A PHARMACEUTICAL COMPOSITION FOR USE IN THE TREATMENT OR PREVENTION OF A C5-RELATED DISEASE AND A METHOD FOR TREATING OR PREVENTING A C5-RELATED DISEASE

Technical Field

[0001]

The present invention relates to dosages and administrations of anti-C5 antibody.

Background Art

[0002]

The complement system plays a central role in the clearance of immune complexes and in immune responses to infectious agents, foreign antigens, virus-infected cells and tumor cells. There are about 25-30 complement proteins, which are found as a complex collection of plasma proteins and membrane cofactors. Complement components achieve their immune defensive functions by interacting in a series of intricate enzymatic cleavages and membrane binding events. The resulting complement cascades lead to the production of products with opsonic, immunoregulatory, and lytic functions.

[0003]

Currently, it is widely accepted that the complement system can be activated through three distinct pathways: the classical pathway, the lectin pathway, and the alternative pathway. These pathways share many components, and while they differ in their initial steps, they converge and share the same terminal complement components (C5 through C9) responsible for the activation and destruction of target cells.

[0004]

The classical pathway is normally activated by the formation of antigen-antibody complexes. Independently, the first step in activation of the lectin pathway is the binding of specific lectins such as mannan-binding lectin (MBL), H-ficolin, M-ficolin, L-ficolin and C-type lectin CL-11. In contrast, the alternative pathway spontaneously undergoes a low level of turnover activation, which can be readily amplified on foreign or other abnormal surfaces (bacteria, yeast, virally infected cells, or damaged tissue). These pathways converge at a point where complement component C3 is cleaved by an active protease to yield C3a and C3b.

[0005]

C3a is an anaphylatoxin. C3b binds to bacterial and other cells, as well as to certain viruses and immune complexes, and tags them for removal from the circulation (the role known as opsonin). C3b also forms a complex with other components to form C5 convertase, which cleaves C5 into C5a and C5b.

[0006]

C5 is a 190 kDa protein found in normal serum at approximately 80 µg/ml (0.4 µM). C5 is glycosylated with about 1.5-3% of its mass attributed to carbohydrate. Mature C5 is a heterodimer of 115 kDa α chain that is disulfide linked to 75 kDa β chain. C5 is synthesized as a single chain precursor protein (pro-C5 precursor) of 1676 amino acids (see, e.g., US Patent No. 6,355,245 and US Patent No. 7,432,356). The pro-C5 precursor is cleaved to yield the β chain as an amino terminal fragment and the α chain as a carboxyl terminal fragment. The alpha chain and the beta chain polypeptide fragments are connected to each other via a disulfide bond and constitute the mature C5 protein.

[0007]

Mature C5 is cleaved into the C5a and C5b fragments during activation of the complement pathways. C5a is cleaved from the α chain of C5 by C5 convertase as an amino terminal fragment comprising the first 74 amino acids of the α chain. The remaining portion of mature C5 is fragment C5b, which contains the rest of the α chain disulfide bonded to the β chain. Approximately 20% of the 11 kDa mass of C5a is attributed to carbohydrate.

[0008]

C5a is another anaphylatoxin. C5b combines with C6, C7, C8 and C9 to form the membrane attack complex (MAC, C5b-9, terminal complement complex (TCC)) at the surface of the target cell. When sufficient numbers of MACs are inserted into target cell membranes, MAC pores are formed to mediate rapid osmotic lysis of the target cells.

[0009]

As mentioned above, C3a and C5a are anaphylatoxins. They can trigger mast cell degranulation, which releases histamine and other mediators of inflammation, resulting in smooth muscle contraction, increased vascular permeability, leukocyte activation, and other inflammatory phenomena including cellular proliferation resulting in hypercellularity. C5a also functions as a chemotactic peptide that serves to attract granulocytes such as neutrophils, eosinophils, basophils and monocytes to the site of complement activation.

[0010]

The activity of C5a is regulated by the plasma enzyme carboxypeptidase N that removes the carboxy-terminal arginine from C5a forming C5a-des-Arg derivative. C5a-des-Arg exhibits only 1% of the anaphylactic activity and polymorphonuclear chemotactic activity of unmodified C5a.

[0011]

While a properly functioning complement system provides a robust defense against infecting microbes, inappropriate regulation or activation of complement has been implicated in the pathogenesis of a variety of disorders including, e.g., rheumatoid arthritis (RA); lupus nephritis; ischemia-reperfusion injury; paroxysmal nocturnal hemoglobinuria (PNH); atypical hemolytic uremic syndrome (aHUS); dense deposit disease (DDD); macular degeneration (e.g., age-related macular degeneration (AMD)); hemolysis, elevated liver enzymes, and low platelets (HELLP) syndrome; thrombotic thrombocytopenic purpura (TTP); spontaneous fetal loss; Pauci-immune vasculitis; epidermolysis bullosa; recurrent fetal loss; multiple sclerosis (MS); traumatic brain injury; and injury resulting from myocardial infarction, cardiopulmonary bypass and hemodialysis (see, e.g., Holers et al., *Immunol. Rev.* 223:300-316 (2008)). Therefore, inhibition of excessive or uncontrolled activations of the complement cascade can provide clinical benefits to patients with such disorders.

[0012]

Paroxysmal nocturnal hemoglobinuria (PNH) is an uncommon blood disorder, wherein red blood cells are compromised and are thus destroyed more rapidly than normal red blood cells. PNH results from the clonal expansion of hematopoietic stem cells with somatic mutations in the PIG-A (phosphatidylinositol glycan class A) gene which is located on the X chromosome. Mutations in PIG-A lead to an early block in the synthesis of glycosylphosphatidylinositol (GPI), a molecule which is required for the anchor of many proteins to cell surfaces. Consequently, PNH blood cells are deficient in GPI-anchored proteins, which include complement-regulatory proteins CD55 and CD59. Under normal circumstances, these complement-regulatory proteins block the formation

of MAC on cell surfaces, thereby preventing erythrocyte lysis. The absence of the GPI-anchored proteins causes complement-mediated hemolysis in PNH.

[0013]

PNH is characterized by hemolytic anemia (a decreased number of red blood cells), hemoglobinuria (the presence of hemoglobin in urine, particularly evident after sleeping), and hemoglobinemia (the presence of hemoglobin in the bloodstream). PNH-afflicted individuals are known to have paroxysms, which are defined here as incidences of dark-colored urine. Hemolytic anemia is due to intravascular destruction of red blood cells by complement components. Other known symptoms include dysphasia, fatigue, erectile dysfunction, thrombosis and recurrent abdominal pain.

[0014]

Eculizumab is a humanized monoclonal antibody directed against the complement protein C5, and the first therapy approved for the treatment of paroxysmal nocturnal hemoglobinuria (PNH) and atypical hemolytic uremic syndrome (aHUS) (see, e.g., Dmytrijuk et al., *The Oncologist* 13(9):993-1000 (2008)). Eculizumab inhibits the cleavage of C5 into C5a and C5b by C5 convertase, which prevents the generation of the terminal complement complex C5b-9. Both C5a and C5b-9 cause the terminal complement-mediated events that are characteristic of PNH and aHUS (see also, WO 2005/074607, WO 2007/106585, WO 2008/069889, and WO 2010/054403).

[0015]

Several reports have described anti-C5 antibodies. For example, WO 95/29697 described an anti-C5 antibody which binds to the α chain of C5 but does not bind to C5a, and blocks the activation of C5, while WO 2002/30985 described an anti-C5 monoclonal antibody which inhibits C5a formation. On the other hand, WO 2004/007553 described

an anti-C5 antibody which recognizes the proteolytic site for C5 convertase on the α chain of C5, and inhibits the conversion of C5 to C5a and C5b. WO 2010/015608 described an anti-C5 antibody which has an affinity constant of at least 1 $x10^7$ M^{-1}.

[0016]

Antibodies (IgGs) bind to neonatal Fc receptor (FcRn), and have long plasma retention times. The binding of IgGs to FcRn is typically observed under acidic conditions (e.g., pH 6.0), and it is rarely observed under neutral conditions (e.g., pH 7.4). Typically, IgGs are nonspecifically incorporated into cells via endocytosis, and return to the cell surfaces by binding to endosomal FcRn under the acidic conditions in the endosomes. Then, IgGs dissociate from FcRn under the neutral conditions in plasma. IgGs that have failed to bind to FcRn are degraded in lysosomes. When the FcRn binding ability of an IgG under acidic conditions is eliminated by introducing mutations into its Fc region, the IgG is not recycled from the endosomes into the plasma, leading to marked impairment of the plasma retention of the IgG. To improve the plasma retention of IgGs, a method that enhances their FcRn binding under acidic conditions has been reported. When the FcRn binding of an IgG under acidic conditions is improved by introducing an amino acid substitution into its Fc region, the IgG is more efficiently recycled from the endosomes to the plasma, and thereby shows improved plasma retention. Meanwhile, it has also been reported that an IgG with enhanced FcRn binding under neutral conditions does not dissociate from FcRn under the neutral conditions in plasma even when it returns to the cell surface via its binding to FcRn under the acidic conditions in the endosomes, and consequently its plasma retention remains unaltered, or rather, is worsened (see, e.g., Yeung et al., *J Immunol.* 182(12): 7663-7671 (2009); Datta-Mannan et al., *J Biol. Chem.* 282(3):1709-1717 (2007); Dall'Acqua et al., *J. Immunol.* 169(9):5171-5180 (2002)).

[0017]

Recently, antibodies that bind to antigens in a pH-dependent manner have been reported (see, e.g., WO 2009/125825 and WO 2011/122011). These antibodies strongly bind to antigens under the plasma neutral conditions and dissociate from the antigens under the endosomal acidic conditions. After dissociating from the antigens, the antibodies become capable once again of binding to antigens when recycled to the plasma via FcRn. Thus, a single antibody molecule can repeatedly bind to multiple antigen molecules. In general, the plasma retention of an antigen is much shorter than that of an antibody that has the above-mentioned FcRn-mediated recycling mechanism. Therefore, when an antigen is bound to an antibody, the antigen normally shows prolonged plasma retention, resulting in an increase of the plasma concentration of the antigen. On the other hand, it has been reported that the above-described antibodies, which bind to antigens in a pH-dependent manner, eliminate antigens from plasma more rapidly than typical antibodies because they dissociate from the antigens within the endosomes during the FcRn-mediated recycling process. WO 2011/111007 also described computer modeling analysis showing that an antibody with pH-dependent binding directed against C5 could extend antigen knockdown.

Citation List

Patent Literature

[0018]

[PTL1] US Patent No. 6,355,245

[PTL2] US Patent No. 7,432,356

[PTL3] WO 2005/074607

[PTL4] WO 2007/106585

[PTL5] WO 2008/069889

[PTL6] WO 2010/054403

[PTL7] WO 95/29697

[PTL8] WO 2002/30985

[PTL9] WO 2004/007553

[PTL10] WO 2010/015608

[PTL11] WO 2009/125825

[PTL12] WO 2011/122011

[PTL13] WO 2011/111007

Non Patent Literature

[0019]

[NPL1] Holers et al., Immunol. Rev. 223:300-316 (2008)

[NPL2] Dmytrijuk et al., The Oncologist 13(9):993-1000 (2008)

[NPL3] Yeung et al., J Immunol. 182(12): 7663-7671 (2009)

[NPL4] Datta-Mannan et al., J Biol. Chem. 282(3):1709-1717 (2007)

[NPL5] Dall'Acqua et al., J. Immunol. 169(9):5171-5180 (2002)

Summary of Invention

[0020]

The invention provides pharmaceutical compositions for use in the treatment or prevention of a C5-related disease and methods for treating or preventing a C5-related disease. The invention also provides dosages and administrations of anti-C5 antibody or pharmaceutical compositions containing the anti-C5 antibody.

[0021]

The inventors of the invention investigated effective dosages and administrations of anti-C5 antibody for use in the treatment and/or prevention of C5-related diseases. As a result, the inventors found that a plasma concentration of anti-C5 antibody rapidly increases to more than a therapeutically efficacious level by intravenously administering the antibody and the concentration is kept at the level by subcutaneously administrating the antibody after the intravenous administration.

[0022]

The inventors also discovered suitable conditions for dosing, including a dose of the antibody, frequency and intervals of the intravenous administrations or subcutaneous administrations, an interval between the intravenous administration and the subcutaneous administration, and switching from other drugs.

[0023]

Specifically, the present invention relates to:

[1] A pharmaceutical composition for use in the treatment or prevention of a C5-related disease, wherein the composition is formulated for subcutaneous injection, comprises an anti-C5 antibody, and is administered subcutaneously, wherein the initial dose of the composition is administered after a dose of another pharmaceutical composition is administered intravenously, wherein the another pharmaceutical composition is formulated for intravenous injection and comprises the anti-C5 antibody.

[2] A pharmaceutical composition for use in the treatment or prevention of a C5-related disease, wherein the composition is formulated for intravenous injection, comprises an anti-C5 antibody, and is administered intravenously, wherein a dose of the composition is administered before the initial dose of another pharmaceutical composition is administered subcutaneously, wherein the another pharmaceutical

composition is formulated for subcutaneous injection and comprises the anti-C5 antibody.

[3] The pharmaceutical composition of [1] or [2], wherein the initial dose of the composition for subcutaneous injection is administered on the same day as or 1 day to 28 days after the dose of the composition for intravenous injection.

[4] The pharmaceutical composition of any one of [1] to [3], wherein the composition for subcutaneous injection is given once every 3 to 42 days.

[5] The pharmaceutical composition of any one of [1] to [4], wherein the composition for the intravenous injection is administered twice or more times and is given once an hour to once every 14 days.

[6] The pharmaceutical composition of any one of [1] to [5], wherein the composition for subcutaneous injection is given at a dose of 17 to 6,000 mg of the antibody.

[7] The pharmaceutical composition of any one of [1] to [6], wherein the composition for intravenous injection is given at a dose of 50 to 5,000 mg of the antibody.

[8] The pharmaceutical composition of any one of [1] to [7], wherein the composition is for use in treatment or prevention of the C5-related disease in a subject who has been treated with at least one pharmacological product for use in treatment or prevention of the disease, wherein the initial dose of the composition for intravenous injection is administered after the final dose of the pharmacological product.

[9] The pharmaceutical composition of [8], wherein the initial dose of the composition for intravenous injection is administered on the third day or after 3 days after the final dose of the pharmacological product.

[10] The pharmaceutical composition of [8] or [9], wherein the pharmacological product comprises an siRNA targeting C5 mRNA, or an anti-C5 antibody which is different from the anti-C5 antibody comprised in the composition.

[11] The pharmaceutical composition of any one of [8] to [10], wherein the pharmacological product comprises eculizumab or its derivative.

[12] A pharmaceutical composition for use in the treatment or prevention of a C5-related disease, wherein the composition is formulated for subcutaneous injection, comprises an anti-C5 antibody, and is administered subcutaneously at a dose of 17 to 6,000 mg of the antibody.

[13] A pharmaceutical composition for use in the treatment or prevention of a C5-related disease, wherein the composition is formulated for intravenous injection, comprises an anti-C5 antibody, and is administered intravenously at a dose of 50 to 5,000 mg of the antibody.

[14] The pharmaceutical composition of any one of [1] to [13], wherein the C5-related disease is any one selected from a group consisting of rheumatoid arthritis (RA); lupus nephritis; ischemia-reperfusion injury; paroxysmal nocturnal hemoglobinuria (PNH); atypical hemolytic uremic syndrome (aHUS); dense deposit disease (DDD); macular degeneration; hemolysis, elevated liver enzymes, and low platelets (HELLP) syndrome; thrombotic thrombocytopenic purpura (TTP); spontaneous fetal loss; Pauci-immune vasculitis; epidermolysis bullosa; recurrent fetal loss; multiple sclerosis (MS); traumatic brain injury; and injury resulting from myocardial infarction, cardiopulmonary bypass or hemodialysis.

[15] A method for treating or preventing a C5-related disease, wherein the method comprises intravenously administrating to a subject a first pharmaceutical

composition which is formulated for intravenous injection and comprises an anti-C5 antibody, and subcutaneously administering to the subject a second pharmaceutical composition which is formulated for subcutaneous injection and comprises the anti-C5 antibody after the administration of the first composition.

[16] The method of [15], wherein the initial dose of the second composition is administered on the same day as or 1 day to 28 days after the final dose of the first composition.

[17] The method of [15] or [16], wherein the second composition is given once every 3 to 42 days.

[18] The method of any one of [15] to [17], wherein the first composition is administered twice or more times and is given once an hour to once every 14 days.

[19] The method of any one of [15] to [18], wherein the second composition is given at a dose of 17 to 6,000 mg of the antibody.

[20] The method of any one of [15] to [19], wherein the first composition is given at a dose of 50 to 5,000 mg of the antibody.

[21] The method of any one of [15] to [20], wherein the method is for treating or preventing the C5-related disease in a subject who has been treated with at least one pharmacological product for use in treatment or prevention of the disease, wherein the initial dose of the first composition is administered after the final dose of the pharmacological product is administered.

[22] The method of [21], wherein the initial dose of the first composition is administered on the third day or after 3 days after the final dose of the pharmacological product is administered.

[23] The method of [21] or [22], wherein the pharmacological product comprises an siRNA targeting C5 mRNA, or an anti-C5 antibody which is different from one comprised in the first composition.

[24] The method of any one of [21] to [23], wherein the pharmacological product comprises eculizumab or its derivative.

[25] A method for treating or preventing C5-related disease, wherein the method comprises administering to a subject a pharmaceutical composition comprising an anti-C5 antibody by subcutaneous injection at a dose of 17 to 6,000 mg of the antibody.

[26] A method for treating or preventing a C5-related disease, wherein the method comprises administering to a subject a pharmaceutical composition comprising an anti-C5 antibody by intravenous injection at a dose of 50 to 5,000 mg of the antibody.

[27] The method of any one of [15] to [26], wherein the C5-related disease is any one selected from the group consisting of rheumatoid arthritis (RA); lupus nephritis; ischemia-reperfusion injury; paroxysmal nocturnal hemoglobinuria (PNH); atypical hemolytic uremic syndrome (aHUS); dense deposit disease (DDD); macular degeneration; hemolysis, elevated liver enzymes, and low platelets (HELLP) syndrome; thrombotic thrombocytopenic purpura (TTP); spontaneous fetal loss; Pauci-immune vasculitis; epidermolysis bullosa; recurrent fetal loss; multiple sclerosis (MS); traumatic brain injury; and injury resulting from myocardial infarction, cardiopulmonary bypass or hemodialysis.

[28] Use of an anti-C5 antibody in the manufacture of a pharmaceutical composition for treating or preventing a C5-related disease, wherein the composition is formulated for subcutaneous injection, comprises the anti-C5 antibody, and is to be administered subcutaneously, wherein an initial dose of the composition is to be

administered after a dose of another pharmaceutical composition is administered intravenously, and wherein the another pharmaceutical composition is formulated for intravenous injection and comprises the anti-C5 antibody.

[29] Use of an anti-C5 antibody in the manufacture of a pharmaceutical composition for treating or preventing a C5-related disease, wherein the composition is formulated for intravenous injection, comprises the anti-C5 antibody, and is to be administered intravenously, wherein a dose of the composition is to be administered before an initial dose of another pharmaceutical composition is administered subcutaneously, wherein the another pharmaceutical composition is formulated for subcutaneous injection and comprises the anti-C5 antibody.

[30] The use of [28] or [29], wherein an initial dose of the composition for subcutaneous injection is to be administered on the same day as or 1 day to 28 days after the dose of the composition for intravenous injection.

[31] The use of any one of [28] to [30], wherein the composition for subcutaneous injection is to be given once every 3 to 42 days.

[32] The use of any one of [28] to [31], wherein the composition for the intravenous injection is to be administered twice or more times and is to be given once an hour to once every 14 days.

[33] The use of any one of [28] to [32], wherein the composition for subcutaneous injection is to be given at a dose of 17 to 6,000 mg of the antibody.

[34] The use of any one of [28] to [33], wherein the composition for intravenous injection is to be given at a dose of 50 to 5,000 mg of the antibody.

[35] The use of any one of [28] to [34], wherein the composition for intravenous injection and the composition for subcutaneous injection are for treating or preventing

the C5-related disease in a subject who has been treated with at least one pharmacological product for use in treatment or prevention of the disease, wherein the initial dose of the composition for intravenous injection is to be administered after the final dose of the pharmacological product.

[36] The use of [35], wherein the initial dose of the composition for intravenous injection is to be administered on the third day or after 3 days after the final dose of the pharmacological product.

[37] The use of [35] or [36], wherein the pharmacological product comprises an siRNA targeting C5 mRNA, or an anti-C5 antibody which is different from the anti-C5 antibody comprised in the composition.

[38] The use of any one of [35] to [37], wherein the pharmacological product comprises eculizumab or its derivative.

[39] Use of an anti-C5 antibody in the manufacture of a first and a second pharmaceutical compositions for treating or preventing a C5-related disease, wherein the first pharmaceutical composition is formulated for intravenous injection, comprises the anti-C5 antibody, and is to be administered intravenously, wherein the second pharmaceutical composition is formulated for subcutaneous injection, comprises the anti-C5 antibody, and is to be administered subcutaneously, and wherein administration of the first and the second pharmaceutical compositions is to be carried out in the order of one or more intravenous doses of the first pharmaceutical composition and one or more subcutaneous doses of the second pharmaceutical composition.

[40] Use of an anti-C5 antibody in the manufacture of a pharmaceutical composition for treating or preventing a C5-related disease, wherein the composition is

formulated for subcutaneous injection, comprises an anti-C5 antibody, and is to be administered subcutaneously at a dose of 17 to 6,000 mg of the antibody.

[41] Use of an anti-C5 antibody in the manufacture of a pharmaceutical composition for treating or preventing a C5-related disease, wherein the composition is formulated for intravenous injection, comprises an anti-C5 antibody, and is to be administered intravenously at a dose of 50 to 5,000 mg of the antibody.

[42] The use of any one of [28] to [41], wherein the C5-related disease is any one selected from the group consisting of rheumatoid arthritis (RA); lupus nephritis; ischemia-reperfusion injury; paroxysmal nocturnal hemoglobinuria (PNH); atypical hemolytic uremic syndrome (aHUS); dense deposit disease (DDD); macular degeneration; hemolysis, elevated liver enzymes, and low platelets (HELLP) syndrome; thrombotic thrombocytopenic purpura (TTP); spontaneous fetal loss; Pauci-immune vasculitis; epidermolysis bullosa; recurrent fetal loss; multiple sclerosis (MS); traumatic brain injury; and injury resulting from myocardial infarction, cardiopulmonary bypass or hemodialysis.

[43] A product for treating or preventing a C5-related disease, comprising (i) a container; (ii) a pharmaceutical composition in the container, wherein the pharmaceutical composition is formulated for subcutaneous injection, comprises an anti-C5 antibody, and is to be administered subcutaneously; and (iii) a document instructing that the pharmaceutical composition is to be administered after a dose of another pharmaceutical composition is administered intravenously, wherein the another pharmaceutical composition is formulated for intravenous injection and comprises the anti-C5 antibody.

[44] A product for treating or preventing a C5-related disease, comprising (i) a container; (ii) a pharmaceutical composition in the container, wherein the pharmaceutical composition is formulated for intravenous injection, comprises an anti-C5 antibody, and is to be administered intravenously; and (iii) a document instructing that the pharmaceutical composition is to be administered before an initial dose of another pharmaceutical composition is administered subcutaneously, wherein the another pharmaceutical composition is formulated for subcutaneous injection and comprises the anti-C5 antibody.

[45] A product for treating or preventing a C5-related disease, comprising (i) a first container; (ii) a first pharmaceutical composition in the first container, wherein the first pharmaceutical composition is formulated for intravenous injection, comprises the anti-C5 antibody, and is to be administered intravenously; (iii) a second container; (iv) a second pharmaceutical composition in the second container, wherein the second pharmaceutical composition is formulated for subcutaneous injection, comprises the anti-C5 antibody, and is to be administered subcutaneously; and (v) a document instructing that the first pharmaceutical composition and the second pharmaceutical composition is to be administered in the order of one or more intravenous doses of the first pharmaceutical composition for intravenous injection and one or more subcutaneous doses of the second pharmaceutical composition for subcutaneous injection.

[46] The product of any one of [43] to [45], wherein the product is for treating or preventing the C5-related disease in a subject who has been treated with at least one pharmacological product for use in treatment or prevention of the disease, wherein the

document further instructs that an initial dose of any one of the pharmaceutical compositions is to be administered after the final dose of the pharmacological product.

[47] A product for treating or preventing a C5-related disease comprising (i) a container; (ii) a pharmaceutical composition in the container, wherein the pharmaceutical composition is formulated for subcutaneous injection and comprises an anti-C5 antibody; and (iii) a document instructing that the pharmaceutical composition is to be administered subcutaneously and at a dose of 17 to 6,000 mg of the antibody.

[48] A product for treating or preventing a C5-related disease comprising (i) a container; (ii) a pharmaceutical composition in the container wherein the pharmaceutical composition is formulated for intravenous injection and comprises an anti-C5 antibody; and (iii) a document instructing that the pharmaceutical composition is to be administered intravenously and at a dose of 50 to 5,000 mg of the antibody.

[49] The product of any one of [43] to [48], wherein the C5-related disease is any one selected from the group consisting of rheumatoid arthritis (RA); lupus nephritis; ischemia-reperfusion injury; paroxysmal nocturnal hemoglobinuria (PNH); atypical hemolytic uremic syndrome (aHUS); dense deposit disease (DDD); macular degeneration; hemolysis, elevated liver enzymes, and low platelets (HELLP) syndrome; thrombotic thrombocytopenic purpura (TTP); spontaneous fetal loss; Pauci-immune vasculitis; epidermolysis bullosa; recurrent fetal loss; multiple sclerosis (MS); traumatic brain injury; and injury resulting from myocardial infarction, cardiopulmonary bypass or hemodialysis.

Brief Description of Drawings

[0024]

[Figure 1] Figure 1 shows the change in plasma 305LO15 concentration in cynomolgus monkeys over time by subcutaneous injection (a) or intravenous injection (b) of antibody 305LO15. The plasma 305LO15 concentration was measured by ELISA. Each point shows a mean value from three males or three females in (a) and a mean value from six males or six females in (b).

[Figure 2] Figure 2 shows the change in plasma 305LO15 concentration in cynomolgus monkeys over time to which monkeys an initial intravenous injection and subsequent subcutaneous maintenance injections of antibody 305LO15 were administered. The plasma 305LO15 concentration was measured by ELISA. Each point shows a mean value from five males or five females.

[Figure 3a] Figure 3a is a graph in which the plasma concentrations of free C5 are plotted against each plasma concentration of the antibody for each indicated dosing amount and route. IV, intravenous injection; SC, subcutaneous injection.

[Figure 3b] Figure 3b is a graph in which the measured complement activities are plotted against each plasma concentration of the antibody for each indicated dosing amount and route. IV, intravenous injection; SC, subcutaneous injection.

[Figure 4a] Figure 4a shows the simulated time-courses of plasma 305LO15 concentration in dosing regimens designed for Part 1 clinical study. IV, intravenous injection; SC, subcutaneous injection.

[Figure 4b] Figure 4b shows the simulated time-course of plasma 305LO15 concentration in dosing regimen designed for Part 2 clinical study. SC, subcutaneous injection; QW, once-every-week administration.

[Figure 4c] Figure 4c shows the simulated time-courses of plasma 305LO15 concentration in dosing regimens designed for Part 3 clinical study. SC, subcutaneous

injection; QW, once-every-week administration; Q2W, once-every-two-week administration; Q4W, once-every-four-week administration.

[Figure 5] Figure 5 shows the profile of immunocomplexes formed with eculizumab (ECZ), human C5 (hC5), and/or 305LO15 analyzed by *in vitro* size-exclusion chromatography at pH7.4 (top) and at pH6.0 (bottom).

[Figure 6] Figure 6 shows the individual observed and simulated plasma 305LO15 concentration-time profiles in healthy subjects who received 75 mg IV infusion (over 60 min). The gray shaded area is the 95% prediction interval of the simulated 305LO15 concentration-time profile. The solid line indicated by an arrow is the simulated median 305LO15 concentration-time profile. The dashed line denotes the 40 μg/mL PD threshold.

[Figure 7] Figure 7 shows the individual observed and simulated plasma 305LO15 concentration-time profiles in healthy subjects who received 125 mg IV infusion (over 60 min). The gray shaded area is the 95% prediction interval of the simulated 305LO15 concentration-time profile. The solid line indicated by an arrow is the simulated median 305LO15 concentration-time profile. The dashed line denotes the 40 μg/mL PD threshold.

[Figure 8] Figure 8 shows the individual observed and simulated plasma 305LO15 concentration-time profiles in healthy subjects who received 100 mg SC. The gray shaded area is a 95% prediction interval of the simulated 305LO15 concentration-time profile. The solid line indicated by an arrow is the simulated median 305LO15 concentration-time profile. For simulations conducted for the single 100 mg SC dosing, a bioavailability of 90% was expected. The dashed line denotes the 40 μg/mL PD threshold.

[Figure 9] Figure 9 shows the relationship between plasma 305LO15 concentrations and hemolytic activity (LIA) following a single IV infusion or SC injection to healthy subjects. Ex vivo LIA dose–response curve was generated using human plasma samples spiked with various concentrations of 305LO15.

[Figure 10a] Figure 10a shows the time profile of serum LDH levels in a PNH patient, patient number 12001, who received an IV dose of 375 mg 305LO15 on day 1, an IV dose of 500 mg 305LO15 on day 8, an IV dose of 1000 mg 305LO15 on day 22, an SC dose of 170 mg 305LO15 on day 36, and an SC dose of 170 mg 305LO15 on day 43 in the Part 2 study.

[Figure 10b] Figure 10b shows the time profile of hemolytic activity (LIA) in the PNH patient, patient number 12001, who received an IV dose of 375 mg 305LO15 on day 1, an IV dose of 500 mg 305LO15 on day 8, an IV dose of 1000 mg 305LO15 on day 22, an SC dose of 170 mg 305LO15 on day 36, and an SC dose of 170 mg 305LO15 on day 43 in the Part 2 study.

Description of Embodiments

[0025]

The techniques and procedures described or referenced herein are generally well understood and commonly employed using conventional methodology by those skilled in the art, such as, for example, the widely utilized methodologies described in Sambrook et al., Molecular Cloning: A Laboratory Manual 3d edition (2001) Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.; Current Protocols in Molecular Biology (F.M. Ausubel, et al. eds., (2003)); the series Methods in Enzymology (Academic Press, Inc.): PCR 2: A Practical Approach (M.J. MacPherson, B.D. Hames and G.R. Taylor eds. (1995)), Harlow and Lane, eds. (1988) Antibodies, A Laboratory Manual, and Animal

Cell Culture (R.I. Freshney, ed. (1987)); Oligonucleotide Synthesis (M.J. Gait, ed., 1984); Methods in Molecular Biology, Humana Press; Cell Biology: A Laboratory Notebook (J.E. Cellis, ed., 1998) Academic Press; Animal Cell Culture (R.I. Freshney), ed., 1987); Introduction to Cell and Tissue Culture (J. P. Mather and P.E. Roberts, 1998) Plenum Press; Cell and Tissue Culture: Laboratory Procedures (A. Doyle, J.B. Griffiths, and D.G. Newell, eds., 1993-8) J. Wiley and Sons; Handbook of Experimental Immunology (D.M. Weir and C.C. Blackwell, eds.); Gene Transfer Vectors for Mammalian Cells (J.M. Miller and M.P. Calos, eds., 1987); PCR: The Polymerase Chain Reaction, (Mullis et al., eds., 1994); Current Protocols in Immunology (J.E. Coligan et al., eds., 1991); Short Protocols in Molecular Biology (Wiley and Sons, 1999); Immunobiology (C.A. Janeway and P. Travers, 1997); Antibodies (P. Finch, 1997); Antibodies: A Practical Approach (D. Catty., ed., IRL Press, 1988-1989); Monoclonal Antibodies: A Practical Approach (P. Shepherd and C. Dean, eds., Oxford University Press, 2000); Using Antibodies: A Laboratory Manual (E. Harlow and D. Lane (Cold Spring Harbor Laboratory Press, 1999); The Antibodies (M. Zanetti and J. D. Capra, eds., Harwood Academic Publishers, 1995); and Cancer: Principles and Practice of Oncology (V.T. DeVita et al., eds., J.B. Lippincott Company, 1993).

[0026]

I. definition

"Administration interval" (an interval between individual administrations) indicates an interval between administration of the nth dose (n is an integer of 1 or greater) and administration of the $(n+1)^{th}$ dose.

[0027]

II. A pharmaceutical composition for subcutaneous injection

A pharmaceutical composition formulated for subcutaneous injection in the present invention is for use in treatment or prevention of a C5-related disease, comprises an anti-C5 antibody, and is administered subcutaneously. The subcutaneous injection can be carried out using ordinary devices and methods to administer a pharmaceutical composition to subcutaneous tissue by injection. Specific devices and methods for the subcutaneous injection also may be selected.

[0028]

Exemplary Anti-C5 Antibodies

The anti-C5 antibody is not limited to a specific embodiment and can be suitably selected from antibodies known as anti-C5 antibody. The term "anti-C5 antibody", or "an antibody that binds to C5" refers to an antibody that is capable of binding C5 with sufficient affinity such that the antibody is useful as a therapeutic agent in targeting C5.

[0029]

In one aspect, the anti-C5 antibodies inhibit activation of C5. In certain embodiments, anti-C5 antibodies prevent the cleavage of C5 to form C5a and C5b, thus preventing the generation of anaphylatoxic activity associated with C5a, as well as preventing the assembly of the C5b-9 membrane attack complex (MAC) associated with C5b. In certain embodiments, anti-C5 antibodies block the conversion of C5 into C5a and C5b by C5 convertase. In certain embodiments, anti-C5 antibodies block access of the C5 convertase to the cleavage site on C5. In certain embodiments, anti-C5 antibodies block hemolytic activity caused by the activation of C5. In further embodiments, anti-C5 antibodies inhibit the activation of C5 via classical pathway and/or alternative pathway.

[0030]

In one aspect, the pharmaceutical compositions of the present invention are useful in treating or preventing a C5-related disease at least in part based on the above-mentioned activities of the anti-C5 antibodies in inhibiting activation of C5.

[0031]

In certain embodiments, C5 activity can be measured as a function of its cell-lysing ability in a subject's body fluids. The cell-lysing ability, or a reduction thereof, of C5 can be measured by methods well known in the art, for example, a conventional hemolytic assay, such as the hemolysis assay described by Kabat and Mayer (eds), Experimental Immunochemistry, 2nd Edition, 135-240, Springfield, IL, CC Thomas (1961), pages 135-139, or a conventional variation of that assay, such as the chicken erythrocyte hemolysis method as described in, e.g., Hillmen et al., *N. Engl. J. Med.* 350(6): 552-559 (2004). In certain embodiments, C5 activity, or inhibition thereof, is quantified using a CH50eq assay. The CH50eq assay is a method for measuring the total classical complement activity in serum. This test is a lytic assay, which uses antibody-sensitized erythrocytes as the activator of the classical complement pathway, and various dilutions of the test serum to determine the amount required to give 50% lysis (CH50). The percentage of hemolysis can be determined, for example, using a spectrophotometer. The CH50eq assay provides an indirect measure of terminal complement complex (TCC) formation, since the TCC themselves are directly responsible for the hemolysis measured. Inhibition of C5 activation can also be detected and/or measured using the methods set forth and exemplified in the working examples. Using assays of these or other suitable types, candidate antibodies capable of inhibiting the activation of C5 can be screened. In certain embodiments, inhibition of C5 activation includes at least a 5%, 10%, 15%, 20%, 25%, 30%, 35%, or 40% or greater decrease in

the C5 activation in an assay as compared to the effect of a negative control under similar conditions. In some embodiments, it refers to inhibition of C5 activation by at least 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95% or greater.

In preferred embodiments, an epitope of the anti-C5 antibody is different from the epitope of eculizumab.

[0032]

In certain embodiments, the anti-C5 antibody for the present invention binds to an epitope within the β chain of C5. In certain embodiments, the anti-C5 antibody binds to an epitope within the MG1-MG2 domain of the β chain of C5. In certain embodiments, the anti-C5 antibody binds to an epitope within a fragment consisting of amino acids 19-180 of the β chain of C5. In certain embodiments, the anti-C5 antibody binds to an epitope within the MG1 domain (amino acids 20-124 of SEQ ID NO: 1) of the β chain of C5. In certain embodiments, the anti-C5 antibody binds to an epitope within a fragment consisting of amino acids 33-124 of the β chain of C5 (SEQ ID NO: 1).

[0033]

In certain embodiments, an anti-C5 antibody for the present invention binds to an epitope within the β chain of C5 which consists of the MG1 domain. In certain embodiments, an anti-C5 antibody binds to an epitope within the β chain (SEQ ID NO: 1) of C5 which comprises at least one fragment selected from the group consisting of amino acids 47-57, 70-76, and 107-110. In certain embodiments, an anti-C5 antibody binds to an epitope within a fragment of the β chain (SEQ ID NO: 1) of C5 which comprises at least one amino acid selected from the group consisting of Thr47, Glu48, Ala49, Phe50, Asp51, Ala52, Thr53, Lys57, His70, Val71, His72, Ser74, Glu76, Val107, Ser108, Lys109, and His110. In certain embodiments, an anti-C5 antibody binds to an epitope within a

fragment of the β chain (SEQ ID NO: 1) of C5 which comprises at least one amino acid selected from the group consisting of Glu48, Asp51, His70, His72, Lys109, and His110. In certain embodiments, binding of an anti-C5 antibody to a C5 mutant is reduced compared to its binding to wild type C5, wherein the C5 mutant has at least one amino acid substitution at a position selected from the group consisting of Glu48, Asp51, His72, and Lys109. In another embodiment, pH-dependent binding (described below) of an anti-C5 antibody to a C5 mutant is reduced compared to its pH-dependent binding to wild type C5, wherein the C5 mutant has at least one amino acid substitution at a position selected from the group consisting of His70, His72, and His110. In a further embodiment, an amino acid at a position selected from Glu48, Asp51, and Lys109 is substituted with alanine, and an amino acid at a position selected from His70, His72, and His110 is substituted with tyrosine in the C5 mutant.

[0034]

In another aspect, the anti-C5 antibodies for the present invention may exhibit pH-dependent binding characteristics. As used herein, the expression "pH-dependent binding" means that the antibody exhibits "reduced binding to C5 at acidic pH as compared to its binding at neutral pH" (for purposes of the present disclosure, both expressions may be used interchangeably). For example, antibodies "with pH-dependent binding characteristics" include antibodies that bind to C5 with higher affinity at neutral pH than at acidic pH. In certain embodiments, the antibodies bind to C5 with at least 2, 3, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 200, 400, 1000, 10000, or more times higher affinity at neutral pH than at acidic pH. In some embodiments, the antibodies bind to C5 with higher affinity at pH7.4 than at pH5.8. In further embodiments, the antibodies bind to C5 with at least 2, 3, 5, 10, 15, 20, 25, 30, 35,

40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 200, 400, 1000, 10000, or more times higher affinity at pH7.4 than at pH5.8.

[0035]

The "affinity" of an antibody for C5, for purposes of the present disclosure, is expressed in terms of the KD of the antibody. The KD of an antibody refers to the equilibrium dissociation constant of an antibody-antigen interaction. The greater the KD value is for an antibody binding to its antigen, the weaker its binding affinity is for that particular antigen. Accordingly, as used herein, the expression "higher affinity at neutral pH than at acidic pH" (or the equivalent expression "pH-dependent binding") means that the KD for the antibody binding to C5 at acidic pH is greater than the KD for the antibody binding to C5 at neutral pH. For example, in the context of the present invention, an antibody is considered to bind to C5 with a higher affinity at neutral pH than at acidic pH if the KD of the antibody binding to C5 at acidic pH is at least 2 times greater than the KD of the antibody binding to C5 at neutral pH. Thus, the anti-C5 antibody for the present invention includes antibodies that bind to C5 at acidic pH with a KD that is at least 2, 3, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 200, 400, 1000, 10000, or more times greater than the KD of the antibody binding to C5 at neutral pH. In another embodiment, the KD value of the antibody at neutral pH can be 10^{-7} M, 10^{-8} M, 10^{-9} M, 10^{-10} M, 10^{-11} M, 10^{-12} M, or less. In another embodiment, the KD value of the antibody at acidic pH can be 10^{-9} M, 10^{-8} M, 10^{-7} M, 10^{-6} M, or greater.

[0036]

In further embodiments an antibody is considered to bind to C5 with a higher affinity at neutral pH than at acidic pH if the KD of the antibody binding to C5 at pH5.8 is at least 2 times greater than the KD of the antibody binding to C5 at pH7.4. In some

embodiments the antibodies bind to C5 at pH5.8 with a KD that is at least 3, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 200, 400, 1000, 10000, or more times greater than the KD of the antibody binding to C5 at pH7.4. In another embodiment, the KD value of the antibody at pH7.4 can be 10^{-7} M, 10^{-8} M, 10^{-9} M, 10^{-10} M, 10^{-11} M, 10^{-12} M, or less. In another embodiment, the KD value of the antibody at pH5.8 can be 10^{-9} M, 10^{-8} M, 10^{-7} M, 10^{-6} M, or greater.

[0037]

The binding properties of an antibody for a particular antigen may also be expressed in terms of the kd of the antibody. The kd of an antibody refers to the dissociation rate constant of the antibody with respect to a particular antigen and is expressed in terms of reciprocal seconds (i.e., sec^{-1}). An increase in kd value signifies weaker binding of an antibody to its antigen. The present invention therefore includes antibodies that bind to C5 with a higher kd value at acidic pH than at neutral pH. The antibodies for the present invention includes antibodies that bind to C5 at acidic pH with a kd that is at least 2, 3, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 200, 400, 1000, 10000, or more times greater than the kd of the antibody binding to C5 at neutral pH. In another embodiment, the kd value of the antibody at neutral pH can be 10^{-2} 1/s, 10^{-3} 1/s, 10^{-4} 1/s, 10^{-5} 1/s, 10^{-6} 1/s, or less. In another embodiment, the kd value of the antibody at acidic pH can be 10^{-3} 1/s, 10^{-2} 1/s, 10^{-1} 1/s, or greater. The antibodies for the present invention also include antibodies that bind to C5 with a higher kd value at pH5.8 than at pH7.4. The antibodies for the present invention include antibodies that bind to C5 at pH5.8 with a kd that is at least 3, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 200, 400, 1000, 10000, or more times greater than the kd of the antibody binding to C5 at pH7.4. In another embodiment, the kd

value of the antibody at pH7.4 can be 10^{-2} 1/s, 10^{-3} 1/s, 10^{-4} 1/s, 10^{-5} 1/s, 10^{-6} 1/s, or less. In another embodiment, the kd value of the antibody at pH5.8 can be 10^{-3} 1/s, 10^{-2} 1/s, 10^{-1} 1/s, or greater.

[0038]

In certain instances, a "reduced binding to C5 at acidic pH as compared to its binding at neutral pH" is expressed in terms of the ratio of the KD value of the antibody binding to C5 at acidic pH to the KD value of the antibody binding to C5 at neutral pH (or vice versa). For example, an antibody may be regarded as exhibiting "reduced binding to C5 at acidic pH as compared to its binding at neutral pH", for purposes of the present invention, if the antibody exhibits an acidic/neutral KD ratio of 2 or greater. In certain exemplary embodiments, the pH5.8/pH7.4 KD ratio for an antibody is 2 or greater. In certain exemplary embodiments, the acidic/neutral KD ratio for an antibody can be 2, 3, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 200, 400, 1000, 10000, or greater. In another embodiment, the KD value of the antibody at neutral pH can be 10^{-7} M, 10^{-8} M, 10^{-9} M, 10^{-10} M, 10^{-11} M, 10^{-12} M, or less. In another embodiment, the KD value of the antibody at acidic pH can be 10^{-9} M, 10^{-8} M, 10^{-7} M, 10^{-6} M, or greater. In further instances an antibody may be regarded as exhibiting "reduced binding to C5 at acidic pH as compared to its binding at neutral pH", for purposes of the present invention, if the antibody exhibits a pH5.8/pH7.4 KD ratio of 2 or greater. In certain exemplary embodiments, the pH5.8/pH7.4 KD ratio for an antibody can be 3, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 200, 400, 1000, 10000, or greater. In another embodiment, the KD value of the antibody at pH7.4 can be 10^{-7} M, 10^{-8} M, 10^{-9} M, 10^{-10} M, 10^{-11} M, 10^{-12} M, or less. In another embodiment, the KD value of the antibody at pH5.8 can be 10^{-9} M, 10^{-8} M, 10^{-7} M, 10^{-6} M, or greater.

[0039]

In certain instances, a "reduced binding to C5 at acidic pH as compared to its binding at neutral pH" is expressed in terms of the ratio of the kd value of the antibody binding to C5 at acidic pH to the kd value of the antibody binding to C5 at neutral pH (or vice versa). For example, an antibody may be regarded as exhibiting "reduced binding to C5 at acidic pH as compared to its binding at neutral pH", for purposes of the present invention, if the antibody exhibits an acidic/neutral kd ratio of 2 or greater. In certain exemplary embodiments, the pH5.8/pH7.4 kd ratio for an antibody is 2 or greater. In certain exemplary embodiments, the acidic/neutral kd ratio for an antibody can be 2, 3, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 200, 400, 1000, 10000, or greater. In further exemplary embodiments, the pH 5.8/pH 7.4 kd ratio for an antibody can be 2, 3, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 200, 400, 1000, 10000, or greater. In another embodiment, the kd value of the antibody at neutral pH can be 10^{-2} 1/s, 10^{-3} 1/s, 10^{-4} 1/s, 10^{-5} 1/s, 10^{-6} 1/s, or less. In a further embodiment, the kd value of the antibody at pH 7.4 can be 10^{-2} 1/s, 10^{-3} 1/s, 10^{-4} 1/s, 10^{-5} 1/s, 10^{-6} 1/s, or less. In another embodiment, the kd value of the antibody at acidic pH can be 10^{-3} 1/s, 10^{-2} 1/s, 10^{-1} 1/s, or greater. In a further embodiment, the kd value of the antibody at pH5.8 can be 10^{-3} 1/s, 10^{-2} 1/s, 10^{-1} 1/s, or greater.

[0040]

As used herein, the expression "acidic pH" means a pH of 4.0 to 6.5. The expression "acidic pH" includes pH values of any one of 4.0, 4.1, 4.2, 4.3, 4.4, 4.5, 4.6, 4.7, 4.8, 4.9, 5.0, 5.1, 5.2, 5.3, 5.4, 5.5, 5.6, 5.7, 5.8, 5.9, 6.0, 6.1, 6.2, 6.3, 6.4, and 6.5. In particular aspects, the "acidic pH" is 5.8.

[0041]

As used herein, the expression "neutral pH" means a pH of 6.7 to about 10.0. The expression "neutral pH" includes pH values of any one of 6.7, 6.8, 6.9, 7.0, 7.1, 7.2, 7.3, 7.4, 7.5, 7.6, 7.7, 7.8, 7.9, 8.0, 8.1, 8.2, 8.3, 8.4, 8.5, 8.6, 8.7, 8.8, 8.9, 9.0, 9.1, 9.2, 9.3, 9.4, 9.5, 9.6, 9.7, 9.8, 9.9, and 10.0. In particular aspects, the "neutral pH" is 7.4.

[0042]

KD values, and kd values, as expressed herein, may be determined using a surface plasmon resonance-based biosensor to characterize antibody-antigen interactions. KD values and kd values can be determined at 25°C or 37°C.

[0043]

In another aspect, an anti-C5 antibody comprises a heavy chain variable domain (VH) sequence having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of SEQ ID NO: 2. In certain embodiments, the VH sequence is the amino acid sequence of SEQ ID NO: 2. In certain embodiments, a VH sequence having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identity contains substitutions (e.g., conservative substitutions), insertions, or deletions relative to the reference sequence, but an anti-C5 antibody comprising that sequence retains the ability to bind to C5. In certain embodiments, a total of 1 to 10 amino acids have been substituted, inserted and/or deleted in SEQ ID NO: 2. In certain embodiments, substitutions, insertions, or deletions occur in regions outside the HVRs (i.e., in the FRs). Optionally, the anti-C5 antibody comprises the VH sequence in SEQ ID NO: 2, including post-translational modifications of that sequence. In a particular embodiment, the VH comprises one, two or three HVRs selected from: (a) a HVR-H1 comprising the amino acid sequence of SEQ ID NO: 3, (b) a HVR-H2

comprising the amino acid sequence of SEQ ID NO: 4, and (c) a HVR-H3 comprising the amino acid sequence of SEQ ID NO: 5.

[0044]

As used herein, the term “HVR” stands for a “hypervariable region” and the term “FR” stands for a “framework”.

[0045]

"Framework" or "FR" refers to variable domain residues other than hypervariable region (HVR) residues. The FR of a variable domain generally consists of four FR domains: FR1, FR2, FR3, and FR4. Accordingly, the HVR and FR sequences generally appear in the following sequence in VH (or VL): FR1-H1(L1)-FR2-H2(L2)-FR3-H3(L3)-FR4.

[0046]

The term "hypervariable region" or "HVR" as used herein refers to each of the regions of an antibody variable domain which are hypervariable in sequence ("complementarity determining regions" or "CDRs") and/or form structurally defined loops ("hypervariable loops") and/or contain the antigen-contacting residues ("antigen contacts"). Generally, antibodies comprise six HVRs: three in the VH (H1, H2, H3), and three in the VL (L1, L2, L3). Exemplary HVRs herein include: (a) hypervariable loops occurring at amino acid residues 26-32 (L1), 50-52 (L2), 91-96 (L3), 26-32 (H1), 53-55 (H2), and 96-101 (H3) (Chothia, J. Mol. Biol. 196:901-917 (1987));(b) CDRs occurring at amino acid residues 24-34 (L1), 50-56 (L2), 89-97 (L3), 31-35b (H1), 50-65 (H2), and 95-102 (H3) (Kabat et al., Sequences of Proteins of Immunological Interest, 5th Ed. Public Health Service, NIH, Bethesda, MD (1991));(c) antigen contacts occurring at amino acid residues 27c-36 (L1), 46-55 (L2), 89-96 (L3), 30-35b (H1), 47-58 (H2), and

93-101 (H3) (MacCallum et al. J. Mol. Biol. 262:732-745 (1996)); and(d) combinations of (a), (b), and/or (c), including HVR amino acid residues 46-56 (L2), 47-56 (L2), 48-56 (L2), 49-56 (L2), 26-35 (H1), 26-35b (H1), 49-65 (H2), 93-102 (H3), and 94-102 (H3). The HVR residues and other residues in the variable domain (e.g., FR residues) are numbered herein according to Kabat et al., supra.

[0047]

In another aspect, an anti-C5 antibody comprises a light chain variable domain (VL) having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of SEQ ID NO: 6. In certain embodiments, the VL sequence is the amino acid sequence of SEQ ID NO: 6. In certain embodiments, a VL sequence having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identity contains substitutions (e.g., conservative substitutions), insertions, or deletions relative to the reference sequence, but an anti-C5 antibody comprising that sequence retains the ability to bind to C5. In certain embodiments, a total of 1 to 10 amino acids have been substituted, inserted and/or deleted in SEQ ID NO: 6. In certain embodiments, the substitutions, insertions, or deletions occur in regions outside the HVRs (i.e., in the FRs). Optionally, the anti-C5 antibody comprises the VL sequence in SEQ ID NO: 6, including post-translational modifications of that sequence. In a particular embodiment, the VL comprises one, two or three HVRs selected from (a) a HVR-L1 comprising the amino acid sequence of SEQ ID NO: 7; (b) a HVR-L2 comprising the amino acid sequence of SEQ ID NO: 8; and (c) a HVR-L3 comprising the amino acid sequence of SEQ ID NO: 9.

[0048]

In another embodiment, the antibody is a full length antibody, e.g., an intact IgG1, IgG2, IgG3 or IgG4 antibody, or an antibody engineered to have regions from two or more IgGs selected from IgG1, IgG2, IgG3, and IgG4 within the region(s) homologous among IgG subclasses by recombination. The antibody may comprise any suitable Fc region comprising a human Fc region sequence (e.g., a human IgG1, IgG2, IgG3 or IgG4 Fc region).

[0049]

Fc region variants

In certain embodiments, an anti-C5 antibody for the present invention comprises an Fc region variant in which one or more amino acid modifications have been introduced into a native sequence Fc region of an antibody. The Fc region variant may comprise a human Fc region sequence (e.g., a human IgG1, IgG2, IgG3 or IgG4 Fc region) comprising an amino acid modification (e.g., a substitution) at one or more amino acid positions.

[0050]

In certain embodiments, the invention contemplates an antibody variant that possesses some but not all effector functions, which make it a desirable candidate for applications in which the half-life of the antibody *in vivo* is important yet certain effector functions (such as complement and ADCC) are unnecessary or deleterious. *In vitro* and/or *in vivo* cytotoxicity assays can be conducted to confirm the reduction/depletion of CDC and/or ADCC activities. For example, Fc receptor (FcR) binding assays can be conducted to ensure that the antibody lacks FcγR binding (hence likely lacking ADCC activity), but retains FcRn binding ability. The primary cells for mediating ADCC, NK cells, express FcγRIII only, whereas monocytes express FcγRI, FcγRII and FcγRIII.

FcR expression on hematopoietic cells is summarized in Table 3 on page 464 of Ravetch and Kinet, *Annu. Rev. Immunol.* 9:457-492 (1991). Non-limiting examples of *in vitro* assays to assess ADCC activity of a molecule of interest is described in US Patent No. 5,500,362 (see, e.g., Hellstrom et al., *Proc. Nat'l Acad. Sci. USA* 83:7059-7063 (1986)) and Hellstrom et al., *Proc. Nat'l Acad. Sci. USA* 82:1499-1502 (1985); US Pat. No. 5,821,337 (see Bruggemann et al., *J. Exp. Med.* 166:1351-1361 (1987)). Alternatively, non-radioactive assays methods may be employed (see, for example, ACTI™ non-radioactive cytotoxicity assay for flow cytometry (CellTechnology, Inc. Mountain View, CA); and CytoTox 96® non-radioactive cytotoxicity assay (Promega, Madison, WI)). Useful effector cells for such assays include peripheral blood mononuclear cells (PBMC) and Natural Killer (NK) cells. Alternatively, or additionally, ADCC activity of the molecule of interest may be assessed *in vivo*, e.g., in an animal model such as that disclosed in Clynes et al., *Proc. Nat'l Acad. Sci. USA* 95:652-656 (1998). C1q binding assays may also be carried out to confirm that the antibody is unable to bind C1q and hence lacks CDC activity. See, e.g., C1q and C3c binding ELISA in WO 2006/029879 and WO 2005/100402. To assess complement activation, a CDC assay may be performed (see, for example, Gazzano-Santoro et al., *J. Immunol. Methods* 202:163 (1996); Cragg et al., *Blood* 101:1045-1052 (2003); and Cragg et al., *Blood* 103:2738-2743 (2004)). FcRn binding and *in vivo* clearance/half-life determinations can also be performed using methods known in the art (see, e.g., Petkova et al., *Int'l. Immunol.* 18(12):1759-1769 (2006)).

[0051]

Antibodies with reduced effector function include those with substitution of one or more of Fc region residues 238, 265, 269, 270, 297, 327 and 329 (US Patent No.

6,737,056). Such Fc mutants include Fc mutants with substitutions at two or more of amino acid positions 265, 269, 270, 297 and 327, including the so-called "DANA" Fc mutant with substitution of residues 265 and 297 to alanine (US Patent No. 7,332,581).

[0052]

Certain antibody variants with improved or diminished binding to FcRs are described. (See, e.g., US Patent No. 6,737,056; WO 2004/056312, and Shields et al., *J. Biol. Chem.* 9(2):6591-6604 (2001).)

[0053]

In certain embodiments, an antibody variant comprises an Fc region with one or more amino acid substitutions which improve ADCC, e.g., substitutions at positions 298, 333, and/or 334 of the Fc region (EU numbering of residues).

[0054]

In some embodiments, alterations are made in the Fc region that result in altered (*i.e.*, either improved or diminished) C1q binding and/or Complement Dependent Cytotoxicity (CDC), e.g., as described in US Patent No. 6,194,551, WO 1999/51642, and Idusogie et al., *J. Immunol.* 164:4178-4184 (2000).

[0055]

Antibodies with increased half-lives and improved binding to the neonatal Fc receptor (FcRn), which is responsible for the transfer of maternal IgGs to the fetus (Guyer et al., *J. Immunol.* 117:587 (1976) and Kim et al., *J. Immunol.* 24:249 (1994)), are described in US2005/0014934 (Hinton et al.). Those antibodies comprise an Fc region with one or more substitutions therein which improve binding of the Fc region to FcRn. Such Fc variants include those with substitutions at one or more of Fc region residues: 238, 256, 265, 272, 286, 303, 305, 307, 311, 312, 317, 340, 356, 360, 362, 376, 378, 380,

382, 413, 424 or 434, e.g., substitution of Fc region residue 434 (US Patent No. 7,371,826).

[0056]

See also Duncan, *Nature* 322:738-40 (1988); US Patent No. 5,648,260; US Patent No. 5,624,821; and WO 1994/29351 concerning other examples of Fc region variants.

[0057]

In certain embodiments, an anti-C5 antibody for the present invention comprises a VH as in any of the embodiments provided above and a heavy chain constant region comprising the amino acid sequence of any one of SEQ ID NOs: 10, 11, 12, 13, 14, and 15. In certain embodiments, an anti-C5 antibody comprises a VL as in any of the embodiments provided above and a light chain constant region comprising the amino acid sequence of any one of SEQ ID NOs: 16, 17, and 18.

[0058]

In certain embodiments, an anti-C5-antibody for the present invention is any one of the antibodies described in WO2016/098356.

[0059]

In certain embodiments, in case that the pharmaceutical composition is administered by subcutaneous injection to a subject repeatedly, the composition is administered once every 3 days to 42 days. An adequate administration interval between subcutaneous injections can be determined within a suitable range according to conditions of a subject or a patient. Specific administration interval between subcutaneous injections is 3 days, 4 days, 5 days, 6 days, 7 days, 8 days, 9 days, 10 days, 11 days, 12 days, 13 days, 14 days, 15 days, 16 days, 17 days, 18 days, 19 days, 20 days,

21 days, 22 days, 23days, 24 days, 25 days, 26 days, 27 days, 28 days, 29 days, 30 days, 31 days, 32 days, 33 days, 34 days, 35 days, 36 days, 37 days, 38 days, 39 days, 40 days, 41 days, or 42 days. In preferred embodiments, the administration interval is 4 days to 35 days. In further preferred embodiments, the administration interval is 5 days to 31 days. In the most preferred embodiments, the administration interval is 7, 14, or 28 days. The pharmaceutical composition may be administered once every week (weekly), once every two weeks (two-weekly or biweekly), or once every four weeks (four-weekly or monthly). The longer interval can reduce patients' burden or suffering.

[0060]

In certain embodiments, the pharmaceutical composition is administered subcutaneously at a dose of 17 to 6,000 mg of the antibody. An adequate dose can be determined within a suitable range according to conditions of a subject or a patient. In case that the pharmaceutical composition is administered subcutaneously to a subject repeatedly, the doses do not have to be the same all the time and may be determined arbitrarily as long as the doses are within the suitable range. For example, the dose may be decreased gradually. In preferred embodiments, the range of the dose is 25 to 4,000 mg of the antibody. In further preferred embodiments, the range of the dose is 50 to 2,000 mg of the antibody. In the most preferred embodiments, the range of the dose is 75 to 1,000 mg of the antibody. Specific preferred doses are 100 mg, 170 mg, 340 mg, and 600mg. Further preferred doses are 170 mg, 340 mg, and 600mg. When the pharmaceutical composition is administered weekly (with about 7 days of interval), a preferred dose is 170 mg. When the pharmaceutical composition is administered two-weekly (with about 14 days of interval), a preferred dose is 340 mg. When the pharmaceutical composition is administered four-weekly (with about 28 days of interval), a preferred

dose is 600 mg. The dosing regimen of 600 mg four-weekly would be particularly preferred, as it can reduce patients' burden or suffering.

[0061]

In certain embodiment, any pharmaceutically acceptable carrier available for an ordinary composition for subcutaneous injection can be included in the composition. A kind of carrier used for subcutaneous injection can be suitably selected from carriers generally known in the art.

[0062]

In certain embodiment, a formulation of the pharmaceutical composition can be any one selected from the group consisting of liquid, semisolid, and solid. The solid composition is usually made by lyophilization from a liquid formulation. The lyophilized composition is usually reconstituted using water or saline solution just before subcutaneous injection.

[0063]

When the composition is administered subcutaneously, a concentration of anti-C5 antibody in a liquid formulation of the composition is determined within an ordinary range in the art. In certain embodiment, a volume of the composition for subcutaneous injection is determined within an ordinary range in the art.

[0064]

In certain embodiments, the C5-related disease is a complement-mediated disease or condition which involves excessive or uncontrolled activation of C5. In certain embodiments, the C5-related disease is at least one selected from a group consisting of rheumatoid arthritis (RA); lupus nephritis; ischemia-reperfusion injury; paroxysmal nocturnal hemoglobinuria (PNH); atypical hemolytic uremic syndrome

(aHUS); dense deposit disease (DDD); macular degeneration; hemolysis, elevated liver enzymes, and low platelets (HELLP) syndrome; thrombotic thrombocytopenic purpura (TTP); spontaneous fetal loss; Pauci-immune vasculitis; epidermolysis bullosa; recurrent fetal loss; multiple sclerosis (MS); traumatic brain injury; and injury resulting from myocardial infarction, cardiopulmonary bypass or hemodialysis; refractory generalized myasthenia gravis (gMG); neuromyelitis optica (NMO). In preferred embodiment, the C5-related disease is at least one selected from a group consisting of PNH, aHUS, gMG and NMO.

[0065]

In certain aspects, an initial dose of the composition for subcutaneous injection is administered after a dose of another pharmaceutical composition is administered intravenously. The pharmaceutical composition for intravenous injection is the same as referred to hereinafter.

[0066]

In certain embodiments, the composition for subcutaneous injection is subcutaneously administered on the same day as or one or more days after a dose of the composition for intravenous injection is intravenously administered. One or more doses of the composition for intravenous injection may be administered to a subject or a patient before administering an initial dose of the composition for subcutaneous injection. In a preferred embodiment, the initial dose of the composition for subcutaneous injection is administered after the final dose of the composition for intravenous injection.

[0067]

In certain embodiments, the initial dose of the composition for subcutaneous injection is administered on the same day as or 1 day to 28 days after a dose of the composition for intravenous injection. Specific interval between the dose of the composition for intravenous injection and the initial dose of the composition for subcutaneous injection is 0 day, 1day, 2 days, 3 days, 4 days, 5 days, 6 days, 7 days, 8 days, 9 days, 10 days, 11 days, 12 days, 13 days, 14 days, 15 days, 16 days, 17 days, 18 days, 19 days, 20 days, 21 days, 22 days, 23days, 24 days, 25 days, 26 days, 27 days, or 28 days. In preferred embodiments, the interval is 1 day to 14 days. In further preferred embodiments, the interval is 3 days to 10 days. In the most preferred embodiments, the interval is 5 days to 8 days.

[0068]

III. A pharmaceutical composition for intravenous injection

A pharmaceutical composition formulated for intravenous injection in the present invention is for use in treatment or prevention of a C5-related disease, comprises an anti-C5 antibody, and is administered intravenously. The intravenous injection can be carried out using ordinary devices and methods to administer a pharmaceutical composition to vein by injection. Specific devices and methods for the intravenous injection also may be selected.

[0069]

The anti-C5 antibody used for intravenous injection can be any antibody as described above for pharmaceutical compositions for subcutaneous injection. The pharmaceutical composition for intravenous injection and the pharmaceutical composition for subcutaneous injection may contain the same anti-C5 antibody or may contain different anti-C5 antibody. In preferred embodiments, the anti-C5 antibody

used for intravenous injection is the same antibody as for subcutaneous injection set forth above.

[0070]

In certain embodiments, the pharmaceutical composition is administered intravenously at a dose of 50 to 5,000 mg of the antibody. An adequate dose can be determined within a suitable range according to conditions of a subject or a patient. In case that the pharmaceutical composition is administered intravenously to a subject repeatedly, the doses do not have to be the same all the time and may be determined arbitrarily as long as the doses are within the suitable range. For example, the dose may be decreased gradually. In preferred embodiments, the range of the dose is 55 to 4,000 mg of the antibody. In further preferred embodiments, the range of the dose is 60 to 2,500 mg of the antibody. In the most preferred embodiments, the range of the dose is 70 to 1,500 mg of the antibody. Specific preferred doses are 75 mg, 125 mg, 150 mg, 300 mg, 375 mg, 500 mg and 1,000 mg. Further preferred doses are 375 mg, 500 mg and 1,000 mg, and among these, the most preferred is 1,000 mg.

[0071]

In certain embodiments, the number of times that the pharmaceutical composition is administered by intravenous injection is not particularly limited and may be once or more times. In preferred embodiments, the number of times is once, twice, or three times. In further preferred embodiments, the number of times is once or twice. In the most preferred embodiment, the number of times is once. The lesser number of times can reduce patients' burden or suffering.

[0072]

In certain embodiments, in case that the pharmaceutical composition is administered intravenously to a subject or a patient repeatedly, the composition is administered once an hour to once every 14 days. An adequate administration interval between intravenous injections can be determined within a suitable range according to conditions of a subject or a patient. Specific administration interval between intravenous injections is one hour, 2 hours, 3 hours, 4 hours, 5 hours, 6 hours, 7 hours, 8 hours, 9 hours, 10 hours, 11 hours, 12 hours, 13 hours, 14 hours, 15 hours, 16 hours, 17 hours, 18 hours, 19 hours, 20 hours, 21 hours, 22 hours, 23hours, 24 hours, 25 hours, 26 hours, 27 hours, 28 hours, 29 hours, 30 hours, 31 hours, 32 hours, 33 hours, 34 hours, 35 hours, 36 hours, 37 hours, 38 hours, 39 hours, 40 hours, 41 hours, 42 hours, 43 hours, 44 hours, 45 hours, 46 hours, 47 hours, 48 hours, 3 days, 4 days, 5 days, 6 days, 7 days, 8 days, 9 days, 10 days, 11 days, 12 days, 13 days, or 14 days. In preferred embodiments, the administration interval is 24 hours to 10 days. In further preferred embodiments, the administration interval is 48 hours to 7 days. In the most preferred embodiments, the administration interval is 3 to 5 days.

[0073]

In certain embodiment, any pharmaceutically acceptable carrier available for an ordinary composition for intravenous injection can be included in the composition. A kind of carrier used for intravenous injection can be suitably selected from carriers generally known in the art.

[0074]

In certain embodiments, a formulation of the pharmaceutical composition can be any one selected from the group consisting of liquid, semisolid, and solid. The sold composition is usually made by lyophilization from a liquid formulation. The lyophilized

composition is usually reconstituted using water or saline solution just before intravenous injection.

[0075]

When the composition is administered intravenously, a concentration of anti-C5 antibody in a liquid formulation of the composition is determined within an ordinary range in the art. In certain embodiment, a volume of the composition for intravenous injection is determined within an ordinary range in the art.

[0076]

In certain aspects, a dose of the composition for intravenous injection is administered before an initial dose of another pharmaceutical composition is administered subcutaneously. The pharmaceutical composition for subcutaneous injection is the same one set forth above.

[0077]

In certain embodiments, the composition for intravenous injection is intravenously administered on the same day as or one or more days before the initial dose of the composition for subcutaneous injection is subcutaneously administered. One or more doses of the composition for intravenous injection may be administered to a subject or a patient before administering an initial dose of the composition for subcutaneous injection. In a preferred embodiment, the final dose of the composition for intravenous injection is administered before the initial dose of the composition for subcutaneous injection.

[0078]

In certain embodiments, the dose of the composition for intravenous injection is administered on the same day as or 1 day to 28 days before the initial dose of the

composition for subcutaneous injection. Specific interval between the dose of the composition for intravenous injection and the initial dose of the composition for subcutaneous injection is 0 day, 1day, 2 days, 3 days, 4 days, 5 days, 6 days, 7 days, 8 days, 9 days, 10 days, 11 days, 12 days, 13 days, 14 days, 15 days, 16 days, 17 days, 18 days, 19 days, 20 days, 21 days, 22 days, 23days, 24 days, 25 days, 26 days, 27 days, or 28 days. In preferred embodiments, the interval is 1 day to 14 days. In further preferred embodiments, the interval is 3 days to 10 days. In the most preferred embodiments, the interval is 5 days to 8 days.

[0079]

IV. Switching from other pharmacological product

In another aspects, the above pharmaceutical compositions for subcutaneous or intravenous injection can be useful for treatment or prevention of a C5-related disease in a subject who has been treated with at least one pharmacological product for use in treatment or prevention of the disease once or more times. For example, the pharmaceutical compositions of the present invention may be useful for treating a patient having a C5-related disease who has received prior treatment with at least one pharmacological product for treating or preventing the disease but is expected to better respond to the treatment by the pharmaceutical compositions of the present invention. In such cases, the medication can be switched from the pharmacological product to the pharmaceutical composition of the present invention. In preferred embodiments, an initial dose of the composition for intravenous injection in the present invention is administered after the final dose of the pharmacological product that has been used in the prior treatment.

[0080]

In certain embodiments, the pharmacological product comprises an siRNA targeting C5 mRNA, or an anti-C5 antibody which is different from the anti-C5 antibody comprised in the above composition for subcutaneous and intravenous injection. In preferred embodiments, the pharmacological product comprises the antibody which is different antibody from ones in the above composition for the subcutaneous and intravenous injection. In the most preferred embodiment, the antibody comprised in the pharmacological product that has been used in the prior treatment is eculizumab or its derivative.

[0081]

In certain embodiments, an initial dose of the composition for intravenous injection in the present invention is administered on the third day or 3 or more days after the final dose of the pharmacological product is administered. Specific interval between the final dose of the pharmacological product and the initial dose of the composition for intravenous injection of the present invention is 3 days, 4 days, 5 days, 6 days, 7 days, 8 days, 9 days, 10 days, 11 days, 12 days, 13 days, 14 days, 15 days, 16 days, 17 days, 18 days, 19 days, 20 days, or 21 days. In preferred embodiments, the composition for intravenous injection of the present invention is administered 7 or more days after the final dose of the pharmacological product. In further preferred embodiments, the composition for intravenous injection of the present invention is administered 14 or more days after the final dose of the pharmacological product. In the most preferred embodiments, the composition for intravenous injection of the present invention is administered 21 or more days after the final dose of the pharmacological product.

[0082]

V. A method for treating or preventing

In other aspects, the invention encompasses methods for treating or preventing a C5-related disease.

[0083]

In one embodiment, the method comprises intravenously administrating a first pharmaceutical composition which is formulated for intravenous injection and comprises an anti-C5 antibody, and subcutaneously administering a second pharmaceutical composition which is formulated for subcutaneous injection and comprises the anti-C5 antibody after the administration of the first composition.

[0084]

Conditions and target diseases of a pharmaceutical composition for subcutaneous or intravenous injection used in the method are the same as those mentioned in the above sections “II. A pharmaceutical composition for intravenous injection” and “III. A pharmaceutical composition for subcutaneous injection”.

[0085]

VI. Vaccination

Infectious disease, e.g. meningococcal infections, can occur in a subject to whom anti-C5 antibody was administered. To prevent such infectious diseases, the subject may be immunized by a vaccine known for treating or preventing the diseases before, after or when the above composition for subcutaneous and intravenous injection is administered.

[0086]

VII. Article of manufacture

In another aspect of the invention, an article of manufacture or product containing materials useful for the treatment or prevention of the C5-related diseases

described above is provided. The article of manufacture or product comprises one or more containers and a label or package insert on or associated with the containers. Suitable containers include, for example, bottles, vials, syringes, IV solution bags, SC solution syringes, etc. The containers may be formed from a variety of materials such as glass or plastic. The container holds a composition which is by itself or combined with another composition effective for treating or preventing the condition and may have a sterile access port (for example the container may be an intravenous solution bag or a vial having a stopper pierceable by a hypodermic injection needle). At least one active agent in the composition is an anti-C5 antibody described above. The label, package insert, or such document indicates that the composition is used for treating the condition of choice.

[0087]

In certain embodiments, the composition contained in the container of the article of manufacture or product is formulated for subcutaneous injection. The article of manufacture or product in this embodiment of the invention may further comprise a package insert indicating that the compositions is to be administered after a dose of another pharmaceutical composition is administered intravenously. Another pharmaceutical composition in this embodiment of the invention comprises the anti-C5 antibody and is formulated for intravenous injection.

[0088]

In certain embodiments, the composition contained in the container of the article of manufacture or product is formulated for intravenous injection. The article of manufacture or product in this embodiment of the invention may further comprise a package insert indicating that the composition is to be administered before an initial

dose of another pharmaceutical composition is administered subcutaneously. Another pharmaceutical composition in this embodiment of the invention comprises the anti-C5 antibody and is formulated for subcutaneous injection.

[0089]

Yet in certain other embodiments, the article of manufacture or product may comprise (a) a first container with a first composition contained therein, wherein the composition comprises the anti-C5 antibody and is formulated for intravenous injection, and (b) a second container with a second composition contained therein, wherein the composition comprises the anti-C5 antibody and is formulated for subcutaneous injection. The article of manufacture in this embodiment of the invention may further comprise a package insert indicating that the compositions can be used to treat a particular condition, such as a C5-related disease, and further indicating that the administration of the first pharmaceutical composition and the second pharmaceutical composition is to be carried out in the order of one or more intravenous administrations of the first pharmaceutical composition for intravenous injection and one or more subcutaneous administrations of the second pharmaceutical composition for subcutaneous injection.

[0090]

Moreover, the article of manufacture or product may further comprise an additional container with a composition contained therein, wherein the composition comprises a further cytotoxic or otherwise therapeutic agent. The article of manufacture in this embodiment of the invention may further comprise a package insert indicating that the compositions can be used to treat a particular condition. Alternatively, or additionally, the article of manufacture may further comprise another

additional container comprising a pharmaceutically-acceptable buffer, such as bacteriostatic water for injection (BWFI), phosphate-buffered saline, Ringer's solution and dextrose solution. It may further include other materials desirable from a commercial and user standpoint, including other buffers, diluents, filters, needles, and syringes.

Examples

[0091]

The following are examples of methods and compositions of the invention. It is understood that various other embodiments may be practiced, given the general description provided above.

Example 1

[0092]

Generation of a C5-antibody

The C5-antibody, 305LO15 in WO2016/098356, was prepared by an ordinary method. Briefly, the genes encoding the heavy chain variable domain (VH) of 305LO15 were combined with the genes encoding a modified human IgG1 heavy chain constant domain (CH) variant SG115 (SEQ ID NO: 13). The genes encoding the light chain variable domain (VL) of 305LO15 were combined with the genes encoding a human light chain constant domain (CL) (SK1, SEQ ID NO: 18). Antibodies were expressed in HEK293 cells co-transfected with the combination of heavy and light chain expression vectors, and were purified by protein A.

Example 2

[0093]

Protein pharmaceuticals, e.g. antibody drugs, are typically administered intravenously because proteins are difficult to concentrate. In fact, eculizumab which is the only approved antibody available for treatment at present is administered intravenously. If subcutaneous injection becomes available, it will reduce patient burden, e.g. ambulant treatment and long time spent in hospital for intravenous injection, since self-injection is possible.

[0094]

To examine the possibility of subcutaneous administration of anti-C5 antibody, 305LO15 disclosed in WO2016/098356 was selected because of its high solubility. It was thought that higher solubility provides the possibility that the protein pharmaceutical can be administered in a smaller volume, allowing subcutaneous injection.

[0095]

To compare pharmacokinetics of 305LO15 after subcutaneous injection or intravenous injection, cynomolgus monkeys were separated into two groups, a subcutaneous injection group and an intravenous injection group. 305LO15 was administered subcutaneously to the animals of the first group (three males and three females) once every week over twenty two times in total at 40 mg/kg of the antibody for each dosing. A plasma 305LO15 concentration of those animals was measured using ELISA analysis at the time points shown in Figure 1a. 305LO15 was administered intravenously to the animals of the second group (six males and six females) once every week over five times in total at 40 mg/kg of the antibody for each dosing. A plasma

305LO15 concentration of the animals was measured by ELISA analysis at the time points shown in Figure 1b.

[0096]

As shown in Figure 1a, the plasma 305LO15 concentration of cynomolgus monkeys increased by subcutaneous injections, verifying that 305LO15 can be administered by subcutaneous injection for therapy. The rate of the initial rise in plasma concentration after the initial dose was lower in the subcutaneous administration (Figure 1a) than in the intravenous injection (Figure 1b).

Example 3

[0097]

Next, it was investigated if the relatively slow rate of the rise in plasma concentration after the initial dose of subcutaneous injection can be compensated by an intravenous injection of 305LO15. To five males and five females of cynomolgus monkeys, the initial dose of 305LO15 was administered intravenously once at the dose of 100 mg/kg of the antibody. After a week from the initial intravenous administration, subcutaneous injection of 305LO15 was started. Twenty six times of the subcutaneous injection were given in total, at the pace of once every week at the dose of 40 mg/kg of the antibody for each dosing.

[0098]

As shown in Figure 2, the rate of the rise in plasma concentration of 305LO15 was faster in the case where the initial dose of the antibody was administered as an intravenous injection before the maintenance doses by subcutaneous injection than in the case of no initial dose of the intravenous injection. The plasma concentration of

305LO15 reached and was maintained at the level comparable to that shown in Figure 1a. It was thought that this faster rise in plasma 305LO15 concentration contributes to faster achievement of therapeutic effect by the drug.

Example 4

[0099]

To estimate efficacious plasma concentration of 305LO15, relationship between plasma concentration of the free antigen (C5) or complement activity and plasma concentration of the antibody (305LO15) was determined in cynomolgus monkeys. 305LO15 was administered intravenously (0.8 mg/kg, 4 mg/kg, or 20 mg/kg) or subcutaneously (4 mg/kg). Plasma concentration of the antibody and free C5 was measured by ELISA. Complement activity was measured by RBC hemolytic assay. As shown in Figure 3, free C5 concentration was suppressed to less than sub μg/mL level when plasma concentration of 305LO15 was maintained at above 40 μg/mL of 305LO15 (Figure 3a) and 40 μg/mL or higher plasma concentration of 305LO15 inhibited complement activity (hemolysis) to less than 20% of baseline (i.e. no antibody is applied) (Figure 3b). It was estimated that complement activity is suppressed to less than 20% of baseline (i.e. no antibody is injected) when 40 μg/mL or higher concentration of 305LO15 is maintained in plasma.

Example 5

[0100]

To determine suitable dosages and administrations for 305LO15 to maintain efficacious plasma 305LO15 concentration for clinical trials, the time course of plasma

305LO15 concentration in humans was predicted by simulation. At first, plasma 305LO15concentrations were measured over time after the antibody was administered to cynomolgus monkeys at 0.8, 4, 20 mg/kg of the antibody for each dosing. Cynomolgus monkey PK parameters were estimated by analyzing the data using a 2-compartment model. Using allometric scaling, human PK parameters were estimated based on the cynomolgus monkey PK parameters. The estimated human PK parameters are shown in Table 1 (abbreviation of parameters in Table 1: BW, body weight; CL, total clearance of drug; F, bioavailability, or a fraction of an administered dose of a drug that becomes systemically available; Ka, absorption rate constant; mAb, monoclonal antibody; PK, pharmacokinetic; Q, intercompartmental clearance; SC, subcutaneous; Vc, volume of distribution for the central compartment; Vp, volume of distribution for the peripheral compartment).

(Table 1)

Parameter	Unit	Cynomolgus Monkey	Human (Allometrically Scaled)	Comment
BW	kg	3.5	70	
F	fraction	0.91	0.75	Not scaled; typical for a high SC bioavailability of a mAb in human*
K_a	per h	0.0485	0.0125	Not scaled; typical value for mAbs in human*
V_c	mL/kg	46.6	46.6	Scaled to body weight with an exponent of 1
V_p	mL/kg	26.1	26.1	Scaled to body weight with an exponent of 1
Cl	mL/h/kg	0.116	0.074	Scaled to body weight with an exponent of 0.85
Q	mL/h/kg	1.16	0.741	Scaled to body weight with an exponent of 0.85

* Dirks NL, Meibohm B. Population pharmacokinetics of therapeutic monoclonal antibodies. Clin Pharmacokinet 2010;49:633–59.

[0101]

Human PK profile was estimated for each cohort expected for phase 1/2 study (Figure 4). Part 1 of the study was designed to include three groups of subjects. The first group is a group of subjects to whom 305LO15 is administered intravenously once at the dose of 75 mg/body. The second group is a group of subjects to whom 305LO15 is administered intravenously once at the dose of 150 mg/body. The third group is a group of subjects to whom 305LO15 is administered subcutaneously once at the dose of 170 mg/body. It was predicted that plasma 305LO15 concentration in subjects who received any one of the above dose is not maintained above the threshold (40 μg/mL) for more than one week (Figure 4a).

[0102]

Part 2 of the study was designed to include a group of subjects to whom 305LO15 is administered intravenously three times (initially at a dose of 300 mg/body, then at 500 mg/body a week after the initial administration, and finally at 1000 mg/body two weeks after the second administration) and, starting from two weeks after the final intravenous administration, 305LO15 is administered subcutaneously once a week at the dose of 170 mg/body. It was predicted that 305LO15 concentration is maintained higher than the PD threshold (40 μg/mL) by the dosing regimen for the Part 2 study (Figure 4b).

[0103]

Part 3 of the study was designed to include three groups of subjects. 305LO15 is initially administered to the subjects of all groups intravenously once at the dose of 500 mg/body. Starting from the day after the initial administration, 305LO15 is administered subcutaneously to subjects of the first group once every week at the dose of 170 mg/body, to subjects of the second group once every two weeks at the dose of

340 mg/body, and to subjects of the third group once every four weeks at the dose of 600 mg/body. It was predicted that 305LO15 concentration is maintained higher than the PD threshold (40 μg/mL) in all of the three groups in the Part 3 study (Figure 4c).

Example 6

[0104]

Eculizumab has been used for treating patients who have PNH or aHUS. In the case that 305LO15 is expected to have a more desirable effect than eculizumab to a PNH or aHUS patient, the drug therapy to the patient may be switched from eculizumab to 305LO15. Meanwhile, there is a possibility that 305LO15 forms immunocomplex with C5 and eculizumab inside the body, because the epitope of 305LO15 is different from that of eculizumab, that is, 305LO15 binds to the β chain of C5, while eculizumab binds to the α chain of C5. Formation of such immunocomplex may cause abnormal activation, and is therefore desirable to be avoided as much as possible.

[0105]

To investigate if 305LO15 forms immunocomplex with eculizumab (ECZ, heavy chain sequence is shown in SEQ ID NO: 19 and light chain sequence is shown in SEQ ID NO: 20) and human C5 (hC5, SEQ ID NO: 21), in vitro analysis was carried out using size-exclusion chromatography (SEC). hC5 was prepared according to the method disclosed in WO2016/098356. ECZ, hC5, and 305LO15 were dialyzed against DPBS pH7.4 for buffer exchange. Then, ECZ and hC5 were mixed and incubated at 37 degrees C for 3-4 hours. After the incubation, 305LO15 was added to this mixture containing ECZ and hC5. Final concentration of ECZ and hC5 was adjusted to 200 μg/mL, and 305LO15 concentration was adjusted to 0, 25, 125, 250 or 1250 μg/mL. After incubating

these mixtures at 37 degrees C for 3-4 hours, SEC analysis was carried out at pH7.4 and pH6.0. Specific SEC conditions were as identified below.

HPLC system: Waters Alliance e2695 HPLC system

Column: TSKgel G4000SWXL

Column temp.: 25 degrees C

Eluent: 50 mM Na-PB/300 mM NaCl, pH7.4 or pH6.0

Flow rate: 0.5 mL/min

Detection: 220 nm UV absorption

Injection: 10 μL

[0106]

As a result, 305LO15 at the concentrations from 125 to 1250 μg/mL did not affect the SEC profile at pH6.0 but affected the profile at pH7.4. More specifically, when 305LO15 was mixed with ECZ and hC5, peaks of the large ECZ/hC5/305LO15 immunocomplexes which were not detected at pH6.0 were detected at pH7.4 (see the peaks indicated by arrows in Figure 5, top), in addition to peaks of the small complexes (see peaks, '2Ag+Ab' and 'Ag+Ab', in Figure 5, top). Only the peaks of the small complexes were detected at pH6.0 (see the peaks of Figure 5, bottom, corresponding to '2Ag+Ab' and 'Ag+Ab'of Figure 5, top). This means that 305LO15 binds to ECZ/hC5 complex at pH7.4, but not at pH6.0, in accordance with the binding characteristics of 305LO15 (binding to hC5 at pH7.4 but not at pH6.0). From these results, it was suggested that simultaneous administration of 305LO15 and ECZ to a subject could result in the formation of large immunocomplexes in plasma.

Example 7

[0107]

Study BP39144, an adaptive phase I/II study, assessed safety, efficacy, pharmacokinetics (PK), and pharmacodynamics (PD) of 305LO15 in healthy volunteers and in patients with paroxysmal nocturnal hemoglobinuria (PNH).

[0108]

Part 1 of Study BP39144 (a randomized, investigator/subject blinded, adaptive, placebo-controlled, parallel group study) evaluated the safety and tolerability of single doses of 305LO15 in healthy volunteers. At each dose-level/cohort, a total of 5 healthy volunteers were randomized to receive a single intravenous (IV; Cohorts 1 and 2) or subcutaneous (SC; Cohort 3) administration of 305LO15 or placebo. The study had an adaptive design, with an ongoing assessment of available safety, tolerability, PK, and PD data prior to the initiation of the next administration. Planned dose levels for cohorts 1, 2 and 3 were 75 mg IV, 150 mg IV, and 170 mg SC, respectively (Figure 4a). Based on preliminary review of the PK and PD data from Cohort 1, doses for Cohorts 2 and 3 were reduced to 125 mg IV (Cohort 2) and 100 mg SC (Cohort 3).

[0109]

For IV infusion, blood samples were collected pre-dose, at the end of IV infusion (1 hour), 2, 6, 12, 24, 48, 72, 96, and 144 hours post-dose, and on days 14, 21, 28, 35, 42, 56, 84, and 91. For SC administration, blood samples were collected pre-dose, at 12, 24, 48, 72, 96 and 144 hours post-dose, and on days 14, 21, 28, 35, 42, 56, 84, and 91.

[0110]

Preliminary safety results from Part 1 of Study BP39144 indicate that 305LO15 was well tolerated at all dose levels evaluated (75 and 125 mg IV and 100 mg SC).

Available clinical safety information raised no significant safety concerns, and showed no evidence suggesting a pattern of adverse events or laboratory test abnormalities. There were no noteworthy findings or trends in vital signs or 12-lead electrocardiograms in any of the treatment groups.

[0111]

Preliminary PK results in serum and simulated concentration-time profiles for 305LO15 are presented in Figure 6, Figure 7, and Figure 8. Based on these data, 305LO15 PK in healthy subjects was in line with the initial predictions of human PK based on cynomolgus monkey PK data (Example 5). Following IV doses, exposure appeared to be dose proportional from 75 to 125 mg. The terminal half-life ($t_{1/2}$) for a typical patient of body weight 70 kg was estimated to be around 25 days. Following SC administrations, preliminary PK results showed that 305LO15 exposure peaked around day 7 and the bioavailability was around 90%. After absorption, the $t_{1/2}$ was comparable with the $t_{1/2}$ of the IV infusion.

[0112]

Preliminary assessment of the exposure-response relationship supported the initial prediction, *i.e.*, approximately 40 μg 305LO15 per mL blood is required to achieve complete complement inhibition (Figure 9).

Example 8

[0113]

Part 2 of Study BP39144 (an open-label, multiple-dose, global multicenter, intra-individual dose-escalation study) evaluated the safety and tolerability for a total duration of 5 months, and the pharmacodynamic effect of multiple doses of 305LO15

on complement activity in treatment naïve PNH patients. In this study six PNH patients were enrolled. The enrolled patients had not been treated with any complement inhibitor or had previously been treated but stopped treatment due to lack of efficacy based on a single missense C5 heterozygous mutation, and showed increased serum LDH levels (>1.5 x ULN) at screening (ULN: upper limit of normal).

[0114]

One (patient number 12001) among six patients was a naïve 47 year-old male Caucasian PNH patient and a candidate for treatment with complement inhibitors. Patient 12001 received three single-ascending IV doses; a single infusion of 375 mg 305LO15 on day 1, followed by a single infusion of 500 mg 305LO15 on day 8, and further followed by a single infusion of 1000 mg 305LO15 on day 22. The first SC administration (170 mg) was initiated on day 36, followed by weekly (QW) SC injections (170 mg) of 305LO15 which will be continued for total treatment duration of 5 months.

[0115]

Preliminary pharmacodynamic results from Patient 12001 are shown in Table 2 and Figure 10. LDH levels, as a pharmacodynamic marker of hemolysis, dropped to levels within the limits of the normal range by day 15, dropped further by day 36, and maintained at the normalized level on day 43 (Table 2 and Figure 10a). As shown in Table 2 and Figure 10b, results of liposome immunoassay (LIA) showed that complement activity was completely inhibited at the end of the infusion on the study-starting day, day 1. A dose of 375 mg 305LO15 on day 1 maintained complete complement inhibition until the next dose of 500 mg 305LO15 on day 8. The dose of 500 mg 305LO15 on day 8 maintained complete complement inhibition until the next

dose of 1000 mg 305LO15 on day 22. The dose of 1000 mg 305LO15 on day 22 maintained complete complement inhibition until the next SC dose of 170 mg 305LO15 on day 36. The SC dose on day 36 maintained complete complement inhibition on day 43.

[0116]

305LO15 was well tolerated with only mild abdominal pain unrelated to the treatment.

(Table 2)

	LDH [U/mL]	LIA [%]
Day 1 pre-dose	2216	80
Day 1 post-dose	Not collected	Complete inhibition
Day 2	2217	Complete inhibition
Day 5	762	Complete inhibition
Day 8 pre-dose	824	Complete inhibition
Day 8 post-dose	634	Complete inhibition
Day 9	Hemolyzed	Complete inhibition
Day 15	474	Complete inhibition
Day 22	455	Complete inhibition
Day 36	414	Complete inhibition
Day 43	461	Complete inhibition

Claims

1. A pharmaceutical composition for use in the treatment or prevention of a C5-related disease, wherein the composition is formulated for subcutaneous injection, comprises an anti-C5 antibody, and is administered subcutaneously, wherein the initial dose of the composition is administered after a dose of another pharmaceutical composition is administered intravenously, wherein the another pharmaceutical composition is formulated for intravenous injection and comprises the anti-C5 antibody.

2. A pharmaceutical composition for use in the treatment or prevention of a C5-related disease, wherein the composition is formulated for intravenous injection, comprises an anti-C5 antibody, and is administered intravenously, wherein a dose of the composition is administered before the initial dose of another pharmaceutical composition is administered subcutaneously, wherein the another pharmaceutical composition is formulated for subcutaneous injection and comprises the anti-C5 antibody.

3. The pharmaceutical composition of claim 1 or 2, wherein the initial dose of the composition for subcutaneous injection is administered on the same day as or 1 day to 28 days after the dose of the composition for intravenous injection.

4. The pharmaceutical composition of any one of claims 1 to 3, wherein the composition for subcutaneous injection is given once every 3 to 42 days.

5. The pharmaceutical composition of any one of claims 1 to 4, wherein the composition for the intravenous injection is administered twice or more times and is given once an hour to once every 14 days.

6. The pharmaceutical composition of any one of claims 1 to 5, wherein the composition for subcutaneous injection is given at a dose of 17 to 6,000 mg of the antibody.

7. The pharmaceutical composition of any one of claims 1 to 6, wherein the composition for intravenous injection is given at a dose of 50 to 5,000 mg of the antibody.

8. The pharmaceutical composition of any one of claims 1 to 7, wherein the composition is for use in treatment or prevention of the C5-related disease in a subject who has been treated with at least one pharmacological product for use in treatment or prevention of the disease, wherein the initial dose of the composition for intravenous injection is administered after the final dose of the pharmacological product.

9. The pharmaceutical composition of claim 8, wherein the initial dose of the composition for intravenous injection is administered on the third day or after 3 days after the final dose of the pharmacological product.

10. The pharmaceutical composition of claim 8 or 9, wherein the pharmacological product comprises an siRNA targeting C5 mRNA, or an anti-C5 antibody which is different from the anti-C5 antibody comprised in the composition.

11. The pharmaceutical composition of any one of claims 8 to 10, wherein the pharmacological product comprises eculizumab or its derivative.

12. A pharmaceutical composition for use in the treatment or prevention of a C5-related disease, wherein the composition is formulated for subcutaneous injection, comprises an anti-C5 antibody, and is administered subcutaneously at a dose of 17 to 6,000 mg of the antibody.

13. A pharmaceutical composition for use in the treatment or prevention of a C5-related disease, wherein the composition is formulated for intravenous injection, comprises an anti-C5 antibody, and is administered intravenously at a dose of 50 to 5,000 mg of the antibody.

14. The pharmaceutical composition of any one of claims 1 to 13, wherein the C5-related disease is any one selected from a group consisting of rheumatoid arthritis (RA); lupus nephritis; ischemia-reperfusion injury; paroxysmal nocturnal hemoglobinuria (PNH); atypical hemolytic uremic syndrome (aHUS); dense deposit disease (DDD); macular degeneration; hemolysis, elevated liver enzymes, and low platelets (HELLP) syndrome; thrombotic thrombocytopenic purpura (TTP); spontaneous fetal loss; Pauci-immune vasculitis; epidermolysis bullosa; recurrent fetal loss; multiple sclerosis (MS); traumatic brain injury; and injury resulting from myocardial infarction, cardiopulmonary bypass or hemodialysis.

SEQUENCE LISTING

<110> CHUGAI SEIYAKU KABUSHIKI KAISHA
F. HOFFMANN-LA ROCHE AG

<120> A pharmaceutical composition for use in the treatment or prevention of a C5-related disease and a method for treating or preventing a C5-related disease

<130> C1-A1713/SG

<160> 21

<170> PatentIn version 3.5

<210> 1
<211> 673
<212> PRT
<213> Homo sapiens

<400> 1

```
Met Gly Leu Leu Gly Ile Leu Cys Phe Leu Ile Phe Leu Gly Lys Thr
1               5                  10                  15

Trp Gly Gln Glu Gln Thr Tyr Val Ile Ser Ala Pro Lys Ile Phe Arg
            20                  25                  30

Val Gly Ala Ser Glu Asn Ile Val Ile Gln Val Tyr Gly Tyr Thr Glu
        35                  40                  45

Ala Phe Asp Ala Thr Ile Ser Ile Lys Ser Tyr Pro Asp Lys Lys Phe
    50                  55                  60

Ser Tyr Ser Ser Gly His Val His Leu Ser Ser Glu Asn Lys Phe Gln
65                  70                  75                  80

Asn Ser Ala Ile Leu Thr Ile Gln Pro Lys Gln Leu Pro Gly Gly Gln
                85                  90                  95

Asn Pro Val Ser Tyr Val Tyr Leu Glu Val Val Ser Lys His Phe Ser
            100                 105                 110

Lys Ser Lys Arg Met Pro Ile Thr Tyr Asp Asn Gly Phe Leu Phe Ile
        115                 120                 125

His Thr Asp Lys Pro Val Tyr Thr Pro Asp Gln Ser Val Lys Val Arg
    130                 135                 140

Val Tyr Ser Leu Asn Asp Asp Leu Lys Pro Ala Lys Arg Glu Thr Val
145                 150                 155                 160

Leu Thr Phe Ile Asp Pro Glu Gly Ser Glu Val Asp Met Val Glu Glu
                165                 170                 175

Ile Asp His Ile Gly Ile Ile Ser Phe Pro Asp Phe Lys Ile Pro Ser
            180                 185                 190
```

```
Asn Pro Arg Tyr Gly Met Trp Thr Ile Lys Ala Lys Tyr Lys Glu Asp
        195                 200                 205

Phe Ser Thr Thr Gly Thr Ala Tyr Phe Glu Val Lys Glu Tyr Val Leu
    210                 215                 220

Pro His Phe Ser Val Ser Ile Glu Pro Glu Tyr Asn Phe Ile Gly Tyr
225                 230                 235                 240

Lys Asn Phe Lys Asn Phe Glu Ile Thr Ile Lys Ala Arg Tyr Phe Tyr
                245                 250                 255

Asn Lys Val Val Thr Glu Ala Asp Val Tyr Ile Thr Phe Gly Ile Arg
            260                 265                 270

Glu Asp Leu Lys Asp Asp Gln Lys Glu Met Met Gln Thr Ala Met Gln
        275                 280                 285

Asn Thr Met Leu Ile Asn Gly Ile Ala Gln Val Thr Phe Asp Ser Glu
    290                 295                 300

Thr Ala Val Lys Glu Leu Ser Tyr Tyr Ser Leu Glu Asp Leu Asn Asn
305                 310                 315                 320

Lys Tyr Leu Tyr Ile Ala Val Thr Val Ile Glu Ser Thr Gly Gly Phe
                325                 330                 335

Ser Glu Glu Ala Glu Ile Pro Gly Ile Lys Tyr Val Leu Ser Pro Tyr
            340                 345                 350

Lys Leu Asn Leu Val Ala Thr Pro Leu Phe Leu Lys Pro Gly Ile Pro
        355                 360                 365

Tyr Pro Ile Lys Val Gln Val Lys Asp Ser Leu Asp Gln Leu Val Gly
    370                 375                 380

Gly Val Pro Val Thr Leu Asn Ala Gln Thr Ile Asp Val Asn Gln Glu
385                 390                 395                 400

Thr Ser Asp Leu Asp Pro Ser Lys Ser Val Thr Arg Val Asp Asp Gly
                405                 410                 415

Val Ala Ser Phe Val Leu Asn Leu Pro Ser Gly Val Thr Val Leu Glu
            420                 425                 430

Phe Asn Val Lys Thr Asp Ala Pro Asp Leu Pro Glu Glu Asn Gln Ala
        435                 440                 445

Arg Glu Gly Tyr Arg Ala Ile Ala Tyr Ser Ser Leu Ser Gln Ser Tyr
    450                 455                 460
```

```
Leu Tyr Ile Asp Trp Thr Asp Asn His Lys Ala Leu Leu Val Gly Glu
465                 470                 475                 480

His Leu Asn Ile Ile Val Thr Pro Lys Ser Pro Tyr Ile Asp Lys Ile
                485                 490                 495

Thr His Tyr Asn Tyr Leu Ile Leu Ser Lys Gly Lys Ile Ile His Phe
            500                 505                 510

Gly Thr Arg Glu Lys Phe Ser Asp Ala Ser Tyr Gln Ser Ile Asn Ile
        515                 520                 525

Pro Val Thr Gln Asn Met Val Pro Ser Ser Arg Leu Leu Val Tyr Tyr
    530                 535                 540

Ile Val Thr Gly Glu Gln Thr Ala Glu Leu Val Ser Asp Ser Val Trp
545                 550                 555                 560

Leu Asn Ile Glu Glu Lys Cys Gly Asn Gln Leu Gln Val His Leu Ser
                565                 570                 575

Pro Asp Ala Asp Ala Tyr Ser Pro Gly Gln Thr Val Ser Leu Asn Met
            580                 585                 590

Ala Thr Gly Met Asp Ser Trp Val Ala Leu Ala Ala Val Asp Ser Ala
        595                 600                 605

Val Tyr Gly Val Gln Arg Gly Ala Lys Lys Pro Leu Glu Arg Val Phe
    610                 615                 620

Gln Phe Leu Glu Lys Ser Asp Leu Gly Cys Gly Ala Gly Gly Gly Leu
625                 630                 635                 640

Asn Asn Ala Asn Val Phe His Leu Ala Gly Leu Thr Phe Leu Thr Asn
                645                 650                 655

Ala Asn Ala Asp Asp Ser Gln Glu Asn Asp Glu Pro Cys Lys Glu Ile
            660                 665                 670

Leu
```

<210> 2
<211> 123
<212> PRT
<213> Artificial Sequence

<220>
<223> An artificially synthesized sequence

<400> 2

Gln Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Arg
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Val His Ser Ser
20 25 30

Tyr Tyr Met Ala Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp
35 40 45

Val Gly Ala Ile Phe Thr Gly Ser Gly Ala Glu Tyr Lys Ala Glu Trp
50 55 60

Ala Lys Gly Arg Val Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
65 70 75 80

Val Leu Thr Met Thr Asn Met Asp Pro Val Asp Thr Ala Thr Tyr Tyr
85 90 95

Cys Ala Ser Asp Ala Gly Tyr Asp Tyr Pro Thr His Ala Met His Tyr
100 105 110

Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
115 120

<210> 3
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> An artificially synthesized sequence

<400> 3

Ser Ser Tyr Tyr Met Ala
1 5

<210> 4
<211> 17
<212> PRT
<213> Artificial Sequence

<220>
<223> An artificially synthesized sequence

<400> 4

Ala Ile Phe Thr Gly Ser Gly Ala Glu Tyr Lys Ala Glu Trp Ala Lys
1 5 10 15

Gly

<210> 5
<211> 13
<212> PRT
<213> Artificial Sequence

<220>
<223> An artificially synthesized sequence

<400> 5

Asp Ala Gly Tyr Asp Tyr Pro Thr His Ala Met His Tyr
1 5 10

<210> 6
<211> 110
<212> PRT
<213> Artificial Sequence

<220>
<223> An artificially synthesized sequence

<400> 6

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1 5 10 15

Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Gly Ile Ser Ser Ser
20 25 30

Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile
35 40 45

Tyr Gly Ala Ser Glu Thr Glu Ser Gly Val Pro Ser Arg Phe Ser Gly
50 55 60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
65 70 75 80

Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Asn Thr Lys Val Gly Ser Ser
85 90 95

Tyr Gly Asn Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
100 105 110

<210> 7
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> An artificially synthesized sequence

<400> 7

Arg Ala Ser Gln Gly Ile Ser Ser Ser Leu Ala
1 5 10

<210> 8
<211> 7
<212> PRT
<213> Artificial Sequence

<220>

<223> An artificially synthesized sequence

<400> 8

Gly Ala Ser Glu Thr Glu Ser
1 5

<210> 9
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> An artificially synthesized sequence

<400> 9

Gln Asn Thr Lys Val Gly Ser Ser Tyr Gly Asn Thr
1 5 10

<210> 10
<211> 325
<212> PRT
<213> Artificial Sequence

<220>
<223> An artificially synthesized sequence

<400> 10

Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Cys Ser Arg
1 5 10 15

Ser Thr Ser Glu Ser Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
20 25 30

Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
35 40 45

Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
50 55 60

Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Lys Thr
65 70 75 80

Tyr Thr Cys Asn Val Asp His Lys Pro Ser Asn Thr Lys Val Asp Lys
85 90 95

Arg Val Glu Ser Lys Tyr Gly Pro Pro Cys Pro Pro Cys Pro Ala Pro
100 105 110

Glu Phe Arg Gly Gly Pro Lys Val Phe Leu Phe Pro Pro Lys Pro Lys
115 120 125

Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val
130 135 140

```
Asp Val Ser Gln Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr Val Asp
145                 150                 155                 160

Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Phe
                165                 170                 175

Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp
            180                 185                 190

Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly Leu
        195                 200                 205

Pro Ser Ser Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg
    210                 215                 220

Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Gln Glu Glu Met Thr Lys
225                 230                 235                 240

Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp
                245                 250                 255

Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys
            260                 265                 270

Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser
        275                 280                 285

Arg Leu Thr Val Asp Lys Ser Arg Trp Gln Glu Gly Asn Val Phe Ser
    290                 295                 300

Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser
305                 310                 315                 320

Leu Ser Leu Ser Leu
                325
```

<210> 11
<211> 325
<212> PRT
<213> Artificial Sequence

<220>
<223> An artificially synthesized sequence

<400> 11

```
Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Cys Ser Arg
1               5                   10                  15

Ser Thr Ser Glu Ser Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
            20                  25                  30

Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
        35                  40                  45
```

```
Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
     50                  55                  60

Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Lys Thr
65                  70                  75                  80

Tyr Thr Cys Asn Val Asp His Lys Pro Ser Asn Thr Lys Val Asp Lys
                85                  90                  95

Arg Val Glu Ser Lys Tyr Gly Pro Pro Cys Pro Pro Cys Pro Ala Pro
            100                 105                 110

Glu Phe Arg Gly Gly Pro Lys Val Phe Leu Phe Pro Pro Lys Pro Lys
        115                 120                 125

Asp Thr Leu Tyr Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val
    130                 135                 140

Asp Val Ser Gln Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr Val Asp
145                 150                 155                 160

Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Phe
                165                 170                 175

Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp
            180                 185                 190

Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly Leu
        195                 200                 205

Pro Ser Ser Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg
    210                 215                 220

Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Gln Glu Glu Met Thr Lys
225                 230                 235                 240

Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp
                245                 250                 255

Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys
            260                 265                 270

Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser
        275                 280                 285

Arg Leu Thr Val Asp Lys Ser Arg Trp Gln Glu Gly Asn Val Phe Ser
    290                 295                 300

Cys Ser Val Met His Glu Ala Leu His Tyr His Val Thr Gln Lys Ser
305                 310                 315                 320
```

Leu Ser Leu Ser Leu
325

<210> 12
<211> 325
<212> PRT
<213> Artificial Sequence

<220>
<223> An artificially synthesized sequence

<400> 12

Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Cys Ser Arg
1 5 10 15

Ser Thr Ser Glu Ser Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
20 25 30

Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
35 40 45

Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
50 55 60

Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Lys Thr
65 70 75 80

Tyr Thr Cys Asn Val Asp His Lys Pro Ser Asn Thr Lys Val Asp Lys
85 90 95

Arg Val Glu Ser Lys Tyr Gly Pro Pro Cys Pro Pro Cys Pro Ala Pro
100 105 110

Glu Phe Arg Gly Gly Pro Lys Val Phe Leu Phe Pro Pro Lys Pro Lys
115 120 125

Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val
130 135 140

Asp Val Ser Gln Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr Val Asp
145 150 155 160

Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Phe
165 170 175

Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp
180 185 190

Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly Leu
195 200 205

Pro Ser Ser Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg
210 215 220

Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Gln Glu Glu Met Thr Lys
225 230 235 240

Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp
245 250 255

Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys
260 265 270

Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser
275 280 285

Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Glu Gly Asn Val Phe Ser
290 295 300

Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser
305 310 315 320

Leu Ser Leu Ser Pro
325

<210> 13
<211> 328
<212> PRT
<213> Artificial Sequence

<220>
<223> An artificially synthesized sequence

<400> 13

Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys
1 5 10 15

Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
20 25 30

Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
35 40 45

Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
50 55 60

Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr
65 70 75 80

Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys
85 90 95

Lys Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys
100 105 110

```
Pro Ala Pro Glu Leu Arg Arg Gly Pro Lys Val Phe Leu Phe Pro Pro
        115                 120                 125

Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys
    130                 135                 140

Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp
145                 150                 155                 160

Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu
                165                 170                 175

Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu
            180                 185                 190

His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn
        195                 200                 205

Lys Gly Leu Pro Ser Ser Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly
    210                 215                 220

Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu
225                 230                 235                 240

Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr
                245                 250                 255

Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn
            260                 265                 270

Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe
        275                 280                 285

Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn
    290                 295                 300

Val Phe Ser Cys Ser Val Leu His Glu Ala Leu His Ala His Tyr Thr
305                 310                 315                 320

Arg Lys Glu Leu Ser Leu Ser Pro
                325
```

<210> 14
<211> 325
<212> PRT
<213> Artificial Sequence

<220>
<223> An artificially synthesized sequence

<400> 14

```
Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Cys Ser Arg
1               5                  10                  15

Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
            20                  25                  30

Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
        35                  40                  45

Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
    50                  55                  60

Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Lys Thr
65                  70                  75                  80

Tyr Thr Cys Asn Val Asp His Lys Pro Ser Asn Thr Lys Val Asp Lys
                85                  90                  95

Arg Val Glu Ser Lys Tyr Gly Pro Pro Cys Pro Pro Cys Pro Ala Pro
            100                 105                 110

Glu Phe Arg Arg Gly Pro Lys Val Phe Leu Phe Pro Pro Lys Pro Lys
        115                 120                 125

Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val
    130                 135                 140

Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp
145                 150                 155                 160

Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr
                165                 170                 175

Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp
            180                 185                 190

Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly Leu
        195                 200                 205

Pro Ser Ser Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg
    210                 215                 220

Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys
225                 230                 235                 240

Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp
                245                 250                 255

Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys
            260                 265                 270
```

Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser
275 280 285

Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser
290 295 300

Cys Ser Val Leu His Glu Ala Leu His Ala His Tyr Thr Arg Lys Glu
305 310 315 320

Leu Ser Leu Ser Pro
325

<210> 15
<211> 325
<212> PRT
<213> Artificial Sequence

<220>
<223> An artificially synthesized sequence

<400> 15

Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Cys Ser Arg
1 5 10 15

Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
20 25 30

Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
35 40 45

Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
50 55 60

Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Lys Thr
65 70 75 80

Tyr Thr Cys Asn Val Asp His Lys Pro Ser Asn Thr Lys Val Asp Lys
85 90 95

Arg Val Glu Ser Lys Tyr Gly Pro Pro Cys Pro Pro Cys Pro Ala Pro
100 105 110

Glu Phe Arg Arg Gly Pro Lys Val Phe Leu Phe Pro Pro Lys Pro Lys
115 120 125

Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val
130 135 140

Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp
145 150 155 160

Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr
165 170 175

```
Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp
            180                 185                 190

Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly Leu
        195                 200                 205

Pro Ser Ser Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg
    210                 215                 220

Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys
225                 230                 235                 240

Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp
                245                 250                 255

Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys
            260                 265                 270

Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser
        275                 280                 285

Arg Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser
    290                 295                 300

Cys Ser Val Leu His Glu Ala Leu His Ala His Tyr Thr Arg Lys Glu
305                 310                 315                 320

Leu Ser Leu Ser Pro
                325
```

<210> 16
<211> 107
<212> PRT
<213> Artificial Sequence

<220>
<223> An artificially synthesized sequence

<400> 16

```
Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu
1               5                   10                  15

Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe
            20                  25                  30

Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln
        35                  40                  45

Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Cys
    50                  55                  60
```

Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu
65 70 75 80

Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser
85 90 95

Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
100 105

<210> 17
<211> 107
<212> PRT
<213> Artificial Sequence

<220>
<223> An artificially synthesized sequence

<400> 17

Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu
1 5 10 15

Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe
20 25 30

Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln
35 40 45

Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser
50 55 60

Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu
65 70 75 80

Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser
85 90 95

Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
100 105

<210> 18
<211> 107
<212> PRT
<213> Artificial Sequence

<220>
<223> An artificially synthesized sequence

<400> 18

Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu
1 5 10 15

Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe
20 25 30

Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln
35 40 45

Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser
50 55 60

Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu
65 70 75 80

Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser
85 90 95

Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
100 105

<210> 19
<211> 448
<212> PRT
<213> Artificial Sequence

<220>
<223> Eculizumab heavy chain

<400> 19

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Ile Phe Ser Asn Tyr
20 25 30

Trp Ile Gln Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
35 40 45

Gly Glu Ile Leu Pro Gly Ser Gly Ser Thr Glu Tyr Thr Glu Asn Phe
50 55 60

Lys Asp Arg Val Thr Met Thr Arg Asp Thr Ser Thr Ser Thr Val Tyr
65 70 75 80

Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Arg Tyr Phe Phe Gly Ser Ser Pro Asn Trp Tyr Phe Asp Val Trp
100 105 110

Gly Gln Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro
115 120 125

Ser Val Phe Pro Leu Ala Pro Cys Ser Arg Ser Thr Ser Glu Ser Thr
130 135 140

Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr
145 150 155 160

```
Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro
                165                 170                 175

Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr
            180                 185                 190

Val Pro Ser Ser Asn Phe Gly Thr Gln Thr Tyr Thr Cys Asn Val Asp
        195                 200                 205

His Lys Pro Ser Asn Thr Lys Val Asp Lys Thr Val Glu Arg Lys Cys
    210                 215                 220

Cys Val Glu Cys Pro Pro Cys Pro Ala Pro Pro Val Ala Gly Pro Ser
225                 230                 235                 240

Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg
                245                 250                 255

Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser Gln Glu Asp Pro
            260                 265                 270

Glu Val Gln Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala
        275                 280                 285

Lys Thr Lys Pro Arg Glu Glu Gln Phe Asn Ser Thr Tyr Arg Val Val
    290                 295                 300

Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr
305                 310                 315                 320

Lys Cys Lys Val Ser Asn Lys Gly Leu Pro Ser Ser Ile Glu Lys Thr
                325                 330                 335

Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu
            340                 345                 350

Pro Pro Ser Gln Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys
        355                 360                 365

Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser
    370                 375                 380

Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp
385                 390                 395                 400

Ser Asp Gly Ser Phe Phe Leu Tyr Ser Arg Leu Thr Val Asp Lys Ser
                405                 410                 415

Arg Trp Gln Glu Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala
            420                 425                 430
```

```
Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Leu Gly Lys
        435                 440                 445


<210>  20
<211>  214
<212>  PRT
<213>  Artificial Sequence

<220>
<223>  Eculizumab light chain

<400>  20

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1               5                   10                  15


Asp Arg Val Thr Ile Thr Cys Gly Ala Ser Glu Asn Ile Tyr Gly Ala
            20                  25                  30


Leu Asn Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile
        35                  40                  45


Tyr Gly Ala Thr Asn Leu Ala Asp Gly Val Pro Ser Arg Phe Ser Gly
    50                  55                  60


Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
65                  70                  75                  80


Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Asn Val Leu Asn Thr Pro Leu
                85                  90                  95


Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala
            100                 105                 110


Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly
        115                 120                 125


Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala
    130                 135                 140


Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln
145                 150                 155                 160


Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser
                165                 170                 175


Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr
            180                 185                 190


Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser
        195                 200                 205
```

Phe Asn Arg Gly Glu Cys
210

<210> 21
<211> 1676
<212> PRT
<213> Homo sapiens

<400> 21

Met Gly Leu Leu Gly Ile Leu Cys Phe Leu Ile Phe Leu Gly Lys Thr
1 5 10 15

Trp Gly Gln Glu Gln Thr Tyr Val Ile Ser Ala Pro Lys Ile Phe Arg
20 25 30

Val Gly Ala Ser Glu Asn Ile Val Ile Gln Val Tyr Gly Tyr Thr Glu
35 40 45

Ala Phe Asp Ala Thr Ile Ser Ile Lys Ser Tyr Pro Asp Lys Lys Phe
50 55 60

Ser Tyr Ser Ser Gly His Val His Leu Ser Ser Glu Asn Lys Phe Gln
65 70 75 80

Asn Ser Ala Ile Leu Thr Ile Gln Pro Lys Gln Leu Pro Gly Gly Gln
85 90 95

Asn Pro Val Ser Tyr Val Tyr Leu Glu Val Val Ser Lys His Phe Ser
100 105 110

Lys Ser Lys Arg Met Pro Ile Thr Tyr Asp Asn Gly Phe Leu Phe Ile
115 120 125

His Thr Asp Lys Pro Val Tyr Thr Pro Asp Gln Ser Val Lys Val Arg
130 135 140

Val Tyr Ser Leu Asn Asp Asp Leu Lys Pro Ala Lys Arg Glu Thr Val
145 150 155 160

Leu Thr Phe Ile Asp Pro Glu Gly Ser Glu Val Asp Met Val Glu Glu
165 170 175

Ile Asp His Ile Gly Ile Ile Ser Phe Pro Asp Phe Lys Ile Pro Ser
180 185 190

Asn Pro Arg Tyr Gly Met Trp Thr Ile Lys Ala Lys Tyr Lys Glu Asp
195 200 205

Phe Ser Thr Thr Gly Thr Ala Tyr Phe Glu Val Lys Glu Tyr Val Leu
210 215 220

Pro His Phe Ser Val Ser Ile Glu Pro Glu Tyr Asn Phe Ile Gly Tyr

225 230 235 240

Lys Asn Phe Lys Asn Phe Glu Ile Thr Ile Lys Ala Arg Tyr Phe Tyr
245 250 255

Asn Lys Val Val Thr Glu Ala Asp Val Tyr Ile Thr Phe Gly Ile Arg
260 265 270

Glu Asp Leu Lys Asp Asp Gln Lys Glu Met Met Gln Thr Ala Met Gln
275 280 285

Asn Thr Met Leu Ile Asn Gly Ile Ala Gln Val Thr Phe Asp Ser Glu
290 295 300

Thr Ala Val Lys Glu Leu Ser Tyr Tyr Ser Leu Glu Asp Leu Asn Asn
305 310 315 320

Lys Tyr Leu Tyr Ile Ala Val Thr Val Ile Glu Ser Thr Gly Gly Phe
325 330 335

Ser Glu Glu Ala Glu Ile Pro Gly Ile Lys Tyr Val Leu Ser Pro Tyr
340 345 350

Lys Leu Asn Leu Val Ala Thr Pro Leu Phe Leu Lys Pro Gly Ile Pro
355 360 365

Tyr Pro Ile Lys Val Gln Val Lys Asp Ser Leu Asp Gln Leu Val Gly
370 375 380

Gly Val Pro Val Thr Leu Asn Ala Gln Thr Ile Asp Val Asn Gln Glu
385 390 395 400

Thr Ser Asp Leu Asp Pro Ser Lys Ser Val Thr Arg Val Asp Asp Gly
405 410 415

Val Ala Ser Phe Val Leu Asn Leu Pro Ser Gly Val Thr Val Leu Glu
420 425 430

Phe Asn Val Lys Thr Asp Ala Pro Asp Leu Pro Glu Glu Asn Gln Ala
435 440 445

Arg Glu Gly Tyr Arg Ala Ile Ala Tyr Ser Ser Leu Ser Gln Ser Tyr
450 455 460

Leu Tyr Ile Asp Trp Thr Asp Asn His Lys Ala Leu Leu Val Gly Glu
465 470 475 480

His Leu Asn Ile Ile Val Thr Pro Lys Ser Pro Tyr Ile Asp Lys Ile
485 490 495

Thr His Tyr Asn Tyr Leu Ile Leu Ser Lys Gly Lys Ile Ile His Phe

500 505 510

Gly Thr Arg Glu Lys Phe Ser Asp Ala Ser Tyr Gln Ser Ile Asn Ile
515 520 525

Pro Val Thr Gln Asn Met Val Pro Ser Ser Arg Leu Leu Val Tyr Tyr
530 535 540

Ile Val Thr Gly Glu Gln Thr Ala Glu Leu Val Ser Asp Ser Val Trp
545 550 555 560

Leu Asn Ile Glu Glu Lys Cys Gly Asn Gln Leu Gln Val His Leu Ser
565 570 575

Pro Asp Ala Asp Ala Tyr Ser Pro Gly Gln Thr Val Ser Leu Asn Met
580 585 590

Ala Thr Gly Met Asp Ser Trp Val Ala Leu Ala Ala Val Asp Ser Ala
595 600 605

Val Tyr Gly Val Gln Arg Gly Ala Lys Lys Pro Leu Glu Arg Val Phe
610 615 620

Gln Phe Leu Glu Lys Ser Asp Leu Gly Cys Gly Ala Gly Gly Gly Leu
625 630 635 640

Asn Asn Ala Asn Val Phe His Leu Ala Gly Leu Thr Phe Leu Thr Asn
645 650 655

Ala Asn Ala Asp Asp Ser Gln Glu Asn Asp Glu Pro Cys Lys Glu Ile
660 665 670

Leu Arg Pro Arg Arg Thr Leu Gln Lys Lys Ile Glu Glu Ile Ala Ala
675 680 685

Lys Tyr Lys His Ser Val Val Lys Lys Cys Cys Tyr Asp Gly Ala Cys
690 695 700

Val Asn Asn Asp Glu Thr Cys Glu Gln Arg Ala Ala Arg Ile Ser Leu
705 710 715 720

Gly Pro Arg Cys Ile Lys Ala Phe Thr Glu Cys Cys Val Val Ala Ser
725 730 735

Gln Leu Arg Ala Asn Ile Ser His Lys Asp Met Gln Leu Gly Arg Leu
740 745 750

His Met Lys Thr Leu Leu Pro Val Ser Lys Pro Glu Ile Arg Ser Tyr
755 760 765

Phe Pro Glu Ser Trp Leu Trp Glu Val His Leu Val Pro Arg Arg Lys

770 775 780

Gln Leu Gln Phe Ala Leu Pro Asp Ser Leu Thr Thr Trp Glu Ile Gln
785 790 795 800

Gly Val Gly Ile Ser Asn Thr Gly Ile Cys Val Ala Asp Thr Val Lys
805 810 815

Ala Lys Val Phe Lys Asp Val Phe Leu Glu Met Asn Ile Pro Tyr Ser
820 825 830

Val Val Arg Gly Glu Gln Ile Gln Leu Lys Gly Thr Val Tyr Asn Tyr
835 840 845

Arg Thr Ser Gly Met Gln Phe Cys Val Lys Met Ser Ala Val Glu Gly
850 855 860

Ile Cys Thr Ser Glu Ser Pro Val Ile Asp His Gln Gly Thr Lys Ser
865 870 875 880

Ser Lys Cys Val Arg Gln Lys Val Glu Gly Ser Ser Ser His Leu Val
885 890 895

Thr Phe Thr Val Leu Pro Leu Glu Ile Gly Leu His Asn Ile Asn Phe
900 905 910

Ser Leu Glu Thr Trp Phe Gly Lys Glu Ile Leu Val Lys Thr Leu Arg
915 920 925

Val Val Pro Glu Gly Val Lys Arg Glu Ser Tyr Ser Gly Val Thr Leu
930 935 940

Asp Pro Arg Gly Ile Tyr Gly Thr Ile Ser Arg Arg Lys Glu Phe Pro
945 950 955 960

Tyr Arg Ile Pro Leu Asp Leu Val Pro Lys Thr Glu Ile Lys Arg Ile
965 970 975

Leu Ser Val Lys Gly Leu Leu Val Gly Glu Ile Leu Ser Ala Val Leu
980 985 990

Ser Gln Glu Gly Ile Asn Ile Leu Thr His Leu Pro Lys Gly Ser Ala
995 1000 1005

Glu Ala Glu Leu Met Ser Val Val Pro Val Phe Tyr Val Phe His
1010 1015 1020

Tyr Leu Glu Thr Gly Asn His Trp Asn Ile Phe His Ser Asp Pro
1025 1030 1035

Leu Ile Glu Lys Gln Lys Leu Lys Lys Lys Leu Lys Glu Gly Met

1040 1045 1050

Leu Ser Ile Met Ser Tyr Arg Asn Ala Asp Tyr Ser Tyr Ser Val
1055 1060 1065

Trp Lys Gly Gly Ser Ala Ser Thr Trp Leu Thr Ala Phe Ala Leu
1070 1075 1080

Arg Val Leu Gly Gln Val Asn Lys Tyr Val Glu Gln Asn Gln Asn
1085 1090 1095

Ser Ile Cys Asn Ser Leu Leu Trp Leu Val Glu Asn Tyr Gln Leu
1100 1105 1110

Asp Asn Gly Ser Phe Lys Glu Asn Ser Gln Tyr Gln Pro Ile Lys
1115 1120 1125

Leu Gln Gly Thr Leu Pro Val Glu Ala Arg Glu Asn Ser Leu Tyr
1130 1135 1140

Leu Thr Ala Phe Thr Val Ile Gly Ile Arg Lys Ala Phe Asp Ile
1145 1150 1155

Cys Pro Leu Val Lys Ile Asp Thr Ala Leu Ile Lys Ala Asp Asn
1160 1165 1170

Phe Leu Leu Glu Asn Thr Leu Pro Ala Gln Ser Thr Phe Thr Leu
1175 1180 1185

Ala Ile Ser Ala Tyr Ala Leu Ser Leu Gly Asp Lys Thr His Pro
1190 1195 1200

Gln Phe Arg Ser Ile Val Ser Ala Leu Lys Arg Glu Ala Leu Val
1205 1210 1215

Lys Gly Asn Pro Pro Ile Tyr Arg Phe Trp Lys Asp Asn Leu Gln
1220 1225 1230

His Lys Asp Ser Ser Val Pro Asn Thr Gly Thr Ala Arg Met Val
1235 1240 1245

Glu Thr Thr Ala Tyr Ala Leu Leu Thr Ser Leu Asn Leu Lys Asp
1250 1255 1260

Ile Asn Tyr Val Asn Pro Val Ile Lys Trp Leu Ser Glu Glu Gln
1265 1270 1275

Arg Tyr Gly Gly Gly Phe Tyr Ser Thr Gln Asp Thr Ile Asn Ala
1280 1285 1290

Ile Glu Gly Leu Thr Glu Tyr Ser Leu Leu Val Lys Gln Leu Arg

1295 1300 1305

Leu Ser Met Asp Ile Asp Val Ser Tyr Lys His Lys Gly Ala Leu
1310 1315 1320

His Asn Tyr Lys Met Thr Asp Lys Asn Phe Leu Gly Arg Pro Val
1325 1330 1335

Glu Val Leu Leu Asn Asp Asp Leu Ile Val Ser Thr Gly Phe Gly
1340 1345 1350

Ser Gly Leu Ala Thr Val His Val Thr Thr Val Val His Lys Thr
1355 1360 1365

Ser Thr Ser Glu Glu Val Cys Ser Phe Tyr Leu Lys Ile Asp Thr
1370 1375 1380

Gln Asp Ile Glu Ala Ser His Tyr Arg Gly Tyr Gly Asn Ser Asp
1385 1390 1395

Tyr Lys Arg Ile Val Ala Cys Ala Ser Tyr Lys Pro Ser Arg Glu
1400 1405 1410

Glu Ser Ser Ser Gly Ser Ser His Ala Val Met Asp Ile Ser Leu
1415 1420 1425

Pro Thr Gly Ile Ser Ala Asn Glu Glu Asp Leu Lys Ala Leu Val
1430 1435 1440

Glu Gly Val Asp Gln Leu Phe Thr Asp Tyr Gln Ile Lys Asp Gly
1445 1450 1455

His Val Ile Leu Gln Leu Asn Ser Ile Pro Ser Ser Asp Phe Leu
1460 1465 1470

Cys Val Arg Phe Arg Ile Phe Glu Leu Phe Glu Val Gly Phe Leu
1475 1480 1485

Ser Pro Ala Thr Phe Thr Val Tyr Glu Tyr His Arg Pro Asp Lys
1490 1495 1500

Gln Cys Thr Met Phe Tyr Ser Thr Ser Asn Ile Lys Ile Gln Lys
1505 1510 1515

Val Cys Glu Gly Ala Ala Cys Lys Cys Val Glu Ala Asp Cys Gly
1520 1525 1530

Gln Met Gln Glu Glu Leu Asp Leu Thr Ile Ser Ala Glu Thr Arg
1535 1540 1545

Lys Gln Thr Ala Cys Lys Pro Glu Ile Ala Tyr Ala Tyr Lys Val

1550 1555 1560

Ser Ile Thr Ser Ile Thr Val Glu Asn Val Phe Val Lys Tyr Lys
1565 1570 1575

Ala Thr Leu Leu Asp Ile Tyr Lys Thr Gly Glu Ala Val Ala Glu
1580 1585 1590

Lys Asp Ser Glu Ile Thr Phe Ile Lys Lys Val Thr Cys Thr Asn
1595 1600 1605

Ala Glu Leu Val Lys Gly Arg Gln Tyr Leu Ile Met Gly Lys Glu
1610 1615 1620

Ala Leu Gln Ile Lys Tyr Asn Phe Ser Phe Arg Tyr Ile Tyr Pro
1625 1630 1635

Leu Asp Ser Leu Thr Trp Ile Glu Tyr Trp Pro Arg Asp Thr Thr
1640 1645 1650

Cys Ser Ser Cys Gln Ala Phe Leu Ala Asn Leu Asp Glu Phe Ala
1655 1660 1665

Glu Asp Ile Phe Leu Asn Gly Cys
1670 1675