

CLAIMS

1. An agonistic antigen-binding molecule comprising:
 - (1) a cancer-specific antigen-binding Fab domain;
 - (2) CD137-binding Fab domain; and
 - (3) an FcRn-binding domain, wherein the FcRn-binding domain is an antibody Fc region having decreased Fc γ receptor-binding activity.
2. The antigen-binding molecule of claim 1, which is an antibody.
3. A pharmaceutical composition comprising as an active ingredient the antigen-binding molecule of claim 1 or 2.
4. The pharmaceutical composition of claim 3, which is a cytotoxicity-inducing composition.
5. A pharmaceutical composition comprising a combination of a first agonistic antigen-binding molecule of claim 1 or 2, and a second antigen-binding molecule that comprises:
 - (1) a cancer-specific antigen-binding domain; and
 - (2) a T cell receptor complex-binding domain.
6. The antigen-binding molecule of claim 1 or claim 2 or the composition of any one of claims 3 to 5 for use in the treatment of cancer.
7. Use of an agonistic antigen-binding molecule comprising:
 - (1) a cancer-specific antigen-binding Fab domain;
 - (2) a CD137-binding Fab domain; and
 - (3) an FcRn-binding domain, wherein the FcRn-binding domain is an antibody Fc region having decreased Fc γ receptor-binding activity,in the manufacture of a medicament for the treatment of cancer.
8. Use of a composition comprising a combination of:
 - a first agonistic antigen-binding molecule comprising:
 - (1) a cancer-specific antigen-binding Fab domain;
 - (2) a CD137-binding Fab domain; and
 - (3) an FcRn-binding domain, wherein the FcRn-binding domain is an antibody Fc region having decreased Fc γ receptor-binding activity,and
 - a second antigen-binding molecule comprising:
 - (1) a cancer-specific antigen-binding domain;
 - (2) a T cell receptor complex-binding domain and

(3) an FcRn-binding domain, wherein the FcRn-binding domain is an antibody Fc region having decreased Fc γ receptor-binding activity, in the manufacture of a medicament for the treatment of cancer.

9. The composition of claim 5 or the use of claim 8, wherein the T cell receptor complex-binding domain is a T cell receptor-binding domain.
10. The composition of claim 5 or the use of claim 8, wherein the T cell receptor complex-binding domain is a CD3-binding domain.
11. The composition of claim 5 or the use of claim 8, wherein the second antigen-binding molecule is an antibody.
12. The composition of any one of claims 3 to 5 or the use of any one of claims 6 to 11, wherein the composition or medicament is cytotoxicity-inducing composition.
13. A kit comprising:
 - a) a pharmaceutical composition comprising as an active ingredient a first agonistic antigen-binding molecule that comprises:
 - (1) a cancer-specific antigen-binding Fab domain;
 - (2) a CD137-binding Fab domain; and
 - (3) an FcRn-binding domain, wherein the FcRn-binding domain is an antibody Fc region having decreased Fc γ receptor-binding activity; and
 - b) a second antigen-binding molecule that comprises:
 - (1) a cancer-specific antigen-binding domain;
 - (2) a T cell receptor complex-binding domain; and
 - (3) an FcRn-binding domain, wherein the FcRn-binding domain is an antibody Fc region having decreased Fc γ receptor-binding activity;

wherein the composition is for use concomitantly with the second antigen-binding molecule in the treatment of cancer.
14. Use of a kit comprising:
 - a) a pharmaceutical composition comprising as an active ingredient a first agonistic antigen-binding molecule that comprises:
 - (1) a cancer-specific antigen-binding Fab domain;
 - (2) a CD137-binding Fab domain; and
 - (3) an FcRn-binding domain, wherein the FcRn-binding domain is an antibody Fc region having decreased Fc γ receptor-binding activity,

in the manufacture of a medicament for the treatment of cancer; and
 - b) a second antigen-binding molecule that comprises:
 - (1) a cancer-specific antigen-binding domain;
 - (2) a T cell receptor complex-binding domain; and

- (3) an FcRn-binding domain, wherein the FcRn-binding domain is an antibody Fc region having decreased Fc γ receptor-binding activity, wherein the composition is to be used concomitantly with the second antigen-binding molecule.
15. Use of a first agonistic antigen-binding molecule comprising:
- (1) a cancer-specific antigen-binding Fab domain;
 - (2) a CD137-binding Fab domain; and
 - (3) an FcRn-binding domain, wherein the FcRn-binding domain is an antibody Fc region having decreased Fc γ receptor-binding activity,
- in the manufacture of a medicament for the treatment of cancer, wherein the medicament is to be used concomitantly with a second antigen-binding molecule that comprises:
- (1) a cancer-specific antigen-binding domain;
 - (2) a T cell receptor complex-binding domain; and
 - (3) an FcRn-binding domain, wherein the FcRn-binding domain is an antibody Fc region having decreased Fc γ receptor-binding activity.
16. The kit of claim 13 or the use of claim 14 or claim 15, wherein the composition or medicament is cytotoxicity-inducing.
17. The kit of claim 13 or the use of any one of claims 14 to 16, wherein the T cell receptor complex-binding domain is a T cell receptor-binding domain.
18. The kit of claim 13 or the use of any one of claims 14 to 16, wherein the T cell receptor complex-binding domain is a CD3-binding domain.
19. The kit of claim 13 or the use of any one of claims 14 to 18, wherein the first antigen-binding molecule and/or the second antigen-binding molecule is an antibody.
20. The kit of claim 13 or the use of any one of claims 14 to 19, wherein the composition or medicament is to be administered either simultaneously with or separately from the second antigen-binding molecule.
21. A kit comprising:
- a) a pharmaceutical composition comprising as an active ingredient a second antigen-binding molecule that comprises:
 - (1) a cancer-specific antigen-binding domain;
 - (2) a T cell receptor complex-binding domain; and
 - (3) an FcRn-binding domain, wherein the FcRn-binding domain is an antibody Fc region having decreased Fc γ receptor-binding activity, in the manufacture of a medicament for the treatment of cancer, and
 - b) a first agonistic antigen-binding molecule that comprises:
 - (1) a cancer-specific antigen-binding Fab domain;

- (2) a CD137-binding Fab domain; and
 - (3) an FcRn-binding domain, wherein the FcRn-binding domain is an antibody Fc region having decreased Fc γ receptor-binding activity,
- wherein the composition is for use concomitantly with the first antigen-binding molecule in the treatment of cancer.

22. Use of a kit comprising:

- a) a pharmaceutical composition comprising as an active ingredient a second antigen-binding molecule that comprises:
 - (1) a cancer-specific antigen-binding domain;
 - (2) a T cell receptor complex-binding domain; and
 - (3) an FcRn-binding domain, wherein the FcRn-binding domain is an antibody Fc region having decreased Fc γ receptor-binding activity,
 in the manufacture of a medicament for the treatment of cancer, and
- b) a first agonistic antigen-binding molecule that comprises:
 - (1) a cancer-specific antigen-binding Fab domain;
 - (2) a CD137-binding Fab domain; and
 - (3) an FcRn-binding domain, wherein the FcRn-binding domain is an antibody Fc region having decreased Fc γ receptor-binding activity,
 wherein the composition is to be used concomitantly with the first agonistic antigen-binding molecule.

23. Use of a second antigen-binding molecule comprising:

- (1) a cancer-specific antigen-binding domain;
- (2) a T cell receptor complex-binding domain; and
- (3) an FcRn-binding domain, wherein the FcRn-binding domain is an antibody Fc region having decreased Fc γ receptor-binding activity, in the manufacture of a medicament for the treatment of cancer, wherein the medicament is to be used concomitantly with a first agonistic antigen-binding molecule that comprises:
 - (1) a cancer-specific antigen-binding Fab domain;
 - (2) a CD137-binding Fab domain; and
 - (3) an FcRn-binding domain, wherein the FcRn-binding domain is an antibody Fc region having decreased Fc γ receptor-binding activity.

24. The kit of claim 21 or the use of claim 22 or claim 23, wherein the composition or medicament is cytotoxicity-inducing composition.

25. The kit of claim 21 or the use of any one of claims 22 to 24, wherein the T cell receptor complex-binding domain is a T cell receptor-binding domain.

26. The kit of claim 21 or the use of any one of claims 22 to 24, wherein the T cell receptor complex-binding domain is a CD3-binding domain.

27. The kit of claim 21 or the use of any one of claims 22 to 26, wherein the first agonistic antigen-binding molecule and/or the second antigen-binding molecule is an antibody.
28. The kit of claim 21 or the use of any one of claims 22 to 27, wherein the composition or medicament is to be administered either simultaneously with or separately from the first agonistic antigen-binding molecule.