


Case 23239BENZIMIDAZOLE DERIVATIVES AS 5-HT₆, 5-HT_{2A}

This invention relates to substituted benzimidazolone and dihydroindolone compounds, and associated compositions, methods for use as therapeutic agents, and methods of preparation thereof.

5 The invention provides compounds of the formula I:



or a pharmaceutically acceptable salt thereof,

wherein:

m is from 0 to 3;

10 n is 1 or 2;

X is:

-NR^a-;

-O-;

-S-;

15 -CR^bR^c-; or

-C(O)-;

wherein:

R^a is hydrogen or alkyl;

R^b is hydrogen, fluoro or alkyl;

20 R^c is hydrogen, fluoro, alkyl, hydroxy; or alkoxy; or

R^b and R^c together form oxo; or


R^b and R^c together with the atom to which they are attached may form a three to six-membered optionally substituted ring that optionally includes a heteroatom selected from O, N and S;

25 Ar is:

optionally substituted aryl; or

- 2 -

optionally substituted heteroaryl;



p is from 1 to 4;

Y is:

5

-O-;

-NR^d-; or-CR^eR^f-;wherein R^d, R^e and R^f each independently is hydrogen or alkyl;each R² is independently:

10

halo;

alkyl;

haloalkyl;

haloalkoxy;

alkoxy;

15

heteroalkyl;

cyano;

-(CH₂)_q-S(O)_r-R^g;-(CH₂)_q-C(=O)-NR^hRⁱ;-(CH₂)_q-SO₂-NR^hRⁱ;

20

-(CH₂)_q-N(R^j)-C(=O)-R^k, or-(CH₂)_q-C(=O)-R^k;

wherein:

q is 0 or 1;

r is from 0 to 2; and

25

R^g, R^h, Rⁱ and R^j each independently is hydrogen or alkyl, and R^k

is

hydrogen, alkyl, alkoxy or hydroxy;

R³ and R⁴ each independently is hydrogen or alkyl;R⁵ and R⁶ each independently is hydrogen or alkyl; andR⁷ and R⁸ each independently is hydrogen or alkyl; or

30

R⁷ and R⁸ together with the nitrogen to which they are attached may form a

four to seven-membered optionally substituted ring that optionally includes an

additional heteroatom selected from O, N and S; or

one of R⁷ and R⁸ and one of R⁵ and R⁶ together with the atoms to which they are attached may form a four to seven-membered optionally substituted ring that optionally includes an additional heteroatom selected from O, N and S; or

one of R⁷ and R⁸ together with R^d and the atoms to which they are attached
5 may form a four to seven-membered optionally substituted ring that optionally includes an additional heteroatom selected from O, N and S; or

one of R⁷ and R⁸ and one of R^e and R^f together with the atoms to which they are attached may form a four to seven-membered ring that optionally includes an additional heteroatom selected from O, N and S.

10 The invention further provides compositions comprising, methods for using, and methods for preparing the aforementioned compounds.

The actions of 5-hydroxytryptamine (5-HT) as a major modulatory neurotransmitter in the brain are mediated through a number of receptor families termed
15 5-HT₁, 5-HT₂, 5-HT₃, 5-HT₄, 5-HT₅, 5-HT₆, and 5-HT₇. Based on a high level of 5-HT₆ receptor mRNA in the brain, it has been stated that the 5-HT₆ receptor may play a role in the pathology and treatment of central nerve system disorders. In particular, 5-HT₂-selective and 5-HT₆ selective ligands have been identified as potentially useful in the treatment of certain CNS disorders such as Parkinson's disease, Huntington's disease,
20 anxiety, depression, manic depression, psychoses, epilepsy, obsessive compulsive disorders, mood disorders, migraine, Alzheimer's disease (enhancement of cognitive memory), sleep disorders, feeding disorders such as anorexia, bulimia and obesity, panic attacks, akathisia, attention deficit hyperactivity disorder (ADHD), attention deficit disorder (ADD), withdrawal from drug abuse such as cocaine, ethanol, nicotine and
25 benzodiazepines, schizophrenia, and also disorders associated with spinal trauma and/or head injury such as hydrocephalus. Such compounds are also expected to be of use in the treatment of certain gastrointestinal (GI) disorders such as functional bowel disorder. See for example, B.L. Roth et al., J. Pharmacol. Exp. Ther., 1994, 268, pages 1403-14120, D. R. Sibley et al., Mol. Pharmacol., 1993, 43, 320-327, A.J. Sleight et al.,
30 Neurotransmission, 1995, 11, 1-5, and A. J. Sleight et al., Serotonin ID Research Alert, 1997, 2(3), 115-8.

While some 5-HT₆ and 5-HT_{2A} modulators have been disclosed, there continues to be a need for compounds that are useful for modulating the 5-HT₆ receptor, the 5-HT_{2A} receptor, or both.

The invention provides benzimidazolone and dihydroindolone compounds, related compositions, methods for use as therapeutic agents, and methods of preparation thereof.

All publications cited in this disclosure are incorporated herein by reference in
5 their entirety.

Unless otherwise stated, the following terms used in this Application, including the specification and claims, have the definitions given below. It must be noted that, as used in the specification and the appended claims, the singular forms "a", "an," and "the" include plural referents unless the context clearly dictates otherwise.

10 "Agonist" refers to a compound that enhances the activity of another compound or receptor site.

"Alkyl" means the monovalent linear or branched saturated hydrocarbon moiety, consisting solely of carbon and hydrogen atoms, having from one to twelve carbon atoms. "Lower alkyl" refers to an alkyl group of one to six carbon atoms (i.e., "C₁-C₆alkyl").

15 Examples of alkyl groups include, but are not limited to, methyl, ethyl, propyl, isopropyl, isobutyl, sec-butyl, tert-butyl, pentyl, n-hexyl, octyl, dodecyl, and the like.

"Alkylene" means a linear saturated divalent hydrocarbon radical of one to six carbon atoms or a branched saturated divalent hydrocarbon radical of three to six carbon atoms, e.g., methylene, ethylene, 2,2-dimethylethylene, propylene, 2-methylpropylene,
20 butylene, pentylene, and the like.

"Alkenylene" means a linear unsaturated divalent hydrocarbon radical of two to six carbon atoms or a branched saturated divalent hydrocarbon radical of three to six carbon atoms, e.g., ethenylene (-CH=CH-), 2,2-dimethylethenylene, propenylene, 2-methylpropenylene, butenylene, pentenylene, and the like.

25 "Alkoxy" means a moiety of the formula -OR, wherein R is an alkyl moiety as defined herein. Examples of alkoxy moieties include, but are not limited to, methoxy, ethoxy, isopropoxy, and the like.

"Aminoalkoxy" means a group -ORR' wherein R' is amino and R is alkylene. Exemplarily aminoalkoxy include aminoethoxy, 2-aminopropoxy, 3-aminopropoxy,
30 and the like. The amino moiety of "aminoalkoxy" may be substituted once or twice with alkyl to provide "alkylaminoalkoxy" and "dialkylaminoalkoxy" respectively.

"Alkylaminoalkoxy" includes methylaminoethoxyl, methylaminopropoxy, ethylaminoethoxy, and the like. "Dialkylaminoalkoxy" includes dimethylaminoethoxyl, dimethylaminopropoxy, N-methyl-N-ethylaminoethoxy, and the like.

35 "Antagonist" refers to a compound that diminishes or prevents the action of another compound or receptor site.

"Aryl" means a monovalent cyclic aromatic hydrocarbon moiety consisting of a mono-, bi- or tricyclic aromatic ring. The aryl group can be optionally substituted as defined herein. Examples of aryl moieties include, but are not limited to, phenyl, naphthyl, naphthalenyl, phenanthryl, fluorenyl, indenyl, pentalenyl, azulenyl, oxydiphenyl, biphenyl, methylenediphenyl, aminodiphenyl, diphenylsulfidyl, diphenylsulfonyl, diphenylisopropylidenyl, benzodioxanyl, benzofuranyl, benzodioxyl, benzopyranyl, benzoxazinyl, benzoxazinonyl, benzopiperadiny, benzopiperazinyl, benzopyrrolidinyl, benzomorpholinyl, methylenedioxyphenyl, ethylenedioxyphenyl, and the like, including partially hydrogenated derivatives thereof.

10 "Arylene" means a divalent aryl radical wherein aryl is as defined herein.

"Arylene" includes, for example, ortho-, meta- and para- phenylene (1,2-phenylene, 1,3-phenylene and 1,4-phenylene respectively), which may be optionally substituted as defined herein.

"Arylalkyl" and "Aralkyl", which may be used interchangeably, mean a radical
15 $-R^aR^b$ where R^a is an alkylene group and R^b is an aryl group as defined herein; e.g., benzyl, phenylethyl, 3-(3-chlorophenyl)-2-methylpentyl, and the like are examples of arylalkyl.

"Cycloalkyl" means a saturated carbocyclic moiety consisting of mono- or bicyclic rings. Cycloalkyl can optionally be substituted with one or more substituents, wherein
20 each substituent is independently hydroxy, alkyl, alkoxy, halo, haloalkyl, amino, monoalkylamino, or dialkylamino, unless otherwise specifically indicated. Examples of cycloalkyl moieties include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, and the like, including partially unsaturated derivatives thereof such as cyclohexenyl, cyclopentenyl, and the like.

25 "Cycloalkylalkyl" means a moiety of the formula $-R'-R''$, where R' is alkylene and R'' is cycloalkyl as defined herein.

"Heteroalkyl" means an alkyl radical as defined herein wherein one, two or three hydrogen atoms have been replaced with a substituent independently selected from the group consisting of $-OR^a$, $-NR^bR^c$, and $-S(O)_nR^d$ (where n is an integer from 0 to 2), with
30 the understanding that the point of attachment of the heteroalkyl radical is through a carbon atom, wherein R^a is hydrogen, acyl, alkyl, cycloalkyl, or cycloalkylalkyl; R^b and R^c are independently of each other hydrogen, acyl, alkyl, cycloalkyl, or cycloalkylalkyl; and when n is 0, R^d is hydrogen, alkyl, cycloalkyl, or cycloalkylalkyl, and when n is 1 or 2, R^d is alkyl, cycloalkyl, cycloalkylalkyl, amino, acylamino, monoalkylamino, or dialkylamino.
35 Representative examples include, but are not limited to, 2-hydroxyethyl, 3-hydroxypropyl, 2-hydroxy-1-hydroxymethylethyl, 2,3-dihydroxypropyl, 1-hydroxymethylethyl, 3-hydroxybutyl, 2,3-dihydroxybutyl, 2-hydroxy-1-methylpropyl, 2-

aminoethyl, 3-aminopropyl, 2-methylsulfonylethyl, aminosulfonylmethyl, aminosulfonylethyl, aminosulfonylpropyl, methylaminosulfonylmethyl, methylaminosulfonylethyl, methylaminosulfonylpropyl, and the like.

“Heteroaryl” means a monocyclic or bicyclic monovalent radical of 5 to 12 ring atoms having at least one aromatic ring containing one, two, or three ring heteroatoms selected from N, O, or S, the remaining ring atoms being C, with the understanding that the attachment point of the heteroaryl radical will be on an aromatic ring. The heteroaryl ring may be optionally substituted as defined herein. Examples of heteroaryl moieties include, but are not limited to, imidazolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, oxadiazolyl, thiadiazolyl, pyrazinyl, thienyl, benzothienyl, thiophenyl, furanyl, pyranal, pyridyl, pyridinyl, pyridazyl, pyrrolyl, pyrazolyl, pyrimidyl, quinoliny, isoquinoliny, benzofuryl, benzothiophenyl, benzothiopyranal, benzimidazolyl, benzooxazolyl, benzooxadiazolyl, benzothiazolyl, benzothiadiazolyl, benzopyranal, indolyl, isoindolyl, triazolyl, triazinyl, quinoxaliny, purinyl, quinazolinyl, quinoliziny, naphthyridiny, pteridiny, carbazolyl, azepiny, diazepiny, acridiny and the like, including partially hydrogenated derivatives thereof.

“Heteroarylene” means a divalent heteroaryl radical wherein heteroaryl is as defined herein. “Heteroarylene” may be optionally substituted as defined herein. “Heteroarylene” includes, for example, indolylene, pyrimidinylene, and the like.

The terms “halo” and “halogen”, which may be used interchangeably, refer to a substituent fluoro, chloro, bromo, or iodo.

“Haloalkyl” means alkyl as defined herein in which one or more hydrogen has been replaced with same or different halogen. Exemplary haloalkyls include $-\text{CH}_2\text{Cl}$, $-\text{CH}_2\text{CF}_3$, $-\text{CH}_2\text{CCl}_3$, perfluoroalkyl (e.g., $-\text{CF}_3$), and the like.

“Heterocycloamino” means a saturated ring wherein at least one ring atom is N, NH or N-alkyl and the remaining ring atoms form an alkylene group.

“Heterocyclyl” means a monovalent saturated moiety, consisting of one to three rings, incorporating one, two, or three or four heteroatoms (chosen from nitrogen, oxygen or sulfur). The heterocyclyl ring may be optionally substituted as defined herein. Examples of heterocyclyl moieties include, but are not limited to, azetidiny, piperidiny, piperazinyl, homopiperazinyl, azepiny, pyrrolidinyl, pyrazolidiny, imidazolidiny, imidazolidinyl, pyridiny, pyridazinyl, pyrimidinyl, oxazolidiny, isoxazolidiny, morpholiny, thiazolidiny, isothiazolidiny, quinuclidiny, quinoliny, isoquinoliny, benzimidazolyl, thiadiazolyldiny, benzothiazolidiny, benzoazolyldiny, dihydrofuryl, tetrahydrofuryl, dihydropyranal, tetrahydropyranal, thiamorpholiny, thiamorpholiny sulfoxide, thiamorpholiny sulfone, dihydroquinoliny,

dihydrisoquinoliny, tetrahydroquinoliny, tetrahydrisoquinoliny, and the like, including partially unsaturated derivatives thereof.

"Heterocyclyloxy" means a group -OR wherein R is heterocyclyl.

"Heterocyclyloxy" includes, by way of example, azetidinyloxy, pyrrolidinyloxy,

5 piperidinyloxy, azepinyloxy and the like.

"Heterocyclylalkoxy" means a group -ORR' wherein R' is heterocyclyl and R is alkylene. Exemplary heterocyclylalkoxy include azetinylmethoxy, pyrrolodinylmethoxy, piperidinylmethoxy, azetinyloxy, pyrrolodinyloxy, piperidinyloxy and the like.

"Optionally substituted", when used in association with "aryl", "arylene", "phenyl",
10 "phenylene", "heteroaryl", heteroarylene or "heterocyclyl", means an aryl, arylene, phenyl, phenylene, heteroaryl, heteroarylene, or heterocyclyl which is optionally substituted independently with one to four substituents, preferably one or two substituents selected from alkyl, cycloalkyl, cycloalkylalkyl, heteroalkyl, hydroxyalkyl, heteroalkylamino, heteroalkoxy, halo, nitro, cyano, hydroxy, alkoxy, amino, acylamino, mono-alkylamino,
15 di-alkylamino, haloalkyl, haloalkoxy, heteroalkyl, -COR (where R is hydrogen, alkyl, phenyl or phenylalkyl), -(CR'R'')_n-COOR (where n is an integer from 0 to 5, R' and R'' are independently hydrogen or alkyl, and R is hydrogen, alkyl, cycloalkyl, cycloalkylalkyl, phenyl or phenylalkyl), or -(CR'R'')_n-CONR^aR^b (where n is an integer from 0 to 5, R' and R'' are independently hydrogen or alkyl, and R^a and R^b are, independently of each
20 other, hydrogen, alkyl, cycloalkyl, cycloalkylalkyl, phenyl or phenylalkyl.

"Leaving group" means the group with the meaning conventionally associated with it in synthetic organic chemistry, i.e., an atom or group displaceable under substitution reaction conditions. Examples of leaving groups include, but are not limited to, halogen, alkane- or arylenesulfonyloxy, such as methanesulfonyloxy,
25 ethanesulfonyloxy, thiomethyl, benzenesulfonyloxy, tosyloxy, and thienyloxy, dihalophosphinoyloxy, optionally substituted benzyloxy, isopropyloxy, acyloxy, and the like.

"Modulator" means a molecule that interacts with a target. The interactions include, but are not limited to, agonist, antagonist, and the like, as defined herein.

30 "Optional" or "optionally" means that the subsequently described event or circumstance may but need not occur, and that the description includes instances where the event or circumstance occurs and instances in which it does not.

"Disease state" means any disease, condition, symptom, or indication.

"Inert organic solvent" or "inert solvent" means the solvent is inert under the
35 conditions of the reaction being described in conjunction therewith, including for example, benzene, toluene, acetonitrile, tetrahydrofuran, N,N-dimethylformamide,

chloroform, methylene chloride or dichloromethane, dichloroethane, diethyl ether, ethyl acetate, acetone, methyl ethyl ketone, methanol, ethanol, propanol, isopropanol, tert-butanol, dioxane, pyridine, and the like. Unless specified to the contrary, the solvents used in the reactions of the present invention are inert solvents.

5 “Pharmaceutically acceptable” means that which is useful in preparing a pharmaceutical composition that is generally safe, non-toxic, and neither biologically nor otherwise undesirable and includes that which is acceptable for veterinary as well as human pharmaceutical use.

 “Pharmaceutically acceptable salts” of a compound means salts that are
10 pharmaceutically acceptable, as defined herein, and that possess the desired pharmacological activity of the parent compound. Such salts include:

 acid addition salts formed with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, and the like; or formed with organic acids such as acetic acid, benzenesulfonic acid, benzoic, camphorsulfonic acid,
15 citric acid, ethanesulfonic acid, fumaric acid, glucoheptonic acid, gluconic acid, glutamic acid, glycolic acid, hydroxynaphthoic acid, 2-hydroxyethanesulfonic acid, lactic acid, maleic acid, malic acid, malonic acid, mandelic acid, methanesulfonic acid, muconic acid, 2-naphthalenesulfonic acid, propionic acid, salicylic acid, succinic acid, tartaric acid, p-toluenesulfonic acid, trimethylacetic acid, and the like; or

20 salts formed when an acidic proton present in the parent compound either is replaced by a metal ion, e.g., an alkali metal ion, an alkaline earth ion, or an aluminum ion; or coordinates with an organic or inorganic base. Acceptable organic bases include diethanolamine, ethanolamine, N-methylglucamine, triethanolamine, tromethamine, and the like. Acceptable inorganic bases include aluminum hydroxide, calcium
25 hydroxide, potassium hydroxide, sodium carbonate and sodium hydroxide.

 The preferred pharmaceutically acceptable salts are the salts formed from acetic acid, hydrochloric acid, sulphuric acid, methanesulfonic acid, maleic acid, phosphoric acid, tartaric acid, citric acid, sodium, potassium, calcium, zinc, and magnesium.

 It should be understood that all references to pharmaceutically acceptable salts
30 include solvent addition forms (solvates) or crystal forms (polymorphs) as defined herein, of the same acid addition salt.

 “Protective group” or “protecting group” means the group which selectively blocks one reactive site in a multifunctional compound such that a chemical reaction can be carried out selectively at another unprotected reactive site in the meaning
35 conventionally associated with it in synthetic chemistry. Certain processes of this invention rely upon the protective groups to block reactive nitrogen and/or oxygen atoms present in the reactants. For example, the terms “amino-protecting group” and “nitrogen

protecting group” are used interchangeably herein and refer to those organic groups intended to protect the nitrogen atom against undesirable reactions during synthetic procedures. Exemplary nitrogen protecting groups include, but are not limited to, trifluoroacetyl, acetamido, benzyl (Bn), benzyloxycarbonyl (carbobenzyloxy, CBZ), p-methoxybenzyloxycarbonyl, p-nitrobenzyloxycarbonyl, tert-butoxycarbonyl (BOC), and the like. The artisan in the art will know how to choose a group for the ease of removal and for the ability to withstand the following reactions.

“Solvates” means solvent addition forms that contain either stoichiometric or non stoichiometric amounts of solvent. Some compounds have a tendency to trap a fixed molar ratio of solvent molecules in the crystalline solid state, thus forming a solvate. If the solvent is water the solvate formed is a hydrate, when the solvent is alcohol, the solvate formed is an alcoholate. Hydrates are formed by the combination of one or more molecules of water with one of the substances in which the water retains its molecular state as H₂O, such combination being able to form one or more hydrate.

“Subject” means mammals and non-mammals. Mammals means any member of the mammalia class including, but not limited to, humans; non-human primates such as chimpanzees and other apes and monkey species; farm animals such as cattle, horses, sheep, goats, and swine; domestic animals such as rabbits, dogs, and cats; laboratory animals including rodents, such as rats, mice, and guinea pigs; and the like. Examples of non-mammals include, but are not limited to, birds, and the like. The term “subject” does not denote a particular age or sex.

“Therapeutically effective amount” means an amount of a compound that, when administered to a subject for treating a disease state, is sufficient to effect such treatment for the disease state. The “therapeutically effective amount” will vary depending on the compound, disease state being treated, the severity of the disease treated, the age and relative health of the subject, the route and form of administration, the judgement of the attending medical or veterinary practitioner, and other factors.

The terms “those defined above” and “those defined herein” when referring to a variable incorporates by reference the broad definition of the variable as well as preferred, more preferred and most preferred definitions, if any.


“Treating” or “treatment” of a disease state includes:

- (i) preventing the disease state, i.e. causing the clinical symptoms of the disease state not to develop in a subject that may be exposed to or predisposed to the disease state, but does not yet experience or display symptoms of the disease state.
- (ii) inhibiting the disease state, i.e., arresting the development of the disease state or its clinical symptoms, or

(iii) relieving the disease state, i.e., causing temporary or permanent regression of the disease state or its clinical symptoms.

The terms “treating”, “contacting” and “reacting” when referring to a chemical reaction means adding or mixing two or more reagents under appropriate conditions to produce the indicated and/or the desired product. It should be appreciated that the reaction which produces the indicated and/or the desired product may not necessarily result directly from the combination of two reagents which were initially added, i.e., there may be one or more intermediates which are produced in the mixture which ultimately leads to the formation of the indicated and/or the desired product.


In general, the nomenclature used in this Application is based on AUTONOMTM v.4.0, a Beilstein Institute computerized system for the generation of IUPAC systematic nomenclature. For convenience, the IUPAC numbering of the positions of representative benzimidazolone and dihydroindolone compounds described herein is shown by the formula:



wherein the 1-position is substituted with arylalkyl or heteroarylalkyl as described below.

Chemical structures shown herein were prepared using ISIS[®] version 2.2. Any open valency appearing on a carbon, oxygen or nitrogen atom in the structures herein indicates the presence of a hydrogen. Where a chiral center is present in a structure but no specific enantiomer is shown, it should be understood that the structure encompasses both enantiomers associated with the chiral center.

The invention provides compounds of the formula I:



or a pharmaceutically acceptable salt thereof,

wherein:

m is from 0 to 3;

n is 1 or 2;

X is:

-NR^a-;
 -O-;
 -S-;
 -CR^bR^c-; or
 -C(O)-;


wherein:

R^a is hydrogen or alkyl;
 R^b is hydrogen, fluoro or alkyl;
 R^c is hydrogen, fluoro, alkyl, hydroxy; or alkoxy; or
 R^b and R^c together form oxo; or
 R^b and R^c together with the atom to which they are attached may
 form a three to six-membered ring that optionally includes a
 heteroatom selected from O, N and S;

Ar is:

optionally substituted aryl; or
 optionally substituted heteroaryl;

R¹ is a group of the formula



p is from 1 to 4;


Y is:

-O-;
 -NR^d-; or
 -CR^eR^f-;

wherein R^d, R^e and R^f each independently is hydrogen or alkyl;

each R² is independently:

halo;
 alkyl;
 haloalkyl;
 haloalkoxy;
 alkoxy;
 heteroalkyl;
 cyano;
 -(CH₂)_q-S(O)_r-R^g;
 -(CH₂)_q-C(=O)-NR^hRⁱ;



wherein:

5 q is 0 or 1;

r is from 0 to 2; and

R^g, R^h, R^i and R^j each independently is hydrogen or alkyl, and R^k

is hydrogen,

alkyl, alkoxy or hydroxy;

10 R³ and R⁴ each independently is hydrogen or alkyl;

R⁵ and R⁶ each independently is hydrogen or alkyl; and

R⁷ and R⁸ each independently is hydrogen or alkyl; or

R⁷ and R⁸ together with the nitrogen to which they are attached may form a

four to seven-membered optionally substituted ring that optionally includes an

15 additional heteroatom selected from O, N and S; or

one of R⁷ and R⁸ and one of R⁵ and R⁶ together with the atoms to which they are attached may form a four to seven-membered ring that optionally includes an additional heteroatom selected from O, N and S; or

one of R⁷ and R⁸ together with R^d and the atoms to which they are attached
20 may form a four to seven-membered optionally substituted ring that optionally includes
an additional heteroatom selected from O, N and S; or

one of R⁷ and R⁸ and one of R^e and R^f together with the atoms to which they are attached may form a four to seven-membered optionally substituted ring that optionally includes an additional heteroatom selected from O, N and S.

25 It should be understood that the scope of this invention encompasses not only the various isomers which may exist but also the various mixture of isomers which may be formed. Furthermore, the scope of the present invention also encompasses solvates and salts of compounds of formula I.

In many embodiments of formula I, n is 1.

30 In many embodiments of formula I, R³ and R⁴ are hydrogen.

In many embodiments of formula I, p is 2 or 3.

In certain embodiments of formula I, Ar is optionally substituted aryl, such as optionally substituted phenyl or optionally substituted naphthyl such as, for example, phenyl or naphthyl optionally substituted independently one, two, three or four times
35 with halo, alkyl, haloalkyl, haloalkoxy, alkoxy, heteroalkyl, cyano, nitro, amino (including alkylamino and dialkylamino), $-(CH_2)_q-S(O)_r-R^g$; $-(CH_2)_q-C(=O)-NR^hR^i$; $-(CH_2)_q-SO_2-NR^hR^i$; $-(CH_2)_q-N(R^j)-C(=O)-R^k$, or $-(CH_2)_q-C(=O)-R^k$, wherein q is 0 or 1, r is

from 0 to 2, R^g , R^h , R^i and R^j each independently is hydrogen or alkyl, and R^k is hydrogen, alkyl, alkoxy or hydroxy. More preferably Ar is optionally substituted phenyl.

In other embodiments of formula I, Ar is optionally substituted heteroaryl. Preferred heteroaryl include pyridinyl, pyrimidinyl, thienyl, furanyl, pyrrolyl, pyrazolyl, imidazolyl, oxazolyl, thiazolyl, quinoline and isoquinoline, and more preferably pyridinyl, pyrimidinyl, thienyl and furanyl..


In certain embodiments of formula I, m is 0 or 1 and R^2 is halo, preferably fluoro or chloro.

In certain embodiments of formula I, m is 0.

In certain embodiments of formula I, m is 1 and R^2 is fluoro or chloro.

In certain embodiments of formula I, m is 2 and each R^2 is independently fluoro or chloro at the 5 and six position of the benzimidazolone or dihydroindolone ring system.

In many embodiments the compounds of the invention may be represented by formula II:




wherein:

s is from 1 to 4;

each R^9 is independently:

- halo;
- alkyl;
- haloalkyl;
- haloalkoxy;
- alkoxy;
- heteroalkyl;
- cyano;
- $-(CH_2)_q-S(O)_r-R^g$;
- $-(CH_2)_q-C(=O)-NR^hR^i$;
- $-(CH_2)_q-SO_2-NR^hR^i$;



wherein:

q is 0 or 1;

5 r is from 0 to 2; and

R^g, R^h, Rⁱ and R^j each independently is hydrogen or alkyl, and R^k
is hydrogen, alkyl, alkoxy or hydroxy; and

m, p, X, Y, R², R³, R⁴, R⁵, R⁶, R⁷ and R⁸ are as defined herein.

In certain embodiments of either of formula I or formula II, X is -NR^a-.

10 In certain embodiments of either of formula I or formula II, X is -O-.

In certain embodiments of either of formula I or formula II, X is -CR^bR^c-.

In certain embodiments of either of formula I or formula II, X is -C(O)-;

In certain embodiments of either of formula I or formula II, Y is -O-.

In certain embodiments of either of formula I or formula II, Y is -NR^d-.

15 In certain embodiments of either of formula I or formula II, Y is -CR^eR^f-.

In certain embodiments of either of formula I or formula II, m is 0 or 1.

In certain embodiments of formula I or formula II, R³ and R⁴ are hydrogen.

In certain embodiments of formula I or formula II, R³ and R⁴ are alkyl, preferably methyl.

20 In certain embodiments of formula I or formula II, one of R³ and R⁴ is hydrogen and the other is alkyl, preferably methyl.

In certain embodiments of formula I or formula II, m is 0 or 1 and R² is halo, preferably fluoro or chloro.

In certain embodiments of formula I or formula II, m is 0.

25 In certain embodiments of formula I or formula II, m is 1 and R² is fluoro or chloro.

In certain embodiments of formula I or formula II, m is 2 and each R² is independently fluoro or chloro at the 5 and six position of the benzimidazolone or dihydroindolone ring system.

30 In certain embodiments of either of formula I or formula II where X is -NR^a-, R^a is hydrogen. In other embodiments of either of formula I or formula II where X is -NR^a-, R^a is alkyl, preferably methyl.

In certain embodiments of either of formula I or formula II where X is -CR^bR^c-, R^b and R^c are hydrogen.

35 In other embodiments of either of formula I or formula II where X is -CR^bR^c-, R^b and R^c are alkyl, preferably methyl.

In still other embodiments of either of formula I or formula II where X is $-\text{CR}^b\text{R}^c-$, one of R^b and R^c is hydrogen and the other is alkyl, preferably methyl.

In yet other embodiments of either of formula I or formula II where X is $-\text{CR}^b\text{R}^c-$, one of R^b and R^c is alkyl, preferably methyl, and the other is hydroxy.

- 5 In other embodiments of either of formula I or formula II where X is $-\text{CR}^b\text{R}^c-$, R^b and R^c together with the atom to which they are attached form a three to six-membered ring that optionally includes a heteroatom selected from O, N and S, and which may optionally be substituted with alkyl. Preferably in such embodiments R^b and R^c together with the atom to which they are attached form a three or four membered carbocyclic ring
- 10 that may optionally be substituted with alkyl.

In certain embodiments of either of formula I or formula II, n is 1, p is 2, and R^3 and R^4 each independently is hydrogen or methyl.

In certain embodiments of either of formula I or formula II, n is 1, p is 2, R^3 and R^4 are hydrogen, and Y is $-\text{O}-$.

- 15 In certain embodiments of either of formula I or formula II, n is 1, p is 2, R^3 and R^4 each independently is hydrogen or methyl, Y is $-\text{O}-$, and R^5 and R^6 are hydrogen.

- In certain embodiments of either of formula I or formula II, n is 1, p is 2, R^3 and R^4 each independently is hydrogen or methyl, Y is $-\text{O}-$, and one of R^7 and R^8 and one of R^5 and R^6 together with the atoms to which they are attached form a four to seven-
- 20 membered ring that may optionally be substituted with alkyl.

In certain embodiments of either of formula I or formula II, n is 1, p is 2, R^3 and R^4 are hydrogen, and Y is $-\text{NR}^d-$.

In certain embodiments of either of formula I or formula II, n is 1, p is 2, R^3 and R^4 are hydrogen Y is $-\text{NR}^d-$, and R^5 and R^6 are hydrogen.

- 25 In certain embodiments of either of formula I or formula II, n is 1, p is 2, R^3 and R^4 each independently is hydrogen or methyl, Y is $-\text{NR}^d-$, and one of R^7 and R^8 and one of R^5 and R^6 together with the atoms to which they are attached form a four to seven-membered ring that may optionally be substituted with alkyl.

- In certain embodiments of either of formula I or formula II, n is 1, p is 2, R^3 and R^4 each independently is hydrogen or methyl, Y is $-\text{NR}^d-$, and one of R^7 and R^8 together with R^d and the atoms to which they are attached form a four to seven-membered ring that may optionally be substituted with alkyl.
- 30

In certain embodiments of either of formula I or formula II, n is 1, p is 2, R^3 and R^4 each independently is hydrogen or methyl, and Y is $-\text{CR}^e\text{R}^f-$.

- 35 In certain embodiments of either of formula I or formula II, n is 1, p is 2, R^3 and R^4 are hydrogen, Y is $-\text{CR}^e\text{R}^f-$, and R^5 , R^6 , R^e and R^f are hydrogen.

In certain embodiments of either of formula I or formula II, n is 1, p is 2, R^3 and R^4 each independently is hydrogen or methyl, Y is $-CR^eR^f-$, and one of R^7 and R^8 and one of R^5 and R^6 together with the atoms to which they are attached form a four to seven-membered ring that may optionally be substituted with alkyl.

5 In certain embodiments of either of formula I or formula II, n is 1, p is 2, R^3 and R^4 each independently is hydrogen or methyl, Y is $-CR^eR^f-$, and one of R^7 and R^8 and one of R^e and R^f together with the atoms to which they are attached form a four to seven-membered ring that may optionally be substituted with alkyl.

In certain embodiments of either of formula I or formula II, n is 1, p is 3, and R^3 and R^4 each independently is hydrogen or methyl.

In certain embodiments of either of formula I or formula II, n is 1, p is 3, R^3 and R^4 each independently is hydrogen or methyl, and Y is $-O-$.

In certain embodiments of either of formula I or formula II, n is 1, p is 3, R^3 and R^4 each independently is hydrogen or methyl, Y is $-O-$, and R^5 and R^6 are hydrogen.

15 In certain embodiments of either of formula I or formula II, n is 1, p is 3, R^3 and R^4 each independently is hydrogen or methyl, Y is $-O-$, and one of R^7 and R^8 and one of R^5 and R^6 together with the atoms to which they are attached form a four to seven-membered ring that may optionally be substituted with alkyl.

In certain embodiments of either of formula I or formula II, n is 1, p is 3, R^3 and R^4 each independently is hydrogen or methyl, and Y is $-NR^d-$.

In certain embodiments of either of formula I or formula II, n is 1, p is 3, R^3 and R^4 each independently is hydrogen or methyl, Y is $-NR^d-$, and R^5 and R^6 are hydrogen.

In certain embodiments of either of formula I or formula II, n is 1, p is 3, R^3 and R^4 each independently is hydrogen or methyl, Y is $-NR^d-$, and one of R^7 and R^8 and one of R^5 and R^6 together with the atoms to which they are attached form a four to seven-membered ring that may optionally be substituted with alkyl.

In certain embodiments of either of formula I or formula II, n is 1, p is 3, R^3 and R^4 each independently is hydrogen or methyl, Y is $-NR^d-$, and one of R^7 and R^8 together with R^d and the atoms to which they are attached form a four to seven-membered ring that may optionally be substituted with alkyl.

In certain embodiments of either of formula I or formula II, n is 1, p is 3, R^3 and R^4 each independently is hydrogen or methyl, and Y is $-CR^eR^f-$.

In certain embodiments of either of formula I or formula II, n is 1, p is 3, R^3 and R^4 each independently is hydrogen or methyl, Y is $-CR^eR^f-$, and R^5 , R^6 , R^e and R^f are hydrogen.

In certain embodiments of either of formula I or formula II, n is 1, p is 3, R^3 and R^4 each independently is hydrogen or methyl, Y is $-CR^eR^f-$, and one of R^7 and R^8 and one

of R^5 and R^6 together with the atoms to which they are attached form a four to seven-membered ring that may optionally be substituted with alkyl.

In certain embodiments of either of formula I or formula II, n is 1, p is 3, R^3 and R^4 each independently is hydrogen or methyl, Y is $-CR^eR^f-$, and one of R^7 and R^8 and one
5 of R^e and R^f together with the atoms to which they are attached form a four to seven-membered ring that may optionally be substituted with alkyl.

In certain embodiments of either of formula I or formula II, n is 1, p is 2, R^3 and R^4 each independently is hydrogen or methyl, and X is $-NR^a-$.

In certain embodiments of either of formula I or formula II, n is 1, p is 2, R^3 and
10 R^4 each independently is hydrogen or methyl, X is $-NR^a-$, and Y is $-O-$.

In certain embodiments of either of formula I or formula II, n is 1, p is 2, R^3 and R^4 each independently is hydrogen or methyl, X is $-NR^a-$, Y is $-O-$, and R^5 and R^6 are hydrogen.

In certain embodiments of either of formula I or formula II, n is 1, p is 2, R^3 and
15 R^4 each independently is hydrogen or methyl, X is $-NR^a-$, Y is $-O-$, and one of R^7 and R^8 and one of R^5 and R^6 together with the atoms to which they are attached form a four to seven-membered ring that may optionally be substituted with alkyl.

In certain embodiments of either of formula I or formula II, n is 1, p is 2, R^3 and R^4 each independently is hydrogen or methyl, X is $-NR^a-$, and Y is $-NR^d-$.

In certain embodiments of either of formula I or formula II, n is 1, p is 2, R^3 and
20 R^4 each independently is hydrogen or methyl, X is $-NR^a-$, Y is $-NR^d-$, and R^5 and R^6 are hydrogen.

In certain embodiments of either of formula I or formula II, n is 1, p is 2, R^3 and R^4 each independently is hydrogen or methyl, X is $-NR^a-$, Y is $-NR^d-$, and one of R^7 and
25 R^8 and one of R^5 and R^6 together with the atoms to which they are attached form a four to seven-membered ring that may optionally be substituted with alkyl.

In certain embodiments of either of formula I or formula II, n is 1, p is 2, R^3 and R^4 are hydrogen, X is $-NR^a-$, Y is $-NR^d-$, and one of R^7 and R^8 together with R^d and the
atoms to which they are attached form a four to seven-membered ring that may
30 optionally be substituted with alkyl.

In certain embodiments of either of formula I or formula II, n is 1, p is 2, R^3 and R^4 each independently is hydrogen or methyl, X is $-NR^a-$, and Y is $-CR^eR^f-$.

In certain embodiments of either of formula I or formula II, n is 1, p is 2, R^3 and R^4 each independently is hydrogen or methyl, X is $-NR^a-$, Y is $-CR^eR^f-$, and R^5 , R^6 , R^e and
35 R^f are hydrogen.

In certain embodiments of either of formula I or formula II, n is 1, p is 2, R^3 and R^4 each independently is hydrogen or methyl, X is $-NR^a-$, Y is $-CR^eR^f-$, and one of R^7 and

R^8 and one of R^5 and R^6 together with the atoms to which they are attached form a four to seven-membered ring that may optionally be substituted with alkyl.

In certain embodiments of either of formula I or formula II, n is 1, p is 2, R^3 and R^4 each independently is hydrogen or methyl, X is $-NR^a$ -, Y is $-CR^eR^f$ -, and one of R^7 and R^8 and one of R^e and R^f together with the atoms to which they are attached form a four to seven-membered ring that may optionally be substituted with alkyl.

In certain embodiments of either of formula I or formula II, n is 1, p is 3, R^3 and R^4 each independently is hydrogen or methyl, and X is $-NR^a$ -.

In certain embodiments of either of formula I or formula II, n is 1, p is 3, R^3 and R^4 each independently is hydrogen or methyl, X is $-NR^a$ -, and Y is $-O$ -.

In certain embodiments of either of formula I or formula II, n is 1, p is 3, R^3 and R^4 each independently is hydrogen or methyl, X is $-NR^a$ -, Y is $-O$ -, and R^5 and R^6 are hydrogen.

In certain embodiments of either of formula I or formula II, n is 1, p is 3, R^3 and R^4 each independently is hydrogen or methyl, X is $-NR^a$ -, Y is $-O$ -, and one of R^7 and R^8 and one of R^5 and R^6 together with the atoms to which they are attached form a four to seven-membered ring that may optionally be substituted with alkyl.

In certain embodiments of either of formula I or formula II, n is 1, p is 3, R^3 and R^4 each independently is hydrogen or methyl, X is $-NR^a$ -, and Y is $-NR^d$ -.

In certain embodiments of either of formula I or formula II, n is 1, p is 3, R^3 and R^4 each independently is hydrogen or methyl, X is $-NR^a$ -, Y is $-NR^d$ -, and R^5 and R^6 are hydrogen.

In certain embodiments of either of formula I or formula II, n is 1, p is 3, R^3 and R^4 each independently is hydrogen or methyl, X is $-NR^a$ -, Y is $-NR^d$ -, and one of R^7 and R^8 and one of R^5 and R^6 together with the atoms to which they are attached form a four to seven-membered ring that may optionally be substituted with alkyl.

In certain embodiments of either of formula I or formula II, n is 1, p is 3, R^3 and R^4 each independently is hydrogen or methyl, X is $-NR^a$ -, Y is $-NR^d$ -, and one of R^7 and R^8 together with R^d and the atoms to which they are attached form a four to seven-membered ring that may optionally be substituted with alkyl.

In certain embodiments of either of formula I or formula II, n is 1, p is 3, R^3 and R^4 each independently is hydrogen or methyl, X is $-NR^a$ -, and Y is $-CR^eR^f$ -.

In certain embodiments of either of formula I or formula II, n is 1, p is 3, R^3 and R^4 each independently is hydrogen or methyl, X is $-NR^a$ -, Y is $-CR^eR^f$ -, and R^5 , R^6 , R^e and R^f are hydrogen.

In certain embodiments of either of formula I or formula II, n is 1, p is 3, R^3 and R^4 each independently is hydrogen or methyl, X is $-NR^a$ -, Y is $-CR^eR^f$ -, and one of R^7 and

R^8 and one of R^5 and R^6 together with the atoms to which they are attached form a four to seven-membered ring that may optionally be substituted with alkyl.

In certain embodiments of either of formula I or formula II, n is 1, p is 3, R^3 and R^4 each independently is hydrogen or methyl, X is $-NR^a-$, Y is $-CR^eR^f-$, and one of R^7 and R^8 and one of R^e and R^f together with the atoms to which they are attached form a four to seven-membered ring.

In certain embodiments of either of formula I or formula II, n is 1, p is 2, R^3 and R^4 each independently is hydrogen or methyl, and X is $-CR^bR^c-$.

In certain embodiments of either of formula I or formula II, n is 1, p is 2, R^3 and R^4 each independently is hydrogen or methyl, X is $-CR^bR^c-$, and Y is $-O-$.

In certain embodiments of either of formula I or formula II, n is 1, p is 2, R^3 and R^4 each independently is hydrogen or methyl, X is $-CR^bR^c-$, Y is $-O-$, and R^5 and R^6 are hydrogen.

In certain embodiments of either of formula I or formula II, n is 1, p is 2, R^3 and R^4 each independently is hydrogen or methyl, X is $-CR^bR^c-$, Y is $-O-$, and one of R^7 and R^8 and one of R^5 and R^6 together with the atoms to which they are attached form a four to seven-membered ring that may optionally be substituted with alkyl.

In certain embodiments of either of formula I or formula II, n is 1, p is 2, R^3 and R^4 each independently is hydrogen or methyl, X is $-CR^bR^c-$, and Y is $-NR^d-$.

In certain embodiments of either of formula I or formula II, n is 1, p is 2, R^3 and R^4 each independently is hydrogen or methyl, X is $-CR^bR^c-$, Y is $-NR^d-$, and R^5 and R^6 are hydrogen.

In certain embodiments of either of formula I or formula II, n is 1, p is 2, R^3 and R^4 each independently is hydrogen or methyl, X is $-CR^bR^c-$, Y is $-NR^d-$, and one of R^7 and R^8 and one of R^5 and R^6 together with the atoms to which they are attached form a four to seven-membered ring that may optionally be substituted with alkyl.

In certain embodiments of either of formula I or formula II, n is 1, p is 2, R^3 and R^4 each independently is hydrogen or methyl, X is $-CR^bR^c-$, Y is $-NR^d-$, and one of R^7 and R^8 together with R^d and the atoms to which they are attached form a four to seven-membered ring that may optionally be substituted with alkyl.

In certain embodiments of either of formula I or formula II, n is 1, p is 2, R^3 and R^4 each independently is hydrogen or methyl, X is $-CR^bR^c-$, and Y is $-CR^eR^f-$.

In certain embodiments of either of formula I or formula II, n is 1, p is 2, R^3 and R^4 each independently is hydrogen or methyl, X is $-CR^bR^c-$, Y is $-CR^eR^f-$, and R^5 , R^6 , R^e and R^f are hydrogen.

In certain embodiments of either of formula I or formula II, n is 1, p is 2, R^3 and R^4 each independently is hydrogen or methyl, X is $-CR^bR^c-$, Y is $-CR^eR^f-$, and one of R^7 and R^8 and one of R^5 and R^6 together with the atoms to which they are attached form a four to seven-membered ring that may optionally be substituted with alkyl.

5 In certain embodiments of either of formula I or formula II, n is 1, p is 2, R^3 and R^4 each independently is hydrogen or methyl, X is $-CR^bR^c-$, Y is $-CR^eR^f-$, and one of R^7 and R^8 and one of R^e and R^f together with the atoms to which they are attached form a four to seven-membered ring that may optionally be substituted with alkyl.

In certain embodiments of either of formula I or formula II, n is 1, p is 3, R^3 and R^4 each independently is hydrogen or methyl, and X is $-CR^bR^c-$.

In certain embodiments of either of formula I or formula II, n is 1, p is 3, R^3 and R^4 each independently is hydrogen or methyl, X is $-CR^bR^c-$, and Y is $-O-$.

In certain embodiments of either of formula I or formula II, n is 1, p is 3, R^3 and R^4 each independently is hydrogen or methyl, X is $-CR^bR^c-$, Y is $-O-$, and R^5 and R^6 are
15 hydrogen.

In certain embodiments of either of formula I or formula II, n is 1, p is 3, R^3 and R^4 each independently is hydrogen or methyl, X is $-CR^bR^c-$, Y is $-O-$, and one of R^7 and R^8 and one of R^5 and R^6 together with the atoms to which they are attached form a four to seven-membered ring that may optionally be substituted with alkyl.

20 In certain embodiments of either of formula I or formula II, n is 1, p is 3, R^3 and R^4 each independently is hydrogen or methyl, X is $-CR^bR^c-$, and Y is $-NR^d-$.

In certain embodiments of either of formula I or formula II, n is 1, p is 3, R^3 and R^4 each independently is hydrogen or methyl, X is $-CR^bR^c-$, Y is $-NR^d-$, and R^5 and R^6 are hydrogen.

25 In certain embodiments of either of formula I or formula II, n is 1, p is 3, R^3 and R^4 each independently is hydrogen or methyl, X is $-CR^bR^c-$, Y is $-NR^a-$, and one of R^7 and R^8 and one of R^5 and R^6 together with the atoms to which they are attached form a four to seven-membered ring.

In certain embodiments of either of formula I or formula II, n is that may optionally be substituted with alkyl, p is 3, R^3 and R^4 each independently is hydrogen or
30 methyl, X is $-CR^bR^c-$, Y is $-NR^d-$, and one of R^7 and R^8 together with R^d and the atoms to which they are attached form a four to seven-membered ring that may optionally be substituted with alkyl.

In certain embodiments of either of formula I or formula II, n is 1, p is 3, R^3 and
35 R^4 each independently is hydrogen or methyl, X is $-CR^bR^c-$, and Y is $-CR^eR^f-$.

In certain embodiments of either of formula I or formula II, n is 1, p is 3, R³ and R⁴ each independently is hydrogen or methyl, X is -CR^bR^c-, Y is -CR^eR^f-, and R⁵, R⁶, R^e and R^f are hydrogen.

In certain embodiments of either of formula I or formula II, n is 1, p is 3, R³ and R⁴ each independently is hydrogen or methyl, X is -CR^bR^c-, Y is -CR^eR^f-, and one of R⁷ and R⁸ and one of R⁵ and R⁶ together with the atoms to which they are attached form a four to seven-membered ring that may optionally be substituted with alkyl.

In certain embodiments of either of formula I or formula II, n is 1, p is 3, R³ and R⁴ each independently is hydrogen or methyl, X is -CR^bR^c-, Y is -CR^eR^f-, and one of R⁷ and R⁸ and one of R^e and R^f together with the atoms to which they are attached form a four to seven-membered ring that may optionally be substituted with alkyl.

In certain embodiments of either of formula I or formula II, n is 1, p is 2, R³ and R⁴ each independently is hydrogen or methyl, and X is -O-.

In certain embodiments of either of formula I or formula II, n is 1, p is 2, R³ and R⁴ each independently is hydrogen or methyl, X is -O-, and Y is -O-.

In certain embodiments of either of formula I or formula II, n is 1, p is 2, R³ and R⁴ each independently is hydrogen or methyl, X is -O-, Y is -O-, and R⁵ and R⁶ are hydrogen.

In certain embodiments of either of formula I or formula II, n is 1, p is 2, R³ and R⁴ each independently is hydrogen or methyl, X is -O-, Y is -O-, and one of R⁷ and R⁸ and one of R⁵ and R⁶ together with the atoms to which they are attached form a four to seven-membered ring that may optionally be substituted with alkyl.

In certain embodiments of either of formula I or formula II, n is 1, p is 2, R³ and R⁴ each independently is hydrogen or methyl, X is -O-, and Y is -NR^d-.

In certain embodiments of either of formula I or formula II, n is 1, p is 2, R³ and R⁴ each independently is hydrogen or methyl, X is -O-, Y is -NR^d-, and R⁵ and R⁶ are hydrogen.

In certain embodiments of either of formula I or formula II, n is 1, p is 2, R³ and R⁴ each independently is hydrogen or methyl, X is -O-, Y is -NR^d-, and one of R⁷ and R⁸ and one of R⁵ and R⁶ together with the atoms to which they are attached form a four to seven-membered ring that may optionally be substituted with alkyl.

In certain embodiments of either of formula I or formula II, n is 1, p is 2, R³ and R⁴ each independently is hydrogen or methyl, X is -O-, Y is -NR^d-, and one of R⁷ and R⁸ together with R^d and the atoms to which they are attached form a four to seven-membered ring that may optionally be substituted with alkyl.

In certain embodiments of either of formula I or formula II, n is 1, p is 2, R³ and R⁴ each independently is hydrogen or methylhydrogen, X is -O-, and Y is -CR^eR^f-.

In certain embodiments of either of formula I or formula II, n is 1, p is 2, R^3 and R^4 each independently is hydrogen or methyl, X is -O-, Y is $-CR^eR^f$ -, and R^5 , R^6 , R^e and R^f are hydrogen.

In certain embodiments of either of formula I or formula II, n is 1, p is 2, R^3 and R^4 each independently is hydrogen or methyl, X is -O-, Y is $-CR^eR^f$ -, and one of R^7 and R^8 and one of R^5 and R^6 together with the atoms to which they are attached form a four to seven-membered ring that may optionally be substituted with alkyl.

In certain embodiments of either of formula I or formula II, n is 1, p is 2, R^3 and R^4 each independently is hydrogen or methyl, X is -O-, Y is $-CR^eR^f$ -, and one of R^7 and R^8 and one of R^e and R^f together with the atoms to which they are attached form a four to seven-membered ring that may optionally be substituted with alkyl.

In certain embodiments of either of formula I or formula II, n is 1, p is 3, R^3 and R^4 each independently is hydrogen or methyl, and X is -O-.

In certain embodiments of either of formula I or formula II, n is 1, p is 3, R^3 and R^4 each independently is hydrogen or methyl, X is -O-, and Y is -O-.

In certain embodiments of either of formula I or formula II, n is 1, p is 3, R^3 and R^4 each independently is hydrogen or methyl, X is -O-, Y is -O-, and R^5 and R^6 are hydrogen.

In certain embodiments of either of formula I or formula II, n is 1, p is 3, R^3 and R^4 each independently is hydrogen or methyl, X is -O-, Y is -O-, and one of R^7 and R^8 and one of R^5 and R^6 together with the atoms to which they are attached form a four to seven-membered ring that may optionally be substituted with alkyl.

In certain embodiments of either of formula I or formula II, n is 1, p is 3, R^3 and R^4 each independently is hydrogen or methyl, X is -O-, and Y is $-NR^d$ -.

In certain embodiments of either of formula I or formula II, n is 1, p is 3, R^3 and R^4 each independently is hydrogen or methyl, X is -O-, Y is $-NR^d$ -, and R^5 and R^6 are hydrogen.

In certain embodiments of either of formula I or formula II, n is 1, p is 3, R^3 and R^4 each independently is hydrogen or methyl, X is -O-, Y is $-NR^a$ -, and one of R^7 and R^8 and one of R^5 and R^6 together with the atoms to which they are attached form a four to seven-membered ring that may optionally be substituted with alkyl.

In certain embodiments of either of formula I or formula II, n is 1, p is 3, R^3 and R^4 each independently is hydrogen or methyl, X is -O-, Y is $-NR^d$ -, and one of R^7 and R^8 together with R^d and the atoms to which they are attached form a four to seven-membered ring that may optionally be substituted with alkyl.

In certain embodiments of either of formula I or formula II, n is 1, p is 3, R^3 and R^4 each independently is hydrogen or methyl, X is -O-, and Y is $-CR^eR^f$ -.

In certain embodiments of either of formula I or formula II, n is 1, p is 3, R³ and R⁴ each independently is hydrogen or methyl, X is -O-, Y is -CR^eR^f-, and R⁵, R⁶, R^e and R^f are hydrogen.

In certain embodiments of either of formula I or formula II, n is 1, p is 3, R³ and R⁴ each independently is hydrogen or methyl, X is -O-, Y is -CR^eR^f-, and one of R⁷ and R⁸ and one of R⁵ and R⁶ together with the atoms to which they are attached form a four to seven-membered ring that may optionally be substituted with alkyl.

In certain embodiments of either of formula I or formula II, n is 1, p is 3, R³ and R⁴ each independently is hydrogen or methyl, X is -O-, Y is -CR^eR^f-, and one of R⁷ and R⁸ and one of R^e and R^f together with the atoms to which they are attached form a four to seven-membered ring that may optionally be substituted with alkyl.

In certain embodiments of either of formula I or formula II, n is 1, p is 2, R³ and R⁴ each independently is hydrogen or methyl, and X is -C(O)-.

In certain embodiments of either of formula I or formula II, n is 1, p is 2, R³ and R⁴ each independently is hydrogen or methyl, X is -C(O)-, and Y is -O-.

In certain embodiments of either of formula I or formula II, n is 1, p is 2, R³ and R⁴ each independently is hydrogen or methyl, X is -C(O)-, Y is -O-, and R⁵ and R⁶ are hydrogen.

In certain embodiments of either of formula I or formula II, n is 1, p is 2, R³ and R⁴ each independently is hydrogen or methyl, X is -C(O)-, Y is -O-, and one of R⁷ and R⁸ and one of R⁵ and R⁶ together with the atoms to which they are attached form a four to seven-membered ring that may optionally be substituted with alkyl.

In certain embodiments of either of formula I or formula II, n is 1, p is 2, R³ and R⁴ each independently is hydrogen or methyl, X is -C(O)-, and Y is -NR^d-.

In certain embodiments of either of formula I or formula II, n is 1, p is 2, R³ and R⁴ each independently is hydrogen or methyl, X is -C(O)-, Y is -NR^d-, and R⁵ and R⁶ are hydrogen.

In certain embodiments of either of formula I or formula II, n is 1, p is 2, R³ and R⁴ each independently is hydrogen or methyl, X is -C(O)-, Y is -NR^d-, and one of R⁷ and R⁸ and one of R⁵ and R⁶ together with the atoms to which they are attached form a four to seven-membered ring that may optionally be substituted with alkyl.

In certain embodiments of either of formula I or formula II, n is 1, p is 2, R³ and R⁴ each independently is hydrogen or methyl, X is -C(O)-, Y is -NR^d-, and one of R⁷ and R⁸ together with R^d and the atoms to which they are attached form a four to seven-membered ring that may optionally be substituted with alkyl.

In certain embodiments of either of formula I or formula II, n is 1, p is 2, R³ and R⁴ each independently is hydrogen or methyl, X is -C(O)-, and Y is -CR^eR^f-.

In certain embodiments of either of formula I or formula II, n is 1, p is 2, R³ and R⁴ each independently is hydrogen or methyl, X is -C(O)-, Y is -CR^eR^f-, and R⁵, R⁶ R^e and R^f are hydrogen.

In certain embodiments of either of formula I or formula II, n is 1, p is 2, R³ and R⁴ each independently is hydrogen or methyl, X is -C(O)-, Y is -CR^eR^f-, and one of R⁷ and R⁸ and one of R⁵ and R⁶ together with the atoms to which they are attached form a four to seven-membered ring that may optionally be substituted with alkyl.

In certain embodiments of either of formula I or formula II, n is 1, p is 2, R³ and R⁴ each independently is hydrogen or methyl, X is -C(O)-, Y is -CR^eR^f-, and one of R⁷ and R⁸ and one of R^e and R^f together with the atoms to which they are attached form a four to seven-membered ring that may optionally be substituted with alkyl.

In certain embodiments of either of formula I or formula II, n is 1, p is 3, R³ and R⁴ each independently is hydrogen or methyl, and X is -C(O)-.

In certain embodiments of either of formula I or formula II, n is 1, p is 3, R³ and R⁴ each independently is hydrogen or methyl, X is -C(O)-, and Y is -O-.

In certain embodiments of either of formula I or formula II, n is 1, p is 3, R³ and R⁴ each independently is hydrogen or methyl, X is -C(O)-, Y is -O-, and R⁵ and R⁶ are hydrogen.

In certain embodiments of either of formula I or formula II, n is 1, p is 3, R³ and R⁴ each independently is hydrogen or methyl, X is -C(O)-, Y is -O-, and one of R⁷ and R⁸ and one of R⁵ and R⁶ together with the atoms to which they are attached form a four to seven-membered ring that may optionally be substituted with alkyl.

In certain embodiments of either of formula I or formula II, n is 1, p is 3, R³ and R⁴ each independently is hydrogen or methyl, X is -C(O)-, and Y is -NR^d-.

In certain embodiments of either of formula I or formula II, n is 1, p is 3, R³ and R⁴ each independently is hydrogen or methyl, X is -C(O)-, Y is -NR^d-, and R⁵ and R⁶ are hydrogen.

In certain embodiments of either of formula I or formula II, n is 1, p is 3, R³ and R⁴ each independently is hydrogen or methyl, X is -C(O)-, Y is -NR^a-, and one of R⁷ and R⁸ and one of R⁵ and R⁶ together with the atoms to which they are attached form a four to seven-membered ring that may optionally be substituted with alkyl.

In certain embodiments of either of formula I or formula II, n is 1, p is 3, R³ and R⁴ each independently is hydrogen or methyl, X is -C(O)-, Y is -NR^d-, and one of R⁷ and R⁸ together with R^d and the atoms to which they are attached form a four to seven-membered ring that may optionally be substituted with alkyl.

In certain embodiments of either of formula I or formula II, n is 1, p is 3, R³ and R⁴ each independently is hydrogen or methyl, X is -C(O)-, and Y is -CR^eR^f-.

In certain embodiments of either of formula I or formula II, n is 1, p is 3, R³ and R⁴ each independently is hydrogen or methyl, X is -C(O)-, Y is -CR^eR^f-, and R⁵, R⁶, R^e and R^f are hydrogen.


5 In certain embodiments of either of formula I or formula II, n is 1, p is 3, R³ and R⁴ each independently is hydrogen or methyl, X is -C(O)-, Y is -CR^eR^f-, and one of R⁷ and R⁸ and one of R⁵ and R⁶ together with the atoms to which they are attached form a four to seven-membered ring that may optionally be substituted with alkyl.

10 In certain embodiments of either of formula I or formula II, n is 1, p is 3, R³ and R⁴ each independently is hydrogen or methyl, X is -C(O)-, Y is -CR^eR^f-, and one of R⁷ and R⁸ and one of R^e and R^f together with the atoms to which they are attached form a four to seven-membered ring that may optionally be substituted with alkyl.

In certain embodiments of formula I, R¹ is azetidiny, pyrrolidiny, piperaziny, piperidiny, azepiny, diazetidiny, diazepiny, aminoalkoxy, azetidinyloxy, pyrrolidinyloxy, piperidinyloxy, azepinyloxy, azetidinylmethoxy, pyrrolidinylmethoxy, piperidinylmethoxy, azepinylmethoxy, or aminoalkyl.

In certain embodiments of formula I, R¹ is piperaziny, piperidiny or pyrrolidiny.

In certain embodiments of formula I, R¹ is a group of the formula:




•

wherein R⁷, R⁸, R¹⁰, R¹¹, R¹² and R¹³ each independently is hydrogen or alkyl.

Preferably R¹⁰, R¹¹, R¹² and R¹³ are hydrogen in such embodiments.

5 In certain embodiments the compounds of the invention may be represented by
formula III:



wherein:

Y is N or CH;

t is 1 or 2;

u is from 1 to 3;

R^{10} , R^{11} , R^{12} and R^{13} each independently is hydrogen or alkyl; and

m, s, X, R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 and R^9 are as defined herein.

5 In certain embodiments of formula III, X is $-NR^a$.

In certain embodiments of formula III, X is $-O$.

In certain embodiments of formula III, X is $-CR^bR^c$.

In certain embodiments of formula III, X is $-C(O)$;

In certain embodiments of formula III, Y is $-NR^d$.

10 In certain embodiments of formula III, Y is $-CR^f$.

In certain embodiments of formula III, m is 0 or 1 and s is 0, 1 or 2.

In certain embodiments of formula III, m is 0 or 1 and R^2 is halo, preferably fluoro or chloro.

In certain embodiments of formula III, m is 0.

15 In certain embodiments of formula III, m is 1 and R^2 is fluoro or chloro.

In certain embodiments of formula III, m is 2 and each R^2 is independently fluoro or chloro at the 5 and six position of the benzimidazolone or dihydroindolone ring system.

In certain embodiments of formula III, R^3 and R^4 are hydrogen.

20 In certain embodiments of formula III, R^3 and R^4 are alkyl, preferably methyl.

In certain embodiments of formula III, one of R^3 and R^4 is hydrogen and the other is alkyl, preferably methyl.

In certain embodiments of formula III where X is $-NR^a$, R^a is hydrogen. In other embodiments of formula III where X is $-NR^a$, R^a is alkyl, preferably methyl.

25 In certain embodiments of formula III where X is $-CR^bR^c$, R^b and R^c are hydrogen.

In other embodiments of formula III where X is $-CR^bR^c$, R^b and R^c are alkyl, preferably methyl.

30 In still other embodiments of formula III where X is $-CR^bR^c$, one of R^b and R^c is hydrogen and the other is alkyl, preferably methyl.

In yet other embodiments of formula III where X is $-CR^bR^c$, one of R^b and R^c is alkyl, preferably methyl, and the other is hydroxy.

35 In other embodiments of formula III where X is $-CR^bR^c$, R^b and R^c together with the atom to which they are attached form a three to six-membered ring that optionally includes a heteroatom selected from O, N and S. Preferably in such embodiments R^b and R^c together with the atom to which they are attached form a three or four membered carbocyclic ring.

In certain embodiments of formula III, Y is -CH-.

In certain embodiments of formula III, Y is -N-.

In certain embodiments of formula III, Y is -CH-, u is 2 and t is 2.

In certain embodiments of formula III, Y is -CH-, u is 1 and t is 2.

5 In certain embodiments of formula III, Y is -N-, u is 2 and t is 2.


In certain embodiments of formula III, Y is -N-, u is 2 and t is 3.

In certain embodiments of formula III, R⁸ is hydrogen.

In certain embodiments of formula III, R¹⁰, R¹¹, R¹² and R¹³ are hydrogen.

In certain embodiments of formula III the subject compounds may be

10 represented by formula IIIa:



wherein Y is N or CH and m, s, t, u, Y, R², R³, R⁴, R⁸, R⁹, R¹⁰, R¹¹, R¹², R¹³ and R^a are as defined herein.

In certain embodiments of formula IIIa, R^a is hydrogen.

15 In certain embodiments of formula IIIa, R⁸ is hydrogen.

In certain embodiments of formula IIIa, Y is -CH-.

In certain embodiments of formula IIIa, Y is -N-.

In certain embodiments of formula IIIa, t is 1 and u is 2.

In certain embodiments of formula IIIa, t is 2 and u is 2.

20 In certain embodiments of formula IIIa, t is 2 and u is 3.

In certain embodiments of formula IIIa, m is 0 or 1 and s is 0, 1 or 2.

In certain embodiments of formula IIIa, R⁸ is hydrogen.

In certain embodiments of formula IIIa, R¹⁰, R¹¹, R¹² and R¹³ are hydrogen.

25 In certain embodiments of formula IIIa, m is 0 or 1 and R² is halo, preferably fluoro or chloro.

In certain embodiments of formula IIIa, m is 0.

In certain embodiments of formula IIIa, m is 1 and R² is fluoro or chloro.


In certain embodiments of formula IIIa, m is 2 and each R² is independently fluoro or chloro at the 5 and six position of the benzimidazolone ring system.

In certain embodiments of formula IIIa, R³ and R⁴ are hydrogen.

In certain embodiments of formula IIIa, R³ and R⁴ are alkyl, preferably methyl.

In certain embodiments of formula IIIa, one of R³ and R⁴ is hydrogen and the other is alkyl, preferably methyl.

5 In other embodiments of formula III the compounds of the invention may be represented by formula IIIb:



wherein Y is N or CH and m, s, t, u, Y, R², R³, R⁴, R⁸, R⁹, R¹⁰, R¹¹, R¹², R¹³, R^b and R^c are as defined herein.

10 In certain embodiments of formula IIIb, R^b and R^c are hydrogen.

In certain embodiments of formula IIIb, R^b and R^c are alkyl, preferably methyl.

In certain embodiments of formula IIIb, one of R^b and R^c is hydrogen and the other is alkyl, preferably methyl.

In certain embodiments of formula IIIb, one of R^b and R^c is hydrogen or alkyl and
15 the other is hydroxyl.

In certain embodiments of formula IIIb, R^b and R^c together form oxo.

In certain embodiments of formula IIIb, Y is -CH-.

In certain embodiments of formula IIIb, Y is -N-.

In certain embodiments of formula IIIb, t is 1 and u is 2.

20 In certain embodiments of formula IIIb, t is 2 and u is 2.

In certain embodiments of formula IIIb, t is 2 and u is 3.

In certain embodiments of formula IIIb, m is 0 or 1 and s is 0, 1 or 2.

In certain embodiments of formula IIIb, R⁸ is hydrogen.

In certain embodiments of formula IIIb, R¹⁰, R¹¹, R¹² and R¹³ are hydrogen.

25 In certain embodiments of formula IIIb, m is 0 or 1 and R² is halo, preferably fluoro or chloro.

In certain embodiments of formula IIIb, m is 0.

In certain embodiments of formula IIIb, m is 1 and R² is fluoro or chloro.


In certain embodiments of formula IIIb, m is 2 and each R^2 is independently fluoro or chloro at the 5 and six position of the dihydroindolone ring system.

In certain embodiments of formula IIIb, R^3 and R^4 are hydrogen.

In certain embodiments of formula IIIb, R^3 and R^4 are alkyl, preferably methyl.

5 In certain embodiments of formula IIIb, one of R^3 and R^4 is hydrogen and the other is alkyl, preferably methyl.

In certain embodiments the compounds of the invention may be represented by formula IV:



10 wherein:

p is 2 or 3; and

m, s, X, Y, R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 and R^9 are as defined herein.

In certain embodiments of formula IV, X is $-NR^a$.

In certain embodiments of formula IV, X is $-O$.

15 In certain embodiments of formula IV, X is $-CR^bR^c$.

In certain embodiments of formula IV, X is $-C(O)$;

In certain embodiments of formula IV, Y is $-NR^d$.

In certain embodiments of formula IV, Y is $-CR^eR^f$.

In certain embodiments of formula IV, m is 0 or 1 and s is 0, 1 or 2.

20 In certain embodiments of formula IV where X is $-NR^a$, R^a is hydrogen. In other embodiments of formula III where X is $-NR^a$, R^a is alkyl, preferably methyl.

In certain embodiments of formula IV where X is $-CR^bR^c$, R^b and R^c are hydrogen.

In other embodiments of formula IV where X is $-CR^bR^c$, R^b and R^c are alkyl, preferably methyl.

In still other embodiments of formula IV where X is $-CR^bR^c$, one of R^b and R^c is hydrogen and the other is alkyl, preferably methyl.

In yet other embodiments of formula IV where X is $-CR^bR^c$, one of R^b and R^c is alkyl, preferably methyl, and the other is hydroxy.

In other embodiments of formula IV where X is $-CR^bR^c-$, R^b and R^c together with the atom to which they are attached form a three to six-membered ring that optionally includes a heteroatom selected from O, N and S. Preferably in such embodiments R^b and R^c together with the atom to which they are attached form a three or four membered

5 carbocyclic ring.

In certain embodiments of formula IV, Y is $-CH_2-$.

In certain embodiments of formula IV, Y is $-NH-$.

In certain embodiments of formula IV, Y is $-O-$.

In certain embodiments of formula IV, p is 2.

10 In certain embodiments of formula IV, p is 3.

In certain embodiments of formula IV, Y is O, p is 2, and one of R^7 and R^8 is hydrogen and the other is alkyl, preferably methyl.

In certain embodiments of formula IV, m is 0 or 1 and R^2 is halo, preferably fluoro or chloro.

15 In certain embodiments of formula IV, m is 0.

In certain embodiments of formula IV, m is 1 and R^2 is fluoro or chloro.


In certain embodiments of formula IV, m is 2 and each R^2 is independently fluoro or chloro at the 5 and six position of the benzimidazolone or dihydroindolone ring system.

20 In certain embodiments of formula IV, R^3 and R^4 are hydrogen.

In certain embodiments of formula IV, R^3 and R^4 are alkyl, preferably methyl.

In certain embodiments of formula IV, one of R^3 and R^4 is hydrogen and the other is alkyl, preferably methyl.

In certain embodiments of formula IV the subject compounds may be
25 represented by formula IVa:



wherein p is 2 or 3 and m, s, p, R^2 , R^3 , R^4 , R^7 , R^8 , R^9 and R^a are as defined herein.

In certain embodiments of formula IVa, R^a is hydrogen.

In certain embodiments of formula IVa, one of R^7 and R^8 is hydrogen and the
30 other is alkyl, preferably methyl.

In certain embodiments of formula IVa, m is 0 or 1 and R^2 is halo, preferably fluoro or chloro.

In certain embodiments of formula IVa, m is 0.

In certain embodiments of formula IVa, m is 1 and R^2 is fluoro or chloro.


5 In certain embodiments of formula IVa, m is 2 and each R^2 is independently fluoro or chloro at the 5 and six position of the benzimidazolone ring system.

In certain embodiments of formula IVa, R^3 and R^4 are hydrogen.

In certain embodiments of formula IVa, R^3 and R^4 are alkyl, preferably methyl.

10 In certain embodiments of formula IVa, one of R^3 and R^4 is hydrogen and the other is alkyl, preferably methyl.

In other embodiments of formula IV the subject compounds may be represented by formul IVb:



wherein p is 2 or 3 and m, s, p, R^2 , R^3 , R^4 , R^7 , R^8 , R^9 and R^a are as defined herein.

15 In certain embodiments of formula IVb, R^b and R^c are hydrogen.

In certain embodiments of formula IVb, R^b and R^c together form oxo.

In certain embodiments of formula IVb, one of R^7 and R^8 is hydrogen and the other is alkyl, preferably methyl.

20 In certain embodiments of formula IVb, m is 0 or 1 and R^2 is halo, preferably fluoro or chloro.

In certain embodiments of formula IVb, m is 0.

In certain embodiments of formula IVb, m is 1 and R^2 is fluoro or chloro.


In certain embodiments of formula IVb, m is 2 and each R^2 is independently fluoro or chloro at the 5 and six position of the dihydroindolone ring system.

25 In certain embodiments of formula IVb, R^3 and R^4 are hydrogen.

In certain embodiments of formula IVb, R^3 and R^4 are alkyl, preferably methyl.

In certain embodiments of formula IVb, one of R^3 and R^4 is hydrogen and the other is alkyl, preferably methyl.

30 In certain embodiments the compounds of the invention may be represented by formula V:



wherein m, s, t, u, X, Y, R², R³, R⁴, R⁸, R⁹, R¹⁰, R¹¹, R¹² and R¹³ are as defined herein.

In certain embodiments of formula V, X is -NR^a-.

5 In certain embodiments of formula V, X is -O-.

In certain embodiments of formula V, X is -CR^bR^c-.

In certain embodiments of formula V, X is -C(O)-;

In certain embodiments of formula V, Y is -NR^d-.

In certain embodiments of formula V, Y is -CR^eR^f-.

10 In certain embodiments of formula V, Y is -O-.

In certain embodiments of formula V, R¹⁰, R¹¹, R¹² and R¹³ are hydrogen.

In certain embodiments of formula V, m is 0 or 1 and s is 0, 1 or 2.

In certain embodiments of formula V, R³ and R⁴ are hydrogen.

In certain embodiments of formula V, R³ and R⁴ are alkyl, preferably methyl.

15 In certain embodiments of formula V, one of R³ and R⁴ is hydrogen and the other is alkyl, preferably methyl.

In certain embodiments of formula V where X is -NR^a-, R^a is hydrogen. In other embodiments of formula III where X is -NR^a-, R^a is alkyl, preferably methyl.

20 In certain embodiments of formula V where X is -CR^bR^c-, R^b and R^c are hydrogen.

In other embodiments of formula V where X is -CR^bR^c-, R^b and R^c are alkyl, preferably methyl.

In still other embodiments of formula V where X is -CR^bR^c-, one of R^b and R^c is hydrogen and the other is alkyl, preferably methyl.

25 In yet other embodiments of formula V where X is -CR^bR^c-, one of R^b and R^c is alkyl, preferably methyl, and the other is hydroxy.

In other embodiments of formula V where X is -CR^bR^c-, R^b and R^c together with the atom to which they are attached form a three to six-membered ring that optionally includes a heteroatom selected from O, N and S. Preferably in such embodiments R^b and

R^c together with the atom to which they are attached form a three or four membered carbocyclic ring.

In certain embodiments of formula V, R^8 is hydrogen.

In certain embodiments of formula V, Y is -O-, t is 2 and tu is 2.

5 In certain embodiments of formula V, Y is -O-, t is 1 and u is 2.

In certain embodiments of formula V, Y is -O-, t is 1 and u is 1.

In certain embodiments of formula V, Y is -O-, t is 2 and u is 3.


In certain embodiments of formula V, m is 0 or 1 and R^2 is halo, preferably fluoro or chloro.

10 In certain embodiments of formula V, m is 0.

In certain embodiments of formula V, m is 1 and R^2 is fluoro or chloro.

In certain embodiments of formula V, m is 2 and each R^2 is independently fluoro or chloro at the 5 and six position of the benzimidazolone or dihydroindolone ring system.

15 In certain embodiments of formula V the compounds of the invention may be represented by formula Va:



wherein m, s, t, u, X, Y, R^2 , R^3 , R^4 , R^8 , R^9 , R^{10} , R^{11} , R^{12} , R^{13} and R^a are as defined herein.

20 In certain embodiments of formula Va, R^a is hydrogen.

In certain embodiments of formula Va, R^8 is hydrogen.

In certain embodiments of formula Va, R^{10} , R^{11} , R^{12} and R^{13} are hydrogen.

In certain embodiments of formula Va, m is 0 or 1 and s is 0, 1 or 2.

In certain embodiments of formula Va, t is 1 and u is 1.

25 In certain embodiments of formula Va, t is 1 and u is 2.

In certain embodiments of formula Va, t is 2 and u is 2.

In certain embodiments of formula Va, t is 2 and u is 3.

In certain embodiments of formula Va, m is 0 or 1 and R^2 is halo, preferably fluoro or chloro.

In certain embodiments of formula Va, m is 0.

In certain embodiments of formula Va, m is 1 and R² is fluoro or chloro.


In certain embodiments of formula Va, m is 2 and each R² is independently fluoro or chloro at the 5 and six position of the benzimidazolone ring system.

5 In certain embodiments of formula Va, R³ and R⁴ are hydrogen.

In certain embodiments of formula Va, R³ and R⁴ are alkyl, preferably methyl.

In certain embodiments of formula Va, one of R³ and R⁴ is hydrogen and the other is alkyl, preferably methyl.

In certain embodiments of formula V the compounds of the invention may be
10 represented by formula Vb:



wherein m, s, t, u, X, Y, R², R³, R⁴, R⁸, R⁹, R¹⁰, R¹¹, R¹², R¹³, R^b and R^c are as defined herein.

In certain embodiments of formula Vb, R^b and R^c are hydrogen.

15 In certain embodiments of formula Vb, R^b and R^c together form oxo.

In certain embodiments of formula Vb, R⁸ is hydrogen.

In certain embodiments of formula Vb, R¹⁰, R¹¹, R¹² and R¹³ are hydrogen.

In certain embodiments of formula Vb, m is 0 or 1 and s is 0, 1 or 2.

In certain embodiments of formula Vb, t is 1 and u is 1.

20 In certain embodiments of formula Vb, t is 1 and u is 2.

In certain embodiments of formula Vb, t is 2 and u is 2.

In certain embodiments of formula Vb, t is 2 and u is 3.

In certain embodiments of formula Vb, m is 0 or 1 and R² is halo, preferably fluoro or chloro.

25 In certain embodiments of formula Vb, m is 0.

In certain embodiments of formula Vb, m is 1 and R² is fluoro or chloro.


In certain embodiments of formula Vb, m is 2 and each R² is independently fluoro or chloro at the 5 and six position of the dihydroindolone ring system.

In certain embodiments of formula Vb, R³ and R⁴ are hydrogen.

In certain embodiments of formula Vb, R^3 and R^4 are alkyl, preferably methyl.

In certain embodiments of formula Vb, one of R^3 and R^4 is hydrogen and the other is alkyl, preferably methyl.

- 5 In certain embodiments the compounds of the invention may be represented by formula VI:



wherein

t is from 1 to 3;

- 10 u is from 0 to 3; and

m , s , X , Y , R^2 , R^3 , R^4 , R^8 , R^9 , R^{10} , R^{11} , R^{12} and R^{13} are as defined herein.

In certain embodiments of formula VI, X is $-NR^a-$.

In certain embodiments of formula VI, X is $-O-$.

In certain embodiments of formula VI, X is $-CR^bR^c-$.

- 15 In certain embodiments of formula VI, X is $-C(O)-$;

In certain embodiments of formula VI, Y is $-NR^d-$.

In certain embodiments of formula VI, Y is $-CR^eR^f-$.

In certain embodiments of formula VI, Y is $-O-$.

In certain embodiments of formula VI, R^{10} , R^{11} , R^{12} and R^{13} are hydrogen.

- 20 In certain embodiments of formula VI, m is 0 or 1 and s is 0, 1 or 2.

In certain embodiments of formula VI, R^3 and R^4 are hydrogen.

In certain embodiments of formula VI, R^3 and R^4 are alkyl, preferably methyl.

In certain embodiments of formula VI, one of R^3 and R^4 is hydrogen and the other is alkyl, preferably methyl.

- 25 In certain embodiments of formula VI where X is $-NR^a-$, R^a is hydrogen. In other embodiments of formula III where X is $-NR^a-$, R^a is alkyl, preferably methyl.

In certain embodiments of formula VI where X is $-\text{CR}^b\text{R}^c-$, R^b and R^c are hydrogen.

In other embodiments of formula VI where X is $-\text{CR}^b\text{R}^c-$, R^b and R^c are alkyl, preferably methyl.

- 5 In still other embodiments of formula VI where X is $-\text{CR}^b\text{R}^c-$, one of R^b and R^c is hydrogen and the other is alkyl, preferably methyl.

In yet other embodiments of formula VI where X is $-\text{CR}^b\text{R}^c-$, one of R^b and R^c is alkyl, preferably methyl, and the other is hydroxy.

- 10 In other embodiments of formula VI where X is $-\text{CR}^b\text{R}^c-$, R^b and R^c together with the atom to which they are attached form a three to six-membered ring that optionally includes a heteroatom selected from O, N and S. Preferably in such embodiments R^b and R^c together with the atom to which they are attached form a three or four membered carbocyclic ring.

In certain embodiments of formula VI, R^8 is hydrogen.

- 15 In certain embodiments of formula VI, Y is $-\text{O}-$, t is 2 and tu is 2.

In certain embodiments of formula VI, Y is $-\text{O}-$, t is 1 and u is 2.

In certain embodiments of formula VI, Y is $-\text{O}-$, t is 1 and u is 1.

In certain embodiments of formula VI, Y is $-\text{O}-$, t is 2 and u is 3.

In certain embodiments of formula VI, Y is $-\text{O}-$, t is 2 and u is 0.


- 20 In certain embodiments of formula V, m is 0 or 1 and R^2 is halo, preferably fluoro or chloro.

In certain embodiments of formula V, m is 0.

In certain embodiments of formula V, m is 1 and R^2 is fluoro or chloro.

- 25 In certain embodiments of formula V, m is 2 and each R^2 is independently fluoro or chloro at the 5 and six position of the benzimidazolone or dihydroindolone ring system.

In certain embodiments of formula VI the compounds of the invention may be represented by formula VIa:



wherein m , s , t , u , X , $\forall$, R^2 , R^3 , R^4 , R^8 , R^9 , R^{10} , R^{11} , R^{12} , R^{13} and R^a are as defined herein.

In certain embodiments of formula VIa, R^a is hydrogen.

5 In certain embodiments of formula VIa, R^8 is hydrogen.

In certain embodiments of formula VIa, R^{10} , R^{11} , R^{12} and R^{13} are hydrogen.

In certain embodiments of formula VIa, m is 0 or 1 and s is 0, 1 or 2.

In certain embodiments of formula VIa, t is 1 and u is 1.

In certain embodiments of formula VIa, t is 1 and u is 2.

10 In certain embodiments of formula VIa, t is 2 and u is 2.

In certain embodiments of formula VIa, t is 2 and u is 3.

In certain embodiments of formula VIa, m is 0 or 1 and R^2 is halo, preferably fluoro or chloro.

In certain embodiments of formula VIa, m is 0.

15 In certain embodiments of formula VIa m is 1 and R^2 is fluoro or chloro.


In certain embodiments of formula VIa, m is 2 and each R^2 is independently fluoro or chloro at the 5 and six position of the benzimidazolone ring system.

In certain embodiments of formula VIa, R^3 and R^4 are hydrogen.

In certain embodiments of formula VIa, R^3 and R^4 are alkyl, preferably methyl.

20 In certain embodiments of formula VIa, one of R^3 and R^4 is hydrogen and the other is alkyl, preferably methyl.

In certain embodiments of formula VI the compounds of the invention may be represented by formula VIb:



wherein m, s, t, u, R², R³, R⁴, R⁸, R⁹, R¹⁰, R¹¹, R¹², R¹³, R^b and R^c are as defined herein.

In certain embodiments of formula VIb, R^b and R^c are hydrogen.

5 In certain embodiments of formula VIb, R^b and R^c together form oxo.

In certain embodiments of formula VIb, R¹⁰, R¹¹, R¹² and R¹³ are hydrogen.

In certain embodiments of formula VIa, R⁸ is hydrogen.

In certain embodiments of formula VIb, m is 0 or 1 and s is 0, 1 or 2.

In certain embodiments of formula VIb, t is 1 and u is 1.

10 In certain embodiments of formula VIb, t is 1 and u is 2.

In certain embodiments of formula VIb, t is 2 and u is 2.

In certain embodiments of formula VIb, t is 2 and u is 3.

In certain embodiments of formula VIb, m is 0 or 1 and R² is halo, preferably fluoro or chloro.

15 In certain embodiments of formula VIb, m is 0.

In certain embodiments of formula VIb m is 1 and R² is fluoro or chloro.


In certain embodiments of formula VIb, m is 2 and each R² is independently fluoro or chloro at the 5 and six position of the dihydroindolone ring system.

In certain embodiments of formula VIb, R³ and R⁴ are hydrogen.

20 In certain embodiments of formula VIb, R³ and R⁴ are alkyl, preferably methyl.

In certain embodiments of formula VIb, one of R³ and R⁴ is hydrogen and the other is alkyl, preferably methyl.

In certain embodiments the compounds of the invention may be represented by formula VII:



wherein:

Y is -N- or -CH- and;

m, s, X, R², R³, R⁴, R⁸ and R⁹ are as defined herein.

5 In certain embodiments of formula VII, X is NH.

In certain embodiments of formula VII, X is CH₂.

In certain embodiments of formula VII, X is C=O.

In certain embodiments of formula VII, m is 0 or 1 and R² is halo, preferably fluoro or chloro.

10 In certain embodiments of formula VII, m is 0.

In certain embodiments of formula VII m is 1 and R² is fluoro or chloro.

In certain embodiments of formula VII, m is 2 and each R² is independently fluoro or chloro at the 5 and six position of the benzimidazolone or dihydroindolone ring system.

15 In certain embodiments of formula VII, s is 0, 1 or 2 and R⁹ is halo.

In certain embodiments of formula VII, R³ and R⁴ are hydrogen.


In certain embodiments of formula VII, R³ and R⁴ are alkyl, preferably methyl.

In certain embodiments of formula VII, one of R³ and R⁴ is hydrogen and the other is alkyl, preferably methyl.

20 In certain embodiments of formula VII, Y is -N-.

In certain embodiments of formula VII, Y is -CH-.

In certain embodiments of formula VII, the subject compounds may be represented by formula VIIa:



wherein:

Y is -N- or -CH- and;

m, s, R², R³, R⁴, R⁸ and R⁹ are as defined herein.

5 In certain embodiments of formula VIIa, m is 0 or 1 and R² is halo, preferably fluoro or chloro.

In certain embodiments of formula VIIa, m is 0.

In certain embodiments of formula VIIa m is 1 and R² is fluoro or chloro.

10 In certain embodiments of formula VIIa, m is 2 and each R² is independently fluoro or chloro at the 5 and six position of the benzimidazolone or dihydroindolone ring system.

In certain embodiments of formula VIIa, s is 0, 1 or 2 and R⁹ is halo.

In certain embodiments of formula VIIa, R⁸ is hydrogen.

In certain embodiments of formula VIIa, R⁸ is methyl.

15 In certain embodiments of formula VIIa, R³ and R⁴ are hydrogen.


In certain embodiments of formula VIIa, R³ and R⁴ are alkyl, preferably methyl.

In certain embodiments of formula VIIa, one of R³ and R⁴ is hydrogen and the other is alkyl, preferably methyl.

In certain embodiments of formula VII, Y is -N-.

20 In certain embodiments of formula VII, Y is -CH-.

In certain embodiments of formula VII, the subject compounds may be represented by formula VIIb:



wherein:

Y is -N- or -CH- and;

m, s, R², R³, R⁴, R⁸ and R⁹ are as defined herein.

5 In certain embodiments of formula VIIb, m is 0 or 1 and R² is halo, preferably fluoro or chloro.

In certain embodiments of formula VIIb, m is 0.

In certain embodiments of formula VIIb m is 1 and R² is fluoro or chloro.

10 In certain embodiments of formula VIIb, m is 2 and each R² is independently fluoro or chloro at the 5 and six position of the benzimidazolone or dihydroindolone ring system.

In certain embodiments of formula VIIb, s is 0, 1 or 2 and R⁹ is halo.

In certain embodiments of formula VIIb, R⁸ is hydrogen.

In certain embodiments of formula VIIb, R⁸ is methyl.

15 In certain embodiments of formula VIIb, R³ and R⁴ are hydrogen.

In certain embodiments of formula VIIb, R³ and R⁴ are alkyl, preferably methyl.

In certain embodiments of formula VIIb, one of R³ and R⁴ is hydrogen and the other is alkyl, preferably methyl.





In certain embodiments of formula VII, Y is -N-.






20 In certain embodiments of formula VII, Y is -CH-.

Where any of R¹, R², R³, R⁴, R⁵, R⁶, R⁷, R⁸, R⁹, R¹⁰, R¹¹, R¹², R¹³, R^a, R^b, R^c, R^d, R^e, R^f, R^g, R^h, Rⁱ, R^j and R^k herein are alkyl or contain an alkyl moiety, such alkyl is preferably lower alkyl, i.e. C₁-C₆alkyl, and more preferably C₁-C₄alkyl.






25 Representative compounds in accordance with the invention are shown in Table 1 together with melting point or mass spectrum M+H, and the experimental examples (described below) associated with each compound. Melting points shown are for the corresponding hydrochloride salts unless indicated otherwise.






TABLE 1






#	Structure	Name (Autonom)	MP °C/ M+H
1		1-Benzyl-4-piperazin-1-yl- 1,3-dihydro- benzoimidazol-2-one	>300 °C
2		1-Benzyl-4-(2- methylamino-ethoxy)-1,3- dihydro-benzoimidazol-2- one	253.5-254.8 °C
3		1-Benzyl-3-methyl-4-(2- methylamino-ethoxy)-1,3- dihydro-benzoimidazol-2- one	203.8-205.1 °C
4		4-(Azetidin-3-ylmethoxy)- 1-benzyl-1,3-dihydro- benzoimidazol-2-one	210.0-216.6 °C






#	Structure	Name (Autonom)	MP °C/ M+H
5		1-Benzyl-4-(3-dimethylamino-propoxy)-1,3-dihydro-benzoimidazol-2-one	186.5-189.5 °C
6		1-Benzyl-4-(pyrrolidin-3-ylmethoxy)-1,3-dihydro-benzoimidazol-2-one	226.0-228.9 °C
7		1-Benzyl-4-(piperidin-4-yloxy)-1,3-dihydro-benzoimidazol-2-one	260.9-263.3 °C
8		4-(Azetidin-3-yloxy)-1-benzyl-1,3-dihydro-benzoimidazol-2-one	181.0-184.0 °C
9		(R)- 1-Benzyl-4-(pyrrolidin-2-ylmethoxy)-1,3-dihydro-benzoimidazol-2-one	239.5-241.8 °C






#	Structure	Name (Autonom)	MP °C/ M+H
10		1-Benzyl-4-piperazin-1-yl- 1,3-dihydro- benzoimidazol-2-one	284.6-285.9 °C
11		1-Benzyl-4-piperazin-1-yl- 1,3-dihydro-indol-2-one	241.0-242.3 °C
12		1-(3-Fluoro-benzyl)-4- piperazin-1-yl-1,3-dihydro- indol-2-one	>300 °C
13		1-(3-Fluoro-benzyl)-4- piperazin-1-yl-1H-indole- 2,3-dione	247.8-248.1 °C
14		4-[1,4]Diazepan-1-yl-1-(3- fluoro-benzyl)-1H-indole- 2,3-dione	354






#	Structure	Name (Autonom)	MP °C/ M+H
15		1-(3-Fluoro-benzyl)-3,3-dimethyl-4-piperazin-1-yl-1,3-dihydro-indol-2-one	354
16		1-(3-fluoro-benzyl)-4-(2-methylamino-ethoxy)-1,3-dihydro-indol-2-one	213.2-213.9 °C
17		1-(3-Fluoro-benzyl)-3-hydroxy-3-methyl-4-piperazin-1-yl-1,3-dihydro-indol-2-one	208.4-209.8 °C
18		4-[1,4]Diazepan-1-yl-1-(3-fluoro-benzyl)-1,3-dihydro-indol-2-one	178.3-181.5 °C
19		1-(3-Fluoro-benzyl)-4-piperidin-4-yl-1,3-dihydro-indol-2-one	221.1-223.7 °C





#	Structure	Name (Autonom)	MP °C/ M+H
20		1-Benzyl-6-chloro-4-(2-methylamino-ethoxy)-1,3-dihydro-benzoimidazol-2-one	188.0-190.1 °C (TFA salt)
21		1-Benzyl-6-fluoro-4-(2-methylamino-ethoxy)-1,3-dihydro-benzoimidazol-2-one	316
22		1-(2-Fluoro-benzyl)-4-(2-methylamino-ethoxy)-1,3-dihydro-benzoimidazol-2-one	251.7-252.5 °C
23		1-(2-Chloro-benzyl)-4-(2-methylamino-ethoxy)-1,3-dihydro-benzoimidazol-2-one	248.4-249.7 °C
24		1-(3-Fluoro-benzyl)-4-(2-methylamino-ethoxy)-1,3-dihydro-benzoimidazol-2-one	219.9-220.6 °C





#	Structure	Name (Autonom)	MP °C/ M+H
25		1-(3-Chloro-benzyl)-4-(2-methylamino-ethoxy)-1,3-dihydro-benzoimidazol-2-one	213.8-214.5 °C
26		1-(2,3-difluoro-benzyl)-4-(2-methylamino-ethoxy)-1,3-dihydro-benzoimidazol-2-one	'238.8-241.6 °C
27		1-Benzyl-4-(2-methylamino-ethyl)-1,3-dihydro-benzoimidazol-2-one	282
28		1-Benzyl-6-fluoro-4-piperazin-1-yl-1,3-dihydro-benzoimidazol-2-one	>300 °C
29		1-(3-Fluoro-benzyl)-6-fluoro-4-piperazin-1-yl-1,3-dihydro-benzoimidazol-2-one	>300 °C





#	Structure	Name (Autonom)	MP °C/ M+H
30		1-(4-Fluoro-benzyl)-6-fluoro-4-(2-methylamino-ethoxy)-1,3-dihydro-benzoimidazol-2-one	271.3-271.9 °C
31		1-(4-Fluoro-benzyl)-4-(2-methylamino-ethoxy)-1,3-dihydro-benzoimidazol-2-one	250.9-251.9 °C
32		1-Benzyl-4-pyrrolidin-3-yl-1,3-dihydro-benzoimidazol-2-one	256.9-257.3 °C
33		1-Benzyl-6-fluoro-4-pyrrolidin-3-yl-1,3-dihydro-benzoimidazol-2-one	267.8-269.9 °C
34		1-(3-Fluoro-benzyl)-6-fluoro-4-(2-methylamino-ethoxy)-1,3-dihydro-benzoimidazol-2-one	312





#	Structure	Name (Autonom)	MP °C/ M+H
35		1-(3-Fluoro-benzyl)-4-piperidin-4-yl-1,3-dihydro-benzoimidazol-2-one	>300 °C
36		1-(3-Fluoro-benzyl)-4-pyrrolidin-3-yl-1,3-dihydro-benzoimidazol-2-one	177.6-179.3 °C
37		1-Benzyl-6-methyl-4-piperidin-4-yl-1,3-dihydro-benzoimidazol-2-one	>300 °C
38		1-(2,3-difluoro-benzyl)-4-pyrrolidin-3-yl-1,3-dihydro-benzoimidazol-2-one	238.0-240.8 °C
39		1-(3-Fluoro-benzyl)-6-methyl-4-piperidin-4-yl-1,3-dihydro-benzoimidazol-2-one	340





#	Structure	Name (Autonom)	MP °C/ M+H
40		6-Fluoro-1-(3-fluoro-benzyl)-4-piperidin-4-yl-1,3-dihydro-benzoimidazol-2-one	>300 °C
41		(S)-1-(1-Phenyl-ethyl)-4-piperazin-1-yl-1,3-dihydro-benzoimidazol-2-one	323
42		(R)-1-(1-Phenyl-ethyl)-4-piperazin-1-yl-1,3-dihydro-benzoimidazol-2-one	323
43		7-Chloro-1-(3-fluoro-benzyl)-4-piperazin-1-yl-1,3-dihydro-benzoimidazol-2-one	361
44		5-Chloro-1-(3-fluoro-benzyl)-4-piperazin-1-yl-1,3-dihydro-benzoimidazol-2-one	361




#	Structure	Name (Autonom)	MP °C/ M+H
45		1-Benzyl-3-methyl-4-piperazin-1-yl-1,3-dihydro-benzoimidazol-2-one	323
46		5,6-Difluoro-1-(3-fluoro-benzyl)-4-piperazin-1-yl-1,3-dihydro-benzoimidazol-2-one	363
47		1-(1-Methyl-1-phenyl-ethyl)-4-piperazin-1-yl-1,3-dihydro-benzoimidazol-2-one	337
48		1-[1-(3-Fluoro-phenyl)-ethyl]-4-piperazin-1-yl-1,3-dihydro-benzoimidazol-2-one	341

#	Structure	Name (Autonom)	MP °C/ M+H
49		1-(2,3-Difluoro-benzyl)-6-fluoro-4-piperazin-1-yl-1,3-dihydro-benzoimidazol-2-one	363
50		6-Fluoro-1-(5-methyl-isoxazol-3-ylmethyl)-4-piperidin-4-yl-1,3-dihydro-benzoimidazol-2-one	331
51		1-(2,3-Difluoro-benzyl)-4-piperazin-1-yl-1,3-dihydro-benzoimidazol-2-one	345
52		6-Fluoro-1-(1-methyl-1-phenyl-ethyl)-4-piperazin-1-yl-1,3-dihydro-benzoimidazol-2-one	355

#	Structure	Name (Autonom)	MP °C/ M+H
53		1-(3-Fluoro-benzyl)-4-pyrrolidin-3-yl-1,3-dihydro-indol-2-one	311
54		1-(2,3-Difluoro-benzyl)-4-piperazin-1-yl-1H-indole-2,3-dione	358
55		1-(2-Fluoro-benzyl)-4-piperazin-1-yl-1H-indole-2,3-dione	232.1-234.3 °C
56		1-(2,3-Difluoro-benzyl)-4-piperazin-1-yl-1,3-dihydro-indol-2-one	>300 °C

#	Structure	Name (Autonom)	MP °C/ M+H
57		1-(3,4-Difluoro-benzyl)-4-piperazin-1-yl-1H-indole-2,3-dione	265.7-267.2 °C
58		1-(3,5-Difluoro-benzyl)-4-piperazin-1-yl-1H-indole-2,3-dione	270.3-273.0
59		1-(3,5-Difluoro-benzyl)-4-piperazin-1-yl-1,3-dihydro-indol-2-one	>300 °C
60		1-(3,4-Difluoro-benzyl)-4-piperazin-1-yl-1,3-dihydro-indol-2-one	>300 °C

#	Structure	Name (Autonom)	MP °C/ M+H
61		1-(3-Chloro-benzyl)-4-piperazin-1-yl-1H-indole-2,3-dione	249.1-251.7 °C
62		1-(2,5-Difluoro-benzyl)-4-piperazin-1-yl-1H-indole-2,3-dione	227.9-230.1 °C
63		1-(2-Fluoro-benzyl)-4-piperazin-1-yl-1,3-dihydroindol-2-one	261.7-263.0 °C
64		1-(3-Chloro-benzyl)-4-piperazin-1-yl-1,3-dihydroindol-2-one	276.1-277.0 °C

#	Structure	Name (Autonom)	MP °C/ M+H
65		5-Chloro-1-(3-fluoro-benzyl)-4-piperazin-1-yl-1,3-dihydro-indol-2-one	360
66		7-Chloro-1-(3-fluoro-benzyl)-4-piperazin-1-yl-1,3-dihydro-indol-2-one	273.4-275.0 °C
67		5,7-Dichloro-1-(3-fluoro-benzyl)-4-piperazin-1-yl-1,3-dihydro-indol-2-one	394

Another aspect of the invention provides a composition comprising a therapeutically effective amount of at least one compound of formula I and a pharmaceutically acceptable carrier.

Yet another aspect of the invention provides a method for treating a central nervous system (CNS) disease state in a subject comprising administering to the subject a therapeutically effective amount of a compound of formula I. The disease state may comprise, for example, psychoses, schizophrenia, manic depressions, neurological disorders, memory disorders, attention deficit disorder, Parkinson's disease, amyotrophic lateral sclerosis, Alzheimer's disease or Huntington's disease.

Still another aspect of the present invention provides a method for treating a disorder of the gastrointestinal tract in a subject comprising administering to the subject a therapeutically effective amount of a compound of formula (I).

Another aspect of the present invention provides a method for producing a compound of formula (I).

Synthesis


Compounds of the present invention can be made by a variety of methods
5 depicted in the illustrative synthetic reaction schemes shown and described below.

The starting materials and reagents used in preparing these compounds generally are either available from commercial suppliers, such as Aldrich Chemical Co., or are prepared by methods known to those skilled in the art following procedures set forth in references such as Fieser and Fieser's Reagents for Organic Synthesis; Wiley & Sons: New
10 York, 1991, Volumes 1-15; Rodd's Chemistry of Carbon Compounds, Elsevier Science Publishers, 1989, Volumes 1-5 and Supplementals; and Organic Reactions, Wiley & Sons: New York, 1991, Volumes 1-40. The following synthetic reaction schemes are merely illustrative of some methods by which the compounds of the present invention can be synthesized, and various modifications to these synthetic reaction schemes can be made
15 and will be suggested to one skilled in the art having referred to the disclosure contained in this Application.

The starting materials and the intermediates of the synthetic reaction schemes can be isolated and purified if desired using conventional techniques, including but not limited to, filtration, distillation, crystallization, chromatography, and the like. Such
20 materials can be characterized using conventional means, including physical constants and spectral data.

Unless specified to the contrary, the reactions described herein preferably are conducted under an inert atmosphere at atmospheric pressure at a reaction temperature range of from about -78 °C to about 150 °C, more preferably from about 0 °C to about
25 125 °C, and most preferably and conveniently at about room (or ambient) temperature, e.g., about 20 °C.

Scheme A below illustrates one synthetic procedure usable to prepare compounds of the invention, wherein m, n, p, Y, R², R³, R⁴, R⁶, R⁷ and R⁸ are as defined herein. Numerous synthetic routes to benzimidazolones are known and may be used in
30 preparation of the subject compounds, and the procedure of Scheme A is only exemplary. Specific examples of the procedure of Scheme A are provided in the following Experimental section.



SCHEME A

In step 1 of Scheme A, difluoronitro compound **a** is treated with amine compound **b** to afford compound **c**. In this reaction Y may be O or NR^d . The reaction of step 1 may be carried out in the presence of potassium carbonate or other weak base under polar solvent conditions. Where one or both of R^7 and R^8 are hydrogen, a suitable amine protection/deprotection strategy may be used, followed by deprotection after step 4 below. The fluoro groups of compound **a** may in certain embodiments be replaced with other leaving groups.

In step 2, an aralkylation reaction is effected by treatment of compound **c** with aralkyl amine **d** to provide aralkylamino compound **e**. Ar may be aryl or heteroaryl as noted above. In many embodiments compound **d** may be a benzyl amine. This reaction may also be carried out in the presence of potassium carbonate or like mild base under polar solvent conditions.


A reduction is carried out in step 3 to reduce the nitro group of compound **e** and provide aniline compound **f**. A mild reducing agent such as sodium dithionite, in polar protic solvent, may be used in this step.

In step 4, a cyclization is achieved by treatment of aniline compound **f** with phosgene or a phosgene equivalent to form benzimidazolone compound **g**.

Benzimidazolone compound **g** is a compound of formula I in accordance with the invention.


Many variations on the procedure of Scheme A are possible, as will be readily apparent to those skilled in the art. In one such variation, compound **b** may be replaced

more simply by a compound of the formula HY-PG wherein PG is a protecting group. After completion of step 4, the protecting group may then be removed and the group



introduced via alkylation. In another such variation, an additional alkylation step may be carried out to introduce an alkyl group at the 3-position of compound **g**. Where either of R^7 and R^8 are hydrogen, a subsequent alkylation step may also be carried out to introduce alkyl for these variables.

Scheme B below provides another synthetic route to the compounds of the invention, wherein m , n , p , Y , R^2 , R^3 , R^4 , R^6 , R^7 and R^8 are as defined herein.



10

SCHEME B


In step 1 of scheme B, indole compound **h** undergoes an N- alkylation by reaction with alkylating agent **i** to form the 1-substituted indole compound **j**. In step 2 compound **j** is treated with amine compound **b** to afford compound **c**. The group X may be O or NR^d as noted above. The presence of potassium carbonate or other weak base under polar solvent conditions may facilitate the reaction of step 2. Where one or both of R^7 and R^8 are hydrogen, a suitable amine protection/deprotection strategy may be used, followed by subsequent deprotection.

In step 3 indole compound **k** undergoes bromination by treatment with N-bromosuccinimide or other bromine source (not shown) to yield 3-bromo indole compound **l**. Bromoindole **l** is then subject to oxidation in step 4 to provide

20

dihydroindolone compound m. Compound m is a compound of formula I in accordance with the invention.

Scheme C illustrates a method for making compounds of formula I wherein R^1 is a pyrrolidinyl group. In Scheme C L is a leaving group such as bromine or other halo, and variables m, n, X, R^2 , R^3 , R^4 , R^6 , R^7 , R^8 , R^{10} and R^{11} are as defined herein.



SCHEME C

In step 1 of Scheme C, compound n is reacted with pyrrole boronic acid compound o to afford a pyrrolyl-substituted compound p. Compound p is then hydrogenated to yield a pyrrolidinyl-substituted compound g. Compound g is a compound of formula I in accordance with the invention. In embodiments where R^8 is hydrogen, suitable protection and deprotection techniques may be used with the method of Scheme C.

More specific details for producing compounds of formula I are described in the Examples section below.

Utility

The compounds of the invention have selective affinity for 5-HT receptors, including the 5-HT₆ the 5-HT_{2A} receptor, or both, and as such are expected to be useful in the treatment of certain CNS disorders such as Parkinson's disease, Huntington's disease, anxiety, depression, manic depression, psychosis, epilepsy, obsessive compulsive disorders, mood disorders, migraine, Alzheimer's disease (enhancement of cognitive memory), sleep disorders, feeding disorders such as anorexia, bulimia, and obesity, panic attacks, akathisia, attention deficit hyperactivity disorder (ADHD), attention deficit disorder (ADD), withdrawal from drug abuse such as cocaine, ethanol, nicotine and benzodiazepines, schizophrenia, and also disorders associated with spinal trauma and/or head injury such as hydrocephalus. Such compounds are also expected to be of use in the treatment of certain GI (gastrointestinal) disorders such functional bowel disorder and irritable bowel syndrome.

Testing

The pharmacology of the compounds of this invention was determined by art recognized procedures. The in vitro techniques for determining the affinities of test compounds at the 5-HT₆ receptor and the 5-HT_{2A} receptor in radioligand binding, FLIPR and functional assays are described below.

5 Administration and Pharmaceutical Composition

The present invention includes pharmaceutical compositions comprising at least one compound of the present invention, or an individual isomer, racemic or non-racemic mixture of isomers or a pharmaceutically acceptable salt or solvate thereof, together with at least one pharmaceutically acceptable carrier, and optionally other therapeutic and/or
10 prophylactic ingredients.

In general, the compounds of the present invention will be administered in a therapeutically effective amount by any of the accepted modes of administration for agents that serve similar utilities. Suitable dosage ranges are typically 1-500 mg daily, preferably 1-100 mg daily, and most preferably 1-30 mg daily, depending upon numerous
15 factors such as the severity of the disease to be treated, the age and relative health of the subject, the potency of the compound used, the route and form of administration, the indication towards which the administration is directed, and the preferences and experience of the medical practitioner involved. One of ordinary skill in the art of treating such diseases will be able, without undue experimentation and in reliance upon
20 personal knowledge and the disclosure of this Application, to ascertain a therapeutically effective amount of the compounds of the present invention for a given disease.

In general, compounds of the present invention will be administered as pharmaceutical formulations including those suitable for oral (including buccal and sub-lingual), rectal, nasal, topical, pulmonary, vaginal, or parenteral (including
25 intramuscular, intraarterial, intrathecal, subcutaneous and intravenous) administration or in a form suitable for administration by inhalation or insufflation. The preferred manner of administration is generally oral using a convenient daily dosage regimen which can be adjusted according to the degree of affliction.

A compound or compounds of the present invention, together with one or more
30 conventional adjuvants, carriers, or diluents, may be placed into the form of pharmaceutical compositions and unit dosages. The pharmaceutical compositions and unit dosage forms may be comprised of conventional ingredients in conventional proportions, with or without additional active compounds or principles, and the unit dosage forms may contain any suitable effective amount of the active ingredient
35 commensurate with the intended daily dosage range to be employed. The pharmaceutical compositions may be employed as solids, such as tablets or filled capsules, semisolids, powders, sustained release formulations, or liquids such as solutions, suspensions,

emulsions, elixirs, or filled capsules for oral use; or in the form of suppositories for rectal or vaginal administration; or in the form of sterile injectable solutions for parenteral use. Formulations containing about one (1) milligram of active ingredient or, more broadly, about 0.01 to about one hundred (100) milligrams, per tablet, are accordingly suitable
5 representative unit dosage forms.

The compounds of the present invention may be formulated in a wide variety of oral administration dosage forms. The pharmaceutical compositions and dosage forms may comprise a compound or compounds of the present invention or pharmaceutically acceptable salts thereof as the active component. The pharmaceutically acceptable
10 carriers may be either solid or liquid. Solid form preparations include powders, tablets, pills, capsules, cachets, suppositories, and dispersible granules. A solid carrier may be one or more substances which may also act as diluents, flavouring agents, solubilizers, lubricants, suspending agents, binders, preservatives, tablet disintegrating agents, or an encapsulating material. In powders, the carrier generally is a finely divided solid which is
15 a mixture with the finely divided active component. In tablets, the active component generally is mixed with the carrier having the necessary binding capacity in suitable proportions and compacted in the shape and size desired. The powders and tablets preferably contain from about one (1) to about seventy (70) percent of the active compound. Suitable carriers include but are not limited to magnesium carbonate,
20 magnesium stearate, talc, sugar, lactose, pectin, dextrin, starch, gelatine, tragacanth, methylcellulose, sodium carboxymethylcellulose, a low melting wax, cocoa butter, and the like. The term "preparation" is intended to include the formulation of the active compound with encapsulating material as carrier, providing a capsule in which the active component, with or without carriers, is surrounded by a carrier, which is in association
25 with it. Similarly, cachets and lozenges are included. Tablets, powders, capsules, pills, cachets, and lozenges may be as solid forms suitable for oral administration.

Other forms suitable for oral administration include liquid form preparations including emulsions, syrups, elixirs, aqueous solutions, aqueous suspensions, or solid form preparations which are intended to be converted shortly before use to liquid form
30 preparations. Emulsions may be prepared in solutions, for example, in aqueous propylene glycol solutions or may contain emulsifying agents, for example, such as lecithin, sorbitan monooleate, or acacia. Aqueous solutions can be prepared by dissolving the active component in water and adding suitable colorants, flavors, stabilizers, and thickening agents. Aqueous suspensions can be prepared by dispersing
35 the finely divided active component in water with viscous material, such as natural or synthetic gums, resins, methylcellulose, sodium carboxymethylcellulose, and other well known suspending agents. Solid form preparations include solutions, suspensions, and

emulsions, and may contain, in addition to the active component, colorants, flavors, stabilizers, buffers, artificial and natural sweeteners, dispersants, thickeners, solubilizing agents, and the like.

The compounds of the present invention may be formulated for parenteral
5 administration (e.g., by injection, for example bolus injection or continuous infusion) and may be presented in unit dose form in ampoules, pre-filled syringes, small volume infusion or in multi-dose containers with an added preservative. The compositions may take such forms as suspensions, solutions, or emulsions in oily or aqueous vehicles, for example solutions in aqueous polyethylene glycol. Examples of oily or nonaqueous
10 carriers, diluents, solvents or vehicles include propylene glycol, polyethylene glycol, vegetable oils (e.g., olive oil), and injectable organic esters (e.g., ethyl oleate), and may contain formulatory agents such as preserving, wetting, emulsifying or suspending, stabilizing and/or dispersing agents. Alternatively, the active ingredient may be in powder form, obtained by aseptic isolation of sterile solid or by lyophilization from solution for
15 constitution before use with a suitable vehicle, e.g., sterile, pyrogen-free water.

The compounds of the present invention may be formulated for topical administration to the epidermis as ointments, creams or lotions, or as a transdermal patch. Ointments and creams may, for example, be formulated with an aqueous or oily base with the addition of suitable thickening and/or gelling agents. Lotions may be
20 formulated with an aqueous or oily base and will in general also containing one or more emulsifying agents, stabilizing agents, dispersing agents, suspending agents, thickening agents, or coloring agents. Formulations suitable for topical administration in the mouth include lozenges comprising active agents in a flavored base, usually sucrose and acacia or tragacanth; pastilles comprising the active ingredient in an inert base such as gelatine and
25 glycerine or sucrose and acacia; and mouthwashes comprising the active ingredient in a suitable liquid carrier.

The compounds of the present invention may be formulated for administration as suppositories. A low melting wax, such as a mixture of fatty acid glycerides or cocoa butter is first melted and the active component is dispersed homogeneously, for example,
30 by stirring. The molten homogeneous mixture is then poured into convenient sized molds, allowed to cool, and to solidify.

The compounds of the present invention may be formulated for vaginal administration. Pessaries, tampons, creams, gels, pastes, foams or sprays containing in addition to the active ingredient such carriers as are known in the art to be appropriate.

35 The compounds of the present invention may be formulated for nasal administration. The solutions or suspensions are applied directly to the nasal cavity by conventional means, for example, with a dropper, pipette or spray. The formulations

may be provided in a single or multidose form. In the latter case of a dropper or pipette, this may be achieved by the patient administering an appropriate, predetermined volume of the solution or suspension. In the case of a spray, this may be achieved for example by means of a metering atomizing spray pump.

5 The compounds of the present invention may be formulated for aerosol administration, particularly to the respiratory tract and including intranasal administration. The compound will generally have a small particle size for example of the order of five (5) microns or less. Such a particle size may be obtained by means known in the art, for example by micronization. The active ingredient is provided in a pressurized
10 pack with a suitable propellant such as a chlorofluorocarbon (CFC), for example, dichlorodifluoromethane, trichlorofluoromethane, or dichlorotetrafluoroethane, or carbon dioxide or other suitable gas. The aerosol may conveniently also contain a surfactant such as lecithin. The dose of drug may be controlled by a metered valve. Alternatively the active ingredients may be provided in a form of a dry powder, for
15 example a powder mix of the compound in a suitable powder base such as lactose, starch, starch derivatives such as hydroxypropylmethyl cellulose and polyvinylpyrrolidone (PVP). The powder carrier will form a gel in the nasal cavity. The powder composition may be presented in unit dose form for example in capsules or cartridges of e.g., gelatine or blister packs from which the powder may be administered by means of an inhaler.

20 When desired, formulations can be prepared with enteric coatings adapted for sustained or controlled release administration of the active ingredient. For example, the compounds of the present invention can be formulated in transdermal or subcutaneous drug delivery devices. These delivery systems are advantageous when sustained release of the compound is necessary and when patient compliance with a treatment regimen is
25 crucial. Compounds in transdermal delivery systems are frequently attached to an skin-adhesive solid support. The compound of interest can also be combined with a penetration enhancer, e.g., Azone (1-dodecylazacycloheptan-2-one). Sustained release delivery systems are inserted subcutaneously into the subdermal layer by surgery or injection. The subdermal implants encapsulate the compound in a lipid soluble
30 membrane, e.g., silicone rubber, or a biodegradable polymer, e.g., polylactic acid.

 The pharmaceutical preparations are preferably in unit dosage forms. In such form, the preparation is subdivided into unit doses containing appropriate quantities of the active component. The unit dosage form can be a packaged preparation, the package containing discrete quantities of preparation, such as packeted tablets, capsules, and
35 powders in vials or ampoules. Also, the unit dosage form can be a capsule, tablet, cachet, or lozenge itself, or it can be the appropriate number of any of these in packaged form.

Other suitable pharmaceutical carriers and their formulations are described in Remington: The Science and Practice of Pharmacy 1995, edited by E. W. Martin, Mack Publishing Company, 19th edition, Easton, Pennsylvania. Representative pharmaceutical formulations containing a compound of the present invention are described in the

5 Examples below.


EXAMPLES

The following preparations and examples are given to enable those skilled in the art to more clearly understand and to practice the present invention. They should not be
10 considered as limiting the scope of the invention, but merely as being illustrative and representative thereof.

Example 1

1-Benzyl-4-piperazin-1-yl-1,3-dihydro-benzoimidazol-2-one


The synthetic procedures described in this Example were carried out according to
15 the process shown in Scheme D.



SCHEME D


Step 1

20 4-(3-Fluoro-2-nitro-phenyl)-piperazine-1-carboxylic acid tert-butyl ester



To a solution of piperazine-1-carboxylic acid tert-butyl ester (0.204 g, 1.1 mmol) in 1 mL dimethylsulfoxide was added potassium carbonate (0.303 g, 2.2 mmol) followed by 1,3-difluoro-2-nitro-benzene (0.159 g, 1 mmol). The solution was stirred 30 minutes
 5 at room temperature, and was then diluted with 50 mL ethyl ether. The organic phase was washed three times with 50 mL water and once with 50 mL brine, dried over sodium sulfated, and concentrated *in vacuo* to give 0.314 g (0.965 mmol, 96.5%) of 4-(3-fluoro-2-nitro-phenyl)-piperazine-1-carboxylic acid tert-butyl ester as a yellow oil. MS: 226 (M-BOC+H)⁺.


10

Step 24-(3-Benzylamino-2-nitro-phenyl)-piperazine-1-carboxylic acid tert-butyl ester

4-(3-Fluoro-2-nitro-phenyl)-piperazine-1-carboxylic acid tert-butyl ester (0.300
 15 g, 0.923 mmol) and benzylamine (0.110 mL, 1.015 mmol) were combined in 1 mL dimethylsulfoxide with potassium carbonate (0.318 g, 2.308 mmol), and the reaction mixture was heated to 120°C for 2 hours. The reaction mixture was poured onto 100 g icewater and extracted twice with 100 mL ethyl acetate. The combined organic phases were washed twice with 50 mL water, once with 50 mL brine, dried over sodium sulfate,
 20 and concentrated *in vacuo*. The residue is purified by flash chromatography (1% to 15% ethyl acetate in hexanes) to afford 0.208 g (0.50 mmol, 54%) of 4-(3-benzylamino-2-nitro-phenyl)-piperazine-1-carboxylic acid tert-butyl ester. MS: 413 (M+H)⁺.

25


Step 34-(2-Amino-3-benzylamino-phenyl)-piperazine-1-carboxylic acid tert-butyl ester



A solution of 4-(3-benzylamino-2-nitro-phenyl)-piperazine-1-carboxylic acid tert-butyl ester (0.200 g, 0.485 mmol) in 5 mL ethanol was added to a solution sodium dithionite (0.565 g, 3.245 mmol) in 10 mL water, and the reaction mixture was heated to 100°C. The mixture was stirred for 5 minutes, cooled to room temperature, and concentrated *in vacuo* to remove ethanol. A yellow solid precipitates and was filtered and dried under vacuum for 18 hours to give 0.173 g (0.45 mmol, 90%) of 4-(2-amino-3-benzylamino-phenyl)-piperazine-1-carboxylic acid tert-butyl ester. MS: 383 (M+H)⁺.

Step 4


4-(1-Benzyl-2-oxo-2,3-dihydro-1H-benzoimidazol-4-yl)-piperazine-1-carboxylic acid tert-butyl ester



To a solution of 4-(2-amino-3-benzylamino-phenyl)-piperazine-1-carboxylic acid tert-butyl ester (0.176 g, 0.459 mmol) in 1 mL dichloromethane was added 1 mL of 2M aqueous sodium carbonate. Phosgene (0.261 mL of a 1.93 M solution in toluene, 0.504 mmol) was added dropwise to the stirring solution over five minutes. Stirring was continued for two hours as a fine white precipitate was formed. The solid was filtered and dried under a stream of nitrogen to give 0.112 g of 4-(1-benzyl-2-oxo-2,3-dihydro-1H-benzoimidazol-4-yl)-piperazine-1-carboxylic acid tert-butyl ester as a fine white powder. MS: 409 (M+H)⁺.

Step 5

1-Benzyl-4-piperazin-1-yl-1,3-dihydro-benzoimidazol-2-one





To a solution of 0.112 g of 4-(1-benzyl-2-oxo-2,3-dihydro-1H-benzoimidazol-4-yl)-piperazine-1-carboxylic acid tert-butyl ester in 15 mL ethyl acetate with 5 mL ethanol is added 3 mL of 2M ethanolic hydrogen chloride. The resulting solution is refluxed for one hour, and on cooling, a solid precipitates. The solid is filtered and dried over night under vacuum to give 82 mg of 1-benzyl-4-piperazin-1-yl-1,3-dihydro-benzoimidazol-2-one hydrochloride. MS: 309 (M+H)⁺.

Similarly prepared, but using 1-methylpiperazine in step 1 instead of piperazine-1-carboxylic acid tert-butyl ester, and omitting step 5, was 1-Benzyl-4-piperazin-1-yl-1,3-dihydro-benzoimidazol-2-one. MS: 323 (M+H)⁺.

Example 2

4-(Azetidin-3-ylmethoxy)-1-benzyl-1,3-dihydro-benzoimidazol-2-one

The synthetic procedures described in this Example were carried out according to the process shown in Scheme E.

**SCHEME E****Step 1****1-Benzyl-2-(benzyloxy)-3-fluoro-4-nitrobenzene**

5

To a suspension of sodium hydride (1.522 g, 38.06 mmol) in anhydrous DMF (150 mL) at room temperature was added benzyl alcohol (3.44 mL, 33.3 mmol) dropwise over 10 minutes, and stirring was continued for another 10 minutes. 2,6-Difluoronitrobenzene (5.046 g, 31.72 mmol) was added to the reaction mixture in one portion. The reaction mixture was stirred for an hour, and 100 mL water and 100 mL ethyl acetate were added. The layers were separated, and the aqueous layer was extracted with 75 mL ethyl acetate. The combined organic layers were washed with 100 mL water


10

and 100 mL brine, dried over sodium sulfate and concentrated *in vacuo*. The resulting residue was purified by flash chromatography (6% to 35% ethyl acetate in hexanes) to give 6.818 g (27.6 mmol, 87%) of 1-benzyloxy-3-fluoro-2-nitro-benzene as a yellow oil. MS: 248 (M+H)⁺.

- 5 Also prepared in a similar manner using (2-hydroxy-ethyl)-methyl-carbamic acid *tert*-butyl ester in place of benzyl alcohol was [2-(3-Fluoro-2-nitro-phenoxy)-ethyl]-methyl-carbamic acid *tert*-butyl ester. MS: 215 (M-BOC+H)⁺.

Step 2

Benzyl-(3-benzyloxy-2-nitro-phenyl)-amine



10


- Potassium carbonate (5.71 g, 41.4 mmol) was added to a solution of 1-benzyloxy-3-fluoro-2-nitro-benzene (6.818 g, 27.6 mmol) and benzylamine (3.32 mL, 30.34 mmol) in tetrahydrofuran, and the resulting suspension was heated to 110°C for 2 hours. After cooling to room temperature, the mixture was poured onto 1 L of ice water and extracted
 15 twice with 150 mL ethyl acetate and twice with 75 mL dichloromethane. The combined organic fractions were dried over sodium sulfate and concentrated *in vacuo* in the presence of 25 g. silica gel. The preloaded silica gel was submitted to flash chromatography (8% to 35% ethyl acetate in hexanes) to give 8.118 g (24.28 mmol, 88%) of benzyl-(3-benzyloxy-2-nitro-phenyl)-amine as an orange solid. MS: 335 (M+H)⁺.

- 20 Also prepared in a similar fashion starting with [2-(3-fluoro-2-nitro-phenoxy)-ethyl]-methyl-carbamic acid *tert*-butyl ester was [2-(3-Benzylamino-2-nitro-phenoxy)-ethyl]-methyl-carbamic acid *tert*-butyl ester. MS: 302 (M-BOC+H)⁺.

Step 3

N*1*-Benzyl-3-benzyloxy-benzene-1,2-diamine

25




A solution of benzyl-(3-benzyloxy-2-nitro-phenyl)-amine (8.116 g, 24.3 mmol) in 500 mL ethanol was added to a solution of sodium thiosulfite (29.0 g, 162.8 g.) in 350 mL water at 100°C while stirring. The reaction mixture was heated for 30 minutes at this temperature, and then cooled and concentrated *in vacuo*. The off-white solid which precipitates was filtered and dried overnight under a stream of nitrogen to give 6.77 g (22.24 mmol, 91.5%) of N¹-Benzyl-3-benzyloxy-benzene-1,2-diamine. MS: 305 (M+H)⁺.

Prepared in a similar fashion, starting with [2-(3-benzylamino-2-nitro-phenoxy)-ethyl]-methyl-carbamic acid tert-butyl ester, was [2-(2-Amino-3-benzylamino-phenoxy)-ethyl]-methyl-carbamic acid tert-butyl ester. MS: 372 (M+H)⁺.

Step 4


1-Benzyl-4-benzyloxy-1,3-dihydro-benzoimidazol-2-one



To a solution of N¹-Benzyl-3-benzyloxy-benzene-1,2-diamine (4.60 g, 15.11 mmol) in 100 mL dichloromethane at 0°C was added triethylamine (4.201 mL, 30.22 mmol) followed by portionwise addition of triphosgene (1.57 g, 5.29 mmol). The resulting solution was stirred for 2 hours at 0°C. The reaction mixture was allowed to warm to room temperature and was then added to 200 mL water. The organic phase was separated, washed with 100 mL 10% aqueous HCl, 100 mL saturated aqueous sodium bicarbonate, and 100 mL brine, dried over sodium sulfate, and concentrated *in vacuo* to give 4.60 g (13.92 mmol, 92%) of 1-benzyl-4-benzyloxy-1,3-dihydro-benzoimidazol-2-one as a tan solid. MS: 331 (M+H)⁺.


Prepared in a similar fashion, starting with [2-(2-Amino-3-benzylamino-phenoxy)-ethyl]-methyl-carbamic acid tert-butyl ester, was [2-(1-Benzyl-2-oxo-2,3-dihydro-1H-benzoimidazol-4-yloxy)-ethyl]-methyl-carbamic acid tert-butyl ester. MS: 398 (M+H)⁺.

5

Step 51-Benzyl-4-hydroxy-1,3-dihydro-benzoimidazol-2-one

To a solution of 1-benzyl-4-benzyloxy-1,3-dihydro-benzoimidazol-2-one (0.875 g, 2.64 mmol) in 100 ml of 1:1 ethyl acetate:THF was added palladium hydroxide (0.270 mg, 1.923 mmol). The reaction mixture was purged with hydrogen gas at 1 atm., allowed to stir at room temperature for 2 hours, and then filtered through celite. The filtrate was concentrated *in vacuo* to give 640 mg (2.64 mmol, quantitative) of 1-benzyl-4-hydroxy-1,3-dihydro-benzoimidazol-2-one as a white solid. MS: 241 (M+H)⁺.

15

Step 63-(1-Benzyl-2-oxo-2,3-dihydro-1H-benzoimidazol-4-yloxymethyl)-azetidine-1-carboxylic acid tert-butyl ester


To a solution of 1-benzyl-4-hydroxy-1,3-dihydro-benzoimidazol-2-one (0.100 g, 0.417 mmol) and 3-hydroxymethyl-azetidine-1-carboxylic acid tert-butyl ester (0.078 g, 0.417 mmol) in 0.5 mL anhydrous THF under nitrogen was added triphenylphosphine (0.109 g, 0.417 mmol) and diisopropyl azodicarboxylate (0.082 mL, 0.417 mmol) dropwise. The reaction mixture was stirred for 24 h, concentrated *in vacuo*, and purified by preparative TLC (3% methanol in dichloromethane) to give 57 mg (0.14 mmol,

20

33.6%) of 3-(1-benzyl-2-oxo-2,3-dihydro-1H-benzoimidazol-4-ylloxymethyl)-azetidine-1-carboxylic acid tert-butyl ester as a clear oil. MS: 410 (M+H)⁺.

Step 7

4-(Azetidin-3-ylmethoxy)-1-benzyl-1,3-dihydro-benzoimidazol-2-one



5

3-(1-Benzyl-2-oxo-2,3-dihydro-1H-benzoimidazol-4-ylloxymethyl)-azetidine-1-carboxylic acid tert-butyl ester 57 mg (0.14 mmol) was dissolved in 0.5 mL ethanol and combined with 0.5 mL 2N ethanolic hydrogen chloride and refluxed for 30 minutes. The white solid which precipitates on cooling is filtered and dried under vacuum for 18 hours to give 23 mg of 4-(azetidin-3-ylmethoxy)-1-benzyl-1,3-dihydro-benzoimidazol-2-one hydrochloride as a white solid. MP: 210.0-216.6 °C. MS: 310 (M+H)⁺.

10

The following compounds were prepared in a similar fashion using the appropriate amino alcohol or N-BOC protected amino alcohol:

1-Benzyl-4-(3-dimethylamino-propoxy)-1,3-dihydro-benzoimidazol-2-one, MP: 186.5-189.5 °C (HCl salt). MS: 326 (M+H)⁺;

15

1-Benzyl-4-(pyrrolidin-3-ylmethoxy)-1,3-dihydro-benzoimidazol-2-one (racemic), MP: 226.0-228.9 °C (HCl salt). MS: 324 (M+H)⁺;

1-Benzyl-4-(piperidin-4-yloxy)-1,3-dihydro-benzoimidazol-2-one, MP: 260.9-263.3 °C (HCl salt). MS: 324 (M+H)⁺; and

20



(R)-1-Benzyl-4-(pyrrolidin-2-ylmethoxy)-1,3-dihydro-benzoimidazol-2-one, MP: 239.5-241.8 °C (HCl salt). MS: 324 (M+H)⁺.

Example 3

4-(Azetidin-3-yloxy)-1-benzyl-1,3-dihydro-benzoimidazol-2-one


The synthetic procedures described in this Example were carried out according to the process shown in Scheme F.

25

**SCHEME F****Step 1****4-(1-Benzhydryl-azetidin-3-yloxy)-1-benzyl-1,3-dihydro-benzoimidazol-2-one**

5

To a suspension of 1-benzyl-4-hydroxy-1,3-dihydro-benzoimidazol-2-one (0.104 g., 0.433 mmol) and 1-benzhydryl-azetidin-3-ol (0.114 g., 0.477 mmol) in 0.5 mL anhydrous THF was added triphenyl phosphine (0.125 g., 0.477 mmol) and diisopropyl azodicarboxylate (0.093 mL, 0.477 mmol). The reaction mixture was refluxed for 2 hours, then concentrated *in vacuo* and purified by preparative TLC (2% methanol in dichloromethane) to give 78 mg (0.17 mmol, 39.5%) of 4-(1-benzhydryl-azetidin-3-yloxy)-1-benzyl-1,3-dihydro-benzoimidazol-2-one as a yellow glass. MS: 462 (M+H)⁺.

Step 2**4-(Azetidin-3-yloxy)-1-benzyl-1,3-dihydro-benzoimidazol-2-one**


15

4-(1-Benzhydryl-azetidin-3-yloxy)-1-benzyl-1,3-dihydro-benzoimidazol-2-one (78 mg, 0.17 mmol) was dissolved in 0.33 mL dichloroethane and cooled to 0°C under nitrogen. α -Chloroethyl chloroformate (0.018 mL, 0.164 mmol) was added dropwise, and the solution was refluxed for 2 hours, concentrated *in vacuo*, and dissolved in 20 mL methanol. The solution was refluxed for one hour, concentrated *in vacuo* and dissolved in 75 mL dichloromethane, and the resulting solution was washed with 75 mL 2M aqueous potassium carbonate and 75 mL water. The organic phase was dried over sodium sulfate, concentrated *in vacuo* and purified by flash (3% methanol in dichloromethane) chromatography to give 42 mg (0.14 mmol, 82%) of 4-(azetidin-3-yloxy)-1-benzyl-1,3-dihydro-benzoimidazol-2-one as a clear oil. MS: 296 (M+H)⁺. The corresponding hydrochloride salt, prepared by recrystallization from EtOH/HCl, gave a melting point of 181.0-184.0 °C.

Example 4

1-Benzyl-3-methyl-4-(2-methylamino-ethoxy)-1,3-dihydro-benzoimidazol-2-one


The synthetic procedures described in this Example were carried out according to the process shown in Scheme G.



SCHEME G

Step 1

[2-(1-Benzyl-3-methyl-2-oxo-2,3-dihydro-1H-benzoimidazol-4-yloxy)-ethyl]-methyl-carbamic acid tert-butyl ester




The [2-(1-benzyl-2-oxo-2,3-dihydro-1H-benzoimidazol-4-yloxy)-ethyl]-methyl-carbamic acid tert-butyl ester used in this example was prepared using the procedure of steps 1-4 of Example 2, but replacing benzyl alcohol in step 1 with (2-Hydroxy-ethyl)-methyl-carbamic acid tert-butyl ester.

- 5 To a solution of [2-(1-benzyl-2-oxo-2,3-dihydro-1H-benzoimidazol-4-yloxy)-ethyl]-methyl-carbamic acid tert-butyl ester (0.100 g, 0.252 mmol) in 2 mL of anhydrous N,N-dimethylformamide was added sodium hydride (11 mg of a 60% dispersion in mineral oil) under nitrogen. The suspension was stirred at room temperature for ten minutes and then methyl iodide (0.018 mL, 0.277 mmol) was added in one portion.
- 10 Stirring was continued for one hour and the reaction mixture was added to 150 mL water, and extracted twice with 75 mL ethyl acetate. The combined organic fraction was washed with 100 mL brine, dried over sodium sulfated and concentrated *in vacuo*. The residue was purified by flash chromatography (18% to 28% ethyl acetate in hexane) to afford 92 mg (0.223 mmol, 88.5%) of [2-(1-benzyl-3-methyl-2-oxo-2,3-dihydro-1H-
- 15 benzoimidazol-4-yloxy)-ethyl]-methyl-carbamic acid tert-butyl ester as a clear oil. MS: 412 (M+H)⁺.

Step 2

1-Benzyl-3-methyl-4-(2-methylamino-ethoxy)-1,3-dihydro-benzoimidazol-2-one




20

- [2-(1-Benzyl-3-methyl-2-oxo-2,3-dihydro-1H-benzoimidazol-4-yloxy)-ethyl]-methyl-carbamic acid tert-butyl ester (92 mg, 0.223 mmol) was dissolved in 1 mL ethanol and combined with 0.5 mL of 2N ethanolic HCl, and the resulting solution was refluxed 30 minutes. On cooling, a white precipitate was observed. The solid was filtered and
- 25 dried for 18 hours under vacuum to give 47 mg of 1-benzyl-3-methyl-4-(2-methylamino-ethoxy)-1,3-dihydro-benzoimidazol-2-one hydrochloride. MP: 203.8-205.1 °C. MS: 312 (M+H)⁺.


- Similarly prepared by the above procedure but omitting step one, was 1-Benzyl-4-(2-methylamino-ethoxy)-1,3-dihydro-benzoimidazol-2-one. MP: 188.7-191.3 °C (HCl
- 30 Salt), MS: 298 (M+H)⁺.

Example 51-Benzyl-4-(2-methoxy-ethyl)-1,3-dihydro-benzoimidazol-2-one

The synthetic procedures described in this Example were carried out according to the process shown in Scheme H.



5

SCHEME HStep 11-(2,2-Dimethoxy-ethyl)-2,3-dinitro-benzene


10

To a solution of 2,3-dinitrotoluene (3.02 g, 16.6 mmol) in 25 mL of DMF was added dimethylformamide dimethylacetal (6.0 mL, 5.4 g, 45 mmol), and the solution was

brought to 140°C and stirred for 16 hours. The solvent was removed under reduced pressure to afford a dark red solid mass. This crude enamine was dissolved in 40 mL of MeOH and 4.0 mL of chlorotrimethylsilane (3.4 g, 31.7 mmol) was added. The solution was brought to reflux and stirred for 16 hours at reflux. The solvent was removed under reduced pressure and the crude material chromatographed directly to afford the 1-(2,2-dimethoxy-ethyl)-2,3-dinitro-benzene (1.85 g, 44%). ¹H NMR (CDCl₃) δ 2.96 (2H, d, *J* = 5.2 Hz), 3.35 (6H, s), 4.49 (1H, t, *J* = 5.2 Hz), 7.60 (apparent t, 1H, *J* = 8.0 Hz), 7.76 (d, 1H, *J* = 6.5 Hz), 8.05 (dd, 1H, *J* = 1.3, 8.0).

Step 2


10 3-(2,2-Dimethoxy-ethyl)-benzene-1,2-diamine




1-(2,2-Dimethoxy-ethyl)-2,3-dinitro-benzene (1.85g, 7.19 mmol) was dissolved in 20 mL of MeOH and 90 mg of 10% by weight Pd/C was added. This mixture was stirred under 1 atmosphere of H₂ for 14 hours at room temperature. The mixture was filtered over celite and purified by chromatography to afford 3-(2,2-dimethoxy-ethyl)-benzene-1,2-diamine (1.04 g, 73%). ¹H NMR (CDCl₃) δ 2.87 (2H, d, *J* = 5.3 Hz), 3.37 (6H, s), 4.50 (1H, t, *J* = 5.3 Hz), 7.60 (6.62-6.66, m, 3H).

Step 3


20 4-(2,2-Dimethoxy-ethyl)-1,3-dihydro-benzoimidazol-2-one




3-(2,2-Dimethoxy-ethyl)-benzene-1,2-diamine (1.03 g, 5.26 mmol) was dissolved in THF and carbonyldiimidazole (935 mg, 5.77 mmol) added. The mixture was stirred at rt for 18 hours. The solvent was removed and the crude material chromatographed to afford 4-(2,2-dimethoxy-ethyl)-1,3-dihydro-benzoimidazol-2-one as a colorless solid (440 mg, 38%). ¹H NMR (CDCl₃) δ 3.03 (2H, d, *J* = 5.2 Hz), 3.40 (6H, s), 4.58 (1H, t, *J* = 5.2 Hz), 6.86-6.89 (m, 1H), 6.96-7.02 (m, 2H), 9.29 (br s, 1H), 10.28 (br s, 1H). MS: 221 (M-H)⁺.

Step 41-Benzyl-4-(2,2-dimethoxy-ethyl)-1,3-dihydro-benzoimidazol-2-one


4-(2,2-Dimethoxy-ethyl)-1,3-dihydro-benzoimidazol-2-one (435 mg, 1.96 mmol) was dissolved along with benzyl bromide (335 mg, 1.96 mmol) in 10 mL of anhydrous DMF. Potassium *tert*-butoxide (1.0 M in THF, 2.2 mL, 2.2 mmol) was added. The reaction mixture was stirred at room temperature for 90 minutes and then partitioned between ether and water. The organic phase was washed with brine and dried over Na_2SO_4 . After filtration and removal of solvent under reduced pressure, the crude material was chromatographed to afford 1-benzyl-4-(2,2-dimethoxy-ethyl)-1,3-dihydro-benzoimidazol-2-one (100 mg, 16%). ^1H NMR (CDCl_3) δ 3.00 (2H, d, $J = 5.0$ Hz), 3.40 (6H, s), 4.55 (1H, t, $J = 5.0$ Hz), 6.75 (d, 1H, $J = 7.5$ Hz), 6.84-6.95 (m, 2H), 7.23-7.35 (m, 5H), 8.95 (br s, 1H). ^{13}C (CDCl_3) δ 36.9, 44.9, 54.4, 105.6, 107.5, 118.8, 121.7, 124.0, 127.9, 128.0, 129.1, 129.2, 130.8, 136.7.

15 Step 5(1-Benzyl-2-oxo-2,3-dihydro-1H-benzoimidazol-4-yl)-acetaldehyde

(1-Benzyl-2-oxo-2,3-dihydro-1H-benzoimidazol-4-yl)-acetaldehyde (90 mg, 0.29 mmol) was dissolved in acetone and *p*-toluene sulfonic acid (10 mg) added. The mixture was stirred for 2 hours and then partitioned between ethyl acetate and water. After washing with saturated sodium bicarbonate, water, and brine, the organic layer was dried over Na_2SO_4 , filtered, and concentrated to afford 1-benzyl-4-(2-hydroxy-ethyl)-1,3-dihydro-benzoimidazol-2-one (80 mg, quant). ^1H NMR (CDCl_3) δ 3.84 (d, 2H, $J = 2.1$ Hz), 5.06 (s, 2H), 6.80-7.02 (M, 3H), 7.20-7.33 (m, 5H), 9.78 (t, 1H, $J = 2.0$ Hz), 10.7 (br s, 1H). MS: 265 ($\text{M}-\text{H}$) $^-$.


Step 61-Benzyl-4-[2-(benzyl-methyl-amino)-ethyl]-1,3-dihydro-benzoimidazol-2-one

To a solution of 1-benzyl-4-(2-hydroxy-ethyl)-1,3-dihydro-benzoimidazol-2-one
 5 (53 mg) and benzylmethylamine (29 mg) in 1.0 mL of methylene chloride was added
 sodium triacetoxy borohydride (51 mg). After one hour the reaction was quenched with
 water, and partitioned with methylene chloride. The crude material was washed with
 brine, dried with sodium sulfate and concentrated to afford 79 mg of 1-benzyl-4-[2-
 (benzyl-methyl-amino)-ethyl]-1,3-dihydro-benzoimidazol-2-one. MS: 372 (M+H)⁺.

10 Step 6[2-(1-Benzyl-2-oxo-2,3-dihydro-1H-benzoimidazol-4-yl)-ethyl]-methyl-
carbamic acid tert-butyl ester

1-Benzyl-4-[2-(benzyl-methyl-amino)-ethyl]-1,3-dihydro-benzoimidazol-2-one
 15 (79 mg) was dissolved in 3 mL of a 1:1 mixture of tetrahydrofuran and ethanol and then
 Pd(OH)₂ was added. The mixture was shaken under 45 psi of H₂ for 24 hours before
 filtering off the catalyst and removing solvent. The crude material was redissolved in 2
 mL of tetrahydrofuran and di-tert-butyl dicarbonate (131 mg) was added. After 16 hours
 the reaction was purified directly by chromatography to afford 31 mg of [2-(1-benzyl-2-
 20 oxo-2,3-dihydro-1H-benzoimidazol-4-yl)-ethyl]-methyl-carbamic acid tert-butyl ester,
 MS: 382 (M+H)⁺.

Step 71-Benzyl-4-(2-methylamino-ethyl)-1,3-dihydro-benzoimidazol-2-one




[2-(1-Benzyl-2-oxo-2,3-dihydro-1H-benzimidazol-4-yl)-ethyl]-methyl-carbamic acid tert-butyl ester (31mg) was dissolved in 2.0 ml of tetrahydrofuran and 100 uL of 2.0 N HCl/EtOH was added. The solution was heated on a steam bath for 30 minutes and then after cooling diethyl ether was slowly added to produce 10 mg of 1-benzyl-4-(2-methylamino-ethyl)-1,3-dihydro-benzimidazol-2-one hydrochloride salt as a colorless powder. MS: 282 (M+H)⁺

Example 6

1-(3-Fluoro-benzyl)-4-piperazin-1-yl-1,3-dihydro-indol-2-one

The synthetic procedures described in this Example were carried out according to the process shown in Scheme I.




SCHEME I

Step 1

4-Bromo-1-(3-fluoro-benzyl)-1H-indole


- 83 -



A solution of 4-bromoindole (3.91g) in 25 ml of EtOH was stirred with KOH (924 mg) for one hour. The solvent was removed under reduced pressure and replaced with 75 mL of acetone. To this solution was added 3-fluorobenzylbromide, and the mixture was stirred at room temperature for 30 minutes before quenching with water. Ethyl acetate was added and the layers separated. After drying with sodium sulfate the organic phase was concentrated and 4-bromo-1-(3-fluoro-benzyl)-1H-indole (4.12 g) was isolated by column chromatography. MS: 305 (M+H)⁺.

Step 2


10 4-[1-(3-Fluoro-benzyl)-1H-indol-4-yl]-piperazine-1-carboxylic acid tert-butyl ester



To a solution of 4-bromo-1-(3-fluoro-benzyl)-1H-indole (750 mg, 25 mmol) in *tert*-butanol was added Boc-piperidine (470 mg, 26 mmol), dicyclohexyl-(2',4',6'-triisopropyl-biphenyl-2-yl)-phosphane (60 mg), potassium carbonate (862 mg), and tris(dibenzylideneacetone)dipalladium (23 mg). The mixture was heated to reflux and stirred for 14 hours. The reaction was cooled and quenched by addition of water, and the resulting mixture was extracted with ethyl acetate. The combined organic phases were dried over Na₂SO₄, filtered, and evaporated to dryness. The resulting crude material was purified by column chromatography to afford 510 mg of 4-[1-(3-fluoro-benzyl)-1H-indol-4-yl]-piperazine-1-carboxylic acid tert-butyl ester. MS: 410 (M+H)⁺.

Step 3


20 4-[3-Bromo-1-(3-fluoro-benzyl)-1H-indol-4-yl]-piperazine-1-carboxylic acid tert-butyl ester



To a solution of 4-[1-(3-fluoro-benzyl)-1H-indol-4-yl]-piperazine-1-carboxylic acid tert-butyl ester (360 mg) in 5 mL of CH_2Cl_2 at 0°C was added N-bromosuccinimide (160 mg). After 30 minutes, water was added and the layers partitioned. The organic phase was dried over sodium sulfate, filtered, and concentrated under reduced pressure. The resulting crude material was purified by column chromatography to afford 4-[3-bromo-1-(3-fluoro-benzyl)-1H-indol-4-yl]-piperazine-1-carboxylic acid tert-butyl ester. 263 mg. MS: 488, 490 ($\text{M}+\text{H}^+$); 510, 512 ($\text{M}+\text{Na}$).

Step 4

4-[1-(3-Fluoro-benzyl)-2-oxo-2,3-dihydro-1H-indol-4-yl]-piperazine-1-carboxylic acid tert-butyl ester




4-[3-Bromo-1-(3-fluoro-benzyl)-1H-indol-4-yl]-piperazine-1-carboxylic acid tert-butyl ester (273 mg) was dissolved in 3 mL of 2-methoxyethanol and 85% phosphoric acid (575 mL) was added. The dark solution was heated to 140°C and stirred for 16 h. The next day the mixture was made basic with 1.0 N NaOH and Et_2O was added. The organic layer was separated, washed with brine, and dried over sodium sulfate. After filtering and removing solvent the crude material was taken up in 3 mL of THF and di-tert-butylidicarbonate (218 mg) was added. The mixture was stirred for 24 hours at room temperature, after which the mixture was partitioned between water and ethyl acetate. The organic layer was dried over sodium sulfate and evaporated under reduced pressure. The resulting residue was purified by column chromatography to provide 115 mg of 4-[1-

(3-fluoro-benzyl)-2,3-dihydro-1H-indol-4-yl]-piperazine-1-carboxylic acid tert-butyl ester, MS: 326 (M-Boc+H)⁺.

Step 5

1-(3-Fluoro-benzyl)-4-piperazin-1-yl-1,3-dihydro-indol-2-one



5

4-[1-(3-Fluoro-benzyl)-2-oxo-2,3-dihydro-1H-indol-4-yl]-piperazine-1-carboxylic acid tert-butyl ester (115 mg) was dissolved in 2 mL of EtOH and 500 μ L of 2.0 N HCl/EtOH was added. The solution was heated to reflux on a steam bath for 30 minutes and then cooled to room temperature. Diethyl ether was added slowly to precipitate 1-(3-Fluoro-benzyl)-4-piperazin-1-yl-1,3-dihydro-indol-2-one as a hydrochloride salt. MS: 326 (M+H)⁺.


10

Example 7

1-(3-Fluoro-benzyl)-3,3-dimethyl-4-piperazin-1-yl-1,3-dihydro-indol-2-one

The synthetic procedures described in this Example were carried out according to the process shown in Scheme J.


15



SCHEME J

Step 1


20 4-[1-(3-Fluoro-benzyl)-3,3-dimethyl-2-oxo-2,3-dihydro-1H-indol-4-yl]-piperazine-1-carboxylic acid tert-butyl ester



4-[1-(3-Fluoro-benzyl)-2-oxo-2,3-dihydro-1H-indol-4-yl]-piperazine-1-carboxylic acid tert-butyl ester (350 mg) was dissolved in 4 mL of DMF and 60% NaH (60 mg) was added. After stirring for 30 minutes methyl iodide was added (1.0 mL), the reaction vessel sealed and the mixture left to stir for 16 hours at room temperature. The reaction was quenched with water and ethyl acetate added. The layers were separated and the organic phase dried with sodium sulfate and concentrated under reduced pressure. The residue was purified by column chromatography to afford 300 mg 4-[1-(3-fluoro-benzyl)-3,3-dimethyl-2-oxo-2,3-dihydro-1H-indol-4-yl]-piperazine-1-carboxylic acid tert-butyl ester, MS: 454 (M+H)⁺.

Step 2

1-(3-Fluoro-benzyl)-3,3-dimethyl-4-piperazin-1-yl-1,3-dihydro-indol-2-one





4-[1-(3-Fluoro-benzyl)-3,3-dimethyl-2-oxo-2,3-dihydro-1H-indol-4-yl]-piperazine-1-carboxylic acid tert-butyl ester (260 mg) was dissolved in 3 mL of EtOH and 0.5 mL of 2.0 N HCl/EtOH was added. After heating for 30 minutes at reflux, the solution was cooled and diethyl ether was added to precipitate out 1-(3-Fluoro-benzyl)-3,3-dimethyl-4-piperazin-1-yl-1,3-dihydro-indol-2-one as a hydrochloride salt (121 mg). MS: 354 (M+H)⁺.

Example 8


1-(3-Fluoro-benzyl)-4-(2-methylamino-ethoxy)-1,3-dihydro-indol-2-one

The synthetic procedures described in this Example were carried out according to the process shown in Scheme K.

**SCHEME K****Step 1****[2-(1H-Indol-4-yloxy)-ethyl]-methyl-carbamic acid tert-butyl ester**

5

To a cooled (0°C) solution of 4-hydroxyindole (2.92 g) and (2-hydroxy-ethyl)-methyl-carbamic acid tert-butyl ester (3.50 g) in 100 mL of THF was added PPh_3 (5.8 g) followed by the dropwise addition of diisopropylazodicarboxylate (4.44 g) over 10 minutes. After 16 h the reaction was worked up in the normal manner and the crude oil was purified by chromatography to afford 4.1 g of [2-(1H-indol-4-yloxy)-ethyl]-methyl-carbamic acid tert-butyl ester. MS: $313 (\text{M}+\text{Na})^+$.


Step 2**[2-[1-(3-Fluoro-benzyl)-1H-indol-4-yloxy]-ethyl]-methyl-carbamic acid tert-butyl ester**

15

[2-(1H-Indol-4-yloxy)-ethyl]-methyl-carbamic acid tert-butyl ester was treated NaOH/EtOH followed by addition of 3-fluorobenzylbromide as described above in Step 1 of Example 5 to afford {2-[1-(3-fluoro-benzyl)-1H-indol-4-yloxy]-ethyl}-methyl-carbamic acid tert-butyl ester. MS: 299 (M-Boc+H)⁺.

5 Step 3


[2-(1-Benzyl-3-bromo-1H-indol-4-yloxy)-ethyl]-methyl-carbamic acid tert-butyl ester



10 {2-[1-(3-Fluoro-benzyl)-1H-indol-4-yloxy]-ethyl}-methyl-carbamic acid tert-butyl ester was treated with N-bromosuccinimide using the procedure of step 3 of Example 6 to provide [2-(1-benzyl-3-bromo-1H-indol-4-yloxy)-ethyl]-methyl-carbamic acid tert-butyl ester.

Step 4


15 [2-(1-Benzyl-2-oxo-2,3-dihydro-1H-indol-4-yloxy)-ethyl]-methyl-carbamic acid tert-butyl ester



20 [2-(1-Benzyl-3-bromo-1H-indol-4-yloxy)-ethyl]-methyl-carbamic acid tert-butyl ester was treated with phosphoric acid using the procedure of step 4 of Example 6 to provide [2-(1-benzyl-2-oxo-2,3-dihydro-1H-indol-4-yloxy)-ethyl]-methyl-carbamic acid tert-butyl ester.

Step 5

1-Benzyl-4-(2-methylamino-ethoxy)-1,3-dihydro-indol-2-one




(1-Benzyl-2-oxo-2,3-dihydro-1H-indol-4-yloxy)-ethyl]-methyl-carbamic acid tert-butyl ester was deprotected using the procedure of step 5 of Example 6 to afford 1-benzyl-4-(2-methylamino-ethoxy)-1,3-dihydro-indol-2-one, MS: 297 (M+H)⁺.

5

Example 9

1-(3-Fluoro-benzyl)-4-piperidin-4-yl-1,3-dihydro-indol-2-one

The synthetic procedures described in this Example were carried out according to the process shown in Scheme L.




10

SCHEME L

Step 1


4-Bromo-1-(3-fluoro-benzyl)-1H-indole-2,3-dione



4-Bromo-1H-indole-2,3-dione (6.38 g) was dissolved in 100 mL of DMF and at 0 °C, and 60% NaH (1.32 g) was added in portions. The mixture was stirred for 20 minutes before adding 3-fluorobenzyl bromide. The solution was stirred for approximately 30 minutes, quenched with water, and partitioned between water and diethyl ether. The organic layer was dried over sodium sulfate and evaporated to dryness under reduced pressure to yield 9.18 g 4-Bromo-1-(3-fluoro-benzyl)-1H-indole-2,3-dione that was sufficiently pure for use in step 2 below. MS: 385 (M+H)⁺.

Step 2


4-[1-(3-Fluoro-benzyl)-2,3-dioxo-2,3-dihydro-1H-indol-4-yl]-3,6-dihydro-2H-pyridine-1-carboxylic acid tert-butyl ester



4-Bromo-1-(3-fluoro-benzyl)-1H-indole-2,3-dione (167 mg) was dissolved in 2.5 mL of dioxane and 4-(4,4,5,5-Tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,6-dihydro-2H-pyridine-1-carboxylic acid tert-butyl ester (154 mg) was added, followed by [1,1'-bis(diphenylphosphino)-ferrocene]dichloropalladium(II), complex with dichloromethane (1:1) (24 mg) and K₂CO₃ (138 mg). The reaction mixture was heated to 80 °C for 18 hours and then cooled and concentrated under reduced pressure. The resulting residue was purified by chromatography to afford 4-[1-(3-Fluoro-benzyl)-2,3-dioxo-2,3-dihydro-1H-indol-4-yl]-3,6-dihydro-2H-pyridine-1-carboxylic acid tert-butyl ester. MS: 337 (M-Boc+H)⁺.

Step 3


4-[1-(3-Fluoro-benzyl)-2-oxo-2,3-dihydro-1H-indol-4-yl]-piperidine-1-carboxylic acid tert-butyl ester



- 5 4-[1-(3-Fluoro-benzyl)-2,3-dioxo-2,3-dihydro-1H-indol-4-yl]-3,6-dihydro-2H-pyridine-1-carboxylic acid tert-butyl ester (930 mg) was dissolved in 17 ml each of hydrazine and ethanol, and the solution was heated at 110 °C for 16 hours. The reaction was cooled and partitioned between water and ethyl acetate, and the organic phase was dried over sodium sulfate and concentrated under reduced pressure. The residue was
- 10 purified by chromatography to afford 146 mg of 4-[1-(3-fluoro-benzyl)-2-oxo-2,3-dihydro-1H-indol-4-yl]-piperidine-1-carboxylic acid tert-butyl ester. MS: 423 (M-H)⁻.

Step 4


1-(3-Fluoro-benzyl)-4-piperidin-4-yl-1,3-dihydro-indol-2-one




- 15 4-[1-(3-Fluoro-benzyl)-2-oxo-2,3-dihydro-1H-indol-4-yl]-piperidine-1-carboxylic acid tert-butyl ester (146 mg) was deprotected as described in step 5 of Example 6 to afford 1-(3-fluoro-benzyl)-4-piperidin-4-yl-1,3-dihydro-indol-2-one hydrochloride as a white powder. MS: 325 (M+H)⁺.

Example 101-(3-Fluoro-benzyl)-3-hydroxy-3-methyl-4-piperazin-1-yl-1,3-dihydro-indol-2-one

The synthetic procedures described in this Example were carried out according to the process shown in Scheme M.

SCHEME MStep 1


4-[1-(3-Fluoro-benzyl)-2,3-dioxo-2,3-dihydro-1H-indol-4-yl]-piperazine-1-carboxylic acid tert-butyl ester



4-Bromo-1-(3-fluoro-benzyl)-1H-indole-2,3-dione (3.05 g) was dissolved in 45 mL of *tert*-butanol and Boc-piperazine (2.04 g), Pd₂(dba)₃ (164 mg), dicyclohexyl-(2',4',6'-triisopropyl-biphenyl-2-yl)-phosphane (217 mg), and potassium carbonate (1.84 g) were added. The mixture was stirred at 120 °C for 4 hours and then cooled to room temperature. The mixture was partitioned between ethyl acetate and water, and the organic layer was separated, dried over sodium sulfate and concentrated under reduced pressure. The residue was purified by chromatography to afford 1.80 g of 4-[1-(3-fluoro-benzyl)-2,3-dioxo-2,3-dihydro-1H-indol-4-yl]-piperazine-1-carboxylic acid tert-butyl ester. MS: 340 (M-Boc+H)⁺.

Step 2


4-[1-(3-Fluoro-benzyl)-3-hydroxy-3-methyl-2-oxo-2,3-dihydro-1H-indol-4-yl]-piperazine-1-carboxylic acid tert-butyl ester



- 5 4-[1-(3-Fluoro-benzyl)-2,3-dioxo-2,3-dihydro-1H-indol-4-yl]-piperazine-1-carboxylic acid tert-butyl ester (330 mg) was dissolved in 4 mL of THF and cooled to -78 °C before adding 1.6 M methyl lithium in hexanes (800 μ L). The solution was stirred and warmed to room temperature, then stirred for an hour at room temperature. The reaction was quenched by addition of water and extracted with ethyl acetate. The combined organic layers were dried over sodium sulfate and evaporated under reduced pressure. The residue was purified by column chromatography afforded 315 mg of 4-[1-(3-fluoro-benzyl)-3-hydroxy-3-methyl-2-oxo-2,3-dihydro-1H-indol-4-yl]-piperazine-1-carboxylic acid tert-butyl ester. MS: 356 (M-Boc+H)⁺.

Step 3


- 15 1-(3-Fluoro-benzyl)-3-hydroxy-3-methyl-4-piperazin-1-yl-1,3-dihydro-indol-2-one



- 20 4-[1-(3-Fluoro-benzyl)-3-hydroxy-3-methyl-2-oxo-2,3-dihydro-1H-indol-4-yl]-piperazine-1-carboxylic acid tert-butyl ester was deprotected in the manner described in step 5 of Example 1 to afford 1-(3-Fluoro-benzyl)-3-hydroxy-3-methyl-4-piperazin-1-yl-1,3-dihydro-indol-2-one (201 mg). MS: 356 (M+H)⁺.

Example 111-Benzyl-4-pyrrolidin-3-yl-1,3-dihydro-benzoimidazol-2-one


The synthetic procedures described in this Example were carried out according to the process shown in Scheme N.



5

SCHEME NStep 1

10

1-Bromo-3-fluoro-2-nitro-benzene


To a suspension of sodium perborate tetrahydrate (135.374 g., 886.4 mmol) in 500 mL acetic acid at 55°C was added dropwise a solution of 2-bromo-6-fluorophenylamine (33.685 g., 177.271 mmol) in 70 mL acetic acid over 1 hour. The reaction mixture was stirred at 55°C for an additional 3 hours, then cooled to 0°C in an ice bath. Insoluble materials were removed by filtration through a plug of celite, which was rinsed

15

with 100 mL acetic acid. The combined acetic acid fractions were added to 3 L ice water with stirring to give a waxy solid which was removed by filtration. The crude solid was dissolved in 250 mL ethyl acetate, washed three times with 200 mL of 10% aqueous hydrogen chloride, 200 mL saturated sodium bicarbonate and 100 mL brine. The solution
5 was concentrated *in vacuo* to give 11. 51 g of 1-bromo-3-fluoro-2-nitro-benzene as a red oil, ^1H NMR (CDCl_3 , 300 MHz) δ : 7.26 (m, 1H), 7.38 (m, 1H), 7.49 (m, 1H)

Step 2


Benzyl-(3-bromo-2-nitro-phenyl)-amine



10 Potassium carbonate (2.58 g, 18.73 mmol) was added to a solution of 1-bromo-3-fluoro-2-nitro-benzene (2.06 g, 9.36 mmol) and benzylamine (1.13 mL, 9.364 mmol) and the resulting suspension was stirred at room temperature for 18 hours. The reaction mixture was poured onto 500 mL of ice water and extracted four times with 100 mL ethyl acetate. The combined organic fractions were dried over sodium sulfate, and
15 concentrated *in vacuo* to give an oil that was purified by recrystallization from 75 mL refluxing ethanol with 3 mL water. 1.83 g of benzyl-(3-bromo-2-nitro-phenyl)-amine was collected by filtration as fine red needles. MS: 308, 309 ($\text{M}+\text{H}$) $^+$.

Step 3

N-1-Benzyl-3-bromo-benzene-1,2-diamine




20

A solution of benzyl-(3-bromo-2-nitro-phenyl)-amine (0.503 g, 1.638 mmol) in 35 mL ethanol was added to a solution of sodium thiosulfite (1.91 g., 10.97 mmol) in 50 mL water at 100°C while stirring. The reaction mixture was heated for 30 minutes at this temperature, and then concentrated *in vacuo* to remove the ethanol. The oil which
25 formed in the resulting water solution was extracted twice with 100 mL dichloromethane. The combined organic fractions were dried over sodium sulfate, then purified by flash

chromatography (2 to 10% gradient ethyl acetate in hexanes) to give 0.327 g. of N-*I*-benzyl-3-bromo-benzene-1,2-diamine as a red oil. MS: 278, 279 (M+H)⁺.

Step 4


5 3-Benzyl-7-bromo-2-oxo-2,3-dihydro-benzoimidazole-1-carboxylic acid tert-butyl ester



A solution of N-*I*-benzyl-3-bromo-benzene-1,2-diamine (0.483 g, 1.742 mmol) in 20 mL dichloromethane was cooled to 0°C under nitrogen. Di-*tert*-butyldicarbonate (1.899 g, 8.712 mmol) was added, followed by 4-dimethylaminopyridine (10 mg, 0.087 mmol), and the reaction mixture was stirred for one hour and allowed to warm to room temperature. The reaction mixture was concentrated in the presence of 0.5 g silica gel, which was submitted to flash chromatography (1 to 10% gradient ethyl acetate in hexanes) to give 0.580 g of 3-benzyl-7-bromo-2-oxo-2,3-dihydro-benzoimidazole-1-carboxylic acid tert-butyl ester as a white solid. MS: 304, 305 (M+H)⁺.

15 Step 5

3-Benzyl-2-oxo-7-(1-triisopropylsilyl-1H-pyrrol-3-yl)-2,3-dihydro-benzoimidazole-1-carboxylic acid tert-butyl ester




To 10 mL of a 9:1 mixture of dimethoxyethane/water was added 3-benzyl-7-bromo-2-oxo-2,3-dihydro-benzoimidazole-1-carboxylic acid tert-butyl ester (0.719 g, 1.783 mmol), 3-boranyl-1-triisopropylsilyl-1H-pyrrole (0.476 g, 1.783 mmol), [1,1'-bis(diphenylphosphino)-ferrocene]dichloropalladium(II) complex with dichloromethane (1:1) (69.7 mg, 0.089 mmol), and cesium carbonate (0.697 g, 2.139 mmol.) The resulting suspension was vacuum-purged with argon and heated to 85°C for 1 hour. The reaction mixture was added to 100 mL of 1:1 water/ethyl acetate and the organic layer was

separated. The combined organic fractions were washed with 100 mL each of water and brine and dried over sodium sulfate. Concentration *in vacuo* gave a crude solid which was purified by flash chromatography (0 to 10% gradient ethyl acetate in hexanes) to give 0.750 g of 3-benzyl-2-oxo-7-(1-triisopropylsilyl-1H-pyrrol-3-yl)-2,3-dihydro-benzoimidazole-1-carboxylic acid tert-butyl ester as a white solid. MS: 446 (M-BOC+H)⁺.

Step 6

3-Benzyl-7-(1-tert-butoxycarbonyl-1H-pyrrol-3-yl)-2-oxo-2,3-dihydro-benzoimidazole-1-carboxylic acid tert-butyl ester



10

To a solution of 3-benzyl-2-oxo-7-(1-triisopropylsilyl-1H-pyrrol-3-yl)-2,3-dihydro-benzoimidazole-1-carboxylic acid tert-butyl ester (0.768 g, 1.40 mmol) in 30 mL 1,4-dioxane was added tetrabutylammonium fluoride (1.548 mL of a 1M solution in tetrahydrofuran, 1.548 mmol) dropwise over 5 minutes. The reaction mixture was stirred for 45 minutes, and was then added to 200 mL of 1:1 water/ethyl acetate. The layers were separated and the aqueous layer was extracted twice with 100 mL ethyl acetate. The combined organic fractions were dried over sodium sulfate and concentrated *in vacuo* and purified by flash chromatography (20 to 40% gradient ethyl acetate in hexanes) to give 0.447 g of crude 3-benzyl-2-oxo-7-(1H-pyrrol-3-yl)-2,3-dihydro-benzoimidazole-1-carboxylic acid tert-butyl ester. This material was dissolved in 7 mL tetrahydrofuran and cooled to 0°C under nitrogen. To this solution was added di-*tert*-butyldicarbonate (0.752 g, 3.45 mmol) and 4-dimethylaminopyridine (14 mg, 0.115 mmol) and stirring was continued for one hour as the reaction mixture was allowed to warm to room temperature. The reaction mixture was concentrated *in vacuo* and the residue was purified by flash chromatography (1 to 10% gradient ethyl acetate in hexanes) to give 0.550 g of 3-benzyl-7-(1-tert-butoxycarbonyl-1H-pyrrol-3-yl)-2-oxo-2,3-dihydro-benzoimidazole-1-carboxylic acid tert-butyl ester, MS: 490 (M + H)⁺.


20

25

Step 7

3-Benzyl-7-(1-tert-butoxycarbonyl-pyrrolidin-3-yl)-2-oxo-2,3-dihydro-benzoimidazole-1-carboxylic acid tert-butyl ester


30



To a Parr vessel flushed with nitrogen was added a 5% dispersion of platinum on charcoal (15 mg) followed by a solution of 3-benzyl-7-(1-tert-butoxycarbonyl-1H-pyrrol-3-yl)-2-oxo-2,3-dihydro-benzoimidazole-1-carboxylic acid tert-butyl ester (0.150 g., 0.307 mmol) in 10 mL methanol. The Parr vessel was vacuum-purged with hydrogen at atmospheric pressure and stirred for 95 hours. The reaction mixture was then vacuum-purged with nitrogen, filtered through a plug of celite to remove catalyst, and concentrated *in vacuo*. The resulting residue was purified by flash chromatography (5 to 10% gradient ethyl acetate in hexanes) to give 80 mg of racemic 3-benzyl-7-(1-tert-butoxycarbonyl-pyrrolidin-3-yl)-2-oxo-2,3-dihydro-benzoimidazole-1-carboxylic acid tert-butyl ester as a clear oil. MS: 516 (M+Na)⁺.

Step 8

1-Benzyl-4-pyrrolidin-3-yl-1,3-dihydro-benzoimidazol-2-one



A solution of 3-benzyl-7-(1-tert-butoxycarbonyl-pyrrolidin-3-yl)-2-oxo-2,3-dihydro-benzoimidazole-1-carboxylic acid tert-butyl ester (80 mg., 0.273 mmol) in 5 mL ethanol was heated to reflux and 2 N ethanolic hydrogen chloride was added (0.5 mL.) The reaction mixture was refluxed for 45 minutes at which point ether (2 mL) was added slowly. A solid precipitate formed on cooling and was removed by filtration to afford 40 mg of 1-benzyl-4-pyrrolidin-3-yl-1,3-dihydro-benzoimidazol-2-one hydrochloride as a tan solid. MS: 294 (M+H)⁺.

Example 12Formulations

Pharmaceutical preparations for delivery by various routes are formulated as shown in the following Tables. "Active ingredient" or "Active compound" as used in the

5 Tables means one or more of the Compounds of Formula I.

Composition for Oral Administration

Ingredient	% wt./wt.
Active ingredient	20.0%
Lactose	79.5%
Magnesium stearate	0.5%

The ingredients are mixed and dispensed into capsules containing about 100 mg

10 each; one capsule would approximate a total daily dosage.

Composition for Oral Administration

Ingredient	% wt./wt.
Active ingredient	20.0%
Magnesium stearate	0.5%
Crosscarmellose sodium	2.0%
Lactose	76.5%
PVP (polyvinylpyrrolidone)	1.0%

The ingredients are combined and granulated using a solvent such as methanol.

15 The formulation is then dried and formed into tablets (containing about 20 mg of active compound) with an appropriate tablet machine.

Composition for Oral Administration

Ingredient	<u>Amount</u>
Active compound	1.0 g
Fumaric acid	0.5 g
Sodium chloride	2.0 g
Methyl paraben	0.15 g
Propyl paraben	0.05 g
Granulated sugar	25.5 g
Sorbitol (70% solution)	12.85 g

Veegum K (Vanderbilt Co.)	1.0 g
Flavoring	0.035 ml
Colorings	0.5 mg
Distilled water	q.s. to 100 ml

The ingredients are mixed to form a suspension for oral administration.

Parenteral Formulation

Ingredient	% wt./wt.
Active ingredient	0.25 g
Sodium Chloride	qs to make isotonic
Water for injection	100 ml

- 5 The active ingredient is dissolved in a portion of the water for injection. A sufficient quantity of sodium chloride is then added with stirring to make the solution isotonic. The solution is made up to weight with the remainder of the water for injection, filtered through a 0.2 micron membrane filter and packaged under sterile conditions.

Suppository Formulation

Ingredient	% wt./wt.
Active ingredient	1.0%
Polyethylene glycol 1000	74.5%
Polyethylene glycol 4000	24.5%

10

The ingredients are melted together and mixed on a steam bath, and poured into molds containing 2.5 g total weight.

Topical Formulation

Ingredients	grams
Active compound	0.2-2
Span 60	2
Tween 60	2
Mineral oil	5
Petrolatum	10
Methyl paraben	0.15
Propyl paraben	0.05
BHA (butylated hydroxy anisole)	0.01
Water	q.s. 100

All of the ingredients, except water, are combined and heated to about 60°C with stirring. A sufficient quantity of water at about 60°C is then added with vigorous stirring to emulsify the ingredients, and water then added q.s. about 100 g.

Nasal Spray Formulations

5 Several aqueous suspensions containing from about 0.025-0.5 percent active compound are prepared as nasal spray formulations. The formulations optionally contain inactive ingredients such as, for example, microcrystalline cellulose, sodium carboxymethylcellulose, dextrose, and the like. Hydrochloric acid may be added to adjust pH. The nasal spray formulations may be delivered via a nasal spray metered pump
10 typically delivering about 50-100 microliters of formulation per actuation. A typical dosing schedule is 2-4 sprays every 4-12 hours.

Example 13

Radioligand binding studies

This example illustrates in vitro radioligand binding studies of compound of
15 formula I.

The binding activity of compounds of this invention in vitro was determined as follows. Duplicate determinations of 5-HT₆ ligand affinity were made by competing for binding of [³H]LSD in cell membranes derived from HEK293 cells stably expressing recombinant human 5-HT₆ receptor. Duplicate determinations of 5-HT_{2A} ligand affinity
20 were made by competing for binding of [³H]Ketanserin (3-(2-(4-(4-fluorobenzoyl)piperidinol)ethyl)-2,4(1H,3H)-quinazolinedione) in cell membranes derived from CHO-K1 cells stably expressing recombinant human 5-HT_{2A} receptor. Membranes were prepared from HEK 293 cell lines by the method described by Monsma et al., Molecular Pharmacology, Vol. 43 pp. 320-327 (1993), and from CHO-K1 cell lines
25 as described by Bonhaus et al., Br J Pharmacol. Jun;115(4):622-8 (1995).

For estimation of affinity at the 5-HT₆ receptor, all determinations were made in assay buffer containing 50 mM Tris-HCl, 10 mM MgSO₄, 0.5 mM EDTA, 1 mM ascorbic acid, pH 7.4 at 37 °C, in a 250 microliter reaction volume. For estimation of affinity at the 5-HT_{2A} receptor all determinations were made in assay buffer containing 50 mM
30 Tris-HCl, 5 mM ascorbic acid, 4 mM CaCl₂, pH 7.4 at 32 °C, in a 250 microliter reaction volume.

Assay tubes containing [³H] LSD or [³H]Ketanserin (5 nM), competing ligand, and membrane were incubated in a shaking water bath for 75 min. at 37 °C (for 5-HT₆) or 60 min. at 32°C (for 5-HT_{2A}), filtered onto Packard GF-B plates (pre-soaked with 0.3%
35 PEI) using a Packard 96 well cell harvester and washed 3 times in ice cold 50 mM Tris-HCl. Bound [³H] LSD or [³H]Ketanserin were determined as radioactive counts per minute using Packard TopCount.

Displacement of [³H]LSD or [³H]Ketanserin from the binding sites was quantified by fitting concentration-binding data to a 4-parameter logistic equation:

$$\text{binding} = \text{basal} + \left(\frac{\text{Bmax} - \text{basal}}{1 + 10^{-\text{Hill}(\log[\text{ligand}] - \log \text{IC}_{50})}} \right)$$

where Hill is the Hill slope, [ligand] is the concentration of competing radioligand and IC₅₀ is the concentration of radioligand producing half-maximal specific binding of radioligand. The specific binding window is the difference between the Bmax and the basal parameters.

Using the procedures of this Example, compounds of Formula I were tested and found to be selective 5-HT₆ antagonists, selective 5-HT_{2A} antagonists, or both. For example, 1-(3,4-Difluoro-benzyl)-4-piperazin-1-yl-1,3-dihydro-indol-2-one exhibited a pKi of approximately 9.48 for the 5-HT_{2A} receptor.

Further test results are shown in the Table below:

#	pKi 5-HT ₆	#	pKi 5-HT ₆
10	9.4	43	9.48
18	8.1	45	8.73
20	8.57	47	9.46
25	8.91	53	8.3
29	9.38	54	9.11
32	8.57	64	9.21
40	9.2	66	8.59


15

While the present invention has been described with reference to the specific embodiments thereof, it should be understood by those skilled in the art that various changes may be made and equivalents may be substituted without departing from the true spirit and scope of the invention. In addition, many modifications may be made to adapt a particular situation, material, composition of matter, process, process step or steps, to the objective spirit and scope of the present invention. All such modifications are intended to be within the scope of the claims appended hereto.

25

Claims:

1. A compound of the formula I:



5 or a pharmaceutically acceptable salt thereof,

wherein:

m is from 0 to 3;

n is 1 or 2;

X is:

- 10 -NR^a-;
 -O-;
 -S-;
 -CR^bR^c-; or
 -C(O)-;

15 wherein:

R^a is hydrogen or alkyl;

R^b is hydrogen, fluoro or alkyl;

R^c is hydrogen, fluoro, alkyl, hydroxy; or alkoxy; or

R^b and R^c together form oxo; or


20 R^b and R^c together with the atom to which they are attached may form a three to six-membered optionally substituted ring that optionally includes a heteroatom selected from O, N and S;

Ar is:

optionally substituted aryl; or

25 optionally substituted heteroaryl;

R¹ is a group of the formula



p is from 1 to 4;

Y is:

- 30 -O-;
 -NR^d-; or

$$-\text{CR}^e\text{R}^f-;$$

wherein R^d, R^e and R^f each independently is hydrogen or alkyl;
each R² is independently:

halo:

5 alkyl;

haloalkyl;

haloalkoxy;

alkoxy;

heteroalkyl;

10 cyano;

$$-(\text{CH}_2)_q-\text{S}(\text{O})_r-\text{R}^g;$$
$$-(\text{CH}_2)_q-\text{C}(=\text{O})-\text{NR}^h\text{R}^i;$$
$$-(\text{CH}_2)_q-\text{SO}_2-\text{NR}^h\text{R}^i;$$
$$-(\text{CH}_2)_q-\text{N}(\text{R}^j)-\text{C}(=\text{O})-\text{R}^k, \text{ or}$$

15 $-(\text{CH}_2)_q-\text{C}(=\text{O})-\text{R}^k;$

wherein:

q is 0 or 1;

r is from 0 to 2; and

R^g, R^h, R^i and R^j each independently is hydrogen or alkyl, and R^k

20 is hydrogen, alkyl, alkoxy or hydroxy;

R³ and R⁴ each independently is hydrogen or alkyl;

R⁵ and R⁶ each independently is hydrogen or alkyl; and

R⁷ and R⁸ each independently is hydrogen or alkyl; or

R⁷ and R⁸ together with the nitrogen to which they are attached may form a

25 four to seven-membered optionally substituted ring that optionally includes an additional heteroatom selected from O, N and S; or

one of R⁷ and R⁸ and one of R⁵ and R⁶ together with the atoms to which they are attached may form a four to seven-membered optionally substituted ring that optionally includes an additional heteroatom selected from O, N and S; or

one of R⁷ and R⁸ together with R^d and the atoms to which they are attached may form a four to seven-membered-ring that optionally includes an additional heteroatom selected from O, N and S; or

one of R⁷ and R⁸ and one of R^e and R^f together with the atoms to which they are attached may form a four to seven-membered optionally substituted ring that

35 optionally includes an additional heteroatom selected from O, N and S.

2. The compound according to claim 1, wherein n is 1.

3. The compound according to claim 1 or 2, wherein R^3 and R^4 are hydrogen.
4. The compound according to any one of claims 1 to 3, wherein Ar is
5 optionally substituted aryl.
5. The compound according to any one of claims 1 to 4, wherein Ar is
optionally substituted phenyl.
- 10 6. The compound according to any one of claims 1 to 5, wherein m is 0 or 1.
7. The compound according to any one of claims 1 to 6, wherein p is 2.
8. The compound according to claim 7, wherein Y is -O-.
- 15 9. The compound according to claim 8, wherein R^5 and R^6 are hydrogen.
10. The compound according to claim 8, wherein one of R^7 and R^8 and one of
 R^5 and R^6 together with the atoms to which they are attached form a four to seven-
20 membered ring.
11. The compound according to claim 7, wherein Y is $-NR^d$.
12. The compound according to claim 11, wherein R^5 and R^6 are hydrogen.
- 25 13. The compound according to claim 12, wherein one of R^7 and R^8 together
with R^d and the atoms to which they are attached form a four to seven-membered ring.
14. The compound according to claim 3, wherein p is 3.
- 30 15. The compound according to claim 14, wherein Y is -O-.
16. The compound according to claim 15, wherein R^5 and R^6 are hydrogen.
- 35 17. The compound according to claim 16, wherein one of R^7 and R^8 and one of
 R^5 and R^6 together with the atoms to which they are attached form a four to seven-
membered ring.


18. The compound according to claim 7, wherein Y is $-\text{CR}^e\text{R}^f-$.

19. The compound according to claim 18, wherein one of R^7 and R^8 and one of R^e and R^f together with the atoms to which they are attached form a five to seven-membered ring.

20. The compound according to any one of claims 1 to 6, wherein R^1 is azepinyl, pyrrolidinyl, piperazinyl, piperidinyl, azepinyl, diazepinyl, aminoalkoxy, azepinyloxy, pyrrolidinyl, piperidinyl, azepinyloxy, or azepinylmethoxy, pyrrolidinylmethoxy, piperidinylmethoxy, azepinylmethoxy, or aminoalkyl.

21. The compound according to any one of claims 1 to 6, wherein:

R^1 is:



; and


$\text{R}^7, \text{R}^8, \text{R}^{10}, \text{R}^{11}, \text{R}^{12}$ and R^{13} each independently is hydrogen or alkyl.

22. The compound according to claim 21, wherein R^{10} , R^{11} , R^{12} and R^{13} are hydrogen.

23. The compound according to claim 21 or 22, wherein Ar is optionally substituted phenyl or isoxazole.

5 24. The compound according to claim 23, wherein the optional substituents on phenyl or isoxazole are selected from one or more halo, alkyl, haloalkyl, haloalkoxy, alkoxy or cyano or mixtures thereof.

25. The compound according to claim 1, wherein said compound is of formula II:



wherein:

s is from 1 to 4;

each R^9 is independently:

halo;

alkyl;

haloalkyl;

haloalkoxy;

alkoxy;

heteroalkyl;

cyano;

$-(CH_2)_q-S(O)_r-R^g$;

$-(CH_2)_q-C(=O)-NR^hR^i$;

$-(CH_2)_q-SO_2-NR^hR^i$;

$-(CH_2)_q-N(R^j)-C(=O)-R^k$, or

$-(CH_2)_q-C(=O)-R^k$;


wherein:

q is 0 or 1;

r is from 0 to 2; and

R^g, R^h, R^i and R^j each independently is hydrogen or alkyl, and R^k is hydrogen, alkyl, alkoxy or hydroxy; and $m, p, X, Y, R^2, R^3, R^4, R^5, R^6, R^7$ and R^8 are as recited in claim 1.


5 26. The compound according to claim 1, wherein said compound is of formula
III:



wherein:

10 t and u each independently is from 1 to 3;
R¹⁰, R¹¹, R¹² and R¹³ each independently is hydrogen or alkyl; and
m, s, X, Y, R², R³, R⁴, R⁸ and R⁹ are as recited in claim 25.


27. The compound according to claim 1, wherein said compound is of the formula IV:



wherein:

p is 2 or 3; and
m, s, X, Y, R^2 , R^3 , R^4 , R^7 , R^8 and R^9 are as recited in claim 25.


28. The compound according to claim 1, wherein said compound is of the formula V:



wherein:

- 5 t and u each independently is from 1 to 3;
 R¹⁰, R¹¹, R¹² and R¹³ each independently is hydrogen or alkyl; and
 m, s, X, Y, R², R³, R⁴, R⁸ and R⁹ are as recited in claim 25.

29. The compound of claim 25, wherein said compound is of the formula VI:



wherein:

- 10 t is from 1 to 3;
 u is from 0 to 3;
 R¹⁰, R¹¹, R¹² and R¹³ each independently is hydrogen or alkyl; and
 15 m, s, X, Y, R², R³, R⁴, R⁸ and R⁹ are as recited in claim 25.

30. The compound of claim 1, wherein said compound is selected from:

- 1-Benzyl-4-piperazin-1-yl-1,3-dihydro-benzoimidazol-2-one;
 1-Benzyl-4-(2-methylamino-ethoxy)-1,3-dihydro-benzoimidazol-2-one;

- 1-Benzyl-3-methyl-4-(2-methylamino-ethoxy)-1,3-dihydro-benzoimidazol-2-one;
- 4-(Azetidin-3-ylmethoxy)-1-benzyl-1,3-dihydro-benzoimidazol-2-one;
- 1-Benzyl-4-(3-dimethylamino-propoxy)-1,3-dihydro-benzoimidazol-2-one;
- 5 1-Benzyl-4-(pyrrolidin-3-ylmethoxy)-1,3-dihydro-benzoimidazol-2-one;
- 1-Benzyl-4-(piperidin-4-yloxy)-1,3-dihydro-benzoimidazol-2-one;
- 4-(Azetidin-3-yloxy)-1-benzyl-1,3-dihydro-benzoimidazol-2-one;
- (R)- 1-Benzyl-4-(pyrrolidin-2-ylmethoxy)-1,3-dihydro-benzoimidazol-2-one;
- 10 1-Benzyl-4-piperazin-1-yl-1,3-dihydro-benzoimidazol-2-one;
- 1-Benzyl-4-piperazin-1-yl-1,3-dihydro-indol-2-one;
- 1-(3-Fluoro-benzyl)-4-piperazin-1-yl-1,3-dihydro-indol-2-one;
- 1-(3-Fluoro-benzyl)-4-piperazin-1-yl-1H-indole-2,3-dione;
- 4-[1,4]Diazepan-1-yl-1-(3-fluoro-benzyl)-1H-indole-2,3-dione;
- 15 1-(3-Fluoro-benzyl)-3,3-dimethyl-4-piperazin-1-yl-1,3-dihydro-indol-2-one;
- 1-(3-fluoro-benzyl)-4-(2-methylamino-ethoxy)-1,3-dihydro-indol-2-one;
- 1-(3-Fluoro-benzyl)-3-hydroxy-3-methyl-4-piperazin-1-yl-1,3-dihydro-indol-2-one;
- 4-[1,4]Diazepan-1-yl-1-(3-fluoro-benzyl)-1,3-dihydro-indol-2-one;
- 20 1-(3-Fluoro-benzyl)-4-piperidin-4-yl-1,3-dihydro-indol-2-one;
- 1-Benzyl-6-chloro-4-(2-methylamino-ethoxy)-1,3-dihydro-benzoimidazol-2-one;
- 1-Benzyl-6-fluoro-4-(2-methylamino-ethoxy)-1,3-dihydro-benzoimidazol-2-one;
- 25 1-(2-Fluoro-benzyl)-4-(2-methylamino-ethoxy)-1,3-dihydro-benzoimidazol-2-one;
- 1-(2-Chloro-benzyl)-4-(2-methylamino-ethoxy)-1,3-dihydro-benzoimidazol-2-one;
- 1-(3-Fluoro-benzyl)-4-(2-methylamino-ethoxy)-1,3-dihydro-benzoimidazol-2-one;
- 30 2-one;
- 1-(3-Chloro-benzyl)-4-(2-methylamino-ethoxy)-1,3-dihydro-benzoimidazol-2-one
- 1-(2,3-difluoro-benzyl)-4-(2-methylamino-ethoxy)-1,3-dihydro-benzoimidazol-2-one;
- 35 1-Benzyl-4-(2-methylamino-ethyl)-1,3-dihydro-benzoimidazol-2-one;
- 1-Benzyl-6-fluoro-4-piperazin-1-yl-1,3-dihydro-benzoimidazol-2-one;


- 1-(3-Fluoro-benzyl)-6-fluoro-4-piperazin-1-yl-1,3-dihydro-benzoimidazol-2-one;
- 1-(4-Fluoro-benzyl)-6-fluoro-4-(2-methylamino-ethoxy)-1,3-dihydro-benzoimidazol-2-one;
- 5 1-(4-Fluoro-benzyl)-4-(2-methylamino-ethoxy)-1,3-dihydro-benzoimidazol-2-one;
- 1-Benzyl-4-pyrrolidin-3-yl-1,3-dihydro-benzoimidazol-2-one;
- 1-Benzyl-6-fluoro-4-pyrrolidin-3-yl-1,3-dihydro-benzoimidazol-2-one;
- 1-(3-Fluoro-benzyl)-6-fluoro-4-(2-methylamino-ethoxy)-1,3-dihydro-
- 10 benzoimidazol-2-one;
- 1-(3-Fluoro-benzyl)-4-piperidin-4-yl-1,3-dihydro-benzoimidazol-2-one;
- 1-(3-Fluoro-benzyl)-4-pyrrolidin-3-yl-1,3-dihydro-benzoimidazol-2-one;
- 1-Benzyl-6-methyl-4-piperidin-4-yl-1,3-dihydro-benzoimidazol-2-one;
- 1-(2,3-difluoro-benzyl)-4-pyrrolidin-3-yl-1,3-dihydro-benzoimidazol-2-
- 15 one;
- 1-(3-Fluoro-benzyl)-6-methyl-4-piperidin-4-yl-1,3-dihydro-benzoimidazol-2-one;
- 6-Fluoro-1-(3-fluoro-benzyl)-4-piperidin-4-yl-1,3-dihydro-benzoimidazol-2-one;
- 20 (*S*)-1-(1-Phenyl-ethyl)-4-piperazin-1-yl-1,3-dihydro-benzoimidazol-2-one;
- (*R*)-1-(1-Phenyl-ethyl)-4-piperazin-1-yl-1,3-dihydro-benzoimidazol-2-one;
- 7-Chloro-1-(3-fluoro-benzyl)-4-piperazin-1-yl-1,3-dihydro-benzoimidazol-2-one;
- 5-Chloro-1-(3-fluoro-benzyl)-4-piperazin-1-yl-1,3-dihydro-benzoimidazol-
- 25 2-one;
- 1-Benzyl-3-methyl-4-piperazin-1-yl-1,3-dihydro-benzoimidazol-2-one;
- 5,6-Difluoro-1-(3-fluoro-benzyl)-4-piperazin-1-yl-1,3-dihydro-benzoimidazol-2-one;
- 1-(1-Methyl-1-phenyl-ethyl)-4-piperazin-1-yl-1,3-dihydro-benzoimidazol-2-
- 30 one;
- 1-[1-(3-Fluoro-phenyl)-ethyl]-4-piperazin-1-yl-1,3-dihydro-benzoimidazol-2-one;
- 1-(2,3-Difluoro-benzyl)-6-fluoro-4-piperazin-1-yl-1,3-dihydro-benzoimidazol-2-one;
- 35 6-Fluoro-1-(5-methyl-isoxazol-3-ylmethyl)-4-piperidin-4-yl-1,3-dihydro-benzoimidazol-2-one;
- 1-(2,3-Difluoro-benzyl)-4-piperazin-1-yl-1,3-dihydro-benzoimidazol-2-one;

6-Fluoro-1-(1-methyl-1-phenyl-ethyl)-4-piperazin-1-yl-1,3-dihydro-benzoimidazol-2-one;

- 1-(3-Fluoro-benzyl)-4-pyrrolidin-3-yl-1,3-dihydro-indol-2-one;
 1-(2,3-Difluoro-benzyl)-4-piperazin-1-yl-1H-indole-2,3-dione;
 5 1-(2-Fluoro-benzyl)-4-piperazin-1-yl-1H-indole-2,3-dione;
 1-(2,3-Difluoro-benzyl)-4-piperazin-1-yl-1,3-dihydro-indol-2-one;
 1-(3,4-Difluoro-benzyl)-4-piperazin-1-yl-1H-indole-2,3-dione;
 1-(3,5-Difluoro-benzyl)-4-piperazin-1-yl-1H-indole-2,3-dione;
 1-(3,5-Difluoro-benzyl)-4-piperazin-1-yl-1,3-dihydro-indol-2-one;
 10 1-(3,4-Difluoro-benzyl)-4-piperazin-1-yl-1,3-dihydro-indol-2-one;
 1-(3-Chloro-benzyl)-4-piperazin-1-yl-1H-indole-2,3-dione;
 1-(2,5-Difluoro-benzyl)-4-piperazin-1-yl-1H-indole-2,3-dione;
 1-(2-Fluoro-benzyl)-4-piperazin-1-yl-1,3-dihydro-indol-2-one;
 1-(3-Chloro-benzyl)-4-piperazin-1-yl-1,3-dihydro-indol-2-one;
 15 5-Chloro-1-(3-fluoro-benzyl)-4-piperazin-1-yl-1,3-dihydro-indol-2-one;
 7-Chloro-1-(3-fluoro-benzyl)-4-piperazin-1-yl-1,3-dihydro-indol-2-one; and
 5,7-Dichloro-1-(3-fluoro-benzyl)-4-piperazin-1-yl-1,3-dihydro-indol-2-one.


31. A method for preparing a compound according to claim 1 wherein X is -NH-,
 20 said method comprising:

reacting a compound of formula f:




with phosgene or a phosgene equivalent,
 to provide a compound of formula g:

- 113 -




wherein m, n, p, Y, Ar, R², R³, R⁴, R⁵, R⁶, R⁷ and R⁸ are as recited in claim 1.

32. A method of making a compound according to claim 21, said method comprising:
- reacting a compound of formula n




- with a pyrrol boronic acid compound of formula o




to form a compound of formula p;

- 114 -



hydrogenating compound **p** to form compound **q**



wherein L is a leaving group, and m, n, X, Ar, R², R³, R⁴, R⁹, R¹⁰ and R¹¹ are as
 5 recited in claim 21.

33. A compound of formula I to VI according to any one of claims 1 to 30, when
 manufactured by a process according to claim 31 or 32.

34. A pharmaceutical composition comprising one or more compounds as
 10 claimed in any one of claims 1 to 30 in admixture with a pharmaceutically acceptable
 carrier.

35. A compound of formula I to VI according to any one of claims 1 to 30 as well
 as its pharmaceutically acceptable salts for the use in the treatment or prevention of
 15 diseases.

36. Use of a compound of formula I to VI according to any of claims 1 to
 30 as well as its pharmaceutically acceptable salts for the manufacture of a
 medicament for the treatment of a central nervous system disease.

20

37. Use according to claim 36, wherein the central nervous disease is selected
 from the group consisting of psychoses, schizophrenia, manic depressions, neurological
 disorders, memory disorders, attention deficit disorder, Parkinson's disease, amyotrophic
 lateral sclerosis, Alzheimer's disease, food uptake disorders, and Huntington's disease.

25

38. The invention as described hereinabove.
