



INHIBITORS OF COGNITIVE DECLINE

This application claims benefit of priority to US provisional patent application serial no. 61/167,984 filed on April 09, 2009, US provisional patent application serial no. 61/308,667 filed on February 26, 2010, and US provisional patent application serial no. 61/309,091 filed on March 01, 2010, each of which is hereby incorporated by reference in its entirety.

SUMMARY

The present invention provides, *inter alia*, compounds of Formula I, II, or III:





or pharmaceutically acceptable salts, wherein constituent members are provided below.

The present invention further provides pharmaceutical compositions comprising a
 5 compound of Formula I, II, or III, or pharmaceutically acceptable salt thereof, and at least one pharmaceutically acceptable carrier.

The present invention further provides methods of inhibiting, treating, and/or abating cognitive decline and/or Alzheimer's disease with a compound of Formula I, II, or III, or pharmaceutically acceptable salt of the same.

10 The present invention further provides methods of inhibiting, treating, or abatement of cognitive decline with a compound of Formula I, II, or III, or pharmaceutically acceptable salt of the same.

The present invention further provides methods of inhibiting, treating, or abatement of one or more of amyloid production, amyloid assembly, amyloid aggregation, amyloid binding
 15 (to cells in the brain such as neuron cells), the activity/effect of Abeta oligomers on neurons, and amyloid deposition (on cells in the brain such as neuron cells) with a compound of Formula I, II, or III, or pharmaceutically acceptable salt of the same.

The present invention further provides compounds of Formula I, II, or III, or pharmaceutically acceptable salts thereof, for use in therapy.

20 The present invention further provides use of the compounds of Formula I, II, or III, or pharmaceutically acceptable salts thereof, for the manufacture/preparation of a medicament for use in therapy.

In some embodiments, methods for preparation of compounds useful for inhibiting,

treating, or abatement of cognitive decline are provided. In a method called "chemical conditioning", certain compounds of the present invention are derived from naturally occurring compounds, such as those found in medicinal plants, like ginger. The chemical conditioning process described herein is applicable to a large variety of biological extracts and may be used to create compound arrays for screening for potential new drug candidates. Further, in general, compounds derived by the chemical conditioning process are chemically stable and structurally diverse, and good candidates for use in drug screenings for pharmaceutical activity. In some embodiments, compounds derived from ginger oil are provided. According to some embodiments of the invention, compounds derived from ginger oil by the chemical conditioning process described herein are provided. In another embodiment, the invention provides a method of preparing an array of chemical compounds from ginger oil.

In some embodiments, the compounds of present invention inhibit, treat, or abate (partially inhibit) binding of the amyloid (including Abeta oligomers) to neurons (such as neurons in the brain) and are useful for the inhibition, treatment, and abatement of cognitive decline and/or Alzheimer's disease. In some embodiments, the compounds of present invention inhibit, treat, or abate one or more of amyloid aggregation, amyloid binding, and amyloid deposition. In some embodiments, the compounds of present invention inhibit, treat, or abate amyloid aggregation. In some embodiments, the compounds of present invention inhibit, treat, or abate amyloid binding. In some embodiments, the compounds of present invention inhibit, treat, or abate amyloid deposition. In some embodiments, the compounds of present invention inhibit, treat, or abate the activity/effect of Abeta oligomers on neurons. In some embodiments, the compounds show activity in a beta-secretase assay and are potentially useful for the inhibition, treatment, and abatement of cognitive decline and Alzheimer's disease. In some embodiments the derivative of ginger oil is a compound in purified and isolated form (for example, with a purity of greater than 80%, 85%, 90%, 95%, 98%, or 99% by weight). The compounds and methods described herein may be used to treat one or more symptoms of cognitive decline and/or Alzheimer's disease such as memory loss, confusion, impaired judgment, personality changes, disorientation, and loss of language skills. Further, the compounds and methods described herein may be useful in inhibiting, treating, and/or abating cognitive decline and/or Alzheimer's disease by restoring long term potentiation, and/or inhibiting, treating, or abatement of one or both of neurodegeneration and general amyloidosis, more specifically, by inhibiting, treating, or abatement of one or more of amyloid production, amyloid assembly, amyloid aggregation, amyloid binding, and amyloid deposition.

DESCRIPTION OF DRAWINGS

Figure 1 shows results of an MTT assay in the presence and absence of a processed product of amyloid precursor protein.

Figure 2 shows inhibition of processed product of amyloid precursor protein-mediated membrane trafficking effect by Compound Example 2.

Figure 3 shows Compound Example 2 inhibiting the memory loss effects of a processed product of amyloid precursor protein.

Figure 4 shows Compound Example 2 inhibiting the membrane trafficking effects of Abeta assemblies isolated from AD patients.

DETAILED DESCRIPTION

Cognitive decline, such as memory loss, confusion, impaired judgment, personality changes, disorientation, and loss of language skills occurs in much of the population as they age, in varying degree. The most common, severe and irreversible form of cognitive decline is Alzheimer's disease, which, at present, is always fatal.

The symptoms of cognitive decline and Alzheimer's disease are thought to stem from the formation of amyloid plaques and neurofibrillary tangles, which are thought to contribute to the degradation of the neurons (nerve cells) in the brain and the subsequent onset of symptoms. Amyloid is a general term for protein fragments that the body produces normally. Beta-amyloid is a fragment of a protein that is snipped from another protein called amyloid precursor protein (APP). In a healthy brain, beta-amyloid protein fragments are broken down and eliminated. In individuals with Alzheimer's disease and other forms of cognitive decline, the fragments accumulate to form hard, insoluble plaques. Neurofibrillary tangles are insoluble twisted fibers that are found inside of the brain's cells. The protein contained in neurofibrillary tangles, i.e., the tau protein, forms a microtubule, which helps transport nutrients and other important substances from one part of the nerve cell to another. In Alzheimer's disease the tau protein is abnormal and the microtubule structures collapse.

Beta-secretase is the enzyme in the human brain responsible for the production of Beta-amyloid, the pathogenic substance responsible for the formation of brain plaques and tangles in the Alzheimer's diseased brain. Beta-amyloid and its oligomers (beta-amyloid oligomers or Abeta oligomers) are also believed to be responsible for early cognitive decline in the pre-Alzheimer's diseased brain. Inhibition of beta-secretase would be expected to lessen beta-amyloid burden in the brain and thus slow cognitive decline, block the formation of amyloid oligomers, the production of plaques and tangles, halt neurodegeneration, and to potentially treat mild cognitive impairment and more serious forms of cognitive impairment


such as Alzheimer's disease.

The gingerols are a series of natural small molecules isolated from ginger, *Zingiber officinale*, and are classified according to their alkyl chain length *e.g.*, [6]-gingerol, [8]-gingerol. Gingerols are known to be relatively unstable under both chemical and biological conditions, forming inactive substances. For example, the beta-hydroxycarbonyl function of the gingerols is vulnerable to oxidation or dehydration to form inactive products, and the gingerols are particularly prone to rapid dehydration under acidic conditions, such that even the pure substance is difficult to store for long periods. Accordingly, simple oral dosing of the gingerols for medicinal action might not be possible due to the acidic environment of the stomach and upper intestinal tract. Further, chemical and biological instability is also likely to be a serious problem for intravenous doses. Accordingly, there is strong need to discover inhibitors of cognitive decline, and in particular, compounds that are useful in the treatment and abatement of cognitive decline and Alzheimer's disease, by methods such as inhibiting amyloid (including Abeta oligomers) production, amyloid (including Abeta oligomers) aggregation, and/or amyloid (including Abeta oligomers) deposition (*i.e.*, plaquing), inhibiting neurodegeneration, and/or restoring long term potentiation, and/or inhibiting the activity/effect of Abeta oligomers on neurons. There is also a need for inhibitors of cognitive decline that are chemically and biologically stable.

Plants have attracted relatively little attention as potentially valuable resources for drug discovery in the area of cognitive decline and Alzheimer's disease. The use of plant extracts to produce unnatural derivatives of compounds of medicinal interest is not generally used. Accordingly, there is also a need for a method of producing compounds of medicinal interest from plant extracts and extracts from other biological sources. In particular, there is also a need to produce and identify compounds derived from plant extracts that are useful in the treatment and abatement of cognitive decline and Alzheimer's disease.

The compounds, compositions, and methods described herein are directed toward these needs and other ends.

Embodiments of the present invention provides, *inter alia*, compounds of Formula I:




6

I

or pharmaceutically acceptable salts thereof wherein:

R¹ is selected from (A1) and (A2):



R², R³, R⁴, R⁵, and R⁶ are each, independently, selected from H, OH, C₁₋₆ alkoxy, C₁₋₆ haloalkoxy, halo, CN, NO₂, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₃₋₇ cycloalkyl, NH₂, NH(C₁₋₄ alkyl), NH(C₃₋₇ cycloalkyl), N(C₁₋₄ alkyl)₂, NHC(O)(C₁₋₄ alkyl), SH, S(C₁₋₆ alkyl), C(O)OR^a, C(O)R^b, C(O)NR^cR^d, OC(O)R^b, OC(O)NR^cR^d, NR^cR^d, NR^cC(O)R^b, NR^cC(O)OR^a, NR^cS(O)₂R^b, NR^cS(O)₂NR^cR^d, S(O)R^b, S(O)₂R^b, and S(O)₂NR^cR^d;

R⁷ is H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, or C₃₋₇ cycloalkyl;

R⁸ is C₁₋₆ alkyl, C₁₋₆ haloalkyl, or C₃₋₇ cycloalkyl;

R⁹ is H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, or C₃₋₇ cycloalkyl;

R¹⁰ is H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, or C₃₋₇ cycloalkyl;

R¹¹ is H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, or C₃₋₇ cycloalkyl;

R¹², R¹³, R¹⁴, R¹⁵, and R¹⁶ are each, independently, selected from H, OH, C₁₋₆ alkoxy, C₁₋₆ haloalkoxy, halo, CN, NO₂, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₃₋₇ cycloalkyl, NH₂, NH(C₁₋₄ alkyl), NH(C₃₋₇ cycloalkyl), N(C₁₋₄ alkyl)₂, NHC(O)(C₁₋₄ alkyl), SH, S(C₁₋₆ alkyl), C(O)OR^{a1}, C(O)R^{b1}, C(O)NR^{c1}R^{d1}, OC(O)R^{b1}, OC(O)NR^{c1}R^{d1}, NR^{c1}R^{d1}, NR^{c1}C(O)R^{b1}, NR^{c1}C(O)OR^{a1}, NR^{c1}S(O)₂R^{b1}, NR^{c1}S(O)₂NR^{c1}R^{d1}, S(O)R^{b1}, S(O)₂R^{b1}, and S(O)₂NR^{c1}R^{d1};

each R^a is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, heterocycloalkylalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl and heterocycloalkyl, wherein each of the C₁₋₆ alkyl, C₁₋₆ haloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, heterocycloalkylalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl and heterocycloalkyl is optionally substituted with 1, 2, 3, 4, or 5

substituents independently selected from OH, CN, amino, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkoxy, aryl, arylalkyl, heteroaryl, heteroarylalkyl, cycloalkyl and heterocycloalkyl;

each R^b is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, arylalkyl,

- 5 heteroarylalkyl, cycloalkylalkyl, heterocycloalkylalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl and heterocycloalkylalkyl, wherein each of the C₁₋₆ alkyl, C₁₋₆ haloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, heterocycloalkylalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl and heterocycloalkylalkyl is
10 optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from OH, amino, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkoxy, aryl, arylalkyl, heteroaryl, heteroarylalkyl, cycloalkyl and heterocycloalkyl;

R^c and R^d are independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, heterocycloalkylalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl,

- 15 heteroaryl, cycloalkyl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl and heterocycloalkylalkyl, wherein each of the C₁₋₆ alkyl, C₁₋₆ haloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, heterocycloalkylalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, heteroaryl, cycloalkyl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl and heterocycloalkylalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from OH,
20 amino, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkoxy, aryl, arylalkyl, heteroaryl, heteroarylalkyl, cycloalkyl and heterocycloalkyl;

or R^c and R^d together with the N atom to which they are attached form a 4-, 5-, 6- or 7-membered heterocycloalkyl group that is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from OH, amino, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkoxy, aryl, arylalkyl, heteroaryl, heteroarylalkyl, cycloalkyl, and
25 heterocycloalkyl;

each R^{a1} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, arylalkyl,

- heteroarylalkyl, cycloalkylalkyl, heterocycloalkylalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl and heterocycloalkyl, wherein each of the C₁₋₆ alkyl, C₁₋₆ haloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, heterocycloalkylalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl and heterocycloalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from OH, CN, amino, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkoxy, aryl, arylalkyl, heteroaryl, heteroarylalkyl, cycloalkyl, and
30 heterocycloalkyl;

each R^{b1} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, heterocycloalkylalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl and heterocycloalkylalkyl, wherein each of the C₁₋₆ alkyl, C₁₋₆ haloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, heterocycloalkylalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl and heterocycloalkylalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from OH, amino, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkoxy, aryl, arylalkyl, heteroaryl, heteroarylalkyl, cycloalkyl, and heterocycloalkyl;

R^{c1} and R^{d2} are independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, heterocycloalkylalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, heteroaryl, cycloalkyl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl and heterocycloalkylalkyl, wherein each of the C₁₋₆ alkyl, C₁₋₆ haloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, heterocycloalkylalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, heteroaryl, cycloalkyl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl and heterocycloalkylalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from OH, amino, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkoxy, aryl, arylalkyl, heteroaryl, heteroarylalkyl, cycloalkyl, and heterocycloalkyl;

or R^{c1} and R^{d1} together with the N atom to which they are attached form a 4-, 5-, 6- or 7-membered heterocycloalkyl group that is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from OH, amino, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkoxy, aryl, arylalkyl, heteroaryl, heteroarylalkyl, cycloalkyl, and heterocycloalkyl; and

m is 0, 1, or 2.

In some embodiments, when R¹ is a moiety of (A1), then two of R², R³, R⁴, R⁵, and R⁶ are independently selected from OH, C₁₋₆ alkoxy, and C₁₋₆ haloalkoxy.

In some embodiments, when R¹ is a moiety of (A1), then at least one of R¹², R¹³, R¹⁴, R¹⁵, and R¹⁶ is other than H.

In some embodiments, two of R², R³, R⁴, R⁵, and R⁶ are independently selected from OH, C₁₋₆ alkoxy, and C₁₋₆ haloalkoxy. In some further embodiments, each of the rest of R², R³, R⁴, R⁵, and R⁶ is H.

In some embodiments, R², R³, R⁴, R⁵, and R⁶ are each, independently, selected from H, OH, C₁₋₆ alkoxy, C₁₋₆ haloalkoxy, halo, CN, NO₂, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₃₋₇ cycloalkyl, NH₂, NH(C₁₋₄ alkyl), NH(C₃₋₇ cycloalkyl), N(C₁₋₄ alkyl)₂, NHC(O)(C₁₋₄ alkyl), SH, S(C₁₋₆ alkyl), C(O)OH, C(O)O(C₁₋₄ alkyl), C(O)(C₁₋₄ alkyl), and C(O)NH(C₁₋₄ alkyl).

In some embodiments, one of R^2 , R^3 , R^4 , R^5 , and R^6 is OH; and one of R^2 , R^3 , R^4 , R^5 , and R^6 is OH, C_{1-6} alkoxy, or C_{1-6} haloalkoxy. In some further embodiments, each of the rest of R^2 , R^3 , R^4 , R^5 , and R^6 is H.

In some embodiments, one of R^2 , R^3 , R^4 , R^5 , and R^6 is OH; and one of R^2 , R^3 , R^4 , R^5 , and R^6 is C_{1-3} alkoxy or C_{1-3} haloalkoxy (In some further embodiments, each of the rest of R^2 , R^3 , R^4 , R^5 , and R^6 is H.). In some further embodiments, one of R^2 , R^3 , R^4 , R^5 , and R^6 is OH; and one of R^2 , R^3 , R^4 , R^5 , and R^6 is methoxy or trihalomethoxy (In some further embodiments, each of the rest of R^2 , R^3 , R^4 , R^5 , and R^6 is H.). In still further embodiments, one of R^2 , R^3 , R^4 , R^5 , and R^6 is OH; and one of R^2 , R^3 , R^4 , R^5 , and R^6 is methoxy (In some further embodiments, each of the rest of R^2 , R^3 , R^4 , R^5 , and R^6 is H.).

In some embodiments, R^4 is OH; and R^5 is methoxy. In some further embodiments, R^4 is OH; R^5 is methoxy; and R^2 , R^3 , and R^6 are each H.

In some embodiments, R^7 is H or C_{1-6} alkyl. In some further embodiments, R^7 is H or C_{1-3} alkyl.

In some embodiments, R^7 is C_{1-3} alkyl. In some further embodiments, R^7 is methyl or ethyl. In still further embodiments, R^7 is methyl.

In some embodiments, R^7 is H.

In some embodiments, R^8 is C_{1-6} alkyl. In some further embodiments, R^8 is C_{1-3} alkyl. In still further embodiments, R^8 is methyl.

In some embodiments, R^9 is H or C_{1-6} alkyl. In some further embodiments, R^9 is H or C_{1-3} alkyl.

In some embodiments, R^9 is H.

In some embodiments, R^9 is C_{1-3} alkyl.

In some embodiments, R^{10} is H or C_{1-6} alkyl. In some further embodiments, R^{10} is H or C_{1-3} alkyl. In still further embodiments, R^{10} is H. In other embodiments, R^{10} is C_{1-3} alkyl.

In some embodiments, R^{11} is H or C_{1-6} alkyl. In some further embodiments, R^{11} is H or C_{1-3} alkyl. In still further embodiments, R^{11} is H. In other embodiments, R^{11} is C_{1-3} alkyl.

In some embodiments, at least one of R^{12} , R^{13} , R^{14} , R^{15} , and R^{16} is other than H.

In some embodiments, R^{12} , R^{13} , R^{14} , R^{15} , and R^{16} are each, independently, selected from H, OH, C_{1-6} alkoxy, C_{1-6} haloalkoxy, halo, CN, NO_2 , C_{1-6} alkyl, C_{1-6} haloalkyl, C_{3-7} cycloalkyl, NH_2 , $NH(C_{1-4}$ alkyl), $NH(C_{3-7}$ cycloalkyl), $N(C_{1-4}$ alkyl)₂, $NHC(O)(C_{1-4}$ alkyl), SH, $S(C_{1-6}$ alkyl), $C(O)OH$, $C(O)O(C_{1-4}$ alkyl), $C(O)(C_{1-4}$ alkyl), and $C(O)NH(C_{1-4}$ alkyl).

In some embodiments, R^{12} , R^{13} , R^{14} , R^{15} , and R^{16} are each, independently, selected from H, halo, CN, NO_2 , C_{1-6} alkyl, C_{1-6} haloalkyl, C_{3-7} cycloalkyl, $C(O)O(C_{1-4}$ alkyl), $C(O)(C_{1-4}$ alkyl), and $C(O)NH(C_{1-4}$ alkyl).

In some embodiments, at least one of R^{12} , R^{13} , R^{14} , R^{15} , and R^{16} is selected from halo, CN, NO_2 , C_{1-6} haloalkyl, $C(O)O(C_{1-4}$ alkyl), $C(O)(C_{1-4}$ alkyl), and $C(O)NH(C_{1-4}$ alkyl).

In some embodiments, at least one of R^{12} , R^{13} , R^{14} , R^{15} , and R^{16} is selected from halo and C_{1-6} haloalkyl. In some further embodiments, at least one of R^{12} , R^{13} , R^{14} , R^{15} , and R^{16} is selected from halo and C_{1-6} haloalkyl, and each of the rest is of R^{12} , R^{13} , R^{14} , R^{15} , and R^{16} is H. In yet further embodiments, one or two of R^{12} , R^{13} , R^{14} , R^{15} , and R^{16} are selected from halo and C_{1-6} haloalkyl, and each of the rest is of R^{12} , R^{13} , R^{14} , R^{15} , and R^{16} is H. In still further embodiments, one of R^{12} , R^{13} , R^{14} , R^{15} , and R^{16} is selected from halo and C_{1-6} haloalkyl, and each of the rest is of R^{12} , R^{13} , R^{14} , R^{15} , and R^{16} is H.

In some embodiments, R^{14} is halo or C_{1-6} haloalkyl (In some further embodiments, each of of R^{12} , R^{13} , R^{15} , and R^{16} is H.). In some further embodiments, R^{14} is halo or C_{1-3} haloalkyl (In some further embodiments, each of of R^{12} , R^{13} , R^{15} , and R^{16} is H.). In still further embodiments, R^{14} is halo or C_1 haloalkyl (In some further embodiments, each of of R^{12} , R^{13} , R^{15} , and R^{16} is H.).

In some embodiments, R^{14} is halo (In some further embodiments, each of of R^{12} , R^{13} , R^{15} , and R^{16} is H.). In some embodiments, R^{14} is Cl or F. In some embodiments, R^{14} is Cl. In some embodiments, R^{14} is F.

In some embodiments, R^{14} is C_{1-6} haloalkyl (In some further embodiments, each of of R^{12} , R^{13} , R^{15} , and R^{16} is H.). In some further embodiments, R^{14} is C_{1-3} haloalkyl. In still further embodiments, R^{14} is C_1 haloalkyl. In yet further embodiments, R^{14} is CF_3 .

In some embodiments, R^{15} is halo or C_{1-6} haloalkyl (In some further embodiments, each of of R^{12} , R^{13} , R^{14} , and R^{16} is H.). In some further embodiments, R^{15} is halo or C_{1-3} haloalkyl (In some further embodiments, each of of R^{12} , R^{13} , R^{14} , and R^{16} is H.). In still further embodiments, R^{15} is halo or C_1 haloalkyl (In some further embodiments, each of of R^{12} , R^{13} , R^{14} , and R^{16} is H.).

In some embodiments, R^{15} is halo. In some embodiments, R^{15} is Cl or F. In some embodiments, R^{15} is Cl. In some embodiments, R^{15} is F.


In some embodiments, R^{15} is C_{1-6} haloalkyl. In some further embodiments, R^{15} is C_{1-3} haloalkyl. In still further embodiments, R^{15} is C_1 haloalkyl. In yet further embodiments, R^{15} is CF_3 .

In some embodiments, R^{14} and R^{15} are each independently halo or C_{1-3} haloalkyl (In some further embodiments, each of of R^{12} , R^{13} , and R^{16} is H.). In some further embodiments, R^{14} and R^{15} are each independently halo or C_1 haloalkyl.

In some embodiments, R^{14} and R^{15} are each independently halo.


In some embodiments, the compound of Formula I is a compound of Formula II:

11




II.

In some embodiments, the compound of Formula II or pharmaceutically acceptable salt thereof is a compound of Formula IIa or IIb:



IIa

or



IIb.

or pharmaceutically acceptable salt thereof.

In some embodiments, the compound of Formula II is a compound of Formula IIa. In some further embodiments, R^{10} and R^{11} are each, independently, selected from H and C_{1-3}

alkyl. In yet further embodiments, R¹⁰ and R¹¹ are each, independently, selected from H and methyl. In still further embodiments, R¹⁰ and R¹¹ are each H.

In some embodiments of the compound of Formula IIa or pharmaceutically acceptable salt thereof, one of R¹⁰ and R¹¹ is selected from H and C₁₋₃ alkyl and the other is H. In some further embodiments, one of R¹⁰ and R¹¹ is C₁₋₃ alkyl. In yet further
5 embodiments, one of R¹⁰ and R¹¹ is methyl.

In some embodiments of the compound of Formula IIa or pharmaceutically acceptable salt thereof, both of R¹⁰ and R¹¹ are selected from C₁₋₃ alkyl. In some further
10 embodiments, both R¹⁰ and R¹¹ are methyl.

In some embodiments of the compound of Formula IIa or pharmaceutically acceptable salt thereof, R¹², R¹³, R¹⁴, R¹⁵, and R¹⁶ are each, independently, selected from H, OH, C₁₋₆ alkoxy, C₁₋₆ haloalkoxy, halo, CN, NO₂, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₃₋₇ cycloalkyl, NH₂, NH(C₁₋₄ alkyl), NH(C₃₋₇ cycloalkyl), N(C₁₋₄ alkyl)₂, NHC(O)(C₁₋₄ alkyl), SH, S(C₁₋₆ alkyl), C(O)OH, C(O)O(C₁₋₄ alkyl), C(O)(C₁₋₄ alkyl), and C(O)NH(C₁₋₄ alkyl).
15

In some embodiments of the compound of Formula IIa or pharmaceutically acceptable salt thereof, R¹², R¹³, R¹⁴, R¹⁵, and R¹⁶ are each, independently, selected from H, halo, CN, NO₂, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₃₋₇ cycloalkyl, C(O)O(C₁₋₄ alkyl), C(O)(C₁₋₄ alkyl), and C(O)NH(C₁₋₄ alkyl).
20

In some embodiments of the compound of Formula IIa or pharmaceutically acceptable salt thereof, R¹², R¹³, R¹⁴, R¹⁵, and R¹⁶ are each, independently, selected from H, halo, CN, NO₂, C₁₋₆ alkyl, C₁₋₆ haloalkyl, and C₃₋₇ cycloalkyl. In some further embodiments, R¹², R¹³, R¹⁴, R¹⁵, and R¹⁶ are each, independently, selected from H, halo, CN, C₁₋₆ alkyl, and C₁₋₆ haloalkyl. In yet further embodiments, R¹², R¹³, R¹⁴, R¹⁵, and R¹⁶ are each, independently, selected from H, halo, C₁₋₆ alkyl, and C₁₋₆ haloalkyl.
25

In some embodiments of the compound of Formula IIa or pharmaceutically acceptable salt thereof, at least one of R¹², R¹³, R¹⁴, R¹⁵, and R¹⁶ is selected from halo and C₁₋₆ haloalkyl, and each of the rest is of R¹², R¹³, R¹⁴, R¹⁵, and R¹⁶ is H. In some further
30 embodiments, one or two of R¹², R¹³, R¹⁴, R¹⁵, and R¹⁶ are selected from halo and C₁₋₆ haloalkyl, and each of the rest is of R¹², R¹³, R¹⁴, R¹⁵, and R¹⁶ is H. In yet further
embodiments, one of R¹², R¹³, R¹⁴, R¹⁵, and R¹⁶ is selected from halo and C₁₋₆ haloalkyl, and each of the rest is of R¹², R¹³, R¹⁴, R¹⁵, and R¹⁶ is H.

In some embodiments of the compound of Formula IIa or pharmaceutically acceptable salt thereof, at least one of R¹², R¹³, R¹⁴, R¹⁵, and R¹⁶ is selected from halo, CN, NO₂, C₁₋₆ haloalkyl, C(O)O(C₁₋₄ alkyl), C(O)(C₁₋₄ alkyl), and C(O)NH(C₁₋₄ alkyl).

In some embodiments of the compound of Formula IIa, at least one of R^{12} , R^{13} , R^{14} , R^{15} , and R^{16} is selected from halo and C_{1-6} haloalkyl.

In some embodiments of the compound of Formula IIa or pharmaceutically acceptable salt thereof, at least one of R^{12} , R^{13} , R^{14} , R^{15} , and R^{16} is selected from halo and C_{1-3} haloalkyl. In some further embodiments, at least one of R^{12} , R^{13} , R^{14} , R^{15} , and R^{16} is selected from halo and C_1 haloalkyl.

In some embodiments of the compound of Formula IIa or pharmaceutically acceptable salt thereof, R^{14} is halo or C_{1-6} haloalkyl. In some further embodiments, R^{14} is halo or C_{1-3} haloalkyl. In still further embodiments, R^{14} is halo or C_1 haloalkyl.

In some embodiments of the compound of Formula IIa or pharmaceutically acceptable salt thereof, R^{14} is halo (In some further embodiments, each of R^{12} , R^{13} , R^{15} , and R^{16} is H.). In some embodiments, R^{14} is Cl or F. In some embodiments, R^{14} is Cl. In some embodiments, R^{14} is F.

In some embodiments of the compound of Formula IIa, R^{14} is C_{1-6} haloalkyl (In some further embodiments, each of R^{12} , R^{13} , R^{15} , and R^{16} is H.). In some further embodiments, R^{14} is C_{1-3} haloalkyl. In still further embodiments, R^{14} is C_1 haloalkyl. In yet further embodiments, R^{14} is CF_3 .

In some embodiments of the compound of Formula IIa or pharmaceutically acceptable salt thereof, R^{14} is halo or C_{1-6} haloalkyl and each of R^{12} , R^{13} , R^{15} , and R^{16} is H.

In some embodiments of the compound of Formula IIa or pharmaceutically acceptable salt thereof, R^{15} is halo or C_{1-6} haloalkyl (In some further embodiments, each of R^{12} , R^{13} , R^{14} , and R^{16} is H.). In some further embodiments, R^{15} is halo or C_{1-3} haloalkyl. In still further embodiments, R^{15} is halo or C_1 haloalkyl.

In some embodiments of the compound of Formula IIa or pharmaceutically acceptable salt thereof, R^{15} is halo. In some embodiments, R^{15} is Cl or F. In some embodiments, R^{15} is Cl. In some embodiments, R^{15} is F.

In some embodiments of the compound of Formula IIa or pharmaceutically acceptable salt thereof, R^{15} is C_{1-6} haloalkyl. In some further embodiments, R^{15} is C_{1-3} haloalkyl. In still further embodiments, R^{15} is C_1 haloalkyl. In yet further embodiments, R^{15} is CF_3 .

In some embodiments of the compound of Formula IIa or pharmaceutically acceptable salt thereof, R^{14} and R^{15} are each independently halo or C_{1-3} haloalkyl (In some further embodiments, each of R^{12} , R^{13} , and R^{16} is H.). In some further embodiments, R^{14} and R^{15} are each independently halo or C_1 haloalkyl. In yet further embodiments, R^{14} and R^{15} are each independently halo.

In some embodiments, the compound of Formula II or pharmaceutically acceptable salt thereof is a compound of Formula IIb or pharmaceutically acceptable salt thereof.

In some embodiments of the compound of Formula IIb or pharmaceutically acceptable salt thereof, R^{12} , R^{13} , R^{14} , R^{15} , and R^{16} are each, independently, selected from H, OH, C_{1-6} alkoxy, C_{1-6} haloalkoxy, halo, CN, NO_2 , C_{1-6} alkyl, C_{1-6} haloalkyl, C_{3-7} cycloalkyl, NH_2 , NH(C_{1-4} alkyl), NH(C_{3-7} cycloalkyl), N(C_{1-4} alkyl)₂, NHC(O)(C_{1-4} alkyl), SH, S(C_{1-6} alkyl), C(O)OH, C(O)O(C_{1-4} alkyl), C(O)(C_{1-4} alkyl), and C(O)NH(C_{1-4} alkyl).

In some embodiments of the compound of Formula IIb or pharmaceutically acceptable salt thereof, R^{12} , R^{13} , R^{14} , R^{15} , and R^{16} are each, independently, selected from H, halo, CN, NO_2 , C_{1-6} alkyl, C_{1-6} haloalkyl, C_{3-7} cycloalkyl, C(O)O(C_{1-4} alkyl), C(O)(C_{1-4} alkyl), and C(O)NH(C_{1-4} alkyl).

In some embodiments of the compound of Formula IIb or pharmaceutically acceptable salt thereof, R^{12} , R^{13} , R^{14} , R^{15} , and R^{16} are each, independently, selected from H, halo, CN, NO_2 , C_{1-6} alkyl, C_{1-6} haloalkyl, and C_{3-7} cycloalkyl. In some further embodiments, R^{12} , R^{13} , R^{14} , R^{15} , and R^{16} are each, independently, selected from H, halo, CN, C_{1-6} alkyl, and C_{1-6} haloalkyl. In yet further embodiments, R^{12} , R^{13} , R^{14} , R^{15} , and R^{16} are each, independently, selected from H, halo, C_{1-6} alkyl, and C_{1-6} haloalkyl.

In some embodiments of the compound of Formula IIb or pharmaceutically acceptable salt thereof, at least one of R^{12} , R^{13} , R^{14} , R^{15} , and R^{16} is selected from halo and C_{1-6} haloalkyl, and each of the rest is of R^{12} , R^{13} , R^{14} , R^{15} , and R^{16} is H. In some further embodiments, one or two of R^{12} , R^{13} , R^{14} , R^{15} , and R^{16} are selected from halo and C_{1-6} haloalkyl, and each of the rest is of R^{12} , R^{13} , R^{14} , R^{15} , and R^{16} is H. In yet further embodiments, one of R^{12} , R^{13} , R^{14} , R^{15} , and R^{16} is selected from halo and C_{1-6} haloalkyl, and each of the rest is of R^{12} , R^{13} , R^{14} , R^{15} , and R^{16} is H.

In some embodiments of the compound of Formula IIb or pharmaceutically acceptable salt thereof, at least one of R^{12} , R^{13} , R^{14} , R^{15} , and R^{16} is selected from halo, CN, NO_2 , C_{1-6} haloalkyl, C(O)O(C_{1-4} alkyl), C(O)(C_{1-4} alkyl), and C(O)NH(C_{1-4} alkyl).

In some embodiments of the compound of Formula IIb, at least one of R^{12} , R^{13} , R^{14} , R^{15} , and R^{16} is selected from halo and C_{1-6} haloalkyl.

In some embodiments of the compound of Formula IIb or pharmaceutically acceptable salt thereof, at least one of R^{12} , R^{13} , R^{14} , R^{15} , and R^{16} is selected from halo and C_{1-3} haloalkyl. In some further embodiments, at least one of R^{12} , R^{13} , R^{14} , R^{15} , and R^{16} is selected from halo and C_1 haloalkyl.

In some embodiments of the compound of Formula IIb or pharmaceutically acceptable salt thereof, R^{14} is halo or C_{1-6} haloalkyl. In some further embodiments, R^{14} is halo or C_{1-3} haloalkyl. In still further embodiments, R^{14} is halo or C_1 haloalkyl.

5 In some embodiments of the compound of Formula IIb or pharmaceutically acceptable salt thereof, R^{14} is halo (In some further embodiments, each of R^{12} , R^{13} , R^{15} , and R^{16} is H.). In some embodiments, R^{14} is Cl or F. In some embodiments, R^{14} is Cl. In some embodiments, R^{14} is F.

10 In some embodiments of the compound of Formula IIb, R^{14} is C_{1-6} haloalkyl (In some further embodiments, each of R^{12} , R^{13} , R^{15} , and R^{16} is H.). In some further embodiments, R^{14} is C_{1-3} haloalkyl. In still further embodiments, R^{14} is C_1 haloalkyl. In yet further embodiments, R^{14} is CF_3 .

In some embodiments of the compound of Formula IIb or pharmaceutically acceptable salt thereof, R^{14} is halo or C_{1-6} haloalkyl and each of R^{12} , R^{13} , R^{15} , and R^{16} is H.

15 In some embodiments of the compound of Formula IIb or pharmaceutically acceptable salt thereof, R^{15} is halo or C_{1-6} haloalkyl (In some further embodiments, each of R^{12} , R^{13} , R^{14} , and R^{16} is H.). In some further embodiments, R^{15} is halo or C_{1-3} haloalkyl. In still further embodiments, R^{15} is halo or C_1 haloalkyl.


20 In some embodiments of the compound of Formula IIb or pharmaceutically acceptable salt thereof, R^{15} is halo. In some embodiments, R^{15} is Cl or F. In some embodiments, R^{15} is Cl. In some embodiments, R^{15} is F.

In some embodiments of the compound of Formula IIb or pharmaceutically acceptable salt thereof, R^{15} is C_{1-6} haloalkyl. In some further embodiments, R^{15} is C_{1-3} haloalkyl. In still further embodiments, R^{15} is C_1 haloalkyl. In yet further embodiments, R^{15} is CF_3 .

25 In some embodiments of the compound of Formula IIb or pharmaceutically acceptable salt thereof, R^{14} and R^{15} are each independently halo or C_{1-3} haloalkyl (In some further embodiments, each of R^{12} , R^{13} , and R^{16} is H.). In some further embodiments, R^{14} and R^{15} are each independently halo or C_1 haloalkyl. In yet further embodiments, R^{14} and R^{15} are each independently halo.

30 In some embodiments, the compound of Formula I is a compound of Formula III:

16



III.

In some embodiments of compounds of Formula III or pharmaceutically acceptable salt thereof, m is 1.

5 In some embodiments of compounds of Formula III or pharmaceutically acceptable salt thereof, m is 0.

In some embodiments of compounds of Formula III or pharmaceutically acceptable salt thereof, at least one of R^2 , R^3 , R^4 , R^5 , and R^6 is selected from OH, C_{1-6} alkoxy, and C_{1-6} haloalkoxy.

10 In some embodiments of compounds of Formula III or pharmaceutically acceptable salt thereof, at least two of R^2 , R^3 , R^4 , R^5 , and R^6 are independently selected from OH, C_{1-6} alkoxy, and C_{1-6} haloalkoxy.

In some embodiments of compounds of Formula III or pharmaceutically acceptable salt thereof, at least one of R^{12} , R^{13} , R^{14} , R^{15} , and R^{16} is other than H.

15 In some embodiments of the compound of Formula III or pharmaceutically acceptable salt thereof, R^{14} and R^{15} are each independently halo or C_{1-3} haloalkyl. In some further embodiments, R^{14} and R^{15} are each independently halo or C_1 haloalkyl.

20 At various places in the present specification, substituents of compounds of the invention are disclosed in groups or in ranges. It is specifically intended that embodiments the invention include each and every individual subcombination of the members of such groups and ranges. For example, the term " C_{1-6} alkyl" is specifically intended to individually

disclose methyl (C_1 alkyl), ethyl (C_2 alkyl), C_3 alkyl, C_4 alkyl, C_5 alkyl, and C_6 alkyl.

For compounds of the invention in which a variable appears more than once, each variable can be a different moiety selected from the Markush group defining the variable. For example, where a structure is described having two R groups that are simultaneously
5 present on the same compound, then the two R groups can represent different moieties selected from the Markush group defined for R.

It is further appreciated that certain features of the invention, which are, for clarity, described in the context of separate embodiments, can also be provided in combination in a single embodiment. Conversely, various features of the invention which are, for brevity,
10 described in the context of a single embodiment, can also be provided separately or in any suitable subcombination.

The term "n-membered" where n is an integer typically describes the number of ring-forming atoms in a moiety where the number of ring-forming atoms is n. For example, pyridine is an example of a 6-membered heteroaryl ring and thiophene is an example of a 5-
15 membered heteroaryl group.

As used herein, the term "alkyl" is meant to refer to a saturated hydrocarbon group which is straight-chained or branched. Example alkyl groups include, but are not limited to, methyl (Me), ethyl (Et), propyl (e.g., n-propyl and isopropyl), butyl (e.g., n-butyl, isobutyl, t-butyl), pentyl (e.g., n-pentyl, isopentyl, neopentyl), and the like. An alkyl group can contain
20 from 1 to about 20, from 2 to about 20, from 1 to about 10, from 1 to about 8, from 1 to about 6, from 1 to about 4, or from 1 to about 3 carbon atoms. The term "alkylene" refers to a divalent alkyl linking group. An example of alkylene is methylene (CH_2).

As used herein, "alkenyl" refers to an alkyl group having one or more double carbon-carbon bonds. Example alkenyl groups include, but are not limited to, ethenyl, propenyl,
25 cyclohexenyl, and the like. The term "alkenylene" refers to a divalent linking alkenyl group.

As used herein, "alkynyl" refers to an alkyl group having one or more triple carbon-carbon bonds. Example alkynyl groups include, but are not limited to, ethynyl, propynyl, and the like. The term "alkynylene" refers to a divalent linking alkynyl group.

As used herein, "haloalkyl" refers to an alkyl group having one or more halogen
30 substituents. Example haloalkyl groups include, but are not limited to, CF_3 , C_2F_5 , CHF_2 , CCl_3 , $CHCl_2$, C_2Cl_5 , CH_2CF_3 , and the like.

As used herein, "aryl" refers to monocyclic or polycyclic (e.g., having 2, 3 or 4 fused rings) aromatic hydrocarbons such as, for example, phenyl, naphthyl, anthracenyl, phenanthrenyl, indanyl, indenyl, and the like. In some embodiments, aryl groups have from
35 6 to about 20 carbon atoms. In some embodiments, aryl groups have from 6 to about 10

carbon atoms.

As used herein, "cycloalkyl" refers to non-aromatic cyclic hydrocarbons including cyclized alkyl, alkenyl, and alkynyl groups that contain up to 20 ring-forming carbon atoms. Cycloalkyl groups can include mono- or polycyclic (e.g., having 2, 3 or 4 fused rings) ring systems as well as spiro ring systems. A cycloalkyl group can contain from 3 to about 15, from 3 to about 10, from 3 to about 8, from 3 to about 6, from 4 to about 6, from 3 to about 5, or from 5 to about 6 ring-forming carbon atoms. Ring-forming carbon atoms of a cycloalkyl group can be optionally substituted by oxo or sulfido. Example cycloalkyl groups include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclopentenyl, cyclohexenyl, cyclohexadienyl, cycloheptatrienyl, norbornyl, norpinyl, norcarnyl, adamantyl, and the like. Also included in the definition of cycloalkyl are moieties that have one or more aromatic rings fused (i.e., having a bond in common with) to the cycloalkyl ring, for example, benzo or thienyl derivatives of pentane, pentene, hexane, and the like (e.g., 2,3-dihydro-1H-indene-1-yl, or 1H-inden-2(3H)-one-1-yl). Preferably, "cycloalkyl" refers to cyclized alkyl groups that contain up to 20 ring-forming carbon atoms. Examples of cycloalkyl preferably include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, adamantyl, and the like.

As used herein, "heteroaryl" groups refer to an aromatic heterocycle having up to 20 ring-forming atoms and having at least one heteroatom ring member (ring-forming atom) such as sulfur, oxygen, or nitrogen. In some embodiments, the heteroaryl group has at least one or more heteroatom ring-forming atoms each independently selected from sulfur, oxygen, and nitrogen. Heteroaryl groups include monocyclic and polycyclic (e.g., having 2, 3 or 4 fused rings) systems. Examples of heteroaryl groups include without limitation, pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl, triazinyl, furyl, quinolyl, isoquinolyl, thienyl, imidazolyl, thiazolyl, indolyl, pyrrol, oxazolyl, benzofuryl, benzothienyl, benzthiazolyl, isoxazolyl, pyrazolyl, triazolyl, tetrazolyl, indazolyl, 1,2,4-thiadiazolyl, isothiazolyl, benzothienyl, purinyl, carbazolyl, benzimidazolyl, indolinyl, and the like. In some embodiments, the heteroaryl group has from 1 to about 20 carbon atoms, and in further embodiments from about 1 to about 5, from about 1 to about 4, from about 1 to about 3, from about 1 to about 2, carbon atoms as ring-forming atoms. In some embodiments, the heteroaryl group contains 3 to about 14, 3 to about 7, or 5 to 6 ring-forming atoms. In some embodiments, the heteroaryl group has 1 to about 4, 1 to about 3, or 1 to 2 heteroatoms.

As used herein, "heterocycloalkyl" refers to non-aromatic heterocycles having up to 20 ring-forming atoms including cyclized alkyl, alkenyl, and alkynyl groups where one or more of the ring-forming carbon atoms is replaced by a heteroatom such as an O, N, or S

atom. Heterocycloalkyl groups can be mono or polycyclic (e.g., both fused and spiro systems). Example "heterocycloalkyl" groups include morpholino, thiomorpholino, piperazinyl, tetrahydrofuranyl, tetrahydrothienyl, 2,3-dihydrobenzofuryl, 1,3-benzodioxole, benzo-1,4-dioxane, piperidinyl, pyrrolidinyl, isoxazolidinyl, isothiazolidinyl, pyrazolidinyl, oxazolidinyl, thiazolidinyl, imidazolidinyl, pyrrolidin-2-one-3-yl, and the like. Ring-forming carbon atoms and heteroatoms of a heterocycloalkyl group can be optionally substituted by oxo or sulfido. For example, a ring-forming S atom can be substituted by 1 or 2 oxo [i.e., form a S(O) or S(O)₂]. For another example, a ring-forming C atom can be substituted by oxo (i.e., form carbonyl). Also included in the definition of heterocycloalkyl are moieties that have one or more aromatic rings fused (i.e., having a bond in common with) to the nonaromatic heterocyclic ring, for example pyridinyl, thiophenyl, phthalimidyl, naphthalimidyl, and benzo derivatives of heterocycles such as indolene, isoindolene, isoindolin-1-one-3-yl, 4,5,6,7-tetrahydrothieno[2,3-c]pyridine-5-yl, 5,6-dihydrothieno[2,3-c]pyridin-7(4H)-one-5-yl, and 3,4-dihydroisoquinolin-1(2H)-one-3yl groups. Ring-forming carbon atoms and heteroatoms of the heterocycloalkyl group can be optionally substituted by oxo or sulfido. In some embodiments, the heterocycloalkyl group has from 1 to about 20 carbon atoms, and in further embodiments from about 3 to about 20 carbon atoms. In some embodiments, the heterocycloalkyl group contains 3 to about 14, 3 to about 7, or 5 to 6 ring-forming atoms. In some embodiments, the heterocycloalkyl group has 1 to about 4, 1 to about 3, or 1 to 2 heteroatoms. In some embodiments, the heterocycloalkyl group contains 0 to 3 double bonds. In some embodiments, the heterocycloalkyl group contains 0 to 2 triple bonds.

As used herein, "halo" or "halogen" includes fluoro, chloro, bromo, and iodo.

As used herein, "alkoxy" refers to an -O-alkyl group. Example alkoxy groups include methoxy, ethoxy, propoxy (e.g., n-propoxy and isopropoxy), t-butoxy, and the like.

As used herein, "haloalkoxy" refers to an -O-haloalkyl group. An example haloalkoxy group is OCF₃. As used herein, "trihalomethoxy" refers to a methoxy group having three halogen substituents. Examples of trihalomethoxy groups include, but are not limited to, -OCF₃, -OCClF₂, -OCCl₃, and the like.

As used herein, "arylalkyl" refers to a C₁₋₆ alkyl substituted by aryl and "cycloalkylalkyl" refers to C₁₋₆ alkyl substituted by cycloalkyl.

As used herein, "heteroarylalkyl" refers to a C₁₋₆ alkyl group substituted by a heteroaryl group, and "heterocycloalkylalkyl" refers to a C₁₋₆ alkyl substituted by heterocycloalkyl.

As used herein, "amino" refers to NH₂.

As used herein, "alkylamino" refers to an amino group substituted by an alkyl group.

As used herein, "dialkylamino" refers to an amino group substituted by two alkyl groups.

As used here, C(O) refers to C(=O).

As used here, C(S) refers to C(=S).

5 As used here, S(O) refers to S(=O).

As used here, S(O)₂ refers to S(=O)₂.

As used herein, the term "optionally substituted" means that substitution is optional and therefore includes both unsubstituted and substituted atoms and moieties. A "substituted" atom or moiety indicates that any hydrogen on the designated atom or moiety can be replaced with a selection from the indicated substituent group, provided that the normal valency of the designated atom or moiety is not exceeded, and that the substitution results in a stable compound. For example, if a methyl group (i.e., CH₃) is optionally substituted, then 3 hydrogen atoms on the carbon atom can be replaced with substituent groups.

15 As used herein, "about" in connection with a numerical value means that the numerical value is approximate and small variations would not significantly affect the practice of the disclosed embodiments. Where a numerical limitation is used, unless indicated otherwise by the context, "about" means the numerical value can vary by $\pm 10\%$ and remain within the scope of the disclosed embodiments.

20 The compounds described in the embodiments herein can be asymmetric (*e.g.*, having one or more stereocenters). All stereoisomers, such as enantiomers and diastereomers, are intended unless otherwise indicated. Compounds of the present invention that contain asymmetrically substituted carbon atoms can be isolated in optically active or racemic forms. Methods on how to prepare optically active forms from optically active starting materials are known in the art, such as by resolution of racemic mixtures or by stereoselective synthesis. Many geometric isomers of olefins, C=N double bonds, and the like can also be present in the compounds described herein, and all such stable isomers are contemplated in the present invention. *Cis* and *trans* geometric isomers of the compounds of the present invention are described and may be isolated as a mixture of isomers or as separated isomeric forms. Where a compound capable of stereoisomerism or geometric isomerism is designated in its structure or name without reference to specific R/S or *cis/trans* configurations, it is intended that all such isomers are contemplated.

35 Resolution of racemic mixtures of compounds can be carried out by any of numerous methods known in the art. An example method includes fractional recrystallization using a chiral resolving acid which is an optically active, salt-forming organic acid. Suitable resolving

agents for fractional recrystallization methods are, for example, optically active acids, such as the D and L forms of tartaric acid, diacetyltartaric acid, dibenzoyltartaric acid, mandelic acid, malic acid, lactic acid or the various optically active camphorsulfonic acids such as β -camphorsulfonic acid. Other resolving agents suitable for fractional crystallization methods include stereoisomerically pure forms of α -methylbenzylamine (e.g., *S* and *R* forms, or diastereomerically pure forms), 2-phenylglycinol, norephedrine, ephedrine, N-methylephedrine, cyclohexylethylamine, 1,2-diaminocyclohexane, and the like.

Resolution of racemic mixtures can also be carried out by elution on a column packed with an optically active resolving agent (e.g., dinitrobenzoylphenylglycine). Suitable elution solvent composition can be determined by one skilled in the art.

Compounds of embodiments the invention also include tautomeric forms. Tautomeric forms result from the swapping of a single bond with an adjacent double bond together with the concomitant migration of a proton. Tautomeric forms include prototropic tautomers which are isomeric protonation states having the same empirical formula and total charge. Example prototropic tautomers include ketone – enol pairs, amide - imidic acid pairs, lactam – lactim pairs, amide - imidic acid pairs, enamine – imine pairs, and annular forms where a proton can occupy two or more positions of a heterocyclic system, for example, 1H- and 3H-imidazole, 1H-, 2H- and 4H- 1,2,4-triazole, 1H- and 2H- isoindole, and 1H- and 2H-pyrazole. Tautomeric forms can be in equilibrium or sterically locked into one form by appropriate substitution.

Compounds of embodiments the invention further include hydrates and solvates, as well as anhydrous and non-solvated forms.

The term, “compound,” as used herein is meant to include all stereoisomers, geometric isomers, tautomers, and isotopes of the structures depicted.

All compounds and pharmaceutically acceptable salts thereof, can be prepared or present together with other substances such as water and solvents (e.g. hydrates and solvates) or can be isolated.

Compounds of embodiments the invention can also include all isotopes of atoms occurring in the intermediates or final compounds. Isotopes include those atoms having the same atomic number but different mass numbers. For example, isotopes of hydrogen include tritium and deuterium.

In some embodiments, the compounds of the invention, or salts thereof, are substantially isolated. By “substantially isolated” is meant that the compound is at least partially or substantially separated from the environment in which is formed or detected.

Partial separation can include, for example, a composition enriched in the compound of the

invention. Substantial separation can include compositions containing at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 90%, at least about 95%, at least about 97%, or at least about 99% by weight of the compound of the invention, or salt thereof. Methods for isolating compounds and their salts are routine in the art.

5 Compounds of embodiments the invention are intended to include compounds with stable structures. As used herein, "stable compound" and "stable structure" are meant to indicate a compound that is sufficiently robust to survive isolation to a useful degree of purity from a reaction mixture, and formulation into an efficacious therapeutic agent.

10 The phrase "pharmaceutically acceptable" is employed herein to refer to those compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio.

15 The expressions, "ambient temperature" and "room temperature," as used herein, are understood in the art, and refer generally to a temperature, *e.g.* a reaction temperature, that is about the temperature of the room in which the reaction is carried out, for example, a temperature from about 20 °C to about 30 °C.

20 Embodiments of the present invention also includes pharmaceutically acceptable salts of the compounds described herein. As used herein, "pharmaceutically acceptable salts" refers to derivatives of the disclosed compounds wherein the parent compound is modified by converting an existing acid or base moiety to its salt form. Examples of pharmaceutically acceptable salts include, but are not limited to, mineral or organic acid salts of basic residues such as amines; alkali or organic salts of acidic residues such as carboxylic acids; and the like. The pharmaceutically acceptable salts of the present invention include
25 the conventional non-toxic salts of the parent compound formed, for example, from non-toxic inorganic or organic acids. The pharmaceutically acceptable salts of the present invention can be synthesized from the parent compound which contains a basic or acidic moiety by conventional chemical methods. Generally, such salts can be prepared by reacting the free acid or base forms of these compounds with a stoichiometric amount of the appropriate base
30 or acid in water or in an organic solvent, or in a mixture of the two; generally, nonaqueous media like ether, ethyl acetate, ethanol, isopropanol, or acetonitrile (ACN) are preferred. Lists of suitable salts are found in *Remington's Pharmaceutical Sciences*, 17th ed., Mack Publishing Company, Easton, Pa., 1985, p. 1418 and *Journal of Pharmaceutical Science*, 66, 2 (1977), each of which is incorporated herein by reference in its entirety.

Synthesis

Compounds of embodiments of the invention, including salts thereof, can be prepared using known organic synthesis techniques and can be synthesized according to any of numerous possible synthetic routes.

5 The reactions for preparing compounds of the invention can be carried out in suitable solvents which can be readily selected by one of skill in the art of organic synthesis. Suitable solvents can be substantially non-reactive with the starting materials (reactants), the intermediates, or products at the temperatures at which the reactions are carried out, e.g., temperatures which can range from the solvent's freezing temperature to the solvent's boiling
10 temperature. A given reaction can be carried out in one solvent or a mixture of more than one solvent. Depending on the particular reaction step, suitable solvents for a particular reaction step can be selected by the skilled artisan.

Preparation of compounds of the invention can involve the protection and deprotection of various chemical groups. The need for protection and deprotection, and the
15 selection of appropriate protecting groups, can be readily determined by one skilled in the art. The chemistry of protecting groups can be found, for example, in T. W. Greene and P. G. M. Wuts, *Protective Groups in Organic Synthesis*, 3rd Ed., Wiley & Sons, Inc., New York (1999), which is incorporated herein by reference in its entirety.


Reactions can be monitored according to any suitable method known in the art. For
20 example, product formation can be monitored by spectroscopic means, such as nuclear magnetic resonance spectroscopy (e.g., ¹H or ¹³C), infrared spectroscopy, spectrophotometry (e.g., UV-visible), mass spectrometry, or by chromatographic methods such as high performance liquid chromatography (HPLC) or thin layer chromatography (TLC).

25 The compounds of embodiments of the invention can be prepared, for example, according to the reaction pathways, synthetic procedures, and techniques described below.

As shown in Scheme 1, benzaldehyde derivative **1-1** can be reacted with acetaldehyde in the presence of either an acid or a base catalyst to afford cinnamic aldehyde **1-2**. Reaction of cinnamic aldehyde **1-2** with an organometallic compound such as a
30 Grignard reagent R⁸MgX¹ [wherein X¹ is halo such as Cl or Br], followed by oxidation of the intermediate alcohol to ketone and by reduction of the C=C bond to a C-C single bond under a hydrogenation condition (for example, in the presence of Pd/C catalyst), affords ketone **1-3**. Reaction of ketone **1-3** with amine **1-4** under reductive amination condition (such as in the presence of a borohydride reducing reagent) affords compound **1-5**.

24


Scheme 1



As shown in Scheme 2, Ketone **2-1** can be reacted with an organometallic compound such as a Grignard reagent R^7MgX^2 [wherein R^7 can be C_{1-6} alkyl or C_{3-7} cycloalkyl; and X^2 can be halo such as Cl or Br] to afford alcohol **2-2a**. Reduction of Ketone **2-1** such as in the presence of a borohydride reducing reagent affords alcohol **2-2b**. The OH group of alcohol **2-2** (wherein R^7 can be H, C_{1-6} alkyl, or C_{3-7} cycloalkyl) can be converted to a better leaving group Lg^1 such as OMs [mesylate or $\text{CH}_3\text{S}(\text{O})_2\text{O}-$] or OTf [triflate or $\text{CF}_3\text{S}(\text{O})_2\text{O}-$], followed by reaction with NH_3 to afford amine **2-4**. Ketone **2-1** can also undergo reductive amination with NH_3 to afford amine **2-5**.


25

Scheme 2



As shown in Scheme 3, amine **3-1** (wherein R^7 can be, e.g., H, C_{1-6} alkyl, or C_{3-7} cycloalkyl) can be reacted with compound **3-2** [Lg^2 can be a leaving group such as triflate group (-OTf) or halo (e.g. Cl or Br)] to afford compound **3-3**.


Scheme 3



As shown in Scheme 4, benzaldehyde derivative **4-1** can be reacted with a methylmetallic compound such as a Grignard reagent $MeMgX^1$ [wherein X^1 is halo such as Cl or Br], followed by oxidation of the intermediate alcohol to ketone to afford ketone **4-2**. Ester compound **4-3** [wherein R' can be alkyl (e.g. methyl or ethyl) or arylalkyl; and Pg^1 can be an amine protecting group (such as tert-butyloxycarbonyl or Boc; benzyloxycarbonyl or Cbz; or benzyl)] can be reduced to an alcohol (in the presence of a reducing reagent such as Lithium aluminium hydride or LAH), followed by conversion of the OH group to a better leaving group Lg^3 such as OMs [mesylate or $CH_3S(O)_2O^-$] or OTf [triflate or $CF_3S(O)_2O^-$], to afford compound **4-4**.

Reaction of compound **4-2** with compound **4-4** in the presence of a strong base (such as lithium diisopropylamide or LDA), followed by hydrogenation (such as in the presence of a Pd/C catalyst) to reduce the $C(O)$ to CH_2 and by removal of the protecting group Pg^1 under suitable conditions (for example, a benzyl group can be removed under hydrogenation condition in the presence of Pd/C; or Boc group can be removed under acidic condition), affords amine **4-5**. Amine **4-5** can be reacted with compound **4-6** [wherein Lg^2 can be a leaving group such as triflate group (-OTf) or halo (e.g. Cl or Br)] to afford compound **4-7**.

Scheme 4



Those skilled in the art can recognize that in all of the schemes described herein, if there are functional (reactive) groups present on a substituent group such as R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , R^9 , R^{10} , R^{11} , R^{12} , R^{13} , R^{14} , R^{15} , R^{16} , etc., further modification can be made if appropriate and/or desired. For example, a CN group can be hydrolyzed to afford an amide

group; a carboxylic acid can be converted to an amide; a carboxylic acid can be converted to an ester, which in turn can be reduced to an alcohol, which in turn can be further modified. For another example, an OH group can be converted into a better leaving group such as mesylate, which in turn is suitable for nucleophilic substitution, such as by CN. For another example, an -S- can be oxidized to -S(O)- and/or -S(O)₂-. For yet another example, unsaturated bond such as C=C or C≡C can be reduced to saturated bond by hydrogenation. In some embodiments, a primary amine or a secondary amine moiety (present on a substituent group such as R¹ – R¹⁶, etc.) can be converted to amide, sulfonamide, urea, or thiourea moiety by reacting it with an appropriate reagent such as an acid chloride, a sulfonyl chloride, an isocyanate, or a thioisocyanate compound. Thus, a compound of Formula I (such as compound **3-3** of Scheme 3) having a substituent which contains a functional group can be converted to another compound of Formula I having a different substituent group.

As used herein, the term “reacting” refers to the bringing together of designated chemical reactants such that a chemical transformation takes place generating a compound different from any initially introduced into the system. Reacting can take place in the presence or absence of solvent.

Chemical Conditioning


In some embodiments, a method of preparing an array of chemical compounds from a biological extract such as ginger oil is provided.

The method of the invention, termed “chemical conditioning” is generally applicable to all biological extracts, in particular, natural plant extracts, common or medicinal. See e.g. US20080193574 and WO2008042755, each of which is incorporated herein by reference in its entirety. Chemical conditioning is a method which produces novel unnatural drug-like compounds from readily available natural materials. In general, the “chemical conditioning” of natural extracts coupled with pre-fractionation of the chemically conditioned extracts facilitates successful biochemical screening of extracts by destroying reactive natural compounds that generate false positive results in biochemical assays. Chemical conditioning produces novel lead-like and drug-like compounds and, the reductive amination protocol described here can produce structurally diverse nitrogen-containing products that are particularly lead-like and drug-like.

In certain embodiments of the present invention, a method of preparing chemical compounds from a biological extract is exemplified in Scheme 5a below. According to the method, first, a biological extract, e.g., a plant extract is provided, the biological extract has one or more biological compounds, each biological compound having one or more reactive

electrophilic groups. Next, the biological compounds in the biological extract are reacted with an amine to incorporate the amine into the biological compounds. Next, the biological compounds having the incorporated amine are reacted with a reducing agent to reduce the intermediate imine and enamine compounds and form one or more nitrogen-containing chemical compounds. Thus, the resultant nitrogen-containing chemical compounds are derivatives of the biological compounds in the biological extract. In some embodiments, the biological compounds in the biological extract are compounds having ketones and aldehydes that are reacted with various amines. This reaction is followed by hydride reduction of the intermediate imines and enamines to provide secondary and tertiary amines. The reaction of ketones and aldehydes with amines, followed by reduction to form imines and enamines is known in the art.

Scheme 5a



R' and R'' represent a variety of substituents that make up a biological compound; and

R* represents a variety of substituent(s) that, together with the nitrogen, make up an amine compound.

The chemical conditioning method described herein employs a biological extract, using many different reagents, to efficiently produce an array of nitrogen-containing chemical compounds. The ready commercial availability of many low molecular weight amines for use as inputs in the reductive amination sequence enables the development of many different and structurally diverse central nervous system druglike mixtures from the same natural extract. Suitable amines for use in the present method are selected from the group consisting of primary amines, secondary amines, cyclic amines, pyrrolidine, and amino acids. Suitable reducing agents for use in the present method are selected from the group of hydride reducing agents including but not limited to sodium borohydride, sodium triacetoxyborohydride, and lithium aluminum hydride.

The method may further comprise quenching the reaction a quenching agent, wherein the quenching agent is selected from but not limited to the group consisting of sodium bicarbonate, sodium carbonate, sodium sulfate, sodium sulfate decahydrate. The method may also further comprise isolating one or more of the nitrogen-containing chemical compounds, in a purified or unpurified form. The resultant nitrogen-containing chemical may then be screened or tested for biological activity.

The process of chemical conditioning by reductive amination described herein destroys reactive electrophiles in the natural extract, including ketones, as in the gingerols, and converts them to chemically stable compounds such as amines. The resulting conditioned extracts contain both natural compounds and novel unnatural nitrogen-
5 containing amine products that are potential drug candidates. In the case of the extracts of gingerol, the nitrogen-containing amine products are potential central nervous system drugs.

For the purpose of this disclosure, the following terms have the following meanings.

The term "biological compound" as used herein refers to a chemical compound that occurs in nature.

10 The term "biological extract" as used herein refers to an extract from a biological sample, such as a plant extract, or other extract from organic matter, containing chemical compounds that occur in nature.


The term "reactive electrophilic group" as used herein refers to an atom or group of atoms that has the ability to react with a nucleophile.

15 The term "nitrogen-containing derivative" as used herein represents those derivatives containing a nitrogen atom, where the nitrogen atom is a substitution another atom, such as oxygen in the parent compound.

In one embodiment, a specific example of the chemical conditioning process is shown in Scheme 5 below. Scheme 5 shows the two-step reductive amination chemical
20 conditioning protocol performed on ginger oil and ginger oleoresin in accordance with one embodiment of the method, wherein ginger oil or gingerol comprising ketone **5-1** are converted to amine **5-4**. According to the method shown in Scheme 5, ginger oil (an extract of ginger containing ketone **5-1** and other molecules occurring in natural ginger) is reacted with amine **5-2** to form compound **5-3**. Then, the resultant compound **5-3** is then reduced,
25 with a reducing agent such as a borohydride, to form the nitrogen-containing compound **5-4** (the reaction crude product also include other chemical compounds).

31


Scheme 5



"conditioned extracts"
nitrogen-containing compounds

In the next step of the method, amine **5-4** is isolated/purified from the extract (the crude reaction product of the 2-step reductive amination). The conditioned extracts can be fractionated by flash chromatography. The fraction that contains amine **5-4** can undergo further purification/isolation according to the methods known to those in the art. Further isolation and characterization of the fraction that contains amine **5-4** may follow. The isolated amine **5-4** is tested for its biological activity such as by those methods described hereinwith.

Some examples of benzylamine **5-2** used in the chemical conditioning process of the invention shown in Scheme 5 include:



New lead compounds generated by this chemical conditioning method can also be prepared to the synthetic methods described hereinwith.

- 5 In some embodiments, the derivatives of ginger oil such as amine **5-4** possess beta-secretase inhibitory activity, and/or inhibit amyloid production, amyloid assembly, the activity/effect of Abeta oligomers on neurons (such as neurons in the brain), amyloid aggregation, amyloid (including amyloid oligomer) binding, or amyloid deposition. These compounds are useful therapeutic agents for the treatment and prevention of cognitive
- 10 decline, amyloid production, neurodegeneration, and Alzheimer's disease.

Methods

- In some embodiments, the compounds of present invention inhibit, treat, or abate (partially inhibit) binding of amyloid (including Abeta oligomers) to neurons (such as neurons
- 15 in the brain) and are useful for the inhibition, treatment, and abatement of cognitive decline and/or Alzheimer's disease. In some embodiments, the compounds of present invention inhibit, treat, or abate (partially inhibit) one or more of amyloid aggregation, amyloid oligomer binding, and amyloid deposition. In some embodiments, the compounds of present invention inhibit, treat, or abate (partially inhibit) amyloid deposition. In some embodiments,
- 20 the compounds of present invention inhibit, treat, or abate (partially inhibit) the activity/effect of Abeta oligomers on neurons (such as neurons in the brain) and are useful for the inhibition, treatment, and abatement of cognitive decline and/or Alzheimer's disease. In some embodiments, the compounds of present invention inhibit, treat, or abate (partially inhibit) the activity/effect of Abeta oligomers on neurons (such as neurons in the brain) via
- 25 disruption of Abeta oligomers, inhibition of Abeta oligomer binding to neurons, and/or counteraction of signal transduction mechanisms of action initiated by Abeta oligomer

binding.

In some embodiments, the compounds show activity in a beta-secretase assay and are useful for the inhibition, treatment, and abatement of cognitive decline and Alzheimer's disease. In some embodiments the derivative of ginger oil is a compound in purified and isolated form (for example, with a purity of greater than 80%, 85%, 90%, 95%, 98%, or 99% by weight). The compounds and methods described herein may be used to treat one or more symptoms of cognitive decline and/or Alzheimer's disease such as memory loss, confusion, impaired judgment, personality changes, disorientation, and loss of language skills. Further, the compounds and methods described herein may be useful in inhibiting, treating, and/or abating cognitive decline and/or Alzheimer's disease by restoring long term potentiation, and/or inhibiting, treating, or abatement of one or both of neurodegeneration and general amyloidosis, more specifically, by inhibiting, treating, or abatement of one or more of amyloid production, amyloid assembly, amyloid aggregation, amyloid (including amyloid oligomer) binding, and amyloid deposition.

In some embodiments, compounds of the invention can inhibit, treat, or abate one or more of amyloid production, amyloid assembly, amyloid aggregation, amyloid oligomer binding, and amyloid deposition. In some embodiments, compounds of the invention can restore long term potentiation, inhibit, treat, or abate one or both of neurodegeneration and general amyloidosis.

In some embodiments, compounds of present invention inhibit, treat, or abate (partially inhibit) one or more of amyloid aggregation, amyloid oligomer binding, and amyloid deposition. In some embodiments, the compounds of present invention inhibit (or partially inhibit) amyloid deposition. In some embodiments, the compounds of present invention inhibit, treat, or abate (partially inhibit) binding of amyloid (including Abeta oligomers) to neurons (such as neurons in the brain). In some embodiments, the compounds of present invention are useful for the inhibition, treatment, and abatement of cognitive decline and/or Alzheimer's disease.

In some embodiments, compounds of the invention can inhibit activity of beta-secretase. In some embodiments, compounds of the invention can be used in methods of inhibiting activity of beta-secretase by contacting the beta-secretase with any one or more of the compounds or compositions described herein.

Another aspect of the present invention pertains to methods of treating cognitive decline and/or Alzheimer's disease in an individual (*e.g.*, patient) by administering to the individual a therapeutically effective amount or dose of a compound of the present invention or a pharmaceutical composition thereof.

Treatment of the diseases/disorders herein includes treating one or more symptoms associated with the diseases/disorders, for example, symptoms of cognitive decline and/or Alzheimer's disease.

As used herein, the term "contacting" refers to the bringing together of indicated
5 moieties in an *in vitro* system or an *in vivo* system. For example, "contacting" a beta-secretase or a neuron cell (or a neuron cell in the presence of one or more of beta-amyloid oligomers) with a compound of the invention includes the administration of a compound of the present invention to an individual or patient, such as a human, having a beta-secretase or a neuron cell, as well as, for example, introducing a compound of the invention into a
10 sample containing a cellular or purified preparation containing the a beta-secretase or a neuron cell (or a neuron cell in the presence of one or more of beta-amyloid oligomers).

As used herein, the term "individual" or "patient," used interchangeably, refers to any animal, including mammals, preferably mice, rats, other rodents, rabbits, dogs, cats, swine, cattle, sheep, horses, or primates, and most preferably humans.

As used herein, the phrase "therapeutically effective amount" refers to the amount of
15 active compound or pharmaceutical agent that elicits the biological or medicinal response that is being sought in a tissue, system, animal, individual or human by a researcher, veterinarian, medical doctor or other clinician.

As used herein, the term "treating" or "treatment" refers to one or more of (1)
20 preventing the disease; for example, preventing a disease, condition or disorder in an individual who may be predisposed to the disease, condition or disorder but does not yet experience or display the pathology or symptomatology of the disease; (2) inhibiting/retarding the disease; for example, inhibiting/retarding a disease, condition or disorder in an individual who is experiencing or displaying the pathology or symptomatology
25 of the disease, condition or disorder; and (3) ameliorating the disease; for example, ameliorating a disease, condition or disorder in an individual who is experiencing or displaying the pathology or symptomatology of the disease, condition or disorder (i.e., reversing the pathology and/or symptomatology) such as decreasing the severity of disease or completely eliminating/curing the disease. As used herein, treating a disease further
30 includes treating one or more symptoms associated with the disease.

Combination Therapies

In certain embodiments, one or more additional pharmaceutical agents for treatment of cognitive decline and/or Alzheimer's disease can be used in combination with the
35 compounds of the present invention for treatment of cognitive decline and/or Alzheimer's

disease. The one or more additional pharmaceutical agents can be administered to a patient simultaneously or sequentially.

Pharmaceutical Formulations and Dosage Forms

5 In certain embodiments, the compounds of the invention can be administered in the form of pharmaceutical compositions. These compositions can be prepared in a manner well known in the pharmaceutical art, and can be administered by a variety of routes, depending upon whether local or systemic treatment is desired and upon the area to be treated. Administration may be topical (including transdermal, epidermal, ophthalmic and to mucous
10 membranes including intranasal, vaginal and rectal delivery), pulmonary (e.g., by inhalation or insufflation of powders or aerosols, including by nebulizer; intratracheal or intranasal), oral or parenteral. Parenteral administration includes intravenous, intraarterial, subcutaneous, intraperitoneal intramuscular or injection or infusion; or intracranial, e.g., intrathecal or intraventricular, administration. Parenteral administration can be in the form of a single bolus
15 dose, or may be, for example, by a continuous perfusion pump. Pharmaceutical compositions and formulations for topical administration may include transdermal patches, ointments, lotions, creams, gels, drops, suppositories, sprays, liquids and powders. Conventional pharmaceutical carriers, aqueous, powder or oily bases, thickeners and the like may be necessary or desirable. Coated condoms, gloves and the like may also be useful.

20 Embodiments of this invention also include pharmaceutical compositions which contain, as the active ingredient, one or more of the compounds of the invention above in combination with one or more pharmaceutically acceptable carriers (excipients). In making the compositions of the invention, the active ingredient is typically mixed with an excipient, diluted by an excipient or enclosed within such a carrier in the form of, for example, a
25 capsule, sachet, paper, or other container. When the excipient serves as a diluent, it can be a solid, semi-solid, or liquid material, which acts as a vehicle, carrier or medium for the active ingredient. Thus, the compositions can be in the form of tablets, pills, powders, lozenges, sachets, cachets, elixirs, suspensions, emulsions, solutions, syrups, aerosols (as a solid or in
30 a liquid medium), ointments containing, for example, up to 10% by weight of the active compound, soft and hard gelatin capsules, suppositories, sterile injectable solutions, and sterile packaged powders.

In preparing a formulation, the active compound can be milled to provide the appropriate particle size prior to combining with the other ingredients. If the active
35 compound is substantially insoluble, it can be milled to a particle size of less than 200 mesh.

If the active compound is substantially water soluble, the particle size can be adjusted by milling to provide a substantially uniform distribution in the formulation, *e.g.* about 40 mesh.

The compounds of the invention may be milled using known milling procedures such as wet milling to obtain a particle size appropriate for tablet formation and for other
5 formulation types. Finely divided (nano particulate) preparations of the compounds of the invention can be prepared by processes known in the art, for example see International Patent Application No. WO 2002/000196.

Some examples of suitable excipients include lactose, dextrose, sucrose, sorbitol, mannitol, starches, gum acacia, calcium phosphate, alginates, tragacanth, gelatin, calcium
10 silicate, microcrystalline cellulose, polyvinylpyrrolidone, cellulose, water, syrup, and methyl cellulose. The formulations can additionally include: lubricating agents such as talc, magnesium stearate, and mineral oil; wetting agents; emulsifying and suspending agents; preserving agents such as methyl- and propylhydroxy-benzoates; sweetening agents; and flavoring agents. The compositions of the invention can be formulated so as to provide quick,
15 sustained or delayed release of the active ingredient after administration to the patient by employing procedures known in the art.

The compositions can be formulated in a unit dosage form, each dosage containing from about 5 to about 1000 mg (1 g), more usually about 100 to about 500 mg, of the active ingredient. The term "unit dosage forms" refers to physically discrete units suitable as unitary
20 dosages for human subjects and other mammals, each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect, in association with a suitable pharmaceutical excipient.

The active compound can be effective over a wide dosage range and can be generally administered in a pharmaceutically effective amount. For example, the dosage of
25 the active compounds of the invention as employed for the treatment of a patient in need thereof (such as an adult human) may range from 0.1 to 3000 mg per day, depending on the route and frequency of administration. Such a dosage corresponds to 0.001 to 50 mg/kg per day. In some embodiments, the dosage of the active compounds of the invention as employed for the treatment of a patient in need thereof (such as an adult human) may range
30 from 1 to 2000 mg per day, from 1 to 1000 mg per day, from 10 to 1000 mg per day, or from 10 to 500 mg per day. It will be understood, however, that the amount of the compound actually administered will usually be determined by a physician, according to the relevant circumstances, including the condition to be treated, the chosen route of administration, the actual compound administered, the age, weight, and response of the individual patient, the
35 severity of the patient's symptoms, and the like.

For preparing solid compositions such as tablets, the principal active ingredient can be mixed with a pharmaceutical excipient to form a solid pre-formulation composition containing a homogeneous mixture of a compound of the present invention. When referring to these pre-formulation compositions as homogeneous, the active ingredient is typically dispersed evenly throughout the composition so that the composition can be readily subdivided into equally effective unit dosage forms such as tablets, pills and capsules. This solid pre-formulation is then subdivided into unit dosage forms of the type described above containing from, for example, about 0.1 to about 1000 mg of the active ingredient of the present invention.

The tablets or pills of the present invention can be coated or otherwise compounded to provide a dosage form affording the advantage of prolonged action. For example, the tablet or pill can comprise an inner dosage and an outer dosage component, the latter being in the form of an envelope over the former. The two components can be separated by an enteric layer which serves to resist disintegration in the stomach and permit the inner component to pass intact into the duodenum or to be delayed in release. A variety of materials can be used for such enteric layers or coatings, such materials including a number of polymeric acids and mixtures of polymeric acids with such materials as shellac, cetyl alcohol, and cellulose acetate.

The liquid forms in which the compounds and compositions of the present invention can be incorporated for administration orally or by injection include aqueous solutions, suitably flavored syrups, aqueous or oil suspensions, and flavored emulsions with edible oils such as cottonseed oil, sesame oil, coconut oil, or peanut oil, as well as elixirs and similar pharmaceutical vehicles.

Compositions for inhalation or insufflation include solutions and suspensions in pharmaceutically acceptable, aqueous or organic solvents, or mixtures thereof, and powders. The liquid or solid compositions may contain suitable pharmaceutically acceptable excipients as described *supra*. In some embodiments, the compositions are administered by the oral or nasal respiratory route for local or systemic effect. Compositions in can be nebulized by use of inert gases. Nebulized solutions may be breathed directly from the nebulizing device or the nebulizing device can be attached to a face masks tent, or intermittent positive pressure breathing machine. Solution, suspension, or powder compositions can be administered orally or nasally from devices which deliver the formulation in an appropriate manner.

The amount of compound or composition administered to a patient will vary depending upon what is being administered, the purpose of the administration, such as prophylaxis or therapy, the state of the patient, the manner of administration, and the like. In

therapeutic applications, compositions can be administered to a patient already suffering from a disease in an amount sufficient to cure or at least partially arrest the symptoms of the disease and its complications. Effective doses will depend on the disease condition being treated as well as by the judgment of the attending clinician depending upon factors such as the severity of the disease, the age, weight and general condition of the patient, and the like.

The compositions administered to a patient can be in the form of pharmaceutical compositions described above. These compositions can be sterilized by conventional sterilization techniques, or may be sterile filtered. Aqueous solutions can be packaged for use as is, or lyophilized, the lyophilized preparation being combined with a sterile aqueous carrier prior to administration. The pH of the compound preparations typically will be between 3 and 11, more preferably from 5 to 9 and most preferably from 7 to 8. It will be understood that use of certain of the foregoing excipients, carriers, or stabilizers will result in the formation of pharmaceutical salts.

The therapeutic dosage of the compounds of the present invention can vary according to, for example, the particular use for which the treatment is made, the manner of administration of the compound, the health and condition of the patient, and the judgment of the prescribing physician. The proportion or concentration of a compound of the invention in a pharmaceutical composition can vary depending upon a number of factors including dosage, chemical characteristics (*e.g.*, hydrophobicity), and the route of administration. For example, the compounds of the invention can be provided in an aqueous physiological buffer solution containing about 0.1 to about 10% w/v of the compound for parenteral administration. Some typical dose ranges are from about 1 g/kg to about 1 g/kg of body weight per day. In some embodiments, the dose range is from about 0.01 mg/kg to about 100 mg/kg of body weight per day. The dosage is likely to depend on such variables as the type and extent of progression of the disease or disorder, the overall health status of the particular patient, the relative biological efficacy of the compound selected, formulation of the excipient, and its route of administration. Effective doses can be extrapolated from dose-response curves derived from *in vitro* or animal model test systems.

The compositions of the invention can further include one or more additional pharmaceutical agents such as a chemotherapeutic, steroid, anti-inflammatory compound, or immunosuppressant, examples of which are listed hereinabove.

Labeled Compounds and Assay Methods

Another aspect of the present invention relates to labeled compounds of the invention (radio-labeled, fluorescent-labeled, etc.) that would be useful not only in radio-imaging but also

in assays, both *in vitro* and *in vivo*, for localizing and quantitating the enzyme in tissue samples, including human, and for identifying ligands by inhibition binding of a labeled compound. Accordingly, the present invention includes enzyme assays that contain such labeled compounds.

Embodiments of the present invention further includes isotopically-labeled compounds of the invention. An "isotopically" or "radio-labeled" compound is a compound of the invention where one or more atoms are replaced or substituted by an atom having an atomic mass or mass number different from the atomic mass or mass number typically found in nature (i.e., naturally occurring). Suitable radionuclides that may be incorporated in compounds of the present invention include but are not limited to ^2H (also written as D for deuterium), ^3H (also written as T for tritium), ^{11}C , ^{13}C , ^{14}C , ^{13}N , ^{15}N , ^{15}O , ^{17}O , ^{18}O , ^{18}F , ^{35}S , ^{36}Cl , ^{82}Br , ^{75}Br , ^{76}Br , ^{77}Br , ^{123}I , ^{124}I , ^{125}I and ^{131}I . The radionuclide that is incorporated in the radio-labeled compounds will depend on the specific application of that radio-labeled compound. For example, for *in vitro* receptor labeling and competition assays, compounds that incorporate ^3H , ^{14}C , ^{82}Br , ^{125}I , ^{131}I , ^{35}S or will generally be most useful. For radio-imaging applications ^{11}C , ^{18}F , ^{125}I , ^{123}I , ^{124}I , ^{131}I , ^{75}Br , ^{76}Br or ^{77}Br will generally be most useful.

It is understood that a "radio-labeled compound" is a compound that has incorporated at least one radionuclide. In some embodiments the radionuclide is selected from ^3H , ^{14}C , ^{125}I , ^{35}S and ^{82}Br .

In some embodiments, the labeled compounds of the present invention contain a fluorescent label.

Synthetic methods for incorporating radio-isotopes and fluorescent labels into organic compounds are well known in the art.

A labeled compound of the invention (radio-labeled, fluorescent-labeled, etc.) can be used in a screening assay to identify/evaluate compounds. For example, a newly synthesized or identified compound (i.e., test compound) which is labeled can be evaluated for its ability to bind a beta-secretase or a neuron cell (or a neuron cell in the presence of one or more of beta-amyloid oligomers) by monitoring its concentration variation when contacting with the beta-secretase or the neuron cell (or the neuron cell in the presence of one or more of beta-amyloid oligomers), through tracking the labeling. For another example, a test compound (labeled) can be evaluated for its ability to reduce binding of another compound which is known to bind to beta-secretase or neuron cell (i.e., standard compound). Accordingly, the ability of a test compound to compete with the standard compound for binding to the beta-secretase or the neuron cell directly correlates to its binding affinity. Conversely, in some other screening assays, the standard compound is labeled and test compounds are unlabeled. Accordingly, the

concentration of the labeled standard compound is monitored in order to evaluate the competition between the standard compound and the test compound, and the relative binding affinity of the test compound is thus ascertained.

5 *Kits*

Embodiments of the present invention also includes pharmaceutical kits useful, for example, in the treatment or prevention of cognitive decline and/or Alzheimer's disease which include one or more containers containing a pharmaceutical composition comprising a therapeutically effective amount of a compound of the invention. Such kits can further include, if
10 desired, one or more of various conventional pharmaceutical kit components, such as, for example, containers with one or more pharmaceutically acceptable carriers, additional containers, etc., as will be readily apparent to those skilled in the art. Instructions, either as inserts or as labels, indicating quantities of the components to be administered, guidelines for administration, and/or guidelines for mixing the components, can also be included in the kit.

15

The invention will be described in greater detail by way of specific examples. The following examples are offered for illustrative purposes, and are not intended to limit the invention in any manner. Those of skill in the art will readily recognize a variety of non-critical parameters which can be changed or modified to yield essentially the same results. Certain
20 compounds of the Examples were found to be inhibit, treat, or abate one or more of amyloid production, amyloid assembly, the activity/effect of Abeta oligomers on neurons (such as neurons in the brain), amyloid aggregation, amyloid oligomer binding, and amyloid deposition according to one or more of the assays provided herein. In some further embodiments, certain compounds of the Examples were found to be inhibit, treat, or abate one or more of
25 the activity/effect of Abeta oligomers on neurons (such as neurons in the brain), amyloid aggregation, amyloid (including amyloid oligomer) binding, and amyloid deposition according to one or more of the assays provided herein.

In some embodiments, the compound of invention has an IC_{50} value of less than 100 μ M, 50 μ M, 20 μ M, 15 μ M, 10 μ M, 5 μ M, 1 μ M, 500 nM, 100 nM, 50 nM, or 10 nM with
30 respect to inhibition of one or more of the activity/effect of Abeta oligomers on neurons (such as neurons in the brain), amyloid aggregation, amyloid (including amyloid oligomer) binding, and amyloid deposition. In some embodiments, the compound of invention has an IC_{50} value of less than 100 μ M, 50 μ M, 20 μ M, 15 μ M, 10 μ M, 5 μ M, 1 μ M, 500 nM, 100 nM, 50 nM, or 10 nM with respect to inhibition the activity/effect of Abeta oligomers on neurons (such as
35 neurons in the brain).

In some embodiments, percentage inhibition of the compound of invention to one or more of the activity/effect of Abeta oligomers on neurons (such as neurons in the brain), amyloid aggregation, amyloid (including amyloid oligomer) binding, and amyloid deposition was measured at a concentration of from 10 nM to 10 μ M. In some embodiments, the percentage inhibition measured is about 1% to about 20%, about 20% to about 50%, about 1% to about 50%, or about 1% to about 80%.

The invention may be appreciated in certain aspects with reference to the following examples, offered by way of illustration, not by way of limitation. Materials, reagents and the like to which reference is made in the following examples are obtainable from commercial sources, unless otherwise noted.

EXAMPLES

Materials And Methods

Ginger oil

The light oil extract from ginger root was obtained by supercritical CO₂ extraction.

Ginger oleoresin

The heavy remainder oil was obtained following extraction of ginger root by supercritical CO₂ extraction.

Example 1

A. Conditioned Extraction of Ginger Oil: Reaction of ginger oil with 4-chlorobenzylamine followed by reduction with sodium borohydride in methanol and by fractioning using column chromatography

Ginger oil (10 g) was dissolved in toluene (250 mL) and 4-chlorobenzylamine (3.4 g) was added. The mixture was maintained under an atmosphere of nitrogen and heated at reflux with removal of water by Dean-Stark distillation for 16 hours. At this time the Dean-Stark trap was removed and the reaction mixture was cooled to 0°C on an ice bath. A solution of sodium borohydride (10 g) in methanol (100 mL) was added portion-wise over 30 minutes with vigorous stirring. When the addition was complete the mixture was heated to reflux for 16 hours. At this time the reaction mixture was cooled to room temperature and poured into saturated aqueous sodium bicarbonate solution (300 mL). The resulting mixture was concentrated by rotary evaporation and the aqueous residue was partitioned between water and chloroform. The chloroform layer was dried over anhydrous sodium sulfate and

then filtered and concentrated. The products were then fractionated using silica gel column chromatography employing a gradient from 100% chloroform to chloroform:methanol (5:1). The product was detected in the relatively polar fractions by thin layer chromatography (TLC). Product-containing fractions were combined and concentrated then dried under high vacuum overnight to provide a light brown oil (0.672 g). ¹H NMR (500 MHz, CDCl₃) δ: 7.30-7.24 (m, 4H), 6.81 (d, *J* = 7.8 Hz, 1H), 6.66-6.62 (m, 2H), 4.25 (br s, 2H), 3.82 (s, 3H), 3.82 (d, *J* = 13.2 Hz, 1H), 3.72 (d, *J* = 13.2 Hz, 1H), 2.73 (m, 1H), 2.66-2.51 (m, 1H), 1.86-1.78 (m, 1H), 1.72-1.63 (m, 1H), 1.62-1.51 (m, 1H), 1.17 (d, *J* = 6.3 Hz, 3H). ¹³C NMR (125 MHz, CDCl₃) δ: 146.6, 143.8, 133.9 132.8, 129.9, 129.7, 128.6, 120.8, 114.5, 110.9, 55.8, 51.9, 50.2, 38.5, 31.9, 31.6, 29.7, 26.9, 22.6, 19.9. Analytic MS(M+H⁺): *m/z* 320.2.

¹H NMR (500 MHz, CD₃OD) δ: 7.10-7.30 (m, 4H), 6.63 (br s, 1H), 6.58 (m, 1H), 6.48 (m, 1H), 3.68 (s, 3H), 3.65 (m, 1H), 3.58 (m, 1H), 2.57 (m, 1H), 2.50 (m, 1H), 2.35 (m, 1H), 1.73 (m, 1H), 1.49 (m, 1H), 1.04 (d, 3H). ¹³C NMR (125 MHz, CD₃OD) δ: 147.5, 145.3, 137.2, 133.3 132.6, 129.9, 128.1, 120.4, 114.7, 111.6, 54.9, 51.3, 49.2, 37.7, 31.5, 18.0.

The weight ratio of Ginger oil to 4-chlorobenzylamine used in the reductive amination is about 3:1 (from 2.7:1 to 3.3:1). The chemical shift measure by ¹H NMR may vary, for example, up to 0.2 ppm. The chemical shift measure by ¹³H NMR may vary, for example, up to 0.5 ppm. The analytical Mass Spectrum may have an experimental error of +/- 0.3.

Purity Determination

The purity of the product was measure by HPLC. The major peak of retention time of 2.22 minutes indicating greater than about 80%, 85%, 90, or 95% of purity. The HPLC conditions used are as follows.

HPLC conditions:

Mobile Phase A: 13.3 mM ammonium formate/6.7 mM formic acid in water

Mobile Phase B: 6 mM ammonium formate/3 mM formic acid in water/CH₃CN (1/9, v/v)

Column: Synergi Fusion-RP 100A Mercury, 2 x 20 mm, 2.5 micron
(Phenomenex Part No 00M-4423-B0_CE)


Gradient Program: RT = 2.22 minutes

Time, minute	% Phase B	Flow rate, ml/min
0	100	0.5
1	100	0.5
2.5	40	0.5

3.4	40	0.5
3.5	100	0.5
4.5	100	0.5

The purity of the product was also measure by ^1H NMR indicating it to be a single compound of a purity of greater than 90% or 95%.



5 The structure of Compound Example 1 (or Example Compound 1) is determined to be as follows.



Compound Example 1

4-(3-(4-chlorobenzylamino)butyl)-2-methoxyphenol

10 **B. Synthesis by reductive amination.**



Vanillylacetone (5.00 g, 25.7 mmol) was dissolved in toluene (250 mL) and 4-chlorobenzylamine (3.82 g, 27.0 mmol) was added. The mixture was maintained under an atmosphere of nitrogen and heated at reflux with removal of water by Dean-Stark distillation for 16 hours. At this time the Dean-Stark trap was removed and the reaction mixture was cooled to 0°C on an ice bath. A solution of sodium borohydride (5 g) in methanol (100 mL) was added portion-wise over 30 minutes with vigorous stirring. When the addition was complete the mixture was heated at reflux for 16 hours. At this time the reaction mixture was cooled to room temperature and poured into saturated aqueous sodium bicarbonate solution (300 mL). The resulting mixture was concentrated by rotary evaporation and the aqueous residue was partitioned between water and chloroform. The chloroform layer was dried over anhydrous sodium sulfate and then filtered and concentrated. The product was then purified using silica gel column chromatography employing a mobile phase of 5% ammonia-methanol in chloroform. Product-containing fractions were combined and concentrated then dried under high vacuum overnight to provide a light brown oil (6.16 g, 75%). ¹H NMR, ¹³C NMR, and Mass Spectrum of the product were substantially the same as those in Example 1, A (made by the Conditioned Extraction method).

Example 2

A. Conditioned Extraction of Ginger Oil: Reaction of ginger oil with 4-trifluoromethylbenzylamine followed by reduction with sodium borohydride in methanol and by fractioning using column chromatography

Ginger oil (10 g) was dissolved in toluene (250 mL) and 4-trifluoromethylbenzylamine (3.5 g) was added. The mixture was maintained under an atmosphere of nitrogen and heated at reflux with removal of water by Dean-Stark distillation for 16 hours. At this time the Dean-Stark trap was removed and the reaction mixture was cooled to 0°C on an ice bath. A solution of sodium borohydride (10 g) in methanol (100 mL) was added portion-wise over 30 minutes with vigorous stirring. When the addition was complete the mixture was heated to reflux for 16 hours. At this time the reaction mixture was cooled to room temperature and poured into saturated aqueous sodium bicarbonate solution (300 mL). The resulting mixture was concentrated by rotary evaporation and the aqueous residue was partitioned between water and chloroform. The chloroform layer was dried over anhydrous sodium sulfate and then filtered and concentrated. The products were then fractionated using silica gel column chromatography employing a gradient from 100% chloroform to chloroform:methanol (5:1). The product was detected in the relatively polar fractions by thin layer chromatography (TLC). Product-containing fractions were combined and concentrated then dried under high

vacuum overnight to provide a light brown oil (0.761 g). ^1H NMR (500 MHz, CDCl_3) δ : 7.57 (d, $J = 7.8$ Hz, 2H), 7.43 (d, $J = 7.9$ Hz, 2H), 6.82 (d, $J = 7.3$ Hz, 1H), 6.65 (m, 2H), 5.16-4.42 (br s, 2H), 3.90 (d, $J = 13.7$ Hz, 1H), 3.84 (s, 3H), 3.80 (d, $J = 13.7$ Hz, 1H), 2.76-2.70 (m, 1H), 2.67-2.55 (m, 2H), 1.84-1.77 (m, 1H), 1.69-1.63 (m, 1H), 1.17 (d, $J = 6.3$ Hz, 3H). ^{13}C NMR (125 MHz, CDCl_3) δ : 146.7, 144.6, 143.9, 134.0, 129.1, 128.4, 127.5, 125.4, 125.3, 123.2, 120.8, 114.6, 111.0, 55.7, 52.1, 50.6, 38.8, 32.0, 20.1. MS (CI) m/z 353 (M^+).

The weight ratio of ginger oil to 4-trifluoromethylbenzylamine used in the reductive amination is about 3:1 (from 2.7:1 to 3.3:1). The chemical shift measure by ^1H NMR may vary, for example, up to 0.2 ppm. The chemical shift measure by ^{13}H NMR may vary, for example, up to 0.5 ppm. The analytical Mass Spectrum may have an experimental error of +/- 0.3.

Purity Determination

The purity of the product was measure by HPLC. The major peak of retention time of 2.22 minutes indicating greater than about 80%, 85%, 90, or 95% of purity. The HPLC conditions used are as follows.

HPLC conditions:

Mobile Phase A: 13.3 mM ammonium formate/6.7 mM formic acid in water

Mobile Phase B: 6 mM ammonium formate/3 mM formic acid in water/ CH_3CN (1/9,

v/v)

Column: Synergi Fusion-RP 100A Mercury, 2 x 20 mm, 2.5 micron
(Phenomenex Part No 00M-4423-B0_CE)


Gradient Program: RT = 2.22 minutes

Time, minute	% Phase B	Flow rate, ml/min
0	100	0.5
1	100	0.5
2.5	40	0.5
3.4	40	0.5
3.5	100	0.5
4.5	100	0.5

The purity of the product was also measure by ^1H NMR indicating it to be a single compound of a purity of greater than 90% or 95%.



The structure of Compound Example 2 (or Example Compound 2) is determined to be as follows.

46



Compound Example 2

4-(3-(4-(trifluoromethyl)benzylamino)butyl)-2-methoxyphenol

5 **B. Synthesis by reductive amination.**

Vanillylacetone (5.00 g, 25.7 mmol) was dissolved in toluene (250 mL) and 4-trifluoromethylbenzylamine (4.73 g, 27.0 mmol) was added. The mixture was maintained under an atmosphere of nitrogen and heated at reflux with removal of water by Dean-Stark distillation for 16 hours. At this time the Dean-Stark trap was removed and the reaction mixture was cooled to 0°C on an ice bath. A solution of sodium borohydride (5 g) in methanol (100 mL) was added portion-wise over 30 minutes with vigorous stirring. When the addition was complete the mixture was heated at reflux for 16 hours. At this time the reaction mixture was cooled to room temperature and poured into saturated aqueous sodium bicarbonate solution (300 mL). The resulting mixture was concentrated by rotary evaporation and the aqueous residue was partitioned between water and chloroform. The chloroform layer was dried over anhydrous sodium sulfate and then filtered and concentrated. The product was then purified using silica gel column chromatography employing a mobile phase of 5% ammonia-methanol in chloroform. Product-containing fractions were combined and concentrated then dried under high vacuum overnight to

provide a light brown oil (6.72 g, 74%). ¹H NMR, ¹³C NMR, and Mass Spectrum of the product were substantially the same as those in Example 2, A (made by the Conditioned Extraction method).

5 **Example AA: Exocytosis assay/MTT assay**

Primary neurons from E18 Sprague-Dawley rat embryos are plated at optimized concentrations in 384 well plates in NB media (Invitrogen). Neurons are maintained in cultures for 3 weeks, with twice weekly feeding of NB media with N₂ supplement (Invitrogen). A test compound is added to cells, followed by addition of Vehicle or Abeta oligomer
10 preparations (1.5 μM), and incubated for 1 to 24 hr at 37 °C in 5% CO₂. MTT reagent (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide) (Roche Molecular Biochemicals) is reconstituted in phosphate buffered saline to 5mg/mL. 10 μL of MTT labeling reagent is added to each well and incubated at 37 °C for 1h, then imaged.

Each assay plate is formatted so that compounds are tested with and without Abeta
15 on each plate. This design eliminates toxic or metabolically active compounds early on in the screening cascade (at the level of the primary screen). Statistical performance of the screening plate layout are assessed, screening will be initiated if the current performance is maintained.

Similar procedures for exocytosis assays/MTT assays can be found in the literature.
20 See e.g., Liu Y, et. al., Detecting bioactive amyloid beta peptide species in Alzheimer's disease. J Neurochem. 2004 Nov;91(3):648-56; Liu Y, and Schubert D. "Cytotoxic amyloid peptides inhibit cellular 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) reduction by enhancing MTT formazan exocytosis." J Neurochem. 1997 Dec;69(6):2285-93; and Liu Y, and Schubert D. "Treating Alzheimer's disease by inactivating bioactive amyloid
25 beta peptide" Curr. Alzheimer Res. 2006 Apr;3(2):129-35.

Experimental controls:

Abeta 1-42 oligomers made according to published methods [See e.g. Dahlgren et al., "Oligomeric and fibrillar species of amyloid-beta peptides differentially affect
30 neuronal viability" J Biol Chem. 2002 Aug 30;277(35):32046-53. Epub 2002 Jun 10.; LeVine H 3rd. "Alzheimer's beta-peptide oligomer formation at physiologic concentrations" Anal Biochem. 2004 Dec 1;335(1):81-90; Shrestha et.al, "Amyloid beta peptide adversely affects spine number and motility in hippocampal neurons" Mol Cell Neurosci. 2006 Nov;33(3):274-82. Epub 2006 Sep 8; Puzzo et al., "Amyloid-beta peptide inhibits activation of the nitric
35 oxide/cGMP/cAMP-responsive element-binding protein pathway during hippocampal

synaptic plasticity" J Neurosci. 2005 Jul 20;25(29):6887-97; Barghorn et al., "Globular amyloid beta-peptide oligomer - a homogenous and stable neuropathological protein in Alzheimer's disease" J Neurochem. 2005 Nov;95(3):834-47. Epub 2005 Aug 31; Johansson et al., Physicochemical characterization of the Alzheimer's disease-related peptides A beta 1-42 Arctic and A beta 1-42wt. FEBS J. 2006 Jun;273(12):2618-30] as well as brain-derived Abeta oligomers (See e.g. Walsh et al., Naturally secreted oligomers of amyloid beta protein potentially inhibit hippocampal long-term potentiation in vivo. Nature (2002). 416, 535-539; Lesne et al., A specific amyloid-beta protein assembly in the brain impairs memory. Nature. 2006 Mar 16;440(7082):352-7; Shankar et al, Amyloid-beta protein dimers isolated directly from Alzheimer's brains impair synaptic plasticity and memory. Nat Med. 2008 Aug;14(8):837-42. Epub 2008 Jun 22) constitute the positive controls. Negative controls include vehicle-treated neurons as well as neurons treated with 28 μ M concentrations of memantine. Memantine produces 50% inhibition of oligomer effects at this dose. These controls, on each plate, serve as normalization tools to calibrate assay performance on a plate-by-plate basis.

Primary neuronal cultures

Optimal cell density is determined based on cellular response to Abeta oligomers using the exocytosis assay as a readout, and immunohistochemical analysis of the relative proportion of glia to neurons in the cultures. Cultures are monitored on a weekly basis with immunohistochemistry and image processing-based quantification to monitor the percentage of the cultures that are neurons vs. glia (Glial cells). Cultures containing more than 20% glia (positive for GFAP) vs. neurons (staining positively with antibodies directed against MAP2) at the screening age of 21 days in vitro (21 DIV) are rejected.

Abeta Oligomer preparations

Human amyloid peptide 1-42 is obtained from California Peptide, with lot-choice contingent upon quality control analysis. Quality controls of oligomer preparations consist of Westerns to determine oligomer size ranges and relative concentrations, and the MTT assay to confirm exocytosis acceleration without toxicity. Toxicity is monitored in each image-based assay via quantification of nuclear morphology visualized with the DNA binding dye DAPI (Invitrogen). Nuclei that are fragmented are considered to be in late stage apoptosis (Majno and Joris '95). Peptide lots producing unusual peptide size ranges or significant toxicity at a standard 1.5 μ M concentration on neurons are rejected. Plate-based controls –The assay optimization will be complete when reformatted plates achieve a minimum of statistically

significant two-fold separation between vehicle and Abeta oligomer-treated neurons ($p < 0.01$, Student's t-test, unequal variance) on a routine basis, with no more than 10% CV between plates, equivalent to its current performance.

5 *Statistical software and Analysis:*

Data handling and analysis are accomplished by Cellomics VTI image analysis software and STORE automated database software. Because of the low dynamic range and neuronal well-to-well variability after three weeks in culture, statistical comparisons are made via pairwise Tukey-Kramer analysis to determine the significance of the separation between
10 compound + Abeta oligomers from Abeta alone, and between compound alone from vehicle. These statistics are more akin to what is seen in animal behavioral testing than the z' statistic that has been used for the past two decades in high throughput screening. The ability of mature primary neurons to more closely approximate the electrophysiologically mediated signal transduction network of the adult brain justifies this screening strategy. Power
15 analysis will be set for a number of replicate screening wells that will minimize false negatives (e.g $N=4$) and shift the burden of distinguishing false positives from actual hits to dose-response confirmation screening. Rank ordering of compounds is done on the basis of secondary assay mechanism of action and physicochemical properties of the compound structures. Certain test compounds significantly reverse the effects of Abeta oligomers but
20 not affect neuronal metabolism.

Compound Example 1 was dosed in the MTT assay and was shown to block the Abeta oligomer-induced acceleration of exocytosis with an EC_{50} of 10 μM , indicating that Compound Example 1 blocks/abate the activity/effect of Abeta oligomer on neuron cells.

25 **Example BB: Binding Assay**

Each test compound was added to a plate followed by an addition of one or more of Abeta 1-42 Oligomers. The plates were fixed with 3.7 % paraformaldehyde in phosphate buffered saline (PBS) for 15 min. The plate was then washed 3x with PBS for 5 min each. The plates were blocked at room temperature for 1 hour in 5 % goat serum and 0.5% Triton
30 X-100 (CAS number: 9002-93-1) in PBS. Primary antibodies (anti-MAP 2 polyclonal, Millipore #AB5622 and anti-Beta Amyloid 6E10 monoclonal, Convance #SIG-39300) were diluted 1:1000 in 5 % goat serum with PBS. Primary antibodies were incubated either overnight at 4 °C or 1 hour at RT. The plate was then washed 3x with PBS for 5 min each. Secondary antibodies (Alexa Fluor 488 polyclonal, Invitrogen # A11008 and Alexa Fluor 647
35 monoclonal, Invitrogen #A21235) were diluted 1:1000 in 5 % goat serum with PBS.

Secondary antibodies were incubated at RT for 1 hr. The plates were washed once with PBS. DAPI (4',6-diamidino-2-phenylindole, Invitrogen) was then applied at 0.03 µg/µL and incubated at RT for 5 min, then washed with PBS. Image process was carried out for analysis.

- 5 Similar procedures for binding assays can be found in the literature. See e.g., Look GC, et. al. Discovery of ADDL--targeting small molecule drugs for Alzheimer's disease. Curr Alzheimer Res. 2007 Dec;4(5):562-7. Review.

Abeta oligomer preparations:

- 10 Human amyloid peptide 1-42 is obtained from California Peptide, with lot-choice contingent upon quality control analysis. Abeta 1-42 oligomers made according to published methods [See e.g. Dahlgren et al., "Oligomeric and fibrillar species of amyloid-beta peptides differentially affect neuronal viability" J Biol Chem. 2002 Aug 30;277(35):32046-53. Epub 2002 Jun 10.; LeVine H 3rd. "Alzheimer's beta-peptide oligomer formation at physiologic concentrations" Anal Biochem. 2004 Dec 1;335(1):81-90; Shrestha et.al, "Amyloid beta peptide adversely affects spine number and motility in hippocampal neurons" Mol Cell Neurosci. 2006 Nov;33(3):274-82. Epub 2006 Sep 8; Puzzo et al., "Amyloid-beta peptide inhibits activation of the nitric oxide/cGMP/cAMP-responsive element-binding protein pathway during hippocampal synaptic plasticity" J Neurosci. 2005 Jul 20;25(29):6887-97;
- 15 Barghorn et al., "Globular amyloid beta-peptide oligomer - a homogenous and stable neuropathological protein in Alzheimer's disease" J Neurochem. 2005 Nov;95(3):834-47. Epub 2005 Aug 31; Johansson et al., Physiochemical characterization of the Alzheimer's disease-related peptides A beta 1-42 Arctic and A beta 1-42wt. FEBS J. 2006 Jun;273(12):2618-30] as well as brain-derived Abeta oligomers (See e.g. Walsh et al., Naturally secreted oligomers of amyloid beta protein potently inhibit hippocampal long-term
- 20 potentiation in vivo. Nature (2002). 416, 535-539; Lesne et al., A specific amyloid-beta protein assembly in the brain impairs memory. Nature. 2006 Mar 16;440(7082):352-7; Shankar et al, Amyloid-beta protein dimers isolated directly from Alzheimer's brains impair synaptic plasticity and memory. Nat Med. 2008 Aug;14(8):837-42. Epub 2008 Jun 22) will
- 25 serve as positive controls. Quality controls of oligomer preparations consist of Westerns to determine oligomer size ranges and relative concentrations, and the MTT assay to confirm exocytosis acceleration without toxicity. Toxicity is monitored in each image-based assay via quantification of nuclear morphology visualized with the DNA binding dye DAPI (Invitrogen). Nuclei that are fragmented are considered to be in late stage apoptosis (Majno and Joris
- 30 Apoptosis, oncosis, and necrosis. An overview of cell death. Am J Pathol 1995;146:3-16).

Peptide lots producing unusual peptide size ranges or significant toxicity at standard concentrations on neurons are rejected.

Image Processing

5 Images were captured and analyzed with the Cellomics VTI automated microscope platform, using the Neuronal Profiling algorithm. For statistical analysis, a Tukey-Kramer pair-wise comparison with unequal variance was used.

Western blots

10 Samples containing Abeta 1-42 were diluted (1:5) in non-reducing lane marker sample buffer (Pierce #1859594). A 30 microliter (μ L) sample was loaded onto an eighteen well precast 4-15% Tris-HCl gel (BIORAD #345-0028). Electrophoresis was performed in a BIO-RAD Criterion precast gel system using Tris-Glycine buffer at 125 volt (V) for 90 minutes. The gels were blotted onto 0.2 μ M nitrocellulose membranes in Tris-Glycine/10%
15 methanol buffer at 30V for 120 minutes. The membranes were boiled for 5 minutes in a PBS solution and blocked over night with TBS/5% milk solution at 4 °C. The membrane was probed with 6E10-HRP (Covance #SIG-39345) diluted to 10 μ g/mL in TBS/1% milk solution for one hour at room temperature. Membrane was washed three times for 40 minutes each with a solution of TBS/0.05% tween-20 and developed with ECL reagent (BIO-RAD #162-
20 0112) for 5 minutes. Image acquisition was performed on an Alpha Innotech FluorChem Q quantitative imaging system and analyzed with AlphaView Q software.

PK studies:

PK studies are performed at CEREP Inc of Redmond WA, according to their standard
25 protocols:: The plasma samples were processed using acetonitrile precipitation and analyzed by HPLC-MS or HPLC-MS/MS. Peak areas were recorded, and the concentrations of the test compound in the unknown plasma samples were determined using the respective calibration curve. The reportable linear range of the assay was determined, along with the lower limit of quantitation (LLQ).

30

NMR spectroscopy and mass spectrometry:

Active fractions were analyzed by 1H NMR (Varian 500 MHz NMR spectrometer) and purified compounds were characterized using a combination 1D and 2D 1H NMR experiments and 13C NMR experiments. Structure proof was obtained using these NMR
35 techniques in combination with low resolution mass spectrometry to determine molecular

weight and high resolution mass spectrometry (Thermo Finnigan LCQ Ion trap) to determine composition-of-matter.

Compound Example 1 was shown to partially block binding of the Abeta oligomer ligand to neurons by 24% according to the binding assay (using imaging processing algorithm).

Example CC: Pharmacokinetic Studies

Pharmacokinetic studies were performed according to the following protocols: The plasma samples were processed using acetonitrile precipitation and analyzed by HPLC-MS or HPLC-MS/MS. Peak areas were recorded, and the concentrations of the test compound in the unknown plasma samples were determined using the respective calibration curve. The reportable linear range of the assay was determined, along with the lower limit of quantitation (LLQ). For example, Compound Example 1 was determined to have a half life of 46 minutes in the plasma of rats when injected intravenously at 1 mg/Kg.

Example DD: A primary neuron-based functional screening assay to detect small molecule Abeta oligomer blockers.

Primary rat neurons grown for at least 3 weeks *in vitro* were chosen as the basis for this screening assay. These neurons express the full complement of synaptic proteins characteristic of neurons in the mature brain, and exhibit a complex network of activity-dependent electrical signaling. Neurons and glia in such cultures have molecular signaling networks exhibiting excellent registration with intact brain circuitry, and for this reason have been used for over two decades as a model system for learning and memory (See e.g. Kaech S, Banker G. Culturing hippocampal neurons. Nat Protoc. 2006;1(5):2406-15. Epub 2007 Jan 11; See also Craig AM, Graf ER, Linhoff MW. How to build a central synapse: clues from cell culture. Trends Neurosci. 2006 Jan;29(1):8-20. Epub 2005 Dec 7. Review). More complex systems such as acute or organotypic brain slices are very useful but not amenable to high throughput screening. Immortalized or transformed neuronal cell lines are amenable to high throughput screening, but do not replicate the electrophysiological state-dependent signaling of primary neuronal cultures and are unlikely to adequately model the subtle alterations in this signaling that are caused by oligomers during the earliest manifestations of the disease state (See e.g. Görtz P, Fleischer W, Rosenbaum C, Otto F, Siebler M. Neuronal network properties of human teratocarcinoma cell line-derived neurons. Brain Res. 2004 Aug 20;1018(1):18-25). For this reason, primary neuronal cultures were

chosen because of their ability to be used in high throughput screens and fidelity to what occurs *in vivo*.

Reduced formazan was first visible in intracellular vesicles (Figure 1A). Example of neurons filled with labeled vesicles following endocytosis of dye and reduction to an insoluble purple product. (Scale bar = 20 microns in Figure 1A). Eventual formazan exocytosis was accelerated via Abeta oligomers in mature hippocampal neurons *in vitro* (Figure 1B).

Example photomicrograph of neurons covered with insoluble purple dye that have been extruded via exocytosis. The dye precipitated in the aqueous environment of the culture and formed needle-shaped crystals on the surface of the neuron. (Figure 1B). Endocytosis rate was altered in the presence of Abeta oligomers. (Figure 1C) Exocytosis rate was altered in the presence of Abeta oligomers (Figure 1D).

Since synaptic and memory deficits, and not widespread cell death, predominate at the earliest stages of Alzheimer's disease, assays that measure these changes can be used to discover small molecule inhibitors of oligomer activity. The MTT assay can be used as a measure of toxicity in cultures. Yellow tetrazolium salts were endocytosed by cells and reduced to insoluble purple formazan in the endosomal pathway. The level of purple formazan was a reflection of the number of actively metabolizing cells in culture, and reduction in the amount of formazan was taken as a measure of cell death or metabolic toxicity in culture. When observed through a microscope, the purple formazan was first visible in intracellular vesicles that fill the cell (Figure. 1A). Over time, the vesicles were exocytosed and the formazan precipitated as needle-shaped crystals on the outer surface of the plasma membrane as the insoluble formazan was exposed to the aqueous media environment (Figure 1B). Cells respond to sublethal levels of Abeta oligomers by selectively accelerating the exocytosis rate of reduced formazan, while leaving endocytosis rate unaffected, which can be seen in mature primary neurons *in vitro* and quantified these morphological shifts via automated microscopy and image processing. At a given point in time following tetrazolium salt addition to the culture well, vehicle-treated cells had the appearance of those in Figure 1A, while Abeta oligomer-treated cells had the appearance of those in Figure 1B. Under these circumstances, there was no overall change in the total amount of reduced formazan, simply a shift in its morphology. This assay is sensitive to low levels of oligomers that do not cause cell death.

Evidence suggests that Abeta oligomer-mediated reduction in neuronal surface receptor expression mediated by membrane trafficking are the basis for oligomer inhibition of electrophysiological measures of synaptic plasticity (LTP) and thus learning and memory (See Kamenetz F, Tomita T, Hsieh H, Seabrook G, Borchelt D, Iwatsubo T, Sisodia S,

Malinow R. APP processing and synaptic function. *Neuron*. 2003 Mar 27;37(6):925-37; and Hsieh H, Boehm J, Sato C, Iwatsubo T, Tomita T, Sisodia S, Malinow R. AMPAR removal underlies Abeta-induced synaptic depression and dendritic spine loss. *Neuron*. 2006 Dec 7;52(5):831-43). Measuring membrane trafficking rate changes induced by oligomers via formazan morphological shifts has been used in cell lines to discover Abeta oligomer-blocking drugs [Maezawa I, Hong HS, Wu HC, Battina SK, Rana S, Iwamoto T, Radke GA, Pettersson E, Martin GM, Hua DH, Jin LW. A novel tricyclic pyrone compound ameliorates cell death associated with intracellular amyloid-beta oligomeric complexes. *J Neurochem*. 2006 Jul;98(1):57-67; Liu Y, Schubert D. Cytotoxic amyloid peptides inhibit cellular 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) reduction by enhancing MTT formazan exocytosis. *J Neurochem*. 1997 Dec;69(6):2285-93; Liu Y, Dargusch R, Banh C, Miller CA, Schubert D. Detecting bioactive amyloid beta peptide species in Alzheimer's disease. *J Neurochem*. 2004 Nov;91(3):648-56; Liu Y, Schubert D. Treating Alzheimer's disease by inactivating bioactive amyloid beta peptide. *Curr Alzheimer Res*. 2006 Apr;3(2):129-35; Rana S, Hong HS, Barrigan L, Jin LW, Hua DH. Syntheses of tricyclic pyrones and pyridinones and protection of Abeta-peptide induced MC65 neuronal cell death. *Bioorg Med Chem Lett*. 2009 Feb 1;19(3):670-4. Epub 2008 Dec 24; and Hong HS, Maezawa I, Budamagunta M, Rana S, Shi A, Vassar R, Liu R, Lam KS, Cheng RH, Hua DH, Voss JC, Jin LW. Candidate anti-Abeta fluorene compounds selected from analogs of amyloid imaging agents. *Neurobiol Aging*. 2008 Nov 18. (Epub ahead of print)] that lower Abeta brain levels in rodents *in vivo* [Hong HS, Rana S, Barrigan L, Shi A, Zhang Y, Zhou F, Jin LW, Hua DH. Inhibition of Alzheimer's amyloid toxicity with a tricyclic pyrone molecule *in vitro* and *in vivo*. *J Neurochem*. 2009 Feb;108(4):1097-1108].

The exocytosis assay was adapted for use with mature primary neuronal cultures grown for 3 weeks *in vitro*. Abeta oligomers caused a dose-dependent decrease in the amount of intracellular vesicles (puncta) filled with reduced purple formazan (Figure. 2A, squares; 3µM dose corresponds to image in Figure 2C) as measured via image processing using a Cellomics VTI automated microscopy system. Increasing the amount of Abeta oligomers eventually resulted in overt toxicity. Thus, the concentration of neuroactive Abeta oligomers was much lower than that causing cell death. This decrease can be blocked by adding stoichiometric amounts of anti-Abeta monoclonal antibody 6E10 (IgG) to the cultures prior to oligomer addition (Figure 2A, circle; the circle corresponds to image in Figure 2D; antibody alone [down triangle] has no effect on the neurons). Several compounds were tested that have been reported to block the effects of Abeta oligomers, including the sugar alcohol scyllo-inositol (AZD-103), the nAChR antagonist hexamethonium bromide, and the

NMDAR antagonists MK-801 and none were active (Fenili et al., '07, Calabrese et al., '06, LeCor et al., '07).

The assay was optimized for performance in 384-well microtiter plates with automated liquid handling robotics for compound formatting and assay plate stamping, routinely achieving statistically significant two-fold separation between vehicle and Abeta oligomer-treated neurons (Student's t-test, unequal variance). Compounds were added to neurons first, then oligomers were added. When configured in this manner the assay was able to detect compounds that act via disruption of oligomers, inhibition of oligomer binding to neurons, or counteraction of signal transduction mechanisms of action initiated by oligomer binding.

Compounds were considered active if they significantly block Abeta-mediated changes in membrane trafficking, but do not significantly affect membrane trafficking when dosed on their own. An example is shown in Figure 2B; Compound Example 2 inhibits oligomer effects on membrane trafficking with an EC₅₀ of 7 μM.

Figure 2A shows dose-dependent decrease of intracellular formazan-filled vesicles (puncta) caused by Abeta 42 oligomer treatment acceleration of exocytosis (squares). Oligomer effects were blocked by anti Abeta IgG (circle and up triangle; circle refers to stoich amount of IgG, i.e., 3 μM of Aβ and 1.5 μM of IgG; up triangle refers to substoich IgG, i.e., 3 μM of Aβ and 0.5 μM of IgG). IgG itself (down triangle) has no effect. Figure 2B shows Example Compound 2, which inhibits oligomer effects on membrane trafficking. Figure 2C shows representative micrographs of 21DIV hippocampal neurons *in vitro* showing oligomer effects membrane trafficking (corresponding to data point 3 μM in Figure 2A); and Figure 2D shows blockade by anti-Abeta antibodies (corresponding to the circle in Figure 2A). Data were the average of 3 experiments. Scale bar = 20 micron in Figure 2D.

Example EE: Fear Conditioning Assay

Compound Example 2 was tested in an animal model of a memory-dependent behavioral task known as fear conditioning. The study protocol was designed based on published protocols (See e.g. Puzzo D, Privitera L, Leznik E, Fà M, Staniszewski A, Palmeri A, Arancio O. Picomolar amyloid-beta positively modulates synaptic plasticity and memory in hippocampus. J Neurosci. 2008 Dec 31;28(53):14537-45.). The formation of contextual memories is dependent upon the integrity of medial temporal lobe structures such as the hippocampus. In this assay mice were trained to remember that a particular salient context (conditioned stimulus; CS) is associated with an aversive event, in this case a mild foot shock (the unconditioned stimulus, US). Animals that show good learning will express an

increase in freezing behavior when placed back into the same context. This freezing is absent in a novel context. Increased freezing in the context indicates strong hippocampal-dependent memory formation in animals. Memory tested in Fear Conditioning is sensitive to elevations of soluble A β . Figure 3 shows the results of administration of Abeta oligomers (bar labeled with "a") during training results in memory deficits when animals are tested 24 later, compared to vehicle administration (bar labeled with "b"). Example Compound 2 was effective at stopping Abeta oligomer mediated effects on membrane trafficking (Figure. 3). When administered to animals prior to Abeta oligomer administration, Example Compound 2 blocked oligomer effects on memory in a dose-dependent manner. The compound completely blocked oligomer-mediated memory deficits at the 2 pmol dose (Figure 3, bar labeled with "d"). This behavioral efficacy demonstrates that the membrane trafficking assay is able to predict which compounds will be efficacious in treating the behavioral memory loss caused by oligomers. The fear condition model for memory was performed as described herein.

Figure 3 shows that Abeta produces significant deficits in memory formation vs. vehicle ($p < 0.05$) in the contextual fear conditioning memory task. Figure 3 shows that the 2 pmol dose of Compound Example 2 + Abeta (200nM) completely blocked the effect of Abeta on memory ($p < 0.05$, one way ANOVA, post hoc comparison with Bonferroni correction). No effect of compound alone was observed (data not shown). No adverse behavioral changes were observed at any dose.

Example FF: Membrane Trafficking Assay:

Abeta assemblies were isolated from patients with Alzheimer's Disease (AD) or from normal patients. The Abeta assemblies were tested for their ability to modulate membrane trafficking. HMW (>100KDa) Abeta assemblies isolated from AD patients do not affect membrane trafficking (not shown). IMW (10-100KDa) Abeta assemblies isolated from AD patients significantly affect membrane trafficking. (Figure 4). IMW Abeta assemblies isolated from Age-matched normal individuals do not affect membrane trafficking (Figure 4). Compound Example 2 has no effect on Abeta assemblies isolated from Age-matched normal individuals. (Figure 4). Compound Example 2 significantly blocked the trafficking effects of AD-brain derived Abeta assemblies. (Figure 4).

Although the present invention has been described in considerable detail with reference to certain preferred versions thereof, other versions are possible. Therefore, the spirit and scope of the invention should not be limited to the description of the preferred

versions described herein.


All features disclosed in the specification, including the abstract and drawings, and all the steps in any method or process disclosed, may be combined in any combination, except combinations where at least some of such features and/or steps are mutually exclusive.

- 5 Each feature disclosed in the specification, including abstract and drawings, can be replaced by alternative features serving the same, equivalent or similar purpose, unless expressly stated otherwise. Thus, unless expressly stated otherwise, each feature disclosed is one example only of a generic series of equivalent or similar features. Various modifications of the invention, in addition to those described herein, will be apparent to those skilled in the art
10 from the foregoing description. Such modifications are also intended to fall within the scope of the appended claims.

Each reference cited in the present application is herein incorporated by reference in its entirety.


What is claimed is:

1. A compound of Formula I:



or pharmaceutically acceptable salt thereof, wherein:

R^1 is selected from (A1) and (A2):



R^2 , R^3 , R^4 , R^5 , and R^6 are each, independently, selected from H, OH, C_{1-6} alkoxy, C_{1-6} haloalkoxy, halo, CN, NO_2 , C_{1-6} alkyl, C_{1-6} haloalkyl, C_{3-7} cycloalkyl, NH_2 , $NH(C_{1-4} \text{ alkyl})$, $NH(C_{3-7} \text{ cycloalkyl})$, $N(C_{1-4} \text{ alkyl})_2$, $NHC(O)(C_{1-4} \text{ alkyl})$, SH, $S(C_{1-6} \text{ alkyl})$, $C(O)OR^a$, $C(O)R^b$, $C(O)NR^cR^d$, $OC(O)R^b$, $OC(O)NR^cR^d$, NR^cR^d , $NR^cC(O)R^b$, $NR^cC(O)OR^a$, $NR^cS(O)_2R^b$, $NR^cS(O)_2NR^cR^d$, $S(O)R^b$, $S(O)_2R^b$, and $S(O)_2NR^cR^d$;

R^7 is H, C_{1-6} alkyl, C_{1-6} haloalkyl, or C_{3-7} cycloalkyl;

R^8 is C_{1-6} alkyl, C_{1-6} haloalkyl, or C_{3-7} cycloalkyl;

R^9 is H, C_{1-6} alkyl, C_{1-6} haloalkyl, or C_{3-7} cycloalkyl;

R^{10} is H, C_{1-6} alkyl, C_{1-6} haloalkyl, or C_{3-7} cycloalkyl;

R^{11} is H, C_{1-6} alkyl, C_{1-6} haloalkyl, or C_{3-7} cycloalkyl;

R^{12} , R^{13} , R^{14} , R^{15} , and R^{16} are each, independently, selected from H, OH, C_{1-6} alkoxy, C_{1-6} haloalkoxy, halo, CN, NO_2 , C_{1-6} alkyl, C_{1-6} haloalkyl, C_{3-7} cycloalkyl, NH_2 , $NH(C_{1-4}$ alkyl), $NH(C_{3-7}$ cycloalkyl), $N(C_{1-4}$ alkyl) $_2$, $NHC(O)(C_{1-4}$ alkyl), SH, $S(C_{1-6}$ alkyl), $C(O)OR^{a1}$, $C(O)R^{b1}$, $C(O)NR^{c1}R^{d1}$, $OC(O)R^{b1}$, $OC(O)NR^{c1}R^{d1}$, $NR^{c1}R^{d1}$, $NR^{c1}C(O)R^{b1}$, $NR^{c1}C(O)OR^{a1}$, $NR^{c1}S(O)_2R^{b1}$, $NR^{c1}S(O)_2NR^{c1}R^{d1}$, $S(O)R^{b1}$, $S(O)_2R^{b1}$, and $S(O)_2NR^{c1}R^{d1}$;

each R^a is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, heterocycloalkylalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, aryl, cycloalkyl, heteroaryl and heterocycloalkyl, wherein each of the C_{1-6} alkyl, C_{1-6} haloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, heterocycloalkylalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, aryl, cycloalkyl, heteroaryl and heterocycloalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from OH, CN, amino, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{1-6} alkoxy, C_{1-6} haloalkoxy, aryl, arylalkyl, heteroaryl, heteroarylalkyl, cycloalkyl and heterocycloalkyl;

each R^b is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, heterocycloalkylalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl and heterocycloalkylalkyl, wherein each of the C_{1-6} alkyl, C_{1-6} haloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, heterocycloalkylalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl and heterocycloalkylalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from OH, amino, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{1-6} alkoxy, C_{1-6} haloalkoxy, aryl, arylalkyl, heteroaryl, heteroarylalkyl, cycloalkyl and heterocycloalkyl;

R^c and R^d are independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, heterocycloalkylalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, aryl, heteroaryl, cycloalkyl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl and heterocycloalkylalkyl, wherein each of the C_{1-6} alkyl, C_{1-6} haloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, heterocycloalkylalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, aryl, heteroaryl, cycloalkyl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl and heterocycloalkylalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from OH, amino, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{1-6} alkoxy, C_{1-6} haloalkoxy, aryl, arylalkyl, heteroaryl, heteroarylalkyl, cycloalkyl and heterocycloalkyl;

or R^c and R^d together with the N atom to which they are attached form a 4-, 5-, 6- or 7-membered heterocycloalkyl group that is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from OH, amino, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{1-6}

alkoxy, C₁₋₆ haloalkoxy, aryl, arylalkyl, heteroaryl, heteroarylalkyl, cycloalkyl, and heterocycloalkyl;

each R^{a1} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, heterocycloalkylalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl and heterocycloalkyl, wherein each of the C₁₋₆ alkyl, C₁₋₆ haloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, heterocycloalkylalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl and heterocycloalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from OH, CN, amino, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkoxy, aryl, arylalkyl, heteroaryl, heteroarylalkyl, cycloalkyl, and heterocycloalkyl;

each R^{b1} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, heterocycloalkylalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl and heterocycloalkylalkyl, wherein each of the C₁₋₆ alkyl, C₁₋₆ haloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, heterocycloalkylalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl and heterocycloalkylalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from OH, amino, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkoxy, aryl, arylalkyl, heteroaryl, heteroarylalkyl, cycloalkyl, and heterocycloalkyl;

R^{c1} and R^{d2} are independently selected from H, C₁₋₁₀ alkyl, C₁₋₆ haloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, heterocycloalkylalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, heteroaryl, cycloalkyl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl and heterocycloalkylalkyl, wherein each of the C₁₋₁₀ alkyl, C₁₋₆ haloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, heterocycloalkylalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, heteroaryl, cycloalkyl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl and heterocycloalkylalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from OH, amino, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkoxy, aryl, arylalkyl, heteroaryl, heteroarylalkyl, cycloalkyl, and heterocycloalkyl;

or R^{c1} and R^{d1} together with the N atom to which they are attached form a 4-, 5-, 6- or 7-membered heterocycloalkyl group that is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from OH, amino, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkoxy, aryl, arylalkyl, heteroaryl, heteroarylalkyl, cycloalkyl, and heterocycloalkyl; and

m is 0, 1, or 2,
with the proviso that

(a) when R^1 is a moiety of (A1), then two of R^2 , R^3 , R^4 , R^5 , and R^6 are independently selected from OH, C_{1-6} alkoxy, and C_{1-6} haloalkoxy; and

(b) when R^1 is a moiety of (A1), then at least one of R^{12} , R^{13} , R^{14} , R^{15} , and R^{16} is other than H.

2. The compound of claim 1 or pharmaceutically acceptable salt thereof, wherein two of R^2 , R^3 , R^4 , R^5 , and R^6 are independently selected from OH, C_{1-6} alkoxy, and C_{1-6} haloalkoxy.

3. The compound of claim 1 or 2 or pharmaceutically acceptable salt thereof, wherein R^2 , R^3 , R^4 , R^5 , and R^6 are each, independently, selected from H, OH, C_{1-6} alkoxy, C_{1-6} haloalkoxy, halo, CN, NO_2 , C_{1-6} alkyl, C_{1-6} haloalkyl, C_{3-7} cycloalkyl, NH_2 , $NH(C_{1-4}$ alkyl), $NH(C_{3-7}$ cycloalkyl), $N(C_{1-4}$ alkyl)₂, $NHC(O)(C_{1-4}$ alkyl), SH, $S(C_{1-6}$ alkyl), $C(O)OH$, $C(O)O(C_{1-4}$ alkyl), $C(O)(C_{1-4}$ alkyl), and $C(O)NH(C_{1-4}$ alkyl).

4. The compound of claim 1 or 2 or pharmaceutically acceptable salt thereof, wherein:
one of R^2 , R^3 , R^4 , R^5 , and R^6 is OH; and
one of R^2 , R^3 , R^4 , R^5 , and R^6 is OH, C_{1-6} alkoxy, or C_{1-6} haloalkoxy.

5. The compound of claim 1 or 2 or pharmaceutically acceptable salt thereof, wherein:
one of R^2 , R^3 , R^4 , R^5 , and R^6 is OH; and
one of R^2 , R^3 , R^4 , R^5 , and R^6 is C_{1-3} alkoxy or C_{1-3} haloalkoxy.

6. The compound of claim 1 or 2 or pharmaceutically acceptable salt thereof, wherein:
one of R^2 , R^3 , R^4 , R^5 , and R^6 is OH; and
one of R^2 , R^3 , R^4 , R^5 , and R^6 is methoxy or trihalomethoxy.


7. The compound of claim 1 or 2 or pharmaceutically acceptable salt thereof, wherein:
one of R^2 , R^3 , R^4 , R^5 , and R^6 is OH; and
one of R^2 , R^3 , R^4 , R^5 , and R^6 is methoxy.

8. The compound of claim 1 or 2 or pharmaceutically acceptable salt thereof, wherein:
 R^4 is OH; and
 R^5 is methoxy.

9. The compound of claim 1 or 2 or pharmaceutically acceptable salt thereof,, wherein:
R⁴ is OH;
R⁵ is methoxy; and
R², R³, and R⁶ are each H.
10. The compound of any one of claims 1-9 or pharmaceutically acceptable salt thereof,
wherein R⁷ is H or C₁₋₆ alkyl.
11. The compound of any one of claims 1-9 or pharmaceutically acceptable salt thereof,
wherein R⁷ is H or C₁₋₃ alkyl.
12. The compound of any one of claims 1-9 or pharmaceutically acceptable salt thereof,
wherein R⁷ is C₁₋₃ alkyl.
13. The compound of any one of claims 1-9 or pharmaceutically acceptable salt thereof,
wherein R⁷ is H.
14. The compound of any one of claims 1-13 or pharmaceutically acceptable salt thereof,
wherein R⁸ is C₁₋₆ alkyl.
15. The compound of any one of claims 1-13 or pharmaceutically acceptable salt thereof,
wherein R⁸ is C₁₋₃ alkyl.
16. The compound of any one of claims 1-13 or pharmaceutically acceptable salt thereof,
wherein R⁸ is methyl.
17. The compound of any one of claims 1-16 or pharmaceutically acceptable salt thereof,
wherein R⁹ is H or C₁₋₆ alkyl.
18. The compound of any one of claims 1-16 or pharmaceutically acceptable salt thereof,
wherein R⁹ is H or C₁₋₃ alkyl.
19. The compound of any one of claims 1-16 or pharmaceutically acceptable salt thereof,
wherein R⁹ is H.


20. The compound of any one of claims 1-16 or pharmaceutically acceptable salt thereof, wherein R^9 is C_{1-3} alkyl.
21. The compound of any one of claims 1-20 or pharmaceutically acceptable salt thereof, wherein R^{10} is H or C_{1-6} alkyl.
22. The compound of any one of claims 1-20 or pharmaceutically acceptable salt thereof, wherein R^{10} is H or C_{1-3} alkyl.
23. The compound of any one of claims 1-20 or pharmaceutically acceptable salt thereof, wherein R^{10} is H.
24. The compound of any one of claims 23 or pharmaceutically acceptable salt thereof, wherein R^{11} is H or C_{1-6} alkyl.
25. The compound of any one of claims 1-23 or pharmaceutically acceptable salt thereof, wherein R^{11} is H or C_{1-3} alkyl.
26. The compound of any one of claims 1-23 or pharmaceutically acceptable salt thereof, wherein R^{11} is H.
27. The compound of any one of claims 1-26 or pharmaceutically acceptable salt thereof, wherein at least one of R^{12} , R^{13} , R^{14} , R^{15} , and R^{16} is other than H.
28. The compound of any one of claims 1-27 or pharmaceutically acceptable salt thereof, wherein R^{12} , R^{13} , R^{14} , R^{15} , and R^{16} are each, independently, selected from H, OH, C_{1-6} alkoxy, C_{1-6} haloalkoxy, halo, CN, NO_2 , C_{1-6} alkyl, C_{1-6} haloalkyl, C_{3-7} cycloalkyl, NH_2 , $NH(C_{1-4}$ alkyl), $NH(C_{3-7}$ cycloalkyl), $N(C_{1-4}$ alkyl) $_2$, $NHC(O)(C_{1-4}$ alkyl), SH, $S(C_{1-6}$ alkyl), $C(O)OH$, $C(O)O(C_{1-4}$ alkyl), $C(O)(C_{1-4}$ alkyl), and $C(O)NH(C_{1-4}$ alkyl).
29. The compound of any one of claims 1-27 or pharmaceutically acceptable salt thereof, wherein R^{12} , R^{13} , R^{14} , R^{15} , and R^{16} are each, independently, selected from H, halo, CN, NO_2 , C_{1-6} alkyl, C_{1-6} haloalkyl, C_{3-7} cycloalkyl, $C(O)O(C_{1-4}$ alkyl), $C(O)(C_{1-4}$ alkyl), and $C(O)NH(C_{1-4}$ alkyl).

30. The compound of any one of claims 1-27 or pharmaceutically acceptable salt thereof, wherein at least one of R^{12} , R^{13} , R^{14} , R^{15} , and R^{16} is selected from halo, CN, NO_2 , C_{1-6} haloalkyl, $\text{C}(\text{O})\text{O}(\text{C}_{1-4}$ alkyl), $\text{C}(\text{O})(\text{C}_{1-4}$ alkyl), and $\text{C}(\text{O})\text{NH}(\text{C}_{1-4}$ alkyl).
31. The compound of any one of claims 1-27 or pharmaceutically acceptable salt thereof, wherein at least one of R^{12} , R^{13} , R^{14} , R^{15} , and R^{16} is selected from halo and C_{1-6} haloalkyl.
32. The compound of any one of claims 1-27 or pharmaceutically acceptable salt thereof, wherein R^{14} is halo.
33. The compound of any one of claims 1-27 or pharmaceutically acceptable salt thereof, wherein R^{14} is C_{1-6} haloalkyl.
34. The compound of any one of claims 1-27 or pharmaceutically acceptable salt thereof, wherein R^{14} and R^{15} are independently halo.
35. The compound of any one of claims 1-34 or pharmaceutically acceptable salt thereof, wherein the compound is a compound of Formula II:




II.

36. The compound of claim 35 or pharmaceutically acceptable salt thereof, wherein the compound is a compound of Formula IIa:



IIa.


37. The compound of claim 35 or pharmaceutically acceptable salt thereof, wherein the compound is a compound of Formula IIb:



IIb.

38. The compound of any one of claims 1-34 or pharmaceutically acceptable salt thereof, wherein the compound is a compound of Formula III:


66



III.

39. The compound of claim 38 or pharmaceutically acceptable salt thereof, wherein m is 1.
40. The compound of claim 38 or pharmaceutically acceptable salt thereof, wherein m is 0.
41. A compound that is 4-(3-(4-chlorobenzylamino)butyl)-2-methoxyphenol or 4-(3-(4-(trifluoromethyl)benzylamino)butyl)-2-methoxyphenol, or pharmaceutically acceptable salt thereof.
42. The compound of claim 41 or pharmaceutically acceptable salt thereof, wherein the compound or pharmaceutically acceptable salt thereof has a purity of greater than 80%, 90%, 95%, or 99% by weight.
43. A pharmaceutical composition comprising a compound of any one of claims 1-42 or pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.


44. A method of preparing a composition comprising a compound of Formula IIa:



IIa

or a pharmaceutically acceptable salt thereof, comprising

- (a) reacting a ginger oil with an amine of Formula XI:



XI

under reactive amination condition to produce the compound of Formula IIa; and

- (b) isolating a composition comprising the compound of Formula IIa or a pharmaceutically acceptable salt thereof, wherein the composition comprises at least 80%, 85%, 90%, or 95% by weight of the compound of Formula IIa or a pharmaceutically acceptable salt thereof,

wherein:

R^{10} is H, C_{1-6} alkyl, C_{1-6} haloalkyl, or C_{3-7} cycloalkyl;

R^{11} is H, C_{1-6} alkyl, C_{1-6} haloalkyl, or C_{3-7} cycloalkyl;

R^{12} , R^{13} , R^{14} , R^{15} , and R^{16} are each, independently, selected from H, OH, C_{1-6} alkoxy, C_{1-6} haloalkoxy, halo, CN, NO_2 , C_{1-6} alkyl, C_{1-6} haloalkyl, C_{3-7} cycloalkyl, NH_2 , $NH(C_{1-4}$ alkyl), $NH(C_{3-7}$ cycloalkyl), $N(C_{1-4}$ alkyl) $_2$, $NHC(O)(C_{1-4}$ alkyl), SH, $S(C_{1-6}$ alkyl), $C(O)OR^{a1}$, $C(O)R^{b1}$, $C(O)NR^{c1}R^{d1}$, $OC(O)R^{b1}$, $OC(O)NR^{c1}R^{d1}$, $NR^{c1}R^{d1}$, $NR^{c1}C(O)R^{b1}$, $NR^{c1}C(O)OR^{a1}$, $NR^{c1}S(O)_2R^{b1}$, $NR^{c1}S(O)_2NR^{c1}R^{d1}$, $S(O)R^{b1}$, $S(O)_2R^{b1}$, and $S(O)_2NR^{c1}R^{d1}$;

each R^{a1} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, heterocycloalkylalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, aryl, cycloalkyl, heteroaryl and heterocycloalkyl, wherein each of the C_{1-6} alkyl, C_{1-6} haloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, heterocycloalkylalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, aryl, cycloalkyl, heteroaryl and heterocycloalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from OH, CN, amino, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{1-6} alkoxy, C_{1-6} haloalkoxy, aryl, arylalkyl, heteroaryl, heteroarylalkyl, cycloalkyl, and heterocycloalkyl;

each R^{b1} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, heterocycloalkylalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl and heterocycloalkylalkyl, wherein each of the C_{1-6} alkyl, C_{1-6} haloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, heterocycloalkylalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl and heterocycloalkylalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from OH, amino, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{1-6} alkoxy, C_{1-6} haloalkoxy, aryl, arylalkyl, heteroaryl, heteroarylalkyl, cycloalkyl, and heterocycloalkyl;

R^{c1} and R^{d2} are independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, heterocycloalkylalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, aryl, heteroaryl, cycloalkyl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl and heterocycloalkylalkyl, wherein each of the C_{1-6} alkyl, C_{1-6} haloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, heterocycloalkylalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, aryl, heteroaryl, cycloalkyl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl and heterocycloalkylalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from OH, amino, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{1-6} alkoxy, C_{1-6} haloalkoxy, aryl, arylalkyl, heteroaryl, heteroarylalkyl, cycloalkyl, and heterocycloalkyl;

or R^{c1} and R^{d1} together with the N atom to which they are attached form a 4-, 5-, 6- or 7-membered heterocycloalkyl group that is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from OH, amino, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{1-6} alkoxy, C_{1-6} haloalkoxy, aryl, arylalkyl, heteroaryl, heteroarylalkyl, cycloalkyl, and heterocycloalkyl; and

m is 0, 1, or 2,

with the proviso that at least one of R^{12} , R^{13} , R^{14} , R^{15} , and R^{16} is other than H.

45. A compound or pharmaceutically acceptable salt thereof formed by a process comprising:
- (a) reacting a ginger oil with 4-chlorobenzylamine under reactive amination condition to produce a crude product, wherein the ratio of the ginger oil to 4-chlorobenzylamine about 3 :1 by weight; and
 - (b) isolating a composition comprising at least 80%, 85%, 90%, or 95% by weight of a compound or a pharmaceutically acceptable salt thereof,
- wherein:
- the compound has an analytical mass spectrum peak of $[M+H]^+$ at about 320.2.
46. The compound of claim 45 or pharmaceutically acceptable salt thereof wherein the 1H NMR (500 MHz, $CDCl_3$) spectrum of the compound comprises the following peaks: 7.30-7.24 (m), 6.81 (d), 6.66-6.62 (m), 4.25 (br s), 3.82 (s), 3.82 (d), 3.72 (d), 2.73 (m), 2.66-2.51 (m), 1.86-1.78 (m), 1.72-1.63 (m), 1.62-1.51 (m), and 1.17 (d).
47. The compound of claim 46 or pharmaceutically acceptable salt thereof wherein the ^{13}C NMR (125 MHz, $CDCl_3$) spectrum of the compound comprises at least 16 peaks selected from the following chemical shifts: 146.6, 143.8, 133.9, 132.8, 129.9, 129.7, 128.6, 120.8, 114.5, 110.9, 55.8, 51.9, 50.2, 38.5, 31.9, 31.6, 29.7, 26.9, 22.6, and 19.9.
48. The compound of claim 45 or pharmaceutically acceptable salt thereof wherein the 1H NMR (500 MHz, CD_3OD) spectrum of the compound comprises the following peaks: 7.10-7.30 (m, 4H), 6.63 (br s, 1H), 6.58 (m, 1H), 6.48 (m, 1H), 3.68 (s, 3H), 3.65 (m, 1H), 3.58 (m, 1H), 2.57 (m, 1H), 2.50 (m, 1H), 2.35 (m, 1H), 1.73 (m, 1H), 1.49 (m, 1H), and 1.04 (d, 3H).
49. The compound of claim 48 or pharmaceutically acceptable salt thereof wherein the ^{13}C NMR (125 MHz, CD_3OD) spectrum of the compound comprises the following peaks at: 147.5, 145.3, 137.2, 133.3, 132.6, 129.9, 128.1, 120.4, 114.7, 111.6, 54.9, 51.3, 49.2, 37.7, 31.5, and 18.0.
50. A compound or pharmaceutically acceptable salt thereof formed by a process comprising:
- (a) reacting a ginger oil with 4-trifluoromethylbenzylamine under reactive amination condition to produce a crude product, wherein the ratio of the ginger oil to 4-trifluoromethylbenzylamine about 3 :1 by weight; and

(b) isolating a composition comprising at least 80%, 85%, 90%, or 95% by weight of a compound or a pharmaceutically acceptable salt thereof,

wherein:

the compound has an analytical mass spectrum peak of $[M+H]^+$ at about 353.

51. The compound of claim 50 or pharmaceutically acceptable salt thereof wherein the 1H NMR (500 MHz, $CDCl_3$) spectrum of the compound comprises the following peaks: 7.57 (d, 2H), 7.43 (d, 2H), 6.82 (d, 1H), 6.65 (m, 2H), 5.16-4.42 (br s, 2H), 3.90 (d, 1H), 3.84 (s, 3H), 3.80 (d, 1H), 2.76-2.70 (m, 1H), 2.67-2.55 (m, 2H), 1.84-1.77 (m, 1H), 1.69-1.63 (m, 1H), and 1.17 (d, 3H).

52. The compound of claim 51 or pharmaceutically acceptable salt thereof wherein the ^{13}C NMR (125 MHz, $CDCl_3$) spectrum of the compound comprises at least 17 peaks selected from the following chemical shifts: 146.7, 144.6, 143.9, 134.0, 129.1, 128.4, 127.5, 125.4, 125.3, 123.2, 120.8, 114.6, 111.0, 55.7, 52.1, 50.6, 38.8, 32.0, 20.1.

53. A method of inhibiting, treating, and/or abatement of cognitive decline and/or Alzheimer's disease in a patient comprising administering to the patient a compound of any one of claims 1-42 or a pharmaceutically acceptable salt thereof.

54. The method of claim 53 wherein the method of inhibiting, treating, and/or abatement of cognitive decline and/or Alzheimer's disease comprises inhibiting, treating, or abatement of one or more symptoms of cognitive decline selected from the group consisting of memory loss, confusion, impaired judgment, personality changes, disorientation, and loss of language skills.

55. The method of claim 53 wherein the method of inhibiting, treating, and/or abatement of cognitive decline and/or Alzheimer's disease comprises one or more of:

- (i) restoration of long term potentiation; and/or
- (ii) inhibiting, treating, and/or abatement of neurodegeneration; and/or
- (iii) inhibiting, treating, and/or abatement of general amyloidosis; and/or
- (iv) inhibiting, treating, and/or abatement of one or more of amyloid production, amyloid assembly, amyloid aggregation, amyloid oligomer binding, and amyloid deposition; and/or
- (v) inhibiting, treating, and/or abatement of the activity/effect of one or more of Abeta oligomers on a neuron cell.

56. The method of claim 53 wherein the method of inhibiting, treating, and/or abatement of cognitive decline and/or Alzheimer's disease comprises inhibiting, treating, and/or abatement of one or more of amyloid production, amyloid assembly, the activity/effect of one or more of Abeta oligomers on a neuron cell, amyloid aggregation, amyloid binding, and amyloid deposition.

57. The method of claim 53 wherein the method of inhibiting, treating, and/or abatement of cognitive decline and/or Alzheimer's disease comprises inhibiting, treating, and/or abatement of one or more of the activity/effect of one or more of Abeta oligomers on a neuron cell, amyloid aggregation, amyloid binding, and amyloid deposition.

58. A method of inhibiting, treating, and/or abatement of cognitive decline and/or Alzheimer's disease in a patient comprising administering to the patient a compound of any one of claims 45-52 or a pharmaceutically acceptable salt thereof.

59. The method of claim 58 wherein the method of inhibiting, treating, and/or abatement of cognitive decline and/or Alzheimer's disease comprises inhibiting, treating, or abatement of one or more symptoms of cognitive decline selected from the group consisting of memory loss, confusion, impaired judgment, personality changes, disorientation, and loss of language skills.

60. The method of claim 58 wherein the method of inhibiting, treating, and/or abatement of cognitive decline and/or Alzheimer's disease comprises one or more of:

- (i) restoration of long term potentiation; and/or
- (ii) inhibiting, treating, and/or abatement of neurodegeneration; and/or
- (iii) inhibiting, treating, and/or abatement of general amyloidosis; and/or
- (iv) inhibiting, treating, and/or abatement of one or more of amyloid production, amyloid assembly, amyloid aggregation, amyloid oligomer binding, and amyloid deposition; and/or
- (v) inhibiting, treating, and/or abatement of the activity/effect of one or more of Abeta oligomers on a neuron cell.

61. The method of claim 58 wherein the method of inhibiting, treating, and/or abatement of cognitive decline and/or Alzheimer's disease comprises inhibiting, treating, and/or abatement of one or more of amyloid production, amyloid assembly, the activity/effect of one

or more of Abeta oligomers on a neuron cell, amyloid aggregation, amyloid binding, and amyloid deposition.

62. The method of claim 58 wherein the method of inhibiting, treating, and/or abatement of cognitive decline and/or Alzheimer's disease comprises inhibiting, treating, and/or abatement of one or more of the activity/effect of one or more of Abeta oligomers on a neuron cell, amyloid aggregation, amyloid binding, and amyloid deposition.