

NUCLEOTIDE AND AMINO ACID SEQUENCES OF OOCYTE FACTORS FOR ALTERING OVARIAN FOLLICULAR GROWTH *IN VIVO* OR *IN VITRO*

The present invention relates to nucleotide and amino acid sequences of oocyte factors for altering ovarian follicular growth *in vivo* or *in vitro*. The present invention also concerns novel homodimeric and heterodimeric polypeptides and their use for altering mammalian ovarian follicular growth *in vivo* or *in vitro*. In particular, the invention broadly concerns active or passive immunisation against these homo- or heterodimeric polypeptides or functional fragments or variants thereof so as to alter follicular growth *in vivo* or *in vitro*.

BACKGROUND OF THE INVENTION

The molecular nature of regulatory molecules responsible for stimulating early phases of ovarian folliculogenesis (i.e. the growth and differentiation of primordial follicles to primary, secondary, and pre-antral follicles) has been poorly understood. On the other hand, follicle-stimulating hormone (FSH) and luteinising hormone are glycoprotein hormones derived from the pituitary and have generally been accepted as the key factors regulating the later stages of ovarian folliculogenesis. Moreover FSH is accepted as the single most important factor for stimulating a greater than normal number of follicles to ovulate, a fact that is well illustrated by the wide use of commercial FSH preparations in ovarian hyperstimulation regimes both in medicine and veterinary medicine. Recent studies have indicated that early folliculogenesis is controlled by intraovarian factors of which the granulosa cell-derived stem cell factor (or c-kit ligand) and the oocyte-derived growth differentiation factor-9 (GDF-9) have gained most attention, because both appear to be essential for early mammalian folliculogenesis.

GDF-9 was first described in 1993 as a novel member of the transforming growth factor beta (TGF- β) superfamily which is specifically expressed in the ovary (McPherron and Lee, 1993). Like other members of the TGF- β family, GDF-9 is encoded as a prepropeptide consisting of a signal peptide, a proregion, and a so called C-terminal mature region, which is cleaved from the precursor peptide by an intracellular protease belonging to a group of furin-like proteases. Growth factors of the TGF- β family are characterised by a common pattern of cysteine residues found in the mature region that is likely to form in all members of the family; this is a rigid intramolecular structure known as the

"cysteine knot" which is composed of six Cys residues forming three characteristic disulphide bridges in a monomer of a TGF- β family member (Daopin et al., 1992; Schlunegger and Grutter, 1992 and 1993; Griffith et al., 1996; Scheufler et al., 1999). Most members of the TGF- β family have a conserved seventh Cys residue that is responsible for the covalent homodimerisation of two identical monomers (homodimers) or for the heterodimerisation of a given TGF- β family member with another distinct member of the family (heterodimers).

In mice, GDF-9 is expressed in oocytes from the primary stage of follicular development until ovulation (McGrath et al., 1995; Laitinen et al., 1998). Using the mouse GDF-9 sequence as a test sequence for the data base searches we identified, a GDF-9 like expressed sequence tag (EST) cDNA derived from a 2-cell mouse embryo library (Laitinen et al., 1998). We showed that the transcript of this novel factor, GDF-9B, which is 55% homologous to GDF-9, is expressed in oocytes of the mouse ovary at the same time as GDF-9 (Laitinen et al., 1998). Using PCR and primers derived from the mouse EST sequence we amplified a fragment of the corresponding gene from human genomic DNA, mapped the gene locus to chromosome Xp11.2, and deduced the human GDF-9B gene structure from isolated cosmid clones (Aaltonen et al., 1999). Interestingly, in the human ovary, GDF-9 mRNA expression begins in primary follicles slightly earlier than that of GDF-9B (Aaltonen et al., 1999). The mouse and human GDF-9B genes have been cloned and the protein encoded by the gene has also been named bone morphogenetic protein 15 (BMP-15) (Dube et al., 1998).

GDF-9 appears to be essential for ovarian folliculogenesis. From the literature it is known that GDF-9 deficient mice (GDF-9 $-/-$) are infertile due to an early arrest in folliculogenesis (Dong et al., 1996). In GDF-9 $-/-$ ovaries folliculogenesis stops at the primary follicle stage when one layer of cuboidal granulosa cells surrounds the oocyte. Even though the oocyte continues to grow, the granulosa cells fail to proliferate and no thecal cell differentiation is associated with follicular enlargement.

The Inverdale fecundity gene (FecX¹) was identified as a major gene affecting the prolificacy of a Romney flock of sheep (Davis et al., 1991). Segregation analyses determined that the gene is carried on the X-chromosome, and that ewes carrying a single copy of the gene (I/+) have litter size about 0.6 lambs larger than noncarrier ewes (+/+). The increase in number of lambs born is directly linked to an altered pattern of follicular development and an increase in ovulation-rate above that in the

wild-type of ~1.0 (Shackell et al., 1993; Davis et al., 1991). By contrast, homozygous carrier ewes having two copies of the gene (I/I) are infertile; due to an ovarian failure condition (Davis et al., 1992). In ovaries of the (I/I) ewe, folliculogenesis stops at the primary follicular stage and the phenotype is not dissimilar to that seen in GDF-9 (-/-) mice (Braw-Tal et al., 1993; McNatty et al., 1995; Smith et al., 1997).

A second prolific Romney flock (Hanna, 1995) with no known connection to the Inverdale flock was also shown to carry an X-linked mutation with similar phenotype to Inverdale. Evidence that the Hanna animals carried a mutation (FecX^H) in the same gene as for Inverdale was obtained when infertile females were produced by mating Inverdale carrier rams with carrier Hanna ewes (Davis et al., 1995). The Hanna line has been maintained at the Invermay AgResearch Centre as a distinct group alongside the Inverdale line.

In New Zealand Patent Application No. 500844 we, the present inventors, identified in Inverdale sheep a nucleotide substitution beyond the mature peptide processing site of the GDF-9B gene which converts the codon GTC (amino acid valine (V)) to GAC (amino acid aspartic acid (D)). We also showed that in Hanna sheep the C nucleotide beyond the mature peptide processing site is converted to a T. This converts a codon CAG (coding for glutamine (Q)) to a codon TAG (coding for termination) thereby resulting in a truncated mature protein. These respective mutations in Inverdale and Hanna are suggested to be the underlying causes for the "streak" ovaries and anovulatory conditions in homozygous Inverdale or Hanna ewes and in the Inverdale cross Hanna ewes.

Previously it has been shown that GDF-9 -/- mice are infertile showing that GDF-9 is important for normal fertility in some mammals. However, with the discovery of the related oocyte-specific factor GDF-9B several novel findings were found by us to support the notion that: (1) GDF-9B is essential for normal folliculogenesis in some mammals; (2) that GDF-9B is critically important for determining the ovulation rate in some mammals and; (3) since GDF-9 and GDF-9B are co-expressed by oocytes, they function co-operatively to enhance both follicular development and ovulation rate. Collectively these novel hypotheses were only made possible by our discovery of the Inverdale and Hanna GDF-9B mutations in sheep.

The inventors have for the first time determined the full gene structure of the sheep GDF-9B gene encoding the wild-type protein and have shown that it is necessary for maintaining normal ovarian folliculogenesis in sheep. The inventors have also identified the full gene structure of the GDF-9B variants in Inverdale and Hanna sheep which cause higher than normal ovulation rates in heterozygous animals and infertility in homozygous animals. It is broadly to the full-length wildtype and mutated GDF-9B sequences and variants thereof and their use in the modulation of mammalian fertility that the present invention is directed.

All references, including any patents or patent applications, cited in this specification are hereby incorporated by reference. No admission is made that any reference constitutes prior art. The discussion of the references states what their authors assert, and the applicants reserve the right to challenge the accuracy and pertinency of the cited documents. It will be clearly understood that, although a number of prior art publications are referred to herein, this reference does not constitute an admission that any of these documents forms part of the common general knowledge in the art, in New Zealand or in any other country.

SUMMARY OF THE INVENTION

Accordingly, to a first aspect, the present invention provides an isolated wildtype GDF-9B nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of:

- a) SEQ ID No:1;
- b) a sequence able to hybridise under stringent conditions to the molecule in (a);
- c) a sequence which is a functional variant or fragment of the molecule in (a);
- d) a sequence complementary to the molecule defined in (a), (b) or (c); and
- e) an anti-sense sequence corresponding to any of the molecules in (a) – (d).

In a second aspect, the present invention provides an isolated full length mutated GDF-9B nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of:

- a) SEQ ID NO: 3 or SEQ ID NO: 5 ;
- b) a sequence able to hybridise under stringent conditions to the molecule(s) in (a);
- c) a sequence which is a functional variant or fragment of the molecule(s) in (a);
- d) a sequence complementary to the molecule(s) defined in (a), (b) or (c); and
- e) an anti-sense sequence corresponding to any of the molecule(s) in (a) – (d).

The nucleic acid molecule may be an RNA, cRNA, genomic DNA or cDNA molecule, and may be single- or double-stranded. The nucleic acid molecule may also optionally comprise one or more synthetic, non-natural or altered nucleotide bases, or combinations thereof.

In a third aspect, the present invention provides an isolated full-length GDF-9B polypeptide comprising an amino acid sequence selected from the group consisting of:

- a) SEQ ID NO:2, SEQ ID NO:4; or SEQ ID NO:6; and
- b) A functional variant or fragment of the sequence(s) in (a).

In a fourth aspect, the present invention provides a homodimeric mature GDF-9B polypeptide having subunits comprising an amino acid sequence selected from the group consisting of SEQ ID NO:2 or a functional fragment or variant of said sequence.

In the fifth aspect, the present invention provides a heterodimeric polypeptide having subunits selected from the group consisting of:

- a) A mature GDF-9B polypeptide comprising an amino acid sequence derived from SEQ ID NO:2 or functional fragment or variant of said sequence; and
- b) A mature GDF-9 polypeptide or a functional variants or fragments thereof.

The present invention further provides in a sixth aspect a method of expressing biologically active processed homodimeric GDF-9B polypeptide comprising the steps of:

- a) generating an expression construct comprising a nucleic acid molecule comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO:1 or a functional fragment or variant of said sequence of the group;
- b) transfecting a suitable cells with said construct;
- c) selecting stable clones; and
- d) isolating and purifying the expressed polypeptide.

In a seventh aspect, the present invention provides a method of expressing biologically-active processed heterodimeric GDF-9B and GDF-9 polypeptides comprising the steps of:

- a) generating an expression construct containing a nucleic acid molecule comprising:
 - (i) a nucleic acid sequence selected from the group consisting of: SEQ ID NO:1 or a functional fragment or variant of said sequence; and
 - (ii) a nucleic acid molecule encoding GDF-9 or a functional fragment or variant thereof;
- b) transfecting suitable cells with said construct;
- c) selecting stable clones; and isolating and purifying the expressed polypeptide

Preferably the cells transfected are vertebrate, however the use of other cell types is envisaged.

The GDF-9 nucleic acid and protein sequences are available in public databases such as GENbank and SWISS-PROT. The accession number for the sheep GDF-9 nucleic acid is AF078545 and for protein is AAC28089.

Also provided by the present invention are recombinant expression vectors which contain a DNA molecule of the invention or functional variant thereof, and hosts transformed with a vector of the invention capable of expressing a polypeptide of the invention.

An additional aspect of the present invention provides a ligand which binds to a polypeptide of the invention. Most usually, the ligand is an antibody. It should be appreciated that the term "antibody" encompasses fragments or analogues of antibodies which retain the ability to bind to a polypeptide of the invention, including but not limited to Fv, F(ab)₂ fragments, ScFv molecules and the like. The antibody may be polyclonal or monoclonal, but is preferably monoclonal. In some embodiments the ligand may be a phage display molecule generated against polypeptides of the present invention, a single cell surface receptor or complex cell surface receptor. The polypeptide or peptide may be present as a monomer, dimer, heterodimer, multimer or a variant thereof.

In an eighth aspect, the invention provides a method for assessing the activity of GDF-9B homodimers and/or GDF-9B/GDF-9 heterodimers, comprising the steps of:

- a) adding an effective amount of a GDF-9B homodimeric polypeptide; and/or a GDF-9B/GDF-9 heterodimeric polypeptide to an ovarian cell or organ culture with or without other ovarian growth factors such as IGF-1 and/or other members of the transforming growth factor superfamily (e.g. activin, BMP2, TGF β 1); and

- b) conducting a bioassay on said cell or organ culture to assess the biological activity of said homodimeric and heterodimeric polypeptides.

In a ninth aspect, the invention provides transgenic animal models useful for demonstrating the effect of systemic production of GDF-9B homodimers and GDF-9B/GDF-9 heterodimers on follicular growth.

In a tenth aspect, the invention provides a method of adenoviral, retroviral and alphaviral transfer of GDF-9B expression cassettes or GDF-9 expression cassettes to host cells or organisms to thereby effect *in vivo* expression of GDF-9B homodimers or GDF-9B/GDF-9 heterodimers, comprising the step of transferring into a recipient cell, organ culture or recipient animal, a recombinant adenovirus including an expression cassette comprising a nucleic acid molecule having a nucleotide sequence selected from the group consisting of SEQ ID NO:1 or a functional fragment or variant of said sequence, said nucleic acid molecule being in operative association with an expression control sequence.

In an eleventh aspect the present invention provides the use of an agent selected from the group consisting of:

- a) a homodimeric polypeptide having subunits comprising GDF-9B or a functional fragment or variant thereof with or without homodimeric polypeptide having subunits comprising GDF-9 polypeptide or a functional fragment or variant thereof;
- b) a heterodimeric polypeptide having subunits comprising GDF-9B and GDF-9 polypeptides, or functional fragments or variants of said GDF-9B or GDF-9 polypeptides;

together with or without supplementary gonadotrophins (e.g. FSH and/or LH) and/or other ovarian growth factors such as IGF-1, kit ligand (stem cell factor), epidermal growth factor or a member of the TGF β superfamily (i.e. an agonist or antagonist) to:

- 1) alter follicular growth in ovaries of a mammal or other vertebrate either *in vivo* or *in vitro*;
or
- 2) alter isolated ovarian cell growth/maturation *in vitro* (e.g. oocyte – cumulus cells and/or granulosa cells)

In yet a twelfth aspect, the invention provides a composition comprising an effective amount of an agent selected from the group consisting of:

- a) a homodimeric polypeptide having subunits comprising a GDF-9B polypeptide or a functional fragment or variant thereof with or without homodimeric polypeptide having subunits comprising GDF-9 polypeptide or a functional fragment or variant thereof;
- b) a heterodimeric polypeptide having subunits comprising a GDF-9B polypeptide and a GDF-9 polypeptide, or functional fragments or variants of said GDF-9B or GDF-9 polypeptides;

together with a pharmaceutically or veterinarily acceptable carrier (including adjuvants) or diluent; and optionally including supplementary gonadotrophins and/or other relevant ovarian growth factor agonists/antagonists.

In a thirteenth aspect the invention provides a method of altering ovarian follicular growth in a female mammal or other female vertebrate, *in vivo* said method comprising the step of transforming mammalian and other vertebrate ovarian host cells with GDF-9B and GDF-9 expression cassettes to allow over-expression of GDF-9B homodimers and GDF-9B/GDF-9 heterodimers.

In a fourteenth aspect the invention provides a method of altering ovarian follicular growth in a female mammal or other female vertebrate, *in vitro* said method comprising the step of transforming mammalian and other vertebrate ovarian host cells with GDF-9B and GDF-9 expression cassettes to allow over-expression of GDF-9B homodimers and GDF-9B/GDF-9 heterodimers.

According to a further aspect the invention provides a nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of SEQ ID NO: 11, SEQ ID NO: 13 and SEQ ID NO: 15, or a functional fragment or variant of said sequence.

According to a further aspect the invention provides a polypeptide comprising an amino acid sequence said from the group consisting of SEQ ID NO: 12 and SEQ ID NO: 16, or a functional fragment or variant of said sequence.

According to another aspect the invention provides a method of altering follicular growth comprising the step of introducing a ligand as claimed in any one of claims 9-15 to:

- i) alter follicular growth in ovaries of a mammal or other vertebrate either *in vivo* or *in vitro*; or
- ii) alter isolated ovarian cell growth/maturation *in vitro*

Preferably said mammal is selected from the group comprising sheep, cattle, goats, deer, pigs, humans, horses, camelids and possums, cats and dogs and any other commercially important species having a GDF-9B gene having a substantial identity to the GDF-9B sequences of the present invention. Said vertebrate is preferably selected from the group comprising chickens, ducks, geese, salmon and any other commercially important species having a substantial identity to the GDF-9B sequences of the present invention.

Brief Description of Drawings

In particular, preferred aspects of the invention will be described in relation to the accompanying drawings, in which:

- Figure 1. Shows the nucleotide sequence of wildtype GDF-9B in sheep.
- Figure 2. Shows a portion of the nucleotide sequence of Figure 1 and illustrates the Inverdale mutation.
- Figure 3. Shows a portion of the nucleotide sequence of Figure 1 and illustrates the Hanna mutation.
- Figure 4. Shows the deduced amino acid sequence of wildtype GDF-9B in sheep.
- Figure 5. Shows a portion of the amino acid sequence of Figure 1 and illustrates the Inverdale mutation.
- Figure 6. Shows a portion of the amino acid sequence of Figure 1 and illustrates the Hanna mutation.
- Figure 7. Show the signal sequence polymorphism for GDF-9B with sheep.

Figure 8. Shows the additional upstream ATG codon present in sheep.

Figure 9. Shows photomicrograph illustrating the localisation of GDF-9B in an oocyte.

DETAILED DESCRIPTION OF THE INVENTION

As discussed above, the primary focus of the invention is the modulation of ovarian follicular growth via GDF-9B homodimers and GDF-9B/GDF-9 heterodimers activity *in vivo* and *in vitro*.

The term "isolated" means substantially separated or purified from contaminating sequences in the cell or organism in which the nucleic acid naturally occurs and includes nucleic acids purified by standard purification techniques as well as nucleic acids prepared by recombinant technology, including PCR technology, and those chemically synthesised.

Preferably, the nucleic acid molecule of SEQ ID NO: 1 is isolated from sheep genomic DNA, and that of SEQ ID NO: 3 and SEQ ID NO: 5 is isolated from DNA of sheep expressing the Inverdale or Hanna phenotype.

It has been noted that a polymorphism may occur in the signal peptide of GDF-9B in sheep (SEQ ID NO: 7, SEQ ID NO: 8). The predicted signal sequence appears to be around 25 amino acids long as predicted using the Signal P programme (Signal P V1.1 server at <http://genome.cbs.dtu.dk/services/SignalP>) (Nielsen et al., 1997) from ATG (Met) to ACA (Thr) in SEQ ID NO: 7 and SEQ ID NO: 8. A 3 base pair deletion is observed in some sheep where one of the two CTT sequences is not present. Therefore some sheep have a shorter signal sequence although most have the full length. From studies of Hanna, Hanna cross Inverdale, Inverdale and wild-type sheep either related or unrelated to Inverdale or Hanna, most were homozygous for two CTT's, although some were heterozygous for one CTT. The prevalence of the shorter version of the signal peptide was found to be high in the Merino breed of sheep but low in Romney breeds indicating that the polymorphism may be related to breed. Most Romney sheep carried the longer signal sequence regardless of their carrier status for Inverdale or Hanna. Although this polymorphism needs to be acknowledged it does not effect or modify the claims made in this invention.

It has also been noted that an additional 'in-frame' ATG start codon is present in the sheep genomic DNA sequence 27 nucleotides upstream from the GDF-9B ATG start codon (SEQ ID NO: 9, SEQ ID NO: 10). This sequence is present in all sheep sequenced, whether wildtype, Inverdale or Hanna carriers, and is clearly independent of the Inverdale and Hanna mutations. It is not known whether this upstream start codon is used during protein translation in sheep (SEQ ID NO: 9, SEQ ID NO: 10). If so it would result in a signal sequence of an additional 9 amino acids. Such a difference between sheep and other mammalian GDF-9B proteins would be unlikely to affect the function of the mature protein as this portion of the molecule is cleaved off in the active mature GDF-9B, but we mention the possibility of an alternative protein translation start site which may be present in sheep. The signal peptide prediction programme (Nielsen et al., 1997) indicates that this additional 9 amino acid sequence may function as a signal peptide and that the likely end site for the signal peptide would still be the Thr (T) amino acid indicated in SEQ ID NO: 7 and SEQ ID NO: 8.

In a further aspect, the present invention provides an isolated polypeptide selected from the amino acid sequences of SEQ ID NO: 2, SEQ ID NO: 4 and SEQ ID NO: 6 or a functional variant thereof which functions to manipulate ovarian follicular growth in a female mammal.

The polypeptide may be produced by expression of a suitable vector comprising the nucleic acid molecule of the invention or a functional variant thereof, in a suitable host cell as would be understood by a person skilled in the art.

The term "variant" as used herein refers to nucleotide and polypeptide sequences wherein the nucleotide or amino acid sequence exhibits substantially 50% or greater homology with the nucleotide or amino acid sequence of SEQ ID NOS: 1-6 respectively, preferably 75% homology and most preferably 90-95% homology to the sequences of the present invention: provided said variant has a biological activity as defined herein. The variant may be arrived at by modification of the native nucleotide or amino acid sequence by such modifications as insertion, substitution or deletion of one or more nucleotides or amino acids or it may be a naturally occurring variant. The term "variant" also includes homologous sequences which hybridise to the sequences of the invention under standard or preferably stringent hybridisation conditions familiar to those skilled in the art. Examples of the in situ hybridisation procedure typically used are described in (Tisdall et al., 1999); (Juengel et al., 2000). Where such a variant is desired, the nucleotide sequence of the native DNA is altered appropriately. This alteration can be made through elective synthesis of the DNA or by

modification of the native DNA by, for example, site-specific or cassette mutagenesis. Preferably, where portions of cDNA or genomic DNA require sequence modifications, site-specific primer directed mutagenesis is employed, using techniques standard in the art.

A "fragment" of a nucleic acid is a portion of the nucleic acid that is less than full length, and comprises at least a minimum sequence capable of hybridising specifically with a nucleic acid molecule according to the invention, or a sequence complementary thereto, under stringent conditions as defined below. A "fragment" of a polypeptide is a portion of the polypeptide which is less than full length, but which still retains the biological activity as defined herein.

The term "biologically active" refers to a polypeptide of the invention which is able to elicit a measurable physiological effect in the ovary or ovarian cell of a mammal or other vertebrate. The physiological effects may be measured by assays such as the incorporation of tritiated thymidine into granulosa cells. An example of such an assay is one in which follicles (1-2.5 mm diameter) are dissected free from ovarian stroma and granulosa cells isolated from theca and oocyte cumulus complexes. The cells are washed and resuspended in fresh media at a final concentration for bioassay of 100,000 viable cells per well and incubated with or without polypeptide for 48 hours. At this time incorporation of tritiated thymidine is measured.

The term "protein (or polypeptide)" refers to a protein encoded by the nucleic acid molecule of the invention, including fragments, mutations and homologs having the same biological activity i.e. ovulation manipulation activity. The polypeptide of the invention can be isolated from a natural source, produced by the expression of a recombinant nucleic acid molecule, or can be chemically synthesised.

The term "ligand" refers to any molecule which can bind to another molecule such as a polypeptides or peptide, and should be taken to include, but not be limited to, antibodies, cell surface receptors or phage display molecules.

In addition, nucleotides and peptides having substantial identity to the nucleotide and amino acid sequences of the invention can also be employed in preferred embodiments. Here "substantial identity" means that two sequences, when optimally aligned such as by the programs GAP or BESTFIT (peptides) using default gap weights, or as measured by computer algorithms BLASTX or

BLASTP, share at least 50%, preferably 75%, and most preferably 90-95% sequence identity. Preferably, residue positions which are not identical differ by conservative amino acid substitutions. For example, the substitution of amino acids having similar chemical properties such as charge or polarity are not likely to effect the properties of a protein. Non-limiting examples include glutamine for asparagine or glutamic acid for aspartic acid.

In a further aspect, the present invention provides in replicable transfer vectors suitable for use in preparing a polypeptide or peptide of the invention. These vectors may be constructed according to techniques well known in the art, or may be selected from cloning vectors available in the art.

The cloning vector may be selected according to the host or host cell to be used. Useful vectors will generally have the following characteristics:

- (a) the ability to self-replicate;
- (b) the possession of a single target for any particular restriction endonuclease; and
- (c) desirably, carry genes for a readily selectable marker such as antibiotic resistance.

Two major types of vector possessing these characteristics are plasmids and bacterial viruses (bacteriophages or phages). Presently preferred vectors are bacterial, insect or mammalian vectors and may include the following: the pUC, pBlueScript, pGEM, PGEX, pBK-CMV, lambda ZAP, lambda GEM, pEFIREP, pUB6/V5/His, pBC1, pADTrack-CMV, pAdenovator, pAdEasy-1, pSFV-PD, pCA3, pBABE, pPIC9, pA0815, pET and pSP series. However, this list should not be seen as limiting the scope of the present invention.

Examples of preferred expression systems are as follows:

1. For an *in vitro* cell expression system, the 293T cell system with a pEFIREP vector (Hobbs S *et al.*, 1998) which confers puromycin resistance may be used. For coexpression of two genes, the aforementioned vector may be modified to change the antibiotic resistance gene to bleomycin resistance. Alternatively, the co-expression of two genes and the selection gene can be achieved by constructing a tricistronic expression vector. A corresponding stably transfected insect cell system can also be used, e.g. the S2 cell system using "DES" vector expression system; www.invitrogen.com.

2. With respect to expressing GDF's in all tissues of transgenic animals, one approach is to use the pUB6/V5-His A vector (www.invitrogen.com) to make the constructs. For tissue-specific expression the rat PEPCK 0.6 kb promoter for liver and kidney expression can be included in the construct by replacing the Ubi-C promoter in the pUB6/V5-His A vector with the PEPCK promoter. For GDF expression in mammary tissue another promoter system would be preferred. For this tissue one approach would be to use the bovine β -lactoglobulin gene promoter and/or the bovine α S1 casein promoter (e.g. pBC1 vector, www.invitrogen.com) to drive the expression of the GDFs into milk. For global over-expression in transgenic animals, the CMV enhanced β -actin promoter (Okabe M, et al.; FEBS Letters 407: 313-319, 1997) or a modified EF1 α -promoter can be used also (Taboit-Dameron F, et al., Transgenic Research 8: 223-235, 1998).

Adenoviruses, retroviruses and alphaviruses are other suitable mammalian expression systems. A typical approach to those skilled in the art is that described by (TC He *et al.*, 1998). With respect to GDF expression the pAd Track-CMV vector or pAdenovator vectors (www.qbiogene.com) can be used to make the construct which is then co-transformed with pAd Easy-1 adenoviral plasmid into *E. coli* to generate a recombinant adenoviral genome which contains a CMV-promoter driven GDF expression cassette. This recombinant adenoviral genome is then transfected into 293T cells to make the virus stock. Alternative methods for generating adenoviruses can also be used for the same purpose (e.g. PCA3 plasmid based gene transfer (www.microbix.com); or COS-TPC method (Miyake S *et al.*, 1996).

3. Non-cytopathogenic Semliki Forest viruses expressing GDF's can be generated using, for example, pSFV-PD vectors as described by Lundstrom et al., Histochem Cell Biol 115: 83-91, 2001. Furthermore, retroviral expression systems based on, for example, pBABE vectors, can be used for expressing GDF's in mammalian cells (Morgenstern, JP and Land, H, 1990; Nucleic Acids Res 18: 3587-3596).
4. Yeast cells (e.g. *Pichia pastoris*, *Saccharomyces cerevisiae*) are another well established expression system to those skilled in the art (C Hadfield, *et al.*, 1993); (MA Romanos *et al.*, 1992). For example, the pPIC9 vector (www.invitrogen.com) can be used in *Pichia pastoris* for the expression of GDF's. For coexpression of two genes, the vector pA0815 (www.invitrogen.com) is a preferred candidate.

5. *Escherichia coli* (*E. coli*) is a standard laboratory expression system in widespread use. For example, the pET expression system (www.novagen.com) can be used to express recombinant mammalian GDF-9 and GDF-9B (steve.lawrence@agresearch.co.nz).

The DNA molecules of the invention may be expressed by placing them in operable linkage with suitable control sequences in a replicable expression vector. Control sequences may include origins of replication, a promoter, enhancer and transcriptional terminator sequences amongst others. The selection of the control sequence to be included in the expression vector is dependent on the type of host or host cell intended to be used for expressing the DNA as would be understood by a person skilled in the art.

The expression vectors useful in the present invention may contain at least one expression control sequence that is operatively linked to the DNA sequence or fragment to be expressed. The control sequence is inserted in the vector in order to control and to regulate the expression of the cloned DNA sequence. Examples of useful expression control sequences are the lac system, the trp system, the tac system, the trc system, major operator and promoter regions of phage lambda, the glycolytic promoters of yeast acid phosphatase, e.g. Pho5, the promoters of the yeast alpha-mating factors, and promoters derived from polyoma, adenovirus, retrovirus, and simian virus, and cytomegalovirus e.g. the early and late promoters of SV40, and other sequences known to control the expression of genes of prokaryotic and eukaryotic cells and their viruses or combinations thereof.

In the construction of a vector it is also an advantage to be able to identify the bacterial clone carrying the vector incorporating the foreign DNA. Such assays include measurable colour changes, antibiotic resistance and the like. In one preferred vector, the β -galactosidase gene is used, which gene is detectable by clones exhibiting a blue phenotype on X-gal plates. This facilitates selection. Once selected, the vectors may be isolated from the culture using standard procedures.

Depending on the host used, transformation and transfection is performed according to standard techniques appropriate to such cells. For prokaryotes or other cells that contain substantial cell walls, the calcium treatment process (Cohen, S N Proceedings, National Academy of Science, USA 69 2110 (1972)) may be employed. For mammalian cells without such cell walls, the calcium phosphate precipitation method of Graeme and Van Der Eb, Virology 52:546 (1978) or liposomal reagents are preferred.

Upon transformation of the selected host with an appropriate vector the polypeptide encoded can be produced, often in the form of a fusion protein, by culturing the host cells. The polypeptide of the invention may be detected by rapid assays as indicated above. The polypeptide is then recovered and purified as necessary. Recovery and purification can be achieved using any procedures known in the art, for example by absorption onto and elution from an anion exchange resin. This method of producing a polypeptide of the invention constitutes a further aspect of the present invention.

Host cells (including whole animal hosts), transformed, transfected or infected with the vectors of the invention also form a further aspect of the present invention.

In addition, a further aspect of the present invention provides for an antibody, antibody fragment, single cell surface receptor or complex cell surface receptor or phage display molecules binding to a polypeptide or peptide of the invention. The polypeptide or peptide may be present as a monomer, dimer, heterodimer, multimer or a variant thereof.

More specifically, the invention provides methods for producing antibodies against the wild-type (SEQ ID NO: 2) or mutated (SEQ ID NO: 4 and SEQ ID NO: 6) GDF-9B polypeptide sequences as monomers or homodimers or as heterodimers in combination with GDF-9. The antibodies can be used for the characterisation of the wild-type endogenous proteins, or peptide fragments and expressed recombinant proteins, or peptide fragments and for passive immunisation of recipient mammals for the modulation of ovarian follicular growth *in vivo*.

It will be appreciated by the reader that a further aspect of the invention contemplates the use of the polypeptides of the invention in the preparation of antisera for the detection of other GDF-9B -like peptides.

Polyclonal antibodies may be produced according to the method used by (Koelle *et al.*, 1991) incorporated herein by reference. Useful antibody production protocols are outlined in US Patent 5,514,578. Monoclonal antibodies may be produced by methods known in the art. These methods include the immunological method described by (Kohler and Milstein, 1975) as well as the recombinant DNA method described by (Huse *et al.*, 1989).

In a further embodiment, the invention provides a method of manipulating ovarian follicular growth in cells in culture and/or transgenic animals. By overexpression of GDF-9B homodimers and GDF-9B/GDF-9 heterodimers, the timing and level of expression of specific ovarian follicular protein genes may be altered in cultured cells or transgenic animals, for example GDF-9B homodimers or GDF-9B/GDF-9 heterodimers encoding sequences may be inserted into a gene cassette under the control of a specific promoter or a promoter that expresses in all cell types (see examples below) (constitutive expression). This cassette also comprises 3' flanking DNA that could stabilise the mRNA and may optionally comprise downstream regulatory sequences. This DNA cassette could be introduced into the genome of mammals by micro injection of the DNA into the pronuclei of eggs (as described in L'Huillier *et al.*, 1996) which are subsequently transferred back to recipient animals and allowed to develop to term. This technique for the production of transgenic animals is described by (Hogan *et al.*, 1996). Transgenic animals may be produced by transfection of cells in culture derived from an embryo, or foetal or adult tissues; followed by nuclear transfer and embryo transfer to recipient animals. Alternatively the gene cassette may be bound to mammalian sperm and delivered to the egg via *in vitro* or *in vivo* fertilisation to produce a non-human transgenic animal. Manipulation of the developmental regulation or the level of expression of GDF-9B homodimers or GDF-9B/GDF-9 heterodimers may be used to alter the level of ovarian follicular protein synthesis or production.

The invention also includes adenovirus-based gene therapy techniques for expressing GDF-9B and GDF-9/GDF-9B in cell cultures, organ cultures and whole experimental animals for manipulating ovarian follicular protein synthesis or production.

Non-limiting examples illustrating the invention will now be provided. It will be appreciated that the above description is provided by way of example only and variations in both the materials and techniques used which are known to those persons skilled in the art are contemplated.

EXAMPLES

EXAMPLE 1

Isolation of wildtype ovine GDF-9B DNA and identification of mutated ovine GDF-9B DNA sequences

Different combinations of oligonucleotide primers derived from human/mouse/rat GDF-9B sequences previously cloned by us were used in PCR on genomic sheep DNA for obtaining fragments of the ovine GDF-9B gene for sequencing. Functional primer pairs were used for obtaining wild-type sheep genomic clones from arrayed libraries and for obtaining cDNA sequence from wild-type ovine ovarian cDNA. The sequence from the full coding regions of Inverdale and Hanna was obtained by sequencing relevant PCR fragments obtained from the respective genomic DNA's. The wildtype GDF-9B sequence is disclosed here as SEQ ID NO: 1, Inverdale as SEQ ID NO: 3, and Hanna as SEQ ID NO: 5.

EXAMPLE 2

Production of specific antibodies and demonstration of natural homodimeric GDF-9B and heterodimeric GDF-9B/GDF-9 proteins in mammalian ovaries.

Antigens for immunising mammals or birds are generated using nucleotide sequences disclosed in the invention as SEQ ID NO: 1, SEQ ID NO: 3, and SEQ ID NO: 5 or variants thereof in operative association with an expression control sequence enabling expression of the protein in *E. coli*. Yet as another approach, anti-peptide antibodies directed against specific peptide sequences from SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6 or variants thereof are generated. Immunoreactivity can be assessed by standard methods (e.g. ELISA) and/or to obtain specific IgGs recognising GDF-9B homodimers and GDF-9B/GDF-9 protein heterodimers from natural sources and cell/tissues overexpressing the respective recombinant proteins.

Previous evidence by the inventors' group (e.g. Aaltonen *et al.* (1999) and Jaatinen *et al.*, 1999) show the presence of GDF-9 mRNA and protein in oocytes of preantral follicles in rodents and primates. Here we show evidence by immunocytochemistry using a mouse antibody for the presence of GDF-9B protein in sheep ovaries (Fig. 9). The immunohistochemical methodology was similar to that described by Tisdall *et al.* (1999) Stem cell factor and c-kit gene expression and protein localisation in the sheep ovary during fetal development, *J Reprod Fert* 116: 277-291. The only exception was that a tyramide signal amplification step was also included in the present study (TSA Biotin System, NEN Life Science Products). The mouse antibody M10 was generated after an *E. coli* derived mature ovine GDF-9B peptide (0.2 mg) was injected in Freund's complete adjuvant (FCA), intraperitoneally (i.p.) and at 2 weekly intervals boosted with 0.1 mg antigen i.p.

and subsequently with 0.05 mg antigen in a Span/Tween/oil mixture and the animal sacrificed 1 week after the final booster and serum collected.

Collectively these and other data show that both the mRNA and protein for GDF-9 and GDF-9B are present in oocytes of mammals (Aaltonen *et al.*, 1999; Galloway *et al.*, 2000).

Evidence that antipeptide antibodies directed against specific peptide sequences affect mammalian ovarian activity is shown in Table 1.

Table 1: Ovulation rate in sheep following administration of plasma containing antibodies to keyhole limpet haemocyanin (KLH) conjugated to GDF-9B peptide

Treatment	Mean Ovulation Rate (range)
Anti-KLH (n=5 animals)	1.6 (1-2)
Anti-GDF-9B peptide (5)	0.2* (0-1)

*p<0.05 (students t-test)

In this study 10 female sheep had their oestrous (i.e. ovarian) cycles synchronised using prostaglandin F_{2α} (Estrumate, 125 µg i.m.). All animals were observed to show synchronised oestrus following markings with vasectomised rams. On day 5 of the subsequent luteal phase, 5 ewes were administered with a pooled plasma recovered from another flock of ewes that had been immunised on 7 consecutive monthly occasions with a 15 mer GDF-9B peptide conjugated to keyhole limpet haemocyanin (KLH). The pooled sera contained high titre antibodies to GDF-9B as assessed by ELISA assay using a full-length *E. coli* expressed GDF-9B as antigen. The other 5 ewes were administered with a pooled plasma to KLH recovered from a separate flock of ewes that had been immunised also on 7 consecutive monthly occasions. The pooled plasma from these animals did not contain detectable GDF-9B antibody.

Antibody levels were measured by an ELISA procedure after the sheep plasmas were diluted 1:50000. The ELISA method involved coating a 96-well plate with 100 ng/well of an *E. coli*

expressed full-length GDF-9B and incubation with 100 μ l of diluted sheep plasma and 100 μ l of assay buffer, after appropriate blocking treatment and successive washes. After incubation with the sheep plasma and several washes, rabbit anti-sheep-HRP was added for 1 h at 37°C. The wells were then washed and developed with o-phenylenediamine plus hydrogen peroxide with development being stopped with sulphuric acid.

The ewes were each given 100 ml of sterile plasma intravenously and 4 days later administered with a second prostaglandin $F_{2\alpha}$ injection to synchronise oestrus. The ovulation rates were examined by laparoscopy 14 days after administration of plasma.

In our previous patent no. 500844 we demonstrated a significant perturbation in ovarian follicular development in mice when the animals were immunised with an *E. coli* derived mature ovine GDF-9B. In this study 10 female mice were immunised intraperitoneally (i.p.) with the *E. coli*-derived mature ovine GDF-9B protein (0.2 mg) in Freund's complete adjuvant (FCA) (0.2 ml), and another 10 female mice were immunised with bovine alpha lactalbumin (0.2 mg) in FCA (0.22 ml) i.p. to serve as controls. Subsequently, 3 booster injections of the appropriate antigens (0.1 mg at first booster and 0.05 mg at second and third booster) were given at 2 week intervals in a Span/Tween/oil mixture and the animals sacrificed 1 week after the final booster.

Here we provide evidence for these GDF-9B immunised mice that the exogenous biological activities of both GDF-9 and GDF-9B are likely to be affected since these animals contained cross-reacting antibodies to both growth factors (Table 2).

Table 2: Mean ($\pm$ s.e.m.) antibody levels in mouse plasma following immunisation with bovine alpha lactalbumin or ovine GDF-9B. The values presented show the absorbance at 490 nm which represents the levels of antibody to GDF-9B or GDF-9

Treatment	GDF-9B	GDF-9
Bovine α -lactalbumin	<0.060	<0.060
GDF-9B	2.177 ($\pm$ 0.163)	0.590 ($\pm$ 0.058)

Thus we assert that procedures that lead to the modulation of endogenous GDF-9 and GDF-9B will alter ovarian function.

Further evidence in support of our claim that modulating homodimeric GDF-9B or GDF-9 and GDF-9B either as homodimeric mixtures or heterodimers is provided by our novel findings following the immunisation of sheep (3-5 per treatment group) against KLH (control), GDF-9B peptide conjugated to KLH or GDF-9 peptide conjugated to KLH. The animals were subjected to 7 consecutive monthly immunisations, the ovaries recovered after slaughter and thereafter the ovarian volumes and numbers of types 1, 1a and 2 follicles were assessed by standard morphometric procedures (Smith *et al.*, 1997). In addition the presence or absence of antral follicles were noted. These results are summarised in Table 3.

Table 3: Mean ovarian volumes and numbers of Types 1-2, 3-4 and antral follicles following immunisation of sheep with KLH, KLH conjugated to GDF-9B peptide or to GDF-9 peptide

Treatment	Ovarian volume (mean range)	Number of Type 1-2 follicles (mean range)	Number of Type 3-4 follicles (mean range)	Number of antral follicles (mean range)
KLH (3)	486 (353-651)	15180 (11623-17433)	63 (61-65)	28 (22-37)
GDF-9B Peptide-KLH (5)	296 (149-488)	8147 (3221-13786)	0	0
GDF-9 Peptide-KLH (2)	284 (248-303)	16278 (10152-25745)	0	4 (0-11)

For these studies the GDF-9B peptide sequence was SEVPGPSREHDGPESC (SEQ ID NO 17) and the GDF-9 peptide sequence was KKPLVPASVNLSEYFC (SEQ ID NO 18). The Romney ewes were injected with 0.4 mg/ewe of KLH or KLH-GDF-9B peptide or KLH-GDF-9 peptide in Freund's complete adjuvant. Subsequently at monthly intervals on 6 occasions, the animals were boosted with further antigen (0.2 mg/ewe on each occasion) in a Span/Tween/oil mixture (s.c.).

The results show that relative to the controls (KLH immunisations): (1) GDF-9-KLH and GDF-9B-KLH immunisations each can inhibit antral follicle development thereby demonstrating that both GDF-9 and GDF-9B are essential for normal follicular development in some mammals.

EXAMPLE 3

Expression of recombinant homodimeric GDF-9B and heterodimeric GDF-9B/GDF-9 protein in mammalian cells *in vitro* and *in vivo*.

Expression constructs generated in plasmid vectors containing nucleotide sequences disclosed in the invention as SEQ ID NO: 1, SEQ ID NO: 3, and SEQ ID NO: 5 or variants thereof in operative association with an expression control sequence (CMV, EF1, and mammary specific promoter sequences) are transfected/transferred to mammalian cells and stable clones are selected using an antibiotic selection marker. For optimising the processing of the expressed recombinant polypeptide sequence the furin processing site is mutated and an auxiliary expression cassette driving the overexpression of the furin protease in the producer cells is introduced.

As an example of an effective mammalian expression system for producing oGDF-9B homodimeric protein we have generated human 293T cell lines transfected with the pEFIREs-P vector containing the rat GDF-9B proregion (Jaatinen et al., Mol Cell Endocrinol. 156:189-93, 1999) fused to an ovine GDF-9B mature region sequence (SEQ ID NO: 1 and NO:2). The furin processing site had been engineered to contain the effectively cleaved RRRR sequence. Cells selected to resist 120-150 ug/ml puromycin were cultured in HamF12/DMEM under serum free conditions for 4 days to produce ovine GDF-9B into the supernatant which was subsequently used in bioassay as described in Example 4.

To gain biochemical evidence for physical heterodimerisation of sheep GDF-9B with GDF-9 the following approach was used. A similar rat/sheep chimeric GDF-9B open reading frame as described above but containing a C-terminal 8 amino acid FLAG epitope was cloned into pSFV-PD Semliki Forest Virus vector and a high titre virus stock of SFV-PD-oGDF-9B-FLAG was generated in BHK cells as described in Lundstrom et al, Histochem Cell Biol 115: 83-91, 2001). A high titre SFV-PD-oGDF-9B-FLAG virus was seen to very effectively infect human 293T cells and cause high expression of oGDF-9B-FLAG in these cells. After infection, during a 4 day culture processed oGDF-9B-FLAG was secreted to medium and the protein was easily visualised as a 18 kd band in Western blot analysis using anti-FLAG M2 antibody. For co-expression of ovine GDF-9B-FLAG with ovine GDF-9 another 293T cell line was generated using a pEFIREs-P vector containing an open reading frame encoding a prepro GDF-9 polypeptide. Parental 293T cells and 293T cells stably

expressing oGDF-9 were infected with equal amounts of SFV-PD-oGDF-9B-FLAG viruses and supernatants were collected after 4 days in culture. One ml of supernatants from uninfected and SFV-PD-oGDF-9B-FLAG infected cells were subjected to immunoprecipitation with 1 µg/ml anti FLAG-M2 antibodies and complexes were recovered with protein G agarose. Eluates were evaluated in Western blots using anti-FLAG-M2 antibodies and anti GDF-9 antibodies. Although anti FLAG M2 antibodies do not react with ovine GDF-9, an immunoreactive 20 kd GDF-9 mature peptide was seen in supernatants of GDF-9 expressing cells that had been infected with SFV-PD-oGDF-9B-FLAG viruses and immunoprecipitated with anti FLAG M2 antibodies. These co-immunoprecipitation experiments indicate a direct physical interaction of recombinantly expressed ovine GDF-9B and GDF-9 and confirm the existence of sheep GDF-9/GDF-9B heterodimers.

Methods are described for over-expressing nucleotide sequences disclosed in the invention as SEQ ID NO: 1, SEQ ID NO: 3, and SEQ ID NO: 5 or variants thereof in several extraovarian sites in transgenic animals for mimicking systemic administration of recombinant homodimeric GDF-9B and heterodimeric GDF-9B/GDF-9 proteins. The relevant GDF encoding sequences are either expressed separately or co-expressed in operative association with expression control sequences. Effects of GDF-9B homodimers and GDF-9B/GDF-9 heterodimers on ovarian follicular growth in these transgenic animals are assessed by morphometric measurements or hormonal assays. This approach provides a general method for altering ovulation rate in transgenic animals. Production of GDF's into milk by mammary gland directed overexpression provides an alternative method of producing large amounts of reagent for medical or pharmaceutical purposes. This approach is not injurious to the health of the transgenic animal as no contraindications have been observed.

A method is described for effectively transferring an expression cassette composed of nucleotide sequences disclosed in the invention as SEQ ID NO: 1, SEQ ID NO: 3, and SEQ ID NO: 5 or variants thereof in operative association with an expression control sequence (CMV promoter) in a recombinant adenovirus into recipient cell or organ cultures or recipient animals for altering follicular growth *in vitro* or *in vivo*. For example, the methods allow systemic infection of recipient mammals with GDF-9B and GDF-9 expression cassette bearing adenoviruses concentrating into the livers of the host and allowing high levels of liver derived recombinant proteins to be released to the circulation. Effects on ovarian follicular growth may be assessed by one or more criteria such as laparoscopy, morphometric measurements or hormonal assays. A similar *in vivo* viral transfer of GDF-9B And GDF-9 sequences and transcriptional units can be achieved with alphaviruses or

retroviruses. These viral approaches offer ways of testing the bioactivity of different GDF-9B and GDF-9 gene constructs *in vivo* and also provide alternative ways of immunising animals against GDF-9B and GDF-9.

EXAMPLE 4

Measurement of the biological activity of homodimeric GDF-9B and heterodimeric GDF-9B/GDF-9 proteins in ovarian cell cultures.

For assessing the biological activity of proteins expressed from the nucleotide sequences disclosed in the invention as SEQ ID NO: 1, SEQ ID NO: 3, and SEQ ID NO: 5 or variants thereof ovarian cell and organ culture models may be used.

An example of assessing the biological activity of homodimeric GDF-9B is shown in Table 4 below whereby a partially purified recombinant (r) ovine (o)GDF-9B extract is assessed by measurement of [³H] thymidine incorporation by isolated ovine granulosa cells during a 48 h incubation at 37°C. The results show that the roGDF-9B caused a 1.9-fold increase in tritiated thymidine incorporation by granulosa cells indicating that the roGDF-9B was biologically active. To obtain the granulosa cells, ovaries were recovered from ewes, follicles (1-2.5 mm diameter) dissected free and the cells isolated and separated from theca and oocyte-cumulus complexes. The cells were washed and resuspended in fresh media at a final concentration for bioassay of 1x10⁵ viable cells per well.

Table 4: Mean $\pm$ s.e.m. increase in [³H] thymidine incorporation in ovine (o) granulosa cells after exposure to roGDF-9B or control media (n=3 separate experiments)

Treatment	[³ H] thymidine incorporation (cpm)
Control	3256 $\pm$ 283
Rec ovine (o)-GDF-9B	6291 $\pm$ 503***

RoGDF-9B was produced by transfected 293T cells and partially purified using heparin-sepharose chromatography and the protein eluted with 0.5 M NaCl and dialysed overnight against tissue culture media. The control for this experiment was media exposed to non-transfected 293T cells and subjected to heparin-sepharose chromatography, NaCl elution and dialysis.

***p<0.001, ANOVA

DISCUSSION

The known human GDF-9B sequence and oligonucleotide primers derived thereof enabled the inventors to determine the sheep GDF-9B genomic and cDNA sequences and to assess the expression of GDF-9B transcripts in sheep ovaries (Galloway et al., 2000).

From literature it appears that GDF-9 is needed for granulosa cell mitogenesis and thecal cell differentiation. Indeed, recombinant rat GDF-9 is able to stimulate rat follicular growth in vitro (Hayashi et al., 1999) as well as the proliferation of rat granulosa cells in culture (Vitt et al., 2000). Recombinant GDF-9 also regulates steroidogenesis and gonadotrophin receptor expression in mouse and rat granulosa cells (Elvin et al., 1999; Vitt et al., 2000). Furthermore, GDF-9 stimulates inhibin B production in human granulosa cells in culture (Vuojolainen et al., in preparation). These recent studies clearly demonstrate that GDF-9 homodimers have potent effects on follicular growth and differentiation in several mammals but before the invention described herein nothing has been known on the possible biological effect of GDF-9B in the ovary.

In our earlier New Zealand patent specification no. 500844 the present inventors showed that the Inverdale gene maps to a sheep X-chromosome region containing genes syntenic to Xp11.2-11.4 in human (Galloway et al., 2000) and determined whether GDF-9B gene is affected in these animals, showing that the Inverdale gene is actually an inactivated form of the sheep GDF-9B gene. In Inverdale animals the T nucleotide at position 92 nucleotides beyond the mature peptide processing site has become an A residue, converting the codon GTC to GAC causing the substitution of the amino acid valine (V) to aspartic acid (D). In all TGF- β family members this very amino acid is either valine, isoleucine or leucine which all represent hydrophobic residues in contrast to the negatively charged aspartic acid. This amino acid substitution causes a change in the surface charge of the very area of the molecule which is involved in the dimerisation process as suggested from the crystal structures of TGF- β 2, BMP-2 and BMP-7 (Schlunegger and Grutter, 1993; Griffith et al., 1996; Scheufler et al., 1999). The invention of New Zealand 500844 also provides evidence for a second GDF-9B gene mutation identified in another flock of sheep, Hanna, having exactly the same heterozygous and homozygous gene carrier phenotypes as Inverdale ewes. Hanna animals present with a C to T nucleotide point mutation at position 67 nucleotides beyond the mature peptide processing site introducing a premature stop codon at the place of a glutamine (Q) residue. This change will cause a major truncation of the mature peptide region causing inactivation of the protein

product. Crosses between the Inverdale and Hanna families lead to 50% infertile females proving that both mutations clearly inactivate the GDF-9B gene product.

REFERENCES

- Aaltonen J, Laitinen M, Vuojolainen K, Jaatinen R, Horelli-Kuitunen N, Seppa L, Louhio H, Tuuri T, Sjoberg J, Butzow R, Hovatta O, Dale L, Ritvos O: Human growth differentiation factor- 9 (GDF-9) and its novel homolog GDF-9B are expressed during early folliculogenesis. *J Clin Endocrinol Metab* 84: 2744-2750, 1999
- Braw-Tal R, McNatty KP, Smith P, Heath DA, Hudson NL, Phillips DJ, McLeod BJ, Davis GH: Ovaries of ewes homozygous for the X-linked Inverdale gene (FecXI) are devoid of secondary and tertiary follicles but contain many abnormal structures. *Biol Reprod* 49:895-907, 1993.
- C Hadfield, KK Raina, K Shashi-Menon, RC Mount (1993) The expression and performance of cloned genes. *Yeast Mycol Res* 9, 897-944
- Daopin S, Piez KA, Ogawa Y, Davies DR: Crystal structure of transforming growth factor-beta 2: an unusual fold for the superfamily. *Science* 257:369-373, 1992.
- Davis GH, McEwan JC, Fennessy PF, Dodds KG, Farquhar PA: Evidence for the presence of a major gene influencing ovulation rate on the X chromosome of sheep. *Biol Reprod* 44:620-624, 1991.
- Davis GH, McEwan JC, Fennessy PF, Dodds KG, McNatty KP, O WS: Infertility due to bilateral ovarian hypoplasia in sheep homozygous (FecXI FecXI) for the Inverdale prolificacy gene located on the X chromosome. *Biol Reprod* 46:636-640, 1992
- Davis GH, McEwan JC, Fennessy PF, Dodds KG: Discovery of the Inverdale gene (FecX). *Proc NZ Soc Anim Prod* 95:289-290, 1995.
- Dong J, Albertini DF, Nishimori K, Kumar TR, Lu N, Matzuk MM: Growth differentiation factor-9 is required during early ovarian folliculogenesis. *Nature* 383:531-535, 1996
- Dube JL, Wang P, Elvin J, Lyons KM, Celeste AJ, Matzuk MM: The bone morphogenetic protein 15 gene is X-linked and expressed in oocytes. *Mol Endocrinol* 12:1809-1817, 1998

- Elvin JA, Clark AT, Wang P, Wolfman NM, Matzuk MM: Paracrine actions of growth differentiation factor-9 in the mammalian ovary. *Mol Endocrinol* 13:1035-1048, 1999
- Galloway SM, McNatty KP, Cambridge LM, Laitinen MPE, Juengel JL, Jokiranta TS, McLaren RJ, Luiro K, Dodds KG, Montgomery GW, Beattie AE, Davis GH, Ritvos O: Mutations in an oocyte-derived growth factor gene (*BMP15*) cause increased ovulation rate and infertility in a dosage-sensitive manner. *Nature Genetics* 25:279-283, 2000.
- Griffith DL, Keck PC, Sampath TK, Rueger DC, Carlson WD: Three-dimensional structure of recombinant human osteogenic protein 1: structural paradigm for the transforming growth factor beta superfamily. *Proc Natl Acad Sci U S A* 93:878-883, 1996.
- Hanna MM: Living with the Inverdale gene (*FecX*) in a Romney flock. *Proc NZ Soc Anim Prod* 55:296-297, 1995.
- Hayashi M, McGee EA, Min G, Klein C, Rose UM, van Duin M, Hsueh AJ: Recombinant growth differentiation factor-9 (GDF-9) enhances growth and differentiation of cultured early ovarian follicles. *Endocrinology* 140:1236-1244, 1999
- Hobbs S, Jitrapakdee S, Wallace JC (1998) Development of a bicistronic vector driven by the human polypeptide chain elongation factor 1 alpha promoter for creation of stable mammalian cell lines that express very high levels of recombinant proteins. *Biochem Biophys Res Commun* 252, 368-372
- Hogan *et al* (1; In: *Manipulating the mouse embryo*, Cold Spring Harbor Laboratory. Press 1996)
- Huse *et al.*, *Science* 246: 1275-1281 (1989).
- Koelle *et al.*, *Cell* 67: 59-77, 1991.
- Kohler and Milstein in *Nature* 256: 495-497 (1975).

- Juengel et al. (2000) Gene expression in abnormal ovarian structures of ewes homozygous for the Inverdale prolificacy gene, *Biol Reprod*, 62, 1467-1478.
- Laitinen M, Vuojolainen K, Jaatinen R, Ketola I, Aaltonen J, Lehtonen E, Heikinheimo M, Ritvos O: A novel growth differentiation factor-9 (GDF-9) related factor is co-expressed with GDF-9 in mouse oocytes during folliculogenesis. *Mech Dev* 78:135-140, 1998
- L'Huillier et al PNAS 93; 6698-6703 (1996)
- MA Romanos, CA Scorer, JJ Clare (1992) Foreign gene expression in yeast: a review. *Yeast* 8, 423-488.
- McGrath SA, Esquela AF, Lee SJ : Oocyte-specific expression of growth differentiation factor-9. *Mol Endocrinol* 9:131-136, 1995.
- McNatty KP, Smith P, Hudson NL, Heath DA, Tisdall DJ, O WS, Braw-Tal R: Development of the sheep ovary during fetal and early neonatal life and the effect of fecundity genes. *J Reprod Fertil Suppl* 49:123-135, 1995.
- McPherron AC, Lee SJ: GDF-3 and GDF-9: two new members of the transforming growth factor-beta superfamily containing a novel pattern of cysteines. *J Biol Chem* 268:3444-3449, 1993
- Miyake S, Makimura M, Kanegae Y, Harada S, Sato Y, Takamori K, Tokuda C, Saito I. Efficient generation of recombinant adenoviruses using adenovirus DNA-terminal protein complex and a cosmid bearing the full-length virus genome. *Proc Natl Acad Sci U S A*. 93:1320-1324, 1996.
- Neilsen H., Engelbrecht J, Brunak S, von Heijne G: Identification of prokaryotic and eukaryotic signal peptides and prediction of their cleavage sites. *Protein Eng* 10:1-6
- Scheufler C, Sebald W, Hulsmeier M: Crystal structure of human bone morphogenetic protein-2 at 2.7 Å resolution. *J Mol Biol* 287:103-115, 1999

- Schlunegger MP, Grutter MG: An unusual feature revealed by the crystal structure at 2.2 Å resolution of human transforming growth factor-beta 2. *Nature* 358:430-434, 1992.
- Schlunegger MP, Grutter MG: Refined crystal structure of human transforming growth factor beta 2 at 1.95 Å resolution. *J Mol Biol* 231:445-458, 1993.
- Shackell GH, Hudson NL, Heath DA, Lun S, Shaw L, Condell L, Blay LR, McNatty KP Plasma gonadotrophin concentrations and ovarian characteristics in Inverdale ewes that are heterozygous for a major gene (*FecX^h*) on the X chromosome that influences ovulation rate. *Biol Reprod* 48:1150-1156, 1993.
- Smith P, O WS, Corrigan KA, Smith T, Lundy T, Davis GH, McNatty KP: Ovarian morphology and endocrine characteristics of female sheep fetuses that are heterozygous or homozygous for the Inverdale prolificacy gene (*fecX1*). *Biol Reprod* 57:1183-1192, 1997
- TC He, S Zhou, LT da Costa, J. Yu, KW Kinzler, B. Vogelstein (1998) A simplified system for generating recombinant adenoviruses. *Proc Natl Acad Sci USA* 95, 2509-2514.
- Tisdall et al. (1999) Stem cell factor and c-kit gene expression and protein localisation in the sheep ovary during fetal development, *J Reprod Fert* 116: 277-291.
- Vitt UA, Hayashi M, Klein C, Hsueh AJW: Growth differentiation factor-9 stimulates proliferation but suppresses the follicle-stimulating hormone-induced differentiation of culture granulosa cells from small antral and preovulatory rat follicles. *Biol Reprod*, 62, 370-377, 2000.
- Vuojolainen K, Bondestam J, Hayashi M, Raivio T, Evans L, Groome NP, Hsueh AJW, Ritvos O: GDF-9 regulates inhibin B production in cultured human granulosa-luteal cells. Manuscript in preparation.

WHAT WE CLAIM IS:

1. An isolated wildtype GDF-9B nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of:
 - a) SEQ ID No:1;
 - b) a sequence able to hybridise under stringent conditions to the molecule in (a);
 - c) a sequence which is a functional variant or fragment of the molecule in (a);
 - d) a sequence complementary to the molecule defined in (a), (b) or (c); and
 - e) an anti-sense sequence corresponding to any of the molecules in (a) – (d).
2. An isolated full length mutated GDF-9B nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of:
 - a) SEQ ID NO: 3 or SEQ ID NO: 5 ;
 - b) a sequence able to hybridise under stringent conditions to the molecule(s) in (a);
 - c) a sequence which is a functional variant or fragment of the molecule(s) in (a);
 - d) a sequence complementary to the molecule(s) defined in (a), (b) or (c); and
 - e) an anti-sense sequence corresponding to any of the molecule(s) in (a) – (d).
3. An isolated full-length GDF-9B polypeptide comprising an amino acid sequence selected from the group consisting of:
 - a) SEQ ID NO:2 , SEQ ID NO:4 ; or SEQ ID NO:6; and
 - b) A functional variant or fragment of the sequence(s) in (a).
4. A homodimeric mature GDF-9B polypeptide having subunits comprising an amino acid sequence selected from the group consisting of SEQ ID NO:2 or a functional fragment or variant of said sequence.
5. A heterodimeric polypeptide having subunits selected from the group consisting of:
 - a) A mature GDF-9B polypeptide comprising an amino acid sequence derived from SEQ ID NO:2 or a functional fragment or variant of said sequence; and
 - b) A mature GDF-9 polypeptide or a functional variant or fragment thereof.
6. A vector comprising the nucleic acid molecule of claim 1 or claim 2.

7. A construct comprising the nucleic acid molecule of claim 1 or claim 2.
8. A host cell which has been transformed by a vector or construct as claimed in claim 6 or claim 7.
9. A ligand which binds to a polypeptide derived from the full-length prepro polypeptide as claimed in claim 3.
10. A ligand which binds to a homodimeric polypeptide as claimed in claim 4.
11. A ligand which binds to a heterodimeric polypeptide as claimed in claim 5.
12. A ligand as claimed in claim 10 or 11, wherein the ligand is an antibody or an antibody fragment comprising the antigen-binding domain.
13. A ligand as claimed in claim 12 wherein the ligand is a monoclonal antibody.
14. A ligand as claimed in claim 10 or 11 wherein the ligand is a phage display molecule.
15. A ligand as claimed in claim 10 or 11 where the ligand is in the form cell surface receptor(s).
16. A method of expressing biologically active processed homodimeric GDF-9B polypeptide comprising the steps of:
 - a) generating an expression construct comprising a nucleic acid molecule comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO:1 or a functional fragment or variant of said sequence of the group;
 - b) transfecting suitable cells with said construct;
 - c) selecting stable clones; and
 - d) isolating and purifying the expressed polypeptide.
17. A method of expressing biologically-active processed heterodimeric GDF-9B and GDF-9 polypeptides comprising the steps of:

- a) generating an expression construct containing a nucleic acid molecule comprising:
 - (i) a nucleic acid sequence selected from the group consisting of: SEQ ID NO:1 or a functional fragment or variant of said sequence; and
 - (ii) a nucleic acid molecule encoding GDF-9 or a functional fragment or variant thereof;
 - b) transfecting suitable cells with said construct;
 - c) selecting stable clones; and
 - d) isolating and purifying the expressed polypeptide.
18. A method of adenoviral, retroviral and alphaviral transfer of GDF-9B expression cassettes or GDF-9 expression cassettes to host cells or organisms to thereby effect *in vivo* expression of GDF-9B homodimers or GDF-9B/GDF-9 heterodimers, comprising the step of transferring into a recipient cell, organ culture or recipient animal, a recombinant adenovirus including an expression cassette comprising a nucleic acid molecule having a nucleotide sequence selected from the group consisting of SEQ ID NO:1 or a functional fragment or variant of said sequence, said nucleic acid molecule being in operative association with an expression control sequence.
19. A transgenic animal which has been transformed by a vector or construct as claimed in claim 6 or 7.
20. A method of altering ovarian follicular growth, *in vivo*, in a female mammal or other female vertebrate, said method comprising the step of transforming mammalian and other vertebrate ovarian host cells with GDF-9B and GDF-9 expression cassettes to allow over-expression of GDF-9B homodimers and GDF-9B/GDF-9 heterodimers.
21. A method of altering ovarian follicular growth, *in vitro*, in a female mammal or other female vertebrate, said method comprising the step of transforming mammalian and other vertebrate ovarian host cells with GDF-9B and GDF-9 expression cassettes to allow over-expression of GDF-9B homodimers and GDF-9B/GDF-9 heterodimers.

22. A method as claimed in claim 19 or 20 wherein said mammal is selected from the group consisting of sheep, cattle, goats, deer, humans, pigs, horses, camelids, and possums, cats and dogs and any other commercially important species.
23. A method as claimed in claims 19 or 20 wherein said vertebrate is selected from the group consisting of chickens, ducks, geese, salmon and any other commercially important species.
24. A composition comprising an effective amount of an agent selected from the group consisting of:
- a) a homodimeric polypeptide having subunits comprising a GDF-9B polypeptide or a functional fragment or variant thereof with or without homodimeric polypeptide having subunits comprising GDF-9 polypeptide or a functional fragment or variant thereof;
 - b) a heterodimeric polypeptide having subunits comprising a GDF-9B polypeptide and a GDF-9 polypeptide, or functional fragments or variants of said GDF-9B or GDF-9 polypeptides;
- together with a pharmaceutically or veterinarily acceptable carrier (including adjuvants) or diluent; and optionally including supplementary gonadotrophins.
25. The use of an agent selected from the group consisting of:
- a) a homodimeric polypeptide having subunits comprising GDF-9B or a functional fragment or variant thereof with or without homodimeric polypeptide having subunits comprising GDF9 polypeptide or a functional fragment or variant thereof;
 - b) a heterodimeric polypeptide having subunits comprising GDF-9B and GDF-9 polypeptides, or functional fragments or variants of said GDF-9B or GDF-9 polypeptides;
- together with or without supplementary gonadotrophins and/or other ovarian growth factors including IGF-1, kit ligand (stem cell factor), epidermal growth factor or TGF β agonists/antagonists to:
- i) alter follicular growth in ovaries of a mammal or other vertebrate either *in vivo* or *in vitro*; or
 - ii) alter isolated ovarian cell growth/maturation *in vitro*.

26. A method for assessing the activity of GDF-9B homodimers and/or GDF-9B/GDF-9 heterodimers, comprising the steps of:
 - a) adding an effective amount of a GDF-9B homodimeric polypeptide; and/or a GDF-9B/GDF-9 heterodimeric polypeptide to an ovarian cell or organ culture with or without other ovarian growth factors (including TGF β agonists/antagonists); and
 - b) conducting a bioassay on said cell or organ culture to assess the biological activity of said homodimeric and heterodimeric polypeptides.
27. A method of altering follicular growth comprising the step of introducing a ligand as claimed in any one of claims 9-15 to:
 - ii) alter follicular growth in ovaries of a mammal or other vertebrate either *in vivo* or *in vitro*; or
 - iii) alter isolated ovarian cell growth/maturation *in vitro*
28. An isolated functional variant polypeptide comprising an amino acid sequence selected from this group consisting of SEQ ID NO: 17.
29. An isolated functional variant polypeptide comprising an amino acid sequence selected from this group consisting of SEQ ID NO: 18.
30. An isolated nucleic acid molecule comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO: 11, SEQ ID NO: 13, and SEQ ID NO: 15, or a functional fragment or variant of said sequences.
31. An isolated polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 12, SEQ ID NO: 14 and SEQ ID NO: 16, or a functional fragment or variant of said sequences.
32. An isolated nucleic acid molecule substantially as described herein with reference to any example and/or drawing thereof.
33. An isolated polypeptide substantially as described herein with reference to any example and/or drawing thereof.

34. A vector or gene construct incorporating an isolated nucleic acid molecule of the present invention substantially as described herein with reference to any example and/or drawing thereof.
35. A ligand which binds to a polypeptide of the present invention substantially as described herein with reference to any example and/or drawing.
36. A homodimeric polypeptide substantially as described herein with reference to any example and /or drawing thereof.
37. A heterodimeric polypeptide substantially as described herein with reference to any example and /or drawing thereof.
38. A method of expressing a biologically active processed homodimeric polypeptide substantially as described herein with reference to any example thereof.
39. A method of expressing a biologically active processed heterodimeric polypeptide substantially as described herein with reference to any example thereof.
40. A method of adenoviral transfer of polypeptides to thereby effect *in vivo* expression of homodimeric or heterodimeric polypeptides of the invention substantially as described herein with reference to any example thereof.
41. A method of altering ovarian follicular growth in a female mammal or other vertebrate substantially as described herein in reference to any example thereof.
42. A composition substantially as described herein with reference to any example and/or drawing thereof.
43. The use of a homodimeric or heterodimeric polypeptide to alter follicular growth substantially as described herein with reference to any example thereof.

44. A method for assessing the biological activity of homodimeric and heterodimeric polypeptides substantially as described herein with reference to any example thereof.

- 8 JAN 2004

28 OCT 2004

45. A use of GDF-9B homodimers and/or GDF-9B/GDF-9 heterodimers in the manufacture of a medicament for altering ovarian follicular growth in a female mammal in vivo.
46. A use of GDF-9B homodimers and/or GDF-9B/GDF-9 heterodimers in the manufacture of a medicament for altering mammalian ovarian follicular growth in vitro.
47. A use as claimed in any one of claims 45-47, wherein said medicament further comprises supplementary gonadotrophins and/or other ovarian growth factors including IGF-1, kit ligand (stem cell factor), epidermal growth factor or TGF β agonist/antagonist.
48. A use of a ligand as claimed in any one of claims 9-15 in the manufacture of a medicament to alter follicular growth in ovaries of a mammal either in vivo or in vitro or to alter isolated mammalian ovarian cell growth/maturation in vitro.
49. A use of an isolated GDF-9B nucleic acid molecule as claimed in any one of claims 1-3 in the manufacture of a medicament for altering mammalian ovarian follicular growth in vivo or in vitro.
50. A use as claimed in any one of claims 45 to 49 wherein said mammal is selected from the group consisting of sheep, cattle, goats, deer, humans, pigs, horses, camelids, possums, cats and dogs and any other commercially important species.