

VACCINE

FIELD OF THE INVENTION

The present invention relates to the field of vaccines for protecting against polio, and in particular to combination vaccines for protecting against polio, diphtheria, tetanus, and pertussis diseases.

BACKGROUND

Combination vaccines (which provide protection against multiple pathogens) are very desirable in order to minimise the number of immunisations required to confer protection against multiple pathogens, to lower administration costs, and to increase acceptance and coverage rates. The well-documented phenomenon of antigenic competition (or interference) complicates the development of multi-component vaccines. Antigenic interference refers to the observation that administering multiple antigens often results in a diminished response to certain antigens relative to the immune response observed when such antigens are administered individually.

Combination vaccines are known which can prevent *Bordetella pertussis*, *Clostridium tetani*, *Corynebacterium diphtheriae*, and optionally inactivated poliovirus (IPV), and/or Hepatitis B virus, and/or Haemophilus type B infection (see for instance WO 93/24148, WO97/00697 and WO2000/030678).

After many years of research the standard dose of polio vaccines accepted as effective within the vaccine community today contains 40 D antigen units of inactivated poliovirus type 1 (Mahoney), 8 D antigen units of inactivated poliovirus type 2 (MEF-1) and 32 D antigen units of inactivated poliovirus type 3 (Saukett) (e.g. Infanrix-IPV™).

The present inventors have surprisingly found that reduced doses of IPV can maintain an adequate or improved level of protection against polio. Such vaccines carry considerable advantages including the ability to provide more doses of IPV vaccines for the individuals in need thereof.

SUMMARY OF THE INVENTION

Accordingly, the present invention provides various reduced-dose IPV vaccines (which may only have IPV components or may have IPV components combined with other antigens).

5 Accordingly, in one aspect the present invention provides an IPV vaccine of the invention comprising inactivated poliovirus type 1 at a dose greater than 10 D-antigen units and less than 20 D-antigen units, e.g. 11, 12, 13, 14, 15, 16, 17, 18 or 19 D-antigen units.

10 In one embodiment, the present invention provides an IPV vaccine of the invention comprising inactivated poliovirus type 3 at a dose of 8-20 D-antigen units, e.g. 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20 D-antigen units.

In another embodiment, the present invention provides an IPV vaccine of the invention comprising inactivated poliovirus type 2 at a dose of 2-4 D-antigen units, e.g. 2, 3 or 4 D-antigen units.

15 In a further embodiment, the present invention provides an IPV vaccine of the invention further comprising diphtheria toxoid and/or tetanus toxoid and/or a pertussis vaccine in the form of killed whole-cell Pw vaccine or acellular pertussis antigens.

20 In a further aspect, the present invention provides an IPV vaccine of the invention which is a thiomersal free DTP-IPV combination vaccine comprising inactivated poliovirus type 1 at a dose between 10 and 36 D-antigen units.

In another embodiment, the present invention provides a thiomersal free DTP-IPV combination vaccine of the invention comprising inactivated poliovirus type 2 at a dose of 2-7 D-antigen units, e.g. 5, 6 or 7 D-antigen units.

25 In another embodiment, the present invention provides a thiomersal free DTP-IPV combination vaccine of the invention comprising inactivated poliovirus type 3 at a dose of 8-29 D-antigen units, e.g. 21, 22, 23, 24, 25, 26, 27, 28 or 29 D-antigen units.

30 In a further embodiment, the vaccines of the present invention may also comprise one or more antigens selected from the group consisting of: Hepatitis B surface antigen, *Haemophilus influenzae* b antigen(s), *Neisseria meningitidis* A antigen(s), *Neisseria meningitidis* C antigen(s), *Neisseria meningitidis* W antigen(s), *Neisseria meningitidis* Y antigen(s), *Neisseria meningitidis* B bleb or antigen(s), Hepatitis A antigen(s) and *Salmonella typhi* antigen(s), in particular capsular saccharide antigens from said bacteria.

Methods of making the vaccines of the invention are also provided.

DEFINITIONS

5 The term “vaccine” is optionally substitutable with the term “immunogenic composition” and vice versa.

 “D-antigen units” (also referred to as “international units” or IU): The D antigenic form of the poliovirus induces protective neutralising antibodies. D antigen units referred to herein (for instance in the vaccines of the invention) are the measured
10 total D antigen units of each unadsorbed bulk IPV antigen type prior to formulation of the final vaccine which are added in each human dose of formulated vaccine (typically 0.5mL final volume). Reliable methods of measuring D-antigen units are well known in the art and are published, for instance, by the European Pharmacopoeia. For instance, D-antigen units may be measured using the ELISA test
15 as described in Example 1 (“D-antigen quantification by ELISA”) below. European Pharmacopoeia provides a test sample (European Pharmacopoeia Biological Reference Preparation – available from Ph. Eur. Secretariat, e.g. Code P 216 0000) for standardisation of such methods between manufacturers (Pharmeuropa Special Issue, Bio 96-2). Thus the D-antigen unit value is well understood in the art.

20 The term “dose” herein is typically one administration of the vaccine of the invention, which is typically one injection. A typical human dose is 0.5mL. Of course various doses may be administered in a vaccine administration schedule.

 The term “IPV” or a vaccine comprising these components herein is intended to mean inactivated polio virus type 1 (e.g. Mahoney, as preferably used, or Brunhilde
25 as marketed by Statens Serum Institut under the name of DiTeKiPol), type 2 (e.g. MEF-1), or type 3 (e.g. Saukett), or a combination of either two or all three of these types. An example of a full (or standard) dose (40-8-32 D antigen units of IPV types 1, 2 and 3 respectively) IPV vaccine for the purposes of this invention could be Poliorix® (GSK Biologicals S.A.). Thus, where it is stated herein that X% of a
30 standard dose of IPV is present in a vaccine of the invention it is meant D-antigen units equating to X% of 40, 8, and/or 32 D-antigen units of IPV types 1, 2 and/or 3 respectively (as measured in each bulk IPV antigen type) are formulated within each dose of said vaccine.

The terms “lipopolysaccharide” (LPS) and “lipooligosaccharide” (LOS) are interchangeable.

The term “saccharide” throughout this specification may indicate polysaccharide or oligosaccharide and includes both. The capsular saccharide antigen
5 may be a full length polysaccharide or it may be extended to bacterial ‘sized-saccharides’ and ‘oligosaccharides’ (which naturally have a low number of repeat units, or which are polysaccharides reduced in size for manageability, but are still capable of inducing a protective immune response in a host) which are well known in the vaccine art (see for instance EP 497525).

10 The term “nucleic acid” herein can comprise single or double stranded deoxyribonucleic acid (DNA) or single or double stranded ribonucleic acid (RNA) or a mixture thereof.

The term “component(s)” from a pathogen or “component(s) affording protection to such a pathogen” within the vaccines of the invention herein is intended
15 to mean one or more antigen(s) from that pathogen.

The terms “around” or “approximately” herein are taken to mean $\pm 10\%$ of the stated value, but should be in keeping with the context of use.

20 DESCRIPTION OF FIGURES

Figure 1. Evolution of the Relative Potency (RP) of DTPwSF-HB-IPV “Method of production 3” with the IPV dose.

The potency of reduced dose IPV of the formulations “Method of production 3” was examined *in vivo* in comparison with reference formulation (Poliorix
25 formulation and DTPaIPVHB). RP of IPV was measured at doses 100%, 50%, 25% and 12.5% of standard IPV dose (40/8/32 D-antigen units for types 1/2/3).

Figure 2. Evolution of the Relative Potency (RP) of DTPwSF-HB-IPV formulation flow-sheet.

30 The potency of reduced dose IPV for both formulations “Method of production 3” and “Method of production 4” was examined *in vivo* in comparison with reference formulations (Poliorix formulation and DTPaIPVHB). RP was measured for both “Method of production 3” and “Method of production 4” at 25% of

the standard IPV dose (40/8/32 D-antigen units for types 1/2/3) in comparison to a placebo with 25% of IPV alone.

Figure 3. Relative potency of IPV types 1, 2 and 3 at time 0 and 8 months.

Relative potency of IPV was measured [relative to DTPaHBIPV (Pediarix) (Figure 3a) or Poliorix (Figure 3b)] to determine whether the Hib component has an effect on IPV potency and to evaluate the stability of IPV over time at different IPV doses.

DETAILED DESCRIPTION

The present invention provides a vaccine (e.g. a combination vaccine) comprising antigens from poliovirus (IPV) and optionally *Corynebacterium diphtheriae* (D), *Clostridium tetani* (T), *Bordetella pertussis* (P) or Hepatitis B.

The antigens of the invention

IPV vaccine components

Vaccines of the invention may be comprised of IPV type 1 or IPV type 2 or IPV type 3, or IPV types 1 and 2, or IPV types 1 and 3, or IPV types 2 and 3, or IPV types 1, 2 and 3.

Methods of preparing inactivated poliovirus (IPV) are well known in the art. In one embodiment, IPV should comprise types 1, 2 and 3 as is common in the vaccine art, and may be the Salk polio vaccine which is inactivated with formaldehyde (see for example, Sutter *et al.*, 2000, *Pediatr. Clin. North Am.* 47:287; Zimmerman & Spann 1999, *Am Fam Physician* 59:113; Salk et al., 1954, *Official Monthly Publication of the American Public Health Association* 44(5):563; Hennesen, 1981, *Develop. Biol. Standard* 47:139; Budowsky, 1991, *Adv. Virus Res.* 39:255).

In one embodiment the IPV is not adsorbed (e.g. before mixing with other components if present). In another embodiment, the IPV component(s) of the invention may be adsorbed onto an aluminium salt such as aluminium hydroxide (e.g. before or after mixing with other components if present). In another embodiment, the IPV component(s) of the invention may be adsorbed onto an aluminium salt such as aluminium phosphate. In a further embodiment the IPV component(s) may be adsorbed onto a mixture of both aluminium hydroxide and aluminium phosphate. If

adsorbed, one or more IPV components may be adsorbed separately or together as a mixture. IPV may be stabilised by a particular drying process as described in WO2004/039417.

Poliovirus may be grown in cell culture. The cell culture may be a VERO cell
5 line or PMKC, which is a continuous cell line derived from monkey kidney. VERO cells can conveniently be cultured microcarriers. Culture of the VERO cells before and during viral infection may involve the use of bovine-derived material, such as calf serum, and this material should be obtained from sources which are free from bovine spongiform encephalitis (BSE). Culture may also involve materials such as
10 lactalbumin hydrolysate. After growth, virions may be purified using techniques such as ultrafiltration, diafiltration, and chromatography. Prior to administration to patients, the viruses must be inactivated, and this can be achieved by treatment with formaldehyde.

Viruses may be grown, purified and inactivated individually, and then
15 combined to give a concentrate bulk mixture for IPV vaccine use or for addition to the adsorbed diphtheria and tetanus antigen and pertussis components for DTPw-IPV or DTPa-IPV comprising vaccines.

Antigens in vaccines of the invention will be present in "immunologically effective amounts" i.e. the administration of that amount to an individual, either in a
20 single dose or as part of a series, is effective for treatment or prevention of disease. Dosage treatment may be a single dose schedule or a multiple dose schedule (e.g. including booster doses).

Standard doses of polio vaccines today tend to contain 40 D antigen units of inactivated poliovirus type 1, 8 D antigen units of inactivated poliovirus type 2 and 32
25 D antigen units of inactivated poliovirus type 3 (e.g. Infanrix-IPV™).

However, the present inventors have surprisingly found that reduced doses of IPV can be used to obtain a good immune response. In one embodiment, an IPV vaccine dose of the present invention may comprise between 10 and 36 D-antigen units of IPV type 1 (e.g. 11-32, 12-28, 13-24, 14-20 or 15-19 D-antigen units). In
30 another embodiment, an IPV vaccine dose of the present invention may comprise IPV type 1 at a dose of 10-20 D-antigen units or a dose greater than 10 D-antigen units and less than 20 D-antigen units. In another embodiment, a vaccine dose of the present invention may comprise 26-49%, 30-45%, 33-40%, 35-37%, or approximately or

exactly one third of a standard 40 D-antigen unit dose of IPV type 1 (equivalent to approximately 10.4-19.6, 12-18, 13.2-16, 14-14.8 or 13.3 D-antigen units). In another embodiment, an IPV vaccine dose of the present invention may comprise 11-32 D-antigen units, 12-28 D-antigen units, 13-24 D-antigen units or 14-20 D-antigen units of IPV type 1.

Alternatively, an IPV vaccine dose of the present invention may comprise 10-19.5 D-antigen units, 12-19 D-antigen units, 14-18.5 D-antigen units, or 15-17 D-antigen units; for instance around or exactly 16 D-antigen units of IPV type 1.

In a further embodiment, the vaccines of the present invention may comprise less than 4 D-antigen units, 2-4 D-antigen units (equivalent to 25-50% of a standard 8 D-antigen unit dose) or around or exactly 3 D-antigen units of IPV type 2 (equivalent to 37.5% of a standard 8 D-antigen unit dose).

In another embodiment, the vaccine of the present invention may comprise approximately or exactly one third of a standard 8 D-antigen unit dose of IPV type 2 (equivalent to approximately 2.7 D-antigen units).

In a further embodiment, the vaccines of the present invention may comprise 2-7 D-antigen units of IPV type 2. In another embodiment, an IPV vaccine dose of the present invention may comprise 3-6 D-antigen units, or 4-5 D-antigen units of IPV type 2.

Alternatively, an IPV vaccine dose of the present invention may comprise 2-4.5 D-antigen units, 2.5-4 D-antigen units or 3-3.5 D-antigen units of IPV type 2.

In a further embodiment the vaccines of the present invention may comprise 8-20 D-antigen units, more than 8 and less than 20 D-antigen units, 9-19 D-antigen units, 10-18 D-antigen units, 11-17 D-antigen units, 12-16 D-antigen units, or 13-15 D-antigen units; for instance around or exactly 14 D-antigen units of IPV type 3 (equivalent to 25-62.5%, 28.125-59.375%, 31.25-46.875% or 43.75% of a standard 32 D-antigen unit dose).

In another embodiment, the vaccine of the present invention may comprise approximately or exactly one third of a standard 32 D-antigen unit dose of IPV type 3 (equivalent to approximately 10.7 D-antigen units).

In a further embodiment, an IPV vaccine dose of the present invention may comprise 8-29 D-antigen units, 9-26 D-antigen units, 10-23 D-antigen units, 11-20 D-antigen units, 12-17 D-antigen units, or 13-14 D-antigen units of IPV type 3.

Alternatively, an IPV vaccine dose of the present invention may comprise 8-19.5 D-antigen units, 9-19 D-antigen units, 10-18.5 D-antigen units, 11-18 D-antigen units, 12-17.5 D-antigen units, 13-17 D-antigen units, or 14-16 D-antigen units; for instance around or exactly 15 D-antigen units.

5

DTP vaccine components

DTP vaccines are well known vaccines to prevent or treat diphtheria, tetanus and *B. pertussis* disease. The vaccines of the invention may comprise diphtheria, tetanus and/or pertussis component(s).

10 The diphtheria antigen is typically a diphtheria toxoid. The preparation of diphtheria toxoids (DT) is well documented. Any suitable diphtheria toxoid may be used. For instance, DT may be produced by purification of the toxin from a culture of *Corynebacterium diphtheriae* followed by chemical detoxification, but is alternatively made by purification of a recombinant, or genetically detoxified analogue of the toxin
15 (for example, CRM197, or other mutants as described in US 4,709,017, US 5,843,711, US 5,601,827, and US 5,917,017). In one embodiment DT is present at an amount of 5-50, 7-30Lf or approximately or exactly 7.5Lf or 25Lf per 0.5mL dose. In a further embodiment DT is present at a low dose of less than 5Lf, or 1-4Lf or approximately or exactly 2Lf per 0.5mL dose. In one embodiment, the diphtheria
20 toxoid of the invention may be adsorbed onto an aluminium salt such as aluminium hydroxide. In another embodiment, the diphtheria toxoid of the invention may be adsorbed onto an aluminium salt such as aluminium phosphate. In a further embodiment the diphtheria toxoid may be adsorbed onto a mixture of both aluminium hydroxide and aluminium phosphate.

25 The tetanus antigen of the invention is typically a tetanus toxoid. Methods of preparing tetanus toxoids (TT) are well known in the art. In one embodiment TT is produced by purification of the toxin from a culture of *Clostridium tetani* followed by chemical detoxification, but is alternatively made by purification of a recombinant, or genetically detoxified analogue of the toxin (for example, as described in EP 209281).
30 Any suitable tetanus toxoid may be used. 'Tetanus toxoid' may encompass immunogenic fragments of the full-length protein (for instance Fragment C – see EP 478602). In one embodiment TT is present at an amount of 2.5-30Lf, 3-20 Lf, 5-15Lf or exactly or approximately 10Lf per 0.5mL dose. In one embodiment, the tetanus toxoid of the invention may be adsorbed onto an aluminium salt such as aluminium

hydroxide. In another embodiment, the tetanus toxoid of the invention may be adsorbed onto an aluminium salt such as aluminium phosphate. In a further embodiment the tetanus toxoid may be adsorbed onto a mixture of both aluminium hydroxide and aluminium phosphate.

5 The pertussis component of the invention may be either acellular (Pa) where purified pertussis antigens are used or whole-cell (Pw) where killed whole cell pertussis is used as the pertussis component. Pw may be inactivated by several known methods, including mercury free methods. Such methods may include heat (e.g. 55-65°C or 56-60°C, for 5-60 minutes or for 10-30 minutes, e.g. 60°C for 30 minutes),
10 formaldehyde (e.g. 0.1% at 37°, 24 hours), glutaraldehyde (e.g. 0.05% at room temperature, 10 minutes), acetone-I (e.g. three treatments at room temperature) or acetone-II (e.g. three treatments at room temperature and fourth treatment at 37°C) inactivation (see for example Gupta *et al.*, 1987, J. Biol. Stand. 15:87; Gupta *et al.*, 1986, Vaccine, 4:185). Methods of preparing killed, whole-cell *Bordetella pertussis*
15 (Pw) suitable for this invention are disclosed in WO 93/24148, as are suitable formulation methods for producing DT-TT-Pw-HepB vaccines. Thiomersal has been used in the past in the preparation of killed whole-cell *Bordetella pertussis* (see below). However, in one embodiment it is not used in the formulation process of the vaccines of the present invention.

20 A Pw dose of 5-50 IOU, 7-40 IOU, 9-35 IOU, 11-30 IOU, 13-25 IOU, 15-21 IOU or around or exactly 20 IOU is typically used.

 Acellular Pa vaccines are also well known, and may comprise 2 or more antigens from: pertussis toxoid (PT), filamentous haemagglutinin (FHA), pertactin (PRN), agglutinogens 2 & 3. In one embodiment, the Pa vaccine comprises PT, FHA
25 and PRN. Kits or vaccines of the invention may comprise PT detoxified by a well known method of formaldehyde treatment or by means of mutations (PT derivative). Substitutions of residues within the S1 subunit of the protein have been found to result in a protein which retains its immunological and protective properties of the PT, but with reduced or no toxicity (EP 322533). The detoxifying mutations discussed in the
30 claims of EP322533 are examples of the DT detoxified mutants of the present invention. Such mutants may be used at doses lower than 20-25µg.

In one embodiment PT is used at an amount of 2-50 μ g, 5-40 μ g, 10-30 μ g or exactly or approximately 25 μ g per 0.5mL dose. In another embodiment PT is used at an amount of exactly or approximately 2.5 or 8 μ g per 0.5mL dose.

5 In one embodiment FHA is used at an amount of 2-50 μ g, 5-40 μ g, 10-30 μ g or exactly or approximately 25 μ g per 0.5mL dose. In another embodiment FHA is used at an amount of exactly or approximately 2.5 or 8 μ g per 0.5mL dose.

In one embodiment PRN is used at an amount of 0.5-20 μ g, 0.8-15 μ g, 2-10 μ g or exactly or approximately 8 μ g per 0.5mL dose. In another embodiment PRN is used at an amount of exactly or around 0.8 or 2.5 μ g per 0.5mL.

10 In one embodiment, the pertussis component of the invention may be adsorbed onto an aluminium salt such as aluminium hydroxide. In another embodiment, the pertussis component of the invention may be adsorbed onto an aluminium salt such as aluminium phosphate. In a further embodiment the pertussis component may be adsorbed onto a mixture of both aluminium hydroxide and aluminium phosphate. For
15 instance in one embodiment at least PRN is adsorbed onto aluminium hydroxide with PT/FHA adsorbed onto aluminium hydroxide, aluminium phosphate or a mixture of both.

Further antigens

20 Vaccine formulations of the invention, optionally also comprising DTP (DTPw or DTPa), can additionally comprise one or more antigens selected from the group consisting of: Hepatitis B surface antigen, *Haemophilus influenzae* b antigen(s), *Neisseria meningitidis* A antigen(s), *Neisseria meningitidis* C antigen(s), *Neisseria meningitidis* W-135 antigen(s), *Neisseria meningitidis* Y antigen(s), *Neisseria meningitidis* B bleb or purified antigen(s), Hepatitis A antigen(s), Salmonella typhi antigen(s) and RTS,S. Typically the capsular saccharide or LOS antigens of these
25 pathogens may be used. Antigens will typically be present at a concentration of at least 1 μ g/mL each, for instance 1-20 μ g/mL, 2-15 μ g/mL, 2.5-10 μ g/mL, 3-8 μ g/mL, or 4-6 μ g/mL. In general, the concentration of any antigen will be sufficient to elicit an
30 immune response against that antigen. It is preferred that the protective efficacy of individual antigens is not removed by combining them, although actual immunogenicity (e.g. ELISA titres) may be reduced.

The further antigen(s) may in one embodiment of the invention be adsorbed onto an aluminium salt such as aluminium hydroxide. In another embodiment, the further antigens of the invention may be adsorbed onto an aluminium salt such as aluminium phosphate. In a further embodiment the further antigens may be adsorbed onto a mixture of both aluminium hydroxide and aluminium phosphate, or may be unadsorbed.

Where a capsular saccharide or LOS antigen is used it may be conjugated to a carrier protein comprising T helper epitopes in order to enhance immunogenicity. The invention may also comprise free "carrier proteins".

As an alternative to using protein antigens in the compositions of the invention, nucleic acid encoding the antigen may be used. Protein components of the compositions of the invention may thus be replaced by nucleic acid (for instance DNA, which may be in the form of a plasmid) that encodes the protein. Similarly, compositions of the invention may comprise proteins which mimic saccharide antigens e.g. mimotopes or anti-idiotypic antibodies. These may replace individual saccharide components, or may supplement them.

Hepatitis B antigen

The preparation of Hepatitis B surface antigen (HBsAg) is well documented. See for example, Hartford *et al.*, 1983, Develop. Biol. Standard 54:125, Gregg *et al.*, 1987, Biotechnology 5:479, EP0226846, EP0299108. It may be prepared as follows. One method involves purifying the antigen in particulate form from the plasma of chronic hepatitis B carriers, as large quantities of HBsAg are synthesised in the liver and released into the blood stream during an HBV infection. Another method involves expressing the protein by recombinant DNA methods. The HBsAg may be prepared by expression in the *Saccharomyces cerevisiae* yeast, pichia, insect cells (e.g. Hi5) or mammalian cells. The HBsAg may be inserted into a plasmid, and its expression from the plasmid may be controlled by a promoter such as the "GAPDH" promoter (from the glyceraldehyde-3-phosphate dehydrogenase gene). The yeast may be cultured in a synthetic medium. HBsAg can then be purified by a process involving steps such as precipitation, ion exchange chromatography, and ultrafiltration. After purification, HBsAg may be subjected to dialysis (e.g. with cysteine). The HBsAg may be used in a particulate form.

As used herein the expression "Hepatitis B surface antigen" or "HBsAg" includes any HBsAg antigen or fragment thereof displaying the antigenicity of HBV surface antigen. It will be understood that in addition to the 226 amino acid sequence of the HBsAg S antigen (see Tiollais *et al.*, 1985, Nature 317:489 and references therein) HBsAg as herein described may, if desired, contain all or part of a pre-S sequence as described in the above references and in EP0278940. In particular, the HBsAg may comprise a polypeptide comprising an amino acid sequence comprising residues 133-145 followed by residues 175-400 of the L-protein of HBsAg relative to the open reading frame on a Hepatitis B virus of ad serotype (this polypeptide is referred to as L*; see EP0414374). HBsAg within the scope of the invention may also include the preS1-preS2 -S polypeptide described in EP0198474 (Endotronics) or analogues thereof such as those described in EP0304578 (McCormick and Jones). HBsAg as herein described can also refer to mutants, for example the "escape mutant" described in WO 91/14703 or EP0511855A1, especially HBsAg wherein the amino acid substitution at position 145 is to arginine from glycine.

The HBsAg may be in particle form. The particles may comprise for example S protein alone or may be composite particles, for example L*, S) where L* is as defined above and S denotes the S-protein of HBsAg. The said particle is advantageously in the form in which it is expressed in yeast.

In one embodiment, HBsAg is the antigen used in EngerixB™ (GlaxoSmithKline Biologicals S.A.), which is further described in WO93/24148.

In one embodiment, HBsAg is present at an amount of 5-20µg, 8-15µg or approximately or exactly 10µg per 0.5mL dose.

Hepatitis B surface antigen may be adsorbed onto aluminium phosphate, which may be done before mixing with the other components (described in WO93/24148). The Hepatitis B component should be substantially thiomersal free (method of preparation of HBsAg without thiomersal has been previously published in EP1307473).

30 ***Haemophilus influenzae b* antigen(s)**

Vaccines comprising antigens from *Haemophilus influenzae* type B have been described in WO97/00697. The vaccines of the invention may use any suitable *Haemophilus influenzae* type B antigen. The antigen may be capsular saccharide

(PRP) from *Haemophilus influenzae* type B conjugated to a carrier protein (Hib). The saccharide is a polymer of ribose, ribitol and phosphate. The Hib antigen may optionally be adsorbed onto aluminium phosphate as described in WO97/00697, or may be unadsorbed as described in WO02/00249 or may not have undergone a specific process of adsorption.

By an antigen being 'unadsorbed onto an aluminium adjuvant salt' herein it is meant for example that an express or dedicated adsorption step for the antigen on fresh aluminium adjuvant salt is not involved in the process of formulating the composition.

Hib may be conjugated to any carrier which can provide at least one T-helper epitope (examples of which are described below), and may be tetanus toxoid, diphtheria toxoid, CRM-197 (diphtheria toxin mutant) or Protein D.

Hib may be lyophilised and may be reconstituted extemporaneously (e.g. with diluent, optionally comprising other antigenic components of the vaccines of the invention).

In one embodiment, Hib is present at an amount of 5-20µg, 8-15µg or approximately or exactly 10µg saccharide per 0.5mL dose.

In a further embodiment, Hib is present at a low dose (e.g. 1-6µg, 2-4µg or around or exactly 2.5µg saccharide) as described in WO 02/00249.

***Neisseria meningitidis* types A, C, W or Y antigens**

The vaccines of the invention may further comprise a capsular saccharide of a bacterium selected from the group consisting of *N. meningitidis* type A (MenA, optionally conjugated to a carrier protein), *N. meningitidis* type C (MenC, optionally conjugated to a carrier protein), *N. meningitidis* type W-135 (MenW, optionally conjugated to a carrier protein), and *N. meningitidis* type Y (MenY, optionally conjugated to a carrier protein).

The vaccines of the invention may comprise one or more antigens from the different strains of *N. meningitidis*, which may be used alone or in any combination of two, three or four components as detailed below:

MenA, MenC, MenW, MenY, or MenA + MenC, MenA + MenW, MenA + MenY, MenC + MenW, MenC + MenY, MenW + MenY or MenA + MenC + MenW, MenA

+ MenC + MenY, MenA + MenW + MenY, MenC + MenW + MenY or MenA + MenC + MenW + MenY.

In one embodiment, the *Neisseria meningitidis* component(s) of the invention may be adsorbed onto an aluminium salt such as aluminium hydroxide. In another embodiment, the *Neisseria meningitidis* component(s) of the invention may be adsorbed onto an aluminium salt such as aluminium phosphate. In a further embodiment the *Neisseria meningitidis* component(s) may be adsorbed onto a mixture of both aluminium hydroxide and aluminium phosphate. In one embodiment the *Neisseria meningitidis* component(s) may be unadsorbed onto an adjuvant, e.g. an aluminium adjuvant salt.

***Neisseria meningitidis* type B bleb or antigen(s)**

The vaccines of the invention may also comprise a MenB component such as an outer membrane vesicle or bleb as described in WO01/09350, WO03/105890, WO04/014417, or WO04/014418 or a conjugated MenB capsular saccharide (or derivative thereof) antigen (e.g. see WO 96/40239) or a free or conjugated L2 or L3 or L2 and L3 meningococcal LOS (as per WO 2004/014417). In one embodiment, the MenB component(s) of the invention may be adsorbed onto an aluminium salt such as aluminium hydroxide. In another embodiment, the MenB component(s) of the invention may be adsorbed onto an aluminium salt such as aluminium phosphate. In a further embodiment the MenB component(s) may be adsorbed onto a mixture of both aluminium hydroxide and aluminium phosphate. In one embodiment the MenB component(s) may be unadsorbed onto an adjuvant, e.g. an aluminium adjuvant salt.

Salmonella typhi antigen(s)

The vaccines of the invention may further comprise the Vi saccharide from *Salmonella typhi*, which may be the registered product Typherix®, described in EP1107787, or a conjugate thereof (e.g. with a carrier protein as described herein). The conjugation process may be carried out as described in WO 2007/000343. In one embodiment, the Vi saccharide(s) of the invention may be adsorbed onto an aluminium salt such as aluminium hydroxide. In another embodiment, the Vi saccharide(s) of the invention may be adsorbed onto an aluminium salt such as aluminium phosphate. In a further embodiment the Vi saccharide(s) may be adsorbed

onto a mixture of both aluminium hydroxide and aluminium phosphate. In one embodiment the Vi saccharide(s) may be unadsorbed onto an adjuvant, e.g. an aluminium adjuvant salt.

5

Hepatitis A antigen(s)

The component affording protection against Hepatitis A may be a killed attenuated Hepatitis A vaccine, for example the product known as Havrix™ (Registered Trade Mark of GlaxoSmithKline Biologicals S.A.) which is a killed attenuated vaccine derived from the HM-175 strain of Hepatitis A virus (HAV) (see “Inactivated Candidate Vaccines for Hepatitis A” by F.E. Andre *et al.*, 1980, Prog. Med. Virol. 37:72 and the product monograph “Havrix” published by SmithKline Beecham Biologicals 1991). Flehmig *et al.* (1990, Prog. Med Virol. 37:56) have reviewed the clinical aspects, virology, immunology and epidemiology of Hepatitis A and discussed approaches to the developments of vaccines against this common viral infection. As used herein the expression “HAV antigen” refers to any antigen capable of stimulating neutralising antibody to HAV in humans. In one embodiment the HAV antigen comprises inactivated attenuated virus particles, or in another embodiment it may be a HAV capsid or HAV viral protein, which may conveniently be obtained by recombinant DNA technology. In one embodiment, the Hepatitis A component of the invention may be adsorbed onto an aluminium salt such as aluminium hydroxide. In another embodiment, the Hepatitis A component of the invention may be adsorbed onto an aluminium salt such as aluminium phosphate. In a further embodiment the Hepatitis A component may be adsorbed onto a mixture of both aluminium hydroxide and aluminium phosphate.

Malarial antigen(s)

The vaccines of the invention may further comprise Malarial antigen(s). The Malarial antigen may be RTS,S (hybrid protein between CS and HBsAg – described in US 6,306,625 and EP 0614465). In one embodiment, RTS,S may be used in the vaccines of the invention in place of HBsAg. Other Malarial antigens may also be used in the vaccines of the invention, including CS protein, RTS, TRAP, 16kD

protein of B 2992, AMA-1, MSP1, optionally including CpG (WO2006/029887, WO98/05355, WO01/00231).

In one embodiment, the Malarial antigen(s) of the invention may be adsorbed onto an aluminium salt such as aluminium hydroxide. In another embodiment, the Malarial antigen(s) of the invention may be adsorbed onto an aluminium salt such as aluminium phosphate. In a further embodiment the Malarial antigen(s) may be adsorbed onto a mixture of both aluminium hydroxide and aluminium phosphate. In one embodiment the Malarial antigen is adjuvanted with an oil-in-water emulsion and/or lipid A derivative (such as MPL) and or sterol (such as cholesterol) and/or tocol (such as α -tocopherol) In another embodiment the Malaria antigen(s) may be unadsorbed onto an adjuvant, e.g. an aluminium adjuvant salt.

Conjugates

Bacterial capsular saccharide conjugates of the invention may comprise any carrier peptide, polypeptide or protein comprising at least one T-helper epitope. The carrier protein(s) used may be selected from the group consisting of: tetanus toxoid, diphtheria toxoid, CRM197, recombinant diphtheria toxin (as described in any of US 4,709,017, WO 93/25210, WO 95/33481, or WO 00/48638), pneumolysin (optionally chemically detoxified, or a detoxified mutant) from *S. pneumoniae* (see e.g. WO 2004/081515 and references referred to therein), OMPC from *N. meningitidis* (EP 0372501), and protein D (PD) from *H. influenzae* (EP 594610). Other carriers may include synthetic peptides (EP 0378881; EP 0427347), heat shock proteins (WO 93/17712; WO 94/03208), pertussis proteins (WO 98/58668; EP 0471177), cytokines (WO 91/01146), lymphokines (WO 91/01146), hormones (WO 91/01146), growth factors (WO 91/01146), artificial proteins comprising multiple human CD4⁺ T cell epitopes from various pathogen-derived antigens (Falugi *et al.*, 2001, Eur. J. Immunol. 31:3816), pneumococcal surface protein PspA (WO 02/091998), iron uptake proteins (WO 01/72337), toxin A or B from *C. difficile* (WO 00/61761), pneumococcal PhtD (WO 00/37105), pneumococcal PhtDE (e.g. WO 01/98334 & WO 03/054007), PhtX, etc.

Saccharides may all be on the same carrier, particularly all saccharides from a particular organism, for instance MenA, MenC, MenW and MenY saccharides may all be conjugated to TT, DT or CRM-197. However, due to the known effect of carrier

suppression, it may be advantageous if in each of the compositions of the invention the saccharide antigens contained therein ('n' antigens) are conjugated to more than one carrier. Thus (n-1) of the saccharides could be carried (separately) on one type of carrier, and 1 on a different carrier, or (n-2) on one, and 2 on two different carriers, etc. For example, in a vaccine containing 4 bacterial saccharide conjugates, 1, 2 or all four could be conjugated to different carriers). Protein D, however, may be used for various (2, 3, 4 or more) saccharides in a composition without a marked carrier suppression effect. Hib may be present as a TT, DT or CRM197 conjugate, and MenA, MenC, MenY and MenW may be either TT, DT, CRM197 or PD conjugates. Vi may be present as a TT, DT or CRM197 conjugate. Protein D is a useful carrier as it provides a further antigen which can provide protection against *H. influenzae*. In one embodiment, all saccharides are conjugated to the same carrier protein.

Vi may be conjugated to a carrier protein for instance by a method using carbodiimide (e.g. EDAC) condensation chemistry (given that the Vi repeat subunit comprises carboxylic acid groups). This could be achieved either by (i) a single carbodiimide reaction between COOH of Vi and NH₂ of protein or (ii) a double carbodiimide reaction which can occur either between COOH of Vi and NH₂ of a homobifunctional linker molecule and COOH of protein and NH₂ of the heterobifunctional linker molecule and NH₂ of protein and COOH of the heterobifunctional linker molecule.

Conjugation may be used in conjunction with free carrier protein(s). In one embodiment, when a given carrier protein is present in both free and conjugated form in a composition of the invention, the unconjugated form is no more than 5% of the total amount of the carrier protein in the composition as a whole, or in another embodiment is present at less than 2% by weight.

The saccharide may be linked to the carrier protein by any known method (for example, by Likhite, U.S. Patent 4,372,945 and by Armor *et al.*, U.S. Patent 4,474,757), with any suitable linker where necessary.

The saccharide will typically be activated or functionalised prior to conjugation. Activation may involve, for example, cyanylating agents such as CDAP (1-cyano-dimethylaminopyridinium tetrafluoroborate) (WO 95/08348 & WO 96/29094). The cyanilation reaction can be performed under relatively mild conditions, which avoids hydrolysis of the alkaline sensitive saccharides. This

synthesis allows direct coupling to a carrier protein. Other suitable techniques use carbodiimides, hydrazides, active esters, norborane, p-nitrobenzoic acid, N-hydroxysuccinimide, S-NHS, EDC or TSTU.

Linkages via a linker group may be made using any known procedure, for example, the procedures described in US 4,882,317 and US 4,695,624. One type of linkage involves reductive amination of the saccharide, coupling the resulting amino group with one end of an adipic acid linker group (EP 0477508, Porro et al., 1985, Mol. Immunol. 22:907, EP 0208375), and then coupling a protein to the other end of the adipic acid linker group. Other linkers include B-propionamido (WO 00/10599), nitrophenyl-ethylamine (Gever *et al.*, 1979, Med. Microbiol. Immunol. 165:171), haloacyl halides (US 4,057,685), glycosidic linkages (US 4,673,574; US 4,761,283; US 4,808,700), 6-aminocaproic acid (US 4,459,286), ADH (US 4,965,338), C4 to C12 moieties (US 4,663,160), etc. As an alternative to using a linker, direct linkage can be used. Direct linkages to the protein may comprise oxidation of the saccharide followed by reductive amination with the protein, as described in, for example US 4,761,283 and US 4,356,170 or a direct CDAP reaction.

After conjugation, free and conjugated saccharides can be separated. There are many suitable methods for this separation, including hydrophobic chromatography, tangential ultrafiltration, diafiltration, etc (see also Lei *et al.*, 2000, Dev Biol. (Basel). 103:259; WO 00/38711; US 6,146,902). In one embodiment, if a vaccine comprises a given saccharide in both free and conjugated forms, the unconjugated form is no more than 20% by weight of the total amount of that saccharide in the composition as a whole (e.g. $\leq 15\%$, $\leq 10\%$, $\leq 5\%$, $\leq 2\%$, $\leq 1\%$).

An amount of saccharide which is capable of conferring protection to a host (an effective amount) can be determined by the skilled person. In one embodiment, each dose will comprise 0.1-100 μg of saccharide, in another embodiment each dose will comprise 0.1-50 μg , in a further embodiment each dose will comprise 0.1-10 μg , in yet another embodiment each dose will comprise 1 to 5 μg .

30 Adjuvants

The vaccines of the invention may include a pharmaceutically acceptable excipient such as a suitable adjuvant. Suitable adjuvants include an aluminium salt such as aluminium hydroxide or aluminium phosphate, but may also be a salt of

calcium, iron or zinc, or may be an insoluble suspension of acylated tyrosine, or acylated sugars, or may be cationically or anionically derivatised saccharides, polyphosphazenes, biodegradable microspheres, monophosphoryl lipid A (MPL), lipid A derivatives (e.g. of reduced toxicity), 3-O-deacylated MPL, quil A, Saponin, QS21, tocol (EP 0382271), Freund's Incomplete Adjuvant (Difco Laboratories, Detroit, MI), Merck Adjuvant 65 (Merck and Company, Inc., Rahway, NJ), AS-2 (Smith-Kline Beecham, Philadelphia, PA), CpG oligonucleotides, bioadhesives and mucoadhesives, microparticles, liposomes, polyoxyethylene ether formulations, polyoxyethylene ester formulations, muramyl peptides or imidazoquinolone compounds (e.g. imiquamod and its homologues). Human immunomodulators suitable for use as adjuvants in the invention include cytokines such as interleukins (e.g. IL-1, IL-2, IL-4, IL-5, IL-6, IL-7, IL-12, etc), macrophage colony stimulating factor (M-CSF), tumour necrosis factor (TNF), granulocyte, macrophage colony stimulating factor (GM-CSF) may also be used as adjuvants.

In one embodiment of the invention, the adjuvant composition of the formulations induces an immune response predominantly of the TH1 type. High levels of TH1-type cytokines (e.g. IFN- γ , TNF α , IL-2 and IL-12) tend to favour the induction of cell mediated immune responses to an administered antigen. Within one embodiment, in which a response is predominantly TH1-type, the level of TH1-type cytokines will increase to a greater extent than the level of TH2-type cytokines. The levels of these cytokines may be readily assessed using standard assays. For a review of the families of cytokines, see Mosmann and Coffman, 1989, Ann. Rev. Immunol. 7:145.

Accordingly, suitable adjuvant systems which promote a predominantly TH1 response include, derivatives of lipid A (e.g. of reduced toxicity), Monophosphoryl lipid A (MPL) or a derivative thereof, particularly 3-de-O-acylated monophosphoryl lipid A (3D-MPL), and a combination of monophosphoryl lipid A, optionally 3-de-O-acylated monophosphoryl lipid A together with an aluminium salt. An enhanced system involves the combination of a monophosphoryl lipid A and a saponin derivative, particularly the combination of QS21 and 3D-MPL as disclosed in WO 94/00153, or a less reactogenic composition where the QS21 is quenched with cholesterol as disclosed in WO 96/33739. A particularly potent adjuvant formulation involving QS21, 3D-MPL and tocopherol in an oil in water emulsion is described in

WO 95/17210. The vaccine may additionally comprise a saponin, which may be QS21. The formulation may also comprise an oil in water emulsion and tocopherol (WO 95/17210). Unmethylated CpG containing oligonucleotides (WO 96/02555) are also preferential inducers of a TH1 response and are suitable for use in the present invention.

The vaccines of the invention may also comprise combinations of aspects of one or more of the adjuvants of the invention identified above.

Any adjuvant of the invention can be adsorbed by or combined with IPV component of the invention.

When referring to aluminium hydroxide or aluminium phosphate, reference is made to all aluminium hydroxide and or aluminium phosphate adjuvants as described by Hem and White (Pharm Biotechnol. 1995;6:249-276).

In one embodiment, aluminium phosphate may also be referred to as aluminium hydroxyphosphate. In another embodiment, aluminium phosphate has a negative charge at a pH of 7.4. Typically, the isoelectric point (pI) of aluminium phosphate is 5-7, or 6-7 or around or exactly 5. In a further embodiment, aluminium phosphate has a molar phosphate : aluminium ratio of 0.3-0.9, or 0.3-0.6, or 0.8-0.9.

In one embodiment, aluminium hydroxide has a positive charge at a pH of 7.4. Typically, the pI of aluminium hydroxide is 8-11, 9-11, 10-11 or around or exactly 11.

Typically, the total aluminium content is 200-1000 μ g, 300-900 μ g, 400-800 μ g, 500-700 μ g or around or exactly 630 μ g Al³⁺ per 0.5 mL dose. This may be all aluminium hydroxide or all aluminium phosphate. Alternatively Al³⁺ content may be from a mixture of aluminium hydroxide and aluminium phosphate in the following ratio: 1:8-8:1, 1:4-4:1, 3:8-8:3, 1:2-2:1 or 1:1 of aluminium phosphate: aluminium hydroxide. In one embodiment a ratio of 12:1-4:1, 11:1-5:1, 10:1-6:1, 9:1-7:1 or 8:1 of aluminium phosphate: aluminium hydroxide is used.

Although most aluminium is provided by preadsorbed antigens before mixture to form a combination vaccine, some aluminium may be added in free form during formulation of the combination vaccine of the invention, e.g. before the pH adjustment step described herein. Typically, free aluminium content per 0.5 mL dose may be 0-300 μ g, 50-250 μ g, 75-200 μ g, 100-150 μ g or around or exactly 115 μ g of Al³⁺. Free Al³⁺ may be all Al(OH)₃ or all AlPO₄, or a mixture of Al(OH)₃ and AlPO₄ in the following ratio (w:w Al³⁺:Al³⁺): 1:1-1:6, 1:1.1-1:5, 1:1.2-1:4, 1:1.3-1:3, 1:1.4-

1:2, e.g. 23/92 or 69/46 or 6:1-1:1, 5:1-1.1:1, 4:1-1.2:1, 3:1-1.3:1, 2:1-1.4:1, e.g. 46/69 or 92/23.

Alternatively certain components of the vaccines of the invention may be not expressly adsorbed onto adjuvant, in particular aluminium salts.

5 IPV may be unadsorbed or adsorbed onto $\text{Al}(\text{OH})_3$ or a mixture of $\text{Al}(\text{OH})_3$ and AlPO_4 . DT may be adsorbed onto $\text{Al}(\text{OH})_3$ or AlPO_4 , TT may be adsorbed onto $\text{Al}(\text{OH})_3$ or AlPO_4 , Pw may be adsorbed onto or mixed with AlPO_4 , PRN may be adsorbed onto $\text{Al}(\text{OH})_3$, FHA may be adsorbed onto $\text{Al}(\text{OH})_3$, PT may be adsorbed onto $\text{Al}(\text{OH})_3$, HB may be adsorbed onto AlPO_4 , Hib may be adsorbed onto AlPO_4 or
 10 unadsorbed, Men ACWY may be adsorbed onto $\text{Al}(\text{OH})_3$ or AlPO_4 or unadsorbed, MenB component may be adsorbed onto $\text{Al}(\text{OH})_3$ or AlPO_4 or unadsorbed, Vi may be adsorbed onto $\text{Al}(\text{OH})_3$ or AlPO_4 or unadsorbed, HepA may be adsorbed onto $\text{Al}(\text{OH})_3$ or AlPO_4 .

Antigens which are preadsorbed onto an aluminium salt can be preadsorbed
 15 individually prior to mixing. In another embodiment, a mix of antigens may be preadsorbed prior to mixing with further adjuvants. In one embodiment, IPV may be adsorbed separately or as a mixture of IPV types 1, 2 and 3 or when mixed with adsorbed D and T components.

The meaning of “adsorbed antigen” is for example taken to mean greater than
 20 20%, 30%, 40%, 50%, 60%, 70%, 80%, or 90% adsorbed.

The meaning of the terms “aluminium phosphate” and “aluminium hydroxide” as used herein includes all forms of aluminium hydroxide or aluminium phosphate which are suitable for adjuvanting vaccines. For example, aluminium phosphate can be a precipitate of insoluble aluminium phosphate (amorphous, semi-crystalline or
 25 crystalline), which can be optionally but not exclusively prepared by mixing soluble aluminium salts and phosphoric acid salts. “Aluminium hydroxide” can be a precipitate of insoluble (amorphous, semi-crystalline or crystalline) aluminium hydroxide, which can be optionally but not exclusively prepared by neutralising a solution of aluminium salts. Particularly suitable are the various forms of aluminium
 30 hydroxide and aluminium phosphate gels available from commercial sources for example, Alhydrogel (aluminium hydroxide, 3% suspension in water) and Adjuphos (aluminium phosphate, 2% suspension in saline) supplied by Brenntag Biosector (Denmark).

Non-immunological components of vaccines of the invention

Vaccines of the invention will typically, in addition to the antigenic and adjuvant components mentioned above, comprise one or more “pharmaceutically acceptable carriers or excipients”, which include any excipient that does not itself induce the production of antibodies harmful to the individual receiving the composition. Suitable excipients are typically large, slowly metabolised macromolecules such as proteins, saccharides, polylactic acids, polyglycolic acids, polymeric amino acids, amino acid copolymers, sucrose (Paoletti *et al.*, 2001, Vaccine, 19:2118), trehalose (WO 00/56365), lactose and lipid aggregates (such as oil droplets or liposomes). Such carriers are well known to those of ordinary skill in the art. The vaccines may also contain diluents, such as water, saline, glycerol, etc. Additionally, auxiliary substances, such as wetting or emulsifying agents, pH buffering substances, and the like, may be present. Sterile pyrogen-free, phosphate buffered physiologic saline is a typical carrier. A thorough discussion of pharmaceutically acceptable excipients is available in reference Gennaro, 2000, Remington: The Science and Practice of Pharmacy, 20th edition, ISBN:0683306472.

Compositions of the invention may be lyophilised or in aqueous form, i.e. solutions or suspensions. Liquid formulations of this type allow the compositions to be administered direct from their packaged form, without the need for reconstitution in an aqueous medium, and are thus ideal for injection. Compositions may be presented in vials, or they may be presented in ready filled syringes. The syringes may be supplied with or without needles. A syringe will include a single dose of the composition, whereas a vial may include a single dose or multiple doses (e.g. 2 doses). In one embodiment the dose is for human. In a further embodiment the dose is for an adult, adolescent, toddler, infant or less than one year old human and may be administered by injection.

Liquid vaccines of the invention are also suitable for reconstituting other vaccines from a lyophilised form. Where a vaccine is to be used for such extemporaneous reconstitution, the invention provides a kit, which may comprise two vials, or may comprise one ready-filled syringe and one vial, with the contents of the syringe being used to reconstitute the contents of the vial prior to injection.

Vaccines of the invention may be packaged in unit dose form or in multiple dose form (e.g. 2 doses). For multiple dose forms, vials are preferred to pre-filled

syringes. Effective dosage volumes can be routinely established, but a typical human dose of the composition for injection has a volume of 0.5mL.

In one embodiment, vaccines of the invention have a pH of between 6.0 and 8.0, in another embodiment vaccines of the invention have a pH of between 6.3 and 6.9, e.g. 6.6 ± 0.2 . Vaccines may be buffered at this pH. Stable pH may be maintained by the use of a buffer. If a composition comprises an aluminium hydroxide salt, a histidine buffer may be used (WO03/009869). The composition should be sterile and/or pyrogen free.

Compositions of the invention may be isotonic with respect to humans.

Vaccines of the invention may include an antimicrobial, particularly when packaged in a multiple dose format. Thiomersal should be avoided as this leads to loss of potency of the IPV component. Other antimicrobials may be used, such as 2-phenoxyethanol or parabens (methyl, ethyl, propyl parabens). Any preservative is preferably present at low levels. Preservative may be added exogenously and/or may be a component of the bulk antigens which are mixed to form the composition (e.g. present as a preservative in pertussis antigens).

In one embodiment, vaccines of the invention are thiomersal free or substantially thiomersal free. By "thiomersal free" or "substantially thiomersal free" it is meant that there is not enough thiomersal present in the final formulation to negatively impact the potency of the IPV component. For instance, if thiomersal is used during the Pw or Hepatitis B surface antigen purification process it should be substantially removed prior to mixture with IPV. Thiomersal content in the final vaccine should be less than $0.025 \mu\text{g}/\mu\text{g}$ protein, $0.02 \mu\text{g}/\mu\text{g}$ protein, $0.01 \mu\text{g}/\mu\text{g}$ protein or $0.001 \mu\text{g}/\mu\text{g}$ protein, for instance $0 \mu\text{g}/\mu\text{g}$ protein. In one embodiment, thiomersal is not added nor used in the purification of any component. See for instance EP1307473 for Hepatitis B and see above for Pw processes where killing is achieved not in the presence of thiomersal.

Vaccines of the invention may comprise detergent e.g. a Tween (polysorbate), such as Tween 80. Detergents are generally present at low levels e.g. $<0.01\%$.

Vaccines of the invention may include sodium salts (e.g. sodium chloride) to give tonicity. The composition may comprise sodium chloride. In one embodiment, the concentration of sodium chloride in the composition of the invention is in the range of 0.1 to 100 mg/mL (e.g. 1-50mg/mL, 2-20mg/mL, 5-15mg/mL) and in a

further embodiment the concentration of sodium chloride is $10 \pm 2 \text{ mg/mL}$ NaCl e.g. about 9 mg/mL .

Vaccines of the invention will generally include a buffer. A phosphate or histidine buffer is typical.

5 Vaccines of the invention may include free phosphate ions in solution (e.g. by the use of a phosphate buffer) in order to favour non-adsorption of antigens. The concentration of free phosphate ions in the composition of the invention is in one embodiment between 0.1 and 10.0 mM , or in another embodiment between 1 and 5 mM , or in a further embodiment about 2.5 mM .

10

Properties of the vaccines of the invention

In one embodiment the vaccines of the invention are formulated as a vaccine for *in vivo* administration to the host in such a way that the individual components of the composition are formulated such that the immunogenicity of individual components is not substantially impaired by other individual components of the composition. By not substantially impaired, it is meant that upon immunisation, an antibody titre against each component is obtained which is more than 60% , 70% , 80% or 90% , or $95-100\%$ of the titre obtained when the antigen is administered in isolation. Thus, in preferred embodiments, no (significantly) detrimental effect occurs to the further components (in terms of protective efficacy) in the combination as compared to their administration in isolation.

20

Vaccine formulations

In one embodiment, the vaccines of the invention are formulated as a vaccine for *in vivo* administration to the host, such that they confer an antibody titre superior to the criterion for seroprotection for each antigenic component for an acceptable percentage of human subjects. This is an important test in the assessment of a vaccine's efficacy throughout the population. Antigens with an associated antibody titre above which a host is considered to be seroconverted against the antigen are well known, and such titres are published by organisations such as WHO. In one embodiment, more than 80% of a statistically significant sample of subjects is seroconverted, in another embodiment more than 90% of a statistically significant sample of subjects is seroconverted, in a further embodiment more than 93% of a

30

statistically significant sample of subjects is seroconverted and in yet another embodiment 96-100% of a statistically significant sample of subjects is seroconverted.

The amount of antigen in each vaccine dose is selected as an amount which induces an immunoprotective response without significant, adverse side effects in typical vaccines. Such amount will vary depending on which specific immunogens are employed. Generally it is expected that each dose will comprise 1-1000 μ g of total immunogen, or 1-100 μ g, or 1-40 μ g, or 1-5 μ g. An optimal amount for a particular vaccine can be ascertained by studies involving observation of antibody titres and other responses in subjects. A primary vaccination course may include 2-3 doses of vaccine, given one to two months apart, e.g. following the WHO recommendations for DTP immunisation (i.e. in first year of life). Booster doses may follow in the second and/or subsequent year(s) of life.

Polio Potency as measured by seroneutralisation test on rats

For the purposes of the invention, the assay for the IPV quantitative evaluation of the vaccine potency of the IPV containing vaccines of the invention should be carried out using a single dose of vaccine and should be done by determining the ratio of test vaccine geometric mean titre (GMT) to reference vaccine GMT and is reported as the relative response (RR) or relative potency (RP). Reference GMT may be the GMT obtained with any IPV vaccine comprising 40-8-32 D-antigen units of IPV types 1-2-3 respectively, and may be the GMT obtained with the known vaccine Poliorix®. Typically, the RP test is carried out by as follows:

The potency of poliovirus Types 1, 2 and 3 is determined on rats by seroneutralisation:

Groups of 10 healthy rats (Sprague-Dawley (OFA) or any beforehand validated strain) are inoculated intramuscularly with dilutions (1/1.25; 1/3.125; 1/7.81) of the test samples or reference material in phosphate buffer saline. If necessary, the dilution range may be extended to 4 dilutions by inoculating undiluted vaccine and the three previous mentioned dilutions. Ten rats inoculated with the diluent are used as negative controls. Rats are observed once a week to detect any abnormal reaction. 20 to 22 days after the inoculation, each animal is deeply anesthetized, and bled and the serum is collected to be analysed by seroneutralisation test.

For the seroneutralisation test, sera are inactivated by incubation at 56°C for 30 minutes in a water bath. Three dilution series of the sera, one for each polio type, are prepared in microplates using the appropriate dilution medium. Plates are stored at + 4°C.

- 5 For the three polio virus types, a predetermined amount of virus (30-300 CCID₅₀) is added to the sera dilutions. The three virus suspensions are diluted taking into account their respective titers. The final dilution is called 'working dilution'. Each working dilution is added to the corresponding microplates. Plates are then sealed and incubated at 37°C ± 1°C for 16 hours. Hep-2 cells are then added and
- 10 microplates are incubated at 37°C ± 1°C for 7 days. The cytopathogenic effect (CPE) of the virus is read using an inverted microscope after Coomassie blue coloration. The presence of anti-poliomyelitis antibodies inhibits the growth of the virus and the appearance of the corresponding CPE. The anti-polio virus titers (type 1, 2 and 3) correspond to the reciprocal of the last dilution without any CPE. In each group,
- 15 animals with neutralising antibodies are recorded and the antibodies titer of each serum sample is determined for the different type of poliovirus. The neutralizing antibody titer is expressed as the log₂ of the inverse of the highest dilution of the serum sample that totally inhibits the cytopathic effect of poliovirus on Hep-2 cells.

- The geometric mean titer per dilution (GMT) and per virus type is also
- 20 determined for each group of rats.

Packaging of vaccines of the invention

- Vaccines of the invention can be packaged in various types of container e.g. in vials, in syringes, etc. A multidose vial will typically comprise a re-sealable plastic
- 25 port through which a sterile needle can be inserted to remove a dose of vaccine, which reseals once the needle has been removed.

- The vaccine may be supplied in various containers (e.g. 2 or 3). The contents of the containers may be mixed extemporaneously before administering to a host in a single injection or may be administered concomitantly at different sites. The dose of
- 30 the vaccine or each vaccine if a kit is administered concomitantly (in two or more containers) will typically be 0.5mL.

In one embodiment of this aspect of the invention there is provided a kit comprising two multi-valent vaccines for conferring protection in a host against

disease caused by poliovirus, *Bordetella pertussis*, *Clostridium tetani*, *Corynebacterium diphtheriae* and optionally one or more of Hepatitis B, *Haemophilus influenza* type B, *Neisseria meningitidis* type A, *Neisseria meningitidis* type C, *Neisseria meningitidis* type W, *Neisseria meningitidis* type Y, *Neisseria meningitidis* type B, *Salmonella typhi*, Hepatitis A or Malaria.

The kit comprises a first container comprising:

- (1) (a) Inactivated polio virus (IPV) of the invention,
- (b) diphtheria toxoid (DT or D) (see above),
- 10 (c) tetanus toxoid (TT or T) (see above),
- (d) killed whole-cell *Bordetella pertussis* (Pw) or 2 or more acellular pertussis components (Pa) (see above),
- (e) optionally Hepatitis B surface antigen (HepB or HB) (see above),
- (f) optionally a conjugate of a carrier protein and the capsular saccharide of *H. influenzae* type B (Hib) (see above),
- 15 (g) optionally either or both conjugates of a carrier protein and a capsular saccharide of a *N. meningitidis* type A (MenA) or *N. meningitidis* type C (MenC) (see above), and

a second container comprising:

- 20 (2A) (a) conjugates of a carrier protein and a capsular saccharide *N. meningitidis* type A (MenA), *N. meningitidis* type C (MenC), *N. meningitidis* type W (MenW) and/or *N. meningitidis* type Y (MenY) (see above for various Men saccharide combinations of the invention), and

- (b) optionally a conjugate of a carrier protein and the capsular saccharide of *H. influenzae* type B (Hib); or
- 25

- (2B) (a) a conjugate of a carrier protein and the capsular saccharide of *H. influenzae* type B (Hib), and

- (b) optionally a conjugate of a carrier protein and Vi saccharide of *Salmonella typhi*

- 30 The kit may optionally comprise a third container comprising:

- (3) (a) optionally Hepatitis B surface antigen

(b) optionally a conjugate of a carrier protein and Vi saccharide of *Salmonella typhi*

The containers may in either case additionally comprise HepA antigen(s) and/or MenB antigen(s) and/or RTS,S and/or *Streptococcus pneumonia* antigen(s).

5 In either case, the same antigen should not be present in both containers.

In one embodiment the first container has in addition to components a), b), c), d) also e), f), g), e)+f), e)+g), f)+g) or e)+f)+g).

10 In one embodiment the vaccine of the first container may be liquid and the vaccine of the second container may be either liquid or lyophilised (e.g. in the presence of a known stabilising excipient such as sucrose or trehalose).

The containers of the kit can be packaged separately or, optionally, packed together. In one embodiment, the kit is provided with a list of instructions for administration of the vaccines in the two or more containers.

15 In one embodiment, where a container in a kit contains a certain saccharide conjugate, the same conjugate is not present in the other containers of the kit.

The inventors believe that a kit provided in the above way may advantageously present the various antigens to a host's immune system in an optimal manner. The kit may provide a medical practitioner with an optimal method of immunising a host with one or more of the following advantages: protective efficacy
20 for all antigens, minimal reactogenicity, minimal carrier suppression interference, minimal adjuvant/antigen interference, or minimal antigen/antigen interference. In such a way, these goals may be achieved with the minimum number (two) administrations, optionally occurring at the same visit to the practitioner.

25 In one embodiment the vaccines of the first and second containers are administered concomitantly at different sites (as described below under "administration of vaccines of the invention), and in an alternative embodiment the inventors envision that the contents of the first and second containers may be mixed (optionally extemporaneously) before administration as a single vaccine.

30 **Preparing vaccines of the invention**

The present invention also provides a method for producing a vaccine formulation comprising the step of mixing the components of the vaccine together with a pharmaceutically acceptable excipient.

In one embodiment of the present invention there is provided a vaccine as herein described for use in a medicament for the treatment or prevention of diseases caused by infection by poliovirus and optionally *Bordetella pertussis*, *Clostridium tetani*, *Corynebacterium diphtheriae*, Hepatitis B virus, *Haemophilus influenzae*,
5 *Neisseria meningitidis* type A, *Neisseria meningitidis* type C, *Neisseria meningitidis* type W, *Neisseria meningitidis* type Y, *Salmonella typhi* or Hepatitis A.

In another embodiment of the invention there is provided a use of the vaccines of the invention in the manufacture of a medicament for the treatment or prevention of diseases caused by infection by poliovirus and optionally *Bordetella pertussis*,
10 *Clostridium tetani*, *Corynebacterium diphtheriae*, Hepatitis B virus, *Haemophilus influenzae*, *Neisseria meningitidis* type A, *Neisseria meningitidis* type C, *Neisseria meningitidis* type W, *Neisseria meningitidis* type Y, *Salmonella typhi* or Hepatitis A.

Additionally, a method of immunising a human host against disease caused by poliovirus and optionally *Bordetella pertussis*, *Clostridium tetani*, *Corynebacterium*
15 *diphtheriae*, Hepatitis B virus, *Haemophilus influenzae*, *Neisseria meningitidis* type A, *Neisseria meningitidis* type C, *Neisseria meningitidis* type W, *Neisseria meningitidis* type Y, *Salmonella typhi* or Hepatitis A, which method comprises administering to the host an immunoprotective dose of the vaccine of the invention is also provided.

20 The amount of antigen in each vaccine dose is selected as an amount which induces an immunoprotective response without significant, adverse side effects in typical vaccines. Such amount will vary depending upon which specific immunogen is employed and how it is presented. In one embodiment each dose will comprise 0.1-100 µg of saccharide, in another embodiment each dose will comprise 0.1-50 µg,
25 in a further embodiment each dose will comprise 0.1-10 µg, in yet another embodiment each dose will comprise 1 to 5 µg saccharide.

In one embodiment, the content of protein antigens in the vaccine will be in the range 1-100µg, in another embodiment the content of the protein antigens in the vaccines will be in the range 5-50µg, in a further embodiment the content of the
30 protein antigens in the vaccines will be in the range 5 - 25µg.

Vaccine preparation is generally described in Vaccine Design [“The subunit and adjuvant approach” (eds Powell M.F. & Newman M.J.) (1995) Plenum Press New York]. Encapsulation within liposomes is described by Fullerton, US Patent

4,235,877. Conjugation of proteins to macromolecules is disclosed, for example by Likhite, US Patent 4,372,945 and by Armor *et al.*, US Patent 4,474,757. Use of Quil A is disclosed by Dalsgaard *et al.*, 1977, Acta Vet Scand. 18:349. 3D-MPL is available from Ribic immunochem, USA and is disclosed in British Patent Application
5 No. 2220211 and US Patent 4,912,094. QS21 is disclosed in US Patent 5,057,540.

In a further embodiment of the invention there is provided a multi-valent vaccine comprising inactivated poliovirus (IPV) of the invention and optionally killed whole-cell *Bordetella pertussis* (Pw), tetanus toxoid (TT), diphtheria toxoid (DT), a conjugate of a carrier protein and the capsular saccharide of *H. influenzae* type B (Hib
10 – optionally conjugated to TT, DT or CRM197), wherein the amount of conjugate per 0.5 mL dose of bulk vaccine is 1-8 µg, and the immunogenicity of the conjugate is equivalent or improved over such compositions comprising larger amounts of conjugate. Optionally, Hepatitis B surface antigen may also be included.

In one embodiment the amount of conjugate per 0.5 mL dose of bulk vaccine
15 is less than 10 µg (of saccharide in the conjugate), in another embodiment the amount of conjugate is 1-7, in another embodiment the amount of conjugate is 2-6 µg, or in a further embodiment about 2.5, 3, 4 or 5 µg.

It will be appreciated that certain components, for example DTPw components, can be combined separately before adding the adsorbed HBsAg or other
20 components.

A method of making vaccines of the invention is also provided comprising the step of mixing IPV type 1, IPV type 2 and/or IPV type 3 with a pharmaceutically acceptable excipient. A typical process for preparing bulk vaccine of the invention with further antigens will add the IPV components to a mixture of the D and T
25 components, i.e. the DT components are mixed with the IPV components. This order of mixing allows the ionic strength and/or pH of the composition to be adjusted (e.g. pH<7) prior to the addition of the Pa or Pw components. Typically, HB pre-adsorbed onto AlPO₄ is added first if included in the composition, followed by the addition of DT pre-adsorbed onto Al(OH)₃ or AlPO₄, followed by the addition of TT pre-
30 adsorbed onto Al(OH)₃ or AlPO₄, followed by the addition of IPV optionally pre-adsorbed onto Al(OH)₃, prior to pH adjustment to e.g. pH5.9-7.2, or pH6-7, or pH6.2-6.8, or pH6.4-6.6, and then the addition of Pw pre-adsorbed onto AlPO₄. Optionally, Hib, Vi, MenA, MenC, MenW, Men Y, MenB and/or HepA antigens may be added at

any point in this process. In one embodiment, Hib, Vi, MenA, MenC, MenW, Men Y, MenB and/or HepA antigens are added prior to pH adjustment. In one embodiment one or more antigens of the invention are adsorbed onto aluminium phosphate or aluminium hydroxide or a mixture of both. In another embodiment the antigens of the invention are mixed with a pharmaceutically acceptable excipient and/or adjuvant(s).

In one embodiment, the vaccine composition of the invention may be prepared in the following order: preadsorbed HBsAg is added, followed by preadsorbed Diphtheria toxoid, followed by preadsorbed tetanus toxoid and IPV, the pH is then adjusted to approximately 6.5 prior to adding preadsorbed Pw.

In another embodiment, the vaccine composition of the invention may be prepared in the following order: preadsorbed tetanus toxoid is added, followed by IPV, followed by preadsorbed HBsAg, followed by preadsorbed Diphtheria toxoid, the pH is then adjusted to approximately 6.5 prior to adding preadsorbed Pw.

In general, the combined vaccine compositions according to any aspect of the invention can be prepared as follows: The IPV, DTPw, HepB, MenA, MenC, MenW, MenY, MenB, Vi, Hepatitis A or other components are pre-adsorbed onto a suitable adjuvant, especially aluminium hydroxide or aluminium phosphate or a mixture of both. After allowing time for complete and stable adsorption of the respective components, the different components are combined under appropriate conditions. The Hib, Vi, MenA, MenC, MenW and/or MenY conjugate(s) may or may not be adsorbed onto aluminium adjuvant salt before being mixed with the DTPw vaccine.

In one embodiment, vaccines of the invention are prepared at between 15°C and 30°C (e.g. between 19°C and 27°C, or at 23±4°C).

Administration of vaccines of the invention

The invention provides a method for raising an immune response in a mammal, comprising the step of administering an effective amount of a vaccine of the invention. The vaccines can be administered prophylactically (i.e. to prevent infection). The immune response is preferably protective and preferably involves antibodies. The method may raise a booster response.

Following an initial vaccination, subjects may receive one or several booster (subsequent) immunisations adequately spaced. Dosing treatment can be a single dose schedule or a multiple dose schedule. Multiple doses may be used in a primary

immunisation schedule and/or in a booster immunisation schedule. A primary dose schedule, which may be in the first year of life, may be followed by a booster dose schedule. Suitable timing between priming doses (e.g. between 4-16 weeks), and between priming and boosting can be routinely determined.

5 In one embodiment, the mammal is a human. Where the vaccine is for prophylactic use, the human is preferably a child (e.g. a toddler or infant) or a teenager; where the vaccine is for therapeutic use, the human is preferably an adult. A vaccine intended for children may also be administered to adults e.g. to assess safety, dosage, immunogenicity, etc.

10 The vaccine preparations of the present invention may be used to protect or treat a mammal susceptible to infection, by means of administering said vaccine directly to a patient. Direct delivery may be accomplished by parenteral administration (intramuscularly, intraperitoneally, intradermally, subcutaneously, intravenously, or to the interstitial space of a tissue); or by rectal, oral, vaginal,
15 topical, transdermal, intranasal, ocular, aural, pulmonary or other mucosal administration. In one embodiment, administration is by intramuscular injection to the thigh or the upper arm. Injection may be via a needle (e.g. a hypodermic needle), but needle free injection may alternatively be used. A typical intramuscular dose is 0.5mL.

20 Bacterial infections affect various areas of the body and so the compositions of the invention may be prepared in various forms. For example, the compositions may be prepared as injectables, either as liquid solutions or suspensions. The composition may be prepared for pulmonary administration e.g. as an inhaler, using a fine powder or spray. The composition may be prepared as a suppository or pessary. The
25 composition may be prepared for nasal, aural or ocular administration e.g. as spray, drops, gel or powder (see e.g. Almeida & Alpar, 1996, J Drug Targeting, 3:455; Bergquist *et al.*, 1998, APMIS, 106:800). Successful intranasal administration of DTP vaccines has been reported (Ryan *et al.*, 1999, Infect. Immun., 67:6270; Nagai *et al.*, 2001, Vaccine, 19:4824).

30 In one embodiment the vaccines of the first and second (and third where applicable) containers are administered concomitantly at different sites, and in an alternative embodiment the inventors envision that the contents of the first and second containers may be mixed (optionally extemporaneously) before administration as a single vaccine.

The invention may be used to elicit systemic and/or mucosal immunity.

One way of checking the efficacy of therapeutic treatment involves monitoring bacterial infection after administration of the composition of the invention. One way of checking efficacy of prophylactic treatment involves monitoring immune responses
5 against the antigens after administration of the composition. Immunogenicity of compositions of the invention can be determined by administering them to test subjects (e.g. children 12-16 months age, or animal models – WO 01/30390) and then determining standard immunological parameters. These immune responses will generally be determined around 4 weeks after administration of the composition, and
10 compared to values determined before administration of the composition. Rather than assessing actual protective efficacy in patients, standard animal and *in vitro* models and correlates of protection for assessing the efficacy of DTP vaccines are well known.

15 The terms “comprising”, “comprise” and “comprises” herein are intended by the inventors to be optionally substitutable with the terms “consisting of”, “consist of” and “consists of”, respectively, in every instance. This does not change the normal meaning of these terms, and is only intended to provide basis for the substitution, not to make them equivalent in meaning.

20

All cited references and publications are incorporated by reference herein.

EXAMPLES


Examples are provided solely for the purposes of illustration and are not intended to limit the scope of the invention.

5 Example 1: Tests on low dose IPV formulations

For all the formulations of the example 1, the antigens are adsorbed by addition of aluminium salt prior to formulation except IPV which is added without adsorption.

The tables below present the adsorption method for D, T, Pw and HBsAg.

10



FINAL COMPOSITION per dose	
Diphtheria	7.5 Lf (+/- 420 Lf/ml)
Al^{3+}	0.075 mg
NaCl	150 mM
pH	6.1 +/- 0.1
Volume	approximately 18 μl

Table 1. Method of productions for Diphtheria toxoid adsorption.

$\text{Al}(\text{OH})_3$
 SUPERFOS type
 |
 T (3.25Lf / 0.070 mg Al^{3+})
 |
 Stirring 15 min. up to 20 min. at R.T.
 |
 Adjust at pH 6.1 +/- 0.1
 |
 Stirring 15 min. up to 20 min. at R.T.
 |
 Check pH 6.1 +/- 0.1
 |
 Stirring 16 hr. up to 24 hr. at R.T.
 |
 NaCl 1500 mM. (ad. 150 mM)
 |
 Stirring 15 min. up to 45 min. at R.T.
 |
 Adjust at pH 6.1 +/- 0.1
 |
 Stirring 15 min. up to 20 min. at R.T.
 |
 Check pH 6.1 +/- 0.1
 |
 Storage minimum 14 days at +2°C / +8°C before formulation.

FINAL COMPOSITION per dose	
Tetanus	3.25 Lf (+/- 360 Lf/ml)
Al^{3+}	0.070 mg
NaCl	150 mM
pH	6.1 +/- 0.1
Volume	approximately 9 μl

Table 2. Method of productions for Tetanus toxoid adsorption.

5

10

AlPO_4
 0.17 mg Al^{3+} /d
 (Brenntag)
 |
 Adjust pH at pH6.5+/- 0.1
 |
 Stirring 15 min. up to 20 min. at room t°
 |
 Check pH 6.5 +/- 0.1
 |
 Pw
 20IOU/d
 |
 Stirring 15 min. up to 45 min. at Room t°
 |
 Measure pH
 |
 Storage at + 2°C - + 8°C


FINAL COMPOSITION per dose			
Antigens		Adjuvant	[Al^{3+}] (mg)
Pw	20 OU	AlPO_4	0.170 mg
Al^{3+}	0.170 mg	AlPO_4	
NaCl	150 mM		
pH	6.8		
Volume	approximately 65 μl		

Table 3. Method of productions for Pw adsorption

5

10

15



FINAL COMPOSITION per dose			
Antigens		Adjuvant	[Al ³⁺] (mg)
HBsAg	10 µg	AlPO ₄	0.200 mg
Al ³⁺	0.200 mg	AlPO ₄	
NaCl	150 mM		
pH	6.1+/-0.1		
Volume	approximately 50 µl		

Table 4. Method of productions for HBsAg adsorption

Several different formulations were tested:

- 5
 - A combination of Diphtheria toxoid, Tetanus toxoid, Pertussis whole cell and Hepatitis B surface antigen : DTP_{W_{SF}}-HB as a reference (DTP_{W_{SF}} means that it is a thiomersal free formulation), formulated with the production method 1 (table 5).
- 10
 - GlaxoSmithKline Biologicals S.A. product Poliorix® (IPV stand-alone not adsorbed) as unadsorbed reference at the standard dose, formulated with the production method 2 (table 5).

- A combination of Diphtheria toxoid, Tetanus toxoid, Pertussis whole cell, Hepatitis B surface antigen and Inactivated polio virus: DTP_{WSF}-HB-IPV with addition of the IPV before Pw, formulated with the production method 3 (table 5).
- A combination of Diphtheria toxoid, Tetanus toxoid, Pertussis whole cell, Hepatitis B surface antigen and Inactivated polio virus: DTP_{WSF}-HB-IPV with addition of the IPV just after T adsorbed. This addition method allows IPV adsorption onto Al(OH)₃. This vaccine is formulated with the production method 4 (table 5).

A placebo containing only aluminium salts, IPV and buffers of the others antigens. As IPV is the only antigen in this placebo, there is no competition for adsorption. Therefore, IPV is completely adsorbed. This vaccine is formulated with the production method 5 (table 5).

The vaccines formulated with production method 2, 3, 4 and 5 were produced with an IPV dose-range between 12.5% and 100% of the standard IPV dose of 40/8/32 IU/0.5mL.

Step	Method of production 1 : DTPw _{SF} -HB
1	Water for injection to reach a final dose volume of 0.5mL
2	Add NaCl 1.5M to reach a final concentration of 150mM
3	Add 115µg of Al ³⁺ as AlPO ₄
4	Add 10µg of HBsAg adsorbed
5	Add 7.5Lf of Diphtheria toxoid adsorbed
6	Add 3.25Lf of Tetanus toxoid adsorbed
7	Stirring
8	Adjust the pH at 6.5+/-0.1
9	Stirring
10	Add 20IOU Pw adsorbed
11	Stirring
12	Store at +2 to +8°C

Step	Method of production 2: IPV standalone		
1	Add IPV at a dose of		
	Type 1	Type 2	Type 3
	40 IU	8 IU	32 IU
	20 IU	4 IU	16 IU
	10 IU	2 IU	8 IU
	5 IU	1 IU	4 IU
2	Add M199 buffer to reach a final volume of 0.5mL		
9	Stirring		
10	Adjust the pH at 6.9+/-0.2		
14	Store at +2 to +8°C		

Step	Method of production 3: DTPw _{SF} -HB-IPV		
1	Water for injection to reach a final dose volume of 0.5mL		
2	Add NaCl 1.5M to reach a final concentration of 150mM		
3	Add 115µg of Al ³⁺ as AlPO ₄		
4	Add 10µg of HBsAg adsorbed		
5	Add 7.5Lf of Diphtheria toxoid adsorbed		
6	Add 3.25Lf of Tetanus toxoid adsorbed		
7	Stirring		
8	Add IPV at a dose of		
	Type 1	Type 2	Type 3
	40 IU	8 IU	32 IU
	20 IU	4 IU	16 IU
	10 IU	2 IU	8 IU
	5 IU	1 IU	4 IU
9	Stirring		
10	Adjust the pH at 6.5+/-0.1		
11	Stirring		
12	Add 20IOU Pw adsorbed		
13	Stirring		
14	Store at +2 to +8°C		

Step	Method of production 4 : DTPw _{SF} -HB-IPV		
1	Water for injection to reach a final dose volume of 0.5mL		
2	Add NaCl 1.5M to reach a final concentration of 150mM		
3	Add 3.25Lf of Tetanus toxoid adsorbed		
4	Add IPV at a dose of		
	Type 1	Type 2	Type 3
	40 IU	8 IU	32 IU
	20 IU	4 IU	16 IU
	10 IU	2 IU	8 IU
	5 IU	1 IU	4 IU
5	Stirring		
6	Add 115µg of Al ³⁺ as AlPO ₄		
7	Add 10µg of HBsAg adsorbed		
8	Add 7.5Lf of Diphtheria toxoid adsorbed		
9	Stirring		
10	Adjust the pH at 6.5+/-0.1		
11	Stirring		
12	Add 20IOU Pw adsorbed		
13	Stirring		
14	Store at +2 to +8°C		
Method of production 5: Placebo			
As per Method of production 3, however all antigens other than IPV have been omitted.			

Table 5. Method of productions per 0.5mL dose

For the method of production 1 formulation: HBsAg, D and T are adsorbed separately on AlPO₄, AlPO₄ and Al(OH)₃ respectively. The three antigens are sequentially added to a suspension containing water, NaCl and free AlPO₄. The mixture is stirred for 60-75 min. Then the pH is adjusted to 6.5 before addition of adsorbed Pw.

For the method of production 3 formulation, the three adsorbed antigens are sequentially added to a suspension containing water, NaCl and free AlPO₄. The mixture is stirred for 60-75 min before IPV addition. The pH is adjusted to 6.5 before addition of Pw antigens.

For the method of production 4 formulation, T antigen is adsorbed on Al(OH)₃. Preadsorbed T antigen is added to a suspension containing water and NaCl, followed by IPV types 1, 2 and 3. The mixture is stirred for 60-75 minutes before addition of free AlPO₄. Preadsorbed HBsAg is then added, followed by preadsorbed D antigen, and the mixture is then stirred for an additional 60-75 minutes. The pH is adjusted to 6.5 before addition of Pw antigens.

The method of production 3 was eventually selected due to ease of manufacture as this protocol only involved one stirring step. During the process of manufacturing the
5 vaccine, thiomersal is not used, and is not added to the final vaccine product.

VB62615

The table below presents the composition of the formulations for a 0.5mL dose.

Description	Al ³⁺ as AlPO ₄ for D ⁺ adsorption	Al ³⁺ as Al(OH) ₃ for D ⁺ adsorption	Al ³⁺ as AlPO ₄ for HBsAg adsorption	Al ³⁺ as AlPO ₄ for Pw adsorption	Free AlPO ₄	Free Al(OH) ₃	Diphtheria toxoid dose	Tetanus toxoid dose	Pertussis dose	HBsAg dose	IPV dose	
											% of standard dose	D-antigen units (*) (T1/T2/T3)
Method of production 1 DTPw _{SF} -HB	75µg	70µg	200µg	170µg	115 µg	NA	7.5Lf	3.25Lf	20IOU	10µg	NA	NA
Method of production 2 IPV	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	100%	40/8/32
											50%	20/4/16
											25%	10/2/8
											12.50%	5/1/4
Method of production 3 DTPw _{SF} -HB-IPV	75µg	70µg	200µg	170µg	115 µg	NA	7.5Lf	3.25Lf	20IOU	10µg	100%	40/8/32
											50%	20/4/16
											25%	10/2/8
											12.50%	5/1/4
Method of production 4 DTPw _{SF} -HB-IPV	75µg	70µg	200µg	170µg	115 µg	NA	7.5Lf	3.25Lf	20IOU	10µg	100%	40/8/32
											50%	20/4/16
											25%	10/2/8
											12.50%	5/1/4
Method of production 5 Placebo	NA	NA	NA	NA	560µg	70µg	NA	NA	NA	NA	100%	40/8/32
											50%	20/4/16
											25%	10/2/8
											12.50%	5/1/4

(*) The D-antigen content is the values targeted for the dilution of the concentrated inactivated polio bulk during the formulation

Table 6. Composition of formulations per 0.5mL dose

Determination of polio potency on rats by seroneutralisation

The potency of the vaccine was determined by a seroneutralisation test after intramuscular inoculation of rats (Sprague-Dawley (OFA) or any beforehand validated strain). Groups of 10 naïve healthy rats were inoculated intramuscularly (0.5 mL) with dilutions of the test samples, reference material in phosphate buffer saline, or diluent (phosphate buffer saline). The ten rats inoculated with the diluent were used as negative controls. Twenty to twenty-two days after the inoculation (immunisation period), each animal was deeply anesthetized prior to blood collection by cardiac puncture. Blood samples were centrifuged (at approximately 800 g), and serums were analysed.

Seroneutralisation test:

Sera were inactivated by incubation at 56°C for 30 minutes. Three dilution series of the sera, one for each polio type, were prepared in microplates using the appropriate dilution medium. For the three polio virus types, a predetermined amount of virus was added to the sera dilutions. The three virus suspensions were diluted taking into account their respective titers. The final dilution is called 'working dilution'. Each working dilution was added to the corresponding microplates. Plates were then sealed and incubated at 37°C $\pm$ 1°C for 16 hours. Hep-2 cells were then added and microplates were incubated at 37°C $\pm$ 1°C for 7 days. The cytopathogenic effect (CPE) of the virus was read using an inverted microscope after Coomassie blue colouration.

The presence of anti-poliomyelitis antibodies inhibits the growth of the virus and the appearance of the corresponding CPE. The anti-polio virus titers (type 1, 2 and 3) correspond to the reciprocal of the last dilution without any CPE.

In each group, animals with neutralising antibodies are recorded and the antibodies titer of each serum sample is determined for the different type of poliovirus. The neutralizing antibody titer is expressed as the log₂ of the inverse of the highest dilution of the serum sample that totally inhibits the cytopathic effect of poliovirus on Hep-2 cells. The geometric mean titer per dilution (GMT) and per virus type is then determined for each group of rats.

For each type of virus, the vaccine dilution and subsequently the amount of D-antigen which induced neutralising antibodies in 50 % of the rats (ED50) was also calculated by probit analysis. The ED50 was expressed in D-antigen units.

In order to quantify the potency relative to that of the reference vaccine (usually Poliorix®, but may be a DTPaHBIPV vaccine such as Pediarix®), the relative potency (RP) defined as the ratio of two equivalent dose responses in a multi-dose test was measured. In this approach, the potency of the test vaccine is calculated by parallel line assay as described in Finney, 1978 (Statistical Method in Biological Assay, Charles Griffin & Company Ltd, London, 1978).

Determination of the Potency Polio Type 1, 2 and 3 by ELISA

The determination of the potency Polio by ELISA is performed in one or two steps depending on whether measurement is being carried out on bulk unadsorbed IPV vs formulated vaccines respectively:

1. Desorption for final adsorbed vaccine (for measuring D-antigen units in formulated vaccines – not required for measurement in unadsorbed IPV antigen bulk);
2. ELISA test for the quantification of D-antigen content of desorbed and unadsorbed vaccine and/ or polio bulk

Desorption step

After centrifugation for 10 minutes of the adsorbed vaccine under test, three successive desorptions are performed, by adding a desorption phosphate buffer to the pellet, mixing and incubating at room temperature. The first and the second desorption periods are of 2 hours, the incubation period for the third extraction being one night at room temperature. The harvests from the three extractions are pooled and diluted with phosphate buffer solution (PBS) without Ca and Mg containing bovine serum albumine (BSA) and Tween 20.

The three poliovirus antigens are quantified by ELISA as described below.

D-antigen quantification by ELISA:

Microtiter plates are coated with specific rabbit anti-polio virus (type 1, 2 or 3) IgG, diluted with carbonate/bicarbonate buffer (pH 9.6), and incubated overnight at 4° C. After washing, the saturating solution (phosphate buffer saline w/o Ca and Mg + 1 % BSA) is added. Blanks (PBS) and serial dilutions of vaccine samples and in-house unadsorbed standard are added in duplicate. The in house trivalent standard preparation contains calibrated type 1, 2 and 3 antigens. The calibrator is the European Pharmacopoeia Biological reference (EPBRP).

For all following steps, the microtiter plates are incubated during 1h30 at 37°C and washed. Rabbit anti-polio virus (type 1, 2 or 3) IgG conjugated to peroxydase, diluted with phosphate buffer (w/o Ca and Mg + Tween 20) containing BSA, is added. The substrate solution, containing the tetramethylbenzidine dissolved in dimethyl sulfoxide (DMSO) and diluted in acetate buffer containing 0.003% H₂O₂, is added, followed by a 15-30 minutes incubation in the dark. The blocking solution, containing H₂SO₄, is then added. Within one hour, the optical density (O.D.) of each well is read using a photometer set at 450 nm with a reference at 620 nm.

The D-antigen concentration in test samples is calculated from the standard curve obtained by plotting the O.D. values against the standard antigen concentrations.

As a supplement to the Potency by ELISA, any unadsorbed IPV antigen may be detected by the completeness method:

Completeness of adsorption to adjuvant unbound Polio Type 1, 2 and 3 by Elisa

Two successive centrifugations are performed. The supernatant is then harvested and tested undiluted in duplicate on microplates by ELISA. Microtiter plates are coated with specific rabbit anti-polio virus (type 1, 2 or 3) IgG, diluted with carbonate/bicarbonate buffer (pH 9.6), and incubated overnight at 4° C. After washing, the saturating solution (phosphate buffer saline w/o Ca and Mg + 1 % BSA) is added. Blanks (PBS), supernatant and in-house unadsorbed standard are added in duplicate.

For all following steps, the microtiter plates are incubated during 1h30 at 37°C and washed. Rabbit anti-polio virus (type 1, 2 or 3) IgG conjugated to peroxydase, diluted

with phosphate buffer (w/o Ca and Mg + Tween 20) containing BSA, is added. The substrate solution, containing the tetramethylbenzidine dissolved in dimethyl sulfoxide (DMSO) and diluted in acetate buffer containing 0.003% H₂O₂, is added, followed by a 15-30 minutes incubation in the dark. The blocking solution, containing H₂SO₄, is then added. Within one hour, the optical density (O.D.) of each well is read using a photometer set at 450 nm with a reference at 620 nm.

The completeness is considered positive (antigen in the supernatant) if the mean OD of sample is higher than the mean OD values of blanks + 3 standard deviations and if the mean OD of sample is higher than 0.1.

In case of positive completeness, the antigen content is measured by ELISA method as described in the second step of the Potency Polio Type 1, 2 and 3 by ELISA.

Method of measuring International Opacity Unit (IOU)

Cell concentration (IOU) can be determined using either visual IRPO (International Reference Preparation of Opacity) standard solution or by absorbance measurement at 660 nm.

The opacity of Single Strain Suspension is then determined by applying the “assigned opacity” equation as follows:

$$AO = LO / KO \times CO;$$

where AO = assigned opacity, LO = live harvest opacity, KO = killed harvest opacity, and CO = concentrate opacity.

RESULTS

Determination of potency polio on rats by seroneutralisation at the standard 40:8:32 dose

Experiments were performed to determine the potency of IPV types 1, 2 and 3. Results are shown in Table 8 below (in the present document, 40:8:32 D-antigen units of IPV types 1, 2 and 3 respectively is equivalent to 100% IPV dose).

Description	Potency IPV (ED50 expressed in IU/dose)		
	Type 1	Type 2	Type 3
<i>Ref. Poliorix</i>	20.78	8.88	40.02
<i>Ref. DTPaHBIPV</i>	3.21	0.57	8.62
DTP_{WSF}-HB-IPV Method of production 3	<1.93	0.64	<2.57

Table 8. Potency of IPV types 1, 2 and 3 in three different vaccine formulations.

The DTP_{WSF}-HB-IPV formulation (100% IPV) presents IPV potencies better than the reference Poliorix and similar or better than the reference DTPaHBIPV.

Evaluation of IPV potency with reduced IPV dosages

The potency is measured by *in vitro* and *in vivo* methods described above.

The potency by Elisa of reduced dose IPV for both formulations of methods of production 3 and 4 was examined *in vitro* and compared with reference DTPaIPVHB as shown in Table 9. Two batches for each formulation were tested for method of production 3.

The percentage of recovery was calculated with regard to the antigen content taken from IPV bulk for each formulation (e.g. 40/8/32 for 100% IPV containing formulation; 20/8/16 for 50 % IPV containing formulation; 10/4/8 for 25 % IPV containing formulation; 5/2/4 for 12.5 IPV containing formulation)

VB62615

Sample	T1		T2		T3	
	Potency Polio (% Recovery)	Completeness (% Recovery)	Potency Polio (% Recovery)	Completeness (% Recovery)	Potency Polio (% Recovery)	Completeness (% Recovery)
DTPaIPVHB Reference	82%	NP	99%	NP	93%	NP
DTPwSF-HB-IPV 1 "Method of Production 3" 100%IPV	46%	47%	94%	<5%	24%	74%
DTPwSF-HB-IPV 2 "Method of Production 3" 100%IPV	80%	<5%	100%	<5.0%	81%	17%
DTPwSF-HB-IPV 1 "Method of Production 3" 50%IPV	48%	31%	98%	<5%	29%	64%
DTPwSF-HB-IPV 2 "Method of Production 3" 50%IPV	71%	<5%	99%	<5%	91%	>5%
DTPwSF-HB-IPV 1 "Method of Production 3" 25%IPV	54%	34%	115%	<5%	33%	71%
DTPwSF-HB-IPV 2 "Method of Production 3" 25%IPV	81%	<5%	115%	<5%	107%	<5%
DTPwSF-HB-IPV "Method of Production 3" 12.5%IPV	50%	24%	110%	<5%	28%	60%
DTPwSF-HB-IPV "Method of Production 4" 100% IPV	41%	48%	93%	<5%	23%	51%
DTPwSF-HB-IPV "Method of Production 4" 50% IPV	47%	37%	98%	<5%	28%	69%
DTPwSF-HB-IPV "Method of Production 4" 25% IPV	51%	28%	80%	<5%	33%	61%
DTPwSF-HB-IPV "Method of Production 4" 12.5% IPV	42%	22%	100%	<5%	40%	58%
Placebo "Method of production 5" 100% IPV	86%	<5%	98%	<5%	107%	<5%
Placebo "Method of production 5" 50% IPV	94%	<5%	110%	<5%	111%	<5%
Placebo "Method of production 5" 25% IPV	80%	<5%	100%	<5%	104%	<5%
Placebo "Method of production 5" 12.5% IPV	72%	<5%	100%	<5%	98%	<5%

Table 9

Table 9 show that the adsorption completeness is similar for all IPV doses. The Type 1 and Type 3 are strongly desorbed (17% - 74%) while the Type 2 stay well adsorbed. The three types are well adsorbed for the placebo formulation for all the IPV doses. The adsorption is similar as for DTPaIPVHB reference vaccine.

There is a variability IPV completeness because of the fact that the completeness quantification method is not validated neither for DTPwHB IPV formulations nor for lower IPV concentrations (<40/8/32 D-antigen Units/0.5 ml).

The relative potency (expressed in comparison with reference poliorix vaccine) of reduced dose IPV for both formulations of methods of production 3 and 4 was examined *in vivo* in comparison with reference formulations as shown in Figures 1 and 2. Two batches for each formulations were tested for method of production 3.

Figure 1 shows that the IPV potency of DTPwSF-HB-IPV with 100% IPV is slightly greater than the potency of IPV in DTPaHBIPV. The IPV potency of DTPwSF-HB-IPV 50% from the formulation of method of production 3 can be seen to be similar to DTPaHBIPV 100%. The IPV potency for DTPwSF-HB-IPV 25% of method of production 3 is slightly lower than for Poliorix®. It was also found that 12.5% of the IPV dose was not sufficient to obtain a good IPV potency.

Figure 2 shows that the IPV potency is similar for the formulation of method of production 3 and the formulation of method of production 4. It is also shown that there is a trend of better potency for the placebo than for DTPwSF-HB-IPV.

These data therefore confirm that a reduced dose of IPV is sufficient to obtain a good potency *in vivo*.

Example 2: Feasibility of using no thiomersal in vaccines of the invention

The Preservative Efficacy Test (PET) allows the demonstration of the antimicrobial activity of the tested vaccine. The test consists in:

- challenging the vaccine preparation, in its final container step, with a prescribed inoculum of suitable micro-organisms,
- storing the inoculated preparation at a prescribed temperature
- withdrawing samples from the container at specified intervals of time and counting the organisms in the taken samples.

The PET testing procedure is described in the European Pharmacopoeia (5.1.3) and in the USP (<51>). According to these guidelines, the antimicrobial activity is evaluated by comparing the reduction in the number of viable micro-organisms with the criteria mentioned in the following table (Table 7)

Microorganisms	Time	Criteria : log reduction			
		EP A	EP B	EP C	USP
<u>Bacteria</u>					
<i>Staphylococcus aureus</i>	6h	2			
<i>Escherichia coli</i>	d1	3	1	Ni*	
<i>Pseudomonas aeruginosa</i>	d7		3	Ni*	1
	d14			3	3
	d28	Nr*	Ni*	Ni*	Ni*
<u>Yeast and moulds</u>					
<i>Candida albicans</i>	d7	2			Ni*
<i>Aspergillus niger</i>	d14		1	Ni*	Ni*
	d28	Ni*	Ni*	Ni*	Ni*

Table 7. EP and USP Criteria

Nr* : not recovered

Ni* : not increased

Example 3: Effect of Hib component on the potency of IPV and stability of IPV over time

Relative potency of IPV was measured as described in Example 1 to determine the effects the Hib component may have on IPV potency and to evaluate the stability of IPV over time at different IPV doses. The vaccines investigated were DTPwHBIPV(40-8-32), DTPwHBIPV with reconstituted Hib and stored for 8 months, DTPwHBIPV(20-4-16), DTPwHBIPV(20-4-16) with reconstituted Hib and stored for 8 months, DTPwHBIPV(20-4-16) and stored for 8 months, DTPwHBIPV(10-2-8) and DTPwHBIPV(10-2-8) with reconstituted Hib and stored for 8 months. RP values were measure relative to DTPaIPVHB (Pediarix) (Figure 3a) or Poliorix (Figure 3b). It was found that the Hib component has no impact on IPV potency. The relative potency of IPV was found to be maintained at 8 months (Figure 3).

Example 4: Effect of AlPO₄/Al(OH)₃ ratio on the visual aspect, the adsorption of D and T and the potency of IPV

Formulations were performed with change of Aluminium composition.

The formulations DTPw_{SF}-HB-IPV usually contains 630µg Aluminium: 560µg Al³⁺ as AlPO₄, 70µg Al³⁺ as Al(OH)₃. Aluminium salts are used to adsorb D, T, Pw and HBsAg. 115µg Al³⁺ of free AlPO₄ is added during the formulation.

Formulations were performed with the following ratios of free Al³⁺:

	Al(OH) ₃ µg Al ³⁺	AlPO ₄ µg Al ³⁺
Lot 1	0	115
Lot 2	23	92
Lot 3	69	46
Lot 4	46	69
Lot 5	92	23
Lot 6	115	0

Table 10. AlPO₄/Al(OH)₃ ratio

Step	Method of production 3: DTPw _{SF} -HB-IPV
1	Water for injection to reach a final dose volume of 0.5mL
2	Add NaCl 1.5M to reach a final concentration of 150mM
3	Add 115µg of Al ³⁺ with at the different ratios Al(OH) ₃ /AlPO ₄
4	Add 10µg of HBsAg adsorbed
5	Add 7.5Lf of Diphtheria toxoid adsorbed
6	Add 3.25Lf of Tetanus toxoid adsorbed
7	Stirring

8	Add IPV at a dose of 40/8/32 IU
9	Stirring
10	Adjust the pH at 6.5+/-0.1
11	Stirring
12	Add 20IOU Pw adsorbed
13	Stirring
14	Store at +2 to +8 °C

Table 11. Method of production for DTPwHB-IPV

Visual aspect was observed and up to ratio 69/46, acceptable aggregation is obtained.

Formulations were performed with the same production method and a dose-range for IPV between 0 and 100% of the regular IPV dose.

The percentage of D and T toxoids adsorption was measured by ELISA. The stability of the adsorption was followed by a treatment of 7 days at 37°C. Results are presented in Table 12 and 14.

		RATIO Al(OH) ₃ /AlPO ₄					
		0/115		23/92		46/69	
		T0	7d37°C	T0	7d37°C	T0	7d37°C
Pw_{SF}	0%	<1%	25%	<1%	6%	/	
	25%	<1%	29%	<1%	15%	<1%	5%
	50%	3%	41%	<1%	26%	<1%	17%
	100%	4%	49%	<1%	23%	<1%	11%

Table 12. Percentage of D toxoid desorption in DTPwHB-IPV with IPV dose-range

		RATIO Al(OH) ₃ /AlPO ₄					
		0/115		23/92		46/69	
		T0	7d37°C	T0	7d37°C	T0	7d37°C
Pw_{SF}	0%	<1%	34%	<1	12%	/	
	25%	<1%	50%	<1	32%	<1%	12%
	50%	5%	61%	<1	51%	<1%	33%
	100%	8%	63%	<1	41%	<1%	29%

Table 13. Percentage of T toxoid desorption in DTPwHB-IPV with IPV dose-range

IPV adsorption was followed. The stability of the adsorption was followed by a treatment of 21 days at 25°C.

IPV		Estimation of Ag not adsorbed					
		RATIO Al(OH) ₃ /AlPO ₄					
		0/115		23/92		46/69	
Dose	Type	T0	21d25°C	T0	21d25°C	T0	21d25°C
0%	N/A	N/A	N/A	N/A	N/A	AGGREGATION	
25%	Type 1	~10-20%	~20-30%	~10-20%	~20-30%	<10%	~20-30%
	Type 2	<10%	<10%	<10%	<10%	<10%	<10%
	Type 3	>30%	>30%	~20-30%	>30%	<10%	>30%
50%	Type 1	>30%	>30%	~10-20%	>30%	~10-20%	>30%

	Type 2	~10-20%	~10-20%	<10%	<10%	<10%	<10%
	Type 3	>30%	>30%	~10-20%	>30%	~10-20%	>30%
100 %	Type 1	>30%	>30%	~10-20%	>30%	~10-20%	>30%
	Type 2	~10-20%	~10-20%	<10%	<10%	<10%	<10%
	Type 3	>30%	>30%	~20-30%	>30%	~20-30%	>30%

Table 14. Percentage of IPV desorption in DTPwHB-IPV with IPV dose-range

The increase of the $\text{Al}(\text{OH})_3$ content in the formulations allows an adsorption improvement for D, T and IPV.

The better adsorption ratio obtained was with the $\text{Al}(\text{OH})_3/\text{AlPO}_4$ ratio of 46/69.

At this ratio:

- The T and D adsorption is complete in T0. Desorption after an accelerated stability study of 7 days at 37 °C present <20% of desorption for D, <30% for T.
- Each IPV type is adsorbed. Desorption of the Type 3 occurs 21 days at 25 °C.

The formulations with the ratio 46/69 were tested in-vivo and compared with Tetravac, Poliorix and a DTPaIPV vaccine.

Sample	ED50		
	Type 1	Type 2	Type 3
DTPw-HB-IPV 100%/HIB Ratio 46/69	<1.93	<0.64	2.57
DTPw-HB-IPV 50%/HIB Ratio 46/69	<1.59	0.46	<1.67
DTPw-HB-IPV 25%/HIB Ratio 46/69	3.08	0.96	3.25
Tetravac	8.53	0.39	9.15
Poliorix	9.52	2.64	15.06
DTPaHBIPV	<5.18	<0.64	15.01

Table 15. In-vivo potencies results

There are no significant differences (ED50) between the DTPw-HB-IPV formulations. DTPaHBIPV, Tetravac and Poliorix give similar results, inferior to the

DTPw-HB-IPV formulations (except for the type 2 for which all the formulations are equivalent).

Example 5: Clinical evaluation of the investigational DTPw-HBV-IPV/Hib vaccine with reduced IPV dosages

A Phase II, feasibility study is planned to assess the immunogenicity, reactogenicity and safety of three different formulations of GSK Biologicals' investigational DTPw-HBV-IPV/Hib vaccine as compared to the commercial DTPw-HBV/Hib and IPV vaccines administered concomitantly.

- **Indication/populations:**

Primary immunization of healthy infants in the first week of life against diphtheria, tetanus, pertussis, hepatitis B, poliomyelitis and *Haemophilus influenzae* type b diseases.

Study groups:

DTPw-HBV-IPV(standard dose)/Hib vaccine

DTPw-HBV-IPV(49% of standard dose)/Hib vaccine

DTPw-HBV-IPV(26% of standard dose)/Hib vaccine

DTPw-HBV/Hib + IPV vaccines

- **Co-primary objectives:**

The co-primary objectives will be assessed in sequential manner: i.e. the second and third objectives will be assessed only if the preceding one has been met.

- ✓ To demonstrate the non-inferiority of the DTPw-HBV-IPV(standard dose)/Hib vaccine to the IPV vaccine co-administered with the DTPw-HBV/Hib vaccine in terms of antibody response to the three poliovirus types, one month after the primary vaccination course.

The objective of non-inferiority will be reached if the upper limit of the standardised asymptotic 95% CI on the difference between groups (DTPw-HBV/Hib + IPV minus DTPw-HBV-IPV(standard dose)/Hib) in terms of seroprotection rates for each of the three poliovirus types is $\leq 10\%$.

- ✓ To demonstrate the non-inferiority of the DTPw-HBV-IPV(49% of standard dose)/Hib vaccine to the IPV vaccine co-administered with the DTPw-HBV/Hib vaccine in terms of antibody response to the three poliovirus types, one month after the primary vaccination course.

The objective of non-inferiority will be reached if the upper limit of the standardised asymptotic 95% CI on the difference between groups (DTPw-HBV/Hib + IPV minus DTPw-HBV-IPV(49% of standard dose)/Hib) in terms of seroprotection rates for each of the three poliovirus types is $\leq 10\%$.

- ✓ To demonstrate the non-inferiority of the DTPw-HBV-IPV(26% of standard dose)/Hib vaccine to the IPV vaccine co-administered with the DTPw-HBV/Hib

vaccine in terms of antibody response to the three poliovirus types, one month after the primary vaccination course.

The objective of non-inferiority will be reached if the upper limit of the standardised asymptotic 95% CI on the difference between groups (DTPw-HBV/Hib + IPV minus DTPw-HBV-IPV(26% of standard dose)/Hib) in terms of seroprotection rates for each of the three poliovirus types is $\leq 10\%$.

- **Secondary objectives:**

Immunogenicity

To assess the immunogenicity the DTPw-HBV-IPV/Hib candidate vaccine in terms of response to all vaccine antigens in comparison with the DTPw-HBV/Hib and IPV vaccines co-administered.

Reactogenicity

To assess the reactogenicity and safety of the study vaccines, in terms of solicited symptoms, unsolicited symptoms and serious adverse events.

- **Vaccination schedule**

Three-dose primary vaccination schedule at 6, 10 and 14 weeks of age. All subjects receive a birth dose of Hepatitis B.

- **Country:**

Philippines

- **Blood sampling:**

Pre-and post-vaccination 3

- **Vaccine formulations:**

Vaccine	Formulation/dose	Presentation	Volume
GSK Biologicals' DTPw-HBV-IPV/Hib IPV component (standard dose) IPV component (49% of standard dose) IPV component (26% of standard dose)	Diphtheria toxoid: not less than 30 IU (7.5 Lf) Tetanus toxoid: not less than 60 IU (3.25 Lf) <i>Bordetella pertussis</i> , killed: not less than 4 IU (20 OU) r-DNA HBsAg: 10 µg Aluminium as salts: 0.66 mg Inactivated Poliovirus type 1: 40 D antigen units Inactivated Poliovirus type 2: 8 D antigen units Inactivated Poliovirus type 3: 32 D antigen units 49% of full standard dose IPV (40-8-32) 26% of full standard dose IPV (40-8-32)	Whitish liquid in monodose vials	0.5 ml of the reconstituted vaccine
	Conjugate of <i>Haemophilus influenzae</i> type b capsular polysaccharide: 2.5 µg and Tetanus toxoid: 5-10 µg Lactose: 12.6 mg Aluminium as salts: 30 µg	Freeze-dried pellet in monodose vials	
GSK Biologicals' DTPw-HBV/Hib (Zilbrix™ Hib)	Diphtheria toxoid: not less than 30 IU (7.5 Lf) Tetanus toxoid: not less than 60 IU (3.25 Lf) <i>Bordetella pertussis</i> , killed: not less than 4 IU (20 OU) r-DNA HBsAg: 10 µg Aluminium as salts: 0.66 mg Thiomersal: 8 µg	Whitish liquid in two-dose vials	1 ml of the reconstituted vaccine
	Conjugate of <i>Haemophilus influenzae</i> type b capsular polysaccharide: 2.5 µg and Tetanus toxoid: 5-10 µg Lactose: 12.6 mg Aluminium as salts: 30 µg	Freeze-dried pellet in two-dose vials	
GSK Biologicals' IPV (Poliorix™)	Inactivated Poliovirus type 1: 40 D antigen units Inactivated Poliovirus type 2: 8 D antigen units Inactivated Poliovirus type 3: 32 D antigen units 2-phenoxyethanol max 2.5 mg Polysorbate max 50 µg Formaldehyde max 100 µg Phosphate buffered saline Contains amino acids for injection q.s, ad 0.5 ml	Whitish liquid in monodose vials	0.5 ml

Table 16. Vaccine formulations


PRE-ADSORPTION OF THE ANTIGENS

The DTPw-HBV-IPV formulation combines diphtheria toxoid, tetanus toxoid, three *Bordetella pertussis* strains, the purified major surface antigen (HBsAg) of the Hepatitis B virus (HBV) and the inactivated polio virus (IPV). These antigens, except IPV, were first pre-adsorbed on aluminium salt before being mixed with aluminium salt, sodium chloride buffer and water for injection.


Adsorption of diphtheria toxoid

The diphtheria purified concentrate was adsorbed on aluminium phosphate in a ratio of 15 Lf Diphtheria toxoid / 0.15 mg Al₃⁺. The two components were stirred for 15 up to 45 minutes at room temperature. The pH was adjusted to pH 5.1 ± 0.1, followed by stirring for 15 up to 45 minutes. The mix was stored for one week at 37°C. After stirring of 15 up to 45 minutes at room temperature, the pH was adjusted to pH 6.1 ± 0.1. The adsorbed concentrate was stored at +2°C -+8°C for at least 7 days before final formulation of DTPw-HB-IPV vaccine. Figure 1 hereafter highlights the adsorption manufacturing process of the pre-adsorbed Diphtheria bulk.


Adsorption flowchart of diphtheria toxoid

**Adsorption of tetanus toxoid**

The purified tetanus concentrate was adsorbed on aluminium hydroxide in a ratio of 3.25 Lf / 0.07 mg Al³⁺. The two components were stirred for 15 up to 20 minutes. The pH was adjusted at pH 6.1 ± 0.1. The mix was stored under stirring for 16 up to 24 hours at room temperature. A sodium chloride solution of 1500 mM of nominal concentration was added (ad 150 mM). After stirring of 15 up to 45 minutes at room temperature, the pH was adjusted to 6.1 ± 0.1. The adsorbed concentrate was stored at +2°C -+8°C for at least 14 days before final formulation of DTPw-HBV-IPV vaccine.

Adsorption flowchart of tetanus toxoid**25 Adsorption of Hepatitis B antigen**

The sterile purified HBsAg bulk was mixed with a sterile suspension of aluminium phosphate in order to obtain a suspension which contains per 10 µg HBsAg, 0.2 mg Al³⁺(as aluminium phosphate), 150 mM NaCl in a final volume of about 50 µl.



Adsorption of Pw antigen

The AlPO_4 solution was transferred aseptically into a sterile vessel. The solution was stirred for 5 to 10 minutes and the pH was adjusted to 6.5 $\pm$ 0.1 with 1M HCl or 0.5M NaOH directly in the vessel. The solution was stirred for 15-20 minutes. The pH was checked (6.5 $\pm$ 0.1) and adjusted if necessary.

Before the adsorption, the pertussis pooled harvest (PPH) was mixed for a minimum of 15 minutes prior to use and then the PPH was added into the sterile vessel containing the AlPO_4 . The suspension was stirred for minimum 15 minutes at room temperature and could be stored overnight at room temperature. If the product was stored overnight at room temperature, it had to be resuspended for minimum 30 minutes before distribution. Samples were taken for testing.

The Pw adsorbed bulk was distributed into sterile glass bottles and stored at 2-8°C.

Flow chart of Pw adsorption

Transfer AlPO_4 in sterile stainless steel vessel



Adjust pH to 6.5 $\pm$ 0.1



Stirring 15-20 minutes at room temperature



Check pH and adjust if necessary (6.5 $\pm$ 0.1)



Add the PPH in the sterile stainless steel vessel



Stirring min. 15 minutes at room temperature



Distribution in glass bottle



Storage of the Pw adsorbed at 2-8 °C

DTPW-HBV-IPV FINAL FORMULATION


The process was done as follows:

- The sodium chloride solution and water were mixed for injections in order to achieve a final concentration of 150 mM NaCl.
- AlPO_4 was added in order to obtain a free Al^{3+} concentration of 0.115 mg/dose
- The adsorbed HEF, diphtheria and tetanus concentrates were added in order to obtain a final concentration of 10 μg of HBsAg, 7.5 Lf diphtheria toxoid and 3.25 Lf tetanus toxoid per 0.5 ml dose.
- IPV was added in order to obtain a final concentration of 40/8/32 or 19.6/3.9/15.7 or 10.4/2.1/8.3 UI/d.
- Stirring gently for 60 up to 120 minutes at room temperature.
- pH was adjusted at 6.5 $\pm$ 0.1
- Stirring for 15 up to 20 minutes at room temperature.
- pH was checked: 6.5 $\pm$ 0.1
- Adsorbed Pw concentrate was added in order to obtain a final concentration of 20 IOU per 0.5 ml dose

- Stirring for 15 to 45 minutes at room temperature.
- pH was measured
- The final bulk was stored between +2°C and +8°C until filling.

5

Formulation flowchart of DTPw-HBV vaccine



30

Example 6: Clinical evaluation of the investigational DTPa-HBV-IPV/Hib vaccine with reduced Hib and IPV dosages

A Phase II, exploratory study is planned to assess the immunogenicity, reactogenicity and safety of 4 different formulations of GSK Biologicals' investigational DTPa-HBV-IPV/Hib vaccine versus the commercial DTPa-HBV-IPV/ Hib vaccine and the commercial DTPw-HBV/Hib and IPV vaccines administered concomitantly.

- **Indication/populations:**
Primary immunization of healthy infants in the first week of life against diphtheria, tetanus, pertussis, hepatitis B, poliomyelitis and *Haemophilus influenzae* type b diseases.

- **Study groups:**

45

DTPa-HBV-IPV(49% of standard dose)/Hib 5µg vaccine
 DTPa-HBV-IPV(49% of standard dose)/Hib 2.5µg vaccine
 DTPa-HBV-IPV(26% of standard dose)/Hib 5µg vaccine

DTPa-HBV-IPV (26% of standard dose)/Hib 2.5µg vaccine

DTPa-HBV-IPV/Hib vaccine

DTPw-HBV/Hib + IPV vaccines

5 • **Primary objectives:**

To assess the immunogenicity of the DTPa-HBV-IPV/Hib candidate vaccines in terms of the response to the PRP and the three polio antigens (polio1, 2 and 3).

• **Secondary objectives:**

10 **Immunogenicity**

To assess the immunogenicity of all study vaccines in terms of response to all vaccine antigens.

Reactogenicity

15 To assess the reactogenicity and safety of the study vaccines, in terms of solicited symptoms, unsolicited symptoms and serious adverse events.

• **Vaccination schedule**

Three-dose primary vaccination schedule as of 6 weeks of age. All subjects receive a birth dose of Hepatitis B.

20

• **Country:**

TBC

• **Blood sampling:**

25 Pre-and post-vaccination 3

• **Vaccine formulations:**

The vaccine is constituted of two parts: a liquid part (DTPa-HB-IPV) and a freeze dried part (Hib).

30

D, T, PT, FHA, PRN and HBsAg are preliminary pre-adsorbed. Water and NaCl are blended with the different antigens. The mixture is stirred to homogenize and pH is adjusted. The final composition of the DTPa-HB-IPV part of the vaccine is presented in the table hereafter.

35

Component	Amount
D toxoid	25Lf
T toxoid	10Lf
PT	25µg
FHA	25µg
PRN	8µg
HBsAg	10µg
IPV type 1	40 or 19.6 or 10.4 IU
IPV type 2	8 or 3.9 or 2.1 IU
IPV type 3	32 or 15.7 or 8.3 IU
Al ³⁺	From 700 to 790µg

Table 17. Composition for one 0.5mL human dose of DTPa-HBV-IPV

- Hib is pre-adsorbed. The Hib pre-adsorbed is mixed with sucrose or lactose prior to freeze drying. The Hib amount will be 2.5 or 5 or 10µg per human dose. Aluminium content will be from 30 to 120µg Al^{3+} as AlPO_4 per human dose.
- 5

Claims

1. A method of making a vaccine (i.e. a combination vaccine) comprising
inactivated poliovirus type 1, comprising the steps of: mixing diphtheria
5 toxoid and tetanus toxoid, followed by the addition of inactivated poliovirus
type 1 at a dose greater than 10 D-antigen units and less than 20 D-antigen
units and optionally IPV type(s) 2 and/or 3 (IPV).
2. A method of making a vaccine (i.e. a combination vaccine) comprising
10 inactivated poliovirus type 1, comprising the steps of: mixing diphtheria
toxoid and tetanus toxoid, followed by the addition of inactivated poliovirus
type 1 at an amount greater than 10 D-antigen units and less than 20 D-antigen
units per 0.5mL and optionally IPV type(s) 2 and/or 3 (IPV).
- 15 3. The method of claim 1 or 2, wherein the vaccine comprises inactivated
poliovirus type 1 at 26-49%, 30-45%, 33-40%, or 35-37% of a standard 40 D-
antigen unit dose.
4. The method of claim 1 or 2, wherein the vaccine comprises inactivated
20 poliovirus type 1 at 26-49%, 30-45%, 33-40%, or 35-37% of a standard 40 D-
antigen unit dose per 0.5mL.
5. The method of claims 1-4, wherein the vaccine additionally comprises
inactivated poliovirus type 3 at a dose of 8-20 D-antigen units, 9-19 D-antigen
25 units, 10-18 D-antigen units, 11-17 D-antigen units, 12-16 D-antigen units, or
13-15 D-antigen units; for instance around or exactly 14 D-antigen units.
6. The method of claims 1-4, wherein the vaccine additionally comprises
inactivated poliovirus type 3 at an amount of 8-20 D-antigen units, 9-19 D-
30 antigen units, 10-18 D-antigen units, 11-17 D-antigen units, 12-16 D-antigen
units, or 13-15 D-antigen units per 0.5mL; for instance around or exactly 14
D-antigen units per 0.5mL.

7. The method of claim 5 or 6, wherein the vaccine comprises inactivated poliovirus type 3 at a dose of greater than 8 and less than 20 D-antigen units.
- 5 8. The method of claim 5 or 6, wherein the vaccine comprises inactivated poliovirus type 3 at an amount of greater than 8 and less than 20 D-antigen units per 0.5mL.
- 10 9. A method of making a vaccine (i.e. a combination vaccine) comprising inactivated poliovirus type 3, comprising the steps of: mixing diphtheria toxoid and tetanus toxoid, followed by the addition of inactivated poliovirus type 3 at a dose of greater than 8 and less than 20 D-antigen units, 9-19 D-antigen units, 10-18 D-antigen units, 11-17 D-antigen units, 12-16 D-antigen units, or 13-15 D-antigen units; for instance around or exactly 14 D-antigen units and optionally IPV type(s) 1 and/or 3 (IPV).
- 15 10. A method of making a vaccine (i.e. a combination vaccine) comprising inactivated poliovirus type 3, comprising the steps of: mixing diphtheria toxoid and tetanus toxoid, followed by the addition of inactivated poliovirus type 3 at an amount of greater than 8 and less than 20 D-antigen units, 9-19 D-antigen units, 10-18 D-antigen units, 11-17 D-antigen units, 12-16 D-antigen units, or 13-15 D-antigen units per 0.5mL; for instance around or exactly 14 D-antigen units per 0.5mL and optionally IPV type(s) 1 and/or 3 (IPV).
- 20 11. The method of claim 9 or 10, wherein the vaccine additionally comprises inactivated poliovirus type 1 at a dose of 10-20 D-antigen units, 11-19 D-antigen units, 12-18 D-antigen units, 13-17 D-antigen units, 14-16 D-antigen units; for instance around or exactly 15 D-antigen units.
- 25 12. The method of claim 9 or 10, wherein the vaccine additionally comprises inactivated poliovirus type 1 at an amount of 10-20 D-antigen units, 11-19 D-antigen units, 12-18 D-antigen units, 13-17 D-antigen units, 14-16 D-antigen units per 0.5mL; for instance around or exactly 15 D-antigen units per 0.5mL.
- 30

13. The method of claims 1-12, wherein the vaccine additionally comprises inactivated poliovirus type 2 at a dose of 2-4 D-antigen units.

5 14. The method of claims 1-12, wherein the vaccine additionally comprises inactivated poliovirus type 2 at an amount of 2-4 D-antigen units per 0.5mL.

10 15. The method of claim 13 or 14, wherein the vaccine comprises inactivated poliovirus type 2 at a dose greater than 2 D-antigen units and less than 4 D-antigen units, or at a dose around or exactly 3 D-antigen units.

15 16. The method of claim 13 or 14, wherein the vaccine comprises inactivated poliovirus type 2 at an amount greater than 2 D-antigen units and less than 4 D-antigen units per 0.5mL, or at an amount of around or exactly 3 D-antigen units per 0.5mL.

17. The method of claims 1-16, wherein the relative potency value of the vaccine is at least 0.85, 0.9, 0.95 or 1 when compared to a vaccine comprising 40 D-antigen units of IPV type 1.

20 18. The method of claims 1-17, wherein the relative potency value of the vaccine is at least 0.85, 0.9, 0.95 or 1 when compared to a vaccine comprising 8 D-antigen units of IPV type 2.

25 19. The method of claims 1-18, wherein the relative potency value of the vaccine is at least 0.85, 0.9, 0.95 or 1 when compared to a vaccine comprising 32 D-antigen units of IPV type 3.

30 20. The method of claims 1-19, comprising the subsequent step of adjusting pH to 6.5 ± 0.5 , or pH5.9-7.2, or pH6-7, or pH6.2-6.8, or pH6.4-6.6.

21. The method of claims 1-20, comprising the subsequent steps of adding a killed whole-cell or acellular pertussis component(s) and a pharmaceutically

acceptable excipient, optionally where free aluminium phosphate is present prior to the addition of the antigens.

5 22. The method of claims 1-21, comprising the additional step of adding Hepatitis B surface antigen prior to pH adjustment or prior to the addition of IPV.

10 23. The method of claims 1-22, comprising the additional step of mixing one or more antigens from a pathogen selected from a list consisting of: *Haemophilus influenzae* b, *Neisseria meningitidis* type A, *Neisseria meningitidis* type C, *Neisseria meningitidis* type W, *Neisseria meningitidis* type Y, *Neisseria meningitidis* type B, *Salmonella typhi* and Hepatitis A, with diphtheria toxoid, tetanus toxoid, inactivated poliovirus types 1, 2 and/or 3 and optionally Hepatitis B surface antigen prior to pH adjustment.

15 24. A method of making a DTP vaccine which is substantially thiomersal free and which comprises one or more IPV components selected from the group consisting of:

- 20 a. IPV type 1 at a dose of 10-36 D-antigen units, 11-32 D-antigen units, 12-28 D-antigen units, 13-24 D-antigen units, 14-20 D-antigen units or 15-19 D-antigen units, or around or exactly 18 D-antigen units;
- b. IPV type 2 at a dose of 2-7 D-antigen units, 3-6 D-antigen units, or 4-5 D-antigen units; and
- 25 c. IPV type 3 at a dose of 8-29 D-antigen units, 9-26 D-antigen units, 10-23 D-antigen units, 11-20 D-antigen units, 12-17 D-antigen units, or 13-14 D-antigen units,

comprising the steps of: mixing diphtheria toxoid and tetanus toxoid, followed by the addition of inactivated poliovirus component(s) .

30 25. The method of claim 24, wherein the vaccine comprises inactivated poliovirus type 1 at a dose of 10-36 D-antigen units, 11-32 D-antigen units, 12-28 D-antigen units, 13-24 D-antigen units, 14-20 D-antigen units or 15-19 D-antigen units, or around or exactly 18 D-antigen units.

26. The method of claim 24 or 25, wherein the vaccine comprises inactivated poliovirus type 2 at a dose of 2-7 D-antigen units, 3-6 D-antigen units, or 4-5 D-antigen units.
- 5 27. The method of claims 24-26, wherein the vaccine comprises inactivated poliovirus type 3 at a dose of 8-29 D-antigen units, 9-26 D-antigen units, 10-23 D-antigen units, 11-20 D-antigen units, 12-17 D-antigen units, or 13-14 D-antigen units.
- 10 28. The method of claims 1-27, wherein the vaccine comprises inactivated poliovirus type 1 adsorbed onto aluminium hydroxide, or aluminium phosphate, or a mixture of both (wherein adsorption takes place either before mixing with antigen other than IPV, or wherein adsorption takes place after mixing with antigen other than IPV).
- 15 29. The method of claim 28, wherein the IPV type 1 is added prior to the pH adjustment step of claim 20.
- 20 30. The method of claims 1-29, wherein the vaccine comprises inactivated poliovirus type 2 adsorbed onto aluminium hydroxide, or aluminium phosphate, or a mixture of both (wherein adsorption takes place either before mixing with antigen other than IPV, or wherein adsorption takes place after mixing with antigen other than IPV).
- 25 31. The method of claim 30, wherein the IPV type 2 is added prior to the pH adjustment step of claim 20.
- 30 32. The method of claims 1-31, wherein the vaccine comprises inactivated poliovirus type 3 adsorbed onto aluminium hydroxide, or aluminium phosphate, or a mixture of both (wherein adsorption takes place either before mixing with antigen other than IPV, or wherein adsorption takes place after mixing with antigen other than IPV).

33. The method of claim 32, wherein the IPV type 3 is added prior to the pH adjustment step of claim 20.
- 5 34. The method of claims 1-33, wherein the vaccine comprises IPV types 1, 2 and 3 adsorbed onto an aluminium salt (wherein adsorption takes place either before mixing with antigen other than IPV, or wherein adsorption takes place after mixing with antigen other than IPV).
- 10 35. The method of claims 1-34, wherein the vaccine comprises IPV types 1, 2 and 3 adsorbed onto aluminium hydroxide or aluminium phosphate or a mixture of both (wherein adsorption takes place either before mixing with antigen other than IPV, or wherein adsorption takes place after mixing with antigen other than IPV).
- 15 36. The method of claims 1-35, wherein the vaccine comprises diphtheria toxoid, optionally adsorbed onto aluminium hydroxide, or aluminium phosphate, or a mixture of both (wherein adsorption takes place either before or after mixing with other antigens).
- 20 37. The method of claim 36, wherein the diphtheria toxoid is added prior to the pH adjustment step of claim 20.
- 25 38. The method of claims 1-37, wherein the vaccine comprises tetanus toxoid, optionally adsorbed onto aluminium hydroxide, or aluminium phosphate, or a mixture of both (wherein adsorption takes place either before or after mixing with other antigens).
- 30 39. The method of claim 38, wherein the tetanus toxoid is added prior to the pH adjustment step of claim 20.
40. The method of claims 1-39, wherein the vaccine comprises killed whole cell *Bordetella pertussis* (Pw), optionally adsorbed onto aluminium hydroxide, or aluminium phosphate, or a mixture of both (wherein adsorption takes place either before or after mixing with other antigens).

41. The method of claim 40, wherein the Pw is added after the pH adjustment step of claim 20.
- 5 42. The method of claim 41, wherein the killed whole-cell *Bordetella pertussis* is adsorbed onto aluminium phosphate (wherein adsorption takes place either before or after mixing with other antigens).
- 10 43. The method of claim 42, wherein the Pw is added after the pH adjustment step of claim 20.
- 15 44. The method of claims 1-43, wherein the vaccine comprises two or more acellular pertussis components (Pa) (e.g. pertussis toxoid (PT), filamentous haemagglutinin (FHA) and pertactin (PRN)), optionally adsorbed onto aluminium hydroxide, or aluminium phosphate, or a mixture of both (wherein adsorption takes place either before or after mixing with antigens other than acellular pertussis antigens).
- 20 45. The method of claim 44, wherein the Pa is added after the pH adjustment step of claim 20.
- 25 46. The method of claim 45, wherein the two or more acellular pertussis components (Pa) are adsorbed onto aluminium phosphate (wherein adsorption takes place either before or after mixing with antigens other than acellular pertussis antigens).
- 30 47. The method of claim 46, wherein the Pa is added after the pH adjustment step of claim 20.
48. The method of claims 1-47, wherein the vaccine comprises Hepatitis B surface antigen, optionally adsorbed onto aluminium phosphate (wherein adsorption takes place either before or after mixing with other antigens).

49. The method of claim 48, wherein the Hepatitis B surface antigen is added prior to the pH adjustment step of claim 20.
50. The method of claims 1-49, wherein the vaccine comprises a conjugate of a carrier protein and the capsular saccharide of *Haemophilus influenzae* type B (Hib), optionally adsorbed onto aluminium phosphate or unadsorbed onto adjuvant (wherein adsorption takes place either before or after mixing with other antigens).
51. The method of claim 50, wherein the Hib is added prior to the pH adjustment step of claim 20.
52. The method of claims 1-51, wherein the vaccine comprises one or more conjugates of a carrier protein and a capsular saccharide of a bacterium selected from the group *Neisseria meningitidis* type A, *Neisseria meningitidis* type C, *Neisseria meningitidis* type W and *Neisseria meningitidis* type Y, optionally adsorbed onto aluminium hydroxide or aluminium phosphate or a mixture of both or unadsorbed onto adjuvant (wherein adsorption takes place either before or after mixing with antigens other than *Neisseria meningitidis* type A, C, W or Y capsular saccharide antigens).
53. The method of claim 52, wherein the one or more conjugates of a carrier protein and a capsular saccharide of a bacterium selected from the group *Neisseria meningitidis* type A, *Neisseria meningitidis* type C, *Neisseria meningitidis* type W and *Neisseria meningitidis* type Y are added prior to the pH adjustment step of claim 20.
54. The method of claims 1-53, wherein the vaccine comprises a *Neisseria meningitidis* type B (MenB) outer membrane vesicle or LOS or a conjugated MenB capsular saccharide, or derivative thereof, optionally adsorbed onto aluminium hydroxide or aluminium phosphate or a mixture of both or unadsorbed onto adjuvant (wherein adsorption takes place either before or after mixing with antigens other than MenB antigens).

55. The method of claim 55, wherein the MenB outer membrane vesicle or LOS or the conjugated MenB capsular saccharide is added prior to the pH adjustment step of claim 20.

5 56. The method of claims 1-55, wherein the vaccine comprises a Vi saccharide from *Salmonella typhi* conjugated to a carrier protein, optionally adsorbed onto aluminium hydroxide or aluminium phosphate or a mixture of both or unadsorbed onto adjuvant (wherein adsorption takes place either before or after mixing with other antigens).

10

57. The method of claim 56, wherein the Vi saccharide from *Salmonella typhi* conjugated to a carrier protein is added prior to the pH adjustment step of claim 20.

15

58. The method of claims 1-57, wherein the vaccine comprises an antigen from Hepatitis A, optionally adsorbed onto aluminium hydroxide or aluminium phosphate or a mixture of both (wherein adsorption takes place either before or after mixing with other antigens).

20

59. The method of claim 58, wherein the antigen from Hepatitis A is added prior to the pH adjustment step of claim 20.

60. The method of claims 1-59, wherein the vaccine comprises DT at an amount of 5-50, 7-30Lf or approximately or exactly 7.5Lf or 25Lf per 0.5mL dose.

25

61. The method of claims 1-59, wherein the vaccine comprises DT at an amount of less than 5Lf, or 1-4Lf or approximately or exactly 2Lf per 0.5mL dose.

62. The method of claims 1-61, wherein the vaccine comprises TT at an amount of 2.5-30Lf, 3-20 Lf, 5-15Lf or exactly or approximately 10Lf per 0.5mL dose.

30

63. The method of claims 1-62, wherein Pa, if present in the vaccine, comprises PT at an amount of 2-50 μ g, 5-40 μ g, 10-30 μ g or exactly or approximately 25 μ g per 0.5mL dose.
- 5 64. The method of claims 1-63, wherein the vaccine comprises Pa which comprises PT at an amount of exactly or approximately 2.5 or 8 μ g per 0.5mL dose.
- 10 65. The method of claims 1-64, wherein the vaccine comprises Pa which comprises FHA at an amount of 2-50 μ g, 5-40 μ g, 10-30 μ g or exactly or approximately 25 μ g per 0.5mL dose.
- 15 66. The method of claims 1-65, wherein the vaccine comprises Pa which comprises FHA at an amount of exactly or approximately 2.5 or 8 μ g per 0.5mL dose
- 20 67. The method of claims 1-66, wherein the vaccine comprises Pa which comprises PRN at an amount of 0.5-20 μ g, 0.8-15 μ g, 2-10 μ g or exactly or approximately 8 μ g per 0.5mL dose.
- 25 68. The method of claims 1-67, wherein the vaccine comprises Pa, which comprises PRN at an amount of exactly or around 0.8 or 2.5 μ g per 0.5mL.
69. The method of claims 1-68, wherein the vaccine comprises Pw at an amount of 5-50 IOU, 7-40 IOU, 9-35 IOU, 11-30 IOU, 13-25 IOU, 15-21 IOU or around or exactly 20 IOU per 0.5mL dose.
- 30 70. The method of claims 1-69, wherein the vaccine comprises HB at an amount of 5-20 μ g, 8-15 μ g or approximately or exactly 10 μ g per 0.5mL
71. The method of claims 1-70, wherein the vaccine comprises Hib at an amount of 5-20 μ g, 8-15 μ g or approximately or exactly 10 μ g per 0.5mL dose.

72. The method of claims 1-70, wherein the vaccine comprises Hib at an amount of 1-6 μ g, 2-4 μ g or around or exactly 2.5 μ g per 0.5mL dose.

5 73. The method of claims 1-72, comprising an additional step of adding an adjuvant.

74. The method of claims 1-73, wherein the vaccine comprises one or more adjuvants selected from the group consisting of: an aluminium salt such as aluminium hydroxide or aluminium phosphate, a salt of calcium, iron or zinc,
10 an insoluble suspension of acylated tyrosine or acylated sugars, cationically or anionically derivatised saccharides, polyphosphazenes, biodegradable microspheres, monophosphoryl lipid A (MPL), lipid A derivatives of reduced toxicity, 3-O-deacylated MPL, quil A, Saponin, QS21, tocol, AS-2, CpG oligonucleotides, bioadhesives, mucoadhesives, microparticles, liposomes,
15 polyoxyethylene ether formulations, polyoxyethylene ester formulations, muramyl peptides, imidazoquinolone compounds, interleukins, macrophage colony stimulating factor (M-CSF), tumour necrosis factor (TNF), granulocyte and macrophage colony stimulating factor (GM-CSF).

20 75. The method of claim 73, wherein the adjuvant is a preferential TH1 inducer.

76. The method of claims 1-75, wherein total aluminium content in the vaccine is 200-1000 μ g, 300-900 μ g, 400-800 μ g, 500-700 μ g or around or exactly 630 μ g Al³⁺ per 0.5mL dose.

25 77. The method of claim 76, wherein Al³⁺ content in the vaccine is all aluminium hydroxide or all aluminium phosphate.

78. The method of claim 76, wherein Al³⁺ content in the vaccine is a mixture of
30 aluminium hydroxide and aluminium phosphate in the following ratio: 1:8-8:1, 1:4-4:1, 3:8-8:3, 1:2-2:1 or 1:1 (w:w Al³⁺:Al³⁺) of aluminium phosphate: aluminium hydroxide.

79. The method of claim 76, wherein Al^{3+} content in the vaccine is a mixture of aluminium hydroxide and aluminium phosphate in the following ratio 12:1-4:1, 11:1-5:1, 10:1-6:1, 9:1-7:1 or 8:1 (w:w Al^{3+} : Al^{3+}) of aluminium phosphate: aluminium hydroxide.

5

80. The method of claims 1-79, comprising a step of adding free aluminium salt adjuvant to the vaccine to the amount of 0-300 μg , 50-250 μg , 75-200 μg , 100-150 μg or around or exactly 115 μg of Al^{3+} per 0.5mL dose.

10

81. The method of claim 80, wherein the free aluminium salt adjuvant is all $\text{Al}(\text{OH})_3$ or all AlPO_4 .

82. The method of claim 80, wherein the free aluminium salt adjuvant is a mixture of $\text{Al}(\text{OH})_3$ and AlPO_4 in the following ratio (w:w Al^{3+} : Al^{3+}): 1:1-1:6, 1:1.1-1:5, 1:1.2-1:4, 1:1.3-1:3, 1:1.4-1:2, e.g. 23/92 or 69/46.

15

83. The method of claim 80, wherein the free aluminium salt adjuvant is a mixture of $\text{Al}(\text{OH})_3$ and AlPO_4 in the following ratio (w:w Al^{3+} : Al^{3+}): 6:1-1:1, 5:1-1.1:1, 4:1-1.2:1, 3:1-1.3:1, 2:1-1.4:1, e.g. 46/69 or 92/23.

20

84. The method of claims 80-83, wherein the free aluminium salt adjuvant is added prior to the pH adjustment step of claim 20.

85. The method of any of claims 1-84, wherein IPV type 1, if present in the vaccine, is the Mahoney strain.

25

86. The method of any of claims 1-85, wherein IPV type 2, if present in the vaccine, is the MEF-1 strain.

30

87. The method of any of claims 1-86, wherein IPV type 3, if present in the vaccine, is the Saukett strain.