

MACROCYCLIC INHIBITORS OF *FLAVIVIRIDAE* VIRUSES

CROSS-REFERENCE TO RELATED APPLICATIONS

5 This application claims the benefit under 35 U.S.C. § 119(e) to U.S. Provisional Application Serial Number 61/657,553, filed on June 8, 2012, the entirety of which is incorporated herein by reference.

FIELD

10 The present application provides novel compounds inhibiting viruses, compositions containing such compounds, and therapeutic methods comprising the administration of such compounds.

BACKGROUND

15 RNA viruses comprising the *Flaviviridae* family include at least three distinguishable genera including *pestiviruses*, *flaviviruses*, and *hepaciviruses* (Calisher, *et al.*, J. Gen. Virol., 1993, 70, 37-43). While *pestiviruses* cause many economically important animal diseases such as bovine viral diarrhea virus (BVDV), classical swine fever virus (CSFV, hog cholera) and border disease of sheep (BDV), their importance in human disease is less well characterized (Moennig, V., *et al.*, Adv. Vir. Res. 1992, 48, 53-98). *Flaviviruses* are responsible for important
20 human diseases such as dengue fever and yellow fever while *hepaciviruses* cause hepatitis C virus infections in humans. Other important viral infections caused by the *Flaviviridae* family include West Nile virus (WNV) Japanese encephalitis virus (JEV), tick-borne encephalitis virus, Junjin virus, Murray Valley encephalitis, St Louis encephalitis, Omsk hemorrhagic fever virus and Zika virus.

25 The hepatitis C virus (HCV) is the leading cause of chronic liver disease worldwide (Boyer, N. *et al.* J Hepatol. 32:98-112, 2000) so a significant focus of current antiviral research is directed toward the development of improved methods of treatment of chronic HCV infections in humans (Di Besceglie, A.M. and Bacon, B. R., Scientific American, Oct.: 80-85, (1999); Gordon, C. P., *et al.*, J. Med. Chem. 2005, 48, 1-20; Maradpour, D., *et al.*, Nat. Rev. Micro.

2007, 5(6), 453-463). A number of HCV treatments are reviewed by Dymock et al. in *Antiviral Chemistry & Chemotherapy*, 11:2; 79-95 (2000). Virologic cures of patients with chronic HCV infection are difficult to achieve because of the prodigious amount of daily virus production in chronically infected patients and the high spontaneous mutability of HCV virus (Neumann, et al.,
5 *Science* 1998, 282, 103-7; Fukimoto, et al., *Hepatology*, 1996, 24, 1351-4; Domingo, et al., *Gene* 1985, 40, 1-8; Martell, et al., *J. Virol.* 1992, 66, 3225-9.

Currently, there are primarily two antiviral compounds, ribavirin, a nucleoside analog, and interferon-alpha (α) (IFN), that are used for the treatment of chronic HCV infections in humans. Ribavirin alone is not effective in reducing viral RNA levels, has significant toxicity,
10 and is known to induce anemia. The combination of IFN and ribavirin has been reported to be effective in the management of chronic hepatitis C (Scott, L. J., et al. *Drugs* 2002, 62, 507-556) but less than half the patients given this treatment show a persistent benefit. Therefore, there is a need to develop more effective anti-HCV therapies.

The macrocycle sanglifehrin and derivatives are immunomodulatory and bind peptidyl-
15 prolyl cis/trans isomerase (PPIase) cyclophilins in a unique manner (WO 97/02285; WO 98/07743; *J. Am. Chem. Soc* 2003, 125, 3849-3859; *J. Org. Chem.* 2000, 65, 9255-9260; *Angew. Chem. Int. Ed.* 1999, 38, 2443-2446). The cyclophilins are peptidyl-prolyl cis/trans isomerases (PPIase) that regulate protein folding *in vivo* and inhibit hepatitis C virus (Lin et al., WO2006/138507). However, none of the sanglifehrins or their derivatives has become available
20 for human anti-viral therapy. Therefore, there is a continuing need to develop macrocyclic sanglifehrins with anti-*Flaviviridae* virus activity and particularly anti-HCV activity.

SUMMARY

In one aspect, there is provided a compound of Formula I:

pharmaceutically acceptable salt, isotope, stereoisomer, mixture of stereoisomers, tautomer, ester or prodrug thereof to a mammal in need thereof. In one example of the embodiment, the treatment results in the reduction of the in viral load or clearance of viral RNA in a patient.

In yet another embodiment, a method for treating *Coronaviridae* viral infection is provided comprising administering a therapeutically effective amount of a compound of Formula I or a pharmaceutically acceptable salt, isotope, stereoisomer, mixture of stereoisomers, tautomer, ester or prodrug thereof to a mammal in need thereof. In one example of the embodiment, the treatment results in the reduction of the in viral load or clearance of viral RNA in a patient.

DETAILED DESCRIPTION

Definitions

Unless stated otherwise, the following terms and phrases as used herein are intended to have the following meanings:

“Alkanoyl” is RC(O)- ; “alkanoyloxy” is RC(O)O- ; and “alkanoylamino” is RC(O)NR'- ; where R is an alkyl group as defined herein, and R' is hydrogen or alkyl.

“Alkenyl” refers to a straight or branched hydrocarbyl group with at least one site of unsaturation, i.e., a carbon-carbon, sp^2 double bond. In some embodiments, alkenyl is a $\text{C}_2\text{-C}_{20}$ alkenyl group, a $\text{C}_2\text{-C}_{10}$ alkenyl group or a $\text{C}_2\text{-C}_6$ alkenyl group. Examples of alkenyl group include, but are not limited to, vinyl ($-\text{CH}=\text{CH}_2$), allyl ($-\text{CH}_2\text{CH}=\text{CH}_2$), cyclopentenyl ($-\text{C}_5\text{H}_7$), and 5-hexenyl ($-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}=\text{CH}_2$).

“Alkenylene” refers to an unsaturated, branched or straight chain or cyclic hydrocarbon radical having two monovalent radical centers derived by the removal of two hydrogen atoms from the same or two different carbon atoms of a parent alkene. For example, and alkenylene group can have 2 to 20 carbon atoms, 2 to 10 carbon atoms, or 2 to 6 carbon atoms. Typical alkenylene radicals include, but are not limited to, 1,2-ethenylene ($-\text{CH}=\text{CH}-$).

“Alkoxy” is RO- where R is alkyl, as defined herein. Non-limiting examples of alkoxy groups include methoxy, ethoxy and propoxy.

“Alkyl” refers to a straight or branched chain hydrocarbyl group. In an embodiment, alkyl has from 1 to 20 carbon atoms (i.e., $\text{C}_1\text{-C}_{20}$ alkyl). In some embodiments, alkyl is a $\text{C}_1\text{-C}_{10}$ alkyl

group or a C₁-C₆ alkyl group. Examples of alkyl groups include, but are not limited to, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *t*-butyl, pentyl, hexyl, heptyl, octyl, nonyl and decyl.

“Alkylene” refers to a saturated, branched or straight chain radical or cyclic hydrocarbon radical having two monovalent radical centers derived by the removal of two hydrogen atoms from the same or two different carbon atoms of a parent alkane. For example, an alkylene group can have 1 to 20 carbon atoms, 1 to 10 carbon atoms, or 1 to 6 carbon atoms. Examples of alkylene radicals include, but are not limited to, methylene (-CH₂-), ethylene (-CH₂CH₂-), propylene (-CH₂CH₂CH₂-) and butylene (-CH₂CH₂CH₂CH₂-).

“Alkynyl” refers to a hydrocarbon containing normal, secondary or tertiary carbon atoms with at least one site of unsaturation, i.e., a carbon-carbon, sp triple bond. For example, an alkynyl group can have 2 to 20 carbon atoms (i.e., C₂-C₂₀ alkynyl), 2 to 12 carbon atoms (i.e., C₂-C₁₂ alkyne,) or 2 to 6 carbon atoms (i.e., C₂-C₆ alkynyl). Examples of alkynyl groups include, but are not limited to, acetylenic (-C≡CH) and propargyl (-CH₂C≡CH).

“Alkylamino” refers to an amino group substituted with one or more alkyl groups. “Mono(alkyl)amino” or “(alkyl)amino” is RNH-, and “di(alkyl)amino” or “(alkyl)₂amino” is R₂N-, where each of the R groups is alkyl as defined herein and are the same or different. Examples of alkylamino groups include, but are not limited to, methylamino, ethylamino, propylamino, butylamino, dimethylamino, diethylamino and methylethylamino.



“Amino” refers to -NH₂.

“Alkynylene” refers to an unsaturated, branched or straight chain or cyclic hydrocarbon radical having two monovalent radical centers derived by the removal of two hydrogen atoms from the same or two different carbon atoms of a parent alkyne. For example, an alkynylene group can have 2 to 20 carbon atoms, 2 to 10 carbon atoms, or 2 to 6 carbon atoms. Typical alkynylene radicals include, but are not limited to, acetylene (-C≡C-), propargylene (-CH₂C≡C-), and 4-pentynylene (-CH₂CH₂CH₂C≡C-).

“Aryl” refers to any monocyclic or bicyclic carbon ring of up to 7 atoms in each ring, wherein at least one ring is aromatic, or an aromatic ring system of 5 to 14 carbons atoms which includes a carbocyclic aromatic group fused with a 5-or 6-membered cycloalkyl group. Examples of aryl groups include, but are not limited to, phenyl, naphthyl, tetrahydronaphthyl and indanyl.

“Arylalkyl” refers to an alkyl as defined herein substituted with an aryl radical.

"Arylene" refers to an aryl as defined above having two monovalent radical centers derived by the removal of two hydrogen atoms from two different carbon atoms of a parent aryl.

Typical arylene radicals include, but are not limited to, phenylene, e.g., , and naphthylene, e.g., .

5 "Arylalkylene" refers to an arylalkyl as defined above having two monovalent radical centers derived by the removal of one hydrogen atom from the aryl radical and the other hydrogen removed from the alkyl radical of the group.

"Cycloalkyl" refers to a hydrocarbonyl group containing at least one saturated or partially unsaturated ring structure, and attached via a ring carbon. Cycloalkyl groups include
10 hydrocarbon mono-, bi-, and poly-cyclic rings, whether fused, bridged, or spiro. In various embodiments, it refers to a saturated or a partially unsaturated C₃-C₁₂ cyclic moiety, examples of which include cyclopropyl, cyclobutyl, cyclopentyl, cyclopentenyl, cyclohexyl, cyclohexenyl, cycloheptyl and cyclooctyl.

"Cycloalkylalkyl" refers to an alkyl moiety substituted with a cycloalkyl group. Examples
15 of cycloalkylalkyl groups include cyclopropylmethyl, cyclobutylmethyl, cyclopentylethyl and cyclohexylmethyl.

"Cycloalkylene" refers to a cycloalkyl, as defined herein, having two monovalent radical centers derived by the removal of two hydrogen atoms from the same or two different carbon atoms of a parent cycloalkyl. Examples of cycloalkylene include, but are not limited to,
20 cyclopropylene, cyclobutylene, cyclopentylene and cyclohexylene.

"Dialkylaminoalkyl" refers to an alkyl moiety substituted with a dialkylamino group, wherein dialkylamino is as defined herein.

"Ester" means any ester of a compound in which any of the -COOH functions of the molecule is replaced by a -C(O)OR function, or in which any of the -OH functions of the
25 molecule are replaced with a -OC(O)R function, in which the R moiety of the ester is any carbon-containing group which forms a stable ester moiety, including but not limited to alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heterocyclyl, heterocyclylalkyl and substituted derivatives thereof.

"Halo" refers to chloro (-Cl), bromo (-Br), fluoro (-F) or iodo (-I).

“Haloalkoxy” refers to alkoxy, as defined herein, substituted with one or more halo radicals.

“Haloalkoxyalkyl” refers to an alkyl moiety substituted with a haloalkoxy group, as defined herein.

5 “Haloalkyl” refers to an alkyl group, in which one or more hydrogen atoms of the alkyl group is replaced with a halogen atom. Examples of haloalkyl groups include, but are not limited to, -CF₃, -CHF₂, -CFH₂ and -CH₂CF₃.

“Heterocycloalkyl” refers to a saturated or partially unsaturated monocyclic, bicyclic or tricyclic group of 2 to 14 ring-carbon atoms and, in addition to ring-carbon atoms, 1 to 4
10 heteroatoms selected from P, N, O and S. The heterocyclic group can be attached through a carbon atom or through a heteroatom, and when substituted, the substituent can be bonded to a carbon atom or a heteroatom. Examples of heterocyclyl include azetidiny, benzoimidazolyl, benzofuranyl, benzofurazanyl, benzopyrazolyl, benzotriazolyl, benzothiophenyl, benzoxazolyl, carbazolyl, carbolinyl, cinnolinyl, furanyl, imidazolyl, indolinyl, indolyl, indolaziny, indazolyl,
15 isobenzofuranyl, isoindolyl, isoquinolyl, isothiazolyl, isoxazolyl, naphthpyridinyl, oxadiazolyl, oxazolyl, oxazoline, isoxazoline, oxetanyl, pyranyl, pyrazinyl, pyrazolyl, pyridazinyl, pyridopyridinyl, pyridazinyl, pyridyl, pyrimidyl, pyrrolyl, quinazolinyl, quinolyl, quinoxalinyl, tetrahydropyranyl, tetrahydrothiopyranyl, tetrahydroisoquinolinyl, tetrazolyl, tetrazolopyridyl, thiadiazolyl, thiazolyl, thienyl, triazolyl, azetidiny, 1,4-dioxanyl, hexahydroazepinyl, piperazinyl,
20 piperidinyl, pyridin-2-onyl, pyrrolidinyl, morpholinyl, thiomorpholinyl, dihydrobenzoimidazolyl, dihydrobenzofuranyl, dihydrobenzothiophenyl, dihydrobenzoxazolyl, dihydrofuranyl, dihydroimidazolyl, dihydroindolyl, dihydroisooxazolyl, dihydroisothiazolyl, dihydrooxadiazolyl, dihydrooxazolyl, dihydropyrazinyl, dihydropyrazolyl, dihydropyridinyl, dihydropyrimidinyl, dihydropyrrolyl, dihydroquinolinyl, dihydrotetrazolyl, dihydrothiadiazolyl, dihydrothiazolyl,
25 dihydrothienyl, dihydrotriazolyl, dihydroazetidiny, methylenedioxybenzoyl, tetrahydrofuranyl, and tetrahydrothienyl and N-oxides thereof.


“Heterocycloalkylene” refers to a heterocycloalkyl, as defined above, having two monovalent radical centers derived by the removal of two hydrogen atoms from the same or two different carbon atoms of a parent heterocycloalkyl group.

30 “Heterocycloalkylalkylene” refers to an heterocycloalkyl as defined above having two monovalent radical centers derived by the removal of one hydrogen atom from the heterocycloalkyl radical and the other hydrogen removed from the alkyl radical of the group.

“Heteroaryl” refers to a monocyclic, bicyclic or tricyclic ring having up to 7 atoms in each ring, wherein at least one ring is aromatic and contains from 1 to 4 heteroatoms in the ring selected from the group consisting of N, O and S. Non-limiting examples of heteroaryl include pyridyl, thienyl, furanyl, pyrimidyl, imidazolyl, pyranyl, pyrazolyl, thiazolyl, thiadiazolyl, isothiazolyl, oxazolyl, isoxazolyl, pyrrolyl, pyridazinyl, pyrazinyl, quinoliny, isoquinoliny, benzofuranyl, dibenzofuranyl, dibenzothiophenyl, benzothienyl, indolyl, benzothiazolyl, benzooxazolyl, benzimidazolyl, isoindolyl, benzotriazolyl, purinyl, thianaphthenyl and pyrazinyl. Attachment of heteroaryl can occur via an aromatic ring, or, if heteroaryl is bicyclic or tricyclic and one of the rings is not aromatic or contains no heteroatoms, through a non-aromatic ring or a ring containing no heteroatoms. “Heteroaryl” is also understood to include the N-oxide derivative of any nitrogen containing heteroaryl.

“Heteroarylalkyl” refers to an alkyl group, as defined herein, in which a hydrogen atom has been replaced with a heteroaryl group.

“Heteroarylene” refers to a heteroaryl, as defined above, having two monovalent radical centers derived by the removal of two hydrogen atoms from the same or two different carbon atoms of a parent heteroaryl group. Non-limiting examples of heteroarylene groups are:



“Hydroxyalkoxy” refers to an alkoxy, as defined herein, substituted with a hydroxyl group (-OH). An example of hydroxyalkoxy is hydroxyethoxy.

“Hydroxyalkyl” refers to an alkyl group substituted with at least one hydroxy group. Examples of hydroxyalkyl groups include, but are not limited to, hydroxymethyl, hydroxyethyl, hydroxypropyl and hydroxybutyl.

“Prodrug” refers to any compound that when administered to a biological system generates the drug substance, i.e., active ingredient, as a result of spontaneous chemical reaction(s), enzyme catalyzed chemical reaction(s), photolysis, and/or metabolic chemical reaction(s). A prodrug is thus a covalently modified analog or latent form of a therapeutically

active compound. Non-limiting examples of prodrugs include ester moieties, quaternary ammonium moieties, glycol moieties, and the like.

The term "optionally substituted" refers to a moiety wherein all substituents are hydrogen or wherein one or more of the hydrogens of the moiety are replaced by non-hydrogen substituents. Multiple substitutions on the same atom are also permitted where chemically feasible (e.g., a dioxo substitution to provide $-S(O)_2-$, geminal substituents, spiro cycloalkyl or heterocycloalkyl rings, etc.). In some embodiments, "one or more" substituents is from one to three substituents.

Embraced herein, where applicable, are permissible isomers such as tautomers, racemates, enantiomers, diastereomers, atropisomers, configurational isomers of double bonds (E- and/or Z-), cis- and trans- configurations in ring substitution patterns, and isotopic variants.

"Pharmaceutically acceptable" means suitable for use in pharmaceutical preparations, generally considered as safe for such use, officially approved by a regulatory agency of a national or state government for such use, or being listed in the U. S. Pharmacopoeia or other generally recognized pharmacopoeia for use in animals, and more particularly in humans.

"Pharmaceutically acceptable carrier" refers to a diluent, adjuvant, excipient, or carrier, or other ingredient which is pharmaceutically acceptable and with which a compound of the invention is administered.

"Pharmaceutically acceptable salt" refers to a salt which may enhance desired pharmacological activity. Examples of pharmaceutically-acceptable salts include acid addition salts formed with inorganic or organic acids, metal salts and amine salts. Examples of acid addition salts formed with inorganic acids include salts with hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid and phosphoric acid. Examples of acid addition salts formed with organic acids such as acetic acid, propionic acid, hexanoic acid, heptanoic acid, cyclopentanepropionic acid, glycolic acid, pyruvic acid, lactic acid, malonic acid, succinic acid, malic acid, maleic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, o-(4-hydroxybenzoyl)-benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, 1,2-ethanedisulfonic acid, 2-hydroxyethane-sulfonic acid, benzenesulfonic acid, p-chlorobenzenesulfonic acid, 2-naphthalenesulfonic acid, p-toluenesulfonic acid, camphorsulfonic acid, 4-methyl-bicyclo[2.2.2]oct-2-ene-1-carboxylic acid, gluco-heptonic acid, 4,4'-methylenebis(3-hydroxy-2-naphthoic) acid, 3-phenylpropionic acid, trimethyl-acetic acid, tertiary butylacetic acid, lauryl sulfuric acid, gluconic acid, glutamic acid, hydroxy-naphthoic

acids, salicylic acid, stearic acid and muconic acid. Examples of metal salts include salts with sodium, potassium, calcium, magnesium, aluminum, iron, and zinc ions. Examples of amine salts include salts with ammonia and organic nitrogenous bases strong enough to form salts with carboxylic acids.

5 A compound of a given formula (e.g. the compound of Formula I, which also includes Formula I-a, I-b, I-c, I-d, I-e or II) is intended to encompass the compounds of the disclosure, and the pharmaceutically acceptable salts, stereoisomers, mixture of stereoisomers or tautomers of such compounds. Additionally, the compounds of the disclosure may possess one or more asymmetric centers, and can be produced as a racemic mixture or as individual
10 enantiomers or diastereoisomers. The number of stereoisomers present in any given compound of a given formula depends upon the number of asymmetric centers present (there are 2^n stereoisomers possible where n is the number of asymmetric centers). The individual stereoisomers may be obtained by resolving a racemic or non-racemic mixture of an intermediate at some appropriate stage of the synthesis or by resolution of the compound by
15 conventional means. The individual stereoisomers (including individual enantiomers and diastereoisomers) as well as racemic and non-racemic mixtures of stereoisomers are encompassed within the scope of the present disclosure, all of which are intended to be depicted by the structures of this specification unless otherwise specifically indicated.

 "Isomers" are different compounds that have the same molecular formula. Isomers
20 include stereoisomers, enantiomers and diastereomers.

 "Stereoisomers" are isomers that differ only in the way the atoms are arranged in space. Stereoisomers include enantiomers and diastereomers.

 "Enantiomers" are a pair of stereoisomers that are non-superimposable mirror images of each other. A 1:1 mixture of a pair of enantiomers is a "racemic" mixture. The term "(±)" is
25 used to designate a racemic mixture where appropriate.

 "Diastereoisomers" are stereoisomers that have at least two asymmetric atoms, but which are not mirror-images of each other.

 The absolute stereochemistry is specified according to the Cahn Ingold Prelog R S system. When the compound is a pure enantiomer the stereochemistry at each chiral carbon
30 may be specified by either R or S. Resolved compounds whose absolute configuration is unknown are designated (+) or (-) depending on the direction (dextro- or laevorotary) that they rotate the plane of polarized light at the wavelength of the sodium D line.

Some of the compounds exist as tautomeric isomers or "tautomers". Tautomeric isomers are in equilibrium with one another. For example, amide containing compounds may exist in equilibrium with imidic acid tautomers. Regardless of which tautomer is shown, and regardless of the nature of the equilibrium among tautomers, the compounds are understood by one of ordinary skill in the art to comprise both amide and imidic acid tautomers. Thus, the amide containing compounds are understood to include their imidic acid tautomers. Likewise, the imidic acid containing compounds are understood to include their amide tautomers.

Any formula or structure provided herein, including Formula I, I-a, I-b, I-c, I-d, I-e or II, is also intended to represent unlabeled forms as well as isotopically labeled forms of the compounds. An "isotope" may have structures depicted by the formulas given herein except that one or more atoms are replaced by an atom having a selected atomic mass or mass number. Examples of isotopes that can be incorporated into compounds of the disclosure include isotopes of hydrogen, carbon, nitrogen, oxygen, phosphorous, fluorine and chlorine, such as, but not limited to ^2H (deuterium, D), ^3H (tritium), ^{11}C , ^{13}C , ^{14}C , ^{15}N , ^{18}F , ^{31}P , ^{32}P , ^{35}S , ^{36}Cl and ^{125}I . Various isotopically labeled compounds of the present disclosure, for example those into which radioactive isotopes such as ^3H , ^{13}C and ^{14}C are incorporated. Such isotopically labelled compounds may be useful in metabolic studies, reaction kinetic studies, detection or imaging techniques, such as positron emission tomography (PET) or single-photon emission computed tomography (SPECT) including drug or substrate tissue distribution assays or in radioactive treatment of patients.

The disclosure also included compounds of Formula I, I-a, I-b, I-c, I-d, I-e or II in which from 1 to n hydrogens attached to a carbon atom is/are replaced by deuterium, in which n is the number of hydrogens in the molecule. Such compounds exhibit increased resistance to metabolism and are thus useful for increasing the half life of any compound of Formula I, I-a, I-b, I-c, I-d, I-e or II when administered to a mammal. See, for example, Foster, "Deuterium Isotope Effects in Studies of Drug Metabolism", Trends Pharmacol. Sci. 5(12):524-527 (1984). Such compounds are synthesized by means well known in the art, for example by employing starting materials in which one or more hydrogens have been replaced by deuterium.

Deuterium labelled or substituted therapeutic compounds of the disclosure may have improved DMPK (drug metabolism and pharmacokinetics) properties, relating to distribution, metabolism and excretion (ADME). Substitution with heavier isotopes such as deuterium may afford certain therapeutic advantages resulting from greater metabolic stability, for example increased in vivo half-life, reduced dosage requirements and/or an improvement in therapeutic

index. An ^{18}F labeled compound may be useful for PET or SPECT studies. Isotopically labeled compounds of this disclosure and prodrugs thereof can generally be prepared by carrying out the procedures disclosed in the schemes or in the examples and preparations described below by substituting a readily available isotopically labeled reagent for a non-isotopically labeled reagent. It is understood that deuterium in this context is regarded as a substituent in the compound of Formula I, I-a, I-b, I-c, I-d, I-e or II.


The concentration of such a heavier isotope, specifically deuterium, may be defined by an isotopic enrichment factor. In the compounds of this disclosure any atom not specifically designated as a particular isotope is meant to represent any stable isotope of that atom. Unless otherwise stated, when a position is designated specifically as "H" or "hydrogen", the position is understood to have hydrogen at its natural abundance isotopic composition. Accordingly, in the compounds of this disclosure any atom specifically designated as a deuterium (D) is meant to represent deuterium.

"Therapeutically-effective amount" refers to an amount of a compound that, when administered to a subject for treating a disease, is sufficient to effect treatment for the disease. "Therapeutically effective amount" can vary depending on the compound, the disease and its severity, the age, the weight, etc. of the subject to be treated.

The term "treating", and grammatical equivalents thereof, when used in the context of treating a disease, means slowing or stopping the progression of a disease, or ameliorating at least one symptom of a disease, more preferably ameliorating more than one symptom of a disease. For example, treatment of a hepatitis C virus infection can include reducing the HCV viral load in an HCV infected human being, and/or reducing the severity of jaundice present in an HCV infected human being.

Compounds

The present application provides a compound represented by Formula I:

**Formula I**

or a pharmaceutically acceptable salt, isotope, stereoisomer, mixture of stereoisomers, tautomer, ester or prodrug thereof, wherein:

- 5 A^1 is (C₂-C₅)alkylene, (C₂-C₅)alkenylene, (C₂-C₅)alkynylene, -O-(C₂-C₄)alkylene, -O-(C₂-C₄)alkenylene, arylene, aryl(C₁-C₂)alkylene, heterocycloalkylene or heterocycloalkyl(C₁-C₂)alkylene, wherein a sp³ carbon atom of A^1 is optionally substituted with one or more (C₁-C₄)alkyl;

A^2 is arylene or heteroarylene, wherein A^2 is optionally substituted with halo;

- 10 X^1 is -O-, -NH- or -N((C₁-C₄)alkyl)-;

R^{1a} and R^{1b} are independently H, (C₁-C₄)alkyl, (C₂-C₄)alkenyl or (C₂-C₄)alkynyl;

R^2 is H, (C₁-C₄)alkyl, (C₂-C₄)alkenyl or (C₂-C₄)alkynyl;

R^{3a} and R^{3b} are independently H or (C₁-C₈)alkyl;


R^{4a} and R^{4b} are independently H, -OH, (C₁-C₄)alkoxy, halo(C₁-C₄)alkoxy or (C₁-C₈)alkyl;

- 15 R^5 is H, (C₁-C₄)alkyl, (C₂-C₄)alkenyl or (C₂-C₄)alkynyl, or

R^5 forms a cyclic moiety along with -N((C₁-C₄)alkyl)- of X^1 or arylene of A^2 ; and

R^6 is H or (C₁-C₄)alkyl.

In certain embodiments, provided is a compound of Formula I:

**Formula I**

or a pharmaceutically acceptable salt, ester or prodrug thereof, wherein:

- 5 A^1 is (C_1-C_5) alkylene, (C_2-C_5) alkenylene, (C_2-C_5) alkynylene, $-O-(C_2-C_4)$ alkylene, $-O-(C_2-C_4)$ alkenylene, arylene, aryl (C_1-C_2) alkylene, heterocycloalkylene or heterocycloalkyl (C_1-C_2) alkylene, wherein a sp^3 carbon atom of A^1 is optionally substituted with one or more (C_1-C_4) alkyl;

A^2 is arylene or heteroarylene, wherein A^2 is optionally substituted with halo;

X^1 is $-O-$, $-NH-$ or $-N((C_1-C_4)alkyl)-$;

- 10 R^{1a} and R^{1b} are independently H, (C_1-C_4) alkyl, (C_2-C_4) alkenyl or (C_2-C_4) alkynyl;

R^2 is H, (C_1-C_4) alkyl, (C_2-C_4) alkenyl or (C_2-C_4) alkynyl;

R^{3a} and R^{3b} are independently H or (C_1-C_8) alkyl;

R^{4a} and R^{4b} are independently H, $-OH$, (C_1-C_4) alkoxy, halo (C_1-C_4) alkoxy or (C_1-C_8) alkyl;


R^5 is H, (C_1-C_4) alkyl, (C_2-C_4) alkenyl or (C_2-C_4) alkynyl, or

- 15 R^5 forms a cyclic moiety along with $-N((C_1-C_4)alkyl)-$ of X^1 or arylene of A^2 ; and

R^6 is H or (C_1-C_4) alkyl.

In one aspect of the embodiment, A^1 is ethenylene, propenylene, butenylene, ethylene, propylene, butylene, oxypropylene, oxypropenylene, pyrazolylenylene, phenylene or pyrimidinylene.


- 20 In another aspect of the embodiment, A^2 is heteroarylene such as isoquinolinylene, phenylene or halophenylene. In one aspect, A^2 is



In another aspect of the embodiment, X^1 is -O- or -NH-; R^{1a} is H; R^{1b} is methyl; R^2 is iso-propyl; R^5 is methyl and R^6 is H or methyl.

In another aspect of the embodiment, R^{3a} is H or methyl; R^{3b} is H; R^{4a} is H, -OH, methoxy, trifluoroethoxy; and R^{4b} is H.

- 5 In another aspect of the embodiment, A^2 is heteroaryl; A^1 is (C_2-C_5) alkylene, (C_2-C_5) alkenylene or (C_2-C_5) alkynylene, wherein A^1 is optionally substituted with one or more (C_1-C_4) alkyl; R^{3a} is H or (C_1-C_8) alkyl; and R^{4a} is H, -OH or (C_1-C_4) alkoxy. In another aspect of the embodiment, A^2 is heteroaryl; A^1 is (C_1-C_5) alkylene, (C_2-C_5) alkenylene or (C_2-C_5) alkynylene, wherein A^1 is optionally substituted with one or more (C_1-C_4) alkyl; R^{3a} is H or (C_1-C_8) alkyl; and R^{4a} is H, -OH or (C_1-C_4) alkoxy. Non-limiting examples of such compounds
- 10 include the following compounds and pharmaceutically acceptable salts thereof.




Formula (I-a)

Compound No.	X^1	A^1	R^{3a}	R^{4a}
a-1	-NH-	$-CH_2CH_2-$	methyl	methoxy
a-2	-NH-	$-CH=CH-$	methyl	methoxy
a-3	-O-	$-CH_2CH_2-$	methyl	methoxy
a-4	-O-	$-CH=CH-$	methyl	methoxy
a-5	-NH-	$-CH_2CH_2-$	methyl	-H
a-6	-NH-	$-CH=CH-$	methyl	-H
a-7	-O-	$-CH_2CH_2-$	methyl	-H
a-8	-O-	$-CH=CH-$	methyl	-H
a-9	-NH-	$-CH=CHCH_2-$	-H	-H
a-10	-NH-	$-CH_2CH_2CH_2-$	-H	-H
a-11	-O-	$-CH=CHCH_2-$	-H	-H
a-12	-O-	$-CH_2CH_2CH_2-$	-H	-H
a-13	-NH-	$-CH=CHCH_2CH_2-$	-H	-H
a-14	-NH-	$-CH_2CH_2CH_2CH_2-$	-H	-H
a-15	-O-	$-CH=CHCH_2CH_2-$	-H	-H
a-16	-O-	$-CH_2CH_2CH_2CH_2-$	-H	-H

wherein the left bond of A¹ linker is attached to A².

In another aspect of the embodiment, A² is arylene; and A¹ is (C₂-C₅)alkylene, (C₂-C₅)alkenylene, (C₂-C₅)alkynylene, -O-(C₂-C₅)alkylene or -O-(C₂-C₄)alkenylene, wherein A¹ is optionally substituted with one or more (C₁-C₄)alkyl; R^{3a} is H or (C₁-C₄)alkyl; and R^{4a} is H, -OH, (C₁-C₄)alkoxy or halo(C₁-C₄)alkoxy. In another aspect of the embodiment, A² is arylene; and A¹ is (C₁-C₅)alkylene, (C₂-C₅)alkenylene, (C₂-C₅)alkynylene, -O-(C₂-C₅)alkylene or -O-(C₂-C₄)alkenylene, wherein A¹ is optionally substituted with one or more (C₁-C₄)alkyl; R^{3a} is H or (C₁-C₄)alkyl; and R^{4a} is H, -OH, (C₁-C₄)alkoxy or halo(C₁-C₄)alkoxy. Non-limiting examples of such compounds include the following compounds and pharmaceutically acceptable salts thereof.




Formula (I-b)

No.	X ¹	A ¹	R ^{1b}	R ^{3a}	R ^{4a}	R ⁵
b-1	-NH-	-CH=CHCH ₂ CH ₂ -	-H	-H	-OH	methyl
b-2	-NH-	-CH ₂ CH ₂ CH ₂ CH ₂ -	-H	-H	-OH	methyl
b-3	-O-	-CH=CHCH ₂ CH ₂ -	-H	-H	-OH	methyl
b-4	-O-	-CH ₂ CH ₂ CH ₂ CH ₂ -	-H	-H	-OH	methyl
b-5	-NH-	-CH=CHCH ₂ CH ₂ -	-H	-H	-OCH ₂ CF ₃	methyl
b-6	-NH-	-CH ₂ CH ₂ CH ₂ CH ₂ -	-H	-H	-OCH ₂ CF ₃	methyl
b-7	-O-	-CH=CHCH ₂ CH ₂ -	-H	-H	-OCH ₂ CF ₃	methyl
b-8	-O-	-CH ₂ CH ₂ CH ₂ CH ₂ -	-H	-H	-OCH ₂ CF ₃	methyl
b-9	-NH-	-CH=CHCH ₂ C(CH ₃) ₂ -	-H	-H	-H	methyl
b-10	-O-	-CH=CHCH ₂ C(CH ₃) ₂ -	-H	-H	-H	methyl
b-11	-NH-	-OCH ₂ CH=CH-	-H	methyl	methoxy	isopropyl
b-12	-O-	-OCH ₂ CH=CH-	-H	methyl	methoxy	isopropyl
b-13	-NH-	-OCH ₂ CH=CH-	-H	methyl	methoxy	methyl
b-14	-O-	-OCH ₂ CH=CH-	-H	methyl	methoxy	methyl
b-15	-NH-	-CH=CHCH ₂ CH ₂ -	methyl	-H	-OH	methyl
b-16	-NH-	-CH ₂ CH ₂ CH ₂ CH ₂ -	methyl	-H	-OH	methyl

wherein the left bond of A¹ linker is attached to A².

Also included are the following compounds and pharmaceutically acceptable salts thereof.




Formula (I-c)

No.	X ¹	A ¹	R ^{3a}	R ^{4a}
c-1	-NH-	-CH=CHCH ₂ CH ₂ -	methyl	methoxy
c-2	-NH-	-CH ₂ CH ₂ CH ₂ CH ₂ -	methyl	methoxy
c-3	-O-	-CH=CHCH ₂ CH ₂ -	methyl	methoxy
c-4	-O-	-CH ₂ CH ₂ CH ₂ CH ₂ -	methyl	methoxy
c-5	-NH-	-CH=CHCH ₂ CH ₂ -	-H	-OH
c-6	-NH-	-CH ₂ CH ₂ CH ₂ CH ₂ -	-H	-OH
c-7	-O-	-CH=CHCH ₂ CH ₂ -	-H	-OH
c-8	-O-	-CH ₂ CH ₂ CH ₂ CH ₂ -	-H	-OH







wherein the left bond of A¹ linker is attached to A².

- 5 In another aspect of the embodiment, A² is phenylene; and A¹ is pyrazolyne, phenylene or pyrimidinylene. Non-limiting examples of such compounds include the following compounds and pharmaceutically acceptable salts thereof.




Formula (I-d)

Compound No.	X ¹	A ¹	R ^{3a}	R ^{4a}
d-1	-O-		-H	-H
d-2	-O-		-H	-H

d-3	-O-		-H	-H
d-4	-O-		-H	-H
d-5	-NH-		-H	-H
d-6	-NH-		-H	-H
d-7	-NH-		-H	-H
d-8	-NH-		-H	-H

wherein the left bond of A¹ linker is attached to A².

In another aspect of the embodiment, A² is halophenylene; and A¹ is -O-(C₂-C₅)alkylene or -O-(C₂-C₄)alkenylene. Non-limiting examples of such compounds include the following compounds and pharmaceutically acceptable salts thereof.






Formula (I-e)


5



Compound No.	X ¹	A ¹	R ^{3a}	R ^{4a}
e-1	-NH-	-OCH ₂ CH=CH-	methyl	-H
e-2	-O-	-OCH ₂ CH=CH-	methyl	-H
e-3	-NH-	-OCH ₂ CH ₂ CH ₂ -	methyl	-H
e-4	-O-	-OCH ₂ CH ₂ CH ₂ -	methyl	-H
e-5	-NH-	-OCH ₂ CH=CH-	methyl	methoxy
e-6	-O-	-OCH ₂ CH=CH-	methyl	methoxy
e-7	-NH-	-OCH ₂ CH ₂ CH ₂ -	methyl	methoxy
e-8	-O-	-OCH ₂ CH ₂ CH ₂ -	methyl	methoxy


wherein the left bond of A¹ linker is attached to A².

In one aspect of the embodiment, A^1 is (C_2-C_5) alkylene or (C_2-C_5) alkenylene; R^5 is

methyl, or R^5 form  along with arylene of A^2 , or R^5 form  or  along with $-N((C_1-C_4)alkyl)-$ of X^1 ; and R^6 is H or methyl. In another aspect of the embodiment,

A^1 is (C_1-C_5) alkylene or (C_2-C_5) alkenylene; R^5 is methyl, or R^5 form  along with


5 arylene of A^2 , or R^5 form  or  along with $-N((C_1-C_4)alkyl)-$ of X^1 ; and R^6 is H or methyl. Non-limiting examples of such compounds include the following compounds and pharmaceutically acceptable salts thereof:



10



and

In another embodiment, there is provided a compound of Formula II:



Formula II

- or a pharmaceutically acceptable salt, isotope, stereoisomer, mixture of stereoisomers,
5 tautomer, ester or prodrug thereof, wherein:



X¹ is -O- or -NH-;

R^{3a} is H or (C₁-C₄)alkyl;

R^{4a} is H, -OH, (C₁-C₄)alkoxy, halo(C₁-C₄)alkoxy or (C₁-C₈)alkyl; and

R⁵ is H or (C₁-C₄)alkyl.

- 15 One skilled in the art will recognize that substituents and other moieties of the compounds of the generic formulae herein should be selected in order to provide a compound which is sufficiently stable to provide a pharmaceutically useful compound which can be formulated into an acceptably stable pharmaceutical composition. Compounds which have such stability are contemplated as falling within the scope of the present invention. It should be


understood by one skilled in the art that any combination of the definitions and substituents described above should not result in an inoperable species or compound.

Preparation of Macrocyclic Compounds

- 5 A compound of the present invention such as those of Formula I, I-a, I-b, I-c, I-d, I-e or II can be prepared according to the schemes described below, but it shall be appreciated that modifications of the illustrated process or other process can also be used. As illustrated in Scheme 1, the macrocyclic compounds **M** are synthesized from the five key components **A-E** by combining them together in sequence with the appropriate use of protecting groups (PG¹-PG⁸) by one skilled in the art. The hashed lines numbered 1-5, hereby referred to as Connection 1, Connection 2, etc., respectively, are the 5 connections for combining Components **A-E**. The order in which the specific connections occur, can vary, and are dependent on the choice of protecting groups and chemistry required. In certain embodiments, Connections 3, 4 or 5 are used as the final macrocyclization step.

15

Scheme 1



PG^x = protecting group as needed

Y¹ and Y² are H, alkyl, halide, triflate, boronic acid, stannane, etc.

Illustratively, Connections 1 through 5 can be performed as described below:

- Connections 1, 2 and 3 are amide bonds. The connections are made between the respective acid and amine using standard peptide coupling agents (EDC/HOBT, DCC, PyBOP, PyBROP, HATU, HBTU, COMU, etc.) known to one skilled in the art. The acid and amine

coupling partners are combined with a coupling agent in an organic solvent, e.g., DMF, dichloromethane, acetonitrile, etc., in the presence of a base, e.g., DIPEA, triethylamine, etc., at RT or slightly elevated temperature. When any of these three steps are chosen as the final macrocyclization step, for example Connection 3, then macrolactamization conditions are preferred. Suitable macrolactamization procedures include, but are not limited to, those found in the following reference: Davies, J. S. J. *Peptide Sci.* **2003**, 9, 471-501.

Connection 4 is typically a carbon-carbon bond or a heteroatom-carbon bond where the heteroatom is O, S or N. When Connection 4 is a carbon-carbon bond, then standard carbon-carbon bond forming procedures typically involving metal mediated cross coupling reactions are preferred. Preferably the carbon-carbon bond is formed using a Heck type coupling between an *sp*² halide group and an terminal alkene, a Suzuki coupling between an *sp*² halide group and a vinyl or aryl boronate, or a ring closing metathesis (RCM) between two alkenes. Stille reactions can also be performed between a vinyl stannane and an aryl or vinyl halide as described in *Journal of American Chemical Society* 2000, 122, 3830 Nicolaou et al. In each of the examples above the aryl or vinyl halide group can also be an aryl or vinyl triflate.

For example, when A²-Y² in **A** contains a terminal alkene, such as -CH=CH₂, and A¹-Y¹ in **B** contains a terminal alkene, or CH₃-CH=CH- then a cross metathesis reaction is performed. The two components are mixed in solvent, e.g., acetonitrile, toluene and a metathesis catalyst, e.g., Grubbs I, Grubbs II or Hoyveda-Grubbs I, Hoyveda-Grubbs II is added followed by heating. If this connection is the final procedure to close the macrocyclic ring, RCM conditions are preferred (e.g., more dilute conditions to avoid dimerization). For relevant RCM conditions and examples see *Journal of American Chemical Society* 2003, 125, 3849 Sedrani et al and *Journal of American Chemical Society* 2000, 122, 3830 Nicolaou et al. A typical RCM procedure includes heating (either conventionally or by microwave) of the acyclic precursor in a solvent such as toluene, or 1,2-dichloroethane, in the presence of a RCM catalyst, e.g., Grubbs I, Grubbs II or Hoyveda-Grubbs I, Hoyveda-Grubbs II.

Alternatively, when Connection 4 is made via a Heck coupling reaction, the vinyl or aryl halide, or the triflate **A** and the alkene component **B** are mixed in a polar solvent, e.g., acetonitrile, or toluene in the presence of a Palladium(II) catalyst, e.g., Palladium(OAc)₂, a phosphine ligand, e.g., P(*o*-toluene)₃, P(*t*-butyl)₃, etc., and a base, e.g., triethylamine. The reaction mixture is heated either conventionally or in a microwave reactor.

Alternatively, when Connection 4 is made via a Suzuki coupling reaction, the vinyl or aryl halide, or the triflate **A** and the vinyl or aryl boronate **B** are mixed in a suitable solvent, e.g.,

cyclopentyl methyl ether, toluene, DMF, DME, etc., in the presence of a Palladium catalyst (e.g., Palladium(II)Cl₂(p-NMe₂Ph)₂ and K₃PO₄ or tetrakis(triphenylphosphine)palladium(0) and a base, such as potassium carbonate). The reaction mixture is heated either conventionally or in a microwave reactor. It is also possible in such a coupling reaction to reverse the reactive functionalities on the two starting materials, such that **A** is an aryl or vinyl boronate and **B** contains a vinyl or aryl halide or triflate.


Alternatively, Connection 4 can be a carbon-oxygen bond and in this case typical alkylation or nucleophilic aromatic substitution conditions can be used between a hydroxyl group and an alkyl halide, or aryl (or heteroaryl) halide. The hydroxyl reagent is mixed with the alkyl or heteroaryl halide (preferably an iodide or bromide), in an inert solvent, e.g., CPME, DMF, THF, etc, in the presence base, e.g., cesium carbonate, cesium hydroxide, sodium hydride, MaHMDS, etc.; and heated.

Alternatively, Connection 4 can be a carbon-nitrogen bond and in this case typical alkylation, nucleophilic aromatic substitution or Buchwald conditions can be used between an amine group and an alkyl halide or heteroaryl halide. For example, the amine and the alkyl or heteroaryl halide are mixed and heated in an inert solvent, e.g., CPME, in the presence base, e.g., cesium carbonate, sodium hydride, etc. An alternative procedure for the carbon-nitrogen connection is to perform a reductive amination between an amine and a carbonyl compound. Typically the amine and aldehyde or ketone are mixed in an inert solvent, e.g., THF, dioxane and treated after a period of time with sodium acetoxy borohydride or alternative reducing agent. Connection 5 is typically an amide (X¹ = -NH or substituted N), or ester (X¹ = O) bond. When forming the amide bond, standard coupling procedures as described for Connections 1-3 can be used. In some embodiments, this is the final step in closing the macrocycle. As such, macrolactamization types of coupling procedures are more effective. Suitable macrolactamization procedures include, but are not limited to, those found in Davies, J. S. *J. Peptide Sci.* **2003**, 9, 471-501.

When forming the ester bond, standard coupling reagents (e.g., EDC, DCC, PyBOP, HATU, COMU) can be used, or when this is the final step in formation of the macrocycle, macrolactonization procedures are preferred (e.g., Shina, Yamaguchi). An exemplary method for the macrolactonization step can be found in *Journal of American Chemical Society* 2002, 124, 4257 Paquette et al or *Chemical Reviews* **2006**, 106(3), 911-939. Typically, the acid and alcohol are mixed in a polar solvent, e.g., DMF, acetonitrile, etc, in the presence of the coupling agent and a base, e.g., DIPEA, DMAP.

The following general schemes provide general examples and sequences for constructing the macrocyclic compound **M** from the common precursors **A-E**.

Scheme 2: Use of RCM method for Connection 4




Compounds **A-E** can be deprotected (PG²-PG⁸) using conditions described in Greene and Wuts, Protective Groups in Organic Synthesis, John Wiley and Sons, Inc. to provide Compounds **1a-1f**.

5 In many cases the optimal protecting groups and their deprotection methods are as follows. For Compound **E**, the typical protecting group PG¹ for the acid is a methyl or trichloroethyl ester. The methyl and trichloroethyl esters can be removed by base, e.g., LiOH in a polar solvent, e.g., aqueous THF, etc. The trichloroethyl ester can also be removed by treatment with zinc and ammonium acetate in a polar solvent, e.g., THF. Typically, PG² and
10 PG⁴ are acid labile groups, e.g., BOC, and are deprotected using HCl in dioxane, or TMSOTf in dioxane, dichloromethane. Typically, PG³ and PG⁵ are ester groups, removed by treatment with alkali metal hydroxide in aqueous THF or dioxane. Typically PG⁶ is an acid labile group, e.g., BOC, for an amine and removed as described for PG², or a silyl ether for a hydroxyl group, and can be removed by treatment with HF.pyridine or TBAF in an organic solvent, e.g.,
15 dichloromethane. Typically PG⁸ is an amine protecting group, e.g., BOC, and is removed as described for PG² or a silyl ether for a hydroxyl group, which can be removed as described for PG⁶, or an acetate protecting group, which can be removed with an alkali metal hydroxide in aqueous THF or dioxane.


Compound **1f** is then coupled to acid **1a** using the conditions described above for
20 Connection 1 to produce compound **1g**. Compound **1g** is then deprotected using conditions described in Greene and Wuts and coupled to **1c** to provide **1h** using the conditions described above for Connection 2. An alternative sequence for generating **1h** begins with coupling amine **1b** to acid **1c** using conditions as described for Connection 2 above, to form **1i**; deprotection of the protecting group PG³ in **1i** using conditions described in Greene and Wuts, and finally
25 coupling with amine **1f** using conditions described for Connection 1 above forms **1h**.

Scheme 3




Compound **1h** is deprotected at PG⁶ using conditions described in Greene and Wuts and
 5 the amine is then coupled to **1d** using the conditions described above for Connection 3 to form
1j. Protecting group PG¹ in Compound **1j** is then removed using conditions described in Greene
 and Wuts, and the acid is then coupled to **1e** using conditions described for Connection 5 to
 form the acyclic intermediate **1k**. An alternative sequence to **1k** is first deprotection of PG¹ and
 10 then coupling to **1e** as described for Connection 5; followed by deprotection of PG⁶ as described
 in Greene and Wuts, followed by coupling to **1d** using the conditions described for Connection 3
 to form **1k**. Acyclic intermediate **1k** is then subjected to the RCM as described above for
 Connection 4 to form the macrocycle **M**.


Scheme 4: Macrolactamization/Macrolactonization Connection 5



- Compound **A** is coupled to Compound **B** using the conditions described above for Connection 4 to generate **2a**. Compound **2a** is then deprotected at PG^7 as described in Greene and Wuts to generate acid **2b**. Acid **2b** is then coupled to the deprotected product of **1h** (prepared from **1h** by deprotection of PG^6 described in Greene and Wuts) to generate the precursor **2c**. Deprotection of **2c** is carried out using conditions described in Greene and Wuts, and then the product is cyclized using the conditions described above for macrolactamization or macrolactonization in Connection 5, to provide Compound **M**.


Scheme 5: Macrolactamization/Macrolactonization Connection 5

- Compound **2a** is deprotected at PG^8 as described in Greene and Wuts and above in
- 5 Connection 5 to generate **3a** which is then coupled, using conditions described above for Connection 5, to the deprotected product of **1h** (prepared from **1h** by deprotection of PG^1 described in Greene and Wuts) to generate the precursor **3b**. Deprotection of **3b** is carried out using conditions described in Greene and Wuts, and then cyclized using the conditions described above for macrolactamization or macrolactonization in Connection 3, to provide
- 10 Compound **M**.

Scheme 6: Further transformation of macrocyclic compounds **M to **M1-M5****

For example, when **M** contains a C=C as a result of RCM, Compound **M** is mixed in a solvent such as ethanol, methanol, etc., in the presence of palladium on carbon catalyst under an atmosphere of hydrogen gas to provide reduced Compound **M1**. The final macrocycle **M** from Schemes 4-6 often contains protecting groups on side-chains that require further removal to generate the final compound **M**. Protecting groups on the R^{4a} , R^{4b} , R^{3a} , R^{3b} , A^1 , A^2 and/or X^1 are removed using conditions described in Greene and Wuts to generate Compound **M2**. Another transformation is click chemistry to produce triazole **M3**. This transformation is performed by treating the alkyne or azide in **M**, in solvent (e.g., DMF) with an alkyne or azide as appropriate in the presence of CuI to form **M3**.


Deprotected compound **M2** can be further transformed after deprotection to additional macrocycle **M**. For example, treatment of **M2** containing an -OH with an alkyl halide in the presence of a base, e.g., cesium carbonate, in a suitable solvent, e.g., DMF, acetonitrile, etc., forms alkylated product **M4**. **M2** containing a ketone group is treated in a suitable solvent (e.g., DMF, methanol, etc.) with an amine followed by the addition of sodium acetoxyborohydride to form amine product **M5**.

Scheme 7: Preparation of acids or esters **B**

Many **B** components containing an acid or ester with a terminal alkene or $\text{CH}_3\text{-CH=C-}$ or vinyl/aryl boronate groups are commercially available or are described in the literature and can

be used in the above schemes directly. In addition, the following schemes are examples of methods that can be used to generate additional **B** components.

Scheme 8: Preparation of acids or esters B



5

In Scheme 8, part 1, a protected acid is treated with a strong deprotonating base, e.g., LDA in an inert solvent, e.g., THF, at -78°C and HMPA. A pre-cooled solution of

dichloroacetylene (prepared by treatment of trichlorethene with potassium hydride and MeOH (catalytic) in THF) is then added to generate the chloro acetylene product. This product is then reduced, for example, by treatment with Cu in acetic acid and THF, to generate the alkyne, which is then further reduced to the alkene, for example, by treatment of an alcoholic solution of the alkyne with a poisoned palladium reducing agent (e.g., Lindlar) in the presence of hydrogen gas. Alternatively, the alkyne is treated with Cp_2ZrHCl in dichloromethane in the presence of pinacolborane to form the vinyl boronate.

In Scheme 8, part 2, a *beta*-keto ester with *alpha* substitution is converted to the vinyl triflate, for example, by treating a THF solution of the *beta*-keto ester with a base, e.g., LDA, in THF at -78°C , followed by addition of $\text{PhN}(\text{Tf})_2$. The triflate product is then treated with pyridine at elevated temperature to form the alkyne. The alkyne is then treated as described above in part 1 to generate the alkene or vinyl boronate products.

In Scheme 8, part 3, a chiral aldol reaction is used. An acyl group is first attached to a chiral auxiliary, e.g., Evans, Oppolzer sultam (see *JACS* 1990, 112, p2767), using the standard amide bond formation conditions as described above for Connection 1-3. The Oppolzer auxiliary product is treated with the aldehyde of choice, TBDMSOTf and base, e.g., triethylamine, in anhydrous solvent, e.g., dichloromethane. The Evans auxiliary is treated with base, e.g., LDA, KHMDS, DIPEA, in organic solvent, e.g., THF, at -78°C and the aldehyde of choice in the presence of a Lewis acid, e.g., TiCl_4 , SnCl_4 , BF_3OEt_2 . Protection of the resulting alcohol from the aldol reaction is performed as described in Greene and Wuts, or alternatively alkylation with an alkyl halide or Meerwein's reagent, i.e., treatment with trimethyloxonium tetrafluoroborate in an inert solvent, e.g., dichloromethane, is performed. The auxiliary is then removed using standard alkali metal hydroxide removal conditions, e.g., LiOH in THF, or LiOH and hydrogen peroxide in THF, to provide the free acids product.

In Scheme 8, part 4, an Evans auxiliary is allylated with an allyl halide as described in *Synlett* 2002, 12, 2039-2040. The product is then isomerized by treatment with RhCl_3 in ethanol and then the auxiliary removed by base and peroxide, e.g., LiOH and H_2O_2 in THF/Water. Alternatively the auxiliary is directly removed by LiOH and H_2O_2 in THF/Water to provide the terminal alkene.

In Scheme 8, part 5, a Horner Wadsworth Emmons reaction is used on an aldehyde (containing a terminal alkene) to generate the alpha-beta unsaturated ester, which is then selectively reduced to the ester. For example, the phosphonate is treated with base, e.g., sodium hydride, in THF at low temperature, followed by addition of the aldehyde and warming to

generate the unsaturated ester. The product is reduced by treatment with magnesium powder in methanol.


In Scheme 8, part 6, an *alpha-beta* unsaturated acid is converted to the unsaturated Evans auxiliary (see *Organic Letters* 2007, 9, p1635) and is treated with an aldehyde to
 5 generate the corresponding alkene product. The hydroxyl group is then protected using methods described in Greene and Wuts and then the auxiliary is removed by treatment with base and peroxide, e.g., LiOH and H₂O₂ in THF/Water. The hydroxyl can also be alkylated by as described above for aldol Scheme 8, part 3.

In Scheme 8, part 7, a ketone is transformed via the nitro olefin as described in *Angew. Chem. Int. Ed.* **2006**, 45 (46), 7736. The nitro olefin is then treated with vinyl magnesium bromide in an inert solvent, e.g., THF, in the presence of a copper(I)salt, e.g., CuI and trimethylsilyl chloride. The nitro alkyl product after addition of the vinyl group is then converted to the acid by treating with sodium nitrite and acetic acid in an inert polar solvent, e.g., DMSO.

Several types of **A** are available commercially or described in the literature where X¹ is O
 15 or NH and Y² is a halide or A²-Y² contains an alkene. The schemes below describe additional general methods for generating **A**.


Scheme 9: Preparation of **A**

1. When Y² = Halogen in **A**



Scheme 9 (cont.): Preparation of A

2. When $Y^2 = H$ and A^2-Y^2 contains a alkene in **A**



In Scheme 9, part 1 (Y^2 is a halogen in **A**); the starting Compound **6a** is typically a commercially available aromatic compound, that contains halogen Y^2 and a group Z that can be transformed to the ketone **6b**. Typical Z groups are halide, acid, or aldehyde, for example.

When Z is an acid, **6a** is treated with a coupling agent, e.g., HATU, EDC in the presence of a base, e.g., DIPEA and the Weinreb amine (Me-NH-OMe) to form the Weinreb amide. The amide is then treated with a nucleophile, e.g., TMS- CF_3 to form the CF_3 substituted ketone **6b** or with a Grignard agent, e.g., MeMgBr in a solvent, e.g., THF, at $-78^\circ C$ to form the methyl ketone **6b**.

When Z is a halogen, then the initial conversion, if required, to a more reactive halogen is performed by treatment with NaI and acetyl chloride in an inert solvent, e.g., acetonitrile. The halogen is then transformed to the ketone by a Stille reaction with an ethoxyvinyl stannane. The halide is treated in an inert solvent, e.g., toluene, with the stannane and a palladium (II) catalyst, e.g., $PdCl_2(PPh_3)_2$, followed by treatment of the product with 2M HCl to afford ketone **6b**. In some cases the formation of an alkyl lithium reagent from the halide group can be performed by treatment with nBuLi at $-78^\circ C$ in THF and then addition of a N-methoxy-N-methyl amide to afford the ketone **6b** (e.g., N-methoxy-N-methylacetamide affords the R^5 is methyl ketone **6b**). A final method to generate the ketone **6b** is through a vinyl group. **6a** is treated with a

vinyltrifluoroborate in the presence of a palladium catalyst, e.g., PdCl₂(dppf) and then the vinyl product is subsequently ozonolysed in a polar solvent, e.g., methanol at low temperature to give an aldehyde. The aldehyde is then reacted with a nucleophile, e.g., TSM-CF₃ or a Grignard reagent, e.g., MeMgBr to afford a secondary alcohol product. The secondary alcohol is then
5 oxidized with Dess Martin Periodinane to give the desired ketone **6b** or can be used as **A** itself.

Chiral alcohol (X¹ is O) and amine (X¹ is NH) **A** are generated using Chiral reduction methods on the ketone **6b**. Chiral alcohol **6c** is formed from **6b** using one of the numerous chiral reduction methods available in the literature. Typically, dichloro(p-cumene)ruthenium(II) dimer and (1R,2R)-(-)-N-p-tosyl-1,2-diphenylethylenediamine are combined in water, and
10 sodium formate and **6b** is added in a water miscible solvent such as tetrahydrofuran. The reaction is then stirred at a temperature between ambient and reflux to produce **6c** where X¹ is O. Alternatively, a chiral CBS reduction can be performed in an inert solvent, e.g., THF at low temperature to also afford the chiral alcohol **6c**. Protection of the OH in **6c** is performed using methods described in Greene and Wuts, typically a TBS ether or acetyl group are used to
15 provide **A** (X¹ is O).

Alternatively, to make chiral **A** (X¹ is NH), ketone **6b** is first converted to a chiral imine (R¹⁰ is chiral group) and then reduced using a variety of methods described in the literature. For example, a chiral sulfinamide is reacted with the ketone **6b** to afford a chiral sulfinimine **6d**, which is then reduced with a suitable reducing agent, typically NaBH₄, or selectride, or a Noyori
20 type reduction as described for the chiral alcohol above, with dichloro(p-cumene)ruthenium(II) dimer and (1R,2R)-(-)-N-p-tosyl-1,2-diphenylethylenediamine. The sulfinamide auxiliary is then removed by treatment with mineral acid, preferably HCl in a suitable organic solvent such as methanol, to afford **6e** where X¹ is NH. Protection of the NH group can then be performed as described in Greene and Wuts to generate **A** (X¹ is NH).

25 In part 2 of Scheme 9, the synthesis of compound **A** where Y² is H and A²-Y² contains -CH=CH₂, a precursor for metathesis and cross coupling reactions is illustrated. Exemplary methods are as follows.

Compound **6c** or **6e** generated in Scheme 9, part 1 is first optionally protected on X¹ using a suitable protecting group as described in Greene and Wuts, and then a vinyl group is
30 introduced by a suitable cross coupling method onto the aryl or sp² halide. For example, a transition metal mediated coupling with a vinyl stannane or vinyl tetrafluoroborate using a suitable palladium catalyst, e.g., PdCl₂(dppf)₂ or PdCl₂(PPh₃)₂ in a suitable organic solvent, e.g., acetonitrile, dichloromethane, etc., with either thermal or microwave heating affords alkene **A**.

Another typical method that can be used to introduce a vinyl group is starting from the ketone **6b** where Y is OH. Initially, triflation of the alcohol is performed by treatment with Tf₂O in the presence of a base, e.g., pyridine. The ketone group is then reduced with a Noyori reduction, or as described above through the sulfonamide, to provide the chiral alcohol or amine. The chiral alcohol or amine is then protected as described in Greene and Wuts, and then the triflate is reacted with a vinyl cross coupling reagent, e.g., vinyl stannane in a Stille coupling, or a vinyltrifluoroborate as described above to introduce the alkene. A further example of alkene generation using **6b** ketone is via introduction of an allyl group. Thus, **6b** where Y is OH is treated in an inert solvent in the presence of a suitable base, e.g., alkali metal carbonate, preferably potassium carbonate with allyl bromide to form **6f**. Compound **6f** is then similar to ketone **6b** and is therefore able to be transformed as described above in part 1 to Compound **A** where X¹ is O or NH with protecting group PG⁸.

Pharmaceutical Formulations

The compounds of this invention are formulated with conventional carriers and excipients, which will be selected in accord with ordinary practice. Tablets will contain excipients, glidants, fillers, binders and the like. Aqueous formulations are prepared in sterile form, and when intended for delivery by other than oral administration generally will be isotonic.

While it is possible for the active ingredients to be administered alone it may be preferable to present them as pharmaceutical formulations. The formulations of the invention, both for veterinary and for human use, comprise at least one active ingredient, together with one or more acceptable carriers and optionally other therapeutic ingredients.

Formulations of the present invention suitable for oral administration may be presented as discrete units such as capsules, cachets or tablets each containing a predetermined amount of the active ingredient; as a powder or granules; as a solution or a suspension in an aqueous or non-aqueous liquid; or as an oil-in-water liquid emulsion or a water-in-oil liquid emulsion. The active ingredient may also be administered as a bolus, electuary or paste.

The effective dose of an active ingredient depends at least on the nature of the condition being treated, toxicity, whether the compound is being used prophylactically (lower doses) or against an active viral infection, the method of delivery, and the pharmaceutical formulation, and will be determined by the clinician using conventional dose escalation studies. The effective dose can be expected to be from about 0.0001 to about 100 mg/kg body weight per day;

typically, from about 0.01 to about 10 mg/kg body weight per day; more typically, from about .01 to about 5 mg/kg body weight per day; most typically, from about .05 to about 0.5 mg/kg body weight per day. For example, the daily candidate dose for an adult human of approximately 70 kg body weight will range from 1 mg to 1000 mg, preferably between 5 mg and 500 mg, and
 5 may take the form of single or multiple doses.

Combination Therapy

The compounds of the present invention may be combined with one or more active agents. Non-limiting examples of suitable active agents to be combined include one or more
 10 interferons, ribavirin or its analogs, HCV NS3 protease inhibitors, NS5a inhibitors, alpha-glucosidase 1 inhibitors, hepatoprotectants, mevalonate decarboxylase antagonists, antagonists of the renin-angiotensin system, other anti-fibrotic agents, endothelin antagonists, nucleoside or nucleotide inhibitors of HCV NS5B polymerase, non-nucleoside inhibitors of HCV NS5B polymerase, HCV NS5A inhibitors, TLR-7 agonists, cyclophilin inhibitors, HCV IRES inhibitors,
 15 pharmacokinetic enhancers and other drugs for treating HCV; or mixtures thereof.

More specifically, one or more compounds to be combined are selected from the group consisting of

- 1) interferons, *e.g.*, pegylated rIFN-alpha 2b (PEG-Intron), pegylated rIFN-alpha 2a (Pegasys), rIFN-alpha 2b (Intron A), rIFN-alpha 2a (Roferon-A), interferon alpha (MOR-22,
 20 OPC-18, Alfaferone, Alfanative, Multiferon, subalin), interferon alfacon-1 (Infergen), interferon alpha-n1 (Wellferon), interferon alpha-n3 (Alferon), interferon-beta (Avonex, DL-8234), interferon-omega (omega DUROS, Biomed 510), albinterferon alpha-2b (Albuferon), IFN alpha XL, BLX-883 (Locteron), DA-3021, glycosylated interferon alpha-2b (AVI-005), PEG-Infergen, PEGylated interferon lambda (PEGylated IL-29), and belerofof;
- 25 2) ribavirin and its analogs, *e.g.*, ribavirin (Rebetol, Copegus), and taribavirin (Virami3);
- 3) HCV NS3 protease inhibitors, *e.g.*, boceprevir (SCH-503034, SCH-7), telaprevir (VX-950), VX-813, TMC-435 (TMC435350), ABT-450, BI-201335, BI-1230, MK-7009, SCH-900518, VBY-376, VX-500, GS-9256, GS-9451, BMS-790052, BMS-605339, PHX-1766, AS-
 30 101, YH-5258, YH5530, YH5531, and ITMN-191 (R-7227);
- 4) alpha-glucosidase 1 inhibitors, *e.g.*, celgosivir (MX-3253), Miglitol, and UT-231B;

- 5) hepatoprotectants, *e.g.*, emericasan (IDN-6556), ME-3738, GS-9450 (LB-84451), silibinin, and MitoQ;
- 6) nucleoside or nucleotide inhibitors of HCV NS5B polymerase, *e.g.*, R1626, R7128 (R4048), IDX184, IDX-102, PSI-7851, BCX-4678, valopicitabine (NM-283), GS-6620 and
5 MK-0608;
- 7) non-nucleoside inhibitors of HCV NS5B polymerase, *e.g.*, filibuvir (PF-868554), ABT-333, ABT-072, BI-207127, VCH-759, VCH-916, JTK-652, MK-3281, VBY-708, VCH-222, A848837, ANA-598, GL60667, GL59728, A-63890, A-48773, A-48547, BC-2329, VCH-796 (nesbuvir), GSK625433, BILN-1941, XTL-2125, and GS-9190;
- 10 8) HCV NS5A inhibitors, *e.g.*, AZD-2836 (A-831), AZD-7295 (A-689), and BMS-790052;
- 9) TLR-7 agonists, *e.g.*, imiquimod, 852A, GS-9524, ANA-773, ANA-975, AZD-8848 (DSP-3025), PF-04878691, and SM-360320;
- 10) cyclophilin inhibitors, *e.g.*, DEBIO-025, SCY-635, and NIM811;
- 15 11) HCV IRES inhibitors, *e.g.*, MCI-067;
- 12) pharmacokinetic enhancers, *e.g.*, BAS-100, SPI-452, PF-4194477, TMC-41629, GS-9350, GS-9585, and roxythromycin;
- 13) other drugs for treating HCV, *e.g.*, thymosin alpha 1 (Zadaxin), nitazoxanide (Alinea, NTZ), BIVN-401 (virostat), PYN-17 (altirex), KPE02003002, actilon (CPG-10101), GS-
20 9525, KRN-7000, civacir, GI-5005, XTL-6865, BIT225, PTX-111, ITX2865, TT-033i, ANA 971, NOc-205, tarvacin, EHC-18, VGX-410C, EMZ-702, AVI 4065, BMS-650032, BMS-791325, Bavituximab, MDX-1106 (ONO-4538), Oglufanide, FK-788, and VX-497 (merimepodib);
- 14) mevalonate decarboxylase antagonists, *e.g.*, statins, HMGCoA synthase inhibitors (*e.g.*, hymeglusin), squalene synthesis inhibitors (*e.g.*, zaragozic acid);
- 25 15) angiotensin II receptor antagonists, *e.g.*, losartan, irbesartan, olmesartan, candesartan, valsartan, telmisartan, eprosartan;
- 16) angiotensin-converting enzyme inhibitors, *e.g.*, captopril, zofenopril, enalapril, ramipril, quinapril, perindopril, lisinopril, benazepril, fosinopril;
- 17) other anti-fibrotic agents, *e.g.*, amiloride; and
- 30 18) endothelin antagonists, *e.g.* bosentan and ambrisentan.

In yet another embodiment, the present application provides a combination therapy comprising a composition of the present invention and a second pharmaceutical composition comprising at least one additional therapeutic agent selected from the group consisting of HIV protease inhibiting compounds, HIV non-nucleoside inhibitors of reverse transcriptase, HIV nucleoside inhibitors of reverse transcriptase, HIV nucleotide inhibitors of reverse transcriptase, HIV integrase inhibitors, gp41 inhibitors, CXCR4 inhibitors, gp120 inhibitors, CCR5 inhibitors, interferons, ribavirin analogs, NS3 protease inhibitors, NS5a inhibitors, alpha-glucosidase 1 inhibitors, cyclophilin inhibitors, hepatoprotectants, non-nucleoside inhibitors of HCV, and other drugs for treating HCV, and combinations thereof.

More specifically, one or more compounds of the present invention may be combined with one or more compounds selected from the group consisting of 1) HIV protease inhibitors, *e.g.*, amprenavir, atazanavir, fosamprenavir, indinavir, lopinavir, ritonavir, lopinavir + ritonavir, nelfinavir, saquinavir, tipranavir, brexanavir, darunavir, TMC-126, TMC-114, mozenavir (DMP-450), JE-2147 (AG1776), AG1859, DG35, L-756423, RO0334649, KNI-272, DPC-681, DPC-684, and GW640385X, DG17, PPL-100, 2) a HIV non-nucleoside inhibitor of reverse transcriptase, *e.g.*, capravirine, emivirine, delaviridine, efavirenz, nevirapine, (+) calanolide A, etravirine, GW5634, DPC-083, DPC-961, DPC-963, Mb-150, and TMC-120, TMC-278 (rilpivirine), efavirenz, BILR 355 BS, VRX 840773, UK-453,061, RDEA806, 3) a HIV nucleoside inhibitor of reverse transcriptase, *e.g.*, zidovudine, emtricitabine, didanosine, stavudine, zalcitabine, lamivudine, abacavir, amdoxovir, elvucitabine, alovudine, Mb-210, racivir ($\pm$ -FTC), D-d4FC, emtricitabine, phosphazide, fozivudine tidoxil, fosaltivudine tidoxil, apricitabine (AVX754), amdoxovir, KP-1461, abacavir + lamivudine, abacavir + lamivudine + zidovudine, zidovudine + lamivudine, 4) a HIV nucleotide inhibitor of reverse transcriptase, *e.g.*, tenofovir, tenofovir disoproxil fumarate + emtricitabine, tenofovir disoproxil fumarate + emtricitabine + efavirenz, and adefovir, 5) a HIV integrase inhibitor, *e.g.*, curcumin, derivatives of curcumin, chicoric acid, derivatives of chicoric acid, 3,5-dicaffeoylquinic acid, derivatives of 3,5-dicaffeoylquinic acid, aurintricarboxylic acid, derivatives of aurintricarboxylic acid, caffeic acid phenethyl ester, derivatives of caffeic acid phenethyl ester, tyrphostin, derivatives of tyrphostin, quercetin, derivatives of quercetin, S-1360, zintevir (AR-177), L-870812, and L-870810, MK-0518 (raltegravir), BMS-707035, MK-2048, BA-011, BMS-538158, GSK364735C, 6) a gp41 inhibitor, *e.g.*, enfuvirtide, sifuvirtide, FB006M, TRI-1144, SPC3, DES6, Locus gp41, CovX, and REP 9, 7) a CXCR4 inhibitor, *e.g.*, AMD-070, 8) an entry inhibitor, *e.g.*, SP01A, TNX-355, 9) a gp120 inhibitor, *e.g.*, BMS-488043 and BlockAide/CR, 10) a G6PD and NADH-oxidase inhibitor, *e.g.*, immunitin, 10) a CCR5 inhibitor, *e.g.*, aplaviroc, vicriviroc, INCB9471, PRO-140, INCB15050,

PF-232798, CCR5mAb004, and maraviroc, 11) an interferon, e.g., pegylated rIFN-alpha 2b, pegylated rIFN-alpha 2a, rIFN-alpha 2b, IFN alpha-2b XL, rIFN-alpha 2a, consensus IFN alpha, interferon, rebif, locteron, AVI-005, PEG-interferon, pegylated IFN-beta, oral interferon alpha, reaféron, intermax alpha, r-IFN-beta, interferon + actimmune, IFN-omega with DUROS, and

5 albuferon, 12) ribavirin analogs, e.g., rebetol, copegus, VX-497, and viramidine (taribavirin) 13) NS5a inhibitors, e.g., A-831, A-689 and BMS-790052, 14) NS5b polymerase inhibitors, e.g., NM-283, valopicitabine, R1626, PSI-6130 (R1656), IDX184, PSI-7851, HCC-796, BILB 1941, MK-0608, NM-107, R7128, VCH-759, PF-868554, GSK625433, and XTL-2125, 15) NS3

10 protease inhibitors, e.g., SCH-503034 (SCH-7), VX-950 (Telaprevir), ITMN-191, and BILN-2065, 16) alpha-glucosidase 1 inhibitors, e.g., MX-3253 (celgosivir) and UT-231B, 17) hepatoprotectants, e.g., IDN-6556, ME 3738, MitoQ, and LB-84451, 18) non-nucleoside inhibitors of HCV, e.g., benzimidazole derivatives, benzo-1,2,4-thiadiazine derivatives, and phenylalanine derivatives, 19) other drugs for treating HCV, e.g., zadaxin, nitazoxanide (alinea), BIVN-401 (virostat), DEBIO-025, VGX-410C, EMZ-702, AVI 4065, bavituximab, oglufanide,

15 PYN-17, KPE02003002, actilon (CPG-10101), KRN-7000, civacir, GI-5005, ANA-975, XTL-ANA 971, NOC-205, tarvacin, EHC-18, and NIM811, 19) pharmacokinetic enhancers, e.g., BAS-100, SPI-452, PF-4194477, TMC-41629, GS-9350, GS-9585, and roxythromycin, 20) RNAse H inhibitors, e.g., ODN-93 and ODN-112, 21) other anti-HIV agents, e.g., VGc-1, PA-457 (bevirimat), ampligen, HRG214, cytolin, polymun, VGX-410, KD247, AMZ 0026, CYT 99007, A-

20 221 HIV, BAY 50-4798, MDX010 (ipilimumab), PBS119, ALG889, and PA-1050040.

In a specific aspect of this embodiment, the additional therapeutic agent is selected from ribavirin, telaprevir, boceprevir and sofosbuvir (GS-7977 (formerly PSI-7977)).

A combination therapy described herein may be administered as a simultaneous or sequential regimen. When administered sequentially, the combination may be administered in

25 two or more administrations.

Co-administration of a compound of the invention with one or more other active agents generally refers to simultaneous or sequential administration of a compound of the invention and one or more other active agents, such that therapeutically effective amounts of the compound of the invention and one or more other active agents are both present in the body of

30 the patient.

Method for treating viral infection

The present application provides a method for treating a *Flaviviridae* viral infection comprising administering a therapeutically effective amount of a compound described herein or a pharmaceutically acceptable salt, isotope, stereoisomer, mixture of stereoisomers, tautomer, ester or prodrug thereof, to a human subject in need thereof.

Also provided is a method for treating a *Coronaviridae* viral infection comprising administering a therapeutically effective amount of a compound described herein or a pharmaceutically acceptable salt, isotope, stereoisomer, mixture of stereoisomers, tautomer, ester or prodrug thereof, to a human subject in need thereof.

In one embodiment, the method of inhibiting or treating a disease comprises administering to an animal a composition comprising an effective amount of one or more compounds of the invention or a pharmaceutically acceptable salt, isotope, stereoisomer, mixture of stereoisomers, tautomer, ester or prodrug thereof and a pharmaceutically acceptable carrier. The composition to be administered may further contain a secondary therapeutic agent as described above.

A method of the present application is particularly suitable for use with humans, but may be used with other animals, particularly mammals, such as, for example, non-human primates, companion animals, farm animals, laboratory animals, and wild and zoo animals.

A method of the present application is particularly useful to treat diseases caused directly or indirectly by *Flaviviridae* virus since the compounds of the present invention have inhibitory activity against those viruses. In some embodiments, therefore, a method of the present invention is used in inhibiting or treating diseases caused by a Hepatitis C virus. In some embodiments, therefore, a method of the present invention is used in inhibiting or treating diseases caused by a Hepatitis B virus. In an aspect, such a method is applied to a patient with a disease caused by the viral infection such as dengue fever, yellow fever, hepatitis C, Japanese encephalitis, Kyasanur forest disease, Murray valley encephalitis, St. Louis encephalitis, tick-borne encephalitis or West Nile encephalitis.

In some embodiments, a sustained virologic response is achieved at about 12 weeks, at about 10 weeks, at about 8 weeks, at about 6 weeks, or at about 4 weeks, or at about 4 months, or at about 5 months, or at about 6 months, or at about 1 year, or at about 2 years.

A method of the present application is also particularly useful to treat diseases caused directly or indirectly by *Coronaviridae* virus since the compounds of the present invention have

inhibitory activity against those viruses. In some embodiments, therefore, a method of the present invention is used in inhibiting or treating diseases caused by a SARS coronavirus. In an aspect, such a method is applied to a patient with a disease caused by the viral infection such as severe acute respiratory syndrome (SARS), cancer, inflammation, obesity, acquired immune deficiency syndrome (AIDS), or cirrhosis.

In another aspect, the compounds disclosed herein can be used for treating cancer. In yet another aspect, the compounds disclosed herein can be used for immunomodulation. In some embodiments, therefore, a method of the present invention comprises adjusting an immune response to a desired level, as in immunopotential, immunosuppression, or induction of immunologic tolerance.

In some embodiments, the compound is administered for about 12 weeks. In further embodiments, the compound is administered for about 12 weeks or less, for about 10 weeks or less, for about 8 weeks or less, for about 6 weeks or less, or for about 4 weeks or less. The compound may be administered once daily, twice daily, once every other day, two times a week, three times a week, four times a week, or five times a week.

In another aspect, the compounds disclosed herein can be used for treating cancer. In yet another aspect, the compounds disclosed herein can be used for immunomodulation. In some embodiments, therefore, a method of the present invention comprises adjusting an immune response to a desired level, as in immunopotential, immunosuppression, or induction of immunologic tolerance.

EXAMPLES

The following examples are merely illustrative, and do not limit this disclosure in any way.


List of Abbreviations and Acronyms

Abbreviation	Meaning
°C	Degree Celsius
Ac	Acetyl
app	Apparent
Aq	Aqueous
Boc	tert-Butoxycarbonyl
Br	Broad
Bu	Butyl
cat	Catalytic
CBS	Corey Bakshi Shibata

CDMT	2-chloro-4,6-dimethoxy-1,3,5-triazine
cm	Centimeter
COMU	(1-Cyano-2-ethoxy-2-oxoethylidenaminoxy)dimethylamino-morpholino-carbenium hexafluorophosphate
CP/Cp	Cyclopentyl
CPME	Cyclopentyl methyl ether
Cy/cHex	Cyclohexyl
d	Doublet
DBU	1,8-Diazabicyclo[5.4.0]undec-7-ene
DCC	N,N'-dicyclohexylcarbodiimide
DCE	Dichloroethane
DCM	Dichloromethane
dd	Doublet of doublets
ddd	Doublet of doublet of doublet
ddt	Doublet of doublet of triplet
DIAD	Diisopropyl azodicarboxylate
DIPEA	N,N-Diisopropylethylamine
DMAP	4-Dimethylaminopyridine
DME	1,2-dimethoxyethane
DMF	Dimethylformamide
DMSO	Dimethylsulfoxide
dppf	1,1'-bis(diphenylphosphino)ferrocene
dq	Doublet of quartet
dt	Doublet of triplet
dtd	Doublet of triplet of doublet
EDC	1-Ethyl-3-(3-dimethylaminopropyl)carbodiimide
Equiv/eq	Equivalents
Et	Ethyl
g	Grams
gen.	Generation
HATU	(Dimethylamino)-N,N-dimethyl(3H-[1,2,3]triazolo[4,5-b]pyridin-3-yloxy)methaniminium hexafluorophosphate
HBTU	2-(1H-benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate
HMDS	Hexamethyldisilazane
HMPA	hexamethylphosphoramide
HOBT	Hydroxybenzotriazole
HPLC	High-performance liquid chromatography
hrs/h	Hours
Hz	Hertz
IC ₅₀	The half maximal inhibitory concentration
Im	imidazole
i-Pr/iPr	Isopropyl
J	Coupling constant
Kg	Kilogram
LCMS	Liquid chromatography-mass spectrometry
LDA	Lithium diisopropylamide
M	Molar
m	Multiplet
m/z	mass-to-charge ratio
M+	Mass peak

Me	Methyl
mg	Milligram
MHz	Megahertz
min	Minute
mL	Milliliter
mM	Millimolar
mm	Millimeter
mmol	Millimole
mol	Mole
Ms	Methanesulfonyl
MW	Microwave
N	normal
nM	Nanomolar
NMM	<i>N</i> -methylmorpholine
NMR	Nuclear magnetic resonance
nPrOH	<i>n</i> -propanol
o-Tol	o-Tolyl
Ph	Phenyl
pTSA	p-Toluenesulfonic acid
Py/pyr	Pyridine
PyAOP	7-azabenzotriazol-1-yloxy)tripyrrolidinophosphonium hexafluorophosphate
PyBop	benzotriazol-1-yl-oxytripyrrolidinophosphonium hexafluorophosphate
q	Quartet
R _f	Retention factor
RT/rt/r.t.	Room temperature
s	Singlet
sat.	Saturated
t	Triplet
TBAF	Tetra- <i>n</i> -butylammonium fluoride
TBDMS/TBS	tert-Butyldimethylsilyl
TBDPS	<i>tert</i> -Butyldiphenylsilyl
<i>t</i> -Bu	<i>tert</i> -butyl
td	Triplet of doublets
TEA	Triethylamine
Tf	Trifluoromethanesulfonyl
TFA	Trifluoroacetic acid
THF	Tetrahydrofuran
TLC	Thin layer chromatography
TMEDA	tetramethylethylenediamine
TMS	Trimethylsilyl
Tr/tr	Retention time
Ts	Tosyl
tt	Triplet of triplet
UV	Ultraviolet
wt.	weight
δ	Chemical shift
μL	Microliter
μM	Micromolar
μmol	Micromole

EXAMPLE 1: (E)-(2R,5S,11S,14S,17R,18R)-18-Hydroxy-14-isopropyl-2,11,17-trimethyl-3,9,12,15,28-pentaaza-tricyclo[21.3.1.1*5,9*]octacos-1(26),21,23(27),24-tetraene-4,10,13,16-tetraone




A solution of 1-((1R,5S)-10,10-dimethyl-3,3-dioxo-3λ⁶-thia-4-azatricyclo[5.2.1.0*1,5*]dec-4-yl)-propan-1-one (3.95 g, 14.55 mmol) in toluene (50 mL) was prepared, then evaporated to dryness. This process was repeated and the resulting white solid was dissolved in anhydrous dichloromethane (16 mL). A small quantity of calcium hydride was added before adding *tert*-butyldimethylsilyl trifluoromethanesulfonate (3.83 mL, 14.5 mmol) and anhydrous triethylamine (2.33 mL, 16.7 mmol). The reaction mixture was stirred at RT ("RT") under a nitrogen atmosphere for 15 hours ("h"). The resulting solution was evaporated to yield a thick paste, which was re-dissolved in anhydrous dichloromethane (15 mL) and added dropwise to a stirred solution of 4-pentenal (2.69 g, 32.0 mmol) and titanium tetrachloride (1 M in dichloromethane, 32 mL, 32 mmol) in anhydrous dichloromethane (20 mL) at -78 °C, under a nitrogen atmosphere. The reaction was stirred at -78 °C for 30 minutes ("min") before diluting with saturated aqueous ammonium chloride solution (100 mL). The layers were separated and the aqueous layer was extracted with dichloromethane (2 x 50 mL). The combined extract was dried over sodium sulfate, filtered and evaporated to give a brown gum. This was purified by silica gel chromatography using *iso*-hexanes/ethyl acetate 4:1 to yield the title compound (3.09 g, 60%) as a colorless gum.

10

15


20



A solution of **1a** (12.0 g, 0.034 mol) in anhydrous dichloromethane (520 mL) was cooled to 0 °C, before adding pyridine (5.5 mL, 0.068 mol) then *tert*-butyldimethylsilyl trifluoromethanesulfonate (9 mL, 0.039 mol). The reaction mixture was stirred at 0 °C for 15 min then allowed to warm to RT and stirred for a further 1.5 h. The reaction mixture was washed


25

with saturated sodium bicarbonate (400 mL). The aqueous wash was back-extracted with dichloromethane (200 mL). The organic layers were combined and washed with dilute brine (200 mL) and 2 M hydrochloric acid (200 mL). The solution was dried over sodium sulfate, filtered and evaporated to give the title product (15.29 g, 96%) as a white solid.



A solution of **1b** (15.29 g, 0.0325 mol) in tetrahydrofuran (300 mL) was prepared and a 2 M aqueous solution of lithium hydroxide (120 mL) was added. The stirred mixture was heated to 60°C for 15 h. The reaction was diluted with 2 M hydrochloric acid (250 mL). The layers were separated and the aqueous was extracted with ethyl acetate (2 x 200 mL). The organic layers were combined, dried over sodium sulfate, filtered and evaporated to give a cream solid (16.7 g). The solid was purified by silica gel chromatography using 3:7 ethyl acetate/ *iso*-hexanes to yield the title product (7.18 g, 81%) as a colorless gum.

10




A solution of (S)-2-*tert*-butoxycarbonylamino-propionic acid (3.28 g, 17.32 mmol) in acetonitrile (160 mL) was cooled to 0°C before addition of *N,N*-diisopropylethylamine (12 mL, 69.3 mmol) then 2-(1H-benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (6.57 g, 17.32 mmol). The reaction mixture was stirred at 0°C for 20 min and a solution of (S)-hexahydropyridazine-3-carboxylic acid 2,2,2-trichloroethyl ester trifluoroacetic acid salt (preparation described in *Angew. Chem. Int. Ed. Engl.* 1999, 38, 2443, 6.49 g, 17.3 mmol) in acetonitrile (80 mL) was added. The reaction was allowed to warm to RT and was stirred for 15 h. The reaction mixture was evaporated then re-dissolved in ethyl acetate (150 mL). The solution was washed with brine (150 mL). The brine was back extracted with ethyl acetate (50 mL). The organic layers were combined, dried over sodium sulfate, filtered and evaporated to give a dark oil. The oil was purified by silica gel chromatography using *iso*-hexanes/ethyl acetate 1:1 to yield the title compound (6.88 g, 92%) as a colorless gum.


15

20

25




A solution of **1d** (6.88 g, 15.9 mmol) in dichloromethane (200 mL) was prepared and trifluoroacetic acid (50 mL) was added. The reaction mixture was stirred at RT for 2 h. TLC showed the reaction to be complete. The solution was evaporated to give a brown oil. This was azeotroped with toluene (50 mL) and the resultant oil was dried under vacuum to give (S)-1-((S)-2-amino-propionyl)-hexahydro-pyridazine-3-carboxylic acid 2,2,2-trichloro-ethyl ester trifluoroacetic acid salt (7.8 g) as a brown gum. A solution of ((S)-1-carbamoyl-2-methylpropyl)-carbamic acid *tert*-butyl ester in acetonitrile (300 mL) was cooled to 0°C before adding *N,N*-diisopropylethylamine (13.8 mL, 79.7 mmol) and 2-(1H-benzotriazole-1-yl)-1,3,3-tetramethyluronium hexafluorophosphate (6.33 g, 16.7 mmol). The reaction was stirred at 0°C for 15 min before adding a solution of the (S)-1-((S)-2-amino-propionyl)-hexahydro-pyridazine-3-carboxylic acid 2,2,2-trichloro-ethyl ester trifluoroacetic acid salt (ca. 15.9 mmol) in acetonitrile (85 mL). The reaction was stirred at 0°C for a further 20 min then allowed to warm to RT and stirred for 15 h. The reaction mixture was evaporated then re-dissolved in ethyl acetate (250 mL). The solution was washed with water (150 mL) then dried over sodium sulfate, filtered and evaporated to give a red oil. This was purified by silica gel chromatography using *iso*-hexanes/ethyl acetate 7:3 then *iso*-hexanes/ethyl acetate 1:1 to yield the title compound (8.2 g, 92%) as a pale orange amorphous solid.



A solution of **1e** (10.0 g, 18.5 mmol) in anhydrous dichloromethane (200 mL) was prepared and trimethylsilyl trifluoromethanesulfonate (5 mL, 27.75 mmol) was added. The reaction mixture was stirred at RT for 2 h, then *N,N*-diisopropylethylamine (13.2 mL, 75.8 mmol) was added and the reaction mixture was evaporated to dryness. The residue was re-dissolved in acetonitrile (200 mL) and a solution of (2R,3R)-3-(*tert*-butyl-dimethyl-silanyloxy)-2-methyl-hept-6-enoic acid (5.04 g, 18.5 mmol) in acetonitrile (60 mL) was added followed by 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide (5.0 g, 26.4 mmol) and 1-hydroxybenzotriazole (4.4 g,

26.4 mmol). The reaction mixture was stirred at RT for 15 h. It was evaporated to give a thick yellow oil. The oil was purified by silica gel chromatography using 1:1 ethyl acetate / *iso*-hexanes then 3:2 ethyl acetate / *iso*-hexanes to yield the title product (8.75 g, 69%) as a white solid.




5

Compound 1g

A solution of (R)-bromo- α -methylbenzylamine (1.023 g, 5.112 mmol) in dichloromethane (20 mL) was subsequently treated with triethylamine (720 μ L, 5.112 mmol) and di-*tert*-butyl dicarbonate (1.784 g, 8.179 mmol). After overnight stirring at RT, the volatiles were removed *in vacuo* and the residue was purified by silica gel chromatography using a 50 g Isolute cartridge eluted with a continuous gradient of *iso*-hexanes/diethyl ether 1:0 to 4:1 to afford the title compound (1.552 g, 100%) as a white solid.


10



Compound 1h

A solution of **1g** (10.26 g, 0.0342 mol.) and tributyl(vinyl)tin (32.5 g, 30 mL, 0.103 mol.) in toluene (175 mL) was purged with nitrogen for 30 min before addition of bis(triphenylphosphine) palladium (II) dichloride (2.38 g, 0.0034 mol.). The stirred mixture was heated to 60°C for 16 h before cooling to RT. The reaction mixture was filtered through hyflo-supercel then evaporated to give a dark coloured oil. The oil was purified by silica gel chromatography using *iso*-hexanes/ethyl acetate 19:1 to yield the title compound (6.95 g, 82%) as a yellow oil.

15




Compound 1i

20

A solution of **1h** (6.95 g, 28.1 mmol.) in 1,4-dioxane (30 mL) was prepared and a solution of hydrogen chloride in 1,4 dioxane (4 M, 60 mL) was added. The reaction mixture was

stirred at RT for 2 h then evaporated to dryness. The resultant solid was re-dissolved in toluene and evaporated. The solid was triturated with diethyl ether, which was removed by decanting. The solid was then dried under vacuum to give the title compound (4.96 g, 96%) as an off-white solid.




A solution of **1f** (8.75 g, 12.75 mmol) in tetrahydrofuran (350 mL) was prepared and zinc powder (8.44 g, 127.5 mmol) was added followed by a 1 M aqueous solution of ammonium acetate (90 mL, 90 mmol). The reaction mixture was vigorously stirred for 16 h then filtered through hyflo-supercel. The solution was cooled over an ice bath before addition of aqueous ammonium chloride solution (350 mL). It was allowed to re-cool before acidifying to pH 1 by addition of 2 M hydrochloric acid. The layers were separated and the aqueous was extracted with ethyl acetate (2 x 250 mL). The organic layers were combined, dried over sodium sulfate filtered and evaporated to give a white solid. The solid was azeotroped with toluene (3 x 200 mL) then dried under vacuum to give a white solid (6.16 g), which was dissolved in acetonitrile (400 mL) before adding (R)-1-(3-vinyl-phenyl)-ethylamine hydrochloride (2.34 g, 12.75 mmol) followed by *N,N*-diisopropylethylamine (8.9 mL, 51 mmol) and 2-(1H-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyl uronium hexafluorophosphate methanaminium (6.8 g, 17.85 mmol). The reaction mixture was stirred at RT for 16 h and then evaporated to give a brown gum. The gum was purified by silica gel chromatography using ethyl acetate then 1:4 acetone/ethyl acetate to yield the title compound (5.51 g, 76%) as a cream solid.


10

15

20




A solution of **1j** (0.50 g, 0.88 mmol) in anhydrous dichloromethane (20 mL) was cooled over an ice bath, before adding pyridine (92 μ L, 1.14 mmol) and *tert*-butyldimethylsilyl trifluoromethanesulfonate (242 μ L, 1.05 mmol). The reaction mixture was stirred at 0°C for 15 min, then allowed to warm to RT and stirred for 1 hour. The reaction mixture was evaporated and the residue was purified by silica gel chromatography using ethyl acetate to yield the title product (477 mg, 80%) as a white solid.



Compound 1k

A solution of **1k** (477 mg, 0.7 mmol) in 1,2-dichloroethane (250 mL) was prepared and Hoveyda-Grubbs 2nd generation catalyst (43 mg, 0.07 mmol) was added. The stirred reaction mixture was heated to 80°C for 1 hour. The reaction mixture was cooled to RT before adding silica gel. The mixture was stirred for 10 min then evaporated and the residue was purified by silica gel chromatography using ethyl acetate to yield the title product (198 mg, 43%) as a white solid.




Compound 1l

A solution of **1l** (198 mg, 0.3 mmol) in tetrahydrofuran (20 mL) was cooled over an ice bath before adding a 1 M solution of tetrabutylammonium fluoride in tetrahydrofuran (1.5 mL, 1.5 mmol). The reaction was allowed to warm to RT and was stirred for 1 hour. The reaction mixture was treated with saturated aqueous sodium bicarbonate solution (20 mL) and extracted with ethyl acetate (2 x 20 mL). The extract was dried over sodium sulfate, filtered and evaporated. The residue was purified by silica gel chromatography using 1:9 acetone / ethyl acetate to yield the title product (150 mg, 92%) as a white solid. ¹H NMR (300 MHz, *d*₆-DMSO) δ 0.83 (d, *J* = 6.5 Hz, 3H), 0.85 (d, *J* = 6.5 Hz, 3H), 1.12-1.21 (m, 4H), 1.28 (d, *J* = 7.1 Hz, 3H), 1.36 (d, *J* = 6.9 Hz, 3H), 1.40-1.90 (m, 7H), 1.92-2.08 (m, 1H), 2.19-2.34 (m, 2H), 2.67-2.80 (m,


1H), 3.56-3.65 (m, 1H), 3.99-4.12 (m, 1H), 4.22 (*br d*, $J = 12.2$ Hz, 1H), 4.75 (d, $J = 11.8$ Hz, 4.87-4.99 (m, 1H), 5.12-5.24 (m, 1H), 5.40 (d, $J = 4.5$ Hz, 1H), 6.14-6.33 (m, 2H), 7.10-7.35 (m, 5H), 7.88 (d, $J = 8.3$ Hz, 1H), 8.55 (d, $J = 8.3$ Hz, 1H). LCMS (m/z) 542.3 [$M+H$], Tr = 1.87 min.

5 **EXAMPLE 2: (E)-(2R,5S,11S,14S,17R,18R)-14-Isopropyl-2,11,17-trimethyl-18-(2,2,2-trifluoro-ethoxy)-3,9,12,15,28-pentaaza-tricyclo[21.3.1.1*5,9*]octacos-1(26),21,23(27),24-tetraene-4,10,13,16-tetraone**



A solution of hexamethyldisilazane (235 μ L, 1.13 mmol) in anhydrous tetrahydrofuran (4 mL) was cooled to -10°C before adding a 2.5 M solution of *n*-butyllithium (380 μ L, 0.94 mmol). The stirred mixture was warmed to 0°C for 10 min before cooling to -78°C . A solution of Compound 1 (102 mg, 0.188 mmol) in anhydrous dimethylformamide (1.5 mL) and anhydrous tetrahydrofuran (1.5 mL) was added dropwise over 2 min followed by the addition of 2,2,2-trifluoroethyl trifluoromethanesulfonate (135 μ L, 0.94 mmol). The reaction mixture was stirred at -78°C under a nitrogen atmosphere then allowed to gradually warm to RT. The reaction mixture was quenched with the addition of a saturated aqueous solution of ammonium chloride (15 mL). The mixture was extracted with ethyl acetate (2 x 15 mL). The extract was dried over sodium sulfate, filtered and evaporated. The residue was purified by silica gel chromatography using 1:9 acetone/ethyl acetate to give a colorless gum (65 mg). The gum was further purified by reverse phase chromatography, using a 10 g C18 cartridge eluted with 2:3 acetonitrile / water. The partially evaporated fractions were extracted with ethyl acetate (2 x 15 mL). The extract was dried over sodium sulfate, filtered and evaporated to yield the title product (17 mg, 30%) as a white solid. ^1H NMR (300 MHz, CDCl_3) 0.92 (d, $J = 6.7$ Hz, 3H), 0.96 (d, $J = 6.9$ Hz, 3H), 1.32 (d, $J = 7.1$ Hz, 3H), 1.39 (d, $J = 7.1$ Hz, 3H), 1.50 (d, $J = 6.7$ Hz, 3H), 1.74-1.85 (m, 1H), 1.86-2.08 (m, 3H), 2.10-2.40 (m, 3H), 2.57-2.78 (m, 2H), 3.19-3.31 (m, 1H), 3.49 (*app t*, $J = 6.7$ Hz, 1H), 3.58-3.79 (m, 4H), 3.81-4.18 (m, 2H), 4.49 (*br d*, $J = 12.3$ Hz, 1H), 5.03-5.17 (m, 1H), 5.32-5.47 (m, 1H), 6.14-6.30 (m, 1H), 6.37-6.72 (m, 4H), 7.06-7.30 (m, 4H). LCMS (m/z) 624.3 [$M+H$], Tr = 2.57 min.


EXAMPLE 3: (2R,5S,11S,14S,17R,18R)-18-Hydroxy-14-isopropyl-2,11,17-trimethyl-3,9,12,15,28-pentaaza-tricyclo[21.3.1.1*5,9*]octacos-1(26),23(27),24-triene-4,10,13,16-tetraone



Compound 3

A solution of Compound 1 (100 mg, 0.184 mmol) in methanol (5 mL) was prepared and 10% palladium on carbon (5 mg) was added. The stirred mixture was placed under a hydrogen atmosphere for 1.5 h. The reaction mixture was filtered through hyflo-supercel then through a 0.2 μ m filter before evaporating to give the title compound (95 mg, 95%) as a white solid. ^1H NMR (300 MHz, CDCl_3) 0.93 (d, $J = 7.1$ Hz, 3H), 0.96 (d, $J = 7.1$ Hz, 3H), 1.32 (d, $J = 6.7$ Hz, 3H), 1.37-2.16 (m, 12H), 1.54 (d, $J = 6.9$ Hz, 3H), 2.34-2.46 (m, 1H), 2.52-2.75 (m, 3H), 3.16-3.27 (m, 1H), 3.43-3.62 (m, 3H), 3.74 (d, $J = 11.8$ Hz, 1H), 3.98-4.06 (m, 1H), 4.50 (d, $J = 13.2$ Hz, 1H), 5.11-5.36 (m, 2H), 6.39-6.50 (m, 2H), 7.70 (d, $J = 8.5$ Hz, 1H), 7.06-7.28 (m, 4H). LCMS (m/z) 544.3 [$\text{M}+\text{H}$], Tr = 1.91 min.


EXAMPLE 4




Compound 4a

To a suspension of sodium hydride (60% in mineral oil, 783 mg, 19.6 mmol) in dry tetrahydrofuran at 0°C , under nitrogen was added dropwise triethyl phosphonoacetate (3.38 mL, 19.6 mmol). The white suspension was stirred for 1 h where it became a solution, then 2,2-dimethyl-4-pentenal (2.42 mL, 17.8 mmol) was slowly added and the resulting green/yellow solution was stirred at 0°C and allowed to warm to RT. After 72 h ethanol (1 mL) was added to the solution, followed by water (100 mL) and the organics were extracted with diethyl ether (2 x 200 mL). The combined organics were washed with water (200 mL), dried over magnesium sulfate, filtered and concentrated *in vacuo* to give a crude residue. This was purified by silica gel


chromatography, using *iso*-hexanes (144 mL), then *iso*-hexanes/diethyl ether 50:50 (72 mL), then diethyl ether (48 mL) to give the title compound (3.20 g, 99%) as an oil.




To **4a** (3.20 g, 17.5 mmol) in dry methanol at RT, under nitrogen was slowly added
 5 magnesium powder (1.28 g, 52.7 mmol) while monitoring the temperature rise. After the addition was complete the mixture was stirred at RT overnight. After this time, to the reaction was added additional magnesium powder (852 mg, 35.1 mmol) and the reaction mixture stirred for 2 h. The mixture was neutralized from pH 10 to pH 7 with 2 M hydrochloric acid and then concentrated *in vacuo* to give a residue. The residue was suspended in ethyl acetate (300 mL) and water (500
 10 mL) and the layers separated. The aqueous phase was then re-extracted with ethyl acetate (200 mL) and the combined organics were dried over magnesium sulfate, filtered and concentrated *in vacuo* to give the title compound (1.96 g, 61%) as a colorless oil.



To **4b** (2.02 g, 11.0 mmol) in a mixture of tetrahydrofuran (35 mL), methanol (9 mL) and
 15 water (9 mL) at RT, was added lithium hydroxide monohydrate (1.38 g, 32.9 mmol) and the mixture was stirred at RT for 1 h. After this time additional lithium hydroxide monohydrate (460 mg, 11.0 mmol) was added and the reaction mixture stirred for 1 h. After this time more lithium hydroxide monohydrate (460 mg, 11.0 mmol) was added and the reaction mixture stirred for 30 min. The reaction mixture was then concentrated *in vacuo* and the residue diluted with water
 20 (200 mL) and the organics extracted with diethyl ether (3 x 50 mL). The aqueous phase was acidified from pH 14 to pH 1 with 2 M hydrochloric acid and the organics extracted with ethyl acetate (3 x 50 mL). The combined ethyl acetate organics were then dried over sodium sulfate, filtered and concentrated *in vacuo* to give the desired product (1.03 g, 60%) as an oil.




A solution of **1e** (1.06 g, 2.00 mmol) in dichloromethane (10 mL) was cooled in an ice-water bath under nitrogen. Trimethylsilyl trifluoromethanesulfonate (0.591 mL, 4.00 mmol) was added dropwise, and the resulting solution was stirred for 1.5 h. The reaction was quenched with *N,N*-diisopropylethylamine (1.4 mL, 8.00 mmol) and the reaction mixture was concentrated *in vacuo* to give (S)-1-[(S)-2-((S)-2-amino-3-methyl-butyrylamino)-propionyl]-hexahydro-pyridazine-3-carboxylic acid 2,2,2-trichloro-ethyl ester a white solid which was used without further purification. To (S)-1-[(S)-2-((S)-2-amino-3-methyl-butyrylamino)-propionyl]-hexahydro-pyridazine-3-carboxylic acid 2,2,2-trichloro-ethyl ester generated in the previous step was added a solution of 4,4-dimethyl-hept-6-enoic acid (343 mg, 2.20 mmol) in acetonitrile (70 mL). To this mixture was added *N,N*-diisopropylethylamine (1.4 mL, 8.03 mmol) and the reaction cooled in an ice-water bath before the addition of 2-(1H-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate methanaminium (1.07 g, 2.80 mmol). The yellow solution was then stirred and allowed to warm to RT overnight. The reaction mixture was quenched with 2 M hydrochloric acid (40 mL) and concentration *in vacuo*. To the residue was added water (200 mL) and the organics were extracted with ethyl acetate (3 x 150 mL). The combined organics were then washed with brine (100 mL) and sodium hydrogen carbonate (200 mL). The organics were then dried over magnesium sulfate, filtered and concentrated *in vacuo* to give a crude yellow oil (2.27 g). This was purified by silica gel chromatography using *iso*-hexanes/ethyl acetate 90:10 (300 mL) then *iso*-hexanes/acetone 50:50 (300 mL) then *iso*-hexanes/acetone 0:100 to give the title compound (1.10 g, 97%) as a colorless oil.




A solution of **4d** (1.10 g, 1.94 mmol) in tetrahydrofuran (40 mL) was prepared and zinc powder (2.79 g, 42.6 mmol) was added followed by a solution of ammonium acetate (2.24 g, 29.0 mmol) in water (10 mL). The reaction mixture was stirred at RT for 24 h. Saturated aqueous potassium hydrogen sulphate (pH 2, 30 mL) and ethyl acetate (50 mL) were added and the suspension filtered through hyflo-supercel washing through with ethyl acetate. The layers were separated. And the aqueous phase was extracted with ethyl acetate (3 x 100 mL). The combined organic layers were washed with brine (250 mL), dried over sodium sulfate,

filtered and evaporated to give a colorless gum. The residue was azeotroped with toluene (3 x 100 mL) to give the title compound (816 mg, 96%) as a white solid.




4e (439 mg, 1.00 mmol) was suspended in acetonitrile (35 mL) and (R)-1-(3-vinyl-phenyl)-ethylamine hydrochloride (202 mg, 1.10 mmol) was added followed by *N,N*-diisopropylethylamine (700 μ L, 5.00 mmol) and the mixture cooled in an ice-water bath before addition of 2-(1H-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate methanaminium (532 mg, 1.40 mmol). The reaction mixture was stirred and allowed to warm to RT over 72 h. To the reaction mixture was added 2 M hydrochloric acid (20 mL) and the mixture concentrated *in vacuo* to give a residue. The aqueous layer was extracted with ethyl acetate (3 x 100 mL) and the combined ethyl acetate layers were washed with brine followed by sodium hydrogen carbonate (100 mL) then dried over sodium sulfate, filtered and concentrated *in vacuo* to give a crude residue. The residue was purified by silica gel chromatography using *iso*-hexanes/ethyl acetate 50:50 then neat ethyl acetate to give the title compound (402 mg, 71%).



To a solution of **4f** (402 mg, 0.71 mmol) in 1,2-dichloroethane (210 mL) was added Hoveyda-Grubbs 2nd generation catalyst (45 mg, 0.07 mmol) and the reaction mixture heated at 80°C for 1 h. After this time the mixture was allowed to cool to RT and concentrated *in vacuo*. The residue was purified by silica gel chromatography using a gradient of ethyl acetate to ethyl acetate/acetone 75:25. Impure product (240 mg) was collected which was further purified by silica gel chromatography using ethyl acetate. After repurification, product (43 mg) was collected containing minor impurities. The impurities were removed via trituration from diethyl ether to


give the title compound (30 mg, 8%) as a white solid. ^1H NMR (300 MHz, CD_3OD) 0.89 (d, J = 6.7 Hz, 3H), 0.92 (d, J = 6.7 Hz, 3H), 0.96 (s, 3H), 0.97 (s, 3H), 1.35 (d, J = 6.9 Hz, 3H), 1.50 (d, J = 7.1 Hz, 3H), 1.59-1.83 (m, 3H), 1.85-2.00 (m, 3H), 2.09 (d, J = 7.1 Hz, 2H), 2.15-2.25 (m, 1H), 2.31-2.44 (m, 1H), 2.78 (td, J = 2.7, 12.7 Hz, 1H), 3.40-3.72 (m, 2H), 4.05-4.17 (m, 1H), 4.34 (br d, J = 13.2 Hz, 1H), 5.05 (q, J = 7.1 Hz, 1H), 5.35 (q, J = 7.1 Hz, 1H), 6.19-6.32 (m, 1H), 6.37 (d, J = 15.8 Hz, 1H), 7.15 (d, J = 7.6 Hz, 1H), 7.21-7.29 (m, 2H), 7.37 (d, J = 7.8 Hz, 1H). LCMS (m/z) 540.3 $[\text{M}+\text{H}]$, Tr = 2.41 min.

EXAMPLE 5




A solution of 6-bromo-3-chloroisoquinoline (8.0 g, 33 mmol) and tributyl-(1-ethoxyvinyl)-tin (14.88 g, 14 mL, 41.2 mmol) in toluene (100 mL) was degassed with nitrogen for 30 min. Bis(triphenylphosphine)palladium(II) dichloride (1.16 g, 1.65 mmol, 5 mol%) was added and the reaction mixture was heated at 60°C for 20 h. The reaction mixture was cooled to RT, the mixture was filtered and the filtrate was evaporated. The residue was purified by silica gel chromatography using a gradient of *iso*-hexanes/ethyl acetate 20:1 to 10:1 to afford the title compound (7.1 g, 92%) as a yellow solid.

15




A solution of **5a** (7.1 g, 30 mmol) in 1,4-dioxane (60 mL) and 2 M hydrochloric acid (30 mL) was stirred at RT for 30 min. The majority of the solvent was evaporated and the residue was partitioned between ethyl acetate and water. The organic extracts were combined, washed with water and brine, dried over sodium sulfate, filtered and evaporated. The residue was triturated with 5% ether in *iso*-hexanes and the resulting solid was collected and dried to afford the title compound (6.0 g, 97%) as a white solid.

20




Compound 5c

A solution of **5b** (1.72 g, 8.3 mmol) in tetrahydrofuran (40 mL) was stirred under nitrogen. Titanium (IV) ethoxide (3.8 g, 3.45 mL, 16.6 mmol, tech. grade) was added followed by (R)-(+)-2-methyl-propanesulfinamide (1.11 g, 9.2 mmol) and the reaction mixture was stirred at 60°C under nitrogen for 18 h. Additional (R)-(+)-2-methyl-propanesulfinamide (190 mg, 1.67 mmol) was added and the reaction mixture was stirred at 65°C for further 2 h. The reaction mixture was cooled to RT and ethyl acetate and brine were added. The suspension was filtered through Celite and the filter pad was washed with ethyl acetate. The ethyl acetate layer was separated, washed with brine, dried over sodium sulfate, filtered and evaporated. The residue was purified by silica gel chromatography using a gradient of *iso*-hexanes/ethyl acetate 7:3 to 3:7 to afford the title compound (2.2 g, 86%) as a yellow solid.




Compound 5d


A mixture of (1S,2R)-(-)-*cis*-1-amino-2-indanol (60 mg, 0.4 mmol), dichloro (*p*-cymene) ruthenium(II) dimer (122 mg, 0.2 mmol) and powdered 4 Å molecular sieves (2 g) was suspended in anhydrous 2-propanol (9 mL) and stirred under nitrogen. The suspension was heated at 90°C for 20 min. The reaction mixture was cooled to 40°C and a solution of **5c** (1.23 g, 4 mmol) in 2-propanol (28 mL) was added followed by a solution of potassium *tert*-butoxide (122 mg, 1.1 mmol) in 2-propanol (10 mL). The reaction mixture was stirred for 2 h at 40°C and then allowed to cool. The mixture was poured directly onto a silica gel cartridge and eluted with ethyl acetate to give, after evaporation, the title compound (1.19 g, 96%) as a brown gum.



A mixture of **5d** (0.66 g, 2.11 mmol), tributyl(vinyl)tin (1.85 mL, 6.35 mmol) and palladium tetrakis(triphenylphosphine) (488 mg, 0.42 mmol) in 1,4-dioxane (10.5 mL) was capped in a microwave vial. The reaction mixture was irradiated and stirred at 160°C for 40 min in a microwave reactor. A second reaction was carried under identical scale and conditions and the reaction mixtures combined and evaporated. The residue was purified by silica gel chromatography using a gradient of 1:3 to 1:0 ethyl acetate /iso-hexanes to afford the title compound (1 g) as a brown gum. ¹H NMR (300 MHz, CDCl₃) δ 1.27 (s, 9H), 1.62 (d, *J* = 7.7 Hz, 3H), 3.53 (*br s*, 1H), 4.69-4.77 (m, 1H), 5.51 (dd, *J* = 10.7, 1.3 Hz, 1H), 6.39 (dd, *J* = 17.4, 1.3 Hz, 1H), 6.95 (dd, *J* = 17.2, 10.7 Hz, 1H), 7.55-7.58 (m, 2H), 7.75 (s, 1H), 7.95 (d, *J* = 8.3 Hz, 1H), 9.20 (s, 1H). LCMS (*m/z*) 303.0 [*M*+*H*], *Tr* = 1.48 min.




5e was suspended in 4 M hydrochloric acid in 1,4-dioxane (17 mL, 68 mmol) and methanol was added (34 mL). The reaction mixture was stirred for 90 min and then evaporated. The residue was passed through a SCX cartridge eluting with methanol and then methanolic ammonia. The basic fraction was collected and evaporated to give the title compound (530 mg, 63% over 2 steps) as a beige solid.



A solution of 1-((1*R*,5*S*)-10,10-dimethyl-3,3-dioxo-3λ⁶-thia-4-azatricyclo[5.2.1.0^{1,5}]dec-4-yl)-propan-1-one (6.0 g, 22.1 mmol) in anhydrous dichloromethane (24 mL) was prepared and *tert*-butyldimethylsilyl trifluoromethanesulfonate (5.0 mL, 22.1 mmol) was added, followed by anhydrous triethylamine (3.54 mL, 25.4 mmol). The reaction mixture was stirred at RT under a nitrogen atmosphere for 15 h. This gave a dark solution that was


evaporated to give an oil. The oil was dissolved in anhydrous dichloromethane (22 mL) and the solution was added dropwise to a solution of crotonaldehyde (3.66 mL, 44.2 mmol) and titanium tetrachloride (1 M in dichloromethane, 44.2 mL, 44.2 mmol) in dichloromethane (22 mL) at -78 °C under a nitrogen atmosphere. The reaction mixture was stirred at -78 °C for 1 hour, before
 5 addition of ammonium chloride solution (30 mL). The stirred mixture was allowed to warm to RT before separating the layers. The aqueous layer was extracted with dichloromethane (2 x 25 mL). The organic layers were combined, dried over sodium sulfate, filtered and evaporated to give a brown oil. The oil was purified by silica gel chromatography using *iso*-hexanes/ethyl acetate 4:1 to yield the title compound (6.7 g, 89%) as a colorless solid.



A solution of **5g** (4.15 g, 12.1 mmol) in anhydrous dichloromethane (80 mL) was prepared and 1,8-bis(dimethylamino)naphthalene (7.78 g, 36.3 mmol) was added followed by trimethyloxonium tetrafluoroborate (3.6 g, 24.2 mmol). The reaction mixture was stirred at RT for 3 h. The reaction mixture was treated with methanol (3 mL) and stirred for 5 min before
 15 adding hydrochloric acid (2 M, 200 mL) and ethyl acetate (250 mL). The mixture was filtered to remove an insoluble solid and the layers were separated. The aqueous layer was extracted with ethyl acetate (2 x 100 mL). The organic layers were combined and washed with brine, dried over sodium sulfate, filtered and evaporated to give the title compound (4.80 g, 100%) as a pale brown solid.




A solution of lithium hydroxide in water (2 M, 50 mL, 100 mmol) was added to a stirred solution of **5h** (4.80 g, 12.1 mmol) in tetrahydrofuran (130 mL). The reaction mixture was heated to 60°C for 15 h. The reaction mixture was cooled to RT, before partially evaporating and adding hydrochloric acid (2 M, 150 mL). The mixture was extracted with ethyl acetate (3 x 50 mL). The
 25 extract was dried over sodium sulfate, filtered and evaporated to give a brown oil (3.5 g). The oil was purified by silica gel chromatography using *iso*-hexanes/diethyl ether 1:1 to give the title compound (1.132 g, 59%) as a colorless oil.



Compound 5j


A cooled (0°C) solution of (S)-1-[(S)-2-((S)-2-*tert*-butoxycarbonylamino-3-methylbutyrylamino)-propionyl]-hexahydro-pyridazine-3-carboxylic acid 2,2,2-trichloro-ethyl ester (2.174 g, 4.088 mmol) in anhydrous dichloromethane (50 mL) was treated with trimethylsilyl trifluoromethanesulfonate (1.2 mL, 6.814 mmol). After 1 h at 0°C, the reaction mixture was
 5 treated with *N,N*-diisopropylethylamine (2.4 mL, 13.628 mmol) and the volatiles were removed *in vacuo* to afford the corresponding amine as a yellow foam. To this amine was added **5i** (539.0 mg, 3.407 mmol), *N,N*-diisopropylethylamine (2.4 mL, 13.628 mmol) and acetonitrile (50 mL). The reaction mixture was cooled to 0°C and treated with 2-(1*H*-7-azabenzotriazol-1-yl)-
 10 1,1,3,3-tetramethyl uronium hexafluorophosphate methanaminium (1.814 g, 4.770 mmol). After overnight stirring at RT the reaction was quenched with 1 M hydrochloric acid (100 mL). The aqueous layer was extracted with ethyl acetate (2x). The organics were combined, dried over sodium sulfate, filtered and the volatiles were removed *in vacuo*. The residue was purified by silica gel chromatography using a 50 g Isolute cartridge eluted with a continuous gradient of *iso*-
 15 hexanes/ethyl acetate 1:0 to 1:4 to afford the title compound (2.193 g, 93%) as a light yellow solid.



Compound 5k


A solution of **5j** (763.4 mg, 1.335 mmol) in tetrahydrofuran (25 mL) was subsequently treated with zinc powder (1.920 g, 29.365 mmol) and a solution of ammonium acetate (1.543 g, 20.025 mmol) in water (5 mL). After overnight stirring the reaction was filtered through Celite and quenched with 2 M hydrochloric acid. The aqueous layer was extracted with ethyl acetate (2x). The organics were combined, dried over sodium sulfate, filtered and the volatiles were

removed *in vacuo*. Residual acetic acid was azeotroped off with toluene to provide the title compound (566.4 mg, 96%) as a light orange solid.



Compound 5I

A solution of **5k** (246 mg, 0.6 mmol), (R)-1-(3-vinyl-isoquinolin-6-yl)-ethylamine dihydrochloride (162 mg, 0.6 mmol) and *N,N*-diisopropylethylamine (387 mg, 0.52 mL, 3 mmol) in acetonitrile (20 mL) was stirred at RT under nitrogen. 2-(1H-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyl uronium hexafluorophosphate methanaminium (320 mg, 0.84 mmol) was added and the reaction mixture was stirred at RT for 4 h. The solvent was evaporated. The residue was diluted with ethyl acetate and saturated sodium hydrogen carbonate solution. A small amount of methanol (5 mL) was added to the suspension to give two clear layers. The organic layer was separated, washed with water and brine, dried over anhydrous sodium sulfate, filtered and evaporated. The residue was purified by silica gel chromatography using methanol/dichloromethane 1:20. The residue was triturated with ether and the resulting solid was collected, washed with ether and dried to afford the title compound (238 mg, 64%) as a pale brown solid.




Compound 5

A solution of **5I** (91 mg, 0.15 mmol) in toluene (45 mL) was stirred at RT under nitrogen. Hoveyda-Grubbs 2nd generation catalyst (10 mg, 0.015 mmol) was added and the reaction mixture was heated at reflux under nitrogen for 2 h. Additional Hoveyda-Grubbs 2nd generation catalyst (10 mg, 0.015 mmol) was added and the reaction mixture was heated at reflux under nitrogen for 6 h. The reaction mixture was cooled to RT, silica gel was added and the reaction mixture was evaporated. The residue was purified by silica gel chromatography using a gradient of ethyl acetate/acetone 20:1 to 5:2. The residue was triturated with ether and the resulting solid

was collected, washed with ether/*iso*-hexanes (1:1) and dried to afford the title compound (23 mg, 26%) as a white solid. ¹H NMR (300 MHz, CDCl₃) δ 0.99 (d, *J* = 6.7 Hz, 3H), 1.04 (d, *J* = 6.7 Hz, 3H), 1.44 (d, *J* = 7.4 Hz, 3H), 1.53 (d, *J* = 6.9 Hz, 3H), 1.61 (d, *J* = 6.9 Hz, 3H), 1.68-2.20 (m, 5H), 2.62-2.74 (m, 2H), 3.35-3.43 (m, 1H), 3.44 (s, 3H), 3.72 (d, *J* = 12.5 Hz, 1H), 3.88-3.92 (m, 2H), 4.23 (dd, *J* = 8.9, 6.0 Hz, 1H), 4.55-4.60 (m, 1H), 5.30-5.37 (m, 1H), 5.80-5.90 (m, 1H), 6.38 (d, *J* = 8.9 Hz, 1H), 6.43 (d, *J* = 7.6 Hz, 1H), 6.93 (d, *J* = 16.3 Hz, 1H), 7.24 (dd, *J* = 16.3, 8.7 Hz, 1H), 7.45 (d, *J* = 8.7 Hz, 1H), 7.56 (d, *J* = 8.5 Hz, 1H), 7.69 (s, 1H), 7.93 (d, *J* = 8.5 Hz, 1H), 9.17 (s, 1H). LCMS (*m/z*) 579.3 [M+H], Tr = 1.40 min.


10

EXAMPLE 6

Compound 6


A solution of Compound 5 (11 mg, 0.019 mmol) in ethyl acetate (10 mL) containing 10% palladium on carbon (10 mg) was hydrogenated at RT and pressure for 3 h. The reaction mixture was filtered through filter aid and the filter pad was washed with ethyl acetate. The filtrate was evaporated and the residue was purified by silica gel chromatography using a gradient of ethyl acetate/acetone 10:1 to 2:1. The residue was triturated with ethyl acetate/ether (1:5) to afford the title compound (2.0 mg, 18%) as a white solid. ¹H NMR (300 MHz, CDCl₃) δ 0.93 (d, *J* = 6.7 Hz, 3H), 0.98 (d, *J* = 6.7 Hz, 3H), 1.39 (d, *J* = 6.9 Hz, 3H), 1.55-2.70 (m, 8H), 1.95-2.50 (m, 5H), 2.64-2.73 (m, 2H), 2.95-3.08 (m, 2H), 3.35-3.42 (m, 2H), 3.53 (s, 3H), 3.61 (d, *J* = 12.1 Hz, 1H), 4.06-4.11 (m, 1H), 4.50-4.57 (m, 1H), 5.25-5.32 (m, 1H), 5.64-5.70 (m, 1H), 6.39 (d, *J* = 8.3 Hz, 1H), 6.47-6.51 (m, 1H), 7.21-7.28 (m, 1H), 7.41 (d, *J* = 8.8 Hz, 1H), 7.62 (s, 1H), 7.69 (s, 1H), 7.92 (d, *J* = 8.8 Hz, 1H), 9.15 (s, 1H). LCMS (*m/z*) 581.3 [M+H], Tr = 1.38 min.

EXAMPLE 7




A mixture of 7-bromo-2-chloro-quinoline (1.46 g, 6.00 mmol) and tributyl(1-ethoxyvinyl)tin (2.71 g, 2.54 mL, 7.50 mmol) in toluene (21 mL) was degassed for 20 min.


- 5 Bis(triphenylphosphine)palladium(II) dichloride (427 mg, 0.60 mmol) was added and the reaction mixture stirred under nitrogen and heated at 80 °C for 19 h before allowing to cool. The volatiles were evaporated and the residue suspended in 1,4-dioxane (15 mL), 2 M aqueous hydrochloric acid (15 mL) was added and the reaction mixture stirred for 45 min and then evaporated to remove the volatiles. The residue was diluted with water and extracted with ethyl
- 10 acetate (2 x) and the combined organic extracts were washed with brine, dried over magnesium sulfate, filtered and evaporated. The product was purified by chromatography using silica gel doped with 10% w/w potassium carbonate eluting using a gradient of *iso*-hexanes/ethyl acetate 1:0 to 9:1 to afford the title compound (836 mg, 56 %) as a yellow solid.




- 15 Dichloro (*p*-cymene) ruthenium(II) dimer (12 mg, 0.019 mmol) and (1R,2R)-(-)-*N*-p-tosyl-1,2-diphenylethylenediamine (17 mg, 0.045 mmol) were suspended in degassed water (7.5 mL) and the mixture was degassed with nitrogen for 15 min. The mixture was stirred at 70°C under nitrogen for 90 min. The resulting turbid orange mixture was allowed to cool to RT. Solid **7a** (948 mg, 3.78 mmol) followed by degassed tetrahydrofuran (7.5 mL) and sodium formate (1.29 g,
- 20 18.9 mmol) were added and the reaction mixture was degassed with nitrogen for 5 min. The reaction mixture was vigorously stirred at 40°C for 3 h and allowed to cool. It was then diluted with ethyl acetate and the mixture washed with water (2 x). The aqueous washes were back-extracted with ethyl acetate and the combined organics washed with brine, dried over magnesium sulfate, filtered and evaporated. The residue was purified by silica gel
- 25 chromatography eluting with *iso*-hexanes/ethyl acetate 2:1 to afford the title compound (814 mg, 85%) as a purple solid.



To a mixture of **7b** (490 mg, 1.94 mmol), *N,N*-dicyclohexylmethylamine (416 mg, 457 μ L, 2.13 mmol) and *tert*-butyl 3-butenate (648 mg, 739 μ L, 4.56 mmol) in toluene (19 mL) was added *bis*(tri-*tert*-butylphosphine)palladium(0) (41 mg, 0.080 mmol) under nitrogen and the reaction mixture stirred and heated under reflux for 5 h then allowed to cool. The mixture was evaporated and then purified by silica gel chromatography using a gradient of *iso*-hexanes/ethyl acetate 4:1 to 3:2 to afford the title compound (367 mg, 60 %) as a yellow oil.




A solution of **1e** (804 mg, 1.51 mmol) in tetrahydrofuran (37.7 mL) was prepared and zinc powder (2.18 g, 33.3 mmol) was added followed by a solution of ammonium acetate (1.75 g, 22.7 mmol) in water (9.4 mL). The reaction mixture was stirred at RT for 72 h. The reaction was filtered through hyflo-supercel washing through with ethyl acetate and saturated aqueous potassium hydrogen sulfate. The mixture was treated with 1 M hydrochloric acid (3 mL) and the layers were separated. The aqueous layer was extracted with ethyl acetate (2 x 100 mL). The organic layers were combined, washed with brine, filtered and evaporated to give a colorless gum. The residue was azeotroped with toluene (3 x 200 mL) to give the title compound (605 mg, quantitative yield) as a white solid.



Compound 7e

To a stirred solution of **7d** (456 mg, 1.14 mmol) and (E)-4-[3-((R)-1-hydroxy-ethyl)-quinoxalin-6-yl]-but-3-enoic acid *tert*-butyl ester (358 mg, 1.14 mmol) in dichloromethane (22 mL) was added 1-hydroxybenzotriazole containing approx. 20% water (270 mg, 1.60 mmol) followed by 4-dimethylaminopyridine (139 mg, 1.14 mmol) and *N*-(3-dimethylaminopropyl)-*N*-ethylcarbodiimide hydrochloride (306 mg, 1.60 mmol). The reaction was stirred for 18 h and then diluted with dichloromethane, washed with saturated ammonium chloride solution (2x), dried over sodium sulfate, filtered and evaporated. The residue was purified by silica gel chromatography using a gradient of *iso*-hexanes/ethyl acetate 3:1 to 0:1 to afford the title compound (335 mg, 45 %) as a white foam.




Compound 7

To a stirred solution of **7e** (309 mg, 0.444 mmol) in dichloromethane (4.5 mL) at 0°C under nitrogen was added trimethylsilyl trifluoromethanesulfonate (346 mg, 359 μ L, 1.56 mmol) and the reaction mixture was allowed to warm to RT over 2.5 h. *N,N*-diisopropylethylamine (164 mg, 221 μ L, 1.27 mmol) was added and the reaction mixture stirred for a further 10 min, evaporated and then suspended in acetonitrile (45 mL). The stirred mixture was cooled to 0°C and 2-(1H-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyl uronium hexafluorophosphate methanaminium (236 mg, 0.622 mmol) and *N,N*-diisopropylethylamine (229 mg, 309 μ L, 1.77 mmol) added. After 90 min the reaction was quenched with a saturated ammonium chloride solution and the mixture evaporated to remove organic volatiles. The residue was diluted with


dichloromethane and the organic layer separated and washed with saturated sodium bicarbonate (2x) and brine then dried over sodium sulfate, filtered and evaporated. The residue was purified by silica gel chromatography eluting with a gradient of ethyl acetate/acetone 1:0 to 9:1. The residue was further purified by reverse phase preparative HPLC using

5 acetonitrile/water 3:7 to afford the title compound (7.6 mg, 3% over 2 steps) as a white solid. ^1H NMR (300 MHz, CD_3OD) δ 1.00 (d, $J = 6.7$ Hz, 3H), 1.01 (d, $J = 6.7$ Hz, 3H), 1.62 (d, $J = 7.1$ Hz, 3H), 1.66-1.75 (m, 2H), 1.77 (d, $J = 6.9$ Hz, 3H), 1.91-2.05 (m, 3H), 2.72-2.82 (m, 1H), 2.98-3.08 (m, 1H), 3.38-3.41 (m, 1H), 3.78-3.84 (m, 1H), 4.25 (d, $J = 10.5$ Hz, 1H), 4.41 (br d, $J = 11.3$ Hz, 1H), 5.68 (q, $J = 7.1$ Hz, 1H), 6.09 (q, $J = 6.9$ Hz, 1H), 6.47 (d, $J = 16.3$ Hz, 1H), 6.55-6.63 (m, 1H), 7.68 (s, 1H), 7.99 (s, 2H), 8.84 (s, 1H). LCMS (m/z) 523.2 $[\text{M}+\text{H}]$, Tr = 1.75 min.

EXAMPLES 8 AND 9




Dichloro (*p*-cymene) ruthenium(II) dimer (3 mg, 0.005 mmol) and (1*R*,2*R*)-(-)-*N*-*p*-tosyl-1,2-diphenylethylenediamine (4.4 mg, 0.012 mmol) were suspended in degassed water (2 mL) and the mixture was degassed with nitrogen for 15 min. The mixture was stirred at 70°C under nitrogen for 90 min. The resulting yellow solution was cooled to RT. 1-(3-Chloro-isoquinolin-6-yl)-ethanone (206 mg, 1 mmol), sodium formate (340 mg, 5 mmol) and degassed tetrahydrofuran (1 mL) were added and the reaction mixture was degassed with nitrogen for 5 min. The reaction mixture was vigorously stirred at 40°C for 2.5 h. The reaction mixture was cooled to RT and was extracted with ethyl acetate. The organic layer was separated, washed with water and brine, dried over sodium sulfate, filtered and evaporated. The residue was purified by silica gel chromatography using a gradient of *iso*-hexanes/ethyl acetate 4:1 to 2:1 to afford the title compound (193 mg, 92%) as a white solid.




1,4-Dioxane (5 mL) was degassed with nitrogen, **8a** (208 mg, 1 mmol), tributyl(vinyl)tin (951 mg, 0.9 mL, 3 mmol) and bis(triphenylphosphine)palladium(II) dichloride (70 mg, 0.1 mmol)

were added and the reaction mixture was heated at 150°C in a microwave reactor for 1 h. Additional tributyl(vinyl)tin (0.3 mL, 1 mmol) and bis(triphenylphosphine)palladium(II) dichloride (70 mg, 0.1 mmol) were added and the reaction mixture was heated at 150°C in a microwave reactor for 1 h. The reaction mixture was cooled to RT and the mixture was filtered through filter aid and the filter pad was washed with ethyl acetate. The filtrate was evaporated and the residue was purified by silica gel chromatography using a gradient of *iso*-hexanes/ethyl acetate 4:1 to 2:1 followed by silica gel chromatography using *iso*-hexanes/ethyl acetate 3:1 to afford the title compound (100 mg, 50%) as a white solid.




Compound 8c

A solution of **1e** (10.6 g, 20 mmol) in dichloromethane (300 mL) was stirred at 0°C under nitrogen. Trimethylsilyl trifluoromethanesulfonate (6.66 g, 5.4 mL, 30 mmol) was added dropwise and the reaction mixture was stirred at 0°C for 1 h. Cold saturated sodium hydrogen carbonate solution (200 mL) was added and the reaction mixture was stirred at 0°C for 15 min. The organic layer was separated, washed with saturated sodium hydrogen carbonate solution and brine, dried over sodium sulfate, filtered and evaporated to afford (S)-1-[(S)-2-((S)-2-amino-3-methyl-butyrylamino)-propionyl]-hexahydro-pyridazine-3-carboxylic acid 2,2,2-trichloro-ethyl ester (20 mmol), which was used crude in the next step. A solution of (S)-1-[(S)-2-((S)-2-amino-3-methyl-butyrylamino)-propionyl]-hexahydro-pyridazine-3-carboxylic acid 2,2,2-trichloro-ethyl ester (20 mmol) in acetonitrile (240 mL) was stirred at 0°C under nitrogen. (E)-Pent-3-enoic acid (2.20 g, 2.2 mL, 22 mmol) and 1-hydroxybenzotriazole hydrate (3.82 g, 20 mmol, wetted with not less than 20 wt.% water) and *N*-(3-dimethylaminopropyl)-*N'*-ethylcarbodiimide hydrochloride (5.38 g, 28 mmol) was added and the reaction mixture was stirred at 0°C for 15 min and then at RT for 20 h. The solvent was evaporated and the residue was partitioned between ethyl acetate and water. The organic extracts were combined, washed with water and brine, dried over anhydrous sodium sulfate, filtered and evaporated. The residue was purified by silica gel chromatography using *iso*-hexanes/ethyl acetate 3:1 to neat ethyl acetate to afford the title compound (9.0 g, 88%) as a white solid.




A solution of **8c** (9.0 g, 17.5 mmol) in tetrahydrofuran (300 mL) was stirred at RT under nitrogen. Zinc powder (25.0 g, 385 mmol) was added followed by a solution of ammonium acetate (20.2 g, 263 mmol) in water (200 mL). The reaction mixture was stirred at RT under nitrogen for 18 h. The reaction mixture was filtered through Celite and the filter pad was washed with water (100 mL) and ethyl acetate (200 mL). The organic layer was separated and the solvent was evaporated to ca. 100 mL and the solution was extracted with water (100 mL). The aqueous layers were combined, saturated ammonium chloride solution (150 mL) was added and the solution was acidified to pH 1 with 2 M aqueous hydrochloric acid. The solution was extracted with ethyl acetate and the organic extracts were combined, washed with brine, dried over anhydrous magnesium sulfate, filtered and evaporated. The residue was co-evaporated with toluene (3 x) to afford (S)-1-[(S)-2-[(S)-3-methyl-2-((E)-pent-3-enoylamino)-butyrylamino]-propionyl]-hexahydro-pyridazine-3-carboxylic acid (5.7 g, 85%) as a white solid which was used in the next reaction. A solution of (S)-1-[(S)-2-[(S)-3-methyl-2-((E)-pent-3-enoylamino)-butyrylamino]-propionyl]-hexahydro-pyridazine-3-carboxylic acid (5.16 g, 13.5 mmol) in dichloromethane (280 mL) and tetrahydrofuran (20 mL) was stirred at RT under nitrogen. (R)-1-(3-Vinyl-isoquinolin-6-yl)-ethanol (2.69 g, 13.5 mmol) was added followed by *N*-(3-dimethylaminopropyl)-*N'*-ethylcarbodiimide hydrochloride (3.63 g, 18.9 mmol) and 4-(dimethylamino)pyridine (1.64 g, 13.5 mmol) and the reaction mixture was stirred at RT for 6 h. Dichloromethane (200 mL) was added and the solution was washed with aqueous citric acid solution (pH 2-3). The organic layer was separated, washed with water and brine, dried over anhydrous sodium sulfate, filtered and evaporated. The residue was purified by silica gel chromatography using a gradient of *iso*-hexanes/ethyl acetate 3:1 to neat ethyl acetate followed by silica gel chromatography using *iso*-hexanes/ethyl acetate 1:8 to afford the title compound (3.91 g, 51%) as a white solid.



A solution of **8d** (1.13 g, 2 mmol) in toluene (600 mL) was stirred at RT under nitrogen for 15 min. Hoveyda-Grubbs 2nd generation catalyst (125 mg, 0.2 mmol) was added and the reaction mixture was heated at reflux under nitrogen for 30 min. Additional Hoveyda-Grubbs 2nd generation catalyst (125 mg, 0.2 mmol) was added and the reaction mixture was heated at reflux for 30 min. The reaction mixture was cooled to RT. The majority of the solvent was evaporated, silica gel was added and the solvent was evaporated. The residue was purified by silica gel chromatography using a gradient of ethyl acetate/acetone 9:1 to 3:2 followed by silica gel chromatography using ethyl acetate/methanol 40:1. The residue was triturated with ethyl acetate/ether (1:4) and the resulting solid was collected, washed with ethyl acetate/ether (1:4) and dried to afford a ~10:1 mixture of Compound 9 and Compound 8, as a pale brown solid (245 mg). A sample of the mixture was purified by reverse phase preparative HPLC to afford Compound 8 (4 mg) as a white solid. ¹H NMR (300 MHz, CD₃OD) δ 1.01 (d, J = 6.5 Hz, 3H), 1.02 (d, J = 6.5 Hz, 3H), 1.37 (d, J = 7.1 Hz, 3H), 1.65 (d, J = 6.5 Hz, 3H), 1.70-2.15 (m, 5H), 2.95-3.05 (m, 1H), 3.70-3.76 (m, 1H), 4.03-4.10 (m, 2H), 4.12 (d, J = 8.3 Hz, 1H), 5.64 (q, J = 7.1 Hz, 1H), 6.10-6.20 (m, 2H), 6.98 (d, J = 11.3 Hz, 1H), 7.65 (d, J = 8.5 Hz, 1H), 7.93 (s, 1H), 8.05-8.10 (m, 2H), 9.21 (s, 1H). LCMS (m/z) 522.3 [M+H], Tr = 1.25 min.


Compound 9, ¹H NMR (300 MHz, CDCl₃) δ 0.97 (d, J = 6.9 Hz, 3H), 0.99 (d, J = 6.7 Hz, 3H), 1.53 (d, J = 6.9 Hz, 3H), 1.71 (d, J = 6.9 Hz, 3H), 1.75-2.70 (m, 6H), 3.20-3.40 (m, 2H), 3.63-3.77 (m, 2H), 4.23-4.29 (m, 1H), 4.53-5.57 (m, 1H), 5.65-5.76 (m, 1H), 6.04 (q, J = 6.7 Hz, 1H), 6.38-6.53 (m, 3H), 6.72 (d, J = 16.3 Hz, 1H), 7.39 (d, J = 8.3 Hz, 1H), 7.55 (s, 1H), 7.72 (s, 1H), 7.92 (d, J = 8.3 Hz, 1H), 9.13 (s, 1H). LCMS (m/z) 522.0 [M+H], Tr = 1.40 min.

EXAMPLE 10




Compound 10a

10a was prepared in the same manner as **4d** using 3-butenic acid instead of 4,4-dimethyl-hept-6-enoic acid in 66% yield.




Compound 10b

10b was prepared in the same manner as **4e** using (S)-1-[(S)-2-((S)-2-but-3-enoylamino-3-methyl-but-3-enoylamino)-propionyl]-hexahydro-pyridazine-3-carboxylic acid 2,2,2-trichloro-ethyl ester instead of (S)-1-[(S)-2-[(S)-2-(4,4-dimethyl-hept-6-enoyloxy)-3-methyl-but-3-enoylamino]-propionyl]-hexahydro-pyridazine-3-carboxylic acid 2,2,2-trichloro-ethyl ester in 84% yield.




Compound 10c

10c was prepared in the same manner as **8d** using (S)-1-[(S)-2-((S)-2-but-3-enoylamino-3-methyl-but-3-enoylamino)-propionyl]-hexahydro-pyridazine-3-carboxylic acid instead of (S)-1-[(S)-2-[(S)-3-methyl-2-((E)-pent-3-enoylamino)but-3-enoylamino]-propionyl]-hexahydro-pyridazine-3-carboxylic acid in 84% yield.




Compound 9 was prepared in the same manner as Compound 5 using (S)-1-[(S)-2-((S)-2-but-3-enoylamino-3-methyl-butyrylamino)-propionyl]-hexahydro-pyridazine-3-carboxylic acid (R)-1-(3-vinyl-isoquinolin-6-yl)-ethyl ester instead of (S)-1-[(S)-2-[(S)-2-((E)-(2R,3R)-3-methoxy-2-methyl-hex-4-enoylamino)-3-methyl-butyrylamino]-propionyl]-hexahydro-pyridazine-3-carboxylic acid [(R)-1-(3-vinyl-isoquinolin-6-yl)-ethyl]-amide in 9% yield.



A solution of Compound 9 (16 mg, 0.03 mmol) in ethyl acetate (5 mL) and tetrahydrofuran (2 mL) containing 10% palladium on carbon (15 mg) was hydrogenated at RT and pressure for 3 h. The reaction mixture was filtered through filter aid and the filter pad was washed with ethyl acetate. The filtrate was evaporated and the residue was triturated with ether (2 x 2 mL) and the resulting solid was dried to afford the title compound (7.6 mg, 48%) as an off-white solid. ¹H NMR (300 MHz, CDCl₃) δ 0.90 (d, *J* = 6.7 Hz, 3H), 0.95 (d, *J* = 6.5 Hz, 3H), 1.50 (d, *J* = 7.1 Hz, 3H), 1.68 (d, *J* = 6.5 Hz, 3H), 1.60-2.40 (m, 9H), 2.70-3.10 (m, 3H), 3.68-3.75 (m, 1H), 3.90 (d, *J* = 11.8 Hz, 1H), 4.32-4.37 (m, 1H), 4.44-4.49 (m, 1H), 5.83-5.92 (m, 1H), 6.05-6.11 (m, 2H), 6.33 (d, *J* = 7.8 Hz, 1H), 7.34 (s, 1H), 7.36 (d, *J* = 8.3 Hz, 1H), 7.80 (s, 1H), 7.93 (d, *J* = 8.3 Hz, 1H), 9.17 (s, 1H). LCMS (*m/z*) 524.3 [*M*+H], Tr = 0.59 min.

EXAMPLES 11 AND 12




Compounds 11a Compound 12b

To a stirred solution of **1e** (10.6 g, 20.0 mmol) in dichloromethane (400 mL), at 0°C
 5 under nitrogen, was added trimethylsilyl trifluoromethanesulfonate (6.67 g, 5.43 mL, 30.0 mmol) and the reaction mixture stirred at 0°C for 2 h. *N,N*-Diisopropylethylamine (10.3 g, 13.9 mL, 80.0 mmol) was added and the mixture allowed to warm to ambient temperature. The volatiles were evaporated and the residue suspended in acetonitrile (250 mL). The stirred mixture was cooled to 0°C under a blanket of nitrogen and then *N,N*-diisopropylethylamine (10.3 g, 13.9 mL, 80
 10 mmol) and 3-butenic acid (1.89 g, 1.86 mL, 4.40 mmol) added, followed by 2-(1H-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyl uronium hexafluorophosphate methanaminium (10.6 g, 28.0 mmol), portionwise. The reaction mixture was allowed to warm to ambient temperature and stirred for 18 h before evaporating. The residue was diluted with ethyl acetate and then washed successively with saturated sodium bicarbonate solution, water, 2 M hydrochloric acid, water
 15 then brine, dried over magnesium sulfate, filtered and evaporated. The residue was purified by silica gel chromatography using a gradient of *iso*-hexanes/ethyl acetate 1:1 to 0:1 to afford a ~1:1 mixture of (S)-1-[(S)-2-((S)-2-but-3-enoylamino-3-methyl-butyrylamino)-propionyl]-hexahydro-pyridazine-3-carboxylic acid 2,2,2-trichloro-ethyl ester and (S)-1-[(S)-2-((E)-but-2-enoylamino)-3-methyl-butyrylamino]-propionyl]-hexahydro-pyridazine-3-carboxylic acid
 20 2,2,2-trichloro-ethyl ester as an orange foam (7.23 g, 72%).

To a stirred solution of a mixture of (S)-1-[(S)-2-((S)-2-but-3-enoylamino-3-methyl-butyrylamino)-propionyl]-hexahydro-pyridazine-3-carboxylic acid 2,2,2-trichloro-ethyl ester and (S)-1-[(S)-2-((E)-but-2-enoylamino)-3-methyl-butyrylamino]-propionyl]-hexahydro-pyridazine-3-carboxylic acid 2,2,2-trichloro-ethyl ester (~1:1, 0.99 g, 2.00 mmol) in
 25 tetrahydrofuran (40 mL) was added zinc powder (2.86 g, 44.0 mmol) and a solution of ammonium acetate (2.31 g, 30.0 mmol) in water (25 mL). The reaction mixture was allowed to warm to ambient temperature and stirred for 18 h, then diluted with ethyl acetate and the mixture filtered. From the filtrate the aqueous layer was separated and diluted with an equal amount of saturated ammonium chloride solution and then acidified to pH 1 with 2 M

hydrochloric acid. The aqueous layer was extracted with ethyl acetate (2 x) and the combined organic layers washed with brine, dried over magnesium sulfate, filtered and evaporated. The residue was azeotroped with toluene (3 x) to give a mixture of the title compounds (~1:1, 466 mg, 63%) as a yellow foam.




A mixture of **11a** and **12b** (~1:1, 250 mg, 0.68 mmol) in acetonitrile (20 mL) was stirred at RT under nitrogen. (R)-1-(3-vinyl-isoquinolin-6-yl)-ethylamine dihydrochloride (136 mg, 0.5 mmol) and *N,N*-diisopropylethylamine (323 mg, 0.44 mL, 2.5 mmol) was added followed by 2-(1H-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyl uronium hexafluorophosphate methanaminium (266 mg, 0.7 mmol) and the reaction mixture was stirred at RT for 18 h. The solvent was evaporated. The residue was diluted with ethyl acetate and the solution was washed with saturated sodium hydrogen carbonate solution, water and brine. The organic extract was dried over anhydrous magnesium sulfate, filtered and evaporated. The residue was purified by silica gel chromatography using a gradient of ethyl acetate to methanol/ethyl acetate 1:5 to afford a ~1:1 mixture of (S)-1-[(S)-2-((S)-2-but-3-enoylamino-3-methyl-butyrylamino)-propionyl]-hexahydro-pyridazine-3-carboxylic acid [(R)-1-(3-vinyl-isoquinolin-6-yl)-ethyl]-amide and (S)-1-[(S)-2-[(S)-2-((E)-but-2-enoylamino)-3-methyl-butyrylamino]-propionyl]-hexahydro-pyridazine-3-carboxylic acid [(R)-1-(3-vinyl-isoquinolin-6-yl)-ethyl]-amide as a white solid (153 mg, 56%) which was used in the next step without further purification.

A solution of a mixture of (S)-1-[(S)-2-((S)-2-but-3-enoylamino-3-methyl-butyrylamino)-propionyl]-hexahydro-pyridazine-3-carboxylic acid [(R)-1-(3-vinyl-isoquinolin-6-yl)-ethyl]-amide and (S)-1-[(S)-2-[(S)-2-((E)-but-2-enoylamino)-3-methyl-butyrylamino]-propionyl]-hexahydro-pyridazine-3-carboxylic acid [(R)-1-(3-vinyl-isoquinolin-6-yl)-ethyl]-amide (~1:1, 150 mg, 0.27 mmol) in toluene (70 mL) was stirred at RT under nitrogen. Hoveyda-Grubbs 2nd generation catalyst (17 mg, 0.027 mmol) was added and the reaction mixture was heated at reflux under nitrogen for 90 min. Additional Hoveyda-Grubbs 2nd generation catalyst (17 mg, 0.027 mmol) was added and the reaction mixture was heated at reflux under nitrogen for 90 min. The reaction mixture was cooled to RT, silica gel was added and the solvent was evaporated. The

residue was purified by silica gel chromatography using a gradient of ethyl acetate to ethyl acetate/acetone 1:1. The residue was further purified by preparative thin layer chromatography using methanol/ethyl acetate 1:4 followed by silica gel chromatography using dichloromethane/methanol 20:1 to give the title compounds. Compound 11 (1.3 mg, 1%) as a white solid: ^1H NMR (300 MHz, CD_3OD) δ 1.01 (d, J = 6.7 Hz, 3H), 1.02 (d, J = 6.7 Hz, 3H), 1.61 (d, J = 7.1 Hz, 3H), 1.66 (d, J = 7.1 Hz, 3H), 1.80-2.05 (m, 5H), 2.71-2.82 (m, 1H), 2.99-3.05 (m, 1H), 3.32-3.43 (m, 1H), 3.64-3.75 (m, 1H), 4.27 (d, J = 10.1 Hz, 1H), 4.40-4.53 (m, 2H), 5.12 (q, J = 7.1 Hz, 1H), 5.62 (q, J = 7.1 Hz, 1H), 6.47-6.56 (m, 1H), 6.62 (d, J = 15.8 Hz, 1H), 7.43 (s, 1H), 7.61 (dd, J = 8.5, 1.6 Hz, 1H), 7.74 (s, 1H), 8.05 (d, J = 8.5 Hz, 1H), 9.13 (s, 1H). LCMS (m/z) 521.3 [M+H], Tr = 0.97 min.

Compound 12 (1.7 mg, 1.2%) as a white solid: ^1H NMR (300 MHz, CD_3OD) δ 0.98 (d, J = 6.7 Hz, 3H), 1.03 (d, J = 6.7 Hz, 3H), 1.47 (d, J = 7.1 Hz, 3H), 1.60 (d, J = 6.9 Hz, 3H), 1.65-2.16 (m, 5H), 2.81-2.89 (m, 1H), 3.61-3.69 (m, 1H), 4.04 (d, J = 6.9 Hz, 1H), 4.30-4.36 (m, 1H), 4.75 (d, J = 11.6 Hz, 1H), 5.23 (q, J = 7.1 Hz, 1H), 5.90 (q, J = 6.9 Hz, 1H), 6.36 (d, J = 13.1 Hz, 1H), 6.92 (d, J = 13.1 Hz, 1H), 7.65 (d, J = 8.5 Hz, 1H), 7.88 (s, 1H), 8.05 (d, J = 8.5 Hz, 1H), 8.46 (s, 1H), 9.16 (s, 1H). LCMS (m/z) 507.2 [M+H], Tr = 0.95 min.


EXAMPLE 13




Compound 13a

A suspension of (S)-1-(7-bromoquinolin-2-yl)-2,2,2-trifluoroethylamine hydrochloride (Asiba Pharmatech, Edison, NJ, USA, 397 mg, 1.16 mmol) in dichloromethane (10 mL) was cooled using an ice bath. Triethylamine (985 μL , 3.48 mmol) was added and the reaction was stirred until a homogenous solution was observed. Di-*tert*-butyl dicarbonate (380 mg, 1.74 mmol) in dichloromethane (5 mL) was then added and the reaction was left to stir overnight. Di-*tert*-butyl dicarbonate (127 mg, 0.58 mmol) was added and the reaction was stirred for a further 6 h. Di-*tert*-butyl dicarbonate (253 mg, 1.16 mmol) was added along with 4 Å molecular sieves and the reaction was left to stir overnight. The reaction was washed with water and brine and the organics were dried over anhydrous sodium sulfate, filtered and concentrated. The residue

was purified by silica gel chromatography using a stepwise gradient from 1:0 to 7:3 *iso*-hexanes/ethyl acetate to afford the title compound (343 mg, 73%) as an orange solid.




To a 3-necked flask, fitted with stirrer bar and thermometer, under an atmosphere of nitrogen and containing magnesium turnings (3.02 g, 0.124 mol) was added anhydrous tetrahydrofuran (105 mL). A single crystal of iodine was added to the reaction followed by 1-chloro-3-methyl-but-2-ene (2 mL, 17.75 mmol). The reaction was stirred for 20 min until all color had left the solution. 1-chloro-3-methyl-but-2-ene (5.57 mL, 49.45 mmol) was added. A gradual temperature increase was observed until the reaction had reached reflux. The reaction was left to stir for 1 hour, allowing the reaction to return to RT. The reaction was then transferred *via* cannula to a flask containing solid carbon dioxide (50 g). The reaction was then stirred for 2 h. A cooling ice bath was added and the reaction was quenched with 2 M hydrochloric acid. The reaction mixture was then extracted with diethyl ether. The organics were then dried over anhydrous sodium sulfate, filtered and concentrated. The residue was purified by silica gel chromatography using a stepwise gradient from 1:0 to 9:1 *iso*-hexanes/ethyl acetate to afford the title compound (2.69 g, 35%) as a pale yellow oil. ¹H NMR (300 MHz, CDCl₃) δ 1.35 (s, 6H), 5.10-5.21 (m, 2H), 6.07 (dd, *J* = 17.6, 10.5 Hz, 1H).




13a (343 mg, 0.846 mmol) and **13b** (106 mg, 0.931 mmol) were placed in a microwave vessel and dissolved in acetonitrile (3 mL). Palladium(II) acetate (19 mg, 0.0846 mmol), tri(*o*-tolyl)phosphine (51 mg, 0.169 mmol) and triethylamine (236 μL, 1.69 mmol) were added and the vessel was sealed before being irradiated in the microwave for 30 min, using fixed hold time, on high absorption at 100°C. The solvent was removed and the residue was taken up in a mixture of water and ethyl acetate. The phases were separated and the aqueous was extracted with ethyl acetate. The combined organics were then dried over anhydrous sodium sulfate, filtered and concentrated. The residue was purified by silica gel chromatography using a stepwise gradient of *iso*-hexanes/ethyl acetate from 1:0 to 6:4. The isolated material was subjected to a

second purification using the same conditions to afford the title compound (142 mg, 38%) as a pale yellow oil.




Compound 13d

- (S)-1-[(S)-2-((S)-2-Amino-3-methyl-butrylamino)-propionyl]-hexahydro-pyridazine-3-carboxylic acid ethyl ester (210 mg, 0.486 mmol) and **13c** (142 mg, 0.324 mmol) were dissolved in *N,N*-dimethylformamide (5 mL), under an atmosphere of nitrogen, and cooled using an ice bath. 2-(1H-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyl uronium hexafluorophosphate methanaminium (203 mg, 0.535 mmol) and *N,N*-diisopropylethylamine (423 μ L, 2.43 mmol) were then added and the reaction was allowed to slowly warm to RT and left to stir overnight.
- The reaction was diluted with water and extracted with ethyl acetate. The combined organics were dried over anhydrous sodium sulfate, filtered and concentrated. The residue was purified by silica gel chromatography using a stepwise gradient of *iso*-hexanes/ethyl acetate from 1:0 to 0:1 to afford the product contaminated with residual solvent. Toluene was added and the solution was concentrated to afford the title compound (79 mg, 29%) as an orange solid.




Compound 13e

- 13d** (79 mg, 0.0925 mmol) was dissolved in tetrahydrofuran (2 mL) and cooled using an ice bath. Methanol (1 mL) and water (1 mL) were then added followed by lithium hydroxide monohydrate (15 mg, 0.37 mmol). The reaction was then left to stir for 1 hour. 1 M hydrochloric acid was added until the solution was pH 2. The solvent was removed and the resultant solid was sequentially azeotroped with methanol, then acetonitrile and finally toluene to afford the title compound (67 mg, 100%) as a yellow solid.




13e (66 mg, 0.0916 mmol) was dissolved in 4 M hydrochloric acid in 1,4-dioxane (2 mL) and left to stir for 30 min. The solvent was removed and the resultant solid was triturated with diethyl ether and dried to afford the title compound (50 mg, 83%) as a pale yellow solid.




13f (50 mg, 0.0761 mmol) was dissolved in dichloromethane (76 mL), under an atmosphere of nitrogen and cooled using an ice bath. 2-(1H-7-Azabenzotriazol-1-yl)-1,1,3,3-tetramethyl uronium hexafluorophosphate methanaminium (58 mg, 0.152 mmol) and *N,N*-diisopropylethylamine (53 μ L, 0.304 mmol) were then added and the reaction was allowed to slowly warm to RT and left to stir overnight. The solvent was removed and the residue was purified by silica gel chromatography using 7:3 *iso*-hexanes/acetone. The residue was re-purified by silica gel chromatography using a stepwise gradient of *iso*-hexanes/acetone from 1:0 to 1:1. The residue was then eluted through an HPLC system fitted with a Phenomenex Gemini 10 μ 110A, 250 x 21.2 mm column using a continuous gradient of acetonitrile/water from 1:4 to 1:0 flow at 20 mL/min to afford the title compound (7 mg, 15%) as a solid. ^1H NMR (300 MHz, CDCl_3) δ 0.86-1.08 (m, 6H), 1.25-1.54 (m, 8H), 1.57-1.81 (m, 5H), 1.88-2.00 (m, 1H), 2.37-2.80 (m, 1H), 3.32-3.65 (m, 1H), 3.69-3.97 (m, 1H), 4.22-4.39 (m, 1H), 4.50-4.68 (m, 1H), 5.74-6.05 (m, 2H), 6.22-6.39 (m, 1H), 6.44-6.54 (m, 1H), 6.63-6.76 (m, 1H), 7.45 (*app* t, J = 8.5 Hz, 1H), 7.56 (*app* t, J = 7.6 Hz, 1H), 7.81 (d, J = 8.3 Hz, 1H), 8.19 (s, 1H), 8.76-9.09 (m, 1H). LCMS (m/z) 603.1 [$\text{M}+\text{H}$], Tr = 2.59 min.

EXAMPLE 14




Dichloro (*p*-cymene) ruthenium(II) dimer (24 mg, 0.040 mmol) and (1*R*,2*R*)-(-)-*N*-p-tosyl-1,2-diphenylethylenediamine (35 mg, 0.096 mmol) were suspended in degassed water (16 mL) and the mixture was degassed with nitrogen for 15 min. The mixture was stirred at 70°C under nitrogen for 90 min. The resulting turbid orange mixture was allowed to cool to RT. Solid 1-(6-bromo-quinolin-3-yl)-ethanone (prepared as in WO2011/063233, 1.92 g, 7.68 mmol) followed by degassed tetrahydrofuran (16 mL) and sodium formate (2.72 g, 40 mmol) were added and the reaction mixture was degassed with nitrogen for 5 min, further degassed tetrahydrofuran (5 mL) was added and the mixture degassed for another minute. The reaction mixture was vigorously stirred at 40°C for 21 h and allowed to cool. It was then diluted with ethyl acetate and the mixture washed with water then brine, dried over magnesium sulfate, filtered and evaporated. The residue was purified by silica gel chromatography using a gradient of *iso*-hexanes/ethyl acetate 7:3 to 0:1 to afford the title compound (1.65 g, 85%) as a brown solid.

15




20

To a mixture of **14a** (356 mg, 1.41 mmol), *N,N*-dicyclohexylmethylamine (275 mg, 301 μ L, 1.41 mmol) and *tert*-butyl 3-butenate (470 mg, 537 μ L, 3.31 mmol) in toluene (14 mL) was added bis(tri-*tert*-butylphosphine)palladium(0) (30 mg, 0.058 mmol) under nitrogen and the reaction mixture stirred and heated under reflux for 90 min then allowed to cool. The mixture was evaporated and then purified by silica gel chromatography using a gradient of *iso*-hexanes/ethyl acetate 4:1 to 2:3 to afford the title compound as a yellow oil (144 mg, 33 %).



Compound 14c

To a stirred solution of (S)-1-[(S)-2-((S)-2-*tert*-butoxycarbonylamino-3-methylbutyrylamino)-propionyl]-hexahydro-pyridazine-3-carboxylic acid (189 mg, 0.473 mmol) and **14b** (148 mg, 0.473 mmol) in dichloromethane (10 mL) was added 1-hydroxybenzotriazole containing approx. 20% water (89 mg, 0.662 mmol) followed by 4-dimethylaminopyridine (58 mg, 0.473 mmol) then *N*-(3-dimethylaminopropyl)-*N'*-ethylcarbodiimide hydrochloride (127 mg, 0.662 mmol). The reaction was stirred for 62 h and then diluted with dichloromethane, washed with saturated ammonium chloride solution (2 x) and brine, dried over magnesium sulfate, filtered and evaporated. Purified by silica gel chromatography using a gradient of *iso*-hexanes/ethyl acetate 3:1 to 2:3 afforded the title compound (126 mg, 38 %) as a white foam.




Compound 14

To a stirred solution of **14c** (126 mg, 0.181 mmol) in dichloromethane (1.8 mL) at 0°C under nitrogen was added trimethylsilyl trifluoromethanesulfonate (141 mg, 147 μ L, 0.635 mmol) and the reaction mixture was allowed to warm to ambient temperature over 1.75 h. Further trimethylsilyl trifluoromethanesulfonate (20 mg, 21 μ L, 0.091 mmol) was added and the reaction mixture was stirred for a further 15 min before the addition of *N,N*-diisopropylethylamine (164 mg, 221 μ L, 1.27 mmol). After a further 10 min of stirring the reaction mixture was evaporated and then suspended in acetonitrile (18.1 mL). The stirred mixture was cooled to 0°C and then 2-(1H-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyl uronium hexafluorophosphate methanaminium (96 mg, 0.253 mmol) and *N,N*-diisopropylethylamine (94 mg, 126 μ L, 0.724 mmol) added. After 45 min the reaction was quenched with saturated ammonium chloride solution (10 mL) and the


mixture evaporated to remove organic volatiles. The residue was diluted with dichloromethane and the aqueous layer separated and extracted with dichloromethane. The organic extracts were washed with saturated sodium bicarbonate, water and then brine, dried over sodium sulfate, filtered and evaporated. The residue was purified by silica gel chromatography eluting with ethyl acetate/acetone 9:1 then by preparative reverse phase HPLC using a gradient of acetonitrile/water 3:7 to 1:1 to afford the title compound (6.5 mg, 7% over 2 steps) as a white solid. ¹H NMR (300 MHz, CD₃OD) δ 1.00 (d, *J* = 6.7 Hz, 3H), 1.03 (d, *J* = 6.7 Hz, 3H), 1.61 (d, *J* = 7.1 Hz, 3H), 1.75 (d, *J* = 6.7 Hz, 3H), 1.91-2.01 (m, 3H), 2.68-2.79 (m, 1H), 3.00 (dd, *J* = 14.5, 3.6 Hz, 1H), 3.35-3.42 (m, 1H), 3.73-3.82 (m, 1H), 4.29 (d, *J* = 9.6 Hz, 1H), 4.41 (d, *J* = 12.9 Hz, 1H), 4.70 (d, *J* = 12.3 Hz, 1H), 5.47 (q, *J* = 7.3 Hz, 1H), 6.15 (q, *J* = 6.6 Hz, 1H), 6.30-6.40 (m, 1H), 6.63 (d, *J* = 16.1 Hz, 1H), 7.55 (s, 1H), 7.82 (dd, *J* = 8.7, 1.5 Hz, 1H), 7.95 (d, *J* = 8.7 Hz, 1H), 8.21 (s, 1H), 8.79 (d, *J* = 1.8 Hz, 1H). LCMS (*m/z*) 522.2 [M+H], Tr = 1.60 min.

EXAMPLE 15




Compound 15a

To a stirred mixture of 3-bromo-quinoline-6-carboxylic acid (prepared as in WO2011/090935, 1.94 g, 7.70 mmol) in tetrahydrofuran (77 mL) was added 2-chloro-4,6-dimethoxy-1,3,5-triazine (2.03 g, 11.5 mmol) and *N*-methylmorpholine (2.34 g, 2.54 mL, 23.1 mmol). The reaction mixture was stirred for 90 min and then *N,O*-dimethylhydroxylamine hydrochloride (751 mg, 7.70 mmol) added in one portion. The reaction mixture was stirred for a further 17 h and further *N,O*-dimethylhydroxylamine hydrochloride (375 mg, 3.85 mmol) was added and then after 5 h a further portion (175 mg, 1.80 mmol). The reaction mixture was stirred for a further hour and then diluted with dichloromethane, washed with water and then saturated ammonium chloride solution. The organic layer was dried over sodium sulfate, filtered and evaporated. The residue was purified by silica gel chromatography eluting with a gradient of *iso*-hexanes/ethyl acetate 4:1 to 7:3 to afford the title compound (1.43 g, 66%) as an off-white solid.




Compound 15b

To a stirred solution of methyl magnesium chloride (3 M in tetrahydrofuran, 4.85 mL, 14.5 mmol) in tetrahydrofuran (10 mL) under nitrogen was added a solution of **15a** (1.43 g, 4.85 mmol) in tetrahydrofuran (20 mL). The reaction mixture was stirred for 1 hour and then quenched with saturated ammonium chloride solution. The mixture was diluted with diethyl ether and water, the organic layer separated and washed with further water and the combined aqueous washes back-extracted with diethyl ether. The combined organic extracts were dried over magnesium sulfate, filtered and evaporated. The residue was purified by silica gel chromatography eluting with *iso*-hexanes/ethyl acetate 4:1 to afford the title compound as a white solid which was taken directly into the next step.




Dichloro (*p*-cymene) ruthenium(II) dimer (9 mg, 0.015 mmol) and (1R,2R)-(-)-*N*-p-tosyl-1,2-diphenylethylenediamine (14 mg, 0.037 mmol) were suspended in degassed water (9 mL) and the mixture was degassed with nitrogen for 15 min. The mixture was stirred at 70°C under nitrogen for 90 min. The resulting turbid orange mixture was allowed to cool to RT. Solid 1-(3-bromo-quinolin-6-yl)-ethanone (771 mg, 3.08 mmol) and sodium formate (2.72g, 40 mmol) followed by degassed tetrahydrofuran (4.5 mL) were added and the reaction mixture was degassed with nitrogen for 5 min. The reaction mixture was vigorously stirred at 40°C for 3 h and allowed to cool. It was then diluted with ethyl acetate and the mixture washed with water then brine, dried over sodium sulfate, filtered and evaporated. The residue was purified by silica gel chromatography using a gradient of *iso*-hexanes/ethyl acetate 3:2 to 1:1 to afford the title compound (616 mg, 79%) as a grey solid.




To a solution of **15c** (560 mg, 2.22 mmol) in toluene (20 mL) and 1,4-dioxane (5 mL) was added *N,N*-dicyclohexylmethylamine (740 μ L, 3.46 mmol) and *tert*-butyl 3-butenate (741 mg, 844 μ L, 5.22 mmol) was added bis(*tri-tert*-butylphosphine)palladium(0) (68 mg, 0.13 mmol) under nitrogen and the reaction mixture stirred and heated under reflux for 6 h then allowed to

cool. The mixture was evaporated and then purified by silica gel chromatography using a gradient of *iso*-hexanes/ethyl acetate 4:1 to 0:1 to give a yellow gum. The gum was suspended in ethyl acetate and washed with saturated ammonium chloride solution (2 x) followed by water then brine, dried over sodium sulfate, filtered and evaporated to afford the title compound (300 mg, 43%) as a yellow oil.



Compound 15e

To a stirred slurry of (S)-1-[(S)-2-((S)-2-*tert*-butoxycarbonylamino-3-methylbutyrylamino)-propionyl]-hexahydro-pyridazine-3-carboxylic acid (325 mg, 0.812 mmol), **15d** (231 mg, 0.738 mmol) and powdered 4Å molecular sieves in dichloromethane (16 mL) was added *N*-(3-dimethylaminopropyl)-*N'*-ethylcarbodiimide hydrochloride (198 mg, 1.03 mmol), 1-hydroxybenzotriazole hydrate containing approx. 20% water (174 mg, 1.03 mmol) followed by 4-dimethylaminopyridine (90 mg, 0.738 mmol). The reaction was stirred under nitrogen for 16 h and then filtered and the solid washed with dichloromethane. The filtrate was washed with saturated ammonium chloride solution (2 x) and water, dried over sodium sulfate, filtered and evaporated. The residue was purified by silica gel chromatography using a gradient of *iso*-hexanes/ethyl acetate 1:1 to 0:1 to afford the title compound (307 mg, 60 %) as a yellow foam.




Compound 15

To a stirred solution of **15e** (150 mg, 0.216 mmol) in dichloromethane (2 mL) at 0°C under nitrogen was added trimethylsilyl trifluoromethanesulfonate (168 mg, 174 µL, 0.755 mmol) dropwise and the reaction mixture was allowed to warm to ambient temperature over 2.25 h.

N,N-Diisopropylethylamine (195 mg, 263 μ L, 1.51 mmol) was added and the reaction mixture stirred for a further 10 min, evaporated and then suspended in acetonitrile (21.6 mL). The stirred mixture was cooled to 0°C and then 2-(1H-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyl uronium hexafluorophosphate methanaminium (115 mg, 0.320 mmol) and *N,N*-diisopropylethylamine (111 mg, 150 μ L, 0.864 mmol) added. After 1 h the reaction was quenched with saturated ammonium chloride solution and the mixture evaporated to remove organic volatiles. The residue was diluted with dichloromethane and the organic layer separated and washed with saturated ammonium chloride (2 x) and brine then dried over sodium sulfate, filtered and evaporated. The residue was purified by preparative reverse phase HPLC using a gradient of acetonitrile/water 3:7 to 11:9 to afford the title compound (4.6 mg, 4% over 2 steps) as an off-white solid. ¹H NMR (300 MHz, CD₃OD) δ 1.01 (*app t*, *J* = 6.7 Hz, 6H), 1.62 (d, *J* = 7.1 Hz, 3H), 1.70 (d, *J* = 6.7 Hz, 3H), 1.75-1.85 (m, 1H), 1.91-2.05 (m, 3H), 2.70-2.80 (m, 1H), 2.98-3.06 (m, 1H), 3.36-3.45 (m, 1H), 3.74-3.84 (m, 1H), 4.26 (d, *J* = 10.5 Hz, 1H), 4.42 (*br d*, *J* = 12.3 Hz, 4.72-4.78 (m, 1H), 5.55 (q, *J* = 7.1 Hz, 1H), 6.09 (q, *J* = 6.5 Hz, 1H), 6.38-6.41 (m, 1H), 6.71 (d, *J* = 16.1 Hz, 1H), 7.68 (dd, *J* = 8.9, 1.9 Hz, 1H), 7.88-8.01 (m, 3H), 8.86 (d, *J* = 2.0 Hz, 1H). LCMS (*m/z*) 522.3 [M+H], Tr = 1.64 min.

EXAMPLE 16




A mixture of 7-bromo-2-chloro-quinoline (1.05 g, 4.32 mmol) and tributyl(1-ethoxyvinyl)tin (1.95 g, 1.83 mL, 5.40 mmol) in toluene (21 mL) was degassed for 20 min. Bis(triphenylphosphine)palladium(II) dichloride (302 mg, 0.432 mmol) was added and the reaction mixture stirred under nitrogen and heated at 80°C for 24 h before allowing to cool. The volatiles were evaporated and the residue suspended in 1,4-dioxane (10 mL) and 2 M aqueous hydrochloric acid (5 mL) was added and the reaction mixture stirred for 30 min and then evaporated to remove organics. The residue was diluted with ethyl acetate and water and the organic layer washed with brine, dried over sodium sulfate, filtered and evaporated. The product was purified on silica gel doped with 10% w/w potassium carbonate eluting using a gradient of


iso-hexanes/ethyl acetate 9:1 to 4:1 to afford the title compound (422 mg, 48 %) as a yellow solid.




Dichloro(*p*-cymene)ruthenium(II) dimer (8.5 mg, 0.014 mmol) and (1*R*,2*R*)-(-)-*N*-*p*-tosyl-1,2-diphenylethylenediamine (12.1 mg, 0.033 mmol) were suspended in degassed water (5.5 mL) and the mixture was degassed with nitrogen for 20 min. The mixture was stirred at 70°C under nitrogen for 90 min. The resulting turbid orange mixture was allowed to cool to RT. Solid **16a** (571 mg, 2.78 mmol) and sodium formate (945 mg, 13.9 mmol) followed by degassed tetrahydrofuran (5.5 mL) were added and the reaction mixture was degassed with nitrogen for 5 min. The reaction mixture was vigorously stirred at 40°C for 4 h and allowed to cool. It was then diluted with ethyl acetate and the mixture washed with water and the aqueous layer back-extracted with ethyl acetate. The combined organic extracts were washed with brine, dried over sodium sulfate, filtered and evaporated. The residue was purified by silica gel chromatography using a gradient of *iso*-hexanes/ethyl acetate 4:1 to 7:3 to afford the title compound (413 mg, 72%) as a beige solid.



To a mixture of **16b** (360 mg, 1.73 mmol), potassium vinyltrifluoroborate (279 mg, 2.08 mmol) and [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II), complex with dichloromethane (27 mg, 0.033 mmol) in dry *n*-propanol (27 mL) was added triethylamine (175 mg, 241 μ L, 1.73 mmol) and the system evacuated and purged with nitrogen (3x). The reaction mixture was stirred and heated at reflux for 3 h before being allowed to cool. The mixture was poured into water and extracted with diethyl ether (2x) and the combined organic extracts dried over magnesium sulfate, filtered and evaporated. The residue was purified by silica gel chromatography using a gradient of *iso*-hexanes/ethyl acetate 7:3 to 3:2 to afford the title compound (270 mg, 78%) as a white solid.




To a stirred mixture of **16c** (100 mg, 0.500 mmol) and (S)-1-[(S)-2-((S)-2-but-3-enoylamino-3-methyl-butyrylamino)-propionyl]-hexahydro-pyridazine-3-carboxylic acid (203 mg, 0.550 mmol) in dichloromethane (10 mL) under nitrogen was added *N*-(3-dimethylaminopropyl)-*N'*-ethylcarbodiimide hydrochloride (134 mg, 0.700 mmol) and 4-dimethylaminopyridine (61 mg, 0.500 mmol). The reaction mixture was stirred for 16 h and then diluted with dichloromethane, washed successively with citric acid solution, water then brine, dried over sodium sulfate, filtered and evaporated. The residue was purified by silica gel chromatography using a gradient of *iso*-hexanes/ethyl acetate 1:1 to 0:1 to afford the title compound (103 mg, 38%) as a brown solid.



16d (60 mg, 0.109 mmol) and Hoveyda-Grubbs 2nd generation catalyst (20.5 mg, 0.0328 mmol) was suspended in 1,2-dichloroethane (6 mL) and the mixture heated in a microwave reactor at 100°C for 1 h. The resulting mixture was combined with an identical reaction carried out using (10 mg, 0.0182) of the (S)-1-[(S)-2-((S)-2-but-3-enoylamino-3-methyl-butyrylamino)-propionyl]-hexahydro-pyridazine-3-carboxylic acid (R)-1-(2-vinyl-quinolin-7-yl)-ethyl ester. The volatiles were evaporated and the residue purified by silica gel chromatography using a gradient of ethyl acetate/acetone 1:0 to 4:1 to give a brown gum. This was further purified by preparative thin layer chromatography using *iso*-hexanes/ethyl acetate 1:3 to afford the title compound (1.8 mg, 3%) as a white solid. ¹H NMR (300 MHz, CDCl₃) δ 0.84-1.02 (m, 6H), 1.26 (app s, 3H), 1.67-1.72 (m, 3H), 1.83-1.96 (m, 1H), 2.00-2.08 (m, 1H), 2.63-2.74 (m, 1H), 3.24 (d, *J* = 4.5 Hz, 1H), 3.50 (d, *J* = 7.6 Hz, 1H), 4.21 (app t, *J* = 10.3 Hz, 1H), 4.53-4.61 (m, 1H), 5.55 (app t, *J* = 7.3


1H), 6.04-6.11 (m, 2H), 6.32 (d, $J = 7.8$ Hz, 1H), 6.62 (d, $J = 16.3$ Hz, 1H), 6.72-6.80 (m, 1H), 7.29-7.33 (m, 1H), 7.45 (d, $J = 8.5$ Hz, 1H), 7.72 (d, $J = 8.3$ Hz, 1H), 7.97-8.02 (m, 1H). LCMS (m/z) 552.2 $[M+H]$, $T_r = 1.86$ min.

5


EXAMPLE 17

Under argon, a solution of diisopropylamine (2.51 g, 24.8 mmol) in tetrahydrofuran (150 mL) was cooled in an ice water bath. A solution of *n*-butyllithium in hexanes (2.5 M, 9.7 mL, 24 mmol) was added dropwise over 2 min, and the resulting solution was stirred for 10 additional min. The solution was then cooled to -78°C in a CO_2 :acetone bath, and methyl 2,2-dimethyl-3-oxobutanoate (3.2 g, 22 mmol) was added dropwise over 30 s. The solution was stirred for an additional 15 min, and *N*-phenyl-bis(trifluoromethanesulfonimide) (8.4 g, 23.5 mmol) was added as a solution in tetrahydrofuran (20 mL) via cannula over 5 min, washing with an additional portion of tetrahydrofuran (10 mL). The resulting solution was stirred for 10 min and was removed from the cold bath. After stirring an additional 1 h, the reaction mixture was concentrated *in vacuo* and diluted with diethyl ether (150 mL). The organic phase was washed with 1 M aqueous sodium hydroxide (1 x 100 mL, 1 x 30 mL), dried over anhydrous magnesium sulfate, filtered, and concentrated to afford the title compound (6.2 g, 100%) as an amber liquid that was used without further purification.


20




A solution of **17a** (6.2 g, 22 mmol) in anhydrous pyridine (11 mL, 140 mmol) was heated to 70°C . After 18.5 h, the temperature was increased to 90°C . After stirring for a total of 72 h, the reaction mixture was partitioned between a stirred mixture of diethyl ether (100 mL) and 3 M aqueous hydrochloric acid (100 mL). The phases were separated, and the organic layer was washed with saturated aqueous sodium bicarbonate (75 mL), dried over magnesium sulfate, filtered, and concentrated to afford the title compound (2.7 g, 97%) as a slightly brown liquid that was used without further purification.



Under argon, bis(cyclopentadienyl)zirconium chloride hydride (290 mg, 1.1 mmol) was cooled in an ice water bath. A solution of **17b** (1.4 g, 11.1 mmol) and pinacolborane (2.4 mL, 16.5 mmol) in dichloromethane (3 mL) was added by cannula, washing with an additional
 5 portion of dichloromethane (2 mL). The resulting mixture was removed from the cold bath and was stirred for 72 h at RT. The reaction was then diluted with ethyl acetate (50 mL), quenched with dropwise water (5 mL), and was further diluted with water (50 mL). The organic and aqueous phases were separated, and the aqueous phase was extracted with ethyl acetate (30 mL). The combined organic phases were dried over anhydrous sodium sulfate, filtered, and
 10 concentrated to afford a crude residue that was purified by silica gel chromatography (5 to 15% ethyl acetate in *iso*-hexanes) to afford the title compound (1.6 g, 57%) as a colorless oil that crystallized on standing at -15°C.




A mixture of 6-bromo-3-chloro-isoquinoline (485 mg, 2.0 mmol), bis(pinacolato)diboron
 15 (560 mg, 2.2 mmol), potassium acetate (392 mg, 4.0 mmol) and [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II), complex with dichloromethane (82 mg, 0.1 mmol) in 1,4-dioxane (4 mL) was heated at 160°C in a microwave for 1 hour. The reaction mixture was filtered through Celite and the filter pad was washed with ethyl acetate. The filtrate was evaporated and the residue was purified by silica gel chromatography using a gradient of
 20 dichloromethane to dichloromethane/methanol 9:1 to afford the title compound (545 mg, 94%).



A solution of **17d** (1.74 g, 6.0 mmol) in tetrahydrofuran (45 mL) was stirred at 0°C under nitrogen. Sodium hydroxide solution (2 M, 9 mL, 18 mmol) and hydrogen peroxide (30%, 2.5
 25 mL) were added dropwise. The reaction mixture was stirred at 0°C for 30 min. Water (30 mL) was added and the solution was acidified to pH 1 with 2 M hydrochloric acid. Sodium


metabisulfite solution (1 M) was added and the mixture was extracted with ethyl acetate. The organic extracts were combined, washed with water and brine. The organic solution was filtered through a hydrophobic frit and the filtrate was evaporated to afford the title compound (662 mg, 61%) as an off-white solid.



To a solution of **17e** (1.25 g, 7.07 mmol) and tributyl-(1-ethoxy-vinyl)-tin (5.09 g, 4.77 mL, 14.1 mmol) in 1,4-dioxane (15 mL) was added bis(triphenylphosphine)palladium(II) dichloride (992 mg, 1.41 mmol) and the reaction mixture was heated at 160°C in a microwave for 30 min. The reaction mixture was filtered through Celite and the filter pad was washed with ethyl acetate. The filtrate was evaporated and the residue was purified by silica gel chromatography using a gradient of *iso*-hexanes/ethyl acetate 9:1 to 3:1 to afford 3-(1-ethoxyvinyl)-isoquinolin-6-ol (670 mg) as a gum which was used crude in the next step. 3-(1-Ethoxyvinyl)-isoquinolin-6-ol (670 mg) was suspended in 1,4-dioxane (4 mL) and 4 M hydrogen chloride in 1,4-dioxane (8 mL) was added and the reaction mixture was stirred at RT for 30 min. The solvent was evaporated to afford the title compound (584 mg, 44% over 2 steps) as a white solid.

10


15




A solution of **17f** (384 mg, 2.05 mmol) and pyridine (0.51 mL, 6.16 mmol) in dichloromethane (30 mL) was stirred in a salt-ice bath for 5 min. Trifluoromethanesulfonic anhydride (0.415 mL, 2.46 mmol) was added dropwise and the mixture was warmed to RT over 20 min. Additional trifluoromethanesulfonic anhydride (0.1 mL, 0.6 mmol) was added and the reaction mixture was stirred at RT for 10 min. Saturated ammonium chloride solution was added and the mixture was extracted with dichloromethane. The organic extracts were combined, dried over sodium sulfate, filtered and evaporated. The residue was purified by silica gel chromatography using *iso*-hexanes/ethyl acetate 19:1 to afford the title compound (300 mg, 44%) as a white solid.

20


25



To dichloro (*p*-cymene) ruthenium (II) dimer (2 mg, 0.003 mmol) in water (2 mL) at RT was added (1*R*,2*R*)-(-)-*N*-*p*-tosyl-1,2-diphenylethylenediamine (3 mg, 0.008 mmol). The system was degassed for 15 min and then heated to 70°C for 1.5 h. The reaction was cooled and a solution of **17g** (224 mg, 0.70 mmol) in degassed anhydrous tetrahydrofuran (1 mL) was added followed by sodium formate (237 mg, 3.50 mmol). The system was degassed for 2 min and then heated at 40°C for 1 hour. After cooling to RT, water was added and the water was extracted with dichloromethane (2 x). The combined organic layers were dried through a hydrophobic frit and concentrated *in vacuo*. The product was purified by silica gel chromatography using *iso*-hexanes/ethyl acetate 1:1 to afford the title compound (220 mg, 97%) as a brown oil.




A round bottom flask was charged with **17h** (100 mg, 0.31 mmol), (E)-2,2-dimethyl-4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-but-3-enoic acid methyl ester (91 mg, 0.36 mmol), bis[(dicyclohexyl)(4-dimethylaminophenyl)phosphine]palladium(II) chloride (13 mg, 0.02 mmol), potassium phosphate tribasic (198 mg, 0.93 mmol) and lithium chloride (40 mg, 0.93 mmol). The system was flushed with nitrogen and cyclopentyl methyl ether (1 mL) and water (0.5 mL) were added. The reaction was heated for 1 hour at 90°C and then cooled to RT. Ethyl acetate was added and the solution was washed with water. The organic layer was dried over magnesium sulfate, filtered and concentrated *in vacuo*. The residue was purified by silica gel chromatography using *iso*-hexanes/ethyl acetate 1:2 to give the title compound (70 mg, 75%) as a yellow oil.




To **17i** (140 mg, 0.47 mmol) in tetrahydrofuran (1 mL), methanol (0.5 mL) and water (0.5 mL) was added lithium hydroxide monohydrate (39 mg, 0.93 mmol) at RT. The reaction was

stirred for 3 h and quenched by adding 2 M aqueous hydrochloric acid (0.5 mL). The reaction was concentrated *in vacuo*, followed by co-evaporation from methanol and then toluene. The ensuing yellow solid was used crude.



Compound 17k

- 5 To **1e** (250 mg, 0.47 mmol) in anhydrous dichloromethane (5 mL) at 0°C and under an atmosphere of nitrogen, was added trimethylsilyl trifluoromethanesulfonate (128 μ L, 0.70 mmol). The reaction mixture was stirred at 0°C for 2 h before adding *N,N*-diisopropylethylamine (252 μ L, 1.41 mmol) and then concentrated *in vacuo*, and co-evaporated with toluene to afford (S)-1-[(S)-2-((S)-2-amino-3-methyl-butyrylamino)-propionyl]-hexahydro-pyridazine-3-carboxylic acid
- 10 2,2,2-trichloro-ethyl ester as a white solid. To **17j** (134 mg, 0.47 mmol) in anhydrous *N,N*-dimethylformamide (2 mL) at RT and under an atmosphere of nitrogen was added *N,N*-diisopropylethylamine (420 μ L, 2.35 mmol) and 2-(1H-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyl uronium hexafluorophosphate methanaminium (250 mg, 0.66 mmol). The solution was stirred at RT for 3 min before adding a solution of (S)-1-[(S)-2-((S)-2-amino-3-methyl-
- 15 butyrylamino)-propionyl]-hexahydro-pyridazine-3-carboxylic acid 2,2,2-trichloro-ethyl ester in anhydrous *N,N*-dimethylformamide (2 mL). The reaction was stirred for 16 h. The reaction was quenched with water and extracted with ethyl acetate (2 x). The combined organic layers were dried over sodium sulfate, filtered and concentrated *in vacuo*. The residue was purified by silica gel chromatography using *iso*-hexanes/acetone 1:1 to give the title compound (220 mg, 67% over 3 steps) as an off white solid.
- 20




Compound 17

To **17k** (220 mg, 0.31 mmol) in tetrahydrofuran (3 mL), methanol (1.5 mL) and water (1.5 mL) was added lithium hydroxide monohydrate (53 mg, 1.26 mmol) at RT. The reaction was

stirred for 1.25 h and quenched by adding 2 M aqueous hydrochloric acid (0.64 mL). The reaction was concentrated *in vacuo*, followed by co-evaporation from methanol (2 x) and then acetonitrile (6 x). The resulting residue was dissolved in anhydrous *N,N*-dimethylformamide (4 mL), and added via syringe pump over 2 h to a suspension of 2-methyl-6-nitrobenzoic anhydride (271 mg, 0.79 mmol), 4-dimethylaminopyridine (288 mg, 2.36 mmol) and powdered 4Å molecular sieves (3 g) in 1,2-dichloroethane (103 mL) at 50°C. Following the addition the reaction was stirred at 50°C for 3 h, cooled to RT and filtered through Celite. The filtrate was concentrated to 1/3 of its volume, diluted with dichloromethane and washed with water (2 x). The organic layer was dried through a hydrophobic frit and concentrated *in vacuo*. The residue was purified by silica gel chromatography using *iso*-hexanes/acetone 1:1 to give an off white solid. The solid was triturated twice with diethyl ether and vacuum dried for 16 h to afford the title compound (65 mg, 38% over 2 steps) as a white powder. ¹H NMR (300 MHz, CDCl₃) 0.93 (d, *J* = 6.7 Hz, 3H), 0.99 (d, *J* = 6.7 Hz, 3H), 1.41 (s, 3H), 1.51 (s, 3H), 1.61 (d, *J* = 7.1 Hz, 3H), 1.76 (d, *J* = 6.7 Hz, 3H), 1.76 (m, 1H), 1.78–2.03 (m, 3H), 2.07–2.17 (m, 1H), 2.62–2.76 (m, 3.62–3.79 (m, 2H), 4.21–4.31 (m, 1H), 4.52–4.62 (m, 1H), 5.82–6.01 (m, 1H), 6.13 (q, *J* = 6.7 Hz, 1H), 6.27 (d, *J* = 16.1 Hz, 1H), 6.30–6.41 (m, 2H), 6.64 (d, *J* = 16.1 Hz, 1H), 7.53 (d, *J* = 8.5 Hz, 1H), 7.71 (s, 1H), 7.82 (s, 1H), 7.90 (d, *J* = 8.5 Hz, 1H), 9.16 (s, 1H). LCMS (*m/z*) 549.9 [M+H], Tr = 5.21 min.

20

EXAMPLE 18


Compound 18

To Compound 17 (40 mg, 0.073 mmol) in ethyl acetate (10 mL) was added 10% palladium on carbon (30 mg) at RT. The system was purged with hydrogen and stirred vigorously for 16 h. The suspension was filtered through Celite and the filtrate concentrated *in vacuo*. The residue was purified by silica gel chromatography using *iso*-hexanes/acetone 1:1 to afford Compound 18 (8 mg, 20%) as a white solid. ¹H NMR (300 MHz, CDCl₃) 0.92 (d, *J* = 6.5 Hz, 3H), 1.00 (d, *J* = 6.5 Hz, 3H), 1.27 (s, 3H), 1.31 (s, 3H), 1.48 (d, *J* = 7.1 Hz, 3H), 1.64–1.79 (m, 1H), 1.75 (d, *J* = 6.7 Hz, 3H), 1.80–2.12 (m, 5H), 2.45 (td, *J* = 12.7, 3.6 Hz, 1H), 2.68–2.93

25

(m, 2H), 3.62–3.78 (m, 2H), 3.86 (d, $J = 12.0$ Hz, 1H), 4.37 (app t, $J = 8.3$ Hz, 1H), 4.50–4.61 (1H), 5.87–6.00 (m, 1H), 6.18 (q, $J = 6.7$ Hz, 1H), 6.38 (d, $J = 8.7$ Hz, 1H), 6.58 (d, $J = 8.3$ Hz, 1H), 7.36 (d, $J = 8.5$ Hz, 1H), 7.42 (s, 1H), 7.64 (s, 1H), 7.83 (d, $J = 8.5$ Hz, 1H), 9.14 (s, 1H). LCMS (m/z) 552.2 [M+H], Tr = 5.11 min.


5

EXAMPLE 19

Compound 19a

To a solution of (S)-1-[(S)-2-((S)-2-*tert*-butoxycarbonylamino-3-methyl-butylamino)-propionyl]-hexahydro-pyridazine-3-carboxylic acid (339 mg, 0.920 mmol) and (R)-1-(3-vinyl-isoquinolin-6-yl)-ethanol (183 mg, 0.920 mmol) in dichloromethane (4.6 mL) were added *N*-(3-dimethylaminopropyl)-*N'*-ethylcarbodiimide hydrochloride (211 mg, 1.10 mmol) and 4-dimethylaminopyridine (56.2 mg, 0.46 mmol) at 23°C under an argon atmosphere. After 21 h, the reaction mixture was purified directly by silica gel flash column chromatography to afford the title compound (224 mg, 42%) as a light tan solid.

15




Compound 19b

To **19a** (800 mg, 1.38 mmol) in anhydrous dichloromethane (12 mL) at 0°C and under an atmosphere of nitrogen, was added trimethylsilyl trifluoromethanesulfonate (374 μ L, 2.07 mmol). The reaction mixture was stirred at 0°C for 2 h before adding *N,N*-diisopropylethylamine (480 2.75 mmol) and then concentrated *in vacuo* to afford a white solid. To the solid was added a solution of (E)-2-ethyl-pent-3-enoic acid (188 mg, 1.65 mmol) in anhydrous acetonitrile (12 mL) followed by *N,N*-diisopropylethylamine (240 μ L, 1.38 mmol) and 2-(1H-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyl uranium hexafluorophosphate methanaminium (733 mg, 1.93 mmol). The

20


reaction was stirred at RT for 3 h and concentrated *in vacuo*. The residue was purified by silica gel chromatography using *iso*-hexanes/ethyl acetate 1:2 then 1:5 to give the title compound (250 mg, 74% over 2 steps) as a viscous yellow oil.



Compound 19

- 5 To **19b** (200 mg, 0.34 mmol) in anhydrous toluene (113 mL) at 70°C and under an atmosphere of nitrogen, was added Hoveyda-Grubbs 2nd generation catalyst (42 mg, 0.07 mmol). The reaction was heated at 120°C for 2 h after which an additional amount of the Hoveyda-Grubbs 2nd generation catalyst (30 mg, 0.05 mmol) was added. Following a further 2 h at 120°C the reaction was cooled to RT and potassium isocyanoacetate (83 mg) in methanol (2
- 10 mL) was added. The reaction was stirred for 1 hour, silica added and then concentrated *in vacuo*. The residue was purified by silica gel chromatography using ethyl acetate/acetone 3:1 to afford a brown solid. This was purified further by reverse phase preparative HPLC to give the title compound (2.2 mg, 1 %) as a white solid. ¹H NMR (300 MHz, CD₃OD) 0.97 (d, *J* = 6.5 Hz, 3H), 1.02 (d, *J* = 6.5 Hz, 3H), 1.11 (t, *J* = 7.4 Hz, 3H), 1.64 (d, *J* = 7.1 Hz, 3H), 1.70 (d, *J* = 6.7
- 15 Hz, 3H), 1.71–2.06 (m, 7H), 2.66–2.82 (m, 1H), 3.13–3.22 (m, 1H), 3.75–3.85 (m, 1H), 4.30–4.48 (m, 2H), 5.51–5.67 (m, 1H), 6.07 (q, *J* = 6.3 Hz, 1H), 6.48–6.57 (m, 1H), 6.61 (d, *J* = 16.5 Hz, 1H), 7.47 (s, 1H), 7.58 (d, *J* = 8.7 Hz, 1H), 7.89 (s, 1H), 8.07 (d, *J* = 8.7 Hz, 1H), 9.15 (s, 1H). LCMS (*m/z*) 550.2 [M+H], Tr = 1.67 min.


20

EXAMPLE 20

Compound 20a

To **19a** (390 mg, 0.67 mmol) in anhydrous dichloromethane (7 mL) at 0°C and under an atmosphere of nitrogen, was added trimethylsilyl trifluoromethanesulfonate (182 μ L, 1.01 mmol).


The reaction mixture was stirred at 0°C for 1 hour before quenching with a saturated aqueous solution of sodium hydrogen carbonate and extracting with ethyl acetate (2 x). The organic layers were dried over sodium sulfate, filtered and concentrated *in vacuo* to afford a white solid. The solid was dissolved in anhydrous acetonitrile (4 mL) and 2-methyl-but-3-enoic acid (81 mg, 0.81 mmol), *N*-(3-dimethylaminopropyl)-*N'*-ethylcarbodiimide hydrochloride (184 mg, 0.94 mmol) and 1-hydroxybenzotriazole monohydrate (103 mg, 0.67 mmol) were added. The reaction was stirred at RT for 16 h and concentrated *in vacuo*. The residue was purified by silica gel chromatography using *iso*-hexanes/ethyl acetate 1:1 then 1:3 to give the title compound (259 mg, 69 %) as a clear viscous oil.




Compound 20

To **20a** (250 mg, 0.44 mmol) in degassed anhydrous toluene (148 mL) was added 2,6-dichlorobenzoquinone (8 mg, 0.04 mmol). The mixture was heated to 105°C and a solution of Hoveyda-Grubbs 2nd generation catalyst (83 mg, 0.13 mmol) in anhydrous toluene (20 mL), was added via syringe pump over 2 h. An additional amount of Hoveyda-Grubbs 2nd generation catalyst (28 mg) was added and the reaction heated to reflux for 1 h. The reaction was cooled to RT, silica gel was added and concentrated *in vacuo*. The residue was purified by silica gel chromatography using ethyl acetate/acetone 1:0 then 3:1 to afford an oil. This was triturated with diethyl ether and a few drops of ethyl acetate to afford a brown solid which was further purified by preparative TLC using ethyl acetate/acetone 5/1 to give the title compound (9 mg, 1 %) as an off-white solid. ¹H NMR (300 MHz, CDCl₃) 0.97 (d, *J* = 6.7 Hz, 3H), 1.01 (d, *J* = 6.7 Hz, 3H), 1.38 (d, *J* = 6.7 Hz, 3H), 1.62 (d, *J* = 7.1 Hz, 3H), 1.70 (d, *J* = 6.9 Hz, 3H), 1.72–1.82 (m, 2H), 1.91–2.01 (m, 2H), 2.03–2.12 (m, 1H), 2.62–2.74 (m, 1H), 3.35 (*app t*, *J* = 7.4 Hz, 1H), 3.59 (d, *J* = 12.5 Hz, 1H), 3.65–3.77 (m, 1H), 4.26 (*app t*, *J* = 8.9 Hz, 1H), 4.51–4.60 (m, 1H), 5.68–5.79 (m, 1H), 6.06 (q, *J* = 6.7 Hz, 1H), 6.16 (d, *J* = 9.2 Hz, 1H), 6.27 (d, *J* = 9.2 Hz, 1H), 6.40 (dd, *J* = 16.0, 7.8 Hz, 1H), 6.71 (d, *J* = 16.0 Hz, 1H), 7.39 (dd, *J* = 8.5, 1.3 Hz, 1H), 7.58 (s, 1H), 7.74 (s, 1H), 7.93 (d, *J* = 8.5 Hz, 1H), 9.16 (s, 1H). LCMS (*m/z*) 536.2 [M+H], Tr = 1.63 min.

EXAMPLE 21




21a was prepared in the same manner as **20a** replacing 2-methyl-but-3-enoic acid with (R)-2-methyl-pent-4-enoic acid (prepared as described in Synlett 2002, No12, 2039-2040, 82 mg, 0.72 mmol), to afford the title compound (280 mg, 67%) as a white foam.




To **21a** (250 mg, 0.43 mmol) in anhydrous toluene (144 mL) at RT and under an atmosphere of nitrogen, was added Hoveyda-Grubbs 2nd generation catalyst (54 mg, 0.09 mmol). The reaction was heated at 120°C for 2 h after which an additional amount of the Hoveyda-Grubbs 2nd generation catalyst (25 mg, 0.04 mmol) was added. Following a further 1 h at 120°C the reaction was cooled to RT and concentrated *in vacuo*. The residue was purified by silica gel chromatography using ethyl acetate/acetone 1:0 then 3:1 to afford a brown solid. This was triturated with diethyl ether/ethyl acetate 3:1 to give the title compound (50 mg, 22%) as an off-white solid. ¹H NMR (300 MHz, CDCl₃) 0.91 (d, *J* = 6.7 Hz, 3H), 0.99 (d, *J* = 6.7 Hz, 3H), 1.29 (d, *J* = 7.1 Hz, 3H), 1.59 (d, *J* = 6.7 Hz, 3H), 1.67 (d, *J* = 6.5 Hz, 3H), 1.70–1.84 (m, 1H), 1.92–2.15 (m, 4H), 2.33–2.48 (m, 1H), 2.63–2.98 (m, 3H), 3.53 (d, *J* = 12.2 Hz, 1H), 3.60–3.72 (m, 1H), 4.42 (dd, *J* = 6.7, 2.2 Hz, 1H), 4.52–4.61 (m, 1H), 5.62–5.74 (m, 1H), 6.11 (q, *J* = 6.7 Hz, 1H), 6.16 (q, *J* = 8.9 Hz, 1H), 6.42 (d, *J* = 8.5 Hz, 1H), 6.53–6.66 (m, 1H), 6.75 (d, *J* = 16.1 Hz, 1H), 7.37 (dd, *J* = 8.7, 1.0 Hz, 1H), 7.80–7.96 (m, 3H), 9.16 (s, 1H). LCMS (*m/z*) 550.3 [M+H], Tr = 1.50 min.

EXAMPLE 22




To a stirred slurry of 7-bromo-2-chloro-quinoline (8.10 g, 33.4 mmol) and sodium iodide (50.0 g, 334 mmol) in acetonitrile (27 mL) was slowly added acetyl chloride (3.56 mL, 50.0 mmol). The flask was stoppered and sealed and heated at 80°C for 3 h before being allowed to cool. The mixture was treated sequentially with 10% w/w aqueous potassium carbonate solution (80 mL), 5% w/w aqueous sodium sulfite solution (80 mL) and saturated aqueous sodium thiosulfate solution (80 mL) and the mixture extracted with dichloromethane (2x). The combined organic extracts were dried over sodium sulfate, filtered and evaporated to give a crude 7-bromo-2-iodo-quinoline. To the quinoline was added tributyl(1-ethoxyvinyl)tin (13.6 mL, 40.1 mmol), 1,4-dioxane (67 mL) and bis(triphenylphosphine)palladium(II) dichloride (2.37 g, 3.34 mmol) and the reaction mixture heated at 100°C for 5 h before allowing to cool. 2 M Aqueous hydrochloric acid (67 mL) was added and the reaction stirred for 1 h. The mixture was filtered and the solids washed with ethyl acetate and the filtrate evaporated to remove organics. The residue was extracted with ethyl acetate (3 x) and the combined organic extracts were dried over sodium sulfate, filtered and evaporated. The product was purified on silica gel doped with 10% w/w potassium carbonate eluting with a gradient of 0 to 6% ethyl acetate in *iso*-hexanes to afford the title compound (5.5 g, 66%) as a white solid.




Dichloro (*p*-cymene) ruthenium(II) dimer (61 mg, 0.100 mmol) and (1*R*,2*R*)-(-)-*N*-*p*-tosyl-1,2-diphenylethylenediamine (88 mg, 0.012 mmol) was suspended in degassed water (40 mL) and the mixture was degassed with nitrogen for 5 min. The mixture was stirred at 70°C under nitrogen for 90 min. The resulting turbid orange mixture was allowed to cool to RT. **22a** (5.00 g, 20 mmol) in degassed tetrahydrofuran (40 mL) was added followed by sodium formate (6.8 g, 100 mmol) and the reaction mixture was degassed with nitrogen for 5 min. The reaction mixture was vigorously stirred at 40°C for 4 h and allowed to cool. It was then diluted with ethyl acetate and water and the organic layer was separated, washed with water and brine, dried over sodium

sulfate, filtered and evaporated. The residue was purified by silica gel chromatography using a gradient of 0% to 30% ethyl acetate in *iso*-hexanes to afford the title compound (4.96 g, 98%) as an off-white solid.




Compound 22c

- 5 To a solution of **22b** (1.00 g, 3.97 mmol) and triethylamine (1.65 mL, 11.9 mmol) in anhydrous dichloromethane at 0°C, was added acetic anhydride (0.75 mL, 7.93 mmol) and 4-(dimethylamino)pyridine (24 mg, 0.197 mmol). The reaction mixture was stirred and allowed to warm to RT. After 1.5 h water (100 mL) was added and the layers separated. The aqueous phase was re-extracted with dichloromethane (2 x 100 mL) and the combined organics were washed with brine (100 mL), dried over magnesium sulfate, filtered and concentrated *in vacuo* to give a crude residue. This was purified by silica gel chromatography using *iso*-hexanes (66 mL) then *iso*-hexanes/ethyl acetate 95:5 (300 mL), then *iso*-hexanes/ethyl acetate 9:1 (1066 mL) to yield the title compound (1.16 g, 99%) as a colorless oil.




Compound 22d


- 15 To a solution of acetic acid **22c** (1.16 g, 3.95 mmol) in anhydrous acetonitrile was added palladium(II) acetate (89 mg, 0.395 mmol), 2,2-dimethyl-but-3-enoic acid (496 mg, 4.35 mmol), tri(*o*-tolyl)phosphine (241 mg, 0.790 mmol) and triethylamine (1.09 mL, 7.90 mmol) then the mixture was heated in the microwave at 100°C for 20 min. The reaction mixture was concentrated *in vacuo* then water (200 mL) was added and the organics extracted with ethyl acetate (2 x 50 mL). The combined organics were dried over sodium sulfate, filtered and concentrated *in vacuo* to give a crude residue. This was purified by silica gel chromatography using *iso*-hexanes/ethyl acetate 95:5 (300 mL), then *iso*-hexanes/ethyl acetate 9:1 (1066 mL) then *iso*-hexanes/ethyl acetate 3:1 (1066 mL) to yield the title compound (792 mg, 61%) as a white solid.



A solution of **1e** (1.29 g, 2.42 mmol) in dichloromethane (12 mL) was cooled in an ice-water bath under nitrogen. Trimethylsilyl trifluoromethanesulfonate (0.715 mL, 4.84 mmol) was added dropwise, and the resulting solution was stirred for 1 h. The reaction was quenched with *N,N*-diisopropylethylamine (1.69 mL, 9.68 mmol) and the reaction mixture was concentrated *in vacuo* to give (S)-1-[(S)-2-((S)-2-amino-3-methyl-butyrylamino)-propionyl]-hexahydro-pyridazine-3-carboxylic acid 2,2,2-trichloro-ethyl ester as a white solid which was used without further purification. To a solution of **22d** (792 mg, 2.42 mmol) in *N,N*-dimethylformamide (10 mL) at 0°C under nitrogen was added *N,N*-diisopropylethylamine (2.11 mL, 12.1 mmol) and 2-(1H-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyl uronium hexafluorophosphate methanaminium (1.01 g, 2.66 mmol). The resulting mixture was stirred at RT for 15 min then re-cooled to 0°C and (S)-1-[(S)-2-((S)-2-amino-3-methyl-butyrylamino)-propionyl]-hexahydro-pyridazine-3-carboxylic acid 2,2,2-trichloro-ethyl ester generated in the previous step, was added as a solution in *N,N*-dimethylformamide (10.7 mL). The reaction mixture was then allowed to warm to RT with stirring. After 1 h the mixture was diluted with ethyl acetate (100 mL) and water (100 mL). The phases were separated and the aqueous was extracted with ethyl acetate (100 mL). The combined organics were washed with brine (100 mL), dried over magnesium sulfate, filtered and concentrated *in vacuo* to give a crude residue. This was purified by silica gel chromatography using *iso*-hexanes/acetone 95:5 (1000 mL) then *iso*-hexanes/acetone 9:1 (1 L) then *iso*-hexanes/acetone 85:15 (1 L) then *iso*-hexanes/acetone 3:1 (1 L) then *iso*-hexanes/acetone 7:3 (1 L) to yield the title compound (1.11 g, 62%) as a colorless solid.



A solution of **22e** (1.11 g, 1.50 mmol) in tetrahydrofuran (14.7 mL) was cooled in an ice-water bath and methanol (7.4 mL), water (7.4 mL), and lithium hydroxide monohydrate (252 mg, 6.0 mmol) were then added. The mixture was stirred for 1 h in the ice-water bath and then quenched with aqueous 1 M hydrochloric acid (6 mL, 6.0 mmol). The resulting solution was concentrated *in vacuo*, and the crude product was concentrated from methanol (4 x 250 mL) then acetonitrile (4 x 250 mL) and toluene (5 x 250 mL). The solid isolated was then left on the freeze dryer overnight to afford the title compound (853 mg, quantitative yield) as a white solid.




Compound 22


Under nitrogen, 2-methyl-6-nitrobenzoic anhydride (2.59 g, 7.51 mmol) and 4-(dimethylamino)pyridine (1.38 g, 11.3 mmol) were dissolved in 1,2-dichloroethane (500 mL) and the resulting solution was heated to 50°C. The crude seco-acid from the previous step, **22f** (853 mg, 1.50 mmol) was dissolved in *N,N*-dimethylformamide (19 mL) and added to the reaction mixture dropwise via syringe pump over 6 h. After the addition was complete, the syringe pump was then rinsed with additional *N,N*-dimethylformamide (3 mL) and the reaction mixture was stirred at 50°C for 40 min. After this time the mixture was concentrated to give a residue. This was purified by silica gel chromatography using *iso*-hexanes (66 mL), then *iso*-hexanes/acetone 4:1 (726 mL), then *iso*-hexanes/acetone 7:3 (726 mL), then *iso*-hexanes/acetone 3:2 (726 mL). This initial column gave impure product (1.10 g) which contained *N,N*-dimethylformamide. To this was added brine (200 mL) and the organics extracted with ethyl acetate (2 x 100 mL). The combined organics were dried over magnesium sulfate, filtered and concentrated *in vacuo* to give impure product (900 mg) which does not contain *N,N*-dimethylformamide. This residue was purified by a second silica gel chromatography using *iso*-hexanes/acetone 95:5 (6 L), then *iso*-hexanes/acetone 93:7 (1.5 L), then *iso*-hexanes/acetone 88:12 (3 L), then *iso*-hexanes/acetone 82:16 until all the product eluted from the column. Two batches of the desired product in 87% purity (batch A) and 77% purity (batch B) were isolated. Batch A was triturated twice with 100% diethyl ether to give the title compound (241 mg) in approximately 90-95% purity. Purification of Batch B by preparative reverse phase HPLC gave the title compound (181 mg) in approximately >95% purity. Batch A and B were combined to give the title compound (422 mg, 51%) as a

white solid. ^1H NMR (500 MHz, CD_3OD) 0.94 (*app* t, $J = 6.7$ Hz, 6H), 1.34 (s, 3H), 1.48 (s, 3H), 1.58 (d, $J = 7.3$ Hz, 3H), 1.62-1.75 (m, 5H), 1.82-1.91 (m, 1H), 1.92-2.03 (m, 2H), 2.66-2.78 (m, 1H), 3.78 (*app* d, $J = 8.5$ Hz, 1H), 4.25-4.33 (m, 1H), 4.34-4.43 (m, 1H), 5.71 (q, $J = 7.3$ Hz, 1H), 5.89 (q, $J = 6.7$ Hz, 1H), 6.24 (d, $J = 16.0$ Hz, 1H), 6.46 (d, $J = 16.0$ Hz, 1H), 7.27 (*br* d, $J = 9.8$ Hz, 1H), 7.38 (d, $J = 8.6$ Hz, 1H), 7.60 (s, 1H), 7.75 (ABq, $\Delta\delta_{\text{AB}} = 0.03$, $J_{\text{AB}} = 8.54$ Hz, 2H), 8.18 (d, $J = 7.9$ Hz, 1H). LCMS (m/z) 550.1 $[\text{M}+\text{H}]$, $\text{Tr} = 2.24$ min.

EXAMPLE 23




A solution of 6-bromo-3-chloroisoquinoline (2.0 g, 8.25 mmol) in *n*-propanol (90 mL) was prepared and potassium vinyltrifluoroborate (1.11 g, 8.25 mmol) was added. The solution was purged with nitrogen for 10 min before addition of [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II), complex with dichloromethane (128 mg, 0.157 mmol) and triethylamine (1.15 mL, 8.25 mmol). The reaction mixture was then purged with nitrogen for a further 3 min before heating to reflux for 1 h. The reaction mixture was then allowed to cool to RT and water was added. The organics were then extracted with diethyl ether (3 x 150 mL). The combined organics were dried over magnesium sulfate, filtered and concentrated *in vacuo* to give a crude residue. This was purified by silica gel chromatography, using *iso*-hexanes/diethyl ether 9:1 to give the title compound (1.26 g, 80%) as an oil.




A suspension of **23a** (1.24 g, 6.54 mmol) in methanol (125 mL) and dichloromethane (125 mL) was cooled to -78°C . The reaction was ozonised until a blue colour persisted (in approximately 15 min), then nitrogen was bubbled through the reaction mixture for 15 min to purge the ozone. The reaction was then treated with solid sodium bicarbonate (549 mg, 6.54 mmol) and dimethyl sulfide (1.31 mL, 1.31 mmol). The mixture was allowed to warm to RT and after 3 h the reaction mixture was concentrated *in vacuo*. Water (200 mL) was added to the residue and the aqueous layer extracted with dichloromethane (3 x 200 mL). The combined


organics were dried over magnesium sulfate, filtered and concentrated *in vacuo* to give the title compound (1.02 g, 81 %) as an oil.



To a solution of **23b** (1.02 g, 5.32 mmol) in tetrahydrofuran (10 mL) at 0°C was added trimethyl(trifluoromethyl)silane solution (3.19 mL, 2 M in tetrahydrofuran, 6.38 mmol) followed by tetrabutylammonium fluoride solution (0.053 mL, 1 M in tetrahydrofuran, 0.053 mmol) and the resulting solution was stirred at 0°C for 1.5 h. The reaction mixture was then allowed to warm to RT and 2 M hydrochloric acid (10 mL) was added and the mixture stirred for 10 min before addition of water (200 mL). The aqueous layer was then extracted with ethyl acetate (3 x 100 mL). The organics were combined, dried over sodium sulfate, filtered and concentrated to give the title compound (1.30 g, 93%) as a solid.




To a solution of **23c** (1.30 g, 4.97 mmol) in dichloromethane (14.7 mL) at 0°C was added Dess-Martin periodinane solution (14.7 mL, 15% in dichloromethane, 6.96 mmol) and the reaction mixture was stirred and allowed to warm to RT. 1 M Aqueous sodium metabisulfite (100 mL) was added and the reaction mixture was stirred for 15 min. To this mixture was added saturated aqueous sodium bicarbonate solution (100 mL) and the aqueous layer was extracted with dichloromethane (3 x 100 mL). The combined organics were washed with saturated aqueous sodium bicarbonate solution (200 mL) followed by brine (100 mL), dried over magnesium sulfate, filtered and concentrated *in vacuo* to give the title compound (1.0 g, 78 %) as a solid.



A mixture of **23d** (686 mg, 2.64 mmol), (R)-2-methyl-2-propanesulfonamide (400 mg, 3.30 mmol) and titanium(IV) isopropoxide (1.95 mL, 6.60 mmol) in tetrahydrofuran (27.4 mL)

was heated at reflux for 16 h then allowed to cool to RT. The reaction mixture was concentrated *in vacuo* and re-dissolved in diethyl ether (27.4 mL) then cooled to -78 °C. To this mixture was added L-selectride® solution (7.93 mL, 1.0 M in tetrahydrofuran, 7.93 mmol) and the reaction mixture stirred at -78 °C for 1 h. Brine (30 mL) was then added and the mixture allowed to warm to RT. More brine (200 mL) and ethyl acetate (300 mL) were added and the layers separated. The aqueous phase was re-extracted with dichloromethane (500 mL). The combined organics were dried over sodium sulfate, filtered and concentrated *in vacuo* to give a crude residue. This was purified by silica gel chromatography, using gradient elution of dichloromethane to dichloromethane/methanol 99:1 to give the title compound (553 mg, 57% yield, 94% d.e.).


10



Compound 23f

To a solution of **23e** (553 mg, 1.52 mmol) in methanol (10 mL) was added 4 M hydrochloric acid solution in 1,4-dioxane (1.52 mL, 6.06 mmol) and the reaction mixture was stirred at RT for 30 min. Additional 4 M hydrochloric acid solution in 1,4-dioxane (1.52 mL, 6.06 mmol) was added and the reaction mixture stirred at RT for 30 min then concentrated *in vacuo* to give the title compound (507 mg, quantitative yield).

15




Compound 23g

A mixture of **23f** (507 mg, 1.52 mmol) and triethylamine (0.84 mL, 6.08 mmol) in dichloromethane (20 mL) was stirred at 0°C to give a solution. Then a solution of di-*tert*-butyl dicarbonate (497 mg, 2.28 mmol) in dichloromethane (4 mL) was added and the reaction mixture was stirred and allowed to warm to RT over 24 h. Water (200 mL) was added and the aqueous layer extracted with dichloromethane (3 x 100 mL), dried over sodium sulfate, filtered and concentrated *in vacuo* to give a residue (816 mg). This residue was dissolved in dichloromethane (20 mL) and cooled to 0°C then triethylamine (0.84 mL, 6.08 mmol) was added followed by a solution of di-*tert*-butyl dicarbonate (497 mg, 2.28 mmol) in dichloromethane (4 mL) and the reaction mixture was stirred and allowed to warm to RT over 72 h. Additional di-

20


25

tert-butyl dicarbonate (162 mg, 0.74 mmol) was added and stirring continued for 30 min then water (100 mL) was added and the aqueous layer extracted with dichloromethane (2 x 50 mL). The combined organics were dried over sodium sulfate, filtered and concentrated *in vacuo* to give a crude residue. This was purified by silica gel chromatography, using gradient elution of
 5 *iso*-hexanes/ethyl acetate 9:1 to *iso*-hexanes/ethyl acetate 1:1 to give the title compound (140 mg, 25%) as an oil.




Compound 23h

To a solution of **23g** (135 mg, 0.37 mmol) in *n*-propanol (6 mL) was added potassium vinyltrifluoroborate (55 mg, 0.412 mmol), [1,1'-bis(diphenylphosphino)
 10 ferrocene]dichloropalladium(II), complex with dichloromethane (5.8 mg, 0.007 mmol) and triethylamine (0.052 mL, 0.37 mmol). The suspension was evacuated and purged with nitrogen 3 times before heating to 100°C for 2 h. After this time additional potassium vinyltrifluoroborate (100 mg, 0.748 mmol) was added and heating continued at 100°C for 16 h. After this time,
 15 additional [1,1'-bis(diphenylphosphino)ferrocene] dichloropalladium(II), complex with dichloromethane (9 mg, 0.011 mmol) was added and the reaction mixture was heated in a microwave reactor at 150°C for 90 min. After this time additional [1,1'-
 bis(diphenylphosphino)ferrocene]dichloropalladium(II), complex with dichloromethane (9 mg, 0.011 mmol) was added and the reaction mixture was heated in a microwave reactor at 150°C
 20 for 10 min. Water (100 mL) was added to the reaction mixture and the aqueous layer was then extracted with diethyl ether (2 x 100 mL). The combined organics were dried over sodium sulfate, filtered and concentrated *in vacuo* to give the title compound (132 mg, quantitative yield).



Compound 23i


To a solution of **23h** (132 mg, 0.37 mmol) in methanol (2 mL) was added 4 M hydrochloric acid in 1,4-dioxane (0.37 mL, 0.75 mmol) and the mixture stirred at RT for 30 min. Additional 4 M hydrochloric acid in 1,4-dioxane (3.0 mL, 6.08 mmol) was added and stirring continued for 2 h then the reaction mixture was concentrated *in vacuo* to give a residue. The residue was then concentrated from diethyl ether (2 mL), followed by acetonitrile (10 mL) and this was repeated twice to give (S)-2,2,2-trifluoro-1-(3-vinyl-isoquinolin-6-yl)-ethylamine hydrochloric acid salt (108 mg, quantitative yield). To a suspension of (S)-2,2,2-trifluoro-1-(3-vinyl-isoquinolin-6-yl)-ethylamine hydrochloric acid salt (108 mg, 0.374 mmol) in acetonitrile (2 mL) at 0°C under nitrogen was added (S)-1-[(S)-2-((S)-2-but-3-enoylamino-3-methyl-butrylamino)-propionyl]-hexahydro-pyridazine-3-carboxylic acid (138 mg, 0.37 mmol), *N*-(3-dimethylaminopropyl)-*N'*-ethylcarbodiimide hydrochloride (100 mg, 0.52 mmol) and 1-hydroxybenzotriazole hydrate (63 mg, 20 wt % in water, 0.37 mmol). The resulting suspension was stirred at 0°C for 15 min before removing from the ice-water bath and stirring at RT for 16 h. *N,N*-Diisopropylethylamine (0.5 mL, 2.87 mmol) was added and the solution was stirred at RT for 2.5 h where additional (S)-1-[(S)-2-((S)-2-but-3-enoylamino-3-methyl-butrylamino)-propionyl]-hexahydro-pyridazine-3-carboxylic acid (69 mg, 0.19 mmol) was added and stirring continued. After 1 h, *N*-(3-dimethylaminopropyl)-*N'*-ethylcarbodiimide hydrochloride (134 mg, 0.70 mmol) was added and the reaction mixture was stirred at RT for 1 hour. After this time, the mixture was concentrated *in vacuo* and the residue diluted with ethyl acetate (60 mL). Water (50 mL) was added and the layers separated. The aqueous phase was re-extracted with ethyl acetate (2 x 10 mL) and the combined organics were then washed with ammonium chloride (2 x 50 mL), dried over sodium sulfate, filtered and concentrated *in vacuo* to give a crude residue. This was purified by silica gel chromatography, using gradient elution of *iso*-hexanes/ethyl acetate 9:1 to *iso*-hexanes/ethyl acetate 4:1 to *iso*-hexanes/ethyl acetate 1:1 and finally *iso*-hexanes/ethyl acetate 0:1. This gave the title compound (124 mg, 55%).



To a solution of **23i** (124 mg, 0.21 mmol) in toluene (69 mL) was added Hoveyda-Grubbs 2nd generation catalyst (13 mg, 0.02 mmol) and the reaction mixture heated at 115°C for


1.5 h. Additional Hoveyda-Grubbs 2nd generation catalyst (13 mg, 0.02 mmol) was added and heating continued at 120°C for 50 min. Additional Hoveyda-Grubbs 2nd generation catalyst (13 mg, 0.02 mmol) was added and heating continued at 120°C for 30 min. After this time the mixture was allowed to cool to RT and concentrated *in vacuo*. The residue was purified by silica gel chromatography using a gradient of ethyl acetate to ethyl acetate/acetone 9:1. Impure product (28.3 mg) was collected which was further purified by preparative TLC eluting with ethyl acetate/acetone 95:5 to afford the title compound (4 mg, 3%) as a white solid. ¹H NMR (300 MHz, CD₃OD) 0.99 (d, *J* = 6.7 Hz, 3H), 1.01 (d, *J* = 6.7 Hz, 3H), 1.29-1.39 (m, 1H), 1.69 (d, *J* = 7.4 Hz, 3H), 1.74-2.01 (m, 5H), 2.73 (td, *J* = 12.9, 2.9 Hz, 1H), 2.99-3.11 (m, 1H), 3.74-3.83 (m, 1H), 4.26 (d, *J* = 10.0 Hz, 1H), 4.41 (*br* d, *J* = 13.2 Hz, 1H), 5.48 (q, *J* = 7.4 Hz, 1H), 5.84 (q, *J* = 8.3 Hz, 1H), 6.58 (d, *J* = 16.3 Hz, 1H), 6.62-6.71 (m, 1H), 7.47 (s, 1H), 7.76 (d, *J* = 8.6 Hz, 1H), 7.89 (s, 1H), 8.13 (d, *J* = 8.6 Hz, 1H), 9.22 (s, 1H). LCMS (*m/z*) 575.2 [M+H], Tr = 1.53 min.

EXAMPLE 24



To a solution of (R)-1-(7-bromo-quinolin-2-yl)-ethanol (295 mg, 1.17 mmol) in anhydrous acetonitrile (12 mL) were added but-3-enoic acid *tert*-butyl ester (0.44 mL, 2.75 mmol), bis(*tert*-butylphosphine)palladium(0) (25 mg, 0.049 mmol) and *N,N*-dicyclohexylmethylamine (0.39 mL, 1.83 mmol) then the mixture was heated at reflux for 2 h. The reaction mixture was concentrated *in vacuo* to give a crude residue. This was purified by silica gel chromatography using *iso*-hexanes/ethyl acetate 95:5 to yield the title compound (348 mg, 95%) as a white solid.


20



To **24a** (363 mg, 1.16 mmol) and (S)-1-[(S)-2-((S)-2-*tert*-butoxycarbonylamino-3-methylbutyrylamino)-propionyl]-hexahydro-pyridazine-3-carboxylic acid (505 mg, 1.26 mmol) in dichloromethane (20 mL) was added *N*-(3-dimethylaminopropyl)-*N'*-ethylcarbodiimide

25

hydrochloride (311 mg, 1.62 mmol) and 4-dimethylaminopyridine (142 mg, 1.16 mmol). The reaction mixture was then stirred at RT for 16 h. To the mixture was added saturated aqueous ammonium chloride solution (100 mL) and the aqueous layer extracted with dichloromethane (2 x 50 mL). The aqueous phase was further washed with dichloromethane (20 mL) and the combined organics washed with brine (50 mL), dried over magnesium sulfate, filtered and concentrated *in vacuo* to give a crude residue. This was purified by silica gel chromatography using *iso*-hexanes/ethyl acetate 4:1 (120 mL) then *iso*-hexanes/ethyl acetate 7:3 (855 mL) then *iso*-hexanes/ethyl acetate 3:2 (540 mL) to yield the title compound (357 mg, 44%) as a white solid.



To a solution of **24b** (342 mg, 0.492 mmol) in dichloromethane (2.5 mL) at 0°C was added trimethylsilyl trifluoromethanesulfonate (0.18 mL, 0.96 mmol) and the pale yellow solution stirred at 0°C for 2.75 h. Additional trimethylsilyl trifluoromethanesulfonate (0.073 mL, 0.45 mmol) was added and stirring continued for 1.33 h. *N,N*-Diisopropylethylamine (0.6 mL, 3.44 mmol) was added and the mixture stirred for 10 min then concentrated *in vacuo* to give intermediate (S)-1-[(S)-2-((S)-2-amino-3-methyl-butyrylamino)-propionyl]-hexahydro-pyridazine-3-carboxylic acid (R)-1-[7-((E)-3-carboxy-propenyl)-quinolin-2-yl]-ethyl ester (266 mg, 0.45 mmol). To a solution of (S)-1-[(S)-2-((S)-2-amino-3-methyl-butyrylamino)-propionyl]-hexahydro-pyridazine-3-carboxylic acid (R)-1-[7-((E)-3-carboxy-propenyl)-quinolin-2-yl]-ethyl ester (266 mg, 0.45 mmol) in acetonitrile (50 mL) at 0°C under nitrogen was added *N,N*-diisopropylethylamine (0.343 mL, 1.971 mmol) and 2-(1H-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyl uronium hexafluorophosphate methanaminium (262 mg, 0.70 mmol). The mixture was stirred at 0°C and allowed to warm to RT over 16 h. To the reaction mixture was added 2 M aqueous hydrochloric acid solution (20 mL) and the mixture concentrated *in vacuo*. The aqueous layer was extracted with dichloromethane/methanol 9:1 (2 x 100 mL). The combined organics were washed with saturated sodium hydrogen carbonate solution (2 x 200 mL), then brine (1 x 100 mL), dried over sodium sulfate, filtered and concentrated *in vacuo* to give a crude residue. This was purified by silica gel chromatography using gradient elution of ethyl acetate/acetone 97:3 to ethyl acetate/acetone 94:6. Collected 45 mg of impure product which was triturated with diethyl ether to give the title compound (25.4 mg, 10%) as a pale yellow solid. ¹H NMR (300 MHz, CD₃OD) δ

15


20

25


30

0.99 (d, $J = 6.48$ Hz, 3H), 1.01 (d, $J = 6.7$ Hz, 3H), 1.64 (d, $J = 7.1$ Hz, 3H), 1.66-1.72 (m, 1H), 1.74 (d, $J = 6.9$ Hz, 3H), 1.85-2.08 (m, 4H), 2.75 (dd, $J = 2.9, 12.9$ Hz, 1H), 2.97-3.07 (m, 1H), 3.35-3.40 (m, 1H), 3.79-3.88 (m, 1H), 4.23 (d, $J = 10.5$ Hz, 1H), 4.43 (br d, $J = 11.2$ Hz, 1H), 5.72 (q, $J = 7.1$ Hz, 1H), 5.94 (q, $J = 6.9$ Hz, 1H), 6.41 (d, $J = 16.7$ Hz, 1H), 6.55 (dt, $J = 4.7, 16.7$ Hz, 1H), 7.40 (d, $J = 8.5$ Hz, 1H), 7.65 (s, 1H), 7.75 (dd, $J = 1.3, 8.7$ Hz, 1H), 7.81 (d, $J = 8.7$ Hz, 1H), 8.21 (d, $J = 8.5$ Hz, 1H). LCMS (m/z) 522.2 [$M+H$], Tr = 1.90 min.


EXAMPLE 25



A cooled (-20°C) solution of (S)-6-bromo-indan-1-ol (1.0462 g, 4.910 mmol, prepared as described in WO 2009/003719) in anhydrous tetrahydrofuran (20 mL) was subsequently treated with triethylamine (2.7 mL, 19.639 mmol) and methanesulfonyl chloride (760 μL , 9.820 mmol). After stirring at -30°C for 2.5 h, a solution of methylamine (2 M in tetrahydrofuran, 25 mL, 50 mmol) was added. After stirring at RT for 22.5 h the reaction mixture was filtered. The white solid was rinsed with diethyl ether. The filtrate was evaporated to dryness to afford the title compound (1.11 g, quantitative yield) as a white solid.




A solution of **25a** (1.11 g, 4.910 mmol) was subsequently treated with di-*tert*-butyl dicarbonate (1.714 g, 7.856 mmol) and triethylamine (690 μL , 4.91 mmol). After stirring for 25.5 h, the volatiles were removed *in vacuo* and the residue was purified by silica gel chromatography using a 50 g Isolute cartridge eluted with a continuous gradient of *iso*-hexanes/ethyl acetate 1:0 to 9:1 to afford the title compound (1.4224 g, 89%) as a white solid as a mixture of rotamers.



Compound 25c


A solution of **25b** (1.4224 g, 4.363 mmol), [1,1-bis(diphenylphosphino)ferrocene] dichloropalladium dichloromethane complex (71.3 mg, 0.087 mmol), potassium vinyltrifluoroborate (701.3 mg, 5.236 mmol) and triethylamine (610 μ L, 4.363 mmol) in anhydrous *n*-propanol (40 mL) was degassed by bubbling nitrogen through for 30 min. The red suspension was then refluxed for 17.5 h. After cooling to RT, the mixture was quenched with water and the aqueous layer was extracted with diethyl ether (2 x). The organics were combined, dried over sodium sulfate, filtered and the volatiles were removed *in vacuo*. The residue was purified by silica gel chromatography using a 50 g Isolute cartridge eluted with a continuous gradient of *iso*-hexanes/diethyl ether 1:0 to 4:1 to afford the title compound (977.3 mg, 82%) as a white solid as a mixture of rotamers.




Compound 25d

A cooled (0°C) solution of **25c** (977.3 mg, 3.575 mmol) in dichloromethane (20 mL) was treated with trifluoroacetic acid (5 mL). After stirring for 30 min at RT the volatiles were removed *in vacuo* and the residual trifluoroacetic acid was azeotroped off with toluene (3 x) to afford the corresponding ammonium salt as a pink solid. A cooled (0°C) solution of the crude ammonium salt, crude (S)-1-[(S)-2-((S)-2-*tert*-butoxycarbonylamino-3-methyl-butyrylamino)-propionyl]-hexahydro-pyridazine-3-carboxylic acid (3.932 mmol) and *N,N*-diisopropylethylamine (2.5 mL, 14.300 mmol) in acetonitrile (60 mL) was treated with 2-(1H-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyl uronium hexafluorophosphate methanaminium (1.903 g, 5.005 mmol). After stirring for 20 h at RT, the reaction was quenched with hydrochloric acid (1 M, 100 mL). The aqueous layer was extracted with ethyl acetate (2 x). The organics were combined, dried over sodium sulfate, filtered and the volatiles were removed *in vacuo*. The residue was purified by silica gel

chromatography using a 100 g Isolute cartridge eluted with a continuous gradient of *iso*-hexanes/ethyl acetate 1:0 to 1:2 to afford the title compound in a mixture which was dissolved in ethyl acetate and washed with aqueous potassium carbonate. The organics were dried over sodium sulfate, filtered and the volatiles were removed *in vacuo* to afford the title compound
 5 (798.9 mg, 36%) as a white solid as a mixture of rotamers.




Compound 25e was prepared in the same manner as **22e** using **25d** instead of **1e** in 27% yield as a complex mixture of rotamers.




10 A solution of **25e** (238.3 mg, 0.390 mmol) in dichloroethane (100 mL) was treated with Hoveyda-Grubbs 2nd generation catalyst (48.9 mg, 0.078 mmol). After stirring at reflux for 1.5 h, the reaction was cooled to RT and the volatiles were removed *in vacuo*. The residue was purified by silica gel chromatography using a 10 g Isolute cartridge eluted *by gravity* with a continuous gradient of ethyl acetate/methanol 1:0 to 95:5 followed by preparative TLC eluted
 15 with ethyl acetate/methanol 97:3 (2 elutions) to provide the final compound (10.2 mg, 5%) as a white solid. ¹H NMR (300 MHz, CD₃CN) δ 0.85-0.97 (m, 6H), 1.21-1.35 (m, 8H), 1.42-1.62 (m, 3H), 1.64-1.77 (m, 2H), 1.80-1.89 (m, 2H), 2.27-2.38 (m, 2H), 2.52 (dd, *J* = 7.6, 3.1 Hz, 1H), 2.76 (s, 3H), 2.82-3.06 (m, 2H), 3.27-3.35 (m, 1H), 3.47 (s, 3H), 3.80-3.91 (m, 1H), 4.06 (dd, *J* = 9.1, 8.2 Hz, 1H), 4.23 (d, *J* = 11.8 Hz, 1H), 4.41 (dd, *J* = 12.9, 3.6 Hz, 1H), 5.44 (*app* pentet, *J* =
 20 8.0 Hz, 1H), 6.10-6.30 (m, 2H), 6.43 (d, *J* = 16.0 Hz, 1H), 6.88 (d, *J* = 8.0 Hz, 1H), 6.96 (d, *J* = 9.4 Hz, 1H), 7.04 (s, 1H), 7.13-7.28 (m, 2H). LCMS (*m/z*) 582.3 [M+H], Tr = 2.57 min.

EXAMPLE 26




Compound 26a

5 A solution of (S)-4-benzyl-3-((R)-2-methyl-pent-4-enoyl)-oxazolidin-2-one (1.65 g, 6.04 mmol, prepared as in *Synlett* 2002, 12, 2039-2040) in ethanol/water (22 mL, 10:1) was treated with rhodium(III) chloride hydrate (31.6 mg, 0.15 mmol). After stirring at 85°C for 24 h, the reaction mixture was cooled to RT and the volatiles were removed *in vacuo*. The residue was purified by silica gel chromatography using a 50 g Isolute cartridge eluted with a continuous gradient of *iso*-hexanes/diethyl ether 1:0 to 4:1 to afford the title compound (892.7 mg, 54%) as a colorless oil.




Compound 26b

A cooled (0°C) solution of **26a** (893 mg, 3.27 mmol) in tetrahydrofuran/water (30 mL, 2:1) was subsequently treated with hydrogen peroxide (1.7 mL, 16.33 mmol, 30% in water) and lithium hydroxide hydrate (274 mg, 6.53 mmol). After stirring for 1.5 h at 0°C, the reaction was
 15 quenched with sodium metabisulfite (6.2 g, 32.66 mmol). After stirring at RT for 40 min, the mixture was acidified with hydrochloric acid (2 M) and the aqueous layer was extracted with dichloromethane (2 x). The organics were combined, filtered through a phase separator and the volatiles were removed *in vacuo*. The residue was purified by silica gel chromatography using a 25 g Isolute cartridge eluted with a continuous gradient of *iso*-hexanes/ethyl acetate 1:0 to 4:1 to
 20 afford the title compound (307.1 mg, 82%) as a colorless oil.




Compound 26c

Compound **26c** was prepared in the same manner as methyl-((R)-6-vinyl-indan-1-yl)-carbamic acid *tert*-butyl ester using (R)-1-(7-bromo-quinolin-2-yl)-ethanol instead of ((R)-6-bromo-indan-1-yl)-methyl-carbamic acid *tert*-butyl ester in quantitative yield.



Compound 26d


- 5 A solution of (S)-1-[(S)-2-((S)-2-*tert*-butoxycarbonylamino-3-methyl-butylamino)-propionyl]-hexahydro-pyridazine-3-carboxylic acid (261.2 mg, 0.653 mmol), **26c** (108.4 mg, 0.544 mmol) and 4-dimethylaminopyridine (79.7 mg, 0.653 mmol) in dichloromethane (10 mL) was treated with *N*-(3-dimethylaminopropyl)-*N'*-ethylcarbodiimide hydrochloride (166.9 mg, 0.870 mmol). After stirring at RT for 17 h, the volatiles were removed *in vacuo* and the residue
- 10 was purified by silica gel chromatography using a 50 g Isolute cartridge eluted with a continuous gradient of *iso*-hexanes/ethyl acetate 1:0 to 1:4 to 0:1 to afford the title compound (139.6 mg, 44%) as a white solid.



Compound 26e

- A cooled (0°C) solution of **26d** (139.6 mg, 0.240 mmol) in anhydrous dichloromethane
- 15 (10 mL) was treated with trimethylsilyl methanesulfonate (90 μ L, 0.480 mmol). After stirring for 1.5 h at 0°C, the reaction was quenched with saturated sodium bicarbonate. The aqueous layer was extracted with dichloromethane. The organics were combined, filtered through a phase separator and the volatiles were removed *in vacuo* to provide the intermediate amine. A solution of this amine, (E)-2-methyl-pent-3-enoic acid (32.9 mg, 0.288 mmol) and 1-
- 20 hydroxybenzotriazole (38.9 mg, 0.288 mmol) in acetonitrile (10 mL) was treated with *N*-(3-dimethylaminopropyl)-*N'*-ethylcarbodiimide hydrochloride (64.4 mg, 0.336 mmol). After stirring at


RT for 17.5 h the volatiles were removed *in vacuo* and the residue was purified by silica gel chromatography using a 25 g Isolute cartridge eluted with a continuous gradient of *iso*-hexanes/ethyl acetate 1:0 to 0:1 to afford the title compound (69.8 mg, 50%) as a colorless oil.



Compound 26

- 5 Compound 26 was prepared in the same manner as compound 25 using **26e** instead of (S)-1-[(S)-2-[(S)-2-((2R,3R)-3-methoxy-2-methyl-hept-6-enoylamino)-3-methyl-butyrylamino]-propionyl]-hexahydro-pyridazine-3-carboxylic acid methyl-((R)-6-vinyl-indan-1-yl)-amide in 4% yield. ¹H NMR (300 MHz, CD₃OD) δ 0.94-1.00 (m, 6H), 1.47 (d, *J* = 7.6 Hz, 3H), 1.63 (d, *J* = 7.1 Hz, 3H), 1.67-1.80 (m, 5H), 1.85-2.07 (m, 3H), 2.76 (td, *J* = 12.1, 3.3 Hz, 1H), 3.79-3.86 (m, 1H),
 10 4.28 (d, *J* = 10.5 Hz, 1H), 4.36-4.46 (m, 1H), 5.73 (q, *J* = 7.1 Hz, 1H), 5.93 (q, *J* = 6.9 Hz, 1H), 6.32-6.51 (m, 2H), 7.41 (d, *J* = 8.5 Hz, 1H), 7.65 (s, 1H), 7.77-7.85 (m, 2H), 8.21 (d, *J* = 8.5 Hz, 1H). LCMS (*m/z*) 536.1 [*M*+*H*], Tr = 1.80 min.


EXAMPLE 27



Compound 27a


- 15 A cooled (-78°C) solution of (of (R)-4-benzyl-3-propionyl-oxazolidin-2-one (3.00 g, 12.9 mmol) in anhydrous tetrahydrofuran (40 mL) was treated with potassium bis(trimethylsilyl)amide (19.3 mL, 19.3 mmol, 1 M in tetrahydrofuran). After stirring for 0.45 h at -78°C, the mixture was treated with allyl bromide (5.6 mL, 64.3 mmol). After stirring for 2 h at -40°C, the reaction was
 20 quenched with 2 M hydrochloric acid. The aqueous was extracted with ethyl acetate (2 x 50 mL). The organics were combined, and the volatiles were removed *in vacuo*. The residue was purified by silica gel chromatography using a 50 g Biotage cartridge eluted with a continuous

gradient of *iso*-hexanes/ethyl acetate 1:0 to 1:1 to afford the title compound (2.73 g, 78%) as a colorless oil.




Compound 27b

- 5 A solution of **27a** (2.73 g, 9.97 mmol) in ethanol/water (22 mL, 10:1) was treated with rhodium(III) chloride hydrate (52 mg, 0.25 mmol). After stirring at 85°C for 3 h, the reaction mixture was cooled to RT and the volatiles were removed *in vacuo*. The residue was purified by silica gel chromatography using a 50 g Biotage cartridge eluted with a continuous gradient of *iso*-hexanes/ethyl acetate 1:0 to 4:1 to afford the title compound (1.76 g, 65%) as a colorless oil.




Compound 27c

- 10 A cooled (0°C) solution of **27b** (1.76 g, 6.44 mmol) in tetrahydrofuran/water (60 mL, 2:1) was subsequently treated with hydrogen peroxide (3.3 mL, 32.2 mmol, 30% in water) and lithium hydroxide hydrate (534 mg, 12.9 mmol). After stirring for 2 h at 0°C, the reaction was quenched with sodium metabisulfite (12.2 g, 64.4 mmol). After stirring at RT for 1 h, the mixture was acidified with 2 M hydrochloric acid and the aqueous layer was extracted with
15 dichloromethane (2 x 100 mL). The organics were combined, filtered and the volatiles were removed *in vacuo*. The residue was purified by silica gel chromatography using a 50 g Biotage cartridge eluted with a continuous gradient of *iso*-hexanes/ethyl acetate 1:0 to 4:1 to afford the title compound (463 mg, 63%) as a colorless oil.



Compound 27d


A cooled (0°C) solution of (S)-1-[(S)-2-[(S)-2-*tert*-butoxycarbonylamino-3-methyl-butrylamino]-propionyl]-hexahydro-pyridazine-3-carboxylic acid (R)-1-(7-vinyl-quinolin-2-yl)-ethyl ester (380 mg, 0.65 mmol) in anhydrous dichloromethane (10 mL) was treated with trimethylsilyl methanesulfonate (237 μ L, 1.31 mmol). After stirring for 1 h at 0 °C, the reaction
 5 was quenched with saturated sodium bicarbonate. The aqueous layer was extracted with dichloromethane. The organics were separated and combined, volatiles were removed *in vacuo* to provide the intermediate amine (305 mg). A solution of this amine, **27c** (90 mg, 0.76 mmol) and 1-hydroxybenzotriazole (103 mg, 0.76 mmol) in acetonitrile (20 mL) was treated with *N*-(3-dimethylaminopropyl)-*N'*-ethylcarbodiimide hydrochloride (170 mg, 0.89 mmol). After stirring at
 10 RT for 16 h the volatiles were removed *in vacuo* and the residue was purified by silica gel chromatography using a 25 g Biotage cartridge eluted with a continuous gradient of *iso*-hexanes/ethyl acetate 1:0 to 0:1 to afford the title compound (236 mg, 65%) as a colorless oil.



Compound 27


Compound 27 was prepared in the same manner as compound 25 using **27d** instead of
 15 (S)-1-[(S)-2-[(S)-2-((2R,3R)-3-methoxy-2-methyl-hept-6-enoylamino)-3-methyl-butrylamino]-propionyl]-hexahydro-pyridazine-3-carboxylic acid methyl-((R)-6-vinyl-indan-1-yl)-amide in 5% yield. ¹H NMR (300 MHz, CD₃OD) δ 0.85-1.00 (m, 6H), 1.30 (d, *J* = 6.9 Hz, 3H), 1.58 (d, *J* = 7.1 Hz, 3H), 1.62-1.80 (m, 6H), 1.83-2.07 (m, 3H), 2.70-2.82 (m, 1H), 3.38-3.48 (m, 1H), 3.76-3.84 (m, 1H), 4.20-4.28 (m, 1H), 4.36-4.47 (m, 1H), 5.62-5.74 (m, 1H), 5.93 (q, *J* = 6.9 Hz, 1H), 6.30-
 20 6.51 (m, 2H), 7.39 (d, *J* = 8.5 Hz, 1H), 7.63 (s, 1H), 7.70-7.80 (m, 2H), 8.13-8.22 (m, 1H). LCMS (m/z) 536.2 [M+H], Tr = 2.15 min.

EXAMPLE 28




Compound 28a


Potassium carbonate (647 mg, 4.68 mmol) and di-*tert*-butyl dicarbonate (716 mg, 3.28 mmol) were added to 2-(3-bromo-phenyl)-piperidine (750 mg, 3.12 mmol) in dichloromethane (10 mL). After overnight stirring at RT water (20 mL) was added to the solution. The resulting biphasic solution was separated into organic and aqueous phases. The aqueous phase was
 5 back extracted with dichloromethane (20 mL). The combined organic phases were dried over anhydrous sodium sulfate, filtered and concentrated to afford the title compound (1.03 g, 97%) as a pale yellow oil.



To a solution of **28a** (1.03 g, 3.03 mmol) and potassium vinyltrifluoroborate (488 mg, 3.64 mmol) in *n*-propanol (30 mL), under an atmosphere of nitrogen was added
 10 1,1'-bis(diphenylphosphino)ferrocenedichloropalladium(II), dichloromethane adduct (49 mg, 0.06 mmol) and triethylamine (306 mg, 422 μ L, 3.03 mmol). The reaction was heated to reflux and left to stir for 3 h before cooling to RT. The reaction mixture poured onto water and the resultant solution was extracted with diethyl ether (3 x 30 mL). The combined organics were dried over
 15 anhydrous sodium sulfate, filtered and concentrated. The resultant residue was purified by silica gel chromatography using a stepwise gradient of *iso*-hexanes/ethyl acetate 1:0 to 4:1 to afford the title compound (652 mg, 75%) as a pale yellow oil.




A solution of (2R, 3R)-1-((1R,5S)-10,10-dimethyl-3,3-dioxo-3 λ 6*-thia-4-aza-tricyclo[5.2.1.0*1,5*]dec-4-yl)-3-hydroxy-2-methyl-hept-6-en-1-one (250 mg, 0.703 mmol) in anhydrous dichloromethane (7 mL) was prepared and trimethyloxonium tetrafluoroborate (208 mg, 1.406 mmol) was added. The reaction mixture was stirred at RT for 15 h. The reaction mixture was treated with methanol (1 mL), then 2 M hydrochloric acid (20 mL) and saturated
 20 brine (20 mL). The mixture was extracted with ethyl acetate (3 x 15 mL) and the extract was dried over sodium sulfate, filtered and evaporated to give a yellow gum. The gum was purified by silica gel chromatography using *iso*-hexanes/ethyl acetate 4:1 to give the title compound
 25 (223 mg, 86%) as a colorless gum.




Compound 28d

A solution of 2 M lithium hydroxide in water (5 mL, 10 mmol) was added to a stirred solution of **28c** (223 mg, 0.60 mmol) in tetrahydrofuran (15 mL). The stirred mixture was heated to 60°C for 15 h. The reaction mixture was partially evaporated before adding 2 M hydrochloric acid (20 mL). The solution was extracted with ethyl acetate (3 x 15 mL). The extract was dried over sodium sulfate, filtered and evaporated to give a yellow gum (209 mg). The gum was purified by silica gel chromatography using *iso*-hexanes/ethyl acetate 3:1 to yield the title compound (68 mg, 66%) as a yellow gum.




Compound 28e

A solution of (S)-1-[(S)-2-((S)-2-*tert*-butoxycarbonylamino-3-methylbutyrylamino)-propionyl]-hexahydropyridazine-3-carboxylic acid 2,2,2-trichloroethyl ester (316 mg, 0.59 mmol) in anhydrous dichloromethane (10 mL) was cooled to 0°C under a nitrogen atmosphere before adding trimethylsilyl trifluoromethanesulfonate (160 μ L, 0.885 mmol). The reaction mixture was stirred at 0°C for 2 h before adding *N,N*-diisopropylethylamine (413 μ L, 2.36 mmol) to afford (S)-1-[(S)-2-((S)-2-amino-3-methyl-butrylamino)-propionyl]-hexahydro-pyridazine-3-carboxylic acid 2,2,2-trichloroethyl ester. The mixture was evaporated and the residue dissolved with (2R,3R)-3-methoxy-2-methyl-hept-6-enoic acid (162 mg, 0.94 mmol.) in acetonitrile (13 mL) and *N*-(3-dimethylaminopropyl)-*N'*-ethylcarbodiimide hydrochloride (250 mg, 1.32 mmol) and 1-hydroxybenzotriazole (220 mg, 1.32 mmol) were added. The reaction was stirred at RT for 15 h then evaporated to give a yellow oil. The oil was purified by silica gel chromatography using ethyl acetate to give the title compound (425 mg, 77%) as a white solid.




Compound 28f

A solution of **28e** (425 mg, 0.725 mmol) in tetrahydrofuran (20 mL) was prepared and zinc powder (0.48 g, 7.25 mmol) was added followed by an aqueous solution of ammonium acetate (1 M, 5 mL, 5 mmol). The reaction mixture was stirred at RT for 15 h. The reaction was filtered through hyflo-supercel washing through with ethyl acetate. The mixture was treated with hydrochloric acid (2 M, 30 mL) and the layers were separated. The aqueous layer was extracted with ethyl acetate (25 mL). The organic layers were combined, washed with brine, filtered and evaporated to give a colorless gum (299 mg) which was used directly without further purification.



Compound 28g


A solution of 2-(3-vinyl-phenyl)-piperidine-1-carboxylic acid *tert*-butyl ester (650 mg, 2.26 mmol) in dichloromethane (15 mL) was cooled to 0°C, before adding trimethylsilyl trifluoromethanesulfonate (569 µL, 3.39 mmol). The reaction mixture was stirred at 0°C for 1 h before adding *N,N*-diisopropylethylamine (1.6 mL, 5.24 mmol) to afford the 2-(3-vinyl-phenyl)-piperidine as a yellow solid. The solid was redissolved, along with **28f** (900 mg, 1.98 mmol) in acetonitrile (20 mL). The solution was cooled to 0°C, before adding 2-(1H-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyl uronium hexafluorophosphate methanaminium (1.05 g, 2.77 mmol) and *N,N*-diisopropylethylamine (1.4 mL, 7.92 mmol). The stirred reaction mixture was allowed to slowly warm to RT. After 2 h, the solvent was evaporated and the remaining residue dissolved in ethyl acetate (30 mL) and washed with water (3 x 30 mL). The organics were dried over anhydrous sodium sulfate, filtered and concentrated. The resultant residue was purified by silica gel chromatography using a stepwise gradient of *iso*-hexanes/ethyl acetate 1:0 to 0:1 to afford the title compound (1.25 g, 100%) as a yellow solid.



Compound 28

To a stirred solution of **28g** (1.05 mg, 1.68 mmol) in 1,2-dichloroethane (550 mL) was added Hoveyda-Grubbs 2nd generation catalyst (105 mg, 0.168 mmol). The solution was heated to 84°C and was left to stir for 2.5 h. The solvent was evaporated and the remaining residue was purified by silica gel chromatography using a stepwise gradient of *iso*-hexanes/ethyl acetate/acetone 1:0:0 to 0:9:1. This material was then subjected to a second round of silica gel chromatography using the same gradient to afford a yellow solid. A final round of silica gel chromatography using neat ethyl acetate and eluting purely by gravity afforded the title compound (175 mg, 18%) as a white solid as a 6:4 mixture of diastereoisomers. ¹H NMR (300 MHz, CDCl₃) δ 0.84 (d, *J* = 6.9 Hz, 3H), 0.91-1.01 (m, 6H), 1.24-1.37 (m, 6H), 1.47-2.03 (m, 14H), 2.06 (s, 2H), 2.14-2.25 (m, 1H), 2.43-2.77 (m, 4H), 3.48 (s, 1H), 3.56 (s, 2H), 3.63-3.86 (m, 1H), 4.00 (*app t*, *J* = 14.7 Hz, 1H), 4.12-4.17 (m, 1H), 4.53-4.65 (m, 1H), 5.45 (q, *J* = 7.2 Hz, 1H), 6.27-6.50 (m, 2H), 6.99-7.10 (m, 2H), 7.14-7.22 (m, 1H), 7.32-7.38 (m, 1H). LCMS (*m/z*) 596.4 [*M*+*H*], Tr = 2.51 min.

EXAMPLES 29 AND 30



Compound 29


Compound 30

Compound 28 (155 mg, 0.26 mmol) was dissolved in a 1:1 mixture of acetonitrile/water to a concentration of 7.8 mg/mL. This solution was then eluted through a reverse phase HPLC system fitted with a Phenomenex Gemini 10μ 110A, 250 x 21.2 mm column using an isocratic 2:3 acetonitrile/water flow at 20 mL/min. The mixture was resolved into the 2 separate

diastereoisomers. On concentration each separate diastereoisomer yielded a white solid. The stereochemistry of each isomer was not determined. First isomer eluted, Compound 29 (15 mg, 10%). ¹H NMR (300 MHz, CDCl₃) δ 0.89-1.05 (m, 8H), 1.20-1.44 (m, 10H), 1.51-1.79 (m, 3H), 1.83-1.99 (m, 4H), 2.03-2.27 (m, 3H), 2.33-2.67 (m, 3H), 2.76 (*app t*, *J* = 12.3 Hz, 1H), 3.09 (*app t*, *J* = 12.3 Hz, 1H), 3.22-3.38 (m, 1H), 3.47 (s, 3H), 3.54-3.62 (m, 1H), 4.17 (d, *J* = 11.7 Hz, 1H), 4.59 (d, *J* = 10.5 Hz, 1H), 5.39-5.53 (m, 1H), 6.01 (s, 1H), 6.09-6.23 (m, 1H), 6.30-6.55 (m, 2H), 6.99-7.23 (m, 4H). LCMS (*m/z*) 596.4 [M+H], Tr = 2.53 min.


Second isomer eluted, Compound 30 (22 mg, 14%). ¹H NMR (300 MHz, CD₃CN) δ 0.98-1.04 (m, 6H), 1.26-1.38 (m, 7H), 1.62-1.80 (m, 5H), 1.83-2.00 (m, 4H), 2.03-2.14 (m, 1H), 2.16-2.25 (m, 2H), 2.36-2.51 (m, 2H), 2.54-2.66 (m, 1H), 2.68-2.82 (m, 1H), 3.03-3.16 (m, 1H), 3.26-3.34 (m, 1H), 3.48 (s, 3H), 3.64-3.73 (m, 1H), 3.76-3.86 (m, 1H), 4.00 (*app t*, *J* = 9.0 Hz, 1H), 4.16 (d, *J* = 12.0 Hz, 1H), 4.55-4.66 (m, 1H), 5.41-5.53 (m, 1H), 5.97-6.06 (m, 1H), 6.10-6.22 (m, 1H), 6.35 (s, 1H), 6.38-6.49 (m, 1H), 7.00-7.11 (m, 2H), 7.14-7.23 (m, 3H). LCMS (*m/z*) 596.3 [M+H], Tr = 2.49 min.

EXAMPLE 31




Compound 31a


A cooled (0°C) solution of (R)-4-benzyl-3-[(R)-2-((S)-1-hydroxy-ethyl)-but-3-enoyl]-oxazolidin-2-one (233.6 mg, 0.807 mmol, prepared as described in *Org. Lett.* 2007, 9, 1635-1638) and imidazole (241.7 mg, 3.551 mmol) in *N,N*-dimethylformamide (2 mL) was treated with *tert*-butyldimethylsilyl chloride (158.2 mg, 1.049 mmol). After stirring for 24 h at RT, the reaction was quenched with saturated ammonium chloride. The aqueous layer was extracted with diethyl ether (2 x 20 mL). The organics were combined, dried over sodium sulfate, filtered and the volatiles were removed *in vacuo*. The residue was purified by silica gel chromatography using a 25 g Isolute cartridge eluted with a continuous gradient of *iso*-hexanes/ethyl acetate 1:0 to 4:1 to afford the title compound (294.6 mg, 90%) as a colorless oil.



A solution of **31a** (294.6 mg, 0.730 mmol), (R)-1-(7-bromo-quinolin-2-yl)-ethanol (184.0 mg, 0.730 mmol), palladium(II) acetate (32.8 mg, 0.146 mmol), tri-(*o*-toluyl)phosphine (44.4 mg, 0.146 mmol) in anhydrous 1,4-dioxane (10 mL) was treated with *N,N*-dicyclohexylmethylamine (250 μ L, 1.168 mmol). After stirring at 100°C for 5 h, the reaction was cooled to RT, diluted with dichloromethane and saturated sodium bicarbonate. The aqueous layer was extracted with dichloromethane. The organics were combined, filtered through a phase separator and the volatiles were removed *in vacuo*. The residue was purified by silica gel chromatography using a 25 g Isolute cartridge eluted with a continuous gradient of *iso*-hexanes/ethyl acetate 1:0 to 7:3 to afford the title compound (230.8 mg, 55%) as a colorless oil.




A cooled (0°C) solution of **31b** (230.8 mg, 0.401 mmol) in tetrahydrofuran/water (15 mL, 2:1) was subsequently treated with hydrogen peroxide (30% aqueous, 210 μ L, 2.005 mmol) and lithium hydroxide hydrate (33.7 mg, 0.803 mmol). After stirring for 2 h at 0°C, the reaction was quenched with sodium metabisulfite (765 mg, 4.1 mmol). After stirring for 3.5 h at RT the volatiles were removed *in vacuo*. The mixture was then diluted with water and the pH was adjusted with potassium carbonate. The aqueous layer was washed with dichloromethane (2 x 20 mL) and acidified with 2 M hydrochloric acid (pH ~1) then extracted with dichloromethane (3 x 30 mL). All the organics were combined, dried over sodium sulfate, filtered and the volatiles were removed *in vacuo*. The residue was purified by silica gel chromatography using a 25 g Isolute cartridge eluted with a continuous gradient of *iso*-hexanes/ethyl acetate 1:0 to 0:1 to afford the title compound (69.8 mg, 42%) as a white solid.



Compound 31d


Compound 31d was prepared in the same manner as **22e** using **31c** instead of (E)-4-[2-((R)-1-acetoxy-ethyl)-quinolin-7-yl]-2,2-dimethyl-but-3-enoic acid in 46% yield.



Compound 31


- 5 Compound 31 was prepared in the same manner as Compound 22 using **31d** instead of (S)-1-[(S)-2-((S)-2-((E)-4-[2-((R)-1-hydroxy-ethyl)-quinolin-7-yl]-2,2-dimethyl-but-3-enoylamino)-3-methyl-butrylamino)-propionyl]-hexahydro-pyridazine-3-carboxylic acid 2,2,2-trichloro-ethyl ester in 10% yield. ¹H NMR (300 MHz, CDCl₃) δ 1.00 (d, J = 6.7 Hz, 3H), 1.01 (d, J = 6.7 Hz, 3H), 1.22-1.44 (m, 4H), 1.64 (d, J = 7.1 Hz, 3H), 1.68-1.77 (m, 4H), 1.89-2.08 (m, 3H), 2.71-2.84 (m, 1H), 3.78-3.86 (m, 1H), 4.11 (dd, J = 8.2, 6.2 Hz, 1H), 4.17-4.25 (m, 2H), 4.38-4.47 (m, 1H), 5.68 (q, J = 7.1 Hz, 1H), 5.94 (q, J = 6.9 Hz, 1H), 6.47 (d, J = 16.5 Hz, 1H), 6.69 (dd, J = 16.5, 5.3 Hz, 1H), 7.40 (d, J = 8.5 Hz, 1H), 7.68 (s, 1H), 7.76 (dd, J = 8.7, 1.3 Hz, 1H), 7.81 (d, J = 8.7 Hz, 1H), 8.21 (d, J = 8.5 Hz, 1H). LCMS (m/z) 566.1 [M+H], Tr = 1.65 min.
- 10

EXAMPLE 32




Compound 32a

To a cooled (0°C) solution of ethyl propiolate (2 mL, 19.735 mmol) in dichloromethane (50 mL) was added dropwise morpholine (1.7 mL, 19.435 mmol). After stirring at RT for 1.5 h, the volatiles were removed *in vacuo* and the residue was purified by silica gel chromatography using a 50 g Isolute cartridge eluted with dichloromethane/methanol 20:1 to afford the title compound (3.5034 g, 97%) as a colorless oil.




Compound 32b

A cooled (0°C) solution of 4-bromoaniline (2.7 g, 15.7 mmol) in water (30 mL) was subsequently treated with concentrated hydrochloric acid (3.5 mL) and sodium nitrite (1.3 g, 18.840 mmol). After 20 min at 0°C, concentrated hydrochloric acid (5.3 mL) and sodium tetrafluoroborate (6.9 g, 62.847 mmol) were added. After 40 min at 0°C, the intermediate diazonium was filtered, washed with water, methanol and diethyl ether (2.1021 g) and was used without further purification. A solution of the diazonium (7.762 mmol) in acetonitrile (50 mL) was treated with **32a** (1.6614 g, 8.970 mmol). After 1 h at RT, silica gel was added. After stirring at RT for 16 h, the volatiles were removed *in vacuo* and the residue was purified by silica gel chromatography using a 100 g Isolute cartridge eluted with a continuous gradient of *iso*-hexanes/ethyl acetate 1:0 to 9:1 to afford the title compound (1.9811 g, 85%) as a highly coloured solid and as a mixture of tautomers.




Compound 32c


A solution of **32b** (1.9811 g, 6.623 mmol) in concentrated sulphuric acid (25 mL) was heated at 100°C for 3 h. After cooling to 0°C, the mixture was diluted with water (150 mL) and a brown solid was filtered off. The filtrate was extracted with diethyl ether (2 x 50 mL), dichloromethane (2 x 50 mL) and ethyl acetate (50 mL). The organics were combined, dried over sodium sulfate, filtered and the volatiles were removed *in vacuo* to provide the title compound (1.5094 g, 90%) as an orange solid that turned dark upon standing.



A cooled (0°C) solution of **32c** (1.5094 g, 5.964 mmol), *N,O*-dimethylhydroxylamine hydrochloride (755.9 mg, 7.753 mmol) and *N,N*-diisopropylethylamine (4.2 mL, 23.856 mmol) in acetonitrile (50 mL) was treated with 2-(1*H*-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyl uronium hexafluorophosphate methanaminium (3.175 g, 8.350 mmol). After stirring for 6 h at RT, the mixture was cooled to 0 °C and quenched with 1 M hydrochloric acid (60 mL). The aqueous layer was extracted with ethyl acetate (3 x 50 mL). The organics were combined, dried over sodium sulfate, filtered and the volatiles were removed *in vacuo*. The residue was purified by silica gel chromatography using a 100 g Isolute cartridge eluted with a continuous gradient of *iso*-hexanes/ethyl acetate 1:0 to 1:1 to afford the title compound (1.4229 g, 81%) as a bright yellow solid.




A cooled (-78°C) solution of **32d** (1.4229 g, 4.805 mmol) in tetrahydrofuran (50 mL) was treated with methylmagnesium chloride (3.2 mL, 9.610 mmol, 3 M in diethyl ether). After 1 h at -78°C and 3 h at 0°C, the reaction was quenched with saturated ammonium chloride (30 mL). The aqueous layer was extracted with ethyl acetate (3 x 50 mL). The organics were combined, dried over sodium sulfate, filtered and the volatiles were removed *in vacuo*. The residue was purified by silica gel chromatography using a 100 g Isolute cartridge eluted with a continuous gradient of *iso*-hexanes/ethyl acetate 1:0 to 4:1 to afford the title compound (751.5 mg, 62%) as a yellow solid.




A solution of **32e** (751.5 mg, 2.993 mmol) in tetrahydrofuran (30 mL) was treated with (S)-(-)-2-methyl-CBS-oxazaborolidine (3.6 mL, 3.592 mmol, 1 M in toluene). After 10 min at RT the mixture was cooled to -60°C and treated with borane-tetrahydrofuran complex (6 mL, 5.986 mmol, 1 M in tetrahydrofuran). After 1.5 h at -55°C to -30°C, the reaction was quenched with methanol (20 mL). After stirring at RT for 16 h, the volatiles were removed *in vacuo*. The residue

was purified by silica gel chromatography using a 50 g Isolute cartridge eluted with a continuous gradient of *iso*-hexanes/ethyl acetate 1:0 to 3:2 to afford the title compound (345.0 mg, 45%) as a yellow solid.




- 5 A solution of **32f** (345.0 mg, 1.363 mmol), palladium II acetate (61.2 mg, 0.273 mmol), tri-(*o*-toluyl)phosphine (83.1 mg, 0.273 mmol), 3-butenic acid *tert*-butyl ester (560 μ L, 3.407 mmol) and *N,N*-dicyclohexylmethylamine (470 μ L, 2.181 mmol) in anhydrous 1,4-dioxane (20 mL) was heated at 100°C for 1.7 h. After cooling to RT the volatiles were removed *in vacuo*. The residue was purified by silica gel chromatography using a 50 g Isolute cartridge eluted with a continuous gradient of *iso*-hexanes/ethyl acetate 1:0 to 2:3 to afford the title compound (244.1 mg, 57%) as a yellow solid.
- 10



- A cooled (0°C) solution of **1e** (422.0 mg, 0.793 mmol) in tetrahydrofuran/water (25 mL, 4:1) was treated with lithium hydroxide hydrate (67.0 mg, 1.587 mmol). After stirring at 0°C for 1.5 h, the reaction was quenched with 1 M hydrochloric acid (20 mL). The aqueous layer was extracted with ethyl acetate (2 x 30 mL). The organics were combined, dried over sodium sulfate, filtered and the volatiles were removed *in vacuo* then residual trichloroethanol was azeotroped off with toluene (3 x) to give the intermediate acid as a white solid which was then combined with **32g** (244.1 mg, 0.793 mmol), 4-dimethylaminopyridine (97.0 mg, 0.793 mmol), *N*-(3-dimethylaminopropyl)-*N'*-ethylcarbodiimide hydrochloride (243.3 mg, 1.269 mmol) and dichloromethane (20 mL). After stirring at RT for 16 h, the reaction was quenched with dilute hydrochloric acid. The aqueous layer was extracted with dichloromethane (30 mL). The organics were combined and filtered through a phase separator and the volatiles were removed *in vacuo*. The residue was purified by silica gel chromatography using a 50 g Isolute cartridge
- 15
- 20

eluted with a continuous gradient of *iso*-hexane/ethyl acetate 1:0 to 1:2 to afford the title compound (164.4 mg, 30% over 2 steps) as a yellow glass.




Compound 32

A cooled (0 °C) solution of **32h** (164.4 mg, 0.236 mmol) in anhydrous dichloromethane (20 mL) was treated with trimethylsilyl methanesulfonate (170 μ L, 0.944 mmol). After stirring for 1 h at 0°C, the reaction was quenched with *N,N*-diisopropylethylamine (330 μ L, 1.888 mmol) and the volatiles were removed *in vacuo*. A cooled (0°C) solution of the crude amino acid in acetonitrile (80 mL) was subsequently treated with *N,N*-diisopropylethylamine (330 μ L, 1.888 mmol) and 2-(1H-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyl uronium hexafluorophosphate methanaminium (125.6 mg, 0.330 mmol). After stirring at RT for 2.5 h, the volatiles were removed *in vacuo*. The residue was purified by silica gel chromatography using a 50 g Isolute cartridge eluted with a continuous gradient of *iso*-hexanes/ethyl acetate 1:0 to 0:1 to afford the final compound (170.7 mg) as a mixture. Reverse phase preparative HPLC which was eluted with a gradient of water/acetonitrile 95:5 to 0:100 provided the title compound (30.6 mg, 25%) as a white solid. ^1H NMR (300 MHz, CD_3OD) δ 1.00 (d, J = 6.7 Hz, 3H), 1.03 (d, J = 6.7 Hz, 3H), 1.61 (d, J = 7.3 Hz, 3H), 1.66-2.07 (m, 8H), 2.68-2.81 (m, 1H), 2.97-3.07 (m, 1H), 3.35-3.47 (m, 1H), 3.76-3.89 (m, 1H), 4.28 (d, J = 9.4 Hz, 1H), 4.37-4.47 (m, 1H), 4.53-4.67 (m, 1H), 5.52 (q, J = 7.1 Hz, 1H), 6.36-6.51 (m, 2H), 6.72 (d, J = 16.0 Hz, 1H), 7.61 (s, 1H), 7.97 (dd, J = 8.9, 1.6 Hz, 1H), 8.06 (s, 1H), 8.38 (d, J = 8.9 Hz, 1H). LCMS (m/z) 523.1 [$\text{M}+\text{H}$], Tr = 1.65 min.


20

EXAMPLES 33 AND 34




Compound 33a

A cooled (0°C) solution of (S)-tetrahydro-pyridazine-1,2,3-tricarboxylic acid 1,2-di-*tert*-butyl ester 3-(2,2,2-trichloro-ethyl) ester (5.5297 g, 11.975 mmol) in tetrahydrofuran/methanol (50 mL, 1:1) was treated with tetrabutylammonium fluoride (23.9 mL, 23.950 mmol, 1 M in tetrahydrofuran). After stirring at RT for 24 h, the volatiles were removed *in vacuo*. The residue was dissolved in diethyl ether and saturated sodium bicarbonate. The aqueous layer was extracted with diethyl ether (50 mL). The organics were combined, dried over sodium sulfate, filtered and the volatiles were removed *in vacuo*. The residue was purified by silica gel chromatography using a 100 g Isolute cartridge eluted with a continuous gradient of *iso*-hexanes/ethyl acetate 1:0 to 9:1 where the mixed fractions were further purified by silica gel chromatography using a 50 g Isolute cartridge eluted with a continuous gradient of *iso*-hexanes/ethyl acetate 1:0 to 4:1 to afford the title compound (3.9765 g, 96%) as a colorless oil and as a mixture of rotamers.




A cooled (-78°C) solution of **33a** (1.4112 g, 4.097 mmol) in anhydrous tetrahydrofuran (10 mL) was treated with lithium bis(trimethylsilyl)amide (6.2 mL, 6.146 mmol, 1 M in tetrahydrofuran). After stirring for 1.25 h at -78°C, the mixture was treated with iodomethane (640 µL, 10.242 mmol). After stirring for 1 h at -78°C, 1 h at 0°C and 1 h at RT, the reaction was quenched with pH 7 buffer at 0°C. The aqueous was extracted with dichloromethane (2 x 50 mL). The organics were combined, dried over sodium sulfate, filtered and the volatiles were removed *in vacuo*. The residue was purified by silica gel chromatography using a 50 g Isolute cartridge eluted with a continuous gradient of *iso*-hexanes/ethyl acetate 1:0 to 4:1 to afford the title compound (760.7 mg, 52%) as a colorless oil and as a mixture of rotamers.




A cooled (0°C) solution of (S)-2-*tert*-butoxycarbonylamino-3-methyl-butyl-benzoic acid (3.020 g, 13.908 mmol), (S)-2-amino-propionic acid methyl ester hydrochloride (1.9413 g, 13.908 mmol) and *N,N*-diisopropylethylamine (17.1 mL, 55.632 mmol) in acetonitrile (50 mL) was treated with

2-(1H-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyl uronium hexafluorophosphate
methanaminium (7.403 g, 19.471 mmol). After stirring at RT for 6 h, the reaction was quenched
with hydrochloric acid (1 M, 100 mL). The aqueous layer was extracted with ethyl acetate (2 x
100 mL). The organics were combined, dried over sodium sulfate, filtered and the volatiles were
5 removed *in vacuo*. The residue was purified by silica gel chromatography using a 100 g Isolute
cartridge eluted with a continuous gradient of *iso*-hexanes/ethyl acetate 1:0 to 3:2 to afford the
title compound (4.0996 g, 97%) as a white solid.




A cooled (0°C) solution of **33c** (317.0 mg, 1.048 mmol) in tetrahydrofuran/water (12 mL,
10 5:1) was treated with lithium hydroxide hydrate (88.0 mg, 2.096 mmol). After stirring at 0°C for
2.5 h, the reaction was quenched with hydrochloric acid (1 M, 30 mL). The aqueous layer was
extracted with dichloromethane (2 x 30 mL). The organics were combined, filtered through a
phase separator and the volatiles were removed *in vacuo* to provide crude acid as a white solid.




A cooled (0°C) solution of **33b** (376.0 mg, 1.049 mmol) in anhydrous dichloromethane
15 (15 mL) was treated with trifluoroacetic acid (5 mL). After stirring at 0°C for 30 min and RT for 2
h, trifluoroacetic acid (4 mL) was added. After 1 h, the volatiles were removed *in vacuo* and the
residual trifluoroacetic acid azeotroped off with toluene (3 x) to provide the bis- trifluoroacetic
acid ammonium salt as an off-white solid. A cooled (0°C) solution of the bis- trifluoroacetic acid
20 ammonium salt and crude (S)-2-((S)-2-*tert*-butoxycarbonylamino-3-methyl-butylamino)-
propionic acid as prepared in the previous step in anhydrous dichloromethane (15 mL) was
subsequently treated with *N*-methylmorpholine (580 µL, 5.240 mmol) and O-(benzotriazol-1-yl)-
N,N,N',N'-tetramethyluronium hexafluorophosphate (596.2 mg, 1.572 mmol). After stirring for 19
h at RT, the reaction was quenched with hydrochloric acid (1 M, 30 mL). The aqueous was
25 extracted with dichloromethane (2 x 30 mL). The organics were combined, washed with
saturated aqueous sodium bicarbonate and filtered through a phase separator. The volatiles
were removed *in vacuo* and the residue was purified by silica gel chromatography using a 50 g

Isolute cartridge eluted with a continuous gradient of *iso*-hexanes/ethyl acetate 1:0 to 1:2 to afford the title compound (373.1 mg, 83% over 2 steps) as a white foam and as a mixture of diastereoisomers.



Compound 33f


- 5 A cooled (0°C) solution of **33e** (373.1 mg, 0.871 mmol) in anhydrous dichloromethane (10 mL) was treated with trimethylsilyl methanesulfonate (310 µL, 1.741 mmol). After stirring for 1 h at 0°C, the reaction was quenched with *N,N*-diisopropylethylamine (610 µL, 3.484 mmol) and the volatiles were removed *in vacuo*. A cooled (0°C) solution of the crude amine and (*E*)-
- 10 (2*R*,3*R*)-3-methoxy-2-methyl-hex-4-enoic acid (137.8 mg, 0.871 mmol) in anhydrous acetonitrile (15 mL) was subsequently treated with *N,N*-diisopropylethylamine (610 µL, 3.484 mmol) and 2-(1*H*-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyl uronium hexafluorophosphate methanaminium (463.7 mg, 1.219 mmol). After stirring at RT for 18 h, the reaction was quenched with hydrochloric acid (1 M, 30 mL). The aqueous layer was extracted with ethyl acetate (2 x 30 mL). The organics were combined, washed with saturated sodium bicarbonate (20 mL), dried over
- 15 sodium sulfate, filtered and the volatiles were removed *in vacuo*. The residue was purified by silica gel chromatography using a 25 g Isolute cartridge eluted with a continuous gradient of *iso*-hexanes/ethyl acetate 1:0 to 1:2 to 1:4 to afford the title compound (332.2 mg, 81%) as a colorless solid and as a mixture of diastereoisomers.



Compound 33g

- 20 Compound **33g** was prepared in the same manner as (*S*)-1-[(*S*)-2-[(*S*)-2-*tert*-butoxycarbonylamino-3-methyl-butyrylamino)-propionyl]-hexahydro-pyridazine-3-carboxylic acid (*R*)-1-[6-[(*E*)-3-*tert*-butoxycarbonyl-propenyl]-cinnolin-3-yl]-ethyl ester using **33f** and (*R*)-1-(3-

vinyl-isoquinolin-6-yl)-ethanol instead of **1e** and (E)-4-[3-((R)-1-hydroxy-ethyl)-cinnolin-6-yl]-but-3-enoic acid *tert*-butyl ester in 30% yield over 2 steps.




- 5 A solution of **33g** (136.2 mg, 0.214 mmol) in toluene (70 mL) was degassed by bubbling N₂ gas through for 20 min. Hoveyda-Grubbs 2nd generation catalyst (27 mg, 0.043 mmol) was added and the mixture was refluxed for 2.5 h. More Hoveyda-Grubbs 2nd generation catalyst (20 mg) was added and after stirring at reflux for 1 h, the mixture was cooled to RT, the volatiles were removed *in vacuo* and the residue was purified by silica gel chromatography using a 25 g
- 10 Isolute cartridge eluted with a continuous gradient of ethyl acetate/acetone 1:0 to 10:1 to afford both diastereoisomers in separate mixtures. Both mixtures were purified by silica gel chromatography using a 10 g Isolute cartridge eluted *by gravity* with a continuous gradient of ethyl acetate/acetone 1:0 to 10:1 to provide the more polar diastereoisomer (19.7 mg, 15%) as a white solid. The less polar diastereomer was further purified by preparative TLC eluted with
- 15 ethyl acetate (4 elutions) and subsequent preparative TLC eluted with *iso*-hexanes/acetone 7:3 (2 elutions) to afford the title compound (7.6 mg, 6%) as a white solid. Relative stereochemistry was not assigned.

More polar diastereoisomer Compound 33: ¹H NMR (300 MHz, CDCl₃) δ 0.83-1.07 (m, 7H), 1.41 (d, *J* = 7.1 Hz, 3H), 1.47 (d, *J* = 6.9 Hz, 3H), 1.50 (s, 3H), 1.70 (d, *J* = 6.7 Hz, 3H), 1.75-1.90 (m, 2H), 1.96-2.15 (m, 3H), 2.64 (dd, *J* = 7.6, 3.1 Hz, 1H), 2.70-2.84 (m, 1H), 3.42 (s, 3H), 3.90 (dd, *J* = 8.7, 3.1 Hz, 1H), 4.00 (*app t*, *J* = 8.0 Hz, 1H), 4.10 (s, 1H), 4.58-4.69 (m, 1H), 6.00 (dd, *J* = 9.4, 7.1 Hz, 1H), 6.22 (q, *J* = 6.2 Hz, 1H), 6.51 (d, *J* = 9.8 Hz, 1H), 6.93 (s, 1H), 7.13-7.23 (m, 1H), 7.34-7.43 (m, 1H), 7.90-7.97 (m, 2H), 8.47 (s, 1H). LCMS (*m/z*) 594.3 [M+H], Tr = 1.87 min.


25 Less polar diastereoisomer Compound 34: ¹H NMR (300 MHz, CD₃CN) δ -0.51- -0.39 (m, 5H), -0.30 (d, *J* = 6.9 Hz, 3H), 0.06 (d, *J* = 7.1 Hz, 3H), 0.15 (d, *J* = 6.9 Hz, 3H), 0.19 (s, 0.25-0.34 (m, 4H), 0.38-0.45 (m, 1H), 0.81-1.02 (m, 2H), 1.14-1.42 (m, 2H), 2.69 (dd, *J* = 8.7, 2.9 Hz, 1H), 3.00-3.10 (m, 2H), 3.45 (s, 3H), 4.32 (*app pentet*, *J* = 7.1 Hz, 1H), 4.76 (q, *J* = 6.7

Hz, 1H), 5.53 (d, $J = 16.3$ Hz, 1H), 5.85 (dd, $J = 16.1, 8.9$ Hz, 1H), 6.20 (dd, $J = 8.5, 1.6$ Hz, 1H), 6.41 (d, $J = 8.9$ Hz, 1H), 6.65-6.75 (m, 2H), 6.91 (s, 1H), 7.11 (d, $J = 7.3$ Hz, 1H), 7.79 (s, 1H). LCMS (m/z) 594.3 [$M+H$], $T_r = 1.83$ min.


5

EXAMPLE 35

A cooled (0°C) solution of 2-fluoro-5-hydroxybenzoic acid (1.0051 g, 6.438 mmol), *N,O*-dimethylhydroxylamine hydrochloride (1.2560 g, 12.856 mmol) and triethylamine (3.6 mL, 25.752 mmol) in dichloromethane (35 mL) was treated with *O*-(benzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium hexafluorophosphate (3.663 g, 9.658 mmol). After stirring at RT for 20 h, the reaction was quenched at 0°C with hydrochloric acid (2 M, 30 mL). The emulsion was filtered on Celite then the aqueous layer was extracted with dichloromethane (50 mL). The organics were combined, filtered through a phase separator and the volatiles were removed *in vacuo*. The residue was purified by silica gel chromatography using a 50 g Isolute cartridge eluted with a continuous gradient of toluene/diethyl ether 1:0 to 1:2 to afford the title compound (260.5 mg, 20%) as a colorless oil.




A solution of **35a** (260.5 mg, 1.308 mmol) in acetone (20 mL) was subsequently treated with potassium carbonate (903.8 mg, 6.539 mmol) and allylbromide (340 μL , 3.924 mmol). After stirring at RT for 24 h, the reaction was quenched with water (20 mL). The aqueous layer was extracted with dichloromethane (2 x 30 mL). The organics were combined, filtered through a phase separator and the volatiles were removed *in vacuo*. The residue was purified by silica gel chromatography using a 25 g Isolute cartridge eluted with a continuous gradient of *iso*-hexanes/ethyl acetate 1:0 to 3:2 to afford the title compound (289.7 mg, 92%) as a colorless oil.




25

A cooled (-78°C) solution of **35b** (289.7 mg, 1.210 mmol) in tetrahydrofuran (10 mL) was treated with methylmagnesium chloride (810 μL , 2.422 mmol, 3 M in diethyl ether). After 1.25 h at -78°C , 2 h at 0°C and 16 h at RT, methylmagnesium chloride (810 μL , 2.422 mmol, 3 M in diethyl ether) was added. After 2.5 h at -78°C and 1.2 h at RT, the reaction was quenched with silica gel and the volatiles were removed *in vacuo*. The residue was purified by silica gel chromatography using a 25 g Isolute cartridge eluted with a continuous gradient of *iso*-hexanes/ethyl acetate 1:0 to 9:1 to afford the title compound (196.7 mg, 84%) as a colorless oil.




A solution of **35c** (196.7 mg, 1.013 mmol) in tetrahydrofuran (15 mL) was treated with (S)-(-)-2-methyl-CBS-oxazaborolidine (1.2 mL, 1.215 mmol, 1 M in toluene). After 10 min at RT the mixture was cooled to -50°C and treated with borane tetrahydrofuran complex (2.1 mL, 2.026 mmol, 1 M in tetrahydrofuran). After 1.5 h at -50°C to -40°C , the reaction was quenched with methanol (6 mL). After stirring at RT for 22 h, the volatiles were removed *in vacuo*. The residue was purified by silica gel chromatography using a 25 g Isolute cartridge eluted with a continuous gradient of *iso*-hexanes/ethyl acetate 1:0 to 4:1 to afford the title compound (179.4 mg, 90%) as a white solid.




A cooled (0°C) solution of **1e** (1.9973 g, 3.755 mmol) in anhydrous dichloromethane (40 mL) was treated with trimethylsilyl trifluoromethanesulfonate (1.4 mL, 7.510 mmol). After 30 min at 0°C , the reaction mixture was treated with *N,N*-diisopropylethylamine (2.6 mL, 15.020 mmol) and the volatiles were removed *in vacuo* to afford the corresponding amine. A cooled (0°C) solution of this amine, (E)-(2R, 3R)-2-methylhex-6-enoic acid (428.6 mg, 3.755 mmol, prepared as described in *Synlett* 2002, 12, pp2039-2040) and *N,N*-diisopropylethylamine (2.6 mL, 15.020 mmol) in acetonitrile (50 mL) was treated with 2-(1H-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate methanaminium (1.999 g, 5.257 mmol). After stirring at RT for 20

h, the reaction was quenched with hydrochloric acid (1 M, 100 mL). The aqueous layer was extracted with ethyl acetate (2 x 50 mL). The organics were combined, washed with saturated sodium bicarbonate, dried over sodium sulfate, filtered and the volatiles were removed *in vacuo*. The residue was purified by silica gel chromatography using a 50 g Isolute cartridge eluted with a continuous gradient of *iso*-hexanes/ethyl acetate 1:0 to 1:4 to afford the title compound
 5 (1.6735 g, 84%) as a brown foam.



A solution of **35e** (534.7 mg, 1.013 mmol) in tetrahydrofuran (20 mL) was subsequently treated with zinc powder (1.457 g, 22.286 mmol) and a solution of ammonium acetate (1.171 g,
 10 15.195 mmol) in water (5 mL). After stirring at RT for 24 h, the mixture was filtered through Celite. The solid was rinsed with saturated potassium bisulfate and ethyl acetate. The pH of the filtrate was adjusted with 2 M hydrochloric acid then the aqueous layer was extracted with ethyl acetate (2 x 50 mL). The organics were combined, dried over sodium sulfate, filtered and the volatiles were removed *in vacuo*. Residual acetic acid was azeotroped off with toluene (3x) to
 15 provide the corresponding acid as a white solid. A solution of the crude acid, (R)-1-(5-allyloxy-2-fluoro-phenyl)-ethanol (179.4 mg, 0.914 mmol) and 4-dimethylaminopyridine (123.7 mg, 1.013 mmol) in dichloromethane (15 mL) was treated with *N*-(3-dimethylaminopropyl)-*N'*-ethylcarbodiimide hydrochloride (310.8 mg, 1.621 mmol). After stirring at RT for 20 h, the reaction was quenched at 0°C with hydrochloric acid (2 M, 15 mL). The aqueous layer was
 20 extracted with ethyl acetate (2x). The organics were combined, dried over sodium sulfate, filtered and the volatiles were removed *in vacuo*. The residue was purified by silica gel chromatography using a 50 g Isolute cartridge eluted with a continuous gradient of *iso*-hexanes/ethyl acetate 1:0 to 1:3 to afford the title compound (310.5 mg, 59% over 2 steps) as a white solid.



Compound 35


A solution of **35f** (310.5 mg, 0.540 mmol) in dichloroethane (180 mL) was treated with Hoveyda-Grubbs 2nd generation catalyst (67.7 mg, 0.108 mmol). After stirring at reflux for 2 h, the reaction was cooled to RT and the volatiles were removed *in vacuo*. The residue was

5 purified by silica gel chromatography using a 50 g Isolute cartridge eluted with a continuous gradient of *iso*-hexanes/ethyl acetate 1:0 to 0:1 then using a 25 g Isolute cartridge eluted *by gravity* with a continuous gradient of *iso*-hexanes/ethyl acetate 1:0 to 0:1 to afford the title compound (109.9 mg) in a mixture. Purification by preparative TLC eluted with *iso*-hexanes/acetone 3:1 (3 elutions) followed by silica gel chromatography using a 25 g Isolute

10 cartridge eluted with a continuous gradient of *iso*-hexanes/acetone 1:0 to 3:2 provided the title compound (58.7 mg, 20%) as a white solid. ¹H NMR (300 MHz, CD₃CN) δ 0.85-0.97 (m, 6H), 1.17-1.22 (m, 3H), 1.28-1.39 (m, 4H), 1.44-1.63 (m, 6H), 1.70-1.82 (m, 1H), 1.83-1.92 (m, 1H), 2.30-2.45 (m, 3H), 3.62-3.73 (m, 2H), 3.89 (*app t*, $J = 8.5$ Hz, 1H), 4.28 (d, $J = 8.0$ Hz, 1H), 4.40-4.60 (m, 2H), 5.32 (*app pentet*, $J = 6.9$ Hz, 1H), 5.62-5.74 (m, 1H), 5.77-5.89 (m, 1H), 6.02

15 (q, $J = 6.7$ Hz, 1H), 6.40 (d, $J = 7.8$ Hz, 1H), 6.76-6.85 (m, 2H), 7.01 (*app t*, $J = 9.4$ Hz, 1H), 7.19 (*br s*, 1H). LCMS (m/z) 547.2 [M+H], Tr = 2.39 min.

EXAMPLE 36




Compound 36


20 To a solution of Compound 35 (44.0 mg, 0.080 mmol) in ethyl acetate (5 mL) was added palladium on carbon (10%, 5 mg). The atmosphere was purged of oxygen. After stirring at RT under an atmosphere of hydrogen for 2.5 h, the volatiles were removed *in vacuo* and the

residue was purified by silica gel chromatography using a 20 g Isolute cartridge eluted with a continuous gradient of *iso*-hexanes/acetone 1:0 to 3:2 followed by preparative TLC eluted with *iso*-hexanes/acetone 3:2 to afford the final compound (19.0 mg, 43%) as a white solid. ¹H NMR (300 MHz, CDCl₃) δ 0.85-0.99 (m, 8H), 1.14-1.22 (m, 4H), 1.26-1.42 (m, 4H), 1.48 (d, *J* = 6.7 Hz, 1H), 1.52-1.84 (m, 8H), 2.28-2.42 (m, 1H), 3.61-3.72 (m, 1H), 3.75-4.10 (m, 4H), 4.25 (d, *J* = 9.4 Hz, 1H), 5.26 (*app* pentet, *J* = 7.1 Hz, 1H), 6.02 (q, *J* = 6.7 Hz, 1H), 6.34 (d, *J* = 8.5 Hz, 1H), 6.75-6.86 (m, 2H), 6.95-7.12 (m, 2H). LCMS (*m/z*) 549.3 [M+H], Tr = 2.54 min.

EXAMPLE 37: (E)-(2R,5S,11S,14S,17R,18R)-2,14-Diisopropyl-18-methoxy-11,17-dimethyl-22-oxa-3,9,12,15,28-pentaaza-tricyclo[21.3.1.1*5,9*]octacos-1(27),19,23,25-tetraene-4,10,13,16-tetraone




37a was prepared in the same manner as **35b** using 3-hydroxybenzaldehyde instead of 2-fluoro-5-hydroxy-*N*-methoxy-*N*-methyl-benzamide in 78% yield.



A cooled (0°C) suspension of **37a** (1.36 g, 8.434 mmol) and magnesium sulfate (5 g) in dichloromethane (40 mL) was treated with D-valinol (870.0 mg, 8.434 mmol). After stirring at 0°C to RT for 23 h, the mixture was filtered and the volatiles were removed *in vacuo*. The residue was dissolved in anhydrous dichloromethane (40 mL) and was subsequently treated with triethylamine (1.3 mL, 9.277 mmol) and a solution of trimethylsilyl chloride (9.3 mL, 9.277 mmol, 1 M in dichloromethane). After stirring at RT for 24 h, the volatiles were removed *in vacuo* and the residue was triturated with diethyl ether/*iso*-hexane (100 mL, 1:1). The white solid was filtered off and the filtrate was evaporated to dryness to provide the intermediate imine. In a cooled (-40°C) 3-neck round-bottom flask, equipped with a nitrogen line and an addition funnel was introduced anhydrous diethyl ether (25 mL) and a solution of *iso*-propyllithium (29 mL, 20.242 mmol, 0.7 M in pentane). To this mixture, a solution of the imine in anhydrous diethyl ether (25 mL) was added dropwise over 15 min. After stirring at -40°C for 2.5 h, the reaction


was quenched with hydrochloric acid (2 M, 50 mL) and the mixture was allowed to warm to RT. The acidic aqueous layer was basified with NaOH pellets at 0°C and then extracted with diethyl ether (2 x). The organics were combined, dried over sodium sulfate, filtered and the volatiles were removed *in vacuo*. The residue was purified by silica gel chromatography using a 50 g Isolute cartridge eluted with a continuous gradient of *iso*-hexanes/ethyl acetate 1:0 to 1:1 to afford the title compound (470.1 mg, 19% over 3 steps) as a colorless oil.



Compound 37c

A solution of **37b** (470.1 mg, 1.613 mmol) in methanol (10 mL) and aqueous methylamine (3 mL, 40 wt% in water) was treated with periodic acid (1.213 g, 5.323 mmol).


After stirring for 24 h at RT more aqueous methylamine (4 mL, 40 wt% in water) and periodic acid (1.213 g, 5.323 mmol) were added. After stirring for 17 h at RT the mixture was filtered over Celite and the solid rinsed with methanol. The volatiles were removed *in vacuo* and the residue was partitioned between water and diethyl ether. The aqueous layer was extracted with diethyl ether, the organics were combined, dried over sodium sulfate, filtered and the volatiles were removed *in vacuo*. The residue was purified by silica gel chromatography using a 25 g Isolute cartridge eluted with ethyl acetate to afford the title compound (191.8 mg, 58%) as a light yellow oil.



Compound 37d


A cooled (0°C) solution of (S)-1-[(S)-2-((S)-2-*tert*-butoxycarbonylamino-3-methylbutyrylamino)-propionyl]-hexahydro-pyridazine-3-carboxylic acid (324.4 mg, 0.810 mmol), **37c** (166.3 mg, 0.810 mmol) and *N,N*-diisopropylethylamine (560 µL, 3.240 mmol) in acetonitrile (15 mL) was treated with 2-(1H-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyl uronium hexafluorophosphate methanaminium (431.2 mg, 1.134 mmol). After stirring at RT for 20 h, the reaction was quenched with hydrochloric acid (2 M, 25 mL) at 0 °C. The aqueous layer was

extracted with ethyl acetate (2 x). The organics were combined, dried over sodium sulfate, filtered and the volatiles were removed *in vacuo*. The residue was purified by silica gel chromatography using a 50 g Isolute cartridge eluted with a continuous gradient of *iso*-hexanes/ethyl acetate 1:0 to 0:1 to afford the title compound (230.5 mg, 48%) as a solid.



Compound 37e was prepared in the same manner as 1-[(S)-2-[(S)-2-((E)-(2R,3R)-3-methoxy-2-methyl-hex-4-enoylamino)-3-methyl-butylamino]-propionyl]-3-methyl-hexahydro-pyridazine-3-carboxylic acid methyl ester using **37d** instead of 1-[(S)-2-[(S)-2-*tert*-butoxycarbonylamino-3-methyl-butylamino]-propionyl]-3-methyl-hexahydro-pyridazine-3-carboxylic acid methyl ester in 30% yield over 2 steps.

10




Compound 37 was prepared in the same manner as **25** using **37e** instead of (S)-1-[(S)-2-[(S)-2-((2R,3R)-3-methoxy-2-methyl-hept-6-enoylamino)-3-methyl-butylamino]-propionyl]-hexahydro-pyridazine-3-carboxylic acid methyl-((R)-6-vinyl-indan-1-yl)-amide in 43% yield. ¹H NMR (300 MHz, *d*₆-DMSO) δ 0.74-0.91 (m, 12H), 1.15 (d, *J* = 7.1 Hz, 3H), 1.28 (d, *J* = 7.3 Hz, 3H), 1.40-1.64 (m, 3H), 1.71-1.88 (m, 3H), 1.98 (*app* sextet, *J* = 6.7 Hz, 1H), 2.66-2.75 (m, 2H), 3.38-3.53 (m, 1H), 3.89 (dd, *J* = 6.7, 2.9 Hz, 1H), 4.10 (*app* t, *J* = 8.9 Hz, 1H), 4.14-4.22 (m, 1H), 4.42 (qd, *J* = 10.0, 4.2 Hz, 1H), 4.57 (*app* t, *J* = 8.2 Hz, 1H), 4.76 (d, *J* = 11.8 Hz, 1H), 5.23 (*app* t, *J* = 7.3 Hz, 1H), 5.69-5.90 (m, 2H), 6.77-6.90 (m, 3H), 7.05 (d, *J* = 9.4 Hz, 1H), 7.16-7.24 (m, 1H), 8.01 (d, *J* = 8.5 Hz, 1H), 8.14 (d, *J* = 8.5 Hz, 1H). LCMS (*m/z*) 586.3 [M+H], Tr = 2.34 min.


15

20


EXAMPLE 38



- To a mixture of (R)-1-(3-bromo-phenyl)-ethanol (201 mg, 1.00 mmol), 4-(2-ethoxy-2-oxoethoxy)benzeneboronic acid (Acros Organics, 224 mg, 1.00 mmol) in 1,2-dimethoxyethane (4 mL) were added potassium carbonate (276 mg, 2.00 mmol) and water (1 mL). The mixture was stirred at RT and tetrakis(triphenylphosphine) palladium(0) (58 mg, 0.05 mmol) was added then the reaction mixture was heated at 100°C in a microwave reactor for 30 min. The reaction mixture was then diluted with ethyl acetate and water. The layers were separated and the organics washed with water and brine, dried over magnesium sulfate, filtered and evaporated.
- The residue was purified by silica gel chromatography using a gradient of *iso*-hexanes/ethyl acetate 9:1 to 7:3 to afford the title compound (230 mg, 76%) as an oil.




- A solution of **38a** (230 mg, 0.77 mmol) in tetrahydrofuran (4 mL) was stirred at 5°C under nitrogen, a solution of lithium hydroxide monohydrate (92 mg, 1.54 mmol) was added and the reaction mixture was stirred at 5°C for 2 h and then at RT overnight. The solution was acidified with 2 M hydrochloric acid and extracted with ethyl acetate (2 x). The organic extracts were combined, washed with water and brine, dried over magnesium sulfate and evaporated to afford the title compound (175 mg, 84%) as a white solid.




- A solution of **1e** (372 mg, 0.70 mmol) in dichloromethane (15 mL) was cooled in an ice-water bath under nitrogen. Trimethylsilyl trifluoromethanesulfonate (0.18 mL, 1.05 mmol) was

added dropwise, and the resulting solution was stirred for 1 hour. Cold saturated aqueous sodium hydrogen carbonate solution (15 mL) was added and the mixture was stirred at 0°C for 15 min. The organic layer was separated, washed with brine, dried over magnesium sulfate and evaporated to afford (S)-1-[(S)-2-((S)-2-amino-3-methyl-butyrylamino)-propionyl]-hexahydro-pyridazine-3-carboxylic acid 2,2,2-trichloro-ethyl ester (0.70 mmol) which was used without further purification. A solution of (S)-1-[(S)-2-((S)-2-amino-3-methyl-butyrylamino)-propionyl]-hexahydro-pyridazine-3-carboxylic acid 2,2,2-trichloro-ethyl ester (0.70 mmol) in acetonitrile (15 mL) was stirred at 0°C under nitrogen. **38b** (175 mg, 0.64 mmol) and 1-hydroxybenzotriazole hydrate (123 mg, 0.64 mmol, wetted with not less than 20 wt. % water) were added, followed by *N*-(3-dimethylaminopropyl)-*N'*-ethylcarbodiimide hydrochloride (173 mg, 0.90 mmol) and the reaction mixture was stirred at RT for 2 h. The solvent was evaporated. The residue was dissolved in ethyl acetate and the solution was washed with water (3 x) followed by brine, dried over magnesium sulfate and evaporated. The residue was purified by silica gel chromatography using a gradient of *iso*-hexanes/ethyl acetate 1:2 to 0:1. The product was triturated with diethyl ether and dried to afford the title compound (367 mg, 83%) as a white solid.




A solution of **38c** (343 mg, 0.50 mmol) in tetrahydrofuran (10 mL) was stirred at RT under nitrogen. Zinc powder (715 mg, 11 mmol) was added followed by a solution of ammonium acetate (578 mg, 7.50 mmol) in water (5 mL). The reaction mixture was stirred at RT under nitrogen for 70 h. The reaction mixture was filtered through Celite and the filter pad was washed with ethyl acetate and 2 M aqueous hydrochloric acid. The filtrate was acidified to pH 2 with 2 M aqueous hydrochloric acid, solid sodium chloride was added to saturate the aqueous layer and the mixture was extracted with ethyl acetate. The ethyl acetate extracts were combined, washed with brine, dried over magnesium sulfate and evaporated. The residue was co-evaporated with toluene (3 x) to afford the title compound (237 mg, 86%) as a white powder.



Compound 38


A solution of **38d** (100 mg, 0.18 mmol) in dichloromethane (180 mL) was stirred at RT under nitrogen. 4-Dimethylaminopyridine (33 mg, 0.27 mmol) and *N*-(3-dimethylaminopropyl)-*N*-ethylcarbodiimide hydrochloride (104 mg, 0.54 mmol) were added and the reaction mixture was stirred at RT for 18 h. The solvent was evaporated. The residue was purified by silica gel chromatography using a gradient of *iso*-hexanes/ethyl acetate 1:2 to 0:1 followed by silica gel chromatography using a gradient of ethyl acetate to ethyl acetate/acetone 4:1. The residue was co-evaporated with dichloromethane then triturated with diethyl ether to afford a solid. The solid was washed with ether and dried to afford the title compound (8 mg, 9%) as a white solid. ¹H NMR (300 MHz, CDCl₃) 0.95 (d, *J* = 6.2 Hz, 3H), 0.96 (d, *J* = 6.5 Hz, 3H), 1.39 (d, *J* = 7.1 Hz, 3H), 1.54-1.80 (m, 5H), 1.82-2.10 (m, 3H), 2.52-2.63 (m, 1H), 3.29-3.62 (m, 2H), 4.07 (*app t*, *J* = 9.6 Hz, 1H), 4.45 (*br d*, *J* = 13.6 Hz, 1H), 4.67 (ABq, $\Delta\delta_{AB}$ = 0.12, *J*_{AB} = 16.0 Hz, 2H), 4.88-5.01 (m, 1H), 5.93-6.00 (m, 2H), 6.57 (d, *J* = 10.3 Hz, 1H), 6.94 (d, *J* = 8.5 Hz, 2H), 7.15-7.53 (m, 6 H). LCMS (*m/z*) 537.2 [*M*+*H*], *Tr* = 2.34 min.

EXAMPLE 39


Compound 39a

A solution of 4-iodo phenol (2.2 g, 10.0 mmol), (R)-2-hydroxy-propionic acid methyl ester (0.95 mL, 10.0 mmol) and triphenylphosphine (2.62 g, 10.0 mmol) was prepared in tetrahydrofuran (40 mL). Diisopropyl azodicarboxylate (2.0 mL, 10 mmol) was added dropwise and the reaction was stirred at -5°C for 1 hour and then at RT for 2 h. The tetrahydrofuran was evaporated and diethyl ether/*iso*-hexanes (1:10, 50 mL) was added. The mixture was stirred at RT for 10 min where a precipitate formed. The filtrate was decanted off and evaporated. The


residue was purified by silica gel chromatography using a gradient of *iso*-hexanes/ethyl acetate 20:1 to 10:1 to afford the title compound (2.02 g, 66%) as an oil.



(R)-1-(3-bromophenyl)-ethanol (1.0 g, 4.97 mmol), bis(pinacolato)diboron (1.39 g, 5.47 mmol), 1,1'-bis(diphenylphosphino)ferrocenedichloropalladium(II), dichloromethane adduct (203 mg, 0.249 mmol) and potassium acetate (976 mg, 9.94 mmol) were dissolved in 1,4-dioxane (10 mL) and the reaction was heated to reflux and left to stir over 3 days. The reaction was allowed to cool to RT before being filtered through a pad of Hyflo. The pad was then washed with ethyl acetate and the combined organics were then concentrated and purified by silica gel chromatography using a stepwise gradient of *iso*-hexanes/ethyl acetate 1:0 to 4:1 to afford the title compound (936 mg, 76%) as a pale yellow oil.




A solution of **39b** (496 mg, 2.00 mmol) and **39a** (612 mg, 2.00 mmol) in 1,2-dimethoxyethane (4 mL) was stirred at RT under nitrogen. A solution of 2 M aqueous sodium carbonate (4 mL) was added followed by tetrakis(triphenylphosphine) palladium(0) (116 mg, 0.1 mmol) and the reaction mixture was heated at 80°C for 1 hour. The reaction mixture was cooled to RT, water was added and the mixture was extracted with ethyl acetate. The ethyl acetate extracts were combined, washed with brine, dried over sodium sulfate and evaporated. The residue was purified by silica gel chromatography using a gradient of *iso*-hexanes/ethyl acetate 17:3 to 1:1 to afford the title compound (253 mg, 42%) as a gum.



A solution of **39c** (250 mg, 0.83 mmol) in tetrahydrofuran (4 mL) was stirred at 5°C under nitrogen. A solution of lithium hydroxide monohydrate (42 mg, 1.00 mmol) in water (1 mL) was

added and the reaction mixture was stirred at 5°C for 1 hour. The reaction mixture was acidified to pH 3 with 2 M aqueous hydrochloric acid and the mixture extracted with ethyl acetate. The organic extracts were separated, washed with water and brine, dried over magnesium sulfate and evaporated to afford the title compound (224 mg, 94%) as a white foam.




A solution of **1e** (478 mg, 0.9 mmol) in dichloromethane (15 mL) was cooled in an ice-water bath under nitrogen. Trimethylsilyl trifluoromethanesulfonate (0.25 mL, 1.35 mmol) was added dropwise, and the resulting solution was stirred for 1 h. Cold saturated aqueous sodium hydrogen carbonate solution (15 mL) was added and the mixture was stirred at 0°C for 15 min.


10 The organic layer was separated, washed with brine, dried over magnesium sulfate and evaporated to afford (S)-1-[(S)-2-((S)-2-amino-3-methyl-butrylamino)-propionyl]-hexahydro-pyridazine-3-carboxylic acid 2,2,2-trichloro-ethyl ester (0.9 mmol) which was used without further purification. A solution of (S)-1-[(S)-2-((S)-2-amino-3-methyl-butrylamino)-propionyl]-hexahydro-pyridazine-3-carboxylic acid 2,2,2-trichloro-ethyl ester (0.9 mmol) in acetonitrile (15

15 mL) was stirred at 0°C under nitrogen. **39d** (224 mg, 0.78 mmol) and 1-hydroxybenzotriazole hydrate (184 mg, 0.96 mmol, wetted with not less than 20 wt. % water) were added, followed by *N*-(3-dimethylaminopropyl)-*N'*-ethylcarbodiimide hydrochloride (225 mg, 1.17 mmol) and the reaction mixture was stirred at RT for 2 h. The solvent was evaporated. The residue was dissolved in ethyl acetate and the solution was washed with water (3 x) followed by brine, dried

20 over magnesium sulfate and evaporated. The residue was purified by silica gel chromatography using a gradient of *iso*-hexanes/ethyl acetate 1:2 to 0:1 to afford the title compound (490 mg, 90%) as a gum.



A solution of **39e** (490 mg, 0.70 mmol) in tetrahydrofuran (15 mL) was stirred at RT under nitrogen. Zinc powder (1.00 g, 15.40 mmol) was added followed by a solution of ammonium acetate (810 mg, 10.50 mmol) in water (8 mL). The reaction mixture was stirred at RT under nitrogen for 24 h. The reaction mixture was filtered through Celite and the filter pad was washed with ethyl acetate and 2 M aqueous hydrochloric acid. The filtrate was acidified to pH 2-3 with 2 M aqueous hydrochloric acid. Solid sodium chloride was added to saturate the aqueous layer and the mixture was extracted with ethyl acetate. The ethyl acetate extracts were combined and washed with brine. The organic extracts were passed through a hydrophobic frit and the filtrate was evaporated. The residue was co-evaporated with ethyl acetate (3 x) then toluene (3 x) to afford the title compound (394 mg, 98%) as a white solid.




A solution of **39f** (100 mg, 0.18 mmol) in dichloromethane (180 mL) was stirred at RT under nitrogen. 4-Dimethylaminopyridine (44 mg, 0.36 mmol) and *N*-(3-dimethylaminopropyl)-*N'*-ethylcarbodiimide hydrochloride (138 mg, 0.72 mmol) were added and the reaction mixture was stirred at RT for 4 h. The solvent was evaporated. Ethyl acetate was added to the residue and the mixture was washed with aqueous citric acid (pH 4) and brine. The organic layer was dried over magnesium sulfate and the solvent was evaporated. The residue was purified by silica gel chromatography using a gradient of *iso*-hexanes/ethyl acetate 3:7 to 0:1. The residue was triturated with diethyl ether/*iso*-hexanes 1:1 to afford the title compound (10 mg, 10%) as a white solid. ¹H NMR (300 MHz, CD₃OD) 0.94 (d, *J* = 6.7 Hz, 3H), 0.95 (d, *J* = 6.9 Hz, 3H), 1.51 (d, *J* = 7.1 Hz, 3H), 1.61 (d, *J* = 6.5 Hz, 3H), 1.67 (d, *J* = 6.9 Hz, 3H), 1.72-2.00 (m, 5H), 2.60-2.69 (m, 1H), 3.54-3.63 (m, 1H), 4.13 (d, *J* = 10.5 Hz, 1H), 4.32 (*br* d, *J* = 12.9 Hz, 1H), 4.57-4.66 (m, 2H), 4.78 (q, *J* = 7.1 Hz, 1H), 5.92 (q, *J* = 6.5 Hz, 1H), 6.91 (d, *J* = 8.9 Hz, 2H), 7.22 (d, *J* = 7.6 Hz, 1H), 7.34-7.61 (m, 5H). LCMS (*m/z*) 551.2 [*M*+*H*], *Tr* = 2.51 min.


EXAMPLE 40




40a was prepared in the same manner as (S)-2-(4-iodophenoxy)-propionic acid methyl ester using (S)-2-hydroxy-propionic acid ethyl ester instead of (R)-2-hydroxy-propionic acid methyl ester in 28% yield.




40b was prepared in the same manner as (S)-2-[3'-((R)-1-hydroxy-ethyl)-biphenyl-4-yloxy]-propionic acid methyl ester using **40a** instead of (S)-2-(4-iodophenoxy)-propionic acid methyl ester in 54% yield.




40c was prepared in the same manner as (S)-2-[3'-((R)-1-hydroxy-ethyl)-biphenyl-4-yloxy]-propionic acid using **40b** instead of (S)-2-[3'-((R)-1-hydroxy-ethyl)-biphenyl-4-yloxy]-propionic acid methyl ester in 71% yield.



40d was prepared in the same manner as (S)-1-[(S)-2-[(S)-2-[(S)-2-[3'-((R)-1-hydroxy-ethyl)-biphenyl-4-yloxy]-propionylamino]-3-methyl-butyrylamino)-propionyl]-hexahydro-pyridazine-3-carboxylic acid 2,2,2-trichloro-ethyl ester using **40c** instead of (S)-2-[3'-((R)-1-hydroxy-ethyl)-biphenyl-4-yloxy]-propionic acid in 71% yield.




40e was prepared in the same manner as ((S)-1-[(S)-2-((S)-2-[(S)-2-[3'-((R)-1-hydroxy-ethyl)-biphenyl-4-yloxy]-propionylamino)-3-methyl-butyrylamino)-propionyl]-hexahydro-pyridazine-3-carboxylic acid using **40d** instead of (S)-1-[(S)-2-((S)-2-[(S)-2-[3'-((R)-1-hydroxy-ethyl)-biphenyl-4-yloxy]-propionylamino)-3-methyl-butyrylamino)-propionyl]-hexahydro-pyridazine-3-carboxylic acid 2,2,2-trichloro-ethyl ester in 83% yield.




A solution of **40e** (102 mg, 0.18 mmol) in dichloromethane (180 mL) was stirred at RT under nitrogen. 4-Dimethylaminopyridine (44 mg, 0.36 mmol) and (7-azabenzotriazol-1-yloxy)tripyrrolidinophosphonium hexafluorophosphate (114 mg, 0.22 mmol) were added and the reaction mixture was stirred at RT for 4 h. The solvent was evaporated. Ethyl acetate was added to the residue and the mixture was washed with aqueous citric acid (pH 4) and brine. The organic layer was separated and dried over magnesium sulfate and evaporated. The residue was purified by silica gel chromatography using a gradient of *iso*-hexanes/ethyl acetate 3:7 to 0:1. The residue was purified by preparative reverse phase HPLC to afford the title compound (2.8 mg, 3%) as a white solid. ¹H NMR (300 MHz, CD₃OD) 1.03 (d, *J* = 6.9 Hz, 3H), 1.05 (d, *J* = 6.9 Hz, 3H), 1.25 (d, *J* = 7.1 Hz, 3H), 1.56 (d, *J* = 6.5 Hz, 3H), 1.57 (d, *J* = 6.3 Hz, 3H), 1.64-2.10 (m, 5H), 2.75-2.83 (m, 1H), 3.60-3.66 (m, 1H), 4.08 (d, *J* = 10.0 Hz, 1H), 4.32 (*br d*, *J* = 12.9 Hz, 1H), 4.74 (q, *J* = 6.5 Hz, 1H), 5.22 (q, *J* = 7.3 Hz, 1H), 5.98 (q, *J* = 6.5 Hz, 1H), 6.86 (d, *J* = 8.5 Hz, 2H), 7.20-7.25 (m, 1H), 7.39-7.51 (m, 5H). LCMS (*m/z*) 551.2 [M+H], Tr = 2.27 min.


EXAMPLE 41




Dichloro (*p*-cymene) ruthenium(II) dimer (31 mg, 0.05 mmol) and (1R,2R)-(-)-*N*-p-tosyl-1,2-diphenylethylenediamine (44 mg, 0.12 mmol) was suspended in degassed water (20 mL) and the mixture was degassed with nitrogen for 10 min. The mixture was stirred at 70°C under nitrogen for 90 min. The resulting yellow solution was cooled to RT. A solution of 1-(6-bromopyridin-2-yl)-ethanone (2.0 mg, 10 mmol) in degassed tetrahydrofuran (10 mL) and sodium formate (3.4 g, 50 mmol) was added and the reaction mixture was degassed with nitrogen for 5 min. The reaction mixture was vigorously stirred at 40°C for 30 min. The reaction mixture was cooled to RT and was extracted with ethyl acetate. The organic extracts were combined, washed with water and brine, dried over sodium sulfate, filtered and evaporated. The residue was purified by silica gel chromatography using a gradient of *iso*-hexanes/ethyl acetate 3:1 to 0:1 to afford the title compound (1.78 g, 89%) as a brown oil.



To a mixture of **41a** (101 mg, 0.50 mmol), 4-(2-ethoxy-2-oxoethoxy)benzeneboronic acid (112 mg, 0.50 mmol) in 1,2-dimethoxyethane (4 mL) was added a solution of potassium carbonate (138 mg, 1.00 mmol) in water (1 mL). Tetrakis(triphenylphosphine)palladium(0) (29 mg, 0.025 mmol) was added and the reaction mixture was heated at 100°C in a microwave reactor for 30 min. The reaction mixture was then diluted with ethyl acetate and water. 2 M Hydrochloric acid was added to adjust the pH of the reaction mixture to pH 3 and the volatiles was evaporated. Methanol was added to the residue and the mixture was filtered through a hydrophobic frit. The filtrate was evaporated and the residue was dried in vacuum to afford the title compound (143 mg, 93%) as a white solid.



- A solution of **1e** (240 mg, 0.45 mmol) in dichloromethane (15 mL) was cooled in an ice-water bath under nitrogen. Trimethylsilyl trifluoromethanesulfonate (0.12 mL, 0.68 mmol) was added dropwise, and the resulting solution was stirred for 1 h. Cold saturated aqueous sodium hydrogen carbonate solution (15 mL) was added and the mixture was stirred at 0°C for 15 min.
- 5 The organic layer was separated, washed with brine, dried over magnesium sulfate and evaporated to afford (S)-1-[(S)-2-((S)-2-amino-3-methyl-butyrylamino)-propionyl]-hexahydro-pyridazine-3-carboxylic acid 2,2,2-trichloro-ethyl ester (0.45 mmol) which was used without further purification. A mixture of (S)-1-[(S)-2-((S)-2-amino-3-methyl-butyrylamino)-propionyl]-hexahydro-pyridazine-3-carboxylic acid 2,2,2-trichloro-ethyl ester (0.45 mmol), **41b** (143 mg, 0.45 mmol) and *N,N*-diisopropylethylamine (0.16 mL, 0.9 mmol) in acetonitrile (15 mL) was
- 10 stirred at RT under nitrogen. 1-Hydroxybenzotriazole hydrate (108 mg, 0.56 mmol, wetted with not less than 20 wt. % water) and *N*-(3-dimethylaminopropyl)-*N'*-ethylcarbodiimide hydrochloride (130 mg, 0.675 mmol) was added and the reaction mixture was stirred at RT for 4 h. *N,N*-Dimethylformamide (2 mL) was added and the reaction mixture was stirred at RT for 22 h.
- 15 Additional *N*-(3-dimethylaminopropyl)-*N'*-ethylcarbodiimide hydrochloride (87 mg, 0.45 mmol) was added and the reaction mixture was stirred at RT for 6 h. The solvent was evaporated. The residue was suspended in a mixture of ethyl acetate and aqueous citric acid solution (pH 3) and the mixture was extracted with ethyl acetate. The organic extracts were combined, washed with water followed by brine. The organic layer was filtered through a hydrophobic frit and the filtrate
- 20 was evaporated. The residue was purified by silica gel chromatography using ethyl acetate to afford the title compound (71 mg, 23%) as a white solid.




Compound 41

- A solution of **41c** (69 mg, 0.10 mmol) in tetrahydrofuran (2 mL) was stirred at 0°C under nitrogen, a solution of lithium hydroxide monohydrate (5 mg, 0.12 mmol) in water (0.5 mL) was
- 25 added and the reaction mixture was stirred at 0°C for 1 h. The solution was acidified to pH 3 with 2 M hydrochloric acid and the solvent was evaporated. The residue was co-evaporated with toluene (3 x) to afford (S)-1-[(S)-2-[(S)-2-(2-{4-[6-((R)-1-hydroxy-ethyl)-pyridin-2-yl]-phenoxy}-acetylamino)-3-methyl-butyrylamino]-propionyl]-hexahydro-pyridazine-3-carboxylic acid (58 mg,


0.1 mmol) as a white solid which was used crude in the next step. A solution of (S)-1-[(S)-2-[(S)-2-(2-{4-[6-((R)-1-hydroxy-ethyl)-pyridin-2-yl]-phenoxy}-acetyl-amino)-3-methyl-butyl-ylamino]-propionyl]-hexahydro-pyridazine-3-carboxylic acid (56 mg, 0.1 mmol) in dichloromethane (100 mL) was stirred at RT under nitrogen. 4-Dimethylaminopyridine (49 mg, 0.4 mmol) was added and the reaction mixture was stirred for 5 min. A solution of (7-azabenzotriazol-1-yl)oxy)tripyrrolidinophosphonium hexafluorophosphate (78 mg, 0.15 mmol) in dichloromethane (20 mL) was added dropwise over 5 min and the reaction mixture was stirred at RT for 2 h. The solvent was evaporated and the residue was purified by silica gel chromatography using ethyl acetate. The residue was purified by preparative reverse phase HPLC to afford the title compound (14 mg, 26%) as a white solid. ¹H NMR (300 MHz, CD₃OD) 0.94 (d, *J* = 6.7 Hz, 3H), 0.98 (d, *J* = 6.5 Hz, 3H), 1.36 (d, *J* = 7.1 Hz, 3H), 1.68 (d, *J* = 6.5 Hz, 3H), 1.72-2.05 (m, 5H), 2.65-2.74 (m, 1H), 3.53-3.60 (m, 1H), 4.14 (d, *J* = 10.7 Hz, 1H), 4.30-4.38 (m, 1H), 4.66 (ABq, $\Delta\delta_{AB}$ = 0.13, *J*_{AB} = 15.8 Hz, 2H), 5.07 (q, *J* = 7.1 Hz, 1H), 5.87 (q, *J* = 6.5 Hz, 1H), 6.89 (d, *J* = 8.9 Hz, 2H), 7.26 (*br d*, *J* = 7.1 Hz, 1H), 7.71-7.78 (m, 2H), 8.06 (d, *J* = 8.9 Hz, 2H). LCMS (*m/z*) 538.2 [M+H], Tr = 2.07 min.

EXAMPLE 42




Compound 42a

42a was prepared in the same manner as (S)-2-[3'-((R)-1-hydroxy-ethyl)-biphenyl-4-yloxy]-propionic acid methyl ester using 3-(4-bromo-pyrazol-1-yl)-propionic acid methyl ester instead of (S)-2-(4-iodo-phenoxy)-propionic acid methyl ester in 28% yield.




Compound 42b

42b was prepared in the same manner as (S)-2-[3'-((R)-1-hydroxy-ethyl)-biphenyl-4-yloxy]-propionic acid using **42a** instead of (S)-2-[3'-((R)-1-hydroxy-ethyl)-biphenyl-4-yloxy]-propionic acid methyl ester in 96% yield.




42c was prepared in the same manner as compound (S)-1-[(S)-2-((S)-2-[3'-((R)-1-hydroxy-ethyl)-biphenyl-4-yloxy]-propionylamino)-3-methyl-butrylamino)-propionyl]-hexahydro-pyridazine-3-carboxylic acid 2,2,2-trichloro-ethyl ester using **42b** instead of (S)-2-[3'-((R)-1-hydroxy-ethyl)-biphenyl-4-yloxy]-propionic acid in 49% yield.




Compound 42 was prepared in the same manner as Compound 41 using **42c** instead of (S)-1-[(S)-2-[(S)-2-(2-[4-[6-((R)-1-hydroxy-ethyl)-pyridin-2-yl]-phenoxy]-acetylamino)-3-methyl-butrylamino]-propionyl]-hexahydro-pyridazine-3-carboxylic acid 2,2,2-trichloro-ethyl ester in 8% yield. ¹H NMR (300 MHz, CD₃OD) 0.94 (d, *J* = 6.7 Hz, 3H), 0.95 (d, *J* = 6.5 Hz, 3H), 1.37 (d, *J* = 7.1 Hz, 3H), 1.62 (d, *J* = 6.5 Hz, 3H), 1.65-2.05 (m, 5H), 2.61-2.69 (m, 1H), 2.88-3.10 (m, 2H), 3.58-3.63 (m, 1H), 3.90-4.00 (m, 1H), 4.01 (d, *J* = 8.5 Hz, 1H), 4.38-4.44 (m, 1H), 4.58-4.67 (m, 1H), 5.31 (q, *J* = 7.1 Hz, 1H), 5.91 (q, *J* = 6.5 Hz, 1H), 7.18 (d, *J* = 7.6 Hz, 1H), 7.31 (*app t*, *J* = 7.6 Hz, 1H), 7.47 (d, *J* = 7.6 Hz, 1H), 7.58 (s, 1H), 7.77 (s, 1H), 7.95 (s, 1H). LCMS (*m/z*) 525.2 [M+H], Tr = 4.51 min.


EXAMPLE 43



Potassium carbonate (663 mg, 4.80 mmol) and tetrakis (triphenylphosphine)palladium (0) (139 mg, 0.12 mmol) were added to a solution of (R)-1-(3-bromophenyl)-ethanol (483 mg, 2.40 mmol) and 4-(2-methoxy carbonyl) benzeneboronic acid (500 mg, 2.40 mmol) in 1,2-dimethoxyethane (5 mL) in a 5 mL microwave vessel. The vessel was sealed before being
 5 heated in the microwave for 20 min, using fixed hold time, on high absorption at 100°C. The reaction mixture was filtered through a pad of Hyflo and the pad was washed with ethyl acetate. The combined organics were then concentrated and the resultant brown oil was purified by silica gel chromatography using a stepwise gradient of *iso*-hexanes/ethyl acetate 1:0 to 3:2 to afford the title compound (416 mg, 61%) as a yellow oil.




43a (416 mg, 1.46 mmol) was dissolved in a mixture of tetrahydrofuran (8 mL) and water (2 mL) and the solution was cooled using an ice bath. Lithium hydroxide monohydrate (175 mg, 2.92 mmol) was added and the solution was allowed to slowly warm to RT overnight. The solution was acidified using 2 M hydrochloric acid and then extracted with dichloromethane (2 x
 15 20 mL). The combined organics were then dried over anhydrous sodium sulfate, filtered and concentrated to afford the title compound (250 mg, 89%) as a white solid.




(S)-1-[(S)-2-((S)-2-Amino-3-methyl-butylamino)-propionyl]-hexahydro-pyridazine-3-carboxylic acid 2,2,2-trichloro-ethyl ester (669 mg, 1.55 mmol) and **43b** (350 mg, 1.29 mmol)
 20 were dissolved in anhydrous acetonitrile (10 mL) and cooled using an ice bath. 2-(1H-7-Azabenzotriazol-1-yl)-1,1,3,3-tetramethyl uronium hexafluorophosphate methanaminium (688 mg, 1.81 mmol) and *N,N*-diisopropylethylamine (899 μ L, 5.16 mmol) were then added and the reaction was allowed to slowly warm to RT and left to stir overnight. The solvent was then removed and the residue was dissolved in ethyl acetate. The solution was then washed with
 25 water (3 x 20 mL) and brine (20 mL), dried over anhydrous sodium sulfate, filtered and

concentrated and the resultant brown oil was purified by silica gel chromatography using a stepwise gradient of *iso*-hexanes/ethyl acetate 1:4 to 0:1 to afford the title compound (314 mg, 36%) as a white solid.



Compound 43d

- 5 **43c** (314 mg, 0.459 mmol) was dissolved in tetrahydrofuran (10 mL). Zinc powder (300 mg, 4.59 mmol) was added followed by ammonium acetate (1 M in water, 3.21 mL, 3.21 mmol). The reaction was then left to stir overnight. The reaction mixture was then filtered through a pad of Hyflo. The pad was then washed with potassium hydrogen sulfate solution and ethyl acetate. The biphasic mixture was further acidified using 2 M hydrochloric acid until the solution was
- 10 approximately pH 1. The phases were separated and the aqueous phase was extracted with ethyl acetate. The combined organic phases were dried over anhydrous sodium sulfate, filtered and concentrated to afford the title compound (212 mg, 84%) as a yellow solid.




Compound 43


- 15 **43d** (100 mg, 0.181 mmol) was dissolved in dichloromethane (181 mL), under an atmosphere of nitrogen. *N*-(3-Dimethylaminopropyl)-*N'*-ethylcarbodiimide hydrochloride (142 0.724 mmol) and 4-dimethylaminopyridine (44 mg, 0.362 mmol) were added and the reaction was left to stir overnight. The solvent was removed and the residue was purified by silica gel chromatography using 100% ethyl acetate to afford the title compound (52 mg, 54%) as a white solid. ¹H NMR (300 MHz, CDCl₃) δ 0.92 (d, *J* = 6.7 Hz, 6H), 1.24-1.29 (m, 3H), 1.60-1.68 (m,
- 20 5H), 1.85-1.98 (m, 2H), 2.37-2.49 (m, 1H), 2.58-2.78 (m, 1H), 2.83-2.93 (m, 1H), 3.16-3.28 (m, 1H), 3.45 (m, 3H), 4.03-4.11 (m, 1H), 4.45-4.54 (m, 1H), 5.06-5.16 (m, 1H), 5.92-6.00 (m, 1H),

6.25 (d, $J = 8.5$ Hz, 1H), 7.19 (s, 1H), 7.22 (s, 2H), 7.39 (*app t*, $J = 7.6$ Hz, 1H), 7.48-7.56 (m, 3H), 7.62 (*br s*, 1H). LCMS (m/z) 535.3 [$M+H$], Tr = 2.43 min.


EXAMPLE 44



Benzyl bromoacetate (2.07 mL, 13.1 mmol) and triphenylphosphine (3.63 g, 13.8 mmol) were dissolved in toluene (60 mL) and left to stir over 3 days. A white solid was collected by filtration and washed with diethyl ether. This afforded the title compound (5.86 g, 93%) as a white solid.




5-bromo-2-pyridine carboxaldehyde (125 mg, 0.666 mmol) and **44a** (655 mg, 1.33 mmol) were dissolved in dichloromethane (5 mL). 1,1,3,3-Tetramethylguanidine (251 μ L, 2 mmol) was then added and the reaction was left to stir for 3.5 h. The reaction was quenched with saturated ammonium chloride solution and the phases were separated. The organic phase was then washed with brine, dried over anhydrous sodium sulfate, filtered and concentrated. The residue was then purified by silica gel chromatography using a stepwise gradient of *iso*-hexanes/ethyl acetate 1:0 to 4:1 to afford the title compound (186 mg, 88%) as a yellow solid.




44b (186 mg, 0.585 mmol), (R)-1-[3-(4,4,5,5-tetramethyl-[1,3,2] dioxaborolan-2-yl)-phenyl]-ethanol (145 mg, 0.585 mmol), cesium carbonate (476 mg, 1.46 mmol) and tetrakis(triphenylphosphine) palladium (0) (66 mg, 0.0585 mmol) were placed in a microwave vessel. 1,2-Dimethoxyethane (2 mL) and water (0.5 mL) were added and the vessel was sealed.


The reaction was heated in the microwave for 30 min, using fixed hold time, on high absorption at 150°C. The reaction mixture was poured onto water and the resultant mixture was extracted with ethyl acetate. The organics were then dried over anhydrous sodium sulfate, filtered and concentrated. The residue was then purified by silica gel chromatography using a stepwise gradient of *iso*-hexanes/ethyl acetate 1:0 to 1:1 to afford the title compound (150 mg, 77%) as an orange gum.



44c (150 mg, 0.417 mmol) was dissolved in ethanol (5 mL). Palladium(II) hydroxide (20% on carbon, wet, 40 mg) was then added followed by ammonium formate (132 mg, 2.09 mmol). The reaction was heated to reflux and left to stir for 30 min. The reaction was allowed to cool to RT and was then filtered through a pad of Hyflo. The pad was washed with ethanol and the combined organics were concentrated to leave the title compound (105 mg, 93%) as an orange gum.




44d (105 mg, 0.387 mmol) and (S)-1-[(S)-2-((S)-2-amino-3-methyl-butylamino)-propionyl]-hexahydro-pyridazine-3-carboxylic acid 2,2,2-trichloro-ethyl ester (200 mg, 0.464 mmol) were dissolved in acetonitrile (5 mL) and cooled using an ice bath. 2-(1H-7-Azabenzotriazol-1-yl)-1,1,3,3-tetramethyl uronium hexafluorophosphate methanaminium (206 mg, 0.542 mmol) and *N,N*-diisopropylethylamine (270 μ L, 1.55 mmol) were then added and the reaction was allowed to slowly warm to RT and left to stir overnight. The solvent was then removed and the residue was dissolved in ethyl acetate. The solution was then washed with water (3 x 20 mL) and brine (20 mL), dried over anhydrous magnesium sulfate, filtered and concentrated and the residue was purified by silica gel chromatography using a stepwise gradient of *iso*-hexanes/ethyl acetate/acetone 3:1:0 to 0:7:3 to afford a brown solid (110 mg). This was further purified by silica gel chromatography using a stepwise gradient of ethyl acetate/acetone 1:0 to 4:1 to afford the title compound (100 mg, 38%) as a yellow solid.



Compound 44f


44e (100 mg, 0.146 mmol) was dissolved in a mixture of tetrahydrofuran (3 mL) and water (1 mL) and cooled using an ice bath. Lithium hydroxide monohydrate (6 mg, 0.153 mmol) was added and the reaction was stirred for 30 min. The solution was acidified to pH 1 using 2 M hydrochloric acid and evaporated to dryness. The residue was then purified by C18 chromatography using a stepwise gradient of acetonitrile/water 0:1 to 1:4 to afford the title compound (25 mg, 31%) as a white solid.




Compound 44

44f (25 mg, 0.045 mmol) was dissolved in dichloromethane (45 mL). *N*-(3-dimethylaminopropyl)-*N'*-ethylcarbodiimide hydrochloride (35 mg, 0.148 mmol) and 4-dimethylaminopyridine (11 mg, 0.009 mmol) were added and the reaction was left to stir for 4 h. The solvent was removed and the residue was purified by silica gel chromatography using a stepwise gradient of ethyl acetate/acetone 1:0 to 3:2 to leave a white solid (6 mg). This was then eluted through an reverse phase HPLC system fitted with a Phenomenex Gemini 10 μ 110A, 250 x 21.2 mm column using an isocratic 2:3 acetonitrile/water flow at 20 mL/min to afford the title compound (1.3 mg, 7%) as a white solid. ¹H NMR (300 MHz, CDCl₃) δ 0.84-1.00 (m, 7H), 1.10 (d, *J* = 6.9 Hz, 3H), 1.62-1.72 (m, 4H), 1.86-2.08 (m, 3H), 2.54-2.71 (m, 2H), 2.90-3.14 (m, 2H), 3.28-3.54 (m, 3H), 3.98 (*app* t, *J* = 9.2 Hz, 1H), 4.43-4.52 (m, 1H), 5.07 (*app* t, *J* = 8.0 Hz, 1H), 5.92-6.00 (m, 1H), 6.10 (d, *J* = 8.0 Hz, 1H), 6.37 (d, *J* = 9.1 Hz, 1H), 7.20 (d, *J* = 8.0 Hz, 1H), 7.26-7.31 (m, 1H), 7.41-7.59 (m, 3H), 7.75 (dd, *J* = 8.0, 2.2 Hz, 1H), 8.78 (d, *J* = 2.4 Hz, 1H). LCMS (*m/z*) 536.0 [M+H]⁺, Tr = 1.47 min.


EXAMPLE 45



6-Bromo-3-pyridinecarboxaldehyde (500 mg, 2.69 mmol) and benzyloxycarbonylmethyltriphenylphosphonium bromide (2.64 g, 5.38 mmol) were dissolved in dichloromethane (15 mL). 1,1,3,3-Tetramethylguanidine (1.01 mL, 8.03 mmol) was then added and the reaction was left to stir for 3.5 h. The reaction was quenched with saturated ammonium chloride solution and the phases were separated. The organic phase was then washed with brine, dried over anhydrous sodium sulfate, filtered and concentrated. The residue was then purified by silica gel chromatography using a stepwise gradient of *iso*-hexanes/ethyl acetate 1:0 to 4:1 to afford the title compound (492 mg, 57%) as a white solid.




45a (382 mg, 1.54 mmol), (R)-1-[3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-phenyl]-ethanol (490 mg, 1.54 mmol), cesium carbonate (1.06 g, 3.85 mmol) and tetrakis(triphenylphosphine) palladium(0) (178 mg, 0.154 mmol) were placed in a microwave vessel. 1,2-Dimethoxyethane (4 mL) and water (1 mL) were added and the vessel was sealed. The reaction was heated in the microwave for 50 min, using fixed hold time, on high absorption at 150°C. The reaction mixture was poured onto water and the resultant mixture was extracted with ethyl acetate. The organics were then dried over anhydrous sodium sulfate, filtered and concentrated. The residue was then purified by silica gel chromatography using a stepwise gradient of *iso*-hexanes/ethyl acetate 1:0 to 1:1 to afford impure product. This was subjected to a second round of purification using the same conditions to afford the title compound (195 mg, 35%) as a yellow solid.



45b (195 mg, 0.543 mmol) was dissolved in ethanol (5 mL). Palladium(II) hydroxide (20% on carbon, wet, 40 mg) was then added followed by ammonium formate (172 mg, 2.72


mmol). The reaction was heated to reflux and left to stir for 30 min. The reaction was allowed to cool to RT and was then filtered through a pad of Hyflo. The pad was washed with ethanol and the combined organics were concentrated to leave the title compound (150 mg, 100%) as a pale yellow solid.



45c (150 mg, 0.553 mmol) and (S)-1-[(S)-2-((S)-2-amino-3-methylbutyrylamino)propionyl]-hexahydro-pyridazine-3-carboxylic acid 2,2,2-trichloro-ethyl ester (287 mg, 0.664 mmol) were dissolved in acetonitrile (5 mL) and cooled using an ice bath. 2-(1H-7-Azabenzotriazol-1-yl)-1,1,3,3-tetramethyl uronium hexafluorophosphate methanaminium (294 mg, 0.774 mmol) and *N,N*-diisopropylethylamine (385 μ L, 2.21 mmol) were then added and the reaction was allowed to slowly warm to RT and left to stir overnight. Ethyl acetate was then poured into the reaction mixture. The resultant solution was then washed with water (3 x 20 mL) and brine (20 mL), dried over anhydrous sodium sulfate, filtered and concentrated and the residue was purified by silica gel chromatography using a stepwise gradient of *iso*-hexanes/ethyl acetate/acetone 3:1:0 to 0:7:3 to afford a solid (334 mg). This was further purified by silica gel chromatography using a stepwise gradient of ethyl acetate/acetone 1:0 to 4:1 to afford the title compound (167 mg, 44%) as a solid.


10

15



45d (160 mg, 0.234 mmol) was dissolved in a mixture of tetrahydrofuran (4 mL) and water (1 mL) and cooled using an ice bath. Lithium hydroxide monohydrate (6 mg, 0.153 mmol) was added and the reaction was stirred for 30 min. The solution was neutralized using 2 M hydrochloric acid and evaporated to dryness. The residue was then purified by C18 chromatography using a stepwise gradient of acetonitrile/water 0:1 to 1:4 to afford the title compound (21 mg, 16%) as a white solid.

20




45e (21 mg, 0.0379 mmol) was dissolved in dichloromethane (38 mL). *N*-(3-dimethylaminopropyl)-*N'*-ethylcarbodiimide hydrochloride (29 mg, 0.152 mmol) and 4-dimethylaminopyridine (9 mg, 0.0758 mmol) were added and the reaction was left to stir for 4 h.

5 The solvent was removed and the residue was purified by silica gel chromatography using 100% ethyl acetate. The resultant material was then eluted through a reverse phase HPLC system fitted with a Phenomenex Gemini 10 μ 110A, 250 x 21.2 mm column using an isocratic 3:7 acetonitrile/water flow at 20 mL/min to afford the title compound (1 mg, 5%). ¹H NMR (300 MHz, CDCl₃) δ 0.91 (d, *J* = 6.5 Hz, 6H), 1.48 (d, *J* = 7.1 Hz, 3H), 1.63-1.72 (m, 7H), 1.81-1.93 (m, 2H), 2.00-2.10 (m, 1H), 2.25-2.38 (m, 1H), 2.54-2.67 (m, 1H), 3.15-3.28 (m, 1H), 3.39-3.59 (m, 2H), 4.03 (*app t*, *J* = 9.6 Hz, 1H), 4.43-4.54 (m, 1H), 5.04-5.16 (m, 1H), 5.80 (d, *J* = 9.8 Hz, 1H), 5.94 (q, *J* = 6.7 Hz, 1H), 7.22-7.27 (m, 1H), 7.43 (*app t*, *J* = 7.8 Hz, 1H), 7.55-7.57 (m, 1H), 7.69 (d, *J* = 8.0 Hz, 1H), 7.92 (d, *J* = 8.3 Hz, 1H), 8.19 (*br s*, 1H), 8.48 (d, *J* = 2.2 Hz, 1H). LCMS (m/z) 536.2 [M+H], Tr = 1.61 min.

15

EXAMPLE 46




To Compound 21 (25 mg, 0.05 mmol) in ethyl acetate (5 mL) at RT was added 10% palladium on carbon (20 mg). The system was purged with hydrogen and stirred for 2 h. The reaction was filtered through Celite and concentrated *in vacuo*. The residue was purified by preparative TLC using ethyl acetate/acetone 5/1 to give the title compound (2.2 mg, 9%) as a white solid. ¹H NMR (300 MHz, CDCl₃) 0.96 (d, *J* = 6.7 Hz, 3H), 0.98 (d, *J* = 6.7 Hz, 3H), 1.24 (*J* = 7.1 Hz, 3H), 1.52 (d, *J* = 6.9 Hz, 3H), 1.67 (d, *J* = 6.7 Hz, 3H), 1.55-2.14 (m, 10H), 2.53-2.80


20

(m, 2H), 2.92–3.01 (m, 2H), 3.45–3.67 (m, 1H), 4.14–4.23 (m, 1H), 4.49–4.60 (m, 1H), 5.39–5.51 (m, 1H), 5.96 (d, $J = 9.4$ Hz, 1H), 6.06 (q, $J = 6.7$ Hz, 1H), 6.43 (d, $J = 8.5$ Hz, 1H), 7.39 (dd, $J = 8.5, 1.1$ Hz, 1H), 7.56 (s, 1H), 7.75 (s, 1H), 7.93 (d, $J = 8.5$ Hz, 1H), 9.16 (s, 1H). LCMS (m/z) = 552.3 [M+H], Tr = 1.25 min.

5

EXAMPLE 47

To a solution of (S)-1-((S)-2-*tert*-butoxycarbonylamino-propionyl)-hexahydro-pyridazine-3-carboxylic acid 2,2,2-trichloro-ethyl ester (320 mg, 0.74 mmol) in dichloromethane (8 mL) was added trifluoroacetic acid (2 mL). The reaction was stirred at RT for 105 min and was then concentrated *in vacuo*. The resulting crude product was dissolved in ethyl acetate (75 mL) and was washed with saturated aqueous sodium bicarbonate. The aqueous phase was extracted with ethyl acetate (2 x 40 mL), and the organics were dried over anhydrous sodium sulfate, filtered, and concentrated to a crude residue that was used without further purification. The residue from the previous step, 1-hydroxybenzotriazole (152 mg, 1.12 mmol) and (S)-2-*tert*-butoxycarbonylamino-3-hydroxy-propionic acid (154 mg, 0.75 mmol) were dissolved in dichloromethane (6 mL). *N,N*-Diisopropylethylamine (190 mg, 1.5 mmol) was added, and the resulting solution was cooled in an ice water bath. *N*-(3-Dimethylaminopropyl)-*N'*-ethylcarbodiimide (149 mg, 0.96 mmol) was added dropwise over 15 s. The reaction was stirred for 18 h, allowing the ice bath to slowly expire. The reaction mixture was then diluted with ethyl acetate (50 mL) and was washed with saturated aqueous sodium bicarbonate and brine. The organic phase was dried over anhydrous sodium sulfate, filtered, and concentrated to afford a crude residue that was purified by silica gel chromatography (75 to 100% ethyl acetate in *iso*-hexanes) to afford the title product (211 mg, 55%) as an oil.



To a solution of **47a** (64 mg, 0.12 mmol) in dichloromethane (1 mL) was added 2,4,6-collidine (98 mg, 1.28 mmol). *tert*-Butyldimethylsilyl trifluoromethanesulfonate (98 mg, 0.37 mmol) was added dropwise over 20 sec. The reaction mixture was stirred for 15 h and was


5 quenched with saturated aqueous sodium bicarbonate (1 mL). The mixture was further diluted with ethyl acetate, water, and 0.1 N aqueous hydrochloric acid to afford an acidic aqueous layer. The phases were separated, and the organics were dried over sodium sulfate, filtered, and concentrated *in vacuo* to afford a crude residue that was used without further purification. The crude product was dissolved in tetrahydrofuran (3 mL). Acetic acid (115 mg, 1.9 mmol) was

10 added in one portion and the resulting solution was stirred for 3.25 h. The reaction was then diluted with ethyl acetate (25 mL) and washed with saturated aqueous sodium bicarbonate (25 mL). The phases were separated, and the aqueous phase was extracted with ethyl acetate (25 mL). The combined organic phases were washed with brine (25 mL) and dried over anhydrous sodium sulfate. The mixture was filtered, and the filtrate was concentrated *in vacuo* to afford

15 crude amine which was used without further purification. The crude amine was dissolved in *N,N*-dimethylformamide (1.5 mL). 2,4,6-Collidine (31 mg, 0.26 mmol) and *trans*-3-pentenoic acid (15.3 mg, 0.152 mmol) were added, and the resulting solution was cooled in an ice water bath. (1-Cyano-2-ethoxy-2-oxoethylidenaminoxy)dimethylamino-morpholino-carbenium

20 hexafluorophosphate (72.4 mg, 0.169 mmol) was added in one portion, and the reaction was stirred for 30 min. The reaction was then removed from the cold bath and warmed to ambient temperature. After 15 h, the reaction was diluted with ethyl acetate (35 mL), saturated aqueous sodium bicarbonate (20 mL), and brine (5 mL). The phases were separated, and the organic layer was washed with 0.1 N aqueous hydrochloric acid (25 mL) and then brine (5 mL). The acidic aqueous layer was extracted with ethyl acetate (25 mL), and the combined organic

25 phases were then dried over anhydrous sodium sulfate, filtered, and concentrated. The resulting crude residue was purified by silica gel chromatography (50 to 80% ethyl acetate in *iso*-hexanes) to afford the title compound (46.9 mg, 62% over 3 steps) as a white foam.




To a solution of **47b** (46.9 mg, 0.0761 mmol) in tetrahydrofuran (1.5 mL) was added water (0.30 mL), ammonium acetate (96 mg, 1.2 mmol), and zinc powder (109 mg, 1.7 mmol). The reaction mixture was stirred vigorously at RT for 17.5 h, at which time the temperature was

5 increased to 35°C. After 25.5 h, additional zinc powder (60 mg, 0.92 mmol) was added and the reaction temperature was increased to 45°C. After 39.5 h, the reaction mixture was filtered through a pad of Celite washing with water and ethyl acetate. The aqueous phase was acidified with 2 M aqueous hydrochloric acid (15 mL), and the phases were separated. The aqueous

10 phase was extracted with ethyl acetate (2 x 20 mL), and the combined organic phases were dried over sodium sulfate, filtered, and concentrated to afford a white solid (34.9 mg, 95%) that was used without further purification. The crude acid (0.072 mmol) was dissolved with (R)-1-(3-vinyl-isoquinolin-6-yl)-ethanol (17.4 mg, 0.087 mmol) in dichloromethane (1.0 mL). 4-Dimethylaminopyridine (13.3 mg, 0.109 mmol) was added followed by *N*-(3-dimethylaminopropyl)-*N'*-ethylcarbodiimide hydrochloride (17 mg, 0.085 mmol). The reaction


15 was stirred for 16.5 h, at which time it was loaded directly onto a silica gel column. Elution with 60 to 100% ethyl acetate in *iso*-hexanes provided the title compound (20 mg, 43%) as an amorphous residue.



A solution of **47c** (19.6 mg, 0.0294 mmol) in toluene (9.4 mL) was sparged with argon for


20 10 min with stirring. Hoveyda- Grubbs 2nd generation - catalyst (2.8 mg, 0.0045 mmol) was then added as a solution in degassed toluene (0.45 mL), and the resulting solution was heated to

105°C. After 25 min, an additional portion of Hoveyda- Grubbs 2nd generation catalyst (1.4 mg, 0.0022 mmol) was added. After 10 additional min, the reaction was cooled to ambient temperature and was concentrated *in vacuo* to ~4.5 mL. The solution was loaded directly onto a silica gel column which was eluted with 80 to 100% ethyl acetate in *iso*-hexanes to afford the title compound (6.9 mg, 38%) as an amorphous solid.




In a polypropylene vial, **47d** (4.7 mg, 0.0075 mmol) was dissolved in tetrahydrofuran (0.90 mL) under argon and the resulting solution was cooled in an ice water bath. HF-pyridine (0.10 mL) was added dropwise. After 10 min, the reaction was quenched by addition to a stirred mixture of ethyl acetate (15 mL) and saturated aqueous sodium bicarbonate (15 mL). The phases were separated, and the aqueous phase was extracted with ethyl acetate (3 x 30 mL). The combined organic phases were dried over sodium sulfate, filtered and concentrated to afford a crude residue. In a polypropylene vial under argon, the aforementioned residue was dissolved in tetrahydrofuran (0.90 mL) and the resulting solution was cooled in an ice water bath. HF-pyridine (0.10 mL) was added dropwise, and the reaction mixture was removed from the cold bath. After 45 min, the reaction was worked up as described above to afford a crude residue. Purification by reverse-phase HPLC (5 to 100% acetonitrile in water, +0.1% trifluoroacetic acid) provided the title compound (1.6 mg, 34%) as an amorphous white solid. ¹H NMR (400 MHz, CD₃OD) 9.54 (s, 1H), 8.38 (d, *J* = 8.7 Hz, 1H), 8.29 (s, 1H), 8.17 (s, 1H), 7.87 (d, *J* = 8.5 Hz, 1H), 6.90–6.75 (m, 2H), 6.18 (q, *J* = 6.9 Hz, 1H), 5.84–5.74 (m, 1H), 4.71–4.64 (m, 1H), 4.50–4.42 (m, 1H), 3.88–3.81 (m, 2H), 3.78 (dd, *J* = 11.3, 6.4 Hz, 1H), 3.52 (dd, *J* = 12.9, 6.2 Hz, 1H), 2.81–2.70 (m, 1H), 2.11–2.04 (m, 1H), 1.99–1.91 (m, 1H), 1.85–1.69 (m, 2H), 1.73 (d, *J* = 6.7 Hz, 3H), 1.60 (d, *J* = 7.1 Hz, 3H). LCMS (*m/z*) 510.2 [*M*+*H*], *Tr* = 2.19 min.

EXAMPLE 48




To a solution of (S)-1-((S)-2-*tert*-butoxycarbonylamino-propionyl)-hexahydro-pyridazine-3-carboxylic acid 2,2,2-trichloro-ethyl ester (320 mg, 0.74 mmol) in dichloromethane (8.8 mL) was added trifluoroacetic acid (2.2 mL). The reaction was stirred at RT for 45 min and was concentrated *in vacuo*. The resulting crude product was twice dissolved in and concentrated from anhydrous toluene (10 mL). The resulting crude residue was used without further purification. The crude residue (ca. 0.74 mmol), 1-hydroxybenzotriazole (153.8 mg, 1.1 mmol), and (2S,3R)-2-*tert*-butoxycarbonylamino-3-hydroxy-butiric acid (161.2 mg, 0.735 mmol) were dissolved in dichloromethane (4 mL). *N,N*-Diisopropylethylamine (192 mg, 1.5 mmol) was added, and the resulting solution was cooled in an ice water bath. *N*-(3-Dimethylaminopropyl)-*N'*-ethylcarbodiimide (132 mg, 0.85 mmol) was added dropwise over 15 s. The reaction was removed from the cold bath and was stirred for 21.5 h. The reaction mixture was then diluted with ethyl acetate (50 mL) and washed with half-saturated aqueous sodium bicarbonate (30 mL). The organic phase was washed with brine, dried over anhydrous sodium sulfate, filtered, and concentrated to afford a crude residue that was purified by silica gel chromatography eluting with 65 to 90 % ethyl acetate in *iso*-hexanes. Impure fractions were repurified by silica gel chromatography and collated with the first product to provide the title compound (230 mg, 58%).




To a solution of **48a** (62.3 mg, 0.12 mmol) in dichloromethane (1 mL) was added 2,4,6-collidine (146 mg, 1.21 mmol). *tert*-Butyldimethylsilyl trifluoromethanesulfonate (94 mg, 0.36 mmol) was added dropwise over 20 s. The reaction mixture was stirred for 15 h and was quenched with saturated aqueous sodium bicarbonate (1 mL). The mixture was further diluted with ethyl acetate (2 mL) and brine (1 mL). The phases were separated, and the aqueous phase was extracted with ethyl acetate (4 x 1.5 mL). The combined organic phases were dried over

sodium sulfate, filtered, and concentrated *in vacuo* to afford a crude residue that was used without further purification. The crude product was dissolved in tetrahydrofuran (3 mL). Acetic acid (104 mg, 1.6 mmol) was added in one portion and the resulting solution was stirred for 3.5 h. The reaction was then diluted with ethyl acetate (30 mL), saturated aqueous sodium bicarbonate (20 mL) and brine (10 mL). The phases were separated, and the aqueous phase was extracted with ethyl acetate (20 mL). The combined organic phases were dried over anhydrous sodium sulfate. The mixture was filtered, and the filtrate was concentrated *in vacuo* to afford crude intermediate amine which was used without further purification. The crude amine was dissolved in *N,N*-dimethylformamide (1.5 mL). 2,4,6-Collidine (28 mg, 0.23 mmol) and *trans*-3-pentenoic acid (14.3 mg, 0.143 mmol) were added, and the resulting solution was cooled in an ice water bath. (1-Cyano-2-ethoxy-2-oxoethylidenaminoxy)dimethylamino-morpholino-carbenium hexafluorophosphate (69 mg, 0.16 mmol) was added in one portion, and the reaction was stirred for 30 min. The reaction was then removed from the cold bath and warmed to ambient temperature. After 15 h, the reaction was diluted with ethyl acetate (25 mL), 0.1 N aqueous hydrochloric acid (30 mL), and brine (5 mL). The phases were separated, and the acidic aqueous layer was extracted with ethyl acetate (25 mL). The combined organic phases were washed with saturated aqueous sodium bicarbonate (25 mL), with brine (5 mL), dried over anhydrous sodium sulfate, filtered, and concentrated. The resulting crude residue was purified by silica gel chromatography (50 to 75% ethyl acetate in *iso*-hexanes) to afford the title compound (74 mg, quantitative yield over 3 steps) as a white foam.




To a solution of **48b** (74 mg, 0.117 mmol) in tetrahydrofuran (2.3 mL) was added water (0.47 mL), ammonium acetate (138 mg, 1.8 mmol), and zinc powder (164 mg, 2.5 mmol). The reaction mixture was stirred vigorously at RT for 17.5 h, at which time the temperature was increased to 35°C. After 25.5 h, additional zinc powder (85 mg, 1.3 mmol) was added and the reaction temperature was increased to 45°C. After 39.5 total hours, the reaction mixture was filtered through a pad of Celite, washing with water and ethyl acetate. The aqueous phase was

acidified with 2 N aqueous hydrochloric acid (15 mL), and the phases were separated. The aqueous phase was extracted with ethyl acetate (2 x 20 mL), and the combined organic phases were dried over anhydrous sodium sulfate, filtered, and concentrated to afford an amorphous white solid (50 mg, 86%) that was used without further purification. The crude product (0.10 mmol) was dissolved along with (R)-1-(3-vinyl-isoquinolin-6-yl)-ethanol (23.9 mg, 0.12 mmol) in dichloromethane (1.0 mL). 4-Dimethylaminopyridine (15 mg, 0.12 mmol) was added followed by *N*-(3-dimethylaminopropyl)-*N'*-ethylcarbodiimide hydrochloride (17 mg, 0.019 mmol). The reaction was stirred for 16 h, at which time it was loaded directly onto a silica gel column. Elution with 50 to 85, and then to 100% ethyl acetate in *iso*-hexanes provided the title compound (40 mg, 58%) as a solid.




A solution of **48c** (26 mg, 0.037 mmol) in toluene (12 mL) was sparged with argon for several min with stirring. Hoveyda-Grubbs 2nd generation catalyst (3.4 mg, 0.0054 mmol) was then added as a solution in degassed toluene (0.40 mL), and the resulting solution was heated to 105°C. After 20 min, an additional portion of Hoveyda-Grubbs 2nd generation catalyst (1.7 mg, 0.0027 mmol) was added as a solution in toluene (0.20 mL). After 40 additional min, the reaction was cooled to RT and was concentrated *in vacuo* to ~3 mL. The solution was loaded directly onto a silica gel column, which was eluted with 70 to 100% ethyl acetate in *iso*-hexanes to afford the title compound (10 mg, 42%) as an amorphous solid.



In a polypropylene vial under argon atmosphere, **48d** (10 mg, 0.016 mmol) was dissolved in tetrahydrofuran (900 μ L). HF·pyridine (~70% as HF, 100 μ L) was added dropwise via syringe, and the resulting solution was stirred for 130 min. An additional aliquot of HF·pyridine (100 μ L) was then added via syringe, and the resulting solution was stirred for an additional 80 min. The reaction mixture was quenched by its careful addition to a stirred mixture of ethyl acetate (50 mL) and saturated aqueous sodium bicarbonate (50 mL). The phases were separated, and the aqueous phase was extracted with ethyl acetate (2 x 30 mL). The combined organic phases were dried over anhydrous sodium sulfate, filtered, and concentrated to afford a crude residue that was purified by reverse phase HPLC (C18, 15 to 100% acetonitrile/water, 0.1% trifluoroacetic acid). Impure fractions from this run were re-purified in the same manner to afford the title compound as the trifluoroacetic acid salt (5.0 mg, 49%) as an amorphous white solid following lyophilization. ^1H NMR (400 MHz, CD_3OD) δ 9.55 (s, 1H), 8.38 (d, J = 8.6 Hz, 1H), 8.22 (s, 1H), 8.15 (s, 1H), 7.87 (dd, J = 8.7, 1.4 Hz, 1H), 6.86–6.75 (m, 2H), 6.15 (q, J = 6.6 Hz, 1H), 5.80–5.70 (m, 1H), 4.47 (d, J = 5.6 Hz, 1H), 4.45–4.38 (m, 1H), 4.09–4.01 (m, 1H), 3.83–3.78 (m, 1H), 3.55 (dd, J = 13.5, 5.6 Hz, 1H), 3.11 (dd, J = 13.2, 4.7 Hz, 1H), 2.78–2.67 (m, 1H), 2.09–1.99 (m, 1H), 1.96–1.89 (m, 1H), 1.82–1.64 (m, 5H), 1.60–1.54 (m, 3H), 1.25 (d, J = 6.3 Hz, 3H). LCMS (m/z) 524.6 [$M+H$], Tr = 2.15 min.


EXAMPLE 49




Compound 49a

A solution of (S)-1-((S)-2-*tert*-butoxycarbonylamino-propionyl)-hexahydro-pyridazine-3-carboxylic acid 2,2,2-trichloro-ethyl ester (865 mg, 2 mmol) in dichloromethane (20 mL) was cooled in an ice water bath. Trimethylsilyl trifluoromethanesulfonate (667 mg, 3 mmol) was added dropwise at 0 °C under argon, and the resulting solution was stirred at RT for 30 min. The reaction mixture was evaporated to dryness and the resulting crude residue was dissolved in anhydrous acetonitrile (25 mL) under argon. The reaction mixture was stirred at 0 °C, (S)-4-

azido-2-(*tert*-butoxycarbonylamino)butanoic acid dicyclohexylamine salt (936 mg, 2.2 mmol) and *N,N*-diisopropylethylamine (1034 mg, 8 mmol) were added followed by 2-(1*H*-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyl uronium hexafluorophosphate methanaminium (1065 mg, 2.8 mmol). The reaction mixture was stirred at RT for 48 h. The solvent was evaporated, the residue was dissolved in ethyl acetate (100 mL) and the solution was washed with 20% water solution of citric acid (2 x 100 mL), water (100 mL) and brine (100 mL), dried over magnesium sulfate, filtered and evaporated. The residue was purified by silica gel chromatography (gradient from 0-40% ethyl acetate + methanol (4/1) in *iso*-hexanes) to afford the title compound (783 mg, 70%) as a white solid after evaporation. $R_f = 0.40$, *iso*-hexanes/ethyl acetate/methanol (6/4/1).




Under argon, (R)-1-(3-chloro-isoquinolin-6-yl)-ethanol (880 mg, 4.23 mmol), (E)-2,2-dimethyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-but-3-enoic acid methyl ester (1.24 g, 4.88 mmol), $\text{PdCl}_2(\text{PCy}_2(p\text{-NMe}_2\text{Ph}))_2$ (bis[(dicyclohexyl)(4-dimethylaminophenyl)phosphine]palladium(II)chloride) 173 mg, 0.21 mmol) and potassium phosphate tribasic (2.64 g, 12.4 mmol) were dissolved in cyclopentyl methyl ether (11.9 mL) and water (5.1 mL). The resulting biphasic mixture was vigorously stirred at 90°C for 3.5 h, at which time the reaction was cooled to ambient temperature and was diluted with ethyl acetate (50 mL) and water (40 mL). The phases were separated, and the aqueous phase was extracted with ethyl acetate (2 x 50 mL). The combined organic phases were dried over anhydrous magnesium sulfate, filtered, and concentrated to afford a crude residue that was purified by silica gel chromatography (25 to 60% ethyl acetate in *iso*-hexanes) to afford the title compound (1.07 g, 85%) as a yellow oil.




To a solution of **49b** (600 mg, 2 mmol) in tetrahydrofuran (8 mL) was added methanol (4 mL), water (4 mL) and lithium hydroxide (96 mg, 4 mmol). The resulting mixture was stirred at RT for 10 h and quenched with 1 M hydrochloric acid (4.2 mL, 4.2 mmol). The resulting solution

was concentrated to a crude residue which was co-distilled twice with tetrahydrofuran (20 mL), twice with anhydrous acetonitrile (20 mL) and twice with anhydrous toluene (20 mL). The resulting white solid was dried under high vacuum overnight and it was used without further purification (735 mg, quantitative yield).



A solution of **49a** (169 mg, 0.302 mmol) in dichloromethane (10 mL) was cooled in an ice water bath. Trimethylsilyl trifluoromethanesulfonate (101 mg, 0.455 mmol) was added dropwise at 0°C under argon, and the resulting solution was stirred at RT for 30 min. The reaction mixture was evaporated to dryness and the resulting crude residue was dissolved in anhydrous acetonitrile (20 mL) under argon. Reaction mixture was stirred at 0°C, (E)-4-[6-((R)-1-hydroxy-ethyl)-isoquinolin-3-yl]-2,2-dimethyl-but-3-enoic acid (123 mg, 0.333 mmol) and *N,N*-diisopropylethylamine (151 mg, 1.220 mmol) were added followed by 2-((1H-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyl uronium hexafluorophosphate methanaminium (161 mg, 0.423 mmol). The reaction mixture was stirred at RT for 2 days. The solvent was evaporated, the residue was dissolved in ethyl acetate (50 mL) and the solution was washed with 20% water solution of citric acid (2 x 50 mL), water (50 mL) and brine (50 mL), dried over magnesium sulfate, filtered and evaporated. The residue was purified by silica gel chromatography (gradient from 0-40% ethyl acetate and methanol mixture (4/1) in *iso*-hexanes) to afford the title compound (198 mg, 90%) as a white solid after evaporation. $R_f = 0.18$, *iso*-hexanes/ethyl acetate/methanol (6/4/1).



To a solution of **49d** (170 mg, 0.23 mmol) in tetrahydrofuran (2 mL) was added methanol (1 mL), water (1 mL) and lithium hydroxide hydrate (7 mg, 0.28 mmol). The mixture was stirred for 2 h at ambient temperature and was quenched with aqueous 1 M hydrochloric acid (0.30 mL, 0.30 mmol). The resulting solution was concentrated to a crude residue which was co-

5 evaporated twice with tetrahydrofuran (5 mL), twice with anhydrous acetonitrile (5 mL) and twice with anhydrous toluene (5 mL). The resulting white solid was dried under high vacuum overnight and used without further purification (151 mg, quantitative yield). Into oven-dried, argon purged flask were placed 2-methyl-6-nitrobenzoic anhydride (317 mg, 0.92 mmol), 4-

10 dimethylaminopyridine (337 mg, 2.76 mmol) and anhydrous 1,2-dichloroethane (300 mL). The resulting solution was heated at 50°C, and the crude seco-acid was added dropwise by syringe as a solution in dry *N,N*-dimethylformamide (5 mL) over 12 h. An additional portion of dry *N,N*-dimethylformamide (2 x 1 mL) was used to complete the quantitative transfer. After stirring for additional 2 h at 50°C, the reaction mixture was transferred to a separatory funnel and washed with water (200 mL, 10 mL of brine was added to support the separation). The aqueous phase


15 was extracted with dichloromethane (100 mL). Combined organic extracts were washed with brine (100 mL) and dried over magnesium sulfate, filtered and concentrated under reduced pressure. The resulting residue was dissolved in ethyl acetate (250 mL) and was washed with water (300 mL, 10 mL of brine was added to support the separation). The aqueous phase was extracted with ethyl acetate (150 mL). Combined organic extracts were washed with water (200

20 mL, 10 mL of brine was added to support the separation). Resulting aqueous phase was extracted with ethyl acetate (150 mL). Combined organic extracts were washed with brine (150 mL) and dried over magnesium sulfate, filtered and concentrated under reduced pressure. The residue was purified by silica gel chromatography (gradient from 0-40% ethyl acetate and methanol mixture (4/1) in *iso*-hexanes) to afford the title compound (76 mg, 57%) as a white

25 solid after evaporation. R_f = 0.51, 10% methanol in dichloromethane. ^1H NMR (400 MHz, CD_3OD) δ 9.03 (s, 1H), 7.95 (d, J = 8.5 Hz, 1H), 7.81 (s, 1H), 7.49 (s, 1H), 7.46 (dd, J = 8.5, 1.6 Hz, 1H), 6.49 (d, J = 16.1 Hz, 1H), 6.40 (d, J = 16.1 Hz, 1H), 5.95 (q, J = 6.6 Hz, 1H), 5.51 (q, J = 7.2 Hz, 1H), 4.67 (dd, J = 8.6, 6.3 Hz, 1H), 4.29 (m, 1H), 3.68 (dd, J = 11.2, 2.7 Hz, 1H), 3.28 (td, J = 6.8, 3.7 Hz, 2H), 2.67–2.56 (m, 1H), 1.88 (m, 2H), 1.85–1.60 (m, 4H), 1.57 (d, J = 6.7


30 Hz, 3H), 1.54 (d, J = 7.3 Hz, 3H), 1.42 (s, 3H), 1.27 (s, 3H). LCMS (m/z) 557.3 ($[\text{M} + \text{H}]$), Tr = 3.15 min.

EXAMPLE 50



Into an oven-dried, argon purged flask, Compound 49 (20 mg, 0.034 mmol), copper(I) iodide (1 mg, 0.005 mmol) and propiolic acid (5 mg, 0.070 mmol) were added. The flask was sealed with septa and repurged with argon three times. Anhydrous *N,N*-dimethylformamide (5 mL) was added and the reaction mixture was repurged with argon three times. This reaction mixture was heated at 70°C for 2 days. After evaporation of the solvent under reduced pressure, the crude residue was dissolved in ethyl acetate (10 mL) and filtered through Celite and the filter pad was washed with ethyl acetate (10 mL). After concentration under reduced pressure, the residue was purified by silica gel chromatography (gradient from 0-40% ethyl acetate and methanol mixture (4/1) in *iso*-hexanes) to afford the title compound (7 mg, 34%) as a white solid. R_f = 0.36, 5% methanol in dichloromethane. ^1H NMR (400 MHz, CD_3OD): δ 9.05 (s, 1H), 8.11 (*br s*, 1H), 8.00 (*br s*, 1H), 7.95 (d, J = 8.8 Hz, 1H), 7.82 (s, 1H), 7.66 (s, 1H), 7.48 (d, J = 10.0 Hz, 1H), 6.50 (d, J = 16.2 Hz, 1H), 6.42 (d, J = 16.2 Hz, 1H), 5.97 (m, 1H), 5.56 (m, 1H), 4.60 (m, 1H), 4.29 (m, 1H), 3.66 (m, 1H), 3.21 (m, 2H), 2.69–2.47 (m, 1H), 1.89 (m, 2H), 1.87–1.55 (m, 4H), 1.53 (d, J = 6.7 Hz, 3H), 1.50 (d, J = 7.1 Hz, 3H), 1.40 (s, 3H), 1.28 (s, 3H). LCMS (m/z) 603.1 $[\text{M}+\text{H}]^+$ T_r = 2.62 min.


EXAMPLE 51




Compound 51a

A solution of (S)-1-((S)-2-*tert*-butoxycarbonylamino-propionyl)-hexahydro-pyridazine-3-carboxylic acid 2,2,2-trichloro-ethyl ester (865 mg, 2 mmol) in dichloromethane (20 mL) was cooled in an ice water bath. Trimethylsilyl trifluoromethanesulfonate (667 mg, 3 mmol) was

added dropwise at 0°C under argon, and the resulting solution was stirred at RT for 30 min. The reaction mixture was evaporated to dryness and the resulting crude residue was dissolved in anhydrous acetonitrile (25 mL) under argon. Reaction mixture was stirred at 0°C, (S)-2-(*tert*-butoxycarbonylamino)pent-4-ynoic acid (469 mg, 2.2 mmol, source: Matrix Scientific, Catalog Number 041479) and *N,N*-diisopropylethylamine (1034 mg, 8 mmol) were added followed by 2-(1H-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyl uronium hexafluorophosphate methanaminium (1065 mg, 2.8 mmol). Reaction mixture was stirred at RT for 48 h. The solvent was evaporated, the residue was dissolved in ethyl acetate (100 mL) and the solution was washed with 20% water solution of citric acid (2 x 100 mL), water (100 mL) and brine (100 mL), dried over magnesium sulfate, filtered and evaporated. The residue was purified by silica gel chromatography (gradient from 0-40% ethyl acetate and methanol (4/1) in *iso*-hexanes) to afford the title compound (738 mg, 70%) as a white solid after evaporation. R_f = 0.30, *iso*-hexanes/ethyl acetate/methanol (6/4/1).




A solution of **51a** (145 mg, 0.275 mmol) in dichloromethane (10 mL) was cooled in an ice water bath. Trimethylsilyl trifluoromethanesulfonate (92 mg, 0.414 mmol) was added dropwise at 0°C under argon, and the resulting solution was stirred at RT for 30 min. The reaction mixture was evaporated to dryness and the resulting crude residue was dissolved in anhydrous acetonitrile (20 mL) under argon. The reaction mixture was stirred at 0°C, (E)-4-[6-((R)-1-hydroxy-ethyl)-isoquinolin-3-yl]-2,2-dimethyl-but-3-enoic acid (112 mg, 0.303 mmol) and *N,N*-diisopropylethylamine (137 mg, 1.101 mmol) were added followed by 2-(1H-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyl uronium hexafluorophosphate methanaminium (147 mg, 0.385 mmol). The reaction mixture was stirred at RT for 2 days. The solvent was evaporated, the residue was dissolved in ethyl acetate (50 mL) and the solution was washed with 20% water solution of citric acid (2 x 50 mL), water (50 mL) and brine (50 mL), dried over magnesium sulfate, filtered and evaporated. The residue was purified by silica gel chromatography (gradient from 0-40% ethyl acetate and methanol (4/1) in *iso*-hexanes) to afford the title compound (163 mg, 85%) as a white solid after evaporation. R_f = 0.29, *iso*-hexanes/ethyl acetate/methanol (6/4/1).



To a solution of **51b** (125 mg, 0.18 mmol) in tetrahydrofuran (2 mL) was added methanol (1 mL), water (1 mL) and lithium hydroxide hydrate (5.2 mg, 0.22 mmol). The mixture was stirred for 2 h at ambient temperature and was quenched with aqueous 1 M hydrochloric acid (0.25 mL, 0.25 mmol). The resulting solution was concentrated to a crude residue which was co-evaporated twice with tetrahydrofuran (5 mL), twice with anhydrous acetonitrile (5 mL) and twice with anhydrous toluene (5 mL). The resulting white solid was dried under high vacuum overnight and it was used without further purification (113 mg, quantitative yield). Into an oven-dried, argon purged flask were placed 2-methyl-6-nitrobenzoic anhydride (248 mg, 0.72 mmol), 4-dimethylaminopyridine (264 mg, 2.16 mmol) and anhydrous 1,2-dichloroethane (200 mL). The resulting solution was heated at 50°C, and the crude seco-acid was added dropwise via syringe as a solution in anhydrous *N,N*-dimethylformamide (5 mL) over 12 h. An additional portion of anhydrous *N,N*-dimethylformamide (2 x 1 mL) was used to complete the quantitative transfer. After stirring for additional 2 h at 50°C, the reaction mixture was transferred to a separatory funnel and washed with water (200 mL, 10 mL of brine was added to support the separation). The aqueous phase was extracted with dichloromethane (100 mL). Combined organic extracts were washed with brine (100 mL) and dried over magnesium sulfate, filtered and concentrated under reduced pressure. The resulting residue was dissolved in ethyl acetate (250 mL) and was washed with water (300 mL, 10 mL of brine was added to support the separation). The aqueous phase was extracted with ethyl acetate (150 mL). Combined organic extracts were washed with water (200 mL, 10 mL of brine was added to support the separation). Resulting aqueous phase was extracted with ethyl acetate (150 mL). Combined organic extracts were washed with brine (150 mL) and dried over magnesium sulfate, filtered and concentrated under reduced pressure. The residue was purified by silica gel chromatography (gradient from 0-40% ethyl acetate + methanol (4/1) in *iso*-hexanes) to afford the title compound (56 mg, 57%) as a white solid after evaporation. R_f = 0.53, 10% methanol in dichloromethane. ^1H NMR (400 MHz, CD_3OD) δ 9.03 (1H), 7.94 (d, J = 8.5 Hz, 1H), 7.79 (s, 1H), 7.54 (s, 1H), 7.45 (d, J = 8.5 Hz, 1H), 6.52 (d, J = 16.1 Hz, 1H), 6.40 (d, J = 16.1 Hz, 1H), 5.95 (q, J = 6.6 Hz, 1H), 5.52 (q, J = 7.2 Hz, 1H), 4.70


(dd, $J = 7.7, 6.7$ Hz, 1H), 4.30 (m, 1H), 4.00 (q, $J = 7.2$ Hz, 1H), 3.72–3.64 (m, 1H), 2.63 (m, 2.55–2.40 (m, 2H), 1.91 (s, 1H), 1.88 (m, 1H), 1.80 (m, 1H), 1.64 (m, 1H), 1.57 (d, $J = 6.7$ Hz, 3H), 1.53 (d, $J = 7.2$ Hz, 3H), 1.42 (s, 3H), 1.27 (s, 3H). LCMS (m/z) 546.2 $[M+H]^+$ Tr = 3.04 min.

5

EXAMPLE 52

A solution of Compound 51 (10 mg, 0.018 mmol) in a mixture of ethyl acetate (4 mL) and tetrahydrofuran (4 mL) containing 5% palladium on calcium carbonate poisoned with lead - Lindlar catalyst (10 mg) was hydrogenated at RT and pressure of hydrogen for 3 h. The reaction mixture was filtered through Celite and the filter pad was washed with tetrahydrofuran (10 mL). The filtrate was evaporated to afford the title compound (10 mg, quantitative yield) as a white solid. $R_f = 0.19$, *iso*-hexanes/ethyl acetate/methanol (6/4/1). ^1H NMR (400 MHz, CD_3OD): δ 9.04 (s, 1H), 7.95 (d, $J = 8.4$ Hz, 1H), 7.82 (s, 1H), 7.54 (s, 1H), 7.46 (d, $J = 8.3$ Hz, 1H), 6.50 (d, $J = 16.1$ Hz, 1H), 6.39 (d, $J = 16.1$ Hz, 1H), 5.95 (q, $J = 6.4$ Hz, 1H), 5.76–5.62 (m, 1H), 5.57 (q, $J = 7.0$ Hz, 1H), 5.02 (d, $J = 17.0$ Hz, 1H), 4.96 (d, $J = 10.1$ Hz, 1H), 4.61–4.54 (m, 1H), 4.29 (m, 1H), 3.68n (m, 1H), 2.61 (m, 1H), 2.42–2.34 (m, 1H), 2.30–2.20 (m, 1H), 1.90 (m, 1H), 1.81 (m, 1H), 1.68–1.60 (m, 2H), 1.57 (d, $J = 6.6$ Hz, 3H), 1.53 (d, $J = 7.2$ Hz, 3H), 1.39 (s, 3H), 1.26 (s, 3H). LCMS (m/z) 548.3 $[M+H]^+$ Tr = 2.85 min.

20


EXAMPLE 53

To a solution of (S)-1-((S)-2-*tert*-butoxycarbonylamino-propionyl)-hexahydro-pyridazine-3-carboxylic acid 2,2,2-trichloro-ethyl ester (300 mg, 0.69 mmol) in dichloromethane (1.84 mL)

was slowly added trifluoroacetic acid (460 μ L, 6.00 mmol) at 0°C under an argon atmosphere. After 3 h, the reaction mixture was concentrated under reduced pressure. The crude residue was dissolved in *N,N*-dimethylformamide (3.45 mL) and (*S*)-*tert*-Butoxycarbonylamino-cyclohexyl-acetic acid (195 mg, 0.760 mmol), 2-(1H-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyl

5 uronium hexafluorophosphate methanaminium (289 mg, 0.760 mmol), and *N,N*-diisopropylethylamine (180 μ L, 1.04 mmol) were sequentially added at 23°C under an argon atmosphere. After 18 h, the reaction mixture was diluted with ethyl acetate (300 mL), and the resulting mixture was washed with brine (4 $\times$ 100 mL), was dried over anhydrous sodium and was concentrated under reduced pressure. The crude residue was purified by silica gel

10 chromatography (12 g Combiflash HP Gold Column, 0-100% ethyl acetate/*iso*-hexanes gradient) to afford the title compound (380 mg, 96%) as a colorless oil.




To a solution of **53a** (220 mg, 0.385 mmol) in dichloromethane (1.92 mL) was added trimethylsilyl trifluoromethanesulfonate (128 mg, 0.587 mmol) at 0°C under an argon


15 atmosphere. After 1.5 h, the reaction mixture was concentrated under reduced pressure. The resulting residue was diluted with acetonitrile (1.92 mL) and 2-(1H-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyl uronium hexafluorophosphate methanaminium (146 mg, 0.385 mmol), *N,N*-diisopropylethylamine (267 μ L, 1.54 mmol), and (*E*)-pent-3-enoic acid (39.4 μ L, 0.385 mmol) were sequentially added at 23°C under an argon atmosphere. After 20 h, the reaction mixture

20 was diluted with dichloromethane (40 mL) and the resulting mixture was washed with saturated aqueous sodium bicarbonate solution (40 mL) and with brine (2 $\times$ 40 mL). The organic layer was separated, was dried over anhydrous sodium sulfate, and was concentrated under reduced pressure. The crude residue was purified by silica gel flash column chromatography (12 g Combiflash HP Gold Column, 0-100% ethyl acetate/*iso*-hexanes gradient) to afford the title


25 compound (180 mg, 84%) as a colorless oil. R_f = 0.75 (ethyl acetate) I_2 /silica stain.



To a solution of **53b** (180 mg, 0.320 mmol) in tetrahydrofuran (5.3 mL) was added zinc powder (418 mg, 6.40 mmol) followed by a solution of ammonium acetate (370 mg, 4.80 mmol) in water (3.5 mL) at 23°C under an argon atmosphere. After 15 h, the reaction mixture was warmed to 45°C. After 2 h, the reaction mixture was allowed to cool to RT and was filtered through a pad of Celite washing with water (10 mL) and ethyl acetate (10 mL). The filtrate layers were split and the aqueous layer was diluted with brine (80 mL) and was acidified to pH 1 with 12 N aqueous hydrogen chloride solution. The aqueous layer was extracted with ethyl acetate (3 × 100 mL), and the combined organic extracts were dried over anhydrous sodium sulfate, and were concentrated under reduced pressure. Residual acetic acid was removed azeotropically via addition of toluene (5 mL) followed by concentration under reduced pressure (3 ×) to afford the title compound (74.5 mg, 55%) as a white solid.




To a solution of **53c** (74.5 mg, 0.176 mmol) and (R)-1-(3-vinyl-isoquinolin-6-yl)-ethanol (42.1 mg, 0.212 mmol) in dichloromethane (3.5 mL) were added *N*-(3-dimethylaminopropyl)-*N'*-ethylcarbodiimide hydrochloride (47.2 mg, 0.246 mmol) and 4-dimethylaminopyridine (11 mg, 88 μmol) at 23°C under an argon atmosphere. After 18 h, the reaction mixture was purified directly by silica gel chromatography (12 g Combiflash HP Gold Column, 0–100% ethyl acetate/*iso*-hexanes gradient) to afford the title compound (67.6 mg, 64%) as a white solid.



To a solution of **53d** (25 mg, 41 μmol) in toluene (8.2 mL) was added the Hoveyda-Grubbs 2nd Generation catalyst (2.5 mg, 4.1 μmol) at 23°C under an argon atmosphere, and the resulting mixture was heated to 110°C. After 2 h, the reaction mixture was quenched with ethyl vinyl ether (300 μL) and the resulting mixture was allowed to cool to 23°C. The reaction mixture was concentrated under reduced pressure and the crude residue was purified by silica gel

chromatography (12 g Combiflash HP Gold Column, 0-100% ethyl acetate/*iso*-hexanes gradient). The fractions containing the desired product were combined and were repurified by preparatory HPLC to afford the title compound (1.7 mg, 7%) as a white powder as a trifluoroacetic acid salt. R_f = 0.40 (ethyl acetate) UV. ^1H NMR (400 MHz, CD_3OD) δ 9.13 (s, 1H), 8.53 (*br s*, 1H), 8.07 (d, J = 7.6 Hz, 1H), 8.05 (d, J = 8.5 Hz, 1H), 7.83 (s, 1H), 7.55 (d, J = 8.3 Hz, 1H), 7.44 (s, 1H), 6.62 (d, J = 16.3 Hz, 1H), 6.55–6.45 (m, 1H), 6.04 (q, J = 6.5 Hz, 1H), 5.50 (q, J = 6.9 Hz, 1H), 4.40 (d, J = 14.1 Hz, 1H), 4.33 (d, J = 9.7 Hz, 1H), 3.78 (d, J = 11.0 Hz, 1H), 3.36 (*app dd*, J = 14.7, 6.7 Hz, 1H), 2.99 (dd, J = 14.5, 4.9 Hz, 1H), 2.90–2.66 (m, 4H), 2.02–1.83 (m, 4H), 1.82–1.70 (m, 3H), 1.68 (d, J = 6.7 Hz, 3H), 1.65 (d, J = 7.3 Hz, 3H), 1.33–1.16 (m, 4H), 1.16–0.94 (m, 2H). HPLC $\text{Tr} = t_R$ (min), 3.091 (Synergi 4u hydro-RP, 50×4.60 mm 4 micron column, 7 min, 2ml/min, 5–100% acetonitrile / water, 0.05% trifluoroacetic acid modifier gradient). LCMS (m/z) 562.3 [$\text{M}+\text{H}$], $\text{Tr} = 2.17$ min.


EXAMPLE 54



Compound 54a


To a solution of (S)-1-((S)-2-*tert*-butoxycarbonylamino-propionyl)-hexahydro-pyridazine-3-carboxylic acid 2,2,2-trichloro-ethyl ester (300 mg, 0.69 mmol) in dichloromethane (1.84 mL) was slowly added trifluoroacetic acid (460 μL , 6.00 mmol) at 0°C under an argon atmosphere. After 2 h, the reaction mixture was concentrated under reduced pressure. The crude residue was dissolved in *N,N*-dimethylformamide (3.45 mL) and (S)-2-*tert*-Butoxycarbonylamino-3-phenyl-propionic acid (201 mg, 0.760 mmol), 2-(1H-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate methanaminium (289 mg, 0.760 mmol), and *N,N*-diisopropylethylamine (180 μL , 1.04 mmol) were sequentially added at 23°C under an argon atmosphere. After 22 h, the reaction mixture was diluted with ethyl acetate (300 mL), and the resulting mixture was washed with brine (3×200 mL), was dried over anhydrous sodium and was concentrated under reduced pressure. The crude residue was purified by silica gel chromatography (24 g Combiflash HP Gold Column, 0-100% ethyl acetate/*iso*-hexanes

gradient) to afford the title compound (319 mg, 96%) as a colorless oil. $R_f = 0.75$ (ethyl acetate) I_2 /silica stain.



Compound 54b


To a solution of **54a** (414 mg, 0.716 mmol) in dichloromethane (3.58 mL) was added trimethylsilyl trifluoromethanesulfonate (238.7 mg, 1.07 mmol) at 0°C under an argon atmosphere. After 1 h, the reaction mixture was concentrated under reduced pressure. The resulting residue was diluted with acetonitrile (3.58 mL) and 2-(1H-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyl uronium hexafluorophosphate methanaminium (272 mg, 0.716 mmol), *N,N*-diisopropylethylamine (498 μ L, 2.86 mmol), and (*E*)-pent-3-enoic acid (73.3 μ L, 0.716 mmol) were sequentially added at 23°C under an argon atmosphere. After 17 h, the reaction mixture was diluted with dichloromethane (50 mL) and the resulting mixture was washed with saturated aqueous sodium bicarbonate solution (50 mL). The organic layer was separated, was dried over anhydrous sodium sulfate, and was concentrated under reduced pressure. The crude residue was purified by silica gel chromatography to afford the title compound (386 mg, 96%) as white solid.



Compound 54c


To a solution of **54b** (443 mg, 0.789 mmol) in tetrahydrofuran (13.2 mL) was added zinc powder (1.03 g, 15.8 mmol) followed by a solution of ammonium acetate (912 mg, 11.8 mmol) in water (8.77 mL) at 23°C under an argon atmosphere. After 17 h, the reaction mixture was warmed to 45°C. After 2 h, the reaction mixture was allowed to cool to RT and was filtered through a pad of Celite washing with water (10 mL) and ethyl acetate (10 mL). The filtrate layers were split and the aqueous layer was diluted with brine (20 mL) and was acidified to pH 2 with 12 N aqueous hydrogen chloride solution. The aqueous layer was extracted with dichloromethane (3 $\times$ 100 mL), and the combined organic extracts were dried over anhydrous sodium sulfate, and were concentrated under reduced pressure. Residual acetic acid was

removed azeotropically via addition of toluene (5 mL) followed by concentration under reduced pressure (3 ×) to afford the title compound (276.2 mg, 81%) as a white solid.



Compound 54d

To a solution of **54c** (275 mg, 0.640 mmol) and (R)-1-(3-vinyl-isoquinolin-6-yl)-ethanol (153 mg, 0.770 mmol) in dichloromethane (3.2 mL) were added *N*-(3-dimethylaminopropyl)-*N'*-ethylcarbodiimide hydrochloride (172 mg, 0.90 mmol) and 4-dimethylaminopyridine (39 mg, 32 μmol) at 23°C under an argon atmosphere. After 23 h, the reaction mixture was purified directly by silica gel chromatography (24 g Combiflash HP Gold Column, 0–100% ethyl acetate/*iso*-hexanes gradient) to afford the title compound (203 mg, 52%) as a white solid.




Compound 54

To a solution of **54d** (170 mg, 278 μmol) in chlorobenzene (56 mL) was added the Hoveyda-Grubbs 2nd Generation catalyst (8.7 mg, 14.0 μmol) at 23°C under an argon atmosphere, and the resulting mixture was heated to 110°C. After 3 h, the reaction mixture was quenched with ethyl vinyl ether (300 μL) and the resulting mixture was allowed to cool to 23°C.

The reaction mixture was concentrated under reduced pressure and the crude residue was purified by silica gel chromatography (12 g Combiflash HP Gold Column, 0–100% ethyl acetate/*iso*-hexanes gradient) to afford the title compound (18.9 mg, 12%) as a tan solid. R_f = 0.25 (ethyl acetate) UV. ¹H NMR (400 MHz, CD₃OD) δ 9.12 (s, 1H), 8.05 (d, J = 8.7 Hz, 1H), 7.84 (s, 1H), 7.56 (d, J = 8.5 Hz, 1H), 7.53 (s, 1H), 7.32–7.12 (m, 5H), 6.66 (d, J = 16.5 Hz, 1H), 6.48 (dt, J = 12.8, 5.6 Hz, 1H), 6.05 (q, J = 6.4 Hz, 1H), 5.53 (q, J = 6.7 Hz, 1H), 4.73 (d, J = 12.2 Hz, 1H), 4.43 (d, J = 11.9 Hz, 1H), 3.80 (*app* t, J = 10.2 Hz, 1H), 3.30–3.22 (m, 1H), 3.09 (dd, J = 14.5, 4.9 Hz, 1H), 2.96–2.69 (m, 3H), 2.04–1.87 (m, 2H), 1.82–1.56 (m, 2H), 1.68 (d, J


= 6.5 Hz, 3H), 1.64 (d, J = 6.9 Hz, 3H). HPLC Tr = 3.060 min. LCMS (m/z) 570.5 [$M+H$], Tr = 2.14 min.

EXAMPLE 55




Compound 55a

To a solution of (S)-1-((S)-2-*tert*-butoxycarbonylamino-propionyl)-hexahydro-pyridazine-3-carboxylic acid 2,2,2-trichloro-ethyl ester (500 mg, 0.16 mmol) in dichloromethane (5.8 mL) was added trimethylsilyl trifluoromethanesulfonate (386 mg, 1.74 mmol) at 0°C under an argon atmosphere. After 1 h, the reaction mixture was concentrated under reduced pressure. The resulting residue was diluted with acetonitrile (5.8 mL) and 2-(1H-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyl uronium hexafluorophosphate methanaminium (485 mg, 1.28 mmol), *N,N*-diisopropylethylamine (302 μ L, 1.74 mmol), *N-tert*-butoxycarbonyl-3-(3-pyridyl)-L-alanine (337 mg, 1.27 mmol) were sequentially added at 23°C under an argon atmosphere. After 20 h, the reaction mixture was concentrated under reduced pressure, and the crude residue was purified by silica gel chromatography (40 g Combiflash HP Gold Column, 0–100% ethyl acetate/*iso*-hexanes gradient) to afford the title compound (570 mg, 85%) as light yellow oil.



Compound 55b


To a solution of **55a** (570 mg, 0.984 mmol) in dichloromethane (5.8 mL) was added trimethylsilyl trifluoromethanesulfonate (386 mg, 1.74 mmol) at 0°C under an argon atmosphere. After 1 h, the reaction mixture was concentrated under reduced pressure to afford the title compound (478 mg) as light yellow oil which was used without further purification.



To a solution of **55b** (120 mg, 0.250 mmol) in acetonitrile (1.25 mL) were sequentially added (E)-4-[6-((R)-1-hydroxy-ethyl)-isoquinolin-3-yl]-2,2-dimethyl-but-3-enoic acid (75 mg, 0.25 mmol), 2-(1H-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyl uronium hexafluorophosphate


5 methanaminium (95 mg, 0.25 mmol), and *N,N*-diisopropylethylamine (173 μ L, 1.00 mmol) at 23°C under an argon atmosphere. *N,N*-Dimethylformamide (100 μ L) was then added to promote solubility of the reagents. After 23 h, the reaction mixture was diluted with saturated aqueous sodium bicarbonate solution (25 mL) and brine (25 mL) and the resulting mixture was extracted with dichloromethane (2 $\times$ 25 mL). The combined organic extracts were dried over anhydrous

10 sodium sulfate and were concentrated under reduced pressure. The crude residue was purified by silica gel chromatography to afford the title compound (76 mg, 41%) as a colorless solid.




To a solution of **55c** (76 mg, 0.10 mmol) in tetrahydrofuran (0.3 mL) and water (0.2 mL) was added lithium hydroxide hydrate (2.4 mg, 0.10 mmol) at 23°C under an argon atmosphere.

15 After 2 h, the reaction mixture was concentrated under reduced pressure to afford the title compound (61 mg, 99%) as a white solid lithium carboxylate salt.



To a solution of **55d** (61 mg, 0.10 mmol) in dichloromethane (50 mL) were added benzotriazol-1-yl-oxytripyrrolidinophosphonium hexafluorophosphate (208 mg, 400 μ mol) and 4-dimethylaminopyridine (366 mg, 3.00 mmol) at 23°C under an argon atmosphere. After 17 h, the reaction mixture was concentrated under reduced pressure and the crude residue was purified by silica gel chromatography (4 g Combiflash HP Gold Column, 0–20% methanol/dichloromethane gradient) to afford the title compound (1.2 mg, 2%) as a colorless solid. R_f = 0.40 (10% methanol in dichloromethane) UV. ^1H NMR (400 MHz, CD_3OD) δ 9.03 (s, 1H), 8.31 (d, J = 1.6 Hz, 1H), 8.27 (dd, J = 4.9, 1.5 Hz, 1H), 7.95 (d, J = 8.5 Hz, 1H), 7.82 (s, 1H), 7.64 (d, J = 8.0 Hz, 1H), 7.53 (s, 1H), 7.46 (dd, J = 8.5, 1.5 Hz, 1H), 7.24 (dd, J = 7.9, 4.9 Hz, 1H), 6.49 (d, J = 16.1 Hz, 1H), 6.37 (d, J = 16.1 Hz, 1H), 5.95 (q, J = 6.5 Hz, 1H), 5.57 (q, J = 7.3 Hz, 1H), 4.65 (d, J = 12.0 Hz, 1H), 4.32 (br d, J = 12.1 Hz, 1H), 3.83 (d, J = 11.8 Hz, 1H), 3.73–3.65 (m, 1H), 3.00 (dd, J = 14.2, 5.0 Hz, 1H), 2.83 (dd, J = 13.9, 9.7 Hz, 1H), 2.64 (br t, J = 11.2 Hz, 1H), 1.95–1.73 (m, 2H), 1.69–1.59 (m, 1H), 1.57 (d, J = 6.7 Hz, 3H), 1.54 (d, J = 7.2 Hz, 3H), 1.33 (s, 3H), 1.16 (s, 3H). HPLC Tr = 4.491 min


EXAMPLE 56




Compound 56a

To a solution of (S)-1-((S)-2-*tert*-butoxycarbonylamino-propionyl)-hexahydro-pyridazine-3-carboxylic acid 2,2,2-trichloro-ethyl ester (791 mg, 1.82 mmol) in dichloromethane (10.0 mL) was slowly added trimethylsilyl trifluoromethanesulfonate (483 μ L, 2.73 mmol) at 0°C under an argon atmosphere. After 45 min, the reaction mixture was concentrated under reduced pressure. The resulting residue was diluted with dichloromethane (10.0 mL) and 2-(1H-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyl uronium hexafluorophosphate methanaminium (761 mg, 2.00 mmol), *N,N*-diisopropylethylamine (1.26 mL, 7.28 mmol), and (S)-2-(*tert*-butoxycarbonylamino)-4-phenylbutanoic acid (Fluka, 560 mg, 2.00 mmol) were sequentially added at 23°C under an argon atmosphere. After 18 h, the reaction mixture was concentrated under reduced pressure. The residue was pre-absorbed on silica and purified by silica gel chromatography (40 g Isco Rf


Gold Column, 0–100% ethyl acetate/*iso*-hexanes gradient) to afford the title compound (799 mg, 74%) as a colorless oil.




To a solution of **56a** (799 mg, 1.34 mmol) in dichloromethane (10.0 mL) was added trimethylsilyl trifluoromethanesulfonate (356 μ L, 2.01 mmol) at 0°C under an argon atmosphere. After 30 min, the reaction mixture was concentrated under reduced pressure. The resulting residue was diluted with acetonitrile (6.0 mL) and 2-(1H-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyl uronium hexafluorophosphate methanaminium (560 mg, 1.47 mmol), *N,N*-diisopropylethylamine (932 μ L, 5.36 mmol), and (E)-pent-3-enoic acid (150 μ L, 1.47 mmol) were sequentially added at 23°C under an argon atmosphere. After 19 h, the reaction mixture was concentrated under reduced pressure. The residue was pre-absorbed on silica and purified by silica gel chromatography to afford the title compound (598 mg, 78%) as a colorless oil.



To a solution of **56b** (167 mg, 0.29 mmol) in tetrahydrofuran (3 mL), methanol (1 mL) and water (1 mL) was added lithium hydroxide hydrate (7.6 mg, 0.32 mmol) at 23°C. After 15 min, the reaction mixture was concentrated to dryness under reduced pressure. The residue was dissolved in water (25 mL) and washed with ethyl acetate. The organic layer was extracted with saturated aqueous sodium bicarbonate solution (25 mL). The combined aqueous layers were acidified with 1 M aqueous hydrochloric acid solution to pH ~2 and extracted with ethyl acetate (3 $\times$ 25 mL), and the combined organic extracts were dried over anhydrous magnesium sulfate, and were concentrated under reduced pressure to afford the title compound (129 mg, 100%) as a white solid.




To a solution of **56c** (129 mg, 0.29 mmol) and (R)-1-(3-vinyl-isoquinolin-6-yl)-ethanol (64 mg, 0.31 mmol) in dichloromethane (3.0 mL) was added *N*-(3-dimethylaminopropyl)-*N'*-ethylcarbodiimide hydrochloride (89 mg, 0.46 mmol) and 4-dimethylaminopyridine (18 mg, 0.14 mmol) at 23°C under an argon atmosphere. After 16 h, the reaction mixture was purified directly by silica gel chromatography to afford the title compound (87 mg, 48%) as a white solid.



To a solution of **56d** (84 mg, 0.13 mmol) in chlorobenzene (27 mL) was added the Hoveyda-Grubbs 2nd Generation catalyst (8 mg, 13 μmol) at 23°C under an argon atmosphere, and the resulting mixture was heated to 110°C. After 2 h, the Hoveyda-Grubbs 2nd Generation catalyst (7.3 mg, 11 μmol) was added under argon at 110°C. The Hoveyda-Grubbs 2nd Generation Catalyst (5 mg, 8 μmol) was then added in the interval of 30 min for three times at which point the reaction was complete. The reaction mixture was quenched with ethyl vinyl ether (1.0 μL) and the resulting mixture was allowed to cool to 23°C. The reaction mixture was concentrated under reduced pressure and the crude residue was purified by silica gel chromatography (24 g Isco Rf Gold Column, 0–100% ethyl acetate/*iso*-hexanes gradient for 10 min and then 100% ethyl acetate for 25 min) to afford the title compound (9.1 mg, 12%) as a pale brown solid. R_f = 0.20 (ethyl acetate). ¹H NMR (400 MHz, CD₃OD) δ 9.00 (s, 1H), 7.93 (d, J = 8.5 Hz, 1H), 7.71 (s, 1H), 7.45 (dd, J = 8.5, 1.5 Hz, 1H), 7.36 (s, 1H), 7.20–7.08 (m, 4H), 7.08–7.00 (m, 1H), 6.55 (d, J = 15.9 Hz, 1H), 6.40 (ddd, J = 15.9, 7.0, 5.1 Hz, 1H), 5.95 (q, J = 6.5 Hz, 1H), 5.40 (q, J = 7.2 Hz, 1H), 4.60 (d, J = 12.1 Hz, 1H), 4.53 (dd, J = 8.2, 6.9 Hz, 1H), 4.30 (d, J = 12.5 Hz, 1H), 3.70–3.63 (m, 1H), 3.31–3.23 (m, 1H), 2.93 (ddd, J = 14.0, 5.1, 1.6


1H), 2.69–2.52 (m, 3H), 2.01–1.73 (m, 4H), 1.71–1.54 (m, 2H), 1.59 (d, $J = 6.7$ Hz, 3H), 1.46 (d, $J = 7.3$ Hz, 3H). HPLC $T_r = 5.108$ min. LCMS (m/z) 584.3 $[M+H]^+$, $T_r = 2.12$ min.

EXAMPLE 57



A solution of (S)-1-[(S)-2-((S)-2-*tert*-butoxycarbonylamino-3-methylbutyrylamino)-propionyl]-hexahydropyridazine-3-carboxylic acid 2,2,2-trichloroethyl ester (1.61 g, 3.03 mmol) in dichloromethane (31 mL) was cooled in an ice water bath under argon. Trimethylsilyl trifluoromethanesulfonate (1.23 g, 5.5 mmol) was added dropwise, and the resulting solution was stirred for 2 h. The reaction was quenched with *N,N*-diisopropylethylamine (1.2 g, 9.2 mmol) and methanol (8.5 mL). The mixture was concentrated *in vacuo* and was redissolved and concentrated from toluene (2 x 25 mL). The resulting crude residue containing (S)-1-[(S)-2-((S)-2-amino-3-methyl-butylamino)-propionyl]-hexahydro-pyridazine-3-carboxylic acid 2,2,2-trichloro-ethyl ester (1.05 g) was used without further purification.


10



To a solution of (E)-4-[6-((R)-1-hydroxy-ethyl)-isoquinolin-3-yl]-2,2-dimethyl-but-3-enoic acid methyl ester (1.05 g, 3.51 mmol) in tetrahydrofuran (8 mL) was added methanol (4 mL), water (4 mL), and lithium hydroxide hydrate (297 mg, 7.08 mmol). The resulting mixture was stirred for 6 h and was quenched with 1 M aqueous hydrochloric acid (7.2 mL, 7.2 mmol). The resulting solution was concentrated to a crude residue that was redissolved and concentrated from anhydrous methanol (50 mL) followed by toluene (50 mL). The resulting yellow solid (1.3 g, 100%) (E)-4-[6-((R)-1-hydroxy-ethyl)-isoquinolin-3-yl]-2,2-dimethyl-but-3-enoic acid was used

20


without further purification. A portion of crude (E)-4-[6-((R)-1-hydroxy-ethyl)-isoquinolin-3-yl]-2,2-dimethyl-but-3-enoic acid (1.05 g, 2.84 mmol) was dissolved in *N,N*-dimethylformamide (12 mL) under argon. *N,N*-diisopropylethylamine (1.85 g, 14.3 mmol) was added followed by 2-(1H-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyl uronium hexafluorophosphate methanaminium (1.42 g, 3.73 mmol). The resulting mixture was stirred for 3 min, at which time crude **57a** was added as a solution in *N,N*-dimethylformamide (8.5 mL), washing with additional *N,N*-dimethylformamide (2 x 5 mL). The reaction was stirred for 40 min and was then diluted with ethyl acetate (200 mL) and water (300 mL). The phases were separated, and the aqueous phase was extracted with ethyl acetate (200 mL). The combined organic phases were washed with water (150 mL), and the second aqueous layer was extracted with ethyl acetate (100 mL). The combined organics were dried over anhydrous sodium sulfate, filtered, and concentrated. The crude residue was purified by silica gel chromatography (35 to 60% acetone in *iso*-hexanes, continuous gradient) to afford the title compound (1.51 g, 71% over 2 steps) as a colorless oil. R_F 0.5 (50% acetone in *iso*-hexanes).



To a solution of **57b** (1.50 g, 2.15 mmol) in tetrahydrofuran (20 mL) was added methanol (10 mL), water (10 mL), and lithium hydroxide hydrate (365 mg, 8.7 mmol). The mixture was stirred for 75 min at ambient temperature and was then quenched with aqueous 1 M hydrochloric acid (8.8 mL, 8.8 mmol). The resulting solution was concentrated *in vacuo*, and the crude product was twice dissolved and concentrated from methanol (40 mL) and suspended and concentrated from acetonitrile (6 x 30 mL) to afford 1.71 g of a colorless solid that was used without further purification. Under argon, 2-methyl-6-nitrobenzoic anhydride (1.85 g, 5.37 mmol) and 4-dimethylaminopyridine (1.97 g, 16.1 mmol) were dissolved in 1,2-dichloroethane (700 mL). The resulting solution was heated to 50°C, and the crude seco-acid was added dropwise via syringe as a solution in *N,N*-dimethylformamide (22 mL) over 6 h. An additional wash with *N,N*-dimethylformamide (1.5 mL) was then added in the same manner over 15 min. After stirring an additional 1.25 h, the reaction mixture was concentrated to a final volume of 200 mL *in vacuo*. The solution was washed with water (250 mL), and the aqueous phase was extracted with


dichloromethane (150 mL). The organic phase was dried over anhydrous magnesium sulfate, filtered, and concentrated. The resulting residue was dissolved in ethyl acetate (200 mL) and was washed with water (150 mL). The aqueous phase was extracted with ethyl acetate (150 mL). The combined organics were washed with water (100 mL), and the second aqueous phase
 5 was extracted with ethyl acetate (100 mL). The combined organics were dried over anhydrous magnesium sulfate, filtered and concentrated. The crude residue was purified by silica gel chromatography (35 to 65% acetone in *iso*-hexanes continuous gradient) to afford 445 mg of the pure title compound as an amorphous white solid along with 389 mg of impure product. The
 10 impure fractions were purified by recrystallization from acetone:*iso*-hexanes to afford an additional 173.5 mg of pure title compound (total: 618.5 mg, 52%). R_F 0.5 (50% acetone in *iso*-hexanes). 1H NMR (400 MHz, CD_3OD) δ 9.13 (s, 1H), 8.04 (d, J = 8.5 Hz, 1H), 7.89 (s, 1H), 7.55 (dd, J = 8.5, 1.6 Hz, 1H), 7.53 (s, 1H), 6.73–6.34 (m, 2H), 6.03 (q, J = 6.6 Hz, 1H), 5.63 (q, J = 7.2 Hz, 1H), 4.67 (d, J = 12.2 Hz, 1H), 4.47–4.21 (m, 2H), 3.81–3.73 (m, 1H), 2.76–2.64 (m, 1H), 2.08–1.81 (m, 3H), 1.80–1.58 (m, 8H), 1.51 (s, 3H), 1.35 (s, 3H), 0.98 (d, J = 6.7 Hz, 3H),
 15 0.93 (d, J = 6.7 Hz, 3H). LCMS (m/z) 550.2 [$M+H$], T_r = 2.74 min.

EXAMPLE 58




A solution of *N,N*-diisopropylamine (2.51 g, 24.8 mmol) in tetrahydrofuran (150 mL)
 20 under argon was cooled in an ice water bath. A solution of *n*-butyllithium in hexanes (2.5 M, 9.7 mL, 24 mmol) was added dropwise over 2 min, and the resulting solution was stirred for 15 additional minutes. The solution was then cooled to $-78^\circ C$ via $CO_{2(s)}$:acetone bath, and ethyl 1-acetylcyclopropanecarboxylate (3.47 g, 22.2 mmol) was added dropwise over 2 min. The solution was stirred for an additional 20 min, and *N*-Phenyl-bis(trifluoromethanesulfonimide) (8.4
 25 g, 23.5 mmol) was added as a solution in tetrahydrofuran (24 mL) via canula over 5 min, washing with additional portions of tetrahydrofuran (2 x 5 mL). The resulting solution was removed from the cold bath. After an additional 30 min, the reaction mixture was concentrated *in vacuo* and was diluted with diethyl ether (200 mL). The organic phase was washed with 1 M aqueous sodium hydroxide (1 x 100 mL, 1 x 30 mL), dried over anhydrous magnesium sulfate,
 30 filtered, and concentrated to afford 6.6 g of a crude oil that was used without further purification. A solution of the crude material from the previous step in anhydrous pyridine (11 mL) was


heated to 90°C. After 16.5 h, the reaction mixture was diluted with diethyl ether (200 mL) and 3 M hydrochloric acid (100 mL) and the phases were separated. The organic phase was washed with 3 M hydrochloric acid (50 mL) and 1 M sodium hydroxide (50 mL), dried over magnesium sulfate, filtered, and concentrated to afford 2.2 g of a crude liquid that was used without further purification. To an argon-flushed vessel containing zirconocene dichloride (410 mg, 1.6 mmol) was added a solution of the crude product from the previous step (2.2 g, ca. 16 mmol) and pinacolborane (3.1 g, 24 mmol) in dichloromethane (8 mL). After 116 h, the stirred reaction mixture was diluted with ethyl acetate (50 mL) and was quenched by dropwise addition of water. The mixture was further diluted with water (50 mL), and the phases were separated. The aqueous phase was extracted with ethyl acetate (2 x 50 mL). The combined organic phases were dried over magnesium sulfate, filtered, and concentrated to a crude residue that was purified by silica gel chromatography (5 to 20% ethyl acetate in *iso*-hexanes, continuous gradient) to afford the title compound (1.26 g, 21% over 3 steps) as an oil that crystallized on standing at -15°C.



To a round-bottomed flask was added (R)-1-(3-chloroisoquinolin-6-yl)ethanol (204 mg, 0.982 mmol), **58a** (314 mg, 1.18 mmol), PdCl₂(PCy₂(p-NMe₂Ph))₂ (bis[(dicyclohexyl)(4-dimethylaminophenyl)phosphine] palladium(II) chloride) (40 mg, 0.049 mmol) and potassium phosphate tribasic (680 mg, 3.2 mmol). The vessel was sealed with a septum cap and was flushed with argon. Cyclopentyl methyl ether (2.8 mL) and water (1.2 mL) were added, and the resulting biphasic mixture was vigorously stirred in an oil bath pre-heated to 90°C. After 6.75 h, the reaction was cooled to ambient temperature and was diluted with ethyl acetate (50 mL) and water (40 mL). The phases were separated, and the aqueous phase was extracted with ethyl acetate (50 mL). The combined organic phases were dried over magnesium sulfate, filtered, and concentrated to afford a crude residue that was purified by silica gel chromatography (35 to 60% ethyl acetate in *iso*-hexanes) to afford the title compound (266 mg, 85%).




To a solution of **58b** (266 mg, 0.854 mmol) in tetrahydrofuran (2 mL) was added methanol (1 mL), water (1 mL), and lithium hydroxide hydrate (70.8 mg, 1.69 mmol). The resulting mixture was stirred for 3 h and was then quenched with 1 M aqueous hydrochloric acid (1.8 mL, 1.8 mmol). Volatiles were removed *in vacuo*, and the resulting solid was suspended in toluene. Volatiles were removed *in vacuo*, and the resulting yellow solid 1-((E)-2-[6-((R)-1-hydroxy-ethyl)-isoquinolin-3-yl]-vinyl)-cyclopropanecarboxylic acid was used without further purification. A solution of (S)-1-[(S)-2-((S)-2-*tert*-butoxycarbonylamino-3-methylbutyrylamino)-propionyl]-hexahydropyridazine-3-carboxylic acid 2,2,2-trichloroethyl ester (486 mg, 0.914 mmol) in dichloromethane (9.4 mL) was cooled in an ice water bath under argon. Trimethylsilyl trifluoromethanesulfonate (370 mg, 1.7 mmol) was added dropwise, and the resulting solution was stirred for 4 h. The reaction was quenched with *N,N*-diisopropylethylamine (360 mg, 2.7 mmol) and methanol (2.5 mL). The mixture was concentrated *in vacuo* and the resulting residue was redissolved and concentrated from toluene (2 x 15 mL). The resulting crude amine was used without further purification. To a solution of crude 1-((E)-2-[6-((R)-1-hydroxy-ethyl)-isoquinolin-3-yl]-vinyl)-cyclopropanecarboxylic acid (ca. 0.854 mmol) in *N,N*-dimethylformamide (3.5 mL) under argon was added *N,N*-diisopropylethylamine (560 mg, 4.3 mmol) and 2-(1H-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyl uronium hexafluorophosphate methanaminium (427 mg, 1.12 mmol). The resulting mixture was stirred for 2 min, at which time crude (S)-1-[(S)-2-((S)-2-amino-3-methyl-butylamino)-propionyl]-hexahydro-pyridazine-3-carboxylic acid 2,2,2-trichloro-ethyl ester from the previous step was added as a solution in *N,N*-dimethylformamide (2.5 mL), washing with two additional portions of *N,N*-dimethylformamide (1.5 mL each). The reaction was stirred for 45 min and was diluted with ethyl acetate (100 mL) and water (150 mL). The phases were separated, and the aqueous phase was extracted with ethyl acetate (100 mL). The combined organic phases were washed with water (75 mL), and the second aqueous layer was extracted with ethyl acetate (75 mL). The combined organics were dried over anhydrous magnesium sulfate, filtered, and concentrated. The crude residue was purified by silica gel chromatography (35 to 60% acetone in *iso*-hexanes, continuous gradient) to afford title compound (467 mg, 78% over 2 steps) as a colorless oil.




To a solution of **58c** (467 mg, 0.67 mmol) in tetrahydrofuran (6 mL) was added methanol (3 mL), water (3 mL), and lithium hydroxide hydrate (113 mg, 2.69 mmol). The mixture was stirred for 1.75 h at ambient temperature and was quenched with aqueous 1 M hydrochloric acid (2.8 mL, 2.8 mmol). The resulting solution was concentrated *in vacuo*, and the crude product was suspended and concentrated from acetonitrile (5 x 20 mL) to afford 530 mg of a pale yellow solid that was used without further purification. Under argon, 2-methyl-6-nitrobenzoic anhydride (283 mg, 0.82 mmol) and 4-dimethylaminopyridine (307 mg, 2.51 mmol) were dissolved in 1,2-dichloroethane (100 mL). The resulting solution was heated to 50°C, and a portion of the crude seco-acid (260 mg, ca. 0.33 mmol) was added dropwise via syringe as a solution in *N,N*-dimethylformamide (3.5 mL) and 1,2-dichloroethane (10 mL) over 6 h. An additional wash of *N,N*-dimethylformamide (1 mL) was then added in the same manner. After stirring an additional 1.25 h, the reaction mixture was concentrated to ~35 mL *in vacuo*. The solution was diluted with ethyl acetate (100 mL) and washed with water (100 mL). The aqueous phase was extracted with ethyl acetate (75 mL), and the combined organic phases were washed with water (50 mL). The second aqueous phase was extracted with ethyl acetate (50 mL), and the combined organics were dried over magnesium sulfate, filtered and concentrated. The crude residue was purified by silica gel chromatography (35 to 60 and then to 100% acetone in *iso*-hexanes) to afford 101 mg of impure product containing the title compound. This material was purified by silica gel chromatography (0 to 5% methanol in ethyl acetate) followed by reverse-phase HPLC (5 to 100% acetonitrile/water + 0.1% trifluoroacetic acid) to afford the title compound as its trifluoroacetic acid salt (37 mg, 17 % over 2 steps). ¹H NMR (400 MHz, CD₃OD) δ 9.55 (s, 1H), 8.37 (d, *J* = 8.6 Hz, 1H), 8.06 (s, 1H), 7.89 (s, 1H), 7.85 (d, *J* = 8.6 Hz, 1H), 6.69 (d, *J* = 16.4 Hz, 1H), 6.54 (d, *J* = 16.4 Hz, 1H), 6.11 (q, *J* = 6.7 Hz, 1H), 5.75–5.67 (m, 1H), 4.46–4.36 (m, 1H), 4.36–4.28 (m, 1H), 3.90–3.78 (m, 1H), 2.80–2.68 (m, 1H), 2.05–1.86 (m, 3H), 1.80–1.68 (m, 5H), 1.66–1.57 (m, 5H), 1.40–1.30 (m, 1H), 1.30–1.21 (m, 1H), 1.03–0.93 (m, 4H), 0.90 (d, *J* = 6.7 Hz, 3H). LCMS (*m/z*) 548.4 [M+H], Tr = 2.79 min.


EXAMPLE 59




- Into an oven dried, argon purged flask were placed oil-free potassium hydride (from 1740 mg of *ca.* 30% dispersion in mineral oil, *ca.* 13 mmol), anhydrous tetrahydrofuran (10 mL) and hexane (1 mL). The flask was repurged with argon and trichloroethylene (900 μ L, 1.32 g, 10 mmol) was added followed by dry methanol (10 μ L, 7.9 mg, 0.25 mmol). This mixture was stirred at RT for two h. After this time, hexane (10 mL) was added and the resulting solution was immediately used in the subsequent step.




- Into an oven dried, argon purged flask tetrahydrofuran (50 mL) was added and the solution was cooled with an ice bath. A 1.8 M solution of lithium diisopropylamide (7.2 mL, 13 mmol) in tetrahydrofuran/heptane/ethylbenzene was added. The resulting solution was cooled to -78°C , and treated dropwise with methyl tetrahydro-2H-pyran-4-carboxylate (1.20 mL, 1.30 g, 9 mmol) followed by hexamethylphosphoramide (1.56 mL, 1.61 g, 9 mmol). The resulting solution was warmed to 0°C , stirred for 20 min., cooled to -78°C , and treated dropwise with pre-cooled (0°C) solution of 1,2-dichloro-ethyne (*ca.* 10 mmol). The reaction mixture was stirred at -78°C for 30 min and then allowed to warm to RT. After 4 h at RT, the reaction mixture was poured into crushed ice and extracted with diethyl ether (200 mL) (5 mL of brine was added to support the separation). The organic phase was separated and washed with water (200 mL). This water phase was extracted with diethyl ether (100 mL). The combined organic fractions were washed with brine (100 mL), dried over magnesium sulfate, filtered through a 2 cm layer of silica gel (silica gel layer was washed with 50 mL of ethyl acetate), and then concentrated under reduced pressure. The crude product was subjected to silica gel chromatography (gradient from 0-15% ethyl acetate in *iso*-hexanes) to afford the title compound (1.22 g, 67%) as a colorless oil. $R_f = 0.48$, 30% ethyl acetate in *iso*-hexanes, phosphomolybdic acid in ethanol.




59b (1.01 g, 5 mmol) and copper powder (1.6 g, 25 mmol) were suspended in tetrahydrofuran (100 mL). Acetic acid (15 mL) was added and the reaction mixture was heated to 60°C for 3 h. After this time, the reaction mixture was poured onto water (copper powder was filtered off with the use of the filtration paper) and extracted with diethyl ether (3 x 50 mL). The combined organic extracts were washed with saturated solution of ammonium chloride (3 x 50 mL), with saturated solution of sodium bicarbonate (2 x 50 mL) and with water (50 mL). This water phase was extracted with diethyl ether (50 mL). Combined organic extracts were washed with brine (50 mL) and dried over magnesium sulfate, filtered through a 2 cm layer of silica gel (silica gel layer was washed with 50 mL of ethyl acetate), and concentrated under reduced pressure. After drying under high vacuum for one day, the title compound was isolated (0.84 g, quantitative yield) as a colorless oil. R_f = 0.37, 30% ethyl acetate in *iso*-hexanes, phosphomolybdic acid in ethanol.




Into an oven dried, argon purged flask were placed **59c** (0.84 g, 5 mmol) and dichloromethane (2mL). This mixture was cooled to 0°C. Pinacolborane (0.96 g, 7.5 mmol) was then added dropwise via syringe. After the mixture was stirred for 1 min., it was transferred by a syringe into another oven dried, argon purged flask, immersed in an ice bath and protected from light, containing zirconocene dichloride (0.13 g, 0.5 mmol). An additional portion of dichloromethane (2 mL) was used to complete the quantitative transfer. After this the mixture was warmed to ambient temperature, it was stirred in the dark for 72 h to achieve full conversion. The reaction mixture was diluted with ethyl acetate (50 mL) and carefully quenched with water (1ml). Water (50 mL) was added and the organic and aqueous phases were separated. The aqueous phase was extracted with ethyl acetate (3 x 40 mL). The combined organic phases were dried over magnesium sulfate, filtered and evaporated to afford the crude residue which was dissolved in *iso*-hexanes (50 mL) and extracted with 5-10% aqueous methanol (3 x 50 mL) and with brine (50 mL). The hexane phase was dried over magnesium sulfate. The title compound was isolated as the white crystalline compound after evaporation (1.42 g, 93%). R_f = 0.38, 30% ethyl acetate in *iso*-hexanes, iodine vapor.




Into an oven dried, argon purged flask were placed (R)-1-(3-chloro-isoquinolin-6-yl)-ethanol (0.83 g, 4 mmol), **59d** (1.40 g, 4.7 mmol), PdCl₂(PCy₂(p-NMe₂Ph))₂ (bis[(dicyclohexyl)(4-dimethylaminophenyl)phosphine] palladium(II) chloride) (173 mg, 0.21 mmol) and potassium phosphate tribasic (2.64 g, 12.4 mmol). The flask was sealed with a septum cap and was re-purged with argon. *N,N*-Dimethylformamide (10 mL) was added and, and the resulting reaction mixture was vigorously stirred in an oil bath pre-heated to 80°C. After 2 h, the reaction was cooled to ambient temperature and was diluted with ethyl acetate (100 mL) and water (100 mL). The phases were separated, and the aqueous phase was extracted with ethyl acetate (2 x 50 mL). The combined organic phases were dried over magnesium sulfate, filtered, and concentrated to afford a crude residue that was purified by silica gel chromatography (gradient from 0-40% ethyl acetate and methanol (4/1) in *iso*-hexanes) to afford the title compound (1.06 g, 78%) as a colorless oil after evaporation. *R_f* = 0.48, *iso*-hexanes/ethyl acetate/methanol (6/4/1).



To a solution of **59e** (1.02 g, 3 mmol) in tetrahydrofuran (8 mL) was added methanol (4 mL), water (4 mL) and lithium hydroxide hydrate (0.15 g, 6.3 mmol). The resulting mixture was stirred at RT for 10 h and quenched with 1 M hydrochloric acid (6.5 mL, 6.5 mmol). The resulting solution was concentrated to a crude residue which was co-evaporated twice with tetrahydrofuran (20 mL), twice with anhydrous acetonitrile (20 mL) and twice with anhydrous toluene (20 mL). The resulting white solid was dried under high vacuum overnight and used without further purification (1.24 g, quantitative yield).




A solution of **1e** (0.53 g, 1 mmol) in dichloromethane (10 mL) was cooled in an ice water bath. Trimethylsilyl trifluoromethanesulfonate (0.69 g, 1.80 mmol) was added dropwise at 0°C under argon, and the resulting solution was stirred at RT for 30 min. The reaction mixture was evaporated to dryness and the resulting crude residue was dissolved in anhydrous acetonitrile (12 mL) under argon. The reaction mixture was stirred at 0°C, **59f** (371 mg, 0.9 mmol) and *N,N*-diisopropylethylamine (517 mg, 4 mmol) were added followed by 2-(1*H*-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyl uronium hexafluorophosphate methanaminium (570 mg, 1.5 mmol). The reaction mixture was stirred at RT for 48 h. The solvent was evaporated, the residue was dissolved in ethyl acetate (50 mL) and the solution was washed with 20% water solution of citric acid (2 x 50 mL), water (50 mL) and brine (50 mL), dried over magnesium sulfate, filtered and evaporated. The residue was purified by silica gel chromatography (gradient from 0-40% ethyl acetate and methanol (4/1) in *iso*-hexanes) to afford the title compound (560 mg, 84%) as a white solid after evaporation. *R*_f = 0.13. *iso*-hexanes/ethyl acetate/methanol (6/4/1).



To a solution of **59g** (371 mg, 0.5 mmol) in tetrahydrofuran (8 mL) was added methanol (4 mL), water (4 mL) and lithium hydroxide hydrate (36 mg, 1.5 mmol). The mixture was stirred for 2 h at ambient temperature and was quenched with aqueous 1 M hydrochloric acid (1.6 mL, 1.6 mmol). The resulting solution was concentrated to a crude residue which was co-evaporated twice with tetrahydrofuran (20 mL), twice with anhydrous acetonitrile (20 mL) and twice with anhydrous toluene (20 mL). The resulting white solid was dried under high vacuum overnight and it was used without further purification (365 mg, quantitative yield). Into oven dried, argon purged flask were placed 2-methyl-6-nitrobenzoic anhydride (258 mg, 0.75 mmol), 4-

dimethylaminopyridine (275 mg, 2.25 mmol) and anhydrous 1,2-dichloroethane (150 mL). The resulting solution was heated at 50°C, and the crude seco-acid was added dropwise via syringe as a solution in dry *N,N*-dimethylformamide (10 mL) over 12 h. An additional portion of dry *N,N*-dimethylformamide (2 x 5 mL) was used to complete the quantitative transfer. After stirring for
5 additional 2 h at 50°C, the reaction mixture was transferred to a separatory funnel and washed with water (100 mL, 5 mL of brine was added to support the separation). The aqueous phase was extracted with dichloromethane (50 mL). Combined organic extracts were washed with brine (50 mL) and dried over magnesium sulfate, filtered and concentrated under reduced pressure. The resulting residue was dissolved in ethyl acetate (100 mL) and was washed with
10 water (100 mL, 5 mL of brine was added to support the separation). The aqueous phase was extracted with ethyl acetate (50 mL). Combined organic extracts were washed with water (100 mL, 5 mL of brine was added to support the separation). Resulting aqueous phase was extracted with ethyl acetate (50 mL). Combined organic extracts were washed with brine (50 mL) and dried over magnesium sulfate, filtered and concentrated under reduced pressure. The
15 residue was purified by silica gel chromatography (gradient from 0-40% ethyl acetate and methanol (4/1) in *iso*-hexanes) to afford the title compound (72 mg, 24%) as a white solid after evaporation. $R_f = 0.42$, 10% methanol in dichloromethane. ^1H NMR (400 MHz, CD_3OD): δ 9.04 (s, 1H), 7.95 (d, $J = 8.5$ Hz, 1H), 7.76 (s, 1H), 7.47 (dd, $J = 8.5, 1.5$ Hz, 1H), 7.31 (s, 1H), 6.43 (m, 2H), 5.93 (q, $J = 6.6$ Hz, 1H), 5.44–5.36 (m, 1H), 4.33–4.23 (m, 2H), 3.84 (dt, $J = 11.6, 4.1$
20 Hz, 1H), 3.76–3.69 (m, 1H), 3.68–3.61 (m, 2H), 3.48 (m, 1H), 2.65–2.56 (m, 1H), 2.16–2.07 (m, 2H), 2.03–1.93 (m, 1H), 1.91–1.82 (m, 2H), 1.79e (m, 1H), 1.67–1.62 (m, 3H), 1.59 (d, $J = 6.6$ Hz, 3H), 1.51 (d, $J = 7.2$ Hz, 3H), 0.90 (d, $J = 6.7$ Hz, 3H), 0.85 (d, $J = 6.7$ Hz, 3H). LCMS (m/z) 592.3 $[\text{M}+\text{H}]^+$ Tr = 3.12 min.


25

EXAMPLE 60


A solution of Compound 59 (10 mg, 0.017 mmol) in a mixture of ethyl acetate (4 mL) and tetrahydrofuran (4 mL) containing 10% palladium on carbon (10 mg) was hydrogenated at RT

and at atmospheric pressure of hydrogen for 3 h. The reaction mixture was filtered through Celite and the filter pad was washed with tetrahydrofuran (10 mL). The filtrate was evaporated to afford the title compound (10 mg, quantitative yield) as a white solid. R_f = 0.47, 10% methanol in dichloromethane. ^1H NMR (400 MHz, CD_3OD): δ 9.02 (s, 1H), 7.94 (d, J = 8.5 Hz, 1H), 7.81 (1H), 7.44 (d, J = 8.5 Hz, 1H), 7.20 (s, 1H), 6.02 (q, J = 6.5 Hz, 1H), 5.78–5.71 (m, 1H), 5.24 (m, 1H), 4.33–4.23 (m, 2H), 3.83–3.61 (m, 6H), 3.52 (m, 1H), 2.71–2.66 (m, 1H), 2.43–2.36 (m, 2.14–2.05 (m, 2H), 2.08–1.91 (m, 1H), 1.90–1.82 (m, 2H), 1.76 (m, 1H), 1.67–1.62 (m, 2H), 1.57 (d, J = 6.6 Hz, 3H), 1.54 (d, J = 7.2 Hz, 3H), 0.96 (d, J = 6.7 Hz, 3H), 0.88 (d, J = 6.7 Hz, 3H). LCMS (m/z) 594.3 [$\text{M}+\text{H}$] $^+$ Tr = 2.72 min.


EXAMPLE 61




A solution of **1e** (1064 mg, 2 mmol) in dichloromethane (30 mL) was cooled in an ice water bath. Trimethylsilyl trifluoromethanesulfonate (666 mg, 3 mmol) was added dropwise at 0°C under argon, and the resulting solution was stirred at RT for 30 min. The reaction mixture was evaporated to dryness and the resulting crude residue was dissolved in anhydrous acetonitrile (25 mL) under argon. The reaction mixture was stirred at 0°C, hex-5-enoic acid (251 mg, 2.2 mmol) and *N,N*-diisopropylethylamine (1034 mg, 8 mmol) were added followed by 2-(1H-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyl uronium hexafluorophosphate methanaminium (1065 mg, 2.8 mmol). The reaction mixture was stirred at RT for 48 h. The solvent was evaporated, the residue was dissolved in ethyl acetate (200 mL) and the solution was washed with 20% water solution of citric acid (2 x 150 mL), water (150 mL) and brine (150 mL), dried over magnesium sulfate, filtered and evaporated. The residue was purified by silica gel chromatography (gradient from 0-40% ethyl acetate and methanol (4/1) in *iso*-hexanes) to afford the title compound (864 mg, 82%) as a white solid after evaporation. R_f = 0.35, *iso*-hexanes/ethyl acetate/methanol (6/4/1).



To a solution of **61a** (830 mg, 1.57 mmol) in tetrahydrofuran (40 mL) were added water (10 mL) and lithium hydroxide hydrate (57 mg, 2.38 mmol). The mixture was stirred for 2 h at ambient temperature and then filtered through a 5 cm layer of DOWEX D50 x 8 resin in H⁺ cycle (resin firstly washed with water). Resin was washed with additional water (50 mL). Filtrates were collected, concentrated under reduced pressure and co-evaporated twice with toluene (10 mL). After drying under high vacuum for one day, the title compound was isolated (590 mg, 95%) as a white solid. $R_f = 0.4$, 30% methanol in dichloromethane.



Into an oven dried, argon purged flask were added **61b** (238 mg, 0.60 mmol) and (R)-1-(3-vinyl-isoquinolin-6-yl)-ethanol (120 mg, 0.6 mmol). The flask was sealed and the reaction mixture was repurged twice with argon. Anhydrous dichloromethane (10 mL) was added and the reaction mixture was repurged twice with argon. *N*-(3-dimethylaminopropyl)-*N'*-ethylcarbodiimide hydrochloride (148 mg, 0.77 mmol) was added followed by 4-dimethylaminopyridine (67 mg, 0.55 mmol). Reaction mixture was quickly repurged twice with argon and was stirred at RT for 12 h. The reaction mixture was diluted with dichloromethane (100 mL) and the solution was washed with 20% water solution of citric acid (2 x 150 mL), water (150 mL) and brine (150 mL), dried over magnesium sulfate, filtered and evaporated. The residue was purified by silica gel chromatography (gradient from 0-40% ethyl acetate and methanol (4/1) in *iso*-hexanes) to afford the title compound (307 mg, 89%) as a white solid after evaporation. $R_f = 0.29$, *iso*-hexanes/ethyl acetate/methanol (6/4/1).




A solution of **61c** (209 mg, 0.362 mmol) in toluene (150 mL) was stirred at RT under argon. Hoveyda-Grubbs 2nd generation catalyst (23 mg, 0.036 mmol) was added and the reaction mixture was heated at reflux under argon for 30 min. Reaction mixture was cooled to

5 RT and ethyl acetate (50 mL) was added. This solution was washed twice with aqueous solution of tris(hydroxymethyl)phosphine (372 mg, 3 mmol in 100 mL of water), with water (2 x 50 mL) and with brine (50 mL). The organic phase was dried over magnesium sulfate, filtered and evaporated. The residue was purified by silica gel chromatography (gradient from 0-40% ethyl acetate and methanol (4/1) in *iso*-hexanes) to afford the title compound (125 mg, 63%) as a

10 white solid after evaporation. $R_f = 0.25$, *iso*-hexanes/ethyl acetate/methanol (6/4/1). ¹H NMR (400 MHz, CD₃OD): δ 9.00 (s, 1H), 7.91 (d, $J = 8.4$ Hz, 1H), 7.77 (s, 1H), 7.41 (d, $J = 8.4$ Hz, 1H), 7.23 (s, 1H), 6.64 (m, 1H), 6.29 (d, $J = 15.4$ Hz, 1H), 5.97 (q, $J = 5.9$ Hz, 1H), 5.53 (q, $J = 7.0$ Hz, 1H), 4.37–4.18 (m, 2H), 3.65 (d, $J = 9.8$ Hz, 1H), 2.65 (t, $J = 12.3$ Hz, 1H), 2.36–2.18 (m, 4H), 1.98 (m, 1H), 1.94–1.73 (m, 3H), 1.71–1.57 (m, 3H), 1.55 (d, $J = 5.6$ Hz, 6H), 0.83 (d, $J =$

15 6.6 Hz, 3H), 0.78 (d, $J = 6.6$ Hz, 3H). LCMS (m/z) 550.2 [M+H]⁺. Tr = 2.55 min.

EXAMPLE 62




A solution of Compound 61 (10 mg, 0.018 mmol) in a mixture of ethyl acetate (4 mL) and tetrahydrofuran (4 mL) containing 10% palladium on carbon (10 mg) was hydrogenated at RT


20 and at atmospheric pressure of hydrogen for 3 h. The reaction mixture was filtered through Celite and the filter pad was washed with tetrahydrofuran (10 mL). The filtrate was evaporated

to afford the title compound (9 mg, 89%) as a white solid. $R_f = 0.16$, *iso*-hexanes/ethyl acetate/methanol (6/4/1). ^1H NMR (400 MHz, CD_3OD) δ 9.02 (s, 1H), 7.93 (d, $J = 8.7$ Hz, 1H), 7.62 (s, 1H), 7.47 (s, 1H), 7.44 (d, $J = 8.7$ Hz, 1H), 6.02–5.93 (m, 1H), 5.39 (m, 1H), 4.03 (m, 1H), 3.69 (m, 1H), 2.84 (m, 2H), 2.27 (s, 1H), 2.10 (m, 1H), 1.97 (m, 1H), 1.85 (m, 2H), 1.76 (m, 2H), 1.62 (m, 4H), 1.54 (d, $J = 5.1$ Hz, 3H), 1.39 (d, $J = 6.6$ Hz, 3H), 1.19 (m, 4H), 0.78 (d, $J = 5.0$ Hz, 3H), 0.73 (d, $J = 5.8$ Hz, 3H). LCMS (m/z) 552.3 $[\text{M}+\text{H}]^+$ Tr = 2.10 min.


EXAMPLE 63




A solution of **1e** (1064 mg, 2 mmol) in dichloromethane (30 mL) was cooled in an ice water bath. Trimethylsilyl trifluoromethanesulfonate (666 mg, 3 mmol) was added dropwise at 0°C under argon, and the resulting solution was stirred at RT for 30 min. The reaction mixture was evaporated to dryness and the resulting crude residue was dissolved in anhydrous acetonitrile (25 mL) under argon. The reaction mixture was stirred at 0°C, hept-6-enoic acid (281 mg, 2.2 mmol) and *N,N*-diisopropylethylamine (1034 mg, 8 mmol) were added followed by 2-(1H-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyl uronium hexafluorophosphate methanaminium (1065 mg, 2.8 mmol). The reaction mixture was stirred at RT for 48 h. The solvent was evaporated, the residue was dissolved in ethyl acetate (200 mL) and the solution was washed twice with 20% water solution of citric acid (2 x 150 mL), water (150 mL) and brine (150 mL), dried over magnesium sulfate, filtered and evaporated. The residue was purified by silica gel chromatography (gradient from 0–40% ethyl acetate + methanol (4/1) in *iso*-hexanes) to afford the title compound (817 mg, 75%) as a white solid after evaporation. $R_f = 0.37$, *iso*-hexanes/ethyl acetate/methanol (6/4/1).



To a solution of **63a** (790 mg, 1.46 mmol) in tetrahydrofuran (40 mL) was added water (10 mL) and lithium hydroxide hydrate (52 mg, 2.19 mmol). The mixture was stirred for 2 h at ambient temperature and then filtered through a 5 cm layer of DOWEX D50 x 8 resin in H⁺ cycle (resin firstly washed with water). Resin was washed with additional water (50 mL). The filtrates were collected, concentrated under reduced pressure and co-evaporated twice with toluene (10 mL). After drying under high vacuum for one day, the title compound was isolated (583 mg, 97%) as a white solid. $R_f = 0.4$, 30% methanol in dichloromethane.



Into an oven dried, argon purged flask were added **63b** (276 mg, 0.67 mmol) and (R)-1-(3-vinyl-isoquinolin-6-yl)-ethanol (134 mg, 0.67 mmol). The flask was sealed and the reaction mixture was repurged twice with argon. Anhydrous dichloromethane (10 mL) was added and the reaction mixture was repurged twice with argon. *N*-(3-Dimethylaminopropyl)-*N'*-ethylcarbodiimide hydrochloride (168 mg, 0.87 mmol) was added followed by 4-dimethylaminopyridine (75 mg, 0.62 mmol). The reaction mixture was quickly repurged twice with argon and was stirred at RT for 12 h. The reaction mixture was diluted with dichloromethane (100 mL) and the solution was washed with 20% water solution of citric acid (2 x 150 mL), water (150 mL) and brine (150 mL), dried over magnesium sulfate, filtered and evaporated. The residue was purified by silica gel chromatography (gradient from 0-40% ethyl acetate and methanol (4/1) in *iso*-hexanes) to afford the title compound (316 mg, 79%) as a white solid after evaporation. $R_f = 0.30$, *iso*-hexanes/ethyl acetate/methanol (6/4/1).




A solution of **63c** (255 mg, 0.431 mmol) in toluene (200 mL) was stirred at RT under argon. Hoveyda-Grubbs 2nd generation catalyst (27 mg, 0.043 mmol) was added and the reaction mixture was heated at reflux under argon for 30 min. Reaction mixture was cooled to

5 RT and ethyl acetate (50 mL) was added. This solution was washed twice with aqueous solution of tris(hydroxymethyl)phosphine (372 mg, 3 mmol in 100 mL of water), twice with water (50 mL) and with brine (50 mL). The organic phase was dried over magnesium sulfate, filtered and evaporated. The residue was purified by silica gel chromatography (gradient from 0-40% ethyl acetate and methanol (4/1) in *iso*-hexanes) to afford the title compound (76 mg, 31%) as a

10 white solid after evaporation. $R_f = 0.20$, *iso*-hexanes/ethyl acetate/methanol (6/4/1). ¹H NMR (400 MHz, CD₃OD): δ 9.01 (s, 1H), 7.93 (d, $J = 8.5$ Hz, 1H), 7.69 (s, 1H), 7.57 (s, 1H), 7.43 (d, $J = 8.5$ Hz, 1H), 6.55 (m, 1H), 5.95 (m, 1H), 5.43 (m, 1H), 5.23 (m, 3H), 4.32 (m, 1H), 4.20 (m, 1H), 3.67 (m, 1H), 2.69 (m, 1H), 2.30 (m, 2H), 2.13 (m, 2H), 1.96 (m, 3H), 1.83 (m, 1H), 1.64 (m, 3H), 1.56 (d, $J = 6.7$ Hz, 3H), 1.44 (d, $J = 7.4$ Hz, 3H), 0.92–0.72 (m, 6H). LCMS (m/z) 564.4

15 [M+H]⁺ Tr = 2.60 min.

EXAMPLE 64




A solution of Compound 63 (15 mg, 0.027 mmol) in a mixture of ethyl acetate (4 mL) and tetrahydrofuran (4 mL) containing 10% palladium on carbon (10 mg) was hydrogenated at RT and at atmospheric pressure of hydrogen for 3 h. The reaction mixture was filtered through Celite and the filter pad was washed with tetrahydrofuran (10 mL). The filtrate was evaporated to afford the title compound (13 mg, 87%) as a white solid. $R_f = 0.11$, *iso*-hexanes/ethyl


20

acetate/methanol (6/4/1). ^1H NMR (400 MHz, CD_3OD) δ 9.02 (s, 1H), 7.95 (d, J = 8.7 Hz, 1H), 7.68 (s, 1H), 7.50 (s, 1H), 7.45 (d, J = 8.7 Hz, 1H), 5.93 (q, J = 6.3 Hz, 1H), 5.20 (m, 1H), 4.21 (m, 1H), 4.07 (d, J = 8.0 Hz, 1H), 3.66 (m, 1H), 2.83 (m, 4H), 2.25 (m, 1H), 2.07 (m, 1H), 1.94 (m, 2H), 1.80e (m, 1H), 1.63 (m, 3H), 1.55 (d, J = 6.6 Hz, 3H), 1.51–1.42 (m, 2H), 1.37 (d, J = 7.0 Hz, 3H), 1.25 (m, 4H), 0.83 (d, J = 6.7 Hz, 6H). LCMS (m/z) 566.3 $[\text{M}+\text{H}]$, Tr = 2.27 min.

EXAMPLE 65




To a solution of (R)-1-(3-chloro-isoquinolin-6-yl)-ethanol (250 mg, 1.21 mmol), and potassium phosphate tribasic (770 mg, 3.63 mmol) in cyclopentyl methy ether (4.5 mL) and water (1.5 mL) preheated to 90 °C under an argon atmosphere were added 3-(2-methoxy-2-oxoethyl)phenylboronic acid, pinacol ester (Combi-Blocks, 387 mg, 1.33 mmol) and (A- C_6Phos) $_2\text{PdCl}_2$ (49 mg, 60 μmol). After 17 h, the reaction was allowed to cool to 23 °C, and was partitioned between dichloromethane (50 mL) and saturated aqueous sodium bicarbonate solution (50 mL). The phases were split and the aqueous layer was extracted with dichloromethane (50 mL). The combined organic layers were dried over anhydrous sodium sulfate, and were concentrated under reduced pressure. The crude residue was purified by silica gel chromatography (24 g Combiflash HP Gold Column, 0–100% ethyl acetate/isohexanes gradient) to afford the title compound (321 mg, 83%) as a faint yellow oil.




To a solution of **65a** (320 mg, 1.00 mmol) in tetrahydrofuran (3 mL) and water (2 mL) was added lithium hydroxide (26 mg, 1.1 mmol) at 23 °C under an argon atmosphere. After 3 h, the reaction mixture was concentrated under reduced pressure and the crude residue was purified by silica gel chromatography (24 g Combiflash HP Gold Column, 0–20%


methanol/dichloromethane gradient) to afford the title compound (64.2 mg, 21%) as a colorless oil. $R_f = 0.5$ (20% methanol in dichloromethane) I_2 /silica stain.



To a solution of **1e** (104 mg, 0.21 mmol) in dichloromethane (1.05 mL) was added trimethylsilyl trifluoromethanesulfonate (70 mg, 1.07 mmol) at 0°C under an argon atmosphere. After 1 h, the reaction mixture was concentrated under reduced pressure. The resulting residue was diluted with acetonitrile (1.05 mL) and 2-(1H-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate methanaminium (79.8 mg, 0.210 mmol), *N,N*-diisopropylethylamine (140 μ L, 0.840 mmol), and **65b** (64 mg, 0.21 mmol) were sequentially added at 23°C under an argon atmosphere. After 18 h, the reaction mixture was concentrated under reduced pressure, and the crude residue was purified by silica gel chromatography to afford the title compound (140 mg, 93%) as a faint yellow oil.




To a solution of **65c** (140 mg, 0.195 mmol) in tetrahydrofuran (0.9 mL) and water (0.3 mL) was added lithium hydroxide hydrate (4.6 mg, 0.195 mmol) at 23°C under an argon atmosphere. After 3 h, the reaction mixture was concentrated under reduced pressure to afford the title compound (131 mg, 99%) as a white solid lithium carboxylate salt.




To a solution of benzotriazol-1-yl-oxytripyrrolidinophosphonium hexafluorophosphate (35.3 mg, 68 μ mol) and 4-dimethylaminopyridine (62.3 mg, 510 μ mol) in dichloromethane (5.7 mL) was added **65d** (10 mg, 17 μ mol) at 23 °C under an argon atmosphere. After 16 h, the reaction mixture was concentrated under reduced pressure and the crude residue was purified by preparative HPLC (Gemini 5u C18 110Å column, 5–100% acetonitrile/water, 0.1% trifluoroacetic acid modifier) to afford the title compound (6.0 mg, 52%) as a white solid trifluoroacetic acid salt. ^1H NMR (400 MHz, CD_3OD) δ 9.70 (s, 1H), 8.58 (s, 1H), 8.48 (d, J = 8.6 Hz, 1H), 8.32 (s, 1H), 7.97–7.87 (m, 2H), 7.79 (d, J = 7.9 Hz, 1H), 7.62 (*app* t, J = 7.7 Hz, 1H), 7.53 (d, J = 7.8 Hz, 1H), 6.21 (q, J = 6.6 Hz, 1H), 5.84–5.71 (m, 1H), 4.45–4.29 (m, 2H), 3.96 (d, J = 15.5 Hz, 1H), 3.75 (dd, J = 11.1, 2.7 Hz, 1H), 3.64 (d, J = 15.5 Hz, 1H), 2.74 (td, J = 12.8, 3.1 Hz, 1H), 2.13–1.96 (m, 2H), 1.91 (*br* d, J = 13.0 Hz, 1H), 1.81–1.65 (m, 2H), 1.67 (d, J = 6.6 Hz, 3H), 1.62 (d, J = 7.1 Hz, 3H), 1.01 (d, J = 6.8 Hz, 3H), 0.94 (d, J = 6.8 Hz, 3H). HPLC Tr = 3.040 min. LCMS (m/z) 572.3 [$M+H$], Tr = 2.07 min.

15

EXAMPLE 66

To a solution of 6-chloropyridin-3-amine (5.00 g, 38.8 mmol) in dioxane (194 mL) was added di-*tert*-butyl dicarbonate (10.2 g, 46.7 mmol) at 23°C under an argon atmosphere, and the resulting mixture was heated to 100°C. After 17 h, the reaction mixture was allowed to cool to 23°C, and was diluted with water (500 mL). The resulting mixture was extracted with ethyl acetate (2 $\times$ 500 mL), and the combined organic extracts were dried over anhydrous sodium sulfate and were concentrated under reduced pressure. The crude residue was purified by silica gel chromatography (120 g Combiflash HP Gold Column, 0–100% ethyl acetate/*iso*-hexanes gradient) to afford the title compound (7.69 g, 87%) as a colorless oil.


20




25

To a solution of **66a** (2.00 g, 8.80 mmol) and tetramethylethylenediamine (2.70 mL, 18.0 mmol) in diethyl ether (44 mL) was added *n*-butyllithium (2.5 M in hexanes, 7.2 mL, 18.0 mmol) at –78°C under an argon atmosphere. After 10 min, the resulting mixture was allowed to warm


to -15°C over a 50 min period. The reaction mixture was cooled to -78°C, and *N,N*-dimethylformamide (1.9 g, 26 mmol) was added via syringe. After 30 min, the reaction mixture was quenched with saturated aqueous ammonium chloride solution (20 mL) and was allowed to warm to 23°C. The resulting mixture was extracted with ethyl acetate (2 × 20 mL), and the combined organic layers were dried over anhydrous sodium sulfate and were concentrated under reduced pressure. The crude residue was purified by silica gel chromatography (120 g Combiflash HP Gold Column, 0–100% ethyl acetate/*iso*-hexanes gradient) to afford the title compound (849 mg, 38%) as colorless oil.



- 10 To a solution of **66b** (258 mg, 1.00 mmol) and (E)-4-(dimethylamino)but-3-en-2-one (452 mg, 4.00 mmol) in 1,4-dioxane (10 mL) was added *p*-toluenesulfonic acid monohydrate (761 mg, 4 mmol) at 23°C under an argon atmosphere and the resulting mixture was heated to 80°C. After 2 h, the reaction mixture was allowed to cool to 23°C and was partitioned between saturated aqueous sodium bicarbonate solution (200 mL) and ethyl acetate (200 mL). The layers were split and the aqueous layer was extracted with ethyl acetate (200 mL). The combined organic layers were dried over anhydrous sodium sulfate and were concentrated under reduced pressure. The crude residue was purified by silica gel chromatography to afford the title compound (107 mg, 51%) as an off-white solid.




- 20 Dichloro (*p*-cymene) ruthenium(II) dimer (5 mg, 8 μmol) and (1*R*,2*R*)-(-)-*N*-*p*-tosyl-1,2-diphenylethylenediamine (7 mg, 19 μmol) were suspended in degassed water (6 mL) and the mixture was degassed with argon for 15 min. The mixture was stirred at 70°C under argon for 90 min. The resulting yellow solution was cooled to RT. **66c** (329 mg, 1.56 mmol), sodium formate (543 mg, 7.98 mmol) and degassed tetrahydrofuran (1 mL) were added and the reaction mixture was degassed for 10 min. The reaction mixture was vigorously stirred at 40°C for 2.5 h. The reaction mixture was cooled to RT and was extracted with ethyl acetate (20 mL). The organic layer was separated, washed with water (20 mL), brine (20 mL), dried over anhydrous magnesium sulfate, and concentrated under reduced pressure. The residue was purified by silica gel chromatography to afford the title compound (180 mg, 54%) as a solid.




To a suspension of potassium phosphate tribasic (550 mg, 2.59 mmol) in cyclopentyl methyl ether (4.5 mL) and water (3 mL) was added **66d** (180 mg, 0.89 mmol) and heated to 90°C. At this temperature, (A-^{ca}Phos)₂PdCl₂ (35 mg, 43 μmol) was added and stirred for 2 min.

- 5 A solution of (E)-2,2-dimethyl-4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-but-3-enoic acid methyl ester (286 mg, 1.12 mmol) in cyclopentyl methyl ether (4.5 mL) was added dropwise and stirred for 19 h at 90°C. The reaction mixture was cooled to RT, diluted with ethyl acetate (40 mL) and water (20 mL). The layers were separated and the aqueous layer was extracted with ethyl acetate (2 × 20 mL). The combined organic layers were dried over anhydrous magnesium sulfate, concentrated and the resulting crude residue was purified via silica gel chromatography
- 10 (24 g SiO₂ Isco Rf Gold Column, 0–100% ethyl acetate/iso-hexanes gradient) to afford the title compound (95 mg, 37%) as a pale brown solid.




- To a solution of **66e** (95 mg, 0.32 mmol) in tetrahydrofuran (1.8 mL), methanol (0.6 mL)
- 15 and water (0.6 mL) was added lithium hydroxide hydrate (15 mg, 0.63 mmol) at 23°C. After 16 h, the resulting mixture was concentrated under reduced pressure and residual solvents were removed azeotropically by addition of toluene (5 mL) followed by concentration under reduced pressure (2 ×) to afford the title compound. This was used in the subsequent amide coupling without further purification.




To a solution of **1e** (212 mg, 0.40 mmol) in dichloromethane (3 mL) was slowly added trimethylsilyl trifluoromethanesulfonate (106 μL, 0.60 mmol) at 0°C under an argon atmosphere.

After 1 h, the resulting mixture was concentrated under reduced pressure and was used in the subsequent amide coupling without further purification. To a solution of **66f** (90.5 mg, 0.32 mmol) in acetonitrile (3 mL), 2-(1H-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyl uronium hexafluorophosphate methanaminium (180 mg, 0.47 mmol), *N,N*-diisopropylethylamine (330 μ L, 1.90 mmol) and (S)-1-[(S)-2-((S)-2-amino-3-methyl-butyrylamino)-propionyl]-hexahydro-pyridazine-3-carboxylic acid 2,2,2-trichloro-ethyl ester (232 mg, 0.4 mmol) in acetonitrile (2 mL) were added sequentially at 23°C. After 16 h, the resulting mixture was concentrated under reduced pressure and the residue was purified directly by silica gel chromatography to afford the title compound (201 mg, 91%) as a solid.



To a solution of **66g** (201 mg, 0.29 mmol) in tetrahydrofuran (2 mL), methanol (0.4 mL) and water (0.4 mL) was added lithium hydroxide hydrate (14 mg, 0.57 mmol) at 23°C. After 40 min, lithium hydroxide hydrate (14 mg, 0.57 mmol) was added at 23°C. After 1 h, the resulting mixture was concentrated under reduced pressure and residual solvents were removed azeotropically by addition of toluene (2 mL) followed by concentration under reduced pressure (3 $\times$) to afford the title compound. This was used in the subsequent macrolactonization without further purification.




To 2-methyl-6-nitrobenzoic anhydride (43 mg, 0.12 mmol) and 4-dimethylaminopyridine (48 mg, 0.39 mmol) was added 1,2-dichloroethane (20 mL) under nitrogen atmosphere and heated to 50°C. At this temperature, **66h** (28 mg, 0.049 mmol) in *N,N*-dimethylformamide (1 mL) was added dropwise via syringe pump over 6 h. An additional wash of *N,N*-dimethylformamide (0.5 mL) was then added in the same manner over 15 min. After stirring an additional 1.25 h, reaction mixture was cool to RT. It was diluted with ethyl acetate (20 mL) and washed with


20

water (2 x 10 mL). The organic layer was dried over anhydrous magnesium sulfate and filtered. The solvent was removed under reduced pressure and the residue was purified by preparative HPLC (Gemini 5u C18 110Å column, 5–100% acetonitrile/water, 0.1% trifluoroacetic acid modifier) to afford the title compound as a white powder trifluoroacetic acid salt. It was washed with saturated solution of sodium bicarbonate to remove acid impurities formed from 2-methyl-6-nitrobenzoic anhydride to afford the title compound (1.5 mg, 5%) as a white solid. ¹H NMR (400 MHz, CD₃OD) δ 9.19 (s, 1H), 8.86 (d, *J* = 2.1 Hz, 1H), 8.67 (d, *J* = 7.5 Hz, 1H), 8.21 (s, 1H), 7.59 (s, 1H), 6.89 (d, *J* = 9.5 Hz, 1H), 6.58 (d, *J* = 16.1 Hz, 1H), 6.46 (d, *J* = 16.1 Hz, 1H), 6.04 (q, *J* = 6.4 Hz, 1H), 5.52–5.41 (m, 1H), 4.26 (d, *J* = 14.4 Hz, 1H), 4.18 (*app t*, *J* = 9.1 Hz, 1H), 3.65 (dd, *J* = 11.4, 2.8 Hz, 1H), 2.58 (td, *J* = 12.9, 3.2 Hz, 1H), 1.91–1.75 (m, 3H), 1.68–1.53 (m, 2H), 1.63 (d, *J* = 6.7 Hz, 3H), 1.51 (d, *J* = 7.3 Hz, 3H), 1.43 (s, 3H), 1.26 (s, 3H), 0.87 (d, *J* = 6.7 Hz, 3H), 0.82 (d, *J* = 6.7 Hz, 3H). LCMS (*m/z*) 551.2 [M+H], Tr = 2.19 min.


EXAMPLE 67




Methyl 2-(3-bromophenyl)-2-methylpropanoate (Pharmabridge, Doylestown, PA, USA (190 mg, 0.74 mmol), 4,4',4'',4''',5,5',5'',5'''-octamethyl-2,2'-bi(1,3,2-dioxaborolane) (206 mg, 0.81 mmol), [1,1'-bis(diphenylphosphino)ferrocene] dichloropalladium(II), complex with dichloromethane (27 mg, 0.04 mmol) and potassium acetate (217 mg, 2.22 mmol) were placed in a screw cap vial and flushed with a vacuum and argon cycle three times. Anhydrous 1,4-dioxane (4 mL) was added under argon and the resulting mixture was heated to 80°C for 19 h. The reaction mixture was cool to RT and was diluted with ethyl acetate (20 mL). Celite (~1 g) was added and filtered through a pad of celite. Solvents were removed under reduced pressure to afford the title compound which was used directly in the subsequent reaction.



To a suspension of potassium phosphate tribasic (445 mg, 2.10 mmol) in cyclopentyl methyl ether (2.5 mL) and water (1.5 mL) was added (R)-1-(3-chloro-isoquinolin-6-yl)-ethanol (180 mg, 0.89 mmol) and the reaction mixture was heated to 90 °C. At this temperature, (A-^{ca}Phos)₂PdCl₂ (28 mg, 35 μmol) was added and the reaction mixture was stirred for 2 min. A solution of **67a** (225 mg, 0.74 mmol) in cyclopentyl methyl ether (2.5 mL) was added dropwise and stirred for 19 h at 90°C. The reaction mixture was cool to RT, diluted with ethyl acetate (30 mL) and water (15 mL). The layers were separated and the aqueous layer was extracted with ethyl acetate (2× 15 mL). The combined organic layers were dried over anhydrous magnesium sulfate, concentrated and the resulting crude residue was purified by silica gel column chromatography to afford the title compound (267 mg, quantitative) as a pale brown solid.




To a solution of **67b** (244 mg, 0.7 mmol) in tetrahydrofuran (10 mL), and water (2.5 mL) was added lithium hydroxide hydrate (18.5 mg, 0.77 mmol) at 23°C. After 48 h, lithium hydroxide hydrate (17 mg, 0.70 mmol) was added to the reaction mixture. After 24 h, the reaction mixture was concentrated under reduced pressure and residual solvents were removed azeotropically by addition of toluene (5 mL) followed by concentration under reduced pressure (2×) to afford the title compound. This was used in the subsequent amide coupling without further purification.




To a solution of **1e** (397 mg, 0.75 mmol) in dichloromethane (5 mL) was slowly added trimethylsilyl trifluoromethanesulfonate (199 μL, 1.12 mmol) at 0°C under an argon atmosphere. After 1 h, the resulting mixture was concentrated under reduced pressure to give (S)-1-[(S)-2-((S)-2-amino-3-methyl-butrylamino)-propionyl]-hexahydro-pyridazine-3-carboxylic acid 2,2,2-trichloro-ethyl ester as a triflate salt. This was used in the subsequent amide coupling without further purification. To a solution of **67c** (234 mg, 0.7 mmol) in acetonitrile (7 mL), 2-(1H-7-

azabenzotriazol-1-yl)-1,1,3,3-tetramethyl uronium hexafluorophosphate methanaminium (532 mg, 1.4 mmol), *N,N*-diisopropylethylamine (730 μ L, 4.2 mmol) and (S)-1-[(S)-2-((S)-2-amino-3-methyl-butylamino)-propionyl]-hexahydro-pyridazine-3-carboxylic acid 2,2,2-trichloro-ethyl ester (435 mg, 0.75 mmol) in acetonitrile (3 mL) were added sequentially at 23°C. After 16 h, the resulting mixture was concentrated under reduced pressure and the residue was purified directly by silica gel chromatography (40 g SiO₂ Isco Rf Gold Column, 0–100% ethyl acetate/*iso*-hexanes gradient) to afford the title compound (230 mg, 44%) as a solid.




To a solution of **67d** (94 mg, 0.13 mmol) in tetrahydrofuran (3 mL), and water (1 mL) was added lithium hydroxide hydrate (3.1 mg, 0.13 mmol) at 23°C. After 48 h, lithium hydroxide hydrate (17 mg, 0.70 mmol) was added to the reaction mixture. After 1 h, the reaction mixture was concentrated under reduced pressure and residual solvents were removed azeotropically by addition of toluene (5 mL) followed by concentration under reduced pressure (2 $\times$) to afford the title compound. This was used in the subsequent macrolactonization without further purification.




To a suspension of **67e** (78 mg, 0.13 mmol) in dichloromethane (42 mL) was added benzotriazol-1-yl-oxytripyrrolidinophosphonium hexafluorophosphate (520 mg, 0.50 mmol) and 4-dimethylaminopyridine (462 mg, 4.0 mmol) at 23°C. After 24 h, the solvent was removed under reduced pressure and the residue was purified by preparative HPLC (Gemini 5u C18 110Å column, 5–100% acetonitrile/water, 0.1% trifluoroacetic acid modifier) to afford the title compound (3.2 mg, 4%) as a white powder and as a trifluoroacetic acid salt. ¹H NMR (400 MHz, CD₃OD) δ 9.65 (s, 1H), 8.69 (s, 1H), 8.49 (d, *J* = 8.7 Hz, 1H), 8.44 (d, *J* = 8.6 Hz, 1H), 8.34 (s, 1H), 7.90 (dd, *J* = 8.7, 1.5 Hz, 1H), 7.83 (d, *J* = 7.7 Hz, 1H), 7.78 (d, *J* = 7.9 Hz, 1H), 7.74 (s,

1H), 7.69 (*app t*, $J = 7.8$ Hz, 1H), 6.82 (d, $J = 8.6$ Hz, 1H), 6.21 (q, $J = 6.5$ Hz, 1H), 5.92–5.84 (1H), 4.41–4.30 (m, 2H), 3.75 (dd, $J = 14.4, 6.0$ Hz, 1H), 2.71 (td, $J = 13.0, 3.0$ Hz, 1H), 2.06–1.99 (m, 1H), 1.96 (dd, $J = 14.2, 7.1$ Hz, 1H), 1.93–1.86 (m, 1H), 1.75 (s, 3H), 1.73–1.59 (m, 2H), 1.68 (d, $J = 6.6$ Hz, 3H), 1.63 (d, $J = 7.1$ Hz, 3H), 1.47 (s, 3H), 0.96 (d, $J = 6.8$ Hz, 3H),
 5 0.84 (d, $J = 6.7$ Hz, 3H). HPLC Tr = 5.319 min. LCMS (m/z) 600.5 [M+H], Tr = 2.71 min.


EXAMPLES 68 AND 69



To a solution of 6-bromo-3-chloroisoquinoline (Frontier Scientific, 1.594 g, 6.57 mmol) and 2,2-dimethylpropane-1,3-diol (684 mg, 6.57 mmol) in cyclopentyl methyl ether (20 mL) was added cesium carbonate (2.354 g, 7.23 mmol) at 23°C. The reaction mixture was heated to 120°C for 18 h. The reaction mixture was cool to RT, diluted with ethyl acetate (50 mL) and washed with water (30 mL), brine (30 mL) and the resulting organic layer was dried over anhydrous magnesium sulfate and concentrated under reduced pressure. The residue was
 15 purified by silica gel chromatography (80 g Isco Rf Gold Column, 0–100% ethyl acetate/*iso*-hexanes gradient) to afford the title compound (461 mg, 23%) as a white solid.




To a solution of **68a** (461 mg, 1.49 mmol) in acetonitrile (10 mL) and water (2.5 mL) was added periodic acid (1.698 g, 7.45 mmol) at 23°C. The reaction mixture was cool to 0°C and chromium trioxide (30 mg, 0.298 mmol) was added in one portion. After 2.5 hour, the reaction mixture was diluted with ethyl acetate (30 mL) and water (30 mL) and the layers were separated. The aqueous layer was extracted with ethyl acetate (30 mL) and combined organic layers were washed with brine, dried over anhydrous magnesium sulfate and concentrated under reduced pressure to afford the title compound which was used without purification (481
 20 mg).




To a solution of **68b** (481 mg, 1.49 mmol) in acetonitrile (10 mL), 2-(1H-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyl uronium hexafluorophosphate methanaminium (1.133

g, 2.98 mmol), *N,N*-diisopropylethylamine (1.55 mL, 8.94 mmol) and (S)-methyl 2-amino-3-methylbutanoate hydrochloride (749 mg, 4.47 mmol) were added sequentially at 23°C. After 16 h, the resulting mixture was concentrated under reduced pressure and the residue was purified directly by silica gel chromatography (24 g Isco Rf Gold Column, 0–100% ethyl acetate/*iso*-hexanes gradient) to afford the title compound (412 mg, 63% over 2 steps) as a brown oil.




To a suspension of **68c** (412 mg, 0.95 mmol) and bis(triphenylphosphine) palladium (II) dichloride in toluene (5 mL) was added tributyl(1-ethoxyvinyl)tin (962 μ L, 2.85 mmol) under argon at 23°C. The reaction mixture was heated to 50°C. After 18 h, water (1 mL) was added at 50°C. After 2 h, the reaction mixture was cool to RT, diluted with ethyl acetate (15 mL) and 1 M solution of potassium fluoride (5 mL) was added. The resulting mixture was vigorously stirred at 23°C. After 2 h, the mixture was filtered through a short pad of Celite and washed with ethyl acetate (10 mL). The filtrate was washed with water (15 mL), brine (15 mL), dried over anhydrous magnesium sulfate and concentrated. This residue was dissolved in 1,4-dioxane (8 mL) and 1 M aqueous hydrochloric acid (1 mL) was added at 23°C. After 5 min, the reaction mixture was quenched with a saturated solution of sodium bicarbonate (2 mL) and concentrated to dryness. The residue was taken up in ethyl acetate (20 mL) and water (20 mL) and layers were separated. The aqueous layer was extracted with ethyl acetate (2 $\times$ 10 mL) and the combined organic layers were washed with brine (20 mL), dried over anhydrous magnesium sulfate and concentrated. The residue was purified by silica gel chromatography to afford the title compound (236 mg, 62%) as a brown gum.




Dichloro (*p*-cymene) ruthenium(II) dimer (2 mg, 3 μ mol) and (1R,2R)-(-)-*N*-*p*-tosyl-1,2-diphenylethylenediamine (2.6 mg, 7 μ mol) were suspended in degassed water (2.5 mL) and the mixture was degassed with argon for 15 min. The mixture was stirred at 70°C under nitrogen for 90 min. The resulting yellow solution was cooled to RT. **68d** (236 mg, 0.59 mmol), sodium formate (200 mg, 2.95 mmol) and degassed tetrahydrofuran (1.25 mL) were added and the reaction mixture was degassed for 10 min. The reaction mixture was vigorously stirred at 40°C for 3.5 h. The reaction mixture was cooled to RT and was extracted with ethyl acetate (15 mL).

The organic layer was separated, washed with water (20 mL), brine (20 mL), dried over anhydrous magnesium sulfate, and concentrated under reduced pressure. The residue was purified by silica gel chromatography to afford the title compound (196 mg, 83%) as a pale brown solid.



To a solution of **68e** (196 mg, 0.487 mmol) in tetrahydrofuran (3 mL), methanol (1 mL) and water (1 mL) was added lithium hydroxide hydrate (23 mg, 0.97 mmol) at 23°C. After 16 h, the reaction mixture was concentrated under reduced pressure. The residue was taken up in ethyl acetate (10 mL) and water (10 mL) and acidified with 1 M aqueous hydrochloric acid solution to pH ~2. The resulting layers were separated. The aqueous layer was extracted with ethyl acetate (2 × 10 mL) and the combined organic layers were washed with brine, dried over anhydrous magnesium sulfate and concentrated under reduced pressure to afford the title compound which was used without purification.

10




To a solution of (S)-1-((S)-2-*tert*-butoxycarbonylamino-propionyl)-hexahydro-pyridazine-3-carboxylic acid 2,2,2-trichloro-ethyl ester (270 mg, 0.62 mmol) in dichloromethane (5 mL) was slowly added trimethylsilyl trifluoromethanesulfonate (165 μ L, 0.93 mmol) at 0°C under an argon atmosphere. After 1 hour, the reaction mixture was concentrated under reduced pressure to afford the triflate salt of (S/R)-1-((S)-2-amino-propionyl)-hexahydro-pyridazine-3-carboxylic acid 2,2,2-trichloro-ethyl ester as yellow oil which was used it without purification. To a solution of **68f** (189 mg, 0.49 mmol) in acetonitrile (7 mL), 2-(1H-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyl uronium hexafluorophosphate methanaminium (278 mg, 0.73 mmol), *N,N*-diisopropylethylamine (730 μ L, 4.2 mmol) and (S/R)-1-((S)-2-amino-propionyl)-hexahydro-pyridazine-3-carboxylic acid 2,2,2-trichloro-ethyl ester (435 mg, 0.62 mmol) in acetonitrile (3 mL) were added sequentially at 23°C. After 16 h, the resulting mixture was concentrated under reduced pressure and the

20

25

residue was purified directly by silica gel chromatography (40 g SiO₂ Isco Rf Gold Column, 0–100% ethyl acetate/*iso*-hexanes gradient) to afford an inseparable diastereomeric mixture of the title compounds (43 mg, 13%) as a colorless residue.




5

Compound 68i Compound 68j

To a solution of a mixture of **68g** and **68h** (43 mg, 61 μ mol) in tetrahydrofuran (4 mL) and water (2 mL) was added lithium hydroxide hydrate (3.2 mg, 0.13 mmol) at 23°C. After 1 h, the reaction mixture was concentrated under reduced pressure and residual solvents were removed azeotropically by addition of toluene (5 mL) followed by concentration under reduced pressure (2 $\times$) to afford the title compounds as a mixture. This was used in the subsequent macrolactonization without further purification. LCMS (*m/z*) 572.2 [*M*+*H*], *Tr* = 2.21 min.

10



Compound 68

Compound 69

To a suspension of **68i** and **68j** (34.8 mg, 61 μ mol) in dichloromethane (20 mL) was added benzotriazol-1-yl-oxytripyrrolidinophosphonium hexafluorophosphate (127 mg, 0.24 mmol) and 4-dimethylaminopyridine (224 mg, 1.83 mmol) at 23°C. After 24 h, the solvent was removed under reduced pressure and the residue was purified by preparative HPLC (Gemini 5u C18 110Å column, 5–100% acetonitrile/water, 0.1% trifluoroacetic acid modifier) to afford both compounds (12 mg, 35%) as a ratio mixture of diastereomers. The diastereomers were separated using Chiral preparative HPLC to afford first eluting (*Tr* = 3.73 min) Compound 68 (2.17 mg, 6%) and the second eluting (*Tr* = 5.713 min)

15


20

Compound 69 (1.85 mg, 5%). Compound 68: ¹H NMR (400 MHz, CD₃OD) δ 8.89 (s, 7.94 (d, *J* = 8.6 Hz, 1H), 7.86 (s, 1H), 7.37 (dd, *J* = 8.6, 1.5 Hz, 1H), 7.28 (d, *J* = 9.2 Hz, 1H),


6.93 (s, 1H), 6.09 (q, J = 6.5 Hz, 1H), 5.54 (q, J = 7.2 Hz, 1H), 4.36–4.26 (m, 3H), 4.21 (d, J = 9.7 Hz, 1H), 3.69 (dd, J = 11.2, 2.6 Hz, 1H), 2.77 (td, J = 12.9, 3.0 Hz, 1H), 2.00–1.84 (m, 2H), 1.83–1.77 (m, 1H), 1.76–1.62 (m, 2H), 1.67 (d, J = 6.7 Hz, 3H), 1.47 (d, J = 7.3 Hz, 3H), 1.38 (s, 3H), 1.27 (s, 3H), 0.99 (d, J = 6.7 Hz, 3H), 0.95 (d, J = 6.7 Hz, 3H). HPLC Tr = 5.866 min. LCMS (m/z) 554.2 [M+H], Tr = 2.44 min.

Compound 69: ^1H NMR (400 MHz, CD_3OD) δ 8.78 (s, 1H), 7.83 (d, J = 8.6 Hz, 1H), 7.74 (s, 1H), 7.24 (dd, J = 8.6, 1.6 Hz, 1H), 7.03 (s, 1H), 5.93 (q, J = 6.7 Hz, 1H), 5.35 (q, J = 6.8 Hz, 1H), 4.31 (d, J = 9.8 Hz, 1H), 4.18 (d, J = 13.3 Hz, 1H), 4.02 – 3.97 (m, 2H), (d, J = 10.0 Hz, 1H), 3.65 (dd, J = 10.8, 2.8 Hz, 1H), 2.71 (td, J = 12.7, 2.8 Hz, 1H), 1.98 – 1.89 (m, 1H), 1.87 – 1.77 (m, 2H), 1.73 – 1.64 (m, 1H), 1.57 (d, J = 6.7 Hz, 3H), 1.49 (d, J = 7.0 Hz, 3H), 1.40 (s, 3H), 1.13 (s, 3H), 0.89 (d, J = 6.7 Hz, 3H), 0.85 (d, J = 6.5 Hz, 3H). HPLC Tr = 5.951 min. LCMS (m/z) 555.2 [M+H], Tr = 2.49 min.

EXAMPLE 70




To 2,7-dibromonaphthalene (1 g, 3.50 mmol) in anhydrous tetrahydrofuran (18 mL), at -78°C and under an atmosphere of nitrogen, was added a solution of *n*-butyllithium (2.5 M in hexanes, 1.5 mL, 3.67 mmol) dropwise. The reaction was stirred at -78°C for 20 min after which *N*-methoxy-*N*-methylacetamide (409 μL , 3.85 mmol) was added. After 15 min, the reaction was warmed to RT and stirred for 30 min. The reaction was quenched with 2 M hydrochloric acid and extracted twice with dichloromethane. The combined organic layers were dried through a hydrophobic frit and concentrated *in vacuo*. The product was purified by silica flash chromatography (*iso*-hexanes/ethyl acetate, 7/1) to afford the title compound (650 mg, 75%) as a colorless solid.




To dichloro (*p*-cymene) ruthenium (II) dimer (8 mg, 0.013 mmol) in water (5 mL) at RT was added (1R, 2R)-(-)-*N*-*p*-tosyl-1,2-diphenylethylenediamine (11.5 mg, 0.031 mmol). The system was degassed for 15 min and then heated to 70°C for 1.5 h. The reaction was cooled


and was added a solution of **70a** (650 mg, 2.61 mmol) in degassed anhydrous tetrahydrofuran (2 mL) followed by sodium formate (874 mg, 13.1 mmol). The reaction was heated at 40°C for 3 h, cooled to RT and extracted twice with dichloromethane. The combined organic layers were dried through a hydrophobic frit and concentrated *in vacuo*. The product was purified by silica gel chromatography (*iso*-hexanes/ethyl acetate, 4/1) to afford the title compound (450 mg, 69%) as a colorless solid.




70b (42 mg, 0.17 mmol) was dissolved in acetonitrile (2 mL) in a microwave vial, to the mixture was added 3-butenic acid (35 mg, 0.41 mmol), palladium (II) acetate (4 mg, 0.017 mmol), tri-(*o*-tolyl)phosphine (10 mg, 0.034 mmol) and triethylamine (0.12 mL). The vial was heated at 120° in the microwave reactor for 15 mins. The reaction mixture was then filtered, the solvent was evaporated and purified with combi-flash column chromatography (Eluent methanol/dichloromethane 1:3) to afford the title compound (35 mg, 81%) as a yellow solid.



70c (40 mg, 0.08 mmol) was dissolved in *N,N*-dimethylformamide (2 mL), to the solution was added 2-(1H-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyl uronium hexafluorophosphate methanaminium (36 mg, 0.1 mmol), the reaction mixture was stirred at RT for 10 mins. Then (S)-1-[(S)-2-((S)-2-amino-3-methylbutyrylamino)-propionyl]-hexahydro-pyridazine-3-carboxylic acid 2,2,2-trichloro-ethyl ester (0.18 mmol) in *N,N*-dimethylformamide (1 mL) was added to above reaction mixture followed by triethylamine (32 mg, 0.32 mmol). The reaction mixture was stirred at RT for 1 h, and then it was diluted with ethyl acetate (20 mL) and washed with brine. The aqueous layer was back extracted with ethyl acetate and the combined organic solvent was evaporated and purified with combi-flash column chromatography (Eluent methanol/dichloromethane 1:10) to afford the title compound (68 mg, 65%) as a yellow solid.




70d (55 mg, 0.08 mmol) was dissolved in the mixture of tetrahydrofuran (2 mL), methanol (1 mL) and water (1 mL). To the solution was added lithium hydroxide hydrate (4 mg, 0.16 mmol). The reaction mixture was stirred at RT for 1 h. Dichloromethane (10 mL) and water (10 mL) were added to the reaction mixture. 1 N Hydrochloric acid was added to the aqueous layer until pH reached 2, the acidic aqueous layer was extracted with dichloromethane (2 x 10 mL). The organic solvent was then evaporated to afford the title compound (37 mg, 84%) as a white solid.




2-Methyl-6-nitrobenzoic anhydride (19 mg, 0.056 mmol) and 4-dimethylaminopyridine (23 mg, 0.185 mmol) were dissolved in 1,2-dichloroethane (18 mL) and the solution was heated at 50°C. To the above solution was added **70e** (20 mg, 0.037 mmol) in *N,N*-dimethylformamide (1 mL) via syringe pump in 10 h. The reaction mixture was stirred at 50°C for 2 h after the completion of addition. The solvent was then evaporated and the residue was purified by reverse phase preparative HPLC (0-100% acetonitrile/water) to afford the title compound (4 mg, 22%) as a white solid. ¹H NMR (400 MHz, CD₃OD) δ 7.79-7.72 (m, 3H), 7.52 (dd, *J* = 1.6, 8.8 Hz, 1H), 7.41 (s, 1H), 7.33 (dd, *J* = 8.4, 1.6 Hz, 1H), 6.48 (d, *J* = 16.4 Hz, 1H), 6.31-6.24 (m, 1H), 6.02 (q, *J* = 7.6 Hz, 1H), 4.64 (d, *J* = 12.4 Hz, 1H), 4.40 (d, *J* = 12.8 Hz, 1H), 4.30 (d, *J* = 10.4 Hz, 1H), 3.78-3.72 (m, 1H), 3.36-3.32 (m, 2H), 2.99-2.93 (m, 1H), 2.77-2.71 (m, 1H), 1.99-1.69 (m, 5H), 1.66 (d, *J* = 6.8 Hz, 3H), 1.58 (d, *J* = 7.2 Hz, 3H), 0.99 (d, *J* = 6.8 Hz, 6H). LCMS (*m/z*) 521.1 [M-H], Tr = 3.18 min.

EXAMPLE 71



Compound 70 (5 mg, 0.01 mmol) was dissolved in ethanol (5 mL) under argon, to the solution was added Pd (10% on activated carbon, 3 mg). The reaction flask was then purged and then charged with H₂ using a balloon. The reaction was filtered through Celite after 2 h, the filtrate was evaporated under reduced pressure and purified by reverse phase preparative HPLC (0-100% acetonitrile/ water) to afford the title compound (3.2 mg, 64%) as a white powder. ¹H-NMR (400 MHz, CD₃OD) δ 8.47 (d, *J* = 6.0 Hz, 1H), 7.97 (d, *J* = 8.8 Hz, 1H), 7.69-7.62 (m, 3H), 7.31 (s, 1H), 7.24-7.19 (m, 2H), 5.97 (dd, *J* = 12.8, 6.0 Hz, 1H), 5.62 (pent, *J* = 7.2 Hz, 1H), 4.76-4.16 (m, 1H), 4.10 (*app t*, *J* = 12.8 Hz, 1H), 4.05-3.69 (m, 1H), 2.85 (*br s*, 1H), 2.73-2.41 (m, 2H), 2.40-2.37 (m, 1H), 2.10-1.93 (m, 1H), 1.92-1.70 (m, 6H), 1.53 (d, *J* = 6.8 Hz, 3H), 1.49 (d, *J* = 7.6 Hz, 3H), 0.99 (d, *J* = 6.8 Hz, 6H). LCMS (*m/z*) 523.140 [M+H], Tr = 3.16 min.


EXAMPLE 72



Into an oven dried, argon purged flask Compound 49 (10 mg, 0.017 mmol), copper(I) iodide (1 mg, 0.005 mmol) and prop-2-yn-1-ol (4 mg, 0.07 mmol) were added. The flask was sealed and repurged with argon three times. Anhydrous *N,N*-dimethylformamide (5 mL) was added and the reaction mixture was repurged with argon three times. This reaction mixture was stirred at RT for 12 h. After evaporation of the solvent under reduced pressure, the crude residue was dissolved in ethyl acetate (10 mL) and filtered through filter aid and the filter pad was washed with ethyl acetate (10 mL). After concentration under reduced pressure, the

residue was purified by silica gel chromatography (gradient from 0–40% ethyl acetate and methanol (4/1) in *iso*-hexanes) to afford the title compound (10 mg, 93%) as a white solid after evaporation. $R_f = 0.33$, 10% methanol in dichloromethane. ^1H NMR (400 MHz, CD_3OD) δ 9.05 (1H), 7.95 (d, $J = 8.6$ Hz, 1H), 7.94 (s, 1H), 7.82 (s, 1H), 7.50 (s, 1H), 7.46 (d, $J = 8.6$ Hz, 1H), 6.49 (d, $J = 16.0$ Hz, 1H), 6.40 (d, $J = 16.0$ Hz, 1H), 5.95 (m, 1H), 5.50 (m, 1H), 4.59–4.53 (m, 3H), 4.36 (m, 2H), 4.30 (m, 1H), 3.68 (m, 1H), 2.80 (m, 1H), 2.60 (m, 1H), 2.23 (m, 1H), 2.14 (m, 1H), 1.90–1.58 (m, 4H), 1.57 (d, $J = 6.6$ Hz, 3H), 1.55 (d, $J = 7.6$ Hz, 3H), 1.41 (s, 3H), 1.27 (s, 3H). LCMS (m/z) 633.4 $[\text{M}+\text{H}]$, $\text{Tr} = 2.66$ min.


10

EXAMPLE 73

To 3-Nitromethylene-oxetane (*Angew. Chem. Int. Ed.* 2006, 45 (46), 7736, 2.5 g, 21.7 mmol) in anhydrous tetrahydrofuran (40 mL) at RT and under an atmosphere of nitrogen was added copper(I) iodide (413 mg, 2.17 mmol) and chlorotrimethylsilane (3.0 mL, 23.9 mmol).

15 The resulting yellow solution was stirred for 5 min and cooled to between -15°C and -11°C with a methanol ice bath. Vinyl magnesium bromide (43.5 mL, 43.5 mmol, 1.0 M in tetrahydrofuran) was slowly added over 3 h via a syringe pump. Following the addition the reaction was quenched with a saturated aqueous solution of ammonium chloride, filtered and extracted with diethyl ether (3x). The combined organics were dried through a hydrophobic frit and


20 concentrated in vacuo. The residue was purified by silica gel chromatography using diethyl ether to give the title compound as a yellow oil (2.0 g, 64%).




To **73a** (600 mg, 4.2 mmol) in anhydrous dimethylsulfoxide (13 mL) was added acetic acid (2.4 mL, 42 mmol) and sodium nitrite (869 mg, 12.6 mmol) and the mixture heated to 35°C

25 for 16 h. The reaction was cooled to RT and diluted with water. The pH was adjusted to pH 3–4 with 10% aqueous hydrochloric acid and the product extracted with diethyl ether (3x) and diethyl ether / ethyl acetate (1:1). The combined organics were dried through a hydrophobic frit and concentrated in vacuo. The residue was purified by silica gel chromatography using *iso*-


hexanes/ethyl acetate (2/1 then 1/1) to afford impure acid. This was dissolved in diethyl ether and extracted with a saturated solution of sodium carbonate. The aqueous was acidified to pH 2 with concentrated hydrochloric acid and extracted with ethyl acetate (3x). The organics were dried through a hydrophobic frit and concentrated in vacuo to give the title compound as a yellow oil (228 mg, 43%).



To **73b** (150 mg, 1.18 mmol) in anhydrous dioxane (1 mL), was added acetic acid (R)-1-(7-bromoquinolin-2-yl)-ethyl ester (347 mg, 1.18 mmol) followed by dicyclohexylmethylamine (0.76 mL, 3.54 mmol), palladium(II) acetate (53 mg, 0.24 mmol) and Tri(*o*-tolyl) phosphine (72 mg, 0.24 mmol). The mixture was heated at 100°C for 1 hour. An additional amount of palladium(II) acetate (26 mg, 0.12 mmol) and Tri(*o*-tolyl) phosphine (36 mg, 0.12 mmol) was added and heating continued at 100°C for a further 45 min. The reaction was cooled to RT and 2M HCl added until pH 3-4 was reached. The product was extracted with ethyl acetate (3x) and ethyl acetate/10% methanol (2x). The combined organics were dried through a hydrophobic frit and concentrated in vacuo to yield the title compound as a brown oil.




A solution of **1e** (3.01 g, 5.66 mmol) in tetrahydrofuran:methanol (1:1, 60 mL) was stirred at 0°C. Tetra-*n*-butylammonium fluoride (1 M in tetrahydrofuran, 11.3 mL, 11.3 mmol) was added and the reaction mixture was stirred at RT for 22 h. The solvent was evaporated and the residue was purified by silica gel chromatography using *iso*-hexane to *iso*-hexane/ethyl acetate 1:1 afford the title compound (2.14 g, 91%) as a white foam.



Compound 73e

To **73d** (365 mg, 0.88 mmol) in anhydrous dichloromethane (15 mL) at 0°C and under an atmosphere of nitrogen, was added trimethylsilyl trifluoromethanesulfonate (239 μ L, 1.32 mmol). The reaction mixture was stirred at 0°C for 1 hour before adding *N,N*-diisopropylethylamine (613 μ L, 3.52 mmol) and then concentrated in vacuo, and co-evaporated with toluene to afford (S)-1-
 5 [(S)-2-((S)-2-amino-3-methyl-butrylamino)-propionyl]-hexahydro-pyridazine-3-carboxylic acid 2,2,2-trichloro-ethyl ester as a white solid. To **73c** (300 mg, 0.88 mmol) in anhydrous acetonitrile (9 mL) at 0°C and under an atmosphere of nitrogen was added *N,N*-diisopropylethylamine (766 μ L, 4.4 mmol) and 2-(1H-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyl
 10 uronium hexafluorophosphate methanaminium (401 mg, 1.06 mmol). The solution was stirred at 0°C for 3 min before adding a solution of (S)-1-[(S)-2-((S)-2-amino-3-methyl-butrylamino)-propionyl]-hexahydro-pyridazine-3-carboxylic acid 2,2,2-trichloro-ethyl ester in anhydrous acetonitrile (2 mL). The reaction was warmed to RT stirred for 2 h. The reaction was quenched
 15 with 1M HCl and extracted with ethyl acetate (3x). The combined organic layers were dried through a hydrophobic frit and concentrated in vacuo. The residue was purified by silica gel chromatography using ethyl acetate to give the title compound as a viscous yellow oil (160 mg, 22%, 2 steps).




Compound 73

To **73e** (160 mg, 0.25 mmol) in tetrahydrofuran (10 mL) and water (2 mL) was added lithium hydroxide monohydrate (53 mg, 1.25 mmol) at 0°C. The reaction was stirred at 0°C for 2
 20 h and quenched by adding 2M aqueous hydrochloric acid (0.63 mL). The reaction was concentrated in vacuo, followed by co-evaporation from toluene/methanol (3x) and then toluene (3x) and dried on a high vacuum for 15 min. The resulting residue was dissolved in anhydrous


tetrahydrofuran (83 mL) and at RT was added *N,N*-diisopropylethylamine (223 μ L, 1.25 mmol), 2-(1H-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyl uronium hexafluorophosphate methanaminium (114 mg, 1.3 mmol) and 4-dimethylaminopyridine (3 mg, 0.03 mmol). The reaction was stirred for 24 h, diluted with ethyl acetate and washed with 1M HCl (1x) and brine (1x). The organic layer was dried through a hydrophobic frit and concentrated in vacuo. The residue was purified by silica gel chromatography using ethyl acetate/methanol 1:0 then 20/1 to give a viscous yellow oil (52 mg). This was further purified by preparative thin layer chromatography (ethyl acetate) to afford the title compound as a white solid which was triturated with diethyl ether, filtered and vacuum dried. (22 mg, 16%, 2 steps). ^1H NMR (300MHz, CDCl_3) 0.96 (d, J = 6.5 Hz, 3H), 1.00 (d, J = 6.5 Hz, 3H), 1.50 (d, J = 7.1 Hz, 3H), 1.69 (d, J = 6.9 Hz, 3H), 1.81 – 2.16 (m, 4H), 2.63 – 2.74 (m, 1H), 3.26 – 3.38 (m 1H), 3.70 – 3.83 (m, 1H), 3.98 (d, J = 12.3 Hz, 1H), 4.28 (t, J = 9.8 Hz, 1H), 4.33 – 4.43 (m, 1H), 4.57 (d, J = 5.8 Hz, 1H), 4.86 (q, J = 6.7 Hz, 2H), 5.13 (d, J = 5.8 Hz, 1H), 5.73 (t, J = 6.9 Hz, 1H), 5.96 (q, J = 6.7 Hz, 1H), 6.24 (d, J = 16.3 Hz, 1H), 6.61 – 6.73 (m, 1H), 7.00 (d, J = 16.3 Hz, 1H), 7.09 – 7.20 (m, 1H), 7.40 (d, J = 8.5 Hz, 1H), 7.58 (s, 1H), 7.81 – 7.93 (m, 2H), 8.22 (d, J = 8.7 Hz, 1H). LCMS (m/z) = 564.2 [M+H], Tr = 1.96 min.

EXAMPLE 74




Compound 57 (20 mg, 0.036 mmol) was dissolved in methanol (5 mL) and added catalytic amount of 10% Pd on carbon. The reaction mixture was stirred under atmosphere hydrogen for 2 h. The catalyst was removed by filtration and the eluent was concentrated under reduced pressure. Purification by reverse phase preparative HPLC gave the title compound (8.4 mg, 42% yield) as a white powder. ^1H NMR (300 MHz, CD_3OD): δ 9.61 (s, 1H), 8.42 (d, J = 9.0 Hz, 1H), 7.99 (s, 1H), 7.89 (d, J = 9.0 Hz, 1H), 7.83 (br s, 1H), 7.29 (d, J = 8.7 Hz, 1H), 6.27–6.21 (m, 1H), 5.92–5.82 (m, 1H), 3.96–3.5 (m, 4H), 3.28–3.19 (m, 1H), 3.08–2.98 (m, 1H), 2.30–2.20 (m, 1H), 2.10–1.90 (m, 4H), 1.78–1.72 (m, 5H), 1.56–1.51 (m, 6H), 1.21 (s, 3H), 0.89 (d, J = 7.2 Hz, 6H). LCMS (m/z): 552.3 [M+H], Tr = 1.74 min.


EXAMPLE 75



A solution of ethyl acetoacetate (20 g, 19.4 mL, 0.154 mol) in a mixture of dioxane (120 mL) and aqueous formaldehyde (37% solution in water, 57.7 mL, 0.77 mol) was stirred at RT. Triethylamine (1.0 M in tetrahydrofuran, 7.7 mL, 7.7 mmol) was added and the reaction mixture was heated at 60 °C for 20 h and then heated at 100 °C for 4 h. The reaction mixture was cooled to RT and was poured into water (1500 mL). The aqueous solution was washed with toluene. The aqueous layer was concentrated to ~50% of the initial volume and extracted with ethyl acetate. The organic extracts were combined and evaporated. The residue was purified by silica gel chromatography using a gradient of *iso*-hexane to *iso*-hexane/ethyl acetate 1:1 followed by silica gel chromatography using a gradient of *iso*-hexane to *iso*-hexane/ethyl acetate 3:2 to afford the title compound (2.33 g, 8%) as a yellow oil.




A solution of **75a** (2.10 g, 11 mmol), 2,2-dimethoxypropane (13.5 mL, 110 mmol) and 4-toluenesulfonic acid hydrate (209 mg, 1.1 mmol) in acetone (8 mL) was stirred at RT for 18 h. Saturated sodium hydrogen carbonate solution was added and the mixture was extracted with ethyl acetate. The organic extracts were combined, washed with water and brine. The organic layer was filtered through a hydrophobic frit and the filtrate was evaporated. The residue was purified by silica gel chromatography using a gradient of *iso*-hexane/ethyl acetate 1:5 to 3:7 to afford the title compound (1.88 g, 74%) as a colorless oil.




A solution of *N,N*-diisopropylethylamine (0.55 mL, 3.9 mmol) in tetrahydrofuran (20 mL) was stirred at -78 °C under nitrogen. *n*-Butyl lithium (1.6 M in hexane, 2.25 mL, 3.6 mmol) was added dropwise and the reaction mixture was warmed to 0 °C. The reaction mixture was stirred

at 0°C for 5 min and then cooled to -78°C. A solution of **75b** (690 mg, 3.0 mmol) in tetrahydrofuran (3 mL) was added dropwise and the reaction mixture was stirred at -78°C for 15 min. A solution of *N*-phenyl-(bistrifluoromethanesulfonamide) (1.18 g, 3.3 mmol) in tetrahydrofuran (10 mL) was added dropwise over 5 min and the reaction mixture was stirred at -78°C for 15 min. The cooling bath was removed and the reaction mixture was warmed to RT and then stirred at RT for 90 min. The solvent was evaporated and diethyl ether (30 mL) was added. The solution was cooled to 5°C and was washed with cold 1 M sodium hydroxide solution (3 x 30 mL). The organic layer was separated, washed with brine, dried over anhydrous sodium sulfate, filtered and the solvent evaporated to afford the title compound (960 mg, 88%) as a yellow oil which was used directly in the next step.



Compound 75d


A solution of **75c** (470 mg, 1.3 mmol) and tri-*n*-butylamine (721 mg, 0.93 mL, 3.9 mmol) in *N,N*-dimethylformamide (3 mL) was stirred at RT under nitrogen. Bis(triphenylphosphine)palladium(II) dichloride (45 mg, 0.065 mmol) and formic acid (120 mg, 0.1 mL, 2.6 mmol) was added and the reaction mixture was heated at 60 °C for 90 min. The reaction mixture was cooled to RT and ethyl acetate and water was added. The organic extract was separated, washed with water (x 5), and brine. The organic solution was filtered through a hydrophobic frit and the filtrate was evaporated. The residue was purified by silica gel chromatography using *iso*-hexane/ethyl acetate 1:9 to afford the title compound (1.88 g, 74%) as a colorless oil.



Compound 75e

A solution of **75d** (150 mg, 0.7 mmol) in tetrahydrofuran (3 mL) was stirred at 5°C under nitrogen. A solution of lithium hydroxide monohydrate (59 mg, 1.4 mmol) in water (1 mL) was added and the reaction mixture was stirred at 5°C for 30 min and then at RT for 5 h. Methanol (0.5 mL) was added to give a clear solution and the reaction mixture was stirred at RT for 22 h. The solvent was evaporated. Water (2 mL) was added to the residue and the solution was acidified to pH 2 with 2 M hydrochloric acid. Brine was added and the mixture was extracted


with ethyl acetate. The organic extracts were combined, washed with brine. The organic extract was separated, washed with water (x 5), and brine. The organic solution was filtered through a hydrophobic frit and the filtrate was evaporated to afford the title compound (117 mg, 90%) as a colorless oil.



A mixture of **75e** (96 mg, 0.5 mmol) and acetic acid (R)-1-(7-bromo-quinolin-2-yl)-ethyl ester (147 mg, 0.5 mmol) in acetonitrile (4 mL) was stirred at RT. Tri(o-tolyl)phosphine (46 mg, 0.15 mmol), palladium(II) acetate (17 mg, 0.075 mmol) and triethylamine (101 mg, 0.14 mL, 1.0 mmol) was added and the reaction mixture was heated in a microwave reactor at 100°C for 20 min. The solvent was evaporated. Water and ethyl acetate was added and the mixture was acidified to pH 3-4 with 2 M hydrochloric acid. The mixture was extracted with ethyl acetate and the organic extracts were combined and washed with brine. The organic solution was filtered through a hydrophobic frit and the filtrate was evaporated. The residue was purified by silica gel chromatography using *iso*-hexane/ethyl acetate 1:1 to ethyl acetate to ethyl acetate/methanol 5:1. The residue was co-evaporated with ethyl acetate and then dichloromethane and dried to afford the title compound (134 mg, 67%) as a yellow gum.

10

15




A solution of (S)-1-[(S)-2-((S)-2-tert-butoxycarbonylamino-3-methyl-butylamino)-propionyl]-hexahydro-pyridazine-3-carboxylic acid methyl ester (150 mg, 0.36 mmol) in dichloromethane (5 mL) was stirred at 0 °C under nitrogen. Trimethylsilyl trifluoromethanesulfonate (160 mg, 0.13 mL, 0.72 mmol) was added and the reaction mixture was stirred at 0°C for 1 hour. *N,N*-Diisopropylethylamine (186 mg, 0.25 mL, 1.44 mmol) was added and the solvent was evaporated to afford (S)-1-[(S)-2-((S)-2-amino-3-methyl-butylamino)-propionyl]-hexahydro-pyridazine-3-carboxylic acid methyl ester (0.36 mmol) as a white solid which was used without further purification. A solution of **75f** (134 mg, 0.33 mmol) in

20


25

acetonitrile was stirred at 0°C under nitrogen. A solution of (S)-1-[(S)-2-((S)-2-amino-3-methylbutyrylamino)-propionyl]-hexahydro-pyridazine-3-carboxylic acid methyl ester (0.36 mmol) and *N,N*-diisopropylethylamine (129 mg, 0.17 mL, 1.0 mmol) in acetonitrile (5 mL) was added followed by 2-(1H-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyl uronium hexafluorophosphate methanaminium (175 mg, 0.46 mmol) and the reaction mixture was stirred at 0°C for 10 min and then at RT for 1 hour. The solvent was evaporated and the residue was aportioned between ethyl acetate and water. The organic extracts were combined and washed with brine. The organic solution was filtered through a hydrophobic frit and the filtrate was evaporated. The residue was purified by silica gel chromatography using *iso*-hexane/ethyl acetate 1:5 to ethyl acetate to afford the title compound (165 mg, 72%) as a yellow solid.



Compound 75h

A solution of **75g** (35 mg, 0.05 mmol) in tetrahydrofuran (0.5 mL) was stirred at 5°C under nitrogen. A solution of lithium hydroxide monohydrate (8.4 mg, 0.2 mmol) in water (0.5 mL) was added followed by methanol (0.5 mL) and the reaction mixture was stirred at 5°C for 1 hour. 1 M hydrochloric acid (0.2 mL) was added and the solvent was evaporated. The residue was co-evaporated with methanol/toluene (1:1, x 2) followed by toluene (x 2). The residue was triturated with diethyl ether (x 3) and dried to afford the title compound (0.05 mmol) as a pale yellow solid which was used crude in the next reaction.




Compound 75

A solution of 2-methyl-6-nitrobenzoic anhydride (86 mg, 0.25 mmol) and 4-dimethylaminopyridine (46 mg, 0.38 mmol) in 1,2-dichloroethane (16 mL) was stirred at RT under nitrogen. A molecular sieves (400 mg) was added and the suspension was heated at

50°C. A solution of crude **75h** (0.05 mmol) in *N,N*-dimethylformamide (2 mL) was added dropwise over 4 h and the reaction mixture was stirred at 50°C for an additional 1 hour. The reaction mixture was cooled to RT and the mixture was filtered through celite. The filter pad was washed with ethyl acetate and the filtrate was evaporated. The residue was diluted with ethyl acetate and the solution was washed with brine (x 3). The organic layer was separated and evaporated. The organic solution was filtered through a hydrophobic frit and the filtrate was evaporated. The residue was purified by preparative HPLC to afford the title compound (4.2 mg, 13%) as a white solid. ¹H NMR (300 MHz, CD₃OD) 1.00 (d, *J* = 6.7 Hz, 3H), 1.01 (d, *J* = 6.6 Hz, 3H), 1.37 (s, 3H), 1.47 (s, 3H), 1.63 (d, *J* = 7.1 Hz, 3H), 1.64-1.70 (m, 2H), 1.73 (d, *J* = 6.9 Hz, 3H), 1.85-2.10 (m, 3H), 2.70-2.80 (m, 1H), 3.78-3.87 (m, 1H), 3.87 (d, *J* = 11.1 Hz, 1H), 4.03-4.07 (m, 1H), 4.26-4.46 (m, 5H), 5.72 (q, *J* = 7.1 Hz, 1H), 5.93 (q, *J* = 6.8 Hz, 1H), 6.38 (s, 2H), 7.42 (d, *J* = 8.5 Hz, 1H), 7.61 (*br s*, 1H), 7.78-7.86 (m, 2H), 8.22 (d, *J* = 8.5 Hz, 1H). LCMS (*m/z*) 622.2 [*M*+*H*], *Tr* = 2.24 min.

EXAMPLE 76



Compound 76

A suspension of crude **75h** (200 mg, 0.25 mmol) in tetrahydrofuran (100 mL) was stirred at RT under nitrogen. *N,N*-diisopropylethylamine (161 mg, 0.22 mL, 1.25 mmol) and 4-dimethylaminopyridine (15 mg, 0.125 mmol) was added and the suspension was stirred for 5 min. 2-(1H-7-Azabenzotriazol-1-yl)-1,1,3,3-tetramethyl uronium hexafluorophosphate methanaminium (133 mg, 0.35 mmol) was added and the reaction mixture was stirred at RT for 2 h. Additional 2-(1H-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyl uronium hexafluorophosphate methanaminium (57 mg, 0.15 mmol) was added and the reaction mixture was stirred at RT for 1 hour. The solvent was evaporated and the residue was diluted with ethyl acetate and 2 M hydrochloric acid. The organic solution was separated and washed with water, saturated sodium hydrogen carbonate solution, and brine. The organic solution was filtered through a hydrophobic frit and the filtrate was evaporated. The residue was purified by silica gel chromatography using

iso-hexane/ethyl acetate 7:3 to ethyl acetate to ethyl acetate/methanol 9:1. The residue was purified by preparative HPLC to afford the title compound (20 mg, 14%) as a white solid. ¹H NMR (500 MHz, CD₃OD) 1.00 (d, *J* = 6.5 Hz, 3H), 1.01 (d, *J* = 6.5 Hz, 3H), 1.68 (d, *J* = 7.1 Hz, 3H), 1.73 (d, *J* = 6.7 Hz, 3H), 1.74-2.05 (m, 5H), 2.72-2.80 (m, 1H), 3.70 (d, *J* = 11.1 Hz, 1H), 3.81-4.07 (m, 4H), 4.35-4.47 (m, 3H), 5.77 (q, *J* = 7.1 Hz, 1H), 5.92 (q, *J* = 6.8 Hz, 1H), 6.43 (s, 2H), 7.40 (d, *J* = 8.5 Hz, 1H), 7.63 (s, 1H), 7.82 (s, 2H), 8.21 (d, *J* = 8.5 Hz, 1H). LCMS (*m/z*) 582.2 [M+H], Tr = 1.66 min.

Biological Examples: Inhibition of Peptidyl-Prolyl Isomerase (PPIase) Activity

The PPIase assay was based on the procedure reported by Janowski et al. (*Anal. Biochem.* 1997, 252, 299). Assay buffer (1980L of a solution containing 35 mM HEPES pH 7.8, 50M DTT, and 0.01% NP40) was pre-equilibrated to 10°C in a quartz cuvette equipped with an overhead stirrer. To this solution was added 10L of compound in DMSO (final concentration: 0.5% DMSO), followed by 5L of a 2M stock solution of cyclophilin A (final concentration: 5 nM). The reaction was initiated with the addition of 5L of 40mM of the tetrapeptide Succ-AAPF-pNA (100M final concentration) dissolved in a solution of 0.5M LiCl in trifluoroethanol. Upon the initiation of the reaction, the absorbance of the peptide substrate was monitored at 330nm for five minutes using a Beckman Coulter DU800 spectrophotometer. Progress curves were fit with a single-exponential decay model to calculate rates. The IC₅₀ values were calculated with a four-parameter logistic fit using GraphPad Prism software.

Cyclophilin A TR-FRET Competitive Binding Assay

Inhibitor potency was measured using a competitive binding assay with a time-resolved fluorescence resonance energy transfer (TR-FRET) readout. To a reaction buffer consisting of 35mM HEPES pH 7.8, 100mM NaCl, 0.01% NP-40 (Pierce), 1mM DTT, and 1% DMSO were added the following: 5 nM of cyclophilin A modified at the N-terminus with an 8×histidine affinity tag (CypA); 150 nM of cyclosporin A modified with a linker attached to a Cy5 fluorophore (CsA-Cy5); 1 nM Eu-labeled anti-(6×His) antibody (Perkin-Elmer); and test compound at one of various concentrations. The total volume of the assay solution was 100L. After a two-hour incubation, the TR-FRET was measured using a Perkin Elmer Envision plate reader (excitation at 340nm, emission measured at 590nm and 665nm). The signal was calculated as the ratio of the emission at 665nm to that at 590nm. An IC₅₀ value was calculated using a four-parameter logistic fit.

When tested, certain compounds of this invention were found to inhibit cyclophilin binding as listed in Table 1 below. The IC₅₀'s are presented as ranges wherein A is ≤ 100 nM, B is 101 to 1000 nM and C is 1001 to 10,000 nM.

Antiviral Activity

5 The antiviral activity of a compound can be measured using standard screening protocols: for example, cell-based Flavivirus immunodetection assay and cell-based Flavivirus cytopathic effect assay as described in U.S. Patent Publication Number US/20130022573, which is hereby incorporated by reference in its entirety.

10 One aspect of the invention relates to methods of inhibiting viral infections, comprising the step of treating a sample or subject suspected of needing such inhibition with a composition of the invention. The antiviral activity of a compound of the invention can be measured using standard screening protocols that are known.

The anti-HCV activity of the compounds of this invention was tested in a human hepatoma Huh-7 cell line harboring a HCV replicon. The assay comprised the following steps:

15 **Step 1 (compound preparation and serial dilution):** Serial dilution was performed in 100% DMSO in a 384-well plate. A solution containing a compound at 225-fold concentration of the starting final serial dilution concentration was prepared in 100% DMSO and 15 µL added to the pre-specified wells in column 3 or 13 of a polypropylene 384-well plate. The rest of the 384-well plate was filled with 10 µL 100% DMSO except for columns 23 and 24, where 10 µL of 500
20 µM a HCV protease inhibitor (ITMN-191) in 100% DMSO was added. The HCV protease inhibitor was used a control of 100% inhibition of HCV replication. The plate was then placed on a Biomek FX Workstation to start the serial dilution. The serial dilution was performed for ten cycles of 3-fold dilution from column 3 to 12 or from column 13 to 22.

25 **Step 2 (cell culture plate preparation and compound addition):** To each well of a black polypropylene 384-well plate, 90 µL of cell media containing 1600 suspended Huh-7 HCV replicon cells was added with a Biotek uFlow Workstation. A volume of 0.4 µL of the compound solution was transferred from the serial dilution plate to the cell culture plate on a Biomek FX Workstation. The DMSO concentration in the final assay condition was 0.44%. The plates were incubated for 3 days at 37°C with 5% CO₂ and 85% humidity.

30 **Step 3 (detection of cytotoxicity and inhibition of viral replication):** a) Assessment of cytotoxicity: The media in the 384-well cell culture plate was aspirated with a Biotek EL405

plate-washer. A volume of 50 μ L of a solution containing 400 nM Calcein AM in 100% PBS was added to each well of the plate with a Biotek uFlow Workstation. The plate was incubated for 30 minutes at RT before the fluorescence signal (emission 490 nm, excitation 520 nm) was measured with a Perkin Elmer Envision Plate Reader.

5 b) Assessment of inhibition of viral replication: The calcein-PBS solution in the 384-well cell culture plate was aspirated with a Biotek EL405 plate-washer. A volume of 20 μ L of Dual-Glo luciferase buffer (Promega, Dual-Glo Luciferase Assay Reagent, cat. #E298B) was added to each well of the plate with a Biotek uFlow Workstation. The plate was incubated for 10 minutes at RT. A volume of 20 μ L of a solution containing 1:100 mixture of Dual-Glo Stop & Glo
10 substrate (Promega, Dual-Glo Luciferase Assay Reagent, cat. #E313B) and Dual-Glo Stop & Glo buffer (Promega, Dual-Glo Luciferase Assay Reagent, cat. #E314B) was then added to each well of the plate with a Biotek uFlow Workstation. The plate was incubated at RT for 10 minutes before the luminescence signal was measured with a Perkin Elmer Envision Plate Reader.

15 **Step 4 (calculation):** The percent cytotoxicity was determined by calcein AM conversion to fluorescent product. The average fluorescent signal from the DMSO control wells were defined as 100% nontoxic. The individual fluorescent signal from testing compound treated well was divided by the average signal from DMSO control wells and then multiplied by 100% to get the percent viability. The percent anti-HCV replication activity was determined by the
20 luminescence signal from the testing well compared to DMSO controls wells. The background signal was determined by the average luminescence signal from the HCV protease inhibitor treated wells and was subtracted from the signal from the testing wells as well as the DMSO control wells. Following 3-fold serial dilutions, the EC₅₀ and CC₅₀ values were calculated by fitting % inhibition at each concentration to the following equation:

25
$$\% \text{ inhibition} = 100\% / [(EC_{50}/[I])^b + 1]$$

 where b is Hill's coefficient. See, for reference, Hill, A. V., *The Possible Effects of the Aggregation of the Molecules of Hæmoglobin on its Dissociation Curves*, J. Physiol. 40: iv-vii. (1910). % inhibition values at a specific concentration, for example 2 μ M, can also be derived from the formula above.

30 When tested, certain compounds of this invention were found to inhibit viral replication as listed in Table 1. The EC₅₀'s are presented as a % inhibition.

Table 1

Example No.	TR-FRET	Replicon 1a % inhibition at 1 μ M	Example No.	TR-FRET	Replicon 1a % inhibition at 1 μ M
1	A	32	39	A	98
2	B	63	40	A	96
3	A	44	41	A	96
4	B	58	42	B	16
5	A	80	43	A	97
6	A	86	44	A	78
7	A	94	45	A	80
8	B	37	46	A	90
9	A	98	47	B	9
10	B	28	48	A	40
11	A	6	49	A	85
12	C	0	50	B	12
13	A	87	51	A	83
14	A	94	52	A	98
15	A	99	53	B	89
16	B	94	54	A	98
17	A	100	55	B	76
18	A	94	56	B	47
19	A	97	57	A	100
20	A	99	58	A	99
21	A	93	59	A	100
22	A	100	60	B	65
23	A	99	61	A	72
24	A	99	62	B	25
25	B	71	63	B	80
26	A	99	64	B	35
27	A	100	65	A	92
28	B	56	66	A	80
29	C	28	67	A	97
30	A	73	68	A	91
31	A	98	69	C	19
32	A	71	70	A	95
33	C	14	71	B	-
34	A	92	72	B	0
35	A	94	73	A	94
36	A	73	74	A	95
37	A	76	75	A	100
38	A	98	76	A	63

The specific pharmacological and biochemical responses observed may vary according to and depending on the particular active compound selected or whether there are present


5 pharmaceutical carriers, as well as the type of formulation and mode of administration

and such expected variations or differences in the results are contemplated in accordance with practice of the present invention.

Although specific embodiments of the present invention are herein illustrated and described in detail, the invention is not limited thereto. The above detailed descriptions are
5 provided as exemplary of the present invention and should not be construed as constituting any limitation of the invention.

WHAT IS CLAIMED IS:

1. A compound of Formula I:



or a pharmaceutically acceptable salt, isotope, stereoisomer, mixture of stereoisomers, tautomer, or ester thereof, wherein:

A¹ is (C₂-C₅)alkylene, (C₂-C₅)alkenylene, (C₂-C₅)alkynylene, -O-(C₂-C₄)alkylene, -O-(C₂-C₄)alkenylene, arylene, aryl(C₁-C₂)alkylene, heterocycloalkylene, pyrazolylene, pyridylene, pyrimidinylene or heterocycloalkyl(C₁-C₂)alkylene, wherein a sp³ carbon atom of A¹ is optionally substituted with one or more (C₁-C₄)alkyl;

A² is arylene or heteroarylene, wherein A² is optionally substituted with halo;

X¹ is -O-, -NH- or -N((C₁-C₄)alkyl)-;

R^{1a} and R^{1b} are independently H, (C₁-C₄)alkyl, (C₂-C₄)alkenyl or (C₂-C₄)alkynyl;

R² is H, (C₁-C₄)alkyl, (C₂-C₄)alkenyl or (C₂-C₄)alkynyl;

R^{3a} and R^{3b} are independently H or (C₁-C₈)alkyl;

R^{4a} and R^{4b} are independently H, -OH, (C₁-C₄)alkoxy, halo(C₁-C₄)alkoxy or (C₁-C₈)alkyl;

R⁵ is H, (C₁-C₄)alkyl, (C₂-C₄)alkenyl or (C₂-C₄)alkynyl, or

R⁵ forms a cyclic moiety along with -N((C₁-C₄)alkyl)- of X¹ or arylene of A²; and

R⁶ is H or (C₁-C₄)alkyl.


2. The compound of Claim 1, wherein A¹ is ethenylene, propenylene, butenylene, ethylene, propylene, butylene, oxypropylene, oxypropenylene, pyrazolylene, phenylene or pyrimidinylene.

3. The compound of any preceding claim, wherein A² is isoquinolinylene, phenylene or halophenylene.


4. The compound of any preceding claim, wherein X¹ is -O- or -NH-; one of R^{1a} and R^{1b} is H and the other is methyl; R² is iso-propyl; R⁵ is methyl and R⁶ is H or methyl.

5. The compound of any preceding claim, wherein R^{3a} is H or methyl; R^{3b} is H; R^{4a} is H, -OH, methoxy, trifluoroethoxy; and R^{4b} is H.

6. The compound of any preceding claim, which is a compound of Formula II:



or a pharmaceutically acceptable salt, isotope, stereoisomer, mixture of stereoisomers, tautomer, or ester thereof, wherein:



X¹ is -O- or -NH-;


R^{3a} is H or (C₁-C₄)alkyl;

R^{4a} is H, -OH, (C₁-C₄)alkoxy, halo(C₁-C₄)alkoxy or (C₁-C₈)alkyl; and

R⁵ is H or (C₁-C₄)alkyl.

7. The compound of any preceding claim, wherein A² is heteroarylene; A¹ is (C₂-C₅)alkylene, (C₂-C₅)alkenylene, (C₂-C₅)alkynylene, wherein A¹ is optionally substituted with one or more (C₁-C₄)alkyl; R^{3a} is H or (C₁-C₈)alkyl; and R^{4a} is H, -OH or (C₁-C₄)alkoxy.


8. The compound of any preceding claim, which is



or a pharmaceutically acceptable salt, isotope, stereoisomer, mixture of stereoisomers, tautomer, or ester thereof.

9. The compound of any preceding claim, wherein A² is arylene; and A¹ is (C₂-C₅)alkylene, (C₂-C₅)alkenylene, (C₂-C₅)alkynylene, -O-(C₂-C₅)alkylene, -O-(C₂-C₄)alkenylene, wherein A¹ is optionally substituted with one or more (C₁-C₄)alkyl; R^{3a} is H or (C₁-C₄)alkyl; and R^{4a} is H, -OH, (C₁-C₄)alkoxy or halo(C₁-C₄)alkoxy.


10. The compound of any preceding claim, which is



or a pharmaceutically acceptable salt, isotope, stereoisomer, mixture of stereoisomers, tautomer, or ester thereof.

11. The compound of any preceding claim, wherein A^2 is arylene; and A^1 is pyrazolylene, phenylene or pyrimidinylene.


12. The compound of any preceding claim, which is



or a pharmaceutically acceptable salt, isotope, stereoisomer, mixture of stereoisomers, tautomer, or ester thereof.


13. The compound of any preceding claim, wherein A^2 is haloarylene; and A^1 is $-O-(C_2-C_5)$ alkylene or $-O-(C_2-C_4)$ alkenylene.


14. The compound of any preceding claim, which is



or a pharmaceutically acceptable salt, isotope, stereoisomer, mixture of stereoisomers, tautomer, or ester thereof.


15. The compound of any preceding claim, wherein A^1 is (C_2-C_5) alkylene or $(C_2-$

$C_5)$ alkenylene; R^5 is methyl, or R^5 form  along with arylene of A^2 , or R^5 form



along with $-N((C_1-C_4)alkyl)-$ of X^1 ; and R^6 is H or methyl.

16. The compound of any preceding claim, which is



or a pharmaceutically acceptable salt, isotope, stereoisomer, mixture of stereoisomers, tautomer, or ester thereof.

17. A pharmaceutical composition comprising a therapeutically effective amount of a compound of any preceding claim or a pharmaceutically acceptable salt, isotope, stereoisomer, mixture of stereoisomers, tautomer, or ester thereof and a pharmaceutically acceptable excipient.

18. The pharmaceutical composition of Claim 17, further comprising at least one additional therapeutic agent selected from the group consisting of interferons, ribavirin, HCV NS3 protease inhibitors, HCV NS5a inhibitors, nucleoside or nucleotide inhibitors of HCV NS5B polymerase, non-nucleoside inhibitors of HCV NS5B polymerase, and TLR-7 agonists; or a mixture thereof.

19. The pharmaceutical composition of Claim 18, wherein the at least one additional therapeutic agent is ribavirin, telaprevir, boceprevir or sofosbuvir.