

target disease, which includes administering the isolated peptide or a composition containing the isolated peptide to a subject in need thereof.

Still another object of the present invention is to provide a use of the isolated peptide or the composition for use in the preparation of a medicament.

### [Technical Solution]

To achieve the above objects, in an example, the present disclosure provides an isolated peptide having activities to a glucagon receptor, a glucagon-like peptide-1 (GLP-1) receptor, and a glucose-dependent insulintropic polypeptide (GIP) receptor.

In one aspect, the present disclosure refers to an isolated peptide having activities to a glucagon receptor, a glucagon-like peptide-1 (GLP-1) receptor, and a glucose-dependent insulintropic polypeptide (GIP) receptor which comprises an amino acid sequence represented by General Formula 1 below:

Xaa1-Xaa2-Xaa3-Gly-Thr-Phe-Xaa7-Ser-Asp-Xaa10-Ser-Xaa12-Xaa13-Xaa14-Xaa15-Xaa16-Xaa17-Xaa18-Xaa19-Xaa20-Xaa21-Phe-Xaa23-Xaa24-Trp-Leu-Xaa27-Xaa28-Xaa29-Xaa30-R1 (General Formula 1, SEQ ID NO: 103)

wherein, in General Formula 1,

Xaa1 is histidine (His, H), 4-imidazoacetyl (CA), or tyrosine (Tyr, Y);

Xaa2 is glycine (Gly, G),  $\alpha$ -methyl-glutamic acid, or aminoisobutyric acid (Aib);

Xaa3 is glutamic acid (Glu, E) or glutamine (Gln, Q);

Xaa7 is threonine (Thr, T) or isoleucine (Ile, I);

Xaa10 is leucine (Leu, L), tyrosine (Tyr, Y), lysine (Lys, K), cysteine (Cys, C), or valine (Val, V);

Xaa12 is lysine (Lys, K), serine (Ser, S), or isoleucine (Ile, I);

Xaa13 is glutamine (Gln, Q), tyrosine (Tyr, Y), alanine (Ala, A), or cysteine (Cys, C);

Xaa14 is leucine (Leu, L), methionine (Met, M), or tyrosine (Tyr, Y);

Xaa15 is cysteine (Cys, C), aspartic acid (Asp, D), glutamic acid (Glu, E), or leucine (Leu, L);

Xaa16 is glycine (Gly, G), glutamic acid (Glu, E), or serine (Ser, S);

Xaa17 is glutamine (Gln, Q), arginine (Arg, R), isoleucine (Ile, I), glutamic acid (Glu, E), cysteine (Cys, C), or lysine (Lys, K);

Xaa18 is alanine (Ala, A), glutamine (Gln, Q), arginine (Arg, R), or histidine (His, H);

Xaa19 is alanine (Ala, A), glutamine (Gln, Q), cysteine (Cys, C), or valine (Val, V);

Xaa20 is lysine (Lys, K), glutamine (Gln, Q), or arginine (Arg, R);

Xaa21 is glutamic acid (Glu, E), glutamine (Gln, Q), leucine (Leu, L), cysteine (Cys, C), or aspartic acid (Asp, D);

Xaa23 is valine (Val, V);

Xaa24 is alanine (Ala, A), glutamine (Gln, Q), cysteine (Cys, C), asparagine (Asn, N), aspartic acid (Asp, D), or glutamic acid (Glu, E);

Xaa27 is valine (Val, V), leucine (Leu, L), or lysine (Lys, K);

Xaa28 is cysteine (Cys, C), lysine (Lys, K), alanine (Ala, A), asparagine (Asn, N), or aspartic acid (Asp, D);

Xaa29 is cysteine (Cys, C), glycine (Gly, G), glutamine (Gln, Q), threonine (Thr, T), glutamic acid (Glu, E), or histidine (His, H);

Xaa30 is cysteine (Cys, C), glycine (Gly, G), lysine (Lys, K), or histidine (His, H), or is absent; and

R1 is cysteine (Cys, C), GKKNDWKHNIT (SEQ ID NO: 106), m-SSGAPPPS-n (SEQ ID NO: 107), or m-SSGQPPPS-n (SEQ ID NO: 108), or is absent;

wherein:

m is -Cys-, -Pro-, or -Gly-Pro-; and

n is -Cys-, -Gly-, -Ser-, or -His-Gly-, or is absent.

In one example, the present disclosure refers to an isolated peptide having activities to a glucagon receptor, a glucagon-like peptide-1 (GLP-1) receptor, and a glucose-dependent insulinotropic polypeptide (GIP) receptor.

In another aspect, the present disclosure refers to a polynucleotide encoding the peptide as disclosed herein.

In yet another aspect, the present disclosure refers to the use of a peptide as disclosed herein in the manufacture of a medicament for preventing or treating a metabolic syndrome.

In yet another example, the present disclosure refers to a pharmaceutical composition for preventing or treating metabolic syndrome comprising a peptide as disclosed herein.

In a specific example, the peptide is an analog of native glucagon with a variation selected from the group consisting of substitution, addition, deletion, modification, and a combination thereof, on at least one amino acid of the native glucagon sequence.

In another specific example, the amino acid sequence with addition is derived from a native GLP-1 amino acid sequence, a native GIP amino acid sequence, or a native exendin-4 amino acid sequence.

In still another specific example, the peptide is an isolated peptide including an amino acid sequence represented by General Formula 1 below:

Xaa1-Xaa2-Xaa3-Gly-Thr-Phe-Xaa7-Ser-Asp-Xaa10-Ser-Xaa12-Xaa13-Xaa14-Xaa15-Xaa16-Xaa17-Xaa18-Xaa19-Xaa20-Xaa21-Phe-Xaa23-Xaa24-Trp-Leu-Xaa27-Xaa28-Xaa29-Xaa30-R1 (General Formula 1, SEQ ID NO: 103)

wherein, in General Formula 1,

Xaa1 is histidine (His, H), 4-imidazoacetyl (CA), or tyrosine (Tyr, Y);

Xaa2 is glycine (Gly, G),  $\alpha$ -methyl-glutamic acid, or Aib (aminoisobutyric acid);

Xaa3 is glutamic acid (Glu, E) or glutamine (Gln, Q);

Xaa7 is threonine (Thr, T) or isoleucine (Ile, I);

Xaa10 is leucine (Leu, L), tyrosine (Tyr, Y), lysine (Lys, K), cysteine (Cys, C), or valine (Val, V);

Xaa12 is lysine (Lys, K), serine (Ser, S), or isoleucine (Ile, I);

Xaa13 is glutamine (Gln, Q), tyrosine (Tyr, Y), alanine (Ala, A), or cysteine (Cys, C);

Xaa14 is leucine (Leu, L), methionine (Met, M), or tyrosine (Tyr, Y);

## **Claims**

1. An isolated peptide having activities to a glucagon receptor, a glucagon-like peptide-1 (GLP-1) receptor, and a glucose-dependent insulintropic polypeptide (GIP) receptor which comprises an amino acid sequence represented by General Formula 1 below:

Xaa1-Xaa2-Xaa3-Gly-Thr-Phe-Xaa7-Ser-Asp-Xaa10-Ser-Xaa12-Xaa13-Xaa14-Xaa15-Xaa16-Xaa17-Xaa18-Xaa19-Xaa20-Xaa21-Phe-Xaa23-Xaa24-Trp-Leu-Xaa27-Xaa28-Xaa29-Xaa30-R1 (General Formula 1, SEQ ID NO: 103)

wherein, in General Formula 1,  
Xaa1 is histidine (His, H), 4-imidazoacetyl (CA), or tyrosine (Tyr, Y);  
Xaa2 is glycine (Gly, G),  $\alpha$ -methyl-glutamic acid, or aminoisobutyric acid (Aib);  
Xaa3 is glutamic acid (Glu, E) or glutamine (Gln, Q);  
Xaa7 is threonine (Thr, T) or isoleucine (Ile, I);  
Xaa10 is leucine (Leu, L), tyrosine (Tyr, Y), lysine (Lys, K), cysteine (Cys, C), or valine (Val, V);  
Xaa12 is lysine (Lys, K), serine (Ser, S), or isoleucine (Ile, I);  
Xaa13 is glutamine (Gln, Q), tyrosine (Tyr, Y), alanine (Ala, A), or cysteine (Cys, C);  
Xaa14 is leucine (Leu, L), methionine (Met, M), or tyrosine (Tyr, Y);  
Xaa15 is cysteine (Cys, C), aspartic acid (Asp, D), glutamic acid (Glu, E), or leucine (Leu, L);  
Xaa16 is glycine (Gly, G), glutamic acid (Glu, E), or serine (Ser, S);  
Xaa17 is glutamine (Gln, Q), arginine (Arg, R), isoleucine (Ile, I), glutamic acid (Glu, E), cysteine (Cys, C), or lysine (Lys, K);  
Xaa18 is alanine (Ala, A), glutamine (Gln, Q), arginine (Arg, R), or histidine (His, H);  
Xaa19 is alanine (Ala, A), glutamine (Gln, Q), cysteine (Cys, C), or valine (Val, V);  
Xaa20 is lysine (Lys, K), glutamine (Gln, Q), or arginine (Arg, R);  
Xaa21 is glutamic acid (Glu, E), glutamine (Gln, Q), leucine (Leu, L), cysteine (Cys, C), or aspartic acid (Asp, D);  
Xaa23 is valine (Val, V);  
Xaa24 is alanine (Ala, A), glutamine (Gln, Q), cysteine (Cys, C), asparagine (Asn, N), aspartic acid (Asp, D), or glutamic acid (Glu, E);  
Xaa27 is valine (Val, V), leucine (Leu, L), or lysine (Lys, K);  
Xaa28 is cysteine (Cys, C), lysine (Lys, K), alanine (Ala, A), asparagine (Asn, N), or aspartic acid (Asp, D);  
Xaa29 is cysteine (Cys, C), glycine (Gly, G), glutamine (Gln, Q), threonine (Thr, T), glutamic acid (Glu, E), or histidine (His, H);  
Xaa30 is cysteine (Cys, C), glycine (Gly, G), lysine (Lys, K), or histidine (His, H), or is absent; and  
R1 is cysteine (Cys, C), GKKNDWKHNIT (SEQ ID NO: 106), m-SSGAPPPS-n

(SEQ ID NO: 107), or m-SSGQPPPS-n (SEQ ID NO: 108), or is absent;

wherein:

m is -Cys-, -Pro-, or -Gly-Pro-; and

n is -Cys-, -Gly-, -Ser-, or -His-Gly-, or is absent.

2. The isolated peptide of claim 1, wherein Xaa14 is leucine or methionine; and Xaa15 is cysteine, aspartic acid, or leucine.

3. The isolated peptide of claim 1,  
wherein, in General Formula 1,  
Xaa2 is glycine,  $\alpha$ -methyl-glutamic acid, or Aib;  
Xaa7 is threonine,  
Xaa10 is tyrosine, cysteine, or valine;  
Xaa12 is lysine or isoleucine;  
Xaa13 is tyrosine, alanine, glutamine, or cysteine;  
Xaa14 is leucine, tyrosine, or methionine;  
Xaa15 is cysteine, leucine, glutamic acid, or aspartic acid;  
Xaa17 is glutamine, arginine, isoleucine, cysteine, glutamic acid, or lysine;  
Xaa18 is alanine, glutamine, arginine, or histidine;  
Xaa19 is alanine, glutamine, valine, or cysteine;  
Xaa20 is lysine, arginine, or glutamine;  
Xaa21 is glutamic acid, glutamine, leucine, cysteine, or aspartic acid;  
Xaa23 is valine;  
Xaa24 is cysteine, alanine, glutamine, asparagine, glutamic acid, or aspartic acid; and  
Xaa27 is leucine or lysine.

4. The isolated peptide of claim 1, wherein the peptide comprises an amino acid sequence represented by General Formula 2 below:

Xaa1-Xaa2-Gln-Gly-Thr-Phe-Thr-Ser-Asp-Xaa10-Ser-Lys-Xaa13-Xaa14-Xaa15-Xaa16-Xaa17-Xaa18-Xaa19-Xaa20-Xaa21-Phe-Xaa23-Xaa24-Trp-Leu-Leu-Xaa28-Xaa29-Xaa30-Xaa31-Ser-Ser-Gly-Gln-Pro-Pro-Pro-Ser-Xaa40 (General Formula 2, SEQ ID NO: 104)

wherein, in General Formula 2,

Xaa1 is 4-imidazoacetyl, histidine, or tyrosine;

Xaa2 is glycine,  $\alpha$ -methyl-glutamic acid, or Aib;

Xaa10 is tyrosine or cysteine;

Xaa13 is alanine, glutamine, tyrosine, or cysteine;

Xaa14 is leucine, methionine, or tyrosine;

Xaa15 is aspartic acid, glutamic acid, or leucine;

Xaa16 is glycine, glutamic acid, or serine;

Xaa17 is glutamine, arginine, isoleucine, glutamic acid, cysteine, or lysine;

Xaa18 is alanine, glutamine, arginine, or histidine;

Xaa19 is alanine, glutamine, cysteine, or valine;  
Xaa20 is lysine, glutamine, or arginine;  
Xaa21 is cysteine, glutamic acid, glutamine, leucine, or aspartic acid;  
Xaa23 is valine;  
Xaa24 is cysteine, alanine, glutamine, asparagine, or glutamic acid;  
Xaa28 is lysine, cysteine, asparagine, or aspartic acid;  
Xaa29 is glycine, glutamine, cysteine, or histidine;  
Xaa30 is cysteine, glycine, lysine, or histidine;  
Xaa31 is proline or cysteine; and  
Xaa40 is cysteine or is absent.

5. The isolated peptide of claim 1,  
wherein, in General Formula 1,  
Xaa2 is glycine,  $\alpha$ -methyl-glutamic acid, or Aib;  
Xaa7 is threonine;  
Xaa10 is tyrosine, cysteine, or valine;  
Xaa12 is lysine or isoleucine;  
Xaa13 is tyrosine, alanine, or cysteine;  
Xaa14 is leucine or methionine;  
Xaa15 is cysteine or aspartic acid;  
Xaa17 is glutamine, arginine, isoleucine, cysteine, or lysine;  
Xaa18 is alanine, arginine, or histidine;  
Xaa19 is alanine, glutamine, or cysteine;  
Xaa20 is lysine or glutamine;  
Xaa21 is glutamic acid, cysteine, or aspartic acid;  
Xaa23 is valine;  
Xaa24 is alanine, glutamine, cysteine, asparagine, or aspartic acid; and  
Xaa27 is leucine or lysine.
6. The isolated peptide of claim 4,  
wherein, in General Formula 2,  
Xaa13 is alanine, tyrosine, or cysteine;  
Xaa15 is aspartic acid or glutamic acid;  
Xaa17 is glutamine, arginine, cysteine, or lysine;  
Xaa18 is alanine, arginine, or histidine;  
Xaa21 is cysteine, glutamic acid, glutamine, or aspartic acid;  
Xaa23 is valine;  
Xaa24 is cysteine, glutamine, or asparagine;  
Xaa28 is cysteine, asparagine, or aspartic acid;  
Xaa29 is glutamine, cysteine, or histidine; and  
Xaa30 is cysteine, lysine, or histidine.
7. The isolated peptide of claim 1,  
wherein, in General Formula 1,  
Xaa2 is  $\alpha$ -methyl-glutamic acid or Aib;

- Xaa7 is threonine;  
 Xaa10 is tyrosine or cysteine;  
 Xaa12 is lysine or isoleucine;  
 Xaa13 is tyrosine, alanine, or cysteine;  
 Xaa14 is leucine or methionine;  
 Xaa15 is cysteine or aspartic acid;  
 Xaa16 is glutamic acid;  
 Xaa17 is arginine, isoleucine, cysteine, or lysine;  
 Xaa18 is alanine, arginine, or histidine;  
 Xaa19 is alanine, glutamine, or cysteine;  
 Xaa20 is lysine or glutamine;  
 Xaa21 is glutamic acid or aspartic acid;  
 Xaa23 is valine;  
 Xaa24 is glutamine, asparagine, or aspartic acid;  
 Xaa27 is leucine; and  
 Xaa28 is cysteine, alanine, asparagine, or aspartic acid.
8. The isolated peptide of claim 1, wherein, in General Formula 1,  
 Xaa1 is histidine or 4-imidazoacetyl;  
 Xaa2 is  $\alpha$ -methyl-glutamic acid or Aib;  
 Xaa3 is glutamine;  
 Xaa7 is threonine;  
 Xaa10 is tyrosine;  
 Xaa12 is isoleucine;  
 Xaa13 is alanine or cysteine;  
 Xaa14 is methionine;  
 Xaa15 is aspartic acid;  
 Xaa16 is glutamic acid;  
 Xaa17 is isoleucine or lysine;  
 Xaa18 is alanine or histidine;  
 Xaa19 is glutamine or cysteine;  
 Xaa20 is lysine;  
 Xaa21 is aspartic acid;  
 Xaa23 is valine;  
 Xaa24 is asparagine;  
 Xaa27 is leucine;  
 Xaa28 is alanine or asparagine;  
 Xaa29 is glutamine or threonine; and  
 Xaa30 is cysteine or lysine, or is absent.
9. The isolated peptide of claim 1, wherein the peptide comprises an amino acid sequence represented by General Formula 3 below:

Xaa1-Xaa2-Gln-Gly-Thr-Phe-Thr-Ser-Asp-Tyr-Ser-Lys-Xaa13-Leu-Asp-Glu-

Xaa17-Xaa18-Xaa19-Lys-Xaa21-Phe-Val-Xaa24-Trp-Leu-Leu-Xaa28-Xaa29-Xaa30-Xaa31-Ser-Ser-Gly-Gln-Pro-Pro-Ser-Xaa40 (General Formula 3, SEQ ID NO: 105),

wherein, in General Formula 3,  
Xaa1 is histidine or tyrosine;  
Xaa2 is  $\alpha$ -methyl-glutamic acid or Aib;  
Xaa13 is alanine, tyrosine, or cysteine;  
Xaa17 is arginine, cysteine, or lysine;  
Xaa18 is alanine or arginine;  
Xaa19 is alanine or cysteine;  
Xaa21 is glutamic acid or aspartic acid;  
Xaa24 is glutamine or asparagine;  
Xaa28 is cysteine or aspartic acid;  
Xaa29 is cysteine, histidine, or glutamine;  
Xaa30 is cysteine or histidine;  
Xaa31 is proline or cysteine; and  
Xaa40 is cysteine or is absent.

10. The isolated peptide of claim 1, wherein R1 is cysteine, GKKNDWKHNIT (SEQ ID NO: 106), CSSGQPPPS (SEQ ID NO: 109), GPSSGAPPPS (SEQ ID NO: 110), GPSSGAPPPSC (SEQ ID NO: 111), PSSGAPPPS (SEQ ID NO: 112), PSSGAPPPSG (SEQ ID NO: 113), PSSGAPPPSHG (SEQ ID NO: 114), PSSGAPPPSS (SEQ ID NO: 115), PSSGQPPPS (SEQ ID NO: 116), or PSSGQPPPS (SEQ ID NO: 117), or is absent.

11. The isolated peptide according to any one of claims 1 to 10, wherein, in General Formulas 1 to 3, the 16th amino acid and the 20th amino acid from the N-terminus together form a ring.

12. The isolated peptide of claim 1, wherein the peptide comprises an amino acid sequence selected from the group consisting of SEQ ID NOS: 6 to 11, 13, 14, and 20 to 102.

13. The isolated peptide according to any one of claims 1 to 10, wherein the peptide has at least one of the following activities of i) to iii):

- i) activation of a GLP-1 receptor;
- ii) activation of a glucagon receptor; and
- iii) activation of a GIP receptor.

14. The isolated peptide according to any one of claims 1 to 10, wherein the peptide has an increased in vivo half-life compared to that of any one of native GLP-1, native glucagon, and native GIP.

15. The isolated peptide according to any one of claims 1 to 10, wherein the C-terminus of the peptide is amidated.

16. A polynucleotide encoding the peptide according to any one of claims 1 to 10.
17. Use of a peptide according to any one of claims 1 to 10 in the manufacture of a medicament for preventing or treating a metabolic syndrome.
18. The use of claim 17, wherein the metabolic syndrome comprises impaired glucose tolerance, hypercholesterolemia, dyslipidemia, obesity, diabetes, hypertension, arteriosclerosis due to dyslipidemia, atherosclerosis, arteriosclerosis, or coronary heart disease.