

0182152

29,845

This Sheet Has Been Replaced



Title: ANTITUMOR ANTIBIOTICS
(LL-E33288 COMPLEX)

SUMMARY OF THE INVENTION

5 This application is a continuation-in-part of co-
pending application S.N. 672,031, filed November 16, 1984.

This invention relates to new antibacterial and
antitumor agents designated LL-E33288 α_1 -Br, LL-E33288 α_1 -I
LL-E33288 α_2 -Br, LL-E33288 α_2 -I, LL-E33288 α_3 -Br, LL-E33288 α_3 -
I, LL-E33288 α_4 -Br, LL-E33288 β_1 -Br, LL-E33288 β_1 -I, LL-E33288
10 β_2 -Br, LL-E33288 γ -I, LL-E33288 γ_1 -Br, LL-E33288 γ_1 -I and LL-
E33288 δ_1 -I, to their production by fermentation, to methods
for their recovery and concentration from crude solutions and
to processes for their purification. The present invention
includes within its scope the antibacterial and antitumor
15 agents in dilute form, as crude concentrates, as a complex of
various or all components, in pure form as individual compo-
nents and novel strains of Micromonospora.

The LL-E33288 antibiotics of this invention are
closely related compounds. The fourteen antibiotics are
20 recovered from fermentation and are initially obtained as a
mixture, hereinafter either the LL-E33888 complex, the the
LL-E33288 Iodo-complex or the LL-E33288 Bromo-complex. In
general, the iodine containing components of the LL-E33288
antibiotics (e.g., α_1 -I, α_2 -I, α_3 -I, β_1 -I, β_2 -I, γ_1 -I and δ_1 -
25 I) are found only in fermentations using media containing
inorganic or organic iodide while the bromine containing
components (e.g., α_1 -Br, α_2 -Br, α_3 -Br, α_4 -Br, β_1 -Br, β_2 -Br
and γ_1 - Br) are found only in fermentations using media
containing inorganic or organic bromide. The ratio of
30 components in the LL-E3388 complex will vary, depending upon

THIS SHEET HAS BEEN
REPLACED

This Sheet Has Been Replaced



-2-

the fermentation of both the bromine and the iodine containing antibiotics, LL-E33288 β_1 and LL-E33288 γ_1 are the major components, together accounting for approximately 90% of the complex. LL-E33288 α_1 , LL-E33288 α_2 , LL-E33288 α_3 , LL-E33288 α_4 -Br, LL-E33288 δ_2 and LL-E33288 δ_1 -I are minor components, together accounting for approximately 10% of the complex.

The LL-E33288 antibiotics are active against gram-positive and gram-negative bacteria. Each of the components were also found to be active in a modification of the Biochemical Induction Assay [Elespuru, R. and Yarmolinsky, M., Environmental Mutagenesis, 1, 65-78 1979)], a test which specifically measures the ability of an agent to directly or indirectly initiate DNA damage. In this assay, both LL-E33288 δ_1 -Br and LL-E33288 γ_1 -Br were active at concentrations lower than 1×10^{-6} mcg/ml.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. I is the ultraviolet absorption spectra of LL-E33288 δ_1 -Br;

FIG. II is the infrared absorption spectrum of LL-E33288 δ_1 -Br;

FIG. III is the proton magnetic resonance spectrum of LL-E33288 δ_1 -Br;

FIG. IV is the carbon 13 magnetic resonance spectrum of LL-E33288 δ_1 -Br;

FIG. V is the ultraviolet absorption spectra of LL-E33288 γ_1 -Br;

FIG. VI is the infrared absorption spectrum of LL-E33288 γ_1 -Br;

FIG. VII is the proton magnetic resonance spectrum of LL-E33288 γ_1 -Br;

FIG. VIII is the carbon 13 magnetic resonance spectrum of LL-E33288 γ_1 -Br;

FIG. IX is the proton magnetic resonance spectrum of LL-E33288 α_2 -I;

FIG. X is the proton magnetic resonance spectrum of LL-E33288 α_3 -I;

THIS SHEET HAS BEEN
REPLACED

FIG. XI is the ultraviolet absorption spectra of LL-E33288 β ₁-I;

FIG. XII is the infrared absorption spectrum of LL-E33288 β ₁-I;

5 FIG. XIII is the protein magnetic resonance spectrum of LL-E33288 β ₁-I;

FIG. XIV is the carbon 13 magnetic resonance spectrum of LL-E33288 β ₁-I;

10 FIG. XV is the ultraviolet absorption spectra of LL-E33288 γ ₁-I;

FIG. XVI is the infrared absorption spectrum of LL-E33288 γ ₁-I;

FIG. XVII is the proton magnetic resonance spectrum of LL-E33288 γ ₁-I; and

15 FIG. XVIII is the carbon 13 magnetic resonance spectrum of LL-E33288 γ ₁-I.

DETAILED DESCRIPTION OF THE INVENTION

The physico-chemical characteristics of LL-E33288 β ₁-Br and LL-E33288 γ ₁-Br are described below:

20 LL-E33288 β ₁-Br

- 1) Approximate elemental analysis: C 48.6; H 5.6; N 2.9; S 9.1; and Br 5.5. (It has been determined by electron spectroscopy for chemical analysis (ESCA) that only the following elements are present: C, 25 H, N, O, S and Br);
- 2) Melting point: 146-150°C (dec.);
- 3) Specific rotation: $[\alpha]_D^{26} = -49 \pm 10^\circ$ (0.1% ethanol);
- 4) Ultraviolet absorption spectra: as shown in Figure I (methanol; acidic methanol; basic methanol);
- 30 5) Infrared absorption spectrum: as shown in Figure II (KBr disc);
- 6) Proton magnetic resonance spectrum: as shown in Figure III (300 MHz, CDCl₃);
- 7) Carbon-13 magnetic resonance spectrum: as shown 35 in Figure IV (75.43 MHz, CDCl₃, ppm from TMS), significant peaks as listed below:

	17.60(q);	17.64(q);	18.9(q);	19.7(q);
	22.4(q);	22.8(q);	23.5(q);	34.3(t);
	36.9(t);	39.2 (t/d);	47.8 (d);	51.7(q);
	52.7(q);	54.6 (t/d);	56.3(q);	57.2(q);
	57.8(d);	61.0 (q/d);	61.7(d);	62.4 (t);
5	66.9(d);	68.4(d);	69.1(d);	69.7(d);
	70.2(d);	71.1(d);	71.9(d);	72.1 (s/t);
	76.1(d);	81.0(d);	83.3(s);	88.2(s);
	97.4(d);	99.7(d);	100.8(s);	102.5(d);
	115.1(s);	123.4(d);	124.4(d);	126.5(d);
	130.2(s);	130.8(s);	144.6(s);	149.3(s);
	149.5(s);	191.7(s);	192.4(s);	

10

8) Molecular weight: 1333/1335 respectively for $^{79}\text{Br}/^{81}\text{Br}$ as determined by FAB-MS; and

9) Molecular formula: $\text{C}_{54}\text{H}_{84}\text{N}_3\text{O}_{22}\text{S}_4\text{Br}$, exact masses at 1258.3699 (^{79}Br) and 1260.3726 (^{81}Br) was determined by high resolution FAB-MS and calculated to be $\text{C}_{52}\text{H}_{81}\text{N}_3\text{O}_{21}\text{S}_3\text{Br}$ ($\text{M}+\text{H}-\text{C}_2\text{H}_4\text{OS}$).

15

LL-E33288 γ_1 -Br

- 1) Ultraviolet absorption spectra: as shown in Figure V (methanol; acidic methanol; basic methanol);
- 20 2) Infrared absorption spectrum: as shown in Figure VI (KBr disc);
- 3) Proton magnetic resonance spectrum: as shown in Figure VII (300 MHz, CDCl_3);
- 25 4) Carbon 13 magnetic resonance spectrum: as shown in Figure VIII (75.43 MHz, CDCl_3 , ppm from TMS), significant peaks as listed below:

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

This Sheet Has Been Replaced

-5-



	14.4	17.6	17.9	19.0
	19.7	-	22.8	-
	-	34.0	37.6	39.5
	42.1	-	51.6	52.7
	54.1	56.3	57.3	-
	59.3	61.1	61.8	61.9
5	67.2	68.18	68.23	69.7
	70.1	70.8	71.1	71.7
	71.8	76.1	-	81.0
	82.9	88.4	-	97.8
	100.0	100.2	101.3	103.0
	115.3	123.0	124.9	126.9
	130.4	131.1	131.8	138.0
10	144.7	-	149.5	149.6
	155.6	192.5	192.9	

- 5) Molecular formula: $C_{53}H_{82}N_3O_{22}S_4Br$ by comparing its UV, IR, 1H NMR, and ^{13}C NMR data to those of LL-E33288 β $_1$ -BR and LL-E33288 γ $_1$ -I; and
- 6) Molecular weight: 1319/1321 respectively for $^{79}Br/^{81}Br$, calculated from its molecular formula.

The physico-chemical characteristics of LL-E33288 α $_1$ -I, LL-E33288 α $_2$ -I, LL-E33288 α $_3$ -I, and LL-E33288 β $_1$ -I and LL-E33288 γ $_1$ -I are described below:

LL-E33288 α $_1$ -I

- 1) Molecular weight: 1145, determined by FAB-MS.

LL-E33288 α $_2$ -I

- 1) Contains and only contains the following elements by electron spectroscopy for chemical analysis (ESCA): C, H, N, O, S, I;
- 2) Molecular weight: 1131, determined by FAB-MS; and
- 3) Proton magnetic resonance spectrum: as shown in Figure IX (300 MHz, $CDCl_3$).

LL-E33288 α $_3$ -I

- 1) Molecular weight: 1066, determined by FAB-MS; and
- 2) Proton magnetic resonance spectrum: as shown in

THIS SHEET HAS BEEN
REPLACED

Figure X (300 MHz, CDCl_3).LL-E33288 β_1 -I

- 5 1) Ultraviolet absorption spectra: as shown in Figure XI (methanol; acidic methanol; basic methanol);
- 2) Infrared absorption spectrum: as shown in Figure XII (KBr disc);
- 3) Proton magnetic resonance spectrum: as shown in Figure XIII (300 MHz, CDCl_3);
- 10 4) Carbon 13 magnetic resonance spectrum: as shown in Figure XIV (75.43 MHz, CDCl_3 , ppm from TMS), significant peaks as listed below:

	-	17.5	17.6	18.9
	-	22.4	22.8	23.4
15	25.4	34.3	36.9	39.2
	-	47.9	51.6	52.8
	54.8	56.3	57.2	57.9
	60.9		61.6	62.2
	67.0	68.4	68.4	69.1
	69.6	70.4	71.1	71.8
	72.2	76.2	-	80.8
	83.3	88.1	93.6	97.4
20	99.6	99.6	-	102.6
	112.4	123.4	124.4	126.4
	-	-	133.4	-
	-	-	-	-
	-	192.3	192.6	-

- 25 5) Molecular formula: $\text{C}_{54}\text{H}_{84}\text{N}_{30}\text{S}_4\text{I}$ by comparing its UV, IR, ^1H NMR and ^{13}C NMR data to those of LL-E33288 β_1 -Br and LL-E33288 γ_1 -I; and
- 6) Molecular weight: 1381, calculated from molecular formula.

30 LL-E33288 γ_1 -I

- 1) Contains and only contains the following elements by electron spectroscopy for chemical analysis (ESCA): C, H, N, O, S, I;
- 35 2) Approximate elemental analysis: C 48.8; H 5.4; N 2.8; S 9.0; I 9.2;



- 3) Molecular weight: 1367, determined by FAB-MS;
 4) Molecular formula: $C_{53}H_{82}N_3O_{22}S_4I$, exact mass for M+H was determined by high resolution FAB-MS to be 1368. 3397 for $C_{53}H_{83}N_3O_{22}S_4I$;
 5) Ultraviolet absorption spectra: as shown in Figure XV (methanol; acidic methanol; basic methanol);
 6) Infrared absorption spectrum: as shown in Figure XVI (KBr disc);
 7) Proton magnetic resonance spectrum: as shown in Figure XVII (300 MHz, $CDCl_3$); and
 8) Carbon 13 magnetic resonance spectrum: as shown in Figure XVIII (75.43 MHz, $CDCl_3$, ppm for TMS) significant peaks as listed below:

15	14.5(q)	17.6(q)	17.6(q)	18.9(q)
	-	-	22.8(q)	-
	25.4(q)	34.1(t)	37.0(t)	39.1(t)
	42.3(t/s)	-	51.5(d)	52.8(q)
	54.8(t)	56.3(q)	57.2(q)	-
	60.4(d)	60.9(q)	61.3(t)	61.7(q)
	67.0(d)	68.4(d)	68.5(d)	69.2(d)
	69.7(d)	70.5(d)	71.1(d)	71.8(d)
20	72.1(s)	75.7(d)	75.8(d)	80.9(d)
	82.8(s)	88.1(s)	93.5(s)	97.3(d)
	99.6(d)	99.7(d)	100.8(s)	102.6(d)
	-	123.4(d)	124.4(d)	126.2(d)
	130.2(s)	131.0(s)	133.4(s)	139.1(s)
	143.0(s)	145.1	150.6(s)	151.5(s)
	154.5	192.0(s)	192.5(s)	

25

The LL-E33288 components are most conveniently separated and identified by high-performance liquid chromatography (HPLC) and by thin-layer chromatography (TLC). It is difficult, although not impossible, to separate the corresponding iodinated and brominated components by HPLC; however, they cannot be distinguished by TLC.

30

This Sheet Has Been Replaced



THIS SHEET HAS BEEN REPLACED

The preferred analytical separation of the LL-E33288-Br components by HPLC uses the following conditions:

Column: "Separalyte C₁₈ 5 m," 4.6 mm x 25 cm
(Analytichem International);

Solvent: Acetonitrile: 0.2M aqueous ammonium acetate (60:40);

Flow rate: 1.5 ml/minute

Detector: Dual wavelength UV at 254 nm and 280 nm;

Sensitivity: 0-0.02 A.U.F.S.

Table IA gives the approximate retention times and volumes of LL-E33288 β ₁-Br, LL-E33288 β ₂-Br, and LL-E33288 γ ₁-Br under these conditions.

TABLE IA

LL-E33288 Components	Retention Time (minutes)	Retention Volume(ml)
β ₁ -Br	5.7	8.6
β ₂ -Br	7.1	10.7
γ ₁ -Br	4.3	6.5

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

The preferred analytical HPLC separation of the iodine containing LL-E33288 components uses the following conditions:

Column: NOVA-PAK C₁₆ Radial-PAK cartridge with
RCM-100 Radial Compression Module
(Millipore, Waters Chromatography Division);

Solvent: Acetonitrile: 0.2M aqueous ammonium
acetate (50:50);

Flow Rate: 1.2 ml/minute;

Detector: Dual wavelength UV at 254 nm and 280 nm;

Sensitivity: 0-0.02 A.U.F.S.

Table IB gives the approximate retention times and volumes of LL-E33288 α_1 -I, LL-E33288 α_2 -I, LL-E33288 α_3 -I, LL-E33288 β_1 -I, LL-E33288 β_2 -I, LL-E33288 γ_1 -I and LL-E33288 δ_1 -I under these conditions.

TABLE IB

LL-E33288 Components	Retention Time (minutes)	Retention Volume(ml)
α_1 -I	11.9	14.3
α_2 -I	9.1	10.9
α_3 -I	1.5	1.8
β_1 -I	4.4	5.3
β_2 -I	5.0	6.0
γ_1 -I	3.6	4.3
δ_1 -I	2.6	3.1

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

The LL-E33288 components are separated and identified by the following TLC system:

Adsorbant: Silica gel 60 F254 pre-coated aluminum sheets, 0.2mm layer thickness, EM Reagents;

Detection: Visualized by quenching effect under short wavelength UV lamp (254 nm), and bioautography using Bacillus subtilis or the modified biochemical induction assay;

Solvent Systems: I, ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate;

II, 3% isopropyl alcohol in ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate;

III, ethyl acetate:methanol (95.5).

Table II gives the approximate R_f values of LL-E33288 components in these three systems:

TABLE II

LL-E33288 Components	R _f Value		
	Solvent System I	Solvent System II	Solvent System III
α_1 -Br, α_1 -I	0.67	0.80	0.79
α_2 -Br, α_2 -I	0.61	0.75	0.73
α_3 -Br, α_3 -I	0.55	0.69	0.61
α_4 -Br	0.49	0.64	0.54
β_2 -Br, β_2 -I	0.32	0.41	0.45
β_1 -Br, β_1 -I	0.24	0.35	0.36
γ_1 -Br, γ_1 -I	0.18	0.28	0.27
δ_1 -I	0.11	0.19	



The new antibacterial and antitumor agents designated LL-E33288 α_1 -Br, LL-E33288 α_2 -Br, LL-E33288 α_3 -Br, LL-E33288 α_4 -Br, LL-E33288 β_1 -Br, LL-E33288 β_2 -Br, LL-E33288 γ_1 -Br, LL-E33288 α_1 -I, LL-E33288 α_2 -I, LL-E33288 α_3 -I, LL-EE288 β_1 -I, LL-E33288 β_2 -I, LL-E33288 γ_1 -I and LL-E33288 δ_1 -I are formed during the cultivation under controlled conditions of a new strain of Micromonospora echinospora ssp. calichensis. This microorganism is maintained in the culture collection of the Medical Research Division, American Cyanamid Company, Pearl River, New York as culture number LL-E33288. A viable culture of this new microorganism has been deposited with the Culture Collection Laboratory, Northern Regional Research Center, U. S. Department of Agriculture, Peoria, Illinois on August 9, 1984, and has been added to its permanent collection. It has been assigned by such depository the strain designation NRRL 15839. Access to such culture, under strain designation NRRL 15839, during pendency of the instant application shall be available to one determined by the Commissioner of Patents and Trademarks to be entitled thereto under 37 C.F.R. §1.14 and 35 U.S.C. §122, and all restrictions on availability to the public of such culture will be irrevocably removed upon grant of a patent on the instant application.

Culture LL-E33288 was isolated from a caliche clay soil sample collected in Texas.

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

This Sheet Has Been Replaced



-12-

The generic assignment of the strain NRRL 15839 to the genus Micromonospora was confirmed morphologically and chemically. The strain produces monospores either singly or in masses on the vegetative hyphae. No aerial hyphae were observed. Electron microscopic examination showed that the spores were warty. Whole cell analysis showed that the strain contained the meso isomer of diaminopimelic acid (DAP). The 3-OH derivative of DAP was present in large (major) amounts. Additionally the strain showed the presence of xylose plus traces of arabinose in its whole cell sugar hydrolysates (whole cell sugar pattern of Type D).

From macromorphological and physiological studies it was concluded that NRRL 15839 can be considered subspecies of M. echinospora (it is closest to M. echinospora ssp. pallida). Data on the morphology of NRRL 15839 are given in Tables A and B. Physiological data are given in Tables C and D.

Table A

Macromorphology Of NRRL 15839
(Colors Are NBS-ISCC)

25	ISP Agar Medium	Spores	Vegetative Mycelium	Soluble Pigments
	Yeast- Malt (ISP 2)	-	Dark orange- yellow (72)	-
30	Oatmeal (ISP 3)	-	Colorless + pale orange-yellow (73)	-
	Inorganic Salts- Starch (ISP 4)	Slight border of black spores	Dark orange- yellow (72) to lt. yellow-brown (76)	Lt. brownish
35	Glycerol- Asparagine (ISP 5)	-	Pale orange-yellow (73) + colorless	-

THIS SHEET HAS BEEN
REPLACED

Table B

Macromorphology of NRRL-15839 on Various Agar
Media Used for Actinomycete Growth (28°C, 2 weeks)

5	<u>Agar Medium</u>	<u>NRRL-15839</u>
	Pablum	Beige veg. Sl. black spores No sol. pig.
10	Yeast Czapeks	Beige veg. No spores No sol. pig.
	Czapek's	Beige veg. Sl. black spores No sol. pig.
15	Yeast Dextrose	Tan veg. Moderate black sp. Sl. dark pig.
	Nutrient	Colorless to tan veg. Sl. black spores No sol. pig.
20	Nutrient- Glycerol	Colorless to light beige veg. No black spores No sol. pig.
25	Bennett's Dextrin	Colorless to beige veg. Sl. black spores Sl. rosy-brown pig.
	Glucose- Asparagine	Colorless to lt. orange-beige veg. No spores No sol. pig.
30		

veg. = vegetative hyphae; pig. = pigment.

This Sheet Has Been Replaced




THIS SHEET HAS BEEN
REPLACED

Table C

Carbohydrate Utilization of NRRL-15839

5	Arabinose	+
	Cellulose	-
	Fructose	+
10	Glucose	+
	Inositol	-
	Mannitol	-
	Raffinose	±
15	Rhamnose	+
	Sucrose	+
	Xylose	+

This Sheet Has Been Replaced



20

Table D

Physiological Reactions of NRRL-15839

25

Hydrolysis of

	Casein	+
	Xanthine	-
	Hypoxanthine	-
	Tyrosine	+
	Adenine	-
30	Gelatin	+
	Potato Starch	+
	Esculin	+

Production of

	Nitrate	
	Reductase	+
30	Phosphatase	W
	Urease	-

THIS SHEET HAS BEEN
REPLACED

-15-

Growth on

Salicin	-
5% NaCl	-
Lysozyme Broth	-

Decarboxylation of

5	Acetate	+
	Benzoate	-
	Citrate	-
	Lactate	-
	Malate	-
	Mucate	-
	Oxalate	-
	Propionate	+
	Pyruvate	+
10	Succinate	-
	Tartrate	-

Acid from

	Adonitol	-
	Arabinose	+
	Cellobiose	+
15	Dextrin	+
	Dulcitol	-
	Erythritol	-
	Fructose	+
	Galactose	V
	Glucose	+
	Glycerol	-
20	Inositol	-
	Lactose	-
	Maltose	+
	Mannitol	-
	Mannose	+
	α -methyl D	-
	Glucoside	-
25	Melibiose	-
	Raffinose	+
	Rhamnose	+
	Salicin	+
	Sorbitol	-
	Sucrose	+
	Trehalose	+
	Xylose	+
30	β -Methyl	-
	D-xyloside	-

Growth at

100	-
420	+
450	+

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

35

+ = positive; - = negative; V = variable;
W = weak.

Derivation of Mutant LL-E33288-R66, NRRL-15975

5 In an effort to improve fermentation yields, the original culture LL-E33288 (NRRL-15839) was plated and 50 single colonies were isolated. These were designated NS1 to NS50 (NS = natural selection).

Fermentation of these isolates showed that those with moderate sporulation were generally better producers of LL-E33288 complex. Selected as representative of this group was isolate NS6.


10 Using isolate NS6 as the starting culture, spore suspensions were prepared and exposed to various mutagens. Single colonies were isolated from a nitroso-guanidine treatment, but none proved to be significantly improved producers of the LL-E33288 complex. From a subsequent series
15 of exposures to ultraviolet irradiation, single colonies were obtained from which isolate UV 610 was selected as a high yielding mutant. Isolate UV 610 was then streaked and sub-isolates 1 to 7 were obtained. Sub-isolate UV 610(3) was selected for further work.

20 Because of the highly potent antibacterial and antineoplastic nature of the LL-E33288 complex it is possible that once a limited concentration of the antibiotic is biosynthesized in the fermentation it may become toxic/inhibitory to the producing culture. Thus, an effort was made
25 to obtain isolates which are resistant to the LL-E33288 antibiotic complex.


Vegetative growth from isolate UV 610(3) was prepared as employed for fermentation and used to inoculate a flask of medium consisting of peptone, dextrose, molasses
30 and water. The medium was supplemented with LL-E33288 β_1 -Br at a concentration of 8 μ g/m. A number of platings were done from this flask and a resistant population was obtained on the seventh day. A total of 97 colonies (R1 to R97) were isolated. Isolate R66 was selected
35 as a potentially improved producer of LL-E33288 β_1 -Br.

The history is represented schematically below.

THIS SHEET HAS BEEN
REPLACED



This Sheet Has Been Replaced



30 The mutant R66 is maintained in the culture collection of the Medical Research Division, American Cyanamid Company, Pearl river, New York as culture number LL-E33288 R66. A viable culture of this new microorganism has been deposited with the Culture Collection Laboratory, Northern Regional Research center, U. S. Department of Agriculture, Peoria, Illinois on June 6, 1985, and has been added to its permanent collection. It has been assigned by such depository the strain designation NRRL 15975. Access to such culture, under strain designation NRRL 15975, during pendency of the instant application

35

THIS SHEET HAS BEEN
REPLACED

shall be available to one determined by the Commissioner of Patents and Trademarks to be entitled thereto under C.F.R. §1.14 and 35 U.S.C. §122, and all restrictions on availability to the public of such culture will be irrevocably removed upon grant of a patent on the instant application.

Morphologically, NRRL-15975 forms fewer spores than NRRL-15839. A comparison of NRRL-15975 with NRRL-15839 is given in Table DD.

Chemically, both NRRL-15839 and NRRL-15975 show the same whole cell sugar patterns (Type D: xylose and traces of arabinose). The whole cell diaminopimelic acid analysis reveals that 15975 does not form the meso isomer but only the 3-hydroxy derivative (NRRL-15839 contains both compounds). This does not change the chemo-taxonomic assignment.

Physiological tests show that NRRL-15839 and NRRL-15975 differ in only two physiological reaction (See Table D). NRRL-15975 is negative for nitrate reductase and positive for utilization of lactate. NRRL-15839 was weakly positive for both, but is now negative after having been maintained on slants for a few months. Thus these characters should be considered variable for this taxon.

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

TABLE DD

Morphological Comparison of NRRL 15839 and NRRL 15975

Agar Medium		NRRL 15839	NRRL 15975
Bennett's-Dextrin	V ¹	Beige-tan	Beige tan
	Sp	Black, copious	None
	SS	None	None
Czapek's	V	Orange-tan	Orange-tan
	Sp	Black, traces	None
	SS	None	None
Yeast Extract-Czapek's	V	Orange tan, flat	Orange tan, convoluted
	Sp	Black, traces	None
	SS	None	Slight yellowish
Potato-Dextrose	V	Very poor growth	Very poor growth
	Sp	None	None
	SS	None	None
Nutrient glycerol	V	Tan	Tan
	Sp	Black, sparse	Black, sparse
	SS	None	Slight brownish
Nutrient	V	Tan	Tan
	Sp	Black, fair	None
	SS	None	None

1 = V = vegetative hyphae; Sp = spores; SS = soluble pigment.

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

5 It is to be understood that for the production
of these new antibacterial and antitumor agents the
present invention is not limited to this particular organ-
isms or to organisms fully answering the above growth
microscopic characteristics which were given for illus-
trative purposes only. In fact, it is desired and intended
to include the use of mutants produced from these organisms
by various means such as exposure to X-radiation, ultra-
violet radiation, N'-methyl-N'-nitro-N-nitrosoguanidine,
10 actinophages and the like.

The in vitro antibacterial activity of LL-
E33288 components was determined against a spectrum of
gram-positive and gram-negative bacteria by a standard
agar dilution method. Mueller-Hinton again containing
15 two-fold decreasing concentrations of the antibiotics
were poured into petri plates. the agar surfaces were
inoculated with 1 to 5 x 10⁴ colong forming units of
bacteria by means of the steers replicating device.
The lowest concentration of LL-E33288 component that
20 inhibited growth of a bacterial strain after about 18
hours of incubation at approximately 35°C was recorded
as the minimal inhibitory concentration (MIC) for that
strain. The results are summarized in Table III.

25
This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED



TABLE III
In vitro Antibacterial Activity of LL-E33288 Components

Organism	Minimal Inhibitory Concentration, mcg/ml			
	β l-Br	β l-I	γ l-Br	γ l-I
<u>Escherichia coli</u>	0.25	0.50	0.50	0.25
"	0.12	0.25	0.25	0.25
"	0.12	0.25	0.25	0.25
<u>Klebsiella pneumoniae</u>	0.25	0.50	0.50	0.25
"	0.12	0.50	0.50	0.25
<u>Enterobacter cloacae</u>	0.5	0.50	0.50	0.50
"	0.25	0.25	0.50	0.50
<u>Serratia marcescens</u>	0.12	0.25	0.50	0.25
"	0.12	0.50	0.25	0.12
<u>Morganella morganii</u>	0.5	1	0.50	0.25
<u>Providencia stuartii</u>	0.25	0.50	1	0.25
<u>Citrobacter diversus</u>	0.12	0.50	0.50	0.25
"	0.12	0.25	0.25	0.12
<u>Acinetobacter sp.</u>	0.06	0.25	0.25	0.12
"	0.12	0.25	0.25	0.06

THIS SHEET HAS BEEN
REPLACED

TABLE III (continued)

Organism	Minimal Inhibitory Concentration, mcg/ml			
	β 1-Br	β 1-I	γ 1-Br	γ 1-I
<u>Pseudomonas aeruginosa</u>	0.5	0.50	1	0.25
"	0.25	0.25	0.50	0.12
<u>Staphylococcus aureus</u>	≤ 0.0025	≤ 0.000031	≤ 0.000031	≤ 0.000031
"	≤ 0.00025	≤ 0.000031	≤ 0.000031	≤ 0.000031
"	≤ 0.00025	≤ 0.000031	≤ 0.000031	≤ 0.000031
"	≤ 0.00025	≤ 0.000031	≤ 0.000031	≤ 0.000031
"	≤ 0.00025	≤ 0.000031	≤ 0.000031	≤ 0.000031
"	≤ 0.00025	≤ 0.000031	≤ 0.000031	≤ 0.000031
"	≤ 0.00025	≤ 0.000031	≤ 0.000031	≤ 0.000031
<u>Staphylococcus epidermidis</u>	≤ 0.00025	≤ 0.000031	≤ 0.000031	≤ 0.000031
"	≤ 0.00025	≤ 0.000031	≤ 0.000031	≤ 0.000031
<u>Enterococcus</u> sp.	0.0038	0.031	0.062	0.0078
<u>Streptococcus faecalis</u>	≤ 0.00025	0.00012	≤ 0.000031	0.00012
<u>Micrococcus luteus</u>	≤ 0.00025	≤ 0.000031	≤ 0.000031	≤ 0.000031
<u>Bacillus subtilis</u>	≤ 0.00025	≤ 0.000031	≤ 0.000031	≤ 0.000031

This Sheet Has Been Replaced

THIS SHEET HAS BEEN
REPLACED

5 Certain in vivo testing systems and protocols have
been developed by the National Cancer Institute for testing
compounds to determine their suitability as anti-neoplastic
agents. These have been reported in "Cancer Chemotherapy
10 Reports", Part III, Vol. 3, No. 2 (1972), Geran, et al. These
protocols have established standardized screening tests
which are generally followed in the field of testing for
antitumor agents. Of these systems, lymphocytic leukemia

10
This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

P388, melanotic melanoma B16, L1210 leukemia and colon 26 adenocarcinoma are particularly significant to the present invention. These neoplasms are utilized for testing as transplantable tumors in mice. Generally, significant anti-tumor activity, shown in these protocols by a percentage increase of mean survival times of the treated animals (T) over the control animals (C), is indicative of similar results in human leukemias and solid tumors.

Lymphocytic Leukemia P388 Test

The animals used were BDF₁ mice, all of one sex, weighing a minimum of 17 g and all within a 3 g weight range. There were 5 or 6 mice per test group. The tumor transplant was by intraperitoneal injection of 0.5 ml of dilute ascitic fluid containing 10⁶ cells of lymphocytic leukemia P388. LL-E33288 antibiotics were tested in the P388 system both as the individual β ₁-Br and γ -Br components and as a complex of all components (Bromo-complex). The test compounds were administered intraperitoneally at a volume of 0.5 ml in 0.2% Klucel in normal saline on days 1, 5 and 9 (relative to tumor inoculation) at the indicated doses. The mice were weighed and the survivors recorded on a regular basis for 30 days. The median survival time and the ratio of survival time for treated (T)/control (C) animals were calculated. The positive control compound was Cisplatin given as an intraperitoneal injection in 0.5 ml of 0.2% Klucel on days 1, 5 and 9 at the indicated doses. The results appear in Table IV.

If $T/C \times 100$ (%) is 125 or over, the tested compound is considered to have significant anti-tumor activity.

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

TABLE IV
Lymphocytic Leukemia P388 Test

Compound	Dose (mg/kg)	Median Survival (Days)	T/C x 100 (%)
LL-E33288 (Bromo-complex)	3.2	16.5	156
	1.6	17.5	165
	0.8	18.5	175
	0.4	19	179
	0.2	16.5	156
Control	-	10.6	-
Positive Control	1.0	21.5	203
	0.25	15	142
	0.06	14.5	137
LL-E-33288 ₁ -Br	0.4	13	105
	0.2	18	145
	0.1	19	153
	0.05	17.5	141
	0.025	18	145
	0.012	14	113
Control	-	12.4	-
Positive Control	1.0	25.5	206
	0.4	19	153
	0.06	15	121

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

TABLE IV (continued)

Compound	Dose (mg/kg)	Median Survival (Days)	T/C x 100 (%)
LL-E33288 γ_1 -Br	0.2	14	113
	0.1	21	169
	0.05	19.5	157
	0.025	18	145
	0.012	14.5	117
Control	-	12.4	-
Positive Control	1.0	25.5	206
	0.4	19	153
	0.06	15	121

Melanotic Melanoma B16

The animals used were BDF₁ mice, all of the same sex, weighing a minimum of 17 g and all within a 3 g weight range. There are normally 6 animals per test group. A 1 g portion of melanotic melanoma B₁₆ tumor was homogenized in 10 ml of cold balanced salt solution and a 0.5 ml aliquot of the homogenate was implanted intraperitoneally into each of the test mice. LL-E33288 antibiotics were tested in the B₁₆ system both as the individual β_1 -Br and γ_1 -Br components and as a complex of all components (Bromo-complex). The test compounds were administered intraperitoneally on days 1 through 9 (relative to tumor inoculation) at various doses. The mice were weighed and survivors recorded on a regular basis for 60 days. The median survival time and the ratio of survival time for treated (T)/control (C) animals were calculated. The positive control compounds were Cisplatin or Adriamycin. The results of this test appear in Table V. If T/C X 100 (%) is 125 or over, the tested compound is considered to have significant anti-tumor activity.



TABLE V
Melanotic Melanoma B₁₆ Test

Compound	Dose (mg/kg)	Median Survival (Days)	T/C x 100 (%)
LL-E33288 (Bromo-complex)	0.8	30	188
	0.4	29.5	184
	0.2	27	169
	0.1	24	150
Control	-	16	-
Cisplatin	0.4	25	156
	0.2	25	156
	0.1	23	144
	0.05	21.5	134
LL-E33288 β_1 -Br	0.05	32	168
	0.025	33.5	176
	0.0125	32	168
Control	-	19	-
Adriamycin	0.8	> 60	> 316
	0.4	> 60	> 316

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

TABLE V (continued)

Compound	Dose (mg/kg)	Median Survival (Days)	T/C x 100 (%)
LL-E33288 γ ₁ -Br	0.05	33.5	176
	0.025	30	159
	0.0125	37	195
Control	-	19	-
Adriamycin	0.8	> 60	> 316
	0.4	> 60	> 316

Lymphocytic Leukemia L1210 Test

The animals used were BDF₁ mice, all of one sex, weighing a minimum of 17 g and all within a 3 g weight range. There were 6 mice in each test group and 18 in control groups. The tumor transplant was by intraperitoneal injection of 0.5 ml of lymphocytic leukemia L1210 at a concentration of 10⁵ cells per mouse. LL-E33288 antibiotics were tested in the L1210 system both as the individual β ₁-Br component and as a complex of all components (Bromo-complex). The test compounds were administered on days 1, 5 and 9 or days 1 through 9 (relative to tumor inoculation) at various doses. The mice were weighed and survivors recorded on a regular basis for 30 days. The median survival time and the ratio of survival time for treated (T)/control (C) animals were calculated. The positive control compound was 1,4-dihydroxy-5,8-bis[[2-(2-hy-



droxyethylamino)ethyl]amino]anthraquinone dihydrochloride or Cisplatin given intraperitoneally at the indicated dose. The results appear in Table VI. If T/C X 100 (%) is 125 or over, the tested compound is considered to have significant anti-tumor activity.

TABLE VI

Lymphocytic Leukemia L1210 Test

Compound	Dose (mg/kg)	Median Survival (Days)	T/C x 100 (%)
LL-E33288 (Bromo-complex)	1.5	29	174
	0.8	28.5	171
	0.4	22.5	135
	0.2	22.5	135
Control	-	16.7	-
Anthraquinone	1.6	30	180
	0.8	30	180
	0.4	30	180
LL-E33288 β_1 -Br	0.2	11.3	136
	0.1	11.4	137
	0.05	11	133
	0.025	11.3	136
Control	-	8.3	-
Cisplatin	5	7.5	90
	2.5	12	145
	1.25	11	133

This Sheet Has Been Replaced



THIS SHEET HAS BEEN REPLACED

This Sheet Has Been Replaced



-30-

Colon 26 Adenocarcinoma Test

The animals used were CD₂F₁ female mice weighing a minimum of 17 g and all within a 3 g weight range. There were 5 or 6 mice per test group with three groups of 5 or 6 animals used as untreated controls for each test. The tumor implant was by intraperitoneal (or subcutaneous) injection of 0.5 ml of a 2% Colon 26 tumor brei in Eagle's MEM medium containing antibiotics. LL-E33288 antibiotics were tested in the Colon 26 system as a complex (Bromo-complex) of all components. The test compounds were administered intraperitoneally on days 1, 5 and 9 (relative to tumor implant doses). The mice were weighed and deaths recorded on a regular basis for 30 days. The median survival times for treated (T)/control (C) animals were calculated. The positive control compound was Cisplatin. The results appear in Table VII. If $T/C \times 100 (\%)$ is 130 or over, the tested compound is considered to have significant anti-tumor activity.

TABLE VIIColon 26 Adenocarcinoma Test

Compound	Dose (mg/kg)	Median Survival (Days)	T/C x 100 (%)
LL-E33288 (Bromo-complex)	1.5	39.5	212
	0.8	32.5	175
	0.4	34	183
	0.2	25.5	137
Control	-	18.6	-
Positive Control	1	29	156
	0.5	38.5	207
	0.25	37.5	202

THIS SHEET HAS BEEN
REPLACED

M5076 Sarcoma

The M5076 reticular cell Sarcoma is propagated as subcutaneous implants in C57B2/6 female mice. In the assays for antitumor activity, BDF₁ mice of either sex were inoculated intraperitoneally with 0.5 ml of a 10% tumor brei. LL-E33288 antibiotics were tested in the M5076 system as a complex (Bromo-complex) of all components. Test compounds were administered intraperitoneally on days 1, 5, 9, 13 and 17 relative to tumor inoculation on day zero. The median survival time in days was determined for each drug dose used on day 60 and the ratio of survival time for treated (T)/control (C) animals were calculated.

The results of this test appear in Table VIII compared to the results obtained with Cisplatin. If T/C x 100 (%) is 125 or over, the tested compound is considered to have significant anti-tumor activity.

TABLE VIIIM5076 Sarcoma

Compound	Dose (mg/kg)	Median Survival (Days)	T/C x 100 (%)
LL-E33288 (Bromo-complex)	1.5	50	175
	0.8	50	175
	0.4	39.5	139
Control	-	28.5	-
Cisplatin	1	30	105
	0.5	44.5	156
	0.25	45	158

This Sheet Has Been Replaced

THIS SHEET HAS BEEN
REPLACED

This Sheet Has Been Replaced



-32-

In the same manner, the following iodo-components were tested for antineoplastic activity.

TABLE IX
Lymphocytic Leukemia P388 Test

5

10

15

20

25

30

35

Compound	Dose (mg/kg)	Median Survival (Days)	T/C x 100 (%)
LL-E33288 γ 1-I (Test 1)	.005	25.5	196
	.0025	22	169
	.00125	18.5	142
	.0006	18	138
	.0003	15.5	119
	.00015	15	115
Control	-	13	-
Positive Control Novantrone®*	1.6	22.5	173
LL-E33288 γ 1-I (Test 2)	.01	11	100
	.005	18	164
	.0025	22.5	205
	.00125	18.5	168
	.0006	16	145
	.0003	14	127
	.00015	14	127
Control	-	11	-
Positive Control Novantrone®	1.6	19	173
	0.8	16	145

*1,4-dihydroxy-5,8-bis[[2-(2-hydroxyethylamino)ethyl]
amino]anthraquinone-2HCl

THIS SHEET HAS BEEN
REPLACED

0182152

-33-

This Sheet Has Been Replaced

TABLE XMelanotic Melanoma B₁₆ Test

Compound	Dose (mg/kg)	Median Survival (Days)	T/C x 100 (%)
LL-E33288 γ 1-I	.0025	26.5	156
	.00125	27	159
	.0006	42.5	250
	.0003	35.5	209
	.00015	33.5	197
	.00007	30.5	179
Control	-	-	-
Adriamycin	0.8	48	282
	0.4	40	235
	0.2	34.5	203

THIS SHEET HAS BEEN
REPLACED

0182152

-34-

TABLE XILymphocytic Leukemia L1210 Test

Compound	Dose (mg/kg)	Median Survival (Days)	T/C x 100 (%)
LL-E33288 γ 1-I	.01	8	89
	.005	14	156
	.0025	11	122
	.0012	10.5	117
	.0006	10	111
Control	-	9	-
Positive Control Novantrone®	3.2	15	167
	1.6	11.5	128
	0.8	12	133
	0.4	11	122

This Sheet Has Been Replaced

THIS SHEET HAS BEEN
REPLACED

0182152

-35-

TABLE XIIColon 26 Adenocarcinoma Test

Compound	Dose (mg/kg)	Median Survival (Days)	T/C x 100 (%)
LL-E33288 γ_1 -I	.01	11	59
	.005	25.5	138
	.0025	27	146
	.00125	22.5	122
	.0006	23.5	127
	.0003	20	108
	.00015	17.5	95
	.00007	17	92
Control	-	18.5	-
Positive Control Cisplatin	2	15.5	84
	1	27.5	149
	0.5	23.5	127

This Sheet Has Been Replaced

THIS SHEET HAS BEEN
REPLACED

General Fermentation Conditions

Cultivation of Micromonospora echinospora NRRL 15839 or NRRL 15975 may be carried out in a wide variety of liquid culture media. Media which are useful for the production of these novel antibacterial and antitumor agents include an assimilable source of carbon, such as starch, sugar, molasses, glycerol, etc.; an assimilable source of nitrogen such as protein, protein hydrolysate, polypeptides, amino acids, corn steep liquor, etc.; and inorganic anions and cations, such as potassium, sodium, ammonium, calcium, sulfate, carbonate, phosphate, chloride, etc. and sources of either bromine (sodium bromide) or iodine (potassium iodide). Trace elements such as boron, molybdenum, copper, etc., are supplied as impurities of other constituents of the media. Aeration in tanks and bottles is supplied by forcing sterile air through or onto the surface of the fermenting medium. Further agitation in tanks is provided by a mechanical impeller. An antifoam agent such as silicone may be added as needed.

General Procedure for the Isolation and Separation of the Antibiotics - LL-E33288

The LL-E33288 antibiotics are recovered from the fermentation broth by extracting the whole mash with an organic solvent such as ethyl acetate or dichloromethane. The antibiotic complex, contained in the organic extract, is further purified by selective precipitation from lower hydrocarbons. The crude LL-E-33288 antibiotic complex thus obtained is further purified and separated into the individual components by a series of column chromatographies using silica gel, Sephadex® LH-20 (Pharmacia Fine Chemicals) and C₁₈ bonded silica.

The invention will be described in greater detail in conjunction with the following non-limiting specific examples.

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

This Sheet Has Been Replaced

Example 1Inoculum Preparation

5 A typical medium used to grow the primary inoculum was prepared according to the following formula:

	Beef extract.....	about 0.3%
	Tryptone.....	about 0.5%
	Dextrose.....	about 0.5%
10	Dextrin.....	about 2.4%
	Calcium carbonate.....	about 0.4%
	Yeast extract.....	about 0.5%
	Water.....qs to.....	100%

15 This medium was adjusted to pH 7.0 and then sterilized. A 100 ml portion of this sterile medium, in a flask, was inoculated with frozen mycelia of the culture

NRRL 15839. The inoculated medium was placed on a rotary shaker and agitated vigorously for 48 hours at 32°C. This incubated medium was then used to inoculate 10
20 liters of the above sterile medium in a 14 liter fermentor. This medium was incubated, with agitation, at 32°C for 48 hours, providing secondary inoculum. This secondary inoculum was then used to inoculate 300 liters of the above sterile medium in a tank and incubated for 48 hours at 30°C
25 while agitated by an impeller driven at 180-200 rpm, providing the tertiary or seed inoculum.

Example 2Tank Fermentation

30 A fermentation medium was prepared according to the following formulation:

	Dextrose.....	about 0.5%
	Sucrose.....	about 1.5%
	Peptone, bacteriological grade.....	about 0.2%
35	Dibasic potassium phosphate.....	about 0.01%

THIS SHEET HAS BEEN
REPLACED

Molasses..... about 0.5%
Calcium carbonate..... about 0.5%
Source of bromine or iodine..... trace amounts
Water.....qs to..... 100%

5 A 2800 liter portion of the above medium was steri-
lized and then inoculated with 300 liters of tertiary (seed)
inoculum prepared as described in Example 1. Aeration was
supplied at the rate of 0.53 liters of sterile air per liter
10 of mash per minute and agitation was supplied by an impeller
driven at 110 rpm. The temperature was maintained at about
28°C and the fermentation was terminated after about 97 hours,
at which time the mash was harvested.

15 The fermentation was monitored for production of
the LL-E33288 antibiotics by antibacterial activity, bio-
chemical induction assay, TLC and HPLC analyses.

20 The whole harvest mash was adjusted to pH 6 and then
extracted with 1/2 mash volume ethyl acetate. The ethyl
acetate extract was concentrated to a syrup which was washed
twice with hexane and filtered through diatomaceous earth.
The diatomaceous earth cake was thoroughly mixed with ethyl
acetate and filtered. The filtrate was concentrated to 3
liters, dried over excess anhydrous sodium sulfate and then
precipitated by the addition of hexane giving about 26.7 g of
crude LL-E33288 complex.

Example 3

Separation of LL-E33288 α_1 -Br, α_2 -Br, α_3 -Br and α_4 -Br from

LL-E33288 β_1 -Br, β_2 -Br and γ_1 -Br

30 The approximately 26.7 g of crude LL-E33288 complex
(Bromo-complex) from Example 2 was divided evenly into three por-
tions and chromatographed on three separate 2.4 x 110 cm silica
gel columns (Silica Woelm®, 32-63 m, Woelm Pharma) packed and
equilibrated with ethyl acetate saturated with 0.1M aqueous
potassium dihydrogen phosphate. The columns were first
35 eluted with the same solvent at a flow rate of 3 ml/minute



for 18 hours, collecting 18 ml fractions. The eluent was changed to ethyl acetate:methanol (95:5) and elution continued for 8 hours. Finally the columns were eluted with ethyl acetate:methanol (90:10) for 10 hours. The fractions were assayed by the modified biochemical induction assay (BIA). The positive fractions were analysed by TLC using silica gel 60 precoated sheets and developed with the solvent system 3% isopropanol in ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate and detected by bioautography using the modified BIA.

Fractions containing LL-E33288 α_1 -Br, α_2 -Br, α_3 -Br and α_4 -Br (LL-E33288 α -Br complex) from the three columns were pooled, concentrated to dryness and the residue was dissolved in ethyl acetate and washed with a small amount of water. The ethyl acetate solution was dried over anhydrous sodium sulfate and precipitated as before to yield about 4.2 g of crude LL-E33288 α -Br complex.

Fractions containing LL-E33288 β_2 -Br, β_1 -Br and γ_1 -Br (LL-E33288 β -Br complex containing γ -Br from the three columns were pooled and worked up as above to yield about 2.0 g of crude LL-E33288 β -Br complex containing γ -Br.

Example 4

Isolation of LL-E33288 β_1 -Br and LL-E33288 γ_1 -Br

An approximately 1.9 g sample of the LL-E33288 β -Br complex containing γ -Br from Example 3 was chromatographed on a 25 x 10 cm Sephadex[®] LH-20 column equilibrated with methanol:water (90:10) at a flow rate of 1.2 ml/minute, collecting 15 ml fractions. The fractions were assayed in the BIA and those active were analysed by TLC as before. Fractions 21-26 containing most of the LL-E33288 β_1 -Br, β_2 -Br and γ_1 -Br were pooled and concentrated to remove methanol and the resulting aqueous mixture was lyophilized to yield about 435 mg of partially purified complex containing approximately 10% of LL-E33288 β_1 -Br, 1% of LL-E33288 β_2 -Br and 4% of LL-E33288 γ_1 -Br.

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED



5 The above partially purified LL-E33288 β -Br complex
containing γ -Br was divided evenly and chromatographed on two
1.5 x 100 cm silica gel columns (Kiesel Gel 60, 40-63 μ m, EM
Products for chromatography) packed and equilibrated with
10 ethyl acetate:methanol (98:2) at a flow rate of 1 ml/minute,
collecting 12 ml fractions. The fractions were assayed and
analysed by TLC as before and those containing primarily LL-
E33288 β_1 -Br were pooled, concentrated and precipitated from
hexane to yield about 26 mg of 80% pure LL-E33288 β_1 -Br. Those
15 fractions containing LL-E33288 γ_1 -Br (chromatographing just
after LL-E33288 β_1 -Br) were pooled and worked up to yield
about 4.5 mg of 30% pure LL-E33288 γ_1 -Br. A few fractions
containing LL-E33288 β_2 -Br (chromatographing just before LL-
E33288 β_1 -Br), were pooled and worked up to yield a trace
amount of LL-E33288 β_2 -Br.

Example 5

Final Purification of LL-E33288 β_1 -Br

20 The approximately 26 mg of 80% pure LL-E33288 β_1 -
Br from Example 4 was combined with other LL-E33288 β_1 -Br
samples of similar purity derived from other fermentations
conducted under identical conditions. A total of about 38 mg
of this combined β_1 -Br was further purified by reverse phase
preparative TLC using Whatman PLKC18F, 100 m precoated TLC
25 plates, developed with methanol:0.1M ammonium acetate buffer
at pH 4.0 (90:10). The band containing LL-E33288 β_1 -Br,
chromatographing at R_f =0.66 and visualized by quenching
effect under short wavelength UV lamp (254 nm), was excised
and the antibiotic was washed off the adsorbant with 10%
isopropyl alcohol in ethyl acetate saturated with 0.1M aque-
30 ous potassium dihydrogen phosphate. The solution was con-
centrated and the residue was dissolved in ethyl acetate and
washed with a small amount of water. The organic solution
containing LL-E33288 β_1 -Br was worked up as before to yield
about 24.5 mg of 90% pure LL-E33288 β_1 -Br. This sample was
35 further purified by preparative TLC on silica gel (Silica

Gel GF precoated plates, 1000 m, Analtech) developed with 3% isopropyl alcohol in ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate. The major quenching band under short wavelength UV lamp (254 nm), chromatographing at $R_f=0.7$, was excised and the antibiotic was washed off the adsorbant with dichloromethane:methanol (80:20). The organic solution containing LL-E33288 β_1 -Br was worked up as before to yield about 18.8 mg of substantially pure LL-E33288 β_1 -Br.

Example 6

Final Purification of LL-E33288 β_1 -Br

The approximately 4.5 mg of 30% pure LL-E33288 γ_1 -Br from Example 4 was combined with other LL-E33288

γ_1 -Br samples of similar purity derived from other fermentations conducted under identical conditions. A total of 18 mg of this combined sample was further purified by preparative TLC on silica gel (Silica Gel GF precoated tapered plates, Analtech) developed with 2% isopropyl alcohol in ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate. The major quenching band under short wavelength UV lamp (254 nm), chromatographing at $R_f=0.5$, was excised and worked up as before to yield about 4.3 mg of substantially pure LL-E33288 γ_1 -Br.

A preferred fermentation medium for production of LL-E33288 Bromo-complex is as follows:

	<u>Ingredient</u>	<u>Percent</u>
	Sucrose	2.0
	Ferrous Sulfate Heptahydrate	0.01
30	Magnesium Sulfate Heptahydrate	0.02
	Calcium Carbonate	0.5
	Peptone	0.2
	Molasses	0.5
	Sodium Bromide	0.05
35	Water qs to	100

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

However, the addition of iodine as potassium iodide, to the fermentation medium provided substantial improvements by:

- 5 1) markedly enhancing vegetative growth in the fermentation;
- 2) increasing zones of inhibition in bioassays versus Escherichia coli #300 and Bacillus subtilis #308;
- 10 3) providing substantial activity at the Rf of LL-E33288₁-I and γ 1-I on the bioautography of TLC plates; and
- 4) enhancement of other components as detected by TLC.

15 The following two media are preferred for the production of LL-E33288 Iodo-complex:

	<u>Ingredient</u>	<u>Percent</u>	
		<u>Media A</u>	<u>Media B</u>
	Sucrose	2.0	2.0
20	Ferrous Sulfate Heptahydrate	0.01	0.01
	Magnesium Sulfate Heptahydrate	0.02	0.02
	Calcium Carbonate*	0.5	0.25
	Peptone**	0.2	0.2
	Molasses	0.5	0.5
25	Potassium iodide	0.05	0.01
	Water qs to	100	100

* Mississippi lime.

30 ** Best results were obtained with MARCOR® bacteriological peptone, but other peptones usable and also polypeptides from meat and casein hydrolyzates.

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

This Sheet Has Been Replaced



-43-

Example 7

A mycelial-spore suspension was prepared by scraping the surface of a slant of culture NRRL-15839 to which 5 ml of sterile distilled water had been added. This suspension was then used to inoculate 100 ml of sterile seed medium of the following formula:

Yeast Extract	0.5%
Beef Extract	0.3%
Tryptose	0.5%
Starch	2.4%
Dextrose	0.5%
Calcium Carbonate	0.4%
Water qs to	100.0%

in a 500 ml flask. This seed flask was incubated at 28°C on a rotary shaker at 200 rpm for 3-4 days, producing Stage I inoculum.


The Stage I inoculum was used to inoculate a Stage II inoculum of the same sterile medium, which was incubated under the same conditions for 2 days.

The Stage II inoculum was then used to inoculate 100 ml of sterile fermentation medium of the formula:

Sucrose	2.0%
Ferrous Sulfate Heptahydrate	0.01%
Magnesium Sulfate Heptahydrate	0.02%
Calcium Carbonate	0.5%
Peptone (MARCOR®)	0.2%
Molasses	0.5%
Potassium Iodide	0.05%
Water qs to	100.0%

This medium was incubated at 28°C on a shaker at 200 rpm for 5 days at which time the mash was harvested.

THIS SHEET HAS BEEN
REPLACED



A concentration of 4 to 20 $\mu\text{g/ml}$ of potassium iodide appears to be optimal, but concentrations of 2 mg/ml do not appear to depress yields.

5 NRRL-15839 can be induced to produce LL-E33288 β_1 -I when potassium iodide is present in the medium, but only at very low levels (0.2-0.3 $\mu\text{g/ml}$) as against 1.5-3.5 $\mu\text{g/ml}$ for the better producing NRRL-15975.

10 Yields of β_1 -I and γ_1 -I in an iodine medium are 2 to 8 times greater than yields of corresponding brominated compounds β_1 -Br and γ_1 -Br in a bromine medium using NRRL-15975.

Example 8

Separation of LL-E33288 α_1 -I, α_2 -I, and α_3 -I from LL-E33288 β_1 -I, β_2 -I, γ_1 -I and δ_1 -I

15 Approximately 41.3 g of crude LL-E33288 complex derived from the processing 7500 liters of a fermentation using NRRL-15975 and medium containing inorganic iodide was divided evenly into two portions and chromatographed on two separate 2.5 x 110 cm silica gel column (Silica Woelm, 32-63 μm)
20 packed and equilibrated with ethyl acetate. The columns were first eluted with ethyl acetate at a flow rate of 4 ml/minute for 4 hours, collecting 20 ml fractions. The eluent was changed to a concave gradient from ethyl acetate saturated with 0.1 M aqueous potassium dihydrogen phosphate to 10% isopropyl alcohol in ethyl acetate saturated with 0.1 M aqueous potassium dihydrogen phosphate
25 over 24 hours. The columns were finally eluted with 10% isopropyl alcohol in ethyl acetate saturated with 0.1 M aqueous potassium dihydrogen phosphate over night. The fractions were assayed in the BIA and those active
30 were analysed by TLC as described in Example 3.

Fractions (86-107) containing LL-E33288 α_3 -I from the two columns were pooled and worked up as before to yield about 2.1 g of crude LL-E33288 α_3 -I.

35

Fractions (182-253) containing LL-E33288 α_1 -I and α_2 -I from the two columns were pooled and worked up to yield about 4.2 g of a crude mixture of LL-E33288 α_1 -I and α_2 -I.

5 Fractions (254-272) containing LL-E33288 β_2 -I and β_1 -I from the two columns were pooled and worked up to yield about 1.2 g of a crude mixture of LL-E33288 β_2 -I and β_1 -I.

10 Fractions (273-302) containing LL-E33288 γ_1 -I from the two columns were pooled and worked up to yield about 1.9 g of 30% pure LL-E33288 γ_1 -I.

15 Fractions (303-340) containing LL-E33288 δ_1 -I from the two columns were pooled and worked up to yield about 1.3 g of partially purified LL-E33288 δ_1 -I.

Example 9

Purification of LL-E33288 γ_1 -I

Approximately 900 mg of the 30% pure LL-E33288 γ_1 -I from Example 8 was chromatographed on a 2.5 x 120 cm sephadex LH-20 column equilibrated with ethyl acetate:dichloromethane:ethanol (2:2:1) at a flow rate of 1 ml/minute, collecting 12 ml fractions. The fractions were assayed and analysed by TLC as before and those containing LL-E33288 γ_1 -I (fractions 24-33) were pooled and worked up to yield 428 mg of 64% pure LL-E33288 γ_1 -I.

25 A 22 mg sample of the above was chromatographed on a 0.8 x 24 cm Sepralyte C₁₈ (35-60 μ m, Analytichem) column equilibrated with acetonitrile:0.2 M aqueous ammonium acetate (55:45) at a flow rate of 2 ml/minute, collecting 12 ml fractions. The fractions were assayed and analysed by TLC as before and those containing pure LL-E33288 γ_1 -I were pooled and worked up to yield 7.7 mg of pure LL-E33288 γ_1 -I.

35 This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

Example 10Purification of LL-E33288 β_1 -I and β_2 -I

Approximately 600 mg of the crude mixture of LL-E33288 β_2 -I and β_1 -I from Example 8 was chromatographed on a 2.5 x 120 cm Sephadex LH-20 column equilibrated with ethyl acetate:dichloromethane:ethanol (2:2:1) at a flow rate of 1 ml/minute, collecting 12 ml fractions. The fractions were assayed and analysed by TLC as before and those containing LL-E33288 β_2 -I and LL-E33288 β_1 -I (fractions 23-31) were pooled and worked up to yield 81 mg of a partially purified mixture of LL-E33288 β_2 -I and β_1 -I.

The sample above was chromatographed on a 1.5 x 90 cm Sephadex LH-20 column equilibrated with hexane:dichloromethane:ethanol (3:1:1) at a flow rate of 0.8 ml/minute, collecting 12 ml fractions. The fractions were assayed and analysed by TLC as before and those containing LL-E33288 β_2 -I (fractions 17-30) and LL-E33288 β_1 -I (fractions 31-38) were pooled separately and worked up to yield 31 mg of partially purified LL-E33288 β_2 -I and 20 mg of 80% pure LL-E33288 β_1 -I.

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

29,845



WHAT IS CLAIMED IS:

1. A compound LL-E33288 α_1 -Br, having the following R_f values in the indicated solvent systems on TLC on silica gel sheets:

- a) ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.67$;
- b) 3% isopropanol in ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.80$; and
- c) ethyl acetate:methanol (95:5), $R_f=0.79$.

2. A compound LL-E33288 α_2 -Br, having the following R_f values in the indicated solvent systems on TLC on silica gel sheets:

- a) ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.61$;
- b) 3% isopropanol in ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.75$; and
- c) ethyl acetate:methanol (95:5), $R_f=0.73$.

3. A compound LL-E33288 α_3 -Br having the following R_f values in the indicated solvent systems on TLC on silica gel sheets:

This Sheet Has Been Replaced

THIS SHEET HAS BEEN
REPLACED



- a) ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.55$;
- b) 3% isopropanol in ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.69$; and
- c) ethyl acetate:methanol (95:5), $R_f=0.61$.

4. A compound LL-E33288 α_4 -Br, having the following R_f values in the indicated solvent systems on TLC on silica gel sheets:

- a) ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.49$;
- b) 3% isopropanol in ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.64$; and
- c) ethyl acetate:methanol (95:5), $R_f=0.54$.

5. A compound LL-E33288 β_1 -Br,

- a) having an approximate elemental analysis:
C 48.6; H 5.6; N 2.9; S 9.1 and Br 5.5;
- b) having a melting point: 146-150°C (dec.);
- c) having a specific rotation: $[\alpha]_D^{26} = -49 \pm 10^\circ$
(0.1%, ethanol);

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED



0182152

-49-

- d) having ultraviolet absorption spectra as shown in Figure I of the drawings;
- e) having an infrared absorption spectrum as shown in Figure II of the drawings;
- f) having a proton magnetic resonance spectrum as shown in Figure III of the drawings;
- g) having a carbon-13 magnetic resonance spectrum as shown in Figure IV of the drawings with significant peaks at:

17.60(q);	17.64(q);	18.9(q);	19.7(q);
22.4(q);	22.8(q);	23.5(q);	34.3(t);
36.9(t);	39.2(t/d);	47.8(d);	51.7(q);
52.7(q);	54.6(d)	56.3(q);	57.2(q);
57.8(d);	61.0(q/d);	61.7(d);	62.4(t);
66.9(d);	68.4(d);	69.1(d);	69.7(d);
70.2(d);	71.1(d);	71.9(d);	72.1(s);
76.1(d);	81.0(d);	83.3(s);	88.2(s);
97.4(d);	99.7(d);	100.8(s);	102.5(d);
115.1(s);	123.4(d);	124.4(d);	126.5(d);
130.2(s);	130.8(s);	144.6(s);	149.3(s);
149.5(s);	191.7(s);	192.4(s);	and

- h) having the following R_f values in the indicated solvent systems on TLC on silica gel sheets:

- i) ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.24$;
- ii) 3% isopropanol in ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.35$;
- iii) ethyl acetate:methanol (95:5), $R_f=0.36$.

This Sheet Has Been Replaced

THIS SHEET HAS BEEN
REPLACED



i) having a molecular weight: 1333/1335, respectively for $^{79}\text{Br}/^{81}\text{Br}$; and

j) having a molecular formula: $\text{C}_{54}\text{H}_{84}\text{N}_3\text{O}_{22}\text{S}_4\text{Br}$.

6. A compound LL-E33288 β_2 -Br, having the following R_f values in the indicated solvent systems on TLC on silica gel sheets:

- a) ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.32$;
- b) 3% isopropanol in ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.41$; and
- c) ethyl acetate:methanol (95:5), $R_f=0.45$.

7. A compound LL-E33288 γ_1 -Br

- a) having ultraviolet absorption spectra as shown in Figure V of the drawings;
- b) having an infrared absorption spectrum as shown in Figure VI of the drawings;
- c) having a proton magnetic resonance spectrum as shown in Figure VII of the drawings; and
- d) having the following R_f values in the indicated solvent systems on TLC on silica gel sheets:
 - i) ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.18$;

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED



ii) 3% isopropanol in ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.28$;

iii) ethyl acetate:methanol (95:5), $R_f=0.27$.

e) having a carbon-13 magnetic resonance spectrum as shown in Figure VIII of the drawings with significant peaks at:

14.4	17.6	17.9	19.0
19.7	-	22.8	-
-	34.0	37.6	39.5
42.1	-	51.6	52.7
54.1	56.3	57.3	-
59.3	61.1	61.8	61.9
67.2	68.18	68.23	69.7
70.1	70.8	71.1	71.7
71.8	76.1	-	81.0
82.9	88.4	-	97.8
100.0	100.2	101.3	103.0
115.3	123.0	124.9	126.9
130.4	131.1	131.8	138.0
144.7	-	149.5	149.6
155.6	192.5	192.9	

f) having a molecular formula: $C_{53}H_{82}N_3O_{22}S_4Br$; and

g) having a molecular weight: 1319/1321, respectively for $^{79}Br/^{81}Br$.

8. A compound LL-E33288 α_1 -I

a) having the following R_f values in the indicated solvent systems on TLC on silica gel sheets:

i) ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.67$;





- ii) 3% isopropanol in ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.80$; and
 - iii) ethyl acetate:methanol (95:5), $R_f=0.80$; and
 - b) having a molecular weight:1145.
9. A compound LL-E33288 α_2 -I
- a) having the following R_f values in the indicated solvent systems on TLC on silica gel sheets:
 - i) ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.61$;
 - ii) 3% isopropanol in ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.75$; and
 - iii) ethyl acetate:methanol (95:5), $R_f=0.73$;
 - b) containing only the following elements: C, H, N, O, S and I;
 - c) having a molecular weight:1131; and
 - d) having a proton magnetic resonance spectrum as shown in Figure IX of the drawings.
10. A compound LL-E33288 α_3 -I
- a) having the following R_f values in the indicated solvent systems on TLC on silica gel sheets:
 - i) ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.55$;





- ii) 3% isopropanol in ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.69$; and
 - iii) ethyl acetate:methanol (95:5), $R_f=0.61$;
- b) having a molecular weight:1066; and
- c) having a proton magnetic resonance spectrum as shown in Figure X of the drawings.

11. A compound LL-E332888₁-I

- a) having the following R_f values in the indicated solvent systems on TLC on silica gel sheets:
- i) ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.24$;
 - ii) 3% isopropanol in ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.35$; and
 - iii) ethyl acetate:methanol (95:5), $R_f=0.36$;
- b) having an ultraviolet absorption spectra as shown in Figure XI of the drawings;
- c) having an infrared absorption spectrum as shown in Figure XII of the drawings;
- d) having a proton magnetic resonance spectrum as shown in Figure XIII of the drawings;
- e) having a carbon 13 magnetic resonance spectrum as shown in Figure XIV of the drawings with significant peaks at



-	17.5	17.6	18.9
-	22.4	22.8	23.4
25.4	34.3	36.9	39.2
-	47.9	51.6	52.8
54.8	56.3	57.2	57.9
60.9		61.6	62.2
67.0	68.4	68.4	69.1
69.6	70.4	71.1	71.8
72.2	76.2	-	80.8
83.3	88.1	93.6	97.4
99.6	99.6	-	102.6
112.4	123.4	124.4	126.4
-	-	133.4	-
-	-	-	-
-	192.3	192.6	-



f) having a molecular formula: $C_{54}H_{84}N_3O_{22}S_4I$; and

g) having a molecular weight: 1381.

12. A compound LL-E33288 β ₂-I having the following R_f values in the indicated solvent systems on TLC on silica gel sheets:

- a) ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.32$.
- b) 3% isopropanol in ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.41$; and
- c) ethyl acetate:methanol (95:5), $R_f=0.45$.

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

13. A compound LL-E33288Y1-I

a) having the following R_f values in the indicated solvent systems on TLC on silica gel sheets:

- i) ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f = 0.18$;
- ii) 3% isopropanol in ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f = 0.28$; and
- iii) ethyl acetate: methanol (95:5), $R_f = 0.27$;

b) containing only the following elements: C, H, N, O, S and I;

c) having an approximate elemental analysis:

C 48.8; H 5.4; N 2.8; S 9.0; and I 9.2;

d) having a molecular weight: 1367;

e) having a molecular formula: $C_{53} H_{82} N_3 O_{22} S_4 I$;

f) having an ultraviolet absorption spectra as shown in Figure XV of the drawings;

g) having an infrared absorption spectrum as shown in Figure XVI of the drawings;

h) having a proton magnetic resonance spectrum as shown in Figure XVII of the drawings; and

i) having a carbon 13 magnetic resonance spectrum as shown in Figure XVIII of the drawings, significant peaks as listed below:

14.5(q)	17.6(q)	17.6(q)	18.9(q)
-	-	22.8(q)	-
25.4(q)	34.1(t)	37.0(t)	39.1(t)
42.3(t/s)	-	51.5(d)	52.8(q)
54.8(t)	56.3(q)	57.2(q)	-
60.4(d)	60.9(q)	61.3(t)	61.7(q)
67.0(d)	68.4(d)	68.5(d)	69.2(d)
69.7(d)	70.5(d)	71.1(d)	71.8(d)
72.1(s)	75.7(d)	75.8(d)	80.9(d)
82.8(s)	88.1(s)	93.5(s)	97.3(d)
99.6(d)	99.7(d)	100.8(s)	102.6(d)
-	123.4(d)	124.4(d)	126.2(d)
130.2(s)	131.0(s)	133.4(s)	139.1(s)
143.0(s)	145.1	150.6(s)	151.5(s)
154.5	192.0(s)	192.5(s)	

This Sheet Has Been Replaced

THIS SHEET HAS BEEN
REPLACED



14.) A compound LL-33288 δ_1 -I having the following R_f values in the indicated solvent systems on TLC on silica gel sheets:

a) ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, R_f = 0.11; and

b) 3% isopropanol in ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, R_f = 0.19.

15. A method of treating bacterial infections in warmblooded animals which comprises administering to said animals an antibacterially effective amount of a compound selected from the group consisting of LL-E33288 α_1 -Br; LL-E33288 α_1 -I; LL-E33288 α_2 -Br; LL-E33288 α_2 -I; LL-E33288 α_3 -Br; LL-E33288 α_3 -I; LL-E33288 α_4 -Br; LL-E33288 β_1 -Br; LL-E33288 β_1 -I; LL-E33288 β_2 -Br; LL-E33288 β_2 -I; LL-E33288 γ_1 -Br; LL-E33288 γ_1 -I; and LL-E33288 δ_1 -I.

16. A method of inhibiting the growth of tumors in a mammal comprising administering to said mammal an effective amount of a compound selected from the group consisting of LL-E33288 α_1 -Br; LL-E33288 α_1 -I; LL-E33288 α_2 -Br; LL-E33288 α_2 -I; LL-E33288 α_3 -Br; LL-E33288 α_3 -I; LL-E33288 α_4 -Br; LL-E33288 β_1 -Br; LL-E33288 β_1 -I; LL-E33288 β_2 -Br; LL-E33288 β_2 -I; LL-E33288 γ_1 -Br; LL-E33288 γ_1 -I; and LL-E33288 δ_1 -I.

17. A method of regressing leukemia in a mammal comprising administering to said mammal an effective amount of a compound selected from the group consisting of LL-E33288 α_1 -Br; LL-E33288 α_1 -I; LL-E33288 α_2 -Br; LL-E33288 α_2 -I; LL-E33288 α_3 -Br; LL-E33288 α_3 -I; LL-E33288 α_4 -Br; LL-E33288 β_1 -Br; LL-E33288 β_1 -I; LL-E33288 β_2 -Br; LL-E33288 β_2 -I; LL-E33288 γ_1 -Br; LL-E33288 γ_1 -I; and LL-E33288 δ_1 -I.

18. A process for producing antibiotics LL-E33288 α_1 -Br; LL-E33288 α_2 -Br; LL-E33288 α_3 -Br; LL-E33288 α_4 -Br; LL-E33288 β_1 -Br; LL-E33288 β_2 -Br and LL-E33288 γ_1 -Br which comprises aerobically fermenting the organism Micromonospora echinospora ssp. calichensis NRRL 15839 or mutants thereof including NRRL 15975 in a liquid medium containing assimilable sources of carbon, nitrogen,



This Sheet Has Been Replaced



-57-

bromine and inorganic salts, until substantial antibiotic activity is imparted to said medium and then recovering the antibiotics therefrom.

19. A process for producing antibiotics LL-E33288 α ₁-Br; LL-E33288 α ₂-Br; LL-E33288 α ₃-Br; LL-E33288 α ₄-Br; LL-E33288 β ₁-Br; LL-E33288 β ₂-Br; and LL-E33288 γ ₁-Br which comprises aerobically fermenting a liquid medium containing assimilable sources of carbon, nitrogen, bromine and inorganic salts; which medium has been inoculated with a viable culture of the organism Micromonospora echinospora ssp. calichensis NRRL 15839 or mutants thereof including NRRL 15975, maintaining said fermentation culture at a temperature of about 24-32°C for a period of approximately 90-200 hours, harvesting the mash and extracting the antibiotics.

20. A process for producing antibiotics LL-E33288 α ₁-I; LL-E33288 α ₂-I; LL-E33288 α ₃-I; LL-E33288 β ₁-I; LL-E33288 β ₂-I; LL-E33288 γ ₁-I; and LL-E33288 δ ₁-I which comprises aerobically fermenting the organism Micromonospora echniospora ssp. calichensis NRRL 15839 or mutants thereof including NRRL 15975 in a liquid medium containing assimilable sources of carbon, nitrogen, iodine and inorganic salts, until substantial antibiotic activity is imparted to said medium and then recovering the antibiotics therefrom.

21. A process for producing antibiotics LL-E33288 α ₁-I; LL-E33288 α ₂-I; LL-E33288 α ₃-I; LL-E33288 β ₁-I; LL-E33288 β ₂-I; LL-E33288 γ ₁-I; and LL-E33288 δ ₁-I which comprises aerobically fermenting a liquid medium containing assimilable sources of carbon, nitrogen, iodine and inorganic salts, which medium has been inoculated with a viable culture of the microorganism Micromonospora echinospora ssp. calichensis NRRL 15839 or mutants thereof including NRRL 15975, maintaining said fermentation culture at a temperature of about 24-32°C for a period of approximately 90-200 hours, harvesting the mash and extracting the antibiotics.

22. A culture containing the microorganism

THIS SHEET HAS BEEN
REPLACED

0182152




-58-

Micromonospora echinospora ssp. calichensis, NRRL 15839, said culture being capable of producing the LL-E33288 complex in recoverable quantity upon aerobic fermentation in an aqueous medium containing assimilable sources of carbon nitrogen, inorganic salts, and either iodine or bromine or both.

23. A culture containing the microorganism Micro-
monospora echinospora ssp. calichensis NRRL 15975, said culture being capable of producing the LL-E33288 complex in recoverable quantity upon aerobic fermentation in an aqueous medium containing assimilable sources of carbon, nitrogen, inorganic salts, and either iodine or bromine or both.

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

ABSTRACTANTITUMOR ANTIBIOTICS (LL-E33288 COMPLEX)

Antibacterial and antitumor agents designated LL-E33288 complex and their production by new strains of *Micromonospora*, *Micromonospora echinospora* ssp. *calichensis*, NRRL-15839 and NRRL-15975, are disclosed.

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

1/18

ULTRAVIOLET ABSORPTION SPECTRA OF LL-E33288 β ,
20 $\mu\text{g}/\text{ml}$ SOLUTION




FIGURE I

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

2/18


INFRARED ABSORPTION SPECTRUM OF LL-E33288 β ,

FIGURE II

This Sheet Has Been Replaced

THIS SHEET HAS BEEN
REPLACED

0182152

3/18

PROTON MAGNETIC RESONANCE SPECTRUM
OF LL-E33288 β .




FIGURE III

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

4/18

CARBON 13 MAGNETIC RESONANCE SPECTRUM
OF LL-E33288 β .




FIGURE IV

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

5/18

ULTRAVIOLET ABSORPTION SPECTRA OF LL-E33288 γ
20 $\mu\text{g/ml}$ SOLUTION




FIGURE V


This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

0182152

INFRARED ABSORPTION SPECTRUM OF LL-E33288 γ



This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

7/18

PROTON MAGNETIC RESONANCE SPECTRUM OF
LL-E33288 γ




FIGURE VII

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

0182152

8/18

^{13}C NMR OF LL-E33288 γ , Br




FIGURE VIII

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

0182152

9/18

PMR OF LL-E33288 α_2 I




FIGURE IX

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

0182152

10 / 18

PMR OF LL-E 33288 α_3 I




FIGURE X

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

0182152

11/18

UV OF LL-E33288 β_1 I




FIGURE XI

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

12/18


IR OF LL-E33288 β , I

FIGURE XII

This Sheet Has Been Replaced

THIS SHEET HAS BEEN
REPLACED

0182152

13/18

PMR OF LL-E33288 β , I





FIGURE XIII

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

0182152

14/18

^{13}C NMR OF LL-E33288 β , I




FIGURE XIV


This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

0182152

15/18



This Sheet Has Been Replaced




THIS SHEET HAS BEEN
REPLACED

0182152

16/18

IR OF LL-E33288Y, I



WAVENUMBERS

FIGURE XVI

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

17 / 18

PMR OF LL-E33288X, I




FIGURE XVII

This Sheet Has Been Replaced

THIS SHEET HAS BEEN
REPLACED

0182152

18 / 18

^{13}C NMR OF LL-E33288X, I




FIGURE XVIII

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED



Title: ANTITUMOR ANTIBIOTICS
(LL-E33288 COMPLEX)

SUMMARY OF THE INVENTION

This application is a continuation-in-part of co-pending application S.N. 672,031, filed November 16, 1984.

This invention relates to new antibacterial and antitumor agents designated LL-E33288 α_1 -Br, LL-E33288 α_1 -I, LL-E33288 α_2 -Br, LL-E33288 α_2 -I, LL-E33288 α_3 -Br, LL-E33288 α_3 -I, LL-E33288 α_4 -Br, LL-E33288 β_1 -Br, LL-E33288 β_1 -I, LL-E33288 β_2 -Br, LL-E33288 γ -I, LL-E33288 γ_1 -Br, LL-E33288 γ_1 -I and LL-E33288 δ_1 -I, to their production by fermentation, to methods for their recovery and concentration from crude solutions and to processes for their purification. The present invention includes within its scope the antibacterial and antitumor agents in dilute form, as crude concentrates, as a complex of various or all components, in pure form as individual components and novel strains of Micromonospora.

The LL-E33288 antibiotics of this invention are closely related compounds. The fourteen antibiotics are recovered from fermentation and are initially obtained as a mixture, hereinafter either the LL-E33888 complex, the the LL-E33288 Iodo-complex or the LL-E33288 Bromo-complex. In general, the iodine containing components of the LL-E33288 antibiotics (e.g., α_1 -I, α_2 -I, α_3 -I, β_1 -I, β_2 -I, γ_1 -I and δ_1 -I) are found only in fermentations using media containing inorganic or organic iodide while the bromine containing components (e.g., α_1 -Br, α_2 -Br, α_3 -Br, α_4 -Br, β_1 -Br, β_2 -Br and γ_1 -Br) are found only in fermentations using media containing inorganic or organic bromide. The ratio of components in the LL-E3388 complex will vary, depending upon

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

18 OCT 1996

-2-

the fermentation of both the bromine and the iodine containing antibiotics, LL-E33288 β ₁ and LL-E33288 γ ₁ are the major components, together accounting for approximately 90% of the complex. LL-E33288 α ₁, LL-E33288 α ₂, LL-E33288 α ₃, LL-E33288 α ₄-Br, LL-E33288 β ₂ and LL-E33288 δ ₁-I are minor components, together accounting for approximately 10% of the complex.

The LL-E33288 antibiotics are active against gram-positive and gram-negative bacteria. Each of the components were also found to be active in a modification of the Biochemical Induction Assay [Elespuru, R. and Yarmolinsky, M., Environmental Mutagenesis, 1, 65-78 1979)], a test which specifically measures the ability of an agent to directly or indirectly initiate DNA damage. In this assay, both LL-E33288 β ₁-Br and LL-E33288 γ ₁-Br were active at concentrations lower than 1×10^{-6} mcg/ml.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. I is the ultraviolet absorption spectra of LL-E33288 β ₁-Br;

FIG. II is the infrared absorption spectrum of LL-E33288 β ₁-Br;

FIG. III is the proton magnetic resonance spectrum of LL-E33288 β ₁-Br;

FIG. IV is the carbon 13 magnetic resonance spectrum of LL-E33288 β ₁-Br;

FIG. V is the ultraviolet absorption spectra of LL-E33288 γ ₁-Br;

FIG. VI is the infrared absorption spectrum of LL-E33288 γ ₁-Br;

FIG. VII is the proton magnetic resonance spectrum of LL-E33288 γ ₁-Br;

FIG. VIII is the carbon 13 magnetic resonance spectrum of LL-E33288 γ ₁-Br;

FIG. IX is the proton magnetic resonance spectrum of LL-E33288 α ₂-I;

FIG. X is the proton magnetic resonance spectrum of LL-E33288 α ₃-I;




FIG. XI is the ultraviolet absorption spectra of LL-E33288 β ₁-I;

FIG. XII is the infrared absorption spectrum of LL-E33288 β ₁-I;

FIG. XIII is the protein magnetic resonance spectrum of LL-E33288 β ₁-I;

FIG. XIV is the carbon 13 magnetic resonance spectrum of LL-E33288 β ₁-I;

FIG. XV is the ultraviolet absorption spectra of LL-E33288 γ ₁-I;

FIG. XVI is the infrared absorption spectrum of LL-E33288 γ ₁-I;

FIG. XVII is the proton magnetic resonance spectrum of LL-E33288 γ ₁-I; and


FIG. XVIII is the carbon 13 magnetic resonance spectrum of LL-E33288 γ ₁-I.

DETAILED DESCRIPTION OF THE INVENTION

The physico-chemical characteristics of LL-E33288 β ₁-Br and LL-E33288 γ ₁-Br are described below:

LL-E33288 β ₁-Br

- 1) Approximate elemental analysis: C 48.6; H 5.6; N 2.9; S 9.1; and Br 5.5. (It has been determined by electron spectroscopy for chemical analysis (ESCA) that only the following elements are present: C, H, N, O, S and Br);
- 2) Melting point: 146-150°C (dec.);
- 3) Specific rotation: $[\alpha]_D^{26} = -49 \pm 10^\circ$ (0.1% ethanol);
- 4) Ultraviolet absorption spectra: as shown in Figure I (methanol; acidic methanol; basic methanol);
- 5) Infrared absorption spectrum: as shown in Figure II (KBr disc);
- 6) Proton magnetic resonance spectrum: as shown in Figure III (300 MHz, CDCl₃);
- 7) Carbon-13 magnetic resonance spectrum: as shown in Figure IV (75.43 MHz, CDCl₃, ppm from TMS), significant peaks as listed below:




-4-

17.60(q);	17.64(q);	18.9(q);	19.7(q);
22.4(q);	22.8(q);	23.5(q);	34.3(t);
36.9(t);	39.2 (t/d);	47.8 (d);	51.7(q);
52.7(q);	54.6 (t/d);	56.3(q);	57.2(q);
57.8(d);	61.0 (q/d);	61.7(d);	62.4 (t);
66.9(d);	68.4(d);	69.1(d);	69.7(d);
70.2(d);	71.1(d);	71.9(d);	72.1 (s/t);
76.1(d);	81.0(d);	83.3(s);	88.2(s);
97.4(d);	99.7(d);	100.8(s);	102.5(d);
115.1(s);	123.4(d);	124.4(d);	126.5(d);
130.2(s);	130.8(s);	144.6(s);	149.3(s);
149.5(s);	191.7(s);	192.4(s);	

- 8) Molecular weight: 1333/1335 respectively for $^{79}\text{Br}/^{81}\text{Br}$ as determined by FAB-MS; and
- 9) Molecular formula: $\text{C}_{54}\text{H}_{84}\text{N}_3\text{O}_{22}\text{S}_4\text{Br}$, exact masses at 1258.3699 (^{79}Br) and 1260.3726 (^{81}Br) was determined by high resolution FAB-MS and calculated to be $\text{C}_{52}\text{H}_{81}\text{N}_3\text{O}_{21}\text{S}_3\text{Br}$ ($\text{M}+\text{H}-\text{C}_2\text{H}_4\text{OS}$).

LL-E33288 γ_1 -Br

- 1) Ultraviolet absorption spectra: as shown in Figure V (methanol; acidic methanol; basic methanol);
- 2) Infrared absorption spectrum: as shown in Figure VI (KBr disc);
- 3) Proton magnetic resonance spectrum: as shown in Figure VII (300 MHz, CDCl_3);
- 4) Carbon 13 magnetic resonance spectrum: as shown in Figure VIII (75.43 MHz, CDCl_3 , ppm from TMS), significant peaks as listed below:



This Sheet Has Been Replaced



14.4	17.6	17.9	19.0
19.7	-	22.8	-
-	34.0	37.6	39.5
42.1	-	51.6	52.7
54.1	56.3	57.3	-
59.3	61.1	61.8	61.9
67.2	68.18	68.23	69.7
70.1	70.8	71.1	71.7
71.8	76.1	-	81.0
82.9	88.4	-	97.8
100.0	100.2	101.3	103.0
115.3	123.0	124.9	126.9
130.4	131.1	131.8	138.0
144.7	-	149.5	149.6
155.6	192.5	192.9	

- 5) Molecular formula: $C_{53}H_{82}N_3O_{22}S_4Br$ by comparing its UV, IR, 1H NMR, and ^{13}C NMR data to those of LL-E33288 β $_1$ -BR and LL-E33288 γ $_1$ -I; and
- 6) Molecular weight: 1319/1321 respectively for $^{79}Br/^{81}Br$, calculated from its molecular formula.

The physico-chemical characteristics of LL-E33288 α $_1$ -I, LL-E33288 α $_2$ -I, LL-E33288 α $_3$ -I, and LL-E33288 β $_1$ -I and LL-E33288 γ $_1$ -I are described below:

LL-E33288 α $_1$ -I

- 1) Molecular weight: 1145, determined by FAB-MS.

LL-E33288 α $_2$ -I

- 1) Contains and only contains the following elements by electron spectroscopy for chemical analysis (ESCA): C, H, N, O, S, I;
- 2) Molecular weight: 1131, determined by FAB-MS; and
- 3) Proton magnetic resonance spectrum: as shown in Figure IX (300 MHz, $CDCl_3$).

LL-E33288 α $_3$ -I

- 1) Molecular weight: 1066, determined by FAB-MS; and
- 2) Proton magnetic resonance spectrum: as shown in

THIS SHEET HAS BEEN
REPLACED




Figure X (300 MHz, CDCl₃).

LL-E33288 β₁-I


- 1) Ultraviolet absorption spectra: as shown in Figure XI (methanol; acidic methanol; basic methanol);
- 2) Infrared absorption spectrum: as shown in Figure XII (KBr disc);
- 3) Proton magnetic resonance spectrum: as shown in Figure XIII (300 MHz, CDCl₃);
- 4) Carbon 13 magnetic resonance spectrum: as shown in Figure XIV (75.43 MHz, CDCl₃, ppm from TMS), significant peaks as listed below:

-	17.5	17.6	18.9
-	22.4	22.8	23.4
25.4	34.3	36.9	39.2
-	47.9	51.6	52.8
54.8	56.3	57.2	57.9
60.9		61.6	62.2
67.0	68.4	68.4	69.1
69.6	70.4	71.1	71.8
72.2	76.2	-	80.8
83.3	88.1	93.6	97.4
99.6	99.6	-	102.6
112.4	123.4	124.4	126.4
-	-	133.4	-
-	-	-	-
-	192.3	192.6	-

- 5) Molecular formula: C₅₄H₈₄N₃O₂₂S₄I by comparing its UV, IR, ¹H NMR and ¹³C NMR data to those of LL-E33288 β₁-Br and LL-E33288 γ₁-I; and
- 6) Molecular weight: 1381, calculated from molecular formula.

LL-E33288 γ₁-I

- 1) Contains and only contains the following elements by electron spectroscopy for chemical analysis (ESCA): C, H, N, O, S, I;
- 2) Approximate elemental analysis: C 48.8; H 5.4; N 2.8; S 9.0; I 9.2;




-7-

- 3) Molecular weight: 1367, determined by FAB-MS;
- 4) Molecular formula: $C_{53}H_{82}N_3O_{22}S_4I$, exact mass for $M+H$ was determined by high resolution FAB-MS to be 1368. 3397 for $C_{53}H_{83}N_3O_{22}S_4I$;
- 5) Ultraviolet absorption spectra: as shown in Figure XV (methanol; acidic methanol; basic methanol);
- 6) Infrared absorption spectrum: as shown in Figure XVI (KBr disc);
- 7) Proton magnetic resonance spectrum: as shown in Figure XVII (300 MHz, $CDCl_3$); and
- 8) Carbon 13 magnetic resonance spectrum: as shown in Figure XVIII (75.43 MHz, $CDCl_3$, ppm for TMS) significant peaks as listed below:

14.5(q)	17.6(q)	17.6(q)	18.9(q)
-	-	22.8(q)	-
25.4(q)	34.1(t)	37.0(t)	39.1(t)
42.3(t/s)	-	51.5(d)	52.8(q)
54.8(t)	56.3(q)	57.2(q)	-
60.4(d)	60.9(q)	61.3(t)	61.7(q)
67.0(d)	68.4(d)	68.5(d)	69.2(d)
69.7(d)	70.5(d)	71.1(d)	71.8(d)
72.1(s)	75.7(d)	75.8(d)	80.9(d)
82.8(s)	88.1(s)	93.5(s)	97.3(d)
99.6(d)	99.7(d)	100.8(s)	102.6(d)
-	123.4(d)	124.4(d)	126.2(d)
130.2(s)	131.0(s)	133.4(s)	139.1(s)
143.0(s)	145.1	150.6(s)	151.5(s)
154.5	192.0(s)	192.5(s)	

The LL-E33288 components are most conveniently separated and identified by high-performance liquid chromatography (HPLC) and by thin-layer chromatography (TLC). It is difficult, although not impossible, to separate the corresponding iodinated and brominated components by HPLC; however, they cannot be distinguished by TLC.



-8-

The preferred analytical separation of the LL-E33288-Br components by HPLC uses the following conditions:

Column: "Separalyte C₁₈ 5 m," 4.6 mm x 25 cm
(Analytichem International);

Solvent: Acetonitrile: 0.2M aqueous ammonium acetate (60:40);

Flow rate: 1.5 ml/minute


Detector: Dual wavelength UV at 254 nm and 280 nm;

Sensitivity: 0-0.02 A.U.F.S.

Table IA gives the approximate retention times and volumes of LL-E33288 β ₁-Br, LL-E33288 β ₂-Br, and LL-E33288 γ ₁-Br under these conditions.

TABLE IA

LL-E33288 Components	Retention Time (minutes)	Retention Volume(ml)
β ₁ -Br	5.7	8.6
β ₂ -Br	7.1	10.7
γ ₁ -Br	4.3	6.5



The preferred analytical HPLC separation of the iodine containing LL-E33288 components uses the following conditions:

Column: NOVA-PAK C₁₆ Radial-PAK cartridge with RCM-100 Radial Compression Module (Millipore, Waters Chromatography Division);

Solvent: Acetonitrile: 0.2M aqueous ammonium acetate (50:50);

Flow Rate: 1.2 ml/minute;


Detector: Dual wavelength UV at 254 nm and 280 nm;

Sensitivity: 0-0.02 A.U.F.S.

Table IB gives the approximate retention times and volumes of LL-E33288 α_1 -I, LL-E33288 α_2 -I, LL-E33288 α_3 -I, LL-E33288 β_1 -I, LL-E33288 β_2 -I, LL-E33288 γ_1 -I and LL-E33288 δ_1 -I under these conditions.

TABLE IB

LL-E33288 Components	Retention Time (minutes)	Retention Volume(ml)
α_1 -I	11.9	14.3
α_2 -I	9.1	10.9
α_3 -I	1.5	1.8
β_1 -I	4.4	5.3
β_2 -I	5.0	6.0
γ_1 -I	3.6	4.3
δ_1 -I	2.6	3.1



The LL-E33288 components are separated and identified by the following TLC system:

Adsorbant: Silica gel 60 F254 pre-coated aluminum sheets, 0.2mm layer thickness, EM Reagents;

Detection: Visualized by quenching effect under short wavelength UV lamp (254 nm), and bioautography using Bacillus subtilis or the modified biochemical induction assay;

Solvent Systems: I, ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate;


II, 3% isopropyl alcohol in ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate;

III, ethyl acetate:methanol (95.5).

Table II gives the approximate R_f values of LL-E33288 components in these three systems:

TABLE II

LL-E33288 Components	R _f Value		
	Solvent System I	Solvent System II	Solvent System III
α ₁ -Br, α ₁ -I	0.67	0.80	0.79
α ₂ -Br, α ₂ -I	0.61	0.75	0.73
α ₃ -Br, α ₃ -I	0.55	0.69	0.61
α ₄ -Br	0.49	0.64	0.54
β ₂ -Br, β ₂ -I	0.32	0.41	0.45
β ₁ -Br, β ₁ -I	0.24	0.35	0.36
γ ₁ -Br, γ ₁ -I	0.18	0.28	0.27
δ ₁ -I	0.11	0.19	



The new antibacterial and antitumor agents designated LL-E33288 α_1 -Br, LL-E33288 α_2 -Br, LL-E33288 α_3 -Br, LL-E33288 α_4 -Br, LL-E33288 β_1 -Br, LL-E33288 δ_2 -Br, LL-E33288 γ_1 -Br, LL-E33288 α_1 -I, LL-E33288 α_2 -I, LL-E33288 α_3 -I, LL-E33288 β_1 -I, LL-E33288 β_2 -I, LL-E33288 γ_1 -I and LL-E33288 δ_1 -I are formed during the cultivation under controlled conditions of a new strain of Micromonospora echinospora ssp. calichensis. This microorganism is maintained in the culture collection of the Medical Research Division, American Cyanamid Company, Pearl River, New York as culture number LL-E33288. A viable culture of this new microorganism has been deposited with the Culture Collection Laboratory, Northern Regional Research Center, U. S. Department of Agriculture, Peoria, Illinois on August 9, 1984, and has been added to its permanent collection. It has been assigned by such depository the strain designation NRRL 15839. Access to such culture, under strain designation NRRL 15839, during pendency of the instant application shall be available to one determined by the Commissioner of Patents and Trademarks to be entitled thereto under 37 C.F.R. §1.14 and 35 U.S.C. §122, and all restrictions on availability to the public of such culture will be irrevocably removed upon grant of a patent on the instant application.

Culture LL-E33288 was isolated from a caliche clay soil sample collected in Texas.

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

-12-

The generic assignment of the strain NRRL 15839 to the genus Micromonospora was confirmed morphologically and chemically. The strain produces monospores either singly or in masses on the vegetative hyphae. No aerial hyphae were observed. Electron microscopic examination showed that the spores were warty. Whole cell analysis showed that the strain contained the meso isomer of diaminopimelic acid (DAP). The 3-OH derivative of DAP was present in large (major) amounts. Additionally the strain showed the presence of xylose plus traces of arabinose in its whole cell sugar hydrolysates (whole cell sugar pattern of Type D).

From macromorphological and physiological studies it was concluded that NRRL 15839 can be considered subspecies of M. echinospora (it is closest to M. echinospora ssp. pallida). Data on the morphology of NRRL 15839 are given in Tables A and B. Physiological data are given in Tables C and D.

Table A

Macromorphology Of NRRL 15839
(Colors Are NBS-ISCC)

ISP Agar Medium	Spores	Vegetative Mycelium	Soluble Pigments
Yeast- Malt (ISP 2)	-	Dark orange- yellow (72)	-
Oatmeal (ISP 3)	-	Colorless - pale orange-yellow (73)	-
Inorganic Salts- Starch (ISP 4)	Slight border of black spores	Dark orange- yellow (72) to lt. yellow-brown (76)	Lt. brownish
Glycerol- Asparagine (ISP 5)	-	Pale orange-yellow (73) - colorless	-




Table B

Macromorphology of NRRL-15839 on Various Agar Media Used for Actinomycete Growth (28°C, 2 weeks)

Agar Medium	NRRL-15839
Pablum	Beige veg. Sl. black spores No sol. pig.
Yeast Czapeks	Beige veg. No spores No sol. pig.
Czapek's	Beige veg. Sl. black spores No sol. pig.
Yeast Dextrose	Tan veg. Moderate black sp. Sl. dark pig.
Nutrient	Colorless to tan veg. Sl. black spores No sol. pig.
Nutrient- Glycerol	Colorless to light beige veg. No black spores No sol. pig.
Bennett's Dextrin	Colorless to beige veg. Sl. black spores Sl. rosy-brown pig.
Glucose- Asparagine	Colorless to lt. orange-beige veg. No spores No sol. pig.

veg. = vegetative hyphae; pig. = pigment.

Table C

Carbohydrate Utilization of NRRL-15839


Arabinose	+
Cellulose	-
Fructose	+
Glucose	+
Inositol	-
Mannitol	-
Raffinose	+
Rhamnose	+
Sucrose	+
Xylose	+

Table D

Physiological Reactions of NRRL-15839

<u>Hydrolysis of</u>	
Casein	+
Xanthine	-
Hypoxanthine	-
Tyrosine	+
Adenine	-
Gelatin	+
Potato Starch	+
Esculin	+
<u>Production of</u>	
Nitrate	
Reductase	+
Phosphatase	W
Urease	-

REPLACEMENT SHEET
18 OCT 1996




Growth on
Salicin -
5% NaCl -
Lysozyme Broth -

Decarboxylation of
Acetate +
Benzoate -
Citrate -
Lactate -
Malate -
Mucate -
Oxalate -
Propionate +
Pyruvate +
Succinate -
Tartrate -

Acid from
Adonitol -
Arabinose +
Cellobiose +
Dextrin +
Dulcitol -
Erythritol -
Fructose +
Galactose V
Glucose +
Glycerol -
Inositol -
Lactose -
Maltose +
Mannitol -
Mannose +
 α -methyl D
Glucoside -
Melibiose -
Raffinose +
Rhamnose +
Salicin +
Sorbitol -
Sucrose +
Trehalose +
Xylose +
B-Methyl
D-xyloside -

Growth at
100 -
420 +
450 +

+ = positive; - = negative; V = variable;
W = weak.



Derivation of Mutant LL-E33288-R66, NRRL-15975

In an effort to improve fermentation yields, the original culture LL-E33288 (NRRL-15839) was plated and 50 single colonies were isolated. These were designated NS1 to NS50 (NS = natural selection).


Fermentation of these isolates showed that those with moderate sporulation were generally better producers of LL-E33288 complex. Selected as representative of this group was isolate NS6.

Using isolate NS6 as the starting culture, spore suspensions were prepared and exposed to various mutagens. Single colonies were isolated from a nitroso-guanidine treatment, but none proved to be significantly improved producers of the LL-E33288 complex. From a subsequent series of exposures to ultraviolet irradiation, single colonies were obtained from which isolate UV 610 was selected as a high yielding mutant. Isolate UV 610 was then streaked and sub-isolates 1 to 7 were obtained. Sub-isolate UV 610(3) was selected for further work.

Because of the highly potent antibacterial and antineoplastic nature of the LL-E33288 complex it is possible that once a limited concentration of the antibiotic is biosynthesized in the fermentation it may become toxic/inhibitory to the producing culture. Thus, an effort was made to obtain isolates which are resistant to the LL-E33288 antibiotic complex.

Vegetative growth from isolate UV 610(3) was prepared as employed for fermentation and used to inoculate a flask of medium consisting of peptone, dextrose, molasses and water. The medium was supplemented with LL-E33288₈₁-Br at a concentration of 8 µg/m. A number of platings were done from this flask and a resistant population was obtained on the seventh day. A total of 97 colonies (R1 to R97) were isolated. Isolate R66 was selected as a potentially improved producer of LL-E33288 ₈₁-Br.

The history is represented schematically below.



E33288 (Wild type)



NS6 (Natural selection)



NTG [NG1-NG120](Nitrosoguanidine)

UV [UV1-UV200](Ultraviolet)

UV [UV201-UV400](Ultraviolet)

UV [UV401-UV702](Ultraviolet)



UV610(3)(Sub-islake 3)




[R1 to R97](Resistant colonies)



R66

The mutant R66 is maintained in the culture collection of the Medical Research Division, American Cyanamid Company, Pearl river, New York as culture number LL-E33288 R66. A viable culture of this new microorganism has been deposited with the Culture Collection Laboratory, Northern Regional Research center, U. S. Department of Agriculture, Peoria, Illinois on June 6, 1985, and has been added to its permanent collection. It has been assigned by such depository the strain designation NRRL 15975. Access to such culture, under strain designation NRRL 15975, during pendency of the instant application



shall be available to one determined by the Commissioner of Patents and Trademarks to be entitled thereto under C.F.R. §1.14 and 35 U.S.C. §122, and all restrictions on availability to the public of such culture will be irrevocably removed upon grant of a patent on the instant application.

Morphologically, NRRL-15975 forms fewer spores than NRRL-15839. A comparison of NRRL-15975 with NRRL-15839 is given in Table DD.

Chemically, both NRRL-15839 and NRRL-15975 show the same whole cell sugar patterns (Type D: xylose and traces of arabinose). The whole cell diaminopimelic acid analysis reveals that 15975 does not form the meso isomer but only the 3-hydroxy derivative (NRRL-15839 contains both compounds). This does not change the chemo-taxonomic assignment.

Physiological tests show that NRRL-15839 and NRRL-15975 differ in only two physiological reactions (See Table D). NRRL-15975 is negative for nitrate reductase and positive for utilization of lactate. NRRL-15839 was weakly positive for both, but is now negative after having been maintained on slants for a few months. Thus these characters should be considered variable for this taxon.





TABLE DD

Morphological Comparison of NRRL 15839 and NRRL 15975

Agar Medium		NRRL 15839	NRRL 15975
Bennett's-Dextrin	V ¹	Beige-tan	Beige tan
	Sp	Black, copious	None
	SS	None	None
Czapek's	V	Orange-tan	Orange-tan
	Sp	Black, traces	None
	SS	None	None
Yeast Extract-Czapek's	V	Orange tan, flat	Orange tan, convoluted
	Sp	Black, traces	None
	SS	None	Slight yellowish
Potato-Dextrose	V	Very poor growth	Very poor growth
	Sp	None	None
	SS	None	None
Nutrient glycerol	V	Tan	Tan
	Sp	Black, sparse	Black, sparse
	SS	None	Slight brownish
Nutrient	V	Tan	Tan
	Sp	Black, fair	None
	SS	None	None

1 = V = vegetative hyphae; Sp = spores; SS = soluble pigment.



It is to be understood that for the production of these new antibacterial and antitumor agents the present invention is not limited to this particular organisms or to organisms fully answering the above growth microscopic characteristics which were given for illustrative purposes only. In fact, it is desired and intended to include the use of mutants produced from these organisms by various means such as exposure to X-radiation, ultra-violet radiation, N'-methyl-N'-nitro-N-nitrosoguanidine, actinophages and the like.

The in vitro antibacterial activity of LL-E33288 components was determined against a spectrum of gram-positive and gram-negative bacteria by a standard agar dilution method. Mueller-Hinton again containing two-fold decreasing concentrations of the antibiotics were poured into petri plates. the agar surfaces were inoculated with 1 to 5×10^4 colong forming units of bacteria by means of the steers replicating device. The lowest concentration of LL-E33288 component that inhibited growth of a bacterial strain after about 18 hours of incubation at approximately 35°C was recorded as the minimal inhibitory concentration (MIC) for that strain. The results are summarized in Table III.

TABLE III
In vitro Antibacterial Activity of LL-E33288 Components


Organism	Minimal Inhibitory Concentration, mcg/ml			
	81-Br	81-I	Y1-Br	Y1-I
<u>Escherichia coli</u>	0.25	0.50	0.50	0.25
"	0.12	0.25	0.25	0.25
"	0.12	0.25	0.25	0.25
<u>Klebsiella pneumoniae</u>	0.25	0.50	0.50	0.25
"	0.12	0.50	0.50	0.25
<u>Enterobacter cloacae</u>	0.5	0.50	0.50	0.50
"	0.25	0.25	0.50	0.50
<u>Serratia marcescens</u>	0.12	0.25	0.50	0.25
"	0.12	0.50	0.25	0.12
<u>Morganella morganii</u>	0.5	1	0.50	0.25
<u>Providencia stuartii</u>	0.25	0.50	1	0.25
<u>Citrobacter diversus</u>	0.12	0.50	0.50	0.25
"	0.12	0.25	0.25	0.12
<u>Acinetobacter sp.</u>	0.06	0.25	0.25	0.12
"	0.12	0.25	0.25	0.06

18 OCT 1996

-22-

TABLE III (continued)

Organism	Minimal Inhibitory Concentration, mcg/ml			
	β 1-Br	β 1-I	γ 1-Br	γ 1-I
<u>Pseudomonas aeruginosa</u>	0.5	0.50	1	0.25
"	0.25	0.25	0.50	0.12
12-4-4 (MP)	≤ 0.0025	≤ 0.000031	≤ 0.000031	≤ 0.000031
ATCC 27853	≤ 0.00025	≤ 0.000031	≤ 0.000031	≤ 0.000031
Smith	≤ 0.00025	≤ 0.000031	≤ 0.000031	≤ 0.000031
SSC 82-31	≤ 0.00025	≤ 0.000031	≤ 0.000031	≤ 0.000031
ATCC 25923	≤ 0.00025	≤ 0.000031	≤ 0.000031	≤ 0.000031
SSC 82-20	≤ 0.00025	≤ 0.000031	≤ 0.000031	≤ 0.000031
SSC 82-26	≤ 0.00025	≤ 0.000031	≤ 0.000031	≤ 0.000031
SSC 82-24	≤ 0.00025	≤ 0.000031	≤ 0.000031	≤ 0.000031
SSC 82-57	≤ 0.00025	≤ 0.000031	≤ 0.000031	≤ 0.000031
<u>Staphylococcus epidermidis</u>	≤ 0.00025	≤ 0.000031	≤ 0.000031	≤ 0.000031
CMC-83-133	≤ 0.00025	≤ 0.000031	≤ 0.000031	≤ 0.000031
ATCC 12228	≤ 0.00025	≤ 0.000031	≤ 0.000031	≤ 0.000031
"	0.0038	0.031	0.062	0.0078
<u>Enterococcus</u> sp.	≤ 0.00025	0.00012	≤ 0.000031	0.00012
<u>Streptococcus faecalis</u>	≤ 0.00025	≤ 0.000031	≤ 0.000031	≤ 0.000031
<u>Micrococcus luteus</u>	≤ 0.00025	≤ 0.000031	≤ 0.000031	≤ 0.000031
<u>Bacillus subtilis</u>	≤ 0.00025	≤ 0.000031	≤ 0.000031	≤ 0.000031
ATCC				



-23-

Certain in vivo testing systems and protocols have been developed by the National Cancer Institute for testing compounds to determine their suitability as anti-neoplastic agents. These have been reported in "Cancer Chemotherapy Reports", Part III, Vol. 3, No. 2 (1972), Geran, et al. These protocols have established standardized screening tests which are generally followed in the field of testing for antitumor agents. Of these systems, lymphocytic leukemia

18 OCT 1996

-24-

P388, melanotic melanoma B16, L1210 leukemia and colon 26 adenocarcinoma are particularly significant to the present invention. These neoplasms are utilized for testing as transplantable tumors in mice. Generally, significant anti-tumor activity, shown in these protocols by a percentage increase of mean survival times of the treated animals (T) over the control animals (C), is indicative of similar results in human leukemias and solid tumors.

Lymphocytic Leukemia P388 Test

The animals used were BDF₁ mice, all of one sex, weighing a minimum of 17 g and all within a 3 g weight range. There were 5 or 6 mice per test group. The tumor transplant was by intraperitoneal injection of 0.5 ml of dilute ascitic fluid containing 10⁶ cells of lymphocytic leukemia P388. LL-E33288 antibiotics were tested in the P388 system both as the individual β ₁-Br and 1-Br components and as a complex of all components (Bromo-complex). The test compounds were administered intraperitoneally at a volume of 0.5 ml in 0.2% Klucel in normal saline on days 1, 5 and 9 (relative to tumor inoculation) at the indicated doses. The mice were weighed and the survivors recorded on a regular basis for 30 days. The median survival time and the ratio of survival time for treated (T)/control (C) animals were calculated. The positive control compound was Cisplatin given as an intraperitoneal injection in 0.5 ml of 0.2% Klucel on days 1, 5 and 9 at the indicated doses. The results appear in Table IV.

If $T/C \times 100 (\%)$ is 125 or over, the tested compound is considered to have significant anti-tumor activity.

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

18 OCT 1996

-25-

TABLE IV

Lymphocytic Leukemia P388 Test

Compound	Dose (mg/kg)	Median Survival (Days)	T/C x 100 (%)
LL-E33288 (Bromo-complex)	3.2	16.5	156
	1.6	17.5	165
	0.8	18.5	175
	0.4	19	179
	0.2	16.5	156
Control	-	10.6	-
Positive Control	1.0	21.5	203
	0.25	15	142
	0.06	14.5	137
LL-E-332888 ₁ -Br	0.4	13	105
	0.2	18	145
	0.1	19	153
	0.05	17.5	141
	0.025	18	145
	0.012	14	113
Control	-	12.4	-
Positive Control	1.0	25.5	206
	0.4	19	153
	0.06	15	121

18 OCT 1996

-26-

TABLE IV (continued)

Compound	Dose (mg/kg)	Median Survival (Days)	T/C x 100 (%)
LL-E33288 γ_1 -Br	0.2	14	113
	0.1	21	169
	0.05	19.5	157
	0.025	18	145
	0.012	14.5	117
Control	-	12.4	-
Positive Control	1.0	25.5	206
	0.4	19	153
	0.06	15	121

Melanotic Melanoma B16

The animals used were BDF₁ mice, all of the same sex, weighing a minimum of 17 g and all within a 3 g weight range. There are normally 6 animals per test group. A 1 g portion of melanotic melanoma B16 tumor was homogenized in 10 ml of cold balanced salt solution and a 0.5 ml aliquot of the homogenate was implanted intraperitoneally into each of the test mice. LL-E33288 antibiotics were tested in the B16 system both as the individual β_1 -Br and γ_1 -Br components and as a complex of all components (Bromo-complex). The test compounds were administered intraperitoneally on days 1 through 9 (relative to tumor inoculation) at various doses. The mice were weighed and survivors recorded on a regular basis for 60 days. The median survival time and the ratio of survival time for treated (T)/control (C) animals were calculated. The positive control compounds were Cisplatin or Adriamycin. The results of this test appear in Table V. If T/C X 100 (%) is 125 or over, the tested compound is considered to have significant anti-tumor activity.




TABLE V
Melanotic Melanoma B₁₆ Test

Compound	Dose (mg/kg)	Median Survival (Days)	T/C x 100 (%)
LL-E33288 (Bromo-complex)	0.8	30	188
	0.4	29.5	184
	0.2	27	169
	0.1	24	150
Control	-	16	-
Cisplatin	0.4	25	156
	0.2	25	156
	0.1	23	144
	0.05	21.5	134
LL-E33288 ₁ -Br	0.05	32	168
	0.025	33.5	176
	0.0125	32	168
Control	-	19	-
Adriamycin	0.8	> 60	> 316
	0.4	> 60	> 316





TABLE V (continued)

Compound	Dose (mg/kg)	Median Survival (Days)	T/C x 100 (%)
LL-E33288 γ ₁ -Br	0.05	33.5	176
	0.025	30	159
	0.0125	37	195
Control	-	19	-
Adriamycin	0.8	> 60	> 316
	0.4	> 60	> 316

Lymphocytic Leukemia L1210 Test

The animals used were BDF₁ mice, all of one sex, weighing a minimum of 17 g and all within a 3 g weight range. There were 6 mice in each test group and 18 in control groups. The tumor transplant was by intraperitoneal injection of 0.5 ml of lymphocytic leukemia L1210 at a concentration of 10⁵ cells per mouse. LL-E33288 antibiotics were tested in the L1210 system both as the individual β ₁-Br component and as a complex of all components (Bromo-complex). The test compounds were administered on days 1, 5 and 9 or days 1 through 9 (relative to tumor inoculation) at various doses. The mice were weighed and survivors recorded on a regular basis for 30 days. The median survival time and the ratio of survival time for treated (T)/control (C) animals were calculated. The positive control compound was 1,4-dihydroxy-5,8-bis[[2-(2-hy-



droxyethylamino)ethyl]amino]anthraquinone dihydrochloride or Cisplatin given intraperitoneally at the indicated dose. The results appear in Table VI. If T/C X 100 (%) is 125 or over, the tested compound is considered to have significant anti-tumor activity.

TABLE VI
Lymphocytic Leukemia L1210 Test

Compound	Dose (mg/kg)	Median Survival (Days)	T/C x 100 (%)
LL-E33288 (Bromo-complex)	1.5	29	174
	0.8	28.5	171
	0.4	22.5	135
	0.2	22.5	135
Control	-	16.7	-
Anthraquinone	1.6	30	180
	0.8	30	180
	0.4	30	180
LL-E33288 β_1 -Br	0.2	11.3	136
	0.1	11.4	137
	0.05	11	133
	0.025	11.3	136
Control	-	8.3	-
Cisplatin	5	7.5	90
	2.5	12	145
	1.25	11	133

18 OCT 1996


-30-

Colon 26 Adenocarcinoma Test

The animals used were CD₂F₁ female mice weighing a minimum of 17 g and all within a 3 g weight range. There were 5 or 6 mice per test group with three groups of 5 or 6 animals used as untreated controls for each test. The tumor implant was by intraperitoneal (or subcutaneous) injection of 0.5 ml of a 2% Colon 26 tumor brei in Eagle's MEM medium containing antibiotics. LL-E33288 antibiotics were tested in the Colon 26 system as a complex (Bromo-complex) of all components. The test compounds were administered intraperitoneally on days 1, 5 and 9 (relative to tumor implant doses). The mice were weighed and deaths recorded on a regular basis for 30 days. The median survival times for treated (T)/control (C) animals were calculated. The positive control compound was Cisplatin. The results appear in Table VII. If $T/C \times 100$ (%) is 130 or over, the tested compound is considered to have significant anti-tumor activity.

TABLE VIIColon 26 Adenocarcinoma Test

Compound	Dose (mg/kg)	Median Survival (Days)	T/C x 100 (%)
LL-E33288 (Bromo-complex)	1.5	39.5	212
	0.8	32.5	175
	0.4	34	183
	0.2	25.5	137
Control	-	18.6	-
Positive Control	1	29	156
	0.5	38.5	207
	0.25	37.5	202



M5076 Sarcoma

The M5076 reticular cell Sarcoma is propagated as subcutaneous implants in C57B2/6 female mice. In the assays for antitumor activity, BDF₁ mice of either sex were inoculated intraperitoneally with 0.5 ml of a 10% tumor brei. LL-E33288 antibiotics were tested in the M5076 system as a complex (Bromo-complex) of all components. Test compounds were administered intraperitoneally on days 1, 5, 9, 13 and 17 relative to tumor inoculation on day zero. The median survival time in days was determined for each drug dose used on day 60 and the ratio of survival time for treated (T)/control (C) animals were calculated.

The results of this test appear in Table VIII compared to the results obtained with Cisplatin. If T/C X 100 (%) is 125 or over, the tested compound is considered to have significant anti-tumor activity.

TABLE VIII

M5076 Sarcoma

Compound	Dose (mg/kg)	Median Survival (Days)	T/C x 100 (%)
LL-E33288 (Bromo-complex)	1.5	50	175
	0.8	50	175
	0.4	39.5	139
Control	-	28.5	-
Cisplatin	1	30	105
	0.5	44.5	156
	0.25	45	158

18 OCT 1996

-32-

In the same manner, the following iodo-components were tested for antineoplastic activity.

TABLE IX
Lymphocytic Leukemia P388 Test

Compound	Dose (mg/kg)	Median Survival (Days)	T/C x 100 (%)
LL-E33288 γ 1-I (Test 1)	.005	>25.5	>196
	.0025	22	169
	.00125	18.5	142
	.0006	18	138
	.0003	15.5	119
	.00015	15	115
Control	-	13	-
Positive Control Novantrone®*	1.6	22.5	173
LL-E33288 γ 1-I (Test 2)	.01	11	100
	.005	18	164
	.0025	22.5	205
	.00125	18.5	168
	.0006	16	145
	.0003	14	127
	.00015	14	127
Control	-	11	-
Positive Control Novantrone®	1.6	19	173
	0.8	16	145

*1,4-dihydroxy-5,8-bis[[2-(2-hydroxyethylamino)ethyl]
amino]anthraquinone-2HCl




TABLE X

Melanotic Melanoma B16 Test

Compound	Dose (mg/kg)	Median Survival (Days)	T/C x 100 (%)
LL-E33288 γ 1-I	.0025	26.5	156
	.00125	27	159
	.0006	42.5	250
	.0003	35.5	209
	.00015	33.5	197
	.00007	30.5	179
Control	-	-	-
Adriamycin	0.8	48	282
	0.4	40	235
	0.2	34.5	203





TABLE XI

Lymphocytic Leukemia L1210 Test


Compound	Dose (mg/kg)	Median Survival (Days)	T/C x 100 (%)
LL-E33288 γ 1-I	.01	8	89
	.005	14	156
	.0025	11	122
	.0012	10.5	117
	.0006	10	111
Control	-	9	-
Positive Control Novantrone®	3.2	15	167
	1.6	11.5	128
	0.8	12	133
	0.4	11	122

TABLE XII

Colon 26 Adenocarcinoma Test



Compound	Dose (mg/kg)	Median Survival (Days)	T/C x 100 (%)
LL-E33288 Y1-I	.01	11	59
	.005	25.5	138
	.0025	27	146
	.00125	22.5	122
	.0006	23.5	127
	.0003	20	108
	.00015	17.5	95
	.00007	17	92
Control	-	18.5	-
Positive Control			
Cisplatin	2	15.5	84
	1	27.5	149
	0.5	23.5	127




General Fermentation Conditions

Cultivation of Micromonospora echinospora NRRL 15839 or NRRL 15975 may be carried out in a wide variety of liquid culture media. Media which are useful for the production of these novel antibacterial and antitumor agents include an assimilable source of carbon, such as starch, sugar, molasses, glycerol, etc.; an assimilable source of nitrogen such as protein, protein hydrolysate, polypeptides, amino acids, corn steep liquor, etc.; and inorganic anions and cations, such as potassium, sodium, ammonium, calcium, sulfate, carbonate, phosphate, chloride, etc. and sources of either bromine (sodium bromide) or iodine (potassium iodide). Trace elements such as boron, molybdenum, copper, etc., are supplied as impurities of other constituents of the media. Aeration in tanks and bottles is supplied by forcing sterile air through or onto the surface of the fermenting medium. Further agitation in tanks is provided by a mechanical impeller. An antifoam agent such as silicone may be added as needed.

General Procedure for the Isolation and Separation of the Antibiotics - LL-E33288

The LL-E33288 antibiotics are recovered from the fermentation broth by extracting the whole mash with an organic solvent such as ethyl acetate or dichloromethane. The antibiotic complex, contained in the organic extract, is further purified by selective precipitation from lower hydrocarbons. The crude LL-E-33288 antibiotic complex thus obtained is further purified and separated into the individual components by a series of column chromatographies using silica gel, Sephadex® LH-20 (Pharmacia Fine Chemicals) and C₁₈ bonded silica.

The invention will be described in greater detail in conjunction with the following non-limiting specific examples.



Example 1

Inoculum Preparation

A typical medium used to grow the primary inoculum was prepared according to the following formula:

Beef extract.....	about 0.3%
Tryptone.....	about 0.5%
Dextrose.....	about 0.5%
Dextrin.....	about 2.4%
Calcium carbonate.....	about 0.4%
Yeast extract.....	about 0.5%
Water.....qs to.....	100%

This medium was adjusted to pH 7.0 and then sterilized. A 100 ml portion of this sterile medium, in a flask, was inoculated with frozen mycelia of the culture

NRRL 15839. The inoculated medium was placed on a rotary shaker and agitated vigorously for 48 hours at 32°C. This incubated medium was then used to inoculate 10 liters of the above sterile medium in a 14 liter fermentor. This medium was incubated, with agitation, at 32°C for 48 hours, providing secondary inoculum. This secondary inoculum was then used to inoculate 300 liters of the above sterile medium in a tank and incubated for 48 hours at 30°C while agitated by an impeller driven at 180-200 rpm, providing the tertiary or seed inoculum.

Example 2

Tank Fermentation

A fermentation medium was prepared according to the following formulation:

Dextrose.....	about 0.5%
Sucrose.....	about 1.5%
Peptone, bacteriological grade.....	about 0.2%
Dibasic potassium phosphate.....	about 0.01%

18 OCT 1996

-38-

Molasses..... about 0.5%
Calcium carbonate..... about 0.5%
Source of bromine or iodine..... trace amounts
Water.....qs to..... 100%

A 2800 liter portion of the above medium was sterilized and then inoculated with 300 liters of tertiary (seed) inoculum prepared as described in Example 1. Aeration was supplied at the rate of 0.53 liters of sterile air per liter of mash per minute and agitation was supplied by an impeller driven at 110 rpm. The temperature was maintained at about 28°C and the fermentation was terminated after about 97 hours, at which time the mash was harvested.

The fermentation was monitored for production of the LL-E33288 antibiotics by antibacterial activity, biochemical induction assay, TLC and HPLC analyses.

The whole harvest mash was adjusted to pH 6 and then extracted with 1/2 mash volume ethyl acetate. The ethyl acetate extract was concentrated to a syrup which was washed twice with hexane and filtered through diatomaceous earth. The diatomaceous earth cake was thoroughly mixed with ethyl acetate and filtered. The filtrate was concentrated to 3 liters, dried over excess anhydrous sodium sulfate and then precipitated by the addition of hexane giving about 26.7 g of crude LL-E33288 complex.

Example 3

Separation of LL-E33288 α_1 -Br, α_2 -Br, α_3 -Br and α_4 -Br from


LL-E33288 β_1 -Br, β_2 -Br and γ_1 -Br

The approximately 26.7 g of crude LL-E33288 complex (Bromo-complex) from Example 2 was divided evenly into three portions and chromatographed on three separate 2.4 x 110 cm silica gel columns (Silica Woelm®, 32-63 m, Woelm Pharma) packed and equilibrated with ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate. The columns were first eluted with the same solvent at a flow rate of 3 ml/minute

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED



for 18 hours, collecting 18 ml fractions. The eluent was changed to ethyl acetate:methanol (95:5) and elution continued for 8 hours. Finally the columns were eluted with ethyl acetate:methanol (90:10) for 10 hours. The fractions were assayed by the modified biochemical induction assay (BIA). The positive fractions were analysed by TLC using silica gel 60 precoated sheets and developed with the solvent system 3% isopropanol in ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate and detected by bioautography using the modified BIA.

Fractions containing LL-E33288 α_1 -Br, α_2 -Br, α_3 -Br and α_4 -Br (LL-E33288 α -Br complex) from the three columns were pooled, concentrated to dryness and the residue was dissolved in ethyl acetate and washed with a small amount of water. The ethyl acetate solution was dried over anhydrous sodium sulfate and precipitated as before to yield about 4.2 g of crude LL-E33288 α -Br complex.

Fractions containing LL-E33288 β_2 -Br, β_1 -Br and γ_1 -Br (LL-E33288 β -Br complex containing γ -Br from the three columns were pooled and worked up as above to yield about 2.0 g of crude LL-E33288 β -Br complex containing γ -Br.

Example 4

Isolation of LL-E33288 β_1 -Br and LL-E33288 γ_1 -Br

An approximately 1.9 g sample of the LL-E33288 β -Br complex containing γ -Br from Example 3 was chromatographed on a 25 x 10 cm Sephadex® LH-20 column equilibrated with methanol:water (90:10) at a flow rate of 1.2 ml/minute, collecting 15 ml fractions. The fractions were assayed in the BIA and those active were analysed by TLC as before. Fractions 21-26 containing most of the LL-E33288 β_1 -Br, β_2 -Br and γ_1 -Br were pooled and concentrated to remove methanol and the resulting aqueous mixture was lyophilized to yield about 435 mg of partially purified complex containing approximately 10% of LL-E33288 β_1 -Br, 1% of LL-E33288 β_2 -Br and 4% of LL-E33288 γ_1 -Br.

18 OCT 1996


-40-

The above partially purified LL-E33288 β -Br complex containing γ -Br was divided evenly and chromatographed on two 1.5 x 100 cm silica gel columns (Kiesel Gel 60, 40-63 μ m, EM Products for chromatography) packed and equilibrated with ethyl acetate:methanol (98:2) at a flow rate of 1 ml/minute, collecting 12 ml fractions. The fractions were assayed and analysed by TLC as before and those containing primarily LL-E33288 β_1 -Br were pooled, concentrated and precipitated from hexane to yield about 26 mg of 80% pure LL-E33288 β_1 -Br. Those fractions containing LL-E33288 γ_1 -Br (chromatographing just after LL-E33288 β_1 -Br) were pooled and worked up to yield about 4.5 mg of 30% pure LL-E33288 γ_1 -Br. A few fractions containing LL-E33288 β_2 -Br (chromatographing just before LL-E33288 β_1 -Br), were pooled and worked up to yield a trace amount of LL-E33288 β_2 -Br.

Example 5

Final Purification of LL-E33288 β_1 -Br

The approximately 26 mg of 80% pure LL-E33288 β_1 -Br from Example 4 was combined with other LL-E33288 β_1 -Br samples of similar purity derived from other fermentations conducted under identical conditions. A total of about 38 mg of this combined β_1 -Br was further purified by reverse phase preparative TLC using Whatman PLKC $_{18}$ F, 100 m precoated TLC plates, developed with methanol:0.1M ammonium acetate buffer at pH 4.0 (90:10). The band containing LL-E33288 β_1 -Br, chromatographing at R_f =0.66 and visualized by quenching effect under short wavelength UV lamp (254 nm), was excised and the antibiotic was washed off the adsorbant with 10% isopropyl alcohol in ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate. The solution was concentrated and the residue was dissolved in ethyl acetate and washed with a small amount of water. The organic solution containing LL-E33288 β_1 -Br was worked up as before to yield about 24.5 mg of 90% pure LL-E33288 β_1 -Br. This sample was further purified by preparative TLC on silica gel (Silica



Gel GF precoated plates, 1000 m, Analtech) developed with 3% isopropyl alcohol in ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate. The major quenching band under short wavelength UV lamp (254 nm), chromatographing at $R_f=0.7$, was excised and the antibiotic was washed off the adsorbant with dichloromethane:methanol (80:20). The organic solution containing LL-E33288 β_1 -Br was worked up as before to yield about 18.8 mg of substantially pure LL-E33288 β_1 -Br.

Example 6

Final Purification of LL-E33288 β_1 -Br

The approximately 4.5 mg of 30% pure LL-E33288 γ_1 -Br from Example 4 was combined with other LL-E33288

γ_1 -Br samples of similar purity derived from other fermentations conducted under identical conditions. A total of 18 mg of this combined sample was further purified by preparative TLC on silica gel (Silica Gel GF precoated tapered plates, Analtech) developed with 2% isopropyl alcohol in ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate. The major quenching band under short wavelength UV lamp (254 nm), chromatographing at $R_f=0.5$, was excised and worked up as before to yield about 4.3 mg of substantially pure LL-E33288 γ_1 -Br.

A preferred fermentation medium for production of LL-E33288 Bromo-complex is as follows:

<u>Ingredient</u>	<u>Percent</u>
Sucrose	2.0
Ferrous Sulfate Heptahydrate	0.01
Magnesium Sulfate Heptahydrate	0.02
Calcium Carbonate	0.5
Peptone	0.2
Molasses	0.5
Sodium Bromide	0.05
Water qs to	100

18 OCT 1996

-42-

However, the addition of iodine as potassium iodide, to the fermentation medium provided substantial improvements by:


- 1) markedly enhancing vegetative growth in the fermentation;
- 2) increasing zones of inhibition in bioassays versus Escherichia coli #300 and Bacillus #308;
- 3) providing substantial activity at the R_f of LL-E33288₁-I and R_f 1-I on the bioautograph of TLC plates; and
- 4) enhancement of other components as detected by TLC.

The following two media are preferred for the production of LL-E33288 Iodo-complex:

<u>Ingredient</u>	<u>Percent</u>	
	<u>Media A</u>	<u>Media B</u>
Sucrose	2.0	2.0
Ferrous Sulfate Heptahydrate	0.01	0.01
Magnesium Sulfate Heptahydrate	0.02	0.02
Calcium Carbonate*	0.5	0.25
Peptone**	0.2	0.2
Molasses	0.5	0.5
Potassium iodide	0.05	0.01
Water qs to	100	100

* Mississippi lime.

** Best results were obtained with MARCOR® bacteriological peptone, but other peptones usable and also polypeptides from meat and casein hydrolyzates.



Example 7

A mycelial-spore suspension was prepared by scraping the surface of a slant of culture NRRL-15839 to which 5 ml of sterile distilled water had been added. This suspension was then used to inoculate 100 ml of sterile seed medium of the following formula:

Yeast Extract	0.5%
Beef Extract	0.3%
Tryptose	0.5%
Starch	2.4%
Dextrose	0.5%
Calcium Carbonate	0.4%
Water qs to	100.0%

in a 500 ml flask. This seed flask was incubated at 28°C on a rotary shaker at 200 rpm for 3-4 days, producing Stage I inoculum.

The Stage I inoculum was used to inoculate a Stage II inoculum of the same sterile medium, which was incubated under the same conditions for 2 days.

The Stage II inoculum was then used to inoculate 100 ml of sterile fermentation medium of the formula:

Sucrose	2.0%
Ferrous Sulfate Heptahydrate	0.01%
Magnesium Sulfate Heptahydrate	0.02%
Calcium Carbonate	0.5%
Peptone (MARCOR®)	0.2%
Molasses	0.5%
Potassium Iodide	0.05%
Water qs to	100.0%

This medium was incubated at 28°C on a shaker at 200 rpm for 5 days at which time the mash was harvested.

18 OCT 1996

-44-

A concentration of 4 to 20 $\mu\text{g/ml}$ of potassium iodide appears to be optimal, but concentrations of 2 mg/ml do not appear to depress yields.

NRRL-15839 can be induced to produce LL-E33288 β_1 -I when potassium iodide is present in the medium, but only at very low levels (0.2-0.3 $\mu\text{g/ml}$) as against 1.5-3.5 $\mu\text{g/ml}$ for the better producing NRRL-15975.

Yields of β_1 -I and γ_1 -I in an iodine medium are 2 to 8 times greater than yields of corresponding brominated compounds β_1 -Br and γ_1 -Br in a bromine medium using NRRL-15975.

Example 8

Separation of LL-E33288 α_1 -I, α_2 -I, and α_3 -I from
LL-E33288 β_1 -I, β_2 -I, γ_1 -I and δ_1 -I

Approximately 41.3 g of crude LL-E33288 complex derived from the processing 7500 liters of a fermentation using NRRL-15975 and medium containing inorganic iodide was divided evenly into two portions and chromatographed on two separate 2.5 x 110 cm silica gel column (Silica Woelm, 32-63 μm) packed and equilibrated with ethyl acetate. The columns were first eluted with ethyl acetate at a flow rate of 4 ml/minute for 4 hours, collecting 20 ml fractions. The eluent was changed to a concave gradient from ethyl acetate saturated with 0.1 M aqueous potassium dihydrogen phosphate to 10% isopropyl alcohol in ethyl acetate saturated with 0.1 M aqueous potassium dihydrogen phosphate over 24 hours. The columns were finally eluted with 10% isopropyl alcohol in ethyl acetate saturated with 0.1 M aqueous potassium dihydrogen phosphate over night. The fractions were assayed in the BIA and those active were analysed by TLC as described in Example 3.

Fractions (86-107) containing LL-E33288 α_3 -I from the two columns were pooled and worked up as before to yield about 2.1 g of crude LL-E33288 α_3 -I.

18 OCT 1996

-45-

Fractions (182-253) containing LL-E33288 α_1 -I and α_2 -I from the two columns were pooled and worked up to yield about 4.2 g of a crude mixture of LL-E33288 α_1 -I and α_2 -I.

Fractions (254-272) containing LL-E33288 β_2 -I and β_1 -I from the two columns were pooled and worked up to yield about 1.2 g of a crude mixture of LL-E33288 β_2 -I and β_1 -I.

Fractions (273-302) containing LL-E33288 γ_1 -I from the two columns were pooled and worked up to yield about 1.9 g of 30% pure LL-E33288 γ_1 -I.

Fractions (303-340) containing LL-E33288 δ_1 -I from the two columns were pooled and worked up to yield about 1.3 g of partially purified LL-E33288 δ_1 -I.

Example 9

Purification of LL-E33288 γ_1 -I

Approximately 900 mg of the 30% pure LL-E33288 γ_1 -I from Example 8 was chromatographed on a 2.5 x 120 cm sephadex LH-20 column equilibrated with ethyl acetate:dichloromethane:ethanol (2:2:1) at a flow rate of 1 ml/minute, collecting 12 ml fractions. The fractions were assayed and analysed by TLC as before and those containing LL-E33288 γ_1 -I (fractions 24-33) were pooled and worked up to yield 428 mg of 64% pure LL-E33288 γ_1 -I.

A 22 mg sample of the above was chromatographed on a 0.8 x 24 cm Septralyte C₁₈ (35-60 μ m, Analytichem) column equilibrated with acetonitrile:0.2 M aqueous ammonium acetate (55:45) at a flow rate of 2 ml/minute, collecting 12 ml fractions. The fractions were assayed and analysed by TLC as before and those containing pure LL-E33288 γ_1 -I were pooled and worked up to yield 7.7 mg of pure LL-E33288 γ_1 -I.


18 OCT 1996

-46-

Example 10Purification of LL-E33288 β_1 -I and β_2 -I

Approximately 600 mg of the crude mixture of LL-E33288 β_2 -I and β_1 -I from Example 8 was chromatographed on a 2.5 x 120 cm Sephadex LH-20 column equilibrated with ethyl acetate:dichloromethane:ethanol (2:2:1) at a flow rate of 1 ml/minute, collecting 12 ml fractions. The fractions were assayed and analysed by TLC as before and those containing LL-E33288 β_2 -I and LL-E33288 β_1 -I (fractions 23-31) were pooled and worked up to yield 81 mg of a partially purified mixture of LL-E33288 β_2 -I and β_1 -I.

The sample above was chromatographed on a 1.5 x 90 cm Sephadex LH-20 column equilibrated with hexane:dichloromethane:ethanol (3:1:1) at a flow rate of 0.8 ml/minute, collecting 12 ml fractions. The fractions were assayed and analysed by TLC as before and those containing LL-E33288 β_2 -I (fractions 17-30) and LL-E33288 β_1 -I (fractions 31-38) were pooled separately and worked up to yield 31 mg of partially purified LL-E33288 β_2 -I and 20 mg of 80% pure LL-E33288 β_1 -I.



-47-

WHAT IS CLAIMED IS:

1. A compound LL-E33288 α ₁-Br, having the following R_f values in the indicated solvent systems on TLC on silica gel sheets:

- a) ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, R_f=0.67;
- b) 3% isopropanol in ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, R_f=0.80; and
- c) ethyl acetate:methanol (95:5), R_f=0.79.



2. A compound LL-E33288 α ₂-Br, having the following R_f values in the indicated solvent systems on TLC on silica gel sheets:

- a) ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, R_f=0.61;
- b) 3% isopropanol in ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, R_f=0.75; and
- c) ethyl acetate:methanol (95:5), R_f=0.73.

3. A compound LL-E33288 α ₃-Br having the following R_f values in the indicated solvent systems on TLC on silica gel sheets:

18 OCT 1996

-48-

- a) ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.55$;
- b) 3% isopropanol in ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.69$; and
- c) ethyl acetate:methanol (95:5), $R_f=0.61$.


4. A compound LL-E33288 α_4 -Br, having the following R_f values in the indicated solvent systems on TLC on silica gel sheets:

- a) ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.49$;
- b) 3% isopropanol in ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.64$; and
- c) ethyl acetate:methanol (95:5), $R_f=0.54$.

5. A compound LL-E33288 β_1 -Br,



- a) having an approximate elemental analysis:
C 48.6; H 5.6; N 2.9; S 9.1 and Br 5.5;
- b) having a melting point: 146-150°C (dec.);
- c) having a specific rotation: $[\alpha]_D^{26} = -49 \pm 10^\circ$
(0.1%, ethanol);




- d) having ultraviolet absorption spectra as shown in Figure I of the drawings;
- e) having an infrared absorption spectrum as shown in Figure II of the drawings;
- f) having a proton magnetic resonance spectrum as shown in Figure III of the drawings;
- g) having a carbon-13 magnetic resonance spectrum as shown in Figure IV of the drawings with significant peaks at:

17.60(q);	17.64(q);	18.9(q);	19.7(q);
22.4(q);	22.8(q);	23.5(q);	34.3(t);
36.9(t);	39.2(t/d);	47.8(d);	51.7(q);
52.7(q);	54.6(d)	56.3(q);	57.2(q);
57.8(d);	61.0(q/d);	61.7(d);	62.4(t);
66.9(d);	68.4(d);	69.1(d);	69.7(d);
70.2(d);	71.1(d);	71.9(d);	72.1(s);
76.1(d);	81.0(d);	83.3(s);	88.2(s);
97.4(d);	99.7(d);	100.8(s);	102.5(d);
115.1(s);	123.4(d);	124.4(d);	126.5(d);
130.2(s);	130.8(s);	144.6(s);	149.3(s);
149.5(s);	191.7(s);	192.4(s);	and

- h) having the following R_f values in the indicated solvent systems on TLC on silica gel sheets:

- i) ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.24$;
- ii) 3% isopropanol in ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.35$;
- iii) ethyl acetate:methanol (95:5), $R_f=0.36$.



-50-

i) having a molecular weight: 1333/1335, respectively for $^{79}\text{Br}/^{81}\text{Br}$; and

j) having a molecular formula: $\text{C}_{54}\text{H}_{84}\text{N}_3\text{O}_{22}\text{S}_4\text{Br}$.

6. A compound LL-E3328882-Br, having the following R_f values in the indicated solvent systems on TLC on silica gel sheets:

- a) ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.32$;
- b) 3% isopropanol in ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.41$; and
- c) ethyl acetate:methanol (95:5), $R_f=0.45$.




7. A compound LL-E3328871-Br

- a) having ultraviolet absorption spectra as shown in Figure V of the drawings;
- b) having an infrared absorption spectrum as shown in Figure VI of the drawings;
- c) having a proton magnetic resonance spectrum as shown in Figure VII of the drawings; and
- d) having the following R_f values in the indicated solvent systems on TLC on silica gel sheets:
 - i) ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.18$;



-51-



ii) 3% isopropanol in ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.28$;

iii) ethyl acetate:methanol (95:5), $R_f=0.27$.

e) having a carbon-13 magnetic resonance spectrum as shown in Figure VIII of the drawings with significant peaks at:

14.4	17.6	17.9	19.0
19.7	-	22.8	-
-	34.0	37.6	39.5
42.1	-	51.6	52.7
54.1	56.3	57.3	-
59.3	61.1	61.8	61.9
67.2	68.18	68.23	69.7
70.1	70.8	71.1	71.7
71.8	76.1	-	81.0
82.9	88.4	-	97.8
100.0	100.2	101.3	103.0
115.3	123.0	124.9	126.9
130.4	131.1	131.8	138.0
144.7	-	149.5	149.6
155.6	192.5	192.9	

f) having a molecular formula: $C_{53}H_{82}N_3O_{22}S_4Br$; and

g) having a molecular weight: 1319/1321, respectively for $^{79}Br/^{81}Br$.

8. A compound LL-E33288 α_1 -I

a) having the following R_f values in the indicated solvent systems on TLC on silica gel sheets:

i) ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.67$;



18 OCT 1996

- ii) 3% isopropanol in ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.80$; and
 - iii) ethyl acetate:methanol (95:5), $R_f=0.80$; and
 - b) having a molecular weight:1145.
9. A compound LL-E33288 α_2 -I
- a) having the following R_f values in the indicated solvent systems on TLC on silica gel sheets:
 - i) ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.61$;
 - ii) 3% isopropanol in ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.75$; and
 - iii) ethyl acetate:methanol (95:5), $R_f=0.73$;
 - b) containing only the following elements: C, H, N, O, S and I;
 - c) having a molecular weight:1131; and
 - d) having a proton magnetic resonance spectrum as shown in Figure IX of the drawings.

10. A compound LL-E33288 α_3 -I
- a) having the following R_f values in the indicated solvent systems on TLC on silica gel sheets:
 - i) ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.55$;

18 OCT 1996



-53-

- ii) 3% isopropanol in ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.69$; and
 - iii) ethyl acetate:methanol (95:5), $R_f=0.61$;
- b) having a molecular weight:1066; and
- c) having a proton magnetic resonance spectrum as shown in Figure X of the drawings.

11. A compound LL-E332888₁-I

- a) having the following R_f values in the indicated solvent systems on TLC on silica gel sheets:
- i) ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.24$;
 - ii) 3% isopropanol in ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.35$; and
 - iii) ethyl acetate:methanol (95:5), $R_f=0.36$;
- b) having an ultraviolet absorption spectra as shown in Figure XI of the drawings;
- c) having an infrared absorption spectrum as shown in Figure XII of the drawings;
- d) having a proton magnetic resonance spectrum as shown in Figure XIII of the drawings;
- e) having a carbon 13 magnetic resonance spectrum as shown in Figure XIV of the drawings with significant peaks at

18 OCT 1996

-54-




-	17.5	17.6	18.9
-	22.4	22.8	23.4
25.4	34.3	36.9	39.2
-	47.9	51.6	52.8
54.8	56.3	57.2	57.9
60.9		61.6	62.2
67.0	68.4	68.4	69.1
69.6	70.4	71.1	71.8
72.2	76.2	-	80.8
83.3	88.1	93.6	97.4
99.6	99.6	-	102.6
112.4	123.4	124.4	126.4
-	-	133.4	-
-	-	-	-
-	192.3	192.6	

f) having a molecular formula: $C_{54}H_{84}N_3O_{22}S_4I$; and

g) having a molecular weight: 1381.

12. A compound LL-E33288^B₂-I having the following R_f values in the indicated solvent systems on TLC on silica gel sheets:

- a) ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.32$.
- b) 3% isopropanol in ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.41$; and
- c) ethyl acetate:methanol (95:5), $R_f=0.45$.



13. A compound LL-E33288Y1-I

a) having the following R_f values in the indicated solvent systems on TLC on silica gel sheets:

- i) ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f = 0.18$;
- ii) 3% isopropanol in ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f = 0.28$; and
- iii) ethyl acetate: methanol (95:5), $R_f = 0.27$;



b) containing only the following elements: C, H, N, O, S and I;

c) having an approximate elemental analysis: C 48.8; H 5.4; N 2.8; S 9.0; and I 9.2;

d) having a molecular weight: 1367;

e) having a molecular formula: $C_{53} H_{82} N_3 O_{22} S_4 I$;

f) having an ultraviolet absorption spectra as shown in Figure XV of the drawings;

g) having an infrared absorption spectrum as shown in Figure XVI of the drawings;

h) having a proton magnetic resonance spectrum as shown in Figure XVII of the drawings; and

i) having a carbon 13 magnetic resonance spectrum as shown in Figure XVIII of the drawings, significant peaks as listed below:

14.5(q)	17.6(q)	17.6(q)	18.9(q)
-	-	22.8(q)	-
25.4(q)	34.1(t)	37.0(t)	39.1(t)
42.3(t/s)	-	51.5(d)	52.8(q)
54.8(t)	56.3(q)	57.2(q)	-
60.4(d)	60.9(q)	61.3(t)	61.7(q)
67.0(d)	68.4(d)	68.5(d)	69.2(d)
69.7(d)	70.5(d)	71.1(d)	71.8(d)
72.1(s)	75.7(d)	75.8(d)	80.9(d)
82.8(s)	88.1(s)	93.5(s)	97.3(d)
99.6(d)	99.7(d)	100.8(s)	102.6(d)
-	123.4(d)	124.4(d)	126.2(d)
130.2(s)	131.0(s)	133.4(s)	139.1(s)
143.0(s)	145.1	150.6(s)	151.5(s)
154.5	192.0(s)	192.5(s)	



18 OCT 1996

14.) A compound LL-33288 δ ₁-I having the following R_f values in the indicated solvent systems on TLC on silica gel sheets:

a) ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, R_f = 0.11; and

b) 3% isopropanol in ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, R_f = 0.19.

15. A method of treating bacterial infections in warmblooded animals which comprises administering to said animals an antibacterially effective amount of a compound selected from the group consisting of LL-E33288 α ₁-Br; LL-E33288 α ₁-I; LL-E33288 α ₂-Br; LL-E33288 α ₂-I; LL-E33288 α ₃-Br; LL-E33288 α ₃-I; LL-E33288 α ₄-Br; LL-E33288 β ₁-Br; LL-E33288 β ₁-I; LL-E33288 β ₂-Br; LL-E33288 β ₂-I; LL-E33288 γ ₁-Br; LL-E33288 γ ₁-I; and LL-E33288 δ ₁-I.


16. A method of inhibiting the growth of tumors in a mammal comprising administering to said mammal an effective amount of a compound selected from the group consisting of LL-E33288 α ₁-Br; LL-E33288 α ₁-I; LL-E33288 α ₂-Br; LL-E33288 α ₂-I; LL-E33288 α ₃-Br; LL-E33288 α ₃-I; LL-E33288 α ₄-Br; LL-E33288 β ₁-Br; LL-E33288 β ₁-I; LL-E33288 β ₂-Br; LL-E33288 β ₂-I; LL-E33288 γ ₁-Br; LL-E33288 γ ₁-I; and LL-E33288 δ ₁-I.

17. A method of regressing leukemia in a mammal comprising administering to said mammal an effective amount of a compound selected from the group consisting of LL-E33288 α ₁-Br; LL-E33288 α ₁-I; LL-E33288 α ₂-Br; LL-E33288 α ₂-I; LL-E33288 α ₃-Br; LL-E33288 α ₃-I; LL-E33288 α ₄-Br; LL-E33288 β ₁-Br; LL-E33288 β ₁-I; LL-E33288 β ₂-Br; LL-E33288 β ₂-I; LL-E33288 γ ₁-Br; LL-E33288 γ ₁-I; and LL-E33288 δ ₁-I.

18. A process for producing antibiotics LL-E33288 α ₁-Br; LL-E33288 α ₂-Br; LL-E33288 α ₃-Br; LL-E33288 α ₄-Br; LL-E33288 β ₁-Br; LL-E33288 β ₂-Br and LL-E33288 γ ₁-Br which comprises aerobically fermenting the organism Micromonospora echinospora ssp. calichensis NRRL 15839 or mutants thereof including NRRL 15975 in a liquid medium containing assimilable sources of carbon, nitrogen,



THIS SHEET HAS BEEN REPLACED



bromine and inorganic salts, until substantial antibiotic activity is imparted to said medium and then recovering the antibiotics therefrom.

19. A process for producing antibiotics LL-E33288 α_1 -Br; LL-E33288 α_2 -Br; LL-E33288 α_3 -Br; LL-E33288 α_4 -Br; LL-E33288 β_1 -Br; LL-E33288 β_2 -Br; and LL-E33288 γ_1 -Br which comprises aerobically fermenting a liquid medium containing assimilable sources of carbon, nitrogen, bromine and inorganic salts; which medium has been inoculated with a viable culture of the organism Micromonospora echinospora ssp. calichensis NRRL 15839 or mutants thereof including NRRL 15975, maintaining said fermentation culture at a temperature of about 24-32°C for a period of approximately 90-200 hours, harvesting the mash and extracting the antibiotics.

20. A process for producing antibiotics LL-E33288 α_1 -I; LL-E33288 α_2 -I; LL-E33288 α_3 -I; LL-E33288 β_1 -I; LL-E33288 β_2 -I; LL-E33288 γ_1 -I; and LL-E33288 δ_1 -I which comprises aerobically fermenting the organism Micromonospora echniospora ssp. calichensis NRRL 15839 or mutants thereof including NRRL 15975 in a liquid medium containing assimilable sources of carbon, nitrogen, iodine and inorganic salts, until substantial antibiotic activity is imparted to said medium and then recovering the antibiotics therefrom.


21. A process for producing antibiotics LL-E33288 α_1 -I; LL-E33288 α_2 -I; LL-E33288 α_3 -I; LL-E33288 β_1 -I; LL-E33288 β_2 -I; LL-E33288 γ_1 -I; and LL-E33288 δ_1 -I which comprises aerobically fermenting a liquid medium containing assimilable sources of carbon, nitrogen, iodine and inorganic salts, which medium has been inoculated with a viable culture of the microorganism Micromonospora echinospora ssp. calichensis NRRL 15839 or mutants thereof including NRRL 15975, maintaining said fermentation culture at a temperature of about 24-32°C for a period of approximately 90-200 hours, harvesting the mash and extracting the antibiotics.

22. A culture containing the microorganism






-58-



Micromonospora echinospora ssp. calichensis, NRRL 15839, said culture being capable of producing the LL-E33288 complex in recoverable quantity upon aerobic fermentation in an aqueous medium containing assimilable sources of carbon nitrogen, inorganic salts, and either iodine or bromine or both.

23. A culture containing the microorganism Micro-monospora echinospora ssp. calichensis NRRL 15975, said culture being capable of producing the LL-E33288 complex in recoverable quantity upon aerobic fermentation in an aqueous medium containing assimilable sources of carbon, nitrogen, inorganic salts, and either iodine or bromine or both.



ABSTRACT

ANTITUMOR ANTIBIOTICS (LL-E33288 COMPLEX)

Antibacterial and antitumor agents designated LL-E33288 complex and their production by new strains of *Micromonospora*, *Micromonospora echinospora* ssp. *calichensis*, NRRL-15839 and NRRL-15975, are disclosed.

FIGURE 1

REPLACEMENT SHEET

18 OCT 1996

1/18

ULTRAVIOLET ABSORPTION SPECTRA OF LL-E33288 β ,
20 $\mu\text{g/ml}$ SOLUTION




FIGURE I

2/18

INFRARED ABSORPTION SPECTRUM OF LL-E33288 β_1




FIGURE II

REPLACEMENT SHEET
18 OCT 1996

3/18

PROTON MAGNETIC RESONANCE SPECTRUM
OF LL-E 33288 *g*.




FIGURE III

REPLACEMENT SHEET

18 OCT 1996

4/18

CARBON 13 MAGNETIC RESONANCE SPECTRUM
OF LL-E33288 β .




FIGURE IV

18 OCT 1996

5/18

ULTRAVIOLET ABSORPTION SPECTRA OF LL-E33288 γ
20 $\mu\text{g}/\text{ml}$ SOLUTION




FIGURE V

REPLACEMENT SHEET
18 OCT 1996

6/18

INFRARED ABSORPTION SPECTRUM OF LL-E33288 γ




FIGURE VI

REPLACEMENT SHEET
18 OCT 1996

7/18

PROTON MAGNETIC RESONANCE SPECTRUM OF
LL-E33288 γ




FIGURE VII



8/18

^{13}C NMR OF LL-E33288 γ , Br




FIGURE VIII

REPLACEMENT SHEET
18 OCT 1996

9/18

PMR OF LL-E33288 α_2 I




FIGURE IX

REPLACEMENT SHEET
18 OCT 1996

10/18

PMR OF LL-E33288 α_3 I




FIGURE X

18 OCT 1996

11/18

UV OF LL-E33288 β_1 I





FIGURE XI

REPLACEMENT SHEET

18 OCT 1996

12/18

IR OF LL-E33288, I



WAVENUMBERS

FIGURE XII

REPLACEMENT SHEET

18 OCT 1996

13/18

PMR OF LL-E33288 β , I




FIGURE XIII

REPLACEMENT SHEET

18 OCT 1996

14/18

^{13}C NMR OF LL-E33288 β_1 I





FIGURE XIV

REPLACEMENT SHEET

18 OCT 1996

15/18

UV OF LL-E33288X, I



REPLACEMENT SHEET
18 OCT 1996

16 / 18

IR OF LL-E33288Y, I




FIGURE XVI

18 OCT 1996

17/18

PMR OF LL-E33288X, I




FIGURE XVII

REPLACEMENT SHEET

18 OCT 1996

18 / 18

^{13}C NMR OF LL-E33288X, I




FIGURE XVIII

08 FEB 1999

-56-



14.) A compound LL-33288 δ ₁-I having the following R_f values in the indicated solvent systems on TLC on silica gel sheets:

a) ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, R_f = 0.11; and

b) 3% isopropanol in ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, R_f = 0.19.

15. Use of a compound selected from the group consisting of LL-E33288 α ₁-Br; LL-E33288 α ₁-I; LL-E33288 α ₂-Br; LL-E33288 α ₂-I; LL-E33288 α ₃-Br; LL-E33288 α ₃-I; LL-E33288 α ₄-Br; LL-E33288 δ ₁-Br; LL-E33288 δ ₁-I; LL-E33288 δ ₂-Br; LL-E33288 δ ₂-I; LL-E33288 γ ₁-Br; LL-E33288 γ ₁-I; and LL-E33288 δ ₁-I for the manufacture of medicaments for treating bacterial infections in warmblooded animals.


16. Use of a compound selected from the group consisting of LL-E33288 α ₁-Br; LL-E33288 α ₁-I; LL-E33288 α ₂-Br; LL-E33288 α ₂-I; LL-E33288 α ₃-Br; LL-E33288 α ₃-I; LL-E33288 α ₄-Br; LL-E33288 δ ₁-Br; LL-E33288 δ ₁-I; LL-E33288 δ ₂-Br; LL-E33288 δ ₂-I; LL-E33288 γ ₁-Br; LL-E33288 γ ₁-I; and LL-E33288 δ ₁-I for the manufacture of medicaments for inhibiting the growth of tumors in a mammal.

17. Use of a compound selected from the group consisting of LL-E33288 α ₁-Br; LL-E33288 α ₁-I; LL-E33288 α ₂-Br; LL-E33288 α ₂-I; LL-E33288 α ₃-Br; LL-E33288 α ₃-I; LL-E33288 α ₄-Br; LL-E33288 δ ₁-Br; LL-E33288 δ ₁-I; LL-E33288 δ ₂-Br; LL-E33288 δ ₂-I; LL-E33288 γ ₁-Br; LL-E33288 γ ₁-I; and LL-E33288 δ ₁-I for the manufacture of medicaments for regressing leukemia in a mammal.

18. A process for producing antibiotics LL-E33288 α ₁-Br; LL-E33288 α ₂-Br; LL-E33288 α ₃-Br; LL-E33288 α ₄-Br; LL-E33288 δ ₁-Br; LL-E33288 δ ₂-Br and LL-E33288 γ ₁-Br which comprises aerobically fermenting the organism Micromonospora echinospora ssp. calichensis NRRL 15839 or mutants thereof including NRRL 15975 in a liquid medium containing assimilable sources of carbon, nitrogen,

This Sheet Has Been Replaced

THIS SHEET HAS BEEN
REPLACED



Title: ANTITUMOR ANTIBIOTICS
(LL-E33288 COMPLEX)

SUMMARY OF THE INVENTION

This application is a continuation-in-part of co-pending application S.N. 672,031, filed November 16, 1984.

This invention relates to new antibacterial and antitumor agents designated LL-E33288 α_1 -Br, LL-E33288 α_1 -I, LL-E33288 α_2 -Br, LL-E33288 α_2 -I, LL-E33288 α_3 -Br, LL-E33288 α_3 -I, LL-E33288 α_4 -Br, LL-E33288 β_1 -Br, LL-E33288 β_1 -I, LL-E33288 β_2 -Br, LL-E33288 β_2 -I, LL-E33288 γ_1 -Br, LL-E33288 γ_1 -I and LL-E33288 δ_1 -I, to their production by fermentation, to methods for their recovery and concentration from crude solutions and to processes for their purification. The present invention includes within its scope the antibacterial and antitumor agents in dilute form, as crude concentrates, as a complex of various or all components, in pure form as individual components and novel strains of Micromonospora.

The LL-E33288 antibiotics of this invention are closely related compounds. The fourteen antibiotics are recovered from fermentation and are initially obtained as a mixture, hereinafter either the LL-E33288 complex, the the LL-E33288 Iodo-complex or the LL-E33288 Bromo-complex. In general, the iodine containing components of the LL-E33288 antibiotics (e.g., α_1 -I, α_2 -I, α_3 -I, β_1 -I, β_2 -I, γ_1 -I and δ_1 -I) are found only in fermentations using media containing inorganic or organic iodide while the bromine containing components (e.g., α_1 -Br, α_2 -Br, α_3 -Br, α_4 -Br, β_1 -Br, β_2 -Br and γ_1 -Br) are found only in fermentations using media containing inorganic or organic bromide. The ratio of components in the LL-E33288 complex will vary, depending upon

18 AUG 1999

14.4	17.6	17.9	19.0
19.7	-	22.8	-
-	34.0	37.6	39.5
42.1	-	51.6	52.7
54.1	56.3	57.3	-
59.3	61.1	61.8	61.9
67.2	68.18	68.23	69.7
70.1	70.8	71.1	71.7
71.8	76.1	-	81.0
82.9	88.4	-	97.8
100.0	100.2	101.3	103.0
115.3	123.0	124.9	126.9
130.4	131.1	131.8	138.0
144.7	-	149.5	149.6
155.6	192.5	192.9	

- 5) Molecular formula: $C_{53}H_{82}N_3O_{22}S_4Br$ by comparing its UV, IR, 1H NMR, and ^{13}C NMR data to those of LL-E33288 δ $_1$ -BR and LL-E33288 γ $_1$ -I; and
- 6) Molecular weight: 1319/1321 respectively for ^{79}Br / ^{81}Br , calculated from its molecular formula.

The physico-chemical characteristics of LL-E33288 α $_1$ -I, LL-E33288 α $_2$ -I, LL-E33288 α $_3$ -I, and LL-E33288 δ $_1$ -I and LL-E33288 γ $_1$ -I are described below:

LL-E33288 α $_1$ -I

- 1) Molecular weight: 1145, determined by FAB-MS.

LL-E33288 α $_2$ -I

- 1) Contains and only contains the following elements by electron spectroscopy for chemical analysis (ESCA): C, H, N, O, S, I;
- 2) Molecular weight: 1131, determined by FAB-MS; and
- 3) Proton magnetic resonance spectrum: as shown in Figure IX (300 MHz, $CDCl_3$).


LL-E33288 α $_3$ -I

- 1) Molecular weight: 1066, determined by FAB-MS; and
- 2) Proton magnetic resonance spectrum: as shown in

18 AUG 1999

The new antibacterial and antitumor agents designated LL-E33288 α_1 -Br, LL-E33288 α_2 -Br, LL-E33288 α_3 -Br, LL-E33288 α_4 -Br, LL-E33288 β_1 -Br, LL-E33288 δ_2 -Br, LL-E33288 γ_1 -Br, LL-E33288 α_1 -I, LL-E33288 α_2 -I, LL-E33288 α_3 -I, LL-E33288 β_1 -I, LL-E33288 β_2 -I, LL-E33288 γ_1 -I and LL-E33288 δ_1 -I are formed during the cultivation under controlled conditions of a new strain of Micromonospora echinospora ssp. calichensis. This microorganism is maintained in the culture collection of the Medical Research Division, American Cyanamid Company, Pearl River, New York as culture number LL-E33288. A viable culture of this new microorganism has been deposited with the Culture Collection Laboratory, Northern Regional Research Center, U. S. Department of Agriculture, Peoria, Illinois on August 9, 1984, and has been added to its permanent collection. It has been assigned by such depository the strain designation NRRL 15839. Access to such culture, under strain designation NRRL 15839, during pendency of the instant application shall be available to one determined by the Commissioner of Patents and Trademarks to be entitled thereto under 37 C.F.R. §1.14 and 35 U.S.C. §122, and all restrictions on availability to the public of such culture will be irrevocably removed upon grant of a patent on the instant application.

Culture LL-E33288 was isolated from a caliche clay soil sample collected in Texas.



11A

According to a first aspect of the invention there is provided a

compound LL-E33288 α_1 -Br, having the following R_f values in the indicated solvent systems on TLC on silica gel sheets:

- a) ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.67$;
- b) 3% isopropanol in ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.80$; and
- c) ethyl acetate:methanol (95:5), $R_f=0.79$.


The invention further provides a

compound LL-E33288 α_2 -Br, having the following R_f values in the indicated solvent systems on TLC on silica gel sheets:

- a) ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.61$;
- b) 3% isopropanol in ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.75$; and
- c) ethyl acetate:methanol (95:5), $R_f=0.73$.

The invention further provides a

compound LL-E33288 α_3 -Br having the following R_f values in the indicated solvent systems on TLC on silica gel sheets:



11B

- a) ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.55$;
- b) 3% isopropanol in ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.69$; and
- c) ethyl acetate:methanol (95:5), $R_f=0.61$.

The invention further provides a compound LL-E33288 α_4 -Br, having the following R_f values in the indicated solvent systems on TLC on silica gel sheets:

- a) ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.49$;
- b) 3% isopropanol in ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.64$; and
- c) ethyl acetate:methanol (95:5), $R_f=0.54$.

The invention further provides a compound LL-E33288 β_1 -Br,

- a) having an approximate elemental analysis: C 48.6; H 5.6; N 2.9; S 9.1 and Br 5.5;
- b) having a melting point: 146-150°C (dec.);
- c) having a specific rotation: $[\alpha]_D^{26} = -49 \pm 10^\circ$ (0.1%, ethanol);

18 AUG 1999

11C

- d) having ultraviolet absorption spectra as shown in Figure I of the drawings;
- e) having an infrared absorption spectrum as shown in Figure II of the drawings;
- f) having a proton magnetic resonance spectrum as shown in Figure III of the drawings;
- g) having a carbon-13 magnetic resonance spectrum as shown in Figure IV of the drawings with significant peaks at:

17.60(q);	17.64(q);	18.9(q);	19.7(q);
22.4(q);	22.8(q);	23.5(q);	34.3(t);
36.9(t);	39.2(t/d);	47.8(d);	51.7(q);
52.7(q);	54.6(d);	56.3(q);	57.2(q);
57.8(d);	61.0(q/d);	61.7(d);	62.4(t);
66.9(d);	68.4(d);	69.1(d);	69.7(d);
70.2(d);	71.1(d);	71.9(d);	72.1(s);
76.1(d);	81.0(d);	83.3(s);	88.2(s);
97.4(d);	99.7(d);	100.8(s);	102.5(d);
115.1(s);	123.4(d);	124.4(d);	126.5(d);
130.2(s);	130.8(s);	144.6(s);	149.3(s);
149.5(s);	191.7(s);	192.4(s);	and

- h) having the following R_f values in the indicated solvent systems on TLC on silica gel sheets:

- i) ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.24$;
- ii) 3% isopropanol in ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.35$;
- iii) ethyl acetate:methanol (95:5), $R_f=0.36$.

18 AUG 1999

11D

i) having a molecular weight: 1333/1335, respectively for $^{79}\text{Br}/^{81}\text{Br}$; and

j) having a molecular formula: $\text{C}_{54}\text{H}_{84}\text{N}_3\text{O}_{22}\text{S}_4\text{Br}$.

The invention further provides a

compound LL-E33288 β -Br, having the following R_f values in the indicated solvent systems on TLC on silica gel sheets:

a) ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.32$;

b) 3% isopropanol in ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.41$; and

c) ethyl acetate:methanol (95:5), $R_f=0.45$.

The invention further provides a

compound LL-E33288 γ -Br


a) having ultraviolet absorption spectra as shown in Figure V of the drawings;

b) having an infrared absorption spectrum as shown in Figure VI of the drawings;

c) having a proton magnetic resonance spectrum as shown in Figure VII of the drawings; and

d) having the following R_f values in the indicated solvent systems on TLC on silica gel sheets:

i) ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.18$;



11E

ii) 3% isopropanol in ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.28$;

iii) ethyl acetate:methanol (95:5), $R_f=0.27$.

e) having a carbon-13 magnetic resonance spectrum as shown in Figure VIII of the drawings with significant peaks at:

14.4	17.6	17.9	19.0
19.7	-	22.8	-
-	34.0	37.6	39.5
42.1	-	51.6	52.7
54.1	56.3	57.3	-
59.3	61.1	61.8	61.9
67.2	68.18	68.23	69.7
70.1	70.8	71.1	71.7
71.8	76.1	-	81.0
82.9	88.4	-	97.8
100.0	100.2	101.3	103.0
115.3	123.0	124.9	126.9
130.4	131.1	131.8	138.0
144.7	-	149.5	149.6
155.6	192.5	192.9	

f) having a molecular formula: $C_{53}H_{82}N_3O_{22}S_4Br$; and

g) having a molecular weight: 1319/1321, respectively for $^{79}Br/^{81}Br$.

The invention further provides a
compound LL-E33288 α_1 -I

a) having the following R_f values in the indicated solvent systems on TLC on silica gel sheets:

i) ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.67$;

18 AUG 1999

11F

ii) 3% isopropanol in ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.80$; and

iii) ethyl acetate:methanol (95:5), $R_f=0.80$; and

b) having a molecular weight:1145.

The invention further provides a
compound LL-E33288 α_2 -I

a) having the following R_f values in the indicated solvent systems on TLC on silica gel sheets:

i) ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.61$;

ii) 3% isopropanol in ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.75$; and

iii) ethyl acetate:methanol (95:5), $R_f=0.73$;

b) containing only the following elements: C, H, N, O, S and I;


c) having a molecular weight:1131; and

d) having a proton magnetic resonance spectrum as shown in Figure IX of the drawings.

The invention further provides a
compound LL-E33288 α_3 -I

a) having the following R_f values in the indicated solvent systems on TLC on silica gel sheets:

i) ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.55$;



11G

- ii) 3% isopropanol in ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.69$; and
- iii) ethyl acetate:methanol (95:5), $R_f=0.61$;

b) having a molecular weight:1066; and

c) having a proton magnetic resonance spectrum as shown in Figure X of the drawings.

The invention further provides a
compound LL-E332888₁-I

a) having the following R_f values in the indicated solvent systems on TLC on silica gel sheets:

i) ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.24$;

ii) 3% isopropanol in ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.35$; and

iii) ethyl acetate:methanol (95:5), $R_f=0.36$;

b) having an ultraviolet absorption spectra as shown in Figure XI of the drawings;

c) having an infrared absorption spectrum as shown in Figure XII of the drawings;

d) having a proton magnetic resonance spectrum as shown in Figure XIII of the drawings;

e) having a carbon 13 magnetic resonance spectrum as shown in Figure XIV of the drawings with significant peaks at



11H


-	17.5	17.6	18.9
-	22.4	22.8	23.4
25.4	34.3	36.9	39.2
-	47.9	51.6	52.8
54.8	56.3	57.2	57.9
60.9		61.6	62.2
67.0	68.4	68.4	69.1
69.6	70.4	71.1	71.8
72.2	76.2	-	80.8
83.3	88.1	93.6	97.4
99.6	99.6	-	102.6
112.4	123.4	124.4	126.4
-	-	133.4	-
-	-	-	-
-	192.3	192.6	-

f) having a molecular formula: $C_{54}H_{84}N_3O_{22}S_4I$; and

g) having a molecular weight: 1381.

The invention further provides a
compound LL-E33288^B₂-I having the
following R_f values in the indicated solvent systems
on TLC on silica gel sheets:


- ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.32$.
- 3% isopropanol in ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f=0.41$; and
- ethyl acetate:methanol (95:5), $R_f=0.45$.



The invention further provides a compound LL-E33288Y1-I

- a) having the following R_f values in the indicated solvent systems on TLC on silica gel sheets:
- ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f = 0.18$;
 - 3% isopropanol in ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, $R_f = 0.28$; and
 - ethyl acetate: methanol (95:5), $R_f = 0.27$;
- b) containing only the following elements: C, H, N, O, S and I;
- c) having an approximate elemental analysis: C 48.8; H 5.4; N 2.8; S 9.0; and I 9.2;
- d) having a molecular weight: 1367;
- e) having a molecular formula: $C_{53} H_{82} N_3 O_{22} S_4 I$;
- f) having an ultraviolet absorption spectra as shown in Figure XV of the drawings;
- g) having an infrared absorption spectrum as shown in Figure XVI of the drawings;
- h) having a proton magnetic resonance spectrum as shown in Figure XVII of the drawings; and
- i) having a carbon 13 magnetic resonance spectrum as shown in Figure XVIII of the drawings, significant peaks as listed below:

14.5(q)	17.6(q)	17.6(q)	18.9(q)
-	-	22.8(q)	-
25.4(q)	34.1(t)	37.0(t)	39.1(t)
42.3(t/s)	-	51.5(d)	52.8(q)
54.8(t)	56.3(q)	57.2(q)	-
60.4(d)	60.9(q)	61.3(t)	61.7(q)
67.0(d)	68.4(d)	68.5(d)	69.2(d)
69.7(d)	70.5(d)	71.1(d)	71.8(d)
72.1(s)	75.7(d)	75.8(d)	80.9(d)
82.8(s)	88.1(s)	93.5(s)	97.3(d)
99.6(d)	99.7(d)	100.8(s)	102.6(d)
-	123.4(d)	124.4(d)	126.2(d)
130.2(s)	131.0(s)	133.4(s)	139.1(s)
143.0(s)	145.1	150.6(s)	151.5(s)
154.5	192.0(s)	192.5(s)	



11J

The invention further provides a compound LL-E33288 δ ₁-I having the following R_f values in the indicated solvent systems on TLC on silica gel sheets:

a) ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, R_f = 0.11; and

b) 3% isopropanol in ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, R_f = 0.19.

The invention further provides the use of a compound

selected from the group consisting of LL-E33288 α ₁-Br; LL-E33288 α ₁-I; LL-E33288 α ₂-Br; LL-E33288 α ₂-I; LL-E33288 α ₃-Br; LL-E33288 α ₃-I; LL-E33288 α ₄-Br; LL-E33288 β ₁-Br; LL-E33288 β ₁-I; LL-E33288 β ₂-Br; LL-E33288 β ₂-I; LL-E33288 γ ₁-Br; LL-E33288 γ ₁-I; and LL-E33288 δ ₁-I for the manufacture of medicaments for treating bacterial infections in warmblooded animals.

The invention further provides the use

of a compound selected from the group consisting of LL-E33288 α ₁-Br; LL-E33288 α ₁-I; LL-E33288 α ₂-Br; LL-E33288 α ₂-I; LL-E33288 α ₃-Br; LL-E33288 α ₃-I; LL-E33288 α ₄-Br; LL-E33288 β ₁-Br; LL-E33288 β ₁-I; LL-E33288 β ₂-Br; LL-E33288 β ₂-I; LL-E33288 γ ₁-Br; LL-E33288 γ ₁-I; and LL-E33288 δ ₁-I for the manufacture of medicaments for inhibiting the growth of tumors in a mammal.

The invention further provides the use

of a compound selected from the group consisting of LL-E33288 α ₁-Br; LL-E33288 α ₁-I; LL-E33288 α ₂-Br; LL-E33288 α ₂-I; LL-E33288 α ₃-Br; LL-E33288 α ₃-I; LL-E33288 α ₄-Br; LL-E33288 β ₁-Br; LL-E33288 β ₁-I; LL-E33288 β ₂-Br; LL-E33288 β ₂-I; LL-E33288 γ ₁-Br; LL-E33288 γ ₁-I; and LL-E33288 δ ₁-I for the manufacture of medicaments for regressing leukemia in a mammal.

The invention further provides a

process for producing antibiotics LL-E33288 α ₁-Br; LL-E33288 α ₂-Br; LL-E33288 α ₃-Br; LL-E33288 α ₄-Br; LL-E33288 β ₁-Br; LL-E33288 β ₂-Br and LL-E33288 γ ₁-Br which comprises aerobically fermenting the organism Micromonospora echinospora ssp. calichensis NRRL 15839 or mutants thereof including NRRL 15975 in a liquid medium containing assimilable sources of carbon, nitrogen,

18 AUG 1999

11K

bromine and inorganic salts, until substantial antibiotic activity is imparted to said medium and then recovering the antibiotics therefrom.

The invention further provides a process for producing antibiotics

LL-E33288 α ₁-Br; LL-E33288 α ₂-Br; LL-E33288 α ₃-Br; LL-E33288 α ₄-Br; LL-E33288 β ₁-Br; LL-E33288 β ₂-Br; and LL-E33288 γ ₁-Br which comprises aerobically fermenting a liquid medium containing assimilable sources of carbon, nitrogen, bromine and inorganic salts; which medium has been inoculated with a viable culture of the organism Micromonospora echinospora ssp. calichensis NRRL 15839 or mutants thereof including NRRL 15975, maintaining said fermentation culture at a temperature of about 24-32°C for a period of approximately 90-200 hours, harvesting the mash and extracting the antibiotics.


The invention further provides a process for producing antibiotics

LL-E33288 α ₁-I; LL-E33288 α ₂-I; LL-E33288 α ₃-I; LL-E33288 β ₁-I; LL-E33288 β ₂-I; LL-E33288 γ ₁-I; and LL-E33288 δ ₁-I which comprises aerobically fermenting the organism Micromonospora echniospora ssp. calichensis NRRL 15839 or mutants thereof including NRRL 15975 in a liquid medium containing assimilable sources of carbon, nitrogen, iodine and inorganic salts, until substantial antibiotic activity is imparted to said medium and then recovering the antibiotics therefrom.

The invention further provides a process for producing antibiotics

LL-E33288 α ₁-I; LL-E33288 α ₂-I; LL-E33288 α ₃-I; LL-E33288 β ₁-I; LL-E33288 β ₂-I; LL-E33288 γ ₁-I; and LL-E33288 δ ₁-I which comprises aerobically fermenting a liquid medium containing assimilable sources of carbon, nitrogen, iodine and inorganic salts, which medium has been inoculated with a viable culture of the microorganism Micromonospora echinospora ssp. calichensis NRRL 15839 or mutants thereof including NRRL 15975, maintaining said fermentation culture at a temperature of about 24-32°C for a period of approximately 90-200 hours, harvesting the mash and extracting the antibiotics.


The invention further provides a culture containing the microorganism



11L

Micromonospora echinospora ssp. calichensis, NRRL 15839, said culture being capable of producing the LL-E33288 complex in recoverable quantity upon aerobic fermentation in an aqueous medium containing assimilable sources of carbon nitrogen, inorganic salts, and either iodine or bromine or both.

The invention further provides a culture containing the microorganism Micro-monospora echinospora ssp. calichensis NRRL 15975, said culture being capable of producing the LL-E33288 complex in recoverable quantity upon aerobic fermentation in an aqueous medium containing assimilable sources of carbon, nitrogen, inorganic salts, and either iodine or bromine or both.



P388, melanotic melanoma B16, L1210 leukemia and colon 26 adenocarcinoma are particularly significant to the present invention. These neoplasms are utilized for testing as transplantable tumors in mice. Generally, significant anti-tumor activity, shown in these protocols by a percentage increase of mean survival times of the treated animals (T) over the control animals (C), is indicative of similar results in human leukemias and solid tumors.

Lymphocytic Leukemia P388 Test

The animals used were BDF₁ mice, all of one sex, weighing a minimum of 17 g and all within a 3 g weight range. There were 5 or 6 mice per test group. The tumor transplant was by intraperitoneal injection of 0.5 ml of dilute ascitic fluid containing 10⁶ cells of lymphocytic leukemia P388. LL-E33288 antibiotics were tested in the P388 system both as the individual β ₁-Br and γ ₁-Br components and as a complex of all components (Bromo-complex). The test compounds were administered intraperitoneally at a volume of 0.5 ml in 0.2% Klucel in normal saline on days 1, 5 and 9 (relative to tumor inoculation) at the indicated doses. The mice were weighed and the survivors recorded on a regular basis for 30 days. The median survival time and the ratio of survival time for treated (T)/control (C) animals were calculated. The positive control compound was Cisplatin given as an intraperitoneal injection in 0.5 ml of 0.2% Klucel on days 1, 5 and 9 at the indicated doses. The results appear in Table IV.

If $T/C \times 100 (\%)$ is 125 or over, the tested compound is considered to have significant anti-tumor activity.

18 AUG 1999

38

Molasses..... about 0.5%
Calcium carbonate..... about 0.5%
Source of bromine or iodine..... trace amounts
Water.....qs to..... 100%

A 2800 liter portion of the above medium was sterilized and then inoculated with 300 liters of tertiary (seed) inoculum prepared as described in Example 1. Aeration was supplied at the rate of 0.53 liters of sterile air per liter of mash per minute and agitation was supplied by an impeller driven at 110 rpm. The temperature was maintained at about 28°C and the fermentation was terminated after about 97 hours, at which time the mash was harvested.

The fermentation was monitored for production of the LL-E33288 antibiotics by antibacterial activity, biochemical induction assay, TLC and HPLC analyses.

The whole harvest mash was adjusted to pH 6 and then extracted with 1/2 mash volume ethyl acetate. The ethyl acetate extract was concentrated to a syrup which was washed twice with hexane and filtered through diatomaceous earth. The diatomaceous earth cake was thoroughly mixed with ethyl acetate and filtered. The filtrate was concentrated to 3 liters, dried over excess anhydrous sodium sulfate and then precipitated by the addition of hexane giving about 26.7 g of crude LL-E33288 complex.

Example 3

Separation of LL-E33288 α_1 -Br, α_2 -Br, α_3 -Br and α_4 -Br from

LL-E33288 β_1 -Br, β_2 -Br and γ_1 -Br

The approximately 26.7 g of crude LL-E33288 complex (Bromo-complex) from Example 2 was divided evenly into three portions and chromatographed on three separate 2.4 x 110 cm silica gel columns (Silica Woelm®, 32-63 μ m, Woelm Pharma) packed and equilibrated with ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate. The columns were first eluted with the same solvent at a flow rate of 3 ml/minute

18 AUG 1999



56

14.) A compound LL-E33288 δ ₁-I having the following R_f values in the indicated solvent systems on TLC on silica gel sheets:

a) ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, R_f = 0.11; and

b) 3% isopropanol in ethyl acetate saturated with 0.1M aqueous potassium dihydrogen phosphate, R_f = 0.19.

15. Use of a compound selected from the group consisting of LL-E33288 α ₁-Br; LL-E33288 α ₁-I; LL-E33288 α ₂-Br; LL-E33288 α ₂-I; LL-E33288 α ₃-Br; LL-E33288 α ₃-I; LL-E33288 α ₄-Br; LL-E33288 δ ₁-Br; LL-E33288 δ ₁-I; LL-E33288 δ ₂-Br; LL-E33288 δ ₂-I; LL-E33288 γ ₁-Br; LL-E33288 γ ₁-I; and LL-E33288 δ ₁-I for the manufacture of medicaments for treating bacterial infections in warmblooded animals.

16. Use of a compound selected from the group consisting of LL-E33288 α ₁-Br; LL-E33288 α ₁-I; LL-E33288 α ₂-Br; LL-E33288 α ₂-I; LL-E33288 α ₃-Br; LL-E33288 α ₃-I; LL-E33288 α ₄-Br; LL-E33288 δ ₁-Br; LL-E33288 δ ₁-I; LL-E33288 δ ₂-Br; LL-E33288 δ ₂-I; LL-E33288 γ ₁-Br; LL-E33288 γ ₁-I; and LL-E33288 δ ₁-I for the manufacture of medicaments for inhibiting the growth of tumors in a mammal.

17. Use of a compound selected from the group consisting of LL-E33288 α ₁-Br; LL-E33288 α ₁-I; LL-E33288 α ₂-Br; LL-E33288 α ₂-I; LL-E33288 α ₃-Br; LL-E33288 α ₃-I; LL-E33288 α ₄-Br; LL-E33288 δ ₁-Br; LL-E33288 δ ₁-I; LL-E33288 δ ₂-Br; LL-E33288 δ ₂-I; LL-E33288 γ ₁-Br; LL-E33288 γ ₁-I; and LL-E33288 δ ₁-I for the manufacture of medicaments for regressing leukemia in a mammal.

18. A process for producing antibiotics LL-E33288 α ₁-Br; LL-E33288 α ₂-Br; LL-E33288 α ₃-Br; LL-E33288 α ₄-Br; LL-E33288 δ ₁-Br; LL-E33288 δ ₂-Br and LL-E33288 γ ₁-Br which comprises aerobically fermenting the organism Micromonospora echinospora ssp. calichensis NRRL 15839 or mutants thereof including NRRL 15975 in a liquid medium containing assimilable sources of carbon, nitrogen,