

Ribonucleic acid (RNA) interactions

The invention relates to a method for analysing ribonucleic acid (RNA) interactions comprising cross-linking base-paired nucleotides of at least one RNA molecule and/or at least one pair of RNA molecules using a tagged reversible cross-linking agent; a kit for analysing ribonucleic acid (RNA) interactions comprising at least said tagged reversible cross-linking agent; a method of studying a subject using the said method and/or kit; and a drug discovery method using the said method and/or kit.

Background of Invention

The ability of an RNA to base pair with itself and with others is crucial for its function *in vivo*. RNA carries information in both its linear sequence and its secondary and tertiary structure. While significant advances have been made to map RNA secondary structures genome-wide, understanding how different parts of an RNA interact to form higher order structures requires considerable pairwise structural information. RNA's ability to interact with other RNAs, such as miRNA-mRNA and lncRNA-mRNA interactions, plays an important role in post-transcriptional gene regulation. However, the global prevalence and dynamics of RNA interaction networks and their impact on gene regulation is still largely unknown. As such, mapping RNA structure and interactomes in different cellular states is crucial to expanding our understanding of RNA function.

To identify which two RNA regions are interacting with each other, we need spatial connectivity information to link nucleotides that are physically pairing. Numerous RNA cross-linkers, including methylene blue, UV and psoralen, have been used to connect far away interacting regions of RNAs to each other. However, the readout for these strategies has typically been slow and tedious. Alternative strategies for identifying pairwise interactions have utilized sequence mutations followed by structure probing to detect base pairing partners within an RNA. These approaches are higher throughput, but are not amenable to studying whole genomes. Recent strategies such as CLASH, Hi-CLIP and RAP have leveraged on high-throughput sequencing to identify subpopulations of RNA interactions that are associated with a specific RNA binding protein or RNA species. A

recent proximity ligation based approach, RPL, has also been used to identify stems in the transcriptome in a non-selective manner. However, RPL does not utilize crosslinking to identify stable interactions and is mostly limited to mapping intramolecular RNA interactions.

We herein disclose a high-throughput methodology, termed Sequencing of Psoralen crosslinked, Ligated, and Selected Hybrids (SPLASH), that maps pairwise RNA interactions *in-vivo* with high sensitivity and specificity, genome-wide. Applying SPLASH to human and yeast transcriptomes permits the diversity and dynamics of thousands of long-range intra and intermolecular RNA-RNA interactions to be studied. This, for example, permitted analysis that highlighted key structural features of RNA classes, including the modular organization of mRNAs, its impact on translation and decay, and the enrichment of long-range interactions in non-coding RNAs. Additionally, intermolecular mRNA interactions were organized into network clusters and were remodelled during cellular differentiation. Also, it allowed identification of hundreds of known and new snoRNA-rRNA binding sites, expanding the knowledge base of rRNA biogenesis. These results highlight the under-explored complexity of RNA interactomes and paves the way to better understand how RNA organization impacts biology.

Statements of Invention

According to a first aspect of the invention there is provided a method for analysing ribonucleic acid (RNA) interactions comprising:

- a. cross-linking base-paired nucleotides of at least one RNA molecule and/or at least one pair of RNA molecules using a tagged, reversible cross-linking agent to produce at least one cross-linked RNA molecule and/or at least one pair of cross-linked RNA molecules;
- b. fragmenting the said cross-linked RNA molecule and/or pair of cross-linked RNA molecules to produce a plurality of fragments comprising at least one cross-linked RNA fragment;
- c. using said tag to extract said cross-linked RNA fragment(s) from said plurality of fragments;

- d. ligating the said cross-linked RNA fragment(s) to produce cross-linked ligated RNA chimera(s);
- e. reversing the cross-linking of the said agent to the said RNA molecule and/or pair of RNA molecules to produce a ligated RNA chimera molecule(s) and/or RNA chimera pair(s);
- f. preparing a sequence library by sequencing the ligated RNA chimera molecule(s) or pair(s); and
- g. analysing the sequence library to determine RNA interactions.

In a preferred method of the invention said RNA is present in a cell and said crosslinking using said tagged, reversible cross-linking agent involves the use of a cellular uptake agent, such as a detergent. Ideally, the detergent is digitonin and preferably used at a concentration of 0.01% or thereabouts. In this embodiment of the invention, said RNA is extracted from said cell prior to performing the fragmentation step of part b.

Those skilled in the art will appreciate that when working the invention part c may be undertaken before or after part b.

In a preferred method of the invention said cross-linking agent comprises a furocoumarin compound, ideally, psoralen. We have found that psoralen intercalates into base-paired regions independently of whether they are formed by the same RNA strand, or between two different RNA strands, enabling SPLASH to interrogate both intra- and inter-molecular RNA interactions.

Psoralen (also called psoralene) is the parent compound in a family of natural products known as furocoumarins. It is structurally related to coumarin by the addition of a fused furan ring, and may be considered as a derivative of umbelliferone. Practising the invention herein described may involve the use of any one or more of these compounds. Advantageously, these furocoumarins are capable of reversibly and/or selectively cross-linking nucleotides.

In yet a further preferred method of the invention said tag of said cross-linking agent comprises a first member of a binding pair. Ideally, said tag is one member of one of the following binding pairs: biotin/streptavidin, antigen/antibody, protein/protein, polypeptide/protein and polypeptide/polypeptide. Accordingly, using said tag to extract said cross-linked RNA fragment from said plurality of fragments involves the use of the other member of said binding pair which may, optionally, be provided on a support.

More preferably still, the cross-linking of said RNA molecule(s) with said cross-linking agent to produce cross-linked RNA molecule(s) is carried out using ultraviolet irradiation at wavelengths in the range of about 300nm to about 400nm. Similarly, reversing the cross-linking of the cross-linked ligated RNA molecule(s) is carried out using ultraviolet irradiation at a different wavelength i.e. in the range of about 200nm to no more than about 300nm.

Preferably, the method step of preparing a sequence library by sequencing the ligated RNA chimera molecule(s) or pair(s) comprises the use of at least one or more of the following techniques: adaptor ligation, reverse transcription, cDNA circularization or polymerase chain reaction (PCR).

In a preferred method of the invention, the step of fragmenting the cross-linked RNA molecule and/or pair of RNA molecules to produce a plurality of fragments comprises producing fragments having an average size in the range of 100 to 500 base pairs in length. Conventional means or agents for fragmenting RNA are used in the method of the invention, such as physical, chemical or enzymatic means including but not limited to acoustic shearing, sonication, hydrodynamic shearing, DNase or ribonuclease treatment, transposase treatment, and heat digestion with a divalent metal cation.

Ideally, when practising the method of the invention, the concentration of cross-linking agent used is calibrated such that it crosslinks at approximately one in every 150 bases.

Ideally, when analysing the sequence library continuous pairwise interactions or those

spaced apart by less than 50 bases are removed, this enables one to focus the analysis on the long-range intramolecular and intermolecular interactions.

In yet a further preferred method of the invention said RNA molecule and/or at least one member of said pair of RNA molecules is ascribed a "circularization score" defined as the average base pair interaction distance within each molecule, normalized by the length of said RNA molecule or the length of said member of said pair of RNA molecules. More ideally still, when analysing the sequence library said RNA molecule and/or said at least one member of said pair of RNA molecules are classified into groups according to their "circularization score".

Reference herein to circularization score is reference to the propensity of RNA to form long-range pairwise interactions which we have found to be related to translation efficiency. Indeed, we have discovered that transcripts with high circularization scores tend to be translated better than those with low circularization scores, moreover, these scores can change as the corresponding RNA, particularly mRNA, undergoes conformational change. For example, mRNAs that shift from having a high circularization score in ES (stem) cells to a low circularization score in RA (differentiated) cells showed a corresponding decrease in translation efficiency and vice versa (Figure 7A). This shows that conformational changes can serve as an underlying mechanism to control translation efficiency during changes in cellular states. For example, one of the chromatin genes, high mobility group 1, HMGA1, exhibited a notable decrease in circularization score and translation efficiency during RA differentiation, consistent with its key role in maintaining ES cell pluripotency (Figure 7B). Corroboratively, protein and mRNA quantification using western blot and qPCR analysis showed that HMGA1 protein levels decrease after 5 days of differentiation, whereas its mRNA levels do not (Figure 7C, D). Furthermore, translation efficiency, measured by ribosome profiling in mouse ES and differentiated cells, showed a corresponding decrease in HMGA1 translation efficiency upon cellular differentiation (Figure S7H), reinforcing the association between structural rearrangement and translation.

In yet a further preferred method of the invention the cell is mammalian, human, bacterial or yeast.

Most typically, analysing the sequence library to determine RNA interactions comprises processing data derived from the sequence library through one or more computational blocks to determine RNA interactions. Most preferably, the one or more computational blocks is/are selected from the group consisting of: a computational block for filtering reads from adaptor RNAs; a computational block for filtering reads from PCR duplicates; a computational block for merging paired-end reads into single reads; a computational block for filtering reads from split alignments less than a predetermined number of base pairs apart; a computational block for filtering reads from splicing related false positives interactions; a computational block for filtering reads of co-transcribed transcripts relating to intermolecular interactions; a computational block for binning and filtering of data relating to interacting RNA pairs; and indeed any combination of the afore blocks.

Ideally, the computational block for filtering reads from split alignments less than a predetermined number of base pairs apart comprises filtering reads from split alignments less than 50 bases pairs apart.

Typically, the invention can be used so that the RNA interactions determined provide useful information relating to, amongst other things, intermolecular RNA interaction, intramolecular RNA interaction, primary RNA structure, secondary RNA structure, tertiary RNA structure, quaternary RNA structure, gene regulation, gene expression, gene translation efficiency, RNA decay rates, metabolites responsive to RNA elements and ribosome biogenesis.

Most advantageously, the method of the invention is indiscriminate in analysing RNA interactions genome-wide and is not limited to analysing RNA interactions associated with a specific RNA binding protein or RNA species.

In yet a further aspect, the invention concerns a kit for analysing ribonucleic acid (RNA)

interactions comprising:

- a tagged, reversible cross-linking agent for reversibly cross-linking base paired nucleotides of at least one RNA molecule and/or at least one pair of RNA molecules to produce at least one cross-linked RNA molecule and/or at least one pair of cross-linked RNA molecules;

- a fragmentation buffer for fragmenting the said cross-linked RNA molecule and/or said pair of cross-linked RNA molecules to produce a plurality of fragments;

- an RNA ligase for ligating the cross-linked RNA fragment(s) to produce cross-linked ligated RNA chimera(s);

- a binding partner for said tag on said agent; and

- optionally, instructions on how to use the kit.

Preferably, the kit further comprising reagents for sequencing the cross-linked ligated RNA chimera(s) to prepare a sequence library. Ideally, the kit comprises at least one of a RNA ligase, reverse transcription primers and DNA polymerase.

Most preferably, the cross-linking agent comprises a furocoumarin compound, such as psoralen.

Additionally, said tag of said cross-linking agent comprises a first member of a binding pair. Ideally, said tag is one member of one of the following binding pairs: biotin/streptavidin, antigen/antibody, protein/protein, polypeptide/protein and polypeptide/polypeptide. Accordingly, using said tag to extract said cross-linked RNA fragment from said plurality of fragments involves the use of said binding partner, or the other member of said binding pair, which may, optionally, be provided on a support.

More preferably still, the kit further comprises an agent to facilitate cellular uptake of the cross-linking agent into a cell such as a detergent, an example of which is a mild detergent such as digitonin, and used at about 0.01%.

According to a further aspect of the invention there is provided a method of studying a subject, the method comprising:

- a. obtaining a cell sample from a subject;
- b. cross-linking base-paired nucleotides of at least one RNA molecule and/or at least one pair of RNA molecules using a tagged, reversible cross-linking agent to produce at least one cross-linked RNA molecule and/or at least one pair of cross-linked RNA molecules;
- c. fragmenting the said cross-linked RNA molecule and/or pair of cross-linked RNA molecules to produce a plurality of fragments comprising at least one cross-linked RNA fragment;
- d. using said tag to extract said cross-linked RNA fragment(s) from said plurality of fragments;
- e. ligating the said cross-linked RNA fragment(s) to produce cross-linked ligated RNA chimera(s);
- f. reversing the cross-linking of the said agent to the said RNA molecule and/or pair of RNA molecule(s) to produce ligated a RNA chimera molecule(s) and/or RNA chimera pair(s);
- g. preparing a sequence library by sequencing the ligated RNA chimera molecule(s) or pair(s);
- h. analysing the sequence library to determine RNA interactions in the cell sample; and
- i. comparing the determined RNA interactions with a set of pre-existing data to attribute a clinical picture to the subject.

In this preferred method of the invention, the method of studying a subject comprises at least one of: diagnosing the subject of a clinical condition, predicting the risk of the subject having a clinical condition, screening the subject for suitability for a particular treatment or determining the efficacy of a drug candidate on the subject.

According to a yet further aspect of the invention there is provided a drug discovery method, the method comprising:

- a. exposing RNA to a drug;
- b. cross-linking base-paired nucleotides of at least one RNA molecule and/or at least one pair of RNA molecules using a tagged, reversible cross-linking agent to produce at least one cross-linked RNA molecule and/or at least one pair of cross-linked RNA molecules;
- c. fragmenting the said cross-linked RNA molecule and/or pair of cross-linked RNA molecules to produce a plurality of fragments comprising at least one cross-linked RNA fragment;
- d. using said tag to extract said cross-linked RNA fragment(s) from said plurality of fragments;
- e. ligating the said cross-linked RNA fragment(s) to produce cross-linked ligated RNA chimera(s);
- f. reversing the cross-linking of the said agent to the said RNA molecule and/or pair of RNA molecules to produce a ligated a RNA chimera molecule(s) and/or RNA chimera pair(s);
- g. preparing a sequence library by sequencing the ligated RNA chimera molecule(s) or pair(s);
- h. analysing the sequence library to determine RNA interactions; and
- i. attributing an efficacy score to the drug based on the determined RNA interactions.

Throughout the description and claims of this specification, the words “comprise” and “contain” and variations of the words, for example “comprising” and “comprises”, mean “including but not limited to” and do not exclude other moieties, additives, components, integers or steps. Throughout the description and claims of this specification, the singular encompasses the plural unless the context otherwise requires. In particular, where the indefinite article is used, the specification is to be understood as contemplating plurality as well as singularity, unless the context requires otherwise.

All references, including any patent or patent application, cited in this specification are hereby incorporated by reference. No admission is made that any reference constitutes prior art. Further, no admission is made that any of the prior art constitutes part of the common general knowledge in the art.

Preferred features of each aspect of the invention may be as described in connection with any of the other aspects.

Other features of the present invention will become apparent from the following examples. Generally speaking, the invention extends to any novel one, or any novel combination, of the features disclosed in this specification (including the accompanying claims and drawings). Thus, features, integers, characteristics, compounds or chemical moieties described in conjunction with a particular aspect, embodiment or example of the invention are to be understood to be applicable to any other aspect, embodiment or example described herein, unless incompatible therewith.

Moreover, unless stated otherwise, any feature disclosed herein may be replaced by an alternative feature serving the same or a similar purpose.

The invention will now be described, by way of example only, with reference to the following figures and tables wherein:-

Figure 1. SPLASH strategy identifies RNA interactions accurately. [A] Schematic of the SPLASH strategy. RNA-RNA interactions are crosslinked *in vivo* using biotinylated psoralen (biopsoralen) and then fragmented. Interacting regions that contain bio-psoralen are enriched by binding to streptavidin beads and ligated together. Chimeric RNAs are then cloned into a cDNA library for deep sequencing. [B] Visualization of chimeric interactions on the secondary structure of the 28S rRNA. The dark grey, light grey and grey lines represent interactions <30Å apart, 30- 60Å apart, and >60Å apart respectively, based on the 80S ribosome crystal structure. [C] Bar chart showing positive predictive value (PPV) and sensitivity in total RNA replicates, based on the 80S rRNA crystal structure, at different cut-offs. The cut-off "All" indicates interactions that exist in at least

one out of the four biological replicates, while “2/4”, “3/4” and “4/4” indicates interactions that are present in at least 2, 3 or 4 replicates. The notation “>8” indicates that the interaction needs to be supported by at least 8 chimeric reads across the libraries. See also Figure 8, 9, Table 1;

Figure 2. Distribution and function of the human RNA interactome. [A, B], Pie charts showing the distribution of intramolecular [A] and intermolecular [B] interactions belonging to different classes of transcripts in four biological replicates of total RNA (Left) and in all polyA(+) enriched RNA samples (Right). [C] Schematic of the circularization score. The circularization score is calculated as the average span of intramolecular interactions normalized by mRNA length. mRNAs with higher circularization score participate in more long-range interactions. [D] Boxplot of circularization scores for rRNAs, lncRNAs and mRNAs. P-values are calculated using Wilcoxon rank sum test. See also Figure 10, 11, Table 2;

Figure 3. Intramolecular interaction patterns and their association with gene regulation. [A] Two-dimensional heat map showing enrichment of intramolecular mRNA interactions based on the location of chimera ends. We aligned transcripts according to their translation start and stop sites and plotted interactions from the last 200 bases of the 5' UTR, the first and last 400 bases of the coding region, and the first 400 bases of the 3' UTR. Enrichment was calculated as $-\log_{10}(p\text{-value})$ based on random sampling across the transcript with 100bp windows. The black dotted lines demarcate boundaries between the 5' UTR, CDS and 3' UTR. [B] Metagene analysis of the frequency of intramolecular interactions along human mRNAs, by aligning mRNAs along their translation start and stop. We plotted interactions that are present in the last 200 bases of the 5' UTR, first and last 400 bases of the coding region and the first 400 bases of the 3' UTR. [C] Boxplot of translation efficiency (Y-axis) in mRNAs with the highest and lowest 20% circularization scores. [D] Boxplot of translation efficiency in mRNAs that have interactions only in the 5' UTRs, versus mRNAs with interactions all over the transcript. [E] Density plot showing the distribution of the left (grey) and right (black) end of a pairwise interaction for top 5% of mRNAs that are translated efficiently (Left plot) and poorly (Right plot) based on ribosome

profiling data. [F] Boxplot of decay rate (Y-axis) in genes that have pairwise interactions confined to the 5' end (Left plot), versus all over the transcript. Pairwise interactions at the 3' end tend to block decay. See also Figure 12, Table 3;

Figure 4. SPLASH identifies known and new human snoRNAs-rRNA interactions.

[A], The black line graph indicates the region that U42B (Top) or U80 (bottom) interacts with 18S or 28S rRNA, respectively, in SPLASH. Light grey bars are the known interaction region for U42B and U80 respectively in the literature. Y-axis indicates the number of chimeric reads that mapped to rRNA. [B] Validation of novel human snoRNA-rRNA interactions. Left: SPLASH data indicates that SNORA32 (Top) interacts with the 5S rRNA, and SNORD83a (Bottom) interacts with the 18S rRNA. Y-axis indicates the number of chimeric reads supporting the interaction. Right: Independent pulldowns of 5S, 18S and 28S rRNA and qPCR analysis of SNORA32 (Top) and SNORD83a (Bottom) in each pulldown confirms the SPLASH data. Y-axis indicates the relative enrichment. [C] SPLASH reads for human U13-rRNA interactions are plotted along the 5' external transcribed spacer, 18S rRNA, and 28S rRNA. The light grey bar in the U13 plot indicates the known position of U13 binding. The grey line indicates the predicted sites for U13-rRNA interaction using the program PLEXY. U13 target sites that are supported by both SPLASH and PLEXY are starred. The Y-axis for PLEXY is in kcal/mol. [D] Model of RNA base pairing between U13 and 28S rRNA. The starred site is a newly identified U13-28S interaction that is supported by a PLEXY prediction. See also Figure 13, Table 4;

Figure 5. SPLASH identifies known and new yeast snoRNAs-rRNAs interactions. [A]

Line graph showing the locations of snR61 target sites on the 25S rRNA that are detected by SPLASH. The position of the known snR61 binding site is marked as a grey bar. The star indicates that the target site that is both identified by SPLASH and predicted by PLEXY. [B] A model showing predicted interactions between snR61 and 25S rRNA. 25S rRNA is shown in blue while snR61 is shown in black. [C] Line graphs showing SPLASH read counts for snR4 binding to 18S rRNA (Top) and 25S rRNA (Bottom). The starred sites indicate sites that are identified in SPLASH data, as well as previously in CLASH data. SPLASH identifies a new snR4 target site in the 18S rRNA, in addition to validating

previously suggested sites [D] snR61, snR4 and snR30 sites are mapped onto the contour map of the yeast 25S rRNA. [E] Line graphs showing SPLASH reads for snR30-18S rRNA interactions in wildtype and Prp43 yeast mutant. [F] Heatmap of snoRNA target sites that are stabilized (left) or lost (right) in the Prp43 mutant as compared to wildtype yeast. The stabilized sites suggest that these snoRNAs might be dependent on Prp43 for release from rRNA. Newly identified target sites that require Prp43 for release are highlighted in red. Stars indicate sites where the snoRNAs have been previously found to bind to Prp43. See also Figure 13, Table 5;

Figure 6. Function and regulation of mRNA interaction modules. [A, B] Barcharts showing enrichment of Thymosin Beta 4, X-Linked (TMSB4X) [A] and Actin (ACTB) [B] interacting genes by qPCR analysis. The following names stands for Eukaryotic Translation Elongation Factor 1 Alpha 1 (EEF1A1), Ribosomal Protein S27 (RPS27), Beta-2-Microglobulin (B2M), Eukaryotic Translation Initiation Factor 5A (EIF5A) and Ribosomal Protein L35 (RPL35). Y-axis indicates $\log_{10}$ enrichment with respect to oligo pulldown against GFP. Error bars depict standard-deviation based on 3 biological replicates. [C] Two-dimensional heatmap showing enrichment of intermolecular interactions based on the location of chimera ends across mRNA pairs. We aligned transcripts according to their translation start and stop sites and plotted interactions from the last 200 bases of the 5' UTR, first and last 400 bases of the coding region, and first 400 bases of the 3' UTR. Enrichment was calculated as $-\log_{10}(p\text{-value})$ based on random sampling across the transcript with 100bp windows. Black dotted lines demarcate the boundaries between 5' UTR, CDS and 3' UTR. [D] Network analysis of lymphoblastoid cells identified a major mRNA-mRNA interaction connected component. [E] Hierarchical clustering based on the density of mRNA-mRNA interactions separates the major component into 9 modules. GO term analysis of each module showed that the modules are enriched for mRNAs with specific functions and subcellular localization patterns. [F] Bar chart of the number of interaction pairs in observed interactions that are both cytoplasmic localized, or with one in the cytoplasm and the other in the nucleus, versus shuffled interaction pairs. Observed interactions between 2 mRNAs are significantly enriched when both RNAs are in the same

cellular compartment, and depleted when they are in different compartments. See also Figure 14, Table 6;

Figure 7. Remodeling of the RNA interactome during human ES cell differentiation.

[A] Boxplot showing changes in translation efficiency in mouse ES and RA cells for conserved human mRNAs with high circularization score in human ES cells and low circularization score in RA cells (Left), and vice versa (Right). mRNAs with a decrease in circularization score typically show a corresponding decrease in translation efficiency. [B] Arc plots of intramolecular interactions in the gene HMGA1, showing a decrease in circularization score after 5 days of RA differentiation. [C] Analysis of HMGA1 and Oct4 mRNA levels by qPCR in human ES cells and ES cells differentiated by retinoic acid (RA) for 5 days. [D] Analysis of HMGA1, Oct4 and GAPDH protein levels by western blotting in human ES and RA differentiated cells. [E, F] Network analysis of mRNA-mRNA intermolecular interactions in ES [E] and RA [F] cells showed that mRNAs are more interconnected in ES cells than in RA cells. [G, H] Hierarchical clustering based on density of mRNA-mRNA interactions identified specific modules in the major connected component of the interaction network. Representative enriched GO terms are shown as labels for each module. ES interaction modules [G] were observed to be more interconnected than RA interaction modules [H]. See also Figure 14, Tables 6 and 7.

Figure 8. Biotinylated psoralen can enter and crosslink RNAs *in vivo*, related to Figure 1.

[A] Structure of biotinylated psoralen. [B] Immunofluorescence images of HeLa cells treated with bio-psoralen and irradiated at 365nm for UV crosslinking. Psoralen is present in both the nucleus and cytoplasm of the cells. 5 min treatment of 0.01% digitonin at 37°C greatly increases the entry of biotinylated-psoralen into the cells. [C] Footprinting analysis on lymphoblastoid cells treated with (lanes 2, 4) and without 200uM psoralen (Lanes 1, 3), in the absence (lanes 1, 2) and presence (lanes 3, 4) of 0.01% digitonin for 5 min. The black arrows indicate positions of reverse transcriptase stoppage due to psoralen crosslinking. Digitonin treatment does not change the pattern of psoralen crosslinking along 18S and 28S rRNA. [D] Titration of the amount of time for UV crosslinking of RNA interactions using psoralen. Dot blot using 2 µg (top) and 0.2 µg

(bottom) of total RNA after crosslinking for 10, 20 and 30 min at UV 365nm. The condition that we chose, 20 min, is boxed in grey dashed lines. [E] Dot blot showing the amount of biotinylated psoralen (bio-psoralen) incorporation into RNA as we increase the concentration of bio-psoralen added to the cells (Top). The biotinylated 20mer (Bottom) serves as a positive control for us to estimate the amount of psoralen incorporation into RNA. [F, G] Dotblot showing that psoralen can enter into *S. cerevisiae* and *E. coli* cells, although a higher concentration of psoralen is needed for a similar level of incorporation as in HeLa cells. [H] Northern blot analysis using probes complementary against U14 (left) and 35S precursor rRNA (right). U14 shows supershift in the presence of bio-psoralen, in Dbp4 mutant cells, confirming that bio-psoralen detects known interactions in the literature.

Figure 9. SPLASH experimental pipeline, related to Figure 1. [A] Quantification of the amount of reverse crosslinking upon irradiating the RNA with UV 254 nm over a time course. The dot blot indicates the amount of RNA that remains crosslinked upon irradiating with UV254 nm. The amount of reverse crosslinking on the Y-axis of the graph is 1- (fraction crosslinked). [B] RNA footprinting analysis of the yeast EFB1 mRNA that has been irradiated with UV 254nm in vitro for 0, 5, 10, 15, 30 min. The bands indicate the stoppage sites by reverse transcription. RNA damage occurs as early as 5 mins after start of irradiation (Left). The percentage of damage is quantified as 100 (the percentage of full length transcripts detected by reverse transcription). The arrow indicates the condition that we used for library preparation (Right). [C] Metal ion hydrolysis retains the biotinylated psoralen on the RNA after fragmentation. Bar graph quantitating the amount of biotinylated psoralen (identified by dot blot) that remains on RNA before fragmentation, after fragmentation using alkaline hydrolysis at pH 9.2, after fragmentation using Mg²⁺ based metal ion hydrolysis. [D] qPCR quantification of the amount of non-specific binding on non-crosslinked EFB1 mRNA that remains bound to the beads after washing using the optimized wash protocol. We observed a 10⁵ fold decrease in non-specific binding in elute RNA compared to its original amount in input RNA after washing. [E] qPCR quantification of the amount of specific binding on bio-psoralen crosslinked TrxB2 mRNA that remains bound to the beads after washing using the optimized wash protocol. We retained about

60% of the mRNA in the elute, as compared to input, after binding and washing. [F] After ligating the two chimeras together, we devised an efficient of the chimeras, and then performed reverse transcription. We then circularized the cDNA and performed PCR amplification to obtain the final cDNA library. [G] Pipeline for analysis of SPLASH libraries, using sequenced 2×150bp paired-end Illumina reads as input.

Figure 10. SPLASH libraries are sensitive and accurate, related to Figure 2. [A] Histogram showing the distribution of the span of chimeras that were found from SPLASH analysis in human lymphoblastoid cells. The median of the distribution is at 300 bases. [B] Barchart showing the sensitivity and specificity of a library without psoralen crosslinking, a psoralen crosslinked library and a biotinylated psoralen crosslinked library (SPLASH) benchmarked against known base pairs on the 28S rRNA. [C] No. of chimeric reads mapped to the 28S rRNA from libraries made with 1X ligase, 0.1X ligase and no ligase SPLASH libraries. Few chimeric reads are identified in the no ligase sample compared to the ligase samples. [D] Correlation analysis between the number of sequencing reads for four lymphoblastoid SPLASH libraries versus solvent accessibility at each base pair, evaluated using the FreeSASA program. Psoralen-biotin crosslinking to 28S rRNA does not show any dependence on solvent accessible surface area. [E] Receiver operating characteristic (ROC) curves for SPLASH data on 28S rRNA (using known base-pairing information as true positives) compared to RNA proximity ligation (RPL) with and without smoothing. [F] Number of known snoRNA-rRNA interactions detected in RPL and SPLASH libraries sequenced to similar depths, based on the human snoRNA database.

Figure 11. Genomic analysis of SPLASH libraries, related to Figure 2. [A] Correlation analysis between coverage of intramolecular (Left) and intermolecular (Right) chimeric interactions in 2 biological replicates of human ES, RA and lymphoblastoid cells. Intramolecular interaction correlations were calculated per interaction window while intermolecular interaction correlations were calculated for each gene pair. Read coverage was normalized by the total number of chimeric reads identified in each library. [B] Pie-charts showing the distribution of intramolecular (top) and intermolecular (bottom) interactions that belong to different classes of transcripts in two biological replicates of

total RNA (Left) and polyA(+) enriched RNA (Right) in yeast. [C] Histogram showing the distribution of the number of RNA partners that an RNA was found to interact with (in lymphoblastoid cells). The median number of interactions was 1 per mRNA.

Figure 12. Analysis and validation of SPLASH intramolecular interactions, related to Figure 3. [A] Box plot of the interaction energy of intramolecular SPLASH chimeras in mRNAs (Left) versus randomly shuffled chimeras (Right). Y-axis shows the RNA hybridization energy (kcal/mol). True chimeras show a lower hybridization energy (calculated by RNA duplex) indicating that they form more stable base pairs. [B] Intramolecular interactions detected in yeast snR86 gene by SPLASH. The interactions suggest the formation of a long hairpin, consistent with the predicted secondary structure of snR86. [C] Top, Intramolecular interactions detected in yeast LSR1 gene by SPLASH. Both the 100-200:1000-1100 and the 400-500:800-900 interactions were consistent with previous crosslinking experiments. Bottom, secondary structure of LSR1 and where the 100-200:1000-1100 interaction is potentially occurring. [D] Top, Intramolecular interactions detected in yeast SCR1 gene by SPLASH. The 0100:400-500 interaction is consistent with previous reports and with the secondary structure of SCR1. [E] Validation of a long range intramolecular interaction between bases 1-100 and 1400-1500 of the yeast mRNA YBR118W. Psoralen and non-psoralen crosslinked yeast total RNA was fragmented and size selected to be between 100-300 bases, before pulldown of bases 1-100 using biotinylated antisense DNA probes. Y-axis indicates log(enrichment) of the fragment 1400-1500 that is pulled down together with bases 1-100 in the presence or absence of psoralen, by qPCR analysis. [F] Average fraction of U among all four nucleotides in human mRNAs, plotted for the last 200 bases of 5' UTR, first and last 400 bases of coding region and first 400 bases of 3' UTR. [G] K-means clustering of the locations of intramolecular interactions (from lymphoblastoid cells), into 5 clusters, show the different patterns of interactions that can occur within an RNA. Top, schematic of the positions of the chimeras along an mRNA. Bottom, heatmap showing the locations of the left (top) and right (bottom) ends of the interaction. The darker the shade [H] Boxplot showing the translation efficiency of the mRNAs in each cluster. mRNAs with more end-

to-end interactions (Groups 4, 5) are translated more efficiently, while efficiently. The p-value was calculated using the Wilcoxon rank sum test.

Figure 13. SPLASH identifies snoRNA-rRNA interactions, related to Figure 4,5. [A] SPLASH identifies the known U14-18S rRNA interaction (top), at the correct region along the 18S rRNA (bottom). Top: U14-rRNA chimeric reads mapped to 5S, 18S and 28S rRNA. Bottom: The black line indicates the number of U14-rRNA chimeric reads mapped along the 18S rRNA, while the light grey bar indicates the known position of U14-18S rRNA interaction. [B] Top and middle: Controls for snoRNA-rRNA pulldowns. Biotinylated antisense oligos against 5S, 18S and 28S rRNAs were used to pull down rRNAs. Y-axis indicates the log(enrichment) of the respective rRNAs normalized to pulldown by antisense probes to GFP. Actin is not selectively enriched in any of the three rRNAs. Bottom, validation of novel SNORA51/ACA51-28S rRNA interaction. Left: ACA51 interacts with 28S rRNA in SPLASH data. Right: ACA51-28S rRNA interaction is experimentally validated by pulldown of 5S, 18S and 28S rRNA. ACA51 binds to 28S with the highest affinity. [C] Model of snR4 interaction with 25S rRNA for an interaction site that is identified by SPLASH and predicted by PLEXY. [D] Prp43 mutant T-123A and WT yeast grown in the presence of galactose and glucose. Prp43 mutant yeast is defective when grown in galactose showing the successful deletion of Prp43.

Figure 14. Properties of intermolecular interactions detected by SPLASH, related to Figure 6,7. [A] Barcharts showing log enrichment of EEF1A1 interacting genes by oligo pulldown of EEF1A1 and qPCR of its interacting genes in human ES cells. Oligo pulldown against GFP was used as negative control. Error bars depict standard-deviation based on 2 biological replicates. The following names stands for Eukaryotic Translation Elongation Factor 1 Alpha 1 (EEF1A1), Ribosomal Protein S27 (RPS27), Thymosin Beta 4, X-Linked (TMSB4X), Ribosomal Protein, Large, P0 (RPLP0), Glyceraldehyde 3-phosphate dehydrogenase (GAPDH), and Actin (ACTB). [B] Boxplot of the interaction energies of intermolecular SPLASH chimeras in mRNAs (Left) versus randomly shuffled chimeras in mRNAs (Right). Y-axis indicates the RNA hybridization energy (kcal/mol). True chimeras show a lower hybridization energy (computed by RNAduplex) indicating that they form more stable base pairs. [C] Metagene analysis of the frequency of intermolecular interactions along lymphoblastoid mRNAs, by aligning mRNAs along their translation start and stop. We plotted interactions that are present in the last 200 bases of 5' UTR, first and

last 400 bases of coding region and first 400 bases of 3' UTR. [D, E] Barcharts showing the number of observed mRNA-mRNA pairs that are correlated in translation efficiency [D] or decay [E] versus random shuffling in each module. [F, G] Two-dimensional heatmap showing enrichment of interactions between one end of a chimera with the other end for all ES [F] and RA [G] mRNAs. We aligned transcripts according to their translation start and stop sites and plotted interactions from the last 200 bases of the 5' UTR, first and last 400 bases of the coding region, and first 400 bases of the 3' UTR. Enrichment was calculated as based on random sampling across the transcript with 100bp windows. Black dotted lines demarcate the boundaries between 5' UTR, CDS and 3' UTR. Globally, the 2D heatmaps resemble the heatmap for lymphoblastoid cells. [H] Bar charts showing translation efficiency of HMGA1, as measured by ribosome profiling in mouse ES and differentiated cells. HMGA1 translation efficiency decreases during differentiation.

Table 1. Evaluation of different protocols for SPLASH, related to figure 1;

Table 2. Information of sequenced SPLASH libraries, related to figure 1;

Table 3. List of common human-human and human yeast interactions, related to Figure 2;

Table 4. List of lymphoblastoid cells snoRNA target sites, related to Figure 4;

Table 5. List of yeast snoRNA target sites, related to Figure 5;

Table 6. GO analysis of network interactions in lymphoblastoid, ES and RA cells, related to Figure 6,7;

Table 7. Probes and qPCR primers used in validation, related to Figure 6.

METHODS & MATERIALS

Cell culture. HeLa cells were grown in Dulbecco's Modified Eagle Medium (DMEM) supplemented with 10% Fetal bovine serum (FBS) and 1% Penicillin Streptomycin (PS). Human lymphoblastoid cells, GM12892, were grown in Roswell Park Memorial Institute (RPMI) supplemented with 20% FBS, 1% PS and 2mM L-Glutamine. hESC line H1 (WA-01, passage 30) was cultured in mTeSR1 (Stem cell technologies) media, on matrigel (BD) coated dishes. For Retinoic Acid (RA) treatment, the cells were seeded at 1:6 ratio and treated with 10uM of RA after 16-24 hrs, and harvested after 5 days of treatment.

Crosslinking and extraction of human, yeast and *E. coli* RNAs. HeLa and GM12892 cells were washed with PBS and treated with 200 μ M of EZ-Link™ Psoralen-PEG3-Biotin (Thermo Fisher Scientific) and 0.01% w/v Digitonin (Sigma) at 37°C for 5 min. *Saccharomyces cerevisiae* (S288C, or W303a) or *Escherichia coli* (*E. coli* K12) were grown to exponential phase (OD=0.6), pelleted and washed in TE buffer and incubated with 2 mM of EZ-Link Psoralen-PEG3-Biotin at 37°C for 10 min in TE. The cells were then spread onto a 10 cm plate and irradiated using 365 nm UV for 20 min on ice. Human and *E. coli* RNAs were isolated by using TRIzol reagent (Invitrogen) while Yeast RNAs were isolated using hot acid phenol extraction.

Fragmentation and enrichment of crosslinked RNA. 20 μ g of RNA were fragmented with RNA fragmentation buffer (9 mM MgCl₂, 225 mM KCl and 150 mM Tris HCl (pH 8.3)) at 95°C for 5 min and size fractionated on a 6% TBE 8M Urea gel. Bases corresponding to 90-110 nt were excised and eluted overnight at 4°C. 1.5 μ g of fragmented RNA was incubated with 100 μ L of Dynabeads® MyOne™ Streptavidin C1 beads (Life Technology), dissolved in 2 mL of fresh Hybridization Buffer (750 mM NaCl, 1% SDS, 50 mM Tris-Cl pH 7.0, 1 mM EDTA, 15% formamide) and 1 ml of supplemented lysis buffer (50 mM Tris-Cl pH 7.0, 10 mM EDTA, 1% SDS) supplemented with Suprase-in (1:200), at 37°C for 30 min. The beads were washed with 1 mL of wash buffer (2x NaCl and Sodium citrate (SSC), 0.5% SDS) at 37°C for 5 min with gentle agitation for five times.

Proximity ligation and reverse crosslinking. Enriched crosslinked samples were washed in cold T4 PNK buffer and treated with 0.5 unit of T4 PNK enzyme (NEB) at 37°C for 4 hours in a 80 µL reaction. We then added fresh 1 mM ATP and 0.5 unit of T4 PNK in a 100 µL reaction, and incubated the reaction for 1 hr at 37 °C. The chimeras were ligated using 2.5 units/µL of T4 RNA ligase I overnight at 16°C, in a 160 µL reaction, and eluted from the beads by incubating at 95°C for 10 min in 100 µL of PK buffer (100mM NaCl, 10 mM TrisCl pH 7.0, 1mM EDTA, 0.5% SDS). Eluted RNA was extracted using TRIzol reagent, and cleaned up using RNeasy Cleanup Kit (Qiagen). We reverse crosslinked the RNA by irradiating at UV 254 nm for 5 min on ice.

3' Adapter ligation. Reverse crosslinked samples were resuspended in 6 µM of 3' adaptors and heat denatured at 80°C for 90 seconds before snap cooling on ice. The 3' adaptors were ligated using T4 RNA ligase 2 KQ at 25°C for 2.5 hours and size fractionated using a 6% TBE 8M Urea gel. RNA corresponding to 110-130 bases were excised and eluted overnight at 4°C.

Reverse transcription (RT). Eluted samples were resuspended in 208 nM of RT primers, heat denatured at 80°C for 2 min and crashed on ice for 1 min. Denatured samples were then incubated at 50°C for 30 min using SuperScript III (Invitrogen) for RT. cDNAs were recovered by degrading RNAs in 100 mM of NaOH, at 98°C for 20 min, and size fractionating on a 6% TBE 8M Urea gel. cDNA of bases 200-220 were excised and eluted overnight at room temperature.

Circularization of cDNA product and PCR. The eluted cDNA samples were recovered by ethanol precipitation, circularized using Circligase II (Epicentre) and purified using DNA Clean & Concentrator™-5 (Zymo). We performed 9-12 cycles of PCR amplification using primers from Primers Set 1 (New England Biolabs) and Q5 DNA polymerase (New England Biolabs). PCR products were run on a 3% GTG Nusieve Agarose (Lonza) and bases 200-300 were gel extracted and purified using DNA gel extraction kit (Qiagen). The

libraries were quantified using Qubit DNA HS Assay (Invitrogen), and sequenced on the Nextseq 500 machine (Illumina).

Human and Yeast Transcriptomes. Human and Yeast sequences were downloaded from the UCSC Genome Browser. Additional sequences belonging to human snoRNAs, snRNAs (extracted from NCBI), tRNAs (extracted from the UCSC Table Browser) and rRNAs were added to the human transcriptome list. Yeast UTR sequences, and noncoding gene sequences including rRNAs, tRNAs, snRNAs, snoRNAs and ncRNAs (Saccharomyces Genome Database) were also added to our transcriptome list.

Processing and detection of chimeric reads. Reads were adapter removed and merged using SeqPrep (version 1.0-7; <https://github.com/jstjohn/SeqPrep>). Merged reads were mapped to the transcriptome (see above) with BWA MEM (Li and Durbin, 2010) (version 0.7.12). Only split alignments that are i) > 50bp apart in transcriptome sequence, ii) not reverse complements of each other, and iii) with mapping quality ≥ 20 are kept for downstream analysis. We further filtered the mapped transcriptome reads by ensuring that i) it could be uniquely mapped back to the human genome (hg19) using the program STAR, ii) does not span annotated splicing junctions, iii) present in at least two out of the four replicates, iii) had a minimum coverage of 2 and iv) if the average coverage in all replicates was at least 2. The final coverage of an interaction site is the average of normalized coverage in all replicates.

Availability. For source code and additional materials see <http://csb5.github.io/splash/>.

SRA accession number. SRP073550

SRA Bioproject ID. PRJNA318958

Cell culture. Human HeLa cells were grown in Dulbecco's Modified Eagle Medium (DMEM) supplemented with 10% Fetal bovine serum (FBS) and 1% Penicillin Streptomycin (PS) and psoralen crosslinked at 80% confluency. Human lymphoblastoid cells, GM12892, were grown in Roswell Park Memorial Institute (RPMI) supplemented with 20% FBS, 1% PS and 2mM L-Glutamine to a concentration of 6×10^5 cells/mL. 20 mL

were used for psoralen crosslinking. The hESC line H1 (WA-01, passage 30) was cultured in mTeSR1 (Stem cell technologies) media, on matrigel (BD) coated dishes. The media was refreshed daily. The hESCs were routinely subcultured with 1 mg/ml Dispase (Stem cell technologies) every 5-7 days. For Retinoic Acid (RA) treatment, the cells were seeded at 1:6 ratio. After 16-24 hrs, the cells were treated with 10uM of RA. The media was refreshed daily and cells were harvested after 5 days of treatment.

SnoRNA immunoprecipitation. SnoRNA enriched samples were obtained by performing immunoprecipitation in IPP150 buffer (6 mM HEPES (pH8.0), 150 mM NaCl, 5 mM MgCl₂, 0.1% Nonidet P-40) with protein A-agarose (Thermo Fischer Scientific) bound anti-TMG (R1131) antibodies. To precipitate TMG cap snoRNAs, total RNA was incubated with protein A-Agarose bound anti-TMG antibodies agarose beads on a rotating wheel for 3 hours at 4°C. The bead bound RNA was digested with proteinase K solution (50 mM Tris-HCl (pH7.5), 5 mM EDTA and proteinase K (2 g/ l) for 30 min at 42°C. The RNA was extracted with phenol-chloroform and concentrated using ethanol precipitation.

3' adapter primer sequence. 5'-CTGTAGGCACCATCAAT-3' (IDT) (SEQ ID NO:1)

Reverse transcription primer sequence. 3' adapter ligated samples were recovered by ethanol precipitation and resuspended in 208 nM of RT primers,

5'AGATCGGAAGAGCGTCGTGTAGGGAAAGAGTGTAGATCTCGGTGGTCGC/iSp18/C
ACTC A/iSp18/TTCAGACGTGTGCTCTTCCGATCTATTGATGGTGCCTACAG-3' (IDT)
(SEQ ID NO:2).

Preparation of control libraries. DMSO and psoralen crosslinked libraries were prepared the same way as the normal libraries, except for the skipping of the enrichment steps by binding to streptavidin beads. As we estimated that around 20 ng of bio-psoralen crosslinked and fragmented RNA is typically bound to streptavidin beads, we used the same amount (20 ng of fragmented, size selected samples) in the subsequent ligation and library preparation steps, using the same conditions as in SPLASH library generation.

Northern blot analysis of U14-18S rRNA interaction. Bio-psoralen crosslinked total RNA was extracted from wild-type, Dbp4p, or Dbp8p metabolic depleted yeast cells, and denatured at 95°C for 5 minutes before separated by the gel electrophoresis (native, 1.2% agarose gel). RNA species that are crosslink by bio-psoralen will co-migrate in the gel. The double stars indicated a supershifted U14-35S rRNA complex, which is accumulated in the Dbp4 mutant. The non-bio-psoralen crosslinked wild-type RNA sample is used as a background control.

Dot blot analysis to detect the presence of biotinylated psoralen on RNA.

Presence of biotinylated psoralen in the cross-linked RNA samples was detected with Chemiluminescent Nucleic Acid Detection Module (Thermo Fisher Scientific) following manufacturer's instructions. 1ug of RNA was dotted on to a Biodyne™ B Nylon Membrane (Thermo Fisher Scientific) and cross-linked to the membrane by baking at 80C for 15 minutes. The membrane was visualized using ChemiDoc™ MP System (BioRad) and quantified using the software Image J.

Calculation of bio-psoralen incorporation into cellular RNAs. Each positive control 20mer oligo contains one biotin molecule. From the number of moles of 20mer oligo and our crosslinked RNAs that are spotted, and intensity of the 20mer oligo by dot blot, we can estimate the amount of incorporation of psoralen in our RNAs.

Western Blotting and qPCR analysis of HMGA1, OCT4 and GAPDH. Human H1 ES cells and ES cells that are differentiated using retinoic acid (RA) for 5 days were lysed using RIPA buffer (150 mM sodium chloride, 1.0% Triton X-100, 0.5% sodium deoxycholate, 0.1% SDS, 50 mM Tris, pH 8.0) supplemented with 1:200 of Protease Inhibitor Cocktail Set III (Merck). Cells were incubated at 4 C for 20 minutes with gentle agitation. The lysate was then clarified by passing through a 25G BD Precision Glide Needle (Becton, Dickinson and Company) for a total of 6 times and centrifuged at 12000 rpm for 30 minutes at 4 C to pellet the insoluble fraction. The supernatant was collected and protein levels were normalized for each sample with Bio-Rad Protein Assay Dye Reagent Concentrate (Bio-Rad). Normalized samples were then size fractionated on a 12% sodium dodecyl sulfate polyacrylamide gel electrophoresis

(SDS-PAGE) gel and transferred onto a Nitrocellulose Membrane (Bio-Rad). Membranes were blocked in 5% Blotting-Grade Blocker in PBST (137 mM NaCl, 2.7 mM KCl, 4.3 mM Na₂HPO₄, 1.47 mM KH₂PO₄, 0.1% Tween 20) and incubated with primary antibodies overnight at 4 C. The membranes were washed and incubated with secondary antibodies conjugated with HRP for 1 hour at room temperature. After washing, the membranes were incubated with Clarity™ Western ECL Substrate (Bio-Rad) and visualized with ChemiDoc™ MP System. The bands were quantified using the software Image J. The following antibodies were used with the dilutions stated: Anti-HMGA1 (cell signaling, #7777) 1:50000, Anti-Oct4 antibody (Abcam, ab19857) 1:10000, Anti-GAPDH Antibody (Merck, MAB374) 1:50000, anti-mouse IgG-HRP antibody (Santa Cruz, sc-2031) 1:5000, and anti-rabbit IgG-HRP antibody (Santa Cruz, sc-2313) 1:5000.

Total RNA was extracted from human ES and RA differentiated cells using the Trizol reagent (Thermo Fisher Scientific) and qPCR analysis was performed using the Brilliant II SYBR Green qRT-PCR 1-Step Master Mix kit, according to manufacturer's instructions. qPCR analysis are normalized to actin control.

Immunofluorescence imaging. HeLa cells were cultured on cover slips and treated with bio-psoralen. *In vivo* treated HeLa cells were rinsed with PBS once and fixed with 4% paraformaldehyde (Sigma) in PBS at room temperature for 30 min. Fixed cells were washed twice with PBS and permeabilized by incubating with 0.1% triton X-100 (Sigma) room temperature for 30 min. Permeabilized cells were rinsed once with PBS and blocked in 8% FBS (Gibco) in PBS for 2 hours at room temperature. 1:1000 of CF488A Streptavidin (Biotium) in blocking buffer was incubated with the cells at room temperature for 2 hours. After the incubation, the cells were washed with 0.1% Tween 20 in PBS and stained with 1:5000 DAPI (Biotium) for 3 min at room temperature and washed with PBS thrice. The prepared cover slips were transferred onto a glass slide coated with Prolong Gold Antifade (Thermo Fisher) and dried overnight at room temperature away from light.

Validation of intermolecular interactions by pulldown and qPCR. 10µg of DNase treated psoralen cross-linked total RNA was diluted in 300µL of water. The samples

were then incubated with 100pM of biotinylated probes, specific to the gene of interest, in fresh Hybridization Buffer (750 mM NaCl, 1% SDS, 50 mM Tris-Cl pH 7.0, 1 mM EDTA, 15% formamide) supplemented with Superscript-in (1:200), and incubated at 37°C overnight with shaking. After hybridization of the probes, 100µL of Dynabeads® MyOne™ Streptavidin C1 beads was used to pull out the RNA complexes. The beads were washed 5 times with wash buffer (0.1x NaCl and Sodium citrate (SSC), 0.5% SDS) that has been pre-warmed to 37 °C. RNA was eluted from the beads by incubating with 100µg of proteinase K (Thermo Scientific) in 95 µL of PK Buffer (100mM NaCl, 10 mM TrisCl pH 7.0, 1mM EDTA, 0.5% SDS) at 50 °C for 45min with end to end shaking, and boiling at 95 °C for 10 minutes. Eluted RNA was recovered by using TRIzol reagent, and cleaned up using RNeasy MinElute Cleanup Kit (Qiagen). The recovered RNA was eluted in 20 µL of nuclease free water, irradiated with 254nm UV for 5 min for reverse crosslinking. The samples were subsequently used for qPCR. Anti-GFP oligos were used as a negative control.

Validation of intramolecular interactions. 100 µg of DNase treated psoralen or DMSO cross-linked total RNA was fragmented with RNA fragmentation buffer (9 mM MgCl₂, 225 mM KCl and 150 mM Tris HCl (pH 8.3) at 95°C for 3 min. After fragmentation, the RNA was size fractionated using a 6% TBE 8M Urea gel and RNA fragments corresponding to 100-300 bases were excised and eluted overnight at 4°C. 5 µg of RNA were used for the hybridization in the same conditions as in the pull down of intermolecular interactions. All probe and qPCR primer sequences for the pull down and qPCR are compiled in Table 7.

Data analysis

Overview of the Computational Pipeline. The SPLASH pipeline automates read processing, mapping, interaction detection and filtering by using the snakemake workflow management system (version 3.4.1 (Koster and Rahmann, 2012)). See Supplementary Figure 3B for a flowchart.

Human Transcriptome. To construct a transcriptome we downloaded all transcripts for hg19 RefSeq genes from the UCSC Table Browser. We then grouped isoforms into genes based on their gene names, and took the longest coding isoform, or if absent,

the longest non-coding isoform as the representative of each gene. To this we added manually curated versions of snoRNAs, snRNAs (extracted from NCBI) and tRNAs (extracted from the UCSC Table Browser) and also replaced the complete repeating rRNA unit (U13369) with the resp. single rRNA sequences (including 5S rRNA and spacers) matching the used PDB structure (see below). This set of sequences was then deduplicated Bbmap's `dedup.sh` (options `absorbcontainment=t minoverlappercent=11 absorbbc=f`; <http://sourceforge.net/projects/bbmap/>). Any non-coding entry that did not belong to either miRNAs, rRNAs, snoRNAs, snRNAs or tRNAs was marked as small non-coding RNA, if its sequence was shorter than 200bp, or lncRNA otherwise.

Yeast Transcriptome. To construct the yeast transcriptome, we extracted the sequences of yeast coding genes from UCSC Table Browser (`sacCer3`, `sgdGene`), and added in UTR sequences to the transcripts based on Nagalakshmi et al (Nagalakshmi et al., 2008). We then supplemented the sequences of noncoding genes, including rRNAs, tRNAs, snRNAs, snoRNAs and ncRNAs downloaded from Saccharomyces Genome Database. Duplicated sequences were then removed to yield the yeast transcriptome used in this study.

Processing of Sequencing Reads. Reads were preprocessed with SeqPrep (version 1.0-7; <https://github.com/jstjohn/SeqPrep>) to remove adapters and merge overlapping paired-end reads into single reads of high quality. To speed this time consuming step up we parallelized the processing by working on split FastQ files. Since the majority of our paired-end reads should overlap, we used only the successfully merged ones for further analysis. Merged reads were mapped to the transcriptome (see above) with BWA MEM (version 0.7.12; and [arXiv:1303.3997v1](https://arxiv.org/abs/1303.3997v1)). We tuned BWA's parameters such that regions of minimum length 20 were detectable (`-T 20`; as opposed to the default 30). Mapped reads were sorted and converted to BAM with samtools (version 1.1). Afterwards, we removed all but the first read aligning to identical start coordinates and with identical CIGAR strings, which aggressively filters potential PCR duplicates (Ramani et al., 2015).

Detection of long range RNA interactions. To detect RNA interactions we scanned the BAM file for primary alignments containing BWA MEM's split alignment (SA) tag. We then discarded split alignments less than 50bp apart. This mainly serves two purposes: 1) these would likely always evaluate as true in our PDB-based evaluation

(see below) because bases are very close in sequence and therefore in structure and 2) we want to focus on the detection of long range RNA interactions. In addition we discarded interacting pairs where either end is mapped as reverse complement (transcriptome mapping) or has a mapping quality below 20. The latter effectively removes ambiguously mapped reads as well as alignments with close second best hits (e.g. pseudogenes).

Removal of splicing related false positives interactions. To deal with false positive interactions caused by splicing events, we remapped split reads from the transcriptome mapping back to the human genome (hg19) using the program STAR, and removed any read that entirely spans an annotated junction, allowing less than 5 bp soft-clip for both ends. The parameters of running STAR are: `--twopassMode Basic --alignSplicedMateMapLminOverLmate 0.1 --outSJfilterOverhangMin 10 6 6 6 --outSJfilterCountUniqueMin 6 1 1 1 --outSJfilterCountTotalMin 6 1 1 1 --outSJfilterDistToOtherSJmin 5 0 5 0 --alignSJDBoverhangMin 3 --alignMatesGapMax 1000000 - -alignIntronMax 1000000 --alignSJstitchMismatchNmax 5 -1 5 5 --outStd SAM --outSAMtype SAM --winAnchorMultimapNmax 9000 --seedPerWindowNmax 1000 --outSAMstrandField None --outSAMmultNmax 1 --outMultimapperOrder Random --outSAMattributes All --outSAMprimaryFlag AllBestScore --outFilterMultimapScoreRange 0 --outFilterMultimapNmax 9000 --outFilterMismatchNmax 2 --outFilterIntronMotifs None --outFilterMatchNminOverLread 0.1 --outFilterScoreMinOverLread 0.1 --alignEndsTypeLocal".--outSAMmultNmax 1 --outMultimapperOrder Random --outSAMattributes All --outSAMprimaryFlag AllBestScore --outFilterMultimapScoreRange 0 --outFilterMultimapNmax 9000 --outFilterMismatchNmax 2 --outFilterIntronMotifs None --outFilterMatchNminOverLread 0.1 --outFilterScoreMinOverLread 0.1 --`

The junction information was downloaded from the ENCODE project database Release 19 (GRCh37.p13).

Evaluation of ribosomal RNA interactions. To evaluate predicted rRNA-rRNA interactions we used the human 80S ribosome (PDB 4V6X), a cryo-EM structure with 5 Angstrom resolution. Each interaction pair window was mapped to the base combination with minimum 3D distance in the PDB structure. For each base we

computed its centroid 3D position and counted a base pair as true, if its respective centroid distance was smaller than 30 Angstrom.

Comparison of sensitivity versus specificity between DMSO, psoralen and bio-psoralen libraries. True base-pairs of 28S rRNA were determined from Petrov et al.(Petrov et al., 2014). Results for RPL was obtained from Ramani et al.(Ramani et al., 2015), and processed as described in their paper and accompanying scripts. The smoothing step was omitted in an alternative analysis to evaluate RPL with minimal post-processing. In both cases the data were then coarse-grained into 100-base windows for direct comparison with SPLASH. The receiver operating characteristic (ROC) curve was then obtained by varying the threshold above which RPL value was deemed to have identified a hit. Similarly, we varied the threshold for SPLASH, systematically increasing the cutoff for identifying hits while still retaining the requirement of having consensus with at least two replicates and total reads of at least 8.

Evaluating the solvent accessibility of bio-psoralen. We consolidated the frequency each base-pair nucleotide appeared in a sequencing read, and estimated the corresponding base-pairs solvent accessible surface area (SASA) as the sum of the SASA of all the nucleotides in the identified base-pair, its preceding and succeeding base-pairs (i.e. total SASA of three consecutive base-pairs). Nucleotide SASA was evaluated using FreeSASA.

Prediction of snoRNA-rRNA interactions. Potential interaction sites of C/D box snoRNAs and the rRNA were predicted with Plexy in conjunction with RNAplex (version 2.1.9). To include weaker interactions the default energy threshold was removed. Interaction interfaces and energies for each predicted interaction were recorded for visualization.

Hybridization energies of RNA interactions. Hybridization energies for 1000 randomly chosen non-rRNA chimeras from human lymphoblastoid cells were computed with RNAduplex (ViennaRNA version 2.1.9). For each observed interaction, we also created a random equivalent, by shuffling the observed sequence preserving dinucleotide content. P-values were computed with Kolmogorov-Smirnov tests.

Visualization. For drawing classical RNA 2D structures we used VARNA (version 3.93). Arc diagrams were plotted using R4RNA.

Classification of RNA classes by circularization score. Circularization score for each mRNA is calculated by taking the average of all pair-wise intramolecular interactions in the RNA, and dividing by RNA length. P-value for boxplots were calculated using Wilcoxon rank sum test.

Association between RNA interactions, translation efficiency and decay. Translation efficiency, obtained from ribosome profiling data (Guo et al., 2010), was calculated for mRNAs with top and bottom 20% of circularization scores. For the association of the location of intramolecular interactions with translation, translation efficiency was calculated for mRNAs with interactions only in the 5' UTRs, versus all other interactions. Translation efficiency for human ES cells and RA cells was estimated from conserved mRNAs using mouse ES and mouse differentiated ribosome profiling data. mRNA decay was calculated for mRNAs with intramolecular interactions present only at the first, and last one third of the transcript, versus all over the transcript.

Two-dimensional RNA interactome maps. To generate a global view of intra-, or intermolecular mRNA-mRNA interaction as a heatmap, we analyzed the last 200 bases of 5'UTR, first and last 400 bases of CDS and the first 400 bases of 3'UTR, centered around the start/codon for each detected transcript. As each bin represents 100 bases along the transcript, we have 14 bins across the 5' UTR, CDS and the 3' UTR region in total. We then calculated the observed interactions on the 14x14 matrix.

We used resampling tests to assess the significance of observed interactions in each bin within the matrix. Specifically, for each interaction, we generated a resampled interaction by randomly picking a pair of positions, weighted by the coverage of non-chimeric reads at the respective positions, from the same transcript as the observed interaction. We then aggregated all of resampled interactions in a 14x14 (or 10x10) matrix. Resampling was repeated 10,000 times. The p-value of observed number of interactions in each bin was calculated from this empirical distribution. Enrichment values as presented as $\log_{10}(\text{p-value})$.

Enrichment of intermolecular mRNA interactions in different cellular compartments. We downloaded the nuclear and cytoplasmic polyA+ RNA-seq data for the GM12892 lymphoblastoid cell line from the GEO database under accession number GSM758560 and GSM765386. The raw reads were mapped to Human

Genome (hg19) by STAR (v2.5.0) and FeatureCounts (v1.4.6) was used to count the number of raw reads for each gene, using GTF file downloaded from Ensembl (vGRCh37.75). We took genes with more than 10 reads in two out of four samples, and used a variance stabilizing transformation algorithm to normalize read counts across different replicates and conditions using DEseq2. The nuclear vs. cytoplasmic enrichment ratio was calculated for each gene by comparing normalized read counts between nuclear and cytoplasmic samples. We defined a gene as either nuclear- or cytoplasmic-enriched if the log2 nuclear vs. cytoplasmic enrichment ratio was greater than 2 or less than -2 respectively. We then used resampling to test the significance of enrichment of inter-molecular interactions (IMIs) among RNAs present in the same cellular compartment. We first grouped interactions based on the cellular compartmentalization of each partner, such as “cytoplasmic RNA – cytoplasmic RNA” and “cytoplasmic RNA – nuclear RNA”. We then sampled the same number of genes from all expressed genes, requiring the distribution of gene expression (estimated from non-chimeric reads, which were mapped without splitting and derived from SPLASH libraries) to be similar to the genes with IMIs. Resampling was repeated 10,000 times. The observed number of IMIs was compared to the number of IMIs from the resampled gene sets for each cellular compartment, and the relative rank of observed IMIs was converted into the enrichment p-value accordingly.

Intermolecular interaction network analysis and correlation with gene regulation. mRNA-mRNA interaction network was constructed by excluding all disconnected edges and extracting modules from the network using the fast-greedy algorithm. We calculated the significance of correlation with gene regulation between pairs of mRNA genes within each of these modules by extracting datasets for gene expression, translation efficiency and decay rates and calculating the pair-wise Pearson correlation for all gene pairs within each module. The significance of correlation was then accessed by permuting the modules 10000 times.

Gene Ontology (GO) enrichment analysis of interaction modules. We used the TopGO package to access the functional enrichment of genes in each individual module in yeast, lymphoblastoid, ES and RA cells, with respect to biological process, molecular function and subcellular components. Genes in each module were compared against all genes with intermolecular interactions detected. The significance

level of enrichment was computed with the “elim” algorithm implemented in TopGO. All reported enrichment terms are based on a false discovery rate threshold of 0.05.

RESULTS

The SPLASH protocol enriches effectively for *in vivo* RNA-RNA hybrids

To develop SPLASH, we used a biotinylated version of the crosslinker psoralen (bio-psoralen, Figure 8A) to identify intramolecular and intermolecular RNA-RNA base pairing. Psoralen enters the cells and intercalates into base paired nucleotides, preferentially crosslinking pyrimidines, especially at Ts and Us, at UV 365nm. The crosslinked RNAs were then extracted, fragmented and enriched for the crosslinked regions using streptavidin beads before undergoing proximity ligation and conversion into a deep sequencing library (Figure 1A). Importantly, the use of bio-psoralen allows the preservation of RNA interactions in living cells, akin to the use of formaldehyde as an *in vivo* crosslinker for detecting protein-DNA and DNA-DNA interactions in chromatin immunoprecipitation (ChIP) and chromatin conformation capture experiments (Hi-C). In particular, the reversibility of bio-psoralen crosslinking at UV 254nm enables reverse transcription across the ligated chimeras during library preparation. The biotin group on bio-psoralen also allows the selective enrichment of crosslinked interaction sites during the library preparation process, increasing the signal of pairwise interactions over the background of non-interacting sites (Figure 1A).

While psoralen has been used to crosslink nucleotides *in vivo*, we observed that the entry of bio-psoralen into human cells was low. To increase the cellular uptake of bio-psoralen, we incubated cells with different concentrations of bio-psoralen, and in the presence of 0.01% digitonin, a mild detergent. Treating human cells with digitonin for 5 min significantly increased the entry of bio-psoralen as determined by immunofluorescence staining (Figure 8B). We confirmed that a brief treatment of digitonin does not change psoralen crosslinking patterns (Figure 8C) and titrated the amount of time it takes to efficiently crosslink RNAs *in vivo* (Figure 8D). As RNA structure probing typically aims for “single hit kinetics”, with few modified molecules per transcript, we titrated the concentration of bio-psoralen used such that it crosslinks at a frequency of approximately one in every 150 bases in the human transcriptome (Figure 8E; Experimental Procedures). Bio-psoralen can also enter and crosslink

RNAs in yeast and *E. coli in vivo*, although a higher concentration is needed (Figure 8F,G). We confirmed that bio-psoralen can crosslink and detect known RNA interactions *in vivo* by performing a northern blot assay to detect known RNA-RNA interactions, such as the U14-18S binding in yeast (Figure 8H).

The reversibility of psoralen crosslinking is key to the success of our library preparation process, however complete reverse crosslinking typically takes about 30 min at UV 254nm, dramatically damaging RNA in the process. We titrated the duration of UV 254nm exposure to the crosslinked RNAs, and identified conditions that maximized the amount of reverse crosslinking while minimizing UV damage (Figure 9A, B). As library preparation processes typically involve multiple steps with different biases, we tested two different library cloning protocols that utilized i) 3' adapter ligation followed by reverse transcription, cDNA circularization and PCR (circularization protocol, Figure 9F), as well as ii) independent 5' and 3' adapter ligations followed by reverse transcription and PCR (RNA ligation protocol) (Table 1). We found that the circularization protocol resulted in less bottlenecking and more efficient capture of chimeric reads than the RNA ligation protocol, while maintaining similar accuracy in capturing real chimeras based on the human 80S ribosome crystal structure (Table 1). We also identified fragmentation conditions that enabled us to fragment crosslinked RNA to ~100 bases without losing the biotin group (Figure 9C), and stringent ligation and wash conditions that enabled us to generate SPLASH libraries with low background noise (Figure 9D,E; Experimental Procedures).

The SPLASH computational pipeline identifies RNA interactions with high specificity

We integrated SPLASH data with a robust computational pipeline that was developed to accurately identify RNA-RNA interactions in the transcriptome. The pipeline stringently removes PCR duplicates, merges paired-end reads and then split maps them along the human and yeast transcriptomes to identify chimeric reads that indicate an RNA-RNA interaction (Experimental Procedures; Figure 9G). Chimeric reads were filtered for splicing artefacts and clustered to identify pairwise interactions (Experimental Procedures). Additionally, all pairwise interactions that are continuous or are spaced less than 50 bases apart were removed, to focus the analysis on the

long-range intramolecular and intermolecular interactions that cannot currently be reliably predicted using computational Experimental Procedures (Experimental Procedures). Overall, our stringent filtering retained 4.6 million chimeric reads (0.4% of all sequenced reads) that identify RNA-RNA interactions across the different transcriptomes. The resulting interactions span a wide range of distances, from 50 to 5000 bases, with a median distance of 300 bases (Figure 10A).

To evaluate sensitivity and precision, intramolecular interactions reported by SPLASH analysis were compared to the crystal structure of the human 80S ribosome. Assessing regions of close spatial proximity in the crystal structure showed that SPLASH predictions provide a good balance between precision (75%) and sensitivity (78%) ($<30\text{\AA}$; Figure 1C; Experimental Procedures). Visualising these interactions on the known secondary structure of the large ribosomal subunit highlights the dense network of long-range RNA interactions that were captured by SPLASH data (Figure 1B). To estimate false discovery rate in reported interactions, we mixed independently crosslinked human and yeast total RNAs in equal amounts to prepare SPLASH libraries. Based on the observed human-yeast pairwise interactions, we estimated a false discovery rate of $<3.7\%$, confirming that SPLASH analysis provides good control over the fraction of false interactions reported overall (Experimental Procedures). To verify that the SPLASH interactions are mediated by psoralen/bio-psoralen crosslinking events, we performed libraries without crosslinking, with psoralen crosslinking and with bio-psoralen crosslinking (SPLASH). Libraries generated with bio-psoralen showed similar levels of high specificity and increased sensitivity as compared to psoralen libraries (Figure S3B), as expected due to enrichment of crosslinking sites by streptavidin beads in SPLASH. In contrast, non-crosslinked libraries showed low overall specificity (17%), confirming that psoralen/bio-psoralen crosslinking is essential to keep interacting RNA partners in close proximity for correct ligations to occur preferentially (Figure S3B).

To further confirm that SPLASH chimeras are enriched for ligation events between crosslinked fragments and not random background ligations, we generated libraries without ligase, with ligase and with $1/10^{\text{th}}$ of the amount of ligase used in SPLASH. Libraries without ligase show a low level of background ligation, indicating that most

pairwise interactions are due to intended proximity ligation events enabled by bio-psoralen crosslinking (Table 1, Figure 10C). Furthermore, correlation analysis between the frequency of crosslinking events and solvent accessibility of a region demonstrates that bio-psoralen crosslinking is largely independent of solvent accessibility (Figure 10D). Finally, benchmarking SPLASH against a recently published proximity ligation based approach indicates that SPLASH has similar precision for detecting intramolecular interactions, while detecting significantly more intermolecular interactions (Figure 10E,F).

Global structure of the yeast and human RNA interactomes

To study RNA interactomes and their dynamics in different organisms, SPLASH was performed on 2-4 biological replicates of human cells, including Hela cells, lymphoblastoid cells, human embryonic stem (ES) cells and retinoic acid (RA) differentiated cells, as well as in wild type and Prp43 helicase mutant *S. cerevisiae* (Table 1,2). In addition, we performed sequencing on total RNA, poly(A)+ enriched, and snoRNA enriched RNA populations in different cell lines to capture RNA-RNA interactions globally and comprehensively. Based on more than two billion Illumina sequencing reads all together, we identified >8,000 intermolecular and >4,000 intramolecular interactions across different cell types (Table 2). We observed a high correlation between biological replicates in the same cell line ($R=0.75-0.9$) confirming that SPLASH data is reproducible (Figure 11A). Overall, 3,497 intramolecular mRNA interactions and 84 lncRNA interactions from 1,311 genes were captured in this study, providing a rich resource for studying human RNA structure and function (Figure 2A). Intermolecular interactions were found to be notably diverse and common, including 990 mRNA-mRNA interactions identified in human cell lines alone (Figure 2B). Similar diversity was captured in thousands of yeast intramolecular and intermolecular interactions, enabling the identification of conserved human-yeast interaction features (Figure 11B, Table 3). The degree distributions of the intermolecular interaction networks were found to have a good fit to a power law distribution for degree less than 10, but were accompanied by a heavy tail with many nodes with large degrees (Figure 11C).

Long-range intramolecular RNA interactions define distinct classes of functional RNAs

To determine if our identified intramolecular interactions are highly stable, we calculated the energy of interactions between true chimeric pairs versus randomly shuffled chimeras with dinucleotide content preserved. Indeed, internal pairwise interactions have lower base-pairing energy compared to the shuffled set ($p < 10^{-6}$, KS test, Figure 12A), indicating that the chimeras are likely to be real. Comparing SPLASH intramolecular interactions with RNAs of known secondary structures, including LSR1, SCR1 and snR86 RNAs in yeast, further confirmed that SPLASH interactions are consistent with known interactions previously found by either biochemical or crosslinking experiments (Figure 12B-D). We also validated a long-range intramolecular interaction in a yeast mRNA, demonstrating the reproducibility of our method (Figure 12E).

To determine if there are differences in the propensity of different classes of RNAs to form long-range pairwise interactions, we calculated a "circularization score", which is an average of interaction distances within a transcript normalized by its length (Figure 2C). Classifying different groups of RNAs according to their circularization score revealed that rRNAs and lncRNAs tend to form more distant interactions than mRNAs (Wilcoxon rank sum test, $p = 0.0045$ and $p = 0.028$ respectively, Figure 2D). Long-range interactions in rRNAs likely contribute to their complex and highly structured conformation. Similar interactions in a subset of lncRNAs may be indicative of structure-mediated functions, such as acting as a scaffold to recruit different protein factors for cellular function.

The structural organization of mRNAs inside cells can impact their regulation and function. Using long-range interactions inferred from SPLASH for the human transcriptome, we constructed two-dimensional heatmaps of enriched interaction sites along a transcript (Figure 3A). The heatmaps bring to light the highly modular nature of an average mRNA in the human cell, with bases in 5' untranslated regions (UTRs), CDS, and 3' UTRs preferentially interacting with other bases in the same domain, and extensions beyond the domain boundaries occurring right at the start and end of the coding region. Sequence composition was found to not correlate with the observed

patterns, with bases near the beginning of mRNAs having very low pyrimidine content, but high levels of psoralen crosslinking (psoralen preferentially crosslink pyrimidines; Figure 12F). Metagene analysis on intramolecular interaction sites aligned along translation start and stop codons also confirmed enrichment for interactions near the start of the mRNA (Figure 3B).

To characterize the impact of interaction domains on mRNA function, mRNAs were grouped according to their propensity to form long-range pairwise interactions (circularization score) and assessed for translation efficiency. This analysis revealed that on average, mRNAs with shorter pairwise interactions are translated less efficiently than mRNAs with longer interactions (Figure 3C). Furthermore, classifying transcripts according to the location of their pairwise interactions revealed that mRNAs with pairwise interactions only in the 5' UTRs tend to be translated slowly, consistent with a model whereby 5' end blocking of mRNAs can reduce translation efficiency (Figure 3D). Sorting transcripts by translation efficiency to detect associated interaction patterns revealed two additional features. Firstly, efficiently translated mRNAs tend to have chimeras that span a longer distance, often connecting the beginning and ends of transcripts (Figure 3E), in support of a circularization model for ribosome recycling and efficient mRNA translation (Wells et al., 1998). Secondly, poorly translated mRNAs tend to contain short chimeras that are clustered near the beginning of the transcript (Figure 3E), highlighting that mRNAs with 5' end interactions are poorly translated. Similar conclusions were obtained when mRNAs were clustered in an unbiased manner based on the location of their interactions (Figure 12G,H). Taken together, our data suggests a transcriptome-wide role for translation inhibition by stable structures near the start codon, as well as increased translation efficiency by end-to-end circularization.

Analysis of mRNA decay information revealed a similar influence of mRNA structure on RNA stability (Figure 3F). Genes with interactions that are confined to the 5' end exhibited the fastest decay rates as compared to control, suggesting that interactions at the 3' end could block the exosome complex during RNA degradation, and emphasizing the importance of structure in post-transcriptional gene regulation.

SPLASH uncovers new rRNA-rRNA and snoRNA-RNA interaction sites

Psoralen intercalates into base-paired regions independent of whether they are formed by the same RNA strand, or between two different RNA strands, enabling SPLASH to interrogate both intra- and intermolecular RNA interactions. As expected, SPLASH captures well-characterized intermolecular interactions corresponding to 5.8S-28S rRNAs, as well as between U4-U6, and U2-U6 snRNAs. In addition, SPLASH analysis identified many known snoRNA-rRNA interactions in the literature, validating the high sensitivity of our approach (Figure 4A, Figure 13A).

SnoRNAs are an important class of non-coding RNAs that guide the maturation of pre-ribosomal RNAs to form the functional ribosome. While the binding regions of some snoRNAs have been identified, the location of many snoRNA-rRNA interactions in the human ribosome remains elusive. Recently, snoRNA-rRNA interactions have been hypothesized to be more widespread than previously appreciated. However, snoRNA target prediction, especially for H/ACA snoRNAs which binds to rRNAs with short complementary stretches, still remains challenging, and experimental strategies such as CLASH have been applied mainly to detecting C/D box snoRNAs with rRNAs in yeast.

To identify snoRNA-rRNA interactions genome-wide in humans, SPLASH was performed on lymphoblastoid cells. Analysis of the trimethylated snoRNA immunoprecipitation libraries, as well as the deeply sequenced total RNA libraries identified 211 human snoRNA-rRNA interactions, corresponding to 78 human snoRNAs (55 C/D box and 23 H/ACA snoRNAs) (Table 4, Experimental Procedures). Based on the human snoRNA database, 122 out of the 211 identified snoRNA-rRNA sites are new, and include target sites for orphan snoRNAs such as SNORA51 (ACA51) and SNORD83. We validated three new snoRNA-rRNA interactions that were captured at different abundances by performing pulldown of 5S, 18S and 28S rRNAs individually and qPCR of the snoRNAs. While SNORA32 was previously thought to only bind to 28S rRNA based on the human snoRNA database, we identified and validated that SNORA32 binds strongly to the 5S rRNA (Figure 4B). SNORD83a is an orphan snoRNA which we identified and validated to bind to the 18S rRNA (Figure 4B). We also validated the weak binding of the orphan snoRNA ACA51 to the

28S rRNA (Figure 13B), suggesting that SPLASH data is accurate even at low chimeric read counts. Predicted snoRNA-rRNA interaction sites from SPLASH were also integrated with a snoRNA prediction program (PLEXY) to refine binding site predictions (Figure 4C,D). Using PLEXY, we narrowed down a new potential U13-28S rRNA binding site to bases 4418-4424 along the 28S rRNA (Figure 4D). Thus, combining high throughput experimental data with snoRNA prediction algorithms can facilitate systematic, high-resolution identification of new snoRNA-rRNA interactions to improve our understanding of ribosome biogenesis.

Beyond human snoRNA-rRNA interactions, SPLASH analysis on two biological replicates of wild-type and Prp43 mutant yeast identified 106 target sites for 39 snoRNAs, including 27 C/D Box and 12 H/ACA snoRNAs (Table 5). For example, we identified the known target site of snR61, a C/D Box snoRNA, as well as two new binding sites on the 25S rRNA (Figure 5A). The snR61 crosslinking site at bases 2800-2900 on the 25S rRNA is also predicted by PLEXY, further refining the location of this new site (Figure 5B). We also identified target sites for the snR4 C/D box snoRNA, which was previously thought to be inactive. snR4-25S interactions were previously reported in CLASH data as low confidence hits that were not reproducibly found in all replicates. Here, we independently identified the same three snR4-25S rRNA interactions sites as in CLASH data, in addition to a new snR4-18S rRNA site, to support the existence of snR4-rRNA interactions (Figure 5C,D, Figure 13C). We also identified a target site for the orphan C/D Box snoRNA snR45 on 25S rRNA, in all 4 biological replicates, indicating that snR45 may play a role in 25S rRNA maturation.

As snoRNA-rRNA interactions are destabilized by helicases upon binding to pre-rRNA, SPLASH analysis in yeast cells that over-express the helicase Prp43 mutant (*prp43-T123A*, Figure 13D) was used to identify additional transient snoRNA-rRNA interactions that are important for rRNA biogenesis. The essential H/ACA box snR30 was previously found to be released from 18S rRNA by the Rok1 helicase and is required for 35S cleavage and release of the 18S rRNA precursor. In our analysis, we identified snR30-18S rRNA interactions in the Prp43 mutant but not in the wildtype (Figure 5E), suggesting that either multiple helicases can work on the same snoRNA substrate(s) to facilitate their release from rRNA or that Prp43 is required for Rok1 to

unwind snR30 from the pre-ribosome. This is consistent with previous reports that both the Dbp4p and Has1p RNA helicases are required for U14 release from the pre-ribosome. Our top interaction sites identified highly confident snoRNA-rRNA basepairs that preferentially accumulated in Prp43 mutant versus wildtype cells (Figure 5F). Many of these accumulated snoRNAs, including snR59, snR60, snR41 and snR55, were previously found to bind directly to Prp43, supporting the hypothesis that Prp43 is important for their release. SPLASH also provides evidence for new rRNA target sites for snR189, snR59, snR40, and snR69 in Prp43 mutant yeast, significantly expanding the list of interactions involved in snoRNA targeting and recycling.

mRNA-mRNA interactions define modules of co-regulated genes

Beyond snoRNA-rRNA interactions, SPLASH analysis identified nearly a thousand mRNA-mRNA interactions. We calculated the folding energies of these intermolecular mRNA interactions to determine whether they are likely to be stable. Intermolecular pairwise interactions exhibit not only lower folding energies than randomly shuffled chimeras with dinucleotide content preserved (median = -27.2 vs -21.85 kcal/mol, KS test, $p < 10^{-15}$), but also lower folding energies compared to intramolecular mRNA interactions (median = -19.7 kcal/mol), indicating that they are likely to be even more stable (Figure 14B). To estimate the true positive rate in intermolecular RNA interaction predictions, we experimentally tested them by using qPCR to determine enrichment of interaction partners after psoralen crosslinking and oligo pulldown (Figure 6A, B; Figure 14A). Overall, 12 out of 13 interaction pairs were validated, indicating high reproducibility and precision (92%) for intermolecular mRNA-mRNA predictions from SPLASH analysis.

To study the distribution of intermolecular interactions along an mRNA, we plotted the interaction density along the length of human mRNAs after aligning the transcripts according to their translation start and stop codons. Interestingly, most intermolecular interactions also occur near the beginning of the transcript (Figure 14C). However, unlike intramolecular interactions whereby RNA interactions tend to occur within the same domain, intermolecular interactions frequently involve the binding of the beginning of one mRNA to another distal region along the second mRNA (Figure 6C).

As a result, intermolecular 2D interaction plots displayed a much more spread-out interaction pattern across the transcript domains, and appear to be less modular.

Network analysis of the human RNA interactome identified a major mRNA interaction cluster that is strongly enriched for genes with RNA binding, metabolic, and translation properties (Figure 6D, Table 6). Hierarchical clustering of the human RNA interactome based on the density of pairwise interactions identified nine modules, showing distinct enrichment for genes with defined functions and subcellular localizations across modules (Figure 6E, Table 6). We observe that mRNAs tend to interact with other mRNAs in the same cellular compartment ($p < 0.05$, Figure 6F), confirming that physical proximity is necessary to drive intermolecular interactions with each other. We also observed that transcripts in mRNA modules can be coordinated in their gene regulation. This was observed for example in module 3 (a large group of translation related genes) which exhibited enrichment for correlated translation efficiency, as well as in module 1 (a group of RNA binding genes) which showed an enrichment for coordinated decay rates compared to controls (Figure 14D,E). These observations highlight the role of intermolecular mRNA interactions as a potential mechanism for coordinating post-transcriptional gene regulation inside cells, with interaction modules serving to refine cellular compartments in enriching for RNA interactions.

Beyond the static picture of the RNA interactome in human cells, the extent to which RNA interactomes are dynamic and rewired during different cellular states is unclear. To investigate the RNA regulatory network governing cellular pluripotency, we performed SPLASH in human ES cells as well as in retinoic acid (RA) differentiated cells. Globally, the intramolecular patterns of RNA interactions for ES and RA cells are highly modular (Figure 14F,G), similar to lymphoblastoid cells, suggesting that the modular pattern of mRNA intramolecular interactions are representative of most RNAs in different human cell types. Based on our previous observation that transcripts with high circularization scores tend to be translated better than those with low circularization scores, we hypothesized that mRNAs that undergo conformational changes can have corresponding changes in translation efficiency. To test this, we calculated the circularization scores for all well expressed genes in both ES and RA cells and identified mRNAs with high circularization scores in ES cells and low

circularization scores in RA cells, and vice versa. Interestingly, mRNAs that shift from having a high circularization score in ES to a low circularization score in RA cells showed a corresponding decrease in translation efficiency and vice versa (Figure 7A). This reaffirms the hypothesis that conformational changes can serve as an underlying mechanism to control translation efficiency during changes in cellular states. One of the chromatin genes, high mobility group 1, HMGA1, exhibited a notable decrease in circularization score and translation efficiency during RA differentiation, consistent with its key role in maintaining ES cell pluripotency (Shah et al., 2012) (Figure 7B). Protein and mRNA quantification using western blot and qPCR analysis showed that HMGA1 protein levels decrease after 5 days of differentiation, whereas its mRNA levels do not (Figure 7C, D). Furthermore, translation efficiency measured by ribosome profiling in mouse ES and differentiated cells showed a corresponding decrease in HMGA1 translation efficiency upon cellular differentiation (Figure S7H), reinforcing the association between structural rearrangement and translation.

Analysis of the intermolecular interactome network in ES and RA cells revealed that mRNAs are more highly interconnected to each other in ES versus RA cells, despite a similar number of detected mRNAs (ES, 277 genes and 402 interactions, RA, 193 genes and 180 interactions; Figure 7E,F). Module analysis of interacting RNAs in ES and RA cells further demonstrated the higher degree of interconnectedness in RNA interactions between ES cell modules when compared to RA cell modules (Figure 7G,H; Table 6). To determine which modules in the ES cell interactome are disrupted during differentiation, we calculated the number of genes that were dissociated from each module upon RA differentiation. We observed that module 3, which is enriched for chromatin remodeling processes, is disrupted during cellular differentiation ($p=0.0088$), consistent with the importance of chromatin remodeling in maintaining pluripotency.

Discussion

The advent of high throughput sequencing has enabled us to obtain a significant amount of sequence information across diverse transcriptomes. However, information in transcriptomes is not limited to their linear sequence and can be encoded in intra- and intermolecular RNA interactions. Studying how RNA molecules pair with

themselves and with others is thus key to understanding their function. The development and application of SPLASH to map pairwise RNA interactions has enabled the generation of transcriptome-wide maps in multiple human and yeast cell types, providing a global view of how transcripts are organized inter- and intramolecularly to impact gene regulation. Its application in different cell states also provides a view of the dynamic interactome and the functional impact of its remodeling during human ES cell differentiation.

Analysis of SPLASH data identified several key features in human interactomes, including the propensity of non-coding RNAs to form longer range interactions than mRNAs, and for mRNAs to adopt a modular configuration where the UTRs tend to interact with themselves and with nearby coding sequences. Interestingly, we do not see this modular pattern in intermolecular mRNA-mRNA interactions, with interactions being spread across the entire transcript. Follow-up experiments are needed to test various hypotheses for this observation, including the role of translation in maintaining mRNA modularity. Additionally, the role of (i) dense RNA interactions near the start codon for inhibiting translation, (ii) long-range end-to-end interactions for promoting efficient translation, and (iii) dense interactions near the 3' end for inhibiting mRNA decay, deserve further investigation. Collectively, our results provide evidence that structural organization of transcripts can play an essential role in gene regulation, and that changes in structural organization to regulate gene expression could be more widespread than previously anticipated.

Intermolecularly, we identified thousands of RNA-RNA interactions in human and yeast cells, including mRNA-rRNA, snoRNA-rRNA, mRNA-mRNA, and mRNA-lncRNA interactions. The majority of our interactions are mRNA-rRNA interactions, which we suspect to be a result of capturing mRNAs during translation. snoRNA-rRNA interactions are critical for ribosome maturation and misregulation of snoRNA abundances has been implicated in diseases such as cancer (Mannoor et al., 2012). Predicting snoRNA-rRNA targets, particularly for H/ACA snoRNAs, can be challenging. In this work, we detected existing and new target sites for 78 human snoRNAs (55 C/D box and 23 H/ACA snoRNAs), as well as for 39 yeast snoRNAs (27 C/D box and 12 H/ACA snoRNAs). The overlap between human and yeast datasets,

as well as between experimental and *in silico* predictions can thus be used to systematically refine and prioritize snoRNA-rRNA interactions for further validation and characterization. In yeast, at least 19 helicases are involved in recycling of snoRNAs after target binding. Our identification of snoRNA-rRNA interactions stabilized in the absence of the Prp43 helicase, highlights an avenue for obtaining additional mechanistic insights for other helicases involved in snoRNA release and ribosome biogenesis.

Mapping of genome-wide RNA interaction networks showed that mRNAs are organized in modules based on connectivity in the interaction network, and mRNAs in the same module are enriched for specific functions and subcellular localizations. These results suggest that RNA interaction modules containing genes of similar functions can be an organizing structure to coordinate translation and decay, and act as a mechanism for gene regulation. Human ES and RA interaction networks also showed that large RNA conformational changes *in vivo* are associated with corresponding changes in translation efficiency, indicating that (i) conformational changes are more widespread than previously appreciated, and (ii) that they could serve as underlying mechanisms for translation changes during ES differentiation. We also observed that the RNA interactome becomes sparser upon differentiation, with fewer mRNAs interacting with each other in differentiated cells, and that a chromatin remodeling associated module was additionally lost during differentiation. Further functional studies disrupting individual interactions in these modules could help understand the robustness of these modules and the key interactions that are involved in the differentiation process.

SUMMARY

In summary, SPLASH expands our understanding of the structural organization of eukaryotic transcriptomes, and helps to define the principles of how RNAs interact with themselves and with other RNAs in gene regulation and ribosome biogenesis. Apart from yeast and human cells, SPLASH is applicable to other organisms (such as *E. coli*) to interrogate RNA interactions under different cellular conditions. Coupled with genome-wide secondary structure mapping and RNA structure modeling, SPLASH data can help refine our current models of RNA structure with *in vivo* information.

SPLASH can also be combined with intermolecular RNA interaction prediction tools, such as snoRNA prediction programs, to improve the accuracy of these predictions. Techniques to enrich specific RNA fractions can be combined with SPLASH to further study rare RNAs. We anticipate that future studies using SPLASH will continue to shed light on the complexity and dynamics of RNA interactions in cellular systems across diverse organisms.

Table 1. Evaluation of different protocols for SPLASH, related to Figure 1.

Condition	No. of reads (merged pairs)	No. of rRNA Chimeric Dup_Removed	PPV
RNA ligation method			
No ligase	1700874	85	0.4824
0.1X ligase	1610716	143	0.5524
1X ligase	1787607	225	0.4889
Circligase method			
No ligase	2310350	117	0.6068
0.1X ligase	1671569	559	0.5510
1X ligase	2386429	1261	0.5234
Wash conditions			
Circligase with Wash buffer I (2X SSC)	2210723	1921	0.5122
Circligase with Wash buffer II (0.1x SSC with 15 % formamide)	2711055	3588	0.4961

Table 2. Information of sequenced SPLASH libraries, related to Figure 1.

Sample	Merged reads	Mapped reads	Mapped reads after PCR dups removal	No. of chimeric reads after filtering	Passed Intra Chimeras	Passed Inter Chimeras
Lymphoblastoid Cells Total RNA Replicate 1	53,747,987	53,309,764	3,228,021	311,079	147,850	163,229
Lymphoblastoid Cells Total RNA Replicate 2	29,859,136	29,865,379	3,081,880	239,724	105,429	134,295
Lymphoblastoid Cells Total RNA Replicate 3	60,332,939	59,565,659	3,482,737	208,630	99,687	108,943
Lymphoblastoid Cells Total RNA Replicate 4	45,400,893	45,667,145	3,923,185	279,368	123,773	155,595
Lymphoblastoid snRNA IP	149,071,830	148,389,870	2,218,555	174,295	68,603	105,692
Lymphoblastoid Cells PolyA Replicate 1	183,913,864	175,803,028	59,637,038	160,800	97919	62881
Lymphoblastoid Cells PolyA Replicate 2	115,234,808	110,846,796	43,847,439	109,274	58016	51258
Lymphoblastoid Cells PolyA Replicate 3	3,963,881	3,814,459	2,751,106	7,211	3819	3392
Lymphoblastoid Cells PolyA Replicate 4	53,371,012	50,589,091	15,197,222	82,921	64919	18002
Human ES PolyA Replicate 1	159,412,735	153,023,928	72,028,001	73,407	34674	38733
Human ES PolyA Replicate 2	68,290,966	66,046,593	21,635,588	42,109	14209	27900
Human RA PolyA Replicate 1	153,884,298	144,714,995	60,325,593	77,245	41090	36155
Human RA PolyA Replicate 2	87,849,979	82,593,496	24,842,241	26,661	12966	13695
Yeast Total RNA replicate 1	12,705,039	12,419,256	2,224,683	26,520	4,063	22,457
Yeast Total RNA replicate 2	29,292,969	28,655,555	5,854,878	39,665	13,635	26,030
Yeast Prp43 mutant Total RNA replicate 1	8,260,248	8,134,607	1,872,711	25,794	8,555	17,239
Yeast Prp43 mutant Total RNA replicate 2	31,719,541	31,194,218	2,609,303	34,530	5,877	28,653
Yeast PolyA replicate 1	16,412,271	16,261,107	9,768,660	167,092	3,910	163,182
Yeast PolyA replicate 2	13,414,403	13,255,714	8,636,175	225,395	3,700	221,695
Biotinylated psoralen libraries replicate 1	12,207,150	11,689,548	2,150,196	36,846	17,080	19,766
Biotinylated psoralen libraries replicate 2	9,114,009	8,876,983	2,052,463	86,823	34,450	52,373
Psoralen libraries replicate 1	32,453,663	32,500,917	1,898,212	112,574	53,980	58,594
Psoralen libraries replicate 2	8,176,811	8,123,956	818,852	18,368	7,681	10,687
DMSO libraries replicate 1	43,587,340	43,411,408	3,446,432	255,787	110,459	145,328
DMSO libraries replicate 2	35,095,430	34,981,150	2,951,955	205,383	92,172	112,911

Table 3. List of common human-human and human yeast interactions

Organism	Type	Human Gene 1	Human Gene 2	Yeast gene1	Yeast gene 2	Region 1	Region 2
Human-yeast	Intermolecular	PKM	EEF1A1	YAL038W	YBR118W		
Human-yeast	Intermolecular	RPL29	GAPDH	YFR032C-A	YGR192C		
Human-yeast	Intermolecular	RPS6	GAPDH	YBR181C	YGR192C		
Human-yeast	Intermolecular	YWHAE	GAPDH	YER177W	YGR192C		
Human-yeast	Intermolecular	TPI1	GAPDH	YDR050C	YGR192C		
Human-yeast	Intermolecular	GAPDH	TPT1	YGR192C	YKL056C		
Human-yeast	Intermolecular	GAPDH	RPL10	YGR192C	YLR075W		
Human-yeast	Intermolecular	RPL10	RPS3	YLR075W	YNL178W		
Human-yeast	Intermolecular	GAPDH	RPL3	YGR192C	YOR063W		
Human-yeast	Intermolecular	GAPDH	RPS12	YGR192C	YOR369C		
Human-human	Intermolecular	ACTB	RPL4				
Human-human	Intermolecular	ATP1A1	TPT1				
Human-human	Intermolecular	BTN2A2	LINC01604				
Human-human	Intermolecular	COX4I1	RPL10				
Human-human	Intermolecular	EDARADD	ENO1				
Human-human	Intermolecular	EEF1A1	EEF1G				
Human-human	Intermolecular	EEF1A1	GAPDH				
Human-human	Intermolecular	EEF1A1	hsnrna-RNU1-1				
Human-human	Intermolecular	EEF1A1	MTRNR2L8				
Human-human	Intermolecular	EEF1A1	RPL10A				
Human-human	Intermolecular	EEF1A1	RPL13				
Human-human	Intermolecular	EEF1A1	RPL18A				
Human-human	Intermolecular	EEF1A1	RPL22				
Human-human	Intermolecular	EEF1A1	RPL3				
Human-human	Intermolecular	EEF1A1	RPL31				
Human-human	Intermolecular	EEF1A1	RPL32				
Human-human	Intermolecular	EEF1A1	RPL35				
Human-human	Intermolecular	EEF1A1	RPL37				
Human-human	Intermolecular	EEF1A1	RPL37A				
Human-human	Intermolecular	EEF1A1	RPL41				
Human-human	Intermolecular	EEF1A1	RPL6				
Human-human	Intermolecular	EEF1A1	RPL7				
Human-human	Intermolecular	EEF1A1	RPL7A				
Human-human	Intermolecular	EEF1A1	RPL9				
Human-human	Intermolecular	EEF1A1	RPLP0				
Human-human	Intermolecular	EEF1A1	RPS15A				
Human-human	Intermolecular	EEF1A1	RPS2				
Human-human	Intermolecular	EEF1A1	RPS23				
Human-human	Intermolecular	EEF1A1	RPS27A				
Human-human	Intermolecular	EEF1A1	RPS3				
Human-human	Intermolecular	EEF1A1	RPS3A				
Human-human	Intermolecular	EEF1A1	RPS6				
Human-human	Intermolecular	EEF1A1	RPS7				
Human-human	Intermolecular	EEF1A1	RPS8				
Human-human	Intermolecular	EEF1A1	TPT1				
Human-human	Intermolecular	EEF1A1	TUBA1B				
Human-human	Intermolecular	EIF5A	PTMA				
Human-human	Intermolecular	ENO1	GAPDH				
Human-human	Intermolecular	FLJ44635	TPT1				
Human-human	Intermolecular	GAPDH	RPL13				
Human-human	Intermolecular	GAS5	U81				
Human-human	Intermolecular	GLTSCR2	PTMA				
Human-human	Intermolecular	GNB2L1	RPS12				
Human-human	Intermolecular	GPX1	RPL10				

Human-	Intermolecula	LRC5A	ZNF45				
Human-	Intermolecula	LYRM7	NOL10				
Human-	Intermolecula	NPM1	RPL18A				
Human-	Intermolecula	PAGR1	TRUB2				
Human-	Intermolecula	PIK3C2B	U80				
Human-	Intermolecula	RAB13	RAB8B				
Human-	Intermolecula	RPL10	RPL35				
Human-	Intermolecula	RPL13A	RPL18A				
Human-	Intermolecula	RPL35	RPLP2				
Human-	Intermolecula	RPL35	RPS28				
Human-	Intermolecula	RPL36A	RPL36AL				
Human-	Intermolecula	RPL37A	RPLP1				
Human-	Intermolecula	RPL3	RPS3				
Human-	Intermolecula	RPL41	hsnma-RNU1-1				
Human-	Intermolecula	RPL41	RPS17				
Human-	Intermolecula	RPL41	RPS3				
Human-	Intermolecula	RPL5	TMSB4X				
Human-	Intermolecula	RPLP1	RPS3				
Human-	Intermolecula	RPS11	RPS15A				
Human-	Intermolecula	RPS20	TMEM70				
Human-	Intermolecula	RPS3	RPS6				
Human-	Intermolecula	TPM3	TRUB2				
Human-	Intermolecula	TPM3	ZNF485				
Human-	Intramolecula	ABI1	ABI1			1000-1100	1100-1200
Human-	Intramolecula	ACTB	ACTB			1400-1500	1700-1800
Human-	Intramolecula	ACTG1	ACTG1			0-100	1900-2000
Human-	Intramolecula	ACTG1	ACTG1			0-100	200-300
Human-	Intramolecula	ACTG1	ACTG1			100-200	200-300
Human-	Intramolecula	ACTG1	ACTG1			1500-1600	1900-2000
Human-	Intramolecula	ACTG1	ACTG1			1600-1700	1700-1800
Human-	Intramolecula	AKR1A1	AKR1A1			0-100	200-300
Human-	Intramolecula	AKR1A1	AKR1A1			300-400	500-600
Human-	Intramolecula	ANAPC11	ANAPC11			200-300	600-700
Human-	Intramolecula	AP2M1	AP2M1			200-300	300-400
Human-	Intramolecula	APEX1	APEX1			100-200	1200-1300
Human-	Intramolecula	ARL6IP1	ARL6IP1			1500-1600	1700-1800
Human-	Intramolecula	ATF4	ATF4			100-200	800-900
Human-	Intramolecula	ATG13	ATG13			1300-1400	1600-1700
Human-	Intramolecula	ATG3	ATG3			1200-1300	2800-2900
Human-	Intramolecula	ATP5A1	ATP5A1			100-200	500-600
Human-	Intramolecula	ATP5B	ATP5B			1000-1100	1100-1200
Human-	Intramolecula	ATP5D	ATP5D			500-600	900-1000
Human-	Intramolecula	ATP5G3	ATP5G3			400-500	1100-1200
Human-	Intramolecula	ATP6V0B	ATP6V0B			900-1000	1700-1800
Human-	Intramolecula	BSG	BSG			200-300	600-700
Human-	Intramolecula	BTF3	BTF3			100-200	300-400
Human-	Intramolecula	C12orf57	C12orf57			300-400	400-500
Human-	Intramolecula	C14orf2	C14orf2			0-100	300-400
Human-	Intramolecula	C19orf70	C19orf70			400-500	600-700
Human-	Intramolecula	CALM2	CALM2			4000-4100	4200-4300
Human-	Intramolecula	CALM2	CALM2			4100-4200	4300-4400
Human-	Intramolecula	CCNB1IP1	CCNB1IP1			400-500	600-700
Human-	Intramolecula	CCNG1	CCNC1			1400-1500	2100-2200
Human-	Intramolecula	CCT2	CCT2			1400-1500	1500-1600
Human-	Intramolecula	CCT8	CCT8			1400-1500	1800-1900
Human-	Intramolecula	CCT8	CCT8			800-900	900-1000
Human-	Intramolecula	CD55	CD55			1300-1400	1500-1600
Human-	Intramolecula	CIRBP	CIRBP			600-700	1000-1100
Human-	Intramolecula	CNN2	CNN2			700-800	800-900

Human-	Intramolecula	COPZ1	COPZ1			0-100	200-300
Human-	Intramolecula	COX4I1	COX4I1			200-300	700-800
Human-	Intramolecula	COX7C	COX7C			0-100	100-200
Human-	Intramolecula	CTNNB1	CTNNB1			2600-2700	3100-3200
Human-	Intramolecula	DNPH1	DNPH1			300-400	500-600
Human-	Intramolecula	DYNC1I2	DYNC1I2			500-600	600-700
Human-	Intramolecula	EDF1	EDF1			400-500	600-700
Human-	Intramolecula	EEF1A1	EEF1A1			0-100	100-200
Human-	Intramolecula	EEF1A1	EEF1A1			0-100	1100-1200
Human-	Intramolecula	EEF1A1	EEF1A1			0-100	1200-1300
Human-	Intramolecula	EEF1A1	EEF1A1			0-100	1300-1400
Human-	Intramolecula	EEF1A1	EEF1A1			0-100	1600-1700
Human-	Intramolecula	EEF1A1	EEF1A1			0-100	800-900
Human-	Intramolecula	EEF1A1	EEF1A1			1000-1100	1100-1200
Human-	Intramolecula	EEF1A1	EEF1A1			1000-1100	1200-1300
Human-	Intramolecula	EEF1A1	EEF1A1			1000-1100	1300-1400
Human-	Intramolecula	EEF1A1	EEF1A1			100-200	1100-1200
Human-	Intramolecula	EEF1A1	EEF1A1			100-200	1200-1300
Human-	Intramolecula	EEF1A1	EEF1A1			100-200	1400-1500
Human-	Intramolecula	EEF1A1	EEF1A1			100-200	1500-1600
Human-	Intramolecula	EEF1A1	EEF1A1			100-200	200-300
Human-	Intramolecula	EEF1A1	EEF1A1			100-200	400-500
Human-	Intramolecula	EEF1A1	EEF1A1			100-200	700-800
Human-	Intramolecula	EEF1A1	EEF1A1			100-200	800-900
Human-	Intramolecula	EEF1A1	EEF1A1			100-200	900-1000
Human-	Intramolecula	EEF1A1	EEF1A1			1100-1200	1200-1300
Human-	Intramolecula	EEF1A1	EEF1A1			1100-1200	1300-1400
Human-	Intramolecula	EEF1A1	EEF1A1			1100-1200	1400-1500
Human-	Intramolecula	EEF1A1	EEF1A1			1100-1200	1600-1700
Human-	Intramolecula	EEF1A1	EEF1A1			1200-1300	1300-1400
Human-	Intramolecula	EEF1A1	EEF1A1			1200-1300	1400-1500
Human-	Intramolecula	EEF1A1	EEF1A1			1200-1300	1500-1600
Human-	Intramolecula	EEF1A1	EEF1A1			1200-1300	1600-1700
Human-	Intramolecula	EEF1A1	EEF1A1			1300-1400	1400-1500
Human-	Intramolecula	EEF1A1	EEF1A1			1300-1400	1500-1600
Human-	Intramolecula	EEF1A1	EEF1A1			1300-1400	1700-1800
Human-	Intramolecula	EEF1A1	EEF1A1			1400-1500	1600-1700
Human-	Intramolecula	EEF1A1	EEF1A1			1400-1500	1700-1800
Human-	Intramolecula	EEF1A1	EEF1A1			1500-1600	1600-1700
Human-	Intramolecula	EEF1A1	EEF1A1			1500-1600	1700-1800
Human-	Intramolecula	EEF1A1	EEF1A1			1600-1700	1700-1800
Human-	Intramolecula	EEF1A1	EEF1A1			200-300	1000-1100
Human-	Intramolecula	EEF1A1	EEF1A1			200-300	1100-1200
Human-	Intramolecula	EEF1A1	EEF1A1			200-300	1200-1300
Human-	Intramolecula	EEF1A1	EEF1A1			200-300	1600-1700
Human-	Intramolecula	EEF1A1	EEF1A1			200-300	300-400
Human-	Intramolecula	EEF1A1	EEF1A1			200-300	400-500
Human-	Intramolecula	EEF1A1	EEF1A1			200-300	500-600
Human-	Intramolecula	EEF1A1	EEF1A1			200-300	900-1000
Human-	Intramolecula	EEF1A1	EEF1A1			300-400	1000-1100
Human-	Intramolecula	EEF1A1	EEF1A1			300-400	1100-1200
Human-	Intramolecula	EEF1A1	EEF1A1			300-400	1200-1300
Human-	Intramolecula	EEF1A1	EEF1A1			300-400	1400-1500
Human-	Intramolecula	EEF1A1	EEF1A1			300-400	1500-1600
Human-	Intramolecula	EEF1A1	EEF1A1			300-400	400-500
Human-	Intramolecula	EEF1A1	EEF1A1			300-400	600-700
Human-	Intramolecula	EEF1A1	EEF1A1			300-400	900-1000
Human-	Intramolecula	EEF1A1	EEF1A1			400-500	1000-1100
Human-	Intramolecula	EEF1A1	EEF1A1			400-500	1100-1200

Human-	Intramolecula	EEF1A1	EEF1A1			400-500	1200-1300
Human-	Intramolecula	EEF1A1	EEF1A1			400-500	1300-1400
Human-	Intramolecula	EEF1A1	EEF1A1			400-500	1400-1500
Human-	Intramolecula	EEF1A1	EEF1A1			400-500	500-600
Human-	Intramolecula	EEF1A1	EEF1A1			400-500	600-700
Human-	Intramolecula	EEF1A1	EEF1A1			400-500	900-1000
Human-	Intramolecula	EEF1A1	EEF1A1			500-600	1100-1200
Human-	Intramolecula	EEF1A1	EEF1A1			500-600	1200-1300
Human-	Intramolecula	EEF1A1	EEF1A1			500-600	600-700
Human-	Intramolecula	EEF1A1	EEF1A1			500-600	700-800
Human-	Intramolecula	EEF1A1	EEF1A1			500-600	800-900
Human-	Intramolecula	EEF1A1	EEF1A1			600-700	1100-1200
Human-	Intramolecula	EEF1A1	EEF1A1			600-700	1200-1300
Human-	Intramolecula	EEF1A1	EEF1A1			600-700	700-800
Human-	Intramolecula	EEF1A1	EEF1A1			600-700	800-900
Human-	Intramolecula	EEF1A1	EEF1A1			600-700	900-1000
Human-	Intramolecula	EEF1A1	EEF1A1			700-800	1000-1100
Human-	Intramolecula	EEF1A1	EEF1A1			700-800	1100-1200
Human-	Intramolecula	EEF1A1	EEF1A1			700-800	1200-1300
Human-	Intramolecula	EEF1A1	EEF1A1			700-800	800-900
Human-	Intramolecula	EEF1A1	EEF1A1			700-800	900-1000
Human-	Intramolecula	EEF1A1	EEF1A1			800-900	1000-1100
Human-	Intramolecula	EEF1A1	EEF1A1			800-900	1100-1200
Human-	Intramolecula	EEF1A1	EEF1A1			800-900	1200-1300
Human-	Intramolecula	EEF1A1	EEF1A1			800-900	900-1000
Human-	Intramolecula	EEF1A1	EEF1A1			900-1000	1000-1100
Human-	Intramolecula	EEF1A1	EEF1A1			900-1000	1100-1200
Human-	Intramolecula	EEF1A1	EEF1A1			900-1000	1200-1300
Human-	Intramolecula	EEF1A1	EEF1A1			900-1000	1300-1400
Human-	Intramolecula	EEF1B2	EEF1B2			0-100	300-400
Human-	Intramolecula	EEF1B2	EEF1B2			300-400	1000-1100
Human-	Intramolecula	EEF1B2	EEF1B2			300-400	900-1000
Human-	Intramolecula	EEF1B2	EEF1B2			500-600	800-900
Human-	Intramolecula	EEF1D	EEF1D			1600-1700	1700-1800
Human-	Intramolecula	EEF1D	EEF1D			200-300	2000-2100
Human-	Intramolecula	EEF1G	EEF1G			1000-1100	1200-1300
Human-	Intramolecula	EEF1G	EEF1G			100-200	300-400
Human-	Intramolecula	EEF1G	EEF1G			1100-1200	1200-1300
Human-	Intramolecula	EEF1G	EEF1G			1200-1300	1300-1400
Human-	Intramolecula	EEF1G	EEF1G			1200-1300	1400-1500
Human-	Intramolecula	EEF1G	EEF1G			300-400	1100-1200
Human-	Intramolecula	EEF1G	EEF1G			300-400	1400-1500
Human-	Intramolecula	EEF1G	EEF1G			300-400	600-700
Human-	Intramolecula	EEF1G	EEF1G			400-500	1200-1300
Human-	Intramolecula	EEF1G	EEF1G			400-500	1400-1500
Human-	Intramolecula	EEF1G	EEF1G			500-600	700-800
Human-	Intramolecula	EEF2	EEF2			1200-1300	2500-2600
Human-	Intramolecula	EEF2	EEF2			1200-1300	3000-3100
Human-	Intramolecula	EEF2	EEF2			1300-1400	1400-1500
Human-	Intramolecula	EEF2	EEF2			1400-1500	2800-2900
Human-	Intramolecula	EEF2	EEF2			200-300	400-500
Human-	Intramolecula	EEF2	EEF2			2700-2800	3000-3100
Human-	Intramolecula	EEF2	EEF2			2800-2900	3000-3100
Human-	Intramolecula	EIF2B2	EIF2B2			1300-1400	1400-1500
Human-	Intramolecula	EIF4A2	EIF4A2			1200-1300	1300-1400
Human-	Intramolecula	EIF4A2	EIF4A2			400-500	500-600
Human-	Intramolecula	EIF4B	EIF4B			2800-2900	3200-3300
Human-	Intramolecula	EIF4B	EIF4B			3000-3100	3100-3200
Human-	Intramolecula	EIF4E	EIF4E			1800-1900	2000-2100

Human-	Intramolecula	EIF4E	EIF4E			1900-2000	2000-2100
Human-	Intramolecula	EIF4H	EIF4H			1200-1300	2200-2300
Human-	Intramolecula	ELP5	ELP5			1100-1200	1800-1900
Human-	Intramolecula	ENO1	ENO1			1500-1600	1600-1700
Human-	Intramolecula	ENO1	ENO1			2000-2100	2200-2300
Human-	Intramolecula	ENO1	ENO1			2300-2400	2400-2500
Human-	Intramolecula	FAM195B	FAM195B			400-500	700-800
Human-	Intramolecula	FTH1	FTH1			700-800	800-900
Human-	Intramolecula	FTL	FTL			100-200	300-400
Human-	Intramolecula	FTL	FTL			500-600	700-800
Human-	Intramolecula	FXR1	FXR1			200-300	600-700
Human-	Intramolecula	FXYD5	FXYD5			0-100	100-200
Human-	Intramolecula	GAPDH	GAPDH			100-200	200-300
Human-	Intramolecula	GAPDH	GAPDH			1100-1200	1300-1400
Human-	Intramolecula	GAPDH	GAPDH			1200-1300	1400-1500
Human-	Intramolecula	GAPDH	GAPDH			300-400	400-500
Human-	Intramolecula	GAPDH	GAPDH			500-600	1100-1200
Human-	Intramolecula	GAPDH	GAPDH			500-600	1200-1300
Human-	Intramolecula	GAPDH	GAPDH			700-800	1000-1100
Human-	Intramolecula	GLUL	GLUL			2700-2800	2900-3000
Human-	Intramolecula	GLUL	GLUL			500-600	900-1000
Human-	Intramolecula	GMPR2	GMPR2			500-600	700-800
Human-	Intramolecula	GNB2L1	GNB2L1			300-400	400-500
Human-	Intramolecula	GNB2L1	GNB2L1			500-600	1000-1100
Human-	Intramolecula	GPX1	GPX1			300-400	600-700
Human-	Intramolecula	H3F3B	H3F3B			1100-1200	1300-1400
Human-	Intramolecula	HMGA1	HMGA1			0-100	300-400
Human-	Intramolecula	HMGA1	HMGA1			0-100	400-500
Human-	Intramolecula	HMGB1	HMGB1			1700-1800	2100-2200
Human-	Intramolecula	HMG2	HMG2			600-700	800-900
Human-	Intramolecula	HN1L	HN1L			2800-2900	2900-3000
Human-	Intramolecula	HNRNPA1	HNRNPA1			1600-1700	1800-1900
Human-	Intramolecula	HNRNPA2B1	HNRNPA2B1			2500-2600	3200-3300
Human-	Intramolecula	HNRNPC	HNRNPC			1000-1100	1300-1400
Human-	Intramolecula	HNRNPD	HNRNPD			1100-1200	1300-1400
Human-	Intramolecula	HNRNPK	HNRNPK			2300-2400	2600-2700
Human-	Intramolecula	HNRNPU	HNRNPU			3100-3200	3300-3400
Human-	Intramolecula	HSD17B4	HSD17B4			100-200	300-400
Human-	Intramolecula	HSD17B4	HSD17B4			300-400	400-500
Human-	Intramolecula		hsnma-RNU1-1hsnma-RNU1-1			0-100	100-200
Human-	Intramolecula	HSP90AA1	HSP90AA1			3000-3100	3500-3600
Human-	Intramolecula	HSPA8	HSPA8			2200-2300	2300-2400
Human-	Intramolecula		HSU13369-5ETHSU13369-5ETS			1800-1900	2700-2800
Human-	Intramolecula		HSU13369-ITS2HSU13369-ITS2			200-300	700-800
Human-	Intramolecula		HSU13369-ITS2HSU13369-ITS2			300-400	700-800
Human-	Intramolecula		HSU13369-ITS2HSU13369-ITS2			400-500	700-800
Human-	Intramolecula	IDH3B	IDH3B			1200-1300	1400-1500
Human-	Intramolecula	INTS6	INTS6			1000-1100	1300-1400
Human-	Intramolecula	IP6K2	IP6K2			0-100	300-400
Human-	Intramolecula	ISCU	ISCU			1500-1600	1600-1700
Human-	Intramolecula	LDHA	LDHA			1400-1500	1900-2000
Human-	Intramolecula	LDHB	LDHB			0-100	300-400
Human-	Intramolecula	LDHB	LDHB			1300-1400	1400-1500
Human-	Intramolecula		LRRC75A-AS1LRRC75A-AS1			0-100	300-400
Human-	Intramolecula		LRRC75A-AS1LRRC75A-AS1			1000-1100	1100-1200
Human-	Intramolecula		LRRC75A-AS1LRRC75A-AS1			1000-1100	1200-1300
Human-	Intramolecula		LRRC75A-AS1LRRC75A-AS1			300-400	600-700
Human-	Intramolecula		LRRC75A-AS1LRRC75A-AS1			400-500	600-700
Human-	Intramolecula	MCM4	MCM4			3800-3900	4300-4400

Human-	Intramolecula	METTL17	METTL17			1400-1500	1500-1600
Human-	Intramolecula	MINOS1	MINOS1			200-300	500-600
Human-	Intramolecula	MORF4L1	MORF4L1			300-400	500-600
Human-	Intramolecula	MRFAP1	MRFAP1			1300-1400	2000-2100
Human-	Intramolecula	MRFAP1	MRFAP1			1300-1400	2100-2200
Human-	Intramolecula	MRFAP1	MRFAP1			600-700	700-800
Human-	Intramolecula	MRPL11	MRPL11			500-600	600-700
Human-	Intramolecula	MTCH1	MTCH1			1200-1300	1300-1400
Human-	Intramolecula	MYL12A	MYL12A			300-400	400-500
Human-	Intramolecula	MYL6	MYL6			500-600	600-700
Human-	Intramolecula	NAP1L1	NAP1L1			1600-1700	2300-2400
Human-	Intramolecula	NAP1L1	NAP1L1			1600-1700	2400-2500
Human-	Intramolecula	NDUFS2	NDUFS2			1500-1600	1800-1900
Human-	Intramolecula	NIPA2	NIPA2			200-300	700-800
Human-	Intramolecula	NONO	NONO			2200-2300	2700-2800
Human-	Intramolecula	NONO	NONO			500-600	700-800
Human-	Intramolecula	OAZ1	OAZ1			700-800	800-900
Human-	Intramolecula	PAICS	PAICS			2000-2100	2700-2800
Human-	Intramolecula	PAICS	PAICS			2100-2200	2700-2800
Human-	Intramolecula	PDK1	PDK1			500-600	600-700
Human-	Intramolecula	PFN1	PFN1			900-1000	1100-1200
Human-	Intramolecula	PGAM1	PGAM1			1100-1200	1200-1300
Human-	Intramolecula	PGK1	PGK1			1600-1700	1700-1800
Human-	Intramolecula	PHPT1	PHPT1			800-900	900-1000
Human-	Intramolecula	PKM	PKM			200-300	600-700
Human-	Intramolecula	PKM	PKM			2200-2300	2600-2700
Human-	Intramolecula	PKM	PKM			2300-2400	2600-2700
Human-	Intramolecula	POLR2F	POLR2F			100-200	300-400
Human-	Intramolecula	PPIA	PPIA			100-200	300-400
Human-	Intramolecula	PPIA	PPIA			600-700	800-900
Human-	Intramolecula	PPP1CC	PPP1CC			400-500	500-600
Human-	Intramolecula	PRDX1	PRDX1			0-100	300-400
Human-	Intramolecula	PRMT1	PRMT1			100-200	200-300
Human-	Intramolecula	PSMB5	PSMB5			700-800	800-900
Human-	Intramolecula	PSMD13	PSMD13			300-400	400-500
Human-	Intramolecula	PTGES3	PTGES3			1100-1200	1500-1600
Human-	Intramolecula	PTMA	PTMA			600-700	900-1000
Human-	Intramolecula	RAC1	RAC1			400-500	500-600
Human-	Intramolecula	RAN	RAN			600-700	900-1000
Human-	Intramolecula	RAP1B	RAP1B			1700-1800	1900-2000
Human-	Intramolecula	RHOA	RHOA			1200-1300	1300-1400
Human-	Intramolecula	RPA2	RPA2			400-500	600-700
Human-	Intramolecula	RPL10A	RPL10A			0-100	200-300
Human-	Intramolecula	RPL10	RPL10			500-600	700-800
Human-	Intramolecula	RPL10	RPL10			600-700	800-900
Human-	Intramolecula	RPL11	RPL11			300-400	500-600
Human-	Intramolecula	RPL11	RPL11			300-400	600-700
Human-	Intramolecula	RPL11	RPL11			400-500	500-600
Human-	Intramolecula	RPL12	RPL12			400-500	500-600
Human-	Intramolecula	RPL13A	RPL13A			0-100	300-400
Human-	Intramolecula	RPL13A	RPL13A			1000-1100	1100-1200
Human-	Intramolecula	RPL13A	RPL13A			900-1000	1100-1200
Human-	Intramolecula	RPL13	RPL13			200-300	400-500
Human-	Intramolecula	RPL13	RPL13			200-300	700-800
Human-	Intramolecula	RPL13	RPL13			200-300	800-900
Human-	Intramolecula	RPL13	RPL13			300-400	400-500
Human-	Intramolecula	RPL13	RPL13			300-400	500-600
Human-	Intramolecula	RPL14	RPL14			0-100	100-200
Human-	Intramolecula	RPL14	RPL14			100-200	300-400

Human-	Intramolecula	RPL15	RPL15			500-600	600-700
Human-	Intramolecula	RPL15	RPL15			500-600	700-800
Human-	Intramolecula	RPL18A	RPL18A			0-100	100-200
Human-	Intramolecula	RPL18A	RPL18A			0-100	200-300
Human-	Intramolecula	RPL18A	RPL18A			0-100	400-500
Human-	Intramolecula	RPL18A	RPL18A			100-200	300-400
Human-	Intramolecula	RPL18A	RPL18A			200-300	300-400
Human-	Intramolecula	RPL18A	RPL18A			200-300	400-500
Human-	Intramolecula	RPL18A	RPL18A			200-300	500-600
Human-	Intramolecula	RPL18A	RPL18A			300-400	500-600
Human-	Intramolecula	RPL18	RPL18			400-500	500-600
Human-	Intramolecula	RPL18	RPL18			500-600	700-800
Human-	Intramolecula	RPL19	RPL19			0-100	500-600
Human-	Intramolecula	RPL19	RPL19			400-500	600-700
Human-	Intramolecula	RPL19	RPL19			400-500	700-800
Human-	Intramolecula	RPL24	RPL24			0-100	400-500
Human-	Intramolecula	RPL24	RPL24			0-100	500-600
Human-	Intramolecula	RPL26	RPL26			100-200	300-400
Human-	Intramolecula	RPL27A	RPL27A			500-600	700-800
Human-	Intramolecula	RPL27A	RPL27A			500-600	800-900
Human-	Intramolecula	RPL27A	RPL27A			600-700	700-800
Human-	Intramolecula	RPL27	RPL27			0-100	100-200
Human-	Intramolecula	RPL27	RPL27			0-100	300-400
Human-	Intramolecula	RPL27	RPL27			300-400	400-500
Human-	Intramolecula	RPL28	RPL28			300-400	500-600
Human-	Intramolecula	RPL29	RPL29			0-100	300-400
Human-	Intramolecula	RPL29	RPL29			0-100	400-500
Human-	Intramolecula	RPL29	RPL29			0-100	500-600
Human-	Intramolecula	RPL29	RPL29			400-500	500-600
Human-	Intramolecula	RPL32	RPL32			200-300	300-400
Human-	Intramolecula	RPL34	RPL34			100-200	300-400
Human-	Intramolecula	RPL37A	RPL37A			0-100	300-400
Human-	Intramolecula	RPL37	RPL37			200-300	300-400
Human-	Intramolecula	RPL37	RPL37			200-300	400-500
Human-	Intramolecula	RPL38	RPL38			0-100	100-200
Human-	Intramolecula	RPL39	RPL39			0-100	300-400
Human-	Intramolecula	RPL39	RPL39			200-300	300-400
Human-	Intramolecula	RPL39	RPL39			200-300	400-500
Human-	Intramolecula	RPL3	RPL3			1000-1100	1100-1200
Human-	Intramolecula	RPL3	RPL3			1000-1100	1200-1300
Human-	Intramolecula	RPL3	RPL3			100-200	600-700
Human-	Intramolecula	RPL3	RPL3			200-300	500-600
Human-	Intramolecula	RPL3	RPL3			300-400	1000-1100
Human-	Intramolecula	RPL3	RPL3			300-400	500-600
Human-	Intramolecula	RPL3	RPL3			300-400	700-800
Human-	Intramolecula	RPL3	RPL3			600-700	1000-1100
Human-	Intramolecula	RPL3	RPL3			600-700	800-900
Human-	Intramolecula	RPL3	RPL3			900-1000	1000-1100
Human-	Intramolecula	RPL3	RPL3			900-1000	1200-1300
Human-	Intramolecula	RPL41	RPL41			0-100	200-300
Human-	Intramolecula	RPL41	RPL41			200-300	400-500
Human-	Intramolecula	RPL41	RPL41			200-300	500-600
Human-	Intramolecula	RPL41	RPL41			300-400	400-500
Human-	Intramolecula	RPL41	RPL41			300-400	500-600
Human-	Intramolecula	RPL4	RPL4			200-300	400-500
Human-	Intramolecula	RPL4	RPL4			300-400	400-500
Human-	Intramolecula	RPL4	RPL4			500-600	1300-1400
Human-	Intramolecula	RPL4	RPL4			600-700	1000-1100
Human-	Intramolecula	RPL4	RPL4			600-700	1300-1400

Human-	Intramolecula	RPL4	RPL4			600-700	800-900
Human-	Intramolecula	RPL4	RPL4			700-800	800-900
Human-	Intramolecula	RPL4	RPL4			700-800	900-1000
Human-	Intramolecula	RPL4	RPL4			800-900	1000-1100
Human-	Intramolecula	RPL4	RPL4			800-900	900-1000
Human-	Intramolecula	RPL5	RPL5			100-200	300-400
Human-	Intramolecula	RPL5	RPL5			300-400	400-500
Human-	Intramolecula	RPL5	RPL5			300-400	500-600
Human-	Intramolecula	RPL5	RPL5			500-600	900-1000
Human-	Intramolecula	RPL5	RPL5			600-700	900-1000
Human-	Intramolecula	RPL6	RPL6			500-600	600-700
Human-	Intramolecula	RPL6	RPL6			900-1000	1000-1100
Human-	Intramolecula	RPL7A	RPL7A			100-200	700-800
Human-	Intramolecula	RPL7A	RPL7A			400-500	500-600
Human-	Intramolecula	RPL7A	RPL7A			500-600	700-800
Human-	Intramolecula	RPL7	RPL7			100-200	500-600
Human-	Intramolecula	RPI 7	RPL7			200-300	300-400
Human-	Intramolecula	RPL7	RPL7			400-500	500-600
Human-	Intramolecula	RPL9	RPL9			200-300	500-600
Human-	Intramolecula	RPL9	RPL9			300-400	400-500
Human-	Intramolecula	RPL9	RPL9			500-600	600-700
Human-	Intramolecula	RPL9	RPL9			500-600	700-800
Human-	Intramolecula	RPLP0	RPLP0			1000-1100	1100-1200
Human-	Intramolecula	RPLP0	RPLP0			100-200	200-300
Human-	Intramolecula	RPLP0	RPLP0			300-400	1000-1100
Human-	Intramolecula	RPLP0	RPLP0			300-400	500-600
Human-	Intramolecula	RPLP0	RPLP0			300-400	800-900
Human-	Intramolecula	RPLP0	RPLP0			300-400	900-1000
Human-	Intramolecula	RPLP0	RPLP0			400-500	600-700
Human-	Intramolecula	RPLP0	RPLP0			400-500	700-800
Human-	Intramolecula	RPLP0	RPLP0			400-500	800-900
Human-	Intramolecula	RPLP0	RPLP0			500-600	700-800
Human-	Intramolecula	RPLP0	RPLP0			600-700	900-1000
Human-	Intramolecula	RPLP0	RPLP0			800-900	900-1000
Human-	Intramolecula	RPLP1	RPLP1			0-100	200-300
Human-	Intramolecula	RPLP1	RPLP1			100-200	200-300
Human-	Intramolecula	RPLP1	RPLP1			100-200	300-400
Human-	Intramolecula	RPLP1	RPLP1			100-200	400-500
Human-	Intramolecula	RPLP1	RPLP1			200-300	400-500
Human-	Intramolecula	RPLP2	RPLP2			100-200	200-300
Human-	Intramolecula	RPS11	RPS11			0-100	400-500
Human-	Intramolecula	RPS11	RPS11			0-100	500-600
Human-	Intramolecula	RPS12	RPS12			0-100	400-500
Human-	Intramolecula	RPS12	RPS12			100-200	300-400
Human-	Intramolecula	RPS12	RPS12			100-200	400-500
Human-	Intramolecula	RPS12	RPS12			200-300	300-400
Human-	Intramolecula	RPS12	RPS12			200-300	400-500
Human-	Intramolecula	RPS13	RPS13			0-100	200-300
Human-	Intramolecula	RPS13	RPS13			200-300	400-500
Human-	Intramolecula	RPS13	RPS13			200-300	500-600
Human-	Intramolecula	RPS13	RPS13			300-400	400-500
Human-	Intramolecula	RPS14	RPS14			300-400	400-500
Human-	Intramolecula	RPS14	RPS14			300-400	500-600
Human-	Intramolecula	RPS15A	RPS15A			0-100	100-200
Human-	Intramolecula	RPS15A	RPS15A			0-100	300-400
Human-	Intramolecula	RPS15A	RPS15A			100-200	200-300
Human-	Intramolecula	RPS15A	RPS15A			200-300	400-500
Human-	Intramolecula	RPS16	RPS16			0-100	100-200
Human-	Intramolecula	RPS16	RPS16			0-100	200-300

Human-	Intramolecula	RPS16	RPS16			0-100	400-500
Human-	Intramolecula	RPS16	RPS16			0-100	500-600
Human-	Intramolecula	RPS16	RPS16			200-300	300-400
Human-	Intramolecula	RPS16	RPS16			200-300	500-600
Human-	Intramolecula	RPS17	RPS17			200-300	500-600
Human-	Intramolecula	RPS17	RPS17			300-400	500-600
Human-	Intramolecula	RPS19	RPS19			400-500	600-700
Human-	Intramolecula	RPS19	RPS19			400-500	700-800
Human-	Intramolecula	RPS19	RPS19			500-600	600-700
Human-	Intramolecula	RPS20	RPS20			100-200	200-300
Human-	Intramolecula	RPS20	RPS20			300-400	400-500
Human-	Intramolecula	RPS21	RPS21			200-300	300-400
Human-	Intramolecula	RPS23	RPS23			200-300	400-500
Human-	Intramolecula	RPS24	RPS24			200-300	300-400
Human-	Intramolecula	RPS25	RPS25			0-100	100-200
Human-	Intramolecula	RPS25	RPS25			0-100	200-300
Human-	Intramolecula	RPS27A	RPS27A			500-600	600-700
Human-	Intramolecula	RPS27A	RPS27A			600-700	700-800
Human-	Intramolecula	RPS27	RPS27			0-100	100-200
Human-	Intramolecula	RPS28	RPS28			0-100	200-300
Human-	Intramolecula	RPS29	RPS29			0-100	100-200
Human-	Intramolecula	RPS29	RPS29			0-100	200-300
Human-	Intramolecula	RPS2	RPS2			0-100	600-700
Human-	Intramolecula	RPS2	RPS2			0-100	800-900
Human-	Intramolecula	RPS2	RPS2			100-200	600-700
Human-	Intramolecula	RPS2	RPS2			200-300	600-700
Human-	Intramolecula	RPS2	RPS2			200-300	700-800
Human-	Intramolecula	RPS2	RPS2			300-400	500-600
Human-	Intramolecula	RPS2	RPS2			300-400	600-700
Human-	Intramolecula	RPS2	RPS2			300-400	700-800
Human-	Intramolecula	RPS2	RPS2			400-500	500-600
Human-	Intramolecula	RPS2	RPS2			600-700	700-800
Human-	Intramolecula	RPS2	RPS2			600-700	800-900
Human-	Intramolecula	RPS2	RPS2			700-800	800-900
Human-	Intramolecula	RPS2	RPS2			700-800	900-1000
Human-	Intramolecula	RPS3A	RPS3A			400-500	1500-1600
Human-	Intramolecula	RPS3	RPS3			0-100	900-1000
Human-	Intramolecula	RPS3	RPS3			100-200	800-900
Human-	Intramolecula	RPS3	RPS3			200-300	300-400
Human-	Intramolecula	RPS3	RPS3			200-300	400-500
Human-	Intramolecula	RPS3	RPS3			300-400	700-800
Human-	Intramolecula	RPS3	RPS3			400-500	600-700
Human-	Intramolecula	RPS3	RPS3			400-500	800-900
Human-	Intramolecula	RPS3	RPS3			500-600	600-700
Human-	Intramolecula	RPS4X	RPS4X			100-200	700-800
Human-	Intramolecula	RPS4X	RPS4X			200-300	600-700
Human-	Intramolecula	RPS5	RPS5			400-500	500-600
Human-	Intramolecula	RPS6	RPS6			0-100	100-200
Human-	Intramolecula	RPS6	RPS6			0-100	700-800
Human-	Intramolecula	RPS6	RPS6			100-200	500-600
Human-	Intramolecula	RPS6	RPS6			100-200	600-700
Human-	Intramolecula	RPS6	RPS6			200-300	400-500
Human-	Intramolecula	RPS6	RPS6			200-300	500-600
Human-	Intramolecula	RPS7	RPS7			200-300	300-400
Human-	Intramolecula	RPS7	RPS7			200-300	400-500
Human-	Intramolecula	RPS8	RPS8			0-100	500-600
Human-	Intramolecula	RPS8	RPS8			200-300	300-400
Human-	Intramolecula	RPS8	RPS8			200-300	600-700
Human-	Intramolecula	RPS8	RPS8			300-400	500-600

Human-	Intramolecula	RPS8	RPS8			300-400	600-700
Human-	Intramolecula	RPS9	RPS9			0-100	600-700
Human-	Intramolecula	RPS9	RPS9			400-500	600-700
Human-	Intramolecula	RSL24D1	RSL24D1			700-800	900-1000
Human-	Intramolecula	SDHC	SDHC			0-100	200-300
Human-	Intramolecula	SDHD	SDHD			300-400	500-600
Human-	Intramolecula	SEC11A	SEC11A			400-500	500-600
Human-	Intramolecula	SEC11A	SEC11A			400-500	600-700
Human-	Intramolecula	SEC61G	SEC61G			300-400	400-500
Human-	Intramolecula	SEPT7	SEPT7			1500-1600	1800-1900
Human-	Intramolecula	SKP1	SKP1			600-700	1300-1400
Human-	Intramolecula	SLC25A3	SLC25A3			0-100	300-400
Human-	Intramolecula	SLC25A3	SLC25A3			500-600	600-700
Human-	Intramolecula	SLC25A6	SLC25A6			500-600	700-800
Human-	Intramolecula	SLC41A3	SLC41A3			1500-1600	2100-2200
Human-	Intramolecula	SNHG16	SNHG16			0-100	200-300
Human-	Intramolecula	SNRPB	SNRPB			800-900	1000-1100
Human-	Intramolecula	SNRPD2	SNRPD2			200-300	400-500
Human-	Intramolecula	SNX3	SNX3			800-900	1100-1200
Human-	Intramolecula	SNX5	SNX5			0-100	300-400
Human-	Intramolecula	SRSF3	SRSF3			800-900	1300-1400
Human-	Intramolecula	STUB1	STUB1			500-600	600-700
Human-	Intramolecula	TCP1	TCP1			900-1000	1100-1200
Human-	Intramolecula	TKT	TKT			400-500	500-600
Human-	Intramolecula	TMBIM6	TMBIM6			1700-1800	2100-2200
Human-	Intramolecula	TMEM147	TMEM147			100-200	300-400
Human-	Intramolecula	TMEM147	TMEM147			200-300	300-400
Human-	Intramolecula	TMEM9	TMEM9			100-200	600-700
Human-	Intramolecula	TMPO	TMPO			1600-1700	1800-1900
Human-	Intramolecula	TMSB4X	TMSB4X			200-300	400-500
Human-	Intramolecula	TMSB4X	TMSB4X			200-300	500-600
Human-	Intramolecula	TMSB4X	TMSB4X			300-400	500-600
Human-	Intramolecula	TOMM20	TOMM20			400-500	500-600
Human-	Intramolecula	TOMM7	TOMM7			0-100	100-200
Human-	Intramolecula	TPI1	TPI1			1200-1300	1300-1400
Human-	Intramolecula	TPT1	TPT1			300-400	1000-1100
Human-	Intramolecula	TPT1	TPT1			400-500	1000-1100
Human-	Intramolecula	TPT1	TPT1			500-600	900-1000
Human-	Intramolecula	TPT1	TPT1			600-700	900-1000
Human-	Intramolecula	TRAPPC5	TRAPPC5			0-100	200-300
Human-	Intramolecula	TRMT112	TRMT112			800-900	900-1000
Human-	Intramolecula	TSFM	TSFM			500-600	600-700
Human-	Intramolecula	UBA52	UBA52			100-200	400-500
Human-	Intramolecula	UBA52	UBA52			200-300	400-500
Human-	Intramolecula	UBB	UBB			300-400	1100-1200
Human-	Intramolecula	UBE2D3	UBE2D3			1400-1500	1600-1700
Human-	Intramolecula	UBL5	UBL5			0-100	100-200
Human-	Intramolecula	UBXN1	UBXN1			1000-1100	1200-1300
Human-	Intramolecula	UFM1	UFM1			300-400	1000-1100
Human-	Intramolecula	USMG5	USMG5			0-100	400-500
Human-	Intramolecula	VDAC2	VDAC2			400-500	600-700
Human-	Intramolecula	VKORC1	VKORC1			500-600	600-700
Human-	Intramolecula	VPS11	VPS11			200-300	400-500
Human-	Intramolecula	YBX1	YBX1			1300-1400	1400-1500
Human-	Intramolecula	YBX1	YBX1			300-400	1000-1100
Human-	Intramolecula	YIF1B	YIF1B			800-900	900-1000
Human-	Intramolecula	YWHAQ	YWHAQ			1000-1100	1800-1900
Human-	Intramolecula	YWHAQ	YWHAQ			1100-1200	1800-1900
Human-	Intramolecula	ZFAND6	ZFAND6			200-300	400-500

Human-	Intramolecula	ZFAS1	ZFAS1			700-800	800-900
Human-	Intramolecula	ZFAS1	ZFAS1			700-800	900-1000
Human-	Intramolecula	ZNF207	ZNF207			900-1000	1000-1100
Human-	Intramolecula	ZNHIT1	ZNHIT1			800-900	900-1000

Table 4. List of lymphoblastoid cells snoRNA target sites

SnoRNA	Start position	End position	Target RNA	Start position	End position	Read count	Notes
ACA13	0	100	human-4V6X-18S	1100	1200	4.25	
ACA22	0	100	human-4V6X-28S	0	100	2.25	
ACA40	0	100	human-4V6X-18S	1100	1200	2.25	
ACA40	0	100	human-4V6X-28S	4500	4600	3.5	
ACA51	0	100	human-4V6X-28S	4600	4700	2.25	
ACA6	0	100	human-4V6X-18S	0	100	4	
HBI-43	0	100	human-4V6X-18S	0	100	2.5	
HBI-43	0	100	human-4V6X-28S	3800	3900	2	
HBII-55	0	100	human-4V6X-18S	1200	1300	2.25	
hTR	400	500	human-4V6X-28S	2200	2300	9	
hTR	200	300	hsnrna-RNU1-1	0	100	5	
mgU12-22/U4-8	200	300	human-4V6X-28S	2200	2300	2	
SNORA1	0	100	human-4V6X-18S	0	100	25.25	
SNORA1	0	100	human-4V6X-18S	1300	1400	6	
SNORA1	0	100	human-4V6X-28S	4500	4600	2.5	
SNORA10	0	100	hsnrna-RNU1-1	0	100	2	
SNORA21	0	100	human-4V6X-28S	4400	4500	2.75	
SNORA28	0	100	human-4V6X-5S	0	100	4.25	
SNORA32	0	100	human-4V6X-5S	0	100	15.25	
SNORA32	0	100	human-4V6X-5S	0	100	4	
SNORA33	100	200	hsnrna-RNU6-1	0	100	3	
SNORA44	0	100	hsnrna-RNU1-1	0	100	2	
SNORA45A	0	100	human-4V6X-18S	0	100	2.25	
SNORA45A	0	100	human-4V6X-18S	400	500	2.25	
SNORA45A	0	100	human-4V6X-18S	800	900	2.25	
SNORA45A	0	100	human-4V6X-18S	1300	1400	6.5	
SNORA45A	100	200	human-4V6X-18S	1300	1400	2	
SNORA45A	100	200	human-4V6X-18S	1400	1500	2	
SNORA45A	0	100	human-4V6X-18S	1400	1500	2	
SNORA45A	0	100	human-4V6X-28S	2400	2500	2.25	
SNORA45A	0	100	human-4V6X-28S	3800	3900	2.5	
SNORA45B	0	100	human-4V6X-18S	1300	1400	4.75	
SNORA58	0	100	human-4V6X-28S	3000	3100	3	
SNORA63	0	100	human-4V6X-28S	4300	4400	3.5	
SNORA63	0	100	human-4V6X-28S	4500	4600	2	
SNORA81	100	200	hsnrna-RNU2-1	0	100	2	
SNORD104	0	100	human-4V6X-28S	1300	1400	3	
SNORD111B	0	100	human-4V6X-28S	3900	4000	3	
SNORD119	0	100	human-4V6X-18S	0	100	2.5	
SNORD11B	0	100	hsnrna-RNU2-1	0	100	3	
SNORD124	0	100	hsnrna-RNU4ATAC	0	100	2	
SNORD12B	0	100	human-4V6X-28S	2900	3000	2	
SNORD15A	100	200	human-4V6X-28S	4600	4700	5.25	
SNORD15A	100	200	human-4V6X-28S	4700	4800	5	
SNORD15B	0	100	human-4V6X-18S	900	1000	2	
SNORD20	0	100	human-4V6X-18S	1700	1800	2	

SNORD21	0	100	human-4V6X-18S	1500	1600	2	
SNORD21	0	100	hsnrna-RNU1-1	100	200	3	
SNORD24	0	100	human-4V6X-28S	2300	2400	2.25	
SNORD24	0	100	human-4V6X-28S	2300	2400	4	
SNORD25	0	100	human-4V6X-18S	1400	1500	2.75	
SNORD25	0	100	human-4V6X-18S	1500	1600	2.75	
SNORD26	0	100	human-4V6X-28S	400	500	5.75	
SNORD26	0	100	human-4V6X-28S	400	500	5	
SNORD26	0	100	human-4V6X-28S	300	400	2	
SNORD26	0	100	human-4V6X-28S	400	500	3	
SNORD27	0	100	human-4V6X-18S	0	100	13	
SNORD27	0	100	human-4V6X-18S	0	100	7	
SNORD28	0	100	human-4V6X-18S	1300	1400	3.75	
SNORD28	0	100	human-4V6X-18S	1400	1500	2	
SNORD32A	0	100	human-4V6X-28S	1500	1600	4	
SNORD32A	0	100	human-4V6X-28S	1500	1600	15	
SNORD45A	0	100	human-4V6X-18S	100	200	16.75	
SNORD45A	0	100	human-4V6X-18S	100	200	7	
SNORD45A	0	100	human-4V6X-18S	100	200	2	
SNORD49A	0	100	human-4V6X-28S	4400	4500	5	
SNORD50A	0	100	human-4V6X-28S	2800	2900	2	
SNORD68	0	100	human-4V6X-28S	2300	2400	2	
SNORD68	0	100	human-4V6X-28S	2700	2800	2	
SNORD76	0	100	human-4V6X-18S	0	100	5.75	
SNORD76	0	100	human-4V6X-28S	2300	2400	4.75	
SNORD83A	0	100	human-4V6X-18S	500	600	2	
SNORD83A	0	100	U16	0	100	2	
SNORD87	0	100	human-4V6X-28S	3700	3800	2.25	
SNORD87	0	100	human-4V6X-28S	3700	3800	3	
SNORD87	0	100	human-4V6X-28S	3700	3800	2	
SNORD91A	0	100	human-4V6X-28S	4600	4700	2	
SNORD91B	0	100	human-4V6X-28S	4600	4700	2.75	
SNORD91B	0	100	human-4V6X-28S	1500	1600	2	
SNORD91B	0	100	human-4V6X-28S	4600	4700	2	
SNORD99	0	100	human-4V6X-28S	2700	2800	2	
SNORD99	0	100	human-4V6X-28S	2800	2900	2	
snR38C	0	100	human-4V6X-28S	5000	5100	2	
U13	0	100	human-4V6X-18S	0	100	2	
U13	0	100	human-4V6X-18S	1700	1800	9	
U13	0	100	human-4V6X-18S	1800	1900	13	
U13	0	100	human-4V6X-28S	4400	4500	12	
U13	0	100	human-4V6X-28S	4500	4600	5.25	
U13	0	100	human-4V6X-18S	0	100	11	
U13	0	100	human-4V6X-18S	100	200	9	
U13	0	100	human-4V6X-18S	400	500	9	
U13	0	100	human-4V6X-18S	600	700	2	
U13	0	100	human-4V6X-18S	700	800	4	
U13	0	100	human-4V6X-18S	900	1000	7	
U13	0	100	human-4V6X-18S	1100	1200	12	
U13	0	100	human-4V6X-18S	1200	1300	5	

U13	0	100	human-4V6X-18S	1300	1400	6	
U13	0	100	human-4V6X-18S	1400	1500	2	
U13	0	100	human-4V6X-18S	1500	1600	5	
U13	0	100	human-4V6X-18S	1600	1700	2	
U13	0	100	human-4V6X-18S	1700	1800	9	
U13	0	100	human-4V6X-18S	1800	1900	16	
U13	0	100	human-4V6X-28S	0	100	7	
U13	0	100	human-4V6X-28S	100	200	2	
U13	0	100	human-4V6X-28S	300	400	4	
U13	0	100	human-4V6X-28S	400	500	14	
U13	0	100	human-4V6X-28S	1300	1400	2	
U13	0	100	human-4V6X-28S	1600	1700	4	
U13	0	100	human-4V6X-28S	1900	2000	2	
U13	0	100	human-4V6X-28S	2400	2500	3	
U13	0	100	human-4V6X-28S	2700	2800	3	
U13	0	100	human-4V6X-28S	2800	2900	8	
U13	0	100	human-4V6X-28S	2900	3000	2	
U13	0	100	human-4V6X-28S	3600	3700	2	
U13	0	100	human-4V6X-28S	3700	3800	3	
U13	0	100	human-4V6X-28S	3800	3900	3	
U13	0	100	human-4V6X-28S	4400	4500	76	
U13	0	100	human-4V6X-28S	4500	4600	10	
U13	0	100	human-4V6X-5.8S	0	100	3	
U13	0	100	U19-2	0	100	2	
U13	0	100	hsnrna-RNU1-1	0	100	88	
U13	0	100	hsnrna-RNU1-1	100	200	17	
U13	0	100	hsnrna-RNU12	100	200	2	
U13	0	100	hsnrna-RNU2-1	0	100	5	
U13	0	100	hsnrna-RNU2-1	100	200	4	
U13	0	100	hsnrna-RNU4-1	0	100	21	
U13	0	100	hsnrna-RNU6-1	0	100	2	
U14A	0	100	human-4V6X-18S	0	100	12.25	
U14A	0	100	human-4V6X-18S	100	200	3.75	
U14A	0	100	human-4V6X-18S	400	500	8.25	
U14A	0	100	human-4V6X-18S	0	100	7	
U14A	0	100	human-4V6X-18S	400	500	5	
U14A	0	100	human-4V6X-18S	500	600	2	
U14B	0	100	human-4V6X-18S	400	500	4	
U14B	0	100	human-4V6X-18S	1300	1400	3.75	
U14B	0	100	human-4V6X-28S	4400	4500	2.25	
U14B	0	100	human-4V6X-18S	400	500	2	
U16	0	100	human-4V6X-18S	500	600	2	
U16	0	100	human-4V6X-18S	400	500	2	
U17a	0	100	human-4V6X-18S	100	200	2.25	
U17a	100	200	human-4V6X-18S	500	600	4.5	Homolog, snR30, found in yeast by SPLASH
U17a	100	200	human-4V6X-18S	1500	1600	2.25	
U17a	100	200	human-4V6X-28S	2800	2900	2.5	

U17a	100	200	human-4V6X-28S	4600	4700	3	Homolog, snR30, found in yeast by SPLASH
U17b	100	200	human-4V6X-18S	400	500	2.5	Homolog, snR30, found in yeast by SPLASH
U17b	100	200	human-4V6X-18S	500	600	4.5	Homolog, snR30, found in yeast by SPLASH
U17b	0	100	human-4V6X-18S	900	1000	2.5	
U17b	100	200	human-4V6X-28S	2300	2400	2.5	
U17b	100	200	human-4V6X-28S	4600	4700	3	Homolog, snR30, found in yeast by SPLASH
U17b	100	200	hsnrna-RNU1-1	0	100	2	
U19-2	0	100	hsnrna-RNU1-1	0	100	2	
U3	0	100	human-4V6X-18S	100	200	12	
U3	0	100	human-4V6X-18S	1300	1400	6	
U3	0	100	human-4V6X-28S	3700	3800	3	
U3	0	100	HSU13369-5ETS	400	500	2	
U3	0	100	hsnrna-RNU1-1	0	100	15	
U3	0	100	hsnrna-RNU1-1	100	200	5	
U31	0	100	human-4V6X-28S	2800	2900	2.25	
U31	0	100	human-4V6X-28S	4100	4200	12.25	
U31	0	100	human-4V6X-28S	4200	4300	17.5	
U31	0	100	human-4V6X-5.8S	0	100	4	
U31	0	100	human-4V6X-5.8S	100	200	4	
U31	0	100	human-4V6X-28S	4200	4300	2	
U3-2	0	100	human-4V6X-18S	1300	1400	8	
U3-2	0	100	human-4V6X-28S	2800	2900	2	
U3-2	0	100	hsnrna-RNU1-1	0	100	3	
U33	0	100	human-4V6X-18S	1300	1400	8.25	
U33	0	100	human-4V6X-18S	1300	1400	10	
U34	0	100	human-4V6X-28S	2800	2900	5.25	
U35A	0	100	human-4V6X-28S	4500	4600	3.25	
U35A	0	100	human-4V6X-28S	2700	2800	2	
U35A	0	100	human-4V6X-28S	4500	4600	3	
U37	0	100	human-4V6X-28S	3700	3800	2.5	
U42B	0	100	human-4V6X-18S	100	200	3	
U45B	0	100	human-4V6X-18S	100	200	2	
U45C	0	100	human-4V6X-18S	100	200	7	
U45C	0	100	human-4V6X-18S	100	200	4	
U45C	0	100	human-4V6X-18S	100	200	2	
U54	0	100	human-4V6X-18S	600	700	2	
U55	0	100	human-4V6X-18S	400	500	2	
U55	0	100	human-4V6X-28S	500	600	2	

U55	0	100	human-4V6X-28S	1400	1500	2	
U55	0	100	human-4V6X-28S	1700	1800	4	
U55	0	100	human-4V6X-28S	4400	4500	4	
U55	0	100	HSU13369-5ETS	2200	2300	2	
U55	0	100	human-4V6X-28S	1400	1500	3	
U55	0	100	human-4V6X-28S	1500	1600	3	
U55	0	100	hsnrna-RNU1-1	0	100	6	
U57	0	100	human-4V6X-18S	0	100	4.5	
U57	0	100	human-4V6X-18S	0	100	4	
U57	0	100	human-4V6X-18S	100	200	4	
U60	0	100	human-4V6X-28S	4300	4400	2	
U60	0	100	human-4V6X-28S	4300	4400	3	
U61	0	100	human-4V6X-18S	1400	1500	2	
U61	0	100	human-4V6X-28S	1500	1600	2	
U64	0	100	human-4V6X-28S	2400	2500	2.5	
U74	0	100	human-4V6X-28S	3800	3900	2.75	
U74	0	100	human-4V6X-28S	3800	3900	6	
U74	0	100	human-4V6X-28S	3800	3900	3	
U80	0	100	human-4V6X-28S	1600	1700	15.25	
U80	0	100	human-4V6X-28S	1600	1700	13	
U80	0	100	human-4V6X-28S	1600	1700	3	
U81	0	100	human-4V6X-28S	300	400	5.5	
U81	0	100	human-4V6X-28S	400	500	9	
U81	0	100	human-4V6X-28S	300	400	7	
U81	0	100	human-4V6X-28S	400	500	6	
U83B	0	100	human-4V6X-18S	400	500	3.25	
U83B	0	100	human-4V6X-28S	2800	2900	3	
U94	0	100	hsnrna-RNU1-1	100	200	2	
U96a	0	100	human-4V6X-5.8S	0	100	3	
U99	0	100	human-4V6X-18S	700	800	3.25	

Table 5. List of yeast snoRNA target sites

SnoRNA	Start position	End position	Target RNA	Start position	End position	Read count	Notes
snR11	0	100	RDN25-2	2900	3000	2	
snR128	0	100	RDN18-1	0	100	95.5	
snR128	0	100	RDN18-1	100	200	33	Known homolog, U14, found in human
snR128	0	100	RDN25-2	1000	1100	2.5	
snR17a	200	300	RDN18-1	400	500	2.5	
snR17a	200	300	RDN18-1	500	600	4	
snR17a	100	200	RDN18-1	1000	1100	2.5	
snR17a	100	200	RDN25-2	1200	1300	7	
snR17a	200	300	RDN25-2	2800	2900	2.5	
snR17a	300	400	RDN25-2	2900	3000	4.5	
snR18	0	100	RDN18-1	800	900	2	
snR18	0	100	RDN18-1	1000	1100	5.5	
snR18	0	100	RDN25-2	600	700	8.5	
snR189	100	200	RDN25-2	1700	1800	23.5	
snR189	100	200	RDN25-2	2800	2900	3	
snR24	0	100	RDN18-1	500	600	4	
snR24	0	100	RDN25-2	1300	1400	4	
snR24	0	100	RDN25-2	1400	1500	575	
snR24	0	100	RDN25-2	3000	3100	2	
snR30	500	600	RDN18-1	400	500	2.5	Homolog, U17, found in human by SPLASH
snR30	300	400	RDN18-1	700	800	5	
snR30	400	500	RDN18-1	1000	1100	7.5	
snR30	500	600	RDN25-2	1100	1200	3	
snR30	400	500	RDN25-2	2900	3000	3	Homolog, U17, found in human by SPLASH
snR30	0	100	RDN25-2	3000	3100	2.5	Homolog, U17, found in human by SPLASH
snR31	0	100	RDN25-2	1100	1200	4.5	
snR32	0	100	RDN25-2	1500	1600	2.5	
snR32	0	100	RDN25-2	2900	3000	3	
snR34	0	100	RDN18-1	800	900	2.5	
snR34	100	200	RDN25-2	2800	2900	6.5	
snR36	0	100	RDN18-1	400	500	3.5	
snR37	0	100	RDN25-2	2900	3000	28	
snR37	0	100	RDN25-2	3000	3100	4.5	
snR38	0	100	RDN18-1	800	900	2	
snR38	0	100	RDN25-2	1300	1400	2.5	
snR38	0	100	RDN25-2	2700	2800	8	

snR38	0	100	RDN25-2	2800	2900	48	
snR38	0	100	RDN25-2	2900	3000	3	
snR39	0	100	RDN25-2	900	1000	3.5	Known homolog, SNORD32A, found in human
snR39B	0	100	RDN18-1	500	600	2.5	
snR39B	0	100	RDN25-2	1200	1300	2.5	
snR39B	0	100	RDN25-2	1300	1400	3.5	
snR39B	0	100	RDN25-2	1700	1800	10	
snR4	100	200	RDN18-1	400	500	6.5	
snR4	0	100	RDN25-2	1000	1100	4.5	
snR4	0	100	RDN25-2	1600	1700	2.5	
snR4	0	100	RDN25-2	1800	1900	3.5	
snR40	0	100	RDN18-1	500	600	15.5	
snR40	0	100	RDN18-1	700	800	4	
snR40	0	100	RDN18-1	800	900	2	
snR40	0	100	RDN18-1	1200	1300	19	
snR40	0	100	RDN25-2	900	1000	10	
snR40	0	100	RDN25-2	2800	2900	4	
snR40	0	100	RDN25-2	3100	3200	2.5	
snR40	0	100	RDN25-2	3200	3300	11	
snR41	0	100	RDN18-1	500	600	15	
snR41	0	100	RDN18-1	1100	1200	65.5	
snR41	0	100	RDN25-2	1800	1900	2	
snR45	100	200	RDN25-2	3100	3200	5.5	
snR47	0	100	RDN18-1	600	700	9	
snR47	0	100	RDN18-1	800	900	2.5	
snR48	0	100	RDN25-2	2700	2800	70	
snR48	0	100	RDN25-2	2800	2900	101	
snR52	0	100	RDN18-1	300	400	9	
snR52	0	100	RDN18-1	400	500	323.5	
snR52	0	100	RDN18-1	500	600	2.5	
snR52	0	100	RDN18-1	800	900	4.5	
snR52	0	100	RDN25-2	1800	1900	5.5	
snR52	0	100	RDN25-2	2800	2900	5.5	
snR52	0	100	RDN25-2	2900	3000	6.5	
snR53	0	100	RDN18-1	700	800	4	
snR54	0	100	RDN18-1	900	1000	6	
snR55	0	100	RDN18-1	1200	1300	199	
snR55	0	100	RDN18-1	1300	1400	2	
snR59	0	100	RDN25-2	1800	1900	14.5	
snR60	0	100	RDN25-2	900	1000	57.5	
snR61	0	100	Q0158	800	900	2.5	
snR61	0	100	RDN25-2	1100	1200	13.5	
snR61	0	100	RDN25-2	1600	1700	4.5	
snR61	0	100	RDN25-2	2800	2900	5	
snR61	0	100	RDN25-2	2900	3000	2	

snR62	0	100	RDN25-2	1800	1900	19.5	
snR62	0	100	RDN25-2	1900	2000	7	
snR69	0	100	RDN25-2	2900	3000	36.5	
snR69	0	100	RDN25-2	3200	3300	2	
snR71	0	100	RDN25-2	1600	1700	4	
snR71	0	100	RDN25-2	2900	3000	16.5	
snR74	0	100	RDN18-1	0	100	11	
snR75	0	100	Q0158	3200	3300	3.5	
snR75	0	100	RDN18-1	400	500	2.5	
snR75	0	100	RDN18-1	500	600	3	
snR77	0	100	RDN18-1	500	600	86	
snR77	0	100	RDN18-1	600	700	5.5	
snR77	0	100	RDN25-2	900	1000	15	
snR77	0	100	RDN25-2	1300	1400	8	
snR79	0	100	RDN18-1	900	1000	103.5	
snR79	0	100	RDN18-1	1000	1100	367.5	
snR80	0	100	RDN18-1	500	600	2.5	
snR80	0	100	RDN25-2	2900	3000	2.5	
snR80	0	100	RDN25-2	3000	3100	7.5	
snR81	100	200	RDN25-2	1900	2000	2	
snR81	0	100	RDN25-2	2900	3000	4.5	
snR83	0	100	RDN18-1	1200	1300	4	
snR86	600	700	RDN18-1	400	500	9	
snR86	600	700	RDN18-1	500	600	2.5	
Y-NME1	200	300	RDN25-2	1500	1600	3.5	

Table 6. Go term analysis of network interactions in lymphoblastoid, ES and RA cells

Cell type	ModID	GO ID	Term	Annotated	Significant	Expected	P-value	GO Type	EnrichFold
Lymphoblastoid	central cluster	00:0006414	translational elongation	69	69	62.15	5.50E-05	BP	1.11
Lymphoblastoid	central cluster	00:0006413	translational initiation	66	66	59.44	1.00E-04	BP	1.11
Lymphoblastoid	central cluster	00:0006415	translational termination	66	66	59.44	1.00E-04	BP	1.11
Lymphoblastoid	central cluster	00:0006614	SRP-dependent cotranslational protein targeting to membrane	66	66	59.44	1.00E-04	BP	1.11
Lymphoblastoid	central cluster	00:0000184	nuclear-transcribed mRNA catabolic process, nonsense-mediated decay	65	65	58.54	0.00012	BP	1.11
Lymphoblastoid	central cluster	00:0032991	macromolecular complex	107	106	95.49	0.00032	CC	1.11
Lymphoblastoid	central cluster	00:0070082	extracellular vesicular exosome	91	88	81.21	0.00049	CC	1.08
Lymphoblastoid	central cluster	00:0016020	membrane	102	97	91.03	0.00203	CC	1.07
Lymphoblastoid	central cluster	00:0005829	cytosol	99	96	88.35	0.00404	CC	1.09
Lymphoblastoid	central cluster	00:0022825	cytosolic large ribosomal subunit	41	41	36.59	0.00436	CC	1.12
Lymphoblastoid	central cluster	00:0003735	structural constituent of ribosome	66	66	58.13	1.30E-05	MF	1.14
Lymphoblastoid	central cluster	00:0044822	poly(A) RNA binding	79	76	59.58	0.0012	MF	1.09
Lymphoblastoid	central cluster	00:0005515	protein binding	102	93	89.84	0.0794	MF	1.04
Lymphoblastoid	central cluster	00:0003723	RNA binding	89	86	78.39	0.0805	MF	1.10
Lymphoblastoid	central cluster	00:0033218	amide binding	14	14	12.33	0.155	MF	1.14
Lymphoblastoid	1	00:0019843	rRNA binding	9	5	1.49	0.0067	MF	3.36
Lymphoblastoid	1	00:0030168	platelet activation	8	4	1.38	0.03	BP	2.90
Lymphoblastoid	1	00:0002576	platelet degranulation	5	3	0.86	0.036	BP	3.49
Lymphoblastoid	1	00:0006887	exocytosis	5	3	0.86	0.036	BP	3.49
Lymphoblastoid	1	00:0016020	membrane	102	21	16.78	0.044	CC	1.25
Lymphoblastoid	2	00:1901575	organic substance catabolic process	88	17	12.38	0.022	BP	1.37
Lymphoblastoid	2	00:0009056	catabolic process	90	17	12.92	0.025	BP	1.36
Lymphoblastoid	2	00:0016052	carbohydrate catabolic process	10	4	1.39	0.034	BP	2.88
Lymphoblastoid	2	00:0044724	single-organism carbohydrate catabolic process	10	4	1.39	0.034	BP	2.88
Lymphoblastoid	2	00:0097285	cell-type specific apoptotic process	6	3	0.83	0.036	BP	3.61
Lymphoblastoid	3	00:0006614	SRP-dependent cotranslational protein targeting to membrane	66	23	13.11	5.10E-05	BP	1.75
Lymphoblastoid	3	00:0003735	structural constituent of ribosome	66	23	13.11	5.10E-05	MF	1.75
Lymphoblastoid	3	00:0006414	translational elongation	66	23	13.71	0.00014	BP	1.68
Lymphoblastoid	3	00:0000184	nuclear-transcribed mRNA catabolic process, nonsense-mediated decay	66	22	12.91	2.00E-04	BP	1.70
Lymphoblastoid	3	00:0019083	viral transcription	66	22	12.91	2.00E-04	BP	1.70
Lymphoblastoid	3	00:0006413	translational initiation	66	22	13.11	0.00027	BP	1.68
Lymphoblastoid	3	00:0022626	cytosolic ribosome	66	22	12.34	0.0031	CC	1.78
Lymphoblastoid	3	00:0044391	ribosomal subunit	66	22	12.53	0.004	CC	1.76
Lymphoblastoid	3	00:0022627	cytosolic small ribosomal subunit	24	10	4.56	0.0044	CC	2.19
Lymphoblastoid	3	00:0003723	RNA binding	89	24	17.68	0.0066	MF	1.36
Lymphoblastoid	3	00:0043232	intracellular non-membrane-bounded organelle	96	28	18.04	0.0111	CC	1.55
Lymphoblastoid	3	00:0043228	non-membrane-bounded organelle	96	28	18.04	0.0111	CC	1.55
Lymphoblastoid	4	00:0042605	peptide antigen binding	5	8	0.6	7.20E-11	MF	13.33
Lymphoblastoid	4	00:0071556	integral component of luminal side of endoplasmic reticulum membrane	10	8	0.63	2.50E-10	CC	12.70
Lymphoblastoid	4	00:0012507	ER to Golgi transport vesicle membrane	11	8	0.7	8.90E-10	CC	11.43
Lymphoblastoid	4	00:0000139	Golgi membrane	12	8	0.76	2.60E-09	CC	10.53
Lymphoblastoid	4	00:0060333	interferon-gamma-mediated signaling pathway	12	8	0.79	3.80E-09	BP	10.13
Lymphoblastoid	4	00:0016045	detection of bacterium	5	5	0.33	4.10E-07	BP	15.15
Lymphoblastoid	4	00:0031901	early endosome membrane	6	5	0.38	1.80E-06	CC	13.16
Lymphoblastoid	4	00:0042612	MHC class I protein complex	6	5	0.38	1.90E-06	CC	13.16

Lymphoblastoid	4	00.0001916	positive regulation of T cell mediated cytotoxicity	6	5	0.4	2.40E-06	BP	12.50
Lymphoblastoid	4	00.0002480	antigen processing and presentation of exogenous peptide antigen via MHC class I, TAP-independent	6	5	0.4	2.40E-06	BP	12.50
Lymphoblastoid	4	00.0002479	antigen processing and presentation of exogenous peptide antigen via MHC class I, TAP-dependent	7	5	0.46	8.20E-06	BP	10.87
Lymphoblastoid	4	00.0005102	receptor binding	9	4	0.6	0.0011	MF	6.67
Lymphoblastoid	4	00.0004872	receptor activity	5	2	0.33	0.0366	MF	6.06

Lymphoblastoid	5	00:0044429	mitochondrial part	9	3	0.51	0.0092	CC	5.88
Lymphoblastoid	5	00:0031975	envelope	10	3	0.57	0.0127	CC	5.26
Lymphoblastoid	5	00:0031967	crannelle envelope	10	3	0.57	0.0127	CC	5.26
Lymphoblastoid	5	00:0097193	intrinsic apoptotic signaling pathway	10	3	0.6	0.014	BP	5.00
Lymphoblastoid	5	00:0010035	response to inorganic substance	5	2	0.3	0.029	BP	6.67
Lymphoblastoid	5	00:0009991	muscle structure development	5	2	0.3	0.029	BP	6.67
Lymphoblastoid	5	00:0061061	response to extracellular stimulus	5	2	0.3	0.029	BP	6.67
Lymphoblastoid	5	00:0016491	oxidoreductase activity	5	2	0.3	0.029	BP	6.67
Lymphoblastoid	5	00:0097190	apoptotic signaling pathway	14	3	0.83	0.038	BP	3.81
Lymphoblastoid	5	00:0031966	mitochondrial membrane	6	2	0.34	0.0366	CC	5.88
Lymphoblastoid	5	00:0019866	organelle inner membrane	6	2	0.34	0.0366	CC	5.88
Lymphoblastoid	5	00:0022857	transmembrane transporter activity	6	2	0.36	0.042	MF	5.56
Lymphoblastoid	5	00:0015075	ion transmembrane transporter activity	6	2	0.36	0.042	MF	5.56
Lymphoblastoid	5	00:0022891	substrate-specific transmembrane transporter activity	6	2	0.36	0.042	MF	5.56
Lymphoblastoid	6	00:0010628	positive regulation of gene expression	5	4	0.53	0.0041	BP	7.55
Lymphoblastoid	6	00:001233	regulation of apoptotic signaling pathway	12	5	1.27	0.00357	BP	3.94
Lymphoblastoid	6	00:0045892	negative regulation of transcription, DNA-templated	8	4	0.85	0.00467	BP	4.71
Lymphoblastoid	6	00:0051347	positive regulation of transcription activity	5	3	0.53	0.00889	BP	5.66
Lymphoblastoid	6	00:0043410	positive regulation of MAPK cascade	5	3	0.53	0.00889	BP	5.66
Lymphoblastoid	7	00:0044430	cytoskeletal part	12	4	1.22	0.021	CC	3.28
Lymphoblastoid	7	00:0015630	microtubule cytoskeleton	7	3	0.71	0.023	CC	4.23
Lymphoblastoid	7	00:0044428	nuclear part	41	8	4.15	0.026	CC	1.93
Lymphoblastoid	8	00:0006812	cation transport	6	2	0.2	0.013	BP	10.00
Lymphoblastoid	8	00:0050801	ion homeostasis	6	2	0.2	0.013	BP	10.00
Lymphoblastoid	8	00:0055082	cellular chemical homeostasis	7	2	0.23	0.017	BP	8.70
Lymphoblastoid	8	00:0043169	metal ion binding	21	3	0.7	0.02	MF	4.29
Lymphoblastoid	8	00:0055887	cation binding	21	3	0.7	0.02	MF	4.29
Lymphoblastoid	8	00:0031226	integral component of plasma membrane	8	2	0.3	0.031	CC	6.67
Lymphoblastoid	8	00:0019725	intrinsic component of plasma membrane	8	2	0.3	0.031	CC	6.67
Lymphoblastoid	8	00:0048878	cellular homeostasis	10	2	0.33	0.036	BP	6.06
Lymphoblastoid	8	00:0044085	chemical homeostasis	10	2	0.33	0.036	BP	6.06
Lymphoblastoid	9	00:0005198	cellular component biogenesis	41	4	1.36	0.019	BP	2.94
Lymphoblastoid	9	00:0005198	structural molecule activity	72	5	2.38	0.023	MF	2.10
Lymphoblastoid	9	00:0061024	membrane organization	75	5	2.48	0.028	BP	2.02
Lymphoblastoid	9	00:0015935	small ribosomal subunit	25	3	0.79	0.028	CC	3.80
Lymphoblastoid	9	00:0016192	vesicle-mediated transport	25	2	0.33	0.036	BP	6.06
ES	central cluster	00:0006413	translational initiation	69	67	58.38	0.0021	BP	1.15
ES	central cluster	00:0006614	SRP-dependent cotranslational protein targeting to membrane	65	63	55	0.0045	BP	1.15
ES	central cluster	00:000184	nuclear-transcribed m RNA catabolic process, nonsense-mediated decay	64	62	54.15	0.0054	BP	1.14
ES	central cluster	00:0019083	viral transcription	63	61	53.31	0.0065	BP	1.14
ES	central cluster	00:0006415	translational termination	63	61	53.31	0.0065	BP	1.14
ES	central cluster	00:0070062	extracellular vesicular exosome	132	122	110.59	9.30E-05	CC	1.10
ES	central cluster	00:0022625	cytosolic large ribosomal subunit	37	37	31	0.0082	CC	1.19
ES	central cluster	00:0005829	cytosol	138	127	115.62	0.0039	CC	1.10
ES	central cluster	00:0005829	ribonucleoprotein complex	93	89	77.92	0.00536	CC	1.14
ES	central cluster	00:0005925	total adhesion	54	51	45.24	0.00973	CC	1.13
ES	central cluster	00:0044822	poly(A) RNA binding	121	114	102.08	1.70E-05	MF	1.12

ES	central cluster	00:0003735	structural constituent of ribosome	65	52	54.84	0.002	MF	1.13
ES	central cluster	00:1901265	nucleoside phosphate binding	56	52	47.24	0.031	MF	1.10
ES	central cluster	00:0000166	nucleotide binding	56	52	47.24	0.031	MF	1.10
ES	central cluster	00:0036094	small molecule binding	61	56	51.46	0.045	MF	1.09
ES	1	00:0006413	translational initiation	69	29	16.2	3.30E-05	BP	1.79
ES	1	00:0006614	SRP-dependent cotranslational protein targeting to membrane	65	27	15.26	0.00011	BP	1.77

ES	1	00:000184	nuclear-transcribed mRNA catabolic process, nonsense-mediated decay	64	26	15.03	0.00026	BP	1.73
ES	1	00:000614	translational elongation	68	27	15.97	0.00029	BP	1.69
ES	1	00:002262	cytosolic ribosome	62	25	14.6	0.00051	CC	1.71
ES	1	00:0019083	viral transcription	63	25	14.79	6.00E-04	BP	1.69
ES	1	00:0003735	structural constituent of ribosome	65	25	15.51	0.0015	MF	1.61
ES	1	00:0015635	small ribosomal subunit	27	12	6.36	0.00943	CC	1.89
ES	1	00:0044822	poly(A) RNA binding	121	37	28.88	0.0107	ME	1.28
ES	1	00:0044391	ribosomal subunit	65	26	15.31	0.01116	CC	1.70
ES	1	00:0005576	extracellular region	136	40	32.03	0.01382	CC	1.25
ES	1	00:0022627	cytosolic small ribosomal subunit	25	11	5.89	0.01446	CC	1.87
ES	1	00:0003723	RNA binding	136	40	32.46	0.0157	MF	1.23
ES	1	00:0003676	nucleic acid binding	164	43	36.76	0.0348	MF	1.17
ES	2	00:0006096	glycolytic process	9	5	1.28	0.0036	BP	3.91
ES	2	00:0046364	monosaccharide biosynthetic process	6	4	0.85	0.0042	BP	4.71
ES	2	00:0006952	defense response	27	9	3.83	0.006	BP	2.35
ES	2	00:0019318	hexose metabolic process	11	5	1.56	0.0106	BP	3.21
ES	2	00:0006008	glucose metabolic process	11	5	1.56	0.0108	BP	3.21
ES	2	00:0005615	extracellular space	16	6	2.16	0.012	CC	2.78
ES	2	00:0005737	cytoplasm	218	34	29.46	0.013	CC	1.15
ES	2	00:0044444	cytoplasmic part	188	31	25.41	0.014	CC	1.22
ES	2	00:0072562	blood microparticle	5	3	0.88	0.019	CC	4.41
ES	2	00:0023023	MHC protein complex binding	5	3	0.7	0.021	MF	4.29
ES	2	00:0023028	MHC class II protein complex binding	5	3	0.7	0.021	MF	4.29
ES	2	00:0016491	oxidoreductase activity	13	5	1.82	0.022	MF	2.75
ES	2	00:0030554	adenyl nucleotide binding	24	7	3.30	0.033	ME	2.08
ES	2	00:0005629	cytosol	138	24	18.55	0.037	CC	1.29
ES	2	00:0048037	cofactor binding	6	3	0.84	0.037	MF	3.57
ES	3	00:0005925	focal adhesion	54	10	4.17	0.024	CC	2.40
ES	3	00:0006339	chromatin remodeling	6	3	0.49	0.078	BP	6.12
ES	3	00:0010628	positive regulation of gene expression	18	5	1.46	0.009	BP	3.42
ES	3	00:0031968	mitochondrial membrane	14	4	1.08	0.0158	CC	3.70
ES	3	00:0008285	negative regulation of cell proliferation	8	3	0.65	0.0197	BP	4.62
ES	3	00:0005740	mitochondrial envelope	15	4	1.16	0.0204	CC	3.45
ES	3	00:0010557	positive regulation of macromolecule biosynthetic process	23	5	1.86	0.0289	BP	2.89
ES	3	00:0016020	membrane	149	16	11.51	0.0269	CC	1.39
ES	3	00:0045893	positive regulation of transcription, DNA-templated	16	4	1.3	0.0303	BP	3.08
ES	4	00:0042273	ribosomal large subunit biogenesis	5	2	0.28	0.027	BP	7.14
ES	4	00:0001880	placenta development	5	2	0.28	0.027	BP	7.14
ES	4	00:0006364	rRNA processing	13	3	0.74	0.03	BP	4.05
ES	4	00:0016072	rRNA metabolic process	13	3	0.74	0.03	BP	4.05
ES	4	00:0034470	ncRNA processing	14	3	0.79	0.037	BP	3.80
ES	4	00:0044822	poly(A) RNA binding	121	10	6.47	0.04	MF	1.55
ES	5	00:00033674	positive regulation of kinase activity	10	3	0.69	0.024	BP	4.35
ES	5	00:0045860	positive regulation of protein kinase activity	10	3	0.69	0.024	BP	4.35
ES	5	00:0020165	MAPK cascade	11	3	0.76	0.032	BP	3.95
ES	5	00:0023014	signal transduction by phosphorylation	11	3	0.76	0.032	BP	3.95
ES	5	00:0043408	regulation of MAPK cascade	11	3	0.76	0.032	BP	3.95

ES	5	00:0030017	sarcomere	5	2	0.39	0.049	CC	5.13
ES	5	00:0044449	contractile fiber part	5	2	0.39	0.049	CC	5.13
ES	6	00:0007600	sensory perception	7	4	0.91	0.0062	BP	4.40
ES	6	00:0043009	chordate embryonic development	13	5	1.68	0.0161	BP	2.98
ES	6	00:0009792	embryo development ending in birth or egg hatching	13	5	1.68	0.0161	BP	2.98
ES	6	00:0030031	cell projection assembly	5	3	0.65	0.0166	BP	4.62

ES	6	00:0048568	embryonic organ development	5	3	0.65	0.0166	BP	4.62
ES	6	00:0003697	single-stranded DNA binding	6	3	0.77	0.029	MF	3.90
ES	6	00:0031252	cell leading edge	7	3	0.69	0.046	CC	3.37
ES	7	00:0043234	protein complex	79	3	3.36	5.00E-04	CC	2.68
ES	7	00:0030001	metal ion transport	7	3	0.28	0.0016	BP	10.71
ES	7	00:0051049	regulation of transport	23	4	0.93	0.0083	BP	4.30
ES	7	00:0019904	protein domain specific binding	13	3	0.53	0.012	ME	5.00
ES	7	00:0012505	endomembrane system	38	5	1.61	0.0125	CC	3.11
ES	7	00:0005794	Golgi apparatus	13	3	0.55	0.013	CC	5.45
ES	7	00:0003013	circulatory system process	5	2	0.2	0.0139	BP	10.00
ES	7	00:0008015	blood circulation	5	2	0.2	0.0139	BP	10.00
ES	7	00:0006308	DNA catabolic process	5	2	0.2	0.0139	BP	10.00
ES	7	00:0005867	transcription factor complex	8	2	0.25	0.0225	CC	8.00
ES	7	00:0005515	protein binding	170	10	7	0.026	MF	1.43
ES	7	00:0005654	nucleoplasm	48	5	2.04	0.0341	CC	2.45
ES	7	00:0019999	enzyme binding	34	4	1.4	0.037	MF	2.86
ES	7	00:0008134	transcription factor binding	9	2	0.37	0.047	MF	5.41
ES	8	00:0009386	cell surface	10	3	0.23	0.00079	CC	13.04
ES	8	00:0015711	organic anion transport	6	2	0.15	0.0071	BP	13.33
ES	8	00:0040011	locomotion	27	3	0.66	0.0187	BP	4.55
ES	8	00:0035770	ribonucleoprotein granule	11	2	0.25	0.02247	CC	8.00
ES	8	00:0036464	cytoplasmic ribonucleoprotein granule	11	2	0.25	0.02247	CC	8.00
ES	8	00:0042330	axis	12	2	0.29	0.0292	BP	6.90
ES	8	00:0016935	chemotaxis	12	2	0.29	0.0292	BP	6.90
ES	8	00:005008	regulation of biological quality	58	4	1.43	0.0303	BP	2.60
ES	8	00:000054	cell junction	62	4	1.44	0.0361	CC	2.78
ES	8	00:0005615	extracellular space	16	2	0.37	0.04648	CC	5.41
ES	9	00:0003729	mRNA binding	9	2	0.33	0.038	ME	6.06
ES	10	00:0031124	mRNA 3'-end processing	6	2	0.15	0.0071	BP	13.33
ES	10	00:0006366	transcription from RNA polymerase II promoter	21	3	0.51	0.009	BP	5.88
ES	10	00:0003723	RNA binding	136	6	3.36	0.029	MF	1.79
ES	10	00:0008283	cell proliferation	32	3	0.78	0.0302	BP	3.85
ES	10	00:0008284	positive regulation of cell proliferation	13	2	0.32	0.0341	BP	6.25
ES	10	00:0006325	chromatin organization	15	2	0.36	0.0449	BP	5.56
ES-RA	central cluster	00:0006414	translational elongation	52	48	35.2	1.10E-06	BP	1.36
ES-RA	central cluster	00:0006413	translational initiation	55	50	37.23	2.10E-06	BP	1.34
ES-RA	central cluster	00:000184	nuclear-transcribed mRNA catabolic process, nonsense-mediated decay	50	46	33.84	3.00E-06	BP	1.36
ES-RA	central cluster	00:0006415	translational termination	50	46	33.84	3.00E-06	BP	1.36
ES-RA	central cluster	00:0019083	viral transcription	49	45	33.16	4.80E-06	BP	1.36
ES-RA	central cluster	00:0070062	extracellular vesicular exosome	79	66	52.37	8.20E-06	CC	1.26
ES-RA	central cluster	00:0005629	cytosol	79	70	52.37	0.00046	CC	1.34
ES-RA	central cluster	00:0022627	cytosolic small ribosomal subunit	20	19	13.28	0.00192	CC	1.43
ES-RA	central cluster	00:0022625	cytosolic large ribosomal subunit	29	26	19.22	0.00203	CC	1.35
ES-RA	central cluster	00:0005925	focal adhesion	37	32	24.53	0.00211	CC	1.30
ES-RA	central cluster	00:0044922	poly(A) RNA binding	75	65	49.54	1.50E-07	MF	1.31
ES-RA	central cluster	00:0003735	structural constituent of ribosome	50	45	33.02	6.30E-06	MF	1.36
ES-RA	central cluster	00:0005515	protein binding	104	79	58.69	0.00037	MF	1.15

ES-RA	central cluster	GO:0003723	RNA binding	85	73	56.14	0.03459	MF	1.30
ES-RA	central cluster	00:0019843	RNA binding	7	7	4.62	0.05118	MF	1.52
ES-RA	1	00:0070062	extracellular vesicular exosome	79	20	12.19	0.00999	CC	1.84
ES-RA	1	GO:0006414	translational elongation	52	16	8.56	0.0012	BP	1.87
ES-RA	1	00:0022626	cytosolic ribosome	49	14	7.66	0.00382	CC	1.85
ES-RA	1	00:0016071	miRNA metabolic process	62	10	10.21	0.0044	BP	1.86

ES-RA	1	00:0044391	ribosomal subunit	50	14	7.71	0.00481	CC	1.82
ES-RA	1	00:0044822	poly(A) RNA binding	75	19	12.5	0.0054	MF	1.52
ES-RA	1	00:0019083	viral transcription	49	14	8.07	0.0076	BP	1.73
ES-RA	1	00:0006614	SRP-dependent cotranslational protein targeting to membrane	49	14	8.07	0.0076	BP	1.73
ES-RA	1	00:0046907	intracellular transport	68	20	11.2	0.0077	BP	1.79
ES-RA	1	00:0003735	structural constituent of ribosome	50	14	8.33	0.0107	MF	1.88
ES-RA	1	00:0022625	cytosolic large ribosomal subunit	29	9	4.47	0.01599	CC	2.01
ES-RA	1	00:0015934	large ribosomal subunit	29	9	4.47	0.01599	CC	2.01
ES-RA	1	00:0005198	structural molecule activity	58	15	9.67	0.0181	MF	1.55
ES-RA	2	00:0043412	macromolecule modification	26	5	1.59	0.0039	BP	3.14
ES-RA	2	00:003924	catalytic activity	31	5	1.91	0.023	MF	2.62
ES-RA	2	00:0006464	cellular protein modification process	24	4	1.46	0.0408	BP	2.74
ES-RA	2	00:0036211	protein modification process	24	4	1.46	0.0408	BP	2.74
ES-RA	2	00:0045892	negative regulation of transcription, DNA-templated	14	3	0.85	0.0418	BP	3.53
ES-RA	2	00:002679	negative regulation of RNA biosynthetic process	14	3	0.85	0.0418	BP	3.53
ES-RA	3	00:0042060	wound healing	8	4	0.88	0.0055	BP	4.55
ES-RA	3	00:0042383	sarcolemma	5	3	0.51	0.0081	CC	5.86
ES-RA	3	00:0048646	anatomical structure formation involved in morphogenesis	9	4	0.99	0.0092	BP	4.04
ES-RA	3	00:0051146	striated muscle cell differentiation	5	3	0.55	0.0098	BP	5.45
ES-RA	3	00:0001101	response to acid chemical	5	3	0.55	0.0098	BP	5.45
ES-RA	3	00:0010035	response to inorganic substance	5	3	0.55	0.0098	BP	5.45
ES-RA	3	00:0065010	extracellular membrane-bounded organelle	79	13	8.13	0.0141	CC	1.60
ES-RA	3	00:0043230	extracellular organelle	79	13	8.13	0.0141	CC	1.60
ES-RA	3	00:0070062	extracellular vesicular exosome	79	13	8.13	0.0141	CC	1.60
ES-RA	3	00:0044421	extracellular region part	80	13	8.23	0.0161	CC	1.58
ES-RA	4	00:0044391	ribosomal subunit	50	14	4.57	0.0055	CC	2.41
ES-RA	4	00:0022626	cytosolic ribosome	49	10	4.48	0.00266	CC	2.23
ES-RA	4	00:0006413	translational initiation	55	11	5.37	0.0027	BP	2.05
ES-RA	4	00:0006614	SRP-dependent cotranslational protein targeting to membrane	49	10	4.78	0.0046	BP	2.09
ES-RA	4	00:0019083	viral transcription	49	10	4.78	0.0046	BP	2.09
ES-RA	4	00:0006415	translational termination	50	10	4.88	0.0055	BP	2.05
ES-RA	4	00:0000184	nuclear-transcribed mRNA catabolic process, nonsense-mediated decay	50	10	4.88	0.0055	BP	2.05
ES-RA	4	00:0003735	structural constituent of ribosome	50	10	4.94	0.0061	MF	2.02
ES-RA	4	00:0003723	RNA binding	85	13	8.4	0.0133	MF	1.55
ES-RA	4	00:0015935	small ribosomal subunit	21	5	1.92	0.02172	CC	2.60
ES-RA	4	00:0006829	cytosol	79	14	7.22	0.02827	CC	1.94
ES-RA	4	00:0030055	cell-substrate junction	37	7	3.38	0.02854	CC	2.07
ES-RA	4	00:003676	nucleic acid binding	104	14	10.27	0.0323	MF	1.36
ES-RA	5	00:0006413	translational initiation	55	8	2.68	0.00011	BP	2.99
ES-RA	5	00:0022627	cytosolic small ribosomal subunit	20	5	0.91	0.0053	CC	5.49
ES-RA	5	00:0019083	viral transcription	49	7	2.39	0.00095	BP	2.93
ES-RA	5	00:0006614	SRP-dependent cotranslational protein targeting to membrane	49	7	2.39	0.00095	BP	2.93
ES-RA	5	00:0000184	nuclear-transcribed mRNA catabolic process, nonsense-mediated decay	50	7	2.44	0.00109	BP	2.87
ES-RA	5	00:0006415	translational termination	50	7	2.44	0.00109	BP	2.87
ES-RA	5	00:0003735	structural constituent of ribosome	50	7	2.47	0.0012	MF	2.83
ES-RA	5	00:0044622	poly(A) RNA binding	75	8	3.7	0.0017	MF	2.16
ES-RA	5	00:0030529	ribonucleoprotein complex	68	8	3.11	0.02842	CC	2.57

ES-RA	5	00:0003729	mRNA binding	6	2	0.3	0.0291	MF	6.67
ES-RA	6	00:0003723	RNA binding	85	5	2.62	0.038	MF	1.91
ES-RA	8	00:0006006	glucose metabolic process	5	3	0.18	0.00027	BP	16.67
ES-RA	8	00:0016051	carbohydrate biosynthetic process	5	3	0.18	0.00027	BP	16.67
ES-RA	8	00:0006091	generation of precursor metabolites and energy	7	3	0.28	0.00092	BP	11.54
ES-RA	8	00:0005886	plasma membrane	28	4	0.96	0.0064	CC	4.17

ES-RA	8	00:0044712	single-organism catabolic process	13	3	0.48	0.00687	BP	5.25
ES-RA	8	00:0044281	small molecule metabolic process	27	4	0.99	0.0071	BP	4.04
ES-RA	8	00:0015629	actin cytoskeleton	5	2	0.17	0.0094	CC	11.76
ES-RA	8	00:0008092	cytoskeletal protein binding	12	2	0.37	0.045	ME	5.41
ES-RA	8	00:0003824	catalytic activity	31	3	0.96	0.049	MF	3.13
ES-RA	9	00:0019843	rRNA binding	7	2	0.26	0.022	MF	7.69
ES-RA	9	00:0048518	positive regulation of biological process	42	4	1.61	0.045	BP	2.48
ES-RA	10	00:0010608	posttranscriptional regulation of gene expression	17	3	0.52	0.0092	BP	5.77
ES-RA	10	00:0006446	regulation of translational initiation	6	2	0.18	0.0107	BP	11.11
ES-RA	10	00:0051248	negative regulation of protein metabolic process	10	2	0.3	0.0304	BP	6.67
ES-RA	10	00:0032269	negative regulation of cellular protein metabolic process	10	2	0.3	0.0304	BP	6.67
ES-RA	10	00:0051129	negative regulation of cellular component organization	10	2	0.3	0.0304	BP	6.67
ES-RA	10	00:0019901	protein kinase binding	11	2	0.34	0.038	MF	5.88
ES-RA	10	00:0019900	kinase binding	12	2	0.37	0.045	MF	5.41
ES-RA	11	00:0010557	positive regulation of macromolecule biosynthetic process	14	2	0.34	0.037	BP	5.88
ES-RA	12	00:0030001	metal ion transport	5	2	0.09	0.0022	BP	22.22
ES-RA	12	00:0006875	cellular metal ion homeostasis	6	2	0.11	0.0033	BP	18.18
ES-RA	12	00:0046872	metal ion binding	30	3	0.56	0.0058	MF	5.36
ES-RA	12	00:0019904	protein domain specific binding	9	2	0.17	0.008	MF	11.76
ES-RA	12	00:0012505	endomembrane system	25	3	0.57	0.0095	CC	5.25
ES-RA	12	00:0022892	substrate-specific transporter activity	10	2	0.19	0.01	MF	10.53
ES-RA	12	00:0005768	endosome	8	2	0.18	0.0105	CC	11.11
ES-RA	12	00:0007154	cell communication	40	3	0.73	0.0137	BP	4.11
ES-RA	12	00:0044700	single organism signaling	40	3	0.73	0.0137	BP	4.11
ES-RA	12	00:0023052	signaling	40	3	0.73	0.0137	BP	4.11
ES-RA	12	00:0005783	endoplasmic reticulum	10	2	0.23	0.0167	CC	8.70
ES-RA	12	00:0005215	transporter activity	13	2	0.24	0.0171	MF	8.33
ES-RA	12	00:0065010	extracellular membrane-bounded organelle	79	4	1.81	0.0398	CC	2.21
ES-RA	12	00:0043230	extracellular organelle	79	4	1.81	0.0398	CC	2.21

Table 7: Probes and qPCR primers used in validation

Type	Gene	Position	
Probe	Human 18S	R1	/5Biosg/ CTGGCAGGATCAACCAGGTA (SEQ ID NO: 3)
		R711	/5Biosg/GGGCGGTGGCTCGCCTCGCG (SEQ ID NO: 4)
		R1661	/5Biosg/TGACCCGCACTTACTGGGAA (SEQ ID NO: 5)
		R1868	/5Biosg/AATGATCCTTCCGCAGGTTCA (SEQ ID NO: 6)
Probe	Human 28S	R1	/5Biosg/ ACGTCTGATCTGAGGTCGCG (SEQ ID NO: 7)
		R1311	/5Biosg/TGGTCCGTGTTTCAAGACGGGT (SEQ ID NO: 8)
		R1737	/5Biosg/CAAGACCTCTAATCATTGCTT (SEQ ID NO: 9)
		R5058	/5Biosg/TGTCGAGGGCTGACTTTCAAT (SEQ ID NO: 10)
Probe	Human 5S	R58	/5Biosg/TGCTTAGCTTCCGAGATCAGA (SEQ ID NO: 11)
		R120	/5Biosg/AAGCCTACAGCACCCGGTATT (SEQ ID NO: 12)
Probe	ACA51	R13	/5Biosg/GTAAGAACACAGCCTGTGGTAAG (SEQ ID NO:13)
		R37	/5Biosg/ TCCTCTTCTATACAGTCAG (SEQ ID NO: 14)
		R60	/5Biosg/ ATATGGGGTAGGTTTACTCT (SEQ ID NO: 15)
Probe	TMSB4X	R52	/5Biosg/GAGGAAAAGCGAAGCGAGGC (SEQ ID NO: 16)
		R241	/5Biosg/GCGAATGCTTGTGGAATGTA (SEQ ID NO: 17)
		R302	/5Biosg/AACTTGATCCAACCTCTTTG (SEQ ID NO: 18)
Probe	EEF1A1	R2	/5Biosg/GGCAAACCCGTTGCGAAAAA (SEQ ID NO: 19)
		R29	/5Biosg/TAGTTTTACGACACCTGTG (SEQ ID NO: 20)
		R1547	/5Biosg/ACCACTGATTAAGAGTGGGG (SEQ ID NO: 21)
Probe	Actin	R435	/5Biosg/ACATGATCTGGGTCATCTTC (SEQ ID NO: 22)
		R488	/5Biosg/GGATAGCACAGCCTGGATAG (SEQ ID NO: 23)
		R745	/5Biosg/ATCTCTTGCTCGAAGTCCAG (SEQ ID NO: 24)
		R823	/5Biosg/TCATTGCCAATGGTGATGAC (SEQ ID NO: 25)
		R1067	/5Biosg/CTCAGGAGGAGCAATGATCT (SEQ ID NO: 26)
		R1400	/5Biosg/CACATTGTGAACTTTGGGGG (SEQ ID NO: 27)
		R1475	/5Biosg/GACTTCCTGTAACAACGCAT (SEQ ID NO: 28)
		R1761	/5Biosg/GTCTCAAGTCAGTGACAGG (SEQ ID NO: 29)
Probe	Yeast YBR118W	R1	/5Biosg/ACCCATGTTTAGTTAATTAT (SEQ ID NO: 30)
		R45	/5Biosg/TCGACATGACCGATAACGAC (SEQ ID NO: 31)
		R97	/5Biosg/ACCACCACACTTGTAATCA (SEQ ID NO: 32)
Probe	GFP Bio control 1		/5Biosg/CACGGATTATTTGCCTGATT (SEQ ID NO: 33)
Probe	GFP Bio control 2		/5Biosg/ATTTTGCGTAACCTATTCGC (SEQ ID NO: 34)
QPCR primer	Human 18S	F1443	TTAGAGGGACAAGTGGCGTT (SEQ ID NO: 35)
		R1513	GGACATCTAAGGGCATCACA (SEQ ID NO: 36)
QPCR primer	Human 28S	F2377	GAGAACTTTGAAGGCCGAAG (SEQ ID NO: 37)
		R2455	CATCTCTCAGGACCGACTGA (SEQ ID NO: 38)

QPCR primer	Human 5S	F25	GCGCCCGATCTCGTCTGATCTC (SEQ ID NO: 39)
		R77	CAGGCGGTCTCCCATCCAAGT (SEQ ID NO: 40)
QPCR primer	ACA51	F20	CAGGCTGTGTTCTTACACTGAC (SEQ ID NO: 41)
		R109	ATGTTCCCCCATTACAATACA (SEQ ID NO: 42)
QPCR primer	SNORA32	F2	GGTCATTACCAAGGCTTTTAG (SEQ ID NO: 43)
		R67	GCAGATAGAAAACCTACTGGG (SEQ ID NO: 44)
QPCR primer	SNORD83a	F19	TCAGAGTGAGCGCTGGGTACAG (SEQ ID NO: 45)
		R63	GGAAGGCAGTAGAGAATGGT (SEQ ID NO: 46)
QPCR primer	TMSB4X	F95	CGATATGGCTGAGATCGAGA (SEQ ID NO: 47)
		R158	CTTTGGAAGGCAGTGGATT (SEQ ID NO: 48)
QPCR primer	EEF1A1	F1010	CTGTCAAGGATGTTTCGTCGT (SEQ ID NO: 49)
		R1105	CTTATTTGGCCTGGATGGTT (SEQ ID NO: 50)
QPCR primer	RPS27	F42	TCGCAAAGGATCTCCTTCAT (SEQ ID NO: 51)
		R89	CCAGGCGTTTCTTCTTG (SEQ ID NO: 52)
QPCR primer	RPLP0	F865	ACTCTGCATTCTCGCTTCCT (SEQ ID NO: 53)
		R960	CTCGTTTGTACCCGTTGATG (SEQ ID NO: 54)
QPCR primer	GAPDH	F787	TGGTATCGTGGAAGGACTCA (SEQ ID NO: 55)
		R899	CCAGTAGAGGCAGGGATGAT (SEQ ID NO: 56)
QPCR primer	Actin	F266	AGAGAGGCATCCTCACCT (SEQ ID NO: 57)
		R353	CACACGCAGCTCATTGTAGA (SEQ ID NO: 58)
QPCR primer	RPL35	F89	GAAGGAGGAGCTGCTGAAAC (SEQ ID NO: 59)
		R174	TCGGATCTTAGAGAGCTTGA (SEQ ID NO: 60)
QPCR primer	B2M	F376	GACTTTGTCACAGCCCAAGA (SEQ ID NO: 61)
		R467	CAAGCAAGCAGAATTTGGAA (SEQ ID NO: 62)
QPCR primer	EIF5A	F1033	GAATCAGAAAGCGGTGGATT (SEQ ID NO: 63)
		R1079	ACCAGACCAGGGATGAGTG (SEQ ID NO: 64)
QPCR primer	Yeast YBR118W	F14	CATGGGTAAAGAGAAGTCTACA (SEQ ID NO: 65)
		R107	GGTTCTCTGTCAATACCACCA (SEQ ID NO: 66)
		F1276	GATTCGCTGTCAGAGACATGA (SEQ ID NO: 67)
		R1349	CAGCCTTGGTAACTTAGCG (SEQ ID NO: 68)
QPCR primer	OCT4	F1764	GAGAAGGATGTGGTCCGAGT (SEQ ID NO: 69)
		R1836	GTGCATAGTCGCTGCTTGAT (SEQ ID NO: 70)

QPCR primer	HMGA1	F609	GCTGGTAGGGAGTCAGAAGG (SEQ ID NO: 71)
		R739	TTGGTTTCCTTCCTGGAGTT (SEQ ID NO:72)
QPCR primer	GAPDH	F1056	TCAAGAAGGTGGTGAAGCAG (SEQ ID NO: 73)
		R1129	CGCTGTTGAAGTCAGAGGAG (SEQ ID NO: 74)

Claims

1. A method of analysing ribonucleic acid (RNA) interactions comprising:
 - a. cross-linking base-paired nucleotides of at least one RNA molecule and/or at least one pair of RNA molecules using a tagged, reversible cross-linking agent to produce at least one cross-linked RNA molecule and/or at least one pair of cross-linked RNA molecules;
 - b. fragmenting the said cross-linked RNA molecule and/or pair of cross-linked RNA molecules to produce a plurality of fragments comprising at least one cross-linked RNA fragment;
 - c. using said tag to extract said cross-linked RNA fragment(s) from said plurality of fragments;
 - d. ligating the said cross-linked RNA fragment(s) to produce cross-linked ligated RNA chimera(s);
 - e. reversing the cross-linking of the said agent to the said RNA molecule and/or pair of RNA molecules to produce a ligated RNA chimera molecule(s) and/or RNA chimera pair(s);
 - f. preparing a sequence library by sequencing the ligated RNA chimera molecule(s) or pair(s); and
 - g. analysing the sequence library to determine RNA interactions.
2. The method according to claim 1 wherein said RNA is present in a cell and said crosslinking using said tagged, reversible cross-linking agent involves the use of a cellular uptake agent, such as a detergent.
3. The method according to claims 1 or 2 wherein part c is undertaken before part b.
4. The method according to any one of claims 1-3 wherein said cross-linking agent comprises a furocoumarin compound.
5. The method according to claim 4 wherein the cross-linking agent comprises psoralen.

6. The method according to any one of claims 1-5 wherein said tag of said cross-linking agent comprises a first member of a binding pair selected from the group comprising: biotin/streptavidin, antigen/antibody, protein/protein, polypeptide/protein and polypeptide/polypeptide.
7. The method according to any one of claims 1-6 wherein the step of cross-linking said RNA molecule(s) with a cross-linking agent to produce cross-linked RNA molecule(s) is carried out using ultraviolet irradiation at wavelengths in the range of about 300nm to about 400nm.
8. The method according to any one of claims 1-7 wherein the step of reversing the cross-linking of the cross-linked ligated RNA molecule(s) is carried out using ultraviolet irradiation at wavelengths in the range of about 200nm to no more than about 300nm.
9. The method according to any one of claims 1-8 wherein the step of preparing a sequence library by sequencing the ligated RNA chimera molecule or pairs comprises the use of at least one or more of the following: adaptor ligation, reverse transcription, cDNA circularization or polymerase chain reaction (PCR).
10. The method according to any one of claims 1-9 wherein the step of fragmenting the cross-linked RNA molecule and/or pair of RNA molecules to produce a plurality of fragments comprises producing fragments having an average size in the range of 100 to 500 base pairs in length.
11. The method according to any one of claims 1-10 wherein the concentration of cross-linking agent used crosslinks at approximately one in every 150 bases.
12. The method according to any one of claims 1-11 wherein continuous pairwise interactions or those spaced apart by less than 50 bases are removed to focus the analysis on the long-range intramolecular and intermolecular interactions.
13. The method according to any one of claims 1-12 wherein said RNA molecule and/or at least one member of said pair of RNA molecules is ascribed a "circularization score" defined as the average base pair interaction distance within

each molecule, normalized by the length of said RNA molecule or the length of said member of said pair of RNA molecules.

14. The method according to claim 13 wherein said RNA molecule and/or said at least one member of said pair of RNA molecules are classified into groups according to their "circularization score".
15. The method according to any one of claims 2-13 wherein the cell is mammalian, human, bacterial or yeast.
16. A kit for analysing ribonucleic acid (RNA) interactions comprising:
 - a tagged, reversible cross-linking agent for reversibly cross-linking base paired nucleotides of at least one RNA molecule and/or at least one pair of RNA molecules to produce at least one cross-linked RNA molecule and/or at least one pair of cross-linked RNA molecules;
 - a fragmentation buffer for fragmenting the cross-linked RNA molecule and/or pair of cross-linked RNA molecules to produce a plurality of fragments;
 - an RNA ligase for ligating the cross-linked RNA fragment(s) to produce cross-linked ligated RNA chimeras;
 - a binding partner for said tag on said agent; and
 - optionally, instructions on how to use the kit.
17. The kit according to claim 16 further comprising reagents for sequencing the cross-linked ligated RNA chimeras to prepare a sequence library.
18. The kit according to claim 17 wherein the reagents for sequencing the ligated RNA chimeras comprises at least one of: a RNA ligase, reverse transcription primers and DNA polymerase.
19. The kit according to any one of claims 16-18 wherein the cross-linking agent comprises a furocoumarin compound.
20. The kit according to claim 19 wherein the cross-linking agent comprises psoralen.

21. The kit according to any one of claims 16-20 wherein said tag of said cross-linking agent comprises a first member of a binding pair selected from the group comprising or consisting of: biotin/streptavidin, antigen/antibody, protein/protein, polypeptide/protein and polypeptide/polypeptide.
22. The kit according to claim 21 wherein said binding partner is the other member of a binding pair selected from the group comprising or consisting of: biotin/streptavidin, antigen/antibody, protein/protein, polypeptide/protein and polypeptide/polypeptide.
23. The kit according to any one of claims 16-21 further comprising a detergent to facilitate cellular uptake of the cross-linking agent into a cell.
24. A method of studying a subject, the method comprising:
- obtaining a cell sample from a subject;
 - cross-linking base-paired nucleotides of at least one RNA molecule and/or at least one pair of RNA molecules using a tagged, reversible cross-linking agent to produce at least one cross-linked RNA molecule and/or at least one pair of cross-linked RNA molecules;
 - fragmenting the said cross-linked RNA molecule and/or pair of cross-linked RNA molecules to produce a plurality of fragments comprising at least one cross-linked RNA fragment;
 - using said tag to extract said cross-linked RNA fragment(s) from said plurality of fragments;
 - ligating the said cross-linked RNA fragment(s) to produce cross-linked ligated RNA chimera(s);
 - reversing the cross-linking of the said agent to the said RNA molecule and/or pair of RNA molecules to produce ligated a RNA chimera molecule(s) and/or RNA chimera pair(s);
 - preparing a sequence library by sequencing the ligated RNA chimera molecule(s) or pair(s);
 - analysing the sequence library to determine RNA interactions in the cell sample; and
 - comparing the determined RNA interactions with a set of pre-existing data to attribute a clinical picture to the subject.

25. The method of claim 23 wherein the method of studying a subject comprises at least one of: diagnosing the subject of a clinical condition, predicting the risk of the subject having a clinical condition, screening the subject for suitability for a particular treatment or determining the efficacy of a drug candidate on the subject.

26. A drug discovery method, the method comprising:

- a. exposing RNA to a drug;
- b. cross-linking base-paired nucleotides of at least one RNA molecule and/or at least one pair of RNA molecules using a tagged, reversible cross-linking agent to produce at least one cross-linked RNA molecule and/or at least one pair of cross-linked RNA molecules;
- c. fragmenting the said cross-linked RNA molecule and/or pair of cross-linked RNA molecules to produce a plurality of fragments comprising at least one cross-linked RNA fragment;
- d. using said tag to extract said cross-linked RNA fragment(s) from said plurality of fragments;
- e. ligating the said cross-linked RNA fragment(s) to produce cross-linked ligated RNA chimera(s);
- f. reversing the cross-linking of the said agent to the said RNA molecule and/or pair of RNA molecules to produce a ligated a RNA chimera molecule(s) and/or RNA chimera pair(s);
- g. preparing a sequence library by sequencing the ligated RNA chimera molecule(s) or pair(s);
- h. analysing the sequence library to determine RNA interactions; and
- i. attributing an efficacy score to the drug based on the determined RNA interactions.