

TREATMENT OF INFLAMMATORY BOWEL DISEASE**FIELD OF THE INVENTION**

- 5 The present invention relates to a method of treating inflammatory bowel disease and a means for carrying out the method.

BACKGROUND OF THE INVENTION

10

Permanent cures for inflammatory bowel disease (IBD), such as Crohn's disease (CD) and ulcerative colitis (UC), are lacking. In Sweden, 1400 new patients are diagnosed with IBD each year, with the peak incidence at 20 years of age, resulting in life
15 long treatment with cortisone and/or immunosuppressive drugs with severe side effects. Furthermore, IBD often results in surgical bowel resection and permanent stoma, which is a major handicap. Ulcerative colitis also increases the risk of cancer in the large bowel.

20

IBD is thought to be an autoimmune disease due to abnormalities in the immunological response to normally harmless mucosal antigens. Based on the cytokines involved the inflammation of CD is characterized as a T_H1 cell mediated
25 immune response. UC is proposed to have a T_H2 cell mediated response, though this has not been clearly shown (Fuss I J et al., 1996. J. Immunol. 157:1261-70; Mullin G E et al., 1996. Inflamm. Bowel Dis. 2:16-26).

30 Cytokines act as regulators of immunological responses as they stimulate differentiation and proliferation of the cellular elements involved. CD4+ helper T cells may differentiate into two major subpopulations of effector cells, T_H1 cells when stimulated by IL-12 and T_H2 cells when stimulated by IL-4. T_H1
35 cells secrete IFN- γ that stimulates phagocyte- and cytolytic T

cell dependent reactions against intracellular microbes such as viruses or intracellular bacteria. T_H2 cells secrete IL-4 and IL-5 that stimulate production of IgE and eosinophil/mast cell mediated defense against parasites. T_H2 also functions to
5 down regulate $TH1$ responses.

OBJECTS OF THE INVENTION

It is an object of the invention to provide a method that
10 cures inflammatory bowel disease or at least keeps it at bay over extended periods of time.

It is another object of the invention to provide a means for use in the method.

15

Further objects will become apparent from the following summary of the invention, a number of figures illustrating it, the description of preferred embodiments thereof as well from the appended claims.

20

SUMMARY OF THE INVENTION

The present invention is based on the insight that inflammatory bowel disease, in particular Crohn's disease
25 and/or ulcerative colitis, can be permanently cured or at least kept in a dormant or inactive state by the administration of regulatory T cells (Treg) collected from sentinel nodes of the patient and expanded *in vitro*. Thereby the inflammation of the bowel is reduced and may be even fully
30 suppressed without having to take recourse to medication such by corticosteroids.

The unresponsiveness of the immune system to orally administered antigens is defined as oral tolerance.
35 Apparently, this mechanism prevents improper immune reaction

against intestinal content and is believed to be mediated by at least two processes: first, the activation of regulatory T cells (Tregs, CD4⁺CD25⁺), which are members of the T helper cell family (CD4⁺), and secondly, the induction of clonal deletion or anergy of T cells. The first process, responsible for the tolerance to low-dose antigen exposure, presumably takes place in the local intestinal lymphoid system and is dependent on the production of the cytokine TGF- β . The second (deletional) process presumably takes place in systemic lymphoid tissue, triggered by filtered antigens, and is similar to the mechanism behind induced tolerance to intravenously administered antigens in the absence of immunological adjuvant. This latter tolerogenic mechanism seems to be more prominent after high-dose antigen exposure.

The activation of T cells is strictly regulated in order to maintain balance in the immune system by avoiding excessive and misdirected immunoresponse as seen in autoimmunity. In patients with Crohn's disease mesenterial lymph nodes are greatly enlarged as a sign of uncontrolled regulation of T cell activation.

As IBD is an inflammation the mesenterial lymph nodes adjacent to the inflammatory site are primarily responsible for the drainage of that site. The first nodes disposed along a direct route from the inflamed area are called sentinel nodes.

According to a first preferred aspect of the invention regulatory T cells in an activated state are collected from a patient's lymph nodes (sentinel nodes) draining bowel segments with IBD, expanded *in vitro*, and re-infused in the patient.

According to a second preferred aspect of the invention regulatory T cells in a non-activated state are collected from a patient's lymph nodes draining healthy bowel segments that

is, bowel segments not affected by IBD, expanded *in vitro*, and re-infused in the patient. The activated as well as the non-activated regulatory T cells so collected can be (further) activated *in vitro* by suitable agents such as cytokines.

5

According to a third preferred aspect of the invention regulatory T cells in an activated state or a non-activated state are collected from lymph nodes draining bowel segments with or without IBD, respectively, and expanded *in vitro*.

- 10 Expansion may be continuous or in intervals, such as in phases interrupted by resting phase(s). The expanded regulatory T cells are activated (stimulated) with cytokine(s) in combination with an antigen extract from an inflammatory bowel segment. Alternatively the regulatory T cells collected from
- 15 lymph nodes draining bowel segments with or without IBD are activated (stimulated) in such manner prior to expansion or between expansion phases. One cytokine, preferably two or more cytokines selected from IL-2, IL-7, IL-10, TGF- β , TSLP, such as a cytokine cocktail comprising IL-2, IL-7, IL-10, TGF- β , in
- 20 particular TGF- β 1, TSLP, are used in combination with the antigen extract from the inflammatory bowel segment to obtain Treg activation. It is also preferred to provide for additional Treg activation by use of anti-CD3, anti-CD28 and rapamycin. After expansion, the expanded and activated
- 25 regulatory T-cells are re-infused in the patient.

Treatment with autologous regulatory T cells has the advantage of reduced or even lacking side effects, and of high efficacy.

- 30 According to a first variant of the third preferred aspect, regulatory T cells in an activated state collected from lymph nodes draining bowel segments with or without IBD from a patient with Crohn's disease, and activated with an antigen extract from the inflammatory site in combination with a low
- 35 dose of the cytokine IL-2 for expansion, preferably in further

combination with antibodies against IFN- γ and/or TNF- α to suppress T_{H1} skewing and T_{H2} expansion. It is also preferred to condition the culture medium in intervals such as from every day to every fifth day, in particular every day for a
5 time period of from one week or more, in particular for a time period of from three to four weeks. Thereby a population of T_{H2} cells is obtained, which is used in autotransfusion to re-establish the T_{H1}/T_{H2} balance in a patient. Expansion can be supported by additional activation with a CD3 antibody. To
10 improve T_{H2} skewing an anti CD28 antibody can additionally be used in the expansion in combination with an anti IL-12 antibody.

According to a second variant of the third preferred aspect,
15 regulatory T cells in an activated state collected from lymph nodes draining bowel segments with or without IBD from a patient with ulcerative colitis disease, are activated with an antigen extract from the inflammatory site in combination with one or more of the cytokines IL-2, IL-12 and IFN- α for
20 expansion, preferably in further combination with anti IL4 and/or anti IL-10 antibodies. It is also preferred to condition the culture medium in intervals such as from every day to every fifth day, in particular every day for a time period of from one week or more, in particular for a time
25 period of from three to four weeks. Additional stimuli such as CD3 and CD28 may be provided, in particular towards the end of the expansion period.

According to a fourth preferred aspect of the invention is
30 provided one or more cytokines in a pharmaceutically acceptable carrier for in-vitro activation of regulatory T cells collected from a patient with IBD, in particular from sentinel nodes affected by IBD.

According to a fifth preferred aspect of the invention is provided a method of obtaining sentinel lymph nodes draining bowel segments with IBD comprising injecting a lymphatic staining agent such as patent blue into lesional areas of the intestine of a patient suffering from IBD, identifying lymphatic tissue stained by the agent, optionally marking the stained tissue by sutures, and resecting the stained tissue. The resected tissue is carefully investigated (inspected, such as by microscopy), whereupon the sentinel lymph nodes draining bowel segments with IBD are excised from the tissue.

According to a sixth preferred aspect of the invention is provided a method of isolating regulatory T cells from a sentinel lymph node draining a bowel segment with IBD, comprising exerting a pressure on the lymph node, collecting the cell suspension thereby expelled from the node, and isolating regulatory T cells from the suspension.

According to a seventh preferred aspect of the invention are provided regulatory T cells isolated from a sentinel lymph node draining a bowel segment with IBD.

Further preferred aspects of the invention are included in the appended claims.

The invention will now be explained in greater detail by reference to preferred but not limiting embodiments of the invention illustrated by a number of figures.

SHORT DESCRIPTION OF THE FIGURES

- Fig. 1 is a diagram showing activated Tregs in a Treg sample identified by the expression of two surface markers;
- Fig. 2 is a diagram showing a functional Treg assay;

- Fig. 3 shows an RT-PCR assay of mRNA extracted from expanded Tregs;
- Fig. 4 is a diagram showing the regulating capacity of a Treg population before and after expansion;
- 5 Fig. 5 is a diagram showing the activation of human immunological cells by an endogenous extract from an inflamed bowel section;
- Fig. 6 is a diagram showing the dose response of similarly activated Tregs derived from sentinel nodes;
- 10 Fig. 7 shows the expression of CCR5 as a marker for T_H2;
- Fig. 8 shows the secretion of IL-4 in Tregs from UC patients;
- Fig. 9 shows the secretion of IFN- γ in Tregs from CD patients.

15

DESCRIPTION OF PREFERRED EMBODIMENTS

EXAMPLE 1. Identification of sentinel nodes draining inflammatory bowel segments

20

The sentinel node technique has been used for a decade to establish the staging of malignant tumours, mainly malignant melanoma and breast cancer. The sentinel node is identified during surgery by depicting the lymph drainage from the cancer

25 using a lymphatic staining agent such as patent blue (CAS 129-17-9 and patent blue V (CAS 3536-49-0; often supplied as a calcium-chelated dimer), which is injected in the lesional areas of the intestine. The sentinel nodes so identified are marked by suture(s) or by other means. Then the intestinal

30 lesions together with unaffected marginal zones and mesentery including vessels and regional lymph nodes are resected in a conventional manner. The resected tissue is carefully investigated and the sentinel nodes identified and removed.

This method is applied to sentinel nodes draining both inflamed and unaffected intestine in patients with ulcerative colitis or Crohn's disease. In addition a venous blood sample is collected from the patient. The peripheral blood sample and
5 the nodes together with specimens from the inflammatory lesions as well as intestinal segments unaffected by the disease are then processed in the laboratory for immunological analysis and T cell expansion. Peripheral blood leukocytes (PBL) are purified by Ficoll-Hypaque (Pharmacia, Amersham).
10 Single cell suspensions of lymph node cells are obtained by gentle pressure using a loose fit glass homogenizer. Antigen extracts from intestinal samples are prepared by homogenizing the tissue fragments in a Dounce homogenizer, followed by 5 min denaturation at 97°C. The cells are then subjected to
15 functional analyses.

EXAMPLE 2. Isolation of T cell populations

Lymphocytes and monocytes are purified from blood samples or
20 buffy coats using Ficoll-Hypaque Plus (Amersham Biosciences, Uppsala, Sweden). The buffy coat or blood sample in question was diluted with PBS and carefully layered onto Ficoll-sodium diatrizoate solution, after which the two-phase system was centrifuged at 400·g for 30 min. The lymphocytes and monocytes
25 collected at the interphase between the Ficoll solution and plasma, whereas erythrocytes and granulocytes gathered at the bottom of the tubes. The lymphocyte layer was removed using a Pasteur pipette, and the cells washed with HBSS to remove excess Ficoll-Hypaque Plus, plasma and platelets. When the
30 cells were not used immediately, they were stored at 37°C in RPMI medium containing 10 % HuS, 1 % PeSt and 1 % glutamine. Conventional CD4⁺ Th cells and CD4⁺CD25⁺ cells were purified using an autoMACS Separator (Miltenyi Biotec, Bergisch Gladbach, Germany). The cells were counted, centrifuged at
35 300·g for 10 min, the supernatant pipetted off completely and

cells dissolved in 90 μL of MACS buffer (0,5 % BSA, 2 mM EDTA, 0,01 % sodium azide in PBS) and 10 μL of Biotin-Antibody Cocktail per 10^7 cells. This labelled all non- CD4^+ cell populations. After 10 min incubation at 4°C , 20 μL of Anti-Biotin MicroBeads per 10^7 cells was added and incubated for an additional 15 min at 4°C . Cells were washed with ~20 volumes of MACS buffer, centrifuged at $300 \cdot g$ for 10 min, the supernatant removed completely and the cells re-suspended in 500 μL of MACS buffer per 10^8 cells. Magnetic separation was carried out in an autoMACS Separator to deplete non- CD4^+ cells from the sample. To isolate the $\text{CD4}^+/\text{CD25}^+$ cells from the CD4^+ population, the CD4^+ cells were counted, centrifuged at $300 \cdot g$ for 10 min, supernatant removed completely and pellet dissolved in 90 μL of MACS buffer and 10 μL of CD25 MicroBeads per 10^7 cells. After 15 min incubation at 4°C , the cells were washed as described above and the CD25^+ cells were isolated by positive selection in the autoMACS Separator. Aliquots from all populations were analyzed by flow cytometry.

20 EXAMPLE 3. Characterization of cells by flow cytometry

Flow cytometry (FACS) was used to investigate the expression of different surface markers on the isolated cell populations. Cells were distributed into 4 mL FACS tubes, washed with 2 mL FACS buffer (2 % FCS, 0,05 % sodium azide in PBS), centrifuged at $300 \cdot g$ for 10 min, the supernatant poured off and the pellet re-suspended in 100 μL of FACS buffer. 7 μL of each antibody to be used was added and incubated for at least 30 min at 4°C whereupon the cells were washed once more with FACS buffer, centrifuged as before and re-suspended in 1 mL of FACS Lysing Solution for cell fixation and erythrocyte lysis, mixed and incubated for 10 min at 4°C before washing and centrifuging them once again as described above. The supernatant was poured off and cells re-suspended in 500 μL of FACS buffer per tube, after which analysis was carried out using a Becton Dickinson FACSCalibur instrument (Franklin Lakes, NJ, USA).

EXAMPLE 4. Proliferation assays

The suppressive ability of the purified CD4⁺CD25⁺ cells was investigated. Round-bottomed 96-well plates were incubated
5 with 25 µL of 5 µg/mL anti-CD3 IgG in PBS at 37°C for 90 min. Wells were washed three times with 200 µL of PBS after which three different cell populations were added. 100 µL of 500,000 cells/mL CD4⁺ cells radiated with 25 Gy were added to each well together with 30 000 CD4⁺CD25⁻ responder cells and 1 µg/mL
10 soluble CD28 (final concentration). CD4⁺CD25⁺ cells were added to the wells in varying amounts to create a number of different CD4⁺CD25⁻/CD4⁺CD25⁺ cell ratios. As a control equal numbers of CD4⁺CD25⁻ cells were added to the responder cells in an additional set of wells.

15 Plates were kept at 37°C for four or five days before 1 µCi of [3H]thymidine was added to each well and the plates were incubated for 18 h after which they were frozen at -20°C. The cells were thawed, and the well content transferred to a glass
20 fibre filter (Wallac, Turku, Finland) by a cell harvester (TOMTEC, Hamden, CT, USA). MeltiLex A - Melt-on scintillator sheets (Wallac, Turku, Finland) were placed on top of the filters and melted at 85°C. Radioactivity was measured using a 1205 Betaplate Liquid Scintillation Counter (Wallac, Turku,
25 Finland).

EXAMPLE 5. TGF-β1 induction of FoxP3

CD4⁺CD25⁻ cells were prepared from buffy coat as described
30 above. PBMCs were prepared from a different donor and radiated (25 Gy). 3·10⁶ CD4⁺ cells were then allo-activated by 1.25·10⁶ PBMCs in a volume of 2 mL/well in 6-well plates with or without 1 ng/mL TGF-β1. The plates were kept at 37°C. Cell samples were taken for flow cytometry analysis and RNA
35 extraction on day 2, 5 and 7.

EXAMPLE 6. RNA extraction and cDNA synthesis

RNA extraction from different cell populations for later use in PCR assays was performed as follows. Cells were lysed in 1 mL of TRIzol (Invitrogen, Carlsbad, CA, USA) per 5-6·10⁶ cells, after which the sample was either immediately frozen at -70°C for storage or incubated 5 min at room temperature before 0.2 mL of chloroform per mL TRIzol reagent was added and the tubes shaken vigorously. After 2-3 min the tubes were centrifuged at 12,000·g for 15 min at 4°C. The transparent, aqueous phase was transferred to a fresh tube, the RNA precipitated with 0.5 mL of isopropanol per mL TRIzol used in a first step and incubated 10 min at room temperature before centrifugation at 12,000·g for 10 min at 4°C. The supernatant was removed and the pellet washed with 1 mL of 75 % ethanol per mL TRIzol used in the first step. Tubes were vortexed and centrifuged at 7,500·g for 5 min at 4°C, after which the pellets were briefly air-dried (5-10 min), dissolved in 10 µL of RNase free water and incubated at 55°C for 10 min. cDNA synthesis was carried out with iScript cDNA Synthesis Kit (Bio-Rad, Hercules, CA, USA) according to the manufacturer's protocol.

EXAMPLE 7. Polymerase Chain Reaction (PCR) and Real Time Quantitative PCR

Synthesized cDNA from the different cell populations was used in PCR. The aim with the PCR was to detect expression of FoxP3, GAPDH and RNA polymerase II (RPII) Primers were obtained from Cybergene AB, Huddinge, Sweden. Sequences: FoxP3, forward - CAGCACATTCCCAGAGTTCCTC, reverse - GCGTGTGAACCACTGGTAGATC; GAPDH, forward - GAAGGTGAAGGTCGGAGTC, reverse - GAAGATGGTGATGGGATTTC; RPII, forward - GCACCACGTCCAATGACAT, reverse - GTGCGGCTGCTTCATAA. For ordinary PCR ThermoPol Reaction Buffer and Taq DNA polymerase

(New England Biolabs, Frankfurt am Main, Germany) was used whereas iQSYBR Green Supermix (Bio-Rad, Hercules, CA, USA) was used in quantitative real time PCR. The reactions were carried out in MyCycler Thermal Cycler or iCycler iQ Real-Time PCR
5 Detection System (Bio-Rad, Hercules, CA, USA) respectively, according to the protocol: 30s at 95°C, 30s at 48-65°C and 30s at 72°C for at least 40 repetitive cycles. For the quantitative real-time PCR reactions a relative quantitation method was applied as reviewed by Ginzinger D G, Exp. Hematol.
10 2002 30:503-12. PCR products were separated on a 2 % agarose gel, and bands detected by UV excitation of incorporated ethidium bromide.

EXAMPLE 8. Investigation of Tregs obtained from sentinel nodes
15 of IBD patients

Sentinel nodes from a total of 20 patients with IBD were harvested by surgery and investigated. Sentinel node acquired lymphocytes (SEALs) were characterized with respect to the
20 expression of surface markers CD4, CD8, CD3, the T cell receptor (TCR), CD25, CD69, CD45RA, CD45RO and CD14. The number of CD4⁺CD25^{hi} Treg population was examined (Figure 1). The number of Treg cells was found to be normal in sentinel nodes draining an inflammatory bowel segment, compared to the
25 number of Tregs in sentinel nodes draining non-affected areas of the intestine.

Tregs (CD4⁺CD25^{hi}) purified from sentinel nodes draining an inflammatory segment and Tregs (CD4⁺CD25^{hi}) purified from a
30 bowel segment not engaged in the inflammation however differed substantially in their regulatory activation of T cells: the Tregs from sentinel nodes draining the inflammatory site performed unsatisfactory in regulating the activation of T cells (Figure 2). In other words, the T cell response

regulating capability of Tregs from sentinel nodes draining affected areas is clearly impaired.

EXAMPLE 9. Expansion of regulatory T cells

5

For Treg induction and expansion CD4⁺CD25⁻ T cells from peripheral blood were stimulated with IL-2, CD28, allogenic feeder cells and TGF- β . After expansion >85% of CD4⁺ T cells expressed high levels of CD25 indicating Treg phenotype as investigated by FACS (not shown). RT-PCR demonstrated that the expanded population contained the Treg hallmark transcript FoxP3, which was not present in the CD4⁺CD25⁻ population prior to expansion (Figure 3).

15 The induced and expanded Treg population was tested functionally to demonstrate its capability to inhibit T cell activation. The expanded CD4⁺CD25⁺ FoxP3⁺ population (Figure 3) exhibited Treg regulatory properties as demonstrated by its inhibition of T helper cell proliferation already at a 1:10 ratio (Figure 4). The number of Tregs seems to be normal in sentinel nodes draining inflammatory bowel segments from patients with IBD; they appear to be inefficient in inhibiting the response of stimulated T cells. By stimulating and expanding naïve T cells efficient T regulatory properties
25 can be accommodated.

EXAMPLE 10. Recognition of antigens derived from the inflammatory site by regulatory T cells derived from sentinel nodes

30

Regulatory T cells derived from sentinel nodes, negative nodes, intestinal epithelia and peripheral blood leukocytes were incubated in different concentrations of intestinal cell homogenates for five to seven days. Proliferation was observed

for Tregs from sentinel nodes stimulated with antigen extract from an inflammatory area (Figure 5).

However, when regulatory T cells derived from sentinel nodes
5 were stimulated with antigen extracts derived from non-inflammatory sites no or very weak responses were seen (not shown). Findings from patients with Crohn's disease were similar to those from patients with ulcerative colitis.

10 Sentinel nodes draining affected areas in the caecum and in the sigmoid colon both recognize and proliferate against the same antigen extract derived from the inflammatory site (Figure 6). The findings indicate that there is a common antigen responsible for clonal expansion of T cells. The high
15 spontaneous proliferation observed is due to *in vivo* clonal T cell expansion in the sentinel node. Regulatory T cells derived from a sentinel node respond in proportion to the dose of antigen extract.

20 Regulatory T cells derived from sentinel nodes display an increased expression of cytokines. CD4⁺ T cells from sentinel nodes are CD45RO⁺/CCR5⁺ in patients with UC suggesting a state of exaggerated Th2 inflammatory response against the inflammatory mucosa (Figure 7).

25 Sentinel nodes from patients with UC secrete increased amounts of IL-4, a Th2 cytokine, upon activation with an antigen extract from the inflammatory site (Figure 8).

30 Regulatory T cells derived from sentinel nodes from patients with Crohn's disease respond with an increased production of the Th1 cytokine IFN- γ upon antigen specific activation (Figure 9).

The findings indicate that T cells derived from sentinel nodes from patients with UC have an exaggerated proliferative response against antigens from the inflammatory site and that the regulatory T cells derived from sentinel nodes react with a T_H2 response as suggested by the increased CCR5 expression (Figure 7) and increased production of T_H2 cytokines such as IL-4 (Figure 8). T cells derived from sentinel nodes from patients with Crohn's disease display an opposite pattern of activation with predominantly production of T_H1 cytokines such as IFN- γ .

Endothelial cells from the ulcer in UC or the inflammatory site in the intestine of patients with UC or Crohn's disease go into apoptosis or necrosis and are then endocytosed by professional antigen presenting cells (APC). APC endocytose disintegrated endothelial cells and migrate via lymphatic vessels to the draining lymph node, the sentinel node, where APC processes and present proteins (antigens) derived from the endothelial cells to T cells. The T cells become activated, undergo clonal expansion and leave the lymph node to seek areas of inflammation where they become effector cells and contribute to the inflammation.

EXAMPLE 11. Treatment of IBD

IBD, in particular Crohn's disease and/or ulcerative colitis, can be treated by harvesting regulatory T cells from sentinel nodes of a patient draining the inflammatory area, expanding them and activating them *in vitro*, and giving them back to the patient by infusion. In particular, *in vitro* expansion of regulatory T cells in an antigen specific manner takes advantage of clonally expanded Tregs accumulated in sentinel nodes. Treg from sentinel nodes from patients with IBD are expanded under GMP conditions and activated using a cytokine cocktail containing IL-2, IL-7, IL-10, TGF- β together with

antigen extract from the inflammatory bowel segment to achieve antigen specific expansion of Tregs. The efficacy of activation is evaluated with RT-PCR, FACS and *in vitro* inhibition assays as described (Figure 4). Additional
5 activation with anti-CD3, anti-CD28 and rapamycin can be used to obtain optimal expansion and functionality of the IBD Tregs. Alternatively, if the Tregs from sentinel nodes draining an inflammatory bowel segment are not satisfactory in regard of their number and/or function, expansion of Tregs
10 from sentinel nodes draining non-inflammatory segments using the same protocols is considered. The rationale behind this alternative approach is that one may consider a non-inflammatory segment being appropriately regulated, thus the Tregs are functional although too few in number to maintain
15 regulation in adjacent bowel segments. Expanded Tregs of either kind are given back as an auto-transfusion to reinstate orderly immunoregulation.

Legends to Figures

20

Figure 1. Tregs identified by the expression of surface markers CD4⁺ and high level of the IL-2 receptor CD25, CD4⁺CD25^{hi}.

Figure 2. Functional Treg assay. Purified Tregs from sentinel
25 nodes draining an inflammatory intestinal segment (open circles) and a non-inflammatory segment (filled circles) were incubated with CD4⁺CD25⁻ T helper cells purified from peripheral blood and activated with anti CD3 and anti CD28. 3H Thymidine was added to the cultures during the final 16 hours.

30 *Figure 3.* Expansion of Treg cells. CD4⁺CD25⁻ cells were activated with IL-2, TGF- β , anti-CD28 and feeder cells in order to stimulate expansion of Treg cells. mRNA was extracted from expanded CD4⁺CD25⁺ Tregs and the FoxP3 transcript was demonstrated by RT-PCR, identifying the expanded population as
35 Treg.

Figure 4. Regulation of stimulated responder T helper cells. A pre-expansion population containing $CD4^+CD25^-$ cells is unable to regulate (filled circles) whereas adequate regulation is obtained by expanded $CD4^+CD25^+$ cells (open circles).

Figure 5. Activation of human immunological cells. Cells obtained from sentinel lymph nodes, non-sentinel lymph nodes, the intestine as well as peripheral blood lymphocytes were activated by an endogenous antigen extract derived from an inflamed bowel section of an ulcerative colitis patient.

Figure 6. Dose response diagram. The dose response of regulatory T cells derived from sentinel nodes draining the caecum and the sigmoid colon, activated by an endogenous antigen extract derived from the same inflammatory bowel section, is shown.

Figure 7. Expression of CCR5 as a marker for T_H2 . Regulatory T cells derived from sentinel nodes from a patient with UC have elevated expression of CCR5 on $CD4^+ CD45RO^+$ memory cells as a sign of $TH2$ activation.

Figure 8. Secretion of the TH_2 cytokine IL-4 in regulatory T cells from patients with UC. Cells from non-sentinel nodes (nSLN) and from sentinel nodes (SLN) were stimulated for 48 h with inflammatory antigen extracts. Secreted IL-4 was measured in cultured supernatants using sandwich ELISA.

Figure 9. Secretion of the T_H1 cytokine IFN- γ by regulatory T cells derived from sentinel nodes from patients with Crohn's disease. Cells from non-sentinel nodes (nSLN) and from sentinel nodes (SLN1 and SLN2) were stimulated for 48 h with an inflammatory antigen extract. Secreted IFN- γ was measured in cultured supernatants using sandwich ELISA.

CLAIMS

1. A composition for treating inflammatory bowel disease comprising expanded T cells, the expanded T cells being obtained by (a) collecting regulatory T cells in an activated or non-activated state from a patient's sentinel lymph nodes draining bowel segments with or without IBD and (b) expanding the collected T cells *in vitro*, said composition being adapted for re-infusing the expanded T cells to the patient.
2. The composition of claim 1, wherein the inflammatory bowel disease is Crohn's disease.
3. The composition of claim 2, wherein the inflammatory bowel disease is ulcerative colitis.
4. The composition of any of claims 1-3, wherein said regulatory T cells are collected in an activated state.
5. The composition of claim 3 or 4, wherein said T cells are collected from lymph nodes draining bowel segments with IBD.
6. The composition of claim 5, wherein the collected T cells are additionally activated by one or more cytokines in combination with an antigen extract obtained from an inflamed bowel segment.
7. The composition of claim 6, wherein activation precedes expansion.
8. The composition of claim 6, wherein activation is concurrent with expansion.
9. The composition of claim 6, wherein expansion is followed by activation.
10. The composition of claim 6, wherein the one or more cytokine is selected from IL-2, IL-7, IL-10, TGF- β 1, TSLP.
11. The composition of claim 6 comprising further activation of collected T cells by anti-CD3, anti-CD28 and/or rapamycin.
12. The composition of any of claims 1-3, wherein said regulatory T cells are collected in a non-activated state.
13. The composition of claim 12, wherein the regulatory T cells are collected from lymph nodes draining bowel segments without IBD.

14. The composition of claim 12 or 13, wherein the collected T cells are activated by one or more cytokines in combination with an antigen extract obtained from an inflamed bowel segment.
15. The composition of claim 14, wherein activation precedes expansion.
- 5 16. The composition of claim 14, wherein activation is concurrent with expansion.
17. The composition of claim 14, wherein expansion is followed by activation.
18. The composition of claim 14, wherein the one of more cytokine is selected from IL-2, IL-7, IL-10, TGF- β 1, TSLP.
- 10 19. The composition of claim 14-18, comprising further activation of collected T cells by anti-CD3, anti CD28 and/or rapamycin.
20. The composition of claim 2, wherein regulatory T cells are collected from lymph nodes draining bowel segments with or without IBD and are activated by contacting them with an antigen extract obtained from the inflammatory site in combination with a low dose of the cytokine IL-2, and expanded.
- 15 21. The composition of claim 20, further comprising activation by contacting said regulatory T cells with an antibody against IFN- γ and/or TNF- α to suppress T_H1 skewing and T_H2 expansion.
- 20 22. The composition of claim 20 or 21, comprising conditioning the culture medium in intervals such as from every day to every fifth day, in particular every day for a time period of from one week or more, in particular for a time period of from three to four weeks.
23. The composition of any of claims 20-22, comprising activation by an anti CD28 antibody in combination with an anti IL-12 antibody to improve T_H2 skewing.
- 25 24. The composition of claim 3, wherein regulatory T cells are collected from lymph nodes draining bowel segments with or without IBD and are activated by contacting them with an antigen extract obtained from the inflammatory site in combination with one or more of the cytokines IL-2, IL12 and IFN- α , and expanded.
- 30

25. The composition of claim 24, wherein additional activation is provided by contacting the regulatory T cells with CD3 and/or CD28.
26. Use of one or more cytokines in a pharmaceutically acceptable carrier for *in-vitro* activation of regulatory T cells collected from a sentinel node of a patient with IBD.
27. The use of claim 26, wherein IBD is Crohn's disease.
28. The use of claim 26, wherein IBD is ulcerative colitis.
29. A method of isolating regulatory T cells from an excised sentinel lymph node draining a bowel segment with IBD, comprising (a) exerting a pressure on the sentinel lymph node, (b) collecting the cell suspension expelled from the node, (c) isolating regulatory T cells from the cell suspension.
30. A regulatory T cell isolated from a sentinel lymph node draining a bowel segment affected by inflammatory bowel disease (IBD).
31. The regulatory T cell of claim 30, wherein the IBD is Crohn's disease.
32. The regulatory T cell of claim 30, wherein the IBD is ulcerative colitis.
33. A method of expanding regulatory T cells useful in the treatment of IBD, comprising (a) providing in a suitable culture medium regulatory T cells collected from a sentinel node of an IBD patient, (b) activating the cells by contacting them with a cytokine; (c) expanding the activated cells.
34. The method of claim 33, wherein some or all activation is carried out during expansion.
35. A composition for re-establishing the T_H1/T_H2 balance in a patient suffering from Crohn's disease comprising expanded T cells, said expanded T cells being obtained by (a) collecting regulatory T cells in an activated state from lymph nodes of the patient draining bowel segments with or without Crohn's disease; (b) activating the T cells so collected with antigen extract from the inflammatory site in combination with a low dose of the cytokine IL-2; (c) expanding the cells; (d) optionally contacting the cells during expansion with antibody against IFN- γ and/or TNF- α to suppress T_H1 skewing and T_H2 expansion, said composition being adapted for re-infusing the expanded T cells to the patient.

- 36. The composition of claim 35, combining conditioning of the culture medium in intervals.
- 37. The composition of claim 36, wherein conditioning is from every day to every fifth day.
- 5 38. The composition of claim 36, comprising stimulating expansion by contacting the cells with a CD3 antibody.
- 39. Use of a cytokine selected from IL-2, IL-7, IL-10, TGF- β 1, TSLP for activating regulatory T cells obtained from a sentinel node of a patient suffering from IBD.
- 10 40. Use of an antigen extract obtained from an inflamed bowel segment of a patient suffering from IBD in combination with a cytokine for activating regulatory T cells obtained from a sentinel node of the patient.
- 41. The use of claim 40, wherein the cytokine is selected from IL-2, IL-7, IL-10, TGF- β 1, TSLP.
- 15 42. The uses of claims 39 and 40-41, respectively in combination.