

Isolated photoprotein mtClytin, and use thereof

The invention relates to the photoprotein mtClytin, to its nucleotide and amino acid sequences and to the activity and use of the photoprotein mtClytin.

Photoproteins

- 5 The phenomenon of the generation of light by living organisms is designated bioluminescence. It is the result of biochemical reactions in cells, in which reactions the chemical energy is emitted in the form of light quanta (what is termed cold emission by means of chemoluminescence). While the light which is produced in this way is monochromatic, since it is emitted in connection with a discrete electron transfer, it can be shifted by secondary luminescent dyes (e.g. fluorescent proteins
- 10 in the case of luminescent jellyfish of the genus *Aequoria*) into spectral regions of longer wavelength.

Bioluminescence has a diversity of biological functions: at an ocean depth of between 200 and 1000 m (mesopelagial), about 90% of all living organisms luminesce. In this case, the luminescent

15 signals are employed for attracting partners, for deception and as a lure. Glowworms and fireflies also use the light signals for seeking partners. On the other hand, the significance of the luminescence of bacteria, fungi and single-cell algae is unclear. It is assumed that it is used for coordinating many single individuals in a large population or else represents a type of biological clock.

- 20 A large number of coelenterates are bioluminescent (Morin et al., 1974). These organisms emit blue or green light. As an isolated protein, aequorin, which is derived from *Aequoria victoria* (Shimomura et al., 1969) and which, in 1962, was the first light-producing protein to be identified, emitted a blue light, and not a green light as observed phenotypically in the case of *Aequoria victoria*. The green fluorescent protein (GFP) which, as a result of being activated by aequorin, causes *Aequoria victoria* to appear phenotypically green was subsequently isolated from this medusa (Johnson et al., 1962; Hastings et al., 1969; Inouye et al., 1994). Other photoproteins which have also been identified and described are clytin (Inouye et al., 1993), mitrocomin (Fagan et al., 1993) and obelin (Illarionov et al., 1995).
- 25

Table 1: Overview of some photoproteins. The table gives the name, the organism from which the protein has been isolated and the identification number (Acc. No.) of the database entry.

Name	Organism	Identification No.
Obelin	Obelia geniculata	AAL86372
Clytin	Clytia gregaria	CAA49754
Aequorin	Aequorea macrodactyla	AAK02061
Aequorin	Aequorea parva	AAK02060
Mitrocomin	Mitrocoma cellularia	AAA29298
Pholasin	Pholas dactylus	AAM18085
?	Symplectoteuthis oualaniensis	AX305029

5 **Table 2:** Overview of some photoproteins. The table gives the organism from which the protein has been isolated, the name of the photoprotein and a selection of patents or applications.

Organism	Fluorescent protein	Patent/Application
Obelia geniculata	Obelin	WO03006497
Clytia gregaria	Clytin	WO03006497
Aequoria victoria	Aequorin	WO200168824 US-0908909 US 6,152,358 JP-0176125
Pholas dactylus	Pholasin	WO0028025 GB-0024357

10 Bioluminescence is nowadays used in technology in a wide variety of ways, e.g. in the form of bioindicators of environmental pollution or in biochemistry for sensitively detecting proteins or for quantifying particular compounds, or as what are termed reporters in connection with investigating gene regulation in the cell.

15 The photoproteins differ not only in their nucleotide and amino acid sequences but also in their biochemical and physical properties.

It has been demonstrated that the physical and biochemical properties of photoproteins can be

altered by altering the amino acid sequences of these proteins. Examples of mutagenized photoproteins are described in the literature (US 6,495,355; US 5,541,309; US 5,093,240; Shimomura et al., 1986).

- 5 The abovementioned photoproteins generate light by oxidizing coelenterazine (Haddock et al., 2001; Jones et al., 1999).

Reporter systems

In general, genes whose gene products can be readily detected using simple biochemical or histochemical methods are termed reporter genes or indicator genes. At least 2 types of reporter
10 gene are distinguished.

1. Resistance genes. This is the term used for genes whose expression confers, on a cell, resistance to antibiotics or other substances whose presence in the growth medium leads to the death of the cell if the resistance gene is absent.
- 15 2. Reporter genes. The products of reporter genes are used in genetic manipulation as fused or unfused indicators. The commonest reporter genes include beta-galactosidase (Alam et al., 1990), alkaline phosphatase (Yang et al., 1997; Cullen et al., 1992), and luciferases and other photoproteins (Shinomura, 1985; Phillips GN, 1997; Snowdowne et al., 1984).

20 The emission of photons in the visible spectral range, with this emission being effected by means of excited emitter molecules, is termed luminescence. In contrast to fluorescence, the energy is not, in this case, supplied from the exterior in the form of radiation of shorter wavelength.

- 25 A distinction is made between chemoluminescence and bioluminescence. A chemical reaction which leads to an excited molecule which itself luminesces when the excited electrons return to the basal state is termed chemoluminescence. If this reaction is catalyzed by an enzyme, the phenomenon is then referred to as being bioluminescence. The enzymes involved in the reaction are generally termed luciferases.

30 Classification of the species *Clytia gregaria*

Cnidaria → Leptomedusae → Campanulariidae → *Clytia gregaria*

The species *Clytia gregaria* belongs to the Cnidaria, specifically to the Medusae. The bioluminescent and fluorescent phenotype, respectively, has already been described in 1998 (Ward et al., 1998).

Isolating the cDNA

In order to investigate the bioluminescence activity of the species *Clytia gregaria*, specimens were caught in the White Sea (Kartesh Biological Station, Russia) and stored in liquid nitrogen. In order to construct the *Clytia gregaria* cDNA libraries, the poly(a)+ RNA was isolated using the “Straight A” isolation method from Novagen (USA).

An RT-PCR was carried out for preparing the cDNA. For this, 1 µg of RNA was incubated with reverse transcriptase (Superscript Gold II) in accordance with the following scheme:

10	PCR	1.	30	seconds	95°C
		2.	6	minutes	68°C
		3.	10	seconds	95°C
		4.	6	minutes	68°C
		17 cycles of step 4 after step 3			

The reaction products were incubated with proteinase K, at 37°C for 30 minutes, in order to inactivate the polymerase, and the cDNA was precipitated with ethanol. The cDNA expression library was constructed using the Clontech (USA) “SMART cDNA” library construction kit in accordance with the manufacturer’s instruction. The cDNA was cloned into the expression vector pTriplEx2 (Clontech; USA). The expression vectors were transformed by electroporation into bacteria of the strain *E. coli* XL1 blue.

The bacteria were plated out on solid LB nutrient medium and incubated at 37°C for 24 hours. A replica plating was then carried out, with the bacteria being transferred to another solid nutrient medium plate using a nitrocellulose filter. The replica plate was in turn incubated at 37°C for 24 hours and the bacterial colonies which had grown were transferred into liquid LB medium. After IPTG (final concentration, 0.1 mM) had been added, the bacteria were incubated at 37°C for 4 hours on a shaker. The bacteria were harvested by centrifugation and the bacterial mass was resuspended, at 0°C, in 0.5 ml of disruption buffer (5 mM EDTA, 20 mM Tris-HCL, pH 9.0). The bacteria were then disrupted by ultrasonication.

After adding coelenterazine (final concentration, 10E-07 M), the lysates were incubated at 4°C for 3 hours. The bioluminescence was then measured in a luminometer after adding calcium chloride (final concentration, 20 mM).

A photoprotein was identified. The photoprotein was designated mtClytin. The photoprotein mtClytin is described in detail below.

With an identity of 87%, the photoprotein mtClytin exhibits the highest homology at the amino acid level with clytin from *Clytia gregaria* and an identity of 77% with obelin from *Obelia geniculata* (shown in Example 8; Figure 8). The homology of 87% - in relation to clytin - occurs at the C-terminal end of the protein, multiple amino acid substitutions being identifiable distributed over the entire protein. At the nucleic acid level, the identity is less than 30% (shown in Example 7; Figure 7). The BLAST method (Altschul et al., 1997) was used for the sequence comparison.

The photoprotein clytin-2 exhibits the highest homology at the amino acid level with clytin from *Clytia gregaria*. However, the sequence exhibits a number of differences in the amino acid sequence, with these differences being depicted in Example 11 (Figure 9). These differences can lead to changes in physicochemical, biochemical and bioluminescent properties. The photoprotein clytin-2 does not possess any signal peptide (as shown in Example 10).

The photoprotein mtClytin possesses a signal peptide which can lead to the photoprotein being translocated into mitochondria. The signal peptide was identified by the computer program MITOPROT (Claros et al., 1996) (shown in Example 10). The signal peptide which was determined by MITOPROT is given in SEQ ID NO: 3. The photoprotein mtClytin is the first photoprotein in which a natural signal peptide for translocation into mitochondria has been identified.

The invention also relates to functional equivalents of mtClytin. Functional equivalents are those proteins which have comparable physicochemical properties and are at least 70% homologous with SEQ ID NO: 2. Preference is given to a homology of at least 80% or 90%. A homology of at least 95% is particularly preferred.

The invention also relates to the functional equivalents of the mtClytin signal peptide. Functional equivalents are those proteins or peptides which have comparable physicochemical properties and are at least 70% homologous to SEQ ID NO: 3. Preference is given to a homology of at least 80% or 90%. A homology of at least 95% is particularly preferred.

The photoprotein mtClytin is suitable for being used as a reporter gene for cellular systems, especially for receptors, for ion channels, for transporters, for transcription factors or for inducible systems.

The mtClytin signal peptide is also suitable for being fused to reporter genes in order to be used as a fused reporter gene for cellular systems, especially for receptors, for ion channels, for transporters, for transcription factors or for inducible systems.

The photoprotein mtClytin is also suitable for being used as a reporter gene by labeling,

identifying and characterizing cell organelles, especially for mitochondria.

The mtClytin signal peptide is also suitable for being fused to peptides or proteins for translocation into cell organelles, especially mitochondria.

5 The photoprotein mtClytin is also suitable for being used as a reporter gene for determining parameters inside and outside cell organelles, especially mitochondria, especially calcium concentrations.

The mtClytin signal peptide is also suitable, as a fusion peptide, for being used as a reporter gene for determining parameters inside and outside cell organelles, especially mitochondria, especially calcium concentrations.

10 The photoprotein mtClytin is suitable for being used as a reporter gene in bacterial and eukaryotic systems, especially in mammalian cells, in bacteria, in yeasts, in baculo and in plants.

The photoprotein mtClytin is suitable for being used as a reporter gene for cellular systems in combination with bioluminescent or chemoluminescent systems, especially systems using luciferases, using oxygenases or using phosphatases.

15 The mtClytin signal peptide is also suitable, as fusion peptide, for being used as a reporter gene for cellular systems in combination with bioluminescent or chemoluminescent systems, especially systems using luciferases, using oxygenases or using phosphatases.

20 The photoprotein mtClytin is suitable for being used as a fusion protein, especially for receptors, for ion channels, for transporters, for transcription factors, for proteinases, for kinases, for phosphodiesterases, for hydrolases, for peptidases, for transferases, for membrane proteins and for glycoproteins.

25 The mtClytin signal peptide is also suitable, as fusion peptide, for being used as a fusion protein, especially for receptors, for ion channels, for transporters, for transcription factors, for proteinases, for kinases, for phosphodiesterases, for hydrolases, for peptidases, for transferases, for membrane proteins and for glycoproteins.

The photoprotein mtClytin is suitable for being immobilized, especially by antibodies, by biotin, or by magnetic or magnetizable supports.

30 The photoprotein mtClytin is suitable for being used as a protein for energy transfer systems, especially FRET (fluorescence resonance energy transfer), BRET (bioluminescence resonance energy transfer), FET (field effect transistors), FP (fluorescence polarization) and HTRF (homogeneous time-resolved fluorescence) systems.

The photoprotein mtClytin is suitable for labeling substrates or ligands, especially for proteases, for kinases or for transferases.

The photoprotein mtClytin is suitable for being expressed in bacterial systems, especially for titer determination, as a substrate for biochemical systems, especially for proteinases and kinases.

- 5 The photoprotein mtClytin is suitable for being used as a label, especially coupled to antibiotics, coupled to enzymes, coupled to receptors or coupled to ion channels and other proteins.

The mtClytin signal peptide is also suitable, as fusion peptide, for being used as a label, especially coupled to antibiotics, coupled to enzymes, coupled to receptors or coupled to ion channels and other proteins.

- 10 The photoprotein mtClytin is suitable for being used as a reporter gene in the search for pharmacological active compounds, especially in HTS (high throughput screening).

The mtClytin signal peptide is also suitable for being used as a reporter gene in the search for pharmacological active compounds, especially in HTS (high throughput screening).

- 15 The photoprotein mtClytin is suitable for being used as a component of detection systems, especially for ELISA (enzyme-linked immunosorbent assay), for immunohistochemistry, for Western blotting or for confocal microscopy.

- 20 The photoprotein mtClytin is suitable for being used as a label for analyzing interactions, especially for protein-protein interactions, for DNA-protein interactions, for DNA-RNA interactions, for RNA-RNA interactions, or for RNA-protein interactions (DNA: desoxyribonucleic acid; RNA: ribonucleic acid).

The photoprotein mtClytin is suitable for being used as a label or fusion protein for expression in transgenic organisms, especially in mice, in rats, in hamsters and other mammals, in primates, in fish, in worms or in plants.

- 25 The mtClytin signal peptide is also suitable, as fusion peptide, for being used as a label or fusion protein for expression in transgenic organisms, especially in mice, in rats, in hamsters and other mammals, in primates, in fish, in worms or in plants.

The photoprotein mtClytin is suitable for being used as a label or fusion protein for analyzing embryonic development.

- 30 The photoprotein mtClytin is suitable for being used as a label by way of a coupling mediator, especially by way of biotin, by way of NHS (N-hydroxysulfosuccinimide) or by way of CN-Br.

The photoprotein mtClytin is suitable for being used as a reporter which is coupled to nucleic acids, especially to DNA or RNA.

The photoprotein mtClytin is suitable for being used as a reporter which is coupled to proteins or peptides.

- 5 The mtClytin signal peptide is also suitable, as fusion peptide, for being used as a reporter which is coupled to proteins or peptides.

The photoprotein mtClytin is suitable for being used as a reporter for measuring intracellular or extracellular calcium concentrations.

The photoprotein mtClytin is suitable for characterizing signal cascades in cellular systems.

- 10 The photoprotein mtClytin which is coupled to nucleic acids or peptides is suitable for being used as a probe, especially for Northern blots, for Southern blots, for Western blots, for ELISA, for nucleic acid sequencings, for protein analyses or for chip analyses.

The photoprotein mtClytin is suitable for being used for labeling pharmacological formulations, especially infectious agents, antibodies or "small molecules".

- 15 The photoprotein mtClytin is suitable for being used for geological investigations, especially for ocean, groundwater and river currents.

The photoprotein mtClytin is suitable for being expressed in expression systems, especially in in-vitro translation systems, in bacterial systems, in yeast systems, in baculo systems, in viral systems and in eukaryotic systems.

- 20 The mtClytin signal peptide is also suitable, as fusion peptide, for being expressed in expression systems, especially in in-vitro translation systems, in bacterial systems, in yeast systems, in baculo systems, in viral systems and in eukaryotic systems.

- 25 The photoprotein mtClytin is suitable for visualizing tissues or cells in connection with surgical interventions, especially in connection with invasive, in connection with noninvasive and in connection with minimally invasive interventions.

The photoprotein mtClytin is also suitable for labeling tumor tissues and other phenotypically altered tissues, especially in connection with histological investigation and in connection with surgical interventions.

The invention also relates to the purification of the photoprotein mtClytin, especially as a wild-type protein, as a fusion protein and as a mutagenized protein.

The invention also relates to the purification of the mtClytin signal peptide, especially as a wild-type protein, as a fusion protein and as a mutagenized protein.

- 5 The invention also relates to the use of the photoprotein mtClytin in the field of cosmetics, especially bath additives, lotions, soaps, body dyes, toothpaste and body powders.

The invention also relates to the use of the photoprotein mtClytin for dyeing, especially dyeing foodstuffs, bath additives, ink, textiles and plastics.

- 10 The invention also relates to the use of the photoprotein mtClytin for dyeing paper, especially greetings cards, paper products, wallpapers and handicraft articles.

The invention also relates to the use of the photoprotein mtClytin for dyeing liquids, especially for water pistols, fountains, beverages and ice.

The invention also relates to the use of the photoprotein mtClytin for producing toys, especially finger dye and makeup.

- 15 The invention relates to nucleic acid molecules which encode the polypeptide which is disclosed by SEQ ID NO: 2.

The invention relates to nucleic acid molecules which encode the polypeptide which is disclosed by SEQ ID NO: 3.

- 20 The invention relates to nucleic acid molecules which encode the polypeptide which is disclosed by SEQ ID NO: 6.

The invention relates to the polypeptide having the amino acid sequence which is disclosed in SEQ ID NO: 2.

The invention relates to the polypeptide having the amino acid sequence which is disclosed in SEQ ID NO: 3.

- 25 The invention relates to the polypeptide having the amino acid sequence which is disclosed in SEQ ID NO: 6.

The invention furthermore relates to nucleic acid molecules which are selected from the group consisting of

- a) nucleic acid molecules which encode a polypeptide which contains the amino acid sequence disclosed by SEQ ID NO: 2;
- b) nucleic acid molecules which contain the sequence depicted by SEQ ID NO: 1;
- c) nucleic acid molecules whose complementary strand hybridizes with a nucleic acid molecule from a) or b) under stringent conditions and which encode a polypeptide which exhibits the biological function of a photoprotein;

A stringent hybridization of nucleic acid molecules can be carried out, for example, in an aqueous solution comprising $0.2 \times \text{SSC}$ ($1 \times$ standard saline citrate = 150 mM NaCl, 15 mM trisodium citrate) at 68°C (Sambrook et al., 1989).

- d) nucleic acid molecules which differ from the nucleic acid molecules mentioned under c) due to the degeneracy of the genetic code;
- e) nucleic acid molecules which exhibit a sequence homology with SEQ ID NO: 1 of at least 95% and whose protein product exhibits the biological function of a photoprotein; and
- f) nucleic acid molecules which exhibit a sequence homology with SEQ ID NO: 1 of at least 65% and whose protein product exhibits the biological function of a photoprotein.

The invention also relates to nucleic acid molecules which exhibit a sequence homology with SEQ ID NO: 1 or SEQ ID NO: 5 of at least 95%, 90%, 85%, 80%, 75%, 70%, 65% or 60% and which encode a polypeptide which possesses the properties of a photoprotein.

- The invention also relates to nucleic acid molecules which exhibit a sequence homology with SEQ ID NO: 4 of at least 95%, 90%, 85%, 80%, 75%, 70%, 65% or 60% and which encode a polypeptide which possesses the properties of a signal or leader peptide.

The invention relates to the abovementioned nucleic acid molecules in which the sequence contains a functional promoter 5' to the photoprotein-encoding sequence or to the leader- or signal-sequence-encoding sequence.

- The invention also relates to nucleic acid molecules as previously described which are constituents of recombinant DNA or RNA vectors.

The invention relates to organisms which harbor such a vector.

The invention relates to oligonucleotides having more than 10 consecutive nucleotides which are identical or complementary to the DNA or RNA sequence of the mtClytin molecules or of the other molecules according to the invention.

5 The invention relates to photoproteins which are encoded by the previously described nucleotide sequences.

The invention relates to methods for expressing the photoprotein polypeptides according to the invention in bacteria, in eukaryotic cells or in *in-vitro* expression systems.

The invention also relates to methods for purifying/isolating a photoprotein polypeptide according to the invention.

10 The invention relates to peptides which have more than 5 consecutive amino acids and which are immunologically recognized by antibodies directed against the photoproteins according to the invention.

The invention relates to the use of the photoprotein-encoding nucleic acids according to the invention as marker genes or reporter genes, in particular for searching for pharmacological active
15 compounds and for diagnostics.

The invention relates to the use of the photoproteins according to the invention or of a photoprotein-encoding nucleic acid according to the invention as labels or reporters or as a marker gene or reporter gene.

20 The invention relates to the use of the photoprotein mtClytin (SEQ ID NO: 2), or to the use of a nucleic acid which encodes the photoprotein mtClytin as a label or reporter, or as a label or reporter gene, in particular for searching for pharmacological active compounds and for diagnostics.

The invention relates to the use of the nucleic acid depicted in SEQ ID NO: 1 as a marker gene or reporter gene, in particular for searching for pharmacological active compounds and diagnostics.

25 The invention relates to the use of the peptide depicted in SEQ ID NO: 6 and its underlying nucleic acid sequence SEQ ID NO: 5 as a marker gene or reporter gene, in particular for searching for pharmacological active compounds and diagnostics.

The invention also relates to polyclonal or monoclonal antibodies which recognize a polypeptide according to the invention.

The invention also relates to monoclonal or polyclonal antibodies which recognize the photoprotein mtClytin (SEQ ID NO: 2) or the photoprotein clytin-2 (SEQ ID NO: 6).

The invention also relates to monoclonal or polyclonal antibodies which recognize the signal peptide of the photoprotein mtClytin (SEQ ID NO: 3).

- 5 The invention furthermore relates to a nucleic acid molecule which is selected from the group consisting of
- a) nucleic acid molecules which encode a polypeptide which contains the amino acid sequence disclosed by SEQ ID NO: 3;
 - b) nucleic acid molecules which contain the sequence depicted by SEQ ID NO: 4;
 - 10 c) nucleic acid molecules whose complementary strand hybridizes with a nucleic acid molecule from a) or b) under stringent conditions and which encode a peptide which exhibits the biological function of a signal or leader peptide;
 - d) nucleic acid molecules which differ from the nucleic acid molecules mentioned under c) due to the degeneracy of the genetic code;
 - 15 e) nucleic acid molecules which exhibit a sequence homology with SEQ ID NO: 4 of at least 95% and encode a peptide which has the biological function of a signal or leader peptide; and
 - f) nucleic acid molecules which exhibit a sequence homology with SEQ ID NO: 4 of at least 65% and encode a peptide which has the biological function of a signal or leader peptide.
- 20 The invention likewise relates to a nucleic acid molecule which is selected from the group consisting of
- a) nucleic acid molecules which encode a polypeptide which contains the amino acid sequence disclosed by SEQ ID NO: 6;
 - b) nucleic acid molecules which contain the sequence depicted by SEQ ID NO: 5;
 - 25 c) nucleic acid molecules whose complementary strand hybridizes with a nucleic acid molecule from a) or b) under stringent conditions and which encode a polypeptide which exhibits the biological function of a photoprotein;
 - d) nucleic acid molecules which differ from the nucleic acid molecules mentioned under c) due to the degeneracy of the genetic code;

- e) nucleic acid molecules which exhibit a sequence homology with SEQ ID NO: 5 of at least 95% and encode a polypeptide which has the biological function of a photoprotein; and
- f) nucleic acid molecules which exhibit a sequence homology with SEQ ID NO: 5 of at least 80% and encode a polypeptide which has the biological function of a photoprotein.

5 The invention also relates to a nucleic acid which is as described in the preceding paragraphs and which contains a functional promoter 5' to the coding sequence.

The invention includes recombinant DNA or RNA vectors which contain the previously described nucleic acids.

10 Organisms which harbor a vector as previously described are likewise in accordance with the invention.

The invention also relates to oligonucleotides having more than 10 consecutive nucleotides which are identical or complementary to a constituent sequence of a nucleic acid molecule as described above.

15 The invention also relates to a polypeptide which is encoded by a nucleic acid sequence as described above.

The invention also relates to a method for expressing the abovementioned polypeptides in bacteria, viral cells, yeasts or eukaryotic cells or in *in-vitro* expression systems.

The invention likewise relates to a method for purifying/isolating a polypeptide according to the invention.

20 The invention likewise relates to peptides having more than 5 consecutive amino acids which are recognized immunologically by antibodies directed against the photoprotein mtClytin.

The invention furthermore relates to peptides having more than 5 consecutive amino acids which are recognized immunologically by antibodies directed against the photoprotein clytin-2.

25 The invention also relates to peptides having more than 5 consecutive amino acids which are recognized immunologically by antibodies directed against the signal or leader peptide disclosed by SEQ ID NO: 3.

The invention also relates to peptides having more than 5 consecutive amino acids which are recognized immunologically by antibodies directed against the photoprotein disclosed by SEQ ID NO: 6 (clytin-2).

The invention relates to the use of a nucleic acid according to the invention as a marker gene or reporter gene.

The invention also relates to the use of a photoprotein according to the invention as a label or reporter.

- 5 The invention furthermore relates to the use, as signal or leader sequence, of a nucleic acid which contains the sequence depicted as SEQ ID NO: 4 or a sequence having 60%, 65%, 70%, 75%, 80%, 85% or 90%, preferably having 95%, sequence identity with SEQ ID NO: 4.

The invention also relates to the use, as signal or leader peptide, of a peptide which contains the sequence depicted as SEQ ID NO: 3 or a sequence having 60%, 65%, 70%, 75%, 80%, 85% or
10 90%, preferably having 95%, sequence identity with SEQ ID NO: 3.

The invention likewise relates to the use, which is described in the two preceding paragraphs, for transporting proteins which are fused to the signal or leader peptide into cell organelles.

The invention also relates to the use which is described in the preceding paragraph, with the cell organelles being mitochondria.

- 15 The invention also relates to the use which is described in the preceding paragraph, with the cell organelles being the endoplasmic reticulum (ER).

The invention furthermore relates to the use of the nucleic acid sequence which is depicted as SEQ ID NO: 4 as a signal or leader sequence.

The invention also relates to the use of the peptide which is depicted as SEQ ID NO: 3 and which
20 contains the depicted sequence as a signal or leader peptide.

The invention likewise relates to the use which is described in the two preceding paragraphs for transporting a protein which is fused to the signal or leader peptide into cell organelles.

The invention also relates to the use which is described in the preceding paragraph, with the cell organelles being mitochondria.

- 25 The invention also relates to the use which is described in the preceding paragraph, with the cell organelles being the endoplasmic reticulum (ER).

The invention likewise relates to the use of the polypeptides according to the invention as reporter proteins in searching for pharmacological active compounds.

Finally, the invention also relates to the use of the nucleic acids according to the invention as

reporter genes in searching for pharmacological active compounds.

Expressing the photoproteins of the invention

The production of a molecule which, after the gene has been introduced into a suitable host cell, enables the foreign gene which is cloned into an expression vector to be transcribed and translated is termed expression. Expression vectors contain the control signals which are required for expressing genes in prokaryotic or eukaryotic cells.

In principle, expression vectors can be constructed in two different ways. In the case of what are termed transcription fusions, the protein encoded by the cloned-in foreign gene is synthesized as an authentic, biologically active protein. For this purpose, the expression vector carries all the 5' and 3' control signals which are required for the expression.

In the case of what are termed translation fusions, the protein encoded by the cloned-in foreign gene is expressed, together with another protein which can be detected readily, as a hybrid protein. The 5' and 3' control signals which are required for the expression, including the start codon and, possibly, a part of the sequences encoding the N-terminal regions of the hybrid protein to be formed, originate from the vector. The additional inserted protein moiety not only in many cases stabilizes the protein, which is encoded by the cloned-in foreign gene, against breakdown by cellular proteases; it can also be used for detecting and isolating the hybrid protein which is formed. The expression can take place either transiently or stably. Suitable host organisms are bacteria, yeasts, viruses or eukaryotic systems.

Purifying the photoproteins of the invention

The isolation of proteins (after they have been overexpressed as well) is frequently termed protein purification. A large number of established methods are available for purifying proteins.

The solid/liquid separation is a basic operation in connection with isolating proteins. This procedural step is required when separating cells from the culture medium, when clarifying the crude extract after having disrupted the cells and removing the cell debris, and when separating off sediments after precipitations, etc. It takes place by means of centrifugation and filtration.

In order to obtain intracellular proteins, the cell wall must be destroyed or rendered permeable.

High-pressure homogenizers or agitator ball mills or glass bead mills are used for this purpose,

depending on the scale and the organism. Mechanical cell integrations and ultrasonic treatment are used, inter alia, on the laboratory scale.

Both in the case of extracellular proteins and in the case of intracellular proteins (following cell
5 disruption), various precipitation methods using salts (in particular ammonium sulfate) or organic solvents (alcohols or acetone) represent rapid and efficient methods for concentrating proteins. When intracellular proteins are being purified, it is desirable to remove the soluble nucleic acids (precipitation with, for example, streptomycin sulfate or protamine sulfate). When extracellular proteins are being isolated, carriers (e.g. starch or kieselguhr) are frequently added before adding
10 the precipitating agents in order to obtain sediments which are easier to handle.

Numerous chromatographic methods and partition methods (absorption chromatography and ion exchange chromatography, gel filtration, affinity chromatography and electrophoreses) are available for high-degree purification. Column chromatography is also used on an industrial scale.
15 Affinity chromatography, which makes possible purification factors of up to several 100s per step, is especially important for the laboratory scale.

Extracellular proteins accrue in relatively dilute solutions. Just like extracellular proteins, they have to be concentrated before being subjected to further use. In addition to the methods which
20 have already been mentioned, ultrafiltration has proved to be of value, on an industrial scale as well.

Inorganic salts which accompany proteins are frequently undesirable in the case of specific applications. They can be removed by, inter alia, gel filtration, dialysis and diafiltration.
25

A large number of proteins are used as dry preparations. Important drying methods are vacuum drying, freeze drying and spray drying.

Nucleotide and amino acid sequences

30 The photoprotein mtClytin is encoded by the following nucleotide sequence (SEQ ID NO: 1):

5`-
gacagataaaaaattcactccttagattattttagtgaataagagaaaaaaggataagaaatcaag
atgcaaagggtttacaaatcgtcttctttccatgtcggttttacgtgcaagatcaagattgcaacgc
acggcaaattttcacaccagcatactcttggctacagattcaaaatacgcggtcaaactcgatcct
5 gattttgcaaattccaaaatggatcaacagacacaaatttatgttcaactttttggacataaacggt
aaggggaaaatcacattagatgaaatcgtctccaaagcttcagacgacatttgtgctaaactggat
gcaacaccagaacagaccaaacgtcaccaggatgctgttgaagcctttttcaagaaaatgggcatg
gattatggtaaagaagttgcattcccagaatttattaagggatgggaagagttggccgaacacgac
ttggaactctggtctcaaaacaaaagtacattgatccgtgaatggggagatgctgttttcgacatt
10 ttcgacaaagacgcaagtggctcaatcagtttagacgaatggaaggcttacggacgaatctctgga
atctgtccatcagacgaagacgctgagaagacggttcaaacattgtgatttggacaacagtggcaaa
cttgatggtgatgagatgaccaggcaacatttaggcttctggtacacattggatccaacttctgat
ggtctttatggcaattttgttcctaagaagcgttcagttaaaaacgctaaacattgttcagttgt
aaaattatattcattttcatttctgtaaaattagttattataaatttgtatcataaattgtatccat
15 gttgtagactaaataagactcggcaaaaaaaaaaaaaaaaaaaaaaaaaaaaaaa -3`.

This yields an amino acid sequence of (SEQ ID NO: 2):

MQRFTNRLLSMSALRARSRLQRTANFHTSILLATDSKYAVKLDPDFANPKWINRHKFMFN
FLDINGKGKITLDEIVSKASDDICAKLDATPEQTKRHQDAVEAFFKKMGMDYGKEVAFPE
FIKGWEELAEHDLELWSQNKSTLIREWGDVDFDKDASGSISLDEWKAYGRISGICPSDE
20 DAEKTFKHCDLDNSGKLDVDEMTRQHLGFWYTLDPSTDGLYGNFVP

The putative signal peptide of the photoprotein mtClytin possesses the following sequence (SEQ ID NO: 3):

MQRFTNRLLSMSALRA

and has the following nucleic acid sequence:

25 5`- atgcaaagggtttacaaatcgtcttctttccatgtcggttttacgtgca -3` (SEQ ID NO 4)

The photoprotein clytin-2 is encoded by the following nucleotide sequence (SEQ ID NO: 5):

5`-
GATCTCAGCTCAACTTGCAATAAGTATCAGATCAAATTTTGCAACTCAAAGCAAATCA
TCAACTTCATCATAATGACTGACACTGCTTCAAAATACGCTGTCAAACCTCAAGACCAA
30 CTTTGAAGATCCAAAATGGGTCAACAGACACAAATTTATGTTCAACTTTTGGACATT
AACGGCAACGGAAAAATCACTTTGGATGAAATTGTCTCCAAAGCTTCGGATGACATTT
GCGCCAAACTTGGAGCTACACCAGCTCAAACCCAACGTCATCAGGAAGCTGTTGAAGC
TTTCTTCAAGAAGATTGGTTTGGATTATGGCAAAGAAGTCGAATTCACAGCTTTCGTTA

ACGGATGGAAAGAACTGGCCAAACATGACTTGAAACTTTGGTCCCAAAACAAGAAAT
CTTTGATCCGCAATTGGGGAGAAGCTGTATTCGACATTTTCGACAAGGACGGAAGTGG
CTCAATCAGTTTGGACGAATGGAAAACATACGGAGGAATCTCTGGAATCTGTCCATCA
GACGAAGACGCTGAAAAGACCTTCAAACATTGCGATTTGGACAACAGTGGCAAACCTT
5 GATGTTGACGAGATGACCAGACAACATTTGGGATTCTGGTACACCTTGGACCCTAACG
CTGATGGTCTTTATGGCAACTTTGTCCCTTAAAAACTTTTTTTGCTGTAAATTCTTTACG
GGTTATTTTTTTCATAATTGTCATTTGATTTTAACTTTGTTTCGGAAAATGAAAAATATT
CTTTATTCAGAAAAAAAAAAAAAAAAAAAAAAAAAAAAA – 3'

This yields an amino acid sequence of (SEQ ID NO: 6):

10 MTDTASKYAVKLKTNFEDPKWVNRHKFMFNFLDINGNGKITLDEIVSKASDDICAKLGAT
PAQTQRHQEAVEAFFKKIGLDYGKEVEFPAFVNGWKELAKHDLKLWSQNKKSLIRNWGE
AVFDIFDKDGSISLDEWKTYGGISGICPSDEDAEKTFKHCDLDNSGKLDVDEMTRQHLG
FWYTLDPNADGLYGNFVP

These sequences are reproduced in the sequence listing.

15 **Brief description of the Figures**

- Figure 1: Figure 1 shows the plasmid map of the vector pTriplEX2-mtClytin.
- Figure 2: Figure 2 shows the plasmid map of the vector pcDNA3-mtClytin.
- Figure 3: Figure 3 shows the result of the bacterial expression of mtClytin and the bioluminescence activity of mtClytin following bacterial expression. (Y = RLU: relative light units; X = dilution; black bar = mtClytin; gray bar = control lysate).
20
- Figure 4: Figure 4 shows the result of the eukaryotic expression of mtClytin and the bioluminescence activity of mtClytin following expression in CHO cells. (Y = RLU: relative light units; X = ATP (logarithmic representation in mol/l)).
- Figure 5: Figure 5 shows the kinetic analysis of the bioluminescence of mtClytin. (Y = RLU: relative light units; X = time [seconds]).
25
- Figure 6: Figure 6 shows the kinetic analysis of the bioluminescence of obelin. (Y = RLU: relative light units; X = time [seconds]).
- Figure 7: Figure 7 shows the alignment of clytin and mtClytin at the amino acid level.
- Figure 8: Figure 8 shows the alignment of clytin and mtClytin at the nucleic acid level.

Figure 9: Figure 9 shows the alignment of clytin, mtClytin and clytin-2 at the amino acid level.

Examples

Example 1

The Clontech plasmid pTriplEx2 was used as vector for preparing the construct which is described below. The derivative of the vector was designated pTriplEx2-mtClytin. The vector pTriplEx2-
5 mtClytin was used for expressing mtClytin in bacterial systems.

Figure 1 shows the plasmid map of the vector pTriplEX2-mtClytin.

Example 2

The Clontech plasmid pcDNA3.1(+) was used as the vector for preparing the construct which is described below. The derivative of the vector was designated pcDNA3-mtClytin. The vector
10 pcDNA3-mtClytin was used for expressing mtClytin in eukaryotic systems.

Figure 2 shows the plasmid map of the vector pcDNA3-mtClytin.

Example 3

Bacterial expression

The bacterial expression was effected in the E. coli strain BL21(DE3) by transforming the bacteria
15 with the expression plasmids pTriplEX2-mtClytin and pTriplEX2. The transformed bacteria were incubated at 37°C for 3 hours in LB medium and the expression was induced for 4 hours by adding IPTG up to a final concentration of 1 mM. The induced bacteria were harvested by centrifugation, resuspended in 50 mM Tris/HCl (pH 9.0) + 5 mM EDTA and disrupted by ultrasonication. The
20 lysate was subsequently centrifuged at 13 000 rpm (16 000 ref) for 15 minutes and the supernatant removed. The supernatant (dilutions 1:5, 1:10; 1:20 and 1:50 with Tris/HCl pH 9.0)) was incubated with coelenterazine (10E-07 M coelenterazine in Tris/HCl pH 9.0) for 3 hours in the dark. The bioluminescence was measured in a luminometer directly after adding 5 mM calcium chloride. The measurement integration time was 40 seconds.

Figure 3 shows the results of measuring the bioluminescence of mtClytin in bacteria.

Example 4

Eukaryotic expression

Constitutive eukaryotic expression was effected in CHO cells by transfecting the cells with the expression plasmids pcDNA3-mtClytin and pcDNA3.1(+) in transient experiments. For this, 10 000 cells per well were plated out, in DMEM-F12 medium, on 96-well microtiter plates and the

plates were incubated overnight at 37°C. Transfection was effected using the Fugene 6 kits (Roche) in accordance with the manufacturer's instructions. The transfected cells were incubated overnight in DMEM-F12 medium at 37°C. The medium was then removed and replaced with 50 µl of coelenterazine (10E-07 M coelenterazine in PBS). The cells were incubated at 37°C for 3 hours and ATP (adenosine triphosphate) was then added to a final concentration of 1 µM. The measurement in a luminometer was started directly after the addition. The integration time was 1 second, with the total measurement time being 60 seconds.

Figure 4 shows the results of measuring the bioluminescence of mtClytin in CHO cells.

Example 5

10 BLAST

Result of a BLAST analysis of mtClytin at the amino acid level:

```
>emb|CAD87655.1| unnamed protein product [Clytia gregaria], Length
= 198, Score = 368 bits (945), Expect = e-101, Identities =
171/195 (87%), Positives = 182/195 (92%)

15 >sp|Q08121|CLYT_CLYGR Clytin precursor (Phialidin), pir||S28860
clytin - hydromedusa (Clytia gregarium), emb|CAA49754.1| clytin
[Clytia gregaria], gb|AAA28293.1| apoclytin, Length = 198, Score =
368 bits (945), Expect = e-101, Identities = 171/195 (87%),
Positives = 182/195 (92%)

20 >emb|CAD87658.1| unnamed protein product [synthetic construct],
Length = 198, Score = 367 bits (943), Expect = e-101, Identities
= 170/195 (87%), Positives = 182/195 (93%)

>sp|Q27709|OBL_OBELO Obelin precursor (OBL), pdb|1EL4|A Chain A,
Structure Of The Calcium-Regulated Photoprotein Obelin, Determined
25 By Sulfur Sas, gb|AAA67708.1| unnamed protein product, Length =
195, Score = 327 bits (837), Expect = 1e-88, Identities = 150/193
(77%), Positives = 170/193 (87%)

>emb|CAD87674.1| unnamed protein product [synthetic construct],
Length = 195, Score = 326 bits (835), Expect = 2e-88, Identities
30 = 149/193 (77%), Positives = 170/193 (87%)
```

```
>emb|CAD87672.1| unnamed protein product [synthetic construct],
Length = 195, Score = 325 bits (834), Expect = 3e-88, Identities
= 149/193 (77%), positives = 170/193 (87%)

>emb|CAD87673.1| unnamed protein product [synthetic construct],
5 Length = 195, Score = 325 bits (833), Expect = 4e-88, Identities
= 149/193 (77%), Positives = 170/193 (87%)

>pdb|1JF0|A Chain A, The Crystal Structure Of Obelin From Obelia
Geniculata At 1.82 A Resolution, gb|AAL86372.1|AF394688_1
apobobelin [Obelia geniculata], Length = 195, Score = 325
10 bits (833), Expect = 4e-88, Identities = 149/193 (77%), Positives
= 168/193 (86%)
```

Example 6

BLAST

Result of a BLAST analysis of mtClytin at the nucleic acid level:

```
15 >emb|AX702125.1| Sequence 23 from Patent WO03006497, Length = 597,
Score = 669 bits (348), Expect = 0.0, Identities = 504/582 (86%)

>emb|AX702119.1| Sequence 17 from Patent WO03006497, Length = 597,
Score = 669 bits (348), Expect = 0.0, Identities = 504/582 (86%)

>emb|X70221.1|CGCLYTIN C.gregaria mRNA for clytin, Length = 747,
20 Score = 669 bits (348), Expect = 0.0, Identities = 504/582 (86%)

>gb|L13247.1|CY1APOCLYT Clytia gregarium apoclytin mRNA, complete
cds, Length = 747, Score = 669 bits (348), Expect = 0.0,
Identities = 504/582 (86%)

>emb|AX702187.1| Sequence 85 from Patent WO03006497, Length = 597,
25 Score = 664 bits (345), Expect = 0.0, Identities = 503/582 (86%)

>emb|AX702185.1| Sequence 83 from Patent WO03006497, Length = 597,
Score = 664 bits (345), Expect = 0.0, Identities = 503/582 (86%)

>emb|AX702183.1| Sequence 81 from Patent WO03006497, Length = 597,
Score = 664 bits (345), Expect = 0.0, Identities = 503/582 (86%)

30 >emb|AX702181.1| Sequence 79 from Patent WO03006497, Length = 597,
Score = 664 bits (345), Expect = 0.0, Identities = 503/582 (86%)
```

>emb|AX702179.1| Sequence 77 from Patent WO03006497, Length = 597,
Score = 664 bits (345), Expect = 0.0, Identities = 503/582 (86%)

>emb|AX702131.1| Sequence 29 from Patent WO03006497, Length = 597,
Score = 664 bits (345), Expect = 0.0, Identities = 503/582 (86%)

5 >emb|AX702129.1| Sequence 27 from Patent WO03006497, Length = 597,
Score = 664 bits (345), Expect = 0.0, Identities = 503/582 (86%)

Example 7

Figure 7 shows the alignment of mtClytin with clytin (*Clytia gregaria*) at the nucleic acid level.

Example 8

10 Figure 8 shows the alignment of mtClytin with clytin (*Clytia gregaria*) at the amino acid level.

Example 9

Kinetic analysis of mtClytin

15 For the kinetic analysis of the bioluminescence of mtClytin, CHO cells were transiently transfected with pcDNA3-mtClytin or pcDNA-obelin or pcDNA3 (without any integrated cDNA). The transfection and measurement were carried out as described in Example 4. The readings were taken for a period of 60 seconds using an integration time of 1 second.

Figures 5 and 6 show the results of the kinetic analysis of mtClytin and obelin.

20 **Example 10**

MITOPROT Analysis

The computer program MITOPROT was used to analyze the mtClytin signal peptide (Claros et al., 1996). The following photoproteins were analyzed: obelin (Q27709), aequorin (P07164), clytin (Q08121) and mtClytin (SEQ ID NO: 2).

Results of the analyses:

Obelin:

Sequence name: OBELIN

5 Input sequence length : 195 aa

VALUES OF COMPUTED PARAMETERS

Net charge of query sequence : -11
Analysed region : 11
10 Number of basic residues in targeting sequence : 3
Number of acidic residues in targeting sequence : 0
Cleavagesite : not predictable
Cleaved sequence : -

15 HYDROPHOBIC SCALE USED

		GES	KD	GVH1	ECS
	H17	: -0.624	0.259	-0.308	0.295
	MesoH	: -1.573	-0.241	-0.642	0.060
	MuHd_075	: 14.019	3.641	4.408	1.523
20	MuHd_095	: 7.994	7.898	3.285	1.838
	MuHd_100	: 13.734	9.836	5.597	2.742
	MuHd_105	: 21.195	11.755	7.339	4.117
	Hmax_075	: -9.450	-2.800	-4.008	1.132
	Hmax_095	: -0.963	1.837	-1.971	1.103
25	Hmax_100	: 0.400	1.300	-1.942	2.240
	Hmax_105	: 10.617	6.067	0.733	3.127

PROBABILITY

of export to mitochondria: 0.1479

30 Aequorin:

Sequence name: AEQUORIN

Input sequence length : 196 aa

VALUES OF COMPUTED PARAMETERS

35 Net charge of query sequence : -13
Analysed region : 3
Number of basic residues in targeting sequence : 0
Number of acidic residues in targeting sequence : 0
Cleavage site : not predictable
40 Cleaved sequence : -

HYDROPHOBIC SCALE USED						
		GES	KD	GVH1	ECS	
	H17	:	0.006	0.794	-0.263	0.368
5	MesoH	:	-1.673	-0.382	-0.703	0.048
	MuHd_075	:	24.326	4.153	5.947	2.450
	MuHd_095	:	12.638	7.213	4.218	1.796
	MuHd_100	:	13.748	8.827	4.477	2.427
	MuHd_105	:	16.581	11.426	5.056	3.453
10	Hmax_075	:	0.438	0.233	-2.490	1.692
	Hmax_095	:	0.525	-1.400	-2.394	0.674
	Hmax_100	:	-0.100	-1.200	-2.292	1.550
	Hmax_105	:	0.500	-0.000	-2.164	1.540

15	PROBABILITY					
	of export to mitochondria: 0.0148					
	<u>Clytin:</u>					
	Sequence name: CLYTIN					
	Input sequence length : 198 aa					
20	-----					
	VALUES OF COMPUTED PARAMETERS					
	Net charge of query sequence : -9					
	Analysed region : 32					
	Number of basic residues in targeting sequence : 6					
25	Number of acidic residues in targeting sequence : 2					
	Cleavage site : not predictable					
	Cleaved sequence : -					

HYDROPHOBIC SCALE USED						
30		GES	KD	GVH1	ECS	
	H17	:	-0.429	0.341	-0.313	0.313
	MesoH	:	-1.778	-0.307	-0.718	0.053
	MuHd_075	:	32.928	17.509	7.351	5.708
	MuHd_095	:	30.874	20.344	9.074	5.834
35	MuHd_100	:	36.596	22.666	10.051	6.762
	MuHd_105	:	39.174	19.336	10.379	7.609
	Hmax_075	:	4.900	7.087	-1.223	3.684
	Hmax_095	:	13.600	10.100	1.251	4.390
	Hmax_100	:	14.000	12.600	1.601	5.060
40	Hmax_105	:	6.650	13.067	-0.468	3.920

PROBABILITY
of export to mitochondria: 0.2047

clytin-2:

5 Sequence name: CLYTIN-2
Input sequence length : 198 aa

VALUES OF COMPUTED PARAMETERS

Net charge of query sequence : -7
10 Analysed region : 16
Number of basic residues in targeting sequence : 3
Number of acidic residues in targeting sequence : 1
Cleavage site : not predictable
Cleaved sequence : -

15 -----

HYDROPHOBIC SCALE USED

		GES	KD	GVH1	ECS
	H17 :	-0.288	0.341	-0.213	0.313
	MesoH :	-1.519	-0.206	-0.681	0.081
20	MuHd_075 :	32.594	15.092	8.192	4.075
	MuHd_095 :	36.090	19.707	8.836	6.716
	MuHd_100 :	38.617	20.269	9.682	6.851
	MuHd_105 :	30.267	16.082	8.229	5.470
	Hmax_075 :	6.533	6.417	-0.793	2.508
25	Hmax_095 :	13.600	10.100	1.251	4.390
	Hmax_100 :	13.600	10.100	1.251	4.390
	Hmax_105 :	13.417	10.150	1.612	3.862

PROBABILITY
30 of export to mitochondria: 0.3974

mtClytin:

Sequence name: mtClytin
Input sequence length : 228 aa

35 VALUES OF COMPUTED PARAMETERS

Net charge of query sequence : -8
Analysed region : 34
Number of basic residues in targeting sequence : 6
Number of acidic residues in targeting sequence : 0
40 Cleavage site : 17

Cleaved sequence : MQRFTNRLLSMSALRA

HYDROPHOBIC SCALE USED

			GES	KD	GVH1	ECS
5	H17	:	-0.135	0.453	-0.343	0.309
	MesoH	:	-1.623	-0.215	-0.701	0.073
	MuHd_075	:	33.394	19.322	8.634	7.593
	MuHd_095	:	34.726	19.634	8.110	8.861
	MuHd_100	:	32.825	16.596	7.376	7.520
10	MuHd_105	:	28.005	19.893	7.410	7.865
	Hmax_075	:	16.683	17.733	2.851	5.763
	Hmax_095	:	13.125	13.388	2.299	4.314
	Hmax_100	:	8.300	11.500	1.845	3.830
	Hmax_105	:	1.700	9.500	-1.171	2.390

15 -----

PROBABILITY

of export to mitochondria: 0.9974

The probability of a translocation of the analyzed peptide into mitochondria increases as the calculated factor approaches 1.

- 20 The analysis of the protein sequences of obelin, aequorin, clytin, clytin-2 and mtClytin has shown that only mtClytin has the features of a protein which can be transported into mitochondria.

Example 11

Figure 9 shows the alignment of mtClytin, clytin (*Clytia gregaria*) and clytin-type2 at the amino acid level.

25

Literature/Patents

US 6,495,355

US 5,541,309

US 5,093,240

30 US-0908909

US 6,152,358

JP-0176125

GB-0024357

WO03006497

35 WO200168824

- Alam J, Cook JL.** Reporter genes: application to the study of mammalian gene transcription. *Anal Biochem.* 1990 Aug 1;188(2):245-54
- Altschul, Stephen F., Thomas L. Madden, Alejandro A. Schäffer, Jinghui Zhang, Zheng Zhang, Webb Miller, and David J. Lipman** (1997); Gapped BLAST and PSI-BLAST: a new generation of protein database search programs; *Nucleic Acids Res.* 25:3389-3402
- Chiesa A, Rapizzi E, Tosello V, Pinton P, de Virgilio M, Fogarty KE, Rizzuto R.** Recombinant aequorin and green fluorescent protein as valuable tools in the study of cell signalling. *Biochem J.* 2001 Apr 1;355(Pt 1):1-12.
- Claros, M.G., Vincens, P.** (1996); Computational method to predict mitochondrially imported proteins and their targeting sequences. *Eur. J. Biochem* 241, 779-786.
- Cullen Bryan R., Malim Michael H.,** Secreted placental alkaline phosphatase as a eukaryotic reporter gene. *Methods in Enzymology.* 216:362ff
- Fagan TF, Ohmiya Y, Blinks JR, Inouye S, Tsuji FI.** Cloning, expression and sequence analysis of cDNA for the Ca(2+)-binding photoprotein, mitrocomin. *FEBS Lett.* 1993 Nov 1;333(3):301-5
- Hastings, J.W. and Morin, J.G.** (1969) Comparative biochemistry of calcium-activated photoproteins from the ctenophore, *Mnemiopsis* and the coelenterates *Aequorea*, *Obelia*, and *Pelagia*. *Biol. Bull.* 137, 402.
- Haddock SH, Rivers TJ, Robison BH.** Can coelenterates make coelenterazine? Dietary requirement for luciferin in cnidarian bioluminescence. *Proc Natl Acad Sci U S A* 2001 Sep 25;98(20):11148-51
- Inouye S, Tsuji FI.** (1994) *Aequorea* green fluorescent protein. Expression of the gene and fluorescence characteristics of the recombinant protein. *FEBS Lett* 1994 Mar 21;341(2-3):277-80
- Inouye S, Tsuji FI.** Cloning and sequence analysis of cDNA for the Ca(2+)-activated photoprotein, clytin. *FEBS Lett.* 1993 Jan 11;315(3):343-6.
- Illarionov BA, Bondar VS, Illarionova VA, Vysotski ES.** Sequence of the cDNA encoding the Ca(2+)-activated photoprotein obelin from the hydroid polyp *Obelia longissima*. *Gene.* 1995 Feb 14;153(2):273-4.

- Jones K, Hibbert F, Keenan M.** Glowing jellyfish, luminescence and a molecule called coelenterazine. *Trends Biotechnol* 1999 Dec;17(12):477-81
- 5 **Johnson, F.H., Shimomura, O., Saiga, Y., Gershman, L.C., Reynolds, G.T., and Waters, J.R.**
(1962) Quantum efficiency of *Cypridina* luminescence, with a note on that of *Aequorea*. *J. Cell. Comp. Physiol.* 60, 85-103.
- 10 **Morin, J.G. and Hastings, J.W.** (1971) Biochemistry of the bioluminescence of colonial hydroids and other coelenterates. *J. Cell. Physiol.* 77, 305-311.
- Phillips GN.** Structure and dynamics of green fluorescent protein. *Curr Opin Struct Biol.* 1997 Dec;7(6):821-7
- 15 **Sambrook, J., Fritsch, E. Maniatis, T.** 1989, Molecular cloning. A laboratory manual Vol 1-3, Cold Spring Harbor, New York : Cold Spring Harbor Laboratory Press
- Shimomura O, Johnson FH.** Properties of the bioluminescent protein aequorin. *Biochemistry* 969 ct;8(10):3991-7
- 20 **Shimomura O.,** Bioluminescence in the sea: photoprotein systems. *Symp Soc Exp Biol.* 1985;39:351-72
- Shimomura O.** Isolation and properties of various molecular forms of aequorin. 25 *Biochem J.* 1986 Mar 1;234(2):271-7.
- Snowdowne KW, Borle AB.** Measurement of cytosolic free calcium in mammalian cells with aequorin. *Am J Physiol.* 1984 Nov;247(5 Pt 1):C396-408.
- 30 **Ward, W.W.** (1998) Biochemical and physical properties of green fluorescent protein. In: *Green Fluorescent Protein: Properties, Applications, and Protocols* (Chalfie, M. and Kain, S., eds) pp. 45-70. Wiley-Liss, Inc.
- Yang Te-Tuan, Sinai Parisa, Kitts Paul A. Kain Seven R.,** Quantification of gene expression with a secreted alkaline phosphatase reporter system. *Biotechnology.* 1997 23(6) 1110ff

Patent Claims

1. A nucleic acid molecule which is selected from the group consisting of
 - a) nucleic acid molecules which encode a polypeptide which contains the amino acid sequence disclosed by SEQ ID NO: 2;
 - 5 b) nucleic acid molecules which contain the sequence depicted by SEQ ID NO: 1;
 - c) nucleic acid molecules whose complementary strand hybridizes with a nucleic acid molecule from a) or b) under stringent conditions and which encode a polypeptide which exhibits the biological function of a photoprotein;
 - 10 d) nucleic acid molecules which differ from the nucleic acid molecules mentioned under c) due to the degeneracy of the genetic code;
 - e) nucleic acid molecules which exhibit a sequence homology with SEQ ID NO: 1 of at least 95% and encode a polypeptide which has the biological function of a photoprotein; and
 - 15 f) nucleic acid molecules which exhibit a sequence homology with SEQ ID NO: 1 of at least 65% and encode a polypeptide which has the biological function of a photoprotein.
2. A nucleic acid molecule which is selected from the group consisting of
 - a) nucleic acid molecules which encode a polypeptide which contains the amino acid sequence disclosed by SEQ ID NO: 3;
 - 20 b) nucleic acid molecules which contain the sequence depicted by SEQ ID NO: 4;
 - c) nucleic acid molecules whose complementary strand hybridizes with a nucleic acid molecule from a) or b) under stringent conditions and which encode a peptide which exhibits the biological function of a signal or leader peptide;
 - 25 d) nucleic acid molecules which differ from the nucleic acid molecules mentioned under c) due to the degeneracy of the genetic code;
 - e) nucleic acid molecules which exhibit a sequence homology with SEQ ID NO: 4 of at least 90% and encode a peptide which has the biological function of a signal or leader peptide; and

- f) nucleic acid molecules which exhibit a sequence homology with SEQ ID NO: 4 of at least 60% and encode a peptide which has the biological function of a signal or leader peptide.
3. A nucleic acid molecule which is selected from the group consisting of
- 5 a) nucleic acid molecules which encode a polypeptide which contains the amino acid sequence disclosed by SEQ ID NO: 6;
- b) nucleic acid molecules which contain the sequence depicted by SEQ ID NO: 5;
- c) nucleic acid molecules whose complementary strand hybridizes with a nucleic acid molecule from a) or b) under stringent conditions and which encode a polypeptide which exhibits the biological function of a photoprotein;
- 10 d) nucleic acid molecules which differ from the nucleic acid molecules mentioned under c) due to the degeneracy of the genetic code;
- e) nucleic acid molecules which exhibit a sequence homology with SEQ ID NO: 5 of at least 95% and encode a polypeptide which has the biological function of a photoprotein; and
- 15 f) nucleic acid molecules which exhibit a sequence homology with SEQ ID NO: 5 of at least 80% and encode a polypeptide which has the biological function of a photoprotein.
4. A nucleic acid as claimed in claim 1, 2 or 3 which contains a functional promoter 5' to the coding sequence.
- 20 5. A recombinant DNA or RNA vector which contains a nucleic acid as claimed in claim 4.
6. An organism which harbors a vector as claimed in claim 5.
- 25
7. A polypeptide which is encoded by a nucleic acid sequence as claimed in claim 1, 2 or 3.
8. A method for expressing the polypeptide as claimed in claim 7 in bacteria, viral systems, yeasts or eukaryotic cells or in *in-vitro* expression systems.

9. A method for purifying/isolating a photoprotein polypeptide as claimed in claim 7.
10. A peptide having more than 5 consecutive amino acids which is recognized immunologically by antibodies directed against the photoprotein mtClytin.
11. A peptide having more than 5 consecutive amino acids which is recognized immunologically by antibodies directed against the photoprotein clytin-2.
12. A peptide having more than 5 consecutive amino acids which is recognized immunologically by antibodies directed against the signal or leader peptide disclosed by SEQ ID NO: 3.
13. The use of a nucleic acid as claimed in claims 1 to 5 as a marker gene or reporter gene.
- 10 14. The use of a photoprotein as claimed in claim 7 as a label or reporter.
15. The use of a nucleic acid which contains the sequence depicted as SEQ ID NO: 4 as a signal or leader sequence.
16. The use of a peptide which contains the sequence depicted as SEQ ID NO: 3 as a signal or leader peptide.
- 15 17. The use as claimed in claim 15 or 16 for transporting a protein which is fused to the signal or leader peptide into cell organelles.
18. The use as claimed in claim 17, wherein the cell organelles are mitochondria or the endoplasmic reticulum (ER).
19. The use of the polypeptides as claimed in claim 7 as reporter proteins in searching for pharmacological active compounds.
- 20 20. The use of the nucleic acids as claimed in claims 1-3 as reporter genes in searching for pharmacological active compounds.

SEQUENCE LISTING

<110> Bayer AG, BHC

<120> Isolated photoprotein mtClytin, and the use thereof

<130> Le A 36 839

<160> 6

<170> PatentIn version 3.1

<210> 1

<211> 912

<212> DNA

<213> Clytia gregaria

<400> 1

```
gacagataaa aaattcactc cttagattat ttagtgaata agagaaaaaa aggataagaa      60
atcaagatgc aaaggtttac aaatcgtctt ctttccatgt cggctttacg tgcaagatca     120
agattgcaac gcacggcaaa ttttcacacc agcatactct tggctacaga ttcaaaatac     180
gcggtcaaac tcgatoctga ttttgcaa at ccaaaatgga tcaacagaca caaatattatg     240
ttcaactttt tggacataaa cggttaagggg aaaatcacat tagatgaaat cgtctccaaa     300
gcttcagacg acatttgtgc taaactggat gcaacaccag aacagaccaa acgtcaccag     360
gatgctgttg aagccttttt caagaaaatg ggcatggatt atggtaaaga agttgcattc     420
ccagaattta ttaagggatg ggaagagttg gccgaacacg acttggaact ctggtctcaa     480
aacaaaagta cattgatccg tgaatgggga gatgctgttt tcgacatttt cgacaaagac     540
gcaagtggct caatcagttt agacgaatgg aaggcttacg gacgaatctc tggaatctgt     600
ccatcagacg aagacgctga gaagacgttc aaacattgtg atttggacaa cagtggcaaa     660
cttgatgttg atgagatgac caggcaacat ttaggcttct ggtacacatt ggatccaact     720
tctgatggtc tttatggcaa ttttgttccc taagaagcgt tcagttaaaa acgctaaaca     780
ttgttcagtt gtaaaattat attcattttc atttcgtaaa attagtattt ataaatttgt     840
atcataaatt gtatccatgt ttagactaa ataagactcg gcaaaaaaaaa aaaaaaaaaa     900
aaaaaaaaaa aa                                                                912
```

<210> 2

<211> 228

<212> PRT

<213> Clytia gregaria

- 2 -

<400> 2

Met Gln Arg Phe Thr Asn Arg Leu Leu Ser Met Ser Ala Leu Arg Ala
 1 5 10 15
 Arg Ser Arg Leu Gln Arg Thr Ala Asn Phe His Thr Ser Ile Leu Leu
 20 25 30
 Ala Thr Asp Ser Lys Tyr Ala Val Lys Leu Asp Pro Asp Phe Ala Asn
 35 40 45
 Pro Lys Trp Ile Asn Arg His Lys Phe Met Phe Asn Phe Leu Asp Ile
 50 55 60
 Asn Gly Lys Gly Lys Ile Thr Leu Asp Glu Ile Val Ser Lys Ala Ser
 65 70 75 80
 Asp Asp Ile Cys Ala Lys Leu Asp Ala Thr Pro Glu Gln Thr Lys Arg
 85 90 95
 His Gln Asp Ala Val Glu Ala Phe Phe Lys Lys Met Gly Met Asp Tyr
 100 105 110
 Gly Lys Glu Val Ala Phe Pro Glu Phe Ile Lys Gly Trp Glu Glu Leu
 115 120 125
 Ala Glu His Asp Leu Glu Leu Trp Ser Gln Asn Lys Ser Thr Leu Ile
 130 135 140
 Arg Glu Trp Gly Asp Ala Val Phe Asp Ile Phe Asp Lys Asp Ala Ser
 145 150 155 160
 Gly Ser Ile Ser Leu Asp Glu Trp Lys Ala Tyr Gly Arg Ile Ser Gly
 165 170 175
 Ile Cys Pro Ser Asp Glu Asp Ala Glu Lys Thr Phe Lys His Cys Asp
 180 185 190
 Leu Asp Asn Ser Gly Lys Leu Asp Val Asp Glu Met Thr Arg Gln His
 195 200 205
 Leu Gly Phe Trp Tyr Thr Leu Asp Pro Thr Ser Asp Gly Leu Tyr Gly
 210 215 220
 Asn Phe Val Pro
 225

<210> 3

<211> 16

<212> PRT

<213> Clytia gregaria

<400> 3

Met Gln Arg Phe Thr Asn Arg Leu Leu Ser Met Ser Ala Leu Arg Ala
 1 5 10 15

<210> 4
 <211> 48
 <212> DNA
 <213> *Clytia gregaria*

<400> 4
 atgcaaaggt ttacaaatcg tcttctttcc atgtcggtt tacgtgca 48

<210> 5
 <211> 791
 <212> DNA
 <213> *Clytia gregaria*

<400> 5
 gatctcagct caacttgcaa taagtatcag atcaaatttt gcaactcaaa gcaaatcatc 60
 aacttcatca taatgactga cactgcttca aaatacgctg tcaaactcaa gaccaacttt 120
 gaagatccaa aatgggtcaa cagacacaaa tttatgttca actttttgga cattaacggc 180
 aacggaaaaa tcactttgga tgaaattgtc tccaaagctt cggatgacat ttgcgcaaaa 240
 cttggagcta caccagctca aacccaacgt catcaggaag ctgttgaagc tttcttcaag 300
 aagattgggt tggattatgg caaagaagtc gaattcccag ctttcgttaa cggatggaaa 360
 gaactggcca aacatgactt gaaactttgg tcccaaaaaca agaaatcttt gatccgcaat 420
 tggggagaag ctgtattcga cattttcgac aaggacggaa gtggctcaat cagtttggac 480
 gaatggaaaa catacggagg aatctctgga atctgtccat cagacgaaga cgctgaaaag 540
 accttcaaac attgcgattt ggacaacagt ggcaaacttg atgttgacga gatgaccaga 600
 caacatttgg gattctggta caccttggac cctaacgctg atggtcttta tggcaacttt 660
 gtcccttaaa aacttttttt gctgtaaatt ctttacgggt tattttttca taattgtcat 720
 ttgattttta ctttgtttcg gaaaatgaaa aatattcttt attcagaaaa aaaaaaaaaa 780
 aaaaaaaaaa a 791

<210> 6
 <211> 198
 <212> PRT
 <213> *Clytia gregaria*

<400> 6
 Met Thr Asp Thr Ala Ser Lys Tyr Ala Val Lys Leu Lys Thr Asn Phe
 1 5 10 15
 Glu Asp Pro Lys Trp Val Asn Arg His Lys Phe Met Phe Asn Phe Leu
 20 25 30
 Asp Ile Asn Gly Asn Gly Lys Ile Thr Leu Asp Glu Ile Val Ser Lys
 35 40 45
 Ala Ser Asp Asp Ile Cys Ala Lys Leu Gly Ala Thr Pro Ala Gln Thr
 50 55 60
 Gln Arg His Gln Glu Ala Val Glu Ala Phe Phe Lys Lys Ile Gly Leu
 65 70 75 80
 Asp Tyr Gly Lys Glu Val Glu Phe Pro Ala Phe Val Asn Gly Trp Lys

85	90	95
Glu Leu Ala Lys His Asp Leu Lys Leu Trp Ser Gln Asn Lys Lys Ser		
100	105	110
Leu Ile Arg Asn Trp Gly Glu Ala Val Phe Asp Ile Phe Asp Lys Asp		
115	120	125
Gly Ser Gly Ser Ile Ser Leu Asp Glu Trp Lys Thr Tyr Gly Gly Ile		
130	135	140
Ser Gly Ile Cys Pro Ser Asp Glu Asp Ala Glu Lys Thr Phe Lys His		
145	150	155
Cys Asp Leu Asp Asn Ser Gly Lys Leu Asp Val Asp Glu Met Thr Arg		
165	170	175
Gln His Leu Gly Phe Trp Tyr Thr Leu Asp Pro Asn Ala Asp Gly Leu		
180	185	190
Tyr Gly Asn Phe Val Pro		
195		