


Description

Recent studies have confirmed that theophylline is probably the most effective bronchodilator which can be administered parenterally and orally for the treatment of asthma. Unfortunately, the therapeutic range for theophylline is fairly narrow and its toxic effects result largely from its action on the central nervous system, cardiovascular system, and gastrointestinal tract. These undesirable pharmacological activities as well as the desired bronchodilation effect may be related to theophylline's ability to inhibit competitively phosphodiesterase (PDE), an enzyme that degrades cyclic AMP. It is desirable, therefore, to discover selective inhibitors of cyclic AMP phosphodiesterase which possess a beneficial bronchodilation effect but lack the adverse reactions of theophylline.

This invention relates to the discovery that certain 5,6-disubstituted-benzothiophene-2-carboxylic acids are potent phosphodiesterase inhibitors which possess bronchodilator activity. 5,6-Dimethoxybenzothiophene-2-carboxylic acid was prepared first by Bew and Clemo, *J. Chem. Soc.*, 1314 (1953) and later by Campaigne and Kreighbaum (*J. Org. Chem.*, 26, 359 (1961)). See also, *Tetrahedron*, 25 (14), 2781 (1969). In *J. Org. Chem.*, 26, 1327—1329 (1961), Campaigne teaches that certain amide derivatives possess analgesic and diuretic activity. Campaigne also teaches 5,6-diethoxybenzothiophene-2-carboxylic acid (*J. Org. Chem.*, 26, 1326 (1961)) although no pharmaceutical utility is disclosed.

In accordance with the invention, a compound represented by Formula (I):



or a pharmaceutically-acceptable salt thereof, in which R_1 is hydrogen or C_1 — C_7 alkyl and R_2 is C_1 — C_7 alkyl or R_1 and R_2 are hydrogen, is useful as an agent capable of producing bronchodilation in mammals. More specifically, this invention provides compounds for treating a mammal suffering from or susceptible to asthma. In such treatment, the mammal is administered a chemotherapeutically-effective amount of a compound of Formula (I), or a pharmaceutically-acceptable salt thereof.

According to a further aspect of the invention, there is provided a pharmaceutical formulation which comprises a compound of Formula (I) as described above associated with a pharmaceutically-acceptable carrier or diluent therefor.

The preferred compounds of this invention are those in which each of R_1 and R_2 is, independently, methyl, ethyl, or propyl. Especially preferred is a compound in which R_1 is ethyl. The most preferred compounds are 5,6-diethoxybenzothiophene-2-carboxylic acid and 5-ethoxy-6-propyloxybenzothiophene-2-carboxylic acid or a pharmaceutically-acceptable salt thereof.

The term " C_1 — C_7 alkyl" refers to straight and branched aliphatic radicals of one to seven carbon atoms such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, amyl, isoamyl, sec-amyl, sec-isoamyl (1,2-dimethylpropyl), tert-amyl (1,1-dimethylpropyl), hexyl, isohexyl (4-methylpentyl), sec-hexyl (1-methylpentyl), 2-methylpentyl, 3-methylpentyl, 1,1-dimethylbutyl, 2,2-dimethylbutyl, 3,3-dimethylbutyl, 1,2-dimethylbutyl, 1,3-dimethylbutyl, 1,2,2-trimethylpropyl, 1,1,2-trimethylpropyl, heptyl, isoheptyl (5-methylhexyl), sec-heptyl (1-methylhexyl), 2,2-dimethylpentyl, 3,3-dimethylbutyl, 4,4-dimethylpentyl, 1,2-dimethylpentyl, 1,3-dimethylpentyl, 1,4-dimethylpentyl, 1,2,3-trimethylbutyl, 1,1,2-trimethylbutyl, or 1,1,3-trimethylbutyl. Straight chain radicals are preferred and methyl, propyl, and especially ethyl are most preferred.

The pharmaceutically-acceptable base addition salts of this invention (i.e. those useful in the chemophary of warm-blooded animals) include salts derived from inorganic bases, such as ammonium and alkali and alkaline earth metal hydroxides, carbonates, bicarbonates, and the like, as well as salts derived from non-toxic basic organic amines, such as aliphatic and aromatic amines, aliphatic diamines, or hydroxy alkylamines. Such bases useful in preparing the salts of this invention may include, for example, sodium hydroxide, sodium acetate, ammonium hydroxide, potassium carbonate, sodium bicarbonate, calcium hydroxide, methyl amine, diethyl amine, ethylenediamine, cyclohexylamine, or ethanolamine. The potassium and sodium salt forms are particularly preferred.

The compounds of Formula (I) are prepared by known methods. These include the cyclization of a 3,4-dialkoxyphenylpropionic acid chloride with thionyl chloride (Bonnin, *et al.*, *Aust. J. Chem.*, 32 (4), 833 (1979), the reaction of a 4,5-dialkoxy-2-nitrobenzaldehyde with methyl thioglycolate followed by hydrolysis of the resulting methyl esters of compounds (I) (Beck, *J. Org. Chem.*, 37 (21), 3224 (1972)), the condensation of a 4,5-alkoxy-2-mercaptobenzaldehyde with a haloacetic acid followed by cyclization (Bew and Clemo, *J. Chem. Soc.*, 1314 (1953)), or preferably the cyclization of a β -(3,4-disubstituted-phenyl)- α -mercaptoacrylic acid (Campaigne and Kreighbaum, *J. Org. Chem.*, 26, 359 (1961); *id.*, 26, 1326 (1961); Chakrabarti, *et al.*, *Tetrahedron*, 25 (14), 2781 (1969)).

In addition, compounds of Formula (I) can also be prepared from other compounds of Formula (I). A dialkoxy compound of Formula (I) can be esterified first by methods known in the art and then mono-dealkylated to provide 5-hydroxy-6-alkoxy-derivatives or di-dealkylated to provide the corresponding 5,6-dihydroxy compound after hydrolysis of the ester group by any of the usual methods. These dealkylations can be performed using methods. These dealkylations can be performed using methods known in the art

such as boron/tribromide in ethylene dichloride, a mixture of methanesulfonic acid/methionine, or ethanethiol/aluminum chloride in ethylene chloride. The resulting hydroxy or dihydroxy ester intermediates can be hydrolyzed to the carboxylic acid compounds of this invention and can be converted also to other compounds of this invention by mono- or di-alkylation in the usual way with the appropriate alkyl halide preferably in a nonreactive solvent containing an acid scavenger, followed by hydrolysis of the ester group.

The compounds may be administered by various routes including the oral, rectal, transdermal, subcutaneous, intravenous, intramuscular, inhalation, or intranasal routes, being employed usually in the form of a pharmaceutical formulation. Such formulations are prepared in a manner well-known in the pharmaceutical art and comprise at least one active compound. Accordingly, the invention includes a pharmaceutical formulation which comprises as an active ingredient, a compound of Formula (I) or a pharmaceutically-acceptable salt thereof, associated with a pharmaceutically-acceptable carrier or diluent therefor.

In making the formulations of the present invention, the active ingredient usually will be mixed with a carrier, or diluted by a carrier, or enclosed within a carrier which may be in the form of a capsule, sachet, paper or other container. When the carrier serves as a diluent, it may be a solid, semi-solid or liquid material which acts as a vehicle, excipient or medium for the active ingredient. Thus, the formulation can be in the form of tablets, pills, powders, lozenges, sachets, cachets, elixirs, suspensions, emulsions, solutions, syrups, aerosols (as a solid or in a liquid medium), ointments containing for example up to 10% by weight of the active compound, soft and hard gelatin capsules, suppositories, sterile injectable solutions and sterile packaged powders.

Examples of suitable carriers are lactose, dextrose, sucrose, sorbitol, mannitol, starches, gum acacia, calcium phosphate, alginates, tragacanth, gelatin, calcium silicate, microcrystalline cellulose, polyvinylpyrrolidone, cellulose, water, syrup, methyl cellulose, methyl- and propyl-hydroxybenzoates, talc, magnesium stearate or mineral oil. The formulations can include additionally lubricating agents, wetting agents, emulsifying and suspending agents, preserving agents, sweetening agents or flavoring agents. The formulations may be formulated to provide quick, sustained or delayed release of the active ingredient after administration to the patient.

Preferably, the compositions are formulated in a unit dosage form, each dosage containing from about 5 to 500 mg (from 5 to 50 mg in the case of parenteral or inhalation administration, and from 25 to 500 mg in the case of oral or rectal administration) of a compound of Formula (I). The term "unit dosage form" refers to physically discrete units suitable as unitary dosages for human subjects and other mammals, each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect, in association with the required pharmaceutical carrier.

The active compounds are effective over a wide dosage range. For example, dosages per day will fall normally within the range of about 0.5 to 300 mg/kg. In the treatment of adult humans, the range of about 1 to 50 mg/kg, in single or divided doses, is preferred. However, the amount of the compound actually administered will be determined by a physician, in the light of the relevant circumstances including the condition to be treated, the choice of compound to be administered, the chosen route of administration, the age, weight, and response of the individual patient, and the severity of the patient's symptoms. The above dosage ranges, therefore, are not intended to limit the scope of the invention in any way.

Preferred pharmaceutical forms of the present invention are capsules, tablets, suppositories, injectable solutions, creams and ointments. Especially preferred are formulations for inhalation or insufflation application, such as an aerosol, and for oral ingestion.

The following examples further illustrate the preparation of the starting materials, intermediates, and compounds used in this invention. The examples are illustrative only and are not intended to limit the scope of the invention in any way.

Example 1

5,6-Diethoxybenzothiophene-2-carboxylic acid

A. Preparation of 5-(3,4-diethoxybenzylidene)rhodanine.

A solution of 10.0 g of 3,4-diethoxybenzaldehyde, 6.86 g of rhodanine, and 14.5 g of fused sodium acetate in 70 ml of acetic acid was heated to reflux for one hour. The hot mixture was poured into one liter of water with stirring. The resulting precipitate was collected by filtration and dried. Recrystallization from ethanol provided 10.22 g of the titled intermediate.

B. Preparation of β -(3,4-diethoxyphenyl)- α -mercaptoacrylic acid.

A solution of 33.0 g of 5-(3,4-diethoxybenzylidene)rhodanine and 23.4 g of sodium hydroxide in 148 ml. of water was heated to 70°C for 30 minutes. The solution was cooled and filtered and the filtrate was acidified by pouring into excess cold 10% hydrochloric acid. The precipitate was collected by filtration affording 25 g of the desired intermediate which was used without further purification.

C. Preparation of 5,6-diethoxybenzothiophene-2-carboxylic acid.

Two grams of β -(3,4-diethoxyphenyl)- α -mercaptoacrylic acid were dissolved in 68.5 ml of dioxane. To the solution were added 2.74 g of iodine. The solution was heated at 60—70°C for 22 hours. The reaction

was then poured into 550 ml of water, decolorized with 11.0 g of sodium bisulfite, and stirred vigorously for several minutes. The crude product was collected by filtration and dissolved in approximately 15 ml of a warm 10% sodium hydroxide solution. The alkaline solution was treated with decolorizing carbon and filtered. The solution was allowed stand at refrigerator temperature overnight. The resulting pink crystals were collected and dissolved in water. Dilute hydrochloric acid was added and the resulting precipitate was collected to give 0.59 g of the desired product. Two recrystallizations from 95% ethanol gave white needles, m.p. 243—245°C.

Analysis: $C_{13}H_{14}O_4S$;

Calc.: C, 58.63; H, 5.30; S, 12.04;

Found: C, 58.84; H, 5.03; S, 12.05.

Examples 2—8

Following the general procedure of Example 1, the following benzothiophene-2-carboxylic acids were prepared from rhodanine and the corresponding benzaldehyde derivatives. Yields are expressed as the percent molar yield from the benzaldehyde.

2. 5-Methoxy-6-ethoxybenzothiophene-2-carboxylic acid, m.p. 236—238°C., 12% yield.

Analysis: $C_{12}H_{12}O_4S$;

Calc.: C, 57.13; H, 4.79; S, 12.71;

Found: C, 57.22; H, 5.05; S, 12.45.

3. 5-Methoxy-6-butyloxybenzothiophene-2-carboxylic acid, 10.5% yield. Mass spectrum: $M^+ = 280$.

Analysis: $C_{14}H_{18}O_4S$;

Calc.: C, 59.98; H, 5.75; S, 11.44;

Found: C, 60.10; H, 5.87; S, 11.22.

4. 5-Methoxy-6-hexyloxybenzothiophene-2-carboxylic acid, m.p. 165—166°C., 13% yield.

Analysis: $C_{16}H_{20}O_4S$;

Calc.: C, 62.31; H, 6.54; S, 10.40;

Found: C, 62.51; H, 6.28; S, 10.13.

5. 5-Methoxy-6-heptyloxybenzothiophene-2-carboxylic acid, m.p. 163—164°C., 9.4% yield.

Analysis: $C_{17}H_{22}O_4S$;

Calc.: C, 63.33; H, 6.88;

Found: C, 63.11; H, 6.96.

6. 5,6-Dihydroxybenzothiophene-2-carboxylic acid, 5.5% yield. Mass spectrum: $M^+ = 210$.

Analysis: $C_9H_6O_4S$;

Calc.: C, 51.43; H, 2.88; S, 15.25;

Found: C, 51.68; H, 3.11; S, 15.01.

7. 5-Hydroxy-6-methoxybenzothiophene-2-carboxylic acid, m.p. 190°C. (decomposition), 3.4% yield. Mass spectrum: $M^+ = 224$.

Analysis: $C_{10}H_8O_4S$;

Calc.: C, 53.56; H, 3.60; S, 14.30;

Found: C, 53.79; H, 3.65; S, 14.09.

8. 5,6-Dimethoxybenzothiophene-2-carboxylic acid, m.p. 257—358°C., 7% yield.

Analysis: $C_{11}H_{10}O_4S$;

Calc.: C, 55.45; H, 4.23; S, 13.46;

Found: C, 55.72; H, 4.25; S, 12.72.

Example 9

5-Ethoxy-6-propyloxybenzothiophene-2-carboxylic acid

A. Preparation of methyl 5-ethoxy-6-propyloxybenzothiophene-2-carboxylate.

A solution of 0.42 g of methyl 5-ethoxy-6-hydroxybenzothiophene-2-carboxylate, 0.25 g of potassium carbonate, a catalytic amount of potassium iodide, and 0.31 g of propyl iodide in methyl ethyl ketone was heated to reflux for 48 hours. The hot solution was filtered and the filtrate was evaporated *in vacuo* to provide the desired intermediate which was used in the subsequent reaction without purification.

B. Preparation of 5-ethoxy-6-propyloxybenzothiophene-2-carboxylic acid.

A solution of 0.45 g of methyl 5-ethoxy-6-propyloxybenzothiophene-2-carboxylate and 0.34 g of potassium hydroxide in ethanol was heated to reflux for two hours. The solution was poured into water, acidified with hydrochloric acid, and extracted with ethyl acetate. The ethyl acetate layer was extracted with 1N sodium hydroxide. The alkaline solution was acidified with hydrochloric acid and was then extracted with ethyl acetate. The organic extract was dried over sodium sulfate and evaporated *in vacuo* to provide

the title product, m.p. 192—194°C., in an overall molar yield of 36% from 5-ethoxy-6-hydroxybenzothiophene-2-carboxylic acid.

Analysis: $C_{14}H_{16}O_4S$;

Calc.: C, 59.98; H, 5.75;

5 Found: C, 59.78; H, 5.99.

Examples 10—11

Following the general procedure of Example 10, the following compounds were prepared from the corresponding hydroxybenzothiophene-2-carboxylic acid and appropriate alkyl halide. Yields are expressed as the percent molar yield from the hydroxybenzothiophene-2-carboxylic acids.

11. 5-Ethoxy-6-methoxybenzothiophene-2-carboxylic acid, m.p. 233—234.5°C., 50% yield.

Analysis: $C_{12}H_{12}O_4S$;

Calc.: C, 57.13; H, 4.79;

15 Found: C, 56.97; H, 4.81.

12. 5-Propyloxy-6-methoxybenzothiophene-2-carboxylic acid, m.p. 172—173°C., 50% yield. Mass spectrum: $M^+ = 266$.

Analysis: $C_{13}H_{14}O_4S$;

Calc.: C, 58.63; H, 5.30; S, 12.04;

20 Found: C, 57.04; H, 5.05; S, 11.85.

The following formulation examples may employ as active compounds any of the pharmaceutical compounds of the invention.

Example 12

Hard gelatin capsules are prepared using the following ingredients:

	Quantity (mg./capsule)
30 Active compound	250
Starch	200
35 Magnesium stearate	10

The above ingredients are mixed and filled in 460 mg. quantities into hard galatin capsules.

Example 13

40 A tablet formula is prepared using the ingredients below:

	Quantity (mg./tablet)
45 Active compound	250
Cellulose, microcrystalline	400
Silicon dioxide, fumed	10
50 Magnesium stearate	5

The components are blended and compressed to form tablets each weighing 665 mg.

Example 14

An aerosol solution is prepared containing the following components:

		Weight %
5	Active ingredient	0.25
	Ethanol	30.00
10	Propellant 11 (trichlorofluoromethane)	10.25
	Propellant 12 (Dichlorodifluoromethane)	29.75
15	Propellant 114 (Dichlorotetrafluoroethane)	29.75

20 The active compound is dissolved in the ethanol and the solution is added to the propellant 11, cooled to -30°C and transferred to a filling device. The required amount is then fed to a container and further filled with the pre-mixed propellants 12 and 114 by means of the cold-filled method or pressure-filled method. The valve units are then fitted to the container.

Example 15

25 Tablets, each containing 60 mg of active ingredient, are made up as follows:

	Active ingredient	60 mg
	Starch	45 mg
30	Microcrystalline cellulose	35 mg
	Polyvinylpyrrolidone (as 10% solution in water)	4 mg
35	Sodium carboxymethyl starch	4.5 mg
	Magnesium stearate	0.5 mg
40	Talc	1 mg
	Total	150 mg

50 The active ingredient, starch and cellulose are passed through a No. 45 mesh U.S. sieve and mixed thoroughly. The solution of polyvinylpyrrolidone is mixed with the resultant powders which are then passed through a No. 14 mesh U.S. sieve. The granules so produced are dried at $50-60^{\circ}\text{C}$ and passed through a No. 18 mesh U.S. sieve. The sodium carboxymethyl starch, magnesium stearate and talc, previously passed through a No. 60 mesh U.S. sieve, are then added to the granules which, after mixing, are compressed on a tablet machine to yield tablets each weighing 150 mg.

Example 16

Capsules each containing 80 mg of medicament, are made as follows:

	Active ingredient	80 mg
55	Starch	59 mg
	Microcrystalline cellulose	59 mg
60	Magnesium stearate	2 mg
	Total	200 mg

65 The active ingredient, cellulose, starch and magnesium stearate are blended, passed through a No. 45 mesh U.S. sieve, and filled into hard gelatin capsules in 200 mg quantities.

Example 17

Suppositories, each containing 225 mg of active ingredient, are made as follows:

5	Active ingredient	225 mg
	Unsaturated or saturated fatty acid glycerides to	2,000 mg

The active ingredient is passed through a No. 60 mesh U.S. sieve and suspended in the fatty acid glycerides previously melted using the minimum heat necessary. The mixture is then poured into a suppository mold of nominal 2 g capacity and allowed to cool.

Example 18

Suspensions, each containing 50 mg of medicament per 5 ml dose, are made as follows:

15	Active ingredient	50 mg
	Sodium carboxymethyl cellulose	50 mg
20	Sugar	1 g
	Methyl paraben	0.05 mg
	Propyl paraben	0.03 mg
25	Flavor	q.v
	Color	q.v
30	Purified water to	5 ml

The medicament is passed through a No. 45 mesh U.S. sieve and mixed with the sodium carboxymethylcellulose, sugar, and a portion of the water to form a suspension. The parabens, flavor and color are dissolved and diluted with some of the water and added, with stirring. Sufficient water is then added to produce the required volume.

A compound of Formula (I), or a pharmaceutically-acceptable salt thereof, possesses useful pharmaceutical properties. The compounds are useful as inhibitors of the enzyme phosphodiesterase and are also useful in blocking anaphylactic responses and bronchoconstriction. The compounds are therefore useful in treating mammals suffering from or susceptible to asthma. This biological activity was demonstrated in the following test system.

Following the general procedure taught by Andersson, *Brit. J. Pharmacol.*, 69, 467 (1980), mixed sex Hartley strain guinea pigs (250—300 g) were sensitized to ovalbumin by a single injection of 1 mcg. of ovalbumin mixed with 50 mg of aluminum hydroxide per animal. These animals were used 21 to 26 days later for aerosol challenge with ovalbumin.

Anaphylaxis was induced by exposure to an aerosol of 10 mg of ovalbumin per milliliter of water for 10 minutes delivered by a Tri-R venturi nebulizer, particle size 2 to 5 micron diameter, at a delivery rate of 0.4 ml of solution per minute. The guinea pigs were placed in specially constructed chamber for exposure to the aerosol. The aerosol was introduced from the bottom of the chamber and exhausted at the top. A baffle above the chamber inlet port provided even distribution of the aerosol throughout the chamber. The animals were supported above the inlet on a wire mesh disc. The chamber was under slight negative pressure to enhance the exhaust of the aerosol which was passed through a calcium chloride trap followed by a super-cooled condenser trap.

Groups of five guinea pigs were treated with the compounds of this invention, clinical anti-asthmatic agents, or placebo (vehicle only) two hours prior to the aerosol challenge. All the animals also received 3 mg/kg pyrilamine orally two hours prior to the aerosol challenge in order to blunt the histamine component of the anaphylaxis. The compounds were administered orally as a suspension in 10% acacia.

Throughout the ten minute period of aerosol challenge, the animals were observed for the symptomology of convulsive cough, convulsive collapse, and death as described in Herxheimer, *J. Physiology*, 117, 251 (1952). The number of animals that responded to the aerosol challenge for each of the above parameters as well as the time for the response to occur were recorded. The data were analyzed by comparing the severity index (the sum of the number of animals that coughed, collapsed, and/or died) between the treated and placebo group of guinea pigs. Thus, the maximum severity index for each group of five guinea pigs is 15. On the average, the severity index for the placebo group was 10—12. The percent inhibition of anaphylaxis was determined by the following formula:

$$\% \text{ inhibition} = \left[1 - \frac{S_d}{S_p} \right] \times 100$$

5 where S_d is the severity index for the drug treated animals and S_p is the severity index of the placebo treated animals. The results of these experiments are summarized in Table 1.

TABLE 1


Antigen induced anaphylaxis in guinea pigs

Compound of Example No.	Percent Inhibition*
1	93%
2	55%
3	23%
4	18%
5	38%
6	40%
7	63%
8	30%
9	90%
10	5%
11	30%

* Average of two experiments. Compounds were given orally at a dose of 50 mg/kg.

30 **Claims for the Contracting States: BE CH DE FR GB IT LI LU NL SE**

1. A pharmaceutical formulation which comprises as an active ingredient, a compound of Formula (I):



in which R_1 is hydrogen or C_1-C_7 alkyl and R_2 is C_1-C_7 alkyl,

or R_1 and R_2 are hydrogen,

or a pharmaceutically-acceptable salt thereof, associated with a pharmaceutically-acceptable carrier or diluent therefor.

2. A formulation as claimed in claim 1 in which R_1 is ethyl.

3. A formulation as claimed in claim 2 in which the compound is 5,6-diethoxybenzothiophene-2-carboxylic acid or a pharmaceutically-acceptable salt thereof.

4. A formulation as claimed in claim 3 in which the compound is in the form of the sodium salt.

5. A formulation as claimed in claim 2 in which the compound is 5-ethoxy-6-propylbenzothiophene-2-carboxylic acid or a pharmaceutically-acceptable salt thereof.

6. A formulation as claimed in claim 3 or claim 4 which is formulated for administration by inhalation or orally.

7. A compound of Formula (I) referred to in claim 1 or a pharmaceutically-acceptable salt thereof for use in chemotherapy.


8. A compound as claimed in claim 7 for therapeutic use as bronchodilator in a mammal.

9. 5,6-diethoxybenzothiophene-2-carboxylic acid or 5-ethoxy-6-propyloxybenzothiophene-2-carboxylic acid, or a pharmaceutically-acceptable salt of either, for use as a bronchodilator.

10. The sodium salt of 5,6-diethoxybenzothiophene-2-carboxylic acid for use as a bronchodilator.

Claims for the Contracting State: AT

1. A method of preparing a formulation for use as a bronchodilator which comprises mixing, diluting or enclosing a compound of Formula (I)



$$\% \text{ inhibition} = \left[1 - \frac{S_d}{S_p} \right] \times 100$$

5 where S_d is the severity index for the drug treated animals and S_p is the severity index of the placebo treated animals. The results of these experiments are summarized in Table 1.

TABLE 1

Antigen induced anaphylaxis in guinea pigs

	Compound of Example No.	Percent Inhibition*
15	1	93%
	2	55%
	3	23%
	4	18%
	5	38%
20	6	40%
	7	63%
	8	30%
	9	90%
	10	5%
25	11	30%

* Average of two experiments. Compounds were given orally at a dose of 50 mg/kg.

30 Claims for the Contracting States: BE CH DE FR GB IT LI LU NL SE

1. A pharmaceutical formulation which comprises as an active ingredient, a compound of Formula (I):



- in which R_1 is hydrogen or C_1-C_7 alkyl and R_2 is C_1-C_7 alkyl,
 40 or R_1 and R_2 are hydrogen,
 or a pharmaceutically-acceptable salt thereof, associated with a pharmaceutically-acceptable carrier or diluent therefor.
2. A formulation as claimed in claim 1 in which R_1 is ethyl.
 3. A formulation as claimed in claim 2 in which the compound is 5,6-diethoxybenzothiophene-2-
 - 45 carboxylic acid or a pharmaceutically-acceptable salt thereof.
 4. A formulation as claimed in claim 3 in which the compound is in the form of the sodium salt.
 5. A formulation as claimed in claim 2 in which the compound is 5-ethoxy-6-propylbenzothiophene-2-
 - carboxylic acid or a pharmaceutically-acceptable salt thereof.
 6. A formulation as claimed in claim 3 or claim 4 which is formulated for administration by inhalation or
 - 50 orally.
 7. A compound of Formula (I) referred to in claim 1 or a pharmaceutically-acceptable salt thereof for use in chemotherapy.
 8. A compound as claimed in claim 7 for therapeutic use as bronchodilator in a mammal.
 9. 5,6-diethoxybenzothiophene-2-carboxylic acid or 5-ethoxy-6-propyloxybenzothiophene-2-carboxylic
 - 55 acid, or a pharmaceutically-acceptable salt of either, for use as a bronchodilator.
 10. The sodium salt of 5,6-diethoxybenzothiophene-2-carboxylic acid for use as a bronchodilator.

Claims for the Contracting State: AT

60 1. A method of preparing a formulation for use as a bronchodilator which comprises mixing, diluting or enclosing a compound of Formula (I)



in which R_1 is hydrogen or C_1-C_7 alkyl and R_2 is C_1-C_7 alkyl, or R_1 and R_2 are hydrogen, or a pharmaceutically-acceptable salt thereof, with or in a carrier.

2. A method of claim 1 wherein R_1 is ethyl.

3. A method of claim 2 in which the compound is 5,6-diethoxybenzothiophene-2-carboxylic acid or a pharmaceutically-acceptable salt thereof.


4. A method of claim 3 wherein the salt is the sodium salt.

5. A method of any one of claims 1 to 4 wherein the formulation is in the form of a unit dosage form.

6. A method of claim 5 wherein the formulation comprises a capsule, a tablet, an aerosol, a suppository or a suspension.

Patentansprüche für die Vertragsstaaten: DE CH DE FR GB IT LI LU NL SE

1. Pharmazeutische Formulierung, dadurch gekennzeichnet, daß sie eine Verbindung der Formel (I)



worin R_1 Wasserstoff oder C_1-C_7 -Alkyl ist und R_2 für C_1-C_7 -Alkyl steht oder R_1 und R_2 Wasserstoff sind, oder ein pharmazeutisch annehmbares Salz hiervon als Wirkstoff in Verbindung mit einem pharmazeutisch annehmbaren Träger oder Verdünnungsmittel hierfür enthält.

2. Formulierung nach Anspruch 1, dadurch gekennzeichnet, daß R_1 Ethyl ist.

3. Formulierung nach Anspruch 2, dadurch gekennzeichnet, daß die Verbindung 5,6-Diethoxybenzothiophen-2-carbonsäure oder ein pharmazeutisch annehmbares Salz hiervon ist.

4. Formulierung nach Anspruch 3, dadurch gekennzeichnet, daß die Verbindung ein Natriumsalz ist.

5. Formulierung nach Anspruch 2, dadurch gekennzeichnet, daß die Verbindung 5-Ethoxy-6-propyloxybenzothiophen-2-carbonsäure oder ein pharmazeutisch annehmbares Salz hiervon ist.

6. Formulierung nach Anspruch 3 oder 4, dadurch gekennzeichnet, daß sie die Form einer durch Inhalation oder auf oralem Weg verabreichbaren Arzneiform hat.

7. Verwendung einer Verbindung der im Anspruch 1 angegebenen Formel (I) oder eines pharmazeutisch annehmbaren Salzes hiervon in der Chemotherapie.


8. Verwendung einer Verbindung nach Anspruch 7 als bronchodilatorisches Therapeutikum bei Säugetieren.

9. Verwendung von 5,6-Diethoxybenzothiophen-2-carbonsäure oder 5-Ethoxy-6-propyloxybenzothiophen-2-carbonsäure oder einem pharmazeutisch annehmbaren Salz hiervon als Bronchodilator.

10. Verwendung des Natriumsalzes von 5,6-Diethoxybenzothiophen-2-carbonsäure als Bronchodilator.

Patentansprüche für den Vertragsstaat: AT

1. Verfahren zur Herstellung einer als Bronchodilator verwendbaren Formulierung, dadurch gekennzeichnet, daß eine Verbindung der Formel (I)



worin R_1 Wasserstoff oder C_1-C_7 -Alkyl ist und R_2 für C_1-C_7 -Alkyl steht oder R_1 und R_2 Wasserstoff sind, oder ein pharmazeutisch annehmbares Salz hiervon mit einem Träger vermischt und/oder verdünnt oder in einen Träger eingeschlossen wird.

2. Verfahren nach Anspruch 1, dadurch gekennzeichnet, daß R_1 Ethyl ist.

3. Verfahren nach Anspruch 2, dadurch gekennzeichnet, daß die Verbindung 5,6-Diethoxybenzothiophen-2-carbonsäure oder ein pharmazeutisch annehmbares Salz hiervon ist.


4. Verfahren nach Anspruch 3, dadurch gekennzeichnet, daß das Salz das Natriumsalz ist.

5. Verfahren nach irgend einem der Ansprüche 1 bis 4, dadurch gekennzeichnet, daß die Formulierung die Form einer Einheitsdosierungsform hat.

6. Verfahren nach Anspruch 5, dadurch gekennzeichnet, daß die Formulierung eine Kapsel, eine Tablette, ein Aerosol, ein Suppositorium oder eine Suspension ist.

Revendications pour les Etats contractants: DE CH DE FR GB IT LI LU NL SE

1. Une formulation pharmaceutique qui comprend comme ingrédient actif un dérivé de formule (I):



dans laquelle:

R_1 représente un atome d'hydrogène ou un radical alkyle en C_1-C_7 ; et
 R_2 représente un radical alkyle en C_1-C_7 , ou R_1 et R_2 représentent un atome d'hydrogène, ou un de leurs sels pharmaceutiquement acceptables, associé à un véhicule ou diluant pharmaceutiquement acceptable de ce dérivé.

2. Une formulation selon la revendication 1, dans laquelle R_1 représente le radical éthyle.

3. Une formulation selon la revendication 2, dans laquelle le dérivé est l'acide 5,6-diéthoxybenzothiophène-2-carboxylique ou un de ses sels pharmaceutiquement acceptables.

4. Une formulation selon la revendication 3, dans laquelle le dérivé se trouve sous la forme de son sel de sodium.

5. Une formulation selon la revendication 2, dans laquelle le dérivé est l'acide 5-éthoxy-6-propyloxybenzothiophène-2-carboxylique ou un de ses sels pharmaceutiquement acceptables.

6. Une formulation selon la revendication 3 ou 4, qui est formulée en vue de l'administration par inhalation ou par voie orale.

7. Un dérivé de formule (I) selon la revendication 1, utilisable en chimiothérapie.


8. Un dérivé selon la revendication 7, destiné à l'utilisation thérapeutique comme bronchodilatateur chez un mammifère.

9. Acide 5,6-diéthoxybenzothiophène-2-carboxylique ou acide 5-éthoxy-6-propyloxybenzothiophène-2-carboxylique ou un sel pharmaceutiquement acceptable de l'un ou l'autre de ces composés, destiné à l'utilisation comme bronchodilatateur.

10. Le sel de sodium de l'acide 5,6-diéthoxybenzothiophène-2-carboxylique destiné à l'utilisation comme bronchodilatateur.

Revendications pour l'Etat contractant: AT

1. Un procédé de préparation d'une formulation utilisable comme bronchodilatateur, qui consiste à mélanger, diluer ou inclure un dérivé de formule (I):



dans laquelle:

R_1 représente un atome d'hydrogène ou un radical alkyle en C_1-C_7 ; et
 R_2 représente un radical alkyle en C_1-C_7 , ou R_1 et R_2 représentent un atome d'hydrogène, ou un de leurs sels pharmaceutiquement acceptables, en présence ou dans un support.

2. Un procédé selon la revendication 1, dans lequel R_1 représente le radical éthyle.

3. Un procédé selon la revendication 2, dans lequel le dérivé est l'acide 5,6-diéthoxybenzothiophène-2-carboxylique ou un de ses sels pharmaceutiquement acceptables.

4. Un procédé selon la revendication 3, dans lequel le sel est le sel de sodium.

5. Un procédé selon l'une quelconque des revendications 1 à 4, dans lequel la formulation se trouve sous la forme de dosage unitaire.

6. Un procédé selon la revendication 5 dans lequel la formulation comprend une gélule, une pastille, un aérosol, un suppositoire ou une suspension.

