

CATIONIC LIPIDS FOR USE IN LIPID NANOPARTICLES

BACKGROUND

Technical Field

The present disclosure generally relates to novel cationic lipids that can be used
5 in combination with other lipid components, such as neutral lipids, cholesterol and
polymer conjugated lipids, to form lipid nanoparticles to facilitate the intracellular
delivery of therapeutic agents, such as nucleic acids (*e.g.*, oligonucleotides, messenger
RNA), both *in vitro* and *in vivo*.

Description of the Related Art

10 There are many challenges associated with the delivery of nucleic acids to affect
a desired response in a biological system. Nucleic acid based therapeutics have
enormous potential but there remains a need for more effective delivery of nucleic acids
to appropriate sites within a cell or organism in order to realize this potential.
Therapeutic nucleic acids include, *e.g.*, messenger RNA (mRNA), antisense
15 oligonucleotides, ribozymes, DNazymes, plasmids, immune stimulating nucleic acids,
antagomir, antimir, mimic, supermir, and aptamers. Some nucleic acids, such as
mRNA or plasmids, can be used to effect expression of specific cellular products as
would be useful in the treatment of, for example, diseases related to a deficiency of a
protein or enzyme. The therapeutic applications of translatable nucleotide delivery are
20 extremely broad as constructs can be synthesized to produce any chosen protein
sequence, whether or not indigenous to the system. The expression products of the
nucleic acid can augment existing levels of protein, replace missing or non-functional
versions of a protein, or introduce new protein and associated functionality in a cell or
organism.
25 Some nucleic acids, such as miRNA inhibitors, can be used to effect expression
of specific cellular products that are regulated by miRNA as would be useful in the
treatment of, for example, diseases related to deficiency of protein or enzyme. The
therapeutic applications of miRNA inhibition are extremely broad as constructs can be

synthesized to inhibit one or more miRNA that would in turn regulate the expression of mRNA products. The inhibition of endogenous miRNA can augment its downstream target endogenous protein expression and restore proper function in a cell or organism as a means to treat disease associated to a specific miRNA or a group of miRNA.

5 Other nucleic acids can down-regulate intracellular levels of specific mRNA and, as a result, down-regulate the synthesis of the corresponding proteins through processes such as RNA interference (RNAi) or complementary binding of antisense RNA. The therapeutic applications of antisense oligonucleotide and RNAi are also extremely broad, since oligonucleotide constructs can be synthesized with any
10 nucleotide sequence directed against a target mRNA. Targets may include mRNAs from normal cells, mRNAs associated with disease-states, such as cancer, and mRNAs of infectious agents, such as viruses. To date, antisense oligonucleotide constructs have shown the ability to specifically down-regulate target proteins through degradation of the cognate mRNA in both *in vitro* and *in vivo* models. In addition, antisense
15 oligonucleotide constructs are currently being evaluated in clinical studies.


 However, two problems currently face the use of oligonucleotides in therapeutic contexts. First, free RNAs are susceptible to nuclease digestion in plasma. Second, free RNAs have limited ability to gain access to the intracellular compartment where the relevant translation machinery resides. Lipid nanoparticles formed from cationic
20 lipids with other lipid components, such as neutral lipids, cholesterol, PEG, PEGylated lipids, and oligonucleotides have been used to block degradation of the RNAs in plasma and facilitate the cellular uptake of the oligonucleotides.

 There remains a need for improved cationic lipids and lipid nanoparticles for the delivery of oligonucleotides. Preferably, these lipid nanoparticles would provide
25 optimal drug:lipid ratios, protect the nucleic acid from degradation and clearance in serum, be suitable for systemic or local delivery, and provide intracellular delivery of the nucleic acid. In addition, these lipid-nucleic acid particles should be well-tolerated and provide an adequate therapeutic index, such that patient treatment at an effective dose of the nucleic acid is not associated with unacceptable toxicity and/or risk to the
30 patient. The present disclosure provides these and related advantages.

BRIEF SUMMARY

In brief, the present disclosure provides lipid compounds, including stereoisomers, pharmaceutically acceptable salts or tautomers thereof, which can be used alone or in combination with other lipid components such as neutral lipids, charged lipids, steroids (including for example, all sterols) and/or their analogs, and/or polymer conjugated lipids to form lipid nanoparticles for the delivery of therapeutic agents. In some instances, the lipid nanoparticles are used to deliver nucleic acids such as antisense and/or messenger RNA. Methods for use of such lipid nanoparticles for treatment or prevention (e.g., vaccination) of various diseases or conditions, such as those caused by infectious entities and/or insufficiency of a protein, are also provided.

In one embodiment, compounds having the following structure (I) are provided:



or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof, wherein a, b, c, d, G¹, G², L¹, L², R^{1a}, R^{1b}, R^{2a}, R^{2b}, R^{3a}, R^{3b}, R^{4a}, R^{4b}, R⁵, R⁶, R⁷, R⁸ and X are as defined herein.

Lipid nanoparticles (LNPs) comprising one or more of the compounds of structure (I) and a therapeutic agent, and pharmaceutical compositions comprising the same, are also provided. In some embodiments, the nanoparticles further comprise one or more components selected from neutral lipids, charged lipids, steroids, and polymer conjugated lipids. Such LNPs are useful for delivery of the therapeutic agent, for example for treatment of a disease or vaccination against a viral pathogen.

These and other aspects of the disclosure will be apparent upon reference to the following detailed description.

DETAILED DESCRIPTION

In the following description, certain specific details are set forth in order to provide a thorough understanding of various embodiments of the disclosure. However, one skilled in the art will understand that the disclosure may be practiced without these
5 details.

The present disclosure is based, in part, upon the discovery of novel cationic (amino) lipids that provide advantages when used in lipid nanoparticles for the *in vivo* delivery of an active or therapeutic agent such as a nucleic acid into a cell of a mammal. In particular, embodiments of the present disclosure provide nucleic acid-lipid
10 nanoparticle compositions comprising one or more of the novel cationic lipids described herein that provide increased activity of the nucleic acid and improved tolerability of the compositions *in vivo*, resulting in a significant increase in the therapeutic index as compared to nucleic acid-lipid nanoparticle compositions previously described. In other embodiments, the disclosed lipids, and lipid
15 nanoparticles comprising the same, have increased safety and/or tolerability when used for delivery of active agents, such as nucleic acids.

In particular embodiments, the present disclosure provides novel cationic lipids that enable the formulation of improved compositions for the *in vitro* and *in vivo* delivery of mRNA and/or other oligonucleotides. In some embodiments, these
20 improved lipid nanoparticle compositions are useful for expression of protein encoded by mRNA. In other embodiments, these improved lipid nanoparticles compositions are useful for upregulation of endogenous protein expression by delivering miRNA inhibitors targeting one specific miRNA or a group of miRNA regulating one target mRNA or several mRNA. In other embodiments, these improved lipid nanoparticle
25 compositions are useful for down-regulating (*e.g.*, silencing) the protein levels and/or mRNA levels of target genes. In some other embodiments, the lipid nanoparticles are also useful for delivery of mRNA and plasmids for expression of transgenes. In yet other embodiments, the lipid nanoparticle compositions are useful for inducing a pharmacological effect resulting from expression of a protein, *e.g.*, increased production
30 of red blood cells through the delivery of a suitable erythropoietin mRNA, or protection

against infection through delivery of mRNA encoding for a suitable antigen or antibody.

The lipid nanoparticles and compositions of embodiments of the present disclosure may be used for a variety of purposes, including the delivery of encapsulated or associated (*e.g.*, complexed) therapeutic agents such as nucleic acids to cells, both *in vitro* and *in vivo*. Accordingly, embodiments of the present disclosure provide methods of treating or preventing diseases or disorders in a subject in need thereof by contacting the subject with a lipid nanoparticle that encapsulates or is associated with a suitable therapeutic agent, wherein the lipid nanoparticle comprises one or more of the novel cationic lipids described herein.

As described herein, embodiments of the lipid nanoparticles of the present disclosure are particularly useful for the delivery of nucleic acids, including, *e.g.*, mRNA, antisense oligonucleotide, plasmid DNA, microRNA (miRNA), miRNA inhibitors (antagomirs/antimirs), messenger-RNA-interfering complementary RNA (micRNA), DNA, multivalent RNA, dicer substrate RNA, complementary DNA (cDNA), etc. Therefore, the lipid nanoparticles and compositions of certain embodiments of the present disclosure may be used to induce expression of a desired protein both *in vitro* and *in vivo* by contacting cells with a lipid nanoparticle comprising one or more novel cationic lipids described herein, wherein the lipid nanoparticle encapsulates or is associated with a nucleic acid that is expressed to produce the desired protein (*e.g.*, a messenger RNA or plasmid encoding the desired protein) or inhibit processes that terminate expression of mRNA (*e.g.*, miRNA inhibitors). In certain embodiments, the protein expressed by the nucleic acid is an antigen, and the LNPs thus induce an immune response (*e.g.*, vaccination). Alternatively, the lipid nanoparticles and compositions of embodiments of the present disclosure may be used to decrease the expression of target genes and proteins both *in vitro* and *in vivo* by contacting cells with a lipid nanoparticle comprising one or more novel cationic lipids described herein, wherein the lipid nanoparticle encapsulates or is associated with a nucleic acid that reduces target gene expression (*e.g.*, an antisense oligonucleotide or small interfering RNA (siRNA)). The lipid nanoparticles and compositions of embodiments of the present disclosure may also be used for co-delivery of different nucleic acids (*e.g.*,

mRNA and plasmid DNA) separately or in combination, such as may be useful to provide an effect requiring co-localization of different nucleic acids (*e.g.*, mRNA encoding for a suitable gene modifying enzyme and DNA segment(s) for incorporation into the host genome).

5 Nucleic acids for use with embodiments of this disclosure may be prepared according to any available technique. For mRNA, the primary methodology of preparation is, but not limited to, enzymatic synthesis (also termed *in vitro* transcription) which currently represents the most efficient method to produce long sequence-specific mRNA. *In vitro* transcription describes a process of template-
10 directed synthesis of RNA molecules from an engineered DNA template comprised of an upstream bacteriophage promoter sequence (*e.g.*, including but not limited to that from the T7, T3 and SP6 coliphage) linked to a downstream sequence encoding the gene of interest. Template DNA can be prepared for *in vitro* transcription from a number of sources with appropriate techniques which are well known in the art
15 including, but not limited to, plasmid DNA and polymerase chain reaction amplification (see Linpinsel, J.L and Conn, G.L., General protocols for preparation of plasmid DNA template and Bowman, J.C., Azizi, B., Lenz, T.K., Ray, P., and Williams, L.D. in RNA *in vitro* transcription and RNA purification by denaturing PAGE in Recombinant and *in vitro* RNA syntheses Methods v. 941 Conn G.L. (ed), New York, N.Y. Humana Press,
20 2012).

 Transcription of the RNA occurs *in vitro* using the linearized DNA template in the presence of the corresponding RNA polymerase and adenosine, guanosine, uridine, and cytidine ribonucleoside triphosphates (rNTPs) under conditions that support polymerase activity while minimizing potential degradation of the resultant mRNA
25 transcripts. *In vitro* transcription can be performed using a variety of commercially available kits including, but not limited to RiboMax Large Scale RNA Production System (Promega), MegaScript Transcription kits (Life Technologies), as well as with commercially available reagents including RNA polymerases and rNTPs. The methodology for *in vitro* transcription of mRNA is well known in the art. (see, *e.g.*,
30 Losick, R., 1972, In vitro transcription, Ann Rev Biochem v.41 409-46; Kamakaka, R. T. and Kraus, W. L. 2001. In Vitro Transcription. Current Protocols in Cell Biology.

2:11.6:11.6.1–11.6.17; Beckert, B. And Masquida, B.,(2010) Synthesis of RNA by In Vitro Transcription in RNA in Methods in Molecular Biology v. 703 (Neilson, H. Ed), New York, N.Y. Humana Press, 2010; Brunelle, J.L. and Green, R., 2013, Chapter Five – In vitro transcription from plasmid or PCR-amplified DNA, Methods in
5 Enzymology v. 530, 101-114; all of which are incorporated herein by reference).

The desired *in vitro* transcribed mRNA is then purified from the undesired components of the transcription or associated reactions (including unincorporated rNTPs, protein enzyme, salts, short RNA oligos, etc.). Techniques for the isolation of the mRNA transcripts are well known in the art. Well known procedures include
10 phenol/chloroform extraction or precipitation with either alcohol (ethanol, isopropanol) in the presence of monovalent cations or lithium chloride. Additional, non-limiting examples of purification procedures which can be used include size exclusion chromatography (Lukavsky, P.J. and Puglisi, J.D., 2004, Large-scale preparation and purification of polyacrylamide-free RNA oligonucleotides, RNA v.10, 889-893), silica-
15 based affinity chromatography and polyacrylamide gel electrophoresis (Bowman, J.C., Azizi, B., Lenz, T.K., Ray, P., and Williams, L.D. in RNA in vitro transcription and RNA purification by denaturing PAGE in Recombinant and in vitro RNA syntheses Methods v. 941 Conn G.L. (ed), New York, N.Y. Humana Press, 2012). Purification can be performed using a variety of commercially available kits including, but not
20 limited to SV Total Isolation System (Promega) and In Vitro Transcription Cleanup and Concentration Kit (Norgen Biotek).

Furthermore, while reverse transcription can yield large quantities of mRNA, the products can contain a number of aberrant RNA impurities associated with undesired polymerase activity which may need to be removed from the full-length
25 mRNA preparation. These include short RNAs that result from abortive transcription initiation as well as double-stranded RNA (dsRNA) generated by RNA-dependent RNA polymerase activity, RNA-primed transcription from RNA templates and self-complementary 3' extension. It has been demonstrated that these contaminants with dsRNA structures can lead to undesired immunostimulatory activity through interaction
30 with various innate immune sensors in eukaryotic cells that function to recognize specific nucleic acid structures and induce potent immune responses. This in turn, can

dramatically reduce mRNA translation since protein synthesis is reduced during the innate cellular immune response. Therefore, additional techniques to remove these dsRNA contaminants have been developed and are known in the art including but not limited to scaleable HPLC purification (see, *e.g.*, Kariko, K., Muramatsu, H., Ludwig, J. And Weissman, D., 2011, Generating the optimal mRNA for therapy: HPLC purification eliminates immune activation and improves translation of nucleoside-modified, protein-encoding mRNA, Nucl Acid Res, v. 39 e142; Weissman, D., Pardi, N., Muramatsu, H., and Kariko, K., HPLC Purification of *in vitro* transcribed long RNA in Synthetic Messenger RNA and Cell Metabolism Modulation in Methods in Molecular Biology v.969 (Rabinovich, P.H. Ed), 2013). HPLC purified mRNA has been reported to be translated at much greater levels, particularly in primary cells and *in vivo*.

A significant variety of modifications have been described in the art which are used to alter specific properties of *in vitro* transcribed mRNA, and improve its utility. These include, but are not limited to modifications to the 5' and 3' termini of the mRNA. Endogenous eukaryotic mRNA typically contain a cap structure on the 5'-end of a mature molecule which plays an important role in mediating binding of the mRNA Cap Binding Protein (CBP), which is in turn responsible for enhancing mRNA stability in the cell and efficiency of mRNA translation. Therefore, highest levels of protein expression are achieved with capped mRNA transcripts. The 5'-cap contains a 5'-5'-triphosphate linkage between the 5'-most nucleotide and guanine nucleotide. The conjugated guanine nucleotide is methylated at the N7 position. Additional modifications include methylation of the ultimate and penultimate most 5'-nucleotides on the 2'-hydroxyl group.

Multiple distinct cap structures can be used to generate the 5'-cap of *in vitro* transcribed synthetic mRNA. 5'-capping of synthetic mRNA can be performed co-transcriptionally with chemical cap analogs (*i.e.*, capping during *in vitro* transcription). For example, the Anti-Reverse Cap Analog (ARCA) cap contains a 5'-5'-triphosphate guanine-guanine linkage where one guanine contains an N7 methyl group as well as a 3'-O-methyl group. However, up to 20% of transcripts remain uncapped during this co-transcriptional process and the synthetic cap analog is not identical to the 5'-cap

structure of an authentic cellular mRNA, potentially reducing translatability and cellular stability. Alternatively, synthetic mRNA molecules may also be enzymatically capped post-transcriptionally. These may generate a more authentic 5'-cap structure that more closely mimics, either structurally or functionally, the endogenous 5'-cap which have enhanced binding of cap binding proteins, increased half-life and reduced susceptibility to 5' endonucleases and/or reduced 5' decapping. Numerous synthetic 5'-cap analogs have been developed and are known in the art to enhance mRNA stability and translatability (see, *e.g.*, Grudzien-Nogalska, E., Kowalska, J., Su, W., Kuhn, A.N., Slepnev, S.V., Darynkiewicz, E., Sahin, U., Jemielity, J., and Rhoads, R.E., Synthetic mRNAs with superior translation and stability properties in Synthetic Messenger RNA and Cell Metabolism Modulation in Methods in Molecular Biology v.969 (Rabinovich, P.H. Ed), 2013).

On the 3'-terminus, a long chain of adenine nucleotides (poly-A tail) is normally added to mRNA molecules during RNA processing. Immediately after transcription, the 3' end of the transcript is cleaved to free a 3' hydroxyl to which poly-A polymerase adds a chain of adenine nucleotides to the RNA in a process called polyadenylation. The poly-A tail has been extensively shown to enhance both translational efficiency and stability of mRNA (see Bernstein, P. and Ross, J., 1989, Poly (A), poly (A) binding protein and the regulation of mRNA stability, Trends Bio Sci v. 14 373-377; Guhaniyogi, J. And Brewer, G., 2001, Regulation of mRNA stability in mammalian cells, Gene, v. 265, 11-23; Dreyfus, M. And Regnier, P., 2002, The poly (A) tail of mRNAs: Bodyguard in eukaryotes, scavenger in bacteria, Cell, v.111, 611-613).

Poly (A) tailing of *in vitro* transcribed mRNA can be achieved using various approaches including, but not limited to, cloning of a poly (T) tract into the DNA template or by post-transcriptional addition using Poly (A) polymerase. The first case allows *in vitro* transcription of mRNA with poly (A) tails of defined length, depending on the size of the poly (T) tract, but requires additional manipulation of the template. The latter case involves the enzymatic addition of a poly (A) tail to *in vitro* transcribed mRNA using poly (A) polymerase which catalyzes the incorporation of adenine residues onto the 3' termini of RNA, requiring no additional manipulation of the DNA template, but results in mRNA with poly(A) tails of heterogeneous length. 5'-capping

and 3'-poly (A) tailing can be performed using a variety of commercially available kits including, but not limited to Poly (A) Polymerase Tailing kit (EpiCenter), mMESSAGE mMACHINE T7 Ultra kit and Poly (A) Tailing kit (Life Technologies) as well as with commercially available reagents, various ARCA caps, Poly (A) polymerase, etc.

5 In addition to 5' cap and 3' poly adenylation, other modifications of the *in vitro* transcripts have been reported to provide benefits as related to efficiency of translation and stability. It is well known in the art that pathogenic DNA and RNA can be recognized by a variety of sensors within eukaryotes and trigger potent innate immune responses. The ability to discriminate between pathogenic and self DNA and RNA has
10 been shown to be based, at least in part, on structure and nucleoside modifications since most nucleic acids from natural sources contain modified nucleosides. In contrast, *in vitro* synthesized RNA lacks these modifications, thus rendering it immunostimulatory which in turn can inhibit effective mRNA translation as outlined above. The introduction of modified nucleosides into *in vitro* transcribed mRNA can be used to
15 prevent recognition and activation of RNA sensors, thus mitigating this undesired immunostimulatory activity and enhancing translation capacity (see, *e.g.*, Kariko, K. And Weissman, D. 2007, Naturally occurring nucleoside modifications suppress the immunostimulatory activity of RNA: implication for therapeutic RNA development, Curr Opin Drug Discov Devel, v.10 523-532; Pardi, N., Muramatsu, H., Weissman, D.,
20 Kariko, K., In vitro transcription of long RNA containing modified nucleosides in Synthetic Messenger RNA and Cell Metabolism Modulation in Methods in Molecular Biology v.969 (Rabinovich, P.H. Ed), 2013; Kariko, K., Muramatsu, H., Welsh, F.A., Ludwig, J., Kato, H., Akira, S., Weissman, D., 2008, Incorporation of Pseudouridine Into mRNA Yields Superior Nonimmunogenic Vector With Increased Translational
25 Capacity and Biological Stability, Mol Ther v.16, 1833-1840). The modified nucleosides and nucleotides used in the synthesis of modified RNAs can be prepared monitored and utilized using general methods and procedures known in the art. A large variety of nucleoside modifications are available that may be incorporated alone or in combination with other modified nucleosides to some extent into the *in vitro* transcribed
30 mRNA (see, *e.g.*, US 2012/0251618). *In vitro* synthesis of nucleoside-modified mRNA

has been reported to have reduced ability to activate immune sensors with a concomitant enhanced translational capacity.

Other components of mRNA which can be modified to provide benefit in terms of translatability and stability include the 5' and 3' untranslated regions (UTR).

- 5 Optimization of the UTRs (favorable 5' and 3' UTRs can be obtained from cellular or viral RNAs), either both or independently, have been shown to increase mRNA stability and translational efficiency of *in vitro* transcribed mRNA (see, *e.g.*, Pardi, N., Muramatsu, H., Weissman, D., Kariko, K., In vitro transcription of long RNA containing modified nucleosides in Synthetic Messenger RNA and Cell Metabolism
10 Modulation in Methods in Molecular Biology v.969 (Rabinovich, P.H. Ed), 2013).

In addition to mRNA, other nucleic acid payloads may be used for this disclosure. For oligonucleotides, methods of preparation include but are not limited to chemical synthesis and enzymatic, chemical cleavage of a longer precursor, *in vitro* transcription as described above, etc. Methods of synthesizing DNA and RNA

- 15 nucleotides are widely used and well known in the art (see, *e.g.*, Gait, M. J. (ed.) Oligonucleotide synthesis: a practical approach, Oxford [Oxfordshire], Washington, D.C.: IRL Press, 1984; and Herdewijn, P. (ed.) Oligonucleotide synthesis: methods and applications, Methods in Molecular Biology, v. 288 (Clifton, N.J.) Totowa, N.J.: Humana Press, 2005; both of which are incorporated herein by reference).

- 20 For plasmid DNA, preparation for use with embodiments of this disclosure commonly utilizes, but is not limited to, expansion and isolation of the plasmid DNA *in vitro* in a liquid culture of bacteria containing the plasmid of interest. The presence of a gene in the plasmid of interest that encodes resistance to a particular antibiotic (penicillin, kanamycin, etc.) allows those bacteria containing the plasmid of interest to
25 selectively grow in antibiotic-containing cultures. Methods of isolating plasmid DNA are widely used and well known in the art (see, *e.g.*, Heilig, J., Elbing, K. L. and Brent, R., (2001), Large-Scale Preparation of Plasmid DNA, Current Protocols in Molecular Biology, 41:II:1.7:1.7.1–1.7.16; Rozkov, A., Larsson, B., Gillström, S., Björnstedt, R. and Schmidt, S. R., (2008), Large-scale production of endotoxin-free plasmids for
30 transient expression in mammalian cell culture, Biotechnol. Bioeng., 99: 557–566; and US 6,197,553 B1). Plasmid isolation can be performed using a variety of commercially

available kits including, but not limited to Plasmid Plus (Qiagen), GenJET plasmid MaxiPrep (Thermo), and PureYield MaxiPrep (Promega) kits as well as with commercially available reagents.

Various exemplary embodiments of the cationic lipids of the present disclosure, lipid nanoparticles and compositions comprising the same, and their use to deliver active (*e.g.*, therapeutic agents), such as nucleic acids, to modulate gene and protein expression, are described in further detail below.

As used herein, the following terms have the meanings ascribed to them unless specified otherwise.

Unless the context requires otherwise, throughout the present specification and claims, the word “comprise” and variations thereof, such as, “comprises” and “comprising” are to be construed in an open and inclusive sense, that is, as “including, but not limited to.”

Reference throughout this specification to “one embodiment” or “an embodiment” means that a particular feature, structure or characteristic described in connection with the embodiment is included in at least one embodiment of the present disclosure. Thus, the appearances of the phrases “in one embodiment” or “in an embodiment” in various places throughout this specification are not necessarily all referring to the same embodiment. Furthermore, the particular features, structures, or characteristics may be combined in any suitable manner in one or more embodiments.

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as is commonly understood by one of skill in the art to which this disclosure belongs. As used in the specification and claims, the singular form “a,” “an” and “the” include plural references unless the context clearly dictates otherwise.

The phrase “induce expression of a desired protein” refers to the ability of a nucleic acid to increase expression of the desired protein. To examine the extent of protein expression, a test sample (*e.g.*, a sample of cells in culture expressing the desired protein) or a test mammal (*e.g.*, a mammal such as a human or an animal) model such as a rodent (*e.g.*, mouse) or a non-human primate (*e.g.*, monkey) model is contacted with a nucleic acid (*e.g.*, nucleic acid in combination with a lipid of the present disclosure). Expression of the desired protein in the test sample or test animal is

compared to expression of the desired protein in a control sample (*e.g.*, a sample of cells in culture expressing the desired protein) or a control mammal (*e.g.*, a mammal such as a human or an animal) model such as a rodent (*e.g.*, mouse) or non-human primate (*e.g.*, monkey) model that is not contacted with or administered the nucleic acid. When the desired protein is present in a control sample or a control mammal, the expression of a desired protein in a control sample or a control mammal may be assigned a value of 1.0. In particular embodiments, inducing expression of a desired protein is achieved when the ratio of desired protein expression in the test sample or the control mammal to the level of desired protein expression in the control sample or the control mammal is greater than 1, for example, about 1.1, 1.5, 2.0, 5.0 or 10.0. When a desired protein is not present in a control sample or a control mammal, inducing expression of a desired protein is achieved when any measurable level of the desired protein in the test sample or the test mammal is detected. One of ordinary skill in the art will understand appropriate assays to determine the level of protein expression in a sample, for example dot blots, northern blots, in situ hybridization, ELISA, immunoprecipitation, enzyme function, and phenotypic assays, or assays based on reporter proteins that can produce fluorescence or luminescence under appropriate conditions.

The phrase “inhibiting expression of a target gene” refers to the ability of a nucleic acid to silence, reduce, or inhibit the expression of a target gene. To examine the extent of gene silencing, a test sample (*e.g.*, a sample of cells in culture expressing the target gene) or a test mammal (*e.g.*, a mammal such as a human or an animal) model such as a rodent (*e.g.*, mouse) or a non-human primate (*e.g.*, monkey) model is contacted with a nucleic acid that silences, reduces, or inhibits expression of the target gene. Expression of the target gene in the test sample or test animal is compared to expression of the target gene in a control sample (*e.g.*, a sample of cells in culture expressing the target gene) or a control mammal (*e.g.*, a mammal such as a human or an animal) model such as a rodent (*e.g.*, mouse) or non-human primate (*e.g.*, monkey) model that is not contacted with or administered the nucleic acid. The expression of the target gene in a control sample or a control mammal may be assigned a value of 100%. In particular embodiments, silencing, inhibition, or reduction of expression of a target

gene is achieved when the level of target gene expression in the test sample or the test mammal relative to the level of target gene expression in the control sample or the control mammal is about 95%, 90%, 85%, 80%, 75%, 70%, 65%, 60%, 55%, 50%, 45%, 40%, 35%, 30%, 25%, 20%, 15%, 10%, 5%, or 0%. In other words, the nucleic acids are capable of silencing, reducing, or inhibiting the expression of a target gene by at least about 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 100% in a test sample or a test mammal relative to the level of target gene expression in a control sample or a control mammal not contacted with or administered the nucleic acid. Suitable assays for determining the level of target gene expression include, without limitation, examination of protein or mRNA levels using techniques known to those of skill in the art, such as, *e.g.*, dot blots, northern blots, *in situ* hybridization, ELISA, immunoprecipitation, enzyme function, as well as phenotypic assays known to those of skill in the art.

An “effective amount” or “therapeutically effective amount” of an active agent or therapeutic agent such as a therapeutic nucleic acid is an amount sufficient to produce the desired effect, *e.g.*, an increase or inhibition of expression of a target sequence in comparison to the normal expression level detected in the absence of the nucleic acid. An increase in expression of a target sequence is achieved when any measurable level is detected in the case of an expression product that is not present in the absence of the nucleic acid. In the case where the expression product is present at some level prior to contact with the nucleic acid, an increase in expression is achieved when the fold increase in value obtained with a nucleic acid such as mRNA relative to control is about 1.05, 1.1, 1.2, 1.3, 1.4, 1.5, 1.75, 2, 2.5, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 40, 50, 75, 100, 250, 500, 750, 1000, 5000, 10000, or greater. Inhibition of expression of a target gene or target sequence is achieved when the value obtained with a nucleic acid such as antisense oligonucleotide relative to the control is about 95%, 90%, 85%, 80%, 75%, 70%, 65%, 60%, 55%, 50%, 45%, 40%, 35%, 30%, 25%, 20%, 15%, 10%, 5%, or 0%. Suitable assays for measuring expression of a target gene or target sequence include, *e.g.*, examination of protein or RNA levels using techniques known to those of skill in the art such as dot blots, northern blots, *in situ* hybridization, ELISA, immunoprecipitation, enzyme function, fluorescence, or luminescence of

suitable reporter proteins, as well as phenotypic assays known to those of skill in the art.


The term “nucleic acid” as used herein refers to a polymer containing at least two deoxyribonucleotides or ribonucleotides in either single- or double-stranded form and includes DNA, RNA, and hybrids thereof. DNA may be in the form of antisense molecules, plasmid DNA, cDNA, PCR products, or vectors. RNA may be in the form of small hairpin RNA (shRNA), messenger RNA (mRNA), antisense RNA, miRNA, micRNA, multivalent RNA, dicer substrate RNA or viral RNA (vRNA), and combinations thereof. Nucleic acids include nucleic acids containing known nucleotide analogs or modified backbone residues or linkages, which are synthetic, naturally occurring, and non-naturally occurring, and which have similar binding properties as the reference nucleic acid. Examples of such analogs include, without limitation, phosphorothioates, phosphoramidates, methyl phosphonates, chiral-methyl phosphonates, 2'-O-methyl ribonucleotides, and peptide-nucleic acids (PNAs). Unless specifically limited, the term encompasses nucleic acids containing known analogues of natural nucleotides that have similar binding properties as the reference nucleic acid. Unless otherwise indicated, a particular nucleic acid sequence also implicitly encompasses conservatively modified variants thereof (*e.g.*, degenerate codon substitutions), alleles, orthologs, single nucleotide polymorphisms, and complementary sequences as well as the sequence explicitly indicated. Specifically, degenerate codon substitutions may be achieved by generating sequences in which the third position of one or more selected (or all) codons is substituted with mixed-base and/or deoxyinosine residues (Batzner et al., *Nucleic Acid Res.*, 19:5081 (1991); Ohtsuka et al., *J. Biol. Chem.*, 260:2605-2608 (1985); Rossolini et al., *Mol. Cell. Probes*, 8:91-98 (1994)). “Nucleotides” contain a sugar deoxyribose (DNA) or ribose (RNA), a base, and a phosphate group. Nucleotides are linked together through the phosphate groups. “Bases” include purines and pyrimidines, which further include natural compounds adenine, thymine, guanine, cytosine, uracil, inosine, and natural analogs, and synthetic derivatives of purines and pyrimidines, which include, but are not limited to, modifications which place new reactive groups such as, but not limited to, amines, alcohols, thiols, carboxylates, and alkylhalides.

The term “gene” refers to a nucleic acid (*e.g.*, DNA or RNA) sequence that comprises partial length or entire length coding sequences necessary for the production of a polypeptide or precursor polypeptide.

“Gene product,” as used herein, refers to a product of a gene such as an RNA transcript or a polypeptide.

The term “lipid” refers to a group of organic compounds that include, but are not limited to, esters of fatty acids and are generally characterized by being poorly soluble in water, but soluble in many organic solvents. They are usually divided into at least three classes: (1) “simple lipids,” which include fats and oils as well as waxes; (2) “compound lipids,” which include phospholipids and glycolipids; and (3) “derived lipids” such as steroids.

A “steroid” is a compound comprising the following carbon skeleton:



Non-limiting examples of steroids include cholesterol, and the like.

A “cationic lipid” refers to a lipid capable of being positively charged. Exemplary cationic lipids include one or more amine group(s) which bear the positive charge. Preferred cationic lipids are ionizable such that they can exist in a positively charged or neutral form depending on pH. The ionization of the cationic lipid affects the surface charge of the lipid nanoparticle under different pH conditions. This charge state can influence plasma protein absorption, blood clearance, and tissue distribution (Semple, S.C., et al., *Adv. Drug Deliv Rev* 32:3-17 (1998)) as well as the ability to form endosomal non-bilayer structures (Hafez, I.M., et al., *Gene Ther* 8:1188-1196 (2001)) critical to the intracellular delivery of nucleic acids.

The term “polymer conjugated lipid” refers to a molecule comprising both a lipid portion and a polymer portion. An example of a polymer conjugated lipid is a pegylated lipid. The term “pegylated lipid” refers to a molecule comprising both a lipid portion and a polyethylene glycol portion. Pegylated lipids are known in the art and

include 1-(monomethoxy-polyethyleneglycol)-2,3-dimyristoylglycerol (PEG-DMG) and the like.

The term “neutral lipid” refers to any of a number of lipid species that exist either in an uncharged or neutral zwitterionic form at a selected pH. At physiological pH, such lipids include, but are not limited to, phosphatidylcholines such as 1,2-Distearoyl-*sn*-glycero-3-phosphocholine (DSPC), 1,2-Dipalmitoyl-*sn*-glycero-3-phosphocholine (DPPC), 1,2-Dimyristoyl-*sn*-glycero-3-phosphocholine (DMPC), 1-Palmitoyl-2-oleoyl-*sn*-glycero-3-phosphocholine (POPC), 1,2-dioleoyl-*sn*-glycero-3-phosphocholine (DOPC), phosphatidylethanolamines such as 1,2-Dioleoyl-*sn*-glycero-3-phosphoethanolamine (DOPE), sphingomyelins (SM), ceramides, steroids such as sterols and their derivatives. Neutral lipids may be synthetic or naturally derived.

The term “charged lipid” refers to any of a number of lipid species that exist in either a positively charged or negatively charged form independent of the pH within a useful physiological range, *e.g.*, pH ~3 to pH ~9. Charged lipids may be synthetic or naturally derived. Examples of charged lipids include phosphatidylserines, phosphatidic acids, phosphatidylglycerols, phosphatidylinositols, sterol hemisuccinates, dialkyl trimethylammonium-propanes, (*e.g.*, DOTAP, DOTMA), dialkyl dimethylaminopropanes, ethyl phosphocholines, dimethylaminoethane carbamoyl sterols (*e.g.*, DC-Chol).

The term “lipid nanoparticle” refers to particles having at least one dimension on the order of nanometers (*e.g.*, 1-1,000 nm) which include one or more of the compounds of structure (I) or other specified cationic lipids. In some embodiments, lipid nanoparticles comprising the disclosed cationic lipids (*e.g.*, compounds of structure (I)) are included in a formulation that can be used to deliver an active agent or therapeutic agent, such as a nucleic acid (*e.g.*, mRNA) to a target site of interest (*e.g.*, cell, tissue, organ, tumor, and the like). In some embodiments, the lipid nanoparticles comprise a compound of structure (I) and a nucleic acid. Such lipid nanoparticles typically comprise a compound of structure (I) and one or more excipient selected from neutral lipids, charged lipids, steroids, and polymer conjugated lipids. In some embodiments, the active agent or therapeutic agent, such as a nucleic acid, may be encapsulated in the lipid portion of the lipid nanoparticle or an aqueous space

enveloped by some or all of the lipid portion of the lipid nanoparticle, thereby protecting it from enzymatic degradation or other undesirable effects induced by the mechanisms of the host organism or cells, *e.g.*, an adverse immune response.

In various embodiments, the lipid nanoparticles have a mean diameter of from
5 about 30 nm to about 150 nm, from about 40 nm to about 150 nm, from about 50 nm to about 150 nm, from about 60 nm to about 130 nm, from about 70 nm to about 110 nm, from about 70 nm to about 100 nm, from about 80 nm to about 100 nm, from about 90 nm to about 100 nm, from about 70 to about 90 nm, from about 80 nm to about 90 nm, from about 70 nm to about 80 nm, or about 30 nm, 35 nm, 40 nm, 45 nm, 50 nm, 55
10 nm, 60 nm, 65 nm, 70 nm, 75 nm, 80 nm, 85 nm, 90 nm, 95 nm, 100 nm, 105 nm, 110 nm, 115 nm, 120 nm, 125 nm, 130 nm, 135 nm, 140 nm, 145 nm, or 150 nm. In some embodiments, the lipid nanoparticles are substantially non-toxic. In certain embodiments, nucleic acids, when present in the lipid nanoparticles, are resistant in aqueous solution to degradation with a nuclease. Lipid nanoparticles comprising
15 nucleic acids and their method of preparation are disclosed in, *e.g.*, U.S. Patent Pub. Nos. 2004/0142025, 2007/0042031 and PCT Pub. Nos. WO 2013/016058 and WO 2013/086373, the full disclosures of which are herein incorporated by reference in their entirety for all purposes.

As used herein, “lipid encapsulated” refers to a lipid nanoparticle that provides
20 an active agent or therapeutic agent, such as a nucleic acid (*e.g.*, mRNA), with full encapsulation, partial encapsulation, or both. In an embodiment, the nucleic acid (*e.g.*, mRNA) is fully encapsulated in the lipid nanoparticle.

As used herein, the term “aqueous solution” refers to a composition comprising water.

25 “Serum-stable” in relation to nucleic acid-lipid nanoparticles means that the nucleotide is not significantly degraded after exposure to a serum or nuclease assay that would significantly degrade free DNA or RNA. Suitable assays include, for example, a standard serum assay, a DNase assay, or an RNase assay.

“Systemic delivery,” as used herein, refers to delivery of a therapeutic product
30 that can result in a broad exposure of an active agent within an organism. Some techniques of administration can lead to the systemic delivery of certain agents, but not

others. Systemic delivery means that a useful, preferably therapeutic, amount of an agent is exposed to most parts of the body. Systemic delivery of lipid nanoparticles can be by any means known in the art including, for example, intravenous, intraarterial, subcutaneous, and intraperitoneal delivery. In some embodiments, systemic delivery of lipid nanoparticles is by intravenous delivery.

“Local delivery,” as used herein, refers to delivery of an active agent directly to a target site within an organism. For example, an agent can be locally delivered by direct injection into a disease site such as a tumor, other target site such as a site of inflammation, or a target organ such as the liver, heart, pancreas, kidney, and the like. Local delivery can also include topical applications or localized injection techniques such as intramuscular, subcutaneous, or intradermal injection. Local delivery does not preclude a systemic pharmacological effect.

“Alkyl” refers to a straight or branched hydrocarbon chain radical consisting solely of carbon and hydrogen atoms, which is saturated, and having, for example, from one to twenty-four carbon atoms (C₁-C₂₄ alkyl), four to twenty carbon atoms (C₄-C₂₀ alkyl), six to sixteen carbon atoms (C₆-C₁₆ alkyl), six to nine carbon atoms (C₆-C₉ alkyl), one to fifteen carbon atoms (C₁-C₁₅ alkyl), one to twelve carbon atoms (C₁-C₁₂ alkyl), one to eight carbon atoms (C₁-C₈ alkyl) or one to six carbon atoms (C₁-C₆ alkyl) and which is attached to the rest of the molecule by a single bond, *e.g.*, methyl, ethyl, *n*-propyl, 1-methylethyl (iso propyl), *n*-butyl, *n*-pentyl, 1,1-dimethylethyl (*t*-butyl), 3-methylhexyl, 2-methylhexyl, and the like. Unless stated otherwise specifically in the specification, an alkyl group is substituted or unsubstituted.

“Alkylene” refers to a straight or branched divalent hydrocarbon chain linking the rest of the molecule to a radical group, consisting solely of carbon and hydrogen, which is saturated, and having, for example, from one to twenty-four carbon atoms (C₁-C₂₄ alkylene), one to fifteen carbon atoms (C₁-C₁₅ alkylene), one to twelve carbon atoms (C₁-C₁₂ alkylene), one to eight carbon atoms (C₁-C₈ alkylene), one to six carbon atoms (C₁-C₆ alkylene), two to four carbon atoms (C₂-C₄ alkylene), one to two carbon atoms (C₁-C₂ alkylene), *e.g.*, methylene, ethylene, propylene, *n*-butylene, and the like. The alkylene chain is attached to the rest of the molecule through a single bond and to the radical group through a single bond. The points of attachment of the alkylene chain to

the rest of the molecule and to the radical group can be through one carbon or any two carbons within the chain. Unless stated otherwise specifically in the specification, an alkylene chain is substituted or unsubstituted.

- “Alkene” and “alkenylene” refer to an alkyl and alkylene, respectively,
 5 comprising at least one carbon-carbon double bond. Alkenes and alkenylenes include the same number of carbon atoms as alkyl and alkylene as defined above, except that alkenes and alkenylenes must include at least two carbons. Unless stated otherwise specifically in the specification, alkenes and alkenylenes are substituted or unsubstituted.
- 10 The term “substituted” used herein means any of the above groups (*e.g.*, alkyl or alkylene) wherein at least one hydrogen atom is replaced by a bond to a non-hydrogen atom such as, but not limited to: a halogen atom such as F, Cl, Br, or I; oxo groups (=O); hydroxyl groups (-OH); carboxyl groups -(CO₂H); C₁-C₁₂ alkyl groups; -(C=O)OR'; -O(C=O)R'; -C(=O)R'; -OR'; -S(O)_xR'; -S-SR'; -C(=O)SR';
 15 -SC(=O)R'; -NR'R'; -NR'C(=O)R'; -C(=O)NR'R'; -NR'C(=O)NR'R'; -OC(=O)NR'R'; -NR'C(=O)OR'; -NR'S(O)_xNR'R'; -NR'S(O)_xR'; and -S(O)_xNR'R', wherein: R' is, at each occurrence, independently H or C₁-C₁₅ alkyl and x is 0, 1 or 2. In some embodiments the substituent is a C₁-C₁₂ alkyl group. In other embodiments, the substituent is a halo group, such as fluoro. In other embodiments, the substituent is an oxo group. In other
 20 embodiments, the substituent is a hydroxyl group. In other embodiments, the substituent is an alkoxy group (-OR'). In other embodiments, the substituent is a carboxyl group. In other embodiments, the substituent is an amine group (-NR'R').

- “Optional” or “optionally” (*e.g.*, optionally substituted) means that the subsequently described event of circumstances may or may not occur, and that the
 25 description includes instances where said event or circumstance occurs and instances in which it does not. For example, “optionally substituted alkyl” means that the alkyl radical may or may not be substituted and that the description includes both substituted alkyl radicals and alkyl radicals having no substitution.

- The disclosure disclosed herein is also meant to encompass all pharmaceutically
 30 acceptable compounds of the compound of structure (I) being isotopically-labelled by having one or more atoms replaced by an atom having a different atomic mass or mass

number. Examples of isotopes that can be incorporated into the disclosed compounds include isotopes of hydrogen, carbon, nitrogen, oxygen, phosphorous, fluorine, chlorine, and iodine, such as ^2H , ^3H , ^{11}C , ^{13}C , ^{14}C , ^{13}N , ^{15}N , ^{15}O , ^{17}O , ^{18}O , ^{31}P , ^{32}P , ^{35}S , ^{18}F , ^{36}Cl , ^{123}I , and ^{125}I , respectively. These radiolabeled compounds could be useful to help determine or measure the effectiveness of the compounds, by characterizing, for example, the site or mode of action, or binding affinity to pharmacologically important site of action. Certain isotopically-labelled compounds of structure (I), (IA) or (IB), for example, those incorporating a radioactive isotope, are useful in drug and/or substrate tissue distribution studies. The radioactive isotopes tritium, *i.e.*, ^3H , and carbon-14, *i.e.*, ^{14}C , are particularly useful for this purpose in view of their ease of incorporation and ready means of detection.

Substitution with heavier isotopes such as deuterium, *i.e.*, ^2H , may afford certain therapeutic advantages resulting from greater metabolic stability, for example, increased *in vivo* half-life or reduced dosage requirements, and hence may be preferred in some circumstances.

Substitution with positron emitting isotopes, such as ^{11}C , ^{18}F , ^{15}O and ^{13}N , can be useful in Positron Emission Topography (PET) studies for examining substrate receptor occupancy. Isotopically-labeled compounds of structure (I) can generally be prepared by conventional techniques known to those skilled in the art or by processes analogous to those described in the Preparations and Examples as set out below using an appropriate isotopically-labeled reagent in place of the non-labeled reagent previously employed.

“Stable compound” and “stable structure” are meant to indicate a compound that is sufficiently robust to survive isolation to a useful degree of purity from a reaction mixture, and formulation into an efficacious therapeutic agent.

“Mammal” includes humans and both domestic animals such as laboratory animals and household pets (*e.g.*, cats, dogs, swine, cattle, sheep, goats, horses, rabbits), and non-domestic animals such as wildlife and the like.

“Pharmaceutically acceptable carrier, diluent or excipient” includes without limitation any adjuvant, carrier, excipient, glidant, sweetening agent, diluent, preservative, dye/colorant, flavor enhancer, surfactant, wetting agent, dispersing agent,

suspending agent, stabilizer, isotonic agent, solvent, or emulsifier which has been approved by the United States Food and Drug Administration as being acceptable for use in humans or domestic animals.

“Pharmaceutically acceptable salt” includes both acid and base addition salts.

5 “Pharmaceutically acceptable acid addition salt” refers to those salts which retain the biological effectiveness and properties of the free bases, which are not biologically or otherwise undesirable, and which are formed with inorganic acids such as, but are not limited to, hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid and the like, and organic acids such as, but not limited to, acetic acid,
10 2,2-dichloroacetic acid, adipic acid, alginic acid, ascorbic acid, aspartic acid, benzenesulfonic acid, benzoic acid, 4-acetamidobenzoic acid, camphoric acid, camphor-10-sulfonic acid, capric acid, caproic acid, caprylic acid, carbonic acid, cinnamic acid, citric acid, cyclamic acid, dodecylsulfuric acid, ethane-1,2-disulfonic acid, ethanesulfonic acid, 2-hydroxyethanesulfonic acid, formic acid, fumaric acid,
15 galactaric acid, gentisic acid, glucoheptonic acid, gluconic acid, glucuronic acid, glutamic acid, glutaric acid, 2-oxo-glutaric acid, glycerophosphoric acid, glycolic acid, hippuric acid, isobutyric acid, lactic acid, lactobionic acid, lauric acid, maleic acid, malic acid, malonic acid, mandelic acid, methanesulfonic acid, mucic acid, naphthalene-1,5-disulfonic acid, naphthalene-2-sulfonic acid, 1-hydroxy-2-naphthoic
20 acid, nicotinic acid, oleic acid, orotic acid, oxalic acid, palmitic acid, pamoic acid, propionic acid, pyroglutamic acid, pyruvic acid, salicylic acid, 4-aminosalicylic acid, sebacic acid, stearic acid, succinic acid, tartaric acid, thiocyanic acid, *p*-toluenesulfonic acid, trifluoroacetic acid, undecylenic acid, and the like.

“Pharmaceutically acceptable base addition salt” refers to those salts which
25 retain the biological effectiveness and properties of the free acids, which are not biologically or otherwise undesirable. These salts are prepared from addition of an inorganic base or an organic base to the free acid. Salts derived from inorganic bases include, but are not limited to, the sodium, potassium, lithium, ammonium, calcium, magnesium, iron, zinc, copper, manganese, aluminum salts and the like. Preferred
30 inorganic salts are the ammonium, sodium, potassium, calcium, and magnesium salts. Salts derived from organic bases include, but are not limited to, salts of primary,

secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines and basic ion exchange resins, such as ammonia, isopropylamine, trimethylamine, diethylamine, triethylamine, tripropylamine, diethanolamine, ethanolamine, deanol, 2-dimethylaminoethanol, 5 2-diethylaminoethanol, dicyclohexylamine, lysine, arginine, histidine, caffeine, procaine, hydrabamine, choline, betaine, benethamine, benzathine, ethylenediamine, glucosamine, methylglucamine, theobromine, triethanolamine, tromethamine, purines, piperazine, piperidine, *N*-ethylpiperidine, polyamine resins and the like. Particularly preferred organic bases are isopropylamine, diethylamine, ethanolamine, 10 trimethylamine, dicyclohexylamine, choline and caffeine.

Often crystallizations produce a solvate of a compound of the disclosure (*i.e.*, a compound of structure (I)). As used herein, the term “solvate” refers to an aggregate that comprises one or more molecules of a compound of the disclosure with one or more molecules of solvent. The solvent may be water, in which case the solvate may be 15 a hydrate. Alternatively, the solvent may be an organic solvent. Thus, the compounds of the present disclosure may exist as a hydrate, including a monohydrate, dihydrate, hemihydrate, sesquihydrate, trihydrate, tetrahydrate and the like, as well as the corresponding solvated forms. Solvates of compound of the disclosure may be true solvates, while in other cases the compound of the disclosure may merely retain 20 adventitious water or be a mixture of water plus some adventitious solvent.

A “pharmaceutical composition” refers to a formulation of a compound of the disclosure and a medium generally accepted in the art for the delivery of the biologically active compound to mammals, *e.g.*, humans. Such a medium includes all pharmaceutically acceptable carriers, diluents or excipients therefor.

25 “Effective amount” or “therapeutically effective amount” refers to that amount of a compound of the disclosure which, when administered to a mammal, preferably a human, is sufficient to effect treatment in the mammal, preferably a human. The amount of a lipid nanoparticle of the disclosure which constitutes a “therapeutically effective amount” will vary depending on the compound, the condition and its severity, 30 the manner of administration, and the age of the mammal to be treated, but can be

determined routinely by one of ordinary skill in the art having regard to his own knowledge and to this disclosure.

“Treating” or “treatment” as used herein covers the treatment of the disease or condition of interest in a mammal, preferably a human, having the disease or condition of interest, and includes:

- (i) preventing the disease or condition from occurring in a mammal, in particular, when such mammal is predisposed to the condition but has not yet been diagnosed as having it;
- (ii) inhibiting the disease or condition, *i.e.*, arresting its development;
- 10 (iii) relieving the disease or condition, *i.e.*, causing regression of the disease or condition; or
- (iv) relieving the symptoms resulting from the disease or condition, *i.e.*, relieving pain without addressing the underlying disease or condition. As used herein, the terms “disease” and “condition” may be used interchangeably or may be different in that the particular malady or condition may not have a known causative agent (so that etiology has not yet been worked out) and it is therefore not yet recognized as a disease but only as an undesirable condition or syndrome, wherein a more or less specific set of symptoms have been identified by clinicians.

The compounds of the disclosure, or their pharmaceutically acceptable salts may contain one or more stereocenters and may thus give rise to enantiomers, diastereomers, and other stereoisomeric forms that may be defined, in terms of absolute stereochemistry, as (*R*)- or (*S*)- or, as (*D*)- or (*L*)- for amino acids. The present disclosure is meant to include all such possible isomers, as well as their racemic and optically pure forms. Optically active (+) and (-), (*R*)- and (*S*)-, or (*D*)- and 25 (*L*)- isomers may be prepared using chiral synthons or chiral reagents, or resolved using conventional techniques, for example, chromatography and fractional crystallization. Conventional techniques for the preparation/isolation of individual enantiomers include chiral synthesis from a suitable optically pure precursor or resolution of the racemate (or the racemate of a salt or derivative) using, for example, chiral high pressure liquid 30 chromatography (HPLC). When the compounds described herein contain olefinic double bonds or other centers of geometric asymmetry, and unless specified otherwise,

it is intended that the compounds include both E and Z geometric isomers. Likewise, all tautomeric forms are also intended to be included.


A “stereoisomer” refers to a compound made up of the same atoms bonded by the same bonds but having different three-dimensional structures, which are not
5 interchangeable. The present disclosure contemplates various stereoisomers and mixtures thereof and includes “enantiomers,” which refers to two stereoisomers whose molecules are non-superimposable mirror images of one another.

A “tautomer” refers to a proton shift from one atom of a molecule to another atom of the same molecule. The present disclosure includes tautomers of any said
10 compounds.

Compounds

In an aspect, the disclosure provides novel lipid compounds which are capable of combining with other lipid components such as neutral lipids, charged lipids, steroids and/or polymer conjugated-lipids to form lipid nanoparticles with therapeutic agents,
15 such as oligonucleotides. Without wishing to be bound by theory, it is thought that these lipid nanoparticles shield the therapeutic agent from degradation in the serum and provide for effective delivery of the therapeutic agent to cells *in vitro* and *in vivo*.

In one embodiment, the compounds have the following structure (I):



(I)

or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof,
wherein:

G^1 and G^2 are each independently C_1 - C_6 alkylene;

L^1 and L^2 are each independently $-O(C=O)-$ or $-(C=O)O-$;

R^{1a} and R^{1b} are, at each occurrence, independently either: (a) H or C₁-C₁₂ alkyl; or (b) R^{1a} is H or C₁-C₁₂ alkyl, and R^{1b} together with the carbon atom to which it is bound is taken together with an adjacent R^{1b} and the carbon atom to which it is bound to form a carbon-carbon double bond;

5 R^{2a} and R^{2b} are, at each occurrence, independently either: (a) H or C₁-C₁₂ alkyl; or (b) R^{2a} is H or C₁-C₁₂ alkyl, and R^{2b} together with the carbon atom to which it is bound is taken together with an adjacent R^{2b} and the carbon atom to which it is bound to form a carbon-carbon double bond;

R^{3a} and R^{3b} are, at each occurrence, independently either (a): H or C₁-C₁₂ 10 alkyl; or (b) R^{3a} is H or C₁-C₁₂ alkyl, and R^{3b} together with the carbon atom to which it is bound is taken together with an adjacent R^{3b} and the carbon atom to which it is bound to form a carbon-carbon double bond;

R^{4a} and R^{4b} are, at each occurrence, independently either: (a) H or C₁-C₁₂ alkyl; or (b) R^{4a} is H or C₁-C₁₂ alkyl, and R^{4b} together with the carbon atom to which it 15 is bound is taken together with an adjacent R^{4b} and the carbon atom to which it is bound to form a carbon-carbon double bond;

R^5 and R^6 are each independently H or methyl;

R^7 is $-O(C=O)R^{10}$, $-(C=O)OR^{10}$, $-NR^9(C=O)R^{10}$ or $-(C=O)NR^9R^{10}$;

R^8 is OH, $-N(R^{11})(C=O)R^{12}$, $-(C=O)NR^{11}R^{12}$, $-NR^{11}R^{12}$, $-(C=O)OR^{12}$ or 20 $-O(C=O)R^{12}$;

R^9 is H or C₁-C₁₅ alkyl;

R^{10} is C₁-C₁₅ alkyl;

R^{11} is H or C₁-C₆ alkyl;


R^{12} is C₁-C₆ alkyl;

25 X is $-(C=O)-$ or a direct bond; and

a, b, c and d are each independently an integer from 1 to 24;

wherein each alkyl and alkylene and is independently optionally substituted.

In other embodiments, the compound has one of the following structures (IA) or (IB):



or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof.

5 In certain embodiments, G¹ is C₂-C₃ alkylene. In different embodiments, G¹ is C₄-C₆ alkylene. For example, in various embodiments, G¹ is C₂, C₃, C₄, C₅ or C₆ alkylene.

In other embodiments, G² is C₂-C₄ alkylene, for example C₂-C₃ alkylene or C₃-C₄ alkylene. In some embodiments, G² is C₂, C₃ or C₄ alkylene.

10 In various different embodiments, X is -(C=O)-, while in different embodiments X is a direct bond.

In any of the foregoing embodiments, R⁷ is -O(C=O)R¹⁰ or -(C=O)OR¹⁰. In certain of these embodiments, R¹⁰ is linear C₁-C₁₅ alkyl, such as linear C₆-C₁₀ alkyl. In other such embodiments, R¹⁰ is methyl or R¹⁰ is branched C₂-C₁₅ alkyl, such as

15 branched C₁₀-C₁₅ alkyl.

In still other of the foregoing embodiments, R⁷ is -NR⁹(C=O) or -(C=O)NR⁹R¹⁰. In some of these embodiments, R⁹ is H. In other of these embodiments, R⁹ and R¹⁰ are each independently C₆-C₁₀ alkyl.



20 In other embodiments, at least one occurrence of R^{1a} and R^{1b}, R^{1a} is H or C₁-C₁₂ alkyl, and R^{1b} together with the carbon atom to which it is bound is taken together with an adjacent R^{1b} and the carbon atom to which it is bound to form a carbon-carbon double bond.


25 In more embodiments, for at least one occurrence of R^{4a} and R^{4b}, R^{4a} is H or C₁-C₁₂ alkyl, and R^{4b} together with the carbon atom to which it is bound is taken together with an adjacent R^{4b} and the carbon atom to which it is bound to form a carbon-carbon double bond.

In still other embodiments, for at least one occurrence of R^{2a} and R^{2b} , R^{2a} is H or C_1 - C_{12} alkyl, and R^{2b} together with the carbon atom to which it is bound is taken together with an adjacent R^{2b} and the carbon atom to which it is bound to form a carbon-carbon double bond.

5 In other embodiments, for at least one occurrence of R^{3a} and R^{3b} , R^{3a} is H or C_1 - C_{12} alkyl, and R^{3b} together with the carbon atom to which it is bound is taken together with an adjacent R^{3b} and the carbon atom to which it is bound to form a carbon-carbon double bond.

10 In various of the foregoing embodiments, R^{1a} , R^{1b} , R^{2a} , R^{2b} , R^{3a} , R^{3b} , R^{4a} and R^{4b} are, at each occurrence, independently H or C_1 - C_{12} alkyl. In other embodiments, R^{2a} , R^{2b} , R^{3a} and R^{3b} are, at each occurrence, H. For example, in certain embodiment R^{1a} and R^{4a} are, at each occurrence, H. In different embodiments, at least one of R^{1b} and R^{4b} is C_1 - C_8 alkyl. For example, in some embodiments C_1 - C_8 alkyl is methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, tert-butyl, n-hexyl or n-octyl.

15 In other more specific embodiments,  or , or both, independently has one of the following structures:



In more embodiments, a, b, c and d are each independently an integer from 2 to 12. In more embodiments, a, b, c and d are each independently an integer from 4 to 10, 5 to 10, 6 to 10, 4 to 9, 5 to 9 or 6 to 9. In still different embodiments, b and c are independently 5, 6, 7, 8, 9 or 10.

In some embodiments, one of R^5 or R^6 is methyl. In other embodiments, each of R^5 and R^6 is methyl.

In some embodiments, R^8 is OH.

In other embodiments, R^8 is $-N(R^{11})(C=O)R^{12}$. In different embodiments, R^8 is $-(C=O)NR^{11}R^{12}$. In more different embodiments, R^8 is $-NR^{11}R^{12}$. In some of these foregoing embodiments, R^{11} and R^{12} are each independently H or C_1 - C_8 alkyl. In other of these embodiments, R^{11} and R^{12} are each independently H or C_1 - C_3 alkyl. For example, in some embodiments the C_1 - C_8 alkyl or C_1 - C_3 alkyl is unsubstituted or substituted with hydroxyl. In other different such embodiments, R^{11} and R^{12} are each methyl.


In other embodiments of the compound of structure (I), R^8 is $-(C=O)OR^{12}$, while in different embodiments R^8 is $-O(C=O)R^{12}$.






In specific embodiments of any of the foregoing compounds, R^8 has one of the following structures:













In various different embodiments, the disclosure provides a compound having one of the structures set forth in Table 1 below, or a pharmaceutically acceptable salt or tautomer thereof.






Table 1. Representative Compounds



No.	Structure
I-1	

No.	Structure
I-2	
I-3	
I-4	
I-5	
I-6	

No.	Structure
I-7	
I-8	
I-9	
I-10	
I-11	

No.	Structure
I-12	
I-13	
I-14	
I-15	
I-16	

No.	Structure
I-17	
I-18	
I-19	
I-20	
I-21	

No.	Structure
I-22	
I-23	

It is understood that any embodiment of the compounds of structure (I), as set forth above, and any specific substituent and/or variable in the compound of structure (I), as set forth above, may be independently combined with other embodiments and/or substituents and/or variables of compounds of structure (I) to form embodiments of the disclosures not specifically set forth above. In addition, in the event that a list of substituents and/or variables is listed for any particular R group, G group, L group or variable a, b, c, d or n, in a particular embodiment and/or claim, it is understood that each individual substituent and/or variable may be deleted from the particular embodiment and/or claim and that the remaining list of substituents and/or variables will be considered to be within the scope of the disclosure.

It is understood that in the present description, combinations of substituents and/or variables of the depicted formulae are permissible only if such contributions result in stable compounds.

In some embodiments, lipid nanoparticles comprising a compound of structure (I) are provided. The lipid nanoparticles optionally include excipients selected from a neutral lipid, a steroid and a polymer conjugated lipid.

In some embodiments, compositions comprising any one or more of the compounds of structure (I) and a therapeutic agent are provided. For example, in some embodiments, the compositions comprise any of the compounds of structure (I) and a therapeutic agent and one or more excipient selected from neutral lipids, steroids and


polymer conjugated lipids. Other pharmaceutically acceptable excipients and/or carriers are also included in various embodiments of the compositions.

In some embodiments, the neutral lipid is selected from DSPC, DPPC, DMPC, DOPC, POPC, DOPE and SM. In some embodiments, the neutral lipid is DSPC. In various embodiments, the molar ratio of the compound to the neutral lipid ranges from about 2:1 to about 8:1.

In various embodiments, the compositions further comprise a steroid or steroid analogue. In certain embodiments, the steroid or steroid analogue is cholesterol. In some of these embodiments, the molar ratio of the compound to cholesterol ranges from about 5:1 to 1:1.

In various embodiments, the polymer conjugated lipid is a pegylated lipid. For example, some embodiments include a pegylated diacylglycerol (PEG-DAG) such as 1-(monomethoxy-polyethyleneglycol)-2,3-dimyristoylglycerol (PEG-DMG), a pegylated phosphatidylethanolamine (PEG-PE), a PEG succinate diacylglycerol (PEG-S-DAG) such as 4-O-(2',3'-di(tetradecanoyloxy)propyl-1-O-(ω -methoxy(polyethoxy)ethyl)butanedioate (PEG-S-DMG), a pegylated ceramide (PEG-cer), or a PEG dialkoxypropylcarbamate such as ω -methoxy(polyethoxy)ethyl-N-(2,3-di(tetradecanoyloxy)propyl)carbamate or 2,3-di(tetradecanoyloxy)propyl-N-(ω -methoxy(polyethoxy)ethyl)carbamate. In various embodiments, the molar ratio of the compound to the pegylated lipid ranges from about 100:1 to about 20:1.

In some embodiments, the composition comprises a pegylated lipid having the following structure (II):



(II)

or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein:

R^8 and R^9 are each independently a straight or branched, saturated or unsaturated alkyl chain containing from 10 to 30 carbon atoms, wherein the alkyl chain is optionally interrupted by one or more ester bonds; and

w has a mean value ranging from 30 to 60.

In some embodiments, R⁸ and R⁹ are each independently straight, saturated alkyl chains containing from 12 to 16 carbon atoms. In other embodiments, the average w ranges from about 42 to 55, for example about 49.

5 In some embodiments of the foregoing composition, the therapeutic agent comprises a nucleic acid. For example, in some embodiments, the nucleic acid is selected from antisense and messenger RNA. In some of the foregoing embodiments, the composition comprises a lipid nanoparticle.


Some related embodiments provide a lipid nanoparticle comprising the
10 compound of any one of the foregoing embodiments (*e.g.*, a compound of structure (I)). In certain embodiments, the lipid nanoparticle further comprises a therapeutic agent (*e.g.*, a nucleic acid such as antisense and messenger RNA).

In some embodiments, the lipid nanoparticle further comprises one or more excipient selected from neutral lipids, steroids and polymer conjugated lipids. In some
15 embodiments, the neutral lipids are selected from DSPC, DPPC, DMPC, DOPC, POPC, DOPE and SM. In more specific embodiments, the neutral lipid is DSPC.

In some more specific embodiments, the molar ratio of the compound to the neutral lipid ranges from about 2:1 to about 8:1. In some embodiments, the steroid is cholesterol. In some embodiments, the molar ratio of the compound to cholesterol
20 ranges from 5:1 to 1:1.

In certain embodiments, the polymer conjugated lipid is pegylated lipid. In certain more specific embodiments, the molar ratio of the compound to pegylated lipid ranges from about 100:1 to about 20:1.

In some embodiments, the pegylated lipid is PEG-DAG, PEG-PE, PEG-S-DAG,
25 PEG-cer or a PEG dialkyoxypropylcarbamate. In other embodiments, the pegylated lipid has the following structure (II):



(II)

or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein:

R⁸ and R⁹ are each independently a straight or branched, saturated or unsaturated alkyl chain containing from 10 to 30 carbon atoms, wherein the alkyl chain
5 is optionally interrupted by one or more ester bonds; and

w has a mean value ranging from 30 to 60.

In some more specific embodiments of structure (II), R⁸ and R⁹ are each independently straight, saturated alkyl chains containing from 12 to 16 carbon atoms. In more specific embodiments, the average w is about 49.

10 In other different embodiments, the disclosure is directed to a method for administering a therapeutic agent to a patient in need thereof, the method comprising preparing or providing any of the foregoing compositions and administering the composition to the patient

For the purposes of administration, embodiments of the compounds of the
15 present disclosure (typically in the form of lipid nanoparticles in combination with a therapeutic agent) may be administered as a raw chemical or may be formulated as pharmaceutical compositions. Pharmaceutical compositions of embodiments of the present disclosure comprise a compound of structure (I) and one or more pharmaceutically acceptable carrier, diluent or excipient. In some embodiments, the
20 compound of structure (I) is present in the composition in an amount which is effective to form a lipid nanoparticle and deliver the therapeutic agent, *e.g.*, for treating a particular disease or condition of interest. Appropriate concentrations and dosages can be readily determined by one skilled in the art.

Administration of the compositions of embodiments of the disclosure can be
25 carried out via any of the accepted modes of administration of agents for serving similar utilities. The pharmaceutical compositions of embodiments of the disclosure may be formulated into preparations in solid, semi-solid, liquid or gaseous forms, such as tablets, capsules, powders, granules, ointments, solutions, suspensions, suppositories, injections, inhalants, gels, microspheres, and aerosols. Typical routes of administering
30 such pharmaceutical compositions include, without limitation, oral, topical, transdermal, inhalation, parenteral, sublingual, buccal, rectal, vaginal, and intranasal.

The term parenteral as used herein includes subcutaneous injections, intravenous, intramuscular, intradermal, intrasternal injection or infusion techniques.

Pharmaceutical compositions of embodiments of the disclosure are formulated so as to allow the active ingredients contained therein to be bioavailable upon administration of the composition to a patient. Compositions that will be administered to a subject or patient in some embodiments take the form of one or more dosage units, where for example, a tablet may be a single dosage unit, and a container of a compound of an embodiment of the disclosure in aerosol form may hold a plurality of dosage units. Actual methods of preparing such dosage forms are known, or will be apparent, to those skilled in this art; for example, see *Remington: The Science and Practice of Pharmacy*, 20th Edition (Philadelphia College of Pharmacy and Science, 2000). In some embodiments, the composition to be administered will, in any event, contain a therapeutically effective amount of a compound of the disclosure, or a pharmaceutically acceptable salt thereof, for treatment of a disease or condition of interest in accordance with the teachings of this disclosure.

A pharmaceutical composition of embodiments of the disclosure may be in the form of a solid or liquid. In one aspect, the carrier(s) are particulate, so that the compositions are, for example, in tablet or powder form. The carrier(s) may be liquid, with the compositions being, for example, oral syrup, injectable liquid or an aerosol, which is useful in, for example, inhalatory administration.

When intended for oral administration, the pharmaceutical composition of certain embodiments is preferably in either solid or liquid form, where semi-solid, semi-liquid, suspension and gel forms are included within the forms considered herein as either solid or liquid.

As a solid composition for oral administration, the pharmaceutical composition of some embodiments may be formulated into a powder, granule, compressed tablet, pill, capsule, chewing gum, wafer or the like form. Such a solid composition will typically contain one or more inert diluents or edible carriers. In addition, one or more of the following may be present: binders such as carboxymethylcellulose, ethyl cellulose, microcrystalline cellulose, gum tragacanth or gelatin; excipients such as starch, lactose or dextrans, disintegrating agents such as alginic acid, sodium alginate,

Primogel, corn starch and the like; lubricants such as magnesium stearate or Sterotex; glidants such as colloidal silicon dioxide; sweetening agents such as sucrose or saccharin; a flavoring agent such as peppermint, methyl salicylate or orange flavoring; and a coloring agent.

5 When the pharmaceutical composition of some embodiments is in the form of a capsule, for example, a gelatin capsule, it may contain, in addition to materials of the above type, a liquid carrier such as polyethylene glycol or oil.

 The pharmaceutical composition of some embodiments may be in the form of a liquid, for example, an elixir, syrup, solution, emulsion or suspension. The liquid may
10 be for oral administration or for delivery by injection, as two examples. When intended for oral administration, preferred composition contain, in addition to a compound of structure (I), one or more of a sweetening agent, preservatives, dye/colorant and flavor enhancer. In a composition intended to be administered by injection, one or more of a surfactant, preservative, wetting agent, dispersing agent, suspending agent, buffer,
15 stabilizer and isotonic agent may be included.

 The liquid pharmaceutical compositions of embodiments of the disclosure, whether they be solutions, suspensions or other like form, may include one or more of the following adjuvants: sterile diluents such as water for injection, saline solution, preferably physiological saline, Ringer's solution, isotonic sodium chloride, fixed oils
20 such as synthetic mono or diglycerides which may serve as the solvent or suspending medium, polyethylene glycols, glycerin, propylene glycol or other solvents; antibacterial agents such as benzyl alcohol or methyl paraben; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid; buffers such as acetates, citrates or phosphates and agents for the adjustment of
25 tonicity such as sodium chloride or dextrose; agents to act as cryoprotectants such as sucrose or trehalose. The parenteral preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic. Physiological saline is a preferred adjuvant. An injectable pharmaceutical composition is preferably sterile.

 The pharmaceutical composition of embodiments of the disclosure may be
30 intended for topical administration, in which case the carrier may suitably comprise a solution, emulsion, ointment or gel base. The base, for example, may comprise one or

more of the following: petrolatum, lanolin, polyethylene glycols, bee wax, mineral oil, diluents such as water and alcohol, and emulsifiers and stabilizers. Thickening agents may be present in a pharmaceutical composition for topical administration. If intended for transdermal administration, the composition may include a transdermal patch or
5 iontophoresis device.

The pharmaceutical composition of embodiments of the disclosure may include various materials, which modify the physical form of a solid or liquid dosage unit. For example, the composition may include materials that form a coating shell around the active ingredients. The materials that form the coating shell are typically inert, and may
10 be selected from, for example, sugar, shellac, and other enteric coating agents. Alternatively, the active ingredients may be encased in a gelatin capsule.

The pharmaceutical composition of embodiments of the disclosure in solid or liquid form may include an agent that binds to the compound of the disclosure and thereby assists in the delivery of the LNP. Suitable agents that may act in this capacity
15 include a monoclonal or polyclonal antibody, or a protein.

The pharmaceutical composition of embodiments of the disclosure may consist of dosage units that can be administered as an aerosol. The term aerosol is used to denote a variety of systems ranging from those of colloidal nature to systems consisting of pressurized packages. Delivery may be by a liquefied or compressed gas or by a
20 suitable pump system that dispenses the active ingredients. Aerosols of LNPs of embodiments of the disclosure may be delivered in single phase, bi-phasic, or tri-phasic systems in order to deliver the active ingredient(s). Delivery of the aerosol includes the necessary container, activators, valves, sub-containers, and the like, which together may form a kit. One skilled in the art, without undue experimentation, may determine
25 preferred aerosols.

The pharmaceutical compositions of embodiments of the disclosure may be prepared by methodology well known in the pharmaceutical art. For example, a pharmaceutical composition intended to be administered by injection can be prepared by combining the lipid nanoparticles of the disclosure with sterile, distilled water or
30 other carrier so as to form a solution. A surfactant may be added to facilitate the formation of a homogeneous solution or suspension. Surfactants are compounds that

non-covalently interact with the compound of the disclosure so as to facilitate dissolution or homogeneous suspension of the compound in the aqueous delivery system.

The compositions of embodiments of the disclosure, or their pharmaceutically acceptable salts, are administered in a therapeutically effective amount, which will vary depending upon a variety of factors including the activity of the specific therapeutic agent employed; the metabolic stability and length of action of the therapeutic agent; the age, body weight, general health, sex, and diet of the patient; the mode and time of administration; the rate of excretion; the drug combination; the severity of the particular disorder or condition; and the subject undergoing therapy.

Compositions of embodiments of the disclosure may also be administered simultaneously with, prior to, or after administration of one or more other therapeutic agents. Such combination therapy includes administration of a single pharmaceutical dosage formulation of a composition of embodiments of the disclosure and one or more additional active agents, as well as administration of the composition of embodiments of the disclosure and each active agent in its own separate pharmaceutical dosage formulation. For example, a composition of embodiments of the disclosure and the other active agent can be administered to the patient together in a single oral dosage composition such as a tablet or capsule, or each agent administered in separate oral dosage formulations. Where separate dosage formulations are used, the compounds of embodiments of the disclosure and one or more additional active agents can be administered at essentially the same time, *i.e.*, concurrently, or at separately staggered times, *i.e.*, sequentially; combination therapy is understood to include all these regimens.

Preparation methods for the above compounds and compositions are described herein below and/or known in the art.


It will be appreciated by those skilled in the art that in the process described herein the functional groups of intermediate compounds may need to be protected by suitable protecting groups. Such functional groups include hydroxy, amino, mercapto and carboxylic acid. Suitable protecting groups for hydroxy include trialkylsilyl or diarylalkylsilyl (for example, *t*-butyldimethylsilyl, *t*-butyldiphenylsilyl or

trimethylsilyl), tetrahydropyranyl, benzyl, and the like. Suitable protecting groups for amino, amidino and guanidino include *t*-butoxycarbonyl, benzyloxycarbonyl, and the like. Suitable protecting groups for mercapto include -C(O)-R'' (where R'' is alkyl, aryl or arylalkyl), *p*-methoxybenzyl, trityl and the like. Suitable protecting groups for carboxylic acid include alkyl, aryl or arylalkyl esters. Protecting groups may be added or removed in accordance with standard techniques, which are known to one skilled in the art and as described herein. The use of protecting groups is described in detail in Green, T.W. and P.G.M. Wutz, *Protective Groups in Organic Synthesis* (1999), 3rd Ed., Wiley. As one of skill in the art would appreciate, the protecting group may also be a polymer resin such as a Wang resin, Rink resin or 2-chlorotrityl-chloride resin.

It will also be appreciated by those skilled in the art, although such protected derivatives of compounds of this disclosure may not possess pharmacological activity as such, they may be administered to a mammal and thereafter metabolized in the body to form compounds of the disclosure which are pharmacologically active. Such derivatives may therefore be described as "prodrugs." All prodrugs of compounds of this disclosure are included within the scope of the disclosure.

Furthermore, compounds of embodiments of the disclosure which exist in free base or acid form can be converted to their pharmaceutically acceptable salts by treatment with the appropriate inorganic or organic base or acid by methods known to one skilled in the art. Salts of compounds of embodiments of the disclosure can be converted to their free base or acid form by standard techniques.


The following General Reaction Scheme 1 illustrates exemplary methods to make compounds of this disclosure, *i.e.*, compounds of structure (I):



(I)

or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof, wherein a, b, c, d, G¹, G², L¹, L², R^{1a}, R^{1b}, R^{2a}, R^{2b}, R^{3a}, R^{3b}, R^{4a}, R^{4b}, R⁵, R⁶, R⁷, R⁸ and X are as defined herein. It is understood that one skilled in the art may be able to make these compounds by similar methods or by combining other methods known to one skilled in the art. It is also understood that one skilled in the art would be able to make, in a similar manner as described below, other compounds of structure (I) not specifically illustrated below by using the appropriate starting components and modifying the parameters of the synthesis as needed. In general, starting components may be obtained from sources such as Sigma Aldrich, Lancaster Synthesis, Inc., Maybridge, Matrix Scientific, TCI, and Fluorochem USA, etc. or synthesized according to sources known to those skilled in the art (see, for example, Advanced Organic Chemistry: Reactions, Mechanisms, and Structure, 5th edition (Wiley, December 2000)) or prepared as described in this disclosure.

GENERAL REACTION SCHEME 1



15

General Reaction Scheme I provides an exemplary method for preparation of a compound of structure (I) (i.e., A5). a, b, c, d, G², L¹, L², R^{1a}, R^{1b}, R^{2a}, R^{2b}, R^{3a}, R^{3b}, R^{4a}, R^{4b}, R⁵, R⁶, R⁸, R⁹ and R¹⁰ in General reaction Scheme 1 are as defined herein, and

Z represents an activated analogue of G¹ sufficient for bond formation with the NH group of A3 (e.g., an alkene or an alkylene terminating in an aldehyde, acid halide, acrylate, etc.). Intermediates and reagents (e.g., A1 and A2) needed for preparation of the compounds according to General Reaction Scheme I can be purchased or prepared
5 according to the examples below or methods known by one of ordinary skill in the art.

It should be noted that various alternative strategies for preparation of compounds of structure (I) are available to those of ordinary skill in the art. For example, other compounds of structure (I) wherein can be prepared according to analogous methods using the appropriate starting material. The use of protecting
10 groups as needed and other modification to the above General Reaction Scheme will be readily apparent to one of ordinary skill in the art.

The following examples are provided for purpose of illustration and not limitation.

EXAMPLE 1

15 LUCIFERASE mRNA IN VIVO EVALUATION USING THE LIPID NANOPARTICLE COMPOSITIONS

Lipid nanoparticles were prepared and tested according to the general procedures described in PCT Pub. Nos. WO 2015/199952 and WO 2017/004143, the full disclosures of which are incorporated herein by reference. Briefly, cationic lipid,
20 DSPC, cholesterol and PEG-lipid were solubilized in ethanol at a molar ratio of about 50:10:38.5:1.5 or about 47.5:10:40.8:1.7. Lipid nanoparticles (LNP) were prepared at a total lipid to mRNA weight ratio of approximately 10:1 to 30:1. The mRNA is diluted to 0.2 mg/mL in 10 to 50 mM citrate or acetate buffer, pH 4. Syringe pumps were used to mix the ethanolic lipid solution with the mRNA aqueous solution at a ratio of about
25 1:5 to 1:3 (vol/vol) with total flow rates above 15 mL/min. The ethanol was then removed and the external buffer replaced with PBS by dialysis. Finally, the lipid nanoparticles were filtered through a 0.2 µm pore sterile filter. Lipid nanoparticle particle size was approximately 55-95 nm diameter, and in some instances approximately 70-90 nm diameter as determined by quasi-elastic light scattering using a
30 Malvern Zetasizer Nano ZS (Malvern, UK).

Studies were performed in 6-8 week old female C57BL/6 mice (Charles River) or 8-10 week old CD-1 (Harlan) mice (Charles River) according to guidelines established by an institutional animal care committee (ACC) and the Canadian Council on Animal Care (CCAC). Varying doses of mRNA-lipid nanoparticle were
5 systemically administered by tail vein injection and animals euthanized at a specific time point (*e.g.*, 4 hours) post-administration. Liver and spleen were collected in pre-weighed tubes, weights determined, immediately snap frozen in liquid nitrogen and stored at -80°C until processing for analysis.

For liver, approximately 50 mg was dissected for analyses in a 2 mL FastPrep
10 tubes (MP Biomedicals, Solon OH). ¼” ceramic sphere (MP Biomedicals) is added to each tube and 500 µL of Glo Lysis Buffer – GLB (Promega, Madison WI) equilibrated to room temperature is added to liver tissue. Liver tissues were homogenized with the FastPrep24 instrument (MP Biomedicals) at 2×6.0 m/s for 15 seconds. Homogenate was incubated at room temperature for 5 minutes prior to a 1:4 dilution in GLB and
15 assessed using SteadyGlo Luciferase assay system (Promega). Specifically, 50 µL of diluted tissue homogenate was reacted with 50 µL of SteadyGlo substrate, shaken for 10 seconds followed by 5 minute incubation and then quantitated using a CentroXS³ LB 960 luminometer (Berthold Technologies, Germany). The amount of protein assayed was determined by using the BCA protein assay kit (Pierce, Rockford IL). Relative
20 luminescence units (RLU) were then normalized to total µg protein assayed. To convert RLU to ng luciferase a standard curve was generated with QuantiLum Recombinant Luciferase (Promega).

The FLuc mRNA (L-6107 or L-7202) from Trilink Biotechnologies will express a luciferase protein, originally isolated from the firefly, *photinus pyralis*. FLuc is
25 commonly used in mammalian cell culture to measure both gene expression and cell viability. It emits bioluminescence in the presence of the substrate, luciferin. This capped and poly-adenylated mRNA was fully substituted with respect to uridine and/or cytidine nucleosides.

EXAMPLE 2

DETERMINATION OF pK_a OF FORMULATED LIPIDS

As described elsewhere, the pK_a of formulated cationic lipids is correlated with the effectiveness of LNPs for delivery of nucleic acids (see Jayaraman et al, *Angewandte Chemie*, International Edition (2012), 51(34), 8529-8533; Semple et al, Nature Biotechnology 28, 172-176 (2010)). The preferred range of pK_a is ~5 to ~7. The pK_a of each cationic lipid was determined in lipid nanoparticles using an assay based on fluorescence of 2-(p-toluidino)-6-naphthalene sulfonic acid (TNS). Lipid nanoparticles comprising cationic lipid/DSPC/cholesterol/PEG-lipid (50/10/38.5/1.5 mol%) in PBS at a concentration of 0.4 mM total lipid were prepared using the in-line process as described in Example 1. TNS was prepared as a 100 μ M stock solution in distilled water. Vesicles were diluted to 24 μ M lipid in 2 mL of buffered solutions containing, 10 mM HEPES, 10 mM MES, 10 mM ammonium acetate, 130 mM NaCl, where the pH ranged from 2.5 to 11. An aliquot of the TNS solution was added to give a final concentration of 1 μ M and following vortex mixing fluorescence intensity was measured at room temperature in a SLM Aminco Series 2 Luminescence Spectrophotometer using excitation and emission wavelengths of 321 nm and 445 nm. A sigmoidal best fit analysis was applied to the fluorescence data and the pK_a was measured as the pH giving rise to half-maximal fluorescence intensity.

20

EXAMPLE 3

DETERMINATION OF EFFICACY OF LIPID NANOPARTICLE FORMULATIONS





CONTAINING VARIOUS CATIONIC LIPIDS USING AN IN VIVO






LUCIFERASE MRNA EXPRESSION RODENT MODEL




Representative compounds of the disclosure shown in Table 2 were formulated using the following molar ratio: 50% cationic lipid / 10% distearoylphosphatidylcholine (DSPC) / 38.5% Cholesterol / 1.5% PEG lipid 2-[2-(ω -methoxy(polyethyleneglycol₂₀₀₀)ethoxy]-N,N-ditetradecylacetamide) or 47.5% cationic lipid / 10% DSPC / 40.7% Cholesterol / 1.8% PEG lipid. Relative activity was determined by measuring luciferase expression in the liver 4 hours following administration via tail vein injection as described in Example 1. The activity was

compared at a dose of 1.0 or 0.5 mg mRNA/kg and expressed as ng luciferase/g liver measured 4 hours after administration, as described in Example 1. Compound numbers in Table 2 refer to the compound numbers of Table 1.

Table 2. Novel Cationic Lipids and Associated Activity

Cmp. No.	pKa	Liver Luc @ 1.0 mg/kg (ng luc/g liver)	Structure
I-1	6.11	24031 ± 4777	
I-2	6.67	122 ± 18* * determined at 0.3 mg/kg 682 ± 147** ** determined at 1.0 mg/kg	
I-4	6.17	46932 ± 17399	
I-5	6.08	11720 ± 2439	


Cmp. No.	pKa	Liver Luc @ 1.0 mg/kg (ng luc/g liver)	Structure
I-6	6.22	$4093 \pm 1036^*$ * determined at 0.3 mg/kg $11290 \pm 6455^{**}$ ** determined at 1.0 mg/kg	
I-14	6.80	$669 \pm 575^*$ * determined at 0.5 mg/kg	
I-15	5.96	21357 ± 15325	
I-16	6.25	66763 ± 15823	
I-17	6.78	26691 ± 13186	

Cmp. No.	pKa	Liver Luc @ 1.0 mg/kg (ng luc/g liver)	Structure
I-18	6.02	40650 ± 11479	
I-19	5.95	32706 ± 4621	
I-20	6.48	4140 ± 1117* * determined at 0.3 mg/kg 17095 ± 8181** ** determined at 1.0 mg/kg	

Cmp. No.	pKa	Liver Luc @ 1.0 mg/kg (ng luc/g liver)	Structure
I-21	6.40	$7723 \pm 1714^*$ * determined at 0.3 mg/kg $20223 \pm 5982^{**}$ ** determined at 1.0 mg/kg	
I-22	6.28	$5291 \pm 1348^*$ * determined at 0.3 mg/kg $17654 \pm 8167^{**}$ ** determined at 1.0 mg/kg	
I-23	6.33	$553 \pm 153^*$ * determined at 0.5 mg/kg	

EXAMPLE 4

SYNTHESIS OF BIS(2-BUTYLOCTYL) 7-((3-(DIMETHYLAMINO)PROPYL)(3-(DIOCTYLAMINO)-3-OXOPROPYL)AMINO)TRIDECANEDIOATE (COMPOUND I-7)

5 Synthesis of acryloyl chloride

Acrylic acid (1.20 g, 16.65 mmol) was dissolved in 20mL anhydrous dichloromethane. Thionyl chloride (1.98 g, 16.65 mmol) was added dropwise while stirring under N_2 , and the reaction mixture was allowed to heat to reflux for 4 hr. After the completion of reaction, the crude product concentrated to yield a pale yellow liquid

10 that used in the next step without further purification.

Synthesis of N,N-Dioctylacrylamide (Intermediate A)

Acryloyl chloride (1.12 g, 12.37 mmol) was added to a cooled solution (0 °C) of dioctylamine in dichloromethane with triethylamine (1 equiv) as a base. The reaction mixture was stirred at 0 °C for 1 h and additional 1 h at room temperature. The reaction mixture was filtered and the solution obtained was washed with hydrochloric acid (1N HCl), then with sat NaHCO₃ solution and brine. The solvent evaporated under reduced pressure to yield crude product (colorless liquid) which was used in the following step without further purification.

Synthesis of bis(2-butyloctyl) 7-oxotridecanedioate

To a solution of 2-butyloctane-1-ol (3.85 g, 20.66 mmol), 7-oxotridecanedioic acid (1.34 g, 5.17 mmol) and 4-dimethylaminopyridine (DMAP) (1.9 g, 15.55 mmol) in anhydrous DCM was added DCC (4.27 g, 20.69 mmol). The resulting mixture was allowed to stir overnight at room temperature. The solid (DCU) was then filtered and washed with DCM. The filtrate was concentrated. The residue (oil/solid) was purified by column chromatography on silica gel (0-5% ethyl acetate in hexane). The desired product was obtained as a colorless oil (2.55 g, 42.86 mmol, 83%).

Synthesis of bis(2-butyloctyl) 7-((3-(dimethylamino)propyl)amino)tridecanedioate

A solution of 3-(dimethylamino)-1-propylamine (0.09 g, 0.88 mmol) and bis(2-butyloctyl) 7-oxotridecanedioate (0.37 g, 0.63 mmol) in DCE was treated with sodium triacetoxyborohydride (0.20 g, 0.94 mmol) and AcOH (55 µL, 0.98 mmol) overnight. The solution was washed with dilute aqueous sodium hydroxide solution (1 N NaOH). The organic phase was washed with brine, dried over anhydrous sodium sulfate, filtered and the solvent removed. The residue was passed down a small pad of silica gel, washed with a mixture of DCM/MeOH/Et₃N (85:15:1). The filtrate was concentrated to give the desired product as a slightly yellow oil (240 mg, 0.35 mmol, 56%).

Synthesis of I-7


An EtOH (10 mL) solution of bis(2-butyloctyl) 7-((3-(dimethylamino)propyl)amino)tridecanedioate (210 mg, 0.30 mmol) and N,N-

Diocetylacrylamide (1.5 eq, 136 mg, 0.46 mmol) was stirred at room temperature overnight. The reaction mixture was heated to reflux for 7 days. Upon completion of the reaction, the solvent was removed. The residue was dissolved in a mixture of hexanes and EtOAc (19:1) and washed with saturated sodium bicarbonate solution, and
5 brine. The extract was dried over sodium sulfate. The dried extract was filtered through a pad of silica gel. The pad was washed with a mixture of hexane/ethyl acetate/triethylamine (80:20:1). The washing was concentrated to give the crude desired product.

The crude product was further purified by flash dry column chromatography on
10 silica gel (MeOH in chloroform, 0 to 5%). This gave the desired product as a colorless oil (30 mg, 0.03 mmol, 10%). ¹HNMR (400 MHz, CDCl₃) δ: 3.96 (d, 5.8 Hz, 4H), 3.30-3.16 (m, 4H), 2.74 (t, 7.2 Hz, 2H), 2.65-2.24 (m, 17H), 1.80-1.44 (m, 12H), 1.43-1.15 (64H), 0.93-0.82 (m, 18H).

EXAMPLE 5

SYNTHESIS OF BIS(2-HEXYLDECYL) 7-((4-(DIHEXYLAMINO)-4-OXOBUTYL)(2-(DIMETHYLAMINO)ETHYL)AMINO)TRIDECANEDIOATE (COMPOUND I-19)

5 Synthesis of N,N-dihexyl-4-oxobutanamide (Intermediate B)

Butyrolactone (2.51 g, 29.15 mmol) and dihexylamine (5.40 g, 29.13 mmol) were heated for 4 days in a pressure flask at 61 °C. The reaction mixture was cooled to room temperature. The crude product was purified by column chromatography on silica gel (0% to 5% of MeOH in DCM) to yield N,N-dihexyl-4-hydroxybutanamide as a slightly yellow oil (6.30 g, 79%).

N,N-dihexyl-4-hydroxybutanamide (3.00 g, 11.05 mmol) was dissolved in DCM and treated with pyridinium chlorochromate (2.38 g, 11.05 mmol) for two hours. Diethyl ether was added and the supernatant filtered through silica gel bed. The solvent was removed from the filtrate and resultant oil dissolved in hexane. The suspension was filtered through silica gel bed and the solvent removed. The crude product (colorless liquid) was used in the following step without further purification.

Synthesis of I-19

A solution of N,N-dihexyl-4-oxobutanamide (0.56 g, 1.97 mmol), and bis(2-hexyldecyl) 7-((2-(dimethylamino)ethyl)amino)tridecanedioate (0.44 g, 0.56 mmol, prepare according to procedures of Example 4) in 1,2-dichloroethane (10 mL) was stirred for 15 min, after which time sodium triacetoxyborohydride (0.41 g, 1.97 mmol) was added in one portion and stirred at room temperature for additional 16 hours. The mixture was concentrated. The residue was taken up in a mixture of hexane and ethyl acetate (96:4) and washed with saturated aqueous NaHCO₃ solution and brine. The organic layer was separated, dried over anhydrous sodium sulphate, filtered and evaporated under reduced pressure to obtain colorless oil. The crude product was purified by flash column chromatography on silica gel (MeOH in chloroform, 0 to 5%) to yield the desired product as a colorless oil (260 mg, 0.25 mmol, 45%). ¹HNMR (400 MHz, CDCl₃) δ: 3.96 (d, 5.8 Hz, 4H), 3.28 (t-like, 7.7 Hz, 2H), 3.20 (t-like, 7.7 Hz, 2H), 2.56-2.47 (m, 2H), 2.44 (t, 6.8 Hz, 2H), 2.39-2.20 (m, 15H), 1.74-1.45 (m, 12H), 1.42-1.15 (72H), 0.93-0.84 (m, 18H).

EXAMPLE 6

SYNTHESIS OF BIS(2-BUTYLOCTYL) 10-((4-(DIHEXYLAMINO)-4-OXOBUTYL)(3-(DIMETHYLAMINO)PROPYL)AMINO)NONADECANEDIOATE (COMPOUND I-21)


Compound I-21 was prepared according to the general procedures of Example 5 to yield 0.05 g of colorless oil, 0.03 mmol, 32%. ¹HNMR (400 MHz, CDCl₃) δ: 3.97 (d, 5.8 Hz, 4H), 3.28 (t-like, 7.6 Hz, 2H), 3.20 (t-like, 7.6 Hz, 2H), 2.43-2.23 (m, 13H), 2.20 (s, 6H), 1.75-1.45 (m, 14H), 1.40-1.12 (m, 68H), 0.93-0.84 (m, 18H).

EXAMPLE 7

SYNTHESIS OF BIS(2-BUTYLOCTYL) 7-((4-(DIHEXYLAMINO)-4-OXOBUTYL)(3-(DIMETHYLAMINO)PROPYL)AMINO)TRIDECANEDIOATE (COMPOUND I-20)

Compound I-20 was prepared according to the general procedures of Example 5 to yield 0.06 g of colorless oil, 0.06 mmol, 41%. ¹HNMR (400 MHz, CDCl₃) δ: 3.96 (d, 5.8 Hz, 4H), 3.27 (t-like, 7.6 Hz, 2H), 3.19 (t-like, 7.6 Hz, 2H), 2.62-2.17 (m, 19H), 1.79-1.43 (m, 14H), 1.42-1.10 (m, 56H), 0.95-0.81 (m, 18H).

EXAMPLE 8

SYNTHESIS OF BIS(2-BUTYLOCTYL) 10-(N-(3-(DIMETHYLAMINO)PROPYL)-6-METHOXY-6-
OXOHEXANAMIDO)NONADECANEDIOATE (COMPOUND I-14)

- 5 Adipoyl chloride (0.12 g, 0.68 mmol) in anhydrous benzene (5 mL) was added via syringe to a solution of bis(2-butyloctyl) 10-((3-(dimethylamino)propyl)amino)nonanedioate (0.26 g, 0.34 mmol, prepared according to Example 4), triethylamine (0.3 mL, 2.5 mmol) and DMAP (5 mg) in benzene (10 mL) at RT over the period of 5 min. The mixture was allowed to stir for 2
- 10 h and then Methanol (0.5 mL) was added to remove excess acyl chloride. The resulting mixture was stirred for another hour and then filtered through a pad of silica gel, washed with a mixture of hexane/EtOAc/Et₃N (70:30:1) and concentrated. The residue was passed down a silica gel column (0-4% MeOH in DCM gradient), yielding compound **I-14** as a colorless oil (0.28 g, 0.30 mmol, 89%). ¹HNMR (400 MHz, CDCl₃) δ: 4.52-4.29 (br., estimated 0.3H, due to slow isomerization about amide bond), 3.96 (d, 5.8 Hz, 4H), 3.65 (s, 3H), 3.59 (quintet-like, 7.0 Hz, 0.7H), 3.14-3.05 (m, 2H), 2.37-2.24 (m, 10H), 2.23-2.18 (m, 6H), 1.73-1.54 (m, 12H), 1.48-1.37 (m, 4H), 1.34-1.14 (m, 52H), 0.93-0.83 (m, 12H).
- 15

EXAMPLE 9

SYNTHESIS OF BIS(2-BUTYLOCTYL) 10-(N-(2-(DIMETHYLAMINO)ETHYL)-6-METHOXY-6-
OXOHEXANAMIDO)NONADECANEDIOATE (COMPOUND I-15)

Compound I-15 was prepared according to the general procedures of Example 8,
5 to yield 0.18 g of colorless oil, 0.20 mmol, 85%. ¹HNMR (400 MHz, CDCl₃) δ: 4.53-
4.30 (br., 0.3H, due to slow isomerization about amide bond), 3.96 (d, 5.8 Hz, 4H), 3.65
(s, 3H), 3.58 (quintet-like, 7 Hz, 0.7H), 3.27-3.15 (m, 2H), 2.46-2.22 (m, 16H), 1.75-
1.54 (m, 10H), 1.50-1.36 (m, 4H), 1.35-1.09 (m, 52H), 0.94-0.82 (m, 12H).

EXAMPLE 10

10 SYNTHESIS OF BIS(2-HEXYLDECYL) 7-(N-(2-(DIMETHYLAMINO)ETHYL)-6-METHOXY-6-
OXOHEXANAMIDO)TRIDECANEDIOATE (COMPOUND I-16)

Compound I-16 was prepared according to the general procedures of Example 8
to yield 0.27 g of colorless oil, 0.29 mmol, 81%. ¹HNMR (400 MHz, CDCl₃) δ: 4.53-
4.30 (br., 0.3H, due to slow isomerization about amide bond), 3.99-3.92 (m, 4H), 3.66
15 (s, 3H), 3.59 (quintet-like, 7.0 Hz, 0.7H), 3.28-3.14 (m, 2H), 2.46-2.20 (m, 16H), 1.75-
1.53 (m, 10H), 1.51-1.36 (m, 4H), 1.35-1.09 (m, 56H), 0.94-0.81 (m, 12H).

EXAMPLE 11

SYNTHESIS OF BIS(2-HEXYLDECYL) 7-(N-(3-(DIMETHYLAMINO)PROPYL)-8-METHOXY-8-
OXOOCTANAMIDO)TRIDECANEDIOATE (COMPOUND I-17)

20 Compound I-17 was prepared according to the general procedures of Example 8
to yield 0.08 g of colorless oil, 0.08 mmol, 75%. ¹HNMR (400 MHz, CDCl₃) δ: 4.53-
4.30 (br., 0.3H, due to slow isomerization about amide bond), 3.99-3.91 (m, 4H), 3.66
(s, 3H), 3.61 (quintet-like, 7.0 Hz, 0.7H), 3.15-3.06 (m, 2H), 2.34-2.23 (m, 10H), 2.22
(s, 6H), 1.75-1.53 (m, 10H), 1.51-1.38 (m, 4H), 1.37-1.15 (m, 62H), 0.93-0.82 (m,
25 12H).


EXAMPLE 12

SYNTHESIS OF BIS(2-HEXYLDECYL) 7-(N-(2-(DIMETHYLAMINO)ETHYL)-8-METHOXY-8-
OXOOCTANAMIDO)TRIDECANEDIOATE (COMPOUND I-18)

Compound I-18 was prepared according to the general procedures of Example 8
5 to yield 0.15 g of colorless oil, 0.16 mmol, 79%. ¹HNMR (400 MHz, CDCl₃) δ: 4.53-
4.30 (br., 0.3H, due to slow isomerization about amide bond), 3.99-3.92 (m, 4H), 3.66
(s, 3H), 3.61 (quintet-like, 7.0 Hz, 0.7H), 3.26-3.14 (m, 2H), 2.47-2.35 (m, 2H), 2.34-
2.20 (m, 14H), 1.73-1.53 (m, 8H), 1.51-1.39 (m, 4H), 1.38-1.14 (m, 62H), 0.93-0.82
(m, 12H).

10

EXAMPLE 13

SYNTHESIS OF BIS(2-BUTYLOCTYL) 10-(N-(3-(DIMETHYLAMINO)PROPYL)-6-METHOXY-6-
OXOHEXANAMIDO)NONADECANEDIOATE (COMPOUND I-1)

Synthesis of Intermediate C

To a solution of acrylic acid (1.1 eq, 8.25 mmol, 594 mg), octanol (1 eq, 975 mg, 7.5 mmol) and DMAP (0.4 eq, 3 mmol, 366 mg) in DCM (15 mL) was added DCC (1.4 eq, 10.5 mmol, 2.16 g). The resulting mixture was stirred at RT for 16h. The reaction mixture was filtered and the filtrate concentrated. The residue was taken up in hexanes (50 mL) and loaded on a column of silica gel. The column was washed with hexane (40 mL). The fractions were combined, re-loaded on the column and eluted with a mixture of hexane and ethyl acetate (ca 99:1 or 98:2, 200 mL). A colorless oil was obtained (986 mg, 71%).

10 Synthesis of Compound I-1

An EtOH (10 mL) solution of bis(2-butyloctyl) 10-((4-(dimethylamino)butyl)amino)nonadecanedioate (1 eq., 220 mg, 0.28 mmol, prepared according to general procedures above), and intermediate C (2.75 eq. 0.77 mmol, 140 mg) in a sealed pressure flask was stirred at RT for 4 days under Ar. The reaction mixture was concentrated. The residue was purified twice by flash dry column chromatography on silica gel (hexane-EtOAc-Et₃N, 95:5:0 to 80:20:1 and MeOH in chloroform, 0 to 5%). The desired product was obtained as colorless oil (68 mg, 0.07 mmol, 25%) ¹HNMR (400 MHz, CDCl₃) δ: 4.03 (t, 6.9 Hz, 2H), 3.97 (d, 5.8 Hz, 4H), 2.69 (t, 7.2 Hz, 2H), 2.38-2.33 (m, 4H), 2.33-2.26 (m, 1H), 2.29 (t, 7.5 Hz, 4H), 2.26-2.22 (m, 2H), 2.21 (s, 6H), 1.61 (quintet-like, 7.0 Hz, 8H), 1.48-1.08 (70H), 0.92-0.86 (m, 15H).

EXAMPLE 14

SYNTHESIS OF BIS(2-BUTYLOCTYL) 10-((5-(DIMETHYLAMINO)PENTYL)(3-(OCTYLOXY)-3-OXOPROPYL)AMINO)NONADECANEDIOATE (COMPOUND I-2)

25 Compound I-2 was prepared according to the general procedures of Example 13 to yield 0.05 g of colorless oil, 0.05 mmol, 10%. ¹HNMR (400 MHz, CDCl₃) δ: 4.03 (t, 6.8 Hz, 2H), 3.96 (d, 5.8 Hz, 4H), 2.69 (t, 7.2 Hz, 2H), 2.38-2.21 (m, 17H), 1.61 (quintet-like, 7.0 Hz, 8H), 1.51-1.10 (m, 72H), 0.93-0.84 (m, 15H).

EXAMPLE 15

SYNTHESIS OF BIS(2-BUTYLOCTYL) 7-((4-(DIMETHYLAMINO)BUTYL)(3-(OCTYLOXY)-3-OXOPROPYL)AMINO)TRIDECANEDIOATE (COMPOUND I-3)

Compound I-3 was prepared according to the general procedures of Example 13 to yield 0.01 g of colorless oil, 0.01 mmol, 11%. ¹HNMR (400 MHz, CDCl₃) δ: 4.03 (t, 6.9 Hz, 2H), 3.96 (d, 5.6 Hz, 4H), 2.69 (t, 7.1 Hz, 2H), 2.38-2.20 (m, 17H), 1.69-1.56 (m, 10H), 1.48-1.09 (m, 56H), 0.92-0.84 (m, 15H).


EXAMPLE 16

SYNTHESIS OF BIS(2-HEXYLDECYL) 7-((4-(DIMETHYLAMINO)BUTYL)(3-(OCTYLOXY)-3-OXOPROPYL)AMINO)TRIDECANEDIOATE (COMPOUND I-4)

Compound I-4 was prepared according to the general procedures of Example 13 to yield 0.05 g of colorless oil, 0.05 mmol, 13%. ¹HNMR (400 MHz, CDCl₃) δ: 4.03 (t, 6.8 Hz, 2H), 3.96 (d, 5.8 Hz, 4H), 2.69 (t, 7.1 Hz, 2H), 2.39-2.21 (m, 17H), 1.66-1.09 (m, 82H), 0.88 (t, 7.0 Hz, 15H).

EXAMPLE 17

SYNTHESIS OF BIS(2-BUTYLOCTYL) 10-((4-(DIMETHYLAMINO)BUTYL)(OCTYL)AMINO)NONADECANEDIOATE (COMPOUND I-23)

5 Synthesis of Compound I-23

A solution of octanal (3.5 eq, 0.90 mmol, 115 mg, 0.141 mL) and bis(2-butyloctyl) 10-((4-(dimethylamino)butyl)amino)nonadecanedioate (200 mg, 0.26 mmol, prepared according to general procedures above) in 1,2-dichloroethane (5 mL) was stirred for 15 min, after which time sodium triacetoxyborohydride (3.5 eq, 0.9 mmol, 190 mg) was added in one portion. Stirring was continued at RT for 16 hours. The reaction mixture was concentrated. The residue was purified twice by flash dry column chromatography on silica gel (hexane-EtOAc-Et₃N, 95:5:0 to 80:20:1 and MeOH in chloroform, 0 to 5%). The desired product was obtained as colorless oil (203 mg, 0.23 mmol, 88%). ¹HNMR (400 MHz, CDCl₃) δ: 3.97 (d, 5.8 Hz, 4H), 2.40-2.18 (m, 17H), 1.69-1.56 (m, 6H), 1.52-1.10 (m, 72H), 0.92-0.86 (m, 15H).

EXAMPLE 18

SYNTHESIS OF BIS(2-ETHYLHEXYL) 10-((4-(DIMETHYLAMINO)BUTYL)(6-((2-
HEXYLDECANOYL)OXY)HEXYL)AMINO)NONADECANEDIOATE (COMPOUND I-22)

Compound I-22 was prepared according to the general procedures of Example
5 17 to yield 0.19 g of colorless oil, 0.19 mmol, 80%. ¹HNMR (400 MHz, CDCl₃) δ: 4.06 (t, 6.7 Hz, 2H), 3.97 (d, 5.6 Hz, 4H), 2.39-2.26 (m, 11H), 2.23 (s, 6H), 1.68-1.10 (m, 83H), 0.94-0.82 (m, 18H).

EXAMPLE 19

SYNTHESIS OF BIS(2-BUTYLOCTYL) 10-((4-(DIMETHYLAMINO)BUTYL)(6-((2-
10 HEXYLDECANOYL)OXY)HEXYL)AMINO)NONADECANEDIOATE (COMPOUND I-5)


Compound I-5 was prepared according to the general procedures of Example 17
to yield 0.04 g of colorless oil, 0.04 mmol, 73%. ¹HNMR (400 MHz, CDCl₃) δ: 4.05 (t, 7.0 Hz, 2H), 3.96 (d, 5., 8 Hz, 4H), 2.38-2.26 (m, 11H), 2.23 (s, 6H), 1.71-1.09 (m, 99H), 0.95-0.82 (m, 18H).

15

The various embodiments described above can be combined to provide further
embodiments. All of the U.S. patents, U.S. patent application publications, U.S. patent
applications, foreign patents, foreign patent applications and non-patent publications
referred to in this specification, including but not limited to U.S. Provisional Patent
20 Application No. 63/052,815, filed July 16, 2020, and U.S. Provisional Patent
Application No. 63/188,996, filed May 14, 2021, are incorporated herein by reference,
in their entirety. Aspects of the embodiments can be modified, if necessary to employ
concepts of the various patents, applications and publications to provide yet further
embodiments. These and other changes can be made to the embodiments in light of the
25 above-detailed description. In general, in the following claims, the terms used should
not be construed to limit the claims to the specific embodiments disclosed in the
specification and the claims, but should be construed to include all possible
embodiments along with the full scope of equivalents to which such claims are entitled.
Accordingly, the claims are not limited by the disclosure.

CLAIMS

1. A compound having a structure of Formula (I):



(I)

or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof,

wherein:

G^1 and G^2 are each independently C_1 - C_6 alkylene;

L^1 and L^2 are each independently $-O(C=O)-$ or $-(C=O)O-$;

R^{1a} and R^{1b} are, at each occurrence, independently either: (a) H or C_1 - C_{12} alkyl; or (b) R^{1a} is H or C_1 - C_{12} alkyl, and R^{1b} together with the carbon atom to which it is bound is taken together with an adjacent R^{1b} and the carbon atom to which it is bound to form a carbon-carbon double bond;

R^{2a} and R^{2b} are, at each occurrence, independently either: (a) H or C_1 - C_{12} alkyl; or (b) R^{2a} is H or C_1 - C_{12} alkyl, and R^{2b} together with the carbon atom to which it is bound is taken together with an adjacent R^{2b} and the carbon atom to which it is bound to form a carbon-carbon double bond;

R^{3a} and R^{3b} are, at each occurrence, independently either (a): H or C_1 - C_{12} alkyl; or (b) R^{3a} is H or C_1 - C_{12} alkyl, and R^{3b} together with the carbon atom to which it is bound is taken together with an adjacent R^{3b} and the carbon atom to which it is bound to form a carbon-carbon double bond;

R^{4a} and R^{4b} are, at each occurrence, independently either: (a) H or C_1 - C_{12} alkyl; or (b) R^{4a} is H or C_1 - C_{12} alkyl, and R^{4b} together with the carbon atom to which it is bound is taken together with an adjacent R^{4b} and the carbon atom to which it is bound to form a carbon-carbon double bond;

R^5 and R^6 are each independently H or methyl;

R^7 is $-O(C=O)R^{10}$, $-(C=O)OR^{10}$, $-NR^9(C=O)R^{10}$ or $-(C=O)NR^9R^{10}$;
 R^8 is OH, $-N(R^{11})(C=O)R^{12}$, $-(C=O)NR^{11}R^{12}$, $-NR^{11}R^{12}$, $-(C=O)OR^{12}$ or
 $-O(C=O)R^{12}$;

R^9 is H or C₁-C₁₅ alkyl;

R^{10} is C₁-C₁₅ alkyl;

R^{11} is H or C₁-C₆ alkyl;


R^{12} is C₁-C₆ alkyl;

X is $-(C=O)-$ or a direct bond; and

a, b, c and d are each independently an integer from 1 to 24;

wherein each alkyl and alkylene and is independently optionally substituted.

2. The compound of claim 1, having one of the following structures (IA) or (IB):



or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof.

3. The compound of claim 1 or 2, wherein G^1 is C₂-C₃ alkylene.
4. The compound of claim 1 or 2, wherein G^1 is C₄-C₆ alkylene.
5. The compound of any one of claims 1-4, wherein G^2 is C₂-C₄ alkylene.
6. The compound of claim 5, wherein G^2 is C₂-C₃ alkylene or C₃-C₄ alkylene.

7. The compound of any one of claims 1-6, wherein X is $-(C=O)-$.
8. The compound of any one of claims 1-6, wherein X is a direct bond.
9. The compound of any one of claims 1-8, wherein R^7 is $-O(C=O)R^{10}$ or $-(C=O)OR^{10}$.
10. The compound of claim 9, wherein R^{10} is linear C_1 - C_{15} alkyl.
11. The compound of claim 10, wherein R^{10} is linear C_6 - C_{10} alkyl.
12. The compound of claim 10, wherein R^{10} is methyl.
13. The compound of claim 10, wherein R^{10} is branched C_2 - C_{15} alkyl.
14. The compound of claim 13, wherein R^{10} is branched C_{10} - C_{15} alkyl.
15. The compound of any one of claims 1-8, wherein R^7 is $-NR^9(C=O)$ or $-(C=O)NR^9R^{10}$.
16. The compound of claim 15, wherein R^9 is H.
17. The compound of claim 15, wherein R^9 and R^{10} are each independently C_6 - C_{10} alkyl.
18. The compound of any one of claims 1-17, wherein for at least one occurrence of R^{1a} and R^{1b} , R^{1a} is H or C_1 - C_{12} alkyl, and R^{1b} together with the carbon atom to which it is bound is taken together with an adjacent R^{1b} and the carbon atom to which it is bound to form a carbon-carbon double bond.

19. The compound of any one of claims 1-18, wherein for at least one occurrence of R^{4a} and R^{4b} , R^{4a} is H or C₁-C₁₂ alkyl, and R^{4b} together with the carbon atom to which it is bound is taken together with an adjacent R^{4b} and the carbon atom to which it is bound to form a carbon-carbon double bond.

20. The compound of any one of claims 1-19, wherein for at least one occurrence of R^{2a} and R^{2b} , R^{2a} is H or C₁-C₁₂ alkyl, and R^{2b} together with the carbon atom to which it is bound is taken together with an adjacent R^{2b} and the carbon atom to which it is bound to form a carbon-carbon double bond.

21. The compound of any one of claims 1-20, wherein for at least one occurrence of R^{3a} and R^{3b} , R^{3a} is H or C₁-C₁₂ alkyl, and R^{3b} together with the carbon atom to which it is bound is taken together with an adjacent R^{3b} and the carbon atom to which it is bound to form a carbon-carbon double bond.


22. The compound of any one of claims 1-17, wherein R^{1a} , R^{1b} , R^{2a} , R^{2b} , R^{3a} , R^{3b} , R^{4a} and R^{4b} are, at each occurrence, independently H or C₁-C₁₂ alkyl.

23. The compound of claim 22, wherein R^{2a} , R^{2b} , R^{3a} and R^{3b} are, at each occurrence, H.

24. The compound of claim 22 or 23, wherein R^{1a} and R^{4a} are, at each occurrence, H.

25. The compound of any one of claims 22-24, wherein at least one of R^{1b} and R^{4b} is C₁-C₈ alkyl.

26. The compound of claim 25, wherein C₁-C₈ alkyl is methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, tert-butyl, n-hexyl or n-octyl.



28. The compound of any one of claims 1-27, wherein a, b, c and d are each independently an integer from 2 to 12

29. The compound of any one of claims 1-27, wherein a, b, c and d are each independently an integer from 4 to 10, 5 to 10, 6 to 10, 4 to 9, 5 to 9 or 6 to 9.


30. The compound of any one of claims 1-29, wherein one of R^5 or R^6 is methyl.

31. The compound of any one of claims 1-30, wherein each of R^5 and R^6 is methyl.

32. The compound of any one of claims 1-31, wherein R^8 is OH.

33. The compound of any one of claims 1-31, wherein R^8 is - $N(R^{11})(C=O)R^{12}$.


34. The compound of any one of claims 1-31, wherein R^8 is $-(C=O)NR^{11}R^{12}$.
35. The compound of any one of claims 1-31, wherein R^8 is $-NR^{11}R^{12}$.
36. The compound of any one of claims 33-35, wherein R^{11} and R^{12} are each independently H or C₁-C₈ alkyl.
37. The compound of any one of claims 33-36, wherein R^{11} and R^{12} are each independently H or C₁-C₃ alkyl.
38. The compound of claim 36 or 37, wherein C₁-C₈ alkyl or C₁-C₃ alkyl is unsubstituted or substituted with hydroxyl.
39. The compound of any one of claims 33-36, wherein R^{11} and R^{12} are each methyl.
40. The compound of any one of claims 1-31, wherein R^8 is $-(C=O)OR^{12}$.
41. The compound of claim 40, wherein R^8 is $-O(C=O)R^{12}$.
42. The compound of any one of claims 1-31, wherein R^8 has one of the following structures:



43. A compound selected from a compound in Table 1.
44. A lipid nanoparticle comprising the compound of any one of claims 1-43 and a therapeutic agent.

45. The lipid nanoparticle of claim 44, further comprising one or more excipient selected from neutral lipids, steroids and polymer conjugated lipids.
46. The lipid nanoparticle of claim 45, wherein the composition comprises one or more neutral lipids selected from DSPC, DPPC, DMPC, DOPC, POPC, DOPE and SM.
47. The lipid nanoparticle of claim 46, wherein the neutral lipid is DSPC.
48. The lipid nanoparticle of any one of claims 44-47, wherein the molar ratio of the compound to the neutral lipid ranges from about 2:1 to about 8:1.
49. The lipid nanoparticle of any one of claims 45-48, wherein the steroid is cholesterol.
50. The lipid nanoparticle of claim 49, wherein the molar ratio of the compound to cholesterol ranges from 5:1 to 1:1.
51. The lipid nanoparticle of any one of claims 45-50, wherein the polymer conjugated lipid is pegylated lipid.
52. The lipid nanoparticle of claim 51, wherein the molar ratio of the compound to pegylated lipid ranges from about 100:1 to about 20:1.
53. The lipid nanoparticle of anyone of claims 51 or 52, wherein the pegylated lipid is PEG-DAG, PEG-PE, PEG-S-DAG, PEG-cer or a PEG dialkoxypolypropylcarbamate.

54. The lipid nanoparticle of any one of claims 51 or 52, wherein the pegylated lipid has the following structure (II):



(II)

or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein:

R⁸ and R⁹ are each independently a straight or branched, alkyl, alkenyl or alkynyl from 10 to 30 carbon atoms, wherein the alkyl, alkenyl or alkynyl is optionally interrupted by one or more ester bonds; and

w has a mean value ranging from 30 to 60.

55. The lipid nanoparticle of claim 54, wherein R⁸ and R⁹ are each independently straight alkyl chain containing from 12 to 16 carbon atoms.

56. The lipid nanoparticle of any one of claims 54 or 55, wherein the average w is about 49.

57. The lipid nanoparticle of any one of claims 44-56, wherein the therapeutic agent comprises a nucleic acid.

58. The lipid nanoparticle of claim 57, wherein the nucleic acid is selected from antisense and messenger RNA.

59. A pharmaceutical composition comprising the lipid nanoparticle of any one of claims 44-58 and a pharmaceutically acceptable diluent or excipient.

60. A method for treating or preventing a disease in a patient in need thereof, the method comprising administering the lipid nanoparticle of any one of claims 44-58, or the pharmaceutical composition of claim 59, to the patient.

61. A method for vaccinating a patient in need thereof against a viral pathogen, the method comprising administering the lipid nanoparticle of any one of claims 44-58, or the pharmaceutical composition of claim 59, to the patient, wherein the therapeutic agent is a viral antigen or a nucleic acid capable of transcribing a viral antigen.