
Claims

1. A catalyst compound represented by the Formula (**II**):



(**II**)

wherein:

M is a group 3, 4, 5, or 6 transition metal or a Lanthanide;

E and E' are each independently O;

each L is independently a Lewis base, selected from the group consisting of ethers, thioethers, amines, nitriles, imines, pyridines, halocarbons, and phosphines;

each X is independently an anionic ligand;

n is 1, 2 or 3;

m is 0, 1, or 2;

n+m is not greater than 4;

each of R^1 and $R^{1'}$ is independently a C_4-C_{40} tertiary hydrocarbyl group;

each of R^2, R^3, R^4, $R^{2'}$, $R^{3'}$, and $R^{4'}$ is independently hydrogen, C_1-C_{40} hydrocarbyl, C_1-C_{40} substituted hydrocarbyl, a heteroatom or a heteroatom-containing group, or one or more of R^2, R^2 and R^3, R^3 and R^4, $R^{2'}$, $R^{2'}$ and $R^{3'}$, $R^{3'}$ and $R^{4'}$ may be joined to form one or more substituted hydrocarbyl rings, unsubstituted hydrocarbyl rings, substituted heterocyclic rings, or unsubstituted heterocyclic rings each having 5, 6, 7, or 8 ring atoms, and where substitutions on the ring can join to form additional rings;any two L groups may be joined together to form a bidentate Lewis base;

an X group may be joined to an L group to form a monoanionic bidentate group;

any two X groups may be joined together to form a dianionic ligand group;

each of R^5, R^6, R^7, R^8, $R^{5'}$, $R^{6'}$, $R^{7'}$, $R^{8'}$, R^{10}, R^{11}, and R^{12} is independently hydrogen, a C_1-C_{40} hydrocarbyl, a C_1-C_{40} substituted hydrocarbyl, a heteroatom or a heteroatom-containing group, or one or more of R^5 and R^6, R^6 and R^7, R^7 and R^8, $R^{5'}$ and $R^{6'}$, $R^{6'}$ and $R^{7'}$, $R^{7'}$ and $R^{8'}$, R^{10} and R^{11}, or R^{11} and R^{12} may be joined to form one or more substituted hydrocarbyl rings,


unsubstituted hydrocarbyl rings, substituted heterocyclic rings, or unsubstituted heterocyclic rings each having 5, 6, 7, or 8 ring atoms, and where substitutions on the ring can join to form additional rings.

2. The catalyst compound claim 1 wherein each X is, independently, selected from the group consisting of substituted or unsubstituted hydrocarbyl radicals having from 1 to 20 carbon atoms, hydrides, amides, alkoxides, sulfides, phosphides, halides, and a combination thereof, or two X's may form a part of a fused ring or a ring system.

3. The catalyst compound of claim 1, wherein each L is independently a Lewis base, selected from ether and thioether groups.

4. The catalyst compound of claim 1, wherein M is Zr or Hf.and both R^1 and $R^{1'}$ are C_4-C_{20} cyclic tertiary alkyls, both R^1 and $R^{1'}$ are adamantan-1-yl or substituted adamantan-1-yl, or both R^1 and $R^{1'}$ are C_6-C_{20} aryls.

5. The catalyst compound of claim 1, wherein the heterocyclic Lewis base is selected from the groups represented by the following formulas:




where each R^{23} is independently selected from hydrogen, C_1-C_{20} alkyls, and C_1-C_{20} substituted alkyls.


6. The catalyst compound of claim 1, wherein M is Zr or Hfand at least on one of the following: (a) R^1 and $R^{1'}$ are C_4-C_{20} cyclic tertiary alkyls or adamantan-1-yl or substituted

adamantan-1-yl; (b) each of R^1, $R^{1'}$, R^3 and $R^{3'}$ are adamantan-1-yl or substituted adamantan-1-yl; and (c) R^1 and $R^{1'}$ are C_4-C_{20} cyclic tertiary alkyls, and both R^7 and $R^{7'}$ are C_1-C_{20} alkyls.


7. The catalyst compound of claim 1 wherein the catalyst compoud is represented by one or more of the following formulas:




Complex 1




Complex 2




Complex 3




Complex 4




Complex 5




Complex 6




Complex 7




Complex 8




Complex 9




Complex 10




Complex 11




Complex 12




Complex 13




Complex 14




Complex 15




Complex 16




Complex 17




Complex 18




Complex 19




Complex 20




Complex 21




Complex 22




Complex 26




Complex 27




Complex 29



Complex 30



Complex 31



Complex 32

8. A process to polymerize olefins comprising: contacting one or more olefins, preferably one or more substituted or unsubstituted C_2 to C_{40} alpha olefins, with a catalyst system including an activator and the catalyst of any one of claims 1 to 7; and obtaining a polymer.

9. The process of claim 8 wherein the process occurs at a temperature of from about 0°C to about 300°C, at a pressure in the range of from about 0.35 MPa to about 10 MPa, and at a time up to 300 minutes.