

Description

Title of Invention: CHICKEN FLAVOR COMPOSITION AND CHICKEN FLAVOR
ENHANCER

5

Technical Field
[0001]

The present invention relates to a chicken flavor composition that contains 1,5-
octadien-3-ol and/or 1,5-octadien-3-one, and to a food, a drink or a seasoning that contains
10 the chicken flavor composition, and further relates to a chicken flavor enhancer that
contains 1,5-octadien-3-ol and/or 1,5-octadien-3-one, and to a food, a drink or a seasoning
that contains the chicken flavor enhancer.

Background Art
15 [0002]

Conventionally, chicken flavor compositions use materials, for example, such as
(1) an animal or a vegetable extract, (2) a yeast extract, (3) an animal or a vegetable protein
hydrolysate, (4) a heating cooking flavor, and (5) a synthetic flavoring, and one or more of
these may be appropriately combined and prepared according to the intended use to obtain
20 a chicken flavor composition (Non Patent Literature 1).

As flavor characteristics desired for a chicken flavor composition, a natural flavor
and a richness characteristic of chicken, and a flavor characteristic that evokes a boiled
flavor, a fried flavor, a roast flavor, or the like that stimulates appetite in harmony with a
dish are required.
25 [0003]

The above-mentioned (2) yeast extract contains relatively larger quantities of an
umami component glutamic acid, and therefore are mainly used to complement umami
(Patent Literatures 1 to 3). It has been investigated to use this expecting an effect on meat
taste and flavor.

30 For example, Patent Literature 4 discloses that a yeast extract containing a free
proline in 8.0% or more of a free amino acid composition can be applied to impart the
characteristic sweetness of animal protein hydrolysates.

[0004]

On the other hand, components that have effects on meat richness and flavor have
35 been investigated. For example, Patent Literature 5 proposes pyrazine compounds as a
seasoning that can impart richness to a food or drink. A seasoning is proposed which
contains, as a pyrazine compound, specifically, one or more compounds selected from a
group consists of 2-methylpyrazine, 2,5-dimethylpyrazine, 2,6-dimethylpyrazine, 2,3,5-

trimethylpyrazine, tetramethylpyrazine, 2,5-diethylpyrazine, 2,6-diethylpyrazine, 2,3-diethyl-5-methylpyrazine, and 2-ethyl-3,5-dimethylpyrazine.

[0005]

Patent Literature 6 proposes, as a stronger and more desirable meat flavor composition, a meat flavor composition characterized in that it contains a flavoring containing at least one or more compound selected from a group consists of (A) pyrroles, (B) pyridines, (C) pyrazines, (D) oxazoles, (E) oxazolines, (F) amines, (G) thiazoles, (H) thiazolines, (I) thiazolidines, (J) thiols, (K) sulfides, (L) thioethers, (M) sulfur-containing carboxylic acids, (N) quinoxalines, and (O) furanones.

10 [0006]

In preparation of a chicken flavor composition by using a synthetic flavoring, an odor analysis data of heat-treated chicken meat and chicken bone are generally used as a reference.

Odor analysis is typically performed by gas chromatography, and the analysis values are generally shown by being sorted in order of odor retention time (hereinafter, also referred to simply as "retention time"), or by being classified according to functional group. Therefore, a flavor preparing person who refers to the odor analysis data often start preparation by selecting various synthetic flavorings referring to the retention time or according to the functional group.

20 [0007]

1,5-Octadien-3-ol is a known compound contained in roast chicken, cauliflower and the like, and is known as a compound which has an odor that evokes a tree aroma or a geranium aroma (Non Patent Literatures 2 and 3). On the other hand, 1,5-octadien-3-one is a known compound contained in beef and the like, and is known to have a metallic odor or a geranium odor (Non Patent Literatures 4 to 8).

25

Prior Art Literatures

Patent Literatures

[0008]

30 Patent Literature 1: JP-A-2005-102549

Patent Literature 2: JP-A-10-327803

Patent Literature 3: WO 99/16860

Patent Literature 4: WO 2008/091519

Patent Literature 5: JP-A-11-313635

35 Patent Literature 6: JP-A-2005-15683

Non Patent Literatures

[0009]

Non Patent Literature 1: Patent Office Publication, Published Collection of Well-Known Prior Arts [Flavor] vol. 2, Food Flavors 3.9.4 Chicken Flavor, published on January 14, 2000

Non Patent Literature 2: Lebensm. Wiss. Technol., 20 (1987), pp. 37-41

5 Non Patent Literature 3: J. Agric. Food Chem., 2002, 50, pp. 6459-6467

Non Patent Literature 4: Journal of food science, Vol.61 (1996), pp.1271-1274

Non Patent Literature 5: Z Lebensm Unters Forsch A (1997) 205: pp.232-238

Non Patent Literature 6: Z Lebensm Unters Forsch A (1997) 204: pp.3-6

Non Patent Literature 7: Z Lebensm Unters Forsch (1988) 186: pp.489-494

10 Non Patent Literature 8: Journal of the Science of Food and Agriculture, Vol.28, 11, pp.1019-1024, November 1977

Summary of Invention

Technical Problem

15 [0010]

However, it is rarely the case that a chicken flavor composition prepared by directly replicating the quantitative analysis data of odor components is usable as a composition with the intended chicken flavor, and usually, satisfying flavor was not able to be obtained.

20 Generally used synthetic flavoring-based chicken flavors are designed flavors that highlight or elaborate the characteristic odor components of chicken by finding the specific odor components of chicken through odor analyses. Specifically, the intended chicken can be recognized by using a chicken flavor composition in combination with a color and a taste of a final product relating chicken, and the desired chicken flavor could not be reproduced by the flavor itself.

25

[0011]

As noted above, in order to prepare desired chicken flavor compositions, they were often composed of, mainly, materials such as (1) an animal or a vegetable extract, (2) a yeast extract, (3) an animal or a vegetable protein hydrolysate, (4) a heating cooking flavor, and (5) a synthetic flavoring. However, because chicken flavor compositions using such materials are of low potency, novel materials are needed and there was room for improvement for the preparation of a composition for obtaining a stronger and more desirable chicken flavor.

30

[0012]

35 Compounds such as 1,5-octadien-3-ol and 1,5-octadien-3-one are known as odor components, as noted above. However, use for imparting or enhancing a chicken flavor that accompanies a perception of boiled flavor, a perception of fried flavor, or a perception of roast flavor, particularly a chicken flavor that accompanies a perception of boiled flavor

was not known.

[0013]

It is accordingly a first object of the present invention to provide a chicken flavor composition and a chicken flavor enhancer that can provide a more desirable chicken
 5 flavor. A further object of the present invention is such as to provide a variety of food, drink or a seasoning that uses the chicken composition or the chicken enhancer, and to impart a chicken flavor to a food, a drink and a seasoning, or enhance it with the use of the composition and the enhancer.

As used in the present specification, "desirable chicken flavor" means that flavor
 10 characteristics (chicken flavor) are more desirable, that evoke such as a flavor of a soup that results from boiling chicken meat or chicken bone or the flavor that results from sauteing chicken (a perception of boiled flavor), a flavor of fried chicken that results from frying chicken meat or chicken bone with oil (a perception of fried flavor), and a flavor of
 15 roast meat that results from roasting chicken meat or chicken bone (a perception of roast flavor).

Solution to Problem

[0014]

The present inventors conducted intensive studies to find a solution to the
 20 foregoing problems, and found that a chicken flavor composition obtained by mixing 1,5-octadien-3-ol and/or 1,5-octadien-3-one surprisingly has a chicken flavor with strong flavor and body, and an excellent roast flavor, resulted in completion of the present invention.

[0015]

25 Specifically, the present invention relates to the following (1) to (12).

- (1) A chicken flavor composition containing at least one of 1,5-octadien-3-ol and 1,5-octadien-3-one.
- (2) The chicken flavor composition according to (1), containing a chicken flavor selected from a boiled flavor, a fried flavor, and a roast flavor.
- 30 (3) The chicken flavor composition according to (1) or (2), in which a total content of the 1,5-octadien-3-ol and the 1,5-octadien-3-one is 0.001 ppm to 1000 ppm by mass.
- (4) A food, a drink or a seasoning, containing the chicken flavor composition according to any one of (1) to (3).
- (5) A method for imparting a chicken flavor to a food, a drink or a seasoning, in
 35 which the chicken flavor composition according to any one of (1) to (3) is added.
- (6) A method for enhancing a chicken flavor of a food, a drink or a seasoning, the method including adding the chicken flavor composition according to any one of (1) to (3).

(7) A chicken flavor enhancer, containing at least one of 1,5-octadien-3-ol and 1,5-octadien-3-one.

(8) The chicken flavor enhancer according to (7), which enhances a chicken flavor selected from a boiled flavor, a fried flavor, and a roast flavor.

5 (9) A food, a drink or a seasoning, containing the chicken flavor enhancer according to (7) or (8).

(10) The food, the drink or the seasoning according to (9), in which a total content of the 1,5-octadien-3-ol and the 1,5-octadien-3-one is 0.0001 ppb to 10 ppm by mass.

10 (11) A method for imparting a chicken flavor to a food, a drink or a seasoning, the method including adding the chicken flavor enhancer according to (7) or (8).

(12) A method for enhancing a chicken flavor of a food, a drink or a seasoning, the method including adding the chicken flavor enhancer according to (7) or (8).

Advantageous Effects of Invention

15 [0016]

The present invention can provide a chicken flavor composition which is useful as a flavor enhancer or the like for food, drink, seasonings or the like, and which has an excellent chicken flavor that accompanies a perception of boiled flavor, a perception of fried flavor, or a perception of roast flavor, by mixing 1,5-octadien-3-ol and/or 1,5-octadien-3-one with a chicken flavor composition.

20

Description of Embodiments

[0017]

The present invention is described below in detail. However, the present invention is not limited to the following embodiment, and may be carried out with arbitrary modifications without deviating from the gist of the present invention.

25

As used in the present specification, "mass%" and "mass part", and "weight%" and "weight part" have the same meaning, respectively, and when simply mention "%", "ppm", and "ppb", they indicate "weight%", "weight ppm", and "weight ppb", respectively.

30 [0018]

The chicken flavor composition according to the present invention contains at least one of 1,5-octadien-3-ol and 1,5-octadien-3-one. At the beginning, the following describes 1,5-octadien-3-ol and 1,5-octadien-3-one.

For 1,5-octadien-3-ol, there exist (3S)-form and (3R)-form due to a single asymmetric carbon within the molecule, and there exist two geometric isomers, (5E)-form and (5Z)-form, depending on cis-trans resulting from the substituents on the double bond. Specifically, there are four kinds, (3S)-(5E)-1,5-octadien-3-ol, (3R)-(5E)-1,5-octadien-3-ol, (3S)-(5Z)-1,5-octadien-3-ol, and (3R)-(5Z)-1,5-octadien-3-ol.

35

For 1,5-octadien-3-one, there exist two geometric isomers, (5E)-form and (5Z)-form, depending on cis-trans resulting from the substituents on the double bond within the molecule. Specifically, there are two kinds, (5E)-1,5-octadien-3-one and (5Z)-1,5-octadien-3-one.

5 All of these 1,5-octadien-3-ol and 1,5-octadien-3-one are in oil (liquid) form at ordinary temperature.

[0019]

The present invention may use any isomer of 1,5-octadien-3-ol and 1,5-octadien-3-one, or may use a mixture of these.

10 These isomers can be distinguished on the basis of the retention time in gas chromatography. In this specification, no distinction is made for these isomers, and they are collectively referred to as "1,5-octadien-3-ol" or "1,5-octadien-3-one".

[0020]

15 As the 1,5-octadien-3-ol and/or 1,5-octadien-3-one, ones obtained by extraction from natural products may be used, or ones obtained by chemical synthesis (for example, ones synthesized according to the methods described in Non Patent Literature 8) may be used. The ones obtained from natural products and obtained by chemical synthesis may be used in a combination.

20 High-purity 1,5-octadien-3-ol and/or 1,5-octadien-3-one can be obtained in large quantities in the case by chemical synthesis.

[0021]

25 The content of the 1,5-octadien-3-ol and the 1,5-octadien-3-on in the chicken flavor composition of the present invention is, by their total content, preferably 0.001 ppm to 1000 ppm, more preferably 0.01 ppm to 100 ppm, and further preferably 0.1 ppm to 10 ppm. With a content of 0.001 ppm or more, the present invention can sufficiently exhibit effects, and with a content of 1000 ppm or less, an odorant note becomes more desirable as a chicken flavor, and therefore they are preferred.

[0022]

30 The chicken flavor composition of the present invention may contain other odor components.

Non-limiting examples of such other odor components include known synthetic flavoring compounds, including: hydrocarbons such as limonene, ocimene, α -pinene, β -pinene, γ -terpinene, sabinene, and myrcene; alcohols such as linalol, citronellol, dihydrolinalol, tetrahydromugol, myrcenol, dihydromyrcenol, tetrahydromyrcenol, ocimenol, terpeneol, 3-thujanol, benzyl alcohol, β -phenylethyl alcohol, α -phenylethyl alcohol, and cis-3-hexenol; aldehydes such as acetaldehyde, n-hexanal, n-heptanal, n-octanal, n-nonanal, cis-3-hexenal, cis-6-nonenal, 3,5,5-trimethylhexanal, decanal, undecanal, 2-methyldecanal, dodecanal, tridecanal, tetradeccanal, trans-2-hexenal, trans-4-

decenal, cis-4-decenal, trans-2-decenal, 10-undecenal, trans-2-undecenal, trans-2-dodecenal, 3-dodecenal, trans-2-tridecenal, 2,4-hexadienal, 2,4-decadienal, 2,4-dodecadienal, 5,9-dimethyl-4,8-decadienal, citral, citronellal, benzaldehyde, cuminaldehyde, vanillin, and ethyl vanillin; ketones such as 2-butanone, 3-heptanone, 3-octanone, 2-nonanone, 2-undecanone, 2-tridecanone, methyl heptenone, dimethyl octenone, geranyl acetone, 2,3,5-trimethyl-4-cyclohexenyl-1-methylketone, 3-methyl-2,4-nonanedione, Nerone (Givaudan, registered trade mark), nootkatone, dihydronootkatone, acetophenone, and 4,7-dihydro-2-isopentyl-2-methyl-1,3-dioxepin; esters such as propyl formate, octyl formate, linalyl formate, citronellyl formate, geranyl formate, neryl formate, terpinyl formate, ethyl acetate, isopropyl acetate, cis-3-hexenyl acetate, trans-2-hexenyl acetate, octyl acetate, nonyl acetate, decyl acetate, dodecyl acetate, dimethyl undecadienyl acetate, ocimenyl acetate, myrcenyl acetate, dihydromyrcenyl acetate, linalyl acetate, citronellyl acetate, geranyl acetate, neryl acetate, tetrahydromugol acetate, lavandulyl acetate, nerolidol acetate, dihydrocuminyl acetate, terpinyl acetate, citryl acetate, nopyl acetate, dihydroterpinyl acetate, 2,4-dimethyl-3-cyclohexenylmethyl acetate, myraldyl acetate, boeticol acetate, decenyl propionate, linalyl propionate, geranyl propionate, neryl propionate, terpinyl propionate, tricyclodecenyl propionate, styralyl propionate, octyl butyrate, neryl butyrate, cinnamyl butyrate, isopropyl isobutyrate, octyl isobutyrate, linalyl isobutyrate, neryl isobutyrate, linalyl isovalerate, terpinyl isovalerate, phenylethyl isovalerate, 2-methylpentyl 2-methyl valerate, methyl 3-hydroxyhexanoate, ethyl 3-hydroxyhexanoate, methyl octanoate, octyl octanoate, linalyl octanoate, methyl nonanoate, methyl undecylenate, linalyl benzoate, methyl cinnamate, isoprenyl angelate, methyl geranate, and triethyl citrate; phenols such as thymol, carvacrol, and β -naphthol isobutyl ether; lactones such as γ -undecalactone, and δ -dodecalactone; fatty acids such as acetic acid, propionic acid, butyric acid, valeric acid, hexanoic acid, octanoic acid, decanoic acid, dodecanoic acid, 2-decenoic acid, and geranic acid; sulfides such as diallyl sulfide, diallyl disulfide, dipropyl disulfide, methyl propyl disulfide, allyl propyl disulfide, and diallyl trisulfide; and nitrogen- or sulfur-containing compounds such as methyl anthranilate, ethyl anthranilate, methyl N-methyl anthranilate, methyl N-2'-methyl pentylidene anthranilate, ligandral, dodecanenitrile, 2-tridecenenitrile, geranyl nitrile, citronellyl nitrile, 3,7-dimethyl-2,6-nonadienonitrile, indole, 5-methyl-3-heptanoneoxime, thiogeraniol, limonenethiol, and P-menthyl-8-thiol, and

chicken meat and/or chicken bone extracts; flavors with chicken flavor obtained by solvent extraction, hydrodistillation, enzyme treatment, heat reactions, and the like; fishery product extracts; plant essential oils such as peppermint oil, spearmint oil, and spice essential oils of such as ginger, pepper, and onion; and oily extracts such as spice extracts, and oleoresins thereof.

These may be arbitrarily combined and mixed to prepare the chicken flavor composition.

[0023]

5 In the chicken flavor composition according to the present invention, the mass ratio of the total content of the 1,5-octadien-3-ol and the 1,5-octadien-3-one to other odor components is preferably 1:9 to 1:99.9 from the standpoint of improving the chicken flavor.

[0024]

10 The chicken flavor composition of the present invention may contain components other than the odor components, exemplified as suitable solvents for dissolving the odor components. Examples thereof include water, ethyl alcohol, glycerine, propylene glycol, triacetin, middle-chain fatty acid glycerine ester, food oils and fats such as food vegetable oils, natural pigments usable for food or drink, vitamins, vegetable resins, and the like.

15 From the standpoint of ease of handling of the composition, these are contained in preferably 0 to 99 mass% of the chicken flavor composition.

[0025]

20 A chicken flavor composition prepared into a liquid form by using the foregoing solvents may be prepared into a powder form by adding carriers such as lactose, dextrin, gum arabic, and cyclodextrin, or into an emulsion with the use of various emulsifiers usable for food or drink. Besides, a composition may be prepared and used in arbitral form such as a paste form, a granule form, and a microcapsule according to its intended use.

[0026]

25 The chicken flavor composition according to the present invention preferably has a chicken flavor selected from a boiled flavor, a fried flavor, or a roast flavor.

[0027]

30 The chicken flavor composition according to the present invention can impart a chicken flavor to a food, a drink or a seasoning by being added thereto. Specifically, a food or drink with an imparted chicken flavor can easily be produced even if the food, the drink or the seasoning does not have any chicken flavor.

A food, a drink or a seasoning with the enhanced chicken flavor can be obtained by adding the chicken flavor composition according to the present invention to a food, a drink or a seasoning that already has a chicken flavor.

[0028]

35 Such food, drink or seasoning (hereinafter, also referred to simply as "chicken flavored food or drink", or "chicken flavored seasoning", respectively) containing the chicken flavor composition according to the present invention are not particularly limited. Preferably, these are food, drink or seasonings relating chicken, such as a chicken flavored

food or drink or a chicken flavored seasoning as represented by soup, instant noodle, snack, and the like.

[0029]

5 A compounding amount of the chicken flavor composition of the present invention to the chicken flavored food or drink or the chicken flavored seasoning varies depending on such as the intended use, and the type and the odorant note strength of the chicken flavor enhancer. For example, a concentration range may be 0.001% to 10%, preferably 0.01 to 3%.

[0030]

10 The present invention also can provide a chicken flavor enhancer containing 1,5-octadien-3-ol and/or 1,5-octadien-3-one. The flavor enhanced by the enhancer is preferably a chicken flavor selected from a boiled flavor, a fried flavor, and a roast flavor.

As the 1,5-octadien-3-ol and/or the 1,5-octadien-3-one here, any of the foregoing geometric isomers may be used, and two or more of them may be used in combination.

15 [0031]

The total of a compounding amount of the 1,5-octadien-3-ol and the 1,5-octadien-3-one in the chicken flavor enhancer of the present invention to the chicken flavored food or drink or the chicken flavored seasoning varies depending on such as the intended use and the type of the chicken flavor enhancer. For example, a concentration range may be 20 0.0001 ppb to 10 ppm, preferably 0.001 ppb to 1 ppm, more preferably 0.01 ppb to 0.1 ppm.

[0032]

As the content of the 1,5-octadien-3-ol and the 1,5-octadien-3-one in the chicken flavor enhancer, in their total content, 0.001 ppm to 1000 ppm is preferred in that the 25 present invention can sufficiently exhibit effects, and an odorant note becomes more desirable as a chicken flavor.

[0033]

30 The chicken flavor enhancer of the present invention may be added to the chicken flavored food or drink or the chicken flavored seasoning to enhance the chicken flavors of them, and may be added to various food, drink or seasonings having no chicken flavor to easily produce a food, a drink or a seasoning that has an imparted chicken flavor.

Examples

[0034]

35 The following describes the present invention in greater detail by using Examples. However, the present invention is in no way limited by the following, and may be altered or modified in many ways without deviating from the scope of the invention. Concerning

to the unit in the formulations described below, composition ratios indicate, unless specifically mentioned, mass ratios.

[0035]

(Example 1) Chicken Flavor Enhancer

- 5 In Example 1, a chicken flavor enhancer was prepared by preparing a 0.1% aqueous solution of 1,5-octadien-3-ol. The 1,5-octadien-3-ol here is (5Z)-1,5-octadien-3-ol, obtained by the synthesis method described in Non Patent Literature 8.

(Example 2) Chicken Flavor Enhancer

- 10 In Example 2, a chicken flavor enhancer was prepared by preparing a 0.1% aqueous solution of 1,5-octadien-3-one. The 1,5-octadien-3-one here is (5Z)-1,5-octadien-3-one, obtained by the synthesis method described in Non Patent Literature 8.

[0036]

(Examples 3 and 4) Chicken Flavor Composition

- 15 Chicken flavor compositions were obtained by mixing each component according to the formulations presented in the Table 1 below. The obtained compositions both had a boiled flavor.

[0037]

[Table 1]

(mass part)		
Component	Example 3	Example 4
Hexanal	7	7
trans-2-Decenal	2	2
2,4-Decadienal	46	46
Dimethyl sulfide 1% ethanol solution	10	10
Linalool	2	2
Terpineol	3	3
1-Octen-3-ol	3	3
Maltol 10% ethanol solution	10	10
Diacetyl 1% ethanol solution	5	5
Acetoin 1% ethanol solution	5	5
Chicken flavor enhancer of Example 1	0.5	
Chicken flavor enhancer of Example 2		0.5
Ethanol	Remainder	Remainder
Total	1000	1000

20

[0038]

(Comparative Example 1) Chicken Flavor Composition

A chicken flavor composition was prepared in the same manner as in Example 3, except that the chicken flavor enhancer was not contained.

[0039]

(Application Example 1) Chicken Flavor Oil

5 To 200 g of chicken carcass was added 1 kg of water and the mixture was heated at 120°C for 4 hours. After cooling, an insoluble solid was removed to obtain 850 g of chicken extract. The extract was concentrated under reduced pressure to mass of 1/3, and the resulting concentrate was, after adding 150 g of canola oil and the chicken flavor composition of Example 3 by 2 mass% with respect to the canola oil, stirred under heat at
10 100°C for 3 hours. The mixture was cooled, and centrifuged to collect the chicken flavor oil layer portion to obtain a present product 1 (chicken flavor oil).

[0040]

(Application Example 2) Chicken Flavor Oil

A present product 2 (chicken flavor oil) was obtained in the same manner as in
15 Application Example 1, except that the chicken flavor composition of Example 4 was used instead of the chicken flavor composition of Example 3.

[0041]

(Comparative Application Example 1) Chicken Flavor Oil

A comparative product 1 (chicken flavor oil) was obtained in the same manner as
20 in Application Example 1, except that the chicken flavor composition of Comparative Example 1 was used instead of the chicken flavor composition of Example 3.

[0042]

The present product 1, the present product 2, and the comparative product 1 were each diluted in 20 times with canola oil, and evaluated for flavor in a sensory evaluation
25 participated by 10 trained panelists. All panelists rated the present product 1 and the present product 2 as more imparting an impression of an enhanced top impact and a distinct edge, and more having an enduring chicken flavor than the comparative product 1.

[0043]

(Test Examples 1-1 to 1-8) Tests on the Effect of 1,5-Octadien-3-ol in Chicken Flavor Oil

30 Chicken flavor enhancers of Test Examples 1-1 to 1-8 were each prepared by preparing 1,5-octadien-3-ol aqueous solutions in the same manner as in Example 1, with the 1,5-octadien-3-ol varied to have the concentrations shown in Table 2. Chicken flavor compositions of Test Examples 1-1 to 1-8 were prepared by each using the chicken flavor enhancers in Example 3 prepared instead of the chicken flavor enhancer of Example 1.

35 [0044]

[Table 2]

Test Examples	1-1	1-2	1-3	1-4	1-5	1-6	1-7	1-8
Concentration (ppm)	0	0.005	0.05	0.5	5	50	500	5000

[0045]

Chicken flavor oils of Test Examples 1-1 to 1-8 were obtained in the same manner as in Application Example 1, except that the chicken flavor compositions of Test Examples 1-1 to 1-8 were used instead of the chicken flavor composition of Example 3.

[0046]

The chicken flavor oils of Test Examples 1-1 to 1-8 were each diluted in 20 times with canola oil, and evaluated for flavor in a sensory evaluation participated by 10 trained panelists. All panelists rated the chicken flavor oils of Test Examples 1-2 to 1-7 as notably having a roundness and a richness with a sense of harmony, and having enduring chicken flavor. The chicken flavor oils of Test Examples 1-4 to 1-6 were particularly found to have a desirable chicken flavor. On the other hand, all panelists rated that the chicken flavor oil of Test Example 1-1 did not sufficiently develop a perception of chicken flavor and that the chicken flavor oil of Test Example 1-8 was undesirable because of the odorant note that was different from a chicken flavor.

[0047]

(Test Examples 2-1 to 2-8) Tests on the Effect of 1,5-Octadien-3-One in Chicken Flavor Oil

Chicken flavor enhancers of Test Examples 2-1 to 2-8 were each prepared by preparing 1,5-octadien-3-one aqueous solutions in the same manner as in Example 2, with the 1,5-octadien-3-one varied to have the concentrations shown in Table 3. Chicken flavor compositions of Test Examples 2-1 to 2-8 were prepared by each using the chicken flavor enhancers in Example 4 prepared instead of the chicken flavor enhancer of Example 2.

[0048]

[Table 3]

Test Examples	2-1	2-2	2-3	2-4	2-5	2-6	2-7	2-8
Concentration (ppm)	0	0.005	0.05	0.5	5	50	500	5000

[0049]

Chicken flavor oils of Test Examples 2-1 to 2-8 were obtained in the same manner as in Application Example 1, except that the chicken flavor compositions of Test Examples 2-1 to 2-8 were used instead of the chicken flavor of Example 4.

[0050]

The chicken flavor oils of Test Examples 2-1 to 2-8 were each diluted in 20 times with canola oil, and evaluated for flavor in a sensory evaluation participated by 10 trained panelists. All panelists rated the chicken flavor oils of Test Examples 2-2 to 2-7 as notably having a roundness and a richness with a sense of harmony, and having enduring

chicken flavor. The chicken flavor oils of Test Examples 2-4 to 2-6 were particularly found to have a desirable chicken flavor. On the other hand, all panelists rated that the chicken flavor oil of Test Example 2-1 did not sufficiently develop a perception of chicken flavor and that the chicken flavor oil of Test Example 2-8 was undesirable because of the odorant note that was different from a chicken flavor.

[0051]

While the present invention has been described in detail with reference to specific embodiments thereof, it will be apparent to one skilled in the art that various changes and modifications can be made therein without departing from the spirit and scope thereof.

This application is based on Japanese Patent Application (No. 2012-220788) filed on October 2, 2012, the contents of which are incorporated herein by reference.

CLAIMS

[Claim 1]

- 5 A chicken flavor composition comprising at least one of 1,5-octadien-3-ol and 1,5-octadien-3-one, wherein a total content of the 1,5-octadien-3-ol and the 1,5-octadien-3-one is 0.25 ppm to 1000 ppm by mass.

[Claim 2]

The chicken flavor composition according to claim 1, comprising a chicken flavor selected from a boiled flavor, a fried flavor, and a roast flavor.