

New Processes For Preparing
Pesticidal Intermediates

5 This invention relates to a process for preparing certain
cyanomethylpropane derivatives and the use of these compounds in
the synthesis of pesticides and pesticide intermediates.

Ethyl 2,3-dicyanopropionate was first prepared and
characterised by Higson and Thorpe (J.Chem.Soc. 82, 1460 (1906))
who obtained the material in good yield (70-81%) by reaction of
10 formaldehyde cyanohydrin with the sodium salt of ethyl cyanoacetate.
Dickinson (J. Am. Chem. Soc. 82, 6132 (1960)) repeated this work.
This method of preparing the dicyanopropionate suffers from a
significant drawback in that it is first necessary to isolate the
intermediate formaldehyde cyanohydrin. This highly water soluble
15 cyanohydrin is obtained by lengthy continuous extraction and has a
limited stability, often decomposing violently upon attempted
distillation. Furthermore, this reaction requires care given the risk of
formation of dimeric side-products. The preparation of
dicyanopropionates has also been described by Whiteley and
20 Marianelli (Synthesis (1978), 392) with the process leading to 2,3-
disubstituted succinodinitriles from the cyanoacetate, an aldehyde (a 1
to 3 carbon alkylaldehyde or benzaldehyde) and potassium cyanide
via 3-substituted-2,3- dicyanopropionates (which were not isolated).
However, the yield decreases dramatically from isobutyraldehyde to
25 acetaldehyde. In the same manner Smith and Horwitz (J. Am. Chem.
Soc. 1949, 71, 3418) described the same reaction with a ketone with a
yield of 70%. This prior art therefore teaches that yields improve
with increasing size of group adjacent to the carbonyl group.

30 In one aspect the present invention seeks to provide a process
for preparing cyanomethyl propane derivatives satisfying one or more
of the following criteria:

- avoiding the use of formaldehyde cyanohydrin;
- avoiding the dimerisation side reaction;

This Sheet Has Been Replaced




THIS SHEET HAS BEEN
REPLACED

CONFIRMATION COPY

- obtaining the required product directly in high yield and with high purity.

Therefore the present invention provides a process for preparing a compound of formula (I):



5

wherein

R represents straight- or branched- chain alkyl having up to 18 carbon atoms; or a salt thereof;

which comprises the reaction of a cyanoacetate of formula (II):-

10



wherein R is as defined above, with a cyanide salt and formaldehyde or a source thereof.

Preferably R represents straight- or branched-chain alkyl having from 1 to 6 carbon atoms and most preferably R represents ethyl.

15

Suitable salts of cyanide include metal salts and organic salts (e.g. tetra-alkylammonium cyanides such as tetrabutylammonium cyanide). Preferably the cyanoacetate of formula (II) reacts with an alkali or alkaline earth metal cyanide salt, with alkali metal cyanide salts being especially suitable for use in the present invention, particularly potassium cyanide or sodium cyanide. The product may conveniently be isolated as the alkaline earth metal or alkali metal salt. Alternatively, the reaction mixture is acidified, for example with a mineral acid such as sulphuric acid or hydrochloric acid, to give the compound of formula (I). Where a compound of formula (I) above is desired (rather than a salt thereof) high yields are generally obtained when the reaction mixture is acidified without the addition of water. Whilst formaldehyde itself can be used in the reaction it is more convenient to use the polymerised form known as paraformaldehyde $[(\text{HCOH})_n]$, available for example from Aldrich Chemical Company.

25

30

The reaction is generally performed using about 1 molar equivalent of a compound of formula (II); about 0.95 to 1.0 molar


equivalents of cyanide salt; and about 1 molar equivalent of formaldehyde compound (based on the formaldehyde content).

5 The reaction may be carried out in the presence of a solvent. Preferably, the reaction is performed in a solvent medium which is usually an alcoholic medium or dimethyl formamide (DMF), N-methyl pyrrolidone (NMP), dioxan, tetrahydrofuran (THF) or dimethoxyethane. Especially preferred solvents are C₁-C₆ alcohols such as methanol or, most preferably, anhydrous ethanol. Although the temperature of the reaction is not critical the reaction will
10 normally be performed from about 0 to about 120°C or at the reflux temperature of the solvent. Generally best results are obtained by introducing the formaldehyde source after the other reactants have been combined.

The reaction generally takes place under substantially
15 anhydrous conditions (it being understood that the reaction proceeds with the formation of one equivalent of water), as in the event of prolonged exposure to aqueous conditions, there is a risk that the ester group of the compound of formula (I) will undergo hydrolysis (due to the basic conditions arising during the reaction) to the
20 corresponding acid of formula (I) (in which R is replaced by hydrogen) and subsequently undergo decarboxylation to give 1,2-dicyanoethane.

The compound of formula (I) is useful in the preparation of pesticidally active compounds, for example as described in European
25 Patent Publication Nos. 0295117 and 0234119, and WO93/06089.

In particular, the process of the invention may form part of an *in situ* preparation of another pesticidal intermediate and in a further aspect the invention provides a process for the preparation of a compound of formula (III):




(III)

wherein R_1 is cyano; W is nitrogen or $-CR_4$; R_2 and R_4 independently represent halogen; and R_3 represents halogen, haloalkyl (preferably trifluoromethyl), haloalkoxy (preferably trifluoromethoxy) or $-SF_5$; which process comprises:


(a) reacting a cyanoacetate of formula (II) as defined above, with a cyanide salt and formaldehyde or a source thereof, to give a compound of formula (I) as defined above; and

(b) reacting the compound of formula (I) thus obtained with the diazonium salt of a compound of formula (IV):



(IV)

wherein W , R_2 and R_3 are as defined above, to give a compound of formula (V):



wherein W, R, R₁, R₂ and R₃ are as defined above, followed by the cyclisation of said compound of formula (V).

5 Compounds of formula (V) above possess a chiral centre giving rise to different enantiomers, and also may exist as different geometric isomers or mixtures thereof. All such forms are embraced by the present invention. In this process, the product of reaction step (a) is generally acidified with an alcoholic solution of a mineral acid, preferably an ethanolic solution of hydrogen chloride. This also
10 ensures that any acid by-product of the reaction step (a) (leading to the corresponding compound of formula (I) in which R is replaced by hydrogen) is re-esterified. For these reasons it is also preferred that in this process, reaction step (a) takes place under substantially
15 anhydrous conditions.

 Reaction step (b) is generally performed in the presence of an inert solvent, for example water, acetonitrile, dichloromethane or DMF, or more preferably an alcoholic solvent (e.g. methanol or ethanol) and is optionally buffered (e.g. with sodium acetate). The
20 diazonium salt of a compound of formula (IV) may be prepared using diazotising agents known in the literature and is conveniently prepared with a molar equivalent of sodium nitrite and a mineral acid (e.g. hydrochloric or sulphuric acid), at a temperature of from about -10°C to about 50°C, more preferably from about 0°C to about 5°C.
25 The diazonium salt of the compound of formula (IV) is generally prepared *in situ* as solvents such as alcohols tend to reduce diazonium salts quickly. In the present reaction, the reaction of the diazonium

salt of the compound of formula (IV) to give a compound of formula (V) above generally occurs faster than the reduction of the diazonium salt.

5 Subsequent hydrolysis, preferably using mild conditions with a base such as aqueous sodium hydroxide, sodium carbonate or ammonia, may be necessary to effect the cyclisation of the compound of formula (V) to a compound of formula (III).

10 The molar ratio of the compounds of formula (II):(IV) is generally from about 1.5:1 to about 1:4, preferably from about 1.3:1 to about 1:1, more preferably about 1.1:1.

Compounds of formula (III) and (IV) above are described in the literature, for example see EP-A1-0295117. Compounds of formula (V) above are novel and thus constitute a further feature of the present invention.

15 The following non-limiting examples illustrate the invention.

Example 1

Preparation of ethyl 2,3-dicyanopropionate.

20 Potassium cyanide (13.0g, 0.2M) was stirred in absolute ethanol and ethyl cyanoacetate (22.6g, 0.2M) and paraformaldehyde (6.0g, 0.2m) were added at ambient temperature. After 5 minutes the white suspension was heated under reflux conditions for 12 minutes, and the orange solution was evaporated to dryness *in vacuo* at below 25°C to give a buff solid. The solid (the potassium salt) was
25 dissolved in water (400ml), acidified to pH 5 with 2M hydrochloric acid solution, giving a red oil. This mixture was extracted with dichloromethane and the extracts dried and evaporated to dryness *in vacuo* to give the title compound as a red oil (23.5g), ¹H NMR (CDCl₃) δ 4.3 (2H, q), 3.95 (1H, t), 3.0 (2H, d), 1.35 (3H, t);
30 identical with an authentic sample. Distilled material had b.p. 132-136°C at 0.5mmHg.

Yield : 77%.

Comparative example according to J. Chem. Soc. 89, 1460 (1906)

A solution of sodium ethoxide [prepared from sodium (25.2g, 1.15M) and absolute ethanol (650ml)] was stirred under an inert atmosphere and treated with ethyl cyanoacetate (127.7ml, 1.2M) over 20 minutes. The solution was cooled to below 10°C and then added slowly to a solution of formaldehyde cyanohydrin (freshly prepared, 70g, 0.2M) in absolute ethanol (200ml) at 5°C over 55 minutes. After standing overnight the mixture was poured into ice-water (1L) and acidified to pH 1-2 with concentrated hydrochloric acid. This was extracted into dichloromethane, dried over anhydrous magnesium sulphate and evaporated to give a dark orange oil (150.6g). This was distilled in vacuo collecting the title compound (73.6g) as a colourless oil, b.p. 144-148°C/1mbar. Yield : 40%

The superiority of the process of the invention is thus clearly demonstrated over this prior art.

Example 2

Process for the preparation of 5-amino-3-cyano-1-(2,6-dichloro-4-trifluoromethyl-phenyl)pyrazole.

Sodium cyanide (20g, 0.408M) and ethyl cyanoacetate (46g, 0.408M) were dissolved in absolute ethanol (300ml) under an inert atmosphere. Paraformaldehyde (12.2g, 0.408M) was added, producing an exotherm, and the temperature was maintained below 50°C. The reaction mixture was then stirred at ambient temperature for between 5 and 7 hours, cooled to between 0 and 5°C, and an ethanolic solution containing hydrogen chloride (0.45M) was added, maintaining the temperature below 5°C. The reaction mixture was left overnight and 111ml of a solution of hydrochloric acid (0.73M) in ethanol was added to the suspension thus obtained at about 5°C. 2,6-Dichloro-4-trifluoromethylaniline (84.44g, 0.367M) was added at this temperature followed by sodium nitrite (35.84g, 0.514M) resulting in the formation of ethyl 2,3-dicyano-2-[2,6-dichloro-4-trifluoromethylphenyl]azo]propionate, which may be isolated by column chromatography, eluting with a pentane/ether solution and/or reverse phase chromatography with an acetonitrile-water solution; or


by removing the ethanol by distillation, dissolving the reaction mixture in toluene, washing the toluene solution with water and evaporating the toluene to dryness. ¹H NMR (CDCl₃) 1.37(t,3H), 3.55(s,2H), 4.43(q,2H), 7.65(s,2H).

5 Ammonia gas (9.6g, 0.56M) was then bubbled into the reaction mixture at 0°C. The ethanol was evaporated from the reaction mixture under reduced pressure and the concentrated liquors were taken up into a mixture of toluene and ethyl acetate. This solution was washed with water and after concentration of the toluene phase at
10 80°C, the solution was cooled to give the title compound as a crystalline solid, and the liquors were then concentrated and cooled to give a second crop of recrystallised product, m.p. 141 - 142°C (combined weight of title compound 87.54g; yield based on aniline starting material = 78%).

15

CLAIMS

1. A process for preparing a compound of formula (I):



5 wherein

R represents straight- or branched- chain alkyl having from 1 to 18 carbon atoms,

or a salt thereof, which comprises the reaction of a cyanoacetate of formula (II):-



10 wherein R is as defined above,

with a cyanide salt, and formaldehyde or a source thereof.

- 15 2. A process according to claim 1 in which the cyanide salt is an alkali or alkaline earth metal salt.

3. A process according to claim 2 wherein the cyanide salt is sodium or potassium cyanide.

20 4. A process according to claim 1, 2 or 3, wherein R represents straight- or branched-chain alkyl having from 1 to 6 carbon atoms.

25 5. A process according to any one of claims 1 to 4, wherein the compound of formula (II) is ethyl cyanoacetate.

6. A process according to any one of claims 1 to 5, wherein the source of formaldehyde is paraformaldehyde.

30 7. A process according to any one of claims 1 to 6, wherein the reaction is performed in presence of a solvent which is an alcoholic medium.



8. A process according to any one of the preceding claims, wherein the reaction is carried out at a temperature from about 0°C to about 120°C.

5

9. A process according to any one of the preceding claims in which the reaction is performed using about 1 molar equivalent of the compound of formula (II); about 0.95 to 1.0 molar equivalents of cyanide salt; and about 1 molar equivalent of formaldehyde compound.

10


10. A process according to any one of the preceding claims in which the reaction mixture is acidified after reaction of the cyanoacetate of formula (II) with the cyanide salt and formaldehyde.

15

11. A process according to any one of the preceding claims, in which the reaction is performed under substantially anhydrous conditions.

20

12. A process for the preparation of a compound of formula (III):



(III)

wherein R₁ is cyano; W is nitrogen or -CR₄; R₂ and R₄ independently represent halogen; and R₃ represents halogen, haloalkyl, haloalkoxy or SF₅; which process comprises:


25

(a) reacting a cyanoacetate of formula (II)




wherein R represents straight- or branched- chain alkyl having from 1 to 18 carbon atoms, with a cyanide salt and formaldehyde or a source thereof, to give a compound of formula (I) as defined in claim 1; and

(b) reacting the compound of formula (I) thus obtained with the diazonium salt of a compound of formula (IV):



(IV)

wherein W, R₂ and R₃ are as defined above; to give a compound of formula (V):



(V)

wherein W, R, R₁, R₂ and R₃ are as defined above, followed by the cyclisation of said compound of formula (V).

13. A process according to claim 12 in which the product of reaction step (a) is treated with an alcoholic solution of an acid, preferably a mineral acid.




14. A process according to claim 12 or 13 in which the molar ratio of the compounds of formula (II):(IV) is from about 1.5:1 to about 1:4, preferably from about 1.3:1 to about 1:1.

5 15. A process according to claim 12, 13 or 14 in which reaction step (a) is performed under substantially anhydrous conditions.

10 16. A process according to any one of claims 12 to 15 in which the cyclisation is effected by hydrolysis of the compound of formula (V).

17. A compound of formula (V):



15 wherein R_1 is cyano; W is nitrogen or $-CR_4$; R_2 and R_4 independently represent halogen; and R_3 represents halogen, haloalkyl, haloalkoxy or SF_5 .

20 18. Ethyl 2,3-dicyano-2-[(2,6-dichloro-4-trifluoromethylphenyl)azo]propionate.