

BROADSPECTRUM SUBSTITUTED BENZIMIDAZOLE SULFONAMIDE HIV
PROTEASE INHIBITORS

5 The present invention relates to substituted-benzimidazole sulfonamides, their use as aspartic protease inhibitors, in particular as broadspectrum HIV protease inhibitors, processes for their preparation as well as pharmaceutical compositions and diagnostic kits comprising them. The present invention also concerns combinations of the present substituted benzimidazole sulfonamides with another anti-retroviral agent. It further relates to their use in assays as reference compounds or as reagents.

10 The virus causing the acquired immunodeficiency syndrome (AIDS) is known by different names, including T-lymphocyte virus III (HTLV-III) or lymphadenopathy-associated virus (LAV) or AIDS-related virus (ARV) or human immunodeficiency virus (HIV). Up until now, two distinct families have been identified, i.e. HIV-1 and
15 HIV-2. Hereinafter, HIV will be used to generically denote these viruses.

One of the critical pathways in a retroviral life cycle is the processing of polyprotein precursors by aspartic protease. For instance the HIV viral *gag-pol* protein is processed by HIV protease. The correct processing of the precursor polyproteins by the aspartic
20 protease is required for the assembly of infectious virions, thus making the aspartic protease an attractive target for antiviral therapy. In particular for HIV treatment, the HIV protease is an attractive target.

HIV protease inhibitors (PIs) are commonly administered to AIDS patients in
25 combination with other anti-HIV compounds such as, for instance nucleoside reverse transcriptase inhibitors (NRTIs), non-nucleoside reverse transcriptase inhibitors (NNRTIs), fusion inhibitors such as T-20 or other protease inhibitors. Despite the fact that these antiretrovirals are very useful, they have a common limitation, namely, the targeted enzymes in the HIV virus are able to mutate in such a way that the known
30 drugs become less effective, or even ineffective against these mutant HIV viruses. Or, in other words, the HIV virus creates an ever increasing resistance against the available drugs.

Resistance of retroviruses, and in particular the HIV virus, against inhibitors is a major
35 cause of therapy failure. For instance, half of the patients receiving anti-HIV combination therapy do not respond fully to the treatment, mainly because of resistance of the virus to one or more drugs used. Moreover, it has been shown that resistant virus is carried over to newly infected individuals, resulting in severely limited therapy options for these drug-naive patients. Therefore, there is a need in the art for new

compounds for retrovirus therapy, more particularly for AIDS therapy. The need in the art is particularly acute for compounds that are active not only on wild type HIV virus, but also on the increasingly more common resistant HIV viruses.

- 5 Known antiretrovirals, often administered in a combination therapy regimen, will eventually cause resistance as stated above. This often may force the physician to boost the plasma levels of the active drugs in order for said antiretrovirals to regain effectivity against the mutated HIV viruses. The consequence of which is a highly undesirable increase in pill burden. Boosting plasma levels may also lead to an
- 10 increased risk of non-compliance with the prescribed therapy. Thus, it is not only important to have compounds showing activity for a wide range of HIV mutants, it is also important that there is little or no variance in the ratio between activity against mutant HIV virus and activity against wild type HIV virus (also defined as fold resistance or FR) over a broad range of mutant HIV strains. As such, a patient may
- 15 remain on the same combination therapy regimen for a longer period of time since the chance that a mutant HIV virus will be sensitive to the active ingredients will be increased.

- Finding compounds with a high potency on the wild type and on a wide variety of
- 20 mutants is also of importance since the pill burden can be reduced if therapeutic levels are kept to a minimum. One way of reducing this pill burden is finding anti-HIV compounds with good bioavailability, i.e. a favourable pharmacokinetic and metabolic profile, such that the daily dose can be minimized and consequently also the number of pills to be taken.

- 25 Another important characteristic of a good anti-HIV compound is that plasma protein binding of the inhibitor has minimal or even no effect on its potency.


- Thus, there is a high medical need for protease inhibitors that are able to combat a
- 30 broad spectrum of mutants of the HIV virus with little variance in fold resistance, have a good bioavailability and experience little or no effect on their potency due to plasma protein binding.

- Up until now, several protease inhibitors are on the market or are being developed.
- 35 One particular core structure (depicted below) has been disclosed in a number of references, such as, WO 95/06030, WO 96/22287, WO 96/28418, WO 96/28463, WO 96/28464, WO 96/28465, WO 99/65870, WO 99/67254, WO 00/76961 and WO 97/18205. The compounds disclosed therein are described as retroviral protease inhibitors.


WO 03/076413

PCT/EP03/50057

-3-




WO 99/67254 discloses 4-substituted-phenyl sulfonamides capable of inhibiting multi-drug resistant retroviral proteases.



- 5 Surprisingly, the substituted-benzimidazole sulfonamides of the present invention are found to have a favourable pharmacological and pharmacokinetic profile. Not only are they active against wild-type HIV virus, but they also show a broad spectrum activity against various mutant HIV viruses exhibiting resistance against known protease inhibitors.


10

The present invention concerns 2-(substituted-amino)-benzimidazole protease inhibitors, having the formula



- and *N*-oxides, salts, stereoisomeric forms, racemic mixtures, prodrugs, esters and
15 metabolites thereof, wherein

R_1 is hydrogen, C_{1-6} alkyl, C_{2-6} alkenyl, aryl C_{1-6} alkyl, aryl C_{2-6} alkenyl, C_{3-7} cycloalkyl, C_{3-7} cycloalkyl C_{1-6} alkyl, aryl, Het¹, Het¹ C_{1-6} alkyl, Het², Het² C_{1-6} alkyl, R_1 may also be a radical of formula



- 20 wherein

- R_9 , R_{10a} and R_{10b} are, each independently, hydrogen, C_{1-4} alkyloxycarbonyl, carboxyl, aminocarbonyl, mono- or di(C_{1-4} alkyl)aminocarbonyl, C_{3-7} cycloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl or C_{1-4} alkyl optionally substituted with aryl, Het^1 , Het^2 , C_{3-7} cycloalkyl, C_{1-4} alkyloxycarbonyl, carboxyl, aminocarbonyl, mono- or di(C_{1-4} alkyl)aminocarbonyl, aminosulfonyl, C_{1-4} alkylS(O)_t, hydroxy, cyano, halogen or optionally mono- or disubstituted amino where the substituents are selected from C_{1-4} alkyl, aryl, aryl C_{1-4} alkyl, C_{3-7} cycloalkyl, C_{3-7} cycloalkyl C_{1-4} alkyl, Het^1 , Het^2 , Het^1C_{1-4} alkyl and Het^2C_{1-4} alkyl; whereby R_9 , R_{10a} and the carbon atoms to which they are attached optionally form a C_{3-7} cycloalkyl radical;
- R_{11a} is hydrogen, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-7} cycloalkyl, aryl, optionally mono- or disubstituted aminocarbonyl, optionally mono- or disubstituted amino C_{1-4} alkylcarbonyloxy, C_{1-4} alkyloxycarbonyl, aryloxycarbonyl, Het^1 oxycarbonyl, Het^2 oxycarbonyl, aryloxycarbonyl C_{1-4} alkyl, aryl C_{1-4} alkyloxycarbonyl, C_{1-4} alkylcarbonyl, C_{3-7} cycloalkylcarbonyl, C_{3-7} cycloalkyl C_{1-4} alkyloxycarbonyl, C_{3-7} cycloalkylcarbonyloxy, carboxyl C_{1-4} alkylcarbonyloxy, C_{1-4} alkylcarbonyloxy, aryl C_{1-4} alkylcarbonyloxy, arylcarbonyloxy, aryloxycarbonyloxy, Het^1 carbonyl, Het^1 carbonyloxy, Het^1C_{1-4} alkyloxycarbonyl, Het^2 carbonyloxy, Het^2C_{1-4} alkylcarbonyloxy, Het^2C_{1-4} alkyloxycarbonyloxy or C_{1-4} alkyl optionally substituted with aryl, aryloxy, Het^2 or hydroxy; wherein the optional substituents on the amino groups are each independently selected from C_{1-4} alkyl, aryl, aryl C_{1-4} alkyl, C_{3-7} cycloalkyl, C_{3-7} cycloalkyl C_{1-4} alkyl, Het^1 , Het^2 , Het^1C_{1-4} alkyl and Het^2C_{1-4} alkyl; whereby R_{11a} may be optionally linked to the remainder of the molecule via a sulfonyl group;
- R_{11b} is hydrogen, C_{3-7} cycloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, aryl, Het^1 , Het^2 or C_{1-4} alkyl optionally substituted with halogen, hydroxy, C_{1-4} alkylS(=O)_t, aryl, C_{3-7} cycloalkyl, Het^1 , Het^2 , optionally mono- or disubstituted amino where the substituents are selected from C_{1-4} alkyl, aryl, aryl C_{1-4} alkyl, C_{3-7} cycloalkyl, C_{3-7} cycloalkyl C_{1-4} alkyl, Het^1 , Het^2 , Het^1C_{1-4} alkyl and Het^2C_{1-4} alkyl;
- Each t independently is zero, 1 or 2;
- R_2 is hydrogen or C_{1-6} alkyl;
- L is -C(=O)-, -O-C(=O)-, -NR₈-C(=O)-, -O-C₁₋₆alkanediyl-C(=O)-, -NR₈-C₁₋₆alkanediyl-C(=O)-, -S(=O)₂-, -O-S(=O)₂-, -NR₈-S(=O)₂-, -NR₈-C₁₋₆alkanediyl-S(=O)₂, whereby either the C(=O) group or the S(=O)₂ group is attached to the NR₂ moiety; whereby the C₁₋₆alkanediyl moiety is optionally substituted with hydroxy, aryl, Het^1 or Het^2 ;

- R₃ is C₁₋₆alkyl, aryl, C₃₋₇cycloalkyl, C₃₋₇cycloalkylC₁₋₄alkyl, or arylC₁₋₄alkyl optionally substituted with C₁₋₄alkyl, halogen, nitro, cyano, polyhaloC₁₋₄alkyl, hydroxy, C₁₋₄alkyloxy, mercapto or C₁₋₄alkylthio;
- R₄ is hydrogen, C₁₋₄alkyloxycarbonyl, carboxyl, aminocarbonyl, mono- or di(C₁₋₄alkyl)aminocarbonyl, C₃₋₇cycloalkyl, C₂₋₆alkenyl, C₂₋₆alkynyl or C₁₋₄alkyl optionally substituted with aryl, Het¹, Het², C₃₋₇cycloalkyl, C₁₋₄alkyloxycarbonyl, carboxyl, aminocarbonyl, mono- or di(C₁₋₄alkyl)aminocarbonyl, aminosulfonyl, C₁₋₄alkylS(=O)₆, hydroxy, cyano, halogen or optionally mono- or disubstituted amino where the substituents are selected from C₁₋₄alkyl, aryl, arylC₁₋₄alkyl, C₃₋₇cycloalkyl, C₃₋₇cycloalkylC₁₋₄alkyl, Het¹, Het², Het¹C₁₋₄alkyl and Het²C₁₋₄alkyl; or
- R₄ may also be -X-R₇ wherein
 X is -O- or -N(R₇)-Y-;
 Y is -C(=O)-, -S(=O)₂-, -O-C(=O)-, -NR₈-C(=O)-, -C(=O)-C(=O)-, -O-S(=O)₂- or -NR₈-S(=O)₂-;
- R₅ is hydrogen, hydroxy, C₁₋₆alkyl, Het¹C₁₋₆alkyl, Het²C₁₋₆alkyl, aminoC₁₋₆alkyl whereby the amino group may optionally be mono- or di-substituted with C₁₋₄alkyl;
- R₆ is hydrogen, C₁₋₆alkyloxy, Het¹, Het¹oxy, Het², Het²oxy, aryl, aryloxy, aryloxy-C₁₋₄alkyl, C₁₋₄alkyloxyaryl, C₁₋₄alkyloxyHet¹, C₁₋₄alkyloxyHet², C₁₋₄alkyloxy-carbonylamino, aminoC₁₋₄alkylamino, amino or aminoC₁₋₄alkyloxy and in case A is other than C₁₋₆alkanediyl, then R₆ may also be C₁₋₆alkyl, Het¹C₁₋₄alkyl, Het¹oxyC₁₋₄alkyl, Het²C₁₋₄alkyl, Het²oxyC₁₋₄alkyl, arylC₁₋₄alkyl, aryloxyC₁₋₄alkyl or aminoC₁₋₄alkyl; whereby each amino group may optionally be mono- or where possible disubstituted with C₁₋₄alkyl;
- R₅ and -A-R₆ taken together with the nitrogen atom to which they are attached may also form Het¹ or Het²;
- R₇ is hydrogen, aryl, C₃₋₇cycloalkyl, Het¹, Het², C₂₋₆alkenyl, C₂₋₆alkynyl or C₁₋₆alkyl optionally substituted with aryl, Het¹, Het², C₃₋₇cycloalkyl, C₁₋₄alkyloxycarbonyl, carboxyl, aminocarbonyl, mono- or di(C₁₋₄alkyl)aminocarbonyl, aminosulfonyl, C₁₋₄alkylS(=O)₆, hydroxy, cyano, halogen or amino optionally mono- or disubstituted where the substituents are selected from C₁₋₄alkyl, aryl, arylC₁₋₄alkyl, C₃₋₇cycloalkyl, C₃₋₇cycloalkylC₁₋₄alkyl, Het¹, Het², Het¹C₁₋₄alkyl and Het²C₁₋₄alkyl;
- R₈ is hydrogen, C₁₋₆alkyl, C₂₋₆alkenyl, arylC₁₋₆alkyl, C₃₋₇cycloalkyl, C₃₋₇cycloalkyl-C₁₋₆alkyl, aryl, Het¹, Het¹C₁₋₆alkyl, Het², Het²C₁₋₆alkyl;
- R₁₂ is hydrogen, -NH₂, -C₁₋₆alkyl or C₁₋₆alkyl-W-R₁₄, wherein said C₁₋₆alkyl is

- optionally substituted with halogen, hydroxy, aryl, heteroaryl, Het¹, Het², or amino wherein said amino is optionally mono- or di-substituted with C₁₋₄ alkyl; or R₁₂ may also be -N(R₅)(AR₆), wherein
- A is C₁₋₆alkanediyl, -C(=O)-, -C(=S)-, -S(=O)₂-, C₁₋₆alkanediyl-C(=O)-, C₁₋₆alkanediyl-C(=S)- or C₁₋₆alkanediyl-S(=O)₂-; wherein if A is C₁₋₆alkanediyl-C(=O)-, C₁₋₆alkanediyl-C(=S)- or C₁₋₆alkanediyl-S(=O)₂-, the NR₅ moiety is attached to the C₁₋₆alkanediyl moiety;
- W is oxy, carbonyl, oxycarbonyl, carbonyloxy, oxycarbonyloxy, amino, aminocarbonyl, carbonylamino or sulphur;
- R₁₃ is hydrogen, C₁₋₆-alkyl, optionally substituted by aryl, Het¹, Het², hydroxy, halogen, amino whereby the amino group may be optionally be mono- or di-substituted with C₁₋₄alkyl;
- R₁₄ is C₁₋₆alkyl, aryl, Het¹ or Het².
- 15 This invention also includes the quaternization of the nitrogen atoms of the present compounds. A basic nitrogen can be quaternized with any agent known to those of ordinary skill in the art including, for instance, lower alkyl halides, dialkyl sulfates, long chain halides and aralkyl halides.
- 20 Whenever the term substituted is used in defining the compounds of formula (I), it is meant to indicate that one or more hydrogens on the atom indicated in the expression using "substituted" is replaced with a selection from the indicated group, provided that the indicated atom's normal valency is not exceeded, and that the substitution results in a chemically stable compound, i.e. a compound that is sufficiently robust to survive
- 25 isolation to a useful degree of purity from a reaction mixture, and formulation into a therapeutic agent.
- As used herein, the term "halo" or "halogen" as a group or part of a group is generic for fluoro, chloro, bromo or iodo.
- 30 The term "C₁₋₄alkyl" as a group or part of a group defines straight and branched chained saturated hydrocarbon radicals having from 1 to 4 carbon atoms, such as, for example, methyl, ethyl, propyl, butyl and 2-methyl-propyl, the like.
- The term "C₁₋₆alkyl" as a group or part of a group defines straight and branched chained saturated hydrocarbon radicals having from 1 to 6 carbon atoms such as the
- 35 groups defined for C₁₋₄alkyl and pentyl, hexyl, 2-methylbutyl, 3-methylpentyl and the like.
- The term "C₁₋₆alkanediyl" as a group or part of a group defines bivalent straight and branched chained saturated hydrocarbon radicals having from 1 to 6 carbon atoms such

as, for example, methylene, ethan-1,2-diyl, propan-1,3-diyl, propan-1,2-diyl, butan-1,4-diyl, pentan-1,5-diyl, hexan-1,6-diyl, 2-methylbutan-1,4-diyl, 3-methylpentan-1,5-diyl and the like.

- 5 The term "C₂₋₆alkenyl" as a group or part of a group defines straight and branched chained saturated hydrocarbon radicals having from 2 to 6 carbon atoms containing at least one double bond such as, for example, ethenyl, propenyl, butenyl, pentenyl, hexenyl and the like.

- 10 The term "C₂₋₆alkynyl" as a group or part of a group defines straight and branched chained saturated hydrocarbon radicals having from 2 to 6 carbon atoms containing at least one triple bond such as, for example, ethynyl, propynyl, butynyl, pentynyl, hexynyl and the like.

The term "C₃₋₇cycloalkyl" as a group or part of a group is generic to cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl or cycloheptyl.

- 15 The term "aryl" as a group or part of a group is meant to include phenyl and naphthyl which both may be optionally substituted with one or more substituents independently selected from C₁₋₆alkyl, C₁₋₆alkyloxy, aminoC₁₋₆alkyl, halogen, hydroxy, optionally mono- or disubstituted amino, nitro, cyano, haloC₁₋₆alkyl, carboxyl, C₁₋₆alkoxy-carbonyl, C₃₋₇cycloalkyl, Het¹, optionally mono- or disubstituted aminocarbonyl, methylthio, methylsulfonyl, and phenyl optionally substituted with one or more
20 substituents selected from C₁₋₆alkyl, C₁₋₆alkyloxy, halogen, hydroxy, optionally mono- or disubstituted amino, nitro, cyano, haloC₁₋₆alkyl, carboxyl, C₁₋₆alkoxycarbonyl, C₃₋₇cycloalkyl, Het¹, optionally mono- or disubstituted aminocarbonyl, methylthio and methylsulfonyl; whereby the optional substituents on any amino function are
25 independently selected from C₁₋₆alkyl, C₁₋₆alkyloxy-A-, Het¹-A-, Het¹C₁₋₆alkyl, Het¹C₁₋₆alkyl-A-, Het¹oxy-A-, Het¹oxyC₁₋₆alkyl-A-, phenyl-A-, phenyloxy-A-, phenyloxyC₁₋₆alkyl-A-, phenylC₁₋₆alkyl-A-, C₁₋₆alkyloxycarbonylamino-A-, amino-A-, aminoC₁₋₆alkyl and aminoC₁₋₆alkyl-A- whereby each of the amino groups may optionally be mono- or where possible di-substituted with C₁₋₆alkyl and whereby A is
30 as defined above.

- The term "haloC₁₋₆alkyl" as a group or part of a group is defined as C₁₋₆alkyl substituted with one or more halogen atoms, preferably, chloro or fluoro atoms, more preferably fluoro atoms. PolyhaloC₁₋₆alkyl as a group or as part of a group is defined as C₁₋₆alkyl substituted with 2 or more halogen atoms. Preferred haloC₁₋₆alkyl group
35 include for instance trifluoromethyl and difluoromethyl.

The term "Het¹" as a group or part of a group is defined as a saturated or partially unsaturated monocyclic, bicyclic or tricyclic heterocycle having preferably 3 to 14 ring members, more preferably 5 to 10 ring members and more preferably 5 to 6 ring

- members, which contains one or more heteroatom ring members selected from nitrogen, oxygen or sulphur and which is optionally substituted on one or more carbon atoms by C₁₋₆alkyl, C₁₋₆alkyloxy, aminoC₁₋₆alkyl, halogen, hydroxy, oxo, optionally mono- or disubstituted amino, optionally mono- or disubstituted aminoalkyl, nitro, cyano, haloC₁₋₆alkyl, carboxyl, C₁₋₆alkoxycarbonyl, C₃₋₇cycloalkyl, optionally mono- or disubstituted aminocarbonyl, methylthio, methylsulfonyl, aryl and a saturated or partially unsaturated monocyclic, bicyclic or tricyclic heterocycle having 3 to 14 ring members; whereby the optional substituents on any amino function are independently selected from C₁₋₆alkyl, C₁₋₆alkyloxy-A-, Het²-A-, Het²C₁₋₆alkyl, Het²C₁₋₆alkyl-A-, Het²oxy-A-, Het²oxyC₁₋₄alkyl-A-, aryl-A-, aryloxy-A-, aryloxyC₁₋₄alkyl-A-, arylC₁₋₆alkyl-A-, C₁₋₆alkyloxycarbonylamino-A-, amino-A-, aminoC₁₋₆alkyl and aminoC₁₋₆alkyl-A- whereby each of the amino groups may optionally be mono- or where possible di-substituted with C₁₋₄alkyl and whereby A is as defined above.
- The term "Het²" as a group or part of a group is defined as an aromatic monocyclic, bicyclic or tricyclic heterocycle having preferably 3 to 14 ring members, more preferably 5 to 10 ring members and more preferably 5 to 6 ring members, which contains one or more heteroatom ring members selected from nitrogen, oxygen or sulphur and which is optionally substituted on one or more carbon atoms by C₁₋₆alkyl, C₁₋₆alkyloxy, aminoC₁₋₆alkyl, halogen, hydroxy, optionally mono- or disubstituted amino, nitro, cyano, haloC₁₋₆alkyl, carboxyl, C₁₋₆alkoxycarbonyl, C₃₋₇cycloalkyl, optionally mono- or disubstituted aminocarbonyl, methylthio, methylsulfonyl, aryl, Het¹ and an aromatic monocyclic, bicyclic or tricyclic heterocycle having 3 to 12 ring members; whereby the optional substituents on any amino function are independently selected from C₁₋₆alkyl, C₁₋₆alkyloxy-A-, Het¹-A-, Het¹C₁₋₆alkyl, Het¹C₁₋₆alkyl-A-, Het¹oxy-A-, Het¹oxyC₁₋₄alkyl-A-, aryl-A-, aryloxy-A-, aryloxyC₁₋₄alkyl-A-, arylC₁₋₆alkyl-A-, C₁₋₆alkyloxycarbonylamino-A-, amino-A-, aminoC₁₋₆alkyl and aminoC₁₋₆alkyl-A- whereby each of the amino groups may optionally be mono- or where possible di-substituted with C₁₋₄alkyl and whereby A is as defined above.
- As used herein, the term (=O) forms a carbonyl moiety with the carbon atom to which it is attached. The term (=O) forms a sulfoxide with the sulfur atom to which it is attached. The term (=O)₂ forms a sulfonyl with the sulfur atom to which it is attached. As used herein, the term (=S) forms a thiocarbonyl moiety with the carbon atom to which it is attached.
- As used herein before, the term "one or more" covers the possibility of all the available C-atoms, where appropriate, to be substituted, preferably, one, two or three.

When any variable (e.g. halogen or C₁₋₄alkyl) occurs more than one time in any constituent, each definition is independent.

5 Lines drawn from substituents into ring systems indicate that the bond may be attached to any of the suitable ring atoms.

The term "prodrug" as used throughout this text means the pharmacologically acceptable derivatives such as esters, amides and phosphates, such that the resulting *in vivo* biotransformation product of the derivative is the active drug as defined in the compounds of formula (I). The reference by Goodman and Gilman (The Pharmacological Basis of Therapeutics, 8th ed, McGraw-Hill, Int. Ed. 1992, "Biotransformation of Drugs", p 13-15) describing prodrugs generally is hereby incorporated. Prodrugs of a compound of the present invention are prepared by modifying functional groups present in the compound in such a way that the modifications are cleaved, either in routine manipulation or *in vivo*, to the parent compound. Prodrugs of the compounds of the present invention include those compounds wherein for instance a hydroxy group, such as the hydroxy group on the asymmetric carbon atom, or an amino group is bonded to any group that, when the prodrug is administered to a patient, cleaves to form a free hydroxyl or free amino, respectively.

20 Typical examples of prodrugs are described for instance in WO 99/33795, WO 99/33815, WO 99/33793 and WO 99/33792 all incorporated herein by reference.

25 Prodrugs are characterized by excellent aqueous solubility, increased bioavailability and are readily metabolized into the active inhibitors *in vivo*.

For therapeutic use, the salts of the compounds of formula (I) are those wherein the counterion is pharmaceutically or physiologically acceptable. However, salts having a pharmaceutically unacceptable counterion may also find use, for example, in the preparation or purification of a pharmaceutically acceptable compound of formula (I). All salts, whether pharmaceutically acceptable or not are included within the ambit of the present invention.

35 The pharmaceutically acceptable or physiologically tolerable addition salt forms which the compounds of the present invention are able to form can conveniently be prepared using the appropriate acids, such as, for example, inorganic acids such as hydrohalic acids, e.g. hydrochloric or hydrobromic acid; sulfuric; nitric; phosphoric and the like acids; or organic acids such as, for example, acetic, propanoic, hydroxyacetic, lactic, pyruvic, oxalic, malonic, succinic, maleic, fumaric, malic, tartaric, citric, methane-

sulfonic, ethanesulfonic, benzenesulfonic, *p*-toluenesulfonic, cyclamic, salicylic, *p*-aminosalicylic, pamoic and the like acids.

- 5 Conversely said acid addition salt forms can be converted by treatment with an appropriate base into the free base form.

- 10 The compounds of formula (I) containing an acidic proton may also be converted into their non-toxic metal or amine addition salt form by treatment with appropriate organic and inorganic bases. Appropriate base salt forms comprise, for example, the ammonium salts, the alkali and earth alkaline metal salts, e.g. the lithium, sodium, potassium, magnesium, calcium salts and the like, salts with organic bases, e.g. the benzathine, N-methyl, -D-glucamine, hydrabamine salts, and salts with amino acids such as, for example, arginine, lysine and the like.

- 15 Conversely said base addition salt forms can be converted by treatment with an appropriate acid into the free acid form.

- 20 The term "salts" also comprises the hydrates and the solvent addition forms which the compounds of the present invention are able to form. Examples of such forms are e.g. hydrates, alcoholates and the like. Generically they also are called solvates.

The *N*-oxide forms of the present compounds are meant to comprise the compounds of formula (I) wherein one or several nitrogen atoms are oxidized to the so-called *N*-oxide.

- 25 The present compounds may also exist in their tautomeric forms. Such forms, although not explicitly indicated in the above formula are intended to be included within the scope of the present invention.


- 30 The term stereochemically isomeric forms of compounds of the present invention, as used hereinbefore, defines all possible compounds made up of the same atoms bonded by the same sequence of bonds but having different three-dimensional structures which are not interchangeable, which the compounds of the present invention may possess. Unless otherwise mentioned or indicated, the chemical designation of a compound encompasses the mixture of all possible stereochemically isomeric forms which said
35 compound may possess. Said mixture may contain all diastereomers and/or enantiomers of the basic molecular structure of said compound. All stereochemically isomeric forms of the compounds of the present invention both in pure form or in admixture with each other are intended to be embraced within the scope of the present invention.

Pure stereoisomeric forms of the compounds and intermediates as mentioned herein are defined as isomers substantially free of other enantiomeric or diastereomeric forms of the same basic molecular structure of said compounds or intermediates. In particular, the term 'stereoisomerically pure' concerns compounds or intermediates having a stereoisomeric excess of at least 80% (i. e. minimum 90% of one isomer and maximum 10% of the other possible isomers) up to a stereoisomeric excess of 100% (i.e. 100% of one isomer and none of the other), more in particular, compounds or intermediates having a stereoisomeric excess of 90% up to 100%, even more in particular having a stereoisomeric excess of 94% up to 100% and most in particular having a stereoisomeric excess of 97% up to 100%. The terms 'enantiomerically pure' and 'diastereomerically pure' should be understood in a similar way, but then having regard to the enantiomeric excess, respectively the diastereomeric excess of the mixture in question.

Pure stereoisomeric forms of the compounds and intermediates of this invention may be obtained by the application of art-known procedures. For instance, enantiomers may be separated from each other by the selective crystallization of their diastereomeric salts with optically active acids. Alternatively, enantiomers may be separated by chromatographic techniques using chiral stationary phases. Said pure stereochemically isomeric forms may also be derived from the corresponding pure stereochemically isomeric forms of the appropriate starting materials, provided that the reaction occurs stereospecifically. Preferably, if a specific stereoisomer is desired, said compound will be synthesized by stereospecific methods of preparation. These methods will advantageously employ enantiomerically pure starting materials.

The diastereomeric racemates of formula (I) can be obtained separately by conventional methods. Appropriate physical separation methods which may advantageously be employed are, for example, selective crystallization and chromatography, e.g. column chromatography.

It is clear to a person skilled in the art that the compounds of formula (I) contain at least one asymmetric center and thus may exist as different stereoisomeric forms. This asymmetric center is indicated with an asterisk (*) in the figure below.



The absolute configuration of each asymmetric center that may be present in the compounds of formula (I) may be indicated by the stereochemical descriptors R and S, this R and S notation corresponding to the rules described in Pure Appl. Chem. 1976, 45, 11-30. The carbon atom marked with the asterisk (*) preferably has the

5 R configuration.

The present invention is also intended to include all isotopes of atoms occurring on the present compounds. Isotopes include those atoms having the same atomic number but different mass numbers. By way of general example and without limitation, isotopes of

10 hydrogen include tritium and deuterium. Isotopes of carbon include C-13 and C-14.

Whenever used hereinafter, the term "compounds of formula (I)", or "the present compounds" or similar term is meant to include the compounds of general formula (I), their *N*-oxides, salts, stereoisomeric forms, racemic mixtures, prodrugs, esters and

15 metabolites, as well as their quaternized nitrogen analogues.

An interesting group of compounds are those compounds of formula (I) wherein L is -C(=O)-, -O-C(=O)-, -NR₈-C(=O)-, -O-C₁₋₆alkanediyl-C(=O)-, -NR₈-C₁₋₆alkanediyl-C(=O)-, -S(=O)₂-, -O-S(=O)₂-, -NR₈-S(=O)₂-,

20 -NR₈-C₁₋₆alkanediyl-S(=O)₂, whereby either the C(=O) group or the S(=O)₂ group is attached to the NR₂ moiety.

A particular group of compounds are those compounds of formula (I) wherein one or more of the following restrictions apply:

25 R₁ is hydrogen, Het¹, Het², aryl, Het¹C₁₋₆alkyl, Het²C₁₋₆alkyl, arylC₁₋₆alkyl, more in particular, R₁ is hydrogen, a saturated or partially unsaturated monocyclic or bicyclic heterocycle having 5 to 8 ring members, which contains one or more heteroatom ring members selected from nitrogen, oxygen or sulfur and which is optionally substituted; phenyl optionally substituted with one or more

30 substituents; an aromatic monocyclic heterocycle having 5 to 6 ring members, which contains one or more heteroatom ring members selected from nitrogen, oxygen or sulfur and which is optionally substituted on one or more carbon atoms; or C₁₋₆alkyl substituted with an aromatic monocyclic heterocycle having 5 to 6 ring members, which contains one or more heteroatom ring members

35 selected from nitrogen, oxygen or sulfur and which is optionally substituted on one or more carbon atoms;

R₂ is hydrogen;

- L is $-C(=O)-$, $-O-C(=O)-$, $-O-C_{1-6}alkanediy-C(=O)-$, more in particular, L is $-C(=O)-$, $-O-C(=O)-$, $-O-CH_2-C(=O)-$, whereby in each case the $C(=O)$ group is attached to the NR_2 moiety;
- R_3 is aryl $C_{1-4}alkyl$, in particular, arylmethyl, more in particular phenylmethyl;
- 5 R_4 is optionally substituted $C_{1-6}alkyl$, in particular $C_{1-6}alkyl$ optionally substituted with aryl, Het^1 , Het^2 , $C_{3-7}cycloalkyl$ or amino optionally mono- or disubstituted where the substituents are selected from $C_{1-4}alkyl$, aryl, Het^1 and Het^2 ;
- R_5 is hydrogen, hydroxy, $C_{1-6}alkyl$, $Het^1C_{1-6}alkyl$, $Het^2C_{1-6}alkyl$, amino $C_{1-6}alkyl$ whereby the amino group may optionally be mono- or di-substituted with
- 10 $C_{1-4}alkyl$;
- R_6 is $C_{1-6}alkyloxy$, Het^1 , Het^1oxy , Het^2 , Het^2oxy , aryl, aryloxy, aryloxy $C_{1-4}alkyl$, $C_{1-4}alkyloxyaryl$, $C_{1-4}alkyloxyHet^1$, $C_{1-4}alkyloxyHet^2$, $C_{1-4}alkyloxycarbonyl-$ amino, amino $C_{1-4}alkylamino$, amino or amino $C_{1-4}alkyloxy$ and in case A is other than $C_{1-6}alkanediy$, then R_6 may also be $C_{1-6}alkyl$, $Het^1C_{1-4}alkyl$,
- 15 $Het^1oxyC_{1-4}alkyl$, $Het^2C_{1-4}alkyl$, $Het^2oxyC_{1-4}alkyl$, aryl $C_{1-4}alkyl$, aryloxy $C_{1-4}alkyl$ or amino $C_{1-4}alkyl$; whereby each amino group may optionally be mono- or where possible disubstituted with $C_{1-4}alkyl$;
- R_5 and $-A-R_6$ taken together with the nitrogen atom to which they are attached may also form Het^1 or Het^2 ;
- 20 R_{12} is H, $-NH_2$, $-N(R_5)(AR_6)$, $-C_{1-6}alkyl$ or $C_{1-6}alkyl-W-R_{14}$, wherein said $C_{1-6}alkyl$ is optionally substituted with halogen, hydroxy, aryl, heteroaryl, Het^1 , Het^2 , or amino wherein said amino is optionally mono- or di-substituted with $C_{1-4}alkyl$;
- A is $C_{1-6}alkanediy$, $-C(=O)-$, $C_{1-6}alkanediy-C(=O)-$;
- W is oxy, carbonyl, oxycarbonyl, carbonyloxy, oxycarbonyloxy, amino,
- 25 aminocarbonyl, carbonylamino or sulphur;
- R_{13} is H, $C_{1-6}alkyl$, optionally substituted by aryl, Het^1 , Het^2 , hydroxy, halogen, amino whereby the amino group may be optionally be mono- or di-substituted with $C_{1-4}alkyl$;
- R_{14} is $C_{1-6}alkyl$, aryl Het^1 or Het^2 .
- 30
- An interesting group of compounds are those compounds of formula (I) wherein one or more of the following restrictions apply
- R_1 is thiazolyl, hexahydrofuro[2,3-b]furan-3-yl, pyrrolidinyl or phenyl which is mono- di- or trisubstituted, wherein the substituents are selected from the group consisting of methyl, nitro, amino, mono or dimethylamino, hydroxy, aminomethyl,
- 35 wherein at least one of said substituents is in ortho vis-à-vis the R_1-L bond.
- L is $-O-C(=O)-$, $-O-CH_2-C(=O)-$, $-CH_2-O-C(=O)-$.

A special group of compounds are those compounds of formula (I) wherein R_1 is Het^1 , aryl, $Het^2C_{1-6}alkyl$; R_2 is hydrogen; L is $-C(=O)-$, $-O-C(=O)-$, $-O-CH_2-C(=O)-$, whereby in each case the $C(=O)$ group is attached to the NR_2 moiety; R_3 is phenylmethyl; and R_4 is $C_{1-6}alkyl$.

5

Another interesting group of compounds are those compounds of formula (I) wherein L is $-O-C_{1-6}alkanediyl-C(=O)-$.

Another interesting group of compounds are those compounds of formula (I) wherein R_{12} is H , $-NH_2$, $-N(R_5)(AR_6)$, $-C_{1-6}alkyl$ or $C_{1-6}alkyl-W-R_{14}$, wherein said $C_{1-6}alkyl$ is optionally substituted with halogen, hydroxy, aryl, heteroaryl, Het^1 , Het^2 , or amino optionally mono- or di-substituted with $C_{1-4}alkyl$;

A is $C_{1-6}alkanediyl$ or $-C(=O)-$; R_5 is hydrogen, methyl, $Het^1C_{1-6}alkyl$, $aminoC_{1-6}alkyl$ whereby the amino group may optionally be mono- or di-substituted with $C_{1-4}alkyl$; R_6 is $C_{1-6}alkyloxy$, Het^1 , amino; and in case $-A-$ is other than $C_{1-6}alkanediyl$ then R_6 may also be $C_{1-6}alkyl$, $Het^1C_{1-4}alkyl$ or $aminoC_{1-4}alkyl$; whereby each of the amino groups may optionally be substituted;

W is oxycarbonyloxy;

R_{14} is Het^1 .

20

Another interesting group of compounds are those compounds of formula (I) wherein R_{12} is $-N(R_5)(AR_6)$;

A is $C_{1-6}alkanediyl$ or $-C(=O)-$; R_5 is hydrogen; R_6 is $C_{1-6}alkyloxy$, Het^1 , $C_{1-4}alkyloxy-aryl$, amino, $aminoC_{1-4}alkylamino$, $aminoC_{1-4}alkyloxy$; and in case $-A-$ is $-C(=O)-$ then R_6 may also be $C_{1-6}alkyl$, $Het^1C_{1-4}alkyl$, or $aminoC_{1-4}alkyl$; whereby each of the amino groups may optionally be substituted with $C_{1-4}alkyl$.

Another interesting group of compounds are those compounds of formula (I) wherein A is $C_{1-6}alkanediyl$ or $-C(=O)-$;

R_5 is hydrogen, $C_{1-6}alkyl$, $Het^1C_{1-6}alkyl$, $Het^2C_{1-6}alkyl$, $aminoC_{1-6}alkyl$ whereby the amino group may optionally be mono- or di-substituted with $C_{1-4}alkyl$; and

in case $-A-$ is $-C(=O)-$ then R_6 is $C_{1-6}alkyloxy$, Het^1 , Het^1oxy or Het^2oxy , aryl, $Het^1C_{1-4}alkyl$, $Het^1oxyC_{1-4}alkyl$, $Het^2C_{1-4}alkyl$, $Het^2oxyC_{1-4}alkyl$, $arylC_{1-4}alkyl$, $aryloxyC_{1-4}alkyl$ or $aminoC_{1-4}alkyl$; and

in case $-A-$ is $C_{1-6}alkanediyl$ then R_6 is amino, $C_{1-6}alkyloxy$, Het^1 , Het^1oxy or Het^2oxy ; and

in case $-A-$ is $C_{1-6}alkanediyl-C(=O)-$ then R_6 is $C_{1-6}alkyloxy$, Het^1 , Het^1oxy or Het^2oxy , aryl, $C_{1-6}alkyl$, $Het^1C_{1-4}alkyl$, $Het^1oxyC_{1-4}alkyl$, $Het^2C_{1-4}alkyl$, $Het^2oxyC_{1-4}alkyl$, $arylC_{1-4}alkyl$, $aryloxyC_{1-4}alkyl$ or $aminoC_{1-4}alkyl$;

35

whereby each of the amino groups in the definition of R_6 may optionally be substituted with one or more substituents selected from C_{1-4} alkyl, C_{1-4} alkylcarbonyl, C_{1-4} alkyloxycarbonyl, aryl, arylcarbonyl, aryloxycarbonyl, Het^1 , Het^2 , aryl C_{1-4} alkyl, Het^1C_{1-4} alkyl or Het^2C_{1-4} alkyl; and

- 5 R_5 and $-A-R_6$ taken together with the nitrogen atom to which they are attached may also form Het^1 whereby Het^1 is substituted by at least an oxo group.

Another interesting group of compounds are those compounds of formula (I) wherein R_{13} is C_{1-4} alkyl preferably methyl, ethyl or isobutyl; optionally substituted with aryl
10 preferably phenyl, Het^1 preferably pyrrolidinyl, Het^2 preferably imidazolyl or pyridinyl, or amino; wherein the amino group is optionally substituted with C_{1-4} alkyl.

Also a special group of compounds are those compounds of formula (I) wherein R_{12} is H, $-NH_2$, $-N(R_5)(AR_6)$, $-C_{1-6}$ alkyl or C_{1-6} alkyl-W- R_{14} , wherein said C_{1-6} alkyl is
15 optionally substituted with hydroxy or amino wherein said amino is optionally mono- or di-substituted with C_{1-4} alkyl;

A is C_{1-6} alkanediyl or $-C(=O)-$; R_5 is hydrogen, methyl, Het^1C_{1-6} alkyl, amino C_{1-6} alkyl whereby the amino group may optionally be mono- or di-substituted with C_{1-4} alkyl; R_6 is C_{1-6} alkyloxy, Het^1 , amino; and in case $-A-$ is other than C_{1-6} alkanediyl then R_6 may
20 also be C_{1-6} alkyl, Het^1C_{1-4} alkyl or amino C_{1-4} alkyl; whereby each of the amino groups may optionally be substituted;

R_{13} is H, C_{1-6} -alkyl, optionally substituted by aryl, Het^1 , Het^2 , hydroxy, halogen, amino whereby the amino group may be optionally be mono- or di-substituted with C_{1-4} alkyl.
 R_{14} is alkyl, aryl Het^1 , Het^2 .

25

Another interesting group of compounds are those compounds of formula (I) wherein R_1 hydrogen, C_{1-6} alkyl, C_{2-6} alkenyl, aryl C_{1-6} alkyl, C_{3-7} cycloalkyl, C_{3-7} cycloalkyl- C_{1-6} alkyl, aryl, Het^1 , Het^1C_{1-6} alkyl, Het^2 , Het^2C_{1-6} alkyl; wherein Het^1 is a saturated or
30 partially unsaturated monocyclic heterocycle having 5 or 6 ring members, which contains one or more heteroatom ring members selected from nitrogen, oxygen or sulfur and which is optionally substituted on one or more carbon atoms.

Another interesting group of compounds are those compounds of formula (I) wherein:
 R_{13} is H, C_{1-6} -alkyl, optionally substituted by aryl, Het^1 , Het^2 , amino whereby the
35 amino group may be optionally be mono- or di-substituted with C_{1-4} alkyl;
 R_{12} is H, $-NH_2$, $-N(R_5)(AR_6)$, $-C_{1-6}$ alkyl or C_{1-6} alkyl-W- R_{14} , wherein said C_{1-6} alkyl is optionally substituted with hydroxy or amino wherein said amino is optionally mono- or di-substituted with C_{1-4} alkyl;

A is C_{1-6} alkanediyl, $-C(=O)-$, C_{1-6} alkanediyl- $C(=O)-$,

- R₅ is hydrogen, hydroxy, C₁₋₆alkyl, Het¹C₁₋₆alkyl, Het²C₁₋₆alkyl, aminoC₁₋₆alkyl whereby the amino group may optionally be mono- or di-substituted with C₁₋₄alkyl;
- R₆ is C₁₋₆alkyloxy, Het¹, Het¹C₁₋₄alkyl, Het¹oxy, Het¹oxyC₁₋₄alkyl, Het², Het²C₁₋₄alkyl, Het²oxy, Het²oxyC₁₋₄alkyl, aryl, aryloxy, aryloxyC₁₋₄alkyl, arylC₁₋₄alkyl, C₁₋₆alkyloxycarbonylamino, amino or aminoC₁₋₄alkyl whereby each amino group may optionally be mono- or where possible di-substituted with C₁₋₄alkyl; and R₆ may also be hydrogen in case -A- is C₁₋₆alkanediyl;
- R₅ and -A-R₆ taken together with the nitrogen atom to which they are attached may also form Het¹ or Het²;
- W is oxy, carbonyl, oxycarbonyl, carbonyloxy, oxycarbonyloxy, amino, aminocarbonyl, carbonylamino or sulphur;
- R₁₄ is alkyl, aryl Het¹ or Het².
- Another interesting group of compounds are those compounds of formula (I) wherein both R₁₂ and R₁₃ are present.
- An interesting group of compounds are those compounds of the formula (I) wherein:
- R₁ is a bicyclic heterocycle comprising at least 7 atoms, of which at least one is O;
- L is -O-C(=O)-, -C(=O)-, -CH₂-O-C(=O)- or -O-CH₂-C(=O)-;
- R₂ is -H;
- R₃ is C₁₋₄ alkylphenyl;
- R₄ is C₄ alkyl;
- R₁₂ is -H; -NH₂-N(R₅)(AR₆), - C₁₋₆alkyl or C₁₋₆alkyl -W-R₁₄, wherein said C₁₋₆alkyl is optionally substituted with hydroxy or amino wherein said amino is optionally mono- or di-substituted with C₁₋₄alkyl;
- A is C₁₋₆alkanediyl or -C(=O)- ;
- R₅ is hydrogen;
- R₆ is C₁₋₆alkyloxy, Het¹, amino; and in case -A- is other than C₁₋₆alkanediyl then R₆ may also be C₁₋₆alkyl, Het¹C₁₋₄alkyl or aminoC₁₋₄alkyl; whereby each of the amino groups may optionally be substituted;
- W is oxycarbonyloxy;
- R₁₄ is Het¹;
- R₁₃ is H, C₁₋₄alkyl, C₁₋₄ alkylaryl, C₁₋₄ alkylHet¹, C₁₋₄ alkylHet².
- Another interesting group of compounds are those compounds of the formula (I) wherein:
- R₁ is a bicyclic heterocycle comprising at least 7 atoms, of which at least one is O;
- L is -O-C(=O)-, -C(=O)-, -CH₂-O-C(=O)- or -O-CH₂-C(=O)-;

WO 03/076413

PCT/EP03/50057

-17-

R₂ is -H;

R₃ is C₁₋₄ alkylphenyl;

R₄ is C₄ alkyl;

R₁₂ is H;

5 R₁₃ is C₁₋₄ alkylaryl, C₁₋₄ alkylHet¹, C₁₋₄ alkylHet².

An interesting group of compounds are those compounds of formula (I) wherein one or more of the following restrictions apply

10 R₁ is thiazolyl, hexahydro furo[2,3-b]furan-3-yl, pyrrolidinyl or phenyl which is mono-di- or trisubstituted, wherein the substituents are selected from the group consisting of methyl, nitro, amino, mono or dimethylamino, hydroxy, aminomethyl, wherein at least one of said substituents is in ortho vis-à-vis the R₁-L bond.

L is -O-C(=O)-, -O-CH₂-C(=O)-, -CH₂-O-C(=O)-;

R₁₂ is hydrogen, methyl, methylcarbonylamino, methyloxycarbonylamino;

15 R₁₃ is hydrogen, methyl, isobutyl, phenylmethyl, phenylethyl.

Another interesting group of compounds are the compounds of formula (I) of structure 1 to 64 as shown in table 1 and 2. An interesting subgroup thereof are the compounds of formula (I) having a number 1-13, 15, 17-19, 21, 22, 24, 37 and 64.

20


An interesting group of compounds are the compounds of formula (I) which contain at least one group selected from thiazolyl, imidazolyl or pyridinyl.

25 The invention also relates to the methods of synthesis of the compounds of the present invention.

The invention further relates to the intermediates in the synthesis of the compounds of the present invention. Interesting intermediates are the intermediates of formula D-2, E-2, F-2, in particular F-2. Another group of interesting intermediates are those 30 intermediates of formula D-4, F-4, in particular F-4. A further interesting group of intermediates are those intermediates of formula F-5, G-5, and H-5, in particular H-5. Also interesting intermediates are those of formula G-5, H-5 and J-7.

35 The invention also relates to a compound of formula (I) obtainable by a process as shown in schemes 1 to 7.

Preferably the compounds of formula (I) have the stereochemistry as indicated in formula (I').



The compounds of formula (I) can generally be prepared using procedures analogous to those procedures described in WO 95/06030, WO 96/22287, WO 96/28418, WO 96/28463, WO 96/28464, WO 96/28465, WO 99/59989 and WO 97/18205.

5


Particular reaction procedures to make the present compounds are described below. In the preparations described below, the reaction products may be isolated from the medium and, if necessary, further purified according to methodologies generally known in the art such as, for example, extraction, crystallization, trituration and chromatography.

10

The following abbreviations are used throughout the synthesis procedures. Boc: t-butyloxycarbonyl, DCM: dichloromethane, DCC: dicyclohexylcarbodiimide; DMF: dimethylformamide; EDCI: 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloric acid; HOBt: 1-hydroxy-1-H-benzotriazole; THF: tetrahydrofurane; mCPBA: metachloroperbenzoic acid. In the following synthesis all substituents R_i , wherein i is an integer from 1 to 14, are as defined *supra*, except if expressly stated otherwise, PG and PG' each independently indicate protecting groups.

15

Scheme 1



20 Compound **A-1** was reacted with chlorosulfonic acid to yield intermediate **B-1**. Those

skilled in the art will appreciate that this intermediate tautomerizes. The depicted **B-1** intermediate is one example intermediate.

Intermediate **D-1** was prepared by reacting an intermediate **C-1**, obtained according to the procedure described in patent WO97/18205 and also depicted in scheme 8, with an intermediate **B-1**, in a reaction-inert solvent such as dichloromethane, in the presence of a base such as triethylamine, at low temperature, for example 0°C.

The aminoterminal protective group in the intermediates may be a protective group known in the art such as *tert*-butyloxycarbonyl group. This protective group may conveniently be replaced by another suitable protective group such as phtalimido, dibenzyl or benzyloxycarbonyl.


Intermediate **D-1** can be deprotected using an acid such as trifluoroacetic acid in a suitable solvent such as dichloromethane, yielding an intermediate **E-1**.


Alternatively, intermediates may be deprotected with a strong acid such as hydrochloric acid in isopropanol, in a suitable solvent such as a mixture of ethanol and dioxane.

The aminoterminal group may subsequently be substituted by procedures known in the art to generate **F-1**. As is indicated *supra*, the compound of formula **F-1** tautomerizes, therefore, the indicated formula is one tautomer.

In a preferred embodiment the protecting group is selected from Fmoc, Acetyl, *tert*-Butyloxycarbonyl, Benzyloxycarbonyl-, Dibenzyl-.

Scheme 2





3-fluoroaniline was nitrated in conditions known in the art to yield intermediate **A-2**, which was further reacted as described in Helvetica Chimica Acta, (1983), 66(1), p 68-75 to give intermediate **B-2**.

Intermediate **D-2** was prepared by reacting an intermediate **C-1**, obtained according to the procedure described in patent WO97/18205 and also depicted in scheme 8, with an intermediate **B-2**, in a reaction-inert solvent such as dichloromethane, in the presence of a base such as triethylamine, at low temperature, for example 0°C.

The aminoterminal protective group in the intermediates may be a protective group known in the art such as *tert*-butoxycarbonyl group. This protective group may conveniently be replaced by another suitable protective group such as phthalimido, dibenzyl or benzyloxycarbonyl.

Nucleophilic substitution of the fluorine atom by a primary amine under heating conditions led to intermediate **E-2**, whose nitro group was reduced to an amino group using a procedure known in the art such as catalytic hydrogenation or transfer hydrogenation, yielding intermediate **F-2**.

Cyclisation of **F-2** with methylorthoformate ($R_{12} = H$) or a carboxylic acid of formula $R_{12}COOH$ (R_{12} differs from H) in an acidic solvent such as hydrochloric acid, or with BrCN led to compounds **G-2**, wherein respectively $R_{12} = H$ (cyclization with methylorthoformate), R_{12} differs from hydrogen (cyclization with $R_{12}COOH$, wherein R_{12} differs from H) or $R_{12} = NH_2$ (cyclization with BrCN).

The aminoterminal protective group PG in the intermediates **G-2** may be a protective group known in the art such as *tert*-butyloxycarbonyl group. This protective group may conveniently be replaced by another suitable protective group such as phthalimido, dibenzyl or benzyloxycarbonyl.

5

Intermediate **G-2** can be deprotected using an acid such as trifluoroacetic acid in a suitable solvent such as dichloromethane to yield intermediate **H-2**.

The aminoterminal group of **H-2** may subsequently be substituted by procedures known in the art to generate the compounds of formula **I-2**.


10

Alternatively, intermediates may be deprotected with a strong acid such as hydrochloric acid in isopropanol, in a suitable solvent such as a mixture of ethanol and dioxane.

15 In a preferred embodiment the protecting group is selected from Fmoc, Acetyl, *tert*-butyloxycarbonyl, benzyloxycarbonyl-, dibenzyl-.

The procedure outlined in scheme 2 generates regioselective substitution of benzimidazol with the R_{12} and R_{13} substituents. If R_{13} is hydrogen, the intermediates and compounds may tautomerize

20

Scheme 3


Intermediate **D-1'**, obtained according to scheme 1, was reacted with an alkaline solution such as sodium hydroxide in methanol and water, to yield intermediate **E-3**, which was further acylated with acyl chlorides R_6COCl , in the presence of a base such as triethylamine, yielding intermediate **F-3**.

25

Intermediate F-3 was then deprotected and further substituted as described in scheme 1 to yield compounds of formula G-3.

The person skilled in the art will recognize that the intermediates in scheme 2 and compounds of formula G-3 may exist as tautomers.

Scheme 4



Reaction of 4-chloro-3-nitrobenzenesulfonic acid with chlorosulfonic acid led to intermediate A-4.

Intermediate D-4 was prepared by reacting an intermediate C-1, obtained according to the procedure described in patent WO97/18205 and also depicted in scheme 8, with an intermediate A-4, in a reaction-inert solvent such as dichloromethane, in the presence of a base such as triethylamine, at low temperature, for example 0°C.

The aminoterminal protective group in the intermediates may be a protective group known in the art such as *tert*-butoxycarbonyl group. This protective group may conveniently be replaced by another suitable protective group such as phthalimido, dibenzyl or benzyloxycarbonyl.

Nucleophilic substitution of the chlorine atom by a primary amine ($R_{13}NH_2$) under heating conditions led to intermediate E-4, whose nitro group was reduced to an amino group using a procedure known in the art such as catalytic hydrogenation or transfer hydrogenation, yielding intermediate F-4.


5

Cyclisation of F-4 with methylorthoformate, a carboxylic acid of formula $R_{12}COOH$ wherein R_{12} differs from hydrogen in an acidic solvent such as hydrochloric acid, or with $BrCN$ led to final compounds G-4.

- 10 Intermediate G-4 can then be deprotected and further substituted according to the procedures as described in scheme 1 to yield final compounds I-4.

This scheme results in regioselective substitution of the benzimidazol moiety with R_{12} and R_{13} respectively. If R_{13} is hydrogen, the intermediates and compounds of scheme 4 may exist as tautomers.

15



Intermediate D-4 was reacted with ammonia in a suitable solvent such as isopropanol, under heating conditions, to yield intermediate E-5. The amino group was then protected (PG') by e.g. a Boc-group, and the nitro group was reduced using conditions known in the art, for example catalytic hydrogenation, yielding compound F-5.

5

Reaction of compound F-5 with an acylchloride of formula RCOCl , in presence of a base such as triethylamine, in an organic solvent such as DCM, led to intermediate G-5, which was further reduced to compound H-5. Compound H-5 can also be obtained directly from F-5 by reductive amination with an aldehyde RCHO , and a reducing agent such as titanium IV isopropoxide, in an organic solvent such as DCM. R is aryl, Het¹, Het² or C₁₋₅alkyl optionally substituted with aryl, Het¹, Het², hydroxy, halogen, amino whereby the amino group may be optionally be mono- or di-substituted with C₁₋₄alkyl. The $-\text{CH}_2\text{-R}$ moiety of intermediates H-5 to K-5 taken together is identical to the definition of R₁₃.

15


After PG' deprotection with, for example, HCl in isopropanol, compound H-5 was cyclized as previously described with methylorthoformate or a carboxylic acid R₁₂COOH wherein R₁₂ differs from hydrogen in an acidic solvent such as hydrochloric acid, or with BrCN to yield compound I-5.

20

Intermediate I-5 can then be deprotected and further substituted according to the methods as described in scheme 1 to yield compounds of formula K-5.

In a preferred embodiment, the amino protecting group PG is dibenzyl, removed by catalytic hydrogenation, in the presence of palladium on charcoal, in an organic solvent such as methanol.

25


Scheme 6

Intermediate **G-6**, obtained as described in scheme 5 from intermediate **F-5** and chloroacetic chloride in the presence of a base such as triethylamine, was substituted by an amine of formula NHRR' to yield compound **H-6**. R and R' are each independently H or C_{1-4} alkyl. PG' preferably is boc.

Compound **H-6** was then reduced to intermediate **I-6**, further deprotected using, for example, HCl in isopropanol, and cyclized as previously described in scheme 2 and 4, to yield compound **J-6**.

Intermediate **J-6** can then be deprotected and further substituted as described in scheme 1 to yield compounds of formula **L-6**.


In an embodiment, the amino protecting group PG is dibenzyl, removed by catalytic hydrogenation, in the presence of palladium on charcoal, in an organic solvent such as methanol.

Scheme 7


Intermediate **H-5** was obtained as described in scheme 5. PG and PG' may be Boc. **H-5** was deprotected using an acid for example HCl in isopropanol, and coupled with a compound of formula R₁-L-(leaving group), in the presence of a base such as triethylamine, in a suitable organic solvent such as DCM, leading to intermediate **I-7**.

Cyclisation of intermediate **I-7** was achieved with thiocarbonylimidazole in a suitable organic solvent such as THF, leading to intermediate **J-7**, which was then reacted with an alkylhalide of formula R-X, wherein R is a C₁₋₆alkyl and X is an halide. An example thereof is methyl iodide, in the presence of a base such as potassium carbonate in DMF.

After oxidation of intermediate **K-7** with a suitable reagent such as mCPBA in DCM, the obtained methylsulfoxide group was substituted by an amine of formula HNR₅AR₆, in the presence of a base such as potassium carbonate in THF, under reflux conditions, yielding compound of formula **M-7**. If R₁₃ is hydrogen, the intermediates and compounds of scheme 7 may exist as tautomers.

Scheme 8


Intermediate **C-1** may be prepared by reacting compound **C-0** with an amine, in a suitable solvent such as isopropanol. It will be clear for the person skilled in the art that starting from a stereospecific intermediate of formula **C-0**, stereoselective intermediates of formula **C-1** will be obtained following reaction with a primary amine.

Example 1: Preparation of compound 8Intermediate **1-e**

25 g of 1H-benzimidazol-2-yl-carbamic acid, methyl ester **1-a** were dissolved in 45 mL of chlorosulfonic acid, at 0°C. The reaction mixture was then warmed up to 50°C and stirred during 4 hours. The reaction mixture was poured onto a mixture of ice and water and was stirred until a white precipitate was formed. The precipitate was next filtered and washed successively with an acidic aqueous solution (pH<2), a basic aqueous solution (pH>10) and a neutral aqueous solution (pH = 7 to 8), and dried in a vacuum oven, yielding 24 g (64%) of the desired intermediate **1-b**, methyl [5-(chlorosulfonyl)-1H-benzimidazol-2-yl]carbamate.


2 g of intermediate **1-c** [2R-hydroxy-3-[(2-methylpropyl)amino]-1S-(phenylmethyl)-propyl]carbamic acid, 1,1-dimethylethyl ester were dissolved in 50 mL of DCM. 1.8 g of triethylamine were then added to the reaction mixture, followed by portion-wise addition of 1.81 g of intermediate **1-b**. After 5 hours stirring at room temperature, the reaction mixture was washed with water, dried over MgSO₄, filtered and evaporated to yield 3 g (85%) of the desired intermediate **1-d**, [(1S,2R)-2-hydroxy-3-[[[2-[(methoxycarbonyl)amino]-1H-benzimidazol-5-yl]sulfonyl]-(2-methylpropyl)amino]-1-(phenylmethyl)propyl]carbamic acid, 1,1-dimethylethyl ester.

- 3 g of intermediate **1-d** were dissolved in a mixture of 8 mL of HCl in isopropanol and 40 mL of ethanol. The reaction mixture was stirred at room temperature overnight, then the solid was filtered off and was redissolved in a mixture DCM/saturated solution of NaHCO₃ in water. The organic layer was dried over MgSO₄ and evaporated to yield 1.4 g (60%) of the intermediate **1-e** [5-[[[(2*R*,3*S*)-3-amino-2-hydroxy-4-phenylbutyl]-(2-methylpropyl)amino]sulfonyl]-1*H*-benzimidazol-2-yl]carbamic acid, methyl ester as a free base.

**Compound 8**

- 1 g of intermediate **1-e** was dissolved in 20 mL of DCM. 0.37 g of 2,6-dimethylphenoxyacetic acid, 0.42 g of DCC and 0.27 g of HOBt were then added to the reaction mixture, which was stirred at room temperature during 4 hours. The reaction mixture was evaporated and the residue dissolved in 20 mL of ethylacetate. The reaction mixture was cooled down to 0°C and the precipitate of dicyclohexylurea was filtered off. The organic layer was then washed with a solution of Na₂CO₃ in water, brine, dried over MgSO₄ and evaporated. The crude compound was purified on silica gel eluting with 5% methanol in DCM, yielding 1.29 g (97%) of the desired final compound [(1*S*,2*R*)-2-hydroxy-3-[[[2-[(methoxycarbonyl)amino]-1*H*-benzimidazol-5-yl]sulfonyl]-(2-methylpropyl)amino]-1-(phenylmethyl)propyl]-(2,6-dimethylphenoxy) acetamide (compound 8).

Example 2: Preparation of Compound 7

**Compound 7**

- 58 g of 3-fluoroaniline and 58 mL of benzaldehyde were heated at 80°C during one hour. 200 mL of sulfuric acid were then added to the reaction mixture, cooled with an ice bath. After removing of the ice bath, the reaction mixture was further stirred at room temperature until complete dissolution of the solid. The reaction mixture was next cooled to 0°C with an ice bath and 36 mL of nitric acid in 120 mL of sulfuric acid were

added dropwise, maintaining the temperature at 0°C. After one hour of stirring at 0°C, the solid was filtered off and poured into a saturated solution of potassium carbonate in water. Ethyl acetate was then added and the two layers were separated. The aqueous phase was extracted two more times with ethyl acetate. The organic phases were
5 collected, dried over MgSO₄ and evaporated. The crude compound was purified on silica gel eluting with 40% ethyl acetate in hexane, yielding 28.65 g (35%) of the desired intermediate **A-2** 3-fluoro-4-nitroaniline.

28.65 g of intermediate **A-2** was dissolved in 230 mL of chlorhydric acid 36%. The
10 reaction mixture was cooled to 0°C with an ice bath and 13.7 g of sodium nitrite was added portion-wise. The reaction was maintained at 0°C during 1.5 hours, then mixed with 145 mL of a SO₂ saturated acetic acid solution, containing 10.5 mL of water and 9.3 g of CuCl₂·2H₂O. After complete addition, the cooling bath was removed and the reaction mixture was stirred at room temperature during one hour, then poured onto ice.
15 The solid was filtered off yielding 37.7 g of the intermediate **B-2** 3-fluoro-4-nitrobenzene sulfonyl chloride.

To a solution of 53 g of intermediate **C-1** (PG = Boc, R₄ = isobutyl) in 500 mL of THF containing 42 mL of triethylamine was added portion-wise 37.7 g of intermediate **B-2**.
20 The reaction mixture was stirred at room temperature overnight then evaporated. The residue was dissolved in ethylacetate and extracted with water, then with a solution of HCl 5% in water and with a K₂CO₃ solution in water. The organic layer was then dried over MgSO₄ and evaporated. The crude compound was purified on silica gel yielding 53 g (65%) of the desired intermediate **2-a** [(1S,2R)-3-[[[3-fluoro-4-nitrophenyl]-sulfonyl](2-methylpropyl)amino]-2-hydroxy-1-(phenylmethyl)propyl] carbamic acid,
25 1,1-dimethylethyl ester.

2 g of intermediate **2-a** were dissolved in 50 mL of DMF and 1.85 mL of isopropylamine were added. The reaction mixture was stirred at 60°C overnight, then
30 concentrated and the residue treated with a mixture of EtOAc and brine. The organic layer was then dried over MgSO₄ and evaporated to yield 2 g (91%) of the desired intermediate **2-b** [(1S,2R)-2-hydroxy-3-[(2-methylpropyl)[[3-(2-methylpropyl)amino-4-nitrophenyl]sulfonyl]amino]-1-(phenylmethyl)propyl]carbamic acid, 1,1-dimethylethyl ester, used without other purification in the next step.

35 2 g of intermediate **2-b** were dissolved in 40 mL of methanol, then 1.5 g of ammonium formate and 0.2 g of palladium on charcoal (10%) were added. The reaction mixture was stirred overnight at 60°C, then 0.5 g of ammonium formate and 0.2 g of palladium on charcoal were added. After three hours, the mixture was filtered on celite and

evaporated. The residue was dissolved in 50 mL of DCM, washed with a solution of Na_2CO_3 in water, then brine, dried over MgSO_4 and evaporated to yield 1.3 g (68%) of intermediate **2-c** [(1*S*,2*R*)-3-[[4-amino-3-[(2-methylpropyl)amino]phenyl]sulfonyl]-(2-methylpropyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]carbamic acid,


5 1,1-dimethylethyl ester, used without other purification in the next step.

1.3 g of intermediate **2-c** was dissolved in 20 mL of ethyl orthoformate. The reaction mixture was stirred at 80°C during 5 hours then concentrated. The residue was dissolved in ethyl acetate and washed with a solution of Na_2CO_3 in water. The organic
10 layer was dried over MgSO_4 and evaporated. The crude compound was purified on silica gel eluting with 0 to 2% methanol in DCM, yielding 0.8 g (70%) of the desired intermediate **2-d** [(1*S*,2*R*)-2-hydroxy-3-[(2-methylpropyl)[[1-(2-methylpropyl)-benzimidazol-6-yl]sulfonyl]amino]-1-(phenylmethyl)propyl] carbamic acid,
15 1,1-dimethylethyl ester.

2.6 g of intermediate **2-d** was dissolved in 100 mL of HCl 5N in isopropanol. The reaction mixture was stirred at room temperature during 2 hours, then concentrated to yield 2.5 g (94%) of the deprotected amine as an HCl salt, N-[(2*R*,3*S*)-3-amino-2-hydroxy-4-phenylbutyl]-N-(2-methylpropyl)[1-(2-methylpropyl) benzimidazol-6-yl]-
20 sulfonamide, hydrochloride (**2-e**).

2.5 g of **2-e** and 1.5 mL of triethylamine were dissolved in 60 mL of DCM. 3.05 g of 1-[(3*R*,3*aS*,6*aR*)-hexahydrofuro[2,3-*b*]furan-3-yl]oxycarbonyloxy]-2,5-pyrrolidinedione were then added and the reaction mixture was stirred at room
25 temperature during 4 hours. The reaction mixture was then washed with a solution of Na_2CO_3 in water, then brine, dried over MgSO_4 and evaporated. The crude compound was purified on silica gel eluting with 5% methanol in DCM, yielding 1.8 g (60%) of the desired final compound [(1*S*,2*R*)-2-hydroxy-3-[(2-methylpropyl)[[1-(2-methylpropyl)benzimidazol-6-yl]sulfonyl]amino]-1-(phenylmethyl)propyl]carbamic acid,
30 [(3*R*,3*aS*,6*aR*)-hexahydrofuro[2,3-*b*]furan-3-yl]ester (compound 7).

Example 3: Preparation of Compound 1



Compound 1


54 mg of intermediate **D-2** (PG = Boc) were dissolved in 20 mL of THF and 11 mg of benzylamine were added. The reaction mixture was stirred at reflux during 4 hours, then concentrated. The crude compound was purified on silica gel eluting with 10% ethylacetate in hexane, yielding 60 mg of intermediate **3-a** [(1S,2R)-2-hydroxy-3-[(2-methylpropyl)][4-nitro-3-[(phenylmethyl)amino]phenyl]sulfonyl]amino]-1-(phenylmethyl)propyl]carbamic acid, 1,1-dimethylethyl ester.

60 mg of intermediate **3-a** was dissolved in 20 mL of methanol. The reaction mixture was then hydrogenated in the presence of 25 mg of palladium on charcoal (10%), at room temperature, overnight. After filtration of the catalyst, the reaction mixture was concentrated to give the intermediate **3-b** [(1S,2R)-3-[[[4-amino-3-[(phenylmethyl)amino]phenyl]sulfonyl](2-methylpropyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]carbamic acid, 1,1-dimethylethyl ester which was used in the next step without further purification.

59 mg of intermediate **3-b** was dissolved in 20 mL of methylorthoformate. The reaction mixture was stirred at 80°C during 5 hours then concentrated. The residue was dissolved in ethyl acetate and washed with a solution of Na₂CO₃ in water. The organic layer was dried over MgSO₄ and evaporated to give the intermediate **3-c** [(1S,2R)-2-hydroxy-3-[(2-methylpropyl)[1-(phenylmethyl)benzimidazol-6-yl]sulfonyl]amino]-1-(phenylmethyl)propyl] carbamic acid, 1,1-dimethylethyl ester which was used in the next step without further purification.

500 mg of intermediate **3-c** was reacted with 15 mL of HCl 5N in isopropanol at room temperature during 2 hours. The reaction mixture was then concentrated to give the intermediate **3-d** as an HCl salt N-[(2R,3S)-3-amino-2-hydroxy-4-phenylbutyl]-N-(2-methylpropyl)[1-(phenylmethyl)benzimidazol-6-yl]sulfonamide, hydrochloride which was used in the next step without further purification.


500 mg of the previous intermediate **3-d** and 85 mg of triethylamine were dissolved in 10 mL of DCM. 271 mg of 1-[(3R,3aS,6aR)-hexahydrofuro[2,3-b]furan-3-yl]oxy-carbonyloxy]-2,5-pyrrolidinedione were then added and the reaction mixture was stirred at room temperature during 4 hours. The reaction mixture was evaporated and the crude compound was purified on silica gel eluting with 5% methanol in DCM, yielding 236 mg (35%) of the desired final compound [(1S,2R)-2-hydroxy-3-[(2-methylpropyl)[1-(phenylmethyl)benzimidazol-6-yl]sulfonyl]amino]-1-(phenylmethyl)propyl]carbamic acid, [(3R,3aS,6aR)-hexahydrofuro[2,3-b]furan-3-yl]ester (compound 1).

Example 4: Preparation of Compound 2**Compound 2**

59 mg of intermediate **F-2** (PG = Boc, R₁₃ = benzyl) and 10 mg of acetic acid were stirred in 5 ml of dioxane/5 ml of 5N HCl for the weekend at 110°C. After cooling, EtOAc was added and the reaction was poured onto a mixture of ice, saturated NaHCO₃ solution and EtOAc. After extraction with EtOAc, the organic layer was dried and evaporated under reduced pressure to give the intermediate **4-a** [(1*S*,2*R*)-2-hydroxy-3-[[[2-methyl-1-(phenylmethyl)]benzimidazol-6-yl]sulfonyl](2-methylpropyl)amino]-1-(phenylmethyl)propyl]carbamic acid, 1,1-dimethylethyl ester which was used in the next step without further purification.

500 mg of intermediate **4-a** was reacted with 15 mL of HCl 5N in isopropanol at room temperature during 2 hours. The reaction mixture was then concentrated to give intermediate **4-b** as an HCl salt N-[(2*R*,3*S*)-3-amino-2-hydroxy-4-phenylbutyl]-N-(2-methylpropyl)[[2-methyl-1-(phenylmethyl)]benzimidazol-6-yl]sulfonamide, hydrochloride which was used in the next step without further purification.

510 mg of intermediate **4-b**, 271 mg of 1-[(3*R*,3*aS*,6*aR*)-hexahydrofuro[2,3-*b*]furan-3-yl]oxycarbonyloxy]-2,5-pyrrolidinedione and 85 mg of triethylamine, were mixed in 30 mL of DCM. The reaction mixture was stirred during 2 hours at room temperature, washed with water then evaporated. The crude compound was purified on silica gel eluting with 5% methanol in DCM, yielding 165 mg (25%) of the final compound [(1*S*,2*R*)-2-hydroxy-3-[[[2-methyl-1-(phenylmethyl)]benzimidazol-6-yl]sulfonyl]-(2-methylpropyl)amino]-1-(phenylmethyl)propyl]carbamic acid, [(3*R*,3*aS*,6*aR*)-hexahydrofuro[2,3-*b*]furan-3-yl]ester (Compound 2).

Example 5: Preparation of Compound 9**Compound 9**

20 g of intermediate **C-1** (PG = dibenzyl, R₄ = isobutyl) were dissolved in 250 mL of DCM. 23 mL of triethylamine were then added to the reaction mixture, followed by 12.7 g of intermediate **1-b**, added portion-wise. After overnight stirring at room temperature, the reaction mixture was washed with water, dried over MgSO₄, filtered and evaporated to yield 11 g (40%) of the desired intermediate **5-a**, [5-[[[(2R,3S)-3-(dibenzylamino)-2-hydroxy-4-phenylbutyl](2-methylpropyl)amino]sulfonyl]-1H-benzimidazol-2-yl]carbamic acid, methyl ester.

11 g of intermediate **5-a** were dissolved in 150 mL of dioxane. Then 18 g of Na₂CO₃ dissolved in 150 mL of water were added to the reaction mixture, which was stirred at 90°C during 24 hours. The reaction mixture was then concentrated and the residue extracted with DCM, then ethylacetate. The residue was then dried by concentration with toluene, then mixed with isopropanol and filtered. The crystals were further dried in a vacuum oven at 50°C, to give 7.76 g (79%) of intermediate **5-b** N-[(2R,3S)-3-(dibenzylamino)-2-hydroxy-4-phenylbutyl]-N-(2-methylpropyl)-[2-aminobenzimidazol-5-yl]sulfonamide.


3 g of intermediate **5-b** and 0.76 mL triethylamine were mixed in 150 mL of THF, at 0°C. Then 0.39 g of acetylchloride dissolved in THF were added to the reaction mixture, which was stirred during 3 hours at room temperature. 0.7 g of 4-dimethylaminopyridine were added and the reaction stirred overnight. 0.39 g of acetylchloride and 0.7 g of 4-dimethylaminopyridine were then added to the reaction mixture which was further stirred until all intermediate **5-b** has reacted. The reaction mixture was then washed with water and extracted with DCM. The organic layer was dried over MgSO₄, filtered and evaporated to yield 1.1 g (34%) of the desired intermediate **5-c** N-[(2R,3S)-3-(dibenzylamino)-2-hydroxy-4-phenylbutyl]-N-(2-methylpropyl)[2-(acetylamino)-benzimidazol-5-yl]sulfonamide.

1.1 g of intermediate **5-c** was dissolved in 200 mL of methanol. Then the reaction mixture was hydrogenated in the presence of 0.5 g of palladium on charcoal (10%), at room temperature, overnight. After filtration of the catalyst, the reaction mixture was hydrogenated again in the presence of newly added palladium on charcoal (10%). This step was again repeated twice then the reaction mixture was concentrated and the crude compound was purified on silica gel eluting with 2% methanol in DCM, yielding 0.2 g (25%) of the desired intermediate N-[(2R,3S)-3-amino-2-hydroxy-4-phenylbutyl]-N-(2-methylpropyl)[2-(acetylamino)benzimidazol-5-yl]sulfonamide (**5-d**).

200 mg of intermediate **5-d** were dissolved in 2 mL of DCM, then 25 mg of 2,6-dimethylphenoxyacetic acid, 27 mg of EDCI and 19 mg of HOBt were added to the

reaction mixture, which was stirred at room temperature overnight then evaporated. The crude compound was purified by preparative HPLC, yielding 27 mg (30%) of the final compound [(1*S*,2*R*)-3-[[[2-(acetylamino)benzimidazol-5-yl]sulfonyl](2-methylpropyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-(2,6-dimethyl phenoxy)acetamide (compound 9).

Example 6: Preparation of Compound 64



Compound 64

- 10 A mixture of 34.20 g of intermediate **C-1** (PG = dibenzyl, R₄ = isobutyl) and 12.48 g of triethylamine in 250 ml of dichloromethane was stirred at 0°C. Then 25.16 g of intermediate **A-4** was added and the reaction was stirred at room temperature. After 1 hour an extra 3 g of intermediate **A-4** was added and the reaction was stirred during weekend. After washing with 5% HCl solution, saturated NaHCO₃ solution and brine,
- 15 the organic layer was separated, dried and evaporated to give the intermediate **6-a** N-[(2*R*,3*S*)-3-(dibenzylamino)-2-hydroxy-4-phenylbutyl]-N-(2-methylpropyl) (4-chloro-3-nitrophenyl)sulfonamide which was used in the next step without further purification.
- 20 A mixture of 11.75 g of intermediate **6-a** and 12 ml of methylamine (40 wt% in H₂O) in 100 ml of methanol was stirred at room temperature for 24 hours. After evaporation under reduced pressure, the product was washed with H₂O and extracted with EtOAc. The organic layer was dried and evaporated under reduced pressure to give the intermediate **6-b** N-[(2*R*,3*S*)-3-(dibenzylamino)-2-hydroxy-4-phenylbutyl]-N-
- 25 (2-methylpropyl)(4-methylamino-3-nitrophenyl)sulfonamide which was used in the next step without further purification.

- A mixture of 11.5 g of intermediate **6-b**, 2 ml of thiophene in diisopropylether (4%) and 1g of palladium on charcoal (10%) in 100 ml of methanol was hydrogenated. After
- 30 filtering over decalite, the filtrate was evaporated under reduced pressure to give the intermediate **6-c** N-[(2*R*,3*S*)-3-(dibenzylamino)-2-hydroxy-4-phenylbutyl]-N-(2-methylpropyl)(3-amino-4-methylaminophenyl)sulfonamide which was used in the next step without further purification.


A mixture of 2 g of intermediate **6-c** and 0.23 g of formic acid was stirred in 100 ml of dioxane/100 ml of 5N HCl for 48 hours at 110°C. After cooling, EtOAc was added and the reaction was poured onto a mixture of ice, saturated NaHCO₃ solution and EtOAc. After extraction with EtOAc, the organic layer was dried and evaporated under reduced pressure. The crude product was purified on silica gel eluting with 1% of methanol in dichloromethane to give 2.23 g of intermediate **6-d** N-[(2*R*,3*S*)-3-(dibenzylamino)-2-hydroxy-4-phenylbutyl]-N-(2-methylpropyl)(1,2-dimethyl benzimidazol-5-yl)-sulfonamide.

- 10 A mixture of 2.23 g of intermediate **6-d**, 1.38 g of ammonium formate and 1.2 g of palladium on charcoal (10%) in 50 ml of ethanol was stirred at 80°C for 1.5 hours. After filtering over decalite, the filtrate was evaporated under reduced pressure. The residue was purified by preparative HPLC to give 0.29 g of intermediate **6-e** N-[(2*R*,3*S*)-3-amino-2-hydroxy-4-phenylbutyl]-N-(2-methylpropyl)(1-methyl benzimidazol-5-yl)sulfonamide, with a yield of 18%.

A mixture of 0.35 g of intermediate **6-e**, 0.22 g of 1-[(3*R*,3*aS*,6*aR*)-hexahydrofuro-[2,3-*b*]furan-3-yl]oxycarbonyloxy-2,5-pyrrolidinedione and 0.12 g of triethylamine in 20 ml of dichloromethane was stirred at room temperature for 48 hours. After washing with saturated NaHCO₃ solution, the organic layer was dried and evaporated under reduced pressure. The residue was purified by preparative HPLC to give 0.105 g of the final compound [(1*S*,2*R*)-3-[[[(1-methylbenzimidazol-5-yl)sulfonyl](2-methylpropyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]carbamic acid, [(3*R*,3*aS*,6*aR*)-hexahydrofuro[2,3-*b*]furan-3-yl]ester, with a yield of 22% (compound 64).

25

Example 7: Preparation of Compound 24



Compound 24

- 20 g of intermediate **D-4** (PG = dibenzyl, R₄ = isobutyl) in 400 ml of ammonia in isopropanol was stirred at 120°C overnight. After evaporation under reduced pressure, the product was washed with H₂O and extracted with dichloromethane. The organic layer was dried and evaporated under reduced pressure to give the intermediate **7-a** N-[(2*R*,3*S*)-3-(dibenzylamino)-2-hydroxy-4-phenylbutyl]-N-(2-methylpropyl)

30

(4-amino-3-nitrophenyl)sulfonamide which was used in the next step without further purification.

5 A mixture of 20 g of intermediate 7-a, 3 ml of thiophene in diisopropylether (4%) and 2 g of palladium on charcoal (10%) in 150 ml of methanol was hydrogenated. After filtering over decalite, the filtrate was evaporated under reduced pressure to give the intermediate 7-b N-[(2*R*,3*S*)-3-(dibenzylamino)-2-hydroxy-4-phenylbutyl]-N-(2-methylpropyl)(3,4-diaminophenyl)sulfonamide which was used in the next step without further purification.

10

A mixture of 1 g of intermediate 7-b and 0.13 g of acetic acid was stirred in 40 ml of dioxane/40 ml of 5N HCl for 48 hours at 110°C. After cooling, EtOAc was added and the reaction was poured onto a mixture of ice, saturated NaHCO₃ solution and EtOAc. After extraction with EtOAc, the organic layer was dried and evaporated under reduced pressure to give the intermediate 7-c N-[(2*R*,3*S*)-3-(dibenzylamino)-2-hydroxy-4-phenylbutyl]-N-(2-methylpropyl)(2-methylbenzimidazol-5-yl)sulfonamide which was used in the next step without further purification.

15

A mixture of 1.55 g of intermediate 7-c and 0.4 g of palladium on charcoal (10%) in 20 ml of methanol was hydrogenated. After filtering over decalite, the filtrate was evaporated under reduced pressure. The crude compound was purified on silica gel eluting with 4% of methanol in dichloromethane to give 0.18 g of intermediate 7-d N-[(2*R*,3*S*)-3-amino-2-hydroxy-4-phenylbutyl]-N-(2-methylpropyl)(2-methylbenzimidazol-5-yl)sulfonamide.


20

A mixture of 0.18 g of intermediate 7-d, 0.11 g of 1-[[[(3*R*,3*aS*,6*aR*)-hexahydrofuro[2,3-*b*]furan-3-yl]oxycarbonyloxy]-2,5-pyrrolidinedione and 46 mg of triethylamine in 20 ml of dichloromethane was stirred at room temperature for 17 hours. After washing with saturated NaHCO₃ solution, the organic layer was dried and evaporated under reduced pressure. The residue was purified on silica gel eluting with 2% of methanol in dichloromethane to give 0.22 g of the final compound [(1*S*,2*R*)-2-hydroxy-3-[[[(2-methylbenzimidazol-5-yl)sulfonyl](2-methylpropyl) amino]-1-(phenylmethyl)propyl]-carbamic acid, [(3*R*,3*aS*,6*aR*)-hexahydrofuro[2,3-*b*]furan-3-yl]ester with a yield of 89% (compound 24).

25


30

35

Example 8: Preparation of compound 20**Intermediate F-5**

37 g of intermediate **F-5** (PG = dibenzyl, PG' = boc, R₄ = isobutyl) and 6.5 g of triethylamine were mixed in 100 mL of DCM. 26.2 g of Boc₂O were then added at 0°C, then the reaction mixture was heated at 50°C during 2 days and evaporated to give 49 g of the desired intermediate **8-a** [4-[[[(2R,3S)-3-(dibenzylamino)-2-hydroxy-4-phenylbutyl](2-methylpropyl)amino]sulfonyl]-2-nitrophenyl]carbamic acid, 1,1-dimethylethyl ester, used without further purification in the next step.

42 g of **8-a** were dissolved in 500 mL of methanol and hydrogenated in the presence of palladium on charcoal (10%) and thiophene. The reaction mixture was then evaporated and purified on silica gel eluting with DCM, yielding 30 g (75%) of intermediate **8-b** [2-amino-4-[[[(2R,3S)-3-(dibenzylamino)-2-hydroxy-4-phenylbutyl](2-methylpropyl)-amino]sulfonyl]phenyl]carbamic acid, 1,1-dimethylethyl ester.

**Compound 20**

1.44 g of intermediate **8-b**, 284 mg of chloroacetyl chloride and 262 mg of triethylamine were mixed in 20 mL of DCM at 0°C. The reaction mixture was then stirred at room temperature overnight, then washed with water, dried over MgSO₄ and evaporated to yield 1.5 g of intermediate **8-c** [2-(chloroacetyl-amino)-4-[[[(2R,3S)-3-(dibenzylamino)-2-hydroxy-4-phenylbutyl](2-methylpropyl)amino] sulfonyl]phenyl] carbamic acid, 1,1-dimethylethyl ester, used without further purification in the next step.

800 mg of **8-c** and 298 mg of pyrrolidine were mixed in 10 mL of THF. The reaction mixture was stirred at room temperature overnight and evaporated. The residue was treated with a mixture of water and DCM. The organic phase was dried over MgSO₄ and evaporated to yield 920 mg of intermediate **8-d** [4-[[[(2R,3S)-3-(dibenzylamino)-2-

hydroxy-4-phenylbutyl](2-methylpropyl)amino] sulfonyl]-2-[[pyrrolidin-1-yl)acetyl]-amino]phenyl]carbamic acid, 1,1-dimethylethyl ester, used without further purification in the next step.

- 5 920 mg of the previous intermediate **8-d** and 1.3 g of acid trifluoroacetic were mixed in 10 mL of DCM. The reaction mixture was stirred at room temperature overnight, then a saturated solution of NaHCO₃ in water was added. The organic layer was separated and evaporated to yield 790 mg of intermediate **8-e** [2-amino-5-[[[(2*R*,3*S*)-3-(dibenzyl-amino)-2-hydroxy-4-phenylbutyl] (2-methylpropyl)amino]sulfonyl]phenyl]-(pyrrolidin-10 1-yl)acetamide, used without further purification in the next step.

- 790 mg of the previous intermediate **8-e** was dissolved in 10 mL of THF. 129 mg of LiAlH₄ was then added and the reaction mixture was stirred at reflux during 3 hours. 2 mL of water and 2 mL of a 20% solution of NaOH in water were then added to the
15 reaction mixture, which was filtered over decalite and evaporated to yield 780 mg of intermediate **8-f** N-[(2*R*,3*S*)-3-(dibenzylamino)-2-hydroxy-4-phenylbutyl]-N-(2-methylpropyl)[4-amino-3-[2-(pyrrolidin-1-yl)ethylamino]phenyl]sulfonamide, used without further purification in the next step.

- 20 780 mg of the previous intermediate **8-f** and 30 mL of HCl 5N in isopropanol were mixed in 30 mL of dioxane. 2 mL of formic acid were then added and the reaction mixture stirred at 100°C during 4 hours. After cooling to room temperature, EtOAc was added and, while maintaining a vigorous stirring, solid K₂CO₃ was added to neutralize the reaction mixture. The organic layer was then separated and evaporated to yield 850
25 mg of intermediate **8-g** N-[(2*R*,3*S*)-3-(dibenzylamino)-2-hydroxy-4-phenylbutyl]-N-(2-methylpropyl)[1-[2-(pyrrolidin-1-yl) ethyl]benzimidazol-6-yl]sulfonamide, used without further purification in the next step.


- 850 mg of the previous intermediate **8-g**, ammonium formate and palladium on
30 charcoal (10%) were mixed in 10 mL of ethanol. The reaction mixture was heated at 80°C and stirred during 2 hours, then filtered on decalite and purified on silica gel, yielding 320 mg of the deprotected compound **8-h** N-[(2*R*,3*S*)-3-amino-2-hydroxy-4-phenylbutyl]-N-(2-methylpropyl)[1-[2-(pyrrolidin-1-yl)ethyl]benzimidazol-6-yl]-sulfonamide.

- 35 320 mg of the previous intermediate **8-h**, 169 mg of 1-[(3*R*,3*aS*,6*aR*)-hexahydrofuro-[2,3-*b*]furan-3-yl]oxycarbonyloxy]-2,5-pyrrolidinedione and 70 mg of triethylamine were mixed in 10 mL of DCM. The reaction mixture was stirred at room temperature for 4 hours, evaporated and purified on silica gel eluting with 8% ammonia in DCM,

yielding 112 mg (27%) of the desired final compound [(1*S*,2*R*)-2-hydroxy-1-(phenylmethyl)-3-[[[1-[2-(pyrrolidin-1-yl)ethyl]benzimidazol-6-yl]sulfonyl](2-methylpropyl)-amino]propyl]carbamic acid, [(3*R*,3*aS*,6*aR*)-hexahydro furo[2,3-*b*]furan-3-yl]ester (compound 25).

5

Example 9: Preparation of compound 40



To 1.5 g of intermediate **F-5** (PG, PG' = Boc, R₄ = isobutyl) dissolved in 25 mL of DCM, were added 321 μ L of phenylacetaldehyde and 1.06 g of sodium triacetoxyborohydride. The mixture was stirred at room temperature overnight, then washed with a saturated solution of Na₂CO₃ in water, then brine, dried over MgSO₄ and evaporated to give 2.1 g (100%) of intermediate **H-5'** (= **H-5** wherein PG, PG' = boc, R₄ = isobutyl, R₁₃ is phenylethyl) as an orange oil.

2.1 g of intermediate **H-5'** were dissolved in 50 mL of dioxane and reacted with 100 mL of HCl 7N in isopropanol, at room temperature, for 16 hours. The reaction mixture was then evaporated and treated with a mixture of DCM and a saturated solution of Na₂CO₃ in water, dried over MgSO₄ and evaporated to give 1.4 g of crude intermediate which was used directly in the next step.

20

1.4 g of the previous intermediate, 677 mg of 1-[(3*R*,3*aS*,6*aR*)-hexahydrofuro[2,3-*b*]furan-3-yl]oxycarbonyloxy]-2,5-pyrrolidinedione and 250 mg of triethylamine were mixed in 25 mL of DCM. The reaction mixture was stirred at room temperature for 2 hours, evaporated and purified on silica gel eluting with 5% methanol in DCM, yielding 900 mg (56%) of the desired compound **I-7'** (= **I-7** wherein R₁₃ is phenylethyl, R_{1-L} is 1-[(3*R*,3*aS*,6*aR*)-hexahydrofuro[2,3-*b*]furan-3-yl]oxycarbonyl, R₄ is isobutyl).

25

800 mg of intermediate **I-7'** and 428 mg of thiocarbonyl diimidazole were dissolved in 10 mL of THF. After stirring at room temperature for 16 hours, 215 mg of thiocarbonyl diimidazole were added to the mixture. After 16 hours, the reaction mixture was evaporated and purified on silica gel eluting with 5% methanol in DCM, yielding 350 mg (41%) of the desired compound **J-7'** (= **J-7** wherein wherein R₁₃ is phenylethyl, R_{1-L} is 1-[(3*R*,3*aS*,6*aR*)-hexahydrofuro[2,3-*b*]furan-3-yl]oxycarbonyl, R₄ is isobutyl).

30

To 350 mg of intermediate **J-7'** were added 31 μ L of MeI and 69 mg of K_2CO_3 in 5 mL of DMF. The mixture was stirred at room temperature for 1 hour, then evaporated, dissolved in DCM and washed with brine, dried over $MgSO_4$ and evaporated to yield
 5 360 mg (100%) of intermediate **K-7'** (= **K-7** wherein R_{13} is phenylethyl, R_{1-L} is 1-[[[(3*R*,3*aS*,6*aR*)-hexahydrofuro[2,3-*b*]furan-3-yl]oxycarbonyl], R_4 is isobutyl, R is methyl) which was used without further purification in the next step.

360 mg of intermediate **K-7'** and 122 mg of mCPBA were dissolved in 5 mL of DCM
 10 and stirred at room temperature for 2 hours. 100 mg of mCPBA were then added to the mixture which was stirred for another hour, then washed with a saturated solution of Na_2CO_3 in water, then brine, dried over $MgSO_4$ and purified on silica gel eluting with 1% methanol in DCM, yielding 140 mg (38%) of the desired intermediate **L-7'** (= **L-7** wherein wherein R_{13} is phenylethyl, R_{1-L} is 1-[[[(3*R*,3*aS*,6*aR*)-hexahydrofuro[2,3-*b*]-
 15 furan-3-yl]oxycarbonyl], R_4 is isobutyl).

To 140 mg of intermediate **L-7'** dissolved in 10 mL of acetonitrile were added 1 mL of 1-(2-Aminoethyl)pyrrolidine. The reaction mixture was stirred 48 hours under reflux, then evaporated and purified by preparative HPLC, yielding 60 mg (40%) of the
 20 desired final compound 40 [(1*S*,2*R*)-2-hydroxy-3-[[[1-(phenethyl)-2-[2-(pyrrolidin-1-yl)ethylamino]]benzimidazol-6-yl]sulfonyl](2-methylpropyl)amino]-1-(phenylmethyl)propyl]carbamic acid, [(3*R*,3*aS*,6*aR*)-hexahydrofuro[2,3-*b*]furan-3-yl]ester. Mass spectral data: m/z = 789 ($M+H$).

25 The compounds of formula (I) may also be converted to the corresponding *N*-oxide forms following art-known procedures for converting a trivalent nitrogen into its *N*-oxide form. Said *N*-oxidation reaction may generally be carried out by reacting the starting material of formula (I) with an appropriate organic or inorganic peroxide. Appropriate inorganic peroxides comprise, for example, hydrogen peroxide, alkali
 30 metal or earth alkaline metal peroxides, e.g. sodium peroxide, potassium peroxide; appropriate organic peroxides may comprise peroxy acids such as, for example, benzenecarboperoxoic acid or halo substituted benzenecarboperoxoic acid, e.g. 3-chloro-benzenecarboperoxoic acid, peroxyalkanoic acids, e.g. peroxyacetic acid, alkylhydroperoxides, e.g. *tert*-butyl hydroperoxide. Suitable solvents are, for example,
 35 water, lower alkanols, e.g. ethanol and the like, hydrocarbons, e.g. toluene, ketones, e.g. 2-butanone, halogenated hydrocarbons, e.g. dichloromethane, and mixtures of such solvents.

An interesting group of intermediates are those intermediates wherein $-A-R_6$ is

hydrogen. Said intermediates may also have pharmacological properties similar to those pharmacological properties of the compounds of formula (I).

5 The present compounds can thus be used in animals, preferably in mammals, and in particular in humans as pharmaceuticals per se, in mixtures with one another or in the form of pharmaceutical preparations.

10 Furthermore, the present invention relates to pharmaceutical preparations which as active constituents contain an effective dose of at least one of the compounds of formula (I) in addition to customary pharmaceutically innocuous excipients and auxiliaries. The pharmaceutical preparations normally contain 0.1 to 90% by weight of a compound of formula (I). The pharmaceutical preparations can be prepared in a manner known per se to one of skill in the art. For this purpose, at least one of a compound of formula (I), together with one or more solid or liquid pharmaceutical
15 excipients and/or auxiliaries and, if desired, in combination with other pharmaceutical active compounds, are brought into a suitable administration form or dosage form which can then be used as a pharmaceutical in human medicine or veterinary medicine.

20 Pharmaceuticals which contain a compound according to the invention can be administered orally using e.g. including suspensions, capsules, tablets, sachets, solutions, suspensions, emulsions; parenterally, using e.g. subcutaneous, intravenous, intramuscular, intrasternal injection or infusion techniques; rectally using e.g. suppositories; intravaginally; by inhalation, or topically, the preferred administration being dependent on the individual case, e.g., the particular course of the disorder to be
25 treated. Oral administration is preferred.

30 The person skilled in the art is familiar on the basis of his expert knowledge with the auxiliaries which are suitable for the desired pharmaceutical formulation. Beside solvents, gel-forming agents, suppository bases, tablet auxiliaries and other active compound carriers, antioxidants, dispersants, emulsifiers, antifoams, flavor corrigents, preservatives, solubilizers, agents for achieving a depot effect, buffer substances or colorants are also useful.

35 Due to their favorable pharmacological properties, particularly their activity against multi-drug resistant HIV protease enzymes, the compounds of the present invention are useful in the treatment of individuals infected by HIV and for the prophylaxis of these individuals.

The prophylaxis treatment can be advantageous in cases where an individual has been subjected to a high risk of exposure to a virus, as can occur when individual has been in

contact with an infected individual where there is a high risk of viral transmission. As an example, prophylactic administration of said compounds would be advantageous in a situation where a health care worker has been exposed to blood from an HIV-infected individual, or in other situations where an individual engaged in high-risk activities that potentially expose that individual to the HIV virus.

In general, the compounds of the present invention may be useful in the treatment of warm-blooded animals infected with viruses whose existence is mediated by, or depends upon, the protease enzyme. Conditions which may be prevented or treated with the compounds of the present invention include, but is not limited to, treating a wide range of states of HIV infection: AIDS, ARC (Aids related complex), both symptomatic and asymptomatic, and actual or potential exposure to HIV. The compounds of the present are also useful for treating progressive generalized lymphadenopathy, Kaposi's syndrome, thrombocytopenia purpurea, AIDS-related neurological conditions such as AIDS dementia complex, multiple sclerosis, tropical paresis, and also anti-HIV antibody positive and HIV-positive conditions, including such conditions in asymptomatic patients. For example, the compounds of this invention are useful in treating infection by HIV after suspected past exposure to HIV by e.g., blood transfusion, exchange of body fluids, bites, accidental needle stick, or exposure to patient blood during surgery. The term prevention includes prophylaxis of HIV infection and prophylaxis of the evolution of HIV infection to AIDS.

The compounds of the present invention or any subgroup thereof may therefore be used as medicines against above-mentioned conditions. Said use as a medicine or method of treatment comprises the systemic administration to HIV-infected subjects of an amount effective to combat the conditions associated with HIV and other pathogenic retroviruses, especially HIV-1. Consequently, the compounds of the present invention can be used in the manufacture of a medicament useful for treating conditions associated with HIV and other pathogenic retroviruses, in particular medicaments useful for treating patients infected with multi-drug resistant HIV virus.

In a preferred embodiment, the invention relates to the use of a compound of formula (I) or any subgroup thereof in the manufacture of a medicament for treating or combating infection or disease associated with multi-drug resistant retrovirus infection in a mammal, in particular HIV-1 infection. Thus, the invention also relates to a method of treating a retroviral infection, or a disease associated with multi-drug resistant retrovirus infection comprising administering to a mammal in need thereof an effective amount of a compound of formula (I) or a subgroup thereof.

In another preferred embodiment, the present invention relates to the use of formula (I) or any subgroup thereof in the manufacture of a medicament for inhibiting a protease of a multi-drug resistant retrovirus in a mammal infected with said retrovirus, in particular HIV-1 retrovirus.

5

In another preferred embodiment, the present invention relates to the use of formula (I) or any subgroup thereof in the manufacture of a medicament for inhibiting multi-drug resistant retroviral replication, in particular HIV-1 replication.

- 10 An advantage of the compounds of the present invention is that the compounds display activity towards HIV strains carrying mutations in the protease gene. It is known in the art that mutants of the HIV protease confer resistance to HIV protease inhibitors. Examples of such mutations comprise those mutations, independently selected from the list comprising mutations at amino acid positions 3, 10, 11, 13, 15, 19, 20, 22, 24, 30,
15 32, 33, 35, 36, 37, 41, 43, 46, 47, 48, 50, 53, 54, 55, 57, 58, 62, 63, 66, 70, 71, 72, 73, 77, 82, 84, 85, 88, 89 or 90 in the HIV protease. The compounds of the present invention may be useful to prevent or delay the onset of mutations in HIV protease, or if the HIV protease contains mutations at the initiation of therapy may prevent or delay the occurrence of additional mutations in the HIV protease.

20

The compounds of the present invention may also find use in inhibiting ex vivo samples containing HIV or expected to be exposed to HIV. Hence, the present compounds may be used to inhibit HIV present in a body fluid sample which contains or is suspected to contain or be exposed to HIV.

25

- Also, the combination of an antiretroviral compound and a compound of the present invention can be used as a medicine. Thus, the present invention also relates to a product containing (a) a compound of the present invention, and (b) another antiretroviral compound, as a combined preparation for simultaneous, separate or
30 sequential use in treatment of retroviral infections, in particular, in the treatment of infections with multi-drug resistant retroviruses. Thus, to combat or treat HIV infections, or the infection and disease associated with HIV infections, such as Acquired Immunodeficiency Syndrome (AIDS) or AIDS Related Complex (ARC), the compounds of this invention may be co-administered in combination with for instance,
35 binding inhibitors, such as, for example, dextran sulfate, suramine, polyanions, soluble CD4, PRO-542, BMS-806; fusion inhibitors, such as, for example, T20, T1249, RPR 103611, YK-FH312, IC 9564, 5-helix, D-peptide ADS-J1; co-receptor binding inhibitors, such as, for example, AMD 3100, AMD-3465, AMD7049, AMD3451 (Bicyclams), TAK220, TAK 779, T-22, ALX40-4C; SHC-C (SCH351125), SHC-D,

PRO-140, RPR103611, AK-602; RT inhibitors, such as, for example, foscarnet and prodrugs; nucleoside RTIs, such as, for example, AZT, 3TC, DDC, DDI, D4T, Abacavir, FTC, DAPD (Amdoxovir), dOTC (BCH-10652), fozivudine, DPC 817; nucleotide RTIs, such as, for example, PMEA, PMPA (tenofovir); NNRTIs, such as, for example, nevirapine, delavirdine, efavirenz, 8 and 9-Cl TIBO (tivrapipe), loviride, TMC-125, dapivirine, MKC-442, UC 781, UC 782, Capravirine, QM96521, GW420867X, GW-3011, GW-4511, GW-4751, DPC 961, DPC963, DPC082, DPC083, TMC-125, calanolide A, SJ-3366, TSAO, 4"-deaminated TSAO, MV150, MV026048, PNU-142721; RNase H inhibitors, such as, for example, SP1093V, PD126338; TAT inhibitors, such as, for example, RO-5-3335, K12, K37; integrase inhibitors, such as, for example, L 708906, L 731988, S-1360; protease inhibitors, such as, for example, amprenavir and prodrug GW908, ritonavir, nelfinavir, saquinavir, indinavir, lopinavir, palinavir, BMS 186316, atazanavir, DPC 681, DPC 684, tipranavir, AG1776, mozenavir, DMP-323, GS3333, KNI-413, KNI-272, L754394, L756425, LG-71350, PD161374, PD173606, PD177298, PD178390, PD178392, PNU 140135, TMC-114, maslinic acid, U-140690, RO-033-4649; glycosylation inhibitors, such as, for example, castanospermine, deoxynojirimycine; entry inhibitors CGP64222.

The compounds of the present invention may also be administered in combination with immunomodulators e.g., broprimine, anti-human alpha interferon antibody, IL-2, methionine enkephalin, interferon alpha, and naltrexone; with antibiotics e.g., pentamidine isothiorate; cytokines (e.g. Th2); modulators of cytokines; chemokines or the receptors thereof (e.g. CCR5); or hormones (e.g. growth hormone) or the receptors thereof; to ameliorate, combat, or eliminate HIV infection and its symptoms. Such combination therapy in different formulations, may be administered simultaneously, sequentially or independently of each other. Alternatively, such combination may be administered as a single formulation, whereby the active ingredients are released from the formulation simultaneously or separately.

The compounds of the present invention may also be administered in combination with modulators of the metabolism following application of the drug to an individual. These modulators include compounds that interfere with the metabolism at cytochromes, such as cytochrome P450. Some modulators inhibit cytochrome P450. It is known that several isoenzymes exist of cytochrome P450, one of which is cytochrome P450 3A4. Ritonavir is an example of a modulator of metabolism via cytochrome P450. Interesting compounds having an effect at cytochrome P450 include those compounds containing a thiazolyl, imidazolyl or pyridinyl moiety. Such combination therapy in different formulations, may be administered simultaneously, separately or

sequentially. Alternatively, such combination may be administered as a single formulation, whereby the active ingredients are released from the formulation simultaneously or separately.

- Such modulator may be administered at the same or different ratio as the compound of the present invention. Preferably, the weight ratio of such modulator vis-à-vis the compound of the present invention (modulator:compound of the present invention) is 1:1 or lower, more preferable the ratio is 1:3 or lower, suitably the ratio is 1:10 or lower, more suitably the ratio is 1:30 or lower.
- 10 The combination may provide a synergistic effect, whereby viral infectivity and its associated symptoms may be prevented, substantially reduced, or eliminated completely. Combinations of the compounds of formula (I) with another HIV protease inhibitor as cytochrome P₄₅₀ inhibitor can act synergistically, in an additive way or antagonistically. This can be assessed in an experimental setting where the potency of
- 15 different ratios of the two HIV-protease inhibitors is measured. Results can be plotted in an isobologram graph according to the method described by Chou and Talalay (Adv. Enzyme Regul. 22: 27-55, 1984). Synergism between two inhibitors would mean a more potent combination therapy, but with no increase in undesired side effects.
- 20 For an oral administration form, compounds of the present invention are mixed with suitable additives, such as excipients, stabilizers or inert diluents, and brought by means of the customary methods into the suitable administration forms, such as tablets, coated tablets, hard capsules, aqueous, alcoholic, or oily solutions. Examples of suitable inert carriers are gum arabic, magnesia, magnesium carbonate, potassium phosphate, lactose,
- 25 glucose, or starch, in particular, corn starch. In this case the preparation can be carried out both as dry and as moist granules. Suitable oily excipients or solvents are vegetable or animal oils, such as sunflower oil or cod liver oil. Suitable solvents for aqueous or alcoholic solutions are water, ethanol, sugar solutions, or mixtures thereof. Polyethylene glycols and polypropylene glycols are also useful as further auxiliaries for
- 30 other administration forms.
- For subcutaneous or intravenous administration, the active compounds, if desired with the substances customary therefor such as solubilizers, emulsifiers or further auxiliaries, are brought into solution, suspension, or emulsion. The compounds of
- 35 formula (I) can also be lyophilized and the lyophilizates obtained used, for example, for the production of injection or infusion preparations. Suitable solvents are, for example, water, physiological saline solution or alcohols, e.g. ethanol, propanol, glycerol, in addition also sugar solutions such as glucose or mannitol solutions, or alternatively

mixtures of the various solvents mentioned.

Suitable pharmaceutical formulations for administration in the form of aerosols or sprays are, for example, solutions, suspensions or emulsions of the compounds of formula (I) or their physiologically tolerable salts in a pharmaceutically acceptable solvent, such as ethanol or water, or a mixture of such solvents. If required, the formulation can also additionally contain other pharmaceutical auxiliaries such as surfactants, emulsifiers and stabilizers as well as a propellant. Such a preparation customarily contains the active compound in a concentration from approximately 0.1 to 50%, in particular from approximately 0.3 to 3% by weight.

In order to enhance the solubility and/or the stability of the compounds of formula (I) in pharmaceutical compositions, it can be advantageous to employ α -, β - or γ -cyclodextrins or their derivatives. Also co-solvents such as alcohols may improve the solubility and/or the stability of the compounds of formula (I) in pharmaceutical compositions. In the preparation of aqueous compositions, addition salts of the subject compounds are obviously more suitable due to their increased water solubility.

Appropriate cyclodextrins are α -, β - or γ -cyclodextrins (CDs) or ethers and mixed ethers thereof wherein one or more of the hydroxy groups of the anhydroglucose units of the cyclodextrin are substituted with C_{1-6} alkyl, particularly methyl, ethyl or isopropyl, e.g. randomly methylated β -CD; hydroxy C_{1-6} alkyl, particularly hydroxyethyl, hydroxypropyl or hydroxybutyl; carboxy C_{1-6} alkyl, particularly carboxymethyl or carboxyethyl; C_{1-6} alkyl-carbonyl, particularly acetyl; C_{1-6} alkyloxycarbonyl C_{1-6} alkyl or carboxy C_{1-6} alkyloxy C_{1-6} alkyl, particularly carboxymethoxypropyl or carboxyethoxypropyl; C_{1-6} alkylcarbonyloxy C_{1-6} alkyl, particularly 2-acetyloxypropyl. Especially noteworthy as complexants and/or solubilizers are β -CD, randomly methylated β -CD, 2,6-dimethyl- β -CD, 2-hydroxyethyl- β -CD, 2-hydroxyethyl- γ -CD, 2-hydroxypropyl- γ -CD and (2-carboxymethoxy)propyl- β -CD, and in particular 2-hydroxypropyl- β -CD (2-HP- β -CD).

The term mixed ether denotes cyclodextrin derivatives wherein at least two cyclodextrin hydroxy groups are etherified with different groups such as, for example, hydroxypropyl and hydroxyethyl.

An interesting way of formulating the present compounds in combination with a cyclodextrin or a derivative thereof has been described in EP-A-721,331. Although the formulations described therein are with antifungal active ingredients, they are equally interesting for formulating the compounds of the present invention. The formulations

described therein are particularly suitable for oral administration and comprise an antifungal as active ingredient, a sufficient amount of a cyclodextrin or a derivative thereof as a solubilizer, an aqueous acidic medium as bulk liquid carrier and an alcoholic co-solvent that greatly simplifies the preparation of the composition. Said
5 formulations may also be rendered more palatable by adding pharmaceutically acceptable sweeteners and/or flavors.

Other convenient ways to enhance the solubility of the compounds of the present invention in pharmaceutical compositions are described in W0-94/05263,
10 W098/42318, EP-A-499,299 and WO 97/44014, all incorporated herein by reference.

More in particular, the present compounds may be formulated in a pharmaceutical composition comprising a therapeutically effective amount of particles consisting of a solid dispersion comprising (a) a compound of formula (I), and (b) one or more
15 pharmaceutically acceptable water-soluble polymers.

The term "a solid dispersion" defines a system in a solid state (as opposed to a liquid or gaseous state) comprising at least two components, wherein one component is dispersed more or less evenly throughout the other component or components. When
20 said dispersion of the components is such that the system is chemically and physically uniform or homogenous throughout or consists of one phase as defined in thermodynamics, such a solid dispersion is referred to as "a solid solution". Solid solutions are preferred physical systems because the components therein are usually readily bioavailable to the organisms to which they are administered.

25 The term "a solid dispersion" also comprises dispersions which are less homogenous throughout than solid solutions. Such dispersions are not chemically and physically uniform throughout or comprise more than one phase.

30 The water-soluble polymer in the particles is conveniently a polymer that has an apparent viscosity of 1 to 100 mPa.s when dissolved in a 2 % aqueous solution at 20°C solution.

Preferred water-soluble polymers are hydroxypropyl methylcelluloses or HPMC.
35 HPMC having a methoxy degree of substitution from about 0.8 to about 2.5 and a hydroxypropyl molar substitution from about 0.05 to about 3.0 are generally water soluble. Methoxy degree of substitution refers to the average number of methyl ether groups present per anhydroglucose unit of the cellulose molecule. Hydroxypropyl molar substitution refers to the average number of moles of propylene oxide which

have reacted with each anhydroglucose unit of the cellulose molecule.

The particles as defined hereinabove can be prepared by first preparing a solid dispersion of the components, and then optionally grinding or milling that dispersion.

- 5 Various techniques exist for preparing solid dispersions including melt-extrusion, spray-drying and solution-evaporation, melt-extrusion being preferred.

- 10 It may further be convenient to formulate the present compounds in the form of nanoparticles which have a surface modifier adsorbed on the surface thereof in an amount sufficient to maintain an effective average particle size of less than 1000 nm. Useful surface modifiers are believed to include those which physically adhere to the surface of the antiretroviral agent but do not chemically bond to the antiretroviral agent.

- 15 Suitable surface modifiers can preferably be selected from known organic and inorganic pharmaceutical excipients. Such excipients include various polymers, low molecular weight oligomers, natural products and surfactants. Preferred surface modifiers include nonionic and anionic surfactants.

- 20 Yet another interesting way of formulating the present compounds involves a pharmaceutical composition whereby the present compounds are incorporated in hydrophilic polymers and applying this mixture as a coat film over many small beads, thus yielding a composition with good bioavailability which can conveniently be manufactured and which is suitable for preparing pharmaceutical dosage forms for oral administration.

- 25 Said beads comprise (a) a central, rounded or spherical core, (b) a coating film of a hydrophilic polymer and an antiretroviral agent and (c) a seal-coating polymer layer.

- 30 Materials suitable for use as cores in the beads are manifold, provided that said materials are pharmaceutically acceptable and have appropriate dimensions and firmness. Examples of such materials are polymers, inorganic substances, organic substances, and saccharides and derivatives thereof.

- 35 Another aspect of the present invention concerns a kit or container comprising a compound of formula (I) in an amount effective for use as a standard or reagent in a test or assay for determining the ability of a potential pharmaceutical to inhibit HIV protease, HIV growth, or both. This aspect of the invention may find its use in pharmaceutical research programs.


The compounds of the present invention can be used in phenotypic resistance

monitoring assays, such as known recombinant assays, in the clinical management of resistance developing diseases such as HIV. A particularly useful resistance monitoring system is a recombinant assay known as the AntivirogramTM. The AntivirogramTM is a highly automated, high throughput, second generation,
 5 recombinant assay that can measure susceptibility, especially viral susceptibility, to the compounds of the present invention. (Hertogs K, de Bethune MP, Miller V *et al.* *Antimicrob Agents Chemother*, 1998; 42(2):269-276, incorporated by reference).

The dose of the present compounds or of the physiologically tolerable salt(s) thereof to
 10 be administered depends on the individual case and, as customary, is to be adapted to the conditions of the individual case for an optimum effect. Thus it depends, of course, on the frequency of administration and on the potency and duration of action of the compounds employed in each case for therapy or prophylaxis, but also on the nature and severity of the infection and symptoms, and on the sex, age, weight and individual
 15 responsiveness of the human or animal to be treated and on whether the therapy is acute or prophylactic. Customarily, the daily dose of a compound of formula (I) in the case of administration to a patient approximately 75 kg in weight is 0.1 mg to 10 g, preferably 1 mg to 1g, more preferably 3 mg to 0.5 g. The dose can be administered in the form of an individual dose, or divided into several, e.g. two, three, or four,
 20 individual doses.


















The following tables list the compounds of formula (I) which were prepared following one of the above reaction schemes. ND means not determined.

Table 1



25
















<u>No</u>	<u>R</u>	<u>R12</u>	<u>R13</u>	<u>pIC₅₀</u> <u>(LAD)</u>
1		H		9.1
2		-CH ₃		9.0

No	R	R12	R13	pIC ₅₀ (LAD)
3		-H		8.7
4		-H		8.7
5		-H		8.7
6		-NHCOOCH ₃	-H	8.6
7		-H		8.6
8		-NHCOOCH ₃	-H	8.5
9		-NHCOCH ₃	-H	8.4
10		-NHCOOCH ₃	-H	8.0
11		-NHCOOCH ₃	-H	8.0
12		-NHCOOCH ₃	-H	7.8
13		-NH ₂	-CH ₃	7.7
14		-H		7.7

WO 03/076413

PCT/EP03/50057

-51-




















No	R	R12	R13	IC ₅₀ (LAI)
15		-H		7.7
16		-H	-CH ₃	7.7
17		-NHCOOCH ₃	-H	7.6
18		-H	-CH ₃	ND
19		-NHCOOCH ₃	-H	7.5
20		-H		7.4
21		-H		7.4
22		-NHCOOCH ₃	-H	7.4
23		-H	-H	7.3
24		-CH ₃	-H	7.3
25		-NHCOOCH ₃	-H	7.1
26		-NHCOOCH ₃	-H	7.0

No	R	R12	R13	pIC ₅₀ (LAD)
27		-NHCOCH ₃	-H	7.3
28		-H		7.5
29		-H	-H	7.1
30		-H		7.1
31		-H	-CH ₃	7.2
32		-H		7.4
33		-H	-CH ₃	7.0
34		-H		7.0
35		-H	-H	7.0
36		-NH ₂	-H	6.5
37			-H	6.9

WO 03/076413

PCT/EP03/50057

-53-

No	R	R12	R13	pIC ₅₀ (LAD)
38		-NHCOCH ₃	-H	6.7
39		-NH ₂		6.9
40				6.4
41		-NH ₂	-CH ₃	6.6
42		-H	-H	7.1
43		-H		6.4
44		-H		6.1
45		-H		6.3
46		-H	-H	6.4
47		-H		6.4
48			-H	6.4

















No	R	R12	R13	pIC ₅₀ (LAD)
49		-H	-H	6.3
50		-H	-H	6.9
51		-H		6.2
52			-H	6.1
53		-NH ₂	-H	6.0
54		-NH ₂	-H	5.8
55			-H	5.8
56		-CH ₂ OH	-H	5.7
57		-NH ₂	-H	5.7
58			-H	5.6
59		-NH ₂	-H	5.3

Table 2

No	R	R12	R13	pIC ₅₀ (LAI)
60		-H		6.5
61		-H		5.7
62		-H		5.7
63		-H		ND
64		-H	-CH ₃	7.6

Antiviral analyses:

The compounds of the present invention were examined for anti-viral activity in a cellular assay. The assay demonstrated that these compounds exhibited potent anti-HIV activity against a wild type laboratory HIV strain (HIV-1 strain LAI). The cellular assay was performed according to the following procedure.

HIV- or mock-infected MT4 cells were incubated for five days in the presence of various concentrations of the inhibitor. At the end of the incubation period, all HIV-infected cells have been killed by the replicating virus in the control cultures in the absence of any inhibitor. Cell viability is measured by measuring the concentration of MTT, a yellow, water soluble tetrazolium dye that is converted to a purple, water insoluble formazan in the mitochondria of living cells only. Upon solubilization of the resulting formazan crystals with isopropanol, the absorbance of the solution is

monitored at 540nm. The values correlate directly to the number of living cells remaining in the culture at the completion of the five day incubation. The inhibitory activity of the compound was monitored on the virus-infected cells and was expressed as IC₅₀ and IC₉₀. These values represent the amount of the compound required to protect 50% and 90%, respectively, of the cells from the cytopathogenic effect of the virus. The toxicity of the compound was measured on the mock-infected cells and was expressed as CC₅₀, which represents the concentration of compound required to inhibit the growth of the cells by 50%. The selectivity index (SI) (ratio CC₅₀/IC₅₀) is an indication of the selectivity of the anti-HIV activity of the inhibitor. Wherever results are reported as e.g. pIC₅₀ or pCC₅₀ values, the result is expressed as the negative logarithm of the result expressed as IC₅₀ or CC₅₀ respectively.

Antiviral spectrum:

Because of the increasing emergence of drug resistant HIV strains, the present compounds were tested for their potency against clinically isolated HIV strains harboring several mutations. These mutations are associated with resistance to protease inhibitors and result in viruses that show various degrees of phenotypic cross-resistance to the currently commercially available drugs such as for instance saquinavir, ritonavir, nelfinavir, indinavir and amprenavir. The viral strains R13025, R13027, R13028, R13029, R13034, R13080, T13127 and R13363 contained mutations as indicated below in Table 3.

Table 3 List of mutations present in the protease gene of the HIV strains (A to H) used.

A	V003I, L010I, V032T, L033M, E035D, S037Y, S037D, M046I, R057R/K, Q058E, L063P, K070T, A071V, I072V, I084V, L089V
B	V003I, L010I, K020R, E035D, M036I, S037D, Q058E, I062V, L063P, A071V, I072M, G073S, V077I, I084V, I085V, L090M
C	V003I, L010I, I015V, L019I, K020M, S037N, R041K, I054V, Q058E, L063P, A071V, I084V, L090M, I093L
D	V003I, L010L/I, I013V, L033I, E035D, M036I, M046L, K055R, R057K, L063P, I066F, A071V, I084V, N088D, L090M
E	V003I, L010I, V011I, A022V, L024I, E035D, M036I, S037T, R041K, I054V, I062V, L063P, A071V, I084V
F	L010F, M046I, M071V, I084V
G	V003I, L010I, V032T, L033M, E035D, S037Y, M046I, I047V, R057R/K, Q058E, L063P, K070T, A071V, I072V, V082I, I084V, L089V
H	V003I, V032I, L035D, M036I, S037N, K043T, M046I, I047V, I050V, K055R, I057K, I062V, L063P, A071L, V082I, I085V, L090M, I093L

Results:

As a measure of the broad spectrum activity of the present compounds, the fold resistance (FR), defined as $FR = IC_{50}(\text{mutant strain})/IC_{50}(\text{HIV-1 strain LAI})$, was determined. Table 4 shows the results of the antiviral testing in terms of fold resistance. As can be seen in this table, the present compounds are effective in inhibiting a broad range of mutant strains: Column A FR value towards mutant A; Column B: FR towards mutant B; Column C: FR towards mutant C; Column D: FR towards mutant D; Column E: FR towards mutant E; Column F: FR towards mutant F; Column G: FR towards mutant G; Column H: FR towards mutant H. The toxicity (column Tox) is expressed as the pCC_{50} value as determined with mock transfected cells. Column WT displays the pIC_{50} value against wild type HIV-LAI strain.

Table 4. Results of the toxicity testing and the resistance testing against strain A to H (expressed as FR). ND indicates not determined

N°	A	B	C	D	E	F	G	H	Tox	WT
1	9.3	3.1	1.1	1.5	1.6	40	275	33	4.2	9.1
2	32	6.3	1.9	6.3	2.9	162	ND	89	4.2	9.0
3	15	2.7	1.0	1.2	0.9	83	112	20	<4	8.7
4	18	3.4	2.0	3.0	2.2	48	224	ND	4.2	8.7
5	ND	1.9	0.9	1.9	3.2	78	110	49	4.1	8.7
6	2.9	2.0	2.7	2.1	1.6	10	60	ND	<4	8.6
7	ND	ND	0.45	1.2	2.1	72	174	ND	<4	8.6
8	2.1	2.0	1.4	1.9	2.3	16.2	89	ND	<4	8.5
9	4.3	1.3	4.4	2.4	2.8	14	107	ND	<4	8.4
11	5.4	1.0	1.5	3.5	1.6	7.1	245	ND	<4	8.0
12	5.5	2.1	9.5	7.9	9.5	49	89	ND	<4	7.8
13	ND	3.4	2.5	3.4	4.5	ND	ND	ND	<4	7.7
15	ND	6.3	2.4	5.4	6.2	269	407	ND	4.2	7.7
16	ND	ND	2.6	5.9	5.4	288	ND	ND	<4.49	7.7
17	0.69	1.1	1.0	ND	0.57	1.3	5.0	ND	4.1	7.6
19	5.6	1.2	1.4	3.6	3.5	83	ND	ND	<4	7.5
21	14	1.1	0.83	1.1	1.1	25	83	ND	<4	7.4
22	0.74	0.78	1.1	0.71	0.72	0.69	4.3	ND	4.1	7.4
23	7.1	2.3	3.5	2.5	4.8	23	ND	ND	<4	7.3
24	2.6	0.85	1.1	0.89	0.83	4.7	35	ND	<4	7.3
25	1.2	ND	ND	ND	1.3	ND	ND	4.6	4.3	7.1

WO 03/076413

PCT/EP03/50057

-58-

Nº	A	B	C	D	E	F	G	H	Tox	WT
26	6.3	ND	ND	ND	6.8	ND	ND	47	4.86	7.0
27	2.7	ND	ND	ND	1.1	ND	ND	2.3	<4	7.3
28	1.5	ND	ND	ND	1.3	ND	ND	2.6	ND	7.5
29	14	ND	ND	ND	13	ND	ND	78	4.21	7.1
30	18	ND	ND	ND	13	ND	ND	65	<4	7.1
31	13	ND	ND	ND	9.5	ND	ND	209	4.24	7.2
32	18	ND	ND	ND	13	ND	ND	41	4.2	7.4
33	60	ND	ND	ND	19	ND	ND	234	<4	7.7
34	32	ND	ND	ND	17	ND	ND	46	<4	7.0
35	91	ND	ND	ND	17	ND	ND	148	<4	7.0
36	4.4	ND	ND	ND	3.1	ND	ND	17	4.19	6.5
37	1.0	0.35	0.95	1.9	1.3	7.9	21	ND	ND	6.9
38	0.12	0.32	0.39	0.19	0.17	0.21	0.60	ND	<4	6.7
39	32	ND	ND	ND	3.5	ND	ND	ND	4.3	6.9
40	10	ND	ND	ND	4.0	ND	ND	7.9	ND	6.4
41	1.4	ND	ND	ND	1.4	ND	ND	1.5	<4	6.6
42	ND	ND	ND	ND	8.7	ND	ND	52	<4	7.1
43	22	ND	ND	ND	12.6	ND	ND	28	4.99	6.4
44	0.29	ND	ND	ND	0.32	ND	ND	ND	<4	6.1
45	9.3	ND	ND	ND	7.4	ND	ND	ND	<4	6.3
46	ND	ND	ND	ND	ND	ND	ND	ND	<4	6.4
47	14	ND	ND	ND	7.8	ND	ND	66	<4	6.4
48	0.37	1.9	0.38	0.35	0.32	0.35	0.85	ND	<4	6.4
49	12	ND	ND	ND	1.0	ND	ND	3.5	<4	6.3
50	1.4	ND	ND	ND	0.83	ND	ND	2.8	<4	6.9
51	2.8	ND	ND	ND	0.89	ND	ND	2.1	ND	6.2
52	4.2	2.9	3.5	3.0	3.4	4.3	ND	ND	<5	6.1
53	ND	ND	ND	ND	ND	ND	ND	ND	4.05	6.0
54	1.6	0.40	0.42	0.46	0.42	1.7	ND	ND	<4	5.8
56	0.28	0.66	0.43	0.18	0.18	0.34	0.83	ND	<4	5.7
57	0.37	0.30	0.31	0.34	0.32	0.30	2.0	ND	<4	5.7
58	1.0	1.0	1.0	0.32	0.37	1.1	1.1	ND	<4	5.6
59	1.5	ND	ND	ND	0.31	ND	ND	0.39	<4	5.3
64	4.3	ND	ND	ND	1.5	ND	ND	13	<4	7.6

Bioavailability:**Caco-2 permeability assay for intestinal absorption**

- The permeability of different compounds is evaluated according to a Caco-2 test protocol as described by Augustijns et al. (Augustijns et al. (1998). *Int. J. of Pharm*, 166, 45-54) whereby, Caco-2 cells at cell passage number between 32 and 45 are grown in 24-well transwell cell culture plates for 21 to 25 days. The integrity of the cell monolayer is checked by measuring the transepithelial electrical resistance (TEER). The test is performed at pH 7.4 and at 100 μ M donor compound concentration.

Aqueous solubility at different pH levels

- The equilibrium solubility in simulated gastrointestinal solutions under thermodynamic conditions is a good measure for the solubility profile of the compound in the stomach and the different parts of the intestine. Simulated gastric fluid (SGF) (without pepsin) is set at pH of 1.5. Simulated intestinal fluids (SIF) (without bile salts) are set at pH 5, pH 6.5, pH 7 and pH 7.5. The experimental protocol uses 96-well flat-bottom microplates in which 1 mg of compound is added per well (stock solution in methanol) and evaporated to dryness. The compounds are resolubilized in SGF and SIF and incubated overnight on a horizontal shaking device at 37°C. After filtration, the compound concentrations are determined by UV-spectrophotometry.

Oral availability in the rat

- The compounds are formulated as a 20 mg/ml solution or suspension in DMSO, PEG400 or cyclodextrin 40% in water. For most experiments in the rat (male and female rats), three dosing groups are formed: 1/ single intraperitoneal (IP) dose at 20 mg/kg using DMSO formulation; 2/ single oral dose at 20 mg/kg using PEG400 formulation and 3/ single oral dose at 20 mg/kg using PEG400 formulation. Blood is sampled at regular time intervals after dosing and drug concentrations in the serum are determined using a LC-MS bioanalytical method. Serum concentrations are expressed in ng/mg. Serum concentration at 30 minutes (30') and at 3 hours (180') can be determined as these values reflect the extent of absorption (30') and the speed of elimination (180'). The rat serum concentration at 30 min and 180 min following IP administration of 20 mg/kg of compound 18 are 2012 ng/ml and 190 ng/ml respectively. The rat serum concentration at 30 min and 180 min following oral administration of 20 mg/kg as a DMSO formulation of compound 18 are 148 ng/ml and 44 ng/ml respectively; whereas in PEG400 said concentrations are respectively 475 ng/ml and 63 ng/ml.

Boosting the systemic bioavailability

- With the described type of compounds (protease-inhibitors), it is known that inhibition of the metabolic degradation processes can markedly increase the systemic availability

- by reducing the first-pass metabolism in the liver and the metabolic clearance from the plasma. This 'boosting' principle can be applied in a clinical setting to the pharmacological action of the drug. This principle can be also explored both in the rat or the dog by simultaneous administration of a compound that inhibits the Cyt-p450
- 5 metabolic enzymes. Known blockers are for example ritonavir and ketoconazole. Dosing a single oral dose of ritonavir at 5 mg/kg in the rat and the dog may result in an increase of the systemic availability.

Protein Binding analyses:

- Human serum proteins like albumin (HSA) or alpha-1 acid glycoprotein (AAG) are
- 10 known to bind many drugs, resulting in a possible decrease in the effectiveness of those compounds. In order to determine whether the present compounds would be adversely effected by this binding, the anti-HIV activity of the compounds was measured in the presence of human serum, thus evaluating the effect of the binding of the protease inhibitors to those proteins.

15 **Film-coated Tablets**

Preparation of Tablet Core


- A mixture of 100 g of active ingredient, *in casu* a compound of formula (I), 570 g lactose and 200 g starch was mixed well and thereafter humidified with a solution of 5 g sodium dodecyl sulfate and 10 g polyvinylpyrrolidone in about 200 ml of water. The
- 20 wet powder mixture was sieved, dried and sieved again. Then there was added 100 g microcrystalline cellulose and 15 g hydrogenated vegetable oil. The whole was mixed well and compressed into tablets, giving 10.000 tablets, each comprising 10 mg of the active ingredient.

Coating

- 25 To a solution of 10 g methylcellulose in 75 ml of denaturated ethanol there was added a solution of 5 g of ethylcellulose in 150 ml of dichloromethane. Then there were added 75 ml of dichloromethane and 2.5 ml 1,2,3-propanetriol. 10 g of polyethylene glycol was molten and dissolved in 75 ml of dichloromethane. The latter solution was added to the former and then there were added 2.5 g of magnesium octadecanoate, 5 g of
- 30 polyvinylpyrrolidone and 30 ml of concentrated color suspension and the whole was homogenated. The tablet cores were coated with the thus obtained mixture in a coating apparatus.

CLAIMS

1. A compound having the formula




or *N*-oxide, salt, stereoisomeric form, racemic mixture, prodrug or ester

5 thereof, wherein

R₁ is hydrogen, C₁₋₆alkyl, C₂₋₆alkenyl, arylC₁₋₆alkyl, arylC₂₋₆alkenyl, C₃₋₇cycloalkyl, C₃₋₇cycloalkylC₁₋₆alkyl, aryl, Het¹, Het¹C₁₋₆alkyl, Het², Het²C₁₋₆alkyl;

R₁ may also be a radical of formula



10 wherein

R₉, R_{10a} and R_{10b} are, each independently, hydrogen, C₁₋₄alkyloxycarbonyl, carboxyl, aminocarbonyl, mono- or di(C₁₋₄alkyl)aminocarbonyl, C₃₋₇cycloalkyl, C₂₋₆alkenyl, C₂₋₆alkynyl or C₁₋₄alkyl optionally substituted with aryl, Het¹, Het², C₃₋₇cycloalkyl, C₁₋₄alkyloxycarbonyl, carboxyl, aminocarbonyl, mono- or di(C₁₋₄alkyl)aminocarbonyl, aminosulfonyl, C₁₋₄alkylS(O)_n, hydroxy, cyano, halogen or optionally mono- or disubstituted amino where the substituents are selected from C₁₋₄alkyl, aryl, arylC₁₋₄alkyl, C₃₋₇cycloalkyl, C₃₋₇cycloalkylC₁₋₄alkyl, Het¹, Het², Het¹C₁₋₄alkyl and Het²C₁₋₄alkyl; whereby R₉, R_{10a} and the carbon atoms to which they are attached optionally form a C₃₋₇cycloalkyl radical;

15

R_{11a} is hydrogen, C₂₋₆alkenyl, C₂₋₆alkynyl, C₃₋₇cycloalkyl, aryl, optionally mono- or disubstituted aminocarbonyl, optionally mono- or disubstituted aminoC₁₋₄alkylcarbonyloxy, C₁₋₄alkyloxycarbonyl, aryloxycarbonyl, Het¹oxycarbonyl, Het²oxycarbonyl, aryloxycarbonylC₁₋₄alkyl, aryl-C₁₋₄alkyloxycarbonyl, C₁₋₄alkylcarbonyl, C₃₋₇cycloalkylcarbonyl, C₃₋₇cycloalkylC₁₋₄alkyloxycarbonyl, C₃₋₇cycloalkylcarbonyloxy, carboxylC₁₋₄alkylcarbonyloxy, C₁₋₄alkylcarbonyloxy, arylC₁₋₄alkylcarbonyloxy, aryloxycarbonyloxy, Het¹carbonyl, Het¹carbonyloxy, Het¹C₁₋₄alkyloxycarbonyl, Het²carbonyloxy, Het²C₁₋₄alkylcarbonyloxy, Het²C₁₋₄alkyloxycarbonyloxy or C₁₋₄alkyl

25

30

- optionally substituted with aryl, aryloxy, Het² or hydroxy; wherein the optional substituents on the amino groups are each independently selected from C₁₋₄alkyl, aryl, arylC₁₋₄alkyl, C₃₋₇cycloalkyl, C₃₋₇cycloalkylC₁₋₄alkyl, Het¹, Het², Het¹C₁₋₄alkyl and Het²C₁₋₄alkyl; whereby R_{11a} may be
- 5 optionally linked to the remainder of the molecule via a sulfonyl group;
- R_{11b} is hydrogen, C₃₋₇cycloalkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, aryl, Het¹, Het² or C₁₋₄alkyl optionally substituted with halogen, hydroxy, C₁₋₄alkylS(=O)_t, aryl, C₃₋₇cycloalkyl, Het¹, Het², optionally mono- or disubstituted amino where the substituents are selected from C₁₋₄alkyl, aryl, arylC₁₋₄alkyl,
- 10 C₃₋₇cycloalkyl, C₃₋₇cycloalkylC₁₋₄alkyl, Het¹, Het², Het¹C₁₋₄alkyl and Het²C₁₋₄alkyl;
- Each t independently is zero, 1 or 2;
- R₂ is hydrogen or C₁₋₆alkyl;
- L is -C(=O)-, -O-C(=O)-, -NR₈-C(=O)-, -O-C₁₋₆alkanediyl-C(=O)-,
- 15 -NR₈-C₁₋₆alkanediyl-C(=O)-, —S(=O)₂-, -O-S(=O)₂-, —NR₈-S(=O)₂-, -NR₈-C₁₋₆alkanediyl-S(=O)₂, whereby either the C(=O) group or the S(=O)₂ group is attached to the NR₂ moiety; whereby the C₁₋₆alkanediyl moiety is optionally substituted with hydroxy, aryl, Het¹ or Het²;
- R₃ is C₁₋₆alkyl, aryl, C₃₋₇cycloalkyl, C₃₋₇cycloalkylC₁₋₄alkyl, or arylC₁₋₄alkyl optionally
- 20 substituted with C₁₋₄alkyl, halogen, nitro, cyano, polyhaloC₁₋₄alkyl, hydroxy, C₁₋₄alkyloxy, mercapto or C₁₋₄alkylthio;
- R₄ is hydrogen, C₁₋₄alkyloxycarbonyl, carboxyl, aminocarbonyl, mono- or di(C₁₋₄alkyl)aminocarbonyl, C₃₋₇cycloalkyl, C₂₋₆alkenyl, C₂₋₆alkynyl or C₁₋₄alkyl optionally substituted with aryl, Het¹, Het², C₃₋₇cycloalkyl, C₁₋₄alkyloxycarbonyl,
- 25 carboxyl, aminocarbonyl, mono- or di(C₁₋₄alkyl)aminocarbonyl, aminosulfonyl, C₁₋₄alkylS(=O)_t, hydroxy, cyano, halogen or optionally mono- or disubstituted amino where the substituents are selected from C₁₋₄alkyl, aryl, arylC₁₋₄alkyl, C₃₋₇cycloalkyl, C₃₋₇cycloalkylC₁₋₄alkyl, Het¹, Het², Het¹C₁₋₄alkyl and Het²C₁₋₄alkyl; or
- 30 R₄ may also be -X-R₇ wherein
- X is -O- or -N(R₇)-Y-;
- Y is -C(=O)-, -S(=O)₂-, -O-C(=O)-, -NR₈-C(=O)-, -C(=O)-C(=O)-, -O-S(=O)₂- or —NR₈-S(=O)₂-;
- R₅ is hydrogen, hydroxy, C₁₋₆alkyl, Het¹C₁₋₆alkyl, Het²C₁₋₆alkyl, aminoC₁₋₆alkyl
- 35 whereby the amino group may optionally be mono- or di-substituted with C₁₋₄alkyl;

- R_6 is hydrogen, C_{1-6} alkyloxy, Het^1 , Het^1oxy , Het^2 , Het^2oxy , aryl, aryloxy, aryloxy- C_{1-4} alkyl, C_{1-4} alkyloxyaryl, C_{1-4} alkyloxy Het^1 , C_{1-4} alkyloxy Het^2 , C_{1-4} alkyloxy-carbonylamino, amino C_{1-4} alkylamino, amino or amino C_{1-4} alkyloxy and in case A is other than C_{1-6} alkanediyl, then R_6 may also be C_{1-6} alkyl, Het^1C_{1-4} alkyl, Het^1oxyC_{1-4} alkyl, Het^2C_{1-4} alkyl, Het^2oxyC_{1-4} alkyl, aryl C_{1-4} alkyl, aryloxy C_{1-4} alkyl or amino C_{1-4} alkyl; whereby each amino group may optionally be mono- or where possible disubstituted with C_{1-4} alkyl;
- R_5 and $-A-R_6$ taken together with the nitrogen atom to which they are attached may also form Het^1 or Het^2 ;
- R_7 is hydrogen, aryl, C_{3-7} cycloalkyl, Het^1 , Het^2 , C_{2-6} alkenyl, C_{2-6} alkynyl or C_{1-6} alkyl optionally substituted with aryl, Het^1 , Het^2 , C_{3-7} cycloalkyl, C_{1-4} alkyloxycarbonyl, carboxyl, aminocarbonyl, mono- or di(C_{1-4} alkyl)aminocarbonyl, aminosulfonyl, C_{1-4} alkyl $S(=O)_t$, hydroxy, cyano, halogen or amino optionally mono- or disubstituted where the substituents are selected from C_{1-4} alkyl, aryl, aryl C_{1-4} alkyl, C_{3-7} cycloalkyl, C_{3-7} cycloalkyl C_{1-4} alkyl, Het^1 , Het^2 , Het^1C_{1-4} alkyl and Het^2C_{1-4} alkyl;
- R_8 is hydrogen, C_{1-6} alkyl, C_{2-6} alkenyl, aryl C_{1-6} alkyl, C_{3-7} cycloalkyl, C_{3-7} cycloalkyl- C_{1-6} alkyl, aryl, Het^1 , Het^1C_{1-6} alkyl, Het^2 , Het^2C_{1-6} alkyl;
- R_{12} is hydrogen, $-NH_2$, $-C_{1-6}$ alkyl or C_{1-6} alkyl- $W-R_{14}$, wherein said C_{1-6} alkyl is optionally substituted with halogen, hydroxy, aryl, heteroaryl, Het^1 , Het^2 , or amino wherein said amino is optionally mono- or di-substituted with C_{1-4} alkyl; or
- R_{12} may also be $-N(R_5)(AR_6)$, wherein
- A is C_{1-6} alkanediyl, $-C(=O)-$, $-C(=S)-$, $-S(=O)_2-$, C_{1-6} alkanediyl $-C(=O)-$, C_{1-6} alkanediyl $-C(=S)-$ or C_{1-6} alkanediyl $-S(=O)_2-$; wherein if A is C_{1-6} alkanediyl $-C(=O)-$, C_{1-6} alkanediyl $-C(=S)-$ or C_{1-6} alkanediyl $-S(=O)_2-$, the NR_5 moiety is attached to the C_{1-6} alkanediyl moiety;
- W is oxy, carbonyl, oxycarbonyl, carbonyloxy, oxycarbonyloxy, amino, aminocarbonyl, carbonylamino or sulphur;
- R_{13} is hydrogen, C_{1-6} -alkyl, optionally substituted by aryl, Het^1 , Het^2 , hydroxy, halogen, amino whereby the amino group may be optionally be mono- or di-substituted with C_{1-4} alkyl;
- R_{14} is C_{1-6} alkyl, aryl, Het^1 or Het^2 .
2. A compound according to claim 1 wherein
- L is $-C(=O)-$, $-O-C(=O)-$, $-NR_8-C(=O)-$, $-O-C_{1-6}$ alkanediyl- $C(=O)-$, $-NR_8-C_{1-6}$ alkanediyl- $C(=O)-$, $-S(=O)_2-$, $-O-S(=O)_2-$, $-NR_8-S(=O)_2-$,

-NR₅-C₁₋₆alkanediyl-S(=O)₂, whereby either the C(=O) group or the S(=O)₂ group is attached to the NR₂ moiety.

3. A compound according to claim 1 or 2 wherein R₁ is hydrogen, C₁₋₆alkyl, C₂₋₆alkenyl, arylC₁₋₆alkyl, C₃₋₇cycloalkyl, C₃₋₇cycloalkylC₁₋₆alkyl, aryl, Het¹, Het¹C₁₋₆alkyl, Het² or Het²C₁₋₆alkyl; wherein Het¹ is a saturated or partially unsaturated monocyclic heterocycle having 5 or 6 ring members, which contains one or more heteroatom ring members selected from nitrogen, oxygen or sulfur and which is optionally substituted on one or more carbon atoms.
4. A compound according to claim 1 or 3 wherein L is -O-CH₂-C(=O)-, -O-C(=O)-, -C(=O)- or -CH₂-O-C(=O)-.
5. A compound according to any one of claims 1 to 4 wherein
 - A is C₁₋₆alkanediyl, -C(=O)- or C₁₋₆alkanediyl-C(=O)-; whereby the point of attachment to the NR₅ moiety is the C₁₋₆alkanediyl group in those moieties containing said group;
 - R₅ is hydrogen, C₁₋₆alkyl, Het¹C₁₋₆alkyl, Het²C₁₋₆alkyl, aminoC₁₋₆alkyl whereby the amino group may optionally be mono- or di-substituted with C₁₋₄alkyl; and
 - in case -A- is -C(=O)- then R₆ is C₁₋₆alkyloxy, Het¹, Het¹oxy or Het²oxy, aryl, Het¹C₁₋₄alkyl, Het¹oxyC₁₋₄alkyl, Het²C₁₋₄alkyl, Het²oxyC₁₋₄alkyl, arylC₁₋₄alkyl, aryloxyC₁₋₄alkyl or aminoC₁₋₄alkyl; and
 - in case -A- is C₁₋₆alkanediyl then R₆ is amino, C₁₋₆alkyloxy, Het¹, Het¹oxy or Het²oxy; and
 - in case -A- is C₁₋₆alkanediyl-C(=O)- then R₆ is C₁₋₆alkyloxy, Het¹, Het¹oxy or Het²oxy, aryl, C₁₋₆alkyl, Het¹C₁₋₄alkyl, Het¹oxyC₁₋₄alkyl, Het²C₁₋₄alkyl, Het²oxyC₁₋₄alkyl, arylC₁₋₄alkyl, aryloxyC₁₋₄alkyl or aminoC₁₋₄alkyl; whereby each of the amino groups in the definition of R₆ may optionally be substituted with one or more substituents selected from C₁₋₄alkyl, C₁₋₄alkyl-carbonyl, C₁₋₄alkyloxycarbonyl, aryl, arylcarbonyl, aryloxycarbonyl, Het¹, Het², arylC₁₋₄alkyl, Het¹C₁₋₄alkyl or Het²C₁₋₄alkyl; and
 - R₅ and -A-R₆ taken together with the nitrogen atom to which they are attached may also form Het¹ whereby Het¹ is substituted by at least an oxo group.
6. A compound according to claim 1 to 4, wherein

- R₁₂ is H, -NH₂, -N(R₅)(AR₆), -C₁₋₆alkyl or C₁₋₆alkyl-W-R₁₄, wherein said C₁₋₆alkyl is optionally substituted with halogen, hydroxy, aryl, heteroaryl, Het¹, Het², or amino optionally mono- or di-substituted with C₁₋₄alkyl;
- A is C₁₋₆alkanediyl or -C(=O)-; R⁵ is hydrogen, methyl, Het¹C₁₋₆alkyl, aminoC₁₋₆alkyl
 5 whereby the amino group may optionally be mono- or di-substituted with C₁₋₄alkyl; R⁶ is C₁₋₆alkyloxy, Het¹, amino; and in case -A- is other than C₁₋₆alkanediyl then R⁶ may also be C₁₋₆alkyl, Het¹C₁₋₄alkyl or aminoC₁₋₄alkyl; whereby each of the amino groups may optionally be substituted;
- W is oxycarbonyloxy;
- 10 R₁₄ is Het¹.
7. A compound according to claim 1 to 4, wherein
 R₁₂ is -N(R₅)(AR₆);
 A is C₁₋₆alkanediyl or -C(=O)-; R₅ is hydrogen; R₆ is C₁₋₆alkyloxy, Het¹, C₁₋₄alkyloxy-
 15 aryl, amino, aminoC₁₋₄alkylamino, aminoC₁₋₄alkyloxy; and in case -A- is -C(=O)- then R₆ may also be C₁₋₆alkyl, Het¹C₁₋₄alkyl, or aminoC₁₋₄alkyl; whereby each of the amino groups may optionally be substituted with C₁₋₄alkyl.
8. A compound according to claim 1 to 4, wherein
 20 R₁₃ is C₁₋₄alkyl preferably methyl, ethyl or isobutyl; optionally substituted with aryl preferably phenyl, Het¹ preferably pyrrolidinyl, Het² preferably imidazolyl or pyridinyl, or amino; wherein the amino group is optionally substituted with C₁₋₄alkyl.
9. A compound according to claim 1 to 4, wherein
 25 R₁₂ is H, -NH₂, -N(R₅)(AR₆), -C₁₋₆alkyl or C₁₋₆alkyl-W-R₁₄, wherein said C₁₋₆alkyl is optionally substituted with hydroxy or amino wherein said amino is optionally mono- or di-substituted with C₁₋₄alkyl;
 A is C₁₋₆alkanediyl or -C(=O)-; R₅ is hydrogen, methyl, Het¹C₁₋₆alkyl, aminoC₁₋₆alkyl
 whereby the amino group may optionally be mono- or di-substituted with C₁₋₄alkyl; R₆
 30 is C₁₋₆alkyloxy, Het¹, amino; and in case -A- is other than C₁₋₆alkanediyl then R₆ may also be C₁₋₆alkyl, Het¹C₁₋₄alkyl or aminoC₁₋₄alkyl; whereby each of the amino groups may optionally be substituted;
- R₁₃ is H, C₁₋₆-alkyl, optionally substituted by aryl, Het¹, Het², hydroxy, halogen, amino
 whereby the amino group may be optionally be mono- or di-substituted with C₁₋₄alkyl.
- 35 R₁₄ is alkyl, aryl Het¹, Het².

-66-

10. A pharmaceutical composition, comprising an effective amount of at least one compound as claimed in any one of claims 1 to 9, and a pharmaceutically tolerable excipient.
11. A substance or composition for use in a method of treating or combating infection or disease associated with multi-drug resistant retrovirus infection in a mammal, and said method comprising administering an effective amount of said substance or composition, said substance or composition comprising at least one compound according to any one of claims 1 to 9, to said mammal.
12. The use of a compound as claimed in any one of claims 1 to 9 in the manufacture of a medicament for treating or combating infection or disease associated with multi-drug resistant retrovirus infection in a mammal.
13. A composition comprising at least (a) a compound of formula (I) as claimed in claims 1 to 9 and, (b) a second antiretroviral agent for the simultaneous, separate or sequential use.
14. Use of a compound according to any one of claims 1 to 9 in the manufacture of a preparation for inhibiting a protease of a multi-drug resistant retrovirus in a mammal infected with said retrovirus.
15. A substance or composition for use in a method of inhibiting a protease of a multi-drug resistant retrovirus in a mammal infected with said retrovirus, said substance or composition comprising a compound according to any one of claims 1 to 9, and said method comprising administering a protease inhibiting amount of said substance or composition to said mammal.