

- 1 -

Compositions and methods for therapy and diagnosis of cancer and cancer metastasis

Despite interdisciplinary approaches and exhaustive use of classical therapeutic procedures, cancers are still among the leading causes of death.

In particular, metastasis is one of the most critical factors responsible for the failure of cancer treatment. Although protein expression profiling, gene array analysis and determination of critical factors in tumor tissue have improved the prognostic classification of tumors, it is still difficult to predict the risk of metastasis analyzing the resected primary tumor. After complete tumor resection survival usually depends on occurrence of metastasis. Presently, it is difficult if not impossible to predict, whether a primary tumor has metastasized or not.

Tumor cells biologically differ substantially from their nonmalignant cells of origin. These differences are due to genetic alterations acquired during tumor development and result, inter alia, also in the formation of qualitatively or quantitatively altered molecular structures in the cancer cells. Tumor-associated structures of this kind are, in particular, genetic products the expression of which is induced or enhanced during the course of malignant transformation.

The factors regulating acquisition of the metastatic phenotype are mostly undefined. It is known that some histopathological parameters are associated with tumor-free survival, for instance tumor stage and

histological grade. However, it is not yet possible to predict the risk of metastasis by quantification of critical factors in tumor tissue.

5 It was the object of the present invention to provide compositions and methods for a diagnosis and therapy of cancer, in particular cancer metastasis. In particular, it was the object of the present invention to provide compositions and methods for a diagnosis of metastatic
10 behaviour of cancer.

These objects are achieved by the subject matter of the claims.

15 The studies presented herein demonstrate that cancers expressing both, TPTE and CXCR4, exhibit a nearly 30 fold increased risk for metastasis, in particular distant metastasis, as compared to tumors lacking at least one of these molecules. A combination of these
20 markers is therefore useful for evaluation of the clinical prognosis of cancer patients and for targeted therapeutical approaches.

Accordingly, the present invention relates to methods
25 which make possible to assess and/or prognose a cancer disease, the metastatic behaviour of a cancer disease and/or the occurrence of a relapse of cancer. In particular, the methods of the invention make possible to assess and/or prognose the occurrence of cancer
30 metastasis, in particular distant metastasis. Preferably, the methods of the invention allow to discriminate malign from benign conditions.

In particular embodiments, the methods of the invention
35 make possible to assess and/or prognose the success of

a cancer therapy which has been administered or will be administered. In particular, the methods of the invention make possible to assess and/or prognose the occurrence of a relapse of cancer following cancer
5 therapy, e.g., by surgery, chemotherapy and/or radiation therapy.

In one aspect the invention relates to a method for diagnosing, monitoring, i.e. determining the
10 regression, progression, course and/or onset of, and/or prognosing cancer, the metastatic behaviour of cancer and/or the presence of a relapse of cancer in a patient, which method comprises quantitatively and/or qualitatively determining the level of expression of
15 TPTE in a biological sample isolated from said patient and quantitatively and/or qualitatively determining the level of expression of CXCR4 in a biological sample isolated from said patient. In a particular preferred embodiment, the invention in this aspect relates to a
20 method of diagnosing whether a patient has cancer metastasis, in particular distant metastasis.

In particular embodiments of the method of the invention, the level of expression of TPTE and the
25 level of expression of CXCR4 are determined in the same sample either concurrently or consecutively. In further embodiments of the method of the invention, the level of expression of TPTE and the level of expression of CXCR4 are determined in different samples wherein said
30 different samples may be the same kind of sample, e.g. both may be a blood sample, taken from the patient at the same or different points in time and from the same or different regions of the body, or may be different kinds of samples, e.g. one is a blood sample and the
35 other is an urine sample.

Preferably a level of expression of TPTE and a level of expression of CXCR4 which is increased compared to the level of expression in a subject without cancer, without a risk for cancer, without metastasis of cancer, without a risk for metastasis of cancer, without a relapse of cancer, and/or without a risk for a relapse of cancer is indicative for cancer or for a potential for cancer, for a metastatic behaviour of cancer or for a potential for a metastatic behaviour of cancer and/or for a relapse of cancer or for a potential for a relapse of cancer.

Preferably, the quantitative and/or qualitative determination of the level of expression of TPTE comprises (i) detecting or determining the amount of a nucleic acid which is selected from the group consisting of (a) a nucleic acid which comprises a nucleic acid sequence selected from the group consisting of SEQ ID NOs: 1, 2, 3, 4, 5, 6, and 7, a part or derivative thereof, (b) a nucleic acid which hybridizes with the nucleic acid of (a) under stringent conditions, (c) a nucleic acid which is degenerate with respect to the nucleic acid of (a) or (b), and (d) a nucleic acid which is complementary to the nucleic acid of (a), (b) or (c), and/or (ii) detecting or determining the amount of a protein or peptide encoded by the nucleic acid under (i) or of a part or derivative thereof, and/or (iii) detecting or determining the amount of an antibody specific for the protein or peptide or for the part or derivative under (ii), and/or (iv) detecting or determining the amount of a T lymphocyte specific for the protein or peptide or for the part or derivative under (ii), optionally in a complex with a MHC molecule, in a biological sample

isolated from a patient. Preferably, the nucleic acid under (i) in said quantitative and/or qualitative determination of the level of expression of TPTE comprises a nucleic acid sequence encoding a protein or peptide comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 8, 9, 10, 11, 12, 13, and 14, a part or derivative thereof and/or the protein or peptide under (ii) in said quantitative and/or qualitative determination of the level of expression of TPTE comprises an amino acid sequence selected from the group consisting of SEQ ID NOs: 8, 9, 10, 11, 12, 13, and 14, a part or derivative thereof.

Preferably, the quantitative and/or qualitative determination of the level of expression of CXCR4 comprises (i) detecting or determining the amount of a nucleic acid which is selected from the group consisting of (a) a nucleic acid which comprises a nucleic acid sequence selected from the group consisting of SEQ ID NOs: 47, and 48, a part or derivative thereof, (b) a nucleic acid which hybridizes with the nucleic acid of (a) under stringent conditions, (c) a nucleic acid which is degenerate with respect to the nucleic acid of (a) or (b), and (d) a nucleic acid which is complementary to the nucleic acid of (a), (b) or (c), and/or (ii) detecting or determining the amount of a protein or peptide encoded by the nucleic acid under (i) or of a part or derivative thereof, and/or (iii) detecting or determining the amount of an antibody specific for the protein or peptide or for the part or derivative under (ii), and/or (iv) detecting or determining the amount of a T lymphocyte specific for the protein or peptide or for the part or derivative under (ii), optionally in a complex with a MHC molecule, in a biological sample

- 6 -

isolated from a patient. Preferably, the nucleic acid under (i) in said quantitative and/or qualitative determination of the level of expression of CXCR4 comprises a nucleic acid sequence encoding a protein or peptide comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 49, and 50, a part or derivative thereof and/or the protein or peptide under (ii) in said quantitative and/or qualitative determination of the level of expression of CXCR4 comprises an amino acid sequence selected from the group consisting of SEQ ID NOs: 49, and 50, a part or derivative thereof.

Means for accomplishing said quantitative and/or qualitative determination of the level of expression are described herein and will be apparent to the skilled person.

According to the invention, detection of a nucleic acid or determining the amount of a nucleic acid may be carried out using a oligo- or polynucleotide probe which hybridizes specifically to said nucleic acid, or may be carried out by selective amplification of said nucleic acid, e.g. by means of PCR amplification. In one embodiment, the probe comprises a sequence of 6-50, in particular 10-30, 15-30 and 20-30, contiguous nucleotides of said nucleic acid and the primers used in said amplification each comprise a sequence of 6-50, in particular 10-30, 15-30 and 20-30, contiguous nucleotides of said nucleic acid.

Preferably, the detection or determination of the amount of said nucleic acid in the method of the invention comprises (i) contacting the biological sample with an agent which binds specifically to the

nucleic acid, and (ii) detecting the formation of or determining the amount of a complex between the agent and the nucleic acid. Preferably, the agent which binds specifically to the nucleic acid is an oligonucleotide
5 or polynucleotide, which hybridizes specifically to said nucleic acid.

According to the invention, detection of a protein or peptide or of a part or derivative thereof or
10 determining the amount of a protein or peptide or of a part or derivative thereof may be carried out using an antibody binding specifically to said protein or peptide or part or derivative thereof. In particular
embodiments, the protein or peptide or part or
15 derivative thereof which is to be detected or the amount of which is to be determined in the methods of the present invention is present in a complex with an MHC molecule.

20 Preferably, the detection or determination of the amount of said protein or peptide or said part or derivative thereof in the method of the invention comprises (i) contacting the biological sample with an agent which binds specifically to the protein or
25 peptide or the part or derivative thereof, and (ii) detecting the formation of or determining the amount of a complex between the agent and the protein or peptide or part or derivative thereof. Preferably, the agent which binds specifically to the protein or peptide or
30 the part or derivative thereof is an antibody binding specifically to said protein or peptide or to said part or derivative thereof.

According to the invention, detection of an antibody or
35 determining the amount of an antibody may be carried

out using a protein or peptide binding specifically to said antibody.

Preferably, the detection or determination of the
5 amount of said antibody in the method of the invention comprises (i) contacting the biological sample with an agent which binds specifically to the antibody, and (ii) detecting the formation of or determining the amount of a complex between the agent and the antibody.
10 Preferably, the agent which binds specifically to the antibody is a protein or peptide binding specifically to said antibody.

According to the invention, detection of a T lymphocyte
15 or determining the amount of a T lymphocyte may be carried out using a cell presenting a complex between a protein or peptide and an MHC molecule for which the T lymphocyte is specific, wherein the cell is preferably an antigen-presenting cell. Detection of or determining
20 the amount of a T lymphocyte may also be carried out by detecting its proliferation, cytokine production, and/or cytotoxic activity which may be triggered by specific stimulation with a complex between a protein or peptide and an MHC molecule for which the T
25 lymphocyte is specific. Detection of or determining the amount of a T lymphocyte may also be carried out with aid of a recombinant MHC molecule or a complex of two or more MHC molecules loaded with one or more proteins or peptides.

30

Preferably, the detection or determination of the amount of said T lymphocyte comprises (i) contacting the biological sample with an agent which binds specifically to the T lymphocyte, and (ii) detecting
35 the formation of or determining the amount of a complex

between the agent and the T lymphocyte. Preferably, the agent which binds specifically to the T lymphocyte is a cell presenting the complex between the protein or peptide or the part or derivative thereof for which the
5 T lymphocyte is specific and an MHC molecule.

An agent which is used for detection or determining the amount in the methods of the invention such as an oligo- or polynucleotide probe, an antibody, a protein
10 or peptide or a cell is preferably labeled in a detectable manner, in particular by a detectable marker or diagnostic substance such as a radioactive marker, fluorescence marker or an enzymic marker.

15 In one embodiment, the method of the invention comprises determining the level of expression in a first sample at a first point in time and in a further sample at a second point in time and a comparison of the two samples. Preferably, a level of expression of
20 TPTE and a level of expression of CXCR4 which is increased in a sample compared to a sample taken earlier from a patient indicates that the patient has developed or is about to develop cancer and/or a metastasis of cancer and/or a relapse of cancer.
25 Preferably, a level of expression of TPTE and a level of expression of CXCR4 which is decreased in a sample compared to a sample taken earlier from a patient indicates regression of cancer and/or a metastasis of cancer in said patient and thus, preferably indicates a
30 successful cancer therapy.

In a further embodiment, the biological sample isolated from the patient is compared to a comparable normal biological sample, e.g. a sample isolated from a
35 healthy individual. Preferably, a level of expression

- 10 -

of TPTE and a level of expression of CXCR4 which is increased in a patient's sample compared to a sample taken from a healthy individual indicates that the patient has developed or is about to develop cancer
5 and/or a metastasis of cancer and/or a relapse of cancer.

The determination of the level of expression of TPTE may also involve the determination of methylation
10 patterns and/or the degree of methylation with a nucleic acid which is selected from the group consisting of (a) a nucleic acid which comprises a nucleic acid sequence selected from the group consisting of SEQ ID NOs: 1, 2, 3, 4, 5, 6, and 7, a
15 part or derivative thereof, (b) a nucleic acid which hybridizes with the nucleic acid of (a) under stringent conditions, (c) a nucleic acid which is degenerate with respect to the nucleic acid of (a) or (b), and (d) a nucleic acid which is complementary to the nucleic acid
20 of (a), (b) or (c), and/or with a nucleic acid comprising a nucleic acid sequence encoding a protein or peptide comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 8, 9, 10, 11, 12, 13, and 14, a part or derivative thereof,
25 preferably within the non-coding region thereof and more preferably within the promoter region thereof.

A degree of methylation which is lower compared to a control, e.g. a subject without cancer, without a risk
30 for cancer, without metastasis of cancer, without a risk for metastasis of cancer, without a relapse of cancer, and/or without a risk for a relapse of cancer, or no methylation preferably is indicative for an increased level of expression of TPTE.

The determination of methylation patterns and/or the degree of methylation can be performed, for example, by using methods on the basis of PCR, with the aid of restriction enzymes or by sequencing. In one preferred
5 embodiment, genomic DNA is selectively amplified following treatment with a bisulfite containing reagent. The oligonucleotides used in such amplification preferably have a sequence binding to the nucleic acid treated with the bisulfite containing
10 reagent and preferably are completely complementary thereto. Preferably the oligonucleotides are adapted to a different degree of methylation of the nucleic acid and bring about amplification products which can be differentiated. A test suitable for this can be as
15 follows: (1) extraction of DNA from tissue samples of patients, for example using paraffin embedded material, (2) treatment of the DNA with bisulfite containing reagents (e.g. as described in Clark S.J. et al., Nucleic Acids Res. 22(15):2990-7, 1994), (3)
20 amplification of DNA by means of PCR and (4) analysis of the amount of sequence specific amplification products (e.g. by means of quantitative PCR, hybridization techniques such as microarray methods).

25 In particular embodiments of the method of the invention, the patient has cancer, is suspected of having cancer or developing cancer, or has a risk for developing cancer. In further embodiments of the method of the invention, the patient has a cancer metastasis,
30 is suspected of having a cancer metastasis or developing a cancer metastasis, or has a risk for developing a cancer metastasis. In particular embodiments of the method of the invention, the patient has already been subjected to cancer therapy such as by
35 tumor resection, radiation therapy and/or chemotherapy,

or it is intended to subject the patient to such therapy.

The method of diagnosing, monitoring, and/or prognosing
5 cancer, the metastatic behaviour of cancer and/or the
presence of a relapse of cancer according to the
invention preferably allows the prognosis of a worsened
course of a disease, whereby among other things
planning of a more aggressive therapy is made possible.
10 This prognostic method also allows to delimit still
benign alterations, e.g. hyperplasias, from tumor
precursors which are already to be appraised as
unfavourable and to anticipate therefore a cancer
disposition already before an invasive tumor has
15 formed.

In a further aspect, the invention relates to a kit
comprising means for quantitatively and/or
qualitatively determining the level of expression of
20 TPTE and means for quantitatively and/or qualitatively
determining the level of expression of CXCR4 in a
biological sample isolated from a patient. Preferably,
the kit is useful in the method for diagnosing,
monitoring and/or prognosing cancer, the metastatic
25 behaviour of cancer and/or the presence of a relapse of
cancer of the invention.

Means for quantitatively and/or qualitatively
determining the level of expression of TPTE and CXCR4
30 are as discussed above.

Preferably, said means for quantitatively and/or
qualitatively determining the level of expression of
TPTE are selected from the group consisting of (i)
35 means for detecting or determining the amount of a

- 13 -

nucleic acid which is selected from the group consisting of (a) a nucleic acid which comprises a nucleic acid sequence selected from the group consisting of SEQ ID NOS: 1, 2, 3, 4, 5, 6, and 7, a
5 part or derivative thereof, (b) a nucleic acid which hybridizes with the nucleic acid of (a) under stringent conditions, (c) a nucleic acid which is degenerate with respect to the nucleic acid of (a) or (b), and (d) a nucleic acid which is complementary to the nucleic acid
10 of (a), (b) or (c), and/or (ii) means for detecting or determining the amount of a protein or peptide encoded by the nucleic acid under (i) or of a part or derivative thereof, and/or (iii) means for detecting or determining the amount of an antibody specific for the
15 protein or peptide or the part or derivative under (ii), and/or (iv) means for detecting or determining the amount of a T lymphocyte specific for the protein or peptide or the part or derivative under (ii), optionally in a complex with a MHC molecule, in a
20 biological sample isolated from a patient. Preferably, the nucleic acid under (i) in said quantitative and/or qualitative determination of the level of expression of TPTE comprises a nucleic acid sequence encoding a protein or peptide comprising an amino acid sequence
25 selected from the group consisting of SEQ ID NOS: 8, 9, 10, 11, 12, 13, and 14, a part or derivative thereof and/or the protein or peptide under (ii) in said quantitative and/or qualitative determination of the level of expression of TPTE comprises an amino acid
30 sequence selected from the group consisting of SEQ ID NOS: 8, 9, 10, 11, 12, 13, and 14, a part or derivative thereof.

Preferably, said means for quantitatively and/or
35 qualitatively determining the level of expression of

- 14 -

CXCR4 are selected from the group consisting of (i) means for detecting or determining the amount of a nucleic acid which is selected from the group consisting of (a) a nucleic acid which comprises a
5 nucleic acid sequence selected from the group consisting of SEQ ID NOs: 47, and 48, a part or derivative thereof, (b) a nucleic acid which hybridizes with the nucleic acid of (a) under stringent conditions, (c) a nucleic acid which is degenerate with
10 respect to the nucleic acid of (a) or (b), and (d) a nucleic acid which is complementary to the nucleic acid of (a), (b) or (c), and/or (ii) means for detecting or determining the amount of a protein or peptide encoded by the nucleic acid under (i) or of a part or
15 derivative thereof, and/or (iii) means for detecting or determining the amount of an antibody specific for the protein or peptide or the part or derivative under (ii), and/or (iv) means for detecting or determining the amount of a T lymphocyte specific for the protein
20 or peptide or the part or derivative under (ii), optionally in a complex with a MHC molecule, in a biological sample isolated from a patient. Preferably, the nucleic acid under (i) in said quantitative and/or qualitative determination of the level of expression of
25 CXCR4 comprises a nucleic acid sequence encoding a protein or peptide comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 49, and 50, a part or derivative thereof and/or the protein or peptide under (ii) in said quantitative and/or
30 qualitative determination of the level of expression of CXCR4 comprises an amino acid sequence selected from the group consisting of SEQ ID NOs: 49, and 50, a part or derivative thereof.

35 Preferably, said means for detecting or determining the

amount of said nucleic acid comprise an agent which binds specifically to the nucleic acid. Preferably, the agent which binds specifically to the nucleic acid is an oligonucleotide or polynucleotide, which hybridizes
5 specifically to said nucleic acid.

Preferably, said means for detecting or determining the amount of said protein or peptide or part or derivative thereof comprise an agent which binds specifically to
10 the protein or peptide or the part or derivative thereof. Preferably, the agent which binds specifically to the protein or peptide or the part or derivative thereof is an antibody binding specifically to said protein or peptide or to said part or derivative
15 thereof.

Preferably, said means for detecting or determining the amount of said antibody comprise an agent which binds specifically to the antibody. Preferably, the agent
20 which binds specifically to the antibody is a protein or peptide binding specifically to said antibody.

Preferably, said means for detecting or determining the amount of said T lymphocyte comprise an agent which
25 binds specifically to the T lymphocyte. Preferably, the agent which binds specifically to the T lymphocyte is a cell presenting the complex between the protein or peptide or the part or derivative thereof for which the T lymphocyte is specific and an MHC molecule.

30

In a further aspect, the invention relates to a pharmaceutical composition comprising (i) an agent which is effective in reducing or inhibiting expression or activity of TPTE and/or which binds to TPTE and has
35 tumor destroying or tumor inhibiting activity, and (ii)

an agent which is effective in reducing or inhibiting expression or activity of CXCR4 and/or which binds to CXCR4 and has tumor destroying or tumor inhibiting activity. The terms "activity of TPTE" and "activity of
5 CXCR4" relate to any activity of TPTE or CXCR4 in a cell or an organism such as enzymatic or regulatory activity, in particular cell migration modulatory activity. Preferably, the agent which binds to TPTE or CXCR4 and has tumor destroying or tumor inhibiting
10 activity is specific for cells expressing or abnormally expressing TPTE or CXCR4, respectively. Preferably, such agent comprises a therapeutic substance.

In certain embodiments of the pharmaceutical
15 composition of the invention, the agent is an antisense nucleic acid which hybridizes selectively with the nucleic acid coding for TPTE and/or hybridizes selectively with the nucleic acid coding for CXCR4. In further embodiments, the agent is a siRNA preferably
20 comprising a sense RNA strand and an antisense RNA strand, wherein the sense and antisense RNA strands form an RNA duplex, and wherein the sense RNA strand comprises a nucleotide sequence substantially identical to a target sequence of about 19 to about 25 contiguous
25 nucleotides in TPTE mRNA and/or in CXCR4 mRNA. In even further embodiments, the agent is an antibody which binds selectively to TPTE and/or CXCR4. The antisense nucleic acid, siRNA and/or antibody discussed above may be combined in the pharmaceutical composition of the
30 invention.

In even a further aspect, the invention relates to a pharmaceutical composition comprising (I) one or more components selected from the group consisting of (i)
35 TPTE or a part or derivative thereof, (ii) a nucleic

- 17 -

acid which codes for TPTE or a part or derivative thereof, (iii) an antibody which binds to TPTE or a part thereof, (iv) an antisense nucleic acid which hybridizes specifically with a nucleic acid coding for TPTE, (v) an siRNA directed against a nucleic acid coding for TPTE, (vi) a host cell which expresses TPTE or a part or derivative thereof, and (vii) isolated complexes between TPTE or a part or derivative thereof and an MHC molecule, and (II) one or more components selected from the group consisting of (i) CXCR4 or a part or derivative thereof, (ii) a nucleic acid which codes for CXCR4 or a part or derivative thereof, (iii) an antibody which binds to CXCR4 or a part or derivative thereof, (iv) an antisense nucleic acid which hybridizes specifically with a nucleic acid coding for CXCR4, (v) an siRNA directed against a nucleic acid coding for CXCR4, (vi) a host cell which expresses CXCR4 or a part or derivative thereof, and (vii) isolated complexes between CXCR4 or a part or derivative thereof and an MHC molecule.

In one embodiment, a nucleic acid coding for TPTE or CXCR4 or a part or derivative thereof is present in the pharmaceutical composition in an expression vector and functionally linked to a promoter.

In a further embodiment, a host cell present in a pharmaceutical composition of the invention secretes TPTE or CXCR4 or the part or derivative thereof, expresses it on the surface and preferably additionally express an MHC molecule which binds to said TPTE or CXCR4 or the part or derivative thereof. In one embodiment, the host cell expresses the MHC molecule endogenously. In a further embodiment, the host cell expresses the MHC molecule and/or TPTE or CXCR4 or the

part or derivative thereof in a recombinant manner. The host cell is preferably nonproliferative. In a preferred embodiment, the host cell is an antigen-presenting cell, in particular a dendritic cell, a
5 monocyte or a macrophage.

In a further embodiment, an antibody present in a pharmaceutical composition of the invention is a monoclonal antibody. In further embodiments, the
10 antibody is a chimeric or humanized antibody, a fragment of a natural antibody or a synthetic antibody.

An antisense nucleic acid present in a pharmaceutical
15 composition of the invention may comprise a sequence of 6-50, in particular 10-30, 15-30 and 20-30, contiguous nucleotides of the nucleic acid coding for TPTE or the part or derivative thereof and/or of the nucleic acid coding for CXCR4 or the part or derivative thereof.

20

In further embodiments, TPTE or CXCR4 or the part or derivative thereof, provided by a pharmaceutical composition of the invention either directly or via expression of a nucleic acid, binds to MHC molecules on
25 the surface of cells, said binding preferably causing a cytolytic response and/or inducing cytokine release.

The antibody contained in the pharmaceutical composition of the invention may be coupled to a
30 therapeutic substance.

In particular embodiments of the siRNA directed against a nucleic acid coding for TPTE contained in the pharmaceutical composition of the invention the target
35 sequence has a nucleic acid sequence selected from the

group consisting of nucleotide positions 3-21 of SEQ ID NO: 15, nucleotide positions 3-21 of SEQ ID NO: 18, nucleotide positions 3-21 of SEQ ID NO: 21, nucleotide positions 3-21 of SEQ ID NO: 24, nucleotide positions 3-21 of SEQ ID NO: 27, nucleotide positions 3-21 of SEQ ID NO: 30, and nucleotide positions 3-21 of SEQ ID NO: 33. In further particular embodiments of the siRNA the sense RNA strand has the sequence of SEQ ID NO: 16 and the antisense RNA strand has the sequence of SEQ ID NO: 17, or the sense RNA strand has the sequence of SEQ ID NO: 19 and the antisense RNA strand has the sequence of SEQ ID NO: 20, or the sense RNA strand has the sequence of SEQ ID NO: 22 and the antisense RNA strand has the sequence of SEQ ID NO: 23, or the sense RNA strand has the sequence of SEQ ID NO: 25 and the antisense RNA strand has the sequence of SEQ ID NO: 26, or the sense RNA strand has the sequence of SEQ ID NO: 28 and the antisense RNA strand has the sequence of SEQ ID NO: 29, or the sense RNA strand has the sequence of SEQ ID NO: 31 and the antisense RNA strand has the sequence of SEQ ID NO: 32, or the sense RNA strand has the sequence of SEQ ID NO: 34 and the antisense RNA strand has the sequence of SEQ ID NO: 35.

A pharmaceutical composition of the invention may comprise a pharmaceutically compatible carrier and/or an adjuvant.

In a further aspect, the invention relates to a method of treating or preventing cancer, metastasis of cancer or relapse of cancer comprising administering to a patient (I) one or more components selected from the group consisting of (i) TPTE or a part or derivative thereof, (ii) a nucleic acid which codes for TPTE or a part or derivative thereof, (iii) an antibody which

- 20 -

binds to TPTE or a part thereof, (iv) an antisense nucleic acid which hybridizes specifically with a nucleic acid coding for TPTE, (v) an siRNA directed against a nucleic acid coding for TPTE, (vi) a host
5 cell which expresses TPTE or a part or derivative thereof, and (vii) isolated complexes between TPTE or a part or derivative thereof and an MHC molecule, and (II) one or more components selected from the group consisting of (i) CXCR4 or a part or derivative
10 thereof, (ii) a nucleic acid which codes for CXCR4 or a part or derivative thereof, (iii) an antibody which binds to CXCR4 or a part or derivative thereof, (iv) an antisense nucleic acid which hybridizes specifically with a nucleic acid coding for CXCR4, (v) an siRNA
15 directed against a nucleic acid coding for CXCR4, (vi) a host cell which expresses CXCR4 or a part or derivative thereof, and (vii) isolated complexes between CXCR4 or a part or derivative thereof and an MHC molecule.

20

The invention also relates to a method of treating or preventing cancer, metastasis of cancer or relapse of cancer comprising administering the pharmaceutical composition of the invention.

25

Preferably, the cancer is a lung tumor, a breast tumor, a prostate tumor, a melanoma, a colon tumor, a gastric tumor, a pancreatic tumor, an ENT tumor, a renal cell carcinoma or a cervical carcinoma, a colon carcinoma or
30 a mammary carcinoma.

Preferably, the cancer, metastasis of cancer or relapse of cancer is characterized by expression or abnormal expression of (i) a nucleic acid which is selected from
35 the group consisting of (a) a nucleic acid which

comprises a nucleic acid sequence selected from the group consisting of SEQ ID NOs: 1, 2, 3, 4, 5, 6, and 7, a part or derivative thereof, (b) a nucleic acid which hybridizes with the nucleic acid of (a) under stringent conditions, (c) a nucleic acid which is degenerate with respect to the nucleic acid of (a) or (b), and (d) a nucleic acid which is complementary to the nucleic acid of (a), (b) or (c), and/or (ii) a protein or peptide encoded by the nucleic acid under (i). Preferably the nucleic acid under (i) comprises a nucleic acid sequence encoding a protein or peptide comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 8, 9, 10, 11, 12, 13, and 14, a part or derivative thereof and/or the protein or peptide under (ii) comprises an amino acid sequence selected from the group consisting of SEQ ID NOs: 8, 9, 10, 11, 12, 13, and 14, a part or derivative thereof.

More preferably, the cancer, metastasis of cancer or relapse of cancer is characterized by further expression or abnormal expression of (i) a nucleic acid which is selected from the group consisting of (a) a nucleic acid which comprises a nucleic acid sequence selected from the group consisting of SEQ ID NOs: 47, and 48, a part or derivative thereof, (b) a nucleic acid which hybridizes with the nucleic acid of (a) under stringent conditions, (c) a nucleic acid which is degenerate with respect to the nucleic acid of (a) or (b), and (d) a nucleic acid which is complementary to the nucleic acid of (a), (b) or (c), and/or (ii) a protein or peptide encoded by the nucleic acid under (i). Preferably the nucleic acid under (i) comprises a nucleic acid sequence encoding a protein or peptide comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 49, and 50, a part or

derivative thereof and/or the protein or peptide under (ii) comprises an amino acid sequence selected from the group consisting of SEQ ID NOs: 49, and 50, a part or derivative thereof.

5

In the methods of the invention the pharmaceutical composition is preferably administered in combination with radiation therapy, chemotherapy or surgery, wherein the chemotherapeutic agent is preferably
10 selected from the group consisting of cisplatin, carboplatin, cyclophosphamide, 5-fluorouracil, adriamycin, daunorubicin, and tamoxifen.

Preferably the subject or patient in the methods of the
15 invention is a human being.

Detailed description of the invention

20 The expression "determining the level of expression" with respect to TPTE or CXCR4 according to the invention relates to the determination of the absence or presence and/or the absolute and/or relative quantification of a gene product of the TPTE gene or
25 CXCR4 gene (nucleic acid and protein/peptide). The expression "determining the level of expression" according to the invention also includes situations wherein no gene product is detected or the amount of said gene product is below the detection limit.

30

Generally all methods suitable to detect and analyse nucleic acids, proteins and/or peptides can be used for determining the level of expression in the methods of the invention. PCR, gene chip/microarray systems,
35 Northern blot, RNase protection assays (RDA) can, for

example, be used for detecting and analyzing nucleic acids. Suitable immunologic methods for detecting and analyzing proteins and/or peptides among others are enzyme linked immuno assays (ELISA), sandwich, direct, 5 indirect, or competitive ELISA assays, enzyme-linked immunospot assays (ELISPOT), radio immuno assays (RIA), flow cytometry assays (FACS = fluorescence activated cell sorting), immunohistochemistry, Western blot, fluorescence resonance energy transfer (FRET) assays, 10 protein-chip assays using for example antibodies, antibody fragments, receptors, ligands, or other binding agents specific for peptides or proteins.

According to the invention, the term "binding" 15 preferably relates to a specific binding. "Specific binding" means that an agent such as an antibody binds stronger to a target such as an epitope for which it is specific compared to the binding to another target. An agent binds stronger to a first target compared to a 20 second target if it binds to the first target with a dissociation constant (K_D) which is lower than the dissociation constant for the second target. Preferably the dissociation constant (K_D) for the target to which the agent binds specifically is more than 10-fold, 25 preferably more than 20-fold, more preferably more than 50-fold, even more preferably more than 100-fold, 200-fold, 500-fold or 1000-fold lower than the dissociation constant (K_D) for the target to which the agent does not bind specifically.

30

According to the invention, a "reference" such as a reference sample or reference organism may be used to correlate and compare the results obtained in the methods of the invention from a test sample or test 35 organism, i.e. a patient. Typically the reference

organism is a healthy organism, in particular an organism which does not suffer from cancer, metastasis of cancer and/or relapse of cancer.

- 5 A "reference value" can be determined from a reference empirically by measuring a sufficiently large number of references. Preferably the reference value is determined by measuring at least 2, preferably at least 3, preferably at least 5, preferably at least 8,
10 preferably at least 12, preferably at least 20, preferably at least 30, preferably at least 50, or preferably at least 100 references.

The term "TPTE" relates to "transmembrane phosphatase
15 with tensin homology" and includes any variants, in particular splice variant, conformations, isoforms and species homologs of TPTE which are naturally expressed by cells or are expressed by cells transfected with the TPTE gene. The expression "determining the level of
20 expression of TPTE" relates to a determination of the level of a nucleic acid of TPTE, such as mRNA, and/or a determination of the level of TPTE protein.

Preferably, a "nucleic acid of TPTE", a "nucleic acid
25 encoding TPTE", a "nucleic acid coding for TPTE" or "TPTE gene" relates to a nucleic acid selected from the group consisting of (a) a nucleic acid which comprises a nucleic acid sequence selected from the group consisting of SEQ ID NOs: 1, 2, 3, 4, 5, 6, and 7, a
30 part or derivative thereof, (b) a nucleic acid which hybridizes with the nucleic acid of (a) under stringent conditions, (c) a nucleic acid which is degenerate with respect to the nucleic acid of (a) or (b), and (d) a nucleic acid which is complementary to the nucleic acid
35 of (a), (b) or (c). The terms may also include mRNA

coding for TPTE. Preferably, "TPTE" protein or simply "TPTE" comprises an amino acid sequence encoded by the afore mentioned nucleic acid, preferably an amino acid sequence selected from the group consisting of SEQ ID
5 NOS: 8, 9, 10, 11, 12, 13, and 14, a part or derivative thereof. One skilled in the art would understand that the cDNA sequence of TPTE as described above would be equivalent to TPTE mRNA, and can be used for the same purpose herein; i.e., the generation of siRNA for
10 inhibiting expression of TPTE.

The term "TPTE" also includes posttranslationally modified variants, isoforms and species homologs of human TPTE which are naturally expressed by cells or
15 are expressed by cells transfected with the TPTE gene.

The term "CXCR4" relates to "chemokine (C-X-C motif) receptor 4" and includes any variants, in particular splice variant, conformations, isoforms and species
20 homologs of CXCR4 which are naturally expressed by cells or are expressed by cells transfected with the CXCR4 gene. The expression "determining the level of expression of CXCR4" relates to a determination of the level of a nucleic acid of CXCR4, such as mRNA, and/or
25 a determination of the level of CXCR4 protein.

Preferably, a "nucleic acid of CXCR4", a "nucleic acid encoding CXCR4", a "nucleic acid coding for CXCR4" or "CXCR4 gene" relates to a nucleic acid selected from
30 the group consisting of (a) a nucleic acid which comprises a nucleic acid sequence selected from the group consisting of SEQ ID NOS: 47, and 48, a part or derivative thereof, (b) a nucleic acid which hybridizes with the nucleic acid of (a) under stringent
35 conditions, (c) a nucleic acid which is degenerate with

respect to the nucleic acid of (a) or (b), and (d) a nucleic acid which is complementary to the nucleic acid of (a), (b) or (c). The terms may also include mRNA coding for CXCR4. Preferably, "CXCR4" protein or simply
5 "CXCR4" comprises an amino acid sequence encoded by the afore mentioned nucleic acid, preferably an amino acid sequence selected from the group consisting of SEQ ID NOs: 49, and 50, a part or derivative thereof. One skilled in the art would understand that the cDNA
10 sequence of CXCR4 as described above would be equivalent to CXCR4 mRNA, and can be used for the same purpose herein; i.e., the generation of siRNA for inhibiting expression of CXCR4.

15 The term "CXCR4" also includes posttranslationally modified variants, isoforms and species homologs of human CXCR4 which are naturally expressed by cells or are expressed by cells transfected with the CXCR4 gene.

20 According to the invention, a nucleic acid is preferably deoxyribonucleic acid (DNA) or ribonucleic acid (RNA). Nucleic acids comprise according to the invention genomic DNA, cDNA, mRNA, recombinantly produced and chemically synthesized molecules.

25 According to the invention, a nucleic acid may be present as a single-stranded or double-stranded and linear or covalently circularly closed molecule.

A degenerate nucleic acid according to the invention is
30 a nucleic acid that differs from a reference nucleic acid in codon sequence due to the degeneracy of the genetic code.

The term "nucleic acid" according to the invention also
35 includes "derivatives" of a nucleic acid. "Derivative"

of a nucleic acid means according to the invention that single or multiple, such as at least 2, at least 4, or at least 6 and preferably up to 3, up to 4, up to 5, up to 6, up to 10, up to 15, or up to 20 nucleotide
5 substitutions, deletions and/or additions are present in said nucleic acid. Furthermore, the term "derivative" also comprises chemical derivatization of a nucleic acid on a nucleotide base, on the sugar or on the phosphate. The term "derivative" also comprises
10 nucleic acids which contain nucleotides and nucleotide analogs not occurring naturally.

The nucleic acids described according to the invention have preferably been isolated. The term "isolated
15 nucleic acid" means according to the invention that the nucleic acid was (i) amplified *in vitro*, for example by polymerase chain reaction (PCR), (ii) recombinantly produced by cloning, (iii) purified, for example by cleavage and gel-electrophoretic fractionation, or (iv)
20 synthesized, for example by chemical synthesis. An isolated nucleic acid is a nucleic acid which is available for manipulation by recombinant DNA techniques.

25 As used herein, the term "RNA" means a molecule comprising at least one ribonucleotide residue. By "ribonucleotide" is meant a nucleotide with a hydroxyl group at the 2'-position of a beta-D-ribo-furanose moiety. The term includes double stranded RNA, single
30 stranded RNA, isolated RNA such as partially purified RNA, essentially pure RNA, synthetic RNA, recombinantly produced RNA, as well as altered RNA that differs from naturally occurring RNA by the addition, deletion, substitution and/or alteration of one or more
35 nucleotides. Such alterations can include addition of

non-nucleotide material, such as to the end(s) of a RNA or internally, for example at one or more nucleotides of the RNA. Nucleotides in RNA molecules can also comprise non-standard nucleotides, such as non-
5 naturally occurring nucleotides or chemically synthesized nucleotides or deoxynucleotides. These altered RNAs can be referred to as analogs or analogs of naturally-occurring RNA.

10 As used herein, the terms "complementarity" or "complementary" means that a nucleic acid can form hydrogen bond(s) with another nucleic acid sequence by either traditional Watson-Crick or other non-traditional types of interaction. In reference to the
15 nucleic molecules described according to the present invention, the binding free energy for a nucleic acid molecule with its complementary sequence is sufficient to allow the relevant function of the nucleic acid to proceed, e.g., RNAi activity. For example, the degree
20 of complementarity between the sense and antisense strand of the siRNA construct can be the same or different from the degree of complementarity between the antisense strand of the siRNA and the target RNA sequence. A percent complementarity indicates the
25 percentage of contiguous residues in a nucleic acid molecule that can form hydrogen bonds (e.g., Watson-Crick base pairing) with a second nucleic acid sequence (e.g., 5, 6, 7, 8, 9, 10 out of 10 being 50%, 60%, 70%, 80%, 90%, and 100% complementary). "Perfectly
30 complementary" means that all the contiguous residues of a nucleic acid sequence will hydrogen bond with the same number of contiguous residues in a second nucleic acid sequence.

35 Preferably, a nucleic acid is "complementary" to

another nucleic acid if the two sequences are capable of hybridizing and forming a stable duplex with one another, with hybridization preferably being carried out under conditions which allow specific hybridization

5 between polynucleotides (stringent conditions). Stringent conditions are described, for example, in Molecular Cloning: A Laboratory Manual, J. Sambrook et al., Editors, 2nd Edition, Cold Spring Harbor Laboratory press, Cold Spring Harbor, New York, 1989 or

10 Current Protocols in Molecular Biology, F.M. Ausubel et al., Editors, John Wiley & Sons, Inc., New York and refer, for example, to hybridization at 65°C in hybridization buffer (3.5 x SSC, 0.02% Ficoll, 0.02% polyvinylpyrrolidone, 0.02% bovine serum albumin,

15 2.5 mM NaH₂PO₄ (pH 7), 0.5% SDS, 2 mM EDTA). SSC is 0.15 M sodium chloride/0.15 M sodium citrate, pH 7. After hybridization, the membrane to which the DNA has been transferred is washed, for example, in 2 x SSC at room temperature and then in 0.1-0.5 x SSC/0.1 x SDS at

20 temperatures of up to 68°C.

Preferably, the degree of complementarity according to the invention is at least 70%, preferably at least 75%, more preferably at least 80%, even more preferably at

25 least 90% or most preferably at least 95%, 96%, 97%, 98% or 99%.

Preferably the degree of identity between a specific nucleic acid sequence described herein and a nucleic

30 acid sequence which is a derivative of said specific nucleic acid sequence, which hybridizes with said specific nucleic acid sequence and/or which is degenerate with respect to said specific nucleic acid sequence will be at least 70%, preferably at least 75%,

35 preferably at least 80%, more preferably at least 85%,

even more preferably at least 90% or most preferably at least 95%, 96%, 97%, 98% or 99%. The degree of identity is preferably given for a region of at least about 30, at least about 50, at least about 70, at least about 90, at least about 100, at least about 200, at least about 300, at least about 400, at least about 500, at least about 1000, at least about 1500, or at least about 2000 nucleotides. In preferred embodiments, the degree of identity is given for the entire length of the reference nucleic acid sequence, such as the nucleic acid sequences given in the sequence listing.

"Sequence similarity" indicates the percentage of amino acids that either are identical or that represent conservative amino acid substitutions. "Sequence identity" between two polypeptide or nucleic acid sequences indicates the percentage of amino acids or nucleotides that are identical between the sequences.

The term "percentage identity" is intended to denote a percentage of nucleotides or of amino acid residues which are identical between the two sequences to be compared, obtained after the best alignment, this percentage being purely statistical and the differences between the two sequences being distributed randomly and over their entire length. Sequence comparisons between two nucleotide or amino acid sequences are conventionally carried out by comparing these sequences after having aligned them optimally, said comparison being carried out by segment or by "window of comparison" in order to identify and compare local regions of sequence similarity. The optimal alignment of the sequences for comparison may be produced, besides manually, by means of the local homology algorithm of Smith and Waterman, 1981, *Adv. App. Math.*

2, 482, by means of the local homology algorithm of Neddleman and Wunsch, 1970, J. Mol. Biol. 48, 443, by means of the similarity search method of Pearson and Lipman, 1988, Proc. Natl Acad. Sci. USA 85, 2444, or by
5 means of computer programs which use these algorithms (GAP, BESTFIT, FASTA, BLAST P, BLAST N and TFASTA in Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Drive, Madison, Wis.).

10 The percentage identity is calculated by determining the number of identical positions between the two sequences being compared, dividing this number by the number of positions compared and multiplying the result obtained by 100 so as to obtain the percentage identity
15 between these two sequences.

As used herein, a nucleic acid sequence "substantially identical" to a target sequence, e.g., a target sequence contained within a target mRNA, is a nucleic
20 acid sequence which is identical to the target sequence, or which differs from the target sequence by one or more nucleotides. Sense strands of siRNAs described herein which comprise nucleic acid sequences substantially identical to a target sequence are
25 characterized in that siRNA comprising such sense strands induce RNAi-mediated degradation of mRNA containing the target sequence. For example, an siRNA can comprise a sense strand which differs from a target sequence by one, two or three or more nucleotides, as
30 long as RNAi-mediated degradation of the target mRNA is induced by the siRNA.

Nucleic acids may, according to the invention, be present alone or in combination with other nucleic
35 acids, which may be homologous or heterologous. In

preferred embodiments, a nucleic acid is functionally linked to expression control sequences which may be homologous or heterologous with respect to said nucleic acid. The term "homologous" means that a nucleic acid is also functionally linked to the expression control sequence naturally and the term "heterologous" means that a nucleic acid is not functionally linked to the expression control sequence naturally.

10 A nucleic acid, such as a nucleic acid expressing RNA and/or protein or peptide, and an expression control sequence are "functionally" linked to one another, if they are covalently linked to one another in such a way that expression or transcription of said nucleic acid is under the control or under the influence of said expression control sequence. If the nucleic acid is to be translated into a functional protein, then, with an expression control sequence functionally linked to a coding sequence, induction of said expression control sequence results in transcription of said nucleic acid, without causing a frame shift in the coding sequence or said coding sequence not being capable of being translated into the desired protein or peptide.

25 The term "expression control sequence" comprises according to the invention promoters, ribosome binding sites, enhancers and other control elements which regulate transcription of a gene or translation of a mRNA. In particular embodiments of the invention, the expression control sequences can be regulated. The exact structure of expression control sequences may vary as a function of the species or cell type, but generally comprises 5'-untranscribed and 5'- and 3'-untranslated sequences which are involved in initiation of transcription and translation, respectively, such as

TATA box, capping sequence, CAAT sequence, and the like. More specifically, 5'-untranscribed expression control sequences comprise a promoter region which includes a promoter sequence for transcriptional
5 control of the functionally linked nucleic acid. Expression control sequences may also comprise enhancer sequences or upstream activator sequences.

According to the invention the term "promoter" or
10 "promoter region" relates to a nucleic acid sequence which is located upstream (5') to the nucleic acid sequence being expressed and controls expression of the sequence by providing a recognition and binding site for RNA-polymerase. The "promoter region" may include
15 further recognition and binding sites for further factors which are involved in the regulation of transcription of a gene. A promoter may control the transcription of a prokaryotic or eukaryotic gene. Furthermore, a promoter may be "inducible" and may
20 initiate transcription in response to an inducing agent or may be "constitutive" if transcription is not controlled by an inducing agent. A gene which is under the control of an inducible promoter is not expressed or only expressed to a small extent if an inducing
25 agent is absent. In the presence of the inducing agent the gene is switched on or the level of transcription is increased. This is mediated, in general, by binding of a specific transcription factor.

30 Promoters which are preferred according to the invention include promoters for SP6, T3 and T7 polymerase, human U6 RNA promoter, and CMV promoter.

According to the invention, the term "expression" is
35 used in its most general meaning and comprises the

production of RNA or of RNA and protein/peptide. It also comprises partial expression of nucleic acids. Furthermore, expression may be carried out transiently or stably.

5

In a preferred embodiment, a nucleic acid molecule is according to the invention present in a vector, where appropriate with a promoter, which controls expression of the nucleic acid. The term "vector" is used here in
10 its most general meaning and comprises any intermediary vehicle for a nucleic acid which enables said nucleic acid, for example, to be introduced into prokaryotic and/or eukaryotic cells and, where appropriate, to be integrated into a genome. Vectors of this kind are
15 preferably replicated and/or expressed in the cells. Vectors comprise plasmids, phagemids, bacteriophages or viral genomes. The term "plasmid" as used herein generally relates to a construct of extrachromosomal genetic material, usually a circular DNA duplex, which
20 can replicate independently of chromosomal DNA.

According to the invention, the term "host cell" relates to any cell which can be transformed or transfected with an exogenous nucleic acid. The term
25 "host cells" comprises according to the invention prokaryotic (e.g. *E. coli*) or eukaryotic cells (e.g. mammalian cells, in particular human cells, yeast cells and insect cells). Particular preference is given to mammalian cells such as cells from humans, mice,
30 hamsters, pigs, goats, or primates. The cells may be derived from a multiplicity of tissue types and comprise primary cells and cell lines. Specific examples comprise keratinocytes, peripheral blood leukocytes, stem cells of the bone marrow and embryonic
35 stem cells. In further embodiments, the host cell is an

antigen-presenting cell, in particular a dendritic cell, monocyte or a macrophage. A nucleic acid may be present in the host cell in the form of a single copy or of two or more copies and, in one embodiment, is
5 expressed in the host cell.

In those cases of the invention in which a MHC molecule presents a protein or peptide, an expression vector may also comprise a nucleic acid sequence coding for said
10 MHC molecule. The nucleic acid sequence coding for the MHC molecule may be present on the same expression vector as the nucleic acid coding for the protein or peptide, or both nucleic acids may be present on different expression vectors. In the latter case, the
15 two expression vectors may be cotransfected into a cell. If a host cell expresses neither the protein or peptide nor the MHC molecule, both nucleic acids coding therefor may be transfected into the cell either on the same expression vector or on different expression
20 vectors. If the cell already expresses the MHC molecule, only the nucleic acid sequence coding for the protein or peptide can be transfected into the cell.

A nucleic acid can be detected or its amount determined
25 by amplification of said nucleic acid. Amplification of a nucleic acid can be done using a pair of amplification primers, i.e. oligonucleotides, which hybridize to the nucleic acid. The primers preferably comprise a sequence of 6-50, in particular 10-30, 15-30
30 and 20-30 contiguous nucleotides of the nucleic acid and are nonoverlapping, in order to avoid the formation of primer dimers. One of the primers will hybridize to one strand of the nucleic acid to be amplified, and the other primer will hybridize to the complementary strand
35 in an arrangement which allows amplification of the

nucleic acid.

"Antisense molecules" or "antisense nucleic acids" may be used for regulating, in particular reducing, expression of a nucleic acid. The term "antisense molecule" or "antisense nucleic acid" refers according to the invention to an oligonucleotide which hybridizes under physiological conditions to DNA comprising a particular gene or to mRNA of said gene, thereby inhibiting transcription of said gene and/or translation of said mRNA. According to the invention, an "antisense molecule" also comprises a construct which contains a nucleic acid or a part thereof in reverse orientation with respect to its natural promoter. An antisense transcript of a nucleic acid or of a part thereof may form a duplex with the naturally occurring mRNA specifying the protein and thus prevent accumulation of or translation of the mRNA into the protein. Another possibility is the use of ribozymes for inactivating a nucleic acid. Antisense oligonucleotides preferred according to the invention have a sequence of 6-50, in particular 10-30, 15-30 and 20-30, contiguous nucleotides of the target nucleic acid and preferably are fully complementary to the target nucleic acid or to a part thereof.

In preferred embodiments, the antisense oligonucleotide hybridizes with an N-terminal or 5' upstream site such as a translation initiation site, transcription initiation site or promoter site. In further embodiments, the antisense oligonucleotide hybridizes with a 3'untranslated region or mRNA splicing site.

According to the invention an oligonucleotide may be an oligoribonucleotide, oligodeoxyribonucleotide, modified

- 37 -

oligoribonucleotide or modified oligo-
deoxyribonucleotide.

In one embodiment, an oligonucleotide consists of
5 ribonucleotides, deoxyribonucleotides or a combination
thereof, with the 5'-end of one nucleotide and the 3'-
end of another nucleotide being linked to one another
by a phosphodiester bond. These oligonucleotides may be
synthesized in the conventional manner or produced
10 recombinantly.

In preferred embodiments, an oligonucleotide is a
"modified" oligonucleotide. Here, the oligonucleotide
may be modified in very different ways, without
15 impairing its ability to bind its target, in order to
increase, for example, its stability. According to the
invention, the term "modified oligonucleotide" means an
oligonucleotide in which (i) at least two of its
nucleotides are linked to one another by a synthetic
20 internucleoside bond (i.e. an internucleoside bond
which is not a phosphodiester bond) and/or (ii) a
chemical group which is usually not found in nucleic
acids is covalently linked to the oligonucleotide.
Preferred synthetic internucleoside bonds are
25 phosphorothioates, alkyl phosphonates,
phosphorodithioates, phosphate esters, alkyl
phosphonothioates, phosphoramidates, carbamates,
carbonates, phosphate triesters, acetamides,
carboxymethyl esters and peptides.

30

The term "modified oligonucleotide" also comprises
oligonucleotides having one or more covalently modified
bases and/or one or more covalently modified sugars.
"Modified oligonucleotides" comprise, for example,
35 oligonucleotides with sugar residues which are

covalently bound to low molecular weight organic groups other than a hydroxyl group at the 3'-position and a phosphate group at the 5'-position. Modified oligonucleotides may comprise, for example, a 2'-O-alkylated ribose residue or another sugar instead of ribose, such as arabinose.

By "small interfering RNA" or "siRNA" as used herein is meant an isolated RNA molecule, preferably greater than 10 nucleotides in length, more preferably greater than 15 nucleotides in length, and most preferably 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, or 30 nucleotides in length that is used to identify a target gene or mRNA to be degraded. A range of 19-25 nucleotides is the most preferred size for siRNAs.

siRNA according to the invention can comprise partially purified RNA, substantially pure RNA, synthetic RNA, or recombinantly produced RNA, as well as altered RNA that differs from naturally-occurring RNA by the addition, deletion, substitution and/or alteration of one or more nucleotides. Such alterations can include addition of non-nucleotide material, such as to the end(s) of the siRNA or to one or more internal nucleotides of the siRNA; modifications that make the siRNA resistant to nuclease digestion (e. g., the use of 2'-substituted ribonucleotides or modifications to the sugar-phosphate backbone); or the substitution of one or more nucleotides in the siRNA with deoxyribonucleotides. Furthermore, siRNA may be modified to increase the stability thereof as described above for modified oligonucleotides, in particular by introducing one or more phosphorothioate linkages.

One or both strands of the siRNA can also comprise a 3'-overhang. As used herein, a "3'-overhang" refers to at least one unpaired nucleotide extending from the 3'-end of an RNA strand. Thus in one embodiment, the siRNA comprises at least one 3'-overhang of from 1 to about 6 nucleotides (which includes ribonucleotides or deoxynucleotides) in length, preferably from 1 to about 5 nucleotides in length, more preferably from 1 to about 4 nucleotides in length, and particularly preferably from about 2 to about 4 nucleotides in length. In the embodiment in which both strands of the siRNA molecule comprise a 3'-overhang, the length of the overhangs can be the same or different for each strand. In a most preferred embodiment, the 3'-overhang is present on both strands of the siRNA, and is 2 nucleotides in length. For example, each strand of the siRNA of the invention can comprise 3'-overhangs of dideoxythymidylic acid ("TT") or diuridylic acid ("uu").

20

In order to enhance the stability of the siRNA, the 3'-overhangs can be also stabilized against degradation. In one embodiment, the overhangs are stabilized by including purine nucleotides, such as adenosine or guanosine nucleotides. Alternatively, substitution of pyrimidine nucleotides by modified analogues, e.g., substitution of uridine nucleotides in the 3'-overhangs with 2'-deoxythymidine, is tolerated and does not affect the efficiency of RNAi degradation. In particular, the absence of a 2'-hydroxyl in the 2'-deoxythymidine significantly enhances the nuclease resistance of the 3'-overhang in tissue culture medium.

The sense and antisense strands of the siRNA can comprise two complementary, single-stranded RNA

35

molecules or can comprise a single molecule in which two complementary portions are base-paired and are covalently linked by a single-stranded "hairpin" area. That is, the sense region and antisense region can be covalently connected via a linker molecule. The linker molecule can be a polynucleotide or non-nucleotide linker. Without wishing to be bound by any theory, it is believed that the hairpin area of the latter type of siRNA molecule is cleaved intracellularly by the "Dicer" protein (or its equivalent) to form a siRNA of two individual base-paired RNA molecules.

As used herein, "target mRNA" refers to an RNA molecule that is a target for downregulation. One skilled in the art would understand that the cDNA sequence is equivalent to the mRNA sequence, and can be used for the same purpose herein, i.e., the generation of siRNA.

As used herein, a gene or mRNA which is "cognate" to human TPTE or CXCR4 is a gene or mRNA from another mammalian species which is homologous to human TPTE or CXCR4.

The mRNA transcribed from the human TPTE or CXCR4 gene can be analyzed for alternative splice forms using techniques well-known in the art. Such techniques include reverse transcription-polymerase chain reaction (RT-PCR), northern blotting and in-situ hybridization.

A technique called "RNase protection" can also be used to identify alternatively spliced TPTE or CXCR4 mRNAs. RNase protection involves transcription of a gene sequence into synthetic RNA, which is hybridized to RNA derived from other cells; for example, cells which are induced to express TPTE. The hybridized RNA is then

- 41 -

incubated with enzymes that recognize RNA:RNA hybrid mismatches. Smaller than expected fragments indicate the presence of alternatively spliced mRNAs. The putative alternatively spliced mRNAs can be cloned and
5 sequenced by methods well known to those skilled in the art.

RT-PCR can also be used to identify alternatively spliced TPTE or CXCR4 mRNAs. In RT-PCR, mRNA from cells
10 known to express TPTE or CXCR4 is converted into cDNA by the enzyme reverse transcriptase, using methods well-known to those of ordinary skill in the art. The entire coding sequence of the cDNA is then amplified via PCR using a forward primer located in the 3'
15 untranslated region, and a reverse primer located in the 5' untranslated region. The amplified products can be analyzed for alternative splice forms, for example by comparing the size of the amplified products with the size of the expected product from normally spliced
20 mRNA, e.g., by agarose gel electrophoresis. Any change in the size of the amplified product can indicate alternative splicing.

The mRNA produced from mutant TPTE or CXCR4 genes can
25 also be readily identified with the techniques described above for identifying alternative splice forms. As used herein, "mutant" TPTE or CXCR4 genes or mRNA include human TPTE or CXCR4 genes or mRNA which differ in sequence from the TPTE or CXCR4 sequences set
30 forth herein. Thus, allelic forms of the TPTE or CXCR4 gene, and the mRNA produced from them, are considered "mutants" for purposes of this invention.

"Reduce" or "inhibit" as used herein means the ability
35 to cause an overall decrease, preferably of 20% or

greater, more preferably of 50% or greater, and most preferably of 75% or greater, in the level, e.g. in the level of protein or mRNA as compared to a reference sample (e.g., a sample not treated with siRNA). This
5 reduction or inhibition of RNA or protein expression can occur through targeted mRNA cleavage or degradation. Assays for protein expression or nucleic acid expression are known in the art and include, for example, ELISA, western blot analysis for protein
10 expression, and northern blotting or RNase protection assays for RNA.

siRNA can be expressed from pol III expression vectors without a change in targeting site, as expression of
15 RNAs from pol III promoters is only believed to be efficient when the first transcribed nucleotide is a purine.

siRNA according to the invention can be targeted to any
20 stretch of approximately 19-25 contiguous nucleotides in any of the target mRNA sequences (the "target sequence"). Techniques for selecting target sequences for siRNA are given, for example, in Tuschl T. et al., "The siRNA User Guide", revised Oct. 11, 2002, the
25 entire disclosure of which is herein incorporated by reference. "The siRNA User Guide" is available on the world wide web at a website maintained by Dr. Thomas Tuschl, Laboratory of RNA Molecular Biology, Rockefeller University, New York, USA, and can be found
30 by accessing the website of the Rockefeller University and searching with the keyword "siRNA". Thus, the sense strand of the present siRNA comprises a nucleotide sequence substantially identical to any contiguous stretch of about 19 to about 25 nucleotides in the
35 target mRNA.

Generally, a target sequence on the target mRNA can be selected from a given cDNA sequence corresponding to the target mRNA, preferably beginning 50 to 100 nt downstream (i.e., in the 3'-direction) from the start codon. The target sequence can, however, be located in the 5'- or 3'-untranslated regions, or in the region nearby the start codon. For example, a suitable target sequence in the TPTE cDNA sequence is selected from the following group of target sequences:

- (i) TCGGTACTTGATAACATTACA (SEQ ID NO: 15)
- (ii) CAGACTTGTGTTATTCTAGCA (SEQ ID NO: 18)
- (iii) CTGAAATATGTTCAACTGCAA (SEQ ID NO: 21)
- 15 (iv) CAGATTGGCAACCAAGACTAA (SEQ ID NO: 24)
- (v) AACCTGCCCACATGTTTCATAT (SEQ ID NO: 27)
- (vi) AATGACAGTCCACAGACAAGT (SEQ ID NO: 30)
- (vii) AAGCTGATAAGAAGGCGGGTT (SEQ ID NO: 33)

20 A preferred siRNA targeting the sequence (i), and which has 3'-overhangs on each strand (overhangs shown in bold), is:

gguacuugauaacauacaTT
25 **AG**ccaugaacuauuguaaugu

A preferred siRNA targeting the sequence (ii), and which has 3'-overhangs on each strand (overhangs shown in bold), is:

30 gacuuguguuauucuagca**TT**
GTcugaacacaauaagaucgu

- 44 -

A preferred siRNA targeting the sequence (iii), and which has 3'-overhangs on each strand (overhangs shown in bold), is:

5 gaaaauauguucaacugcaa**TT**
 GAcuuuauacaaguugacguu

A preferred siRNA targeting the sequence (iv), and which has 3'-overhangs on each strand (overhangs shown in bold), is:

10 gauuggcaaccaagacuaa**TT**
 GTcuaaccguugguucugauu

15 A preferred siRNA targeting the sequence (v), and which has 3'-overhangs on each strand (overhangs shown in bold), is:

 cccugccacauguucauau**TT**
20 **TT**gggacgguguacaaguaa

A preferred siRNA targeting the sequence (vi), and which has 3'-overhangs on each strand (overhangs shown in bold), is:

25 ugacaguccacagacaagu**TT**
 TTacugucaggugucuguuca

A preferred siRNA targeting the sequence (vii), and which has 3'-overhangs on each strand (overhangs shown in bold), is:

 gcugauaagaaggcggguu**TT**
30 **TT**cgacuaauucuuccgccc

35

In the above list, all deoxyribonucleotides in a nucleic acid sequence are represented by capital letters (e.g., deoxythymidine is "T"), and ribonucleotides in a nucleic acid sequence are represented by lower case letters (e.g., uridine is "u").

It is understood that the target sequences given herein are with reference to the human TPTE cDNA, and thus these sequences contain deoxythymidines represented by "T". One skilled in the art would understand that, in the actual target sequence of the TPTE mRNA, the deoxythymidines would be replaced by uridines ("u"). Likewise, a target sequence contained within an siRNA of the invention would also contain uridines in place of deoxythymidines.

siRNA can be obtained using a number of techniques known to those of skill in the art. For example, siRNA can be chemically synthesized or recombinantly produced using methods known in the art, such as the *Drosophila* in vitro system described in U.S. published application 2002/0086356 of Tuschl et al., the entire disclosure of which is herein incorporated by reference.

25

Preferably, siRNA is chemically synthesized using appropriately protected ribonucleoside phosphoramidites and a conventional DNA/RNA synthesizer. siRNA can be synthesized as two separate, complementary RNA molecules, or as a single RNA molecule with two complementary regions.

Alternatively, siRNA can also be expressed from recombinant circular or linear DNA plasmids using any suitable promoter. Suitable promoters for expressing

35

siRNA of the invention from a plasmid include, for example, the U6 or H1 RNA pol III promoter sequences and the cytomegalovirus promoter.

5 Selection of other suitable promoters is within the skill in the art. The recombinant plasmids of the invention can also comprise inducible or regulatable promoters for expression of the siRNA in a particular tissue or in a particular intracellular environment.

10

The siRNA expressed from recombinant plasmids can either be isolated from cultured cell expression systems by standard techniques, or can be expressed intracellularly. The use of recombinant plasmids to
15 deliver siRNA to cells in vivo is discussed in more detail below. siRNA can be expressed from a recombinant plasmid either as two separate, complementary RNA molecules, or as a single RNA molecule with two complementary regions.

20

Selection of plasmids suitable for expressing siRNA, methods for inserting nucleic acid sequences for expressing the siRNA into the plasmid, and methods of delivering the recombinant plasmid to the cells of
25 interest are within the skill in the art.

siRNA can also be expressed from recombinant viral vectors intracellularly in vivo. The recombinant viral vectors comprise sequences encoding the siRNA and any
30 suitable promoter for expressing the siRNA sequences. The recombinant viral vectors can also comprise inducible or regulatable promoters for expression of the siRNA in a particular tissue or in a particular intracellular environment. The use of recombinant viral
35 vectors to deliver siRNA to cells in vivo is discussed

in more detail below. siRNA can be expressed from a recombinant viral vector either as two separate, complementary RNA molecules, or as a single RNA molecule with two complementary regions.

5

According to the invention the term "peptide" refers to substances comprising two or more, preferably 3 or more, preferably 4 or more, preferably 6 or more, preferably 8 or more, preferably 10 or more, preferably
10 13 or more, preferably 16 more, preferably 20 or more and up to preferably 8, 10, 20, 30, 40 or 50, in particular 100 amino acids joined covalently by peptide bonds. The term "protein" refers to large peptides, preferably to peptides with more than 100 amino acid
15 residues, but in general the terms "peptides" and "proteins" are synonyms and are used in this application interchangeably.

Preferably, the proteins and peptides described
20 according to the invention have been isolated. The terms "isolated protein" or "isolated peptide" mean that the protein or peptide has been separated from its natural environment. An isolated protein or peptide may be in an essentially purified state. The term
25 "essentially purified" means that the protein or peptide is essentially free of other substances with which it is associated in nature or *in vivo*.

Such proteins and peptides may be used, for example, in
30 producing antibodies and in an immunological or diagnostic assay. Proteins and peptides described according to the invention may be isolated from biological samples such as tissue or cell homogenates and may also be expressed recombinantly in a
35 multiplicity of pro- or eukaryotic expression systems.

For the purposes of the present invention, "derivatives" of a protein or peptide or of an amino acid sequence comprise amino acid insertion variants, amino acid deletion variants and/or amino acid substitution variants.

Amino acid insertion variants comprise amino- and/or carboxy-terminal fusions and also insertions of single or two or more amino acids in a particular amino acid sequence. In the case of amino acid sequence variants having an insertion, one or more amino acid residues are inserted into a particular site in an amino acid sequence, although random insertion with appropriate screening of the resulting product is also possible.

Amino acid deletion variants are characterized by the removal of one or more amino acids from the sequence.

Amino acid substitution variants are characterized by at least one residue in the sequence being removed and another residue being inserted in its place. Preference is given to the modifications being in positions in the amino acid sequence which are not conserved between homologous proteins or peptides and/or to replacing amino acids with other ones having similar properties such as hydrophobicity, hydrophilicity, electronegativity, volume of the side chain and the like (conservative substitution). Conservative substitutions, for example, relate to the exchange of one amino acid with another amino acid listed below in the same group as the amino acid to be substituted:

1. small aliphatic, nonpolar or slightly polar residues: Ala, Ser, Thr (Pro, Gly)

2. negatively charged residues and their amides: Asn, Asp, Glu, Gln
3. positively charged residues: His, Arg, Lys
4. large aliphatic, nonpolar residues: Met, Leu, Ile,
5 Val (Cys)
5. large aromatic residues: Phe, Tyr, Trp.

Owing to their particular part in protein architecture, three residues are shown in brackets. Gly is the only
10 residue without a side chain and thus imparts flexibility to the chain. Pro has an unusual geometry which greatly restricts the chain. Cys can form a disulfide bridge.

15 Preferably the degree of similarity, preferably identity between a specific amino acid sequence described herein and an amino acid sequence which is a derivative of said specific amino acid sequence will be at least 70%, preferably at least 80%, even more
20 preferably at least 90% or most preferably at least 95%, 96%, 97%, 98% or 99%. The degree of similarity or identity is given preferably for a region of at least about 10, at least about 20, at least about 40, at least about 60, at least about 80, at least about 100,
25 at least about 150, at least about 200, at least about 300, at least about 400 or 500 amino acids. In preferred embodiments, the degree of similarity or identity is given for the entire length of the reference amino acid sequence.

30

The amino acid variants described above may be readily prepared with the aid of known peptide synthesis techniques such as, for example, by solid phase synthesis (Merrifield, 1964) and similar methods or by
35 recombinant DNA manipulation. The manipulation of DNA

sequences for preparing proteins and peptides having substitutions, insertions or deletions, is described in detail in Sambrook et al. (1989), for example.

- 5 According to the invention, "derivatives" of proteins and peptides also comprise single or multiple substitutions, deletions and/or additions of any molecules associated with the protein or peptide, such as carbohydrates, lipids and/or proteins or peptides.
- 10 The term "derivative" also extends to all functional chemical equivalents of said proteins and peptides which do not only contain amino acid constituents but also non-amino acid constituents such as sugar and phosphate structures and extends also to substances
- 15 containing bonds such ester, thioether and disulfide bonds.

According to the invention, a part or fragment of a protein or peptide preferably has a functional property

20 of the protein or peptide from which it has been derived. Such functional properties comprise the interaction with antibodies, and the interaction with other peptides or proteins. A particular property is the ability to form a complex with MHC molecules and,

25 where appropriate, generate an immune response, preferably by stimulating cytotoxic or T helper cells. A part or fragment of a protein or peptide preferably comprises a sequence of at least 6, in particular at least 8, at least 10, at least 12, at least 15, at

30 least 20, at least 30 and preferably up to 8, in particular up to 10, up to 12, up to 15, up to 20, up to 30 or up to 50, consecutive amino acids of the protein or peptide.

35 A part or a fragment of a nucleic acid coding for a

protein or peptide preferably relates to the part of the nucleic acid, which codes at least for the protein or peptide and/or for a part or a fragment of said protein or peptide, as defined above. A part or
5 fragment of a nucleic acid coding for a protein or peptide is preferably that part of the nucleic acid corresponding to the open reading frame.

Antisera which contain specific antibodies specifically
10 binding to the target protein can be prepared by various standard processes; see, for example, "Monoclonal Antibodies: A Practical Approach" by Philip Shepherd, Christopher Dean ISBN 0-19-963722-9; "Antibodies: A Laboratory Manual" by Ed Harlow, David
15 Lane, ISBN: 0879693142 and "Using Antibodies: A Laboratory Manual: Portable Protocol NO" by Edward Harlow, David Lane, Ed Harlow ISBN 0879695447. Thereby it is also possible to generate affine and specific antibodies which recognize complex membrane proteins in
20 their native form (Azorsa et al., J. Immunol. Methods 229: 35-48, 1999; Anderson et al., J. Immunol. 143: 1899-1904, 1989; Gardsvoll, J. Immunol. Methods 234: 107-116, 2000). This is in particular relevant for the preparation of antibodies which are to be used
25 therapeutically, but also for many diagnostic applications. In this respect, it is possible to immunize with the whole protein, with extracellular partial sequences as well as with cells which express the target molecule in physiologically folded form.

30

Monoclonal antibodies are traditionally prepared using the hybridoma technology. (for technical details see: "Monoclonal Antibodies: A Practical Approach" by Philip Shepherd, Christopher Dean ISBN 0-19-963722-9;
35 "Antibodies: A Laboratory Manual" by Ed Harlow, David

Lane ISBN: 0879693142; "Using Antibodies: A Laboratory Manual: Portable Protocol NO" by Edward Harlow, David Lane, Ed Harlow ISBN: 0879695447).

5 It is known that only a small part of an antibody molecule, the paratope, is involved in binding of the antibody to its epitope (cf. Clark, W.R. (1986), *The Experimental Foundations of Modern Immunology*, Wiley & Sons, Inc., New York; Roitt, I. (1991), *Essential*
10 *Immunology*, 7th Edition, Blackwell Scientific Publications, Oxford). The pFc' and Fc regions are, for example, effectors of the complement cascade but are not involved in antigen binding. An antibody from which the pFc' region has been enzymatically removed or which
15 has been produced without the pFc' region, referred to as F(ab')₂ fragment, carries both antigen binding sites of a complete antibody. Similarly, an antibody from which the Fc region has been enzymatically removed or which has been produced without said Fc region,
20 referred to as Fab fragment, carries one antigen binding site of an intact antibody molecule. Furthermore, Fab fragments consist of a covalently bound light chain of an antibody and part of the heavy chain of said antibody, referred to as Fd. The Fd
25 fragments are the main determinants of antibody specificity (a single Fd fragment can be associated with up to ten different light chains, without altering the specificity of the antibody) and Fd fragments, when isolated, retain the ability to bind to an epitope.

30

Located within the antigen-binding part of an antibody are complementary-determining regions (CDRs) which interact directly with the antigen epitope and framework regions (FRs) which maintain the tertiary
35 structure of the paratope. Both the Fd fragment of the

- 53 -

heavy chain and the light chain of IgG immunoglobulins contain four framework regions (FR1 to FR4) which are separated in each case by three complementary-determining regions (CDR1 to CDR3). The CDRs and, in particular, the CDR3 regions and, still more particularly, the CDR3 region of the heavy chain are responsible to a large extent for antibody specificity.

Non-CDR regions of a mammalian antibody are known to be able to be replaced by similar regions of antibodies with the same or a different specificity, with the specificity for the epitope of the original antibody being retained. This made possible the development of "humanized" antibodies in which nonhuman CDRs are covalently linked to human FR and/or Fc/pFc' regions to produce a functional antibody.

As another example, WO 92/04381 describes the production and use of humanized murine RSV antibodies in which at least part of the murine FR regions have been replaced with FR regions of a human origin. Antibodies of this kind, including fragments of intact antibodies with antigen-binding capability, are often referred to as "chimeric" antibodies.

25

According to the invention, the term "antibody" also includes $F(ab')_2$, Fab, Fv, and Fd fragments of antibodies, chimeric antibodies, in which the Fc and/or FR and/or CDR1 and/or CDR2 and/or light chain-CDR3 regions have been replaced with homologous human or nonhuman sequences, chimeric $F(ab')_2$ -fragment antibodies in which the FR and/or CDR1 and/or CDR2 and/or light chain-CDR3 regions have been replaced with homologous human or nonhuman sequences, chimeric Fab-fragment antibodies in which the FR and/or CDR1 and/or

CDR2 and/or light chain-CDR3 regions have been replaced with homologous human or nonhuman sequences, and chimeric Fd-fragment antibodies in which the FR and/or CDR1 and/or CDR2 regions have been replaced with
5 homologous human or nonhuman sequences. The term "antibody" also comprises "single-chain" antibodies.

Antibodies may also be coupled to specific diagnostic substances for displaying cells and tissues expressing
10 particular proteins or peptides.

Diagnostic substances include any label that functions to: (i) provide a detectable signal; (ii) interact with a second label to modify the detectable signal provided
15 by the first or second label, e.g. FRET (Fluorescence Resonance Energy Transfer); (iii) affect mobility, e.g. electrophoretic mobility, by charge, hydrophobicity, shape, or other physical parameters, or (iv) provide a capture moiety, e.g., affinity, antibody/antigen, or
20 ionic complexation. Suitable as label are structures, such as fluorescent labels, luminescent labels, chromophore labels, radioisotopic labels, isotopic labels, preferably stable isotopic labels, isobaric labels, enzyme labels, particle labels, in particular
25 metal particle labels, magnetic particle labels, polymer particle labels, small organic molecules such as biotin, ligands of receptors or binding molecules such as cell adhesion proteins or lectins, label-sequences comprising nucleic acids and/or amino acid
30 residues which can be detected by use of binding agents, etc. Diagnostic substances comprise, in a nonlimiting manner, barium sulfate, iocetamic acid, iopanoic acid, calcium ipodate, sodium diatrizoate, meglumine diatrizoate, metrizamide, sodium tyropanoate
35 and radio diagnostic, including positron emitters such

as fluorine-18 and carbon-11, gamma emitters such as iodine-123, technetium-99m, iodine-131 and indium-111, nuclides for nuclear magnetic resonance, such as fluorine and gadolinium.

5

Antibodies may also be coupled to specific therapeutic substances.

According to the invention, the term "therapeutic substance" means any molecule which may exert a therapeutic effect. According to the invention, a therapeutic substance is preferably selectively guided to a diseased cell and includes anticancer agents, radioactive iodine-labeled compounds, toxins, cytostatic or cytolytic drugs, etc. Anticancer agents comprise, for example, aminoglutethimide, azathioprine, bleomycin sulfate, busulfan, carmustine, chlorambucil, cisplatin, cyclophosphamide, cyclosporine, cytarabidine, dacarbazine, dactinomycin, daunorubin, doxorubicin, taxol, etoposide, fluorouracil, interferon- α , lomustine, mercaptopurine, methotrexate, mitotane, procarbazine HCl, thioguanine, vinblastine sulfate and vincristine sulfate. Other anticancer agents are described, for example, in Goodman and Gilman, "The Pharmacological Basis of Therapeutics", 8th Edition, 1990, McGraw-Hill, Inc., in particular Chapter 52 (Antineoplastic Agents (Paul Calabresi and Bruce A. Chabner). Toxins may be proteins such as pokeweed antiviral protein, cholera toxin, pertussis toxin, ricin, gelonin, abrin, diphtheria exotoxin or *Pseudomonas* exotoxin. Toxin residues may also be high energy-emitting radionuclides such as cobalt-60.

The term "major histocompatibility complex" or "MHC"

relates to a complex of genes present in all vertebrates. MHC proteins or molecules are involved in signaling between lymphocytes and antigen presenting cells in normal immune reactions by binding peptides and presenting them for recognition by T cell receptors (TCR). MHC molecules bind peptides within an intracellular processing compartment and present these peptides on the surface of antigen presenting cells for recognition by T cells. The human MHC region also termed HLA is located on chromosome 6 and includes the class I and class II region. In one preferred embodiment of all aspects of the invention an MHC molecule is an HLA molecule.

The term "patient" or "subject" means according to the invention a human being, a nonhuman primate or another animal, in particular a mammal such as a cow, horse, pig, sheep, goat, dog, cat or a rodent such as a mouse and rat. In a particularly preferred embodiment, the patient is a human being.

"Abnormal expression" means according to the invention that expression is altered, preferably increased, compared to the state in a healthy individual.

25

According to the invention the term "increased" or "increased amount" preferably refers to an increase by at least 10%, in particular at least 20%, at least 50% or at least 100%. The amount of a substance is also increased in a test sample such as a biological sample compared to a reference sample if it is detectable in the test sample but absent or not detectable in the reference sample.

35 According to the invention, the term "disease" refers

to any pathological state, including, in particular, cancer, wherein the term "cancer" according to the invention comprises leukemias, seminomas, melanomas, teratomas, lymphomas, neuroblastomas, gliomas, rectal
5 cancer, endometrial cancer, kidney cancer, adrenal cancer, thyroid cancer, blood cancer, skin cancer, cancer of the brain, cervical cancer, intestinal cancer, liver cancer, colon cancer, stomach cancer, intestine cancer, head and neck cancer,
10 gastrointestinal cancer, lymph node cancer, esophagus cancer, colorectal cancer, pancreas cancer, ear, nose and throat (ENT) cancer, breast cancer, prostate cancer, cancer of the uterus, ovarian cancer and lung cancer and the metastases thereof. Examples thereof are
15 lung carcinomas, mamma carcinomas, prostate carcinomas, colon carcinomas, renal cell carcinomas, cervical carcinomas, or metastases of the cancer types or tumors described above. The term cancer according to the invention also comprises cancer metastases.

20

By "tumor" is meant an abnormal group of cells or tissue that grows by a rapid, uncontrolled cellular proliferation and continues to grow after the stimuli that initiated the new growth cease. Tumors show
25 partial or complete lack of structural organization and functional coordination with the normal tissue, and usually form a distinct mass of tissue, which may be either benign or malignant.

30 By "metastasis" is meant the spread of cancer cells from its original site to another part of the body. The formation of metastasis is a very complex process and depends on detachment of malignant cells from the primary tumor, invasion of the extracellular matrix,
35 penetration of the endothelial basement membranes to

enter the body cavity and vessels, and then, after being transported by the blood, infiltration of target organs. Finally, the growth of a new tumor at the target site depends on angiogenesis. Tumor metastasis
5 often occurs even after the removal of the primary tumor because tumor cells or components may remain and develop metastatic potential. In one embodiment, the term "metastasis" according to the invention relates to "distant metastasis" which relates to a metastasis
10 which is remote from the primary tumor and the regional lymph node system.

The term "relapse" relates to the return of signs and symptoms of a disease after a patient has enjoyed a
15 remission, e.g. after therapy such as tumor resection, chemotherapy and/or radiation therapy. In particular, the term "relapse" relates to the reappearance of cancer after a disease-free period. For example, after treatment a patient with cancer went into remission
20 with no sign or symptom of the tumor, remained in remission for some time, but then suffered a relapse and has to be treated once again for cancer.

According to the invention, a biological sample may be
25 a tissue sample, including bodily fluids, and/or a cellular sample and may be obtained in the conventional manner such as by tissue biopsy, including punch biopsy, and by taking blood, bronchial aspirate, sputum, urine, feces or other body fluids. According to
30 the invention, the term "biological sample" also includes fractions of biological samples.

The terms "T cell" and "T lymphocyte" are used interchangeably herein and include T helper cells and
35 cytolytic T cells which comprise cytotoxic T cells.

The pharmaceutical compositions and methods of treatment described according to the invention may also be used for immunization or vaccination to prevent a disease described herein.

In the present methods, nucleic acids can be administered to the subject either as naked nucleic acids, in conjunction with a delivery reagent, or as a recombinant plasmid or viral vector which may express the nucleic acid. The invention also provides for administering nucleic acids in vivo by using target-controlled liposomes.

For example vectors derived from adenovirus (AV); adeno-associated virus (AAV); retroviruses (e. g, lentiviruses (LV), Rhabdoviruses, murine leukemia virus); herpes virus, and the like can be used. The tropism of the viral vectors can be modified by pseudotyping the vectors with envelope proteins or other surface antigens from other viruses, or by substituting different viral capsid proteins, as appropriate.

Liposomes can aid in the delivery of the nucleic acid to a particular tissue, such as tumor tissue, and can also increase the blood half-life of the nucleic acid. Liposomes suitable for use in the invention are formed from standard vesicle-forming lipids, which generally include neutral or negatively charged phospholipids and a sterol, such as cholesterol. The selection of lipids is generally guided by consideration of factors such as the desired liposome size and half-life of the liposomes in the blood stream. A variety of methods are known for preparing liposomes, for example as described

in Szoka et al. (1980), Ann. Rev. Biophys. Bioeng. 9 : 467; and U.S. Pat. Nos. 4,235,871, 4,501,728, 4,837,028, and 5,019,369, the entire disclosures of which are herein incorporated by reference.

5

In particular embodiments, preference is given to directing nucleic acids to particular cells. In such embodiments, a carrier used for administering a nucleic acid to a cell (e.g. a retrovirus or a liposome) may have a bound target control molecule. For example, a molecule such as an antibody specific for a surface membrane protein on the target cell or a ligand for a receptor on the target cell may be incorporated into or attached to the nucleic acid carrier. Preferred antibodies comprise antibodies which bind selectively an antigen associated with tumor cells. If administration of a nucleic acid via liposomes is desired, proteins binding to a surface membrane protein associated with endocytosis may be incorporated into the liposome formulation in order to make target control and/or uptake possible. Such proteins comprise capsid proteins or fragments thereof which are specific for a particular cell type, antibodies to proteins which are internalized, proteins addressing an intracellular site, and the like.

RNAi-mediated degradation of the target mRNA can be detected by measuring levels of the target mRNA or protein, using standard techniques for isolating and quantifying mRNA such as Northern blot or dot blotting techniques, or quantitative RT-PCR or protein such as ELISA or Western blot.

The therapeutic compositions of the invention may be administered in pharmaceutically compatible

preparations. Such preparations may usually contain pharmaceutically compatible concentrations of salts, buffer substances, preservatives, carriers, supplementing immunity-enhancing substances such as
5 adjuvants, e.g. CpG oligonucleotides, cytokines, chemokines, saponin, GM-CSF and/or RNA and, where appropriate, other therapeutically active compounds.

The therapeutically active compounds of the invention
10 may be administered via any conventional route, including by injection or infusion. The administration may be carried out, for example, orally, intravenously, intraperitoneally, intramuscularly, subcutaneously or transdermally.

15 Suitable techniques for delivering nucleic acids to cells include administration of the nucleic acid to a subject by gene gun, electroporation, nanoparticles, micro-encapsulation, and the like, or by parenteral and
20 enteral administration routes.

Suitable enteral administration routes include oral, rectal, or intranasal delivery.

25 Suitable parenteral administration routes include intravascular administration (e. g. intravenous bolus injection, intravenous infusion, intra-arterial bolus injection, intra-arterial infusion and catheter instillation into the vasculature); peri- and intra-
30 tissue administration (e. g., peri-tumoral and intra-tumoral injection); subcutaneous injection or deposition including subcutaneous infusion (such as by osmotic pumps); direct (e.g., topical) application at or near the site of the diseased area, for example by a
35 catheter or other placement device (e.g., a

suppository, or an implant comprising a porous, non-porous, or gelatinous material); and inhalation.

The compositions of the invention are administered in effective amounts. An "effective amount" refers to the amount which achieves a desired reaction or a desired effect alone or together with further doses. In the case of treatment of a particular disease or of a particular condition, the desired reaction preferably relates to inhibition of the course of the disease. This comprises slowing down the progress of the disease and, in particular, interrupting or reversing the progress of the disease. The desired reaction in a treatment of a disease or of a condition may also be delay of the onset or a prevention of the onset of said disease or said condition. As used herein, an "effective amount" of the siRNA is preferably an amount sufficient to cause RNAi-mediated degradation of the target mRNA in a subject.

20

An effective amount of a composition of the invention will depend on feelers such as the condition to be treated, the severeness of the disease, the individual parameters of the patient, including age, physiological condition, size and weight, the duration of treatment, the type of an accompanying therapy (if present), the specific route of administration and similar factors.

The compositions of the invention can be administered to a subject in combination with another therapeutic method designed to treat the pathology. For example, they can be administered in combination with therapeutic methods currently employed for treating cancer or preventing tumor metastasis (e. g., radiation therapy, chemotherapy, and surgery). For treating

tumors, the compositions of the invention are preferably administered to a subject in combination with radiation therapy, or in combination with chemotherapeutic agents such as cisplatin, carboplatin, cyclophosphamide, 5-fluorouracil, adriamycin, daunorubicin or tamoxifen.

The pharmaceutical compositions of the invention are preferably sterile and contain an effective amount of the therapeutically active substance to generate the desired reaction or the desired effect.

The doses administered of the compositions of the invention may depend on various parameters such as the type of administration, the condition of the patient, the desired period of administration, etc. In the case that a reaction in a patient is insufficient with an initial dose, higher doses (or effectively higher doses achieved by a different, more localized route of administration) may be used.

The pharmaceutical compositions of the invention are generally administered in pharmaceutically compatible amounts and in pharmaceutically compatible compositions. The term "pharmaceutically compatible" refers to a nontoxic material which does not interact with the action of the active component of the pharmaceutical composition. Preparations of this kind may usually contain salts, buffer substances, preservatives, carriers and, where appropriate, other therapeutically active compounds. When used in medicine, the salts should be pharmaceutically compatible. However, salts which are not pharmaceutically compatible may be used for preparing pharmaceutically compatible salts and are included in

the invention. Pharmacologically and pharmaceutically compatible salts of this kind comprise in a nonlimiting way those prepared from the following acids: hydrochloric, hydrobromic, sulfuric, nitric, phosphoric, maleic, acetic, salicylic, citric, formic, malonic, succinic acids, and the like. Pharmaceutically compatible salts may also be prepared as alkali metal salts or alkaline earth metal salts, such as sodium salts, potassium salts or calcium salts.

10

A pharmaceutical composition of the invention may comprise a pharmaceutically compatible carrier. According to the invention, the term "pharmaceutically compatible carrier" refers to one or more compatible solid or liquid fillers, diluents or encapsulating substances, which are suitable for administration to humans. The term "carrier" refers to an organic or inorganic component, of a natural or synthetic nature, in which the active component is combined in order to facilitate application. The components of the pharmaceutical composition of the invention are usually such that no interaction occurs which substantially impairs the desired pharmaceutical efficacy.

25 The pharmaceutical compositions of the invention may contain suitable buffer substances such as acetic acid in a salt, citric acid in a salt, boric acid in a salt and phosphoric acid in a salt.

30 The pharmaceutical compositions may, where appropriate, also contain suitable preservatives such as benzalkonium chloride, chlorobutanol, paraben and thimerosal.

The pharmaceutical compositions are usually provided in a uniform dosage form and may be prepared in a manner known per se. Pharmaceutical compositions of the invention may be in the form of capsules, tablets, 5 lozenges, solutions, suspensions, syrups, elixirs or in the form of an emulsion, for example.

Compositions suitable for parenteral administration usually comprise a sterile aqueous or nonaqueous 10 preparation of the active compound, which is preferably isotonic to the blood of the recipient. Examples of compatible carriers and solvents are Ringer solution and isotonic sodium chloride solution. In addition, usually sterile, fixed oils are used as solution or 15 suspension medium.

The present invention is described in detail by the figures and examples below, which are used only for illustration purposes and are not meant to be limiting. 20 Owing to the description and the examples, further embodiments which are likewise included in the invention are accessible to the skilled worker.

25 Figures:

Figure 1. Selective transcription of TPTE in malignant tissues and cancer cell lines.

a, b, Analysis of TPTE mRNA expression by (a) 30 conventional RT-PCR and (b) quantitative Real-Time RT-PCR in normal human tissues, TPTE positive tumor specimens and cancer cell lines. c, Western blot analysis of protein lysates from normal tissues and cancer cell lines with constitutive TPTE expression. 35 Controls were NIH3T3 cells transfected with TPTE cDNA

- 66 -

(+) or a control plasmid (-). **d**, Immunohistochemical staining of testis and malignant tissues for TPTE. Blocking with the recombinant protein fragment used for immunization (+) as compared to the buffer control (-)
5 confirmed specificity of the polyclonal antiserum pAK2091. **e**, (left) Induction of *TPTE* mRNA expression in TPTE negative BT-549 breast cancer cells treated with the methylation inhibitor dAC as shown by real time RT-PCR analysis. (right) Relative amounts of TPTE
10 transcripts in DNA methyltransferase knockout variants of HCT116 cells as compared to wild type cells.

Figure 2. TPTE is a phosphoinositol 3'-phosphatase localized at the plasmamembrane.

15 **a**, In vitro phosphatase assay with recombinant proteins using $PI(3,4,5)P_3$ and $PI(4,5)P_2$ as substrates. **b**, Colocalization of TPTE-eGFP fluorescence and pAK2091 staining for verification of the specificity of the polyclonal rabbit antiserum. **c**, Immunofluorescence
20 analysis of cancer cell lines displaying a constitutive expression of TPTE. **d**, Localization of endogenous TPTE in filo- and pseudopodia of PC-3 prostate cancer cells; arrows, accumulation of TPTE at the lateral margins of cell protrusions; asterisks, tips of protrusions are
25 free of TPTE. **e**, Colocalization of endogenously expressed TPTE in PC-3 cells with PIP_2 visualized by staining of PLC- $\delta 1$ -PH-eGFP transfected cells with anti-TPTE. **f**, Transfection of TPTE but not of the catalytically inactive mutant TPTE_{C338S} into NIH3T3-her2
30 cells reduces constitutive PIP_3 signaling and thus leads to cytoplasmic redistribution of AKT-PH-eGFP.

Figure 3. TPTE establishes a growth-factor dependent phenotype in cancer cells.

The influence of TPTE on AKT phosphorylation, cell proliferation and resistance to apoptosis induced by growth factor deprivation was analyzed in siRNA transfected TPTE positive tumor cell lines (a-d) as well as in transformed cells ectopically expressing TPTE or the catalytically inactive mutant TPTE_{C338S} and eGFP as controls (e-i). **a**, Western blot analysis of siRNA transfected cells. TPTE siRNA specifically suppresses the respective phosphatase. Note that PTEN protein levels are not affected by TPTE siRNA. **b**, suppression of TPTE by TPTE siRNA (+) but not scrambled siRNA duplexes (-) leads to an increase of cellular phospho-AKT levels. **c**, **d**, As documented by proliferation rates (**c**) and apoptotic fraction (**d**) of MDA-MB-435 breast cancer and MelJuso melanoma cells cultured for 48h in medium supplemented with various concentrations of serum, TPTE downregulation uncouples cell proliferation and survival from dependency on external growth factors. Similar data were obtained for MDA-MB-231 and PC-3 cells. RFU stands for relative fluorescence unit. **e**, Western blot analysis of HER-2/neu expression and AKT phosphorylation in wild type and HER2/neu transformed NIH3T3 fibroblasts (NIH3T3-her2). **f**, TPTE but not a catalytically inactive variant reduces cellular PIP₃ levels in Her2/neu transformed fibroblasts. Cellular PIP₃ levels were quantified from lysates of serum deprived cells as described in example 1 using a PIP₃ specific ELISA. **g**, AKT phosphorylation in NIH3T3-her2 cells transfected with TPTE-eGFP and controls. **h**, Stable expression of catalytically active TPTE-eGFP abrogates autonomous growth of NIH3T3-her2 cells as determined by flowcytometric cell cycle analysis. **i**, tumor growth kinetics after s.c.

inoculation of stably transfected NIH3T3-her2 cells in immunocompromised mice.

Figure 4. TPTE promotes cell chemotaxis.

5 **a**, Transwell migration assay using PDGF-BB (200ng/ml) and SDF-1 α /CXCL12* (200ng/ml) as chemoattractants 48h after transfection of cells with siRNA oligos; (*MDA-MB-435 cells do not express CXCR4, the receptor for CXCL12). **b, c** Chemotaxis of NIH-3T3-her2 transfectants
10 analyzed in transwell migration assay using FCS or various concentrations of PDGF-BB as chemoattractants. **d**, Morphological characteristics of serum-free cultured NIH3T3-her2 cells stably transfected with either TPTE-eGFP or control vectors. **e, f, g** impact of single and
15 combined siRNA mediated knock down of PTEN and TPTE on tumor cell chemotaxis, cellular PIP₃ levels and pAKT levels (** note that PC-3 cells do not express PTEN).

Figure 5. TPTE is essential for metastatic spread of tumor cells.

20 **a**, In vivo tumor cell extravasation assay using breast cancer cells transfected with siRNA and labeled with a fluorophore. Cells extravasated into the lungs were documented by fluorescence microscopy and counted 6h
25 after injection. **b**, Experimental metastasis assay based on injection of TPTE siRNA or control siRNA treated breast cancer and melanoma cells into the tail vein of NOD/SCID (MDA-MB-231) or nude (MCF-7, MelJuso) mice. Metastatic tumor load was determined five weeks after
30 inoculation by quantitative PCR using oligonucleotides specific for human microsatellite DNA. **c**, Representative lungs and HE-stained lung tissue sections obtained by an independent experiment from nude mice four weeks after inoculation with siRNA
35 transfected MDA-MB-231 cells. Note that experiments

with PTEN siRNA transfected MDA-MB-231 cells resulted in similar reduction of metastatic tumor load. d, Venn diagram for illustration of metastasis rates in groups of patients. Patients were grouped according to the status of TPTE and CXCR4 expression in the primary tumor samples. The statistics illustrate the numbers of cases with metastasis versus the total number of patients in the respective group.

10 Example 1: Materials and Methods

Tissues and cell lines

This study was approved by the local ethical review board ("Ethikkommission der Ärztekammer des Landes Rheinland-Pfalz"). Recombinant DNA work was done with the official permission and according to the rules of the state government of Rheinland-Pfalz. Tissues were obtained as human surplus materials during routine diagnostic or therapeutic procedures and were stored at -80 °C until use. If not otherwise stated, cell lines were obtained from commercial providers. For demethylation studies cells were split to 20-30% confluency and cultured with 2µM or 10µM 5-Aza-2'-deoxycytidine (5-Aza-dC) (Sigma-Aldrich) for 72h. Colon cancer cell lines HCT116^{WT}, HCT116^{DNMT1-/-}, HCT116^{DNMT3b-/-} and HCT116^{DKO} were kindly provided by Bert Vogelstein.

RNA-Isolation, RT-PCR and real-Time RT-PCR

RNA extraction, first-strand cDNA synthesis, RT-PCR and real-time RT-PCR were performed as previously described (Koslowski, M. et al., Cancer Res. 62, 6750-6755 (2002), Koslowski, M. et al., Cancer Res. 64, 5988-5993 (2004)). All homologous members and pseudogenes have been aligned, to design specific primer pairs for TPTE.

- 70 -

Specificity was confirmed by sequencing of randomly selected amplification products. For end-point analysis TPTE-specific oligonucleotides (sense 5'-TGG ATG TCA CTC TCA TCC TTG-3'; antisense 5'-CCA TAG TTC CTG TTC TAT CTG-3', 63°C annealing) were used in a 35 cycle RT-PCR. Real-time quantitative expression analysis was performed in triplicates using TPTE-specific oligonucleotides (sense 5'-GAG TCT ACA ATC TAT GCA GTG-3'; antisense 5'-CCA TAG TTC CTG TTC TAT CTG-3', 63°C annealing) in a 40 cycle PCR. After normalization to 18sRNA (sense 5'-CGA TGC TCT TAG CTG AGT GTC-3'; antisense 5'-TAA CCA GAC AAA TCG CTC CAC-3', 65°C annealing) TPTE transcripts in tumor samples were quantified relative to normal tissues using $\Delta\Delta CT$ calculation.

Antisera, immunochemistry and Western blot

Polyclonal antiserum pAK2091 raised against the n-terminus (aa 1-51) of TPTE was generated by a custom antibody service (SeqLab). Immunohistochemistry was performed on formalin-fixed and paraffin-embedded tissue sections after antigen retrieval by boiling the slides for 15 minutes in citrate buffer (pH 6) followed by a cooling period of 15 minutes at room temperature. For Western blot analysis 60 μ g of total protein extracted from cells lysed with Triton-X was used. Extracts were diluted in reducing sample buffer (Roth), subjected to SDS-PAGE and subsequently electrotransferred onto PVDF membrane (Pall). For immunostaining antibodies reactive to HER2/neu (Abcam), pAKT (Cell Signaling), AKT (Cell Signaling) and beta-Actin (Abcam) were used followed by horseradish-peroxidase conjugated goat anti-mouse and goat anti-rabbit secondary antibodies (Dako). Detection of the

- 71 -

anti-TPTE pAK2091 primary antibody was performed using the anti-rabbit Envision+ System (Dako) according to the manufacturer's instructions.

5 Expression of eGFP-tagged TPTE in eukaryotic cells

The open reading frame of TPTE was amplified (sense primer 5'-GAG AGA AAG CTT CCA CCA TGA ATG AAA GTC CTG ATC CCA CTG ACC T-3', antisense primer 5'-GAG AGA AAG CTT GAT CGG ATC CAG CTA CAA CAT CAC TGC AAG TC-3')
10 introducing two HindIII sites. The amplified fragment was ligated into vectors pEGFP-C1 and pEGFP-N3 (BD Biosciences). Variant TPTE_{C338S} carrying a mutation in the active site of the phosphatase domain was generated by PCR-mediated site directed mutagenesis.

15

Immunofluorescence and colocalization studies

Cells expressing TPTE either endogenously or heterologously upon transfection with vector constructs were grown on slides for 12-24 h and were fixed with 2%
20 paraformaldehyde/0.1% saponin/PBS. Indirect immunofluorescence staining for TPTE was performed with pAK2091 polyclonal rabbit antiserum and a fluorescence-tagged secondary anti-rabbit IgG antibody. To analyze colocalization of TPTE with F-Actin we stained
25 permeabilised, fixed cells with rhodamine-phalloidin (Molecular Probes). Expression plasmids for eGFP-tagged PH-domains of PLC- δ 1 and AKT for visualization of membrane bound PIP₂ and PIP₃ by protein-protein interaction were kindly provided by Mario. J. Rebecchi
30 and Julian Downward, respectively. Coverslips were mounted on slides in Slow-Fade (Molecular Probes) and analysed by immunofluorescence microscopy.

35

Immunopurification of eGFP-fusion proteins

Cells expressing eGFP-fusion proteins were lysed in buffer containing 1% Triton X-100 and protease inhibitors (8 μ M Leupeptin, 3.3 μ M Chymostatin, 2.9 μ M⁵ Pepstatin A, 1 mM AEBSF-Hydrochloride). Lysates were spin down for 5 min at 4°C. For preclearing the lysates were incubated with protein A Sepharose CL-4B (Sigma-Aldrich) for 1 h at 4°C. The precleared lysates were incubated with anti-eGFP antibody (Delta Biolabs) for 2¹⁰ h at 4°C followed by incubation with protein A Sepharose CL-4B for 1 h at 4°C and precipitated by centrifugation for 2 min. Immune complexes were washed with IP buffer (50 mM HEPES (pH 7.5), 150 mM NaCl), and resuspended in reaction buffer (100 mM HEPES (pH 7.5),¹⁵ 150 mM NaCl, 10 mM DTT). Proteins were separated by SDS-PAGE and analyzed by immunoblotting.

In vitro phosphatase assay

To measure phosphatase activity PTEN and TPTE fusion²⁰ proteins were immunoprecipitated and incubated in reaction buffer containing 110 μ M water-soluble phosphatidylinositol phosphate (Echelon) for 90 min at 37°C. The amount of phosphate released from the substrate was determined using a malachite green assay²⁵ (Echelon). After 15 min colour development the absorbance of the samples was measured at 620 nm on a Tecan Safire reader. Each sample was analyzed as triplicate.

30 siRNA Duplexes

siRNA duplexes were designed following common rules and were purchased from Ambion. The TPTE siRNA duplex (sense 5'-r(GGU ACU UGA UAA CAU UAC A)dTdT-3', antisense 5'-r(UGU AAU GUU AUC AAG UAC C)dGdA-3')³⁵ targeted nucleotides 1722-1742 of the TPTE mRNA

sequence (NM_013315). As control a scrambled siRNA duplex (sense 5'-r(UAA CUG UAU AAU CGA CUA G)dTdT-5', antisense 5'-r(CUA GUC GAU UAU ACA GUU A)dGdA-3') was used. For TPTE silencing studies cells were transfected with 100nm siRNA duplex using RNAiFect transfection reagent (Qiagen) according to the manufacturer's instructions. All functional assays were conducted 24h after transfection with siRNA duplexes. All results were reproduced with a second set of TPTE siRNA duplexes (sense 5'-r(GAU UGG CAA CCA AGA CUA A)dTdT-3', antisense 5'-r(UAA GUC UUG GUU GCC AAU C)dTdG-3') targeting nucleotides 2487-2505. Duplexes for PTEN silencing (5'-r(GGC GUA UAC AGG AAC AAU A)dTdT-3', antisense 5'-r(UAU UGU UCC UGU AUA CGC C)dTdT-3') were directed against nucleotides 1161-1179 (NM_000314).

Cell migration

Cell migration assays were conducted in transwell chambers with 8.0 μ m pore membranes (BD Biosciences) with cells cultured in serum-free medium for 12h prior to the experiments. For siRNA experiments cells were transferred to serum-free conditions 24 h after transfection with siRNA duplexes as described above. 4×10^4 cells in 400 μ l serum-free culture medium were added to the upper chamber. The bottom chambers contained 800 μ l culture medium supplemented with either FCS, PDGF-BB (Sigma-Aldrich) or SDF-1 α /CXCL12 (R&D Systems) as chemoattractants. After 24 hours cells that had migrated to the bottom side of the membrane were fixed in ice-cold methanol; membranes were excised, placed on microscope slides and mounted with Hoechst (Dako) for fluorescence microscopy. Cells in five random visual fields (100x magnification) were counted for each membrane. All experiments were done in

triplicates. Effects on chemokinesis of cells was analyzed using the same experimental setup with (i) no chemoattractant added to the upper and lower chamber and (ii) with chemoattractant added to both the upper
5 and lower chamber.

Cell proliferation analysis

24 h after transfection with siRNA duplexes 1×10^4 cells were cultured in medium supplemented with varying
10 concentrations of FCS for 48 h. Proliferation was analyzed by measuring the incorporation of BrdU into newly synthesized DNA strands using the DELFIA cell proliferation Kit (Perkin Elmer) according to the manufacturer's instructions on a Wallac Victor2 multi-
15 label counter (Perkin Elmer).

Cell cycle analysis and apoptosis

Cells were cultured in medium supplemented with FCS in varying concentrations, harvested after 48 h and
20 stained with propidiumiodide prior to flowcytometric DNA content analysis. Apoptotic cells and cells in S/G2/M phases of the cell cycle were quantified using CellQuest-Software (Becton Dickinson).

25 In vivo tumor growth analysis and experimental metastasis assay

For analysis of in vivo tumor growth 5×10^6 cells (NIH3T3-her2, NIH3T3-her2-eGFP, NIH3T3-her2-TPTE-eGFP, and NIH3T3-her2-TPTE_{C338S}-eGFP) were injected
30 subcutaneously into the flanks of NOD/SCID mice (5 animals per group). Tumors were measured periodically with a caliper rule, and the tumor volume was calculated ($V = a \times b \times c / 2$). For assessment of tumor cell extravasation 1×10^6 cells labeled with CFSE (Vybrant

CFDA SE Cell Tracer Kit; Molecular probes) were injected into the tail vein of NOD/SCID mice (3 animals per group). Mice were sacrificed after 6 h and Hoechst 33258 labeled cryosections (20 μ M) of the lungs were
5 analyzed for extravasated tumor cells by fluorescence microscopy. Tumor cells in 50 random visual fields per lung were counted.

Real-time PCR was used for quantification of the tumor
10 load in the lungs of NOD/SCID mice (4 animals per group) five weeks after i.v. injection of 2×10^6 MDA-MB-231 cells. DNA was extracted using QIAamp DNA Mini Kit (Qiagen) and a 226 bp fragment of the alpha-satellite region of the human chromosome 17 (sense 5'-CAG CTG ACT
15 AAA CAG AAG CAG-3'; antisense 5'-GAG TTG AAT GCA GTC ATC ACA G-3') was amplified from 1 μ g DNA. The tumor load was quantified by referring to a standard curve generated by a serial dilution of MDA-MB-231 cells in NIH3T3 mouse fibroblasts.

20

Statistical Analysis

Statistical analysis of *TPTE* and *CXCR4* expression in tumors in relation to the metastatic rate of the patients was performed using SPSS software (Fisher's
25 exact test).

Example 2: *TPTE* is ectopically expressed in human tumors

TPTE mRNA expression was investigated in a large set of
30 normal and neoplastic tissue specimens. *TPTE* expression is confined to testis and transcript amounts are below detection limit of highly sensitive RT-PCR in all other normal tissue specimens (Fig. 1a,b). In contrast, strong *TPTE* expression was detected in 59 of 155 (38%)
35 tumor samples across different cancer types including

- 76 -

malignant melanoma (50%), breast carcinomas (47%) and lung carcinomas (55%) as well as in a large set of cancer cell lines (62%) (Tab. 1).

Table 1 Expression of *TPTE* in human tissues and cell lines analyzed by RT-PCR and Real-Time PCR.

Normal tissues	Positive/tested
Testis	3/3
Small intestine	0/2
Colon	0/3
Liver	0/2
Lung	0/3
Lymph node	0/2
Stomach	0/2
Spleen	0/2
Adrenal gland	0/1
Kidney	0/3
Esophagus	0/1
Ovary	0/2
Thymus	0/1
Skin	0/2
Breast	0/3
Pancreas	0/2
PBMC's, resting	0/3
PBMC's, proliferating	0/3
Prostate	0/2
Thyroid	0/2
Endometrium	0/3
Cerebellum	0/1
Brain	0/2
Tumour tissues	Positive/tested
Breast cancer	17/36 (47%)
Lung cancer	25/45 (55%)
Malignant melanoma	9/18 (50%)
Colon cancer	0/20
Prostate cancer	3/8
Ovarian cancer	2/7
Cervical cancer	1/6
Tumour cell lines	Positive/tested
Breast cancer cell lines	4/5
Lung cancer cell lines	2/6
Melanoma cell lines	5/8
Prostate cancer cell lines	2/2

- 77 -

Cloning and sequencing of amplification products from all cell lines tested and from arbitrarily selected tumor derived samples verified them as transcripts derived from TPTE on chromosome 21p11.

5

A polyclonal rabbit antibody (pAK2091) against the N-terminus (aa 1-51) of TPTE was used to verify expression data at the protein level. In accordance with the predicted size of TPTE, a 65 kDa band was
10 detected by Western blot analysis in testicular tissue, in a number of tumor cell lines typed positive for constitutive TPTE expression by RT-PCR, as well as in cells transfected with TPTE-cDNA confirming specificity of the antibody (Fig. 1c, left). In agreement with RT-
15 PCR data, normal somatic tissues scored negative in Western blot for TPTE, whereas TPTE RT-PCR positive cancer tissues contain significant amounts of TPTE protein (Figure 1c, right).

20 Immunohistochemistry with pAK2091 on testicular tissue showed specific immunoreactivity in type II spermatocytes and prespermatids in agreement with in situ hybridisation data described recently for the mouse orthologue (Wu, Y. et al., *J. Biol. Chem.* 276,
25 21745-21753 (2001)) (Fig. 1d). Tissue specimens obtained from cancers of lung, breast and prostate as well as malignant melanomas displayed a tumor cell-specific staining in immunohistochemistry. In contrast, adjacent stromal and non-neoplastic epithelial cells
30 (Fig. 1d) as well as patient matched normal tissues were not reactive (not shown). Having established TPTE as a molecular tumor marker, the mechanism responsible for its ectopic activation in cancer cells was investigated. DNA methylation at CpG-rich promoters has

been reported to be the primary mechanism for silencing of a subset of germline-specific genes in somatic tissues. Genomic demethylation, in turn, appears to be sufficient for aberrant activation of these genes in tumor cells (Koslowski, M. et al., *Cancer Res.* **64**, 5988-5993 (2004), De Smet, C. et al., *Mol. Cell Biol.* **19**, 7327-7335 (1999)). Sequence analysis of the *TPTE* promoter revealed a classical CpG island extending from upstream of the first exon over the first exon and intron. Due to the presence of nearly identical promoter sequences on chromosome 7 and chromosome 20, locus-specific bisulfite sequencing for direct analysis of the methylation status of the *TPTE* promoter in tumor cells could not be applied. Therefore, the effects of global methylation alterations on *TPTE* expression were studied. *TPTE* transcription was robustly induced upon treatment of several non-expressing cancer cell lines with the DNA methylation inhibitor 5-Aza-2'-deoxycytidine (dAC) (Fig. 1e). Methylation-dependent regulation of *TPTE* transcription was further evaluated in wild type HCT116 colon cancer cells and descendants with disrupted DNA methyltransferase (DNMT) genes. HCT116WT cells as well as the *DNMT3b*^{-/-} and *DNMT1*^{-/-} single knockout variants, which are known to display almost normal or only moderately reduced global DNA methylation (Rhee, I. et al., *Nature* **416**, 552-556 (2002)), do not express *TPTE*. In contrast, HCT116DKO cells lacking both methyltransferases and exhibiting vastly diminished overall DNA methylation resulted in a robust expression of *TPTE* (Fig. 1e). Both assays independently confirmed that DNA methylation is necessary for *TPTE* silencing and that genomic demethylation as frequently observed in tumors (Ehrlich, M., *Oncogene* **21**, 5400-5413 (2002),

Feinberg, A.P. & Vogelstein, B., *Nature* **301**, 89-92 (1983)) is sufficient for its activation.

Example 3: TPTE is a plasma membrane PIP₃-phosphatase

5 TPTE contains a phosphatase as well as a lipid-binding C2 domain, which have been shown to be essential and sufficient for the lipid phosphatase activity of its homologue PTEN (Lee, J.O. et al., *Cell* **99**, 323-334 (1999)). Whereas a lipid phosphatase activity with
10 substrate-specificity for PIP₃ and PI(3,4)P₂ has previously been shown for the mouse orthologue of TPTE (Wu, Y. et al., *J. Biol. Chem.* **276**, 21745-21753 (2001)) in vitro, no enzymatic activity was detected for the human counterpart (Walker, S.M. et al., *Biochem. J.* **360**,
15 277-283 (2001)), leaving PTEN as the only so far known human PIP₃-phosphatase. Since the latter study used recombinant protein of bacterial origin, enzymatic activity of human TPTE with eucaryotically produced protein was reassessed. The phosphatase and C2 domains
20 of TPTE and PTEN fused to eGFP were expressed in HEK-293 cells, the proteins purified by immunoprecipitation with anti-eGFP antibody coupled protein A beads and used in a malachite green assay. Equimolar amounts of immunoprecipitates obtained from cells transfected with
25 eGFP or with TPTE^{C338S}-eGFP, a TPTE variant mutated at a site critical for the putative phosphatase activity, served as controls to rule out contamination with copurifying phosphatases. Surprisingly, it was found that TPTE but not the respective controls release
30 phosphate specifically from PIP₃ at a rate comparable to PTEN (Fig. 2a). This finding together with the aberrant activation of TPTE in human cancers indicates that TPTE is involved in phosphoinositide-mediated plasma membrane signalling events in tumor cells.

TPTE-negative cells transfected with TPTE-eGFP as well as cancer cell lines, which express TPTE constitutively were stained with anti-TPTE antibody and investigated by immunofluorescence microscopy. In addition to the previously described localization in the Golgi apparatus and endoplasmatic reticulum (Wu,Y. et al., *J. Biol. Chem.* **276**, 21745-21753 (2001)), a major portion of TPTE was found at the plasma membrane (Fig. 2b, 2c). TPTE appears accentuated at membrane ruffles and at lateral margins of membrane protrusions including pseudopodia and filopodia, but not at the tips of such structures (Fig. 2d). To dissect the spatial association of TPTE with plasma membrane phosphoinositides colocalisation studies with pleckstrin domain-eGFP fusion proteins were performed using the PLC- δ 1-PH (phospholipase C- δ 1 pleckstrin homology) (Tall,E.G. et al., *Curr. Biol.* **10**, 743-746 (2000)) and the AKT-PH (Watton,S.J. & Downward,J., *Curr. Biol.* **9**, 433-436 (1999)) domains, which selectively bind to either PIP(4,5)P₂ (PIP₂) or 3'-phosphorylated phospholipids, respectively. Remarkably, staining of cells coexpressing TPTE cDNA and eGFP-tagged PH-domains with pAK2091 demonstrated an almost complete overlap of TPTE with PLC- δ 1-PH-eGFP (Fig. 2f) but not with AKT-PH-eGFP, establishing that TPTE colocalizes with PIP₂ (Fig. 2e).

A trafficking assay, which allows to indirectly determine membrane PIP₃ levels (Halet,G., *Biol. Cell* **97**, 501-518 (2005)), demonstrated that cotransfection of TPTE cDNA but not TPTE_{C338S}-cDNA together with AKT-PH-eGFP resulted in a complete redistribution of AKT-PH-eGFP from the plasma membrane to the cytosol in fibroblasts with PI3K overactivation due to HER-2/neu transformation (Fig. 2f), proving that TPTE decreases

plasma membrane PIP_3 levels. Altogether these observations prove that TPTE metabolizes PIP_3 and imply that TPTE may be involved in spatial control of plasma membrane phosphoinositides in tumor cells.

5

Example 4: Use of siRNA in silencing TPTE expression

The effects of small interfering RNA (siRNA) induced gene silencing of TPTE in breast cancer, prostate cancer and malignant melanoma cell lines, which
10 endogenously express the tumor-associated phosphatase TPTE were analyzed. Quantitative RT-PCR and Western blot demonstrated that TPTE specific siRNA duplexes induce a robust knockdown of TPTE transcripts and protein without affecting cellular PTEN levels (Fig.
15 3a).

First, levels of Ser473 phosphorylated AKT (pAKT) were quantified as a measure of cellular PIP_3 signalling. siRNA mediated downregulation of TPTE resulted in a
20 substantial upregulation of cellular pAKT in all tumor lines tested (Fig. 3b) establishing that TPTE counteracts PI3K signaling in cancer cells. Upregulation of pAKT by TPTE silencing was most prominent in PC-3 cells, which lack PTEN, suggesting
25 that endogenous TPTE may at least partially compensate for loss of PTEN activity in tumor cells. Most importantly, upregulation of pAKT upon TPTE silencing translates into a reduced growth factor-dependency of the respective tumor cells giving rise to sustained
30 proliferation rates (Fig. 3c) and protection from apoptosis even under serum starvation in all TPTE positive tumor lines tested (Fig. 3d).

In order to clarify the effects directly mediated by

the phosphatase activity of TPTE, Her2/neu transformed fibroblasts stably transfected with either TPTE or the catalytically inactive TPTE_{C338S} variant were used.

5 Her2/neu transformed fibroblasts (NIH3T3-her2) display permanent AKT activation (Fig. 3e) resulting from constitutively PI3K overactivation associated with increased cellular PIP₃ levels (Fig. 3f). As a consequence, these cells are resistant to apoptosis and
10 sustain proliferation under growth-factor starvation (Fig. 3h). Expression of TPTE but not mutated TPTE_{C338S} downregulates cellular PIP₃ (Fig. 3f), reduces pAKT levels (Fig. 3g), resets proliferation and survival autonomy and induces strictly serum-dependent
15 proliferation and rapid onset of a G0/G1 cell cycle block upon growth factor removal (Fig. 3h). Notably, growth of NIH3T3-her2 cells expressing TPTE in immunocompromised mice was markedly reduced compared to controls lacking the phosphatase activity, but were
20 still tumorigenic (Fig. 3g). These findings demonstrate that TPTE counteracts upstream oncogene-induced PI3K overactivation by metabolizing PIP₃ and renders tumor cell growth and survival dependent on external growth factors without abrogating tumorigenicity.

25

Example 5: TPTE promotes tumor cell chemotaxis

TPTE specific siRNA duplexes but not control duplexes reduced tumor cell migration towards PDGF or SDF-1/CXCL12 gradients in all tumor cell lines tested in
30 transwell migration assays and chemokine based invasion assays (Fig. 4a)

To exclude siRNA off target activity these findings were confirmed with a second set of TPTE specific siRNA

duplexes and controls. Moreover, it was observed that TPTE but not its catalytically inactive mutant variant potentiates HER-2/neu effects on cell migration. The increased baseline migration rate of NIH3T3-her2 cells (Dittmar, T. et al., *FASEB J.* 16, 1823-1825 (2002)) owing to transformation by this oncogene is further augmented upon coexpression of TPTE (Fig. 4b). Such double positive cells migrate efficiently even towards lowest gradients of chemoattractants (Fig. 4c) indicating that a combination of PI3K overactivation and TPTE expression promotes both chemokine sensing and efficient chemotactic migration. In line with this, expression of TPTE but not TPTE_{C338S}, results in profound morphological changes, i.e. the transition from a rounded cell shape to a polarized, polymorphic phenotype with pseudo- and filopodia (Fig. 4d). As shown for constitutively expressing cancer cell lines (Fig 2d), TPTE is strongly enriched in these protrusions suggesting that the lipid phosphatase is directly involved in the generation of filopodial extensions.

The chemotaxis promoting activity observed for TPTE is surprising in particular in light of previous data for PTEN, which exhibits the same catalytic PIP₃ phosphatase activity as TPTE, but was reported to inhibit migration (Tamura, M. et al., *Science* 280, 1614-1617 (1998)). However, these studies were based on tumor cells transfected with PTEN cDNA. In contrast, a recent report (Li, Z. et al., *Nat. Cell Biol.* 7, 399-404 (2005), Meili, R., Sasaki, A.T. & Firtel, R.A., *Nat. Cell Biol.* 7, 334-335 (2005)) which made use of siRNA for PTEN knockdown demonstrated clearly that PTEN is

essential for SDF-1 mediated chemotaxis in transformed Jurkat cells.

Robust reduction of PTEN expression by specific siRNA
5 duplexes resulted in marked and selective reduction of
chemotaxis (Fig. 4e) but not of chemokinesis in all
investigated PTEN positive tumor cell lines. The lack
of effects of PTEN siRNA in PTEN deficient PC-3 cells
ruled out that the observed inhibition on chemotactic
10 migration is mediated by siRNA off target activity.
Importantly, inhibition of both phosphatases resulted
in nearly complete abrogation of chemotaxis. Analysis
of cellular PIP_3 levels in siRNA treated cells
demonstrate that abrogation of both, TPTE and PTEN by
15 combined siRNA results in a more profound upregulation
of cellular PIP_3 as compared to the increase of PIP_3
levels in single siRNA treated cells (Fig. 4f). This
together with the observation, that inhibition of both
phosphatases resulted in a more intense increase of
20 cellular pAKT (Fig 4g) underlines that activities of
TPTE and PTEN are additive for promotion of tumor cell
chemotaxis and reduction of PIP_3 /AKT signaling.

Example 6: TPTE promotes metastatic spread

25 Chemotaxis mediated by growth factor receptors like EGF
and PDGF or chemokine receptors such as CXCR4 and CCR7
promotes tumor invasion and metastasis (Muller, A. et
al., *Nature* **410**, 50-56 (2001), Staller, P. et al.,
Nature **425**, 307-311 (2003)). To investigate the impact
30 of TPTE in metastasis, tumor cell extravasation, which
is a critical step for metastatic dissemination of
cancer cells mediated by chemotaxis (Chambers, A.F.,
Groom, A.C. & MacDonald, I.C., *Nat. Rev. Cancer* **2**, 563-
572 (2002)) was studied. siRNA treated, fluorophore-

labeled MDA-MB-231 or MCF-7 breast cancer cells were injected into the tail vein of NOD/SCID mice. Six hours later animals were sacrificed and the number of tumor cells extravasated into the lungs was quantified in whole mount lung sections by fluorescence microscopy. For both tumor cell lines siRNA mediated knockdown of TPTE significantly reduced the number of extravasated cells (Fig. 5a). Quantification of submacroscopic metastatic tumor lesions in the lungs of mice several weeks after inoculation with metastases forming breast cancer (MDA-MB-231, MCF-7) or malignant melanoma (MelJuso) cells by human microsatellite specific PCR demonstrated a 100-1000 fold reduction of the tumor load in animals which received tumor cells transiently transfected with TPTE siRNA (Fig. 5b).

Independently, experimental metastases assays with MDA-MB-231 in nude mice giving rise to macroscopic lesions confirmed these striking findings and proved a crucial role of TPTE for metastatic dissemination (Fig. 5c).

Example 7: TPTE and CXCR4 are markers for tumor metastasis

It was assessed whether the strong promigratory and metastasis promoting activity of TPTE is of relevance for the metastatic spread of tumors. To this end, independently collected samples from 34 breast cancer patients from a thoroughly characterized cohort (Ahr, A. et al., *Lancet* 359, 131-132 (2002)) and 24 non-small cell lung cancer specimens were typed for TPTE expression by real-time RT-PCR. There was no significant correlation between TPTE expression and tumor stage or differentiation grade. However, TPTE positive tumors displayed significantly more regional

lymphatic metastasis (76%) and distant metastasis (21%) at the time of diagnosis than TPTE negative primaries (37% and 0%) (Tab. 2a). CXCR4 expression is also a metastasis predicting marker for various cancers.

5 Therefore, the same set of samples was tested for CXCR4 expression. Indeed, CXCR4 positive cancers (n=23) showed a significantly higher rate of distant metastasis (26%) as compared to CXCR4 negative cancers (3%). Importantly, TPTE and CXCR4 expression do not

10 correlate and both molecules represent independent metastasis predictors (Fig. 5c). Cancers with combined expression of TPTE and CXCR4 exhibit a highly increased metastasis rate (60%) whereas tumors lacking either TPTE, CXCR4 or both molecules display an even reduced

15 risk for metastasis (2%, $p < 0.00005$, Tab. 2c) indicating that co-expression of both molecules is important for metastatic spread of cancer, in particular breast and lung cancer.

20 **Table 2a TPTE expression correlates with metastatic spread.**

Status TPTE- Expression	metastatic site	
	lymph nodes	distant
TPTE positive n=33	25 (76%)	7 (21%)
TPTE negative n=25	9 (37%)	0 (0%)
Total n=58	$P < 0.003$	$P < 0.02$

Table 2b CXCR4 expression correlates with distant metastasis.

Status CXCR4- Expression	metastatic site	
	lymph nodes	distant
CXCR4 positive N=23	14 (61%)	6 (26%)
CXCR4 negative n=35	20 (57%)	1 (3%)
Total n=58	P = 0.5	P < 0.02

5 Table 2c Simultaneous expression of TPTE and CXCR4 highly correlates with distant metastasis.

Status TPTE and CXCR4- Expression	metastatic site	
	lymph nodes	distant
TPTE positive AND CXCR4 positive n=10	7 (70%)	6 (60%)
TPTE negative OR CXCR4 negative n=48	27 (56%)	1 (2%)
Total n=58	P = 0.33	P < 0.00005

Claims

1. A method for diagnosing, monitoring and/or prognosing the metastatic behaviour of cancer regarding the occurrence of distant metastasis in a patient, which method comprises quantitatively and/or qualitatively determining the level of expression of TPTE in a biological sample isolated from said patient and quantitatively and/or qualitatively determining the level of expression of CXCR4 in a biological sample isolated from said patient, wherein a level of expression of TPTE and a level of expression of CXCR4 which is increased compared to the level of expression in a subject without metastasis of cancer or without a risk for metastasis of cancer is indicative for a metastatic behaviour of cancer regarding the occurrence of distant metastasis or for a potential for a metastatic behaviour of cancer regarding the occurrence of distant metastasis.

20

2. The method of claim 1 wherein said quantitative and/or qualitative determination of the level of expression of TPTE comprises

(i) detecting or determining the amount of a nucleic acid which is selected from the group consisting of:

(a) a nucleic acid which comprises a nucleic acid sequence selected from the group consisting of SEQ ID NOs: 1, 2, 3, 4, 5, 6, and 7, or a derivative thereof having at least 70% sequence identity,

(b) a nucleic acid which hybridizes with the nucleic acid of (a) under stringent conditions,

(c) a nucleic acid which is degenerate with respect to the nucleic acid of (a) or (b), and

(d) a nucleic acid which is complementary to the nucleic acid of (a), (b) or (c), and/or

(ii) detecting or determining the amount of a protein or peptide encoded by the nucleic acid under (i),
5 and/or

(iii) detecting or determining the amount of an antibody specific for the protein or peptide under (ii), and/or

(iv) detecting or determining the amount of a T
10 lymphocyte specific for the protein or peptide under (ii) or a part thereof comprising a sequence of at least 6 consecutive amino acids of said protein or peptide, optionally in a complex with a MHC molecule, in a biological sample isolated from a patient.

15

3. The method of claim 1 or 2 wherein said quantitative and/or qualitative determination of the level of expression of CXCR4 comprises

(i) detecting or determining the amount of a nucleic
20 acid which is selected from the group consisting of:

(a) a nucleic acid which comprises a nucleic acid sequence selected from the group consisting of SEQ ID NOS: 47, and 48, or a derivative thereof having at least 70% sequence identity,

25 (b) a nucleic acid which hybridizes with the nucleic acid of (a) under stringent conditions,

(c) a nucleic acid which is degenerate with respect to the nucleic acid of (a) or (b), and

(d) a nucleic acid which is complementary to
30 the nucleic acid of (a), (b) or (c), and/or

(ii) detecting or determining the amount of a protein or peptide encoded by the nucleic acid under (i), and/or

- 90 -

(iii) detecting or determining the amount of an antibody specific for the protein or peptide under (ii), and/or

5 (iv) detecting or determining the amount of a T lymphocyte specific for the protein or peptide under (ii) or a part thereof comprising a sequence of at least 6 consecutive amino acids of said protein or peptide, optionally in a complex with a MHC molecule, in a biological sample isolated from a patient.

10

4. The method of claim 2 or 3, wherein the nucleic acid under (i) in said quantitative and/or qualitative determination of the level of expression of TPTE comprises a nucleic acid sequence encoding a protein or peptide comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 8, 9, 10, 11, 12, 13, and 14, or a derivative thereof having at least 70% sequence identity and/or the protein or peptide under (ii) in said quantitative and/or qualitative
15 determination of the level of expression of TPTE comprises an amino acid sequence selected from the group consisting of SEQ ID NOs: 8, 9, 10, 11, 12, 13, and 14, or a derivative thereof having at least 70% sequence identity.

20
25

5. The method of claim 3 or 4, wherein the nucleic acid under (i) in said quantitative and/or qualitative determination of the level of expression of CXCR4 comprises a nucleic acid sequence encoding a protein or peptide comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 49, and 50, or a derivative thereof having at least 70% sequence identity and/or the protein or peptide under (ii) in
30

- 91 -

said quantitative and/or qualitative determination of the level of expression of CXCR4 comprises an amino acid sequence selected from the group consisting of SEQ ID NOs: 49, and 50, or a derivative thereof having at least 70% sequence identity.

6. The method of any one of claims 2 to 5, wherein the detection or determination of the amount of said nucleic acid comprises

- 10 (i) contacting the biological sample with an agent which binds specifically to the nucleic acid, and
(ii) detecting the formation of or determining the amount of a complex between the agent and the nucleic acid.

15

7. The method of any one of claims 2 to 5, wherein the detection or determination of the amount of said protein or peptide comprises

- 20 (i) contacting the biological sample with an agent which binds specifically to the protein or peptide, and
(ii) detecting the formation of or determining the amount of a complex between the agent and the protein or peptide.

25

8. The method of any one of claims 2 to 5, wherein the detection or determination of the amount of said antibody comprises

- 30 (i) contacting the biological sample with an agent which binds specifically to the antibody, and
(ii) detecting the formation of or determining the amount of a complex between the agent and the antibody.

- 92 -

9. The method of any one of claims 2 to 5, wherein the detection or determination of the amount of said T lymphocyte comprises

(i) contacting the biological sample with an agent which binds specifically to the T lymphocyte, and

(ii) detecting the formation of or determining the amount of a complex between the agent and the T lymphocyte.

10. The method of claim 6, wherein the agent which binds specifically to the nucleic acid is an oligonucleotide or polynucleotide, which hybridizes specifically to said nucleic acid.

11. The method of claim 7, wherein the agent which binds specifically to the protein or peptide is an antibody binding specifically to said protein or peptide.

12. The method of claim 8, wherein the agent which binds specifically to the antibody is a protein or peptide binding specifically to said antibody.

13. The method of claim 9, wherein the agent which binds specifically to the T lymphocyte is a cell presenting the complex between the protein or peptide or the part thereof comprising a sequence of at least 6 consecutive amino acids of said protein or peptide for which the T lymphocyte is specific and an MHC molecule.

30

14. The method of any one of claims 1-13 wherein said monitoring comprises determining regression, course or onset of said metastatic behaviour of cancer.

15. The method of claim 14, which comprises determining the level of expression in a first sample at a first point in time and in a further sample at a second point in time and a comparison of the two samples.

16. The method of any one of claims 1-15 wherein said patient has cancer, metastasis of cancer and/or a relapse of cancer or is suspected of or has a risk for falling ill with cancer, metastasis of cancer and/or a relapse of cancer.

17. The method of any one of claims 6-16, wherein said agent is labeled in a detectable manner.

18. The method of any one of claims 1-17, wherein said biological sample comprises body fluid and/or body tissue.

20

19. A kit for quantitatively and/or qualitatively determining the level of expression of TPTE and the level of expression of CXCR4 for diagnosing, monitoring and/or prognosing the metastatic behaviour of cancer regarding the occurrence of distant metastasis comprising means for quantitatively and/or qualitatively determining the level of expression of TPTE and means for quantitatively and/or qualitatively determining the level of expression of CXCR4 in a biological sample isolated from a patient, wherein if said means are means for detection of a nucleic acid or determining the amount of a nucleic acid, said detection or determining of the amount is by

- 94 -

amplification of said nucleic acid.

20. The kit of claim 19 wherein said means for quantitatively and/or qualitatively determining the level of expression of TPTE are selected from the group consisting of

(i) means for detecting or determining the amount of a nucleic acid which is selected from the group consisting of:

10 (a) a nucleic acid which comprises a nucleic acid sequence selected from the group consisting of SEQ ID NOS: 1, 2, 3, 4, 5, 6, and 7, or a derivative thereof having at least 70% sequence identity,

(b) a nucleic acid which hybridizes with the nucleic acid of (a) under stringent conditions,

(c) a nucleic acid which is degenerate with respect to the nucleic acid of (a) or (b), and

(d) a nucleic acid which is complementary to the nucleic acid of (a), (b) or (c), and/or

20 (ii) means for detecting or determining the amount of a protein or peptide encoded by the nucleic acid under (i), and/or

(iii) means for detecting or determining the amount of an antibody specific for the protein or peptide under

25 (ii), and/or

(iv) means for detecting or determining the amount of a T lymphocyte specific for the protein or peptide under (ii) or a part thereof comprising a sequence of at least 6 consecutive amino acids of said protein or peptide, optionally in a complex with a MHC molecule, in a biological sample isolated from a patient.

30

21. The kit of claim 19 or 20 wherein said means for

- 95 -

quantitatively and/or qualitatively determining the level of expression of CXCR4 are selected from the group consisting of

(i) means for detecting or determining the amount of
5 a nucleic acid which is selected from the group consisting of:

(a) a nucleic acid which comprises a nucleic acid sequence selected from the group consisting of SEQ ID NOs: 47, and 48, or a derivative thereof having at
10 least 70% sequence identity,

(b) a nucleic acid which hybridizes with the nucleic acid of (a) under stringent conditions,

(c) a nucleic acid which is degenerate with respect to the nucleic acid of (a) or (b), and

15 (d) a nucleic acid which is complementary to the nucleic acid of (a), (b) or (c), and/or

(ii) means for detecting or determining the amount of a protein or peptide encoded by the nucleic acid under (i), and/or

20 (iii) means for detecting or determining the amount of an antibody specific for the protein or peptide under (ii), and/or

(iv) means for detecting or determining the amount of a T lymphocyte specific for the protein or peptide
25 under (ii) or a part thereof comprising a sequence of at least 6 consecutive amino acids of said protein or peptide, optionally in a complex with a MHC molecule, in a biological sample isolated from a patient.

30 22. The kit of claim 20 or 21, wherein the nucleic acid under (i) in said quantitative and/or qualitative determination of the level of expression of TPTE comprises a nucleic acid sequence encoding a protein or

- 96 -

peptide comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 8, 9, 10, 11, 12, 13, and 14, or a derivative thereof having at least 70% sequence identity and/or the protein or peptide under

5 (ii) in said quantitative and/or qualitative determination of the level of expression of TPTE comprises an amino acid sequence selected from the group consisting of SEQ ID NOs: 8, 9, 10, 11, 12, 13, and 14, or a derivative thereof having at least 70%

10 sequence identity.

23. The kit of claim 21 or 22, wherein the nucleic acid under (i) in said quantitative and/or qualitative determination of the level of expression of CXCR4

15 comprises a nucleic acid sequence encoding a protein or peptide comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 49, and 50, or a derivative thereof having at least 70% sequence identity and/or the protein or peptide under (ii) in

20 said quantitative and/or qualitative determination of the level of expression of CXCR4 comprises an amino acid sequence selected from the group consisting of SEQ ID NOs: 49, and 50, or a derivative thereof having at least 70% sequence identity.

25

24. The kit of any one of claims 20 to 23, wherein said means for detecting or determining the amount of said nucleic acid comprise an agent which binds specifically to the nucleic acid.

30

25. The kit of any one of claims 20 to 24, wherein said means for detecting or determining the amount of said protein or peptide comprise an agent which binds

specifically to the protein or peptide.

26. The kit of any one of claims 20 to 25, wherein said means for detecting or determining the amount of
5 said antibody comprise an agent which binds specifically to the antibody.

27. The kit of any one of claims 20 to 26, wherein said means for detecting or determining the amount of
10 said T lymphocyte comprise an agent which binds specifically to the T lymphocyte.

28. The kit of claim 24, wherein the agent which binds specifically to the nucleic acid is an
15 oligonucleotide or polynucleotide, which hybridizes specifically to said nucleic acid.

29. The kit of claim 25, wherein the agent which binds specifically to the protein or peptide is an
20 antibody binding specifically to said protein or peptide.

30. The kit of claim 26, wherein the agent which binds specifically to the antibody is a protein or
25 peptide binding specifically to said antibody.

31. The kit of claim 27, wherein the agent which binds specifically to the T lymphocyte is a cell
presenting the complex between the protein or peptide
30 or the part thereof comprising a sequence of at least 6 consecutive amino acids of said protein or peptide for which the T lymphocyte is specific and an MHC molecule.