

RNA-GUIDED GENOME RECOMBINEERING AT KILOBASE SCALE**CROSS-REFERENCE TO RELATED APPLICATIONS**

[0001] This application claims the benefit of U.S. Provisional Application No. 62/984,618, filed March 3, 2020, and U.S. Provisional Application No. 63/146,447, filed February 5, 2021, the contents of each are incorporated herein by reference.

FIELD

[0002] The present invention relates to RNA-guided recombineering-editing systems using phage recombination enzymes as well as methods, vectors, nucleic acid compositions, and kits thereof.

BACKGROUND

[0003] The Clustered Regularly Interspaced Short Palindromic Repeats (CRISPR) system, originally found in bacteria and archaea as part of the immune system to defend against invading viruses, forms the basis for genome editing technologies that can be programmed to target specific stretches of a genome or other DNA for editing at precise locations. While various CRISPR-based tools are available, the majority are geared towards editing short sequences. Long-sequence editing is highly sought after in the engineering of model systems, therapeutic cell production and gene therapy. Prior studies have developed technologies to improve Cas9-mediated homology-5 directed repair (HDR), and tools leveraging nucleic acid modification enzymes with Cas9, e.g., prime-editing, demonstrated editing up to 80 base-pairs (bp) in length. Despite these progresses, there are continued demands for large-scale mammalian genome engineering with high efficiency and fidelity.

SUMMARY

[0004] Provided herein are systems and methods that facilitate nucleic acid editing in a manner that allows large-scale nucleic acid editing with high accuracy and low off-target errors. These systems and methods employ a combination of microbial recombination components with CRISPR recombination components.

[0005] For example, disclosed herein are systems comprising a protein, a nucleic acid molecule comprising a guide RNA sequence that is complementary to a target DNA sequence, and a microbial recombination protein. The microbial recombination protein may be, for example, RecE, RecT, lambda exonuclease (Exo), Bet protein (betA, redB), exonuclease gp6, single-stranded DNA-binding protein

gp2.5, or a derivative or variant thereof. In some embodiments, the system further comprises donor DNA. In some embodiments, the target DNA sequence is a genomic DNA sequence in a host cell.

[0006] In some embodiments, the system further comprises a recruitment system comprising at least one aptamer sequence and an aptamer binding protein functionally linked to the microbial recombination protein as part of a fusion protein. In some embodiments, the aptamer sequence is an RNA aptamer sequence or a peptide aptamer sequence. In some embodiments, the RNA aptamer sequence is part of the nucleic acid molecule. In some embodiments, the nucleic acid molecule comprises two RNA aptamer sequences. In some embodiments, the microbial recombination protein is functionally linked to the aptamer binding protein as a fusion protein. In some embodiments, the binding protein comprises a MS2 coat protein, a lambda N22 peptide, or a functional derivative, fragment, or variant thereof. In some embodiments, the fusion protein further comprises a linker and/or a nuclear localization sequence.

[0007] Disclosed herein are compositions comprising a nucleic acid sequence encoding a fusion protein comprising a microbial recombination protein functionally linked to an aptamer binding protein. The microbial recombination protein may be RecE, RecT, lambda exonuclease (Exo), Bet protein (betA, redB), exonuclease gp6, single-stranded DNA-binding protein gp2.5, or a derivative or variant thereof. The compositions may further comprise one or both of a polynucleotide comprising a nucleic acid sequence encoding a Cas protein and a nucleic acid molecule comprising a guide RNA sequence that is complementary to a target DNA sequence. In some embodiments, the nucleic acid molecule further comprises at least one RNA aptamer sequence. In some embodiments, the polynucleotide comprising a nucleic acid sequence encoding a Cas protein further comprises a sequence encoding at least one peptide aptamer sequence.

[0008] Also disclosed herein are vectors comprising a nucleic acid sequence encoding a fusion protein comprising a microbial recombination protein functionally linked to an aptamer binding protein. The microbial recombination protein may be RecE, RecT, lambda exonuclease (Exo), Bet protein (betA, redB), exonuclease gp6, single-stranded DNA-binding protein gp2.5, or a derivative or variant thereof. The vectors may further comprise one or both of a polynucleotide comprising a nucleic acid sequence encoding a Cas protein and a nucleic acid molecule comprising a guide RNA sequence that is complementary to a target DNA sequence. In some embodiments, the nucleic acid molecule further comprises at least one RNA aptamer sequence. In some embodiments, the polynucleotide comprising a nucleic acid sequence encoding a Cas protein further comprises a sequence encoding at least one peptide aptamer sequence.

[0009] In some embodiments, the RecE and RecT recombination protein is derived from *E. coli*. In some embodiments, the RecE, or derivative or variant thereof, comprises an amino acid sequence with at least 70% similarity to amino acid sequences selected from the group consisting of SEQ ID NOs: 1-8. In some embodiments, the RecT, or derivative or variant thereof, comprises an amino acid sequence with at least 70% similarity to amino acid sequences selected from the group consisting of SEQ ID NO: 9.

[0010] In some embodiments, the Cas protein is Cas9 or Cas12a. In some embodiments, the Cas protein is a catalytically dead. In some embodiments, the Cas9 protein is wild-type *Streptococcus pyogenes* Cas9 or a wild-type *Staphylococcus aureus* Cas9. In some embodiments, the Cas9 protein is a Cas9 nickase (e.g., wild-type *Streptococcus pyogenes* Cas9 with an amino acid substitution at position 10 of D10A).

[0011] Also disclosed is a eukaryotic cell comprising the systems or vectors disclosed herein.

[0012] Further disclosed herein are methods of altering a target genomic DNA sequence in a host cell. The methods comprise contacting the systems, compositions, or vectors described herein with a target DNA sequence (e.g., introducing the systems, compositions, or vectors described herein into a host cell comprising a target genomic DNA sequence). Kits containing one or more reagents or other components useful, necessary, or sufficient for practicing any of the methods are also disclosed herein.

[0013] Other aspects and embodiments of the disclosure will be apparent in light of the following detailed description and accompanying figures.

BRIEF DESCRIPTION OF THE DRAWINGS

[0014] FIG. 1A and FIG. 1B are the reconstructed RecE (FIG. 1A) and RecT (FIG. 1B) phylogenetic trees with eukaryotic recombination enzymes from yeast and human.

[0015] FIG. 2A is a phylogenetic tree and length distribution of RecE/RecT homologs. FIG. 2B is the metagenomics distribution of RecE/T. FIG. 2C is a schematic showing central models disclosed herein. FIG. 2D are graphs of the genome knock-in efficiency of RecE/T homologs.

[0016] FIG. 3A and 3B are graphs of the high-throughput sequencing (HTS) reads of homology directed repair (HDR) at the *EMX1* (FIG. 3A) locus and the *VEGFA* (FIG. 3B) locus. FIGS. 3C-3D are graphs of the mKate knock-in efficiency at *HSP90AA1* (FIG. 3C), *DYNLT1* (FIG. 3D), and *AAVSI* (FIG. 3E) loci in HEK293T cells. FIG. 3F is images of mKate knock-in efficiency in HEK293T cells with RecT. FIG. 3G is a schematic of an exemplary *AAVSI* knock-in strategy and chromatogram trace from RecT knock-in group. FIG. 3H is schematics and graphs of the recruitment control experiment and

corresponding knock-in efficiency. All results are normalized to NR. (NC, no cutting; NR, no recombinator).

[0017] FIGS. 4A-4C are graphs of the relative mKate knock-in efficiencies to the NE group at *HSP90AA1* (FIG. 4A), *DYNLT1* (FIG. 4B), and *AAVSI* (FIG. 4C) loci in HEK293T cells. (NC, no cutting control group. NR, no recombinator control group.) FIG. 4D is an image of an exemplary agarose gel of junction PCR that validates mKate knock-in at *AAVSI* locus. FIG. 4E and 4F are graphs of the absolute and (FIG. 4E) and relative (FIG. 4F) LOV knock-in efficiencies at *AAVSI* locus.

[0018] FIGS. 5A-5D are graphs of the genomic knock-in efficiencies at different loci across cell lines A549 (FIG. 5A), HepG2 (FIG. 5B), HeLa (FIG. 5C), and hESCs (H9) (FIG. 5D). FIG. 5E is images of mKate knock-ins in hESCs. FIG. 5F and 5G are genomic-wide off-target site (OTS) counts (FIG. 5F) and OTS chromosomal distribution (FIG. 5G) of REDITv1 tools.

[0019] FIGS. 6A-6D are graphs of the relative mKate knock-in efficiency at the *AAVSI* locus and the *DYNLT1* locus in A549 cell line (FIG. 6A), the *DYNLT1* locus and the *HSP90AA1* locus in HepG2 cell line (FIG. 6B), the *DYNLT1* locus and the *HSP90AA1* locus in Hela cell line (FIG. 6C), and the *HSP90AA1* locus and the *OCT4* locus in hES-H9 cell line (FIG. 6D). (NC, no cutting control group. NR, no recombinator control group. All data normalized to NR group.) FIG. 6E is representative FACS results of *HSP90AA1* mKate knock-in in hES-H9 cells.

[0020] FIGS. 7A-7D are graphs of the absolute mKate knock-in efficiencies of different homology arm lengths at the *DYNLT1* (FIG. 7A) and *HSP90AA1* (FIG. 7B) loci and the no recombinator controls for *DYNLT1* (FIG. 7C) and *HSP90AA1* (FIG. 7D).

[0021] FIGS. 8A-8E are graphs of the indel rates of the top 3 predicted off-target loci associated with sgEMX1 (FIGS. 8A-8C) or sgVEGFA (FIGS. 8D-8E) in the REDITv1 system.

[0022] FIG. 9A is a schematic of select embodiments of REDITv2N and corresponding knock-in efficiencies in HEK293T cells. FIG. 9B and 9C are graphs of genomic-wide off-target site (OTS) counts (FIG. 9B) and OTS chromosomal distribution (FIG. 9C) comparing REDITv2N against REDITv1. FIG. 9D is a schematic of select embodiments of REDITv2D and corresponding knock-in efficiencies. FIG. 9E is a graph of editing efficiency of REDITv1, REDITv2N, and REDITv2D under serum starvation conditions. FIG. 9F is the knock-in efficiencies of REDITv3 in hESCs. FIG. 9G is images of mKate knock in using REDITv3 in hESCs.

[0023] FIG. 10A and 10B are schematics and graphs of the relative mKate knock-in efficiencies of select embodiments of REDITv2N (FIG. 10A) and REDITv2D (FIG. 10B) at the *DYNLT1* locus and the *HSP90AA1* locus.

[0024] FIGS. 11A-11D are images of agarose gels showing junction PCR of mKate knock-in at the *DYNLT1* locus and the *HSP90AA1* locus for a select REDITv2N system.

[0025] FIG. 12A and 12B are graphs of the genomic distribution of detected off-target cleavages of select embodiments of REDITv2 (FIG. 12A) and REDITv2N (FIG. 12B). A pileup includes alignments that have two or more reads overlapping with each other. Flanking pairs include alignments that show up on opposite strands within 200bp upstream of each other. Target matched includes alignments that match to a treated target in the upstream sequence (up to 6 mismatches, including 1 mismatch in the PAM, are allowed in the target sequence). FIG. 12C is a graph of the HTS HDR and indel reads at *EMXI* locus for REDITv2N system.

[0026] FIG. 13A is an image of an agarose gel showing junction PCR of mKate knock-ins at the *DYNLT1* locus for REDITv2D system.

[0027] FIGS. 14A-14C are graphs of the mKate knock-in efficiencies at the *HSP90AA1* locus in REDITv2 (FIG. 14A), REDITv2N (FIG. 14B) and REVITv2D (FIG. 14C) when treated with different FBS concentrations. FIGS. 14D-14F are graphs of the mKate knock-in efficiencies at the *HSP90AA1* locus in REDITv2 (FIG. 14D), REDITv2N (FIG. 14E) and REVITv2D (FIG. 14F) when treated with different serum FBS concentrations.

[0028] FIG. 15 is images of the nuclear localization of RecE_587 and RecT following EGFP fusion to the REDITv1 systems. Nuclei were stained with NucBlue Live Ready Probes Reagent.

[0029] FIG. 16A and 16B are the relative mKate knock-in efficiencies at *HSP90AA1* and *DYNLT1* loci following fusion of different nuclear localization sequences to either the N- or C-terminus of RecT and RecE_587. FIG. 16C and 16D are graphs of the absolute mKate knock-in efficiencies of the constructs from FIGS. 16A and 16B for the *DYNLT1* locus (FIG. 16C) and the *HSP90AA1* locus (FIG. 16D).

[0030] FIGS. 17A-17D are graphs of the relative (FIGS. 17A and 17B) and absolute (FIGS. 17C and 17D) mKate knock-in efficiencies for the *DYNLT1* locus (FIGS. 17A and 17C) and the *HSP90AA1* locus (FIGS. 17B and 17D) following fusion new NLS sequences as well as optimal linkers to REDITv2 and REDITv3 variants. The REDITv2 versions using REDITv2N (D10A or H840A) and REDITv2D (dCas9) are indicated in the horizontal axis, along with the number of guides used. The different colors represent the different control groups and REDIT versions.

[0031] FIG. 18 is a graph of the relative editing efficiency of REDITv3N system at *HSP90AA1* locus in hES-H9 cells.

[0032] FIG. 19A is a diagram of an exemplary saCas9 expression vector. FIGS. 19B-19E are graphs of the relative mKate knock-in efficiencies at the *AAVS1* locus (FIG. 19D) and *HSP90AA1* locus (FIG. 19E) of different effectors in saCas9 system and the respective absolute efficiencies (FIG. 19B and 19C, respectively). NC, no cutting control group. NR, no recombinator control group.

[0033] FIG. 20A is a schematic of RecT truncations. FIGS. 20B and 20C are graphs of the relative mKate knock-in efficiencies at the *DYNLT1* locus for wild-type *Streptococcus pyogenes* Cas9 and *Streptococcus pyogenes* Cas9n(D10A) with single- and double-nicking.

[0034] FIG. 21A is a schematic of RecE_587 truncations. FIGS. 21B and 21C are graphs of the relative mKate knock-in efficiencies at the *DYNLT1* locus for wild-type *Streptococcus pyogenes* Cas9 and *Streptococcus pyogenes* Cas9n(D10A) with single- and double-nicking.

[0035] FIGS. 22A and 22B are graphs of comparison of efficiency to perform recombineering-based editing with various exonucleases (FIG. 22A) and single-strand DNA annealing protein (SSAP) (FIG. 22B) from naturally occurring recombineering systems, including NR (no recombinator) as negative control. The gene-editing activity was measured using mKate knock-in assay at genomic loci (*DYNLT1* and *HSP90AA1*). The data shown are percentage of successful mKate knock-in using human HEK293 cells, each experiments were performed in triplicate (n=3).

[0036] FIGS. 23A-23E show a compact recruitment system using boxB and N22. The REDIT recombinator proteins were fused to N22 peptide and within the sgRNA was boxB, the short cognizant sequence of N22 peptide (FIG. 23A). FIGS. 23B-23E are graphs of the gene-editing efficiency using mKate knock-in assay, with wildtype SpCas9, with side-by-side comparisons to the MS2-MCP recruitment system. FIGS. 23B and 23D are absolute mKate knock-in efficiency at *DYNLT1*, *HSP90AA1* loci and FIGS. 23C and 23E are relative efficiencies. The data shown are percentage of successful mKate knock-in using HEK293 human cells, each experiments were performed in triplicate (n=3).

[0037] FIGS. 24A-24C show a SunTag recruitment system. The REDIT recombinator proteins were fused to scFV antibody and the GCN4 peptide in tandem fashion (10 copies of GCN4 peptide separated by linkers) was fused to the Cas9 protein (FIG. 24A). An mKate knock-in experiment (FIG. 24B) with the *DYNLT1* locus was used to measure the gene-editing knock-in efficiency (FIG. 24C). All data are measurements of gene-editing efficiency using mKate knock-in assay, with wildtype SpCas9. Absolute mKate knock-in efficiency at *DYNLT1* are shown in the bottom right corner of each flow cytometry plot,

where the control is without recombinator (NR), which included scFV fused to GFP protein as negative control, all experiments done in HEK293 human cells.

[0038] FIGS. 25A and 25B exemplify REDIT with a Cas12A system. A Cpf1/Cas12a based REDIT system via the SunTag recruitment design was created (FIG. 25A) for two different Cpf1/Cas12a proteins. Using the mKate knock-in assay, the efficiencies at two endogenous loci (*DYNLT1* and *AASI*) were measured. (FIG. 25B). Shown are absolute mKate knock-in efficiency as measured by mKate⁺ cell percentage using HEK293 human cells, each experiment was performed in triplicate (n=3), where the negative control is without recombinator (NR).

[0039] FIGS. 26A and 26B are the measurements of precision recombineering activity via mKate knock-in gene-editing assay using RecE and RecT homologs at the *DYNLT1* locus (A) and the *HSP90AA1* locus (B). Shown are absolute mKate knock-in efficiency as measured by mKate⁺ cell percentage using HEK293 human cells, each experiments were performed in triplicate (n=3), where the negative control is without recombinator (NR) and no cutting (NC). The original RecE and RecT from *E. coli* were also included as positive controls.

[0040] FIGS. 27A and 27B is a schematic showing the SunTag-based recruitment of SSAP RecT to Cas9-gRNA complex for gene-editing (FIG. 27A) and a graph quantifying the editing efficiencies of SunTag compared to MS2-based strategies (FIG. 27B).

[0041] FIGS. 28A-28C show comparisons of REDIT with alternative HDR-enhancing gene-editing approaches. FIG. 28A is schematics showing alternative HDR-enhancing approaches via fusing functional domains, CtIP or Geminin (Gem), to Cas9 protein (left) and when combined with REDIT (right). FIG. 28B is an alternative small-molecule HDR-enhancing approach through cell cycle control. Nocodazole was used to synchronize cells at the G2/M boundary (left) according to the timeline shown (right). FIG. 28C is comparisons of gene-editing efficiencies using REDIT and alternative HDR-enhancing tools, Cas9-HE (CtIP fusion), Cas9-Gem (Geminin fusion), and Nocodazole (noc), along with combination of REDIT with these methods (Cas9-HE/Cas9-Gem/noc+REDIT). Donor DNAs have 200 + 400 bp (*DYNLT1*) or 200 + 200bp (*HSP90AA1*) of HAs. All assays performed with no donor, NTC and Cas9 (no enhancement) controls. #*P* < 0.05, compared to REDIT; ##*P* < 0.01, compared to REDIT.

[0042] FIGS. 29A-29D show template design guideline, junction precision, and capacity of REDIT gene-editing methods. FIG. 29A is graphs of a homology arm (HA) length test comparing different template designs of HDR donors (longer HAs) or NHEJ/MMEJ donors (zero/shorter HAs) using REDIT and Cas9 references. Top and bottom are two genomic loci tested using mKate knock-in assay. FIG. 29B

is a design of an exemplary junction profiling assay through isolation of knock-in clones, followed by genomic PCR using primers (fwd, rev) binding outside donor to avoid template amplification. Paired Sanger sequencing of the PCR products reveal homologous and non-homologous edits at the 5'- and 3'- junctions. FIG. 29C is a graph of the percentage of colonies with indicated junction profiles from the Sanger sequencing of knock-in clones as in FIG. 29B. Editing methods and donor DNA are listed at the bottom (HA lengths indicated in bracket). FIG. 29D is a graph of knock-in efficiencies using a 2-kb cassette to insert dual-GFP/mKate tags to validate REDIT methods with Cas9. HA lengths of donor DNAs indicated at the bottom.

[0043] FIGS. 30A-30C show GISseq results indicating that REDIT is an efficient method with the ability to insert kilobase-length sequences with less unwanted editing events. FIG. 30A is a schematic showing the design, procedures, and analysis steps for GIS-seq to measure genome-wide insertion sites of the knock-in cassettes. High-molecular-weight (HMW) genomic DNA purification was needed to remove potential contamination from donor DNAs. Donor DNAs had 200 bp HAs each side. FIG. 30B is representative GIS-seq results showing plus/minus reads at on-target locus *DYNLT1*. The expected 2A-mKate knock-in site before the stop codon of the last exon are the center of the trimmed reads (reads clipped to remove 2A-mKate cassette). The template mutations help to avoid gRNA targeting and distinguish genomic and edited reads are labeled. FIG. 30C is a summary of top GIS-seq insertion sites comparing Cas9dn and REDITdn groups, showing the expected on-target insertion site (highlighted) and reduced number of identified off-target insertion sites when using REDITdn. (Left) *DYNLT1* and (Right) *ACTB* loci with MLE calculated from the distribution of filtered and trimmed GIS-seq reads.

[0044] FIGS. 31A-31F show the dependence of REDIT gene-editing on endogenous DNA repair and applying REDIT methods for human stem cell engineering. FIG. 31A is a model showing the editing process and major repair pathways involved when using REDIT or Cas9 for gene-editing, the HDR pathway are highlighted for chemical perturbation (inhibition of RAD51). Donor DNAs with 200 + 200 bp HAs are used for all inhibitor experiments. FIGS. 31B and 31C are graphs showing the relative knock-inefficiency of REDIT tools compared with Cas9 reference treated with RAD51 inhibitor B02 and RI-1, or vehicle-treated, for the wtCas9-based REDIT and Cas9 (FIG. 31B) and for Cas9 nickase-based REDITdn and Cas9dn (FIG. 31C). All conditions were measured with 1-kb knock-in assay at two genomic loci (*DYNLT1* and *HSP90AA1*). FIG. 31D are graphs of knock-in efficiencies in hESCs (H9) using REDIT and REDITdn tested across three genomic loci, compared with corresponding Cas9 and Cas9dn references. FIGS. 31E and 31F are flow cytometry plots of mKate knock-in results in hESCs

using REDIT, REDITdn with Cas9, Cas9dn, and NTC controls. Donor DNAs in the hESC experiments have 200 + 200 bp HAs across all loci tested.

[0045] FIGS. 32A-32B show chemical perturbations to dCas9 REDIT. Gene editing efficiencies were determined when treated with mammalian DNA repair pathway inhibitors (Mirin, RI-1, and B02) with (FIG. 32A) and without (FIG. 32B) cell cycle inhibitor (Thy, doubly Thymidine) blocking. Statistical analyses are from t-test results with 1% FDR via a two-stage step-up method.

[0046] FIGS. 33A and 33B are schematics of the DNA components (gene-editing vectors and template DNA) and tail vein injection of mice, respectively.

[0047] FIGS. 34A-34C are results from the tail vein injection of mice with gene-editing vectors. FIG. 34A is a schematic and gel electrophoresis of PCR analysis of liver hepatocytes from the injected mice. FIG. 34B is the Sanger sequencing results of the PCR amplicon (SEQ ID NO: 162). FIG. 34C is a schematic of next-generation sequencing and a graph of the quantification of knock-in junction errors.

[0048] FIGS. 35A and 35B are schematics of the DNA components (gene-editing and control vector) and adeno-associated virus (AAV) treatment, respectively. FIG. 35C is fluorescent images of lungs from AAV treated mice and graphs of corresponding quantitation of tumor number.

DETAILED DESCRIPTION OF THE INVENTION

[0049] The present disclosure is directed to a system and the components for DNA editing. In particular, the disclosed system based on CRISPR targeting and homology directed repair by phage recombination enzymes. The system results in superior recombination efficiency and accuracy at a kilobase scale.

1. Definitions

[0050] To facilitate an understanding of the present technology, a number of terms and phrases are defined below. Additional definitions are set forth throughout the detailed description.

[0051] The terms “comprise(s),” “include(s),” “having,” “has,” “can,” “contain(s),” and variants thereof, as used herein, are intended to be open-ended transitional phrases, terms, or words that do not preclude the possibility of additional acts or structures. The singular forms “a,” “and” and “the” include plural references unless the context clearly dictates otherwise. The present disclosure also contemplates other embodiments “comprising,” “consisting of” and “consisting essentially of,” the embodiments or elements presented herein, whether explicitly set forth or not.

[0052] For the recitation of numeric ranges herein, each intervening number there between with the same degree of precision is explicitly contemplated. For example, for the range of 6-9, the numbers 7 and 8 are contemplated in addition to 6 and 9, and for the range 6.0-7.0, the number 6.0, 6.1, 6.2, 6.3, 6.4, 6.5, 6.6, 6.7, 6.8, 6.9, and 7.0 are explicitly contemplated.

[0053] Unless otherwise defined herein, scientific, and technical terms used in connection with the present disclosure shall have the meanings that are commonly understood by those of ordinary skill in the art. For example, any nomenclature used in connection with, and techniques of, cell and tissue culture, molecular biology, immunology, microbiology, genetics and protein and nucleic acid chemistry and hybridization described herein are those that are well known and commonly used in the art. The meaning and scope of the terms should be clear; in the event, however of any latent ambiguity, definitions provided herein take precedent over any dictionary or extrinsic definition. Further, unless otherwise required by context, singular terms shall include pluralities and plural terms shall include the singular.

[0054] The terms “complementary” and “complementarity” refer to the ability of a nucleic acid to form hydrogen bond(s) with another nucleic acid sequence by either traditional Watson-Crick base-pairing or other non-traditional types of pairing. The degree of complementarity between two nucleic acid sequences can be indicated by the percentage of nucleotides in a nucleic acid sequence which can form hydrogen bonds (e.g., Watson-Crick base pairing) with a second nucleic acid sequence (e.g., 50%, 60%, 70%, 80%, 90%, and 100% complementary). Two nucleic acid sequences are “perfectly complementary” if all the contiguous nucleotides of a nucleic acid sequence will hydrogen bond with the same number of contiguous nucleotides in a second nucleic acid sequence. Two nucleic acid sequences are “substantially complementary” if the degree of complementarity between the two nucleic acid sequences is at least 60% (e.g., 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97%, 98%, 99%, or 100%) over a region of at least 8 nucleotides (e.g., 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 30, 35, 40, 45, 50, or more nucleotides), or if the two nucleic acid sequences hybridize under at least moderate, preferably high, stringency conditions. Exemplary moderate stringency conditions include overnight incubation at 37° C in a solution comprising 20% formamide, 5×SSC (150 mM NaCl, 15 mM trisodium citrate), 50 mM sodium phosphate (pH 7.6), 5×Denhardt’s solution, 10% dextran sulfate, and 20 mg/ml denatured sheared salmon sperm DNA, followed by washing the filters in 1×SSC at about 37-50° C., or substantially similar conditions, e.g., the moderately stringent conditions described in Sambrook et al., *infra*. High stringency conditions are conditions that use, for example (1) low ionic strength and high temperature for washing, such as 0.015 M sodium chloride/0.0015 M sodium citrate/0.1% sodium dodecyl sulfate (SDS) at 50° C,

(2) employ a denaturing agent during hybridization, such as formamide, for example, 50% (v/v) formamide with 0.1% bovine serum albumin (BSA)/0.1% Ficoll/0.1% polyvinylpyrrolidone (PVP)/50 mM sodium phosphate buffer at pH 6.5 with 750 mM sodium chloride and 75 mM sodium citrate at 42° C., or (3) employ 50% formamide, 5×SSC (0.75 M NaCl, 0.075 M sodium citrate), 50 mM sodium phosphate (pH 6.8), 0.1% sodium pyrophosphate, 5×Denhardt's solution, sonicated salmon sperm DNA (50 µg/ml), 0.1% SDS, and 10% dextran sulfate at 42° C., with washes at (i) 42° C. in 0.2×SSC, (ii) 55° C. in 50% formamide, and (iii) 55° C. in 0.1×SSC (preferably in combination with EDTA). Additional details and an explanation of stringency of hybridization reactions are provided in, e.g., Sambrook et al., *Molecular Cloning: A Laboratory Manual*, 3rd ed., Cold Spring Harbor Press, Cold Spring Harbor, N.Y. (2001); and Ausubel et al., *Current Protocols in Molecular Biology*, Greene Publishing Associates and John Wiley & Sons, New York (1994).

[0055] A cell has been “genetically modified,” “transformed,” or “transfected” by exogenous DNA, e.g., a recombinant expression vector, when such DNA has been introduced inside the cell. The presence of the exogenous DNA results in permanent or transient genetic change. The transforming DNA may or may not be integrated (covalently linked) into the genome of the cell. In prokaryotes, yeast, and mammalian cells for example, the transforming DNA may be maintained on an episomal element such as a plasmid. With respect to eukaryotic cells, a stably transformed cell is one in which the transforming DNA has become integrated into a chromosome so that it is inherited by daughter cells through chromosome replication. This stability is demonstrated by the ability of the eukaryotic cell to establish cell lines or clones that comprise a population of daughter cells containing the transforming DNA. A “clone” is a population of cells derived from a single cell or common ancestor by mitosis. A “cell line” is a clone of a primary cell that is capable of stable growth in vitro for many generations.

[0056] As used herein, a “nucleic acid” or a “nucleic acid sequence” refers to a polymer or oligomer of pyrimidine and/or purine bases, preferably cytosine, thymine, and uracil, and adenine and guanine, respectively. The present technology contemplates any deoxyribonucleotide, ribonucleotide, or peptide nucleic acid component, and any chemical variants thereof, such as methylated, hydroxymethylated, or glycosylated forms of these bases, and the like. The polymers or oligomers may be heterogenous or homogenous in composition and may be isolated from naturally occurring sources or may be artificially or synthetically produced. In addition, the nucleic acids may be DNA or RNA, or a mixture thereof, and may exist permanently or transitionally in single-stranded or double-stranded form, including homoduplex, heteroduplex, and hybrid states. In some embodiments, a nucleic acid or nucleic acid

sequence comprises other kinds of nucleic acid structures such as, for instance, a DNA/RNA helix, peptide nucleic acid (PNA), morpholino nucleic acid (see, e.g., Braasch and Corey, *Biochemistry*, 41(14): 4503-4510 (2002)) and U.S. Pat. No. 5,034,506, incorporated herein by reference), locked nucleic acid (LNA; see Wahlestedt et al., *Proc. Natl. Acad. Sci. U.S.A.*, 97: 5633-5638 (2000), incorporated herein by reference), cyclohexenyl nucleic acids (see Wang, *J. Am. Chem. Soc.*, 122: 8595-8602 (2000), incorporated herein by reference), and/or a ribozyme. Hence, the term “nucleic acid” or “nucleic acid sequence” may also encompass a chain comprising non-natural nucleotides, modified nucleotides, and/or non-nucleotide building blocks that can exhibit the same function as natural nucleotides (e.g., “nucleotide analogs”); further, the term “nucleic acid sequence” as used herein refers to an oligonucleotide, nucleotide or polynucleotide, and fragments or portions thereof, and to DNA or RNA of genomic or synthetic origin, which may be single or double-stranded, and represent the sense or antisense strand. The terms “nucleic acid,” “polynucleotide,” “nucleotide sequence,” and “oligonucleotide” are used interchangeably. They refer to a polymeric form of nucleotides of any length, either deoxyribonucleotides or ribonucleotides, or analogs thereof.

[0057] A “peptide” or “polypeptide” is a linked sequence of two or more amino acids linked by peptide bonds. The peptide or polypeptide can be natural, synthetic, or a modification or combination of natural and synthetic. Polypeptides include proteins such as binding proteins, receptors, and antibodies. The proteins may be modified by the addition of sugars, lipids or other moieties not included in the amino acid chain. The terms “polypeptide” and “protein,” are used interchangeably herein.

[0058] As used herein, the term “percent sequence identity” refers to the percentage of nucleotides or nucleotide analogs in a nucleic acid sequence, or amino acids in an amino acid sequence, that is identical with the corresponding nucleotides or amino acids in a reference sequence after aligning the two sequences and introducing gaps, if necessary, to achieve the maximum percent identity. Hence, in case a nucleic acid according to the technology is longer than a reference sequence, additional nucleotides in the nucleic acid, that do not align with the reference sequence, are not taken into account for determining sequence identity. Methods and computer programs for alignment are well known in the art, including BLAST, Align 2, and FASTA.

[0059] A “vector” or “expression vector” is a replicon, such as plasmid, phage, virus, or cosmid, to which another DNA segment, e.g., an “insert,” may be attached or incorporated so as to bring about the replication of the attached segment in a cell.

[0060] The term “wild-type” refers to a gene or a gene product that has the characteristics of that gene or gene product when isolated from a naturally occurring source. A wild-type gene is that which is most frequently observed in a population and is thus arbitrarily designated the “normal” or “wild-type” form of the gene. In contrast, the term “modified,” “mutant,” or “polymorphic” refers to a gene or gene product that displays modifications in sequence and or functional properties (e.g., altered characteristics) when compared to the wild-type gene or gene product. It is noted that naturally occurring mutants can be isolated; these are identified by the fact that they have altered characteristics when compared to the wild-type gene or gene product.

2. RNA-guided CRISPR Recombineering System

[0061] In bacteria and archaea, CRISPR/Cas systems provide immunity by incorporating fragments of invading phage, virus, and plasmid DNA into CRISPR loci and using corresponding CRISPR RNAs (“crRNAs”) to guide the degradation of homologous sequences. Each CRISPR locus encodes acquired “spacers” that are separated by repeat sequences. Transcription of a CRISPR locus produces a “pre-crRNA,” which is processed to yield crRNAs containing spacer-repeat fragments that guide effector nuclease complexes to cleave dsDNA sequences complementary to the spacer. Three different types of CRISPR systems are known, type I, type II, or type III, and classified based on the Cas protein type and the use of a proto-spacer-adjacent motif (PAM) for selection of proto-spacers in invading DNA. The endogenous type II systems comprise the Cas9 protein and two noncoding crRNAs: trans-activating crRNA (tracrRNA) and a precursor crRNA (pre-crRNA) array containing nuclease guide sequences (also referred to as “spacers”) interspaced by identical direct repeats (DRs). tracrRNA is important for processing the pre-crRNA and formation of the Cas9 complex. First, tracrRNAs hybridize to repeat regions of the pre-crRNA. Second, endogenous RNaseIII cleaves the hybridized crRNA-tracrRNAs, and a second event removes the 5’ end of each spacer, yielding mature crRNAs that remain associated with both the tracrRNA and Cas9. Third, each mature complex locates a target double stranded DNA (dsDNA) sequence and cleaves both strands using the nuclease activity of Cas9.

[0062] CRISPR/Cas gene editing systems have been developed to enable targeted modifications to a specific gene of interest in eukaryotic cells. CRISPR/Cas gene editing systems are commonly based on the RNA-guided Cas9 nuclease from the type II prokaryotic clustered regularly interspaced short palindromic repeats (CRISPR) adaptive immune system. Engineering CRISPR/Cas systems for use in eukaryotic cells typically involves reconstitution of the crRNA-tracrRNA-Cas9 complex. In human cells, for example, the Cas9 amino acid sequence may be codon-optimized and modified to include an

appropriate nuclear localization signal, and the crRNA and tracrRNA sequences may be expressed individually or as a single chimeric molecule via an RNA polymerase II promoter. Typically, the crRNA and tracrRNA sequences are expressed as a chimera and are referred to collectively as “guide RNA” (gRNA) or single guide RNA (sgRNA). Thus, the terms “guide RNA,” “single guide RNA,” and “synthetic guide RNA,” are used interchangeably herein and refer to a nucleic acid sequence comprising a tracrRNA and a pre-crRNA array containing a guide sequence. The terms “guide sequence,” “guide,” and “spacer,” are used interchangeably herein and refer to the about 20 nucleotide sequence within a guide RNA that specifies the target site. In CRISPR/Cas9 systems, the guide RNA contains an approximate 20-nucleotide guide sequence followed by a protospacer adjacent motif (PAM) that directs Cas9 via Watson-Crick base pairing to a target sequence.

[0063] In some embodiments, the disclosure provides a system for RNA-guided recombineering utilizing tools from CRISPR gene editing systems. The system comprises: a Cas protein, a nucleic acid molecule comprising a guide RNA sequence that is complementary to a target DNA sequence and a microbial recombination protein.

[0064] Cas protein families are described in further detail in, e.g., Haft et al., *PLoS Comput. Biol.*, 1(6): e60 (2005), incorporated herein by reference. The Cas protein may be any Cas endonucleases. In some embodiments, the Cas protein is Cas9 or Cas12a, otherwise referred to as Cpf1. In one embodiment, the Cas9 protein is a wild-type Cas9 protein. The Cas9 protein can be obtained from any suitable microorganism, and a number of bacteria express Cas9 protein orthologs or variants. In some embodiments, the Cas9 is from *Streptococcus pyogenes* or *Staphylococcus aureus*. Cas9 proteins of other species are known in the art (see, e.g., U.S. Patent Application Publication 2017/0051312, incorporated herein by reference) and may be used in connection with the present disclosure. The amino acid sequences of Cas proteins from a variety of species are publicly available through the GenBank and UniProt databases.

[0065] In some embodiments, the Cas9 protein is a Cas9 nickase (Cas9n). Wild-type Cas9 has two catalytic nuclease domains facilitating double-stranded DNA breaks. A Cas9 nickase protein is typically engineered through inactivating point mutation(s) in one of the catalytic nuclease domains causing Cas9 to nick or enzymatically break only one of the two DNA strands using the remaining active nuclease domain. Cas9 nickases are known in the art (see, e.g., U.S. Patent Application Publication 2017/0051312, incorporated herein by reference) and include, for example, *Streptococcus pyogenes* with point mutations at D10 or H840. In select embodiments, the Cas9 nickase is *Streptococcus pyogenes* Cas9n (D10A).

[0066] In some embodiments, the Cas protein is a catalytically dead Cas. For example, catalytically dead Cas9 is essentially a DNA-binding protein due to, typically, two or more mutations within its catalytic nuclease domains which renders the protein with very little or no catalytic nuclease activity. *Streptococcus pyogenes* Cas9 may be rendered catalytically dead by mutations of D10 and at least one of E762, H840, N854, N863, or D986, typically H840 and/or N863 (see, e.g., U.S. Patent Application Publication 2017/0051312, incorporated herein by reference). Mutations in corresponding orthologs are known, such as N580 in *Staphylococcus aureus* Cas9. Oftentimes, such mutations cause catalytically dead Cas proteins to possess no more than 3% of the normal nuclease activity.

[0067] In some embodiments, the system comprises a nucleic acid molecule comprising a guide RNA sequence complementary to a target DNA sequence. The guide RNA sequence, as described above, specifies the target site with an approximate 20-nucleotide guide sequence followed by a protospacer adjacent motif (PAM) that directs Cas9 via Watson-Crick base pairing to a target sequence.

[0068] The terms “target DNA sequence,” “target nucleic acid,” “target sequence,” and “target site” are used interchangeably herein to refer to a polynucleotide (nucleic acid, gene, chromosome, genome, etc.) to which a guide sequence (e.g., a guide RNA) is designed to have complementarity, wherein hybridization between the target sequence and a guide sequence promotes the formation of a Cas9/CRISPR complex, provided sufficient conditions for binding exist. In some embodiments, the target sequence is a genomic DNA sequence. The term “genomic,” as used herein, refers to a nucleic acid sequence (e.g., a gene or locus) that is located on a chromosome in a cell. The target sequence and guide sequence need not exhibit complete complementarity, provided that there is sufficient complementarity to cause hybridization and promote formation of a CRISPR complex. A target sequence may comprise any polynucleotide, such as DNA or RNA. Suitable DNA/RNA binding conditions include physiological conditions normally present in a cell. Other suitable DNA/RNA binding conditions (e.g., conditions in a cell-free system) are known in the art; see, e.g., Sambrook, referenced herein and incorporated by reference. The strand of the target DNA that is complementary to and hybridizes with the DNA-targeting RNA is referred to as the “complementary strand” and the strand of the target DNA that is complementary to the “complementary strand” (and is therefore not complementary to the DNA-targeting RNA) is referred to as the “noncomplementary strand” or “non-complementary strand.”

[0069] The target genomic DNA sequence may encode a gene product. The term “gene product,” as used herein, refers to any biochemical product resulting from expression of a gene. Gene products may be RNA or protein. RNA gene products include non-coding RNA, such as tRNA, rRNA, micro RNA

(miRNA), and small interfering RNA (siRNA), and coding RNA, such as messenger RNA (mRNA). In some embodiments, the target genomic DNA sequence encodes a protein or polypeptide.

[0070] In some embodiments, for instance, when the system includes a Cas9 nickase or a catalytically dead Cas 9, two nucleic acid molecules comprising a guide RNA sequence may be utilized. The two nucleic acid molecules may have the same or different guide RNA sequences, thus complementary to the same or different target DNA sequence. In some embodiments, the guide RNA sequences of the two nucleic acid molecules are complementary to a target DNA sequences at opposite ends (e.g., 3' or 5') and/or on opposite strands of the insert location.

[0071] In some embodiments, the system further comprises a recruitment system comprising at least one aptamer sequence and an aptamer binding protein functionally linked to the microbial recombination protein as part of a fusion protein.

[0072] In some embodiments, the aptamer sequence is an RNA aptamer sequence. In some embodiments, the nucleic acid molecule comprising the guide RNA also comprises one or more RNA aptamers, or distinct RNA secondary structures or sequences that can recruit and bind another molecular species, an adaptor molecule, such as a nucleic acid or protein. The RNA aptamers can be naturally occurring or synthetic oligonucleotides that have been engineered through repeated rounds of in vitro selection or SELEX (systematic evolution of ligands by exponential enrichment) to bind to a specific target molecular species. In some embodiments, the nucleic acid comprises two or more aptamer sequences. The aptamer sequences may be the same or different and may target the same or different adaptor proteins. In select embodiments, the nucleic acid comprises two aptamer sequences.

[0073] Any RNA aptamer/ aptamer binding protein pair known may be selected and used in connection with the present disclosure (see, e.g., Jayasena, S.D., *Clinical Chemistry*, 1999. 45(9): p. 1628-1650; Gelinas, et al., *Current Opinion in Structural Biology*, 2016. 36: p. 122-132; and Hasegawa, H., *Molecules*, 2016; 21(4): p. 421, incorporated herein by reference).

[0074] A number of RNA aptamer binding, or adaptor, proteins exist, including a diverse array of bacteriophage coat proteins. Examples of such coat proteins include but are not limited to: MS2, Q β , F2, GA, fr, JP501, M12, R17, BZ13, JP34, JP500, KU1, M11, MX1, TW18, VK, SP, FI, ID2, NL95, TW19, AP205, ϕ Cb5, ϕ Cb8r, ϕ Cb12r, ϕ Cb23r, 7s and PRR1. In some embodiments, the RNA aptamer binds MS2 bacteriophage coat protein or a functional derivative, fragment or variant thereof. MS2 binding RNA aptamers commonly have a simple stem-loop structure, classically defined by a 19 nucleotide RNA molecule with a single bulged adenine on the 5' leg of the stem (Witherall G.W., et al., (1991) *Prog.*

Nucleic Acid Res. Mol. Biol., 40, 185–220, incorporated herein by reference). However, a number of vastly different primary sequences were found to be able to bind the MS2 coat protein (Parrott AM, et al., *Nucleic Acids Res.* 2000;28(2):489–497, Buenrostro JD, et al. *Natura Biotechnology* 2014; 32, 562-568, and incorporated herein by reference). Any of the RNA aptamer sequence known to bind the MS2 bacteriophage coat protein may be utilized in connection with the present disclosure. In select embodiments, the MS2 RNA aptamer sequence comprises: AAC AUGAGGAUCACCCAUGUCUGCAG (SEQ ID NO:145), AGCAUGAGGAUCACCCAUGUCUGCAG (SEQ ID NO:146), or AGCGUGAGGAUCACCCAUGCCUGCAG (SEQ ID NO:147).

[0075] N-proteins (Nut-utilization site proteins) of bacteriophages contain arginine-rich conserved RNA recognition motifs of ~20 amino acids, referred to as N peptides. The RNA aptamer may bind a phage N peptide or a functional derivative, fragment or variant thereof. In some embodiments, the phage N peptide is the lambda or P22 phage N peptide or a functional derivative, fragment or variant thereof.

[0076] In select embodiments, the N peptide is lambda phage N22 peptide, or a functional derivative, fragment or variant thereof. In some embodiments, the N22 peptide comprises an amino acid sequence with at least 70% similarity to the amino acid sequence GNARTRRRERRAEKQAQWKAAN (SEQ ID NO: 149). N22 peptide, the 22 amino acid RNA-binding domain of the λ bacteriophage antiterminator protein N (λ N-(1–22) or λ N peptide), is capable of specifically binding to specific stem-loop structures, including but not limited to the BoxB stem-loop. See, for example Cilley and Williamson, *RNA* 1997; 3(1):57-67, incorporated herein by reference. A number of different BoxB stem-loop primary sequences are known to bind the N22 peptide and any of those may be utilized in connection with the present disclosure. In some embodiments, the N22 peptide RNA aptamer sequence comprises a nucleotide sequence with at least 70% similarity to an RNA sequence selected from the group consisting of GCCCUGAAAAAGGGC (SEQ ID NO: 150), GCCCUGAAGAAGGGC (SEQ ID NO: 151), GCGCUGAAAAAGCGC (SEQ ID NO: 152), GCCCUGACAAAGGGC (SEQ ID NO: 153), and GCGCUGACAAAGCGC (SEQ ID NO: 154). In some embodiments, the N22 peptide RNA aptamer sequence is selected from the group consisting of SEQ ID NOs: 150-154.

[0077] In select embodiments, the N peptide is the P22 phage N peptide, or a functional derivative, fragment or variant thereof. A number of different BoxB stem-loop primary sequences are known to bind the P22 phage N peptide and variants thereof and any of those may be utilized in connection with the present disclosure. See, for example Cocozaki, Ghattas, and Smith, *Journal of Bacteriology* 2008; 190(23):7699-7708, incorporated herein by reference. In some embodiments, the P22 phage N peptide

comprises an amino acid sequence with at least 70% similarity to the amino acid sequence GNAKTRRHERRRKLAIERDTI (SEQ ID NO: 155). In some embodiments, the P22 phage N peptide RNA aptamer sequence comprises a sequence with at least 70% similarity to an RNA sequence selected from the group consisting of GCGCUGACAAAGCGC (SEQ ID NO: 156) and CCGCCGACAACGCGG (SEQ ID NO: 157). In some embodiments, the P22 phage N peptide RNA aptamer sequence is selected from the group consisting of SEQ ID NOs: 156-157, UGCGCUGACAAAGCGCG (SEQ ID NO: 158) or ACCGCCGACAACGCGGU (SEQ ID NO: 159).

[0078] In some embodiments, the aptamer sequence is a peptide aptamer sequence. The peptide aptamers can be naturally occurring or synthetic peptides that are specifically recognized by an affinity agent. Such aptamers include, but are not limited to, a c-Myc affinity tag, an HA affinity tag, a His affinity tag, an S affinity tag, a methionine-His affinity tag, an RGD-His affinity tag, a 7× His tag, a FLAG octapeptide, a strep tag or strep tag II, a V5 tag, or a VSV-G epitope. Corresponding aptamer binding proteins are well-known in the art and include, for example, primary antibodies, biotin, affimers, single domain antibodies, and antibody mimetics.

[0079] An exemplary peptide aptamer includes a GCN4 peptide (Tanenbaum et al., *Cell* 2014; 159(3):635-646, incorporated herein by reference). Antibodies, or GCN4 binding protein can be used as the aptamer binding proteins.

[0080] In some embodiments, the peptide aptamer sequence is conjugated to the Cas protein. The peptide aptamer sequence may be fused to the Cas in any orientation (e.g., N-terminus to C-terminus, C-terminus to N-terminus, N-terminus to N-terminus). In select embodiments, the peptide aptamer is fused to the C-terminus of the Cas protein.

[0081] In some embodiments, between 1 and 24 peptide aptamer sequences may be conjugated to the Cas protein. The aptamer sequences may be the same or different and may target the same or different aptamer binding proteins. In select embodiments, 1 to 24 tandem repeats of the same peptide aptamer sequence are conjugated to the Cas protein. In preferred embodiments between 4 and 18 tandem repeats are conjugated to the Cas protein. The individual aptamers may be separated by a linker region. Suitable linker regions are known in the art. The linker may be flexible or configured to allow the binding of affinity agents to adjacent aptamers without or with decreased steric hindrance. The linker sequences may provide an unstructured or linear region of the polypeptide, for example, with the inclusion of one or more glycine and/or serine residues. The linker sequences can be at least about 2, 3, 4, 5, 6, 7, 8, 9, 10 or more amino acids in length.

[0082] In some embodiments, the fusion protein comprises a microbial recombination protein functionally linked to an aptamer binding protein. The microbial recombination protein may be RecE, RecT, lambda exonuclease (Exo), Bet protein (betA, redB), exonuclease gp6, single-stranded DNA-binding protein gp2.5, or a derivative or variant thereof.

[0083] In select embodiments, the microbial recombination protein is RecE or RecT, or a derivative or variant thereof. Derivatives or variants of RecE and RecT are functionally equivalent proteins or polypeptides which possess substantially similar function to wild type RecE and RecT. RecE and RecT derivatives or variants include biologically active amino acid sequences similar to the wild-type sequences but differing due to amino acid substitutions, additions, deletions, truncations, post-translational modifications, or other modifications. In some embodiments, the derivatives may improve translation, purification, biological half-life, activity, or eliminate or lessen any undesirable side effects or reactions. The derivatives or variants may be naturally occurring polypeptides, synthetic or chemically synthesized polypeptides or genetically engineered peptide polypeptides. RecE and RecT bioactivities are known to, and easily assayed by, those of ordinary skill in the art, and include, for example exonuclease and single-stranded nucleic acid binding, respectively.

[0084] The RecE or RecT may be from a number of microbial organisms, including *Escherichia coli*, *Pantoea breeneri*, Type-F symbiont of *Plautia stali*, *Providencia* sp. MGF014, *Shigella sonnei*, *Pseudobacteriovorax antillogorgiicola*, among others. In preferred embodiments, the RecE and RecT protein is derived from *Escherichia coli*.

[0085] In some embodiments, the fusion protein comprises RecE, or a derivative or variant thereof. The RecE, or derivative or variant thereof, may comprise an amino acid sequence selected from the group consisting of SEQ ID NOs: 1-8. The RecE, or derivative or variant thereof, may comprise an amino acid sequences with at least 70% (e.g., 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100%) similarity to amino acid sequences selected from the group consisting of SEQ ID NOs: 1-8. In select embodiments, the RecE, or derivative or variant thereof, comprises an amino acid sequences with at least 90% similarity to amino acid sequences selected from the group consisting of SEQ ID NOs: 1-8. In exemplary embodiments, the RecE, or derivative or variant thereof, comprises an amino acid sequences with at least 90% similarity to amino acid sequences selected from the group consisting of SEQ ID NOs: 1-3.

[0086] In some embodiments, the fusion protein comprises RecT, or a derivative or variant thereof. The RecT, or derivative or variant thereof, may comprise an amino acid sequence selected from the group

consisting of SEQ ID NOs: 9-14. The RecT, or derivative or variant thereof, may comprise an amino acid sequences with at least 70% (e.g., 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100%) similarity to amino acid sequences selected from the group consisting of SEQ ID NOs: 9-14. In select embodiments, the RecT, or derivative or variant thereof, comprises an amino acid sequences with at least 90% similarity to amino acid sequences selected from the group consisting of SEQ ID NOs: 9-14. In exemplary embodiments, the RecT, or derivative or variant thereof, comprises an amino acid sequences with at least 90% similarity to amino acid sequences selected from the group consisting of SEQ ID NO: 9.

[0087] Truncations may be from either the C-terminal or N-terminal ends, or both. For example, as demonstrated in Example 6 below, a diverse set of truncations from either end or both provided a functional product. In some embodiments, one or more (2, 3, 4, 5, 10, 20, 30, 40, 50, 60, 100, 120 or more) amino acids may be truncated from the C-terminal, N-terminal ends as compared to the wild-type sequence.

[0088] In the fusion protein, the microbial recombination protein may be linked to either terminus of the aptamer binding protein in any orientation (e.g., N-terminus to C-terminus, C-terminus to N-terminus, N-terminus to N-terminus). In select embodiments, the microbial recombination protein N-terminus is linked to the aptamer binding protein C-terminus. Thus, the overall fusion protein from N- to C-terminus comprises the aptamer binding protein (N- to C-terminus) linked to the microbial recombination protein (N- to C-terminus).

[0089] In some embodiments, the fusion protein further comprises a linker between the microbial recombination protein and the aptamer binding protein. The linkers may comprise any amino acid sequence of any length. The linkers may be flexible such that they do not constrain either of the two components they link together in any particular orientation. The linkers may essentially act as a spacer. In select embodiments, the linker links the C-terminus of the microbial recombination protein to the N-terminus of the aptamer binding protein. In select embodiments, the linker comprises the amino acid sequence of the 16-residue XTEN linker, SGSETPGTSESATPES (SEQ ID NO: 15) or the 37-residue EXTEN linker, SASGGSSGGSSGSETPGTSESATPESGGSSGGSGGS (SEQ ID NO: 148).

[0090] In some embodiments, the fusion protein further comprises a nuclear localization sequence (NLS). The nuclear localization sequence may be at any location within the fusion protein (e.g., C-terminal of the aptamer binding protein, N-terminal of the aptamer binding protein, C-terminal of the microbial recombination protein). In select embodiments, the nuclear localization sequence is linked to

the C-terminus of the microbial recombination protein. A number of nuclear localization sequences are known in the art (see, e.g., Lange, A., et al., J Biol Chem. 2007; 282(8): 5101-5105, incorporated herein by reference) and may be used in connection with the present disclosure. The nuclear localization sequence may be the SV40 NLS, PKKKRKV (SEQ ID NO:16); the Ty1 NLS, NSKKRSLEDNETEIKVSRDTWNTKNMRSLEPPRSKKRIH (SEQ ID NO: 17); the c-Myc NLS, PAAKRVKLD (SEQ ID NO:18); the biSV40 NLS, KRTADGSEFESPKKKRKV (SEQ ID NO: 19); and the Mut NLS, PEKKRRRPSGSVPVLARSPPKAGKSSCI (SEQ ID NO: 20). In select embodiments, the nuclear localization sequence is the SV40 NLS, PKKKRKV (SEQ ID NO: 16).

[0091] The Cas protein and the fusion protein are desirably included in a single composition alone, in combination with each other, and/or the polynucleotide(s) (e.g., a vector) comprising the guide RNA sequence and the aptamer sequence. The Cas protein and/or the fusion protein may or may not be physically or chemically bound to the polynucleotide. The Cas protein and/or the microbial recombination protein can be associated with a polynucleotide using any suitable method for protein-protein linking or protein-virus linking known in the art.

[0092] The disclosure further provides compositions and vectors comprising a polynucleotide comprising a nucleic acid sequence encoding a fusion protein comprising a microbial recombination protein functionally linked to an RNA aptamer binding protein.

[0093] The compositions or vectors may further comprise at least one or both of a polynucleotide comprising a nucleic acid sequence encoding a Cas protein and a nucleic acid molecule comprising a guide RNA sequence that is complementary to a target DNA sequence. In some embodiments, the nucleic acid molecule comprising a guide RNA sequence further comprises at least one RNA aptamer sequence. In some embodiments, the polynucleotide comprising a nucleic acid sequence encoding a Cas protein further comprises a sequence encoding at least one peptide aptamer sequence.

[0094] Descriptions of the nucleic acid molecule comprising a guide RNA sequence, the aptamer sequences, the Cas proteins, the microbial recombination proteins, and the aptamer binding proteins set forth above in connection with the inventive system also are applicable to the polynucleotides of the recited compositions and vectors.

[0095] The nucleic acid sequence encoding the Cas protein and/or the nucleic acid sequence encoding a fusion protein comprising a microbial recombination protein functionally linked to an aptamer binding protein can be provided to a cell on the same vector (e.g., in *cis*) as the nucleic acid molecule comprising the guide RNA sequence and/or the RNA aptamer sequence. In such embodiments, a unidirectional

promoter can be used to control expression of each nucleic acid sequence. In another embodiment, a combination of bidirectional and unidirectional promoters can be used to control expression of multiple nucleic acid sequences.

[0096] In other embodiments, a nucleic acid sequence encoding the Cas protein, the nucleic acid sequence encoding a fusion protein comprising a microbial recombination protein functionally linked to an aptamer binding protein, and the nucleic acid molecule comprising the guide RNA sequence and/or the RNA aptamer sequence can be provided to a cell on separate vectors (e.g., in *trans*). Each of the nucleic acid sequences in each of the separate vectors can comprise the same or different expression control sequences. The separate vectors can be provided to cells simultaneously or sequentially.

[0097] The vector(s) comprising the nucleic acid sequences encoding the Cas protein and encoding a fusion protein comprising a microbial recombination protein functionally linked to an aptamer binding protein can be introduced into a host cell that is capable of expressing the polypeptide encoded thereby, including any suitable prokaryotic or eukaryotic cell. As such, the disclosure provides an isolated cell comprising the vector or nucleic acid sequences disclosed herein. Preferred host cells are those that can be easily and reliably grown, have reasonably fast growth rates, have well characterized expression systems, and can be transformed or transfected easily and efficiently. Examples of suitable prokaryotic cells include, but are not limited to, cells from the genera *Bacillus* (such as *Bacillus subtilis* and *Bacillus brevis*), *Escherichia* (such as *E. coli*), *Pseudomonas*, *Streptomyces*, *Salmonella*, and *Envinia*. Suitable eukaryotic cells are known in the art and include, for example, yeast cells, insect cells, and mammalian cells. Examples of suitable yeast cells include those from the genera *Kluyveromyces*, *Pichia*, *Rhizosporidium*, *Saccharomyces*, and *Schizosaccharomyces*. Exemplary insect cells include Sf-9 and HIS (Invitrogen, Carlsbad, Calif.) and are described in, for example, Kitts et al., *Biotechniques*, 14: 810-817 (1993); Lucklow, *Curr. Opin. Biotechnol.*, 4: 564-572 (1993); and Lucklow et al., *J. Virol.*, 67: 4566-4579 (1993), incorporated herein by reference. Desirably, the host cell is a mammalian cell, and in some embodiments, the host cell is a human cell. A number of suitable mammalian and human host cells are known in the art, and many are available from the American Type Culture Collection (ATCC, Manassas, Va.). Examples of suitable mammalian cells include, but are not limited to, Chinese hamster ovary cells (CHO) (ATCC No. CCL61), CHO DHFR-cells (Urlaub et al., *Proc. Natl. Acad. Sci. USA*, 97: 4216-4220 (1980)), human embryonic kidney (HEK) 293 or 293T cells (ATCC No. CRL1573), and 3T3 cells (ATCC No. CCL92). Other suitable mammalian cell lines are the monkey COS-1 (ATCC No. CRL1650) and COS-7 cell lines (ATCC No. CRL1651), as well as the CV-1 cell line (ATCC No. CCL70). Further

exemplary mammalian host cells include primate, rodent, and human cell lines, including transformed cell lines. Normal diploid cells, cell strains derived from *in vitro* culture of primary tissue, as well as primary explants, are also suitable. Other suitable mammalian cell lines include, but are not limited to, mouse neuroblastoma N2A cells, HeLa, HEK, A549, HepG2, mouse L-929 cells, and BHK or HaK hamster cell lines. Methods for selecting suitable mammalian host cells and methods for transformation, culture, amplification, screening, and purification of cells are known in the art.

3. Methods of Altering Target DNA

[0098] The disclosure also provides a method of altering a target DNA. In some embodiments, the method alters genomic DNA sequence in a cell, although any desired nucleic acid may be modified. When applied to DNA contained in cells, the method comprises introducing the systems, compositions, or vectors described herein into a cell comprising a target genomic DNA sequence. Descriptions of the nucleic acid molecule comprising a guide RNA sequence, the Cas proteins, the microbial recombination proteins, the recruitment systems, and polynucleotides encoding thereof, the cell, the target genomic DNA sequence, and components thereof, set forth above in connection with the inventive system are also applicable to the method of altering a target genomic DNA sequence in a cell. The systems, composition or vectors may be introduced in any manner known in the art including, but not limited to, chemical transfection, electroporation, microinjection, biolistic delivery via gene guns, or magnetic-assisted transfection, depending on the cell type.

[0099] Upon introducing the systems described herein into a cell comprising a target genomic DNA sequence, the guide RNA sequence binds to the target genomic DNA sequence in the cell genome, the Cas protein associates with the guide RNA and may induce a double strand break or single strand nick in the target genomic DNA sequence and the aptamer recruits the microbial recombination proteins to the target genomic DNA sequence through the aptamer binding protein of the fusion protein, thereby altering the target genomic DNA sequence in the cell. When introducing the compositions, or vectors described herein into the cell, the nucleic acid molecule comprising a guide RNA sequence, the Cas9 protein, and the fusion protein are first expressed in the cell.

[0100] In some embodiments, the cell is in an organism or host, such that introducing the disclosed systems, compositions, vectors into the cell comprises administration to a subject. The method may comprise providing or administering to the subject, *in vivo*, or by transplantation of *ex vivo* treated cells, systems, compositions, vectors of the present system.

[00101] A “subject” may be human or non-human and may include, for example, animal strains or species used as “model systems” for research purposes, such a mouse model as described herein. Likewise, subject may include either adults or juveniles (e.g., children). Moreover, subject may mean any living organism, preferably a mammal (e.g., human or non-human) that may benefit from the administration of compositions contemplated herein. Examples of mammals include, but are not limited to, any member of the Mammalian class: humans, non-human primates such as chimpanzees, and other apes and monkey species; farm animals such as cattle, horses, sheep, goats, swine; domestic animals such as rabbits, dogs, and cats; laboratory animals including rodents, such as rats, mice and guinea pigs, and the like. Examples of non-mammals include, but are not limited to, birds, fish, and the like. In one embodiment of the methods and compositions provided herein, the mammal is a human.

[00102] As used herein, the terms “providing”, “administering,” “introducing,” are used interchangeably herein and refer to the placement of the systems of the disclosure into a subject by a method or route which results in at least partial localization of the system to a desired site. The systems can be administered by any appropriate route which results in delivery to a desired location in the subject.

[00103] The phrase “altering a DNA sequence,” as used herein, refers to modifying at least one physical feature of a DNA sequence of interest. DNA alterations include, for example, single or double strand DNA breaks, deletion, or insertion of one or more nucleotides, and other modifications that affect the structural integrity or nucleotide sequence of the DNA sequence. The modifications of a target sequence in genomic DNA may lead to, for example, gene correction, gene replacement, gene tagging, transgene insertion, nucleotide deletion, gene disruption, gene mutation, gene knock-down, and the like.

[00104] In some embodiments, the systems and methods described herein may be used to correct one or more defects or mutations in a gene (referred to as “gene correction”). In such cases, the target genomic DNA sequence encodes a defective version of a gene, and the system further comprises a donor nucleic acid molecule which encodes a wild-type or corrected version of the gene. Thus, in other words, the target genomic DNA sequence is a “disease-associated” gene. The term “disease-associated gene,” refers to any gene or polynucleotide whose gene products are expressed at an abnormal level or in an abnormal form in cells obtained from a disease-affected individual as compared with tissues or cells obtained from an individual not affected by the disease. A disease-associated gene may be expressed at an abnormally high level or at an abnormally low level, where the altered expression correlates with the occurrence and/or progression of the disease. A disease-associated gene also refers to a gene, the mutation or genetic variation of which is directly responsible or is in linkage disequilibrium with a gene(s) that is responsible

for the etiology of a disease. Examples of genes responsible for such “single gene” or “monogenic” diseases include, but are not limited to, adenosine deaminase, α -1 antitrypsin, cystic fibrosis transmembrane conductance regulator (CFTR), β -hemoglobin (HBB), oculocutaneous albinism II (OCA2), Huntingtin (HTT), dystrophin myotonia-protein kinase (DMPK), low-density lipoprotein receptor (LDLR), apolipoprotein B (APOB), neurofibromin 1 (NF1), polycystic kidney disease 1 (PKD1), polycystic kidney disease 2 (PKD2), coagulation factor VIII (F8), dystrophin (DMD), phosphate-regulating endopeptidase homologue, X-linked (PHEX), methyl-CpG-binding protein 2 (MECP2), and ubiquitin-specific peptidase 9Y, Y-linked (USP9Y). Other single gene or monogenic diseases are known in the art and described in, e.g., Chial, H. *Rare Genetic Disorders: Learning About Genetic Disease Through Gene Mapping, SNPs, and Microarray Data*, *Nature Education* 1(1):192 (2008), incorporated herein by reference; Online Mendelian Inheritance in Man (OMIM); and the Human Gene Mutation Database (HGMD).

[00105] In another embodiment, the target genomic DNA sequence can comprise a gene, the mutation of which contributes to a particular disease in combination with mutations in other genes. Diseases caused by the contribution of multiple genes which lack simple (e.g., Mendelian) inheritance patterns are referred to in the art as a “multifactorial” or “polygenic” disease. Examples of multifactorial or polygenic diseases include, but are not limited to, asthma, diabetes, epilepsy, hypertension, bipolar disorder, and schizophrenia. Certain developmental abnormalities also can be inherited in a multifactorial or polygenic pattern and include, for example, cleft lip/palate, congenital heart defects, and neural tube defects.

[00106] In another embodiment, the method of altering a target genomic DNA sequence can be used to delete nucleic acids from a target sequence in a cell by cleaving the target sequence and allowing the cell to repair the cleaved sequence in the absence of an exogenously provided donor nucleic acid molecule. Deletion of a nucleic acid sequence in this manner can be used in a variety of applications, such as, for example, to remove disease-causing trinucleotide repeat sequences in neurons, to create gene knock-outs or knock-downs, and to generate mutations for disease models in research.

[00107] The term “donor nucleic acid molecule” refers to a nucleotide sequence that is inserted into the target DNA (e.g., genomic DNA). As described above the donor DNA may include, for example, a gene or part of a gene, a sequence encoding a tag or localization sequence, or a regulating element. The donor nucleic acid molecule may be of any length. In some embodiments, the donor nucleic acid molecule is between 10 and 10,000 nucleotides in length. For example, between about 100 and 5,000 nucleotides in length, between about 200 and 2,000 nucleotides in length, between about 500 and 1,000 nucleotides in

length, between about 500 and 5,000 nucleotides in length, between about 1,000 and 5,000 nucleotides in length, or between about 1,000 and 10,000 nucleotides in length,

[00108] The disclosed systems and methods overcome challenges encountered during conventional gene editing, including low efficiency and off-target events, particularly with kilobase-scale nucleic acids. In some embodiments, the disclosed systems and methods improve the efficiency of gene editing. For example, the disclosed systems and methods can have a 2- to 10-fold increase in efficiency over conventional CRISPR-Cas9 systems and methods, as shown in Examples 2, 3, and 5. In some embodiments, the improvement in efficiency is accompanied by a reduction in off-target events. The off-target events may be reduced by greater than 50% compared to conventional CRISPR-Cas9 systems and methods, for example, a reduction of off-target events by about 90% is shown in Example 3. Another aspect of increasing the overall accuracy of a gene editing system is reducing the on-target insertion-deletions (indels), a byproduct of HDR editing. In some embodiments, the disclosed systems and methods reduce the on-target indels by greater than 90% compared to conventional CRISPR-Cas9 systems and methods, as shown in Example 3.

[00109] The disclosure further provides kits containing one or more reagents or other components useful, necessary, or sufficient for practicing any of the methods described herein. For example, kits may include CRISPR reagents (Cas protein, guide RNA, vectors, compositions, etc.), recombineering reagents (recombination protein-aptamer binding protein fusion protein, the aptamer sequence, vectors, compositions, etc.) transfection or administration reagents, negative and positive control samples (e.g., cells, template DNA), cells, containers housing one or more components (e.g., microcentrifuge tubes, boxes), detectable labels, detection and analysis instruments, software, instructions, and the like.

[00110] Any element of any suitable CRISPR/Cas gene editing system known in the art can be employed in the systems and methods described herein, as appropriate. CRISPR/Cas gene editing technology is described in detail in, for example, U.S. Patent Nos. 8,546,553; 8,697,359; 8,771,945; 8,795,965; 8,865,406; 8,871,445; 8,889,356; 8,889,418; 8,895,308; 8,9066,616; 8,932,814; 8,945,839; 8,993,233; 8,999,641; 9,115,348; 9,149,049; 9,493,844; 9,567,603; 9,637,739; 9,663,782; 9,404,098; 9,885,026; 9,951,342; 10,087,431; 10,227,610; 10,266,850; 10,601,748; 10,604,771; and 10,760,064; and U.S. Patent Application Publication Nos. US2010/0076057; US2014/0113376; US2015/0050699; US2015/0031134; US2014/0357530; US2014/0349400; US2014/0315985; US2014/0310830; US2014/0310828; US2014/0309487; US2014/0294773; US2014/0287938; US2014/0273230; US2014/0242699; US2014/0242664; US2014/0212869; US2014/0201857; US2014/0199767;

US2014/0189896; US2014/0186919; US2014/0186843; and US2014/0179770, each incorporated herein by reference.

[00111] The following examples further illustrate the invention but should not be construed as in any way limiting its scope.

EXAMPLES

Materials and Methods

[00112] *RecE/T Homolog Screening* RefSeq non-redundant protein database was downloaded from NCBI on October 29, 2019. The database was searched with *E. coli* Rac prophage RecT (NP_415865.1) and RecE (NP_415866.1) as queries using position-specific iterated (PSI)-BLAST¹ to retrieve protein homologs. Hits were clustered with CD-HIT2 and representative sequences were selected from each cluster for multiple alignment with MUSCLE³. Then, FastTree4 was used for maximum likelihood tree reconstruction with default parameters. A diverse set of RecET homologs were selected, synthesized by GenScript, and cloned into pMPH_MCP vectors for testing.

[00113] *Plasmids construction* pX330, pMPH and pU6-(BbsI)_CBh-Cas9-T2A-BFP plasmids were obtained from Addgene. Tested effector DNA fragments were ordered from IDT, Genewiz, and GenScript. The fragments were Gibson assembled into the backbones using NEBuilder HiFi DNA Assembly Master Mix (New England BioLabs). All sgRNAs (Table 1) were inserted into backbones using Golden Gate cloning. All constructs were sequence-verified with Sanger sequencing of prepped plasmids.

Table 1. Sequence for sgRNAs

Primer Name	Genomic Target	Sequence
sp-EMX1	<i>EMX1</i>	GTCACCTCCAATGACTAGGG (SEQ ID NO:21)
sp-VEGFA	<i>VEGFA</i>	GGTGAGTGAGTGTGTGCGTG (SEQ ID NO:22)
sp-DYNLT1	<i>DYNLT1</i>	AAGGCCATAGGCTGGACTGC (SEQ ID NO:23)
sp-HSP90AA1	<i>HSP90AA1</i>	GTAGACTAATCTCTGGCTGA (SEQ ID NO:24)
sp-OCT4	<i>OCT4</i>	TCTCCCATGCATTCAAAGTG (SEQ ID NO:25)
sp-AAVS1	<i>AAVS1</i>	ACCCACAGTGGGGCCACTA (SEQ ID NO:26)
nsp-EMX1-guide1	<i>EMX1</i>	GTCACCTCCAATGACTAGGG (SEQ ID NO:27)
nsp-EMX1-guide2	<i>EMX1</i>	GTCACCTCCAATGACTAGGG (SEQ ID NO:28)

nsp-DYNLT1-guide1	<i>DYNLT1</i>	AAGGCCATAGGCTGGACTGC (SEQ ID NO:29)
nsp-DYNLT1-guide2	<i>DYNLT1</i>	GGCACTGACGATGCAGTACA (SEQ ID NO:30)
nsp-HSP90AA1-guide1	<i>HSP90AA1</i>	GTAGACTAATCTCTGGCTGA (SEQ ID NO:31)
nsp-HSP90AA1-guide2	<i>HSP90AA1</i>	TCGTCATCTCCTTCAAGGGG (SEQ ID NO:32)
nsp-OCT4-guide1	<i>OCT4</i>	ATGCATGGGAGAGCCCAGAG (SEQ ID NO:33)
nsp-OCT4-guide2	<i>OCT4</i>	GCCTGCCCTTCTAGGAATGG (SEQ ID NO:34)

[00114] *Cell culture* Human Embryonic Kidney (HEK) 293T, HeLa and HepG2 were maintained in Dulbecco's Modified Eagle's Medium (DMEM, Life Technologies), with 10% fetal bovine serum (FBS, HyClone), 100 U/mL penicillin, and 100 µg/mL streptomycin (Life Technologies) at 37 °C with 5% CO₂.

[00115] hES-H9 cells were maintained in mTeSR1 medium (StemCell Technologies) at 37 °C with 5% CO₂. Culture plates were pre-coated with Matrigel (Corning) 12 hours prior to use, and cells were supplemented with 10 µM Y27632 (Sigma) for the first 24 hours after passaging. Culture media was changed every 24 hours.

[00116] *Transfection* HEK293T cells were seeded into 96-well plates (Corning) 12-24 hours prior to transfection at a density of 30,000 cells/well, and 250 ng of total DNA was transfected per well. HeLa and HepG2 cells were seeded into 48-well plates (Corning) one day prior to transfection at a density of 50,000 and 30,000 cells/well respectively, and 400 ng of total DNA was transfected per well. Transfections were performed with Lipofectamine 3000 (Life Technologies) following the manufacturer's instructions.

[00117] *Electroporation* For hES-H9 related transfection experiments, P3 Primary Cell 4D-Nucleofector™ X Kit S (Lonza) was used following the manufacturer's protocol. For each reaction, 300,000 cells were nucleofected with 4 µg total DNA using the DC100 Nucleofector Program.

[00118] *Fluorescence-activated cell sorting (FACS)* mKate knock-in efficiency was analyzed on a CytoFLEX flow cytometer (Beckman Coulter; Stanford Stem Cell FACS Core). 72 hours after transfection, cells were washed once with PBS and dissociated with TrypLE Express Enzyme (Thermo Fisher Scientific). Cell suspension was then transferred to a 96-well U-bottom plate (Thermo Fisher Scientific) and centrifuged at 300xG for 5 minutes. After removing the supernatant, pelleted cells were resuspended with 50 µl 4% FBS in PBS, and cells were sorted within 30 minutes of preparation.

[00119] *RFLP* HEK293T cells were transfected with plasmid DNA and PCR templates and harvested after 72 hours for genomic DNA using the QuickExtract DNA Extraction Solution (Biosearch

Technologies) following the manufacturer's protocol. The target genomic region was amplified using specific primers outside of the homology arms of the PCR template. PCR products were purified with Monarch PCR & DNA Cleanup Kit (New England BioLabs). 300 ng of purified product was digested with BsrGI (EMX1, New England BioLabs) or XbaI (VEGFA, NEB), and the digested products were analyzed on a 5% Mini-PROTEAN TBE gel (Bio-Rad).

[00120] *Next-Generation Sequencing Library Preparation* 72 hours after transfection, genomic DNA was extracted using QuickExtract DNA Extraction Solution (Biosearch Technologies). 200 ng total DNA was used for NGS library preparation. Genes of interest were amplified using specific primers (Table 2) for the first round PCR reaction. Illumina adapters and index barcodes were added to the fragments with a second round PCR using the primers listed in Table 2. Round 2 PCR products were purified by gel electrophoresis on a 2% agarose gel using the Monarch DNA Gel Extraction Kit (NEB). The purified product was quantified with Qubit dsDNA HS Assay Kit (Thermo Fisher) and sequenced on an Illumina MiSeq according to the manufacturer's instructions.

Table 2. Sequence for primers used for PCR template, RFLP and NGS

Primer Name	Usage	Genomic Target	Sequence
EMX1-PCR-F	PCR template	<i>EMX1</i>	CATTCTGCCTCTCTGTATGGAAAAGAGC (SEQ ID NO:35)
EMX1-PCR-R	PCR template	<i>EMX1</i>	CCCATTGAACTACCTGGGCCTGATTC (SEQ ID NO:36)
VEGFA-PCR-F	PCR template	<i>VEGFA</i>	AGGTTTGAATCATCACGCAGGC (SEQ ID NO:37)
VEGFA-PCR-R	PCR template	<i>VEGFA</i>	ATTCAAGTGGGGAATGGCAAGC (SEQ ID NO:38)
DYNLT1-PCR-100bp-F	PCR template	<i>DYNLT1</i>	TGCCGTAAATGCTGCTCTCT (SEQ ID NO:39)
DYNLT1-PCR-200bp-F	PCR template	<i>DYNLT1</i>	AGACTTGCCAAGGTTCTTTGTG (SEQ ID NO:40)
DYNLT1-PCR-400bp-F	PCR template	<i>DYNLT1</i>	AGTGACCTGTGTAATTATGCAGAAG (SEQ ID NO:41)
DYNLT1-PCR-100bp-R	PCR template	<i>DYNLT1</i>	TGAAAGTGCCACAAAACAAAGAGA (SEQ ID NO:42)
DYNLT1-PCR-200bp-R	PCR template	<i>DYNLT1</i>	AAGACAAGTGGCAACGCAG (SEQ ID NO:43)
DYNLT1-PCR-400bp-R	PCR template	<i>DYNLT1</i>	CGTTTATGATACTATGCAGACTATGAAGAA C (SEQ ID NO:44)
HSP90AA1-PCR-100bp-F	PCR template	<i>HSP90AA1</i>	ATGAAGATGACCCTACTGCTGAT (SEQ ID NO:45)

HSP90AA1-PCR-200bp-F	PCR template	<i>HSP90AA1</i>	TACTGTCTTGAAAGCAGATAGAAACC (SEQ ID NO:46)
HSP90AA1-PCR-600bp-R	PCR template	<i>HSP90AA1</i>	GCAGCAAAGAAACACCTGGA (SEQ ID NO:47)
HSP90AA1-PCR-100bp-R	PCR template	<i>HSP90AA1</i>	GTTGTCATGCCATACAGACTTTTT (SEQ ID NO:48)
HSP90AA1-PCR-200bp-R	PCR template	<i>HSP90AA1</i>	AGCATTACTAGCTCTGCTTTAGTG (SEQ ID NO:49)
HSP90AA1-PCR-600bp-R	PCR template	<i>HSP90AA1</i>	TCCACAAGACTGGGTCTGAG (SEQ ID NO:50)
OCT4-PCR-F	PCR template	<i>OCT4</i>	GCGACTATGCACAACGAGAGG (SEQ ID NO:51)
OCT4-PCR-R	PCR template	<i>OCT4</i>	AAGTGTGTCTATCTACTGTGTCCCAG (SEQ ID NO:52)
AAVS1-PCR-F	PCR template	<i>AAVS1</i>	GATGCTCTTTCCGGAGCACT (SEQ ID NO:53)
AAVS1-PCR-R	PCR template	<i>AAVS1</i>	GCCAAGGACTCAAACCCAGAA (SEQ ID NO:54)
EMX1-RFLP-F	RFLP	<i>EMX1</i>	TGGTGGATTTCGGACTACCCT (SEQ ID NO:55)
EMX1-RFLP-R	RFLP	<i>EMX1</i>	TTCGGACTGGAACCGTCAGC (SEQ ID NO:56)
VEGFA-RFLP-F	RFLP	<i>VEGFA</i>	AGACGTTCCCTTAGTGCTGGC (SEQ ID NO:57)
VEGFA-RFLP-R	RFLP	<i>VEGFA</i>	AAAAGTTTCAGTGCGACGCC (SEQ ID NO:58)
DYNLT1 KI PCR-F	Junction PCR	<i>DYNLT1</i>	AGGAGGTCCCATCAGATGCT (SEQ ID NO:59)
HSP90AA1 KI PCR-F	Junction PCR	<i>HSP90AA1</i>	GGCTGGACAGCAAACATGGA (SEQ ID NO:60)
AAVS1 KI PCR-F	Junction PCR	<i>AAVS1</i>	GATGCTCTTTCCGGAGCACT (SEQ ID NO:61)
Junction PCR universal-R	Junction PCR	<i>mKate</i>	TTGCTGCCGTACATGAAGCTG (SEQ ID NO:62)
EMX1-NGS-F	NGS	<i>EMX1</i>	CCATCTCATCCCTGCGTGTCTCCAGAAGAAGGGCTCCCATCAC (SEQ ID NO:63)
EMX1-NGS-R	NGS	<i>EMX1</i>	CCTCTCTATGGGCAGTCGGTGATgAGCAGCAAGCAGCACTCTG (SEQ ID NO:64)
VEGFA-NGS-F	NGS	<i>VEGFA</i>	CCATCTCATCCCTGCGTGTCTCCAGCGTCTTCGAGAGTGAGG (SEQ ID NO:65)
VEGFA-NGS-R	NGS	<i>VEGFA</i>	CCTCTCTATGGGCAGTCGGTGATgTTGGAATCCTGGAGTGACCC (SEQ ID NO:66)
EMX-OT1-F	Off Target	<i>EMX1 OT-1</i>	CCATCTCATCCCTGCGTGTCTCCACAAAA GCTCCACATGCTAGGA (SEQ ID NO:67)
EMX-OT1-R	Off Target	<i>EMX1 OT-1</i>	CCTCTCTATGGGCAGTCGGTGATgGCTGACTTTGGGCTCCTTCT (SEQ ID NO:68)

EMX-OT2-F	Off Target	<i>EMX1 OT-2</i>	<u>CCATCTCATCCCTGCGTGTCTCCACACAC</u> TCCCAGGATCTCA (SEQ ID NO:69)
EMX-OT2-R	Off Target	<i>EMX1 OT-2</i>	<u>CCTCTCTATGGGCAGTCGGTGATgAATGT</u> CAGCTGAAGCAGGCT (SEQ ID NO:70)
EMX-OT3-F	Off Target	<i>EMX1 OT-3</i>	<u>CCATCTCATCCCTGCGTGTCTCCGGCTAC</u> CCTGACAACTGCTT (SEQ ID NO:71)
EMX-OT3-R	Off Target	<i>EMX1 OT-3</i>	<u>CCTCTCTATGGGCAGTCGGTGATgAGGAC</u> AGACATGACAAGGCA (SEQ ID NO:72)
VEGFA-OT1-F	Off Target	<i>VEGFA OT-1</i>	<u>CCATCTCATCCCTGCGTGTCTCCGCAGGC</u> AAGCTGTCAAGGGT (SEQ ID NO:73)
VEGFA-OT1-R	Off Target	<i>VEGFA OT-1</i>	<u>CCTCTCTATGGGCAGTCGGTGATgCCCTC</u> ACACCCACACCCTCA (SEQ ID NO:74)
VEGFA-OT2-F	Off Target	<i>VEGFA OT-2</i>	<u>CCATCTCATCCCTGCGTGTCTCCGGAGG</u> GGTGTCACTGTTCTG (SEQ ID NO:75)
VEGFA-OT2-R	Off Target	<i>VEGFA OT-2</i>	<u>CCTCTCTATGGGCAGTCGGTGATgCAAAT</u> TGCGCCATAGCTGGG (SEQ ID NO:76)
VEGFA-OT3-F	Off Target	<i>VEGFA OT-3</i>	<u>CCATCTCATCCCTGCGTGTCTCTGAGCG</u> CTCTTCGTCTTTCC (SEQ ID NO:77)
VEGFA-OT3-R	Off Target	<i>VEGFA OT-3</i>	<u>CCTCTCTATGGGCAGTCGGTGATgGCCAG</u> GAACACAGGAATGCTA (SEQ ID NO:78)

[00121] *High-throughput Sequencing Data Analysis* Processed (demultiplexed, trimmed, and merged) sequencing reads were analyzed to determine editing outcomes using CRISPPResso2⁵ by aligning sequenced amplicons to reference and expected HDR amplicons. The quantification window was increased to 10 bp surrounding the expected cut site to better capture diverse editing outcomes, but substitutions were ignored to avoid inclusion of sequencing errors. Only reads containing no mismatches to the expected amplicon were considered for HDR quantification; reads containing indels that partially matched the expected amplicons were included in the overall reported indel frequency.

[00122] *Statistical Analysis* Unless otherwise stated, all statistical analysis and comparison were performed using t-test, with 1% false-discovery-rate (FDR) using two-stage step-up method of Benjamini, Krieger and Yekutieli (Benjamini, Y., et. al, Biometrika 93, 491–507 (2006), incorporated herein by reference). All experiments were performed in triplicates unless otherwise noted to ensure sufficient statistical power in the analysis.

[00123] *Determination of editing at predicted Cas9 off-target sites* To evaluate RecT/RecE off-target editing activity at known Cas9 off-target sites, same genomic DNA extracts for knock-in analysis were used as template for PCR amplification of top predicted off-targets sites (high scored as predicted CRISPOR, a web-based analysis tool) for the EMX1, VEGFA guides, primer sequences are listed in Table 2.

[00124] *iGUIDE Off-target Analysis* Genome-wide, unbiased off-target analysis was performed following the iGUIDE pipeline (Nobles, C.L., *et al. Genome Biol* 20, 14 (2019), incorporated herein by reference) based on Guide-seq invented previously (Tsai, S., *et al. Nat Biotechnol* 33, 187–197 (2015), incorporated herein by reference). HEK293T cells were transfected in 20uL Lonza SF Cell Line Nucleofector Solution on a Lonza Nucleofector 4-D with program DS-150 according to the manufacturer's instructions. 300ng of gRNA-Cas9 plasmids (or 150ng of each gRNACas9n plasmid for the double nickase), 150ng of the effector plasmids, and 5pmol of double stranded oligonucleotides (dsODN) were transfected. Cells were harvested after 72hrs for genomic DNA using Agencourt DNAdvance reagent kit. 400ng of purified gDNA which was then fragmented to an average of 500bp and ligated with adaptors using NEBNext Ultra II FS DNA Library Prep kit following manufacturer's instructions. Two rounds of nested anchored PCR from the oligo tag to the ligated adaptor sequence were performed to amplify targeted DNA, and the amplified library was purified, size-selected, and sequenced using Illumina Miseq V2 PE300. Sequencing data was analyzed using the published iGUIDE pipeline, with the addition of a downsampling step which ensures an unbiased comparison across samples.

EXAMPLE 1

[00125] In contrast to mammals, convenient recombineering-edit tools are available for bacteria, e.g., the phage lambda Red and RecE/T. Microbial recombineering has two major steps: template DNA is chewed back by exonucleases (Exo), then the single-strand annealing protein (SSAP) supports homology directed repair by the template, optionally facilitated by nuclease inhibitor. A system for RNA-guided targeting of RecE/T recombineering activities was developed and achieved kilobase (kb) human gene-editing without DNA cutting.

[00126] Candidate microbial systems with recombineering activities were surveyed. Two lines of reasoning guided the search: 1) Orthogonality: prioritizing proteins with minimal resemblance to mammalian repair enzymes; 2) Parsimony: focusing on systems with fewest interdependent components. Three protein families were identified: lambda Red, RecE/T, and phage T7 gp6 (Exo) and gp2.5 (SSAP) recombination machinery. Based on phylogenetic reconstruction, RecE/T proteins were determined to be the most distant from eukaryotic recombination proteins and among the most compact (FIG. 1). Thus, RecE/T systems were utilized for downstream analysis.

[00127] The NCBI protein database was systematically searched for RecE/T homologs. To develop a portable tool, evolutionary relationships and lengths were examined (FIG. 2A). Co-occurrence analysis

revealed that most RecE/T systems have only one of the two proteins (FIG. 2B). As prophage integration could be imprecise, the 11% of species harboring both homologs were prioritized as evidence for intact functionality.

[00128] The top 12 candidates were codon-optimized and MS2 coat protein (MCP) fusions were constructed to recruit these RecE/T homologs, hereafter termed “recombinator”, to wild-type *Streptococcus pyogenes* Cas9 (wtCas9) via MS2 RNA aptamers. To understand their respective molecular effects as Exo and SSAP, each was tested independently (FIG. 2C). Initial results revealed *Escherichia coli* RecE/T proteins (simplified as RecE and RecT) as promising candidates, as determined by genome knock-in assays (FIG. 2D). While RecT is only 269 amino acid (AA) long, RecE was truncated from AA587 (RecE_587) and the carboxy terminus domain (RecE_CTD) based on functional studies (Muyrers, J.P., Genes Dev. (2000); 14, 1971-1982, incorporated herein by reference).

[00129] To validate RecE/T recombineering in human cells, homology directed repair (HDR) was measured at five genomic sites with two templates. While the RecE variants (RecE_587, RecE_CTD) demonstrated variable increases in knock-in efficiency, RecT significantly enhanced HDR in all cases, replacing ~16bp sequences at *EMXI* and *VEGFA*, and knocking-in ~1kb cassette at *HSP90AA1*, *DYNLT1*, *AAVSI* (FIGS. 3A-E, FIG. 4). These results were verified using imaging (FIG. 3F) and junction sites were sequenced using Sanger sequencing to confirm precise insertion (FIG. 3G). To test if these activities are truly sequence-specific, a no-recruitment control with the PP7 coat protein (PCP) that recognizes PP7 aptamers not MS2 aptamers was employed. RecE had activities without recruitment, whereas RecT showed efficiency increases in a recruitment-dependent manner (FIG. 3H). Without being bound by theory, this may be explained by RecE exonuclease activity acting promiscuously (FIG. 2C). The RecE/T recombineering-edit (REDIT) tools was termed as REDITv1, with REDITv1_RecT as the preferred variant.

EXAMPLE 2

[00130] Three tests on REDITv1 were performed to explore: 1) activity across cell types, 2) optimal designs of HDR template, and 3) specificity. REDITv1 activity was robust across multiple genomic sites in HEK, A549, HepG2, and HeLa cells (FIGS. 5A-C, FIGS. 6A-C). Noticeably, in human embryonic stem cells (hESCs), REDITv1 exhibited consistent increases of kilobase knock-in efficiency at *HSP90AA1* and *OCT4*, with up to 3.5-fold improvement relative to Cas9-HDR (FIGS. 5D-E, FIGS. 6D-E). Different template designs were also tested. REDITv1 performed efficient kilobase editing using HA

length as short as 200bp total, with longer HA supporting higher efficiency. It achieved up to 10% efficiency (without selection) for kb-scale knock-in, a 5-fold increase over Cas9-HDR and significantly higher than the 1~2% typical efficiency (FIG. 7). Lastly, the accuracy of REDITv1 accuracy was determined using deep sequencing of predicted off-target sites (OTs) and GUIDE-seq. Although REDITv1 did not increase off-target effects, detectable OTs remained at previously reported sites for *EMXI* and *VEGFA* (FIGS. 5F-G, FIG. 8). In short, REDITv1 showcased kilobase-scale genome recombineering but retained the off-target issues, with REDITv1_RecT having the highest efficiency.

EXAMPLE 3

[00131] To alleviate unwanted edits, a version of REDIT with non-cutting Cas9 nickases (Cas9n) was assessed. A similar strategy was previously employed (Ran, F.A., et al., Cell (2013), 154: 1380-1389, incorporated herein by reference) to address off-target issues but had low HDR efficiency. REDIT was tested to determine if this system could overcome the limitation of endogenous repair and promote nicking-mediated recombination. Indeed, the nickase version demonstrated higher efficiencies, with the best results from Cas9n(D10A) with single- and double-nicking. This Cas9n(D10A) variant was designated REDITv2N (FIG. 9A). A 5%~10% knock-in without selection was observed using REDITv2N double-nicking, comparable to REDITv1 using wtCas9 (FIG. 9A, FIG. 10A). Junction sequencing confirmed the precision of knock-in for all targets (FIG. 11). This result represented 6- to 10-fold improvement over Cas9n-HDR. Even with single-nicking REDITv2N, a ~2% efficiency for 1kb knock-in was observed, a level considerably higher than the 0.46% HDR efficiency in previous report (Cong, L. et al., Science. 339, 819-823, incorporated herein by reference) using regular single-nicking Cas9n and a less-challenging 12-bp knock-in template (FIG. 9A).

[00132] The off-target activity of REDITv2N was investigated using GUIDE-seq. Results showed minimal off-target cleavage and a reduction of OTs by ~90% compared to REDITv1 (FIG. 9B). Specifically, for *DYNLT1*-targeting guides, the most abundant *KIF6* OTS was significantly enriched in REDITv1 group but disappeared when using REDITv2N (FIG. 9C). REDITv2N was highly accurate (FIGS. 9B-C, FIG. 12).

[00133] Another byproduct of HDR editing is on-target insertion-deletions (indels). They could drastically lower yields of gene-editing, especially for long sequences. Indel formation was measured in an *EMXI* knock-in experiment using deep sequencing. REDITv2N increased HDR to the same efficiency

as its counterpart using wtCas9 (FIG. 12C, top), with a reduction of unwanted on-target indels by 92% (FIG. 12C, bottom).

[00134] Concepts from GUIDE-seq, LAM-PCR, and TLA were used to develop an NGS-based assay to identify genome-wide insertion sites (GIS), or GIS-seq (FIG. 30A). Using GIS-seq, NGS read clusters/peaks representing knock-in insertion sites were obtained (FIG. 30B), showing representative reads from the on-target site). GIS-seq was applied to *DYNLT1* and *ACTB* loci to measure the knock-in accuracy. Sequencing results indicated that, when considering sites with high confidence based on maximum likelihood estimation, REDIT had less off-target insertion sites identified compared with Cas9 (FIG. 30C). Together, the clonal Sanger sequencing of knock-in junctions (FIGS. 9C and 12), GUIDE-seq analysis (FIG. 9B), and GIS seq results (FIGS. 30A-30C) indicated that REDIT can be an efficient method with the ability to insert kilobase-length sequences with less unwanted editing events.

EXAMPLE 4

[00135] REDIT was examined for long sequence editing ability in the absence of any nicking/cutting of the target DNA. Remarkably, when using catalytically dead Cas9 (dCas9) to construct REDITv2D, an exact genomic knock-in of a kilobase cassette was observed in human cells (FIG. 9D, top, FIG. 13). While REDITv2D has lower efficiency than REDITv2N, it achieved programmable DNA-damage-free editing at kilobase-scale with 1~2% efficiency and no selection (FIG. 9D, FIG. 10B). It was hypothesized that two processes could be contributing to the REDITv2D recombineering. One possibility was via dCas9 unwinding. If dCas9 could unwind DNA as it induces sequence-specific formation of loop, a double-binding with two dCas9s would be expected to promote genome accessibility to RecE/T. However, a significant increase upon delivering two guide RNAs was not observed (FIG. 9D, bottom). Another possibility was that the unwinding of DNA during cell cycle permitted RecE/T to access the target region mediated by dCas9 binding. A 1kb knock-in was performed with different REDIT tools at varying serum levels (10% regular, 2% reduced, and no serum). As serum starvation arrests cell proliferation, the results indicated that the cell cycle correlated positively with REDITv2D recombineering (FIG. 9E). Upon no-serum treatment, HDR efficiency only dropped in REDITv2D(dCas9) group, whereas REDITv1(wtCas9) and REDITv2N(D10A) were not affected (FIG. 9E, FIG. 14), supporting that DNA unwinding permitted RecE/T to access the target region.

EXAMPLE 5

[00136] Microscopy analysis revealed incomplete nuclei-targeting of REDITv1, particularly REDITv1_RecT (FIG. 15). Hence, different designs of protein linkers and nuclear localization signals (NLSs) were tested (FIG. 15A). The extended XTEN-linker with C-terminal SV40-NLS was identified as a preferred configuration, termed REDITv3 (FIG. 16). REDITv3 further achieved a 2- to 3- fold increase of HDR efficiencies over REDITv2 across genome targets and Cas9 variants (wtCas9, Cas9n, dCas9) (FIG. 17).

[00137] Finally, REDITv3 was utilized in hESCs to engineer kilobase knock-in alleles in human stem cells. REDITv3N single- and double-nicking designs resulted in 5-fold and 20-fold increased HDR efficiencies over no-recombinator controls, respectively (FIG. 9F). The efficacy and fidelity were confirmed via a combination of assays described for previous REDIT versions (FIGS. 9F-G, FIG. 18). Additionally, REDITv3 works effectively with *Staphylococcus aureus* Cas9 (SaCas9), a compact CRISPR system suitable for *in vivo* delivery (FIG. 19).

EXAMPLE 6

[00138] To further investigate RecT and RecE_587 variants, both RecT and RecE_587 were truncated at various lengths as shown in FIG. 20A and FIG. 21A, respectively. The resulting efficiencies were measured using an mKate knock-in assay, with both wildtype SpCas9 and Cas9n(D10A) with single- and double-nicking at the *DYNLT1* locus (FIGS. 20B-C and FIGS. 21B-C, respectively). Efficiencies of the no recombination group are shown as the control.

[00139] The truncated versions of both RecT and RecE_587 retained significant recombineering activity when used with different Cas9s. In particular, compared with the full-length RecT(1-269aa), the new truncated versions such as RecT(93-264aa) are over 30% smaller yet they preserved essentially the full activities of RecT in stimulating recombination in eukaryotic cells. Similarly, compared with the full-length RecE(1-280aa), truncated versions such as RecE_587(120-221aa) and RecE_587(120-209aa) are over 60% smaller but still retained high recombination activities in human cells. These truncated versions demonstrated the potential to further engineer minimal-functional recombineering enzymes using RecE and RecT protein variants, but also provide valuable compact recombineering tools for human genome editing that is ideal for *in vitro*, *ex vivo*, and *in vivo* delivery given their small size.

[00140] Overall, REDIT harnessed the specificity of CRISPR genome-targeting with the efficiency of RecE/RecT recombineering. The disclosed high-efficiency, low-error system makes a powerful addition

to existing CRISPR toolkits. The balanced efficiency and accuracy of REDITv3N makes it an attractive therapeutic option for knock-in of large cassette in immune and stem cells.

EXAMPLE 7

[00141] The reconstructed RecE and RecT phylogenetic trees with eukaryotic recombination enzymes from yeast and human (FIGS. 1A and 1B) show the evolutionary distance of the proteins based on sequence homology. The dotted boxes indicate the full-length *E. coli* RecB and *E. coli* RecE protein. The catalytic core domain of *E. coli* RecB and *E. coli* RecE protein (solid boxes) was used for the comparison. The gene-editing activities of these families of recombineering proteins were measured using the MS2-MCP recruitment system, where sgRNA bearing MS2 stem-loop is used with recombineering proteins fused to the MCP protein via peptide linker and with nuclear-localization signals.

[00142] Three exonuclease proteins were used: the exonuclease from phage Lambda, the RecE587 core domain of *E. coli* RecE protein, and the exonuclease (gene name gp6) from phage T7 (FIG. 22A). The gene-editing activity was measured using mKate knock-in assay at genomic loci (DYNLT1 and HSP90AA1).

[00143] Similar measurements were made testing the genome editing efficiencies of three single-strand DNA annealing proteins (SSAPs) from the same three species of microbes as the exonucleases, namely Bet protein from phage Lambda, RecT protein from *E. coli*, and SSAP (gene name gp2.5) from phage T7 (FIG. 22B).

[00144] From these results, the genome recombineering activities of all three major family of phage/microbial recombination systems was systematically measured and validated in eukaryotic cells (lambda phage exonuclease and beta proteins; *E. coli* prophase RecE and RecT proteins, T7 phage exonuclease gp6 and single-strand binding gp2.5 proteins). All six proteins from three systems achieved efficient gene editing to knock-in kilobase-long sequences into mammalian genome across two genomic loci. Overall, the exonucleases showed ~3-fold higher recombination efficiency (up to 4% mKate genome knock-in) when compared with no-recombinator controls. The single-strand annealing proteins (SSAP) showed higher activities, with 4-fold to 8-fold higher gene-editing activities over the control groups. This demonstrated the general applicability and validity that microbial recombination proteins in the exonuclease and SSAP families could be engineered via the Cas9-based fusion protein system to achieve highly efficient genome recombination in mammalian cells.

EXAMPLE 8

[00145] In order to demonstrate the generalizability of REDIT protein design, alternative recruitment systems were developed and tested. For a more compact REDIT system, the REDIT recombinator proteins were fused to N22 peptide and at the same time the sgRNA included boxB, the short cognizant sequence of N22 peptide, replacing MCP within the sgRNA (FIG. 23A). This boxB-N22 system demonstrated comparable editing efficiencies at the two genomic sites tested as shown in FIGS. 23B-23E with side-by-side comparisons of the MS2-MCP recruitment system.

[00146] A REDIT system using SunTag recruitment, a protein-based recruitment system, was developed (FIGS. 24A and 27A). Because SunTag is based on fusion protein design, the sgRNA or guideRNAs are the same as wild-type CRISPR system. Specifically, the REDIT recombinator proteins were fused to scFV antibody peptide (replacing MCP), and the GCN4 peptide was fused in tandem fashion (10 copies of GCN4 peptide separated by linkers) to the Cas9 protein. Thus, the scFV-REDIT could be recruited to the Cas9 complex via affinity of GCN4 to scFV.

[00147] mKate knock-in experiments (FIG. 24B and 27B) were used to measure the editing efficiencies at the *DYNLT1* locus and the HSP90AA1 locus, respectively. This SunTag-based REDIT system demonstrated significant increase of gene-editing knock-in efficiency at the *DYNLT1* genomic sites tested. In addition, the SunTag design significantly increased HRD efficiencies to ~2-fold better than Cas9 but did not achieve increases as high as the MS2-aptamer.

EXAMPLE 9

[00148] In order to demonstrate the generalizability of REDIT protein design and develop versatile REDIT system applicable to a range of CRISPR enzymes, Cpf1/Cas12a based REDIT system using the SunTag recruitment design was developed (FIG. 25A). Two different Cpf1/Cas12a proteins were tested (*Lachnospiraceae bacterium ND2006*, LbCpf1 and *Acidaminococcus sp. BV3L6*) using the mKate knock-in assay as previously shown (FIG. 25B).

[00149] These results showed that the microbial recombination proteins (exonuclease and single-strand annealing proteins) could be engineered using alternative designs such as the SunTag recruitment system to perform genome editing in eukaryotic cells. These protein-based recruitment system does not require the usage of RNA aptamers or RNA-binding proteins, instead, they took advantage of fusion protein domains directly connecting to the CRISPR enzymes to recruit REDIT proteins.

[00150] In addition to the flexibility in recruitment system design, these results using Cpf1/Cas12a-type CRISPR enzymes also demonstrated the general adaptability of REDIT proteins to various CRISPR systems for genome recombination. Cpf1/Cas12a enzymes have different catalytic residues and DNA-recognition mechanisms from the Cas9 enzymes. Hence, the REDIT recombination proteins (exonucleases and single-strand annealing proteins) could function independent from the specific choices of the CRISPR enzyme components (Cas9, Cpf1/Cas12a, and others). This proved the generalizability of the REDIT system and open up possibility to use additional CRISPR enzymes (known and unknown) as components of REDIT system to achieve accurate genome editing in eukaryotic cells.

EXAMPLE 10

[00151] Fifteen different species of microbes having RecE/RecT proteins were selected for a screen of various RecE and RecT proteins across the microbial kingdom (Table 3). Each protein was codon-optimized and synthesized. As previously described for *E. coli* RecE/RecT based REDIT systems, each protein was fused via E-XTEN linker to the MCP protein with additional nuclear localization signal. mKate knock-in gene-editing assay was used to measure efficiencies at *DYNLT1* locus (FIG. 26A, Table 4) and *HSP90AA1* locus (FIG. 26B, Table 4). The homologs demonstrated the ability to enable and enhance precision gene-editing.

Table 3: RecE and RecT protein homologs

Homolog	Source	Protein
T1	<i>Pantoea stewartii</i>	RecT
E1	<i>Pantoea stewartii</i>	RecE
T2	<i>Pantoea brenneri</i>	RecT
E2	<i>Pantoea brenneri</i>	RecE
T3	<i>Pantoea dispersa</i>	RecT
E3	<i>Pantoea dispersa</i>	RecE
T4	Type-F symbiont of <i>Plautia stali</i>	RecT
E4	Type-F symbiont of <i>Plautia stali</i>	RecE
T5	<i>Providencia stuartii</i>	RecT
E5	<i>Providencia stuartii</i>	RecE
T6	<i>Providencia</i> sp. MGF014	RecT
E6	<i>Providencia</i> sp. MGF014	RecE
T7	<i>Providencia alcalifaciens</i> DSM 30120	RecT
E7	<i>Providencia alcalifaciens</i> DSM 30120	RecE
T8	<i>Shewanella putrefaciens</i>	RecT
E8	<i>Shewanella putrefaciens</i>	RecE

T9	Bacillus sp. MUM 116	RecT
E9	Bacillus sp. MUM 116	RecE
T10	Shigella sonnei	RecT
E10	Shigella sonnei	RecE
T11	Salmonella enterica	RecT
E11	Salmonella enterica	RecE
T12	Acetobacter	RecT
E12	Acetobacter	RecE
T13	Salmonella enterica subsp. enterica serovar Javiana str. 10721	RecT
E13	Salmonella enterica subsp. enterica serovar Javiana str. 10721	RecE
T14	Pseudobacteriovorax antillogorgiicola	RecT
E14	Pseudobacteriovorax antillogorgiicola	RecE
T15	Photobacterium sp. JCM 19050	RecT
E15	Photobacterium sp. JCM 19050	RecE

Table 4: mKate Knock-In Gene-Editing Efficiencies

	DYNLT1		HSP90AA1	
	Mean mKate+ (%)	SEM	Mean mKate+ (%)	SEM
NC	1.2100	0.0802	1.7333	0.1245
NR	2.0500	0.1442	4.0100	0.2166
EcRecE 587	5.1767	0.0897	3.7067	0.1784
EcRecT	9.9467	1.0143	6.5467	0.4646
Homolog_T1	11.7333	0.4667	8.0733	0.8752
Homolog_E1	5.7333	0.8503	7.6567	0.4556
Homolog_T2	12.0000	0.5292	6.9233	0.4594
Homolog_E2	7.4533	0.8553	6.4867	0.4359
Homolog_T3	11.9000	1.3013	7.1200	0.2730
Homolog_E3	2.0533	0.1020	6.7467	0.1565
Homolog_T4	10.4433	0.7331	5.7567	0.8704
Homolog_E4	5.7200	0.4744	6.2567	0.3339
Homolog_T5	10.8267	0.9445	6.4300	0.3262
Homolog_E5	4.4667	0.7116	6.0233	0.4366
Homolog_T6	9.0533	0.3548	6.2500	0.4100
Homolog_E6	5.4100	0.5981	5.9300	0.4708
Homolog_T7	5.6467	0.7383	5.3700	0.4795
Homolog_E7	4.4733	0.2444	5.7367	0.2105
Homolog_T8	5.0400	0.5599	5.7133	0.4886

Homolog E8	4.6567	0.3088	7.0533	0.4388
Homolog T9	8.1300	0.3523	6.2000	0.2511
Homolog E9	5.3233	0.5233	5.6900	0.4903
Homolog T10	8.5333	0.1601	5.5900	0.2237
Homolog E10	4.4000	1.0149	3.5900	0.1442
Homolog T11	9.8467	1.4374	4.9233	0.4074
Homolog E11	7.0567	1.5872	3.1167	0.2010
Homolog T12	8.5900	0.5401	5.2733	0.2935
Homolog E12	5.2633	0.3374	6.0800	0.5164
Homolog T13	9.9567	0.3324	5.7200	0.4267
Homolog E13	5.6333	0.2360	5.6900	0.3729
Homolog T14	6.7700	0.7022	4.7200	0.3612
Homolog E14	6.0167	0.4890	5.7100	0.1793
Homolog T15	7.8033	0.7075	5.2333	0.2302
Homolog E15	5.0700	0.5543	6.0500	0.5696

EXAMPLE 11

[00152] Next, to benchmark the RecT-based REDIT design, it was compared with three categories of existing HDR-enhancing tools (FIGS. 28A and 28B): DNA repair enzyme CtIP fusion with the Cas9 (Cas9-HE), a fusion of the functional domain (amino acids 1 to 110) of human Geminin protein with the Cas9 (Cas9-Gem), and a small-molecule enhancers of HDR via cell cycle control, Nocodazole. Across endogenous targets tested, the RecT-based REDIT design had favorable performance compared with three alternative strategies (FIG. 28C). Furthermore, the RecT-based REDIT design, which putatively acted through activity independently from the other approaches, may synergize with existing methods. To test this hypothesis, RecT-based REDIT design was combined with three different approaches (conveniently through the MS2-aptamer) (FIG. 28A, right). The RecT-based REDIT design could indeed further enhance the HDR-promoting activities of the tested tools (FIG. 28C).

EXAMPLE 12

[00153] The effect of template HA lengths on the editing efficiency of REDIT was quantified when using the canonical HDR donor bearing HAs of at least 100 bp on each side (FIG. 29A, left). Higher HDR rates were observed for both Cas9 and RecT groups with increasing HA lengths, and REDIT effectively stimulated HDR over Cas9 using HA lengths as short as ~100bp each side. When supplied with a longer

template bearing 600–800 bp total HA, RecT achieved over 10% HDR efficiencies for kb-scale knock-in without selection, significantly higher than the 2–3% efficiency when only using Cas9. Recent reports identified that using donor DNAs with shorter HAs (usually between 10 and 50 bp) could significantly stimulate knock-in efficiencies thanks to the high repair activities from the Microhomology-mediated end joining (MMEJ) pathway. Knock-in efficiencies of the REDIT-based method were compared with Cas9, using donor DNA with 0bp (NHEJ-based), 10bp or 50bp (MMEJ-based) HAs. The results demonstrated that short-HA donors leveraging MMEJ mechanisms yielded higher editing efficiencies compared with HDR donors (FIG. 29A, right). At the same time, REDIT was able to enhance the knock-in efficiencies as long as there is HA present (no effect for the 0bp NHEJ donor). This effect is particularly significant with The 10 bp donors in which there was a significant effect, were chosen for further characterization and comparison with the HDR donors.

[00154] The knock-in cells were clonally isolated and the target genomic region was amplified using primers binding completely outside of the donor DNAs for colony Sanger sequencing (FIG. 29B). Junction sequencing analysis (~48 colonies per gene per condition) revealed varying degrees of indels at the 5' and 3' knock-in junctions, including at single or both junctions (FIG. 29C). Overall, HDR donors had better precision than MMEJ donors, and REDIT modestly improved the knock-in yield compared with Cas9, though junction indels were still observed.

[00155] Furthermore, the efficiencies of REDIT and Cas9 were compared when making different lengths of editing. For longer edits, 2-kb knock-in cassettes were used (FIG. 29D), and for shorter edits single-stranded oligo donors (ssODN) were used. When the knock-in sequence length was increased to ~2-kb using a dual-mKate/GFP template, REDIT maintained its HDR-promoting activity compared with Cas9 across endogenous targets tested (FIG. 29D). For ssODN tests, at two well-established loci *EMXI* and *VEGFA*, REDIT and Cas9 were used to introduce 12–16-bp exogenous sequences. As ssODN templates are short (<100 bp HAs on each side), next-generation sequencing (NGS) was used to quantify the editing events. Comparable levels of indels were observed between Cas9 and REDIT with improved HDR efficiencies using REDIT.

EXAMPLE 13

[00156] The sensitivity of REDIT's ability to promote HDR in the presence or absence of two distinctive pharmacological inhibitors of RAD51, B02 and RI-1 (FIG. 31A). As expected, for Cas9-based editing, RAD51 inhibition significantly lowered HDR efficiencies (FIGS. 31B, 31C, and 32A).

Intriguingly, RAD51 inhibition decreased REDIT and REDITdn efficiencies only moderately, as both REDIT/REDITdn methods maintained significantly higher knock-in efficiencies compared with Cas9/Cas9dn under RAD51 inhibition.

[00157] Mirin, a potent chemical inhibitor of DSB repair, which has also been shown to prevent MRN complex formation, MRN-dependent ATM activation, and inhibit Mre11 exonuclease activity was also used. When treating cells with Mirin, only the editing efficiencies of Cas9 reference experiments were affected by the Mirin treatment, whereas the REDIT versions were essentially the same as vehicle-treated groups across all genomic targets (FIG. 32A).

[00158] To test if cell cycle inhibition affected recombination, cells were chemically synchronized at the G1/S boundary using double Thymidine blockage (DTB). REDIT versions had reduced editing efficiencies under DTB treatment, though it maintained higher editing efficiencies under DNA repair pathway inhibition, compared with Cas9 reference experiments, when Mirin RI-1, or B02 were combined with DTB treatment (FIG. 32B).

[00159] To validate REDIT in different contexts, REDIT was applied in human embryonic stem cells (hESCs) to test their ability to engineer long sequences in non-transformed human cells. Robust stimulation of HDR was observed across all three genomic sites (*HSP90AA1*, *ACTB*, *OCT4/POU5F1*) using REDIT and REDITdn (FIGS. 31D and 31E). Of note, REDIT and REDITdn editing used donor DNAs with 200-bp HAs on each side and achieved up to over 5% efficiency for kb-scale gene-editing without selection compared with ~1% efficiency using non-REDIT methods. Additionally, REDIT improved knock-in efficiencies in A549 (lung-derived), HepG2 (liver-derived), and HeLa (cervix-derived) cells, demonstrating up to ~15% kb-scale genomic knock-in without selection. This improvement was up to 4-fold higher than the Cas9 groups, supporting the potential of using REDIT methods in different cell types.

EXAMPLE 14

[00160] *In vivo* use of dCas9-EcRecT (SAFE-dCas9) was tested using cleavage free dCas9 editor via hydrodynamic tail vein injection. The gene editing vectors and template DNA used are shown in FIG. 33A. A gene editing vector (60 µg) and template DNA (60 µg) were injected via hydrodynamic tail vein injection to deliver the components to the mouse. Successful gene editing of liver hepatocytes was monitored by transgene-encoded protein expression from the albumin locus. A schematic of the experimental procedure is shown in FIG. 33B.

[00161] At approximately seven days after injection, the perfused mice livers were dissected. The lobes of the liver were homogenized and processed to extract liver genomic DNA from the primary hepatocytes. The extracted genomic DNA was used for three different downstream analyses: 1) PCR using knock-in-specific primers and agarose gel electrophoresis (FIG. 34A); 2) Sanger sequencing of the knock-in PCR product (FIG. 34B); 3) high-throughput deep sequencing of the knock-in junction to confirm and quantify the accuracy of gene-editing using SAFE-dCas9 *in vivo* (FIG. 34C). Each downstream analysis confirmed knock-in success with .

[00162] In addition, *in vivo* use was tested using adeno-associated virus (AAV) delivery into LTC mice lungs. LTC mice include three genome alleles: 1) Lkb1 (flox/flox) allele allows Lkb1-KO when expressing Cre; 2) R26(LSL-TdTom) allele allows detection of AAV-transduced cells via TdTom red fluorescent protein; and 3) H11(LSL-Cas9) allele allows expression of Cas9 in AAV-transduced cells. Schematics of the REDI gene editing vector and Cas9 control vectors are shown in FIG. 35A. As shown in FIG. 35B, successful gene editing using the gene editing vector leads to Kras alleles that drive tumor growth in the lung of the treated mice.

[00163] Approximately fourteen weeks after the AAV injection, perfused mice lungs were dissected. Fixed lung tissue was used for imaging analysis to identify tumor formation from successful gene-editing (FIG. 35C). Quantification of the surface tumor number via imaging analysis showed increased gene-editing efficiencies and total number of tumors in the REDIT treated mice (FIG. 35C).

Escherichia coli RecE amino acid sequence (SEQ ID NO:1):

MSTKPLFLLRKAKKSSGEPDVVLWASNDFESTCATLDYLIVKSGKKLSSYFKAVATNFPVVNDL
PAEGEIDFTWSERYQLSKDSMTWELKPGAAPDNAHYQGNTNVNGEDMTEIEENMLLPISGQELP
IRWLAQHGSEKPVTHVSRDGLQALHIARAEELPAVTALAVSHKTSLLDPLEIRELHKLVRDTRDKV
FPNPGNSNLGLITAFFEAYLNADYTDRGLLTKIEWMKGNRVSHITRTASGANAGGGNLTDREGGF
VHDLTSLARDVATGVLARSMDLDIYNLHPAHAKRIEIIAENKPPFSVFRDKFITMPGGLDYSGAI
VVASVKEAPIGIEVIPAHVTEYLNKVL TETDHANPDPEIVDIACGRSSAPMPQRVTEEGKQDDEEK
PQPSGTTAVEQGEAETMEPDATEHHQDTQPLDAQSQVNSVDAKYQELRAELHEARKNIPSKNPV
DDDKLLAASRGEFVDGISDPNDPKWVGIGQTRDCVYQNPETEKTSPPDMNQPEPVVQGEPEIAC
NACGQTGGDNCPDCGAVMGDATYQETFEESQVEAKENDPEEMEGAEHPHNENAGSDPHRDC
SDETGEVADPVIVEDIPIGIYYGISNENYHAGPGISKSQLDIADTPALYLWRKNAPVDTTKTKTL
DLGTAHFHCRVLEPEEFSNRFIVAPEFNRRRTNAGKEEEKAFLECASTGKTVITAEGRKIELMYQS
VMALPLGQWLVESAGHAESSIYWEDPETGILCRCPDKIPEFWIMDVKTTADIQRFKTAYYDY
RYHVQDAFYSDGYEAQFGVQPTFVFLVASTTIECGRYPVEIFMMGEEAKLAGQQEYHRNLRTLA
DCLNTDEWPA IKTLSPRWAKEYAND

Escherichia coli RecE_587 amino acid sequence (SEQ ID NO:2):

ADPVIVEDIPIGIYYGISNENYHAGPGVSKSQLDDIADTPALYLWRKNAPVDTTKTKTLD

LGTAHFHCRVLEPEEFSNRFIVAPEFNRRNTNSGKEEEKAFLECASTGKTVITAEGRKIEL
MYQSVMALPLGQWLVESAGHAESSIYWEDPETAILECRCPDKIPEFWIMDVKTTADI
QRFKTAYYDYRYHVQDAFYSDGYEAQFGVQPTFVFLVASTTIECGRYPVEIFMMGEEA
KLAGQLEYHRNLRTLADCLNTDEWPAIKTLSPRWAKEYAND*

Escherichia coli CTD_RecE amino acid sequence (SEQ ID NO:3):

GISNENYHAGPGVSKSQLDDIADTPALYLWRKNAPVDTTKTKTLDLGTAFHCRVLEPEE
FSNRFIVAPEFNRRNTNSGKEEEKAFLECASTGKTVITAEGRKIELMYQSVMALPLGQW
LVESAGHAESSIYWEDPETAILECRCPDKIPEFWIMDVKTTADIQRFKTAYYDYRYHV
QDAFYSDGYEAQFGVQPTFVFLVASTTIECGRYPVEIFMMGEEAKLAGQLEYHRNLRTL
ADCLNTDEWPAIKTLSPRWAKEYAND*

Pantoea brenneri RecE amino acid sequence (SEQ ID NO:4):

MQPGIYYDISNEDYHRGAGISKSQDDIAISPAIYQWRKHAPVDEEKTAALDLGTALHCL
LLEPDEFKSRFQIGPEVNRRTTAGKEKEKEFIERCEAEGITPITHDDNRKLLMRDSALAH
PIARWMLEAQGNAEASIWNRDAGVLSRCRPDKIITEFNWCVDVKSTADIMKFQKDF
YSYRYHVQDAFYSDGYESHFETPTFAFLAVSTSIDCGRYPVQVFIMDQQAADAGRAE
YKRNIHTFAECLSRNEWPGIATLSLPFWAKELRNE

Type-F symbiont of Plautia stali RecE amino acid sequence (SEQ ID NO:5):

MQPGIYYDISNEDYHGGPGISKSQDDIAISPAIYQWRKHAPVDEEKTAALDLGTALHCL
LLEPDEFKSRFEIGPEVNRRTTAGKEKEKEFMERCEAEGVTPITHDDNRKLLMRDSAM
AHPIARWMLEAQGNAEASIWNRDGTGVLSCRPDKIITDFNWCVDVKSTADIIFQKD
FYSYRYHVQDAFYSDGYESHFDEPTFAFLAVSTSIDCGRYPVQVFIMDQQAADAGRAE
YKRNIHTFAECLSRNEWPGIATLSLPYWAKELRNE

Providencia sp. MGF014 RecE amino acid sequence (SEQ ID NO:6):

MKEGIYYNISNEDYHNGLGISKSQDLINEMPAEYIWSKEAPVDEEKIKPLEIGTALHCLL
LEPDEYHKRYKIGPDVNRRTNVGKEKEKEFFDMCEKEGITPITHDDNRKLLMIMRDSALA
HPIAKWCLEADGVSESIYWTDKETDVLRCRCPDRIITAHNYIIDVKSSGDIEKFDYEEYN
YRYHVQDAFYSDGYKEVTGITPTFLFLVSTKIDCGKYPVRTYVMSEEAASAGRTAYK
HNLLTYAECLKTDEWAGIRTLSPRWAKELRNE

Shigella sonnei RecE amino acid sequence (SEQ ID NO:7):

DRGLLTKEWRKGNRVSRIIRTAGANAGGGNLTDRGEGFVHDLTSLARDIATGVLARS
MDVDIYNLHPAHAKRIEIIAENKPPFSVFRDKFITMPGGLDYSRIVVASVKEAPIGIEVI
PAHVTAAYLNKVLTTETDHANPDPEIVDIACGRSSAPMPQRVTEEGKQDDEEKLQPSGTTA
DEQGEAETMEPDATKHHQDTQPLDAQSQVNSVDAKYQELRAELHEARKNIPSKNPVDA
DKLLAASRGFVVDGISDPNDPKWVKGIQTRDSVYQNPETEKTSPDMKQPEPVVQQEPE
IAFNACGQTGGDNCPDCGAVMGDATYQETFDENQVEAKENDPEEMEGAEPHNENA
GSDPHRDCSDETGEVADPVIVEDIPIGIYYGISNENYHAGPGVSKSQLDDIADTPALYLW
RKNAPVDTTKTKTLDLGTAFHCRVLEPEEFSNRFIVAPEFNRRNTNAGKEEEKAFLECA
STGKMVITAEGRKIELMYQSVMALPLGQWLVESAGHAESSIYWEDPETGILRCRCPDK
IPEFWIMDVKTTADIQRFKTAYYDYRYHVQDAFYSDGYEAQFGVQPTFVFLVASTTIE
CGRYPVEIFMMGEEAKLAGQLEYHRNLRTLADCLNTDEWPAIKTLSPRWAKEYAND

Pseudobacteriovorax antillogorgiicola RecE amino acid sequence (SEQ ID NO:8):

MSKLSNLKVSNSDVDTLRIRMKEGVYRDLPYESYHQSPGYSKTSLCQIDKAPIYLKTKV
 PQKSTKSLNIGTAFHEAMEGVFKDKYVVHPDPGVNKTTSWKDFVKRYPKHMPLKRSE
 YDQVLAMYDAARSYRPFQKYHLSRGFYESSFYWHDAVTNSLIKCRPDYITPDGMSVIDF
 KTTVDPSPKGFQYQAYKYHYVSAALTLEGIEAVTGIRPKEYLFLAVSNSAPYLTALYR
 ASEKEIALGDHFIRSLTLKTCLESGKWPGQEILELGLPFGSLKELREEQEVEDEFME
 LVG

Escherichia coli RecT amino acid sequence (SEQ ID NO:9):

MTKQPPIAKADLQKTQGNRAPAAVKNSDVISFINQPSMKEQLAAALPRHMTAERMIRIA
 TTEIRKVPALGNCDTMSFVSAIVQCSQLGLEPGSALGHAYLLPFGNKNEKSGKKNVQLII
 GYRGMIDLARRSGQIASLSARVVREGDEFSEFGLDEKLIHRPGENEDAPVTHVYAVAR
 LKDGGTQFEVMTRKQIELVRSLSKAGNNGPWVTHWEEMAKKTAIRRLFKYLPVSIEIQR
 AVSMDEKEPLTIDPADSSVLTGEYSVIDNSEE*

Pantoea brenneri RecT amino acid sequence (SEQ ID NO:10):

MSNQPPIASADLQKTQQSKQVANKTPEQTLVGFMNQPMKSQLAAALPRHMTADRMIRI
 RIVTTEIRKTPQLAQCDQSSFIGAVVQCSQLGLEPGSALGHAYLLPFGNGRSKSGQSNVQ
 LIIGYRGMIDLARRSGQIVSLSARVVRADEFSEFYGLDENLVHRPGENEDAPITHVYAV
 ARLKDGGTQFEVMTVKQVEKVKAQSKASSNGPWVTHWEEMAKKTVIRRLFKYLPVSI
 EMQKAVVLDEKAESDQDQNASVLSAEYSVLESGDEATN

Type-F symbiont of Plautia stali RecT amino acid sequence (SEQ ID NO:11):

MSNQPPIASADLQKTQQSKQVANKTPEQTLVGFMNQPMKSQLAAALPRHMTADRMIRI
 RIVTTEIRKTPALATCDQSSFIGAVVQCSQLGLEPGSALGHAYLLPFGNGRSKSGQSNVQ
 LIIGYRGMIDLARRSGQIVSLSARVVRADEFSEFYGLDENLVHRPGDNEDAPITHVYAV
 ARLKDGGTQFEVMATAKQVEKVKAQSKASSNGPWVTHWEEMAKKTVIRRLFKYLPVSI
 EMQKAVVLDEKAESDQDQNASVLSAEYSVLEGDGGE

Providencia sp. MGF014 RecT amino acid sequence (SEQ ID NO:12):

MSNPPLAQSDLQKTQGTEVKVKTQDQQLIQFINQPSMKAQLAAALPRHMTTPDRMIRIVT
 TEIRKTPALATCDMQSFVGAVVQCSQLGLEPGNALGHAYLLPFGNGKAKSGQSNVQLII
 GYRGMIDLARRSNQIISARTVRQGDNFHFYGLNEDLTHTPSENEDESPITHVYAVARL
 KDGGVQFEVMTYNQVEKVRASSKAGQNGPWVSHWEEMAKKTVIRRLFKYLPVSIEMQ
 KAVVLDEKAEANVDQENATIFEGEYEEVGTGDN

Shigella sonnei RecT amino acid sequence (SEQ ID NO:13):

MTKQPPIAKADLQKTQENRAPAAIKNNDVISFINQPSMKEQLAAALPRHMTAERMIRIA
 TTEIRKVPALGNCDTMSFVSAIVQCSQLGLEPGSALGHAYLLPFGNKNEKSGKKNVQLII
 GYRGMIDLARRSGQIASLSARVVREGDEFNFEFGLDEKLIHRPGENEDAPVTHVYAVAR
 LKDGGTQFEVMTRRQIELVRSQSKAGNNGPWVTHWEEMAKKTAIRRLFKYLPVSIEIQR
 AVSMDEKEPLTIDPADSSVLTGEYSVIDNSEE

Pseudobacteriovorax antillogorgiicola RecT amino acid sequence (SEQ ID NO:14):

MGHLVSKTEQDYIKQHYAKGATDQEFHFVCRARGLNPAANQIYFVKYRSKDGPAK
 PAFILSIDSLRLIAHRTGDYAGCSEPIFTDGGKACTVTVRRNLKSGETGNFSGMAFYDEQ
 VQKNGRPTSFWQSKPRTMLEKCAEAKALRKAFPQDLGQFYIREEMPPQYDEPIQVHK
 PKALEEPRFSKSDLRRKGLNRKLSALGVDPSRFDEVATFLDGTDPRELQKLKLWLKE
 AGYGVNQ

SV40 NLS amino acid sequence (SEQ ID NO:16):

PKKKRKV

Ty1 NLS amino acid sequence (SEQ ID NO:17):

NSKKRSLEDNETEIKVSRDTWNTKNMRSLEPPRSKKRIH

c-Myc NLS amino acid sequence (SEQ ID NO:18):

PAAKRVKLD

biSV40 NLS amino acid sequence (SEQ ID NO:19):

KRTADGSEFESPKKKRKV

Mut NLS amino acid sequence (SEQ ID NO:20):

PEKKRRRPSGSPVVLARSPPKAGKSSCI

Template DNA sequences (underlining marks the replaced or inserter editing sequences)**EMX1 HDR template sequence (SEQ ID NO:79):**

CATTCTGCCTCTCTGTATGGAAAAGAGCATGGGGCTGGCCCGTGGGGTGGTGTCCAC
 TTTAGGCCCTGTGGGAGATCATGGGAACCCACGCAGTGGGTcataggctctctcattactacat
 ccactctgtgaagaagcattatgatctctctctagaaaCTCGTAGAGTCCCATGTCTGCCGGCTTCCAGAG
 CCTGCACTCCTCCACCTTGGCTTGGCTTTGCTGGGGCTAGAGGAGCTAGGATGCACA
 GCAGCTCTGTGACCCTTTGTTTGAGAGGAACAGGAAAACACCCTTCTCTCTGGCCC
 ACTGTGTCCTCTTCTGCCCTGCCATCCCCTTCTGTGAATGTTAGACCCATGGGAGCA
 GCTGGTCAGAGGGGACCCCGGCCTGGGGCCCCTAACCTATGTAGCCTCAGTCTTCC
 CATCAGGCTCTCAGCTCAGCCTGAGTGTTGAGGCCCCAGTGGCTGCTCTGGGGGCCT
 CCTGAGTTTCTCATCTGTGCCCCCTCCCTCCCTGGCCCAGGTGAAGGTGTGGTTCCAG
 AACCGGAGGACAAAGTACAAACGGCAGAAGCTGGAGGAGGAAGGGCCTGAGTCCG
 AGCAGAAGAAGAAGGGCTCCCATCACATCAACCGGTGGCGCATTGCCACGAAGCAG
 GCCAATGGGGAGGACATCGATGTCACCTCCAATGACTCGGATGTACACGGTCTGCA
 ACCACAAACCCACGAGGGCAGAGTGCTGCTTGCTGCTGGCCAGGCCCTGCGTGGG
 CCCAAGCTGGACTCTGGCCACTCCCTGGCCAGGCTTTGGGGAGGCCTGGAGTCATGG
 CCCACAGGGCTTGAAGCCCGGGGCCGCGCATTGACAGAGGGACAAGCAATGGGCTG
 GCTGAGGCCTGGGACCACTTGGCCTTCTCCTCGGAGAGCCTGCCTGCCTGGGCGGGC
 CCGCCCCGCCACCGCAGCCTCCCAGCTGCTCTCCGTGTCTCCAATCTCCCTTTTGT
 TTTGATGCATTTCTGTTTTAATTTATTTTCCAGGCACCACTGTAGTTTAGTGATCCCCAGT
 GTCCCCCTTCCCTATGGGAATAATAAAAGTCTCTCTCTTAATGACACGGGCATCCAG

CTCCAGCCCCAGAGCCTGGGGTGGTAGATTCCGGCTCTGAGGGCCAGTGGGGGCTG
GTAGAGCAAACGCGTTCAGGGCCTGGGAGCCTGGGGTGGGGTACTGGTGGAGGGGG
TCAAGGGTAATTCATTAACCTCCTCTCTTTTGTGGGGGACCCTGGTCTCTACCTCCAG
CTCCACAGCAGGAGAAACAGGCTAGACATAGGGAAGGGCCATCCTGTATCTTGAGG
GAGGACAGGCCAGGTCTTTCTTAACGTATTGAGAGGTGGGAATCAGGCCAGGTA
GTTCAATGGG

VEGFA HDR template sequence (SEQ ID NO:80):

AGGTTTGAATCATCACGCAGGCCCTGGCCTCCACCCGCCCCACCAGCCCCCTGGCC
TCAGTTCCCTGGCAACATCTGGGGTTGGGGGGGCAGCAGGAACAAGGGCCTCTGTC
TGCCAGCTGCCTCCCCCTTTGGGTTTTGCCAGACTCCACAGTGCATACGTGGGCTC
CAACAGGTCCTCTCCCTCCAGTCACTGACTAACCCCGGAACCACACAGCTTCCCG
TTctcagctccacaaacttggtccaaattcttctccctgggaagcatccctggacacttccaaaggacccagtcactccagcctgtg
gctgccgctcactttgatgtctgcaggccagatgagggctccagatggcacattgtcagagggacacactgtggccctgtgcccagccct
gggtctctgtacatgaagcaactccagtcaccaaatatgtagctgtttgggaggtcagaataaggggtccaggagcaaaactccccacc
ccctttccaaagccattccctcttagccagagccgggtgtgcagacggcagtcactagggggcgctcgccaccacaggggaagctg
ggtgaatggagcgagcagcgtcttcgagagtgaggacgtgtgtctgtgtgggtgagtgtgtgCgcACTCTAGAGgtgtCg
Tgttgagggcggttgagcgggggagaagccaggggtcactccaggttccaatagatctgtgtgtccctctccccaccgtccctgtccg
gctctccgcttccctgccccctcaatattcctagcaaaagagggaacggctctcaggccctgtccgcacgtaacctcactttctgtccct
cctcgccaatgccccggggcggtgtctgtgacagagtttccggggcggtggttaattttcaggctgtgaaccttggtgggggtcga
gcttcccccttattggggcggtGCGGGCCAGGCTTCACTGAGCGTCCGCAGAGCCCGGGCCCCGA
GCCGCGTGTGAAGGGCTGAGGCTCGCTGTccccgcccccgggcgggcgggggcggggtcccg
cgggcgggAGCCATGCGCCCCCCCCtttttttAAAAGTCGGCTGGTAGCGGGGAGGATCGC
GGAGGCTTGGGGCAGCCGGGTAGCTCGGAGGTCGTGGCGCTGGGGGCTAGCACCAG
CGTCTGTCTGGGAGGGCGCAGCGGTTAGGTGGACCGGTCAGCGGACTCACCGGCCAG
GGCGCTCGGTGCTGGAATTTGATATTCATTGATCCGGGttttatccctcttcttttctaaacattttttA
AAACTGTATTGTTTCTCGTTTAAATTTATTTTGTCTTGCCATTCCCCACTTGAAT

DYNLT1 HDR template sequence (SEQ ID NO:81):

AGTGACCTGTGTAATTATGCAGAAGAATGGAGCTGGATTACACACAGCAAGTTCCTGCTTCT
GGGACAGCTCTACTGACGGTATGATTTTTCATTCATGTTTGTGAAGTTTTGTTGTGTGAATAT
ATGACTGGAAGTTTCTATCTTTGAATGCAATGCATGTTTATCACCTTTTAAAACATTTAATA
ATAGACTTGCCAAGGTTCTTTGTGTAGCATAGAGATGGGTACTTGAATGTTGGCCTTATTGTG
AGTAAAACGTCGTCCCCCAGCTTTCCCTGCCGTAAATGCTGCTCTCTTCCCTCCCGCAGGGAG
CTGCACTGTGCGATGGGAGAATAAGACCATGTACTGCATCGTCAGTGCCTTCGGACTGTCTA
TTGGAAGCGGAGCTACTAAGCTTACGCTGCTGAAGCAGGCTGGAGACGTGGAGGAGAACCC
TGGAACCTgccaccatggtgagcgagctgattaaggagaacatgcacatgaagctgtacatggaggccaccgtgaacaaccaccactcaagtgc
acatccgagggcggaaggcaagccctacgagggcaccagaccatgagaatcaaggcggtcagggcgccctctcccttcgcttcgacatcctgg
ctaccagcttcacatcgccagcaaaacctcatcaaccacaccagggcatccccacttcttaagcagtccttccccgagggcttcacatgggagag
agtcaccacatacgaagatggggcggtgctgaccgctacccaggacaccagcctccaggacggctgcctcatctacaacgtcaagatcagaggggtg
aacttcccacccaacggccctgtgatgcagaagaaaacactcggtgggagggcctccaccgagacactgtaccccgctgacggcggtggaaggca
gagccgacatggccctgaagctcgtggcgggggccacctgatctgcaaccttaagaccacatacagatccaagaaacccgctaagaacctcaagatg
cccggcgctactatgtggacaggagactggaagaatcaaggaggccgacaagagacatacgtcgagcagcacgaggtggctgtggccagatact
gagacctccctagcaactggggcacaacttaattccTAACCaGcGTCCtGCCTATGGCCTTTCTCCTTTTGTCTCT
AGTTCATCCTCTAACCACCAGCCATGAATTCAGTGAACCTTTTTCTCATTCTCTTTGTTTTGTG
GCACTTTCACAATGTAGAGGAAAAAACCAATGACCGCACTGTGATGTGAATGGCACCGAA

GTCAGATGAGTATCCCTGTAGGTCACCTGCAGCCTGCGTTGCCACTTGTCTTAACTCTGAATA
TTTCATTTCAAAGGTGCTAAAATCTGAAATCTGCTAGTGTGAACTTGTCTACTCTCTGAAA
TGATTCAAATACACTAATTTTCCATACTTTATACTTTTGTGTTAGAATAAATTATTCAAATCTAA
AGTCTGTTGTGTTCTTCATAGTCTGCATAGTATCATAAACG

[0100] HSP90AA1 HDR template sequence (SEQ ID NO:82):

GCAGCAAAGAAACACCTGGAGATAAACCTGACCATTCCATTATTGAGACCTTAAGGCAAA
AGGCAGAGGCTGATAAGAACGACAAGTCTGTGAAGGATCTGGTCATCTTGCTTTATGAACT
GCGCTCCTGTCTTCTGGCTTCAGTCTGGAAGATCCCCAGACACATGCTAACAGGATCTACAG
GATGATCAAACCTTGGTCTGGGTAAGCCTTATACTATGTAATGTTAAAAAGAAAATAAACACA
CGTGACATTGAAGAAAATGGTGAACCTTTCAGTTATCCAACTTGGAGCACCTTGTCTGCTT
GCTGCTTGGAGGTATTAAGTATGTTTTAGGGATAAGTAAGGTCTTACAAGAGCAAAGAAAT
GAAATTGAGACTCATATGTCCTGTAATACTGTCTTGAAAGCAGATAGAAACCAAGAGTATTA
CCCTAATAGCTGGCTTTAAGAAATCTTTGTAATATGAGGATTTTATTTTGGAAACAGGTATTG
ATGAAGATGACCCTACTGCTGATGATACCACTGCTGCTGTAACCTGAAGAAATGCCACCCCTT
GAAGGAGATGACGACACATCACGCATGGAAGAAGTAGACGGAAGCGGAGCTACTAAGCTTCA
GCCTGCTGAAGCAGGCTGGAGACGTGGAGGAGAACCCTGGACCTgtgagcgagctgattaaggagaaacatg
cacatgaagctgtacatggaggccaccgtgaacaaccaccacttcaagtgcacatccgagggcgaaggcaagccctacgagggcaccagaccatg
agaatcaaggcggcgcagggcgccctctcccttcgcttcgacatcctggctaccagcttcatgtacggcagcaaaccttcatcaaccacaccag
ggcatccccgacttctttaagcagtccttccccgagggcttcacatggagagagtcaccacatacgaagatggggcgtgctgaccgctaccagggac
accagctccaggagcgctgcctcatctacaacgtcaagatcagaggggtgaacttccatccaacggccctgtgatgcagaagaaacactcggtcgt
ggagggcctccaccgagacactgtaccccgctgacggcgccctgggaaggcagagccgacatggccctgaagctcgtggcgggggccacctgactg
caaccttaagaccacatacagatccaagaaccgctaagaacctcaagatgccggcgctctactatgtggacagagactggaagaatcaaggagg
ccgacaagagacatacgtcgagcagcagaggtggctgtggccagatactgcgacctccctagcaaaactggggcacaaacttaattccTAAATC
TgTGGCTGAGGGATGACTTACCTGTTCACTACTCTACAATTCCTCTGATAATATATTTTCAAG
GATGTTTTTCTTTATTTTGTAAATATTAAGTCTGTATGGCATGACAACCTACTTTAAGGG
GAAGATAAGATTTCTGTCTACTAAGTGATGCTGTGATACCTTAGGCACTAAAGCAGAGCTAG
TAATGCTTTTTGAGTTTCATGTTGGTTTATTTTCACAGATTGGGGTAACGTGCACTGTAAAGAC
GTATGTAACATGATGTTAACTTTGTGGTCTAAAGTGTTTAGCTGTCAAGCCGGATGCCTAAGT
AGACCAAATCTTGTTATTGAAGTGTTCTGAGCTGTATCTTGATGTTTAGAAAAGTATTCGTTA
CATCTTGATAGGATCTACTTTTTGAACCTTTTCATTCCCTGTAGTTGACAATTCTGCATGTACTAG
TCCTCTAGAAATAGGTTAACTGAAGCAACTTGATGGAAGGATCTCTCCACAGGGCTTGTTT
TCCAAAGAAAAGTATTGTTTGGAGGAGCAAAGTTAAAGCCTACCTAAGCATATCGTAAAG
CTGTTCAAAAATAACTCAGACCCAGTCTTGTGGA

[0101] AAVS1 HDR template sequence (SEQ ID NO:83):

gatgcttttcggagcacttctctcgccgctgcaccacgtgatgtcctctgagcggatcctccccgtgctggtcctctccggcactctctcctcctc
acccaacccatgcccgtcttctcctgctgggttccctttctctcctctctggggcctgtgccatctctcgttcttaggatggccttcccgacggatgtctcc
cttgcgtcccgctcccttctgtaggcctgcatcaccggtttttctggacaaccccaagtaccccgctccttggttagccacctctccatcctcttg
cttcttctgctggacacccgttctcctgtggatcgggtcactctcactcttctcatttgggcagctccctaccccttacctctctagtctgtgtagctc
ttccagccccctgtcatggcatctccaggggtccgagagctcagctagtcttcttctccaacccggggccctatgtccacttcaggacagcatgtttctg
cctccagggatcctgtgtcccgagctgggaccaccttatattccagggccggtaattgtggtctgtggttctgggtactttatctgtccctccacccac
agtggggcaagcttctgacctcttctctctccacagggcctcgagagatctggcagcggaGGAAGCGGAGCTACTAAGCTTCAAG
CCTGCTGAAGCAGGCTGGAGACGTGGAGGAGAACCCTGGACCTgtgagcgagctgattaaggagaaacatgca
catgaagctgtacatggaggccaccgtgaacaaccaccacttcaagtgcacatccgagggcgaaggcaagccctacgagggcaccagaccatgag
aatcaaggcggcgcagggcgccctctcccttcgcttcgacatcctggctaccagcttcatgtacggcagcaaaccttcatcaaccacaccaggg

catccccgacttctttaagcagtccttccccgagggcctcacatgggagagagtcaccacatacgaagatgggggctgctgaccgctaccagggacac
cagcctccaggagcggctgcctcatctacaacgtcaagatcagaggggtgaacttcccatccaacggcctgtgatgcagaagaaaacactcggctggg
aggcctccaccgagacactgtaccccgctgacggcgccctggaaggcagagccgacatggcctgaagctcgtggcgggggccacgtgatctgca
accttaagaccacatacagatccaagaaacccgctaagaacctcaagatccccggcgtctactatgtggacaggagactgaaagaatcaaggagggc
gacaaagagacatacgtcagcagcagcagagtggtgctgtggccagatactcgcacctccctagcaaaactggggcacaacttaattccTAaactaggg
acaggattggtgacagaaaagccccatccttaggcctcctcctctagtcctctgataattgggttaacccccacctcctgttaggcagattccttatctgt
gacacacccccatttctggagccatctctccttggcagaaccttaaggtttgcttacgatggagccagagaggatcctgggaggagagcttggca
gggggtgggagggaagggggggatgctgacctgccgggttcagtgccaccctgcgctaccctctccagaacctgagctgctctgacgcggct
gtctggtgcgtttcactgatcctgggtgctgcagcttcttacacttcccaaggaggagaagcagtttgaaaaacaaatcagaataagttggctcgtagttct
aactttggctcttcacctttctagtcctcaatttatattgtcctcggcgtcagttttacctgtgagataaggccagtagccagccccgctcctggcagggctg
tggtgaggaggggggtgtccgtgtgaaaactccctttgtgagaatgtgtgcgtcctaggtgttcaccaggtcgtggccgctctactccctttctcttcc
atccttctttccttaagagtccccagtgctatctgggacatattcctcgccagagcaggggtcccgttccttaagccctgctctgggcttctgggttga
tccttggc

OCT4 HDR template sequence (SEQ ID NO:84):

GCGACTATGCACAACGAGAGGATTTTGAGGCTGCTGGGTCTCCTTTCTCAGGGGGACCACTG
 TCCTTTCTCTGGCCCCAGGGCCCCATTTTGGTACCCAGGCTATGGGAGCCCTCACTTCACT
 GCACTGTACTCCTCGGTCCCTTTCCTTGAGGGGGAAGCCTTTCCCCCTGTCTCCGTCACCACT
 CTGGGCTCTCCCATGCATTCAAATGGAAGCGGAGCTACTAACTTCAGCCTGCTGAAGCAGGC
TGGAGACGTGGAGGAGAACCCTGGACCTgccaccatgtgagcgcgagctgattaaaggagaacctgcacatgaagctgtacat
ggagggcaccgtgaacaaccaccacttcaagtgcacatccgaggcggaaggcaagccctacgagggcaccagaccatgagaatcaaggcggtgcg
agggcgccctctcccccttcgccttcgacatcctggctaccagcttcatgtacggcagcaaaaccttcatcaaccacaccagggcatccccgacttctt
aagcagtccttccccgagggcttcacatgggagagagtcaccacatacgaagatggggcgctgctgaccgctaccagggacaccagcctccaggagc
gtgcctcatctacaacctcaagatcagaggggtgaacttccatccaacggcctgtgatgcagaagaaaacactcggctgggagggcctccaccgag
acactgtaccccgctgacggcgccctggaaggcagagccgacatggcctgaagctcgtggcgggggccacctgactgcaaccttaagaccacat
acagatccaagaaacccgctaagaacctcaagatgccggcgctactatgtggacaggagactggaaagaatcaaggagggcgacaaagagacata
cgtcagcagcagcagagtggtgctgtggccagatactcgcacctccctagcaaaactggggcacaacttaattccTAaTGACTAGGAATGG
 GGGACAGGGGGAGGGGAGGAGCTAGGGAAAGAAAACCTGGAGTTTGTGCCAGGGTTTTTGG
 GATTAAGTTCTTCATTCATAAGGAAGGAATTGGGAACACAAAGGGTGGGGGCAGGGGAGT
 TTGGGGCAACTGGTTGGAGGGAAGGTGAAGTTCAATGATGCTCTTGATTTTAATCCCACATC
 ATGTATCACTTTTTTCTTAATAAAGAAGCCTGGGACACAGTAGATAGACACACTT

Pantoea stewartii RecT DNA (SEQ ID NO:85):

AGCAACCAGCCCCCTATCGCCTCCGCCGATCTGCAGAAGGCCAACACCGGCAAGCAGGTGG
 CCAATAAGACCCCTGAGCAGACACTGGTGGGCTTCATGAATCAGCCAGCAATGAAGAGCCA
 GCTGGCCGCCGCCCTGCCAAGGCACATGACAGCCGATCGGATGATCAGAATCGTGACCACA
 GAGATCCGCAAGACCCCCGCCCTGGCCACATGCGACCAGAGCTCCTTCATCGGCGCCGTGGT
 GCAGTGTTCTCAGCTGGGCCTGGAGCCTGGCAGCGCCCTGGGCCACGCCTACCTGCTGCCAT
 TTGGCAACGGCCGGAGCAAGTCCGGACAGTCCAATGTGCAGCTGATCATCGGCTATAGAGG
 CATGATCGATCTGGCCCGGAGATCTGGCCAGATCGTGTCTCTGAGCGCCAGGGTGGTGC GCG
 CAGACGATGAGTTCTCCTTTGAGTACGGCCTGGATGAGAACCTGATCCACCGGCCAGGCGAG
 AATGAGGACGCACCCATCACCCACGTGTATGCAGTGGCAAGACTGAAGGACGGAGGCACCC
 AGTTCGAAGTGATGACAGTGAAGCAGATCGAGAAGGTGAAGGCCAGTCCAAGGCCTCTAG
 CAACGGACCCTGGGTGACCCACTGGGAGGAGATGGCCAAGAAAACCGTGATCAGGCGCCTG
 TTTAAGTACCTGCCCCTGAGCATCGAGATGCAGAAGGCCGTGATCCTGGATGAGAAGGCCG

AGTCTGACGTGGATCAGGACAATGCCTCCGTGCTGTCTGCCGAGTATAGCGTGCTGGACGGC
TCCTCTGAGGAG

Pantoea stewartii RecE DNA (SEQ ID NO:86):

CAGCCCGGCGTGTACTATGACATCTCCAACGAGGAGTATCACGCCGGCCCTGGCATCAGCAA
GTCCCAGCTGGACGACATCGCCGTGTCCCCAGCCATCTTCCAGTGGAGAAAGTCTGCCCCCG
TGGACGATGAGAAAACCGCCGCCCTGGACCTGGGCACAGCCCTGCACTGCCTGCTGCTGGA
GCCTGATGAGTTCTCCAAGAGGTTTATGATCGGCCAGAGGTGAACCGGAGAACCAATGCC
GGCAAGCAGAAGGAGCAGGACTTCCTGGATATGTGCGAGCAGCAGGGCATCACCCCTATCA
CACACGACGATAACCGGAAGCTGAGACTGATGAGGGACTCTGCCTTTGCCACCCAGTGGCC
AGATGGATGCTGGAGACAGAGGGCAAGGCCGAGGCCTCTATCTACTGGAATGACAGGGATA
CACAGATCCTGAGCAGGTGCCGCCCGACAAGCTGATCACCGAGTTCTCTTGGTGCGTGGAC
GTGAAGAGCACAGCCGACATCGGCAAGTTCCAGAAGGACTTCTACAGCTATCGCTACCACGT
GCAGGACGCCTTCTATTCCGATGGCTACGAGGCCAGTTTTGCGAGGTGCCAACCTTCGCCT
TTCTGGTGGTGAGCTCCTCTATCGATTGTGGCCGGTATCCCGTGCAGGTGTTTATCATGGACC
AGCAGGCAAAGGATGCAGGAAGGGCCGAGTATAAGCGGAACCTGACCACATACGCCGAGT
GCCAGGCAAGGAATGAGTGGCCTGGCATCGCCACACTGAGCCTGCCTTACTGGGCCAAGGA
GATCCGGAATGTG

Pantoea brenneri RecT DNA (SEQ ID NO:87):

AGCAACCAGCCCCCTATCGCCTCCGCCGATCTGCAGAAAACCCAGCAGTCCAAGCAGGTGG
CCAACAAGACCCCCTGAGCAGACACTGGTGGGCTTCATGAATCAGCCAGCAATGAAGAGCCA
GCTGGCCGCCGCCCTGCCAAGGCACATGACCGCCGATCGGATGATCAGAATCGTGACCACA
GAGATCCGCAAGACACCACAGCTGGCCCAGTGCAGCCAGAGCTCCTTCATCGGCGCCGTGGT
GCAGTGTCTCAGCTGGGCCTGGAGCCTGGCAGCGCCCTGGGCCACGCCTACCTGCTGCCAT
TTGGCAACGGCCGGTCCAAGTCTGGCCAGAGCAATGTGCAGCTGATCATCGGCTATAGAGGC
ATGATCGATCTGGCCCGGAGATCCGGACAGATCGTGAGCCTGTCCGCCAGGGTGGTGCGCGC
AGACGATGAGTTCTCTTTTGTAGTACGGCCTGGATGAGAACCTGGTGCACCGGCCAGGCGAGA
ATGAGGACGCACCCATCACCCACGTGTATGCAGTGGCAAGACTGAAGGACGGAGGCACCCA
GTTCAAGTGATGACAGTGAAGCAGGTGGAGAAGGTGAAGGCCAGTCCAAGGCCTCTAGC
AATGGCCCCTGGGTGACCCACTGGGAGGAGATGGCCAAGAAAACCGTGATCAGGCGCCTGT
TTAAGTACCTGCCCGTGAGCATCGAGATGCAGAAGGCCGTGGTGCTGGATGAGAAGGCCGA
GTCTGACGTGGATCAGGACAACGCCTCTGTGCTGAGCGCCGAGTATCCGTGCTGGAGTCTG
GCGACGAGGCCACAAAT

Pantoea brenneri RecE DNA (SEQ ID NO:88):

CAGCCTGGCATCTACTATGACATCAGCAACGAGGATTATCACAGGGGAGCAGGCATCAGCA
AGTCCCAGCTGGACGACATCGCCATCTCCCCAGCCATCTACCAAGTGGAGAAAGCAGCCCCC
GTGGACGAGGAGAAAACCGCCGCCCTGGATCTGGGCACAGCCCTGCACTGCCTGCTGCTGG
AGCCTGACGAGTTCTCTAAGAGGTTTCAGATCGGCCAGAGGTGAACCGGAGAACCACAGC
CGGCAAGGAGAAGGAGAAGGAGTTCATCGAGCGGTGCGAGGCAGAGGGAATCACCCCAAT
CACACACGACGATAATAGGAAGCTGAAGCTGATGAGGGATTCCGCCCTGGCCCACCCAATC
GCAAGGTGGATGCTGGAGGCACAGGGAACGCAGAGGCCTCTATCTATTGGAATGACAGAG
ATGCCGGCGTGCTGAGCAGGTGCCGCCCGACAAGATCATCACCGAGTTCAACTGGTGCGTG
GACGTGAAGTCCACAGCCGACATCATGAAGTTCCAGAAGGACTTCTACTTTACAGATACCA
CGTGACAGGACGCCCTTCTATTCCGATGGCTACGAGTCTCACTTTCACGAGACACCCACATTG

CCTTTCTGGCCGTGTCTACCAGCATCGACTGCGGCAGGTATCCTGTGCAGGTGTTTATCATGG
ACCAGCAGGCAAAGGATGCAGGAAGGGCCGAGTACAAGAGAAACATCCACACCTTCGCCGA
GTGTCTGAGCAGGAATGAGTGGCCTGGCATCGCCACACTGTCCCTGCCTTTTTGGGCCAAGG
AGCTGCGCAATGAG

Pantoea dispersa RecT DNA (SEQ ID NO:89):

TCCAACCAGCCACCTCTGGCCACCGCAGATCTGCAGAAAACCCAGCAGTCTAACCAGGTGGC
CAAGACCCCTGAGCAGACACTGGTGGGCTTCATGAATCAGCCAGCAATGAAGAGCCAGCTG
GCCGCCGCCCTGCCAAGGCACATGACCGCCGATCGGATGATCAGAATCGTGACCACAGAGA
TCCGAAGACACCCGCCCTGGCCAGTGCAGACCAGAGCTCCTTCATCGGAGCAGTGGTGCAG
TGTAAGCAGCTGGGCCTGGAGCCTGGCTCCGCCCTGGGCCACGCCTACCTGCTGCCATTTGG
CAACGGCCGGTCCAAGTCTGGCCAGAGCAATGTGCAGCTGATCATCGGCTATAGAGGCATG
ATCGATCTGGCCCGGAGATCCGGACAGATCGTGAGCCTGTCCGCCAGGGTGGTGCAGCGCAG
ACGATGAGTTCTCTTTTGAGTACGGCCTGGATGAGAACCTGATCCACCGGCCAGGCGACAAT
GAGTCCGCCCCCATCACCCACGTGTATGCAGTGGCAAGACTGAAGGACGGAGGCACCCAGT
TCGAAGTGATGACAGCCAAGCAGGTGGAGAAGGTGAAGGCCAGTCCAAGGCCTCTAGCAA
CGGACCCTGGGTGACCCACTGGGAGGAGATGGCCAAGAAAACCGTGATCAGGCGCCTGTTT
AAGTACCTGCCCCGTGAGCATCGAGATGCAGAAGGCCGTGGTGCTGGACGAGAAGGCCGAGA
GCGACGTGGATCAGGACAATGCCTCTGTGCTGAGCGCCGAGTATTCCGTGCTGGAGTCTGGC
ACAGGCGAG

Pantoea dispersa RecE DNA (SEQ ID NO:90):

GAGCCAGGCATCTACTATGACATCAGCAACGAGGCCTACCACTCCGGCCCCGGCATCAGCA
AGTCCCAGCTGGACGACATCGCCAGGAGCCCTGCCATCTTCCAGTGGCGCAAGGACGCCCA
GTGGATACCGAGAAAACCAAGGCCCTGGACCTGGGCACCGATTTCCTACTGCGCCGTGCTGG
AGCCAGAGAGGTTTGCAGACATGTATCGCGTGGGCCCTGAAGTGAATCGGAGAACCACAGC
CGGCAAGGCCGAGGAGAAGGAGTTCTTTGAGAAGTGTGAGAAGGATGGAGCCGTGCCCATC
ACCCACGACGATGCACGGAAGGTGGAGCTGATGAGAGGCTCCGTGATGGCCACCCTATCG
CCAAGCAGATGATCGCAGCACAGGGACACGAGAGGCCTCTATCTACTGGCACGACGAGAG
CACAGGCAACCTGTGCCGGTGTAGACCCGACAAGTTTATCCCTGATTGGAATTGGATCGTGG
ACGTGAAAACACAGCCGATATGAAGAAGTTCAGGCGCGAGTTTACGATCTGCGGTATCAC
GTGCAGGACGCCTTCTACACCGATGGCTATGCCGCCAGTTTGGCGAGCGGCCTACCTTCGT
GTTTGTGGTGACATCCACCACAATCGACTGCGGCAGATACCCACCGAGGTGTTCTTTCTGG
ATGAGGAGACAAAGGCCGCCGGCAGGTCTGAGTACCAGAGCAACCTGGTGACCTATTCCGA
GTGTCTGTCTCGCAATGAGTGGCCAGGCATCGCCACACTGTCTCTGCCCCACTGGGCCAAGG
AGCTGAGGAACGTG

Type-F symbiont of Plautia stali RecT DNA (SEQ ID NO:91):

TCCAACCAGCCCCCTATCGCCTCTGCCGATCTGCAGAAAACCCAGCAGTCTAAGCAGGTGGC
CAACAAGACCCCTGAGCAGACACTGGTGGGCTTCATGAATCAGCCAGCAATGAAGTCCAG
CTGGCCGCCGCCCTGCCAAGGCACATGACAGCCGATCGGATGATCAGAATCGTGACCACAG
AGATCCGCAAGACCCCCGCCCTGGCCACATGCGACCAGAGCTCCTTCATCGGAGCAGTGGTG
CAGTGTAGCCAGCTGGGCCTGGAGCCTGGCTCCGCCCTGGGCCACGCCTACCTGCTGCCATT
TGGAACGGCCGGTCCAAGTCTGGCCAGTCTAATGTGCAGCTGATCATCGGCTATAGAGGCA
TGATCGACCTGGCCCGGAGAAGCGGACAGATCGTGAGCCTGTCCGCCAGGGTGGTGCAGCGC

AGACGATGAGTTCTCCTTTGAGTACGGCCTGGATGAGAACCTGATCCACCGGCCAGGCGATA
ATGAGGACGCCCCCATCACCCACGTGTATGCAGTGGCAAGACTGAAGGACGGAGGCACCCA
GTTCAAGTGATGACAGCCAAGCAGGTGGAGAAGGTGAAGGCCAGAGCAAGGCCTCTAGC
AACGGACCCTGGGTGACCCACTGGGAGGAGATGGCCAAGAAAACCGTGATCAGGCGCCTGT
TTAAGTACCTGCCCGTGAGCATCGAGATGCAGAAGGCCGTGGTGCTGGATGAGAAGGCCGA
GAGCGACGTGGATCAGGACAATGCCTCTGTGCTGAGCGCCGAGTATTCCGTGCTGGAGGGC
GACGGCGGCGAG

Type-F symbiont of *Plautia stali* RecE DNA (SEQ ID NO:92):

CAGCCTGGCATCTACTATGACATCAGCAACGAGGATTATCACGGCGGCCCTGGCATCAGCAA
GTCCCAGCTGGACGACATCGCCATCTCCCCAGCCATCTACCAGTGGAGGAAGCACGCCCCCG
TGGACGAGGAGAAAACCGCCGCCCTGGATCTGGGCACAGCCCTGCACTGCCTGCTGCTGGA
GCCTGACGAGTTCTCTAAGAGATTTGAGATCGGCCAGAGGTGAACCGGAGAACCACAGCC
GGCAAGGAGAAGGAGAAGGAGTTCATGGAGAGGTGTGAGGCAGAGGGAGTGACCCCTATC
ACACACGACGATAATCGGAAGCTGAGACTGATGAGGGATAGCGCAATGGCCCACCCAATCG
CCAGATGGATGCTGGAGGCACAGGGAAACGCAGAGGCCTCTATCTATTGGAATGACAGGGA
TACCGGCGTGCTGAGCAGGTGCCGCCCCGACAAGATCATCACCGACTTCAACTGGTGCGTGG
ACGTGAAGTCCACAGCCGACATCATCAAGTTCCAGAAGGACTTTTACTCTTATCGCTACCAC
GTGCAGGACGCCTTCTATTCCGATGGCTACGAGTCTCACTTTGACGAGACACCAACATTCGC
CTTTCTGGCCGTGTCTACAAGCATCGATTGCGGCCGGTATCCCGTGCAAGGTGTTTCATCATGGA
CCAGCAGGCAAAGGATGCAGGAAGGGCCGAGTACAAGCGGAACATCCACACCTTTGCCGAG
TGTCTGAGCCGCAATGAGTGGCCTGGCATCGCCACACTGTCCCTGCCTTACTGGGCCAAGGA
GCTGCGGAATGAG

Providencia stuartii RecT DNA (SEQ ID NO:93):

AGCAACCCACCTCTGGCCCAGGCAGACCTGCAGAAAACCCAGGGGCACAGAGGTGAAGGAGA
AAACCAAGGATCAGATGCTGGTGGAGCTGATCAATAAGCCTTCCATGAAGGCACAGCTGGC
CGCCGCCCTGCCAAGGCACATGACACCCGACCGGATGATCAGAATCGTGACCACAGAGATC
AGAAAGACCCCCGCCCTGGCCACATGCGATATGCAGAGCTTCGTGGGAGCAGTGGTGCAGT
GTTCCCAGCTGGGCCTGGAGCCTGGCAACGCCCTGGGACACGCCTACCTGCTGCCTTTTGGC
AACGGCAAGTCTAAGAGCGGCCAGTCTAATGTGCAGCTGATCATCGGCTATCGGGGCATGAT
CGACCTGGCCCGGAGAAGCGGCCAGATCGTGTCCATCTCTGCCAGGACCGTGCGCCAGGGC
GATAACTTCCACTTTGAGTACGGCCTGAACGAGAATCTGACCCACGTGCCTGGCGAGAATGA
GGACTCTCCAATCACACACGTGTACGCAGTGGCAAGGCTGAAGGATGGAGGCGTGCAAGTTC
GAAGTGATGACCTATAACCAGATCGAGAAGGTGCGCGCCAGCTCCAAGGCAGGACAGAATG
GACCCTGGGTGAGCCACTGGGAGGAGATGGCCAAGAAAACCGTGATCAGGCGCCTGTTCAA
GTACCTGCCCCGTGTCTATCGAGATGCAGAAGGCCGTGATCCTGGACGAGAAGGCCGAGGCC
AACATCGATCAGGAGAATGCCACCATCTTTGAGGGCGAGTATGAGGAAGTGGGCACAGACG
GCAAG

Providencia stuartii RecE DNA (SEQ ID NO:94):

GAGGGCATCTACTATAACATCAGCAATGAGGACTACCACAACGGCCTGGGCATCTCCAAGTC
TCAGCTGGATCTGATCAATGAGATGCCTGCCGAGTATATCTGGTCCAAGGAGGCCCGTGG
ACGAGGAGAAGATCAAGCCTCTGGAGATCGGCACCGCCCTGCACTGCCTGCTGCTGGAGCC
AGACGAGTACCACAAGAGATATAAGATCGGCCCCGATGTGAACCGGAGAACAAATGCCGGC
AAGGAGAAGGAGAAGGAGTTCTTTGATATGTGCGAGAAGGAGGGCATCACCCCCATCACAC

ACGACGATAACCGGAAGCTGATGATCATGAGAGACTCTGCCCTGGCCCACCCTATCGCCAAG
TGGTGTCTGGAGGCCGATGGCGTGAGCGAGAGCTCCATCTACTGGACCGACAAGGAGACAG
ATGTGCTGTGCAGGTGTGCGCCAGACCGCATCATCACCGCCCACAACCTACATCGTGGATGTG
AAGTCTAGCGGCGACATCGAGAAGTTCGATTACGAGTACTACAACCTACAGATACCACGTGC
AGGACGCCTTTTACTCCGATGGCTATAAGGAGGTGACCGGCATCACCCCTACATTCTGTTC
TGGTGGTGTCTACCAAGATCGACTGCGGCAAGTACCCCGTGCGGACCTACGTGATGAGCGAG
GAGGCAAAGTCCGCCGGAAGGACCGCCTACAAGCACAACCTGCTGACCTATGCCGAGTGTG
TGAAAACCGATGAGTGGGCCGGCATCAGGACACTGTCTCTGCCCAGATGGGCAAAGGAGCT
GCGGAATGAG

Providencia sp. MGF014 RecT DNA (SEQ ID NO:95):

TCTAACCCCCCTCTGGCCCAGAGCGACCTGCAGAAAACCCAGGGCACAGAGGTGAAGGTGA
AAACCAAGGATCAGCAGCTGATCCAGTTCATCAATCAGCCTTCTATGAAGGCACAGCTGGCC
GCCGCCCTGCCAAGGCACATGACACCCGACCGGATGATCAGAATCGTGACCACAGAGATCA
GAAAGACCCCCGCCCTGGCCACATGCGATATGCAGTCTTCGTGGGCGCCGTGGTGCAGTGT
TCTCAGCTGGGCCTGGAGCCTGGCAACGCCCTGGGACACGCCTACCTGCTGCCTTTTGGCAA
CGGCAAGGCCAAGTCCGGCCAGTCTAATGTGCAGCTGATCATCGGCTATCGGGGCATGATCG
ACCTGGCCCGGAGATCCAACCAGATCATCTCTATCAGCGCCAGGACCGTGCGCCAGGGCGAT
AACTTCCACTTTGAGTACGGCCTGAATGAGGACCTGACCCACACACCTAGCGAGAATGAGG
ATTCCCAATCACCCACGTGTACGCAGTGGCAAGGCTGAAGGACGGAGGCGTGCAGTTTGA
AGTGATGACATATAACCAGGTGGAGAAGGTGCGCGCCAGCTCCAAGGCAGGACAGAATGGA
CCCTGGGTGAGCCACTGGGAGGAGATGGCCAAGAAAACCGTGATCAGGCGCCTGTTCAAGT
ACCTGCCCCGTGTCCATCGAGATGCAGAAGGCAGTGGTGTGCTGGACGAGAAGGCAGAGGCCAA
CGTGATCAGGAGAATGCCACCATCTTTGAGGGCGAGTATGAGGAAGTGGGCACAGATGGC
AAT

Providencia sp. MGF014 RecE DNA (SEQ ID NO:96):

AAGGAGGGCATCTACTATAACATCAGCAATGAGGACTACCACAACGGCCTGGGCATCTCCA
AGTCTCAGCTGGATCTGATCAATGAGATGCCTGCCGAGTATATCTGGTCCAAGGAGGCCCCC
GTGGACGAGGAGAAGATCAAGCCTCTGGAGATCGGCACCGCCCTGCACTGCCTGCTGCTGG
AGCCAGACGAGTACCACAAGAGATATAAGATCGGCCCCGATGTGAACCGGAGAACAAATGT
GGGCAAGGAGAAGGAGAAGGAGTTCTTTGATATGTGCGAGAAGGAGGGCATCACCCCCATC
ACACACGACGATAACCGGAAGCTGATGATCATGAGAGACTCTGCCCTGGCCCACCCTATCGC
CAAGTGGTGTCTGGAGGCCGATGGCGTGAGCGAGAGCTCCATCTACTGGACCGACAAGGAG
ACAGATGTGCTGTGCAGGTGTCGCCCAGACCGCATCATCACCGCCCACAACCTACATCATCGA
TGTGAAGTCTAGCGGCGACATCGAGAAGTTCGATTACGAGTACTACAACCTACAGATACCACG
TGCAGGACGCCTTTTACTCCGATGGCTATAAGGAGGTGACCGGCATCACCCCTACATTCTG
TTTCTGGTGGTGTCTACCAAGATCGACTGCGGCAAGTACCCCGTGCGGACCTACGTGATGAG
CGAGGAGGCAAAGTCCGCCGGAAGGACCGCCTACAAGCACAACCTGCTGACCTATGCCGAG
TGTCTGAAAACCGATGAGTGGGCCGGCATCAGGACACTGTCTCTGCCCAGATGGGCAAAGG
AGCTGCGGAATGAG

Shewanella putrefaciens RecT DNA (SEQ ID NO:97):

CAGACCGCACAGGTGAAGCTGAGCGTGCCCCACCAGCAGGTGTACCAGGACAACCTTCAATT
ATCTGAGCTCCCAGGTGGTGGGCCACCTGGTGGATCTGAACGAGGAGATCGGCTACCTGAAC

CAGATCGTGTTTAATTCTCTGAGCACCGCCTCTCCCCTGGACGTGGCAGCACCTTGGAGCGT
GTACGGCCTGCTGCTGAACGTGTGCCGGCTGGGCCTGTCCCTGAATCCAGAGAAGAAGCTGG
CCTATGTGATGCCCTCCTGGTCTGAGACAGGCGAGATCATCATGAAGCTGTACCCCGGCTAT
AGGGGCGAGATCGCCATCGCCTCTAACTTCAATGTGATCAAGAACGCCAATGCCGTGCTGGT
GTATGAGAACGATCACTTCCGCATCCAGGCAGCAACCGGCGAGATCGAGCACTTTGTGACA
AGCCTGTCCATCGACCCTAGGGTGC GCGGAGCATGCAGCGGAGGCTACTGTCGGTCCGTGCT
GATGGATAATAACAATCCAGATCTCTTATCTGAGCATCGAGGAGATGAACGCCATCGCCCAGA
ATCAGATCGAGGCCAACATGGGCAATACCCCTTGGAATCCATCTGGCGGACAGAGATGAA
TAGAGTGGCCCTGTACCGGAGAGCAGCAAAGGACTGGAGGCAGCTGATCAAGGCCACCCCA
GAGATCCAGTCCGCCCTGTCTGATACAGAGTAT

Shewanella putrefaciens RecE DNA (SEQ ID NO:98):

GGCACCGCCCTGGCCCAGACAATCAGCCTGGACTGGCAGGATACCATCCAGCCAGCATACA
CAGCCTCCGGCAAGCCTAACTTCCTGAATGCCCAGGGCGAGATCGTGGAGGGCATCTACACC
GATCTGCCTAATTCCGTGTATCACGCCCTGGACGCACACAGCTCCACCGGCATCAAGACATT
CGCCAAGGGCCGCCACCACTACTTTCGGCAGTATCTGTCTGACGTGTGCCGGCAGAGAACAA
AGCAGCAGGAGTACACCTTCGACGCCGGCACCTACGGCCACATGCTGGTGCTGGAGCCAGA
GAACTTCCACGGCAACTTCATGAGGAACCCCGTGCCTGACGATTTTCCAGACATCGAGCTGA
TCGAGAGCATCCCACAGCTGAAGGCCGCCCTGGCCAAGAGCAACCTGCCCCTGTCCGGAGC
AAAGGCCGCCCTGATCGAGAGACTGTACGCCTTCGACCCATCCCTGCCCCTGTTTGAGAAGA
TGAGGGAGAAGGCCATCACCGACTATCTGGATCTGCGCTACGCCAAGTATCTGCGGACCGAC
GTGGAGCTGGATGAGATGGCCACATTCTACGGCATCGATACCTCTCAGACACGGGAGAAGA
AGATCGAGGAGATCCTGGCCATCTCTCCTAGCCAGCCAATCTGGGAGAAGCTGATCAGCCAG
CACGTGATCGACCACATCGTGTGGGACGATGCCATGAGGGTGGAGAGATCCACCAGGGCCC
ACCCTAAGGCAGACTGGCTGATCTCTGATGGCTATGCCGAGCTGACAATCATCGCAAGGTGC
CCAACCACCGGCCCTGCTGCTGAAGGTGCGGTTTGACTGGCTGAGGAATGATGCCATCGGCGT
GGACTTCAAGACCACACTGTCTACCAACCCACAAAGTTTGGCTACCAGATCAAGGACCTGC
GGTATGATCTGCAGCAGGTGTTCTACTGTTATGTGGCCAATCTGGCCGGCATCCCTGTGAAG
CACTTCTGCTTTGTGGCCACCGAGTACAAGGACGCCGATAACTGTGAGACATTTGAGCTGTC
TCACAAGAAAGTGATCGAGAGCACCGAGGAGATGTTGACCTGCTGGATGAGTTTAAGGAG
GCCCTGACCTCCGGCAATTGGTATGGCCACGACAGGTCCCGCTCTACATGGGTCATCGAGGT
G

Bacillus sp. MUM 116 RecT DNA (SEQ ID NO:99):

AGCAAGCAGCTGACCACAGTGAATACCCAGGCCGTGGTGGGCACATTCTCCCAGGCCGAGC
TGGATACCCTGAAGCAGACAATCGCCAAGGGCACCACAAACGAGCAGTTCGCCCTGTTTGTG
CAGACCTGCGCCAACCTCTAGGCTGAATCCATTTCTGAACCACATCCACTGTATCGTGTATAA
CGGCAAGGAGGGGCGCCACCATGAGCCTGCAGATCGCAGTGGAGGGCATCCTGTACCTGGCA
CGCAAGACAGACGGCTATAAGGGCATCGAGTGCCAGCTGATCCACGAGAATGACGAGTTCA
AGTTTGATGCCAAGTCCAAGGAGGTGGATCACCAGATCGGATTCCCCAGGGGCAACGTGAT
CGGAGGATATGCAATCGCAAAGAGGGAGGGCTTTGACGATGTGGTGGTGTGCTGATGGAGTCT
AACGAGGTGGACCACATGCTGAAGGGCCGGAATGGCCACATGTGGAGAGACTGGTTCAACG
ATATGTTTAAGAAGCACATCATGAAGCGGGCCGCCAAGCTGCAGTACGGCATCGAGATCGC
AGAGGACGAGACAGTGAGCAGCGGACCTAGCGTGGATAATATCCCAGAGTATAAGCCACAG
CCCCGGAAGGACATCACACCCAACCAGGACGTGATCGATGCCCCCCTCAGCAGCCTAAGC
AGGACGATGAGGCCGCCAAGCTGAAGGCCGCCAGATCTGAGGTGAGCAAGAAGTTCAAGAA

GCTGGGCATCGTGAAGGAGGATCAGACCGAGTACGTGGAGAAGCACGTGCCTGGCTTCAAG
GGCACACTGTCCGACTTTATCGGCCTGTCTCAGCTGCTGGATCTGAATATCGAGGCCAGGA
GGCCAGTCCGCCGACGGCGATCTGCTGGAC

Bacillus sp. MUM 116 RecE DNA (SEQ ID NO:100):

ACCTACGCCGCCGACGAGACACTGGTGCAGCTGCTGCTGTCCGTGGATGGCAAGCAGCTGCT
GCTGGGAAGGGGCTGAAGAAGGGCAAGGCCAGTACTATATCAATGAGGTGCCATCTAAG
GCCAAGGAGTTCGAGGAGATCCGGGACCAGCTGTTTGACAAGGATCTGTTTCATGTCCCTGTT
TAACCCCTCTTACTTCTTTACCCTGCACTGGGAGAAGCAGAGGGCCATGATGCTGAAGTATG
TGACAGCCCCCGTGTCTAAGGAGGTGCTGAAGAATCTGCCTGAGGCCAGTCCGAGGTGCTG
GAGAGATACCTGAAGAAGCACTCTCTGGTGGATCTGGAGAAGATCCACAAGGACAACAAGA
ATAAGCAGGATAAGGCCTATATCTCTGCCAGAGCAGGACCAACACACTGAAGGAGCAGCT
GATGCAGCTGACCGAGGAGAAGCTGGACATCGATTCCATCAAGGCCGAGCTGGCCACATC
GACATGCAGGTCATCGAGCTGGAGAAGCAGATGGATACAGCCTTCGAGAAGAACCAGGCCT
TTAATCTGCAGGCCAGATCAGGAATCTGCAGGACAAGATCGAGATGAGCAAGGAGCGGTG
GCCCTCCCTGAAGAACGAAGTGATCGAGGATACCTGCCGGACATGCAAGCGGCCCTGGAC
GAGGATAGCGTGGAGGCCGTGAAGGCCGACAAGGATAATCGGATCGCCGAGTACAAGGCCA
AGCACAACCTCCCTGGTGTCTCAGAGAAATGAGCTGAAGGAGCAGCTGAACACCATCGAGTA
TATCGACGTGACAGAGCTGAGAGAGCAGATCAAGGAGCTGGATGAGTCCGGACAGCCTCTG
AGGGAGCAGGTGCGCATCTACAGCCAGTATCAGAATCTGGACACCCAGGTGAAGTCCGCCG
AGGCAGACGAGAACGGCATCCTGCAGGATCTGAAGGCCTCTATCTTCATCCTGGATAGCATC
AAGGCCTTTAGGGGCAAGGAGGCCGAGATGCAGGCCGAGAAGGTGCAGGCCCTGTTACCA
CACTGAGCGTGCCTGTTTAAAGCAGAATAAGGGCGACGGCGAGATCAAGCCAGATTTTCA
GATCGAGATGAACGACAAGCCCTATCGGACCCTGAGCCTGTCCGAGGGCATCCGGGCAGGC
CTGGAGCTGCGGGACGTGCTGAGCCAGCAGTCCGAGCTGGTGACCCCTACATTCGTGGATAA
TGCCGAGTCTATCACCAGCTTCAAGCAGCCAAACGGCCAGCTGATCATCAGCCGGGTGGTGG
CAGGACAGGAGCTGAAGATCGAGGCCGTGAGCGAG

Shigella sonnei RecT DNA (SEQ ID NO:101):

ACCAAGCAGCCCCCTATCGCCAAGGCCGACCTGCAGAAAACCCAGGAGAACAGGGCACCAG
CAGCCATCAAGAACAAATGATGTGATCTCCTTTATCAATCAGCCCTCTATGAAGGAGCAGCTG
GCCGCCGCCCTGCCTAGGCACATGACCGCCGAGAGGATGATCCGCATCGCCACCACAGAGA
TCCGCAAGGTGCCTGCCCTGGGCAACTGCGACACAATGAGCTTCGTGAGCGCCATCGTGCAG
TGTAAGCAGCTGGGCCTGGAGCCAGGCTCCGCCCTGGGCCACGCTACCTGCTGCCCTTCGG
CAACAAGAATGAGAAGTCCGGCAAGAAGAATGTGCAGCTGATCATCGGCTATAGGGGCATG
ATCGATCTGGCCCCGAGATCTGGCCAGATCGCCTCTCTGAGCGCCAGAGTGTTGCGGGAGG
GCGACGAGTTCAACTTTGAGTTCGGCCTGGATGAGAAGCTGATCCACCGGCCTGGCGAGAA
TGAGGACGCCCCAGTGACCCACGTGTACGCAGTGGCCAGACTGAAGGATGGCGGCACCCAG
TTTGAAGTGATGACAAGGCGCCAGATCGAGCTGGTGAGGTCCCAGTCTAAGGCCGGCAACA
ATGGCCCTTGGGTGACCCACTGGGAGGAGATGGCCAAGAAAACCGCCATCCGGAGACTGTT
CAAGTACCTGCCAGTGTCTATCGAGATCCAGCGCGCCGTGAGCATGGACGAGAAGGAGCCA
CTGACCATCGACCCCGCCGATAGCTCCGTGCTGACAGGCGAGTATTCTGTGATCGATAACAG
CGAGGAG

Shigella sonnei RecE DNA (SEQ ID NO:102):

GATCGCGGCCTGCTGACAAAGGAGTGGAGGAAGGGAAACCGGGTGAGCCGGATCACCAGG
ACAGCCAGCGGAGCAAACGCAGGAGGAGGAAATCTGACCGACAGAGGCGAGGGCTTCGTG
CACGATCTGACAAGCCTGGCCCCGCGACATCGCAACCGGCGTGTCTGGCCCCGAGCATGGACG
TGGACATCTACAACCTGCACCCTGCCCACGCCAAGAGGATCGAGGAGATCATCGCCGAGAA
TAAGCCCCCTTTTCAGCGTGTTTAGAGACAAGTTTATCACAATGCCAGGCGGCCTGGACTACT
CCAGGGCCATCGTGGTGGCCTCTGTGAAGGAGGCCCCAATCGGCATCGAAGTGATCCCCGCC
CACGTGACCGCCTATCTGAACAAGGTGCTGACCGAGACAGACCACGCCAATCCAGATCCCCG
AGATCGTGGACATCGCATGCGGCAGAAGCTCCGCCCTATGCCACAGAGGGTGACCGAGGA
GGGCAAGCAGGACGATGAGGAGAAGCTGCAGCCTTCTGGCACCACAGCAGATGAGCAGGG
AGAGGCAGAGACAATGGAGCCAGACGCCACAAAGCACCACCAGGATACCCAGCCTCTGGAC
GCCCAGAGCCAGGTGAACAGCGTGGATGCCAAGTATCAGGAGCTGAGAGCCGAGCTGCACG
AGGCCAGGAAGAATCCCTTCCAAGAATCCAGTGGACGCAGATAAGCTGCTGGCCGCCTC
TCGCGGCGAGTTTCGTGGACGGCATCAGCGACCCAAACGATCCCAAGTGGGTGAAGGGCATC
CAGACACGGGATTCCGTGTACCAGAATCAGCCTGAGACAGAGAAAACAGCCCCGACATGA
AGCAGCCAGAGCCTGTGGTGCAGCAGGAGCCTGAGATCGCCTTCAACGCCTGCGGACAGAC
CGGCGGCGACAATTGCCCAGATTGTGGCGCCGTGATGGGCGATGCCACCTATCAGGAGACA
TTTGACGAGGAGAACCAGGTGGAGGCCAAGGAGAATGATCCTGAGGAGATGGAGGGCGCC
GAGCACCACACAACGAGAATGCCGGCAGCGACCCCCACAGAGACTGTTCCGATGAGACAG
GCGAGGTGGCCGATCCCGTGATCGTGGAGGACATCGAGCCTGGCATCTACTATGGCATCAGC
AACGAGAATTACCACGCAGGCCCCGGCGTGTCCAAGTCTCAGCTGGACGACATCGCCGACA
CACCTGCCCTGTATCTGTGGAGGAAGAACGCCCCAGTGGATACCACAAAGACCAAGACACT
GGACCTGGGCACCGCATTCCTACTGCCGCGTGTCTGGAGCCAGAGGAGTTCAGCAATCGGTTTA
TCGTGGCCCCCGAGTTCAACCGGAGAACAAATGCCGGCAAGGAGGAGGAGAAGGCCTTTCT
GATGGAGTGTGCCTCCACAGGCAAGATGGTCATCACCGCCGAGGAGGGCAGAAAGATCGAG
CTGATGTACCAGTCTGTGATGGCACTGCCACTGGGACAGTGGCTGGTGGAGAGCGCCGGAC
ACGCAGAGTCTAGCATCTATTGGGAGGACCCCGAGACAGGCATCCTGTGCAGGTGTGCCCC
GACAAGATCATCCCTGAGTTCCACTGGATCATGGACGTGAAAACCACAGCCGACATCCAGC
GGTTCAAGACAGCCTACTATGATTACAGGTATCACGTGCAGGATGCCTTCTACTCCGACGGC
TATGAGGCCAGTTTGGCGTGCAGCCACCTTCGTGTTTCTGGTGGCCTCTACCACAATCGAG
TGCGGCAGATACCCCGTGGAGATCTTTATGATGGGAGAGGAGGCAAAGCTGGCCGGACAGC
TGGAGTATACCGCAACCTGCGGACACTGGCCGATTGTCTGAATACCGACGAGTGGCCAGCC
ATCAAGACCCTGTCCCTGCCCAGATGGGCAAAGGAGTACGCCAACGAC

Salmonella enterica RecT DNA (SEQ ID NO:103):

ACCAAGCAGCCCCCTATCGCCAAGGCCGACCTGCAGAAAACCCAGGGAAACAGGGCACCTG
CAGCAGTGAATGACAAGGATGTGCTGTGCGTGATCAACAGCCCTGCCATGAAGGCACAGCT
GGCCGCCGCCCTGCCAAGGCACATGACCGCCGAGAGGATGATCCGCATCGCCACCACAGAG
ATCAGGAAGGTGCCAGAGCTGCGCAACTGCGACAGCACCAGCTTCATCGGCGCCATCGTGC
AGTGTTCCTCAGCTGGGCCTGGAGCCCCGCGAGCGCCCTGGGCCACGCCTACCTGCTGCCTTTT
GGCAATGGCAAGGCCAAGAACGGCAAGAAGAATGTGCAGCTGATCATCGGCTATCGGGGCA
TGATCGATCTGGCCCCGAGATCTGGCCAGATCATCTCCCTGAGCGCCAGAGTGGTGCGGGAG
TGTGACGAGTTCTCCTACGAGCTGGGCCTGGATGAGAAGCTGGTGCACCGGCCAGGCGAGA
ACGAGGACGCACCCATCACCCACGTGTATGCCGTGGCCAAGCTGAAGGATGGCGGCGTGCA
GTTTGAAGTGATGACCAAGAAGCAGGTGGAGAAGGTGAGAGATACACACTCCAAGGCCGCC
AAGAATGCCGCCTCTAAGGGCGCCAGCTCCATCTGGGACGAGCACTTCGAGGATATGGCCA
AGAAAACCGTGATCCGGAAGCTGTTTAAGTACCTGCCCGTGAGCATCGAGATCCAGAGAGC

CGTGAGCATGGACGGCAAGGAGGTGGAGACAATCAACCCAGACGACATCAGCGTGATCGCC
GGCGAGTATTCCGTGATCGATAATCCCGAGGAG

Salmonella enterica RecE DNA (SEQ ID NO:104):

GATCGCGGCCTGCTGACAAAGGAGTGGAGGAAGGGAAACCGGGTGAGCCGGATCACCAGG
ACAGCCAGCGGAGCAAACGCAGGAGGAGGAAATCTGACCGACAGAGGCGAGGGCTTCGTG
CACGATCTGACAAGCCTGGCCCGCGACGTGGCAACCGGCGTGCTGGCCCGGAGCATGGACG
TGGACATCTACAACCTGCACCCTGCCCACGCCAAGAGGGTGGAGGAGATCATCGCCGAGAA
TAAGCCCCCTTTCAGCGTGTTTAGAGACAAGTTTATCACAATGCCTGGCGGCCTGGACTACT
CCAGGGCCATCGTGGTGGCCTCTGTGAAGGAGGCCCCCTATCGGCATCGAAGTGATCCCAGCC
CACGTGACCGAGTATCTGAACAAGGTGCTGACCGAGACAGACCACGCCAATCCAGATCCCCG
AGATCGTGGACATCGCATGCGGCAGAAGCTCCGCCCTATGCCACAGAGGGTGACCGAGGA
GGGCAAGCAGGACGATGAGGAGAAGCCCCAGCCTTCTGGAGCTATGGCCGACGAGCAGGCA
ACCGCAGAGACAGTGGAGCCAAACGCCACAGAGCACCACCAGAATACCCAGCCCCCTGGATG
CCCAGAGCCAGGTGAACTCCGTGGACGCCAAGTATCAGGAGCTGAGAGCCGAGCTGCAGGA
GGCCAGGAAGAACATCCCCTCCAAGAATCCTGTGGACGCGAGATAAGCTGCTGGCCGCCTCTC
GCGGCGAGTTTCGTGGATGGCATCAGCGACCCTAACGATCCAAAGTGGGTGAAGGGCATCCA
GACACGGGATTCCGTGTACCAGAATCAGCCCCGAGACAGAGAAGATCTCTCCTGACGCCAAG
CAGCCAGAGCCCGTGGTGCAGCAGGAGCCCGAGACAGTGTGCAACGCCTGTGGACAGACCG
GCGGCGACAATTGCCCTGATTGTGGCGCCGTGATGGGCGACGCCACATATCAGGAGACATTC
GGCGAGGAGAATCAGGTGGAGGCCAAGGAGAAGGACCCCGAGGAGATGGAGGGGAGCAGAG
CACCTCACAAACGAGAATGCCGGCAGCGACCCACACAGAGACTGTTCCGATGAGACAGGCG
AGGTGGCCGATCCAGTGATCGTGGAGGACATCGAGCCTGGCATCTACTATGGCATCAGCAAC
GAGAATTACCACGCAGGCCCCCGCGTGTCCAAGTCTCAGCTGGACGACATCGCCGACACAC
CCGCCCTGTATCTGTGGAGGAAGAACGCCCTGTGGATACCACAAAGACCAAGACACTGGA
CCTGGGCAACGCATTCCACTGCCGCGTGCTGGAGCCTGAGGAGTTCAGCAATCGGTTTATCG
TGGCCCCAGAGTTCAACCGGAGAACAATGCCGGCAAGGAGGAGGAGAAGGCCTTTCTGAT
GGAGTGTGCCTCCACCGGCAAGACAGTGATCACCGCCGAGGAGGGCAGAAAGATCGAGCTG
ATGTACCAGTCTGTGATGGCACTGCCTCTGGGACAGTGGCTGGTGGAGAGCGCCGGACACGC
AGAGTCTAGCATCTATTGGGAGGACCCCGAGACAGGCATCCTGTGCAGGTGTGCCCCAGAC
AAGATCATCCCCGAGTTCCACTGGATCATGGACGTGAAAACACAGCCGACATCCAGCGGTT
CAAGACAGCCTACTATGATTACAGGTATCACGTGCAGGATGCCTTCTACTCCGACGGCTATG
AGGCCAGTTTGGCGTGCAGCCAACCTTCGTGTTTCTGGTGGCCTCTACCACAGTGGAGTGC
GGCAGATACCCCGTGGAGATCTTTATGATGGGAGAGGAGGCAAAGCTGGCCGGACAGCAGG
AGTATCACCGCAACCTGCGGACACTGGCCGATTGTCTGAATACCGACGAGTGGCCTGCCATC
AAGACCCTGTCCCTGCCACGGTGGGCCAAGGAGTACGCCAACGAC

Acetobacter RecT DNA (SEQ ID NO:105):

AACGCCCCCAGAAAGCAGAATACCAGAGCCGCCGTGAAGAAGATCAGCCCTCAGGAGTTCG
CCGAGCAGTTTGGCGCCATCATCCCACAGGTGAAGTCCGTGCTGCCCCCCACGTGACCTTC
GAGAAGTTTGAAGCGGTGGTGAGACTGGCCGTGCGGAAGAACCCTGACCTGCTGACATGCT
CCCCAGCCTCTCTGTTTCATGGCATGTATCCAGGCAGCCTCCGACGGCCTGCTGCCTGATGGA
AGGGAGGGAGCAATCGTGAGCCGGTGGAGCTCCAAGAAGAGCTGCAACGAGGCCTCCTGGA
TGCCAATGGTGGCCGGCCTGATGAAGCTGGCCCCGAACAGCGGCGACATCGCCAGCATCTCT
AGCCAGGTGGTGTTCGAGGGCGAGCACTTTAGAGTGGTGCTGGGCGACGAGGAGGATCG
AGCACGAGCGCGATCTGGGCAAGACCGGCGGCAAGATCGTGGCAGCCTACGCCGTGGCAAG

GCTGAAGGACGGCAGCGATCCAATCCGCGAGATCATGTCCTGGGGCCAGATCGAGAAGATC
AGAAACACAAATAAGAAGTGGGAGTGGGGACCCTGGAAGGCCTGGGAGGACGAGATGGCC
AGAAAGACCGTGATCCGGAGACTGGCCAAGAGACTGCCCATGTCTACAGATAAGGAGGGAG
AGAGGCTGCGCAGCGCCATCGAGAGGATCGACTCCCTGGTGGACATCTCTGCCAACGTGGA
CGCACCTCAGATCGCAGCAGACGATGAGTTTGGCGCCGCCGCCACGGCGTGGAGCCACAG
CAGATCGCAGCACCTGACCTGATCGGCCGCTGGCCCAGATGCAGTCCCTGGAGCAGGTGCA
GGACATCGAGCCCCAGGTGTCTACGCCATCCAGGAGGCCGACAAGAGGGGCGACAGCGAT
ACAGCCAATGCCCTGGATGCCGCCCTGCAGAGCGCCCTGTCCCGCACCTCTACAGCCAAGGA
GGAGGTGCCTGCC

Acetobacter RecE DNA (SEQ ID NO:106):

GTGATCTCTAAGAGCGGCATCTACGACCTGACCAACGAGCAGTATCACGCCGATCCTTGCCC
AGAGATGTCCCTGAGCTCCTCTGGAGCCAGGGACCTGCTGAGCTCCTGTCTGCCAAGTTCA
TCGCCGCCAAGCAGCTGCCACAGCAGAATAAGAGGTGCTTTGACATCGGCTCTGCCGGACAC
CTGATGGTGTCTGGAGCCACACCTGTTTCGACCAGAAGGTGTGCGAGATCAAGCACCCCTGATTG
GCGCACAAAGGCAGCAAAGGAGGAGCGGGACGCCGCTACGCCGAGGGAAGAATCCCCCT
GCTGAGCCGCGAGGTGGAGGACATCAGGGCAATGCACTCCGTGGTGTGGAGAGATTCTCTG
GGAGCCAGGGCCTTCAGCGGAGGCAAGGCAGAGCAGTCCCTGGTGTGGCGCGACGAGGAGT
TTGGCATCTGGTGCCGGCTGCGGCCCGATTACGTGCCTAACAATGCCGTGCGGATCTTCGAC
TATAAGACCGCCACAAACGGCTCCCCCGATGCCTTTATGAAGGAGATCTACAATCGGGGCTA
TCACCAGCAGGCCGCTGGTATCTGGACGGATATGAGGCAGTGACCGGCCACAGGCCACGC
GAGTTCTGGTTTGTGGTGCAGGAGAAAACCGCCCCCTTCTGCTGTCTTTCTTTCAGATGGAT
GAGATGAGCCTGGAGATCGGCCGGACCCTGAACAGACAGGCCAAGGGCATCTTTGCCTGGT
GCCTGCGCAACAATTGTTGGCCAGGCTATCAGCCCGAGGTGGATGGCAAGGTGAGATTCTTT
ACCACATCTCCCCCTGCCTGGCTGGTGAGGGAGTACGAGTTTAAGAATGAGCACGGCGCCTA
TGAGCCACCCGAGATCAAGCGGAAGGAGGTGGCC

Salmonella enterica subsp. enterica serovar Javiana str. 10721 RecT DNA (SEQ ID NO:107):

CCAAAGCAGCCCCCTATCGCCAAGGCAGACCTGCAGAAAACCCAGGGAGCACGGACCCCAA
CAGCAGTGAAGAACAATAACGATGTGATCTCCTTTATCAATCAGCCTTCTATGAAGGAGCAG
CTGGCCGCCGCCCTGCCAAGGCACATGACCGCCGAGCGGATGATCAGAATCGCCACCACAG
AGATCAGGAAGGTGCCCGCCCTGGGCGACTGCGATACAATGTCTTTTGTGAGCGCCATCGTG
CAGTGTAGCCAGCTGGGCCTGGAGCCTGGCGGCGCCCTGGGCCACGCCTACCTGCTGCCTTT
CGGCAATCGGAACGAGAAGTCCGGCAAGAAGAATGTGCAGCTGATCATCGGCTATAGAGGC
ATGATCGACCTGGCCCGGAGATCCGGACAGATCGCCAGCCTGTCCGCCAGGGTGGTGCGCG
AGGGCGACGATTTCTCTTTTGTAGTTCGGCCTGGAGGAGAAGCTGGTGCACAGGCCAGGCGA
GAACGAGGACGCCCCCGTGACCCACGTGTACGCAGTGGCACGCCTGAAGGATGGAGGCACC
CAGTTTGAAGTGATGACACGGAAGCAGATCGAGCTGGTGAGAGCCAGTCTAAGGCCGGCA
ATAACGGCCCTTGGGTGACCCACTGGGAGGAGATGGCCAAGAAAACCGCCATCAGGCGCCT
GTTCAAGTACCTGCCCCTGAGCATCGAGATCCAGAGGGCCGTGAGCATGGATGAGAAGGAG
AACTGACAATCGACCCAGCCGATGCCAGCGTGATCACCGGCGAGTATTCCGTGGTGGAGA
ATGCCGGCGTGGAGGAGAACGTGACAGCC

Salmonella enterica subsp. enterica serovar Javiana str. 10721 RecE DNA (SEQ ID NO:108):

TACTATGACATCCCAAACGAGGCCTACCACGCAGGCCCCGGCGTGTCTAAGAGCCAGCTGG
ACGACATCGCCGATACCCCGCCATCTATCTGTGGCGGAAGAATGCCCTGTGGACACCGAG

AAAACCAAGTCCCTGGATACCGGCACAGCCTTCCACTGCAGGGTGCTGGAGCCAGAGGAGT
TCAGCAAGCGGTTTCATCATCGCCCCGAGTTCAACCGGAGAACCTCCGCGGCAAGGAGGA
GGAGAAAACCTTCCTGGAGGAGTGTACCCGGACAGGCAGAACCGTGCTGACAGCCGAGGAG
GGCAGGAAGATCGAGCTGATGTACCAGTCCGTGATGGCACTGCCACTGGGACAGTGGCTGG
TGGAGTCTGCCGGCTACGCCGAGAGCTCCGTGTATTGGGAGGACCCTGAGACAGGCATCCT
GTGCCGGTGTAGACCCGATAAGATCATCCCTGAGTTCCACTGGATCATGGACGTGAAAACCA
CAGCCGACATCCAGAGGTTTCGCACCGCCTACTATGACTACAGATACCACGTGCAGGACGCC
TTCTACTCTGATGGCTATAGAGCCCAGTTTGGCGAGATCCCTACATTCTGTGTTTCTGGTGGCC
AGCACACAGCAGAGTGCGGCAGATACCCCGTGGAGATCTTTATGATGGGAGAGGACGCAA
AGCTGGCCGGACAGCGCGAGTATAGGCGCAATCTGCAGACCCTGGCCGAGTGTCTGAACAA
TGATGAGTGGCCTGCCATCAAGACACTGTCTCTGCCACGGTGGGCCAAGGAGAACGCCAAT
GCC

Pseudobacteriovorax antillogorgiicola RecT DNA (SEQ ID NO:109):

GGCCACCTGGTGAGCAAGACCGAGCAGGATTACATCAAGCAGCACTATGCCAAGGGCGCCA
CAGACCAGGAGTTCGAGCACTTTATCGGCGTGTGCAGGGCCAGAGGCCTGAACCCAGCCGC
CAATCAGATCTACTTCGTGAAGTATCGGTCCAAGGATGGACCAGCAAAGCCAGCCTTTATCC
TGTCTATCGACAGCCTGAGGCTGATCGCACACCCGACCCGGCGATTACGCAGGATGCTCTGAG
CCCATCTTCACAGACGGCGGCAAGGCCTGTACCGTGACAGTGCGGAGAAACCTGAAGAGCG
GCGAGACAGGCAATTTCTCCGGCATGGCCTTTTATGACGAGCAGGTGCAGCAGAAGAACGG
CCGGCCTACCTCCTTTTGGCAGTCTAAGCCAAGAACAATGCTGGAGAAGTGTGCAGAGGCAA
AGGCCCTGAGGAAGGCCTTCCCTCAGGATCTGGGCCAGTTTACATCAGAGAGGAGATGCCC
CCTCAGTATGACGAGCCTATCCAGGTGCACAAGCCAAAGGCCCTGGAGGAGCCCAGGTTCA
GCAAGTCCGATCTGTCCAGGCGCAAGGGCCTGAACAGGAAGCTGTCTGCCCTGGGAGTGGA
CCCCAGCCGCTTCGATGAGGTGGCCACCTTTCTGGACGGCACACCTGATCGCGAGCTGGGCC
AGAAGCTGAAGCTGTGGCTGAAGGAGGCCGGCTACGGCGTGAATCAG

Pseudobacteriovorax antillogorgiicola RecE DNA (SEQ ID NO:110):

AGCAAGCTGTCCAACCTGAAGGTGTCTAATAGCGACGTGGATACACTGAGCCGGATCAGAA
TGAAGGAGGGCGTGTATCGGGACCTGCCAATCGAGAGCTACCACCAGTCCCCCGGCTATTCT
AAGACCAGCCTGTGCCAGATCGATAAGGCCCTATCTACCTGAAAACCAAGGTGCCACAGA
AGTCCACAAAGTCTCTGAACATCGGCACCGCCTTCCACGAGGCTATGGAGGGCGTGTTTAAG
GACAAGTATGTGGTGCACCCCGATCCTGGCGTGAATAAGACCACAAAGTCTTGGAAGGACTT
CGTGAAGAGGTATCCTAAGCACATGCCACTGAAGCGCAGCGAGTACGACCAGGTGCTGGCC
ATGTACGATGCCGCCCCGTCTTATAGACCTTTTTCAGAAGTACCACCTGAGCCGGGGCTTCTA
CGAGAGCTCCTTTTATTGGCACGATGCCGTGACAAACAGCCTGATCAAGTGCAGACCCGACT
ATATCACCCCTGATGGCATGAGCGTGATCGACTTCAAGACCACAGTGGACCCAGCCCCAAG
GGCTTTCAGTACCAGGCCTACAAGTATCACTACTACGTGAGCGCCGCCCTGACCCTGGAGGG
AATCGAGGCAGTGACCGGCATCAGGCCAAAGGAGTACCTGTTCTGCGCGTGTCCAATTCTG
CCCCATACCTGACCGCCCTGTATCGCGCCTCTGAGAAGGAGATCGCCCTGGGCGACCACTTT
ATCCGGCGGAGCCTGCTGACCCTGAAAACCTGTCTGGAGTCTGGCAAGTGGCCCGGCCTGCA
GGAGGAGATCCTGGAGCTGGGCCTGCCTTTCTCCGGCCTGAAGGAGCTGAGAGAGGAGCAG
GAGGTGGAGGATGAGTTTATGGAGCTGGTGGCC

Photobacterium sp. JCM 19050 RecT DNA (SEQ ID NO:111):

AACACCGACATGATCGCCATGCCCCCTTCTCCAGCCATCAGCATGCTGGACACAAGCAAGCT
GGATGTGATGGTGCGGGCAGCAGAGCTGATGTCCCAGGCCGTGGTCATGGTGCCCGACCACT
TCAAGGGCAAGCCAGCCGATTGCCTGGCAGTGGTCATGCAGGCAGACCAGTGGGGCATGAA
CCCCTTTACCGTGGCCCAGAAAACCCACCTGGTGAGCGGCACCCTGGGATACGAGTCCCAGC
TGGTGAATGCCGTGATCAGCTCCTCTAAGGCCATCAAGGGCCGGTTCCACTATGAGTGGTCT
GATGGCTGGGAGAGACTGGCCGGCAAGGTGCAGTACGTGAAGGAGTCTCGGCAGAGAAAG
GGCCAGCAGGGCAGCTATCAGGTGACCGTGGCCAAGCCAACATGGAAGCCAGAGGACGAGC
AGGGCCTGTGGGTGCGGTGTGGAGCCGTGCTGGCCGGAGAGAAGGACATCACATGGGGCCC
TAAGCTGTACCTGGCCAGCGTGCTGGTGCAGAACAGCGAGCTGTGGACCACAAAGCCCTAC
CAGCAGGCCGCCTATACCGCCCTGAAGGATTGGTCCCGCCTGTATACACCTGCCGTGATGCA
GGGCTCTATGACCGGCAAGAGCTGGTCCCTGACAGGCAGGCTGATCAGCCCCCGC

Photobacterium sp. JCM 19050 RecE DNA (SEQ ID NO:112):

GCCGAGCGGGTGAGAACCTATCAGCGGGACGCCGTGTTTCGCACACGAGCTGAAGGCCGAGT
TTGATGAGGCCGTGGAGAACGGCAAGACCGGCGTGACACTGGAGGACCAGGCCAGGGCCAA
GAGGATGGTGCACGAGGCCACCACAAACCCCGCCTCTCGGAATTGGTTCAGATACGACGGA
GAGCTGGCCGCATGCGAGAGGAGCTATTTTTGGCGCGATGAGGAGGCAGGCCTGGTGCTGA
AGGCCAGGCCTGACAAGGAGATCGGCAACAATCTGATCGATGTGAAGTCCATCGAGGTGCC
AACCGACGTGTGCGCCTGTGATCTGAACGCCTATATCAATCGGCAGATCGAGAAGAGAGGC
TACCACATCTCCGCCGCCACTATCTGTCTGGCACAGGCAAGGACCGCTTCTTTTGGATCTTC
ATCAATAAGGTGAAGGGCTACGAGTGGGTGGCAATCGTGGAGGCCTCTCCCCTGCACATCG
AGCTGGGCACCTATGAGGTGCTGGAGGGCCTGCGGAGCATCGCCAGCTCCACAAAGGAGGC
AGATTACCCAGCACCTCTGTCCCACCCTGTGAACGAGAGAGGCATCCCACAGCCCCTGATGT
CTAATCTGAGCACATACGCCATGAAGAGGCTGGAGCAGTTTCGCGAGCTG

Providencia alcalifaciens DSM 30120 RecT DNA (SEQ ID NO:113):

AAGGCACAGCTGGCCGCCGCCCTGCCTAAGCACATCACCAGCGACCGGATGATCAGAATCG
TGTCCACCGAGATCAGAAAGACCCCATCTCTGGCCAACTGCGACATCCAGAGCTTCATCGGC
GCCGTGGTGCAGTGTTCTCAGCTGGGCCTGGAGCCAGGCAACGCCCTGGGACACGCCTACCT
GCTGCCCTTTGGCAATGGCAAGTCCGACAACGGCAAGTCTAATGTGCAGCTGATCATCGGCT
ATCGGGGCATGATCGATCTGGCCCGGAGAAGCGGCCAGATCATCTCTATCAGCGCCAGGAC
CGTGCGCCAGGGCGACAACCTTCCACTTTGAGTACGGCCTGAACGAGAATCTGACCCACATCC
CCGAGGGCAATGAGGACTCCCCTATCACACACGTGTACGCAGTGGCACGGCTGAAGGATGA
GGGCGTGCAAGTTCGAAGTGATGACATATAACCAGATCGAGAAGGTGAGAGATAGCTCCAAG
GCCGGCAAGAATGGCCCCTGGGTGACCCACTGGGAGGAGATGGCCAAGAAAACCGTGATCA
GGCGCCTGTTTAAGTACCTGCCCGTGAGCATCGAGATGCAGAAGGCCGTGATCCTGGACGAG
AAGGCCGAGGCCAATATCGAGCAGGATCACTCCGCCATCTTCGAGGCCGAGTTTGAGGAGG
TGACTCTAACGGCAAT

Providencia alcalifaciens DSM 30120 RecE DNA (SEQ ID NO:114):

AACGAGGGCATCTACTATGACATCTCTAATGAGGACTATCACCACGGCCTGGGCATCTCTAA
GAGCCAGCTGGATCTGATCGACGAGAGCCCCGCCGATTTTCATCTGGCACCGGGATGCCCCTG
TGGACAACGAGAAAACCAAGGCCCTGGATTTTGGCACAGCCCTGCACTGCCTGCTGCTGGAG
CCAGACGAGTTCCAGAAGAGGTTTCGCATCGCCCCCGAGGTGAACCGGAGAACAATGCCG
GCAAGGAGCAGGAGAAGGAGTTCTGGAGATGTGCGAGAAGGAGAATATCACCCCCATCAC
AAACGAGGATAATAGGAAGCTGTCTCTGATGAAGGACAGCGCAATGGCCCCACCCTATCGCC

CGCTGGTGTCTGGAGGCCAAGGGCATCGCCGAGAGCTCCATCTATTGGAAGGACAAGGATA
 CAGACATCCTGTGCCGGTGTAGACCAGACAAGCTGATCGAGGAGCACCCTGGCTGGTGA
 TGTGAAGTCCACCGCCGACATCCAGAAGTTCGAGCGGTCTATGTACGAGTATAGATACCACG
 TGCAGGATTCCTTTTATTCTGACGGCTACAAGAGCCTGACAGGCGAGATGCCCCGTGTTCTG
 TTCCTGGCCGTGTCCACCGTGATCAACTGCGGCAGATACCCCGTGCAGGCTGTTCTGCTGGA
 CGAGCAGGCAAAGTCCGTGGGACGGATCACCTATAAGCAGAATCTGTTTACATACGCCGAG
 TGTCTGAAAACCGACGAGTGGGCCGGCATCAGAACCCTGAGCCTGCCCTCCTGGGCAAAGG
 AGCTGAAGCACGAGCACACCACAGCCTCT

Pantoea stewartii RecT Protein (SEQ ID NO:115):

MSNQPIASADLQKANTGKQVANKTPEQTLVGFMNQPMKSQLAAALPRHMTADRMIRIVTTEI
 RKTPALATCDQSSFIGAVVQCSQLGLEPGSALGHAYLLPFGNGRSKSGQSNVQLIIGYRGMIDLA
 RRSQIVSLSARVVRADDEFSFEYGLDENLIHRPGENEDAPITHVYAVARLKDGGTQFEVMTVK
 QIEKVKAQSKASSNGPWVTHWEEMAKKTVIRRLFKYLPVSIEMQKAVILDEKAESDQDQNAS
 VLSAEYSVLDGSSEE

Pantoea stewartii RecE Protein (SEQ ID NO:116):

MQPGVYYDISNEEYHAGPGISKSQLDDIAVSPAIFQWRKSAPVDDEKTAALDLGTALHCLLLEPD
 EFSKRFMIGPEVNRRTNAGKQKEQDFLDMCEQQGITPITHDDNRKLRLMRDSAFAPHPVARWML
 ETEGKAEASIYWNDRDTQILSRCRPDKLITEFSWCVDVKSTADIGKFQKDFYSYRYHVQDAFYSD
 GYEAQFCEVPTFAFLVVSSSIDCGRYPVQVFIMDQQAQKADAGRAEYKRNLTITYAECQARNEWPGI
 ATLSLPYWAKEIRNV

Pantoea brenneri RecT Protein (SEQ ID NO:117):

MSNQPIASADLQKTQSKQVANKTPEQTLVGFMNQPMKSQLAAALPRHMTADRMIRIVTTEI
 RKTPQLAQCDQSSFIGAVVQCSQLGLEPGSALGHAYLLPFGNGRSKSGQSNVQLIIGYRGMIDLA
 RRSQIVSLSARVVRADDEFSFEYGLDENLVHRPGENEDAPITHVYAVARLKDGGTQFEVMTVK
 QVEKVKAQSKASSNGPWVTHWEEMAKKTVIRRLFKYLPVSIEMQKAVVLDEKAESDQDQDNA
 SVLSAEYSVLESGDEATN

Pantoea brenneri RecE Protein (SEQ ID NO:118):

MQPGIYYDISNEDYHRGAGISKSQLDDIAISPAIYQWRKHAPVDEEKTAALDLGTALHCLLLEPD
 EFSKRFQIGPEVNRRTTAGKEKEKEFIERCEAEGITPITHDDNRKLKLMRDSALAHPIARWML
 QGNAEASIYWNDRDAGVLSRCRPDKIITEFNWCVDVKSTADIMKFQKDFYSYRYHVQDAFYSD
 GYESHFHETPTFAFLAVSTSIDCGRYPVQVFIMDQQAQKADAGRAEYKRNHTFAECLSRNEWPGIA
 TLSLPFWAKELRNE

Pantoea dispersa RecT Protein (SEQ ID NO:119):

MSNQPLATADLQKTQQSNQVAKTPEQTLVGFMNQPMKSQLAAALPRHMTADRMIRIVTTEI
 RKTPALAQCDQSSFIGAVVQCSQLGLEPGSALGHAYLLPFGNGRSKSGQSNVQLIIGYRGMIDLA
 RRSQIVSLSARVVRADDEFSFEYGLDENLIHRPGDNESAPITHVYAVARLKDGGTQFEVMTAK
 QVEKVKAQSKASSNGPWVTHWEEMAKKTVIRRLFKYLPVSIEMQKAVVLDEKAESDQDQDNA
 SVLSAEYSVLESGTGE

Pantoea dispersa RecE Protein (SEQ ID NO:120):

MEPGIYYDISNEAYHSGPGISKSQLDDIARSPAIFQWRKDAPVDTEKTKALDLGTDFHCAVLEPER
FADMYRVGPEVNRRTTAGKAEKEFFFEKCEKDGAVPITHDDARKVELMRGSVMAHPIAKQMIA
AQGHAEASIYWHEDESTGNLCRCRDPKFI PDWNWIVDVKTADMKKFRREFYDLRYHVQDAFYT
DGYAAQFGERPTFVFVVTSTTIDCGRYPTEVFFLDEETKAAGRSEYQSNLVTYSECLSRNEWPGI
ATLSLPHWAKELRNV

Type-F symbiont of *Plautia stali* RecT Protein (SEQ ID NO:121):

MSNQPIASADLQKTQQSKQVANKTPEQTLVGFMNQPAMKSQLAAALPRHMTADRMIRIVTTEI
RKTPALATCDQSSFIGAVVQCSQLGLEPGSALGHAYLLPFGNGRSKSGQSNVQLIIGYRGMIDLA
RRSGQIVSL SARVVRADDEFSEYGLDENLIHRPGDNEDAPITHVYAVARLKDGGTQFEVMTAK
QVEKVKAQSKASSNGPWVTHWEEMAKKTVIRRLFKYLPVSIEMQKAVVLDEKAESDQDQNA
SVLSAEYSVLEGDGGE

Type-F symbiont of *Plautia stali* RecE Protein (SEQ ID NO:122):

MQPGIYYDISNEDYHGGPGISKSQLDDIAISPAIQWRKHAPVDEEKTAALDLGTALHCLLLEPDE
FSKRFEIGPEVNRRTTAGKEKEKEFMERCEAEGVTPITHDDNRKLRLMRDSAMAHPIARWMLA
QGNAEASIYWNRDRTGVL SRCRPDKIITDFNWCVDVKSTADIIFQKDFYSYRYHVQDAFYSDG
YESHFDETPTF AFLAVSTSIDCGRYPVQVFIMDQQA K DAGRAEYKRNIHTFAECLSRNEWPGIAT
LSLPYWAKELRNE

Providencia stuartii RecT Protein (SEQ ID NO:123):

MSNPPLAQADLQKTQGTEVKEKTKDQMLVELINKPSMKAQLAAALPRHMTDPDRMIRIVTTEIRK
TPALATCDMQSFVGAVVQCSQLGLEPGNALGHAYLLPFGNGKSKSGQSNVQLIIGYRGMIDLAR
RSGQIVSISARTVRQGNDFHFEYGLNENLTHVPGENEDSPITHVYAVARLKDGGVQFEVMTYNQI
EKVRASSKAGQNGPWVSHWEEMAKKTVIRRLFKYLPVSIEMQKAVILDEKAEANIDQENATIFE
GEYEEVGTDGK

Providencia stuartii RecE Protein (SEQ ID NO:124):

EGIYYNISNEDYHNGLGISKSQLDLINEMPAEYIWSKEAPVDEEKIKPLEIGTALHCLLLEPDEYH
KRYKIGPDVNRRTNAGKEKEKEFFDMCEKEGITPITHDDNRKLMIMRDSALAHPIAKWCLEADG
VSESIYWTDKETDVLRCRCPDRIITAHNYIVDVKSSGDIEKFDY EYNYRYHVQDAFYSDGYKE
VTGITPTFLFLVSTKIDCGKYPVRTYVMSEEAKSAGRTAYKHNLITYAECLKTDEWAGIRTL SL
PRWAKELRNE

Providencia sp. MGF014 RecT Protein (SEQ ID NO:125):

MSNPPLAQSDLQKTQGTEVKVKTQQLIQFINQPSMKAQLAAALPRHMTDPDRMIRIVTTEIRKT
PALATCDMQSFVGAVVQCSQLGLEPGNALGHAYLLPFGNGKAKSGQSNVQLIIGYRGMIDLARR
SNQIISISARTVRQGNDFHFEYGLNEDLTHTPSENEDSPITHVYAVARLKDGGVQFEVMTYNQVE
KVRASSKAGQNGPWVSHWEEMAKKTVIRRLFKYLPVSIEMQKAVVLDEKAEANVDQENATIFE
GEYEEVGTDGN

Providencia sp. MGF014 RecE Protein (SEQ ID NO:126):

MKEGIYYNISNEDYHNGLGISKSQLDLINEMPAEYIWSKEAPVDEEKIKPLEIGTALHCLLLEPDE
YHKRYKIGPDVNRRTNVGKEKEKEFFDMCEKEGITPITHDDNRKLMIMRDSALAHPIAKWCLEA
DGVSESIYWTDKETDVLRCRCPDRIITAHNYIIDVKSSGDIEKFDY EYNYRYHVQDAFYSDGY

KEVTGITPTFLFLVVSTKIDCGKYPVRTYVMSEEAKSAGRTAYKHNLITYAECLKTDEWAGIRTL
SLPRWAKELRNE

Shewanella putrefaciens RecT Protein (SEQ ID NO:127):

MQTAQVKLSVPHQQVYQDNFNLYSSQVVGHVLDLNEEIGYLNQIVFNSLSTASPLDVAAPWSV
YGLLLNVCRLGLSLNPEKKLAYVMPSWSETGEIIMKLYPGYRGEIAIASNFNVIKNANAVLVYEN
DHFRIQAATGEIEHFVTSLSIDPRVRGACSGGYCRSVLMDNTIQISYLSIEEMNAIAQNQIEANMG
NTPWNSIWRTEMNRVALYRRAAKDWRQLIKATPEIQSALSDTEY

Shewanella putrefaciens RecE Protein (SEQ ID NO:128):

MGTALAQTTISLDWQDTIQPAYTASGKPNFLNAQGEIVEGIYTDLPNSVYHALDAHSSTGIKTFK
GRHHYFRQYLSDVCRQRTKQQEYTFDAGTYGHMLVLEPENFHGNFMRNPVPDDFPDIELIESIPQ
LKAALAKSNLPVSGAKAALIERLYAFDPSLPLEFKMREKAITDYLDLRYAKYLRTDVELDEMAT
FYGIDTSQTREKKIEELAISPSQPIWEKLISQHVHDHIVWDDAMRVERSTRAHPKADWLISDGYAE
LTIIARCPTTGLLLKVRFDWLRNDAIGVDFKTTLSTNPTKFGYQIKDLRYDLQQVFYCYVANLAG
IPVKHFCFVATEYKDADNCETFELSHKKVIESTEEMFDLLDEFKEALTSGNWDHRSRSTWVIE
V

Bacillus sp. MUM 116 RecT Protein (SEQ ID NO:129):

MSKQLTTVNTQAVVGTFSSQAEIDLKQTIAGTTNEQFALFVQTCANSRLNPFLNHHHCIVYNGK
EGATMSLQIAVEGILYLARKTDGYKGIECQLIHENDEFKFDKAKSKEVDHQIGFPRGNVIGGYAIA
KREGFDDVVLMESNEVDHMLKGRNGHMRDWFNDMFKKHIMKRAAKLQYGIEIAEDETSS
GPSVDNIPEYKPKPRKDITPNQDVIDAPPQQPKQDDEAAKLKAARSEVSKKFKKLGIKEDQTEY
VEKHVPGFKGTLSDFIGLSQLLDLNIEAQEAQSADGDLDD

Bacillus sp. MUM 116 RecE Protein (SEQ ID NO:130):

MTYAADETLVQLLLSVDGKQLLLGRGLKKGKAQYYINEVPSKAKEFEEIRDQLFDKDLFMSLFN
PSYFFTLHWKQRAMMLKYVTAPVSKEVLKLNPEAQSEVLERYLKKHSLVDLEKIHKDNKNKQ
DKAYISAQSRTNTLKEQLMQLTEEKLDIDSIKAEALAHIDMQVIELEKQMDTAFEKNQAFNLQAQI
RNLQDKIEMSKERWPSLKNEVIEDTCRTCKRPLDEDSVEAVKADKDNRIAEYKAKHNSLVSRN
ELKEQLNTIEYIDVTELREKQELDESQPLREQVRIYSQYQNLDTQVKSADENGILQDLKASIF
ILDSIKAFRGKEAEMQAQKQALFTTSLVRLFKQNGDGEIKPDFEIEMNDKPYRTLTLSEGIRAG
LELRDVLSSQSELVPTPTFDNAESITSFKQPNGQLIISRVVAGQELKIEAVSE

Shigella sonnei RecT Protein (SEQ ID NO:131):

MTKQPPIAKADLQKTQENRAPAAIKNNDVISFINQPSMKEQLAAALPRHMTAERMIRIATTEIRK
VPALGNCDTMSFVSAIVQCSQLGLEPGSALGHAYLLPFGNKNEKSGKKNVQLIIGYRGMIDLARR
SGQIASLSARVVREGDEFNFEFGLDEKLIHRPGENEDAPVTHVYAVARLKDGGTQFEVMTRRQIE
LVRSQSKAGNNGPWVTHWEEMAKKTAIRRLFKYLPVSIEIQRAVSMDEKEPLTIDPADSSVLTGE
YSVIDNSEE

Shigella sonnei RecE Protein (SEQ ID NO:132):

DRGLLTKEWRKGNRVSRI RTASGANAGGGNLDRGEGFVHDLTSLARDIATGVLARSMDVDI
YNLHPAHAKRIEIIAENKPPFSVFRDKFITMPGGLDYSRAIVVASVKEAPIGIEVIPAHVTAYLNK
VLTETDHANPDPEIVDIACGRSSAPMPQVRVTEEGKQDDEEKLQPSGTTADEQGEAETMEPDATK
HHQDTQPLDAQSQVNSVDAKYQELRAELHEARKNIPSKNPVDADKLLAASRGFVVDGISDPNDP

KWVKGITRDSVYQNQPETEKTS PDMKQPEPVVQQEPEIAFNACGQTGGDNCPDCGAVMGDAT
YQETFDEENQVEAKENDPEEMEGAEPHNENAGSDPHRDCSDETGEVADPVIVEDIEPGIYYGIS
NENYHAGPGVSKSQLDDIADTPALYLWRKNAPVDTTTKTKTLDLGTAFHCRVLEPEEFSNRFIVAP
EFNRRTNAGKEEEKAFLEMECASTGKMVITAEEGRKIELMYQSVMALPLGQWLVESAGHAESSIY
WEDPETGILCRCRPDKIPEFWIMDVKTTADIQRFKTAYYDYRYHVQDAFYSDGYEAQFGVQP
TFVFLVASTTIECGRYPVEIFMMGEEAKLAGQLEYHRNLRTLADCLNTDEWPAIKTSLPRWAKE
YAND

Salmonella enterica RecT Protein (SEQ ID NO:133):

MTKQPPIAKADLQKTQGNRAPAAVNDKDVLCVINSPAMKAQLAAALPRHMTAERMIRIATTEIR
KVPELRNCDSTSFIGAIVQCSQLGLEPGSALGHAYLLPFGNGKAKNGKKNVQLIIGYRGMIDLAR
RSGQIISLSARVVRECDEFSEYELGLDEKLVHRPGENEDAPITHVYAVAKLKDGGVQFEVMTKKQ
VEKVRDTHSKAAKNAASKGASSIWDEHFEDMAKKTVIRKLFKYLPVSIEIQRAVSMGKEVETI
NPDDISVIAGEYSVIDNPEE

Salmonella enterica RecE Protein (SEQ ID NO:134):

DRGLLTKEWRKGNRVSRITRTASGANAGGGNLTDRGEGFVHDLTSLARDVATGVLARSMDVDI
YNLHPAHAKRVEEIIAENKPPFSVFRDKFITMPGGLDYSAIVVASVKEAPIGIEVIPAHVTEYLNK
VLTTEDHANPDPEIVDIACGRSSAPMPQVRTEEGKQDDEEKQPSGAMADEQATAETVEPNATE
HHQNTQPLDAQSQVNSVDAKYQELRAELQEARKNIPSKNPVDADKLLAASRGEFVDGISDPNDP
KWVKGITRDSVYQNQPETEKISPDQKQPEPVVQQEPEVCNACGQTGGDNCPDCGAVMGDAT
YQETFGEENQVEAKEKDPEEMEGAEPHNENAGSDPHRDCSDETGEVADPVIVEDIEPGIYYGIS
NENYHAGPGVSKSQLDDIADTPALYLWRKNAPVDTTTKTKTLDLGTAFHCRVLEPEEFSNRFIVA
PEFNRRTNAGKEEEKAFLEMECASTGKTVITAEEGRKIELMYQSVMALPLGQWLVESAGHAESSI
YWEDPETGILCRCRPDKIPEFWIMDVKTTADIQRFKTAYYDYRYHVQDAFYSDGYEAQFGVQ
PTFVFLVASTTVECGRYPVEIFMMGEEAKLAGQQEYHRNLRTLADCLNTDEWPAIKTSLPRWA
KEYAND

Acetobacter RecT Protein (SEQ ID NO:135):

MNAPQKQNTAAVKKISPQFAEQFAAIIPQVKSVPALHVTFEKFERVVRLAVRKNPDLLTCSA
SLFMACIQAAASDGLLPDREGAIVSRWSSKKSCNEASWMPMVAGLMKLARNSGDIASISSQVVF
EGEHFRVVLGDEERIEHERDLGKTGGKIVAAAYAVARLKDGSPIREIMSWGQIEKIRNTNKKWE
WGPWKAWEDEMARKTVIRRLAKRLPMSTDKEGERLRSIERIDSLVDISANVDAPQIAADDEFA
AAAHGVEPQQAAPDLIGRLAQMQSLEQVQDIEPQVSHAIQEADKRGDSDTANALDAALQSALS
RTSTAKEEVPA

Acetobacter RecE Protein (SEQ ID NO:136):

MVISKSGIYDLTNEQYHADPCPEMSLSSSGARDLLSSCPAKFIAAKQLPQQNKRCFDIGSAGHLM
VLEPHLFDQKVCEIKHPDWRTKAAKEERDAAYAEGRIPLLSREVDIRAMHSVVWRDSL GARAF
SGGKAEQSLVWRDEEFGIWCRLRPDYVPNNVRIFDYKTATNGSPDAFMKEIYNRGYHQQA
AWYLDGYEAVTGHRPREFWFVQEKTAFFLLSFFQMDEMSLEIGRTLNRQAKGIFAWCLRNNCWP
GYQPEVDGKVRFFTTSPPAWL VREYEFKNEHGAYEPPEIKRKEVA

Salmonella enterica subsp. enterica serovar Javiana str. 10721 RecT Protein (SEQ ID NO:137):

MPKQPPIAKADLQKTQGARTPTAVKNNNDVISFINQPSMKEQLAAALPRHMTAERMIRIATTEIR
KVPALGDCDTMSFVSAIVQCSQLGLEPGGALGHAYLLPFGNRNEKSGKKNVQLIIGYRGMIDLA

RRSGQIASLSARVVREGDDFSFEFGLEEKLVHRPGENEDAPVTHVYAVARLKDGGTQFEVMTRK
QIELVRAQSKAGNNGPWVTHWEEMAKKTAIRRLFKYLPVSIQRAVSMDEKETLTIDPADASVI
TGEYSVVENAGVEENVTA

Salmonella enterica subsp. enterica serovar Javiana str. 10721 RecE Protein (SEQ ID NO:138):
MYDIPNEAYHAGPGVSKSQLDDIADTPAIYLWRKNAPVDTEKTKSLDTGTAFHCRVLEPEEFS
KRFIIAPEFNRRTSAGKEEEKTFLEECTRTGRTVLTAEGRKIELMYQSVMALPLGQWLVESAGY
AESSVYWEDPETGILCRCPDKIIEPFHWIMDVKTADTADIRFRTAYDYRYHVQDAFYSDGYRA
QFGEIPTFVFLVASTTAECGRYPVEIFMMGEDAKLAGQREYRRNLQTLAECLNDEWPAIKTSL
PRWAKENANA

Pseudobacteriovorax antillogorgiicola RecT Protein (SEQ ID NO:139):
MGHLVSKTEQDYIKQHYAKGATDQEFHFVCRARGLNPAANQIYFVKYRSKDGPAKPAFILSI
DSLRLIAHRTGDYAGCSEPIFTDGGKACTVTVRNLKSGETGNFSGMAFYDEQVQKNGRPTSF
WQSKPRTMLEKCAEAKALRKAFPQDLGQFYIREEMPPQYDEPIQVHKPKALEEPRFSKSDLRRK
GLNRKLSALGVDPSRFEDEVATFLDGTDPRELQKLLWLKEAGYGVNQ

Pseudobacteriovorax antillogorgiicola RecE Protein (SEQ ID NO:140):
MSKLSNLKVSNSDVTLSRIRMKEGVYRDLPIESYHQSPGYSKTSLCQIDKAPIYLLTKVPQKSTK
SLNIGTAFHEAMEGVFKDKYVVPDPGVNKTTSKWKDFVKRYPKHMPLKRSEYDQVLAMYDA
ARSYRPFQKYHLRSGFYESSFYWHDAVTNSLIKCRPDYITPDGMSVIDFKTTVDPSPKGFQYQAY
KYHYVSAALTLEGIEAVTGIRPKEYLFLAVSNSAPYLTALYRASEKEIALGDHFIRRSLLTLKTC
LESGKWPGQLQEEILELGLPFSGLKELREEQEVEDEFMELVG

Photobacterium sp. JCM 19050 RecT Protein (SEQ ID NO:141):
MNTDMIAMPPSPAISMLDTSKLDVMVRAAELMSQAVVMVPDHFKGKPADCLAVVMQADQWG
MNPFTVAQKTHLVSGTLGYESQLVNAVSSSKAIKGRFHYEWSGWERLAGKVQYVKESRQRK
GQQGSYQVTVAKPTWKPEDEQGLWVRGAVLAGEKDITWGPKLYLASVLVRNSELWTTKPYQ
QAAYTALKDWSRLYTPAVMQGSMTGKSWSLTGRLISPR

Photobacterium sp. JCM 19050 RecE Protein (SEQ ID NO:142):
MAERVRTYQRDAVFAHELKAEFDEAVENGKTGVTLEDQARAKRMVHEATTNPASRNWFRYDG
ELAACERSYFWRDEEAGLVLKARPDKEIGNNLDVKSIEVPTDVCACDLNAYINRQIEKRGYHIS
AAHYLSGTGKDRFFWIFINKVKGYEWWAIVEASPLHIELGTYEVLGLRSIASSTKEADYPAPLSH
PVNERGIPQPLMSNLSTYAMKRLEQFREL

Providencia alcalifaciens DSM 30120 RecT Protein (SEQ ID NO:143):
MKAQLAAALPKHITSRDMIRIVSTEIRKTPSLANCDIQSFIGAVVQCSQLGLEPGNALGHAYLLPF
GNGKSDNGKSNVQLIIGYRGMIDLARRSGQIISARTVRQGNFHFYGLNENLTHIPEGNEDSPI
THVYAVARLKDEGVQFEVMTYNQIEKVRDSSKAGKNGPWVTHWEEMAKKTVIRRLFKYLPVSI
EMQKAVILDEKAEANIEQDHSIFAEEFEVDSNGN

Providencia alcalifaciens DSM 30120 RecE Protein (SEQ ID NO:144):
MNEGIYYDISNEDYHHGLGISKSQLDLIDESPADFIWHRDAPVDNEKTKALDFGTALHCLLLEPD
EFQKRFRIAPEVNRRTNAGKEQEKEFLEMCEKENITPITNEDNRKLSLMKDSAMAHPIARWCLEA
KGIAESSIYWKDKDITDILCRCPDKLIEHHWLVDVKSTADIQKFERSMYEYRYHVQDSFYSDG

YKSLTGEMPVFVFLAVSTVINCGRYPVRVFLDEQAKSVGRITYKQNLFTYAECLKTDEWAGIR
TSLSPSWAKELKHEHTTAS

Mouse Albumin knock-in sense template (SEQ ID NO: 160)

CACCTTCAGATTTTCCTGTAACGATCGGGAAGTGGCATCTTCAGGGAGTAGctgacctcttctctctcc
cacaggATCCTGGAGCCACCCGCAGTTCGAAAAGCTCAGTGAAGAGAAGAACAAAAAGCAGCA
TATTACAGTTAGTTGTCTTCATCAATCTTTAAATATGTTGTGTGGTTTTTCTCTCCCTGTTTCC
AC

Mouse Albumin knock-in anti-sense template (SEQ ID NO: 161)

GTGGAAACAGGGAGAGAAAAACACACAACATATTTAAAGATTGATGAAGACAACCTAACTG
TAATATGCTGCTTTTTGTTCTTCTCTTCACTGAGCTTTTCGAACTGCGGGTGGCTCCAGGATcct
gtgggaggaagagaagaggtcagCTACTCCCTGAAGATGCCAGTTCCCGATCGTTACAGGAAAATCTGAA
GGTG

(SEQ ID NO: 162)

ACTTTGAGTGTAGCAGAGAGGAACCATTTGCCACCTTCAGATTTTCCTGTAACGATCGGGAAC
TGGCATCTTCAGGGAGTAGCTGACCTCTTCTCTCTCTCCACAGGATCCTGGAGCCACC

[0102] All references, including publications, patent applications, and patents, cited herein are hereby incorporated by reference to the same extent as if each reference were individually and specifically indicated to be incorporated by reference and were set forth in its entirety herein.

[0103] Preferred embodiments of this invention are described herein, including the best mode known to the inventors for carrying out the invention. Variations of those preferred embodiments may become apparent to those of ordinary skill in the art upon reading the foregoing description. The inventors expect skilled artisans to employ such variations as appropriate, and the inventors intend for the invention to be practiced otherwise than as specifically described herein. Accordingly, this invention includes all modifications and equivalents of the subject matter recited in the claims appended hereto as permitted by applicable law. Moreover, any combination of the above-described elements in all possible variations thereof is encompassed by the invention unless otherwise indicated herein or otherwise clearly contradicted by context.

CLAIMS

What is claimed is:

1. A system comprising:
 - a Cas protein;
 - a nucleic acid molecule comprising a guide RNA sequence that is complementary to a target DNA sequence; and
 - a microbial recombination protein,wherein the microbial recombination protein is selected from the group consisting of RecE, RecT, lambda exonuclease, Bet protein, exonuclease gp6, single-stranded DNA-binding protein gp2.5, or a derivative or variant thereof.
2. The system of claim 1, further comprising a recruitment system comprising:
 - at least one aptamer sequence; and
 - an aptamer binding protein functionally linked to the microbial recombination protein as part of a fusion protein.
3. The system of claim 2, wherein the at least one aptamer sequence is an RNA aptamer sequence or a peptide aptamer sequence.
4. The system of claim 3, wherein the nucleic acid molecule comprises the at least one RNA aptamer sequence.
5. The system of claim 4, wherein the nucleic acid molecule comprises two RNA aptamer sequences.
6. The system of claim 5, wherein the two RNA aptamer sequences comprise the same sequence.
7. The system of any of claims 2-6, wherein the aptamer binding protein comprises a MS2 coat protein, or a functional derivative or variant thereof.
8. The system of any of claims 2-6, wherein the aptamer binding protein comprises phage N peptide, or a functional derivative or variant thereof.

9. The system of claim 3, wherein the at least one peptide aptamer sequence is conjugated to the Cas protein.
10. The system of claim 9, wherein the at least one peptide aptamer sequence comprises between 1 and 24 peptide aptamer sequences.
11. The system of claim 9 or 10, wherein the aptamer sequences comprise the same sequence.
12. The system of any of claims 2-3 or 9-11, wherein the aptamer sequence comprises a GCN4 peptide sequence.
13. The system of any of claims 2-12, wherein the microbial recombination protein N-terminus is linked to the aptamer binding protein C-terminus.
14. The system of any of claims 2-13, wherein the fusion protein further comprises a linker between the microbial recombination protein and the aptamer binding protein.
15. The system of claim 14, wherein the linker comprises the amino acid sequence of SEQ ID NO: 15.
16. The system of any of claims 2-15, wherein the fusion protein further comprises a nuclear localization sequence.
17. The system of claim 16, wherein the nuclear localization sequence comprises the amino acid sequence of SEQ ID NO: 16.
18. The system of claim 16 or claim 17, wherein the nuclear localization sequence is on the microbial recombination protein C-terminus.
19. The system of any of claims 1-18, wherein the RecE or RecT recombination protein is derived from *E. coli*.

20. The system of any of claims 1-19, wherein the microbial recombination protein comprises RecE, or derivative or variant thereof.
21. The system of any of claims 1-20, wherein the RecE, or derivative or variant thereof, comprises an amino acid sequence with at least 70% similarity to amino acid sequences selected from the group consisting of SEQ ID NOs: 1-8.
22. The system of any of claims 1-21, wherein the RecE, or derivative or variant thereof, comprises an amino acid sequence with at least 70% similarity to amino acid sequences selected from the group consisting of SEQ ID NOs: 1-3.
23. The system of any of claims 1-19, wherein the fusion protein comprises RecT, or derivative or variant thereof.
24. The system of any of claims 1-19 or 23, wherein the RecT, or derivative or variant thereof, comprises an amino acid sequence with at least 70% similarity to amino acid sequences selected from the group consisting of SEQ ID NOs: 9-14.
25. The system of any of claims 1-19 or 23-24, wherein the RecT, or derivative or variant thereof, comprises an amino acid sequence with at least 70% similarity to amino acid sequences selected from the group consisting of SEQ ID NO: 9.
26. The system of any of claims 1-25, wherein the Cas protein is catalytically dead.
27. The system of any of claims 1-26, wherein the Cas protein is Cas9 or Cas12a.
28. The system of any of claims 27, wherein the Cas9 protein is wild-type *Streptococcus pyogenes* Cas9 or a wild-type *Staphylococcus aureus* Cas9.
29. The system of any of claims 27-28, wherein the Cas9 protein is a Cas9 nickase.

30. The system of claim 29, wherein the Cas9 nickase is wild-type *Streptococcus pyogenes* Cas9 with an amino acid substitution at position 10 of D10A.
31. The system of any of claims 1-30, further comprising donor nucleic acid.
32. The system of any of claims 1-31, wherein the target DNA sequence is a genomic DNA sequence in a host cell.
33. A composition comprising:
a polynucleotide comprising a nucleic acid sequence encoding a fusion protein comprising a microbial recombination protein functionally linked to an aptamer binding protein,
wherein the microbial recombination protein is RecE, RecT, lambda exonuclease, Bet protein, exonuclease gp6, single-stranded DNA-binding protein gp2.5, or a derivative or variant thereof.
34. The composition of claim 33, further comprising at least one of:
a polynucleotide comprising a nucleic acid sequence encoding a Cas protein; and
a nucleic acid molecule comprising a guide RNA sequence that is complementary to a target DNA sequence.
35. The composition of claim 34, wherein the nucleic acid molecule further comprises at least one RNA aptamer sequence.
36. The composition of claim 34, wherein the polynucleotide comprising a nucleic acid sequence encoding a Cas protein further comprises a sequence encoding at least one peptide aptamer sequence.
37. A vector comprising a polynucleotide comprising a nucleic acid sequence encoding a fusion protein comprising a microbial recombination protein functionally linked to an aptamer binding protein,
wherein the microbial recombination protein is RecE, RecT, lambda exonuclease, Bet protein, exonuclease gp6, single-stranded DNA-binding protein gp2.5, or a derivative or variant thereof.
38. The vector of claim 37, further comprising at least one of:

a polynucleotide comprising a nucleic acid sequence encoding a Cas protein; and
a nucleic acid molecule comprising a guide RNA sequence that is complementary to a target DNA sequence.

39. The vector of claim 38, wherein the nucleic acid molecule further comprises at least one RNA aptamer sequence.

40. The vector of claim 38, wherein the polynucleotide comprising a nucleic acid sequence encoding a Cas protein further comprises a sequence encoding at least one peptide aptamer sequence.

41. A eukaryotic cell comprising the system of any one of claims 1-32, the composition of any one of claims 33-36, or the vector of any of claims 37-40.

42. A method of altering a target genomic DNA sequence in a cell, comprising introducing the system of any one of claims 1-32, the composition of any one of claims 33-36, or the vector of any one of claims 37-40 into a cell comprising a target genomic DNA sequence.

43. The method of claim 42, wherein the cell is a mammalian cell.

44. The method of claim 42 or claim 43, wherein the cell is a human cell.

45. The method of any one of claims 42-44, wherein the cell is a stem cell.

46. The method of any one of claims 42-45, wherein the target genomic DNA sequence encodes a gene product.

47. The method of any one of claims 42-46, wherein the introducing into a cell comprises administering to a subject.

48. The method of claim 47, wherein the subject is a human.

49. The method of claim 47 or 48, wherein the administering comprises in vivo administration.

50. The method of claim 47 or 48, wherein the administering comprises transplantation of ex vivo treated cells comprising the system, composition, or vector.

51. Use of the system of any one of claims 1-32, the composition of any one of claims 33-36, or the vector of any one of claims 37-40 for the alteration of a target DNA sequence in a cell.