

PDK-1/Akt Signaling Inhibitors

Cross-Reference to Related Applications

[0001] This application claims priority to US Provisional Patent Application No. 60/508,619, titled "PDK-1/Akt Signaling Inhibitors" filed October 3, 2003 and US Provisional Patent Application No. 60/509,814, titled "PDK-1/Akt Signaling Inhibitors" filed October 8, 2003; the entirety of each of these is incorporated herein by reference.

Statement Regarding Federally Funded Research


[0002] This invention was made, at least in part, with government support under National Institutes of Health Grant CA94829 and Army Grant DAMD 17-02-1-0117. The government may have certain rights in this invention.

Background of the Invention









[0003] The phosphoinositide (PI) 3-kinase/PDK-1/Akt signaling cascade represents a convergence point for a plethora of receptor tyrosine kinase and cytokine-mediated pathways that regulate cell proliferation and survival, and offers a framework to account for the ability of many extracellular trophic factors to maintain cell survival. Dysregulation of this signaling cascade due to constitutive growth factor-receptor activation and/or PTEN mutations results in Akt up-regulation, which subsequently, promotes tumor invasiveness, angiogenesis, and progression. Thus, PDK-1/Akt signaling inhibitors are of translational relevance for development into useful chemotherapeutic or chemopreventive agents. There exists a need for development of new compounds that are potent PDK-1/Akt signaling inhibitors. There further exists a need for the development of chemotherapeutic agents and chemopreventative agents based on PDK-1/Akt signaling inhibition.

Summary of the Invention

[0004] Provided are compounds of formula I:



wherein X is selected from alkyl and haloalkyl; Ar is selected from the group consisting of phenyl, biphenyl, naphthyl, anthryl, phenanthryl, and fluorenyl; R is selected from the group

consisting of $-\text{CN}$, $-\text{CH}_2\text{CN}$, $-\text{CH}_2\text{CH}_2\text{CN}$, $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CN}$, $-\text{CONH}_2$, , , , , , , , , and . Formula I also includes pharmaceutically acceptable salts thereof, metabolism products, and prodrugs thereof.

Brief Description of the Figures


[0005] **Figure 1** shows dose-dependent effects of compounds **70** (left panels) and **71** (right panels) on cell viability of PC-3 cells and on the cell growth in nine representative human tumor cell lines.

[0006] **Figure 2** shows the synthesis of compounds **37-72** using the 1,1,1-trifluoro-4-hydroxy-4-phenanthren-2-yl-but-3-en-2-one as a common precursor.










Detailed Description of the Invention

[0007] Provided is a new class of PDK-1/Akt signaling inhibitors. The compounds described herein are useful for inducing apoptosis in undesirable proliferating cells, such as cancer cells. The compounds described herein are also useful in promoting wound healing and preventing scarring. The compounds described herein further have application in preventing restenosis. The compounds described herein further have application in organ transplantation.

[0008] The compounds described herein can be shown in the general formula I:



wherein X is selected from alkyl and haloalkyl; Ar is selected from the group consisting of phenyl, biphenyl, naphthyl, anthryl, phenanthryl, and fluorenyl; R is selected from the group


consisting of $-\text{CN}$, $-\text{CH}_2\text{CN}$, $-\text{CH}_2\text{CH}_2\text{CN}$, $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CN}$, $-\text{CONH}_2$, , , , , , , , , and . Stated otherwise, R is selected from nitrile, acetonitrile,

ethylnitrile, propylnitrile, carboxamide, amidine, pyrazole, oxime, hydrazone, acetamidine, acetamide, guanidine, and urea. Formula I also includes pharmaceutically acceptable salts thereof, metabolism products, and prodrugs thereof.


[0009] In some embodiments, X is C_1 to C_4 haloalkyl. In some embodiments, X is CF_3 . In some embodiments, Ar may be substituted at any substitutable position with one or more radicals, such as, but not limited to halo, C_1 - C_4 alkyl, C_1 - C_4 haloalkyl, azido, C_1 - C_4 azidoalkyl, aryl, alkaryl, haloaryl, haloalkaryl, and combinations thereof. In some embodiments, Ar is selected from 2-naphthyl, 4-biphenyl, 9-anthryl, 2-fluorenyl, 4-azidophenyl, 4-azidomethylphenyl, 4-(2-azidoethyl)phenyl, 4-(3-azidopropyl)phenyl, 4-(4-azidobutyl)phenyl, 4-(4-azidophenyl)phenyl, 4-(4-azidomethylphenyl)phenyl, 4-methylphenyl, 4-ethylphenyl, 4-propylphenyl, 4-butylphenyl, 4-(2-bromoethyl)phenyl, 4-(3-bromopropyl)phenyl, 4-(4-bromobutyl)phenyl, 4-(trifluoromethyl)phenyl, 4-(4-methylphenyl)phenyl, 4-(4-bromomethylphenyl)phenyl, 4-(4-butylphenyl)phenyl, 4-(4-*tert*-butylphenyl)phenyl, 2-chlorophenyl, 4-chlorophenyl, 2,4-dichlorophenyl,

3,4-dichlorophenyl, 2,5-dichlorophenyl, 2,4-dimethylphenyl, 2,5-dimethylphenyl, 3,4-dimethylphenyl, 3,5-dimethylphenyl, 4-(4-chlorophenyl)phenyl, 4-(3,5-dichlorophenyl)phenyl, 4-(2,3-dichlorophenyl)phenyl, 4-(3,5-dimethylphenyl)phenyl, 4-(2,4,5-trichlorophenyl)phenyl, 4-(4-trifluoromethylphenyl)phenyl, 2-phenanthrenyl, 3-indolyl, 2-pyrrolyl, and 4-(benzyl)phenyl. In some embodiments, Ar is selected from 4-(2-bromoethyl)phenyl, 4-(3-bromopropyl)phenyl, 4-(2-azidoethyl)phenyl; 4-(3-azidopropyl)phenyl, 4-butylphenyl, 4-*t*-butylphenyl, 2-naphthalenyl, 3-indolyl, 4-biphenyl, 4'-chloro[1,1'-biphenyl]-4-yl, 3',5'-dichloro[1,1'-biphenyl]-4-yl, 2',3'-dichloro[1,1'-biphenyl]-4-yl, 4'-methyl[1,1'-biphenyl]-4-yl, 4'-trifluoromethyl[1,1'-biphenyl]-4-yl, 4'-bromomethyl[1,1'-biphenyl]-4-yl, 3',5'-dimethyl[1,1'-biphenyl]-4-yl, 4'-butyl[1,1'-biphenyl]-4-yl, 4'-*tert*-butyl[1,1'-biphenyl]-4-yl, 4-(phenylmethyl)phenyl, 9H-fluoren-2-yl, 9-anthracenyl, 2-phenanthrenyl, 9-phenanthrenyl. In some embodiments, Ar is 2-phenanthrenyl. In some embodiments, R is selected from aminoacetamide and guanidine.

[0010] Another embodiment described herein is that of formula II:




wherein X is selected from alkyl and haloalkyl; R is selected -CN, -CH₂CN, -CH₂CH₂CN,




otherwise, R is selected from nitrile, acetonitrile, ethylnitrile, propylnitrile, carboxamide, amidine, pyrazole, oxime, hydrazone, acetamidine, acetamide, guanidine, and urea. In some embodiments, X is C₁ to C₄ haloalkyl, and in some embodiments, X is CF₃. In some

embodiments, R is aminoacetamide or guanidine. Formula II also includes pharmaceutically acceptable salts thereof, metabolism products, and prodrugs thereof.

[0011] Another embodiment described herein is that of formula III:




wherein R is selected from the group consisting of -CN, -CH₂CN, -CH₂CH₂CN,




otherwise, R is selected from nitrile, acetonitrile, ethylnitrile, propylnitrile, carboxamide, amidine, pyrazole, oxime, hydrazone, acetamidine, acetamide, guanidine, and urea. In some embodiments, R is aminoacetamide or guanidine. Formula II also includes pharmaceutically acceptable salts thereof, metabolism products, and prodrugs thereof.


[0012] Some additional compounds of formula III include the following groups for R:



[0013] In another embodiment, the compounds are that of formula IV or V:



IV



V

[0014] Provided also are methods of using the compounds of formulae I - V to induce apoptosis in the undesirable proliferating cells in subjects in need of such treatment. The method involves treating the subject in need of such treatment with a therapeutically effective amount of a compound of formulae I - V or derivative, metabolites, or pharmaceutically acceptable salts thereof.

[0015] The compounds and methods described herein are useful for, but not limited to treating, inhibiting, or delaying the onset of cancers. The compounds and methods are also useful in the treatment of precancers and other incidents of undesirable cell proliferation. The compounds of formula I, II, III, IV or V, or combinations thereof, are administered to a subject that has been diagnosed with or is at risk of developing a disorder characterized by undesirable cell proliferation. The compounds and methods are useful for treating cancers including, but not limited to, leukemia, melanoma, non-small cell lung cancer, colon cancer, cancers of the central nervous system, ovarian cancer, breast cancer, kidney cancer, and prostate cancer. Furthermore, they are useful in the slowing the growth of these cancers in individuals with precancers, as well as individuals prone to or having a genetic predisposition to these disorders.

[0016] The compounds are useful in methods of inducing apoptosis in unwanted rapidly proliferating cells, the method comprising introducing a therapeutically effective amount of a compound of formula I, II, III, IV, or V to the unwanted rapidly proliferating cells. In accordance with this method, the unwanted rapidly proliferating cells may be cancer cells. The cancer cells may be selected from the group consisting of leukemia, melanoma, non-small cell lung cancer, colon cancer, cancers of the central nervous system, ovarian cancer, breast cancer, kidney cancer, and prostate cancer.

[0017] The compounds are further useful for preventing restenosis in a subject who has undergone an angioplasty or stent procedure comprising administering a therapeutically effective amount of a compound of formula I, II, III, IV, or V, or combinations thereof, or pharmaceutically acceptable salts and/or metabolites thereof to the subject who has undergone an angioplasty or stent procedure.

[0018] The following terms used herein include, but are not limited to the following definitions.

[0019] The term “PDK-1/Akt signaling inhibitor” signifies that a specific compound or combination of compounds is capable of disrupting the PDK-1/Akt signaling pathway, as measured versus a blank, regardless of whether *in vivo* or *in vitro*. One method is set forth in the examples below, though other methods now known or later developed may also be used.

[0020] The term “treatment” as used herein, encompasses the administration and/or application of one or more compounds described herein, to a subject, for the purpose of providing prevention of or management of, and/or remedy for a condition. “Treatment” for the purposes of this disclosure, may, but does not have to, provide a cure; rather, “treatment” may be in the form of management of the condition. When the compounds described herein are used to treat unwanted proliferating cells, including cancers, “treatment” includes partial or total destruction of the undesirable proliferating cells with minimal destructive effects on normal cells. A desired mechanism of treatment of unwanted rapidly proliferating cells, including cancer cells, at the cellular level is apoptosis.

[0021] The term “prevention” as used herein includes either preventing or slowing the onset of a clinically evident unwanted cell proliferation altogether or preventing or slowing the onset of a preclinically evident stage of unwanted rapid cell proliferation in individuals at risk. Also intended to be encompassed by this definition is the prevention or slowing of metastasis of malignant cells or to arrest or reverse the progression of malignant cells. This includes prophylactic treatment of those at risk of developing precancers and cancers. Also encompassed by this definition is the prevention or slowing of restenosis in subjects that have undergone angioplasty or a stent procedure.

[0022] The terms “therapeutically effective” and “pharmacologically effective” are intended to qualify the amount of each agent which will achieve the goal of improvement in disease severity and the frequency of incidence over treatment with the compounds described herein.

Therapeutically effective or pharmacologically effective amounts may readily be determined by those skilled in the art.

[0023] The term “subject” for purposes of treatment includes any human or animal subject who has been diagnosed with, has symptoms of, or is at risk of developing a disorder characterized by unwanted, rapid cell proliferation. Such disorders include, but are not limited to cancers and precancers. For methods of prevention the subject is any human or animal subject. To illustrate, for purposes of prevention, a subject may be a human subject who is at risk of or is genetically predisposed to obtaining a disorder characterized by unwanted, rapid cell proliferation, such as cancer. The subject may be at risk due to exposure to carcinogenic agents, being genetically predisposed to disorders characterized by unwanted, rapid cell proliferation, and so on. Besides being useful for human treatment, the compounds described herein are also useful for veterinary treatment of mammals, including companion animals and farm animals, such as, but not limited to dogs, cats, horses, cows, sheep, and pigs.

[0024] **Dosage and Administration**

[0025] Preliminary animal studies have shown that these compounds can be orally absorbed, can generate average serum concentrations several-fold higher than TGI, and more importantly, incur little toxicity to the animals after daily oral administration for one month (data not shown)

[0026] The compounds of the present invention can be formulated into suitable pharmaceutical preparations such as tablets, capsules, or elixirs for oral administration or in sterile solutions or suspensions for parenteral administration. The therapeutic agents described herein can be formulated into pharmaceutical compositions using techniques and procedures well known in the art.

[0027] The PDK-1/Akt signaling inhibitor described herein is compounded with a physiologically acceptable vehicle, carrier, excipient, binder, preservative, stabilizer, flavor, etc., in a unit dosage form as called for by accepted pharmaceutical practice. The amount of active substance in those compositions or preparations is such that a suitable dosage in the range indicated is obtained. In some embodiments, the dosage may be between 0.1 to 1000 mg of the PDK-1/Akt signaling inhibitor. In some embodiments, the compositions can be formulated in a unit dosage form, each dosage containing from 1 to 500 mg. In other embodiments, the dosage may be from 10 to 100 mg of the active ingredient. The term “unit dosage form” refers to physically discrete units suitable as unitary dosages for human subjects and other mammals,

each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect, in association with a suitable pharmaceutical excipient. The dosage may depend on many factors, such as the age and size of the subject, the condition being treated, the severity of the condition, and other factors known to those skilled in the art. Taking those factors into account, dosages can be determined by those skilled in the art.

[0028] To prepare compositions, one or more of the therapeutic agents employed in the methods of the invention are mixed with a suitable pharmaceutically acceptable carrier. Upon mixing or addition of the therapeutic agent(s), the resulting mixture may be a solution, suspension, emulsion, or the like. These may be prepared according to methods known to those skilled in the art. The form of the resulting mixture depends upon a number of factors, including the intended mode of administration and the solubility of the compound in the selected carrier or vehicle. The effective concentration is sufficient for inducing apoptosis in undesired cells, such as cancer cells, and may be empirically determined.

[0029] Pharmaceutical carriers or vehicles suitable for administration of the present therapeutic agents include any such carriers suitable for the particular mode of administration. In addition, the active materials can also be mixed with other active materials that do not impair the desired action, or with materials that supplement the desired action, or have another action. The present therapeutic agents may be formulated as the sole pharmaceutically active ingredient in the composition or may be combined with other active ingredients. Derivatives of the present therapeutic agents, such as salts or prodrugs, may also be used in formulating effective pharmaceutical compositions.

[0030] The present therapeutic agents may be prepared with carriers that protect them against rapid elimination from the body, such as time-release formulations or coatings. Such carriers include controlled release formulations, such as, but not limited to, microencapsulated delivery systems. The active compound can be included in the pharmaceutically acceptable carrier in an amount sufficient to exert a therapeutically useful effect in the absence of undesirable side effects on the patient treated.

[0031] The active ingredient may be administered at once, or may be divided into a number of smaller doses to be administered at intervals of time. It is understood that the precise dosage and duration of treatment is a function of the disease being treated and may be determined empirically using known testing protocols or by extrapolation from in vivo or in vitro test data.

It is to be noted that concentrations and dosage values may also vary with the severity of the condition to be alleviated. It is to be further understood that for any particular subject, specific dosage regimens should be adjusted over time according to the individual need and the professional judgment of the person administering or supervising the administration of the compositions, and that the concentration ranges set forth herein are exemplary only and are not intended to limit the scope or practice of the claimed compositions.

[0032] Oral compositions will generally include an inert diluent or an edible carrier and may be compressed into tablets or enclosed in gelatin capsules. For the purpose of oral therapeutic administration, the active compound or compounds can be incorporated with excipients and used in the form of tablets, capsules, or troches. Pharmaceutically compatible binding agents and adjuvant materials can be included as part of the composition.

[0033] The tablets, pills, capsules, troches, and the like can contain any of the following ingredients or compounds of a similar nature: a binder such as, but not limited to, gum tragacanth, acacia, corn starch, or gelatin; an excipient such as microcrystalline cellulose, starch, or lactose; a disintegrating agent such as, but not limited to, alginic acid and corn starch; a lubricant such as, but not limited to, magnesium stearate; a glidant, such as, but not limited to, colloidal silicon dioxide; a sweetening agent such as sucrose or saccharin; and a flavoring agent such as peppermint, methyl salicylate, or fruit flavoring.

[0034] The compounds of formulae I - V may trigger cell death by a number of different mechanisms, however, in most embodiments, the compounds of formulae I - V are able to induce apoptosis in unwanted, proliferative cells. The term "apoptosis" refers to the process of programmed cell death. In every person hundreds of thousands of old or damaged cells die each day by the process of apoptosis and are replaced in the ebb and flow of maintaining a constant number of living cells in the body. Old and damaged cells die in response to a signal triggered on the cell surface for the targeted cell to self destruct. Apoptosis is distinguished from other mechanisms of cell death, such as necrosis, which results in inflammation including swelling, redness, pain and tenderness. Apoptosis does not stimulate such reactions. In apoptosis, the cells shrivel up, break into pieces and the contents are quietly removed by methods that do not induce inflammation. For these reasons, it is highly desirable to induce apoptosis, rather than necrosis, in rapidly proliferating cells, such as cancer cells. However, mutations in some cancer cells confer resistance of these cells to apoptosis. The compounds of formulae I - V have been found to induce apoptosis even in cancer cells which, because of mutations, are otherwise

resistant to apoptosis. Apoptosis can be distinguished from other treatment mechanisms by methods such as microscopy, which are known in the art.

[0035] The terms “proliferative cells,” “proliferating cells,” “rapidly proliferating cells,” “undesirable proliferating cells,” “undesirable rapidly proliferating cells,” “unwanted rapidly proliferating cells,” and the like, refer to cancer cells, precancer cells, and other unwanted, rapidly dividing cells in a subject.

MATERIALS AND METHODS

[0036] **Materials** 4-[5-(4-methylphenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl] benzenesulfonamide was extracted from capsules obtained from Amerisource Health (Malvern, PA) with ethyl acetate followed by recrystallization from a mixture of ethyl acetate and hexane. The Cell Death Detection ELISA kit was purchased from Roche Diagnostics (Mannheim, Germany). Rabbit polyclonal antibodies against Akt and phospho-⁴⁷³ Ser Akt were obtained from Cell Signaling Technologies (Beverly, MA). Mouse monoclonal anti-poly(ADPribose) polymerase (PARP) antibody was provided by Pharmingen (Sari Diego, CA). The PDK-1 kinase assay kit was purchased from Upstate (Lake Placid, NY). Other chemical and biochemicals were obtained from Sigma-Aldrich (St. Louis, MO) unless otherwise mentioned. Nuclear magnetic resonance spectra (¹H NMR) were measured on Bruker 250 MHz. Chemical shifts (δ) are reported in parts per million (ppm) relative to TMS peak with CDCl₃ as solvent unless otherwise mentioned. High-resolution electrospray ionization mass spectrometry analyses were performed with a 3-Tesla Finnigan FTMS-2000 Fourier Transform mass spectrometer.

Synthesis of Chemicals

[0037] The compounds listed in Table 1 were prepared and tested as indicated below. The chemical names, proton nuclear magnetic resonance (¹H NMR) and high-resolution mass spectrometry (HRMS) data are summarized below. The procedures used to synthesize compounds 1 - 36 are described in the Examples, below.

Table 1 Nomenclatures, ¹H NMR (proton nuclear magnetic resonance), and HRMS (high resolution mass spectrometry) characterizations of compounds 1 - 36.

Compound	Description
1	4-[5-(4-(2-bromoethyl)phenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide
	¹ H-NMR δ 3.16 (t, <i>J</i> = 6.4, 2.0Hz, 2H), 3.60 (t, <i>J</i> = 6.4, 2.0 Hz, 2H), 4.90 (s, 2H), 6.75 (s, 1H), 7.13 (d, <i>J</i> = 8.0Hz, 2H), 7.20 (d, <i>J</i> = 8.0 Hz, 2H), 7.47 (d, <i>J</i> = 8.5Hz, 2H), 7.91 (d, <i>J</i> = 8.5Hz, 2H)
	C ₁₈ H ₁₅ BrF ₃ N ₃ O ₂ S; HRMS (M + Na ⁺): theoretical mass, 495.9913; actual mass, 495.9943
2	4-[5-(4-(3-bromopropyl)phenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide
	¹ H-NMR δ 2.16 (m, 2H), 2.81 (t, <i>J</i> = 7.1 Hz, 2H), 3.41 (t, <i>J</i> = 6.4 Hz, 2H), 5.08 (s, 2H), 6.76 (s, 1H), 7.15 (d, <i>J</i> = 8.2 Hz, 2H), 7.25 (d, <i>J</i> = 8.2 Hz, 2H), 7.47 (d, <i>J</i> = 8.5 Hz, 2H), 7.90 (d, <i>J</i> = 8.5 Hz, 2H)
	C ₁₉ H ₁₇ BrF ₃ N ₃ O ₂ S; HRMS (M + Na ⁺): theoretical mass, 510.0069; actual mass, 510.0042
3	4-[5-(4-(2-azidoethyl)phenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide
	¹ H-NMR δ 2.90 (t, <i>J</i> = 6.8 Hz, 2H), 3.51 (t, <i>J</i> = 6.8 Hz, 2H), 5.49 (s, 2H), 6.76 (s, 1H), 7.17 (d, <i>J</i> = 8.3 Hz, 2H), 7.24 (d, <i>J</i> = 8.3 Hz, 2H), 7.42 (d, <i>J</i> = 8.7 Hz, 2H), 7.85 (d, <i>J</i> = 8.7, 2.0 Hz, 2H)
	C ₁₈ H ₁₅ F ₃ N ₆ O ₂ S; HRMS (M + Na ⁺): theoretical mass, 459.0821; actual mass, 459.0817
4	4-[5-(4-(3-azidopropyl)phenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide
	¹ H-NMR δ 1.83 (m, 2H), 2.64 (t, <i>J</i> = 7.5 Hz, 2H), 3.20 (t, <i>J</i> = 7.5 Hz, 2H), 5.31 (br s, 2H), 6.67 (s, 1H), 7.07 (m, 4H), 7.35 (dd, <i>J</i> = 7.5, 2.0 Hz, 2H), 7.79 (d, <i>J</i> = 7.5, 2.0 Hz, 2H)
	C ₁₉ H ₁₇ F ₃ N ₆ O ₂ S; HRMS (M + Na ⁺): theoretical mass, 473.0978; actual mass, 473.0946
5	4-[5-(4-butylphenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide
	¹ H-NMR δ 0.93 (t, <i>J</i> = 7.2 Hz, 3H), 1.36 (m, 2H), 1.64 (m, 2H), 2.63 (t, <i>J</i> = 7.6 Hz, 2H), 5.54 *sm 2H), 6.76 (s, 1H), 7.15 (d, <i>J</i> = 8.3 Hz, 2H), 7.20 (d, <i>J</i> = 8.3 Hz, 2H), 7.45 (dt, <i>J</i> = 8.8, 2.0 Hz, 2H), 7.88 (dt, <i>J</i> = 8.8, 2.0 Hz, 2H)
	C ₂₀ H ₂₀ F ₃ N ₃ O ₂ S; HRMS (M + Na ⁺): theoretical mass, 446.1120; actual mass, 446.1149
6	4-[5-(4- <i>t</i> -butylphenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide
	¹ H-NMR δ 1.33 (s, 9H), 4.90 (s, 2H), 6.53 (s, 1H), 7.32 (dd, <i>J</i> = 9.7 Hz, 4H), 7.42 (d, <i>J</i> = 8.8 Hz, 2H), 8.02 (d, <i>J</i> = 8.8 Hz, 2H)
	C ₂₀ H ₂₀ F ₃ N ₃ O ₂ S; HRMS (M + Na ⁺): theoretical mass, 446.1120; actual mass, 446.1118
7	4-[5-(2-naphthalenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide
	¹ H-NMR δ 5.47 (s, 2H), 6.89 (s, 1H), 7.18 (dd, <i>J</i> = 8.6, 1.6 Hz, 1H), 7.42 (bd, <i>J</i> = 8.6 Hz, 2H), 7.51-7.55 (m, 2H), 7.78-7.83 (m, 6H)
	C ₂₀ H ₁₄ F ₃ N ₃ O ₂ S; HRMS (M + Na ⁺): theoretical mass, 440.0651; actual mass, 440.0657
8	4-[5-(3-indolyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide
	¹ H-NMR δ (acetone-d ₆) 6.69 (br s, 1H), 7.03-7.08 (m, 2H), 7.19 (t, <i>J</i> = 7.2 Hz, 1H), 7.40 (d, <i>J</i> = 7.8 Hz, 1H), 7.50 (d, <i>J</i> = 7.8 Hz, 1H), 7.67 (d, <i>J</i> = 8.7 Hz, 2H), 7.92 (d, <i>J</i> = 8.7 Hz, 2H)
	C ₁₈ H ₁₃ F ₃ N ₄ O ₂ S; HRMS (M + Na ⁺): theoretical mass, 429.0603; actual mass, 429.0606
9	4-[5-(4-biphenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide
	¹ H-NMR δ 4.81 (s, 2H), 6.75 (s, 1H), 7.23 (d, <i>J</i> = 8.5 Hz, 2H), 7.34-7.56 (m, 5H), 7.56 (m, 4H), 7.86 (d, <i>J</i> = 8.5 Hz, 2H)
	C ₂₂ H ₁₆ F ₃ N ₃ O ₂ S; HRMS (M + Na ⁺): theoretical mass, 466.0807; actual mass, 466.0811

Compound	Description
10	4-[5-(4'-chloro[1,1'-biphenyl]-4-yl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide
	¹ H-NMR δ 6.42 (s, 2H), 6.83 (s, 1H), 7.30 (d, <i>J</i> = 8.2 Hz, 2H), 7.40-7.59 (m, 8H), 7.92 (d, <i>J</i> = 8.2 Hz, 2H)
	C ₂₂ H ₁₅ ClF ₃ N ₃ O ₂ S; HRMS (M + Na ⁺): theoretical mass, 500.0418; actual mass, 500.0432
11	4-[5-(3',5'-dichloro[1,1'-biphenyl]-4-yl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide
	¹ H-NMR δ 4.85 (s, 2H), 6.82 (s, 1H), 7.30 (d, <i>J</i> = 8.8 Hz, 2H), 7.36 (s, 1H), 7.37-7.57 (m, 6H), 7.93 (d, <i>J</i> = 8.8 Hz, 2H)
	C ₂₂ H ₁₄ Cl ₂ F ₃ N ₃ O ₂ S; HRMS (M + Na ⁺): theoretical mass, 534.0028; actual mass, 534.0016
12	4-[5-(2',3'-dichloro[1,1'-biphenyl]-4-yl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide
	¹ H-NMR δ 4.85 (s, 2H), 6.76 (s, 1H), 7.18-7.25 (m, 3H), 7.35-7.49 (m, 6H), 7.88 (d, <i>J</i> = 8.6 Hz, 2H)
	C ₂₂ H ₁₄ Cl ₂ F ₃ N ₃ O ₂ S; HRMS (M + Na ⁺): theoretical mass, 534.0028; actual mass, 533.9999
13	4-[5-(2',4',5'-trichloro[1,1'-biphenyl]-4-yl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide
	¹ H-NMR δ 4.86 (s, 2H), 6.77 (s, 1H), 7.25 (dt, <i>J</i> = 8.6, 2.0 Hz, 2H), 7.37 (dt, <i>J</i> = 8.6, s.0 Hz, 2H), 7.39 (s, 1H), 7.46 (dt, <i>J</i> = 8.8, 2.0 Hz, 2H), 7.54 (s, 1H), 7.88 (dt, <i>J</i> = 8.9, 1.2 Hz, 2H)
	C ₂₂ H ₁₃ Cl ₃ F ₃ N ₃ O ₂ S; HRMS (M + Na ⁺): theoretical mass, 567.9638; actual mass, 567.9679
14	4-[5-(4'-methyl[1,1'-biphenyl]-4-yl)-3-(trifluoromethyl)-1H-1-yl]benzenesulfonamide
	¹ H-NMR δ 2.32 (s, 3H), 4.57 (s, 2H), 6.72 (s, 1H), 7.18-7.21 (m, 4H), 7.39-7.52 (m, 6H), 7.84 (d, <i>J</i> = 8.9 Hz, 2H)
	C ₂₃ H ₁₈ F ₃ N ₃ O ₂ S; HRMS (M + Na ⁺): theoretical mass, 480.0964; actual mass, 480.0961
15	4-[5-(4'-trifluoromethyl[1,1'-biphenyl]-4-yl)-3-(trifluoromethyl)-1H-1-yl]benzenesulfonamide
	¹ H-NMR δ 5.19 (s, 2H), 6.86 (s, 1H), 7.36 (d, <i>J</i> = 8.0 Hz, 2H), 7.53 (d, <i>J</i> = 8.5 Hz, 2H), 7.65 (m, 6H), 7.92 (d, <i>J</i> = 8.5 Hz, 2H)
	C ₂₃ H ₁₅ F ₆ N ₃ O ₂ S; HRMS (M + Na ⁺): theoretical mass, 534.0681; actual mass, 534.0677
16	4-[5-(4'-bromomethyl[1,1'-biphenyl]-4-yl)-3-(trifluoromethyl)-1H-1-yl]benzenesulfonamide
	¹ H-NMR δ 3.92 (s, 2H), 4.93 (s, 2H), 6.66 (s, 1H), 7.03-7.26 (m, 8H), 7.38 (d, <i>J</i> = 8.6 Hz, 2H), 7.82 (d, <i>J</i> = 8.6 Hz, 2H)
	C ₂₃ H ₁₇ BrF ₃ N ₃ O ₂ S; HRMS (M + Na ⁺): theoretical mass, 558.0069; actual mass, 558.0112
17	4-[5-(3',5'-dimethyl[1,1'-biphenyl]-4-yl)-3-(trifluoromethyl)-1H-1-yl]benzenesulfonamide
	¹ H-NMR δ 2.40 (s, 6H), 5.38 (br s, 2H), 6.83 (s, 1H), 7.05 (s, 1H), 7.25 (m, 4H), 7.50 (dd, <i>J</i> = 6.7, 1.7 Hz, 2H), 7.59 (dd, <i>J</i> = 6.7, 1.7 Hz, 2H), 7.92 (dd, <i>J</i> = 6.7, 1.7 Hz, 2H)
	C ₂₄ H ₂₀ F ₃ N ₃ O ₂ S; HRMS (M + Na ⁺): theoretical mass, 494.1120; actual mass, 494.1119
18	4-[5-(4'-butyl[1,1'-biphenyl]-4-yl)-3-(trifluoromethyl)-1H-1-yl]benzenesulfonamide
	¹ H-NMR δ 0.96 (t, <i>J</i> = 7.5 Hz, 3H), 1.41 (m, 2H), 1.66 9m, wH), 2.68 (t, <i>J</i> = 7.5 Hz, 2H), 5.20 (br s, 2H), 6.84 (s, 1H), 7.29 (dd, <i>J</i> = 8.2, 2.0 Hz, 4H), 7.53 (dt, <i>J</i> = 8.2, 2.0 Hz, 4H), 7.62 (d, <i>J</i> = 8.5 Hz, 2H), 7.93 (d, <i>J</i> = 8.5 Hz, 2H)
	C ₂₆ H ₂₄ F ₃ N ₃ O ₂ S; HRMS (M + Na ⁺): theoretical mass, 522.1433; actual mass, 522.1466
19	4-[5-(4'-tert-butyl[1,1'-biphenyl]-4-yl)-3-(trifluoromethyl)-1H-1-yl]benzenesulfonamide
	¹ H-NMR δ 1.35 (s, 9H), 4.87 (s, 2H), 6.59 (s, 1H), 7.44-7.57 (m, 6H), 7.58 (d, <i>J</i> = 7.5 Hz, 2H), 7.92 (d, <i>J</i> = 8.7 Hz, 2H), 8.12 (d, <i>J</i> = 7.5 Hz, 2H)
	C ₂₆ H ₂₄ F ₃ N ₃ O ₂ S; HRMS (M + Na ⁺): theoretical mass, 522.1433; actual mass, 522.1401
20	4-[5-(4-(phenylmethyl)phenyl)-3-(trifluoromethyl)-1H-1-yl]benzenesulfonamide
	¹ H-NMR δ 3.71 (s, 2H), 4.74 (s, 2H), 6.52 (s, 1H), 6.91-7.11 (m, 9H), 7.27 (d, <i>J</i> = 8.9 Hz, 2H), 7.69 (d, <i>J</i> = 8.9 Hz, 2H)
	C ₂₃ H ₁₈ F ₃ N ₃ O ₂ S; HRMS (M + Na ⁺): theoretical mass, 480.0964; actual mass, 580.0938

Compound	Description
21	4-[5-(9H-fluoren-2-yl)-3-(trifluoromethyl)-1H-1-yl]benzenesulfonamide
	¹ H-NMR δ 3.88 (s, 2H), 4.64 (s, 2H), 6.68 (s, 1H), 7.26-7.38 (m, 4H), 7.56 (d, <i>J</i> = 8.7 Hz, 2H), 7.74-7.81 (m, 3H), 7.90 (d, <i>J</i> = 8.7 Hz, 2H)
	C ₂₃ H ₁₆ F ₃ N ₃ O ₂ S; HRMS (M + Na ⁺): theoretical mass, 478.0807; actual mass, 478.0771
22	4-[5-(9-anthracenyl)-3-(trifluoromethyl)-1H-1-yl]benzenesulfonamide
	¹ H-NMR δ 4.63 (s, 2H), 6.93 (s, 1H), 7.33 (d, <i>J</i> = 6.8 Hz, 2H), 7.45-7.55 (m, 8H), 8.04 (d, <i>J</i> = 6.8 Hz, 2H), 8.60 (s, 1H)
	C ₂₄ H ₁₆ F ₃ N ₃ O ₂ S; HRMS (M + Na ⁺): theoretical mass, 490.0807; actual mass 490.0769
23	4-[5-(2-phenanthrenyl)-3-(trifluoromethyl)-1H-1-yl]benzenesulfonamide
	¹ H-NMR δ (600 MHz) 4.89 (s, 2H), 6.92 (s, 1H), 7.37 (d, <i>J</i> = 8.5, 1.4 Hz, 1H), 7.51 (d, <i>J</i> = 8.6 Hz, 2H), 7.54-7.69 (m, 3H), 7.80 (d, <i>J</i> = 8.8 Hz, 1H), 7.86-7.92 (m, 4H), 8.64 (d, <i>J</i> = 8.4 Hz, 2H)
	C ₂₄ H ₁₆ F ₃ N ₃ O ₂ S; HRMS (M + Na ⁺): theoretical mass, 490.0807; actual mass, 490.0805
24	4-[5-(9-phenanthrenyl)-3-(trifluoromethyl)-1H-1-yl]benzenesulfonamide
	¹ H-NMR δ 4.76 (s, 2H), 6.90 (s, 1H), 7.43-7.84 (m, 11H), 8.72 (t, <i>J</i> = 7.8 Hz, 2H)
	C ₂₄ H ₁₆ F ₃ N ₃ O ₂ S; HRMS (M + Na ⁺): theoretical mass, 490.0807; actual mass, 490.0833
25	4-[5-(2-phenanthrenyl)-3-(trifluoromethyl)-1H-1-yl]benzenecarboxamide
	¹ H-NMR δ 5.75-6.05 (br d, 2H), 7.0 (s, 1H), 7.50 (dd, <i>J</i> = 8.5, 1.4 Hz, 1H), 7.55 (d, <i>J</i> = 8.5 Hz, 2H), 7.77 (m, 3H), 7.88 (m, 3H), 7.90 (m, 2H), 8.72 (m, 2H)
	C ₂₅ H ₁₆ F ₃ N ₃ O; HRMS (M + Na ⁺): theoretical mass, 454.0038; actual mass, 454.1142
26	4-[5-(2-phenanthrenyl)-3-(trifluoromethyl)-1H-1-yl]benzonitrile
	¹ H-NMR δ 6.91 (s, 1H), 7.46 (s, 1H), 7.50 (d, <i>J</i> = 2.0 Hz, 2H), 7.63-7.79 (m, 5H), 7.83 (d, <i>J</i> = 2.0 Hz, 2H), 7.92 (m, 1H), 8.64 (d, <i>J</i> = 8.4 Hz, 2H)
	C ₂₅ H ₁₄ F ₃ N ₃ O; HRMS (M + Na ⁺): theoretical mass, 436.1032; actual mass, 436.1032
27	4-[5-(2-phenanthrenyl)-3-(trifluoromethyl)-1H-1-yl]-N-hydroxy-benzimidine
	¹ H-NMR δ 7.10 (s, 1H), 7.34 (dd, <i>J</i> = 4.0, 0.9 Hz, 1H), 7.36 (d, <i>J</i> = 0.9 Hz, 1H), 7.37 (d, <i>J</i> = 0.9 Hz, 1H), 7.42-7.45 (m, 3H), 7.46 (d, <i>J</i> = 0.8 Hz, 1H), 7.51-7.52 (m, 2H), 7.53 (d, <i>J</i> = 0.9 Hz, 1H), 7.57 (s, 1H), 7.89 (s, 1H), 7.91 (s, 1H)
	C ₂₅ H ₁₇ F ₃ N ₃ O; HRMS (M + Na ⁺): theoretical mass, 469.1220; actual mass, 469.1247
28	5-(2-phenanthrenyl)-3-(trifluoromethyl)-4-(1H-1-tetrazol-5-ylphenyl)-1H-pyrazole
	¹ H-NMR δ 6.82 (s, 1H), 7.28 (d, <i>J</i> = 1.8 Hz, 1H), 7.38 (d, <i>J</i> = 8.7 Hz, 2H), 7.48-7.74 (m, 5H), 7.74 (d, <i>J</i> = 2.5 Hz, 2H), 7.95 (d, <i>J</i> = 8.7 Hz, 2H), 8.47 (d, <i>J</i> = 8.7 Hz, 2H)
	C ₂₅ H ₁₅ F ₃ N ₆ ; HRMS (M + Na ⁺): theoretical mass, 479.1202; actual mass, 479.1225
29	4-[5-(2-phenanthrenyl)-3-(trifluoromethyl)-1H-1-pyrazol-1-yl]-benzaldehyde oxime
	¹ H-NMR δ 6.81 (s, 1H), 7.27-7.30 (m, 3H), 7.47 (d, <i>J</i> = 8.7 Hz, 2H), 7.52-7.57 (m, 4H), 7.68 (d, <i>J</i> = 8.8 Hz, 2H), 7.75-7.79 (m, 2H), 8.48-8.53 (m, 2H)
	C ₂₅ H ₁₆ F ₃ N ₃ O; HRMS (M + Na ⁺): theoretical mass, 454.1137; actual mass, 454.1106
30	4-[5-(2-phenanthrenyl)-3-(trifluoromethyl)-1H-1-pyrazol-1-yl]-benzaldehyde hydrazone
	¹ H-NMR δ 6.81 (s, 1H), 7.27-7.30 (m, 2H), 7.33 (d, <i>J</i> = 1.8 Hz, 1H), 7.42 (d, <i>J</i> = 8.6 Hz, 1H), 7.53-7.55 (m, 2H), 7.57-7.60 (m, 2H), 7.68 (d, <i>J</i> = 8.9 Hz, 2H), 7.75 (d, <i>J</i> = 1.7 Hz, 1H), 7.80 (s, 1H), 8.48-8.55 (m, 2H)
	C ₂₅ H ₁₇ F ₃ N ₄ ; HRMS (M + Na ⁺): theoretical mass, 453.1297; actual mass, 453.1302
31	4-[5-(2-phenanthrenyl)-3-(trifluoromethyl)-1H-1-pyrazol-1-yl]-phenyl}-acetonitrile
	¹ H-NMR δ 3.77 (s, 2h), 6.93 (s, 1h), 7.29-7.43 (m, 4h), 7.66-7.86 (m, 6h), 8.65 (t, <i>J</i> = 7.0 Hz, 3H)
	C ₂₆ H ₁₆ F ₃ N ₃ ; HRMS (M + Na ⁺): theoretical mass, 450.1151; actual mass, 450.1188

Compound	Description
32	2-{4-[5-(2-phenanthrenyl)-3-(trifluoromethyl)-1 <i>H</i> -pyrazol-1-yl]-phenyl}- <i>N</i> -hydroxy-acetamidine
	¹ H-NMR δ 3.30 (s, 1H), 3.38 (s, 1H), 6.83 (s, 1H), 7.20-7.41 (m, 4H), 7.59-7.89 (m, 6H), 8.55-8.60 (m, 3H)
	C ₂₆ H ₁₉ F ₃ N ₄ O; HRMS (M + Na ⁺): theoretical mass, 461.1580; actual mass, 461.1584
33	5-(2-phenanthrenyl)-3-(trifluoromethyl)-4-(1 <i>H</i> -tetrazol-5-ylmethylphenyl)-1 <i>H</i> -pyrazole
	¹ H-NMR δ 4.45 (s, 2H), 7.15 (s, 1H), 7.42 (s, 4H), 7.53 (d, <i>J</i> = 6.9 Hz, 1H), 7.66-7.76 (m, 3H), 7.89 (d, <i>J</i> = 7.2 Hz, 1H), 8.01 (m, 2H), 8.78 (t, <i>J</i> = 6.9 Hz, 2H)
	C ₂₆ H ₁₇ F ₃ N ₆ ; HRMS (M + Na ⁺): theoretical mass, 493.1335; actual mass, 493.1359
34	2-amino- <i>N</i> -{4-[5-(2-phenanthrenyl)-3-(trifluoromethyl)-1 <i>H</i> -pyrazol-1-yl]-phenyl}acetamide
	¹ H-NMR δ 3.48 (s, 2H), 6.92 (s, 1H), 7.35 (d, <i>J</i> = 8.8 Hz, 2H), 7.42 (dd, <i>J</i> = 8.6, 1.7 Hz, 1H), 7.62-7.72 (m, 5H), 7.79 (d, <i>J</i> = 8.8 Hz, 1H), 7.85-7.94 (m, 2H), 8.62 (t, <i>J</i> = 8.5 Hz, 2H), 9.56 (br s 1H)
	C ₂₆ H ₁₉ F ₃ N ₄ O; HRMS (M + Na ⁺): theoretical mass, 483.1403; actual mass, 483.1389
35	4-[5-(2-phenanthrenyl)-3-(trifluoromethyl)-1 <i>H</i> -pyrazol-1-yl]-phenyl-guanidine
	¹ H-NMR δ 6.90 (s, 1H), 7.19 (d, <i>J</i> = 8.7 Hz, 2H), 7.34 (dd, <i>J</i> = 8.7, 2.0 Hz, 1H), 7.39 (d, <i>J</i> = 8.7 Hz, 2H), 7.61-7.67 (m, 3H), 7.79 (d, <i>J</i> = 9.0 Hz, 1H), 7.84-7.91 (m, 3H), 8.62 (d, <i>J</i> = 8.3 Hz, 2H), 9.95 (s, 1H)
	C ₂₅ H ₁₈ F ₃ N ₅ ; HRMS (M + H): theoretical mass, 446.1587 (M+H); actual mass, 446.1596 (M+H)
36	4-[5-(2-phenanthrenyl)-3-(trifluoromethyl)-1 <i>H</i> -pyrazol-1-yl]-phenyl-urea
	¹ H-NMR δ 6.98 (s, 1H), 7.19 (dt, <i>J</i> = 8.9, 2.1 Hz, 2H), 7.34-7.42 (m, 3H), 7.51-7.62 (m, 4H), 7.70 (d, <i>J</i> = 9.0 Hz, 1H), 7.81-7.85 (m, 2H), 8.59-8.64 (m, 2H)
	C ₂₅ H ₁₇ F ₃ N ₄ O; HRMS (M + Na ⁺): theoretical mass, 469.1252; actual mass, 469.1199

[0038] **Cell Culture** PC-3 (p53^{-/-}) human androgen-nonresponsive prostate cancer cells were purchased from the American Type Tissue Collection (Manassas, VA). Cells were cultured in RPMI 1640 medium (Gibco, Grand Island, NY) supplemented with 10% fetal bovine serum (FBS; Gibco) at 37°C in a humidified incubator containing 5% CO₂.

[0039] **Cell viability analysis** The effect of 4-[5-(4-methylphenyl)-3-(trifluoromethyl)-1*H*-pyrazol-1-yl]benzenesulfonamide and its derivatives on PC-3 cell viability was assessed by using the MTT {[3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl-2*H*-tetrazolium bromide]} assay in six replicates. Cells were grown in 10 % FBS-supplemented RPMI 1640 medium in 96-well, flat bottomed plates for 24 h, and were exposed to various concentrations of compounds 1 - 36 dissolved in DMSO (final concentration ≤0.1 %) in 1 % serum-containing RPMI 1640 medium for different time intervals. Controls received DMSO vehicle at a concentration equal to that in drug-treated cells. The medium was removed, replaced by 200 μl of 0.5 mg/ml of MTT in 10 % FBS-containing RPMI-1640 medium, and cells were incubated in the CO₂ incubator at 37°C for 2 h. Supernatants were removed from the wells, and the reduced MTT dye was solubilized in 200 μL/well DMSO. Absorbance at 570 nm was determined on a plate reader.

[0040] **Cell proliferation** PC-3 cells were seeded into six-well plates at 50,000 cells/well in 10 % FBS-containing RPMI 1640 medium. Following a 24 h attachment period, cells were treated in triplicate with the indicated concentration of compounds 1 - 36 or DMSO vehicle in 10 % FBS-containing RPMI-1640 medium. At different time intervals, cells were harvested by trypsinization, and numerated using a Coulter counter model Z1 D/T (Beckman Coulter, Fullerton, CA).

[0041] **Apoptosis analysis** Two methods were used to assess drug-induced apoptotic cell death: detection of DNA fragmentation by the Cell Death Detection ELISA kit (Roche Diagnostics, Mannheim, Germany) and Western blot analysis of poly-(ADP-ribose)polymerase (PARP) cleavage. The ELISA was performed according to the manufacturer's instructions, and is based on the quantitative determination of cytoplasmic histone-associated DNA fragments in the form of mononucleosomes or oligonucleosomes generated after induced apoptotic death. In brief, 4×10^5 PC-3 cells were cultured in a T-25 flask for 24 h before treatment. Cells were treated with the DMSO vehicle or the test agent at the indicated concentrations for 6 - 24 h, collected, and cell lysates equivalent to 2×10^3 PC-3 cells were used in the ELISA. For the PARP cleavage assay, drug-treated cells were collected 4 - 8 h post-treatment, washed with ice-cold PBS, and resuspended in lysis buffer containing 20 mM Tris-HCl, pH 8, 137 mM NaCl, 1 mM CaCl_2 , 10 % glycerol, 1 % Nonidet P-40, 0.5 % deoxycholate, 0.1 % SDS, 100 μM 4-(2-aminoethyl)benzenesulfonyl fluoride, leupeptin at 10 $\mu\text{g/mL}$, and aprotinin at 10 $\mu\text{g/mL}$. Soluble cell lysates were collected after centrifugation at 10,000 g for 5 min. Equivalent amounts of proteins (60-100 μg) from each lysate were resolved in 8 % SDS-polyacrylamide gels. Bands were transferred to nitrocellulose membranes, and analyzed by immunoblotting with anti-PARP antibody.

[0042] **Immunoblotting.** The general procedure for the Western blot analysis of Akt and phospho-Akt is described as follows. Cells were washed in PBS, resuspended in SDS sample buffer sonicated by an ultrasonic sonicator for 5 sec, and boiled for 5 min. After brief centrifugation, equivalent protein concentrations from the soluble fractions were resolved in 10 % SDS-polyacrylamide gels on a Minigel apparatus, and transferred to a nitrocellulose membrane using a semi-dry transfer cell. The transblotted membrane was washed three times with TBS containing 0.05 % Tween 20 (TBST). After blocking with TBST containing 5 % nonfat milk for 60 min., the membrane was incubated with the primary antibody at 1:1,000 dilution in TBST-5 % low fat milk at 4°C for 12 h, and was then washed three times with TBST.

The membrane was probed with goat anti-rabbit IgG-HRP conjugates (1:1,000) for 1 h at room temperature, and was washed with TBST three times. The immunoblots were visualized by enhanced chemiluminescence.


[0043] **PDK-1 kinase activity** This *in vitro* assay was performed using a PDK-1 kinase assay kit (Upstate, Lake Placid, NY) according to the vendor's instructions. This cell-free assay is based on the ability of recombinant PDK-1, in the presence of DMSO vehicle or the test agent, to activate its downstream kinase serum- and glucocorticoid-regulated kinase (SGK.) which, in turn, phosphorylates the Akt/SGK-specific peptide substrate RPRAATF with $[\gamma\text{-}^{32}\text{P}]\text{-ATP}$. The $[\text{}^{32}\text{P}]$ -phosphorylated peptide substrate was then separated from the residual $[\gamma\text{-}^{32}\text{P}]\text{-ATP}$ using P81 phosphocellulose paper and quantitated by a scintillation counter after three washes with 0.75 % phosphoric acid. The reported values represent the means of two independent determinations.

[0044] **Immunoprecipitated Akt kinase assay** Akt immunoprecipitation was carried out according to a modified, published procedure. PC-3 cells were treated with DMSO vehicle or the test agents at the indicated concentrations for 2 h and then lysed at 4°C for 1 h in buffer A containing 50 mM Tris-HCl, pH 7.5, 1% Triton X-100, 1 mM EDTA, 1 mM EGTA, 50 mM sodium fluoride, 10 mM sodium β -glycerophosphate, 0.1 % 2-mercaptoethanol, 0.1 mM phenylmethylsulfonyl fluoride, and 1 $\mu\text{g/mL}$ each of aprotinin, pepstatin, and leupeptin. Cell lysates were centrifuged at 10,000 g for 5 min, and the supernatant was treated with anti-Akt at 4°C for 60 min., followed by protein G-agarose beads for additional 60 min. The immunoprecipitate was used to analyze Akt kinase activity by using the Akt/SGK-specific peptide substrate RPRAATF as described above. Values represented the means of two independent determinations.

















[0045] **Statistical analysis** Each experiment was performed in triplicate, unless otherwise mentioned. All experiments were carried out at least two times on different occasions. Where appropriate, the data are presented as the mean $\pm$ 95 % confidence interval.









[0046] The structure and potency in inhibiting PDK-1 kinase activity and PC-3 cell growth of 24 representative derivatives are summarized in Table 2.

Table 2 Structures and potency for inhibiting recombinant PDK-1 kinase activity and for inducing apoptotic death in PC-3 cells for 4-[5-(4-methylphenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl] benzenesulfonamide and



compounds 37-60 (1-24)

Number	Ar	IC ₅₀ (μM)	
		PDK-1	PC-3
Comparative Compound		48	30
37		42	18
38		38	17
39		32	17
40		34	18
41		20	9
42		34	18
43		24	11
44		65	31
45		21	11
46		22	9
47		18	10
48		23	10
49		9	5
50		15	8
51		18	8
52		20	11

Number	Ar	IC ₅₀ (μM)	
		PDK-1	PC-3
53		17	9
54		32	15
55		32	15
56		15	8
57		16	9
58		12	7
59		9	5
60		42	23

[0047] These compounds, except the indole derivative **44**, showed improved PDK-1 inhibitory and anti-proliferative activities vis-à-vis 4-[5-(4-methylphenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl] benzenesulfonamide. Additionally, none of these compounds displayed measurable COX-2 inhibitory activity (data not shown). A general increase in PDK-1 inhibitory activity was noted with increasing bulkiness of the aromatic ring, *i.e.*, tricyclic aromatic rings (**57-59**) > substituted biphenyl (**45-55**) > substituted phenyl (**37-42**). These data suggested that the aromatic system bound to a large, hydrophobic region of the enzyme pocket. Among the 24 analogues examined, compound **59** represented the optimal derivative with IC₅₀ values of 9 μM and 5 μM for inhibiting PDK-1 activity and PC-3 cell viability, respectively, as reported in Table 2. These IC₅₀ values corresponded to a five- to six-fold improvement over the activities of 4-[5-(4-methylphenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl] benzenesulfonamide (48 μM and 30 μM, respectively). However, compound **60** exhibited a decrease, compared with compound **59**

in PDK-1 inhibitory activity, which might be attributable to steric hindrance imposed by an unfavorable orientation of the tricyclic aromatic ring.

[0048] There existed a correlation between PDK-1 and PC-3 growth inhibition potency in all compounds examined, suggesting the mechanistic relevance of PDK-1 inhibition to the antiproliferative effect. Overall, the IC₅₀ value for inhibiting PC-3 cell proliferation was approximately one half of that of PDK-1 inhibition. This discrepancy might arise from a mechanistic synergy between PDK-1 inhibition and concomitant Akt dephosphorylation by protein phosphatase 2A (PP2A) in Aug-treated cells, resulting in augmented Akt deactivation. To examine this premise, PC-3 cells were treated with different concentrations of compound 59 for 2 h, and the consequent effect on AM was assessed by two independent assays: immunoprecipitated Akt kinase activity and Akt phosphorylation status. Both assays gave consistent results.

[0049] According to the kinase assay, the IC₅₀ of compound 59 for inhibiting intracellular Akt activation was 5 μM. Neither compound 59 nor other the other compounds displayed a direct inhibitory effect on immunoprecipitated Akt activity. Meanwhile, Western blot analysis shows that treatment of PC-3 cells with compound 59 at 5 μM and above led to significant Akt dephosphorylation.


[0050] The inhibition of PDK-1/Akt signaling led to apoptotic death in PC-3 cells in 1 % FBS-containing RPMI 1640 medium in a dose-dependent manner, as was evidenced by DNA fragmentation and PARP cleavage. The dose of compound 59 required to induce 50 % PC-3 cell death at 24 h was 5 μM. The IC₅₀ values for compound 59 to induce PC-3 cell death was consistent with that of inhibiting Akt activation in drug-treated cells. Furthermore, the effect of compound 59 on PC-3 cell proliferation was examined in 10 % FBS-supplemented RPMI 1640 medium. Compound 59 at 1 μM showed substantial anti-proliferative activity. Together, these data clearly indicated the *in vitro* efficacy of compound 59 in PC-3 growth inhibition.

[0051] The modeling showed that compound 59 was docked into the ATP-binding domain that is located within a deep cleft between the two lobes of PDK-1. Although compound 59 competed with ATP for binding, the mode of binding for compound 59 was found to be somewhat different from that of ATP. While the benzenesulfonamide moiety occupied the adenine-binding motif, the planar pyrazole moiety was perpendicular to the ribose ring. This arrangement positioned the adjacent phenanthrene ring behind the trisphosphate-binding pocket.




The phenanthrene ring formed hydrophobic interactions with an apolar region formed by residues 88-96 encompassing part of two adjacent β sheets joined by a glycine-rich loop.

[0052] Structures of twelve representative derivatives, their potency against PDK-1, and their ability to cause apoptotic death in PC-3 cells are summarized in Table 3.

Table 3 Structures and potency for inhibiting recombinant PDK-1 kinase activity and for inducing apoptotic death in PC-3 cells for compounds 25 - 36. The general structures of these compounds is shown at top.



Number	R	IC ₅₀ (μM)	
		PDK-1	PC-3
61	-CONH ₂	12	7
62	-CN	45	30
63		40	25
64		52	32
65		25	14
66	-CH=N-NH ₂	16	10
67	-CH ₂ CN	42	25
68		15	8
69		45	27

Number	R	IC ₅₀ (μM)	
		PDK-1	PC-3
70		5	5
71		2	3
72		40	24

[0053] Among these derivatives, compound **70** and **71** exhibited IC₅₀ values for PDK-1 inhibition of 5 μM and 2 μM, respectively, which represented two- and five-fold increases in potency over compound **59**. Compounds **70** and **71** contained side chains of 2-aminoacetamide (-NHC(O)CH₂NH₂) and guanidine (-NHC(=NH)NH₂), respectively. Like compound **59**, they exhibited no appreciable direct inhibition on immunoprecipitated Akt kinase activity, nor was any measurable COX-2 inhibitory activity detected at concentrations up to 50 μM. Exposure of PC-3 cells to either agent, even at 1 μM, resulted in a substantial decrease in the phospho-Akt level. This improvement in potency reflected a strengthening of the hydrogen bonding in the protein-ligand interactions for these derivatives. This premise was supported by the modeled docking of compound **71** into the ATP-binding site. The guanidino group of compound **71** resembled the partial structure of ATP's purine ring, which allowed the formation of hydrogen bonds with Ser160 and Ala162 as depicted by the docking model.

[0054] **Cellular effects of PDK-1/Akt signaling inhibitors** Both compounds **70** and **71** induced apoptotic death in PC-3 cells in 1 % FBS-containing medium in a dose-dependent manner, as was demonstrated by DNA fragmentation and PARP cleavage. These agents exhibited higher potency than compound **59** in apoptosis induction at concentrations greater than 2.5 μM. Moreover, these derivatives were submitted to the Developmental Therapeutic Program (DTP) at the National Cancer Institute (NCI) for screening against sixty human tumor cell lines, representing leukemia, melanoma, and cancers of the lung, colon, brain, ovary, breast, prostate, and kidney. Dose-response data of one representative cell line from each class of tumor cells after two-day exposure in 5% FBS-containing medium are shown in Fig. 1C, which include: 1, RPMI-8226 leukemia cells; 2, NCI-H322M non-small cell lung cancer cells; 3, HT29 colon

cancer cells; 4, U251 CNS cancer cells; 5, SK-MEL-28 melanoma cancer cells; 6, SK-OV-3 ovarian cancer cells; 7, RXF 393 renal cancer cells; 8, PC-3 prostate cancer cells; 9, MDA-MB-231 breast cancer cells. Many of these cell lines were responsive to the growth inhibitory effect of both agents at concentrations as low as 0.1 μ M

[0055] In the sixty cell line assay, three dose response parameters for each cell line were calculated based on growth inhibition curves. These parameters include G150 (concentration resulting in 50 % growth inhibition), TGI (concentration resulting in total growth inhibition), and LC50 (concentration resulting in a 50 % reduction in the measured protein level at the end of drug treatment as compared to that at the beginning). The means of these parameters among the sixty different cell lines for compounds **70** and **71** after two-day treatment were as follows, respectively, G150: 1.1 and 1.2 μ M; TGI: 3.2 and 2.9 μ M; LC50: 24 and 8.5 μ M. These data clearly demonstrate the *in vitro* efficacy of compounds **70** and **71**. Both agents were able to completely suppress cell growth in a diverse range of tumor cell lines at the 3 - 5 μ M therapeutic range.

[0056] In light of the conserved role of PDK-1/Akt signaling in cancer cell survival and proliferation, this pathway represents a therapeutically relevant target for developing orally bioavailable, small-molecule inhibitors.

[0057] *In silico* docking of compound **59** into the ATP-binding pocket showed that the molecule was anchored into the ATP binding domain, in part, through hydrogen bonding between the sulfonamide and the amide of Ala162. Ala162 has also been reported to play a key role in anchoring other ligands such as ATP¹⁷ and UCN-01 to PDK-1. Together, these data suggest that the sulfonamide moiety of compound **59** might be amenable to alterations for optimizing potency.


[0058] Accordingly, replacement of the sulfonamide function with 2-aminoacetamide (-NHC(O)CH₂NH₂) and guanidine [-NHC(=NH)NH₂] led to compounds **70** and **71**, respectively, both of which exhibited improved PDK-1 inhibition with IC₅₀ values of 5 and 2 μ M, respectively. Docking of compound **71** into the ATP binding site revealed the existence of an additional hydrogen bond between the guanidine moiety and the backbone-oxygen of Ser160, suggesting that the enhancement in potency might be attributable to an increase in hydrogen bonding. The effect of these side chains on ligand binding, however, is subtle, as illustrated by the structure-activity relationship summarized in Table 3.

[0059] The high potency of compounds **70** and **71** in PDK-1 inhibition was reflected in their abilities to effectively block Akt activation and to induce apoptotic cell death in PC-3 cells at low μM concentrations (Fig. 1A, B). More importantly, due to the conserved role of PDK-1/Akt signaling in cell proliferation and survival, these agents were potent in inhibiting cell growth in serum-containing medium in all 60 human tumor cell lines examined, with mean GI50 (50% cell growth inhibition) values of 1.2 μM and 1.3 μM , respectively, and TGI (total growth inhibition) values of 3.2 μM and 2.9 μM , respectively. Our preliminary animal studies have shown that these compounds can be orally absorbed, can generate average serum concentrations several-fold higher than TGI, and more importantly, incur little toxicity to the animals after daily oral administration for one month (data not shown).

[0060] Testing of *in vivo* efficacy against different tumor xenografts in nude mice is currently under way in this laboratory. In addition, toxicological and pharmacological testing of these agents will be undertaken under the Rapid Access to Intervention Development (RAID) program at NCI.


General Synthetic Procedures for Compounds 37-60

[0061] All chemical reagent and organic solvents were purchased from Aldrich (St. Louis, MO) unless otherwise mentioned. Compounds **1** – **24** were synthesized according to a two-step general procedure described in Scheme 1, in which Ar represents the respective aromatic ring structures.



Compound **59** is used here as an example to illustrate the synthesis of the group of compounds (Scheme 2). Other compounds followed the same procedures *via* precursors and the respective intermediates with different aromatic ring structures (compounds **I** and **II**).

Scheme 2



[0062] **EXAMPLE 1 Synthesis of the 1,1,1-Trifluoro-4-hydroxy-4-phenanthren-2-yl-but-3-en-2-one Precursor** (step 1). To a suspension of sodium hydride (NaH; 0.13 g, 5.4 mmol) in 5 mL of anhydrous tetrahydrofuran (THF) was added ethyl trifluoroacetate (CF_3COOEt ; 0.64 g, 4.5 mmol) under argon. After stirring at 25°C for 10 minutes, 2-acetylphenanthrene (1 g, 4.5 mmol) in 5 mL of THF was added dropwise to the solution. The mixture became clear and orange-hued within 30 minutes, and after stirring for an additional 2 hours, was concentrated under vacuum. The residue was suspended in water, and extracted with ethyl acetate (15 mL) twice. The organic phase was separated, dried over sodium sulfate, and concentrated to dryness under vacuum to give the product (yellow solid; 1.29 g, 90% yield). The product was used directly without further purification.

[0063] **EXAMPLE 2 Synthesis of Compound 59** (step 2). 4-Hydrazinobenzene-1-sulfonamide hydrochloride (1.1 g; 4.9 mmol) was added to a stirred solution of 1,1,1-trifluoro-4-hydroxy-4-phenanthren-2-yl-but-3-en-2-one (1.29 g, 4.1 mmol) in 40 mL of ethanol. The mixture was refluxed for 12 hours, cooled to room temperature, and concentrated to dryness under vacuum. The residue was dissolved in ethyl acetate, and washed with water. The organic layer was dried over sodium sulfate, and concentrated under vacuum. The crude product was purified by silica gel flash chromatography to yield **59** (1.52 g, 80% yield).

[0064] **EXAMPLES 3 – 14 Syntheses of Compounds 61-72** Compounds **61-72** were synthesized using 1,1,1-trifluoro-4-hydroxy-4-phenanthren-2-yl-but-3-en-2-one, product of the aforementioned step 1, as a common precursor (Scheme 3, Error! Reference source not found..)

[0065] **EXAMPLE 3 4-[5-(2-Phenanthracenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]-benzenecarboxamide (61)** (step 3). (4-Carbamoylphenyl)-hydrazine hydrochloride (0.92 g, 4.9 mmol) was added to a stirred solution of 1,1,1-trifluoro-4-hydroxy-4-phenanthren-2-yl-but-3-en-2-one (1.29 g, 4.1 mmol) in 40 mL of ethanol at 25°C. The mixture was refluxed for 12 hours, cooled to room temperature and concentrated to dryness under vacuum. The residue

was dissolved in ethyl acetate, and washed with water. The organic layer was dried over sodium sulfate, and concentrated under vacuum. The crude product was purified by silica gel flash chromatography (ethyl acetate-hexane, 1:1), - yielding **61** (1 g, 60 % yield).

[0066] **EXAMPLE 4** **4-[5-(2-Phenanthrenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]-benzonitrile (62)** (step 4). To a stirred solution of 1,1,1-trifluoro-4-hydroxy-4-phenanthren-2-yl-but-3-en-2-one (2.45 g, 7.7 mmol) in 60 mL of ethanol was added 4-cyanophenylhydrazine hydrochloride (2.53 g, 15 mmol) at 25°C. The mixture was stirred under reflux for 12 hours, cooled to room temperature and concentrated to dryness under vacuum. The residue was dissolved in methylene chloride, and washed with water. The organic layer was dried over sodium sulfate, and concentrated under vacuum. The crude product was purified by silica gel flash chromatography (ethyl-acetate-hexane, 1:4) to afford **62** (2.7 g, 85 % yield).

[0067] **EXAMPLE 5** **4-[5-(2-Phenanthrenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]-N-hydroxybenzamidine (63)** (step 5). Hydroxylamine hydrochloride (25 mg, 0.36 mmol) was added to a suspension of Na metal (8.3 mg, 0.36 mmol) in methanol (3 mL). The mixture was stirred at room temperature for 10 minutes, and compound **62** (1224 mg, 0.3 mmol) was added. The mixture was refluxed for 2 hours, then stirred at 25°C for an additional 16 hours, and concentrated under vacuum. The residue was purified by silica gel flash chromatography (ethyl acetate-hexane, 1:4 to 1:1) to give **63** (120 mg, 76 % yield).

[0068] **EXAMPLE 6** **5-(2-Phenanthrenyl)-3-(trifluoromethyl)-4-(1H-tetrazol-5-ylphenyl)-1H-pyrazole (64)** (step 6). A mixture containing compound **62** (125 mg, 0.3 mmol), NH₄Cl (123.7 mg), and NaN₃ (58.5 mg, 0.9 mmol) in 5 mL of 10 % HCl was added, and extracted with 20 mL of methylene chloride, twice. The organic phase was dried over sodium sulfate, and concentrated to dryness under vacuum. The crude product was purified by silica gel flash chromatography (ethyl acetate-hexane 1:4) to give **64** (96 mg, 70 % yield).

[0069] **EXAMPLE 7** **4-5[-(2-Phenanthrenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]-benzaldehyde oxime (65)** (step 7). DIBAL-H (3.1 mL, 3.1 mmol, 1.0 M in hexane) was added dropwise to a solution of compound **62** (0.417 g, 1.1 mmol) in 5 mL THF at -40°C. The mixture was stirred for 8 hours, poured into 5 mL of 10 % acetic acid, and stirred for 30 minutes. The organic layer was dried over sodium sulfate, and concentrated to dryness under vacuum. The crude product was purified by silica gel flash chromatography (ethyl acetate-hexane, 1:4) to give an aldehyde intermediate (141 mg, 0.34 mmol) that was immediately added to a solution

containing hydroxylamide hydrochloride (211 mg) and K_2CO_3 in 5 mL of ethanol. The mixture was stirred under reflux for 16 hours. After removal of solvent, the residue was extracted with CH_2Cl_2 and washed with water.

[0070] **EXAMPLE 8** **4-[5-(2-Phenanthrenyl)-3-(trifluoromethyl)-1*H*-pyrazol-1-yl]-benzaldehyde hydrazone (66)** (step 8). Compound **66** (124 mg, 85 % yield) was synthesized in the same manner as **65** except that hydrazine monohydrate (153 mg, 3.1 mmol) was used instead of hydroxylamine hydrochloride.

[0071] **EXAMPLE 9** **{4-[5-(2-Phenanthrenyl)-3-(trifluoromethyl)-1*H*-pyrazol-1-yl]-phenyl}-acetonitrile (67)** (step 9). (a) Preparation of (4-Hydrazinophenyl)acetonitrile hydrochloride. A solution of sodium nitrite (3.15 g, 45.7 mmol) in water (20 mL) was added dropwise to a cooled ($-15^\circ C$), stirred suspension of 4-aminobenzonitrile (5 g, 42.3 mmol) in a concentrated hydrogen chloride solution (55 mL) at such a rate as to maintain a temperature below $-10^\circ C$. After the addition was finished, the reaction mixture was quickly filtered to remove solids, and the filtrate was added in portions to a cooled ($-20^\circ C$), stirred solution of $SnCl_2 \cdot 2H_2O$ (47.7 g, 0.21 mol) in a concentrated hydrogen chloride solution (37 mL) at such a rate as to keep the temperature below $-10^\circ C$. After stirring the solution for an additional 15 minutes, the solid was collected, washed with diethyl ether (4×25 mL), and dried to give (4-hydrazinophenyl)acetonitrile hydrochloride (5.6 g, 78 %). (b) Compound **67**. A mixture of (4-hydrazinophenyl)acetonitrile hydrochloride (0.32 g, 1 mmol) and 1,1,1-trifluoro-4-hydroxy-4-phenanthren-2-yl-but-3-en-2-one (0.18 g, 1.1 mmol) in ethanol (20 mL) was stirred under reflux for 24 hours, cooled to room temperature, concentrated to dryness under vacuum, and dissolved in ethyl acetate. The organic layer was dried over magnesium sulfate, and concentrated to dryness under vacuum. The crude product was purified by silica gel column chromatography (hexane-ethyl acetate, 2:1) to give compound **67** (0.35 g, 81 % yield).

[0072] **EXAMPLE 10** **2-{4-[5-(2-Phenanthrenyl)-3-(trifluoromethyl)-1*H*-pyrazol-1-yl]-phenyl}-*N*-hydroxy-acetamidine (68)** (step 10). A solution of compound **67** (0.43 g, 1 mmol) and hydroxylamine hydrochloride (0.075 g, 1.1 mmol) in ethanol (10 mL) was stirred under reflux for 8 hours, and concentrated to dryness under vacuum. The residue was dissolved in water, brought to pH 8-9 by addition of saturated $NaHCO_3$ solution, and extracted with ethyl acetate. The organic layer was dried over magnesium sulfate, and concentrated to dryness under vacuum. The crude product was recrystallized in diethyl ether-hexane to give compound **68** (0.32 g, 71 % yield).

[0073] **EXAMPLE 11** **5-(2-Phenanthrenyl)-3-(trifluoromethyl)-4-(1*H*-tetrazol-5-ylmethylphenyl)-1*H*-pyrazole (69)** (step 11). A mixture containing compound **67** (0.43 g, 1 mmol), sodium azide (0.08 g, 1.2 mmol), and triethylamine hydrochloride (0.12 g, 1.2 mmol) in toluene (5 mL) was stirred at 100 °C for 5 hours, cooled to room temperature, and extracted with water (10 mL). To the aqueous phase was added dropwise a 36 % hydrogen chloride solution to salt out the resulting tetrazole **69**. After filtration, the solid was dried under vacuum, yielding compound **33** (0.39 g, 84 % yield).

[0074] **EXAMPLES 12-14 1-(4-Nitrophenyl)-5-phenyl-3-(trifluoromethyl)-1*H*-pyrazole (III)** (step 12). To a solution of 1,1,1-trifluoro-4-hydroxy-4-phenanthren-2-yl-but-3-en-2-one (1.29 g, 4.1 mmol) in 40 mL of ethanol was added 4-nitrophenylhydrazine hydrochloride (0.93 g, 4.9 mmol) under stirring, refluxed for 1 hour, cooled to room temperature, and concentrated to dryness under vacuum. The residue was dissolved in ethyl acetate, and washed with water. The organic phase was dried over magnesium sulfate, and concentrated to dryness under vacuum. The crude product was purified by silica gel column chromatography to afford compound **III** (0.88 g, 50 % yield).

[0075] **4-[5-(2-Phenanthrenyl)-3-(trifluoromethyl)-1*H*-pyrazol-1-yl]phenylamine (IV)** (step 13). To a solution of compound **III** (0.88 g, 2 mmol) in 20 mL ethanol was added platinum oxide (27 mg, 0.12 mmol), stirred under H₂ at 55 psi for 12 hours, filtered to remove the catalyst, and concentrated to dryness under vacuum. The crude product was purified by silica gel chromatography to yield compound **IV** (0.57 g, 70 % yield).

[0076] **EXAMPLE 12 2-Amino-*N*-{4-[5-(2-phenanthrenyl)-3-(trifluoromethyl)-1*H*-pyrazol-1-yl]-phenyl}-acetamide (70)** (steps 14 and 15). To a solution of *t*-butoxycarbonyl (tBOC)-glycine (0.25 g, 1.4 mmol) and compound **IV** (0.57 g, 1.4 mmol) in 10 mL of tetrahydrofuran was added 1-[3-(dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride (0.41 g, 2.1 mmol), stirred at 25 °C for 12 hours, and concentrated to dryness under vacuum in a rotary evaporator. The residue was suspended in water, and the product was extracted with ethyl acetate. The organic phase was dried over magnesium sulfate, and concentrated to dryness under vacuum to give compound **V** (0.67 g, 85 % yield). Compound **V** (0.67 g, 1.2 mmol) was dissolved in 8 mL of ethyl acetate containing 0.7 mL of concentrated HCl solution, stirred at room temperature for 2 hours, and concentrated to dryness under vacuum. The crude product was purified by silica gel column chromatography to yield compound **70** as a white powder (0.49 g, 90 %).

[0077] **EXAMPLE 13 4-[5-(2-Phenanthrenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]-phenyl-guanidine (71)** (step 16). To a solution of compound **IV** (0.57 g, 1.4 mmol) in 7 mL of ethanol was added cyanamide (89 mg, 2.1 mmol) and 1.5 mL of 1N HCl. The mixture was refluxed for 24 hours, and concentrated to dryness under vacuum. The product was purified by silica gel column chromatography to give compound **71** as a white solid (0.25 g, 40 %yield).

[0078] **EXAMPLE 14 4-[5-(2-Phenanthrenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]phenyl urea (72)** (step 17). Into a 250 mL round bottom flask containing acetic acid (50 mL), water (12 mL), and ethanol (20 mL) was added compound **IV** (2.25 g, 5.6 mmol), followed by sodium isocyanate (0.74 g, 11.2 mmol). The reaction was stirred for 1.5 hours, and then neutralized with the addition of 1 N sodium hydroxide followed by sodium hydroxide pellets until the pH had changed to 7.0. The product was separated and then washed with 100 mL of water, dried with magnesium sulfate, and then solvent was removed to obtain the crude product. Purification was performed by silica gel chromatography with (hexane-ethyl acetate, 3:2 to hexane-acetone, 1:3) to afford compound **72**.

[0079] **EXAMPLE 15 Preparation of Additional Compounds** The compounds in Table 4 were prepared using the methods of the Examples above.




Table 4 Additional Compounds

Compound	R	Nomenclature
73		4-(5-Phenanthren-2-yl-3-trifluoromethyl-pyrazol-1-yl)-N-(4-sulfamoyl-phenyl)-benzamide
74		N-[4-(5-Phenanthren-2-yl-3-trifluoromethyl-pyrazol-1-yl)-phenyl]-4-sulfamoyl-benzamide
75		2-Guanidino-N-[4-(5-phenanthren-2-yl-3-trifluoromethyl-pyrazol-1-yl)-phenyl]-acetamide
76		2-[4-(5-Phenanthren-2-yl-3-trifluoromethyl-pyrazol-1-yl)-phenylamino]-acetamide
77		N-{[4-(5-Phenanthren-2-yl-3-trifluoromethyl-pyrazol-1-yl)-phenylcarbamoyl]-methyl}-benzamide

[0080] **EXAMPLE 16 Screening of compounds 70 and 71 against several cancer cell lines**

The human tumor cell lines of the cancer screening panel are grown in RPMI 1640 medium containing 5% fetal bovine serum and 2 mM L-glutamine. For a typical screening experiment, cells are inoculated into 96 well microtiter plates in 100 μ L at plating densities ranging from 5,000 to 40,000 cells/well depending on the doubling time of individual cell lines. After cell inoculation, the microtiter plates are incubated at 37°C, 5 % CO₂, 95 % air and 100 % relative humidity for 24 h prior to addition of experimental drugs.

[0081] After 24 h, two plates of each cell line are fixed *in situ* with TCA, to represent a measurement of the cell population for each cell line at the time of drug addition (Tz). Experimental drugs are solubilized in dimethyl sulfoxide at 400-fold the desired final maximum test concentration and stored frozen prior to use. At the time of drug addition, an aliquot of frozen concentrate is thawed and diluted to twice the desired final maximum test concentration with complete medium containing 50 μ g/ml gentamicin. Additional four, 10-fold or 1/2 log serial dilutions are made to provide a total of five drug concentrations plus control. Aliquots of 100 μ l of these different drug dilutions are added to the appropriate microtiter wells already containing 100 μ l of medium, resulting in the required final drug concentrations.

[0082] Following drug addition, the plates are incubated for an additional 48 h at 37°C, 5 % CO₂, 95 % air, and 100 % relative humidity. For adherent cells, the assay is terminated by the addition of cold TCA. Cells are fixed *in situ* by the gentle addition of 50 μ l of cold 50 % (w/v) TCA (final concentration, 10 % TCA) and incubated for 60 minutes at 4°C. The supernatant is discarded, and the plates are washed five times with tap water and air dried. Sulforhodamine B (SRB) solution (100 μ l) at 0.4 % (w/v) in 1 % acetic acid is added to each well, and plates are incubated for 10 minutes at room temperature. After staining, unbound dye is removed by washing five times with 1 % acetic acid and the plates are air dried. Bound stain is subsequently solubilized with 10 mM trizma base, and the absorbance is read on an automated plate reader at a wavelength of 515 nm. For suspension cells, the methodology is the same except that the assay is terminated by fixing settled cells at the bottom of the wells by gently adding 50 μ l of 80 % TCA (final concentration, 16 % TCA). Using the seven absorbance measurements [time zero, (Tz), control growth, (C), and test growth in the presence of drug at the five concentration levels (Ti)], the percentage growth is calculated at each of the drug concentrations levels. Percentage growth inhibition is calculated as:

$$[(Ti - Tz) / (C - Tz)] \times 100 \text{ for concentrations for which } Ti \geq Tz$$

$$[(Ti-Tz)/Tz] \times 100 \text{ for concentrations for which } Ti < Tz$$

[0083] Three dose response parameters are calculated for each experimental agent. Growth inhibition of 50 % (GI_{50}) is calculated from $[(Ti-Tz)/(C-Tz)] \times 100 = 50$, which is the drug concentration resulting in a 50% reduction in the net protein increase (as measured by SRB staining) in control cells during the drug incubation. The drug concentration resulting in total growth inhibition (TGI) is calculated from $Ti = Tz$. The LC_{50} (concentration of drug resulting in a 50% reduction in the measured protein at the end of the drug treatment as compared to that at the beginning) indicating a net loss of cells following treatment is calculated from $[(Ti-Tz)/Tz] \times 100 = -50$. Values are calculated for each of these three parameters if the level of activity is reached; however, if the effect is not reached or is exceeded, the value for that parameter is expressed as greater or less than the maximum or minimum concentration tested.

[0084] The methods described were used to test compounds **70** and **71** on a panel of sixty cell lines under a screening service provided by the Developmental Therapeutics Program at the National Institutes of Health. Shown in Figure 1C are 1, RPMI-8226 leukemia cells; 2, NCI-H322M non-small cell lung cancer cells; 3, HT29 colon cancer cells; 4, U251 CNS cancer cells; 5, SK-MEL-28 melanoma cancer cells; 6, SK-OV-3 ovarian cancer cells; 7, RXF 393 renal cancer cells; 8, PC-3 prostate cancer cells; 9, MDA-MB-231 breast cancer cells.

[0085] Results of testing compounds **70** and **71** against the full 60 cell lines are shown in the tables below. The testing was done by the National Cancer Institute Developmental Therapeutics Program. The results shown are *in vitro* testing results.

Table 5. Results of testing Compound 70 against 60 cancer cell lines.

Panel/Cell Line	Time Zero	Ctrl	Mean Optical Densities				Log10 Concentration				Percent Growth				GI50	TGI	LC50
			-8.0	-7.0	-6.0	-5.0	-4.0	-3.0	-2.0	-1.0	0.0	1.0	2.0	3.0			
Leukemia																	
CCRF-CEM	0.295	1.099	0.893	0.946	0.523	0.168	0.194	74	81	28	-43	-34	3.87E-07	2.49E-06	>1.00E-04		
K-562	0.297	1.467	1.435	1.022	0.681	0.209	0.247	97	62	33	-30	-17	2.57E-07	3.34E-06	>1.00E-04		
MOLT-4	0.313	1.048	0.964	0.951	0.614	0.180	0.296	89	87	41	-42	-6	6.36E-07	3.10E-06	>1.00E-04		
RPMI-8226	0.332	0.945	0.766	0.701	0.547	0.164	0.229	85	72	42	-51	-31	5.37E-07	2.84E-06			
SR	0.364	0.902	0.816	0.796	0.605	0.265	0.363	84	80	45	-27	.	7.11E-07	4.17E-06	>1.00E-04		
Non-Small Cell Lung Cancer																	
AS49/ATCC	0.363	1.279	1.294	1.149	0.831	0.155	0.252	102	86	51	-57	-31	1.02E-06	2.96E-06			
EKVV	0.557	0.860	0.891	0.860	0.786	0.156	0.193	110	100	75	-72	-65	1.49E-06	3.25E-06	7.08E-06		
HOP-62	0.510	1.395	1.310	1.330	1.075	0.155	0.217	90	93	64	-70	-57	1.27E-06	3.01E-06	7.13E-06		
HOP-92	1.070	1.526	1.509	1.474	1.465	0.095	0.267	96	89	87	-91	-75	1.61E-06	3.07E-06	5.87E-06		
NCI-H226	1.190	1.594	1.593	1.586	1.429	0.505	0.614	100	98	59	-58	-48	1.20E-06	3.21E-06			
NCI-H23	0.457	1.655	1.672	1.673	1.255	0.097	0.156	101	102	67	-79	-66	1.30E-06	2.87E-06	6.33E-06		
NCI-H322M	0.626	2.465	1.363	1.562	1.420	0.100	0.269	40	51	43	-84	-57	2.18E-06	5.40E-06			
NCI-H460	0.319	1.843	1.799	1.499	0.830	0.173	0.213	97	77	34	-46	-33	4.21E-07	2.64E-06	>1.00E-04		
NCI-H522	0.766	1.549	1.626	1.718	1.528	0.260	0.486	110	122	97	-66	-37	1.95E-06	3.94E-06			
Colon Cancer																	
COLO 205	0.221	1.412	1.286	1.252	0.916	0.104	0.239	89	87	58	-53	1	1.19E-06				
HCT-116	0.114	0.708	0.751	0.665	0.452	0.104	0.146	107	93	57	-9	5	1.27E-06		>1.00E-04		
HCT-15	0.279	0.875	0.760	0.599	0.561	0.149	0.278	81	54	47	-47	.	3.65E-07	3.19E-06	>1.00E-04		
HT29	0.215	1.453	1.355	1.299	0.801	0.075	0.159	92	88	47	-65	-26	8.58E-07	2.64E-06			
KM12	0.535	2.184	2.151	2.044	1.425	0.146	0.264	98	91	54	-73	-51	1.07E-06	2.67E-06	6.62E-06		
SW-620	0.187	1.202	1.197	1.198	0.736	0.127	0.136	100	100	54	-32	-28	1.12E-06	4.23E-06	>1.00E-04		
CNS Cancer																	
SF-268	0.376	1.142	1.016	0.930	0.789	0.252	0.280	84	72	54	-33	-26	1.11E-06	4.16E-06	>1.00E-04		
SF-295	0.421	1.224	1.173	1.190	0.915	0.225	0.287	94	96	62	-47	-32	1.28E-06	3.71E-06	>1.00E-04		
SNB-19	0.533	1.721	1.562	1.655	1.230	0.219	0.324	87	94	59	-59	-39	1.18E-06	3.15E-06			
U251	0.391	1.301	1.298	1.139	0.810	0.144	0.320	100	82	46	-63	-18	7.75E-07	2.64E-06			
Melanoma																	
LOX IMVI	0.353	1.166	1.204	0.974	0.742	0.176	0.352	105	76	48	-50	.	8.36E-07	3.07E-06			
MALME-3M	0.666	1.165	1.123	1.128	1.102	0.218	0.290	92	93	87	-67	-56	1.74E-06	3.67E-06	7.73E-06		
M14	0.315	0.986	0.975	0.974	0.916	0.211	0.256	98	98	90	-33	-19	2.10E-06	5.37E-06	>1.00E-04		
SK-MEL-2	0.456	0.883	0.943	0.956	0.885	0.289	0.330	114	117	100	-37	-28	2.33E-06	5.40E-06	>1.00E-04		
SK-MEL-28	0.530	1.482	1.426	1.320	1.130	0.071	0.218	94	83	63	-87	-59	1.22E-06	2.64E-06	5.69E-06		
SK-MEL-5	0.549	2.147	1.995	1.958	1.659	0.136	0.364	91	88	69	-75	-34	1.36E-06	3.02E-06			
UACC-62	0.675	1.600	1.649	1.546	1.407	0.137	0.364	105	94	79	-80	-46	1.53E-06	3.15E-06			
Ovarian Cancer																	
IGROV1	0.462	1.060	1.044	1.103	0.902	0.235	0.302	97	107	74	-49	-35	1.56E-06	1.97E-06	>1.00E-04		
OVCA-3	0.460	0.929	0.879	0.849	0.816	0.140	0.301	89	83	76	-70	-35	1.51E-06	3.32E-06			
OVCA-4	0.530	1.121	1.114	1.073	1.086	0.093	0.172	99	92	94	-82	-68	1.78E-06	3.41E-06	6.55E-06		
OVCA-5	0.512	1.507	1.480	1.540	1.412	0.087	0.149	97	103	90	-83	-71	1.71E-06	3.32E-06	6.45E-06		
OVCA-8	0.400	1.101	1.078	1.114	0.755	0.269	0.306	97	102	51	-33	-24	1.02E-06	4.05E-06	>1.00E-04		
SK-OV-3	0.399	1.623	1.514	1.597	1.262	0.083	0.302	91	98	71	-79	-24	1.37E-06	2.96E-06			
Renal Cancer																	
786-0	0.652	1.958	2.012	1.897	1.440	0.181	0.299	104	95	60	-72	-54	1.20E-06	2.85E-06	6.79E-06		
A498	0.045	2.261	2.230	2.598	2.201	0.660	0.433	98	126	95	-30	-54	2.30E-06	5.75E-06	6.69E-06		
ACHN	0.374	1.026	0.985	1.037	0.829	0.150	0.313	94	102	70	-60	-16	1.42E-06	3.45E-06			
CAKI-1	0.597	1.862	1.692	1.802	1.126	0.147	0.335	87	95	42	-75	-44	7.03E-07	2.27E-06			
RXP 393	0.403	0.801	0.766	0.760	0.628	0.177	0.441	91	75	56	-56	10	1.14E-06				
SN12C	0.584	1.233	1.209	1.194	1.093	0.203	0.308	96	94	78	-65	-47	1.58E-06	3.51E-06			
TK-10	0.672	1.496	1.512	1.492	1.304	0.138	0.200	102	100	77	-80	-70	1.48E-06	3.10E-06	6.47E-06		
UO-31	0.443	1.782	1.572	1.563	1.356	0.151	0.285	84	84	68	-64	-36	1.37E-06	3.29E-06			
Prostate Cancer																	
PC-3	0.242	0.996	0.910	0.763	0.593	0.055	0.108	89	69	47	-77	-55	7.05E-07	2.38E-06	6.02E-06		
DU-145	0.327	0.961	0.993	1.018	0.895	0.078	0.132	105	109	90	-76	-60	1.73E-06	3.47E-06	6.95E-06		
Breast Cancer																	
MCF7	0.452	1.672	1.566	1.541	0.982	0.196	0.255	91	89	43	-57	-44	7.18E-07	2.72E-06			
NCI/ADR-RES	0.556	1.904	1.883	1.878	1.217	0.229	0.212	98	98	49	-59	-62	9.54E-07	2.45E-06	8.28E-06		
MDA-MB-231/ATCC	0.642	0.988	0.988	0.889	0.843	0.173	0.306	100	71	58	-73	-52	1.15E-06	2.77E-06	6.67E-06		
HS 578T	0.534	1.186	1.221	1.179	1.110	0.351	0.333	105	99	88	-34	-38	2.05E-06	5.25E-06	>1.00E-04		
MDA-MB-435	0.404	1.469	1.502	1.447	1.126	0.127	0.347	103	98	68	-69	-14	1.35E-06	3.14E-06			
BT-549	0.492	0.957	0.912	0.891	0.785	0.143	0.195	90	86	63	-71	-60	1.25E-06	2.95E-06	6.98E-06		
T-47D	0.423	1.019	0.921	1.071	0.935	0.205	0.238	84	109	86	-52	-44	1.82E-06	4.21E-06			

Table 6. Results of testing Compound 71 against 60 cancer cell lines.


Panel/Cell Line	Time Zero	Ctrl	Log10 Concentration					Percent Growth					GIS6	TOI	LC50
			-8.0	-7.0	-6.0	-5.0	-4.0	-3.0	-2.0	-1.0	0.0	1.0			
Leukemia															
CCRF-CEM	0.295	0.949	0.808	0.811	0.718	0.002	0.140	78	79	65	-99	-53	1.23E-06	2.48E-06	5.00E-06
K-562	0.297	1.401	1.338	1.283	0.438	0.126	0.119	94	89	13	-58	-60	3.26E-07	1.52E-06	7.77E-06
MO1T-4	0.313	0.957	0.833	0.736	0.461	0.143	0.143	81	65	23	-54	-54	2.33E-07	1.98E-06	8.75E-06
RFMT-8226	0.332	0.682	0.536	0.559	0.513	0.016	0.176	58	65	52	-95	-47	1.03E-06	2.25E-06	.
SR	0.364	0.888	0.688	0.583	0.254	0.234	0.287	62	42	-30	-36	-21	3.86E-08	3.79E-07	>1.00E-04
Non-Small Cell Lung Cancer															
A549/ATCC	0.363	1.175	1.140	1.145	1.121	-0.032	0.135	96	96	93	-100	-63	1.68E-06	3.04E-06	5.51E-06
EKVX	0.557	0.841	0.713	0.737	0.711	-0.008	0.145	55	63	54	-100	-74	1.06E-06	2.24E-06	4.74E-06
HOP-62	0.510	1.288	1.285	1.267	1.255	0.011	0.329	100	97	96	-98	-36	1.72E-06	3.12E-06	.
HOP-92	1.070	1.647	1.567	1.541	1.361	0.089	0.767	86	82	50	-92	-28	1.01E-06	2.26E-06	.
NCI-H23	0.457	1.669	1.585	1.515	1.679	-0.041	0.151	93	87	101	-100	-67	1.79E-06	3.18E-06	5.64E-06
NCI-H322M	0.626	1.181	0.988	1.049	1.031	-0.052	0.257	65	76	73	-100	-59	1.36E-06	2.64E-06	5.14E-06
NCI-H460	0.319	1.792	1.654	1.658	1.650	0.011	0.257	91	91	90	-97	-20	1.64E-06	3.04E-06	.
NCI-H522	0.766	1.671	1.509	1.558	1.471	0.037	0.439	82	87	78	-95	-43	1.45E-06	2.82E-06	.
Colon Cancer															
COLO 205	0.221	1.196	1.146	1.028	0.930	0.009	0.151	95	83	73	-96	-32	1.36E-06	2.69E-06	.
HCT-116	0.114	0.714	0.607	0.633	0.525	-0.022	0.058	82	86	68	-100	-49	1.29E-06	2.55E-06	.
HCT-15	0.279	0.893	0.669	0.779	0.797	-0.003	0.082	64	81	84	-100	-71	1.54E-06	2.87E-06	5.36E-06
HT29	0.215	1.333	1.276	1.238	1.135	-0.054	0.003	95	91	82	-100	-99	1.50E-06	2.83E-06	5.32E-06
KM12	0.535	1.957	1.867	1.832	1.777	0.013	0.307	94	91	87	-98	-43	1.59E-06	2.97E-06	.
SW-620	0.187	1.073	0.923	0.993	1.060	-0.048	0.015	83	91	98	-100	-92	1.75E-06	3.13E-06	5.60E-06
CNS Cancer															
SP-268	0.376	1.016	0.893	0.939	0.886	0.148	0.278	81	88	80	-61	-26	1.63E-06	3.69E-06	.
SP-295	0.421	1.107	1.033	0.931	0.987	-0.025	0.195	89	74	84	-100	-54	1.53E-06	2.86E-06	5.35E-06
SNB-19	0.532	1.483	1.385	1.385	1.351	0.063	0.317	90	90	85	-88	-40	1.61E-06	3.12E-06	.
U251	0.391	1.156	1.125	1.059	1.123	0.026	0.484	96	87	96	-93	12	1.74E-06	.	.
Melanoma															
LOX IMVI	0.353	1.164	0.948	1.033	0.935	0.008	0.165	73	84	72	-98	-53	1.34E-06	2.65E-06	5.22E-06
MALME-3H	0.666	0.984	0.832	0.838	0.826	0.089	0.333	52	54	50	-87	-50	1.01E-06	2.33E-06	1.00E-04
M14	0.315	0.935	0.772	0.823	0.816	-0.004	0.210	74	82	81	-100	-33	1.48E-06	2.80E-06	.
SK-MEL-2	0.456	0.830	0.775	0.819	0.780	-0.010	0.324	85	97	87	-100	-29	1.57E-06	2.91E-06	.
SK-MEL-28	0.510	1.508	1.439	1.408	1.367	-0.002	0.248	93	90	86	-100	-53	1.56E-06	2.89E-06	5.38E-06
SK-MEL-5	0.549	1.640	1.611	1.762	1.812	0.143	0.363	97	111	116	-74	-34	2.22E-06	4.07E-06	.
UACC-62	0.675	1.477	1.323	1.338	1.301	-0.041	0.272	81	83	78	-100	-60	1.44E-06	2.74E-06	5.24E-06
Ovarian Cancer															
IGROV1	0.462	1.091	0.997	0.983	0.806	-0.039	0.216	55	83	55	-100	-53	1.07E-06	2.26E-06	4.75E-06
OVCAR-3	0.400	0.868	0.833	0.792	0.770	-0.043	0.131	91	81	76	-100	-72	1.40E-06	2.70E-06	5.20E-06
OVCAR-4	0.530	1.123	1.011	0.999	0.945	-0.001	0.179	81	79	70	-100	-66	1.31E-06	2.58E-06	5.08E-06
OVCAR-5	0.512	1.592	1.538	1.437	1.559	-0.033	0.474	95	86	97	-100	-8	1.73E-06	3.11E-06	.
OVCAR-8	0.400	0.968	0.923	0.946	0.934	-0.001	0.126	92	96	94	-100	-69	1.69E-06	3.05E-06	5.52E-06
SK-OV-3	0.399	1.327	1.261	1.245	1.093	0.062	0.278	93	91	75	-85	-30	1.43E-06	2.94E-06	.
Renal Cancer															
786-O	0.652	1.912	1.862	1.668	1.663	0.046	0.458	96	96	80	-93	-30	1.49E-06	2.91E-06	.
A498	0.945	2.207	2.413	2.087	1.869	1.903	0.540	116	91	73	76	-43	1.65E-05	4.35E-05	>1.00E-04
ACHN	0.374	1.026	0.898	0.902	0.964	0.034	0.291	80	81	90	-91	-22	1.67E-06	3.15E-06	.
CAKI-1	0.597	1.767	1.735	1.642	1.804	-0.067	0.315	97	89	103	-100	-47	1.83E-06	3.22E-06	.
RXFP 393	0.403	0.829	0.582	0.629	0.629	0.156	0.385	42	53	53	-61	-4	.	2.91E-06	.
SN12C	0.584	1.115	1.050	0.998	1.045	-0.014	0.224	88	78	87	-100	-62	1.57E-06	2.91E-06	5.40E-06
TK-10	0.672	1.134	1.032	1.087	0.998	-0.021	0.245	78	90	71	-100	-64	1.32E-06	2.59E-06	5.09E-06
VO-31	0.443	1.467	1.375	1.407	1.332	-0.030	0.288	93	94	87	-100	-35	1.57E-06	2.92E-06	.
Prostate Cancer															
PC-3	0.242	1.068	0.862	0.846	0.857	-0.022	0.179	75	73	75	-100	-26	1.38E-06	2.67E-06	.
DU-145	0.327	0.835	0.694	0.767	0.760	-0.074	-0.019	72	87	85	-100	-100	1.55E-06	2.89E-06	5.37E-06
Breast Cancer															
HCP7	0.452	1.540	1.467	1.340	1.507	0.073	0.229	93	82	97	-84	-49	1.82E-06	3.44E-06	.
NCI/ADR-RES	0.556	1.942	1.920	1.882	1.874	0.490	0.294	98	96	95	-12	-47	2.64E-06	7.73E-06	>1.00E-04
MDA-MB-231/ATCC	0.642	0.884	0.756	0.830	0.709	0.015	0.389	47	78	27	-98	-39	.	1.66E-06	.
HS 578T	0.534	2.167	1.099	1.165	1.127	0.168	0.333	89	100	94	-69	-38	1.86E-06	3.78E-06	.
MDA-MB-415	0.404	1.498	1.376	1.382	1.444	-0.073	0.295	89	89	95	-100	-27	1.70E-06	3.07E-06	.
T-47D	0.423	0.750	0.541	0.578	0.529	0.109	0.198	36	47	32	-76	-53	<1.00E-08	1.98E-06	5.72E-06

[0086] The examples described herein are meant to be illustrative of the synthesis and applications of the compounds described. The examples are not meant to limit the scope of the invention described herein.

The invention claimed is:

The invention claimed is:

1. A compound of formula I




wherein

X is selected from the group consisting of alkyl and haloalkyl;

Ar is selected from the group consisting of phenyl, biphenyl, naphthyl, anthryl, phenanthryl, and fluorenyl;

R is selected from the group consisting of -CN, -CH₂CN, -CH₂CH₂CN, -CH₂CH₂CH₂CN -



or metabolites or pharmaceutically acceptable salts thereof.

2. The compound of claim 1 wherein X is C₁ to C₄ haloalkyl.
3. The compound of claim 2 wherein X is CF₃.
4. The compound of claim 1, wherein Ar is substituted at any substitutable position with one or more radicals selected from the group consisting of halo, C₁-C₄ alkyl, C₁-C₄ haloalkyl, azido, C₁-C₄ azidoalkyl, aryl, alkylaryl, haloaryl, haloalkylaryl, and combinations thereof.
5. The compound of claim 1 wherein Ar is selected from the group consisting of 2-naphthyl, 4-biphenyl, 9-anthryl, 2-fluorenyl, 4-azidophenyl, 4-azidomethylphenyl,

4-(2-azidoethyl)phenyl, 4-(3-azidopropyl)phenyl, 4-(4-azidobutyl)phenyl, 4-(4-azidophenyl)phenyl, 4-(4-azidomethylphenyl)phenyl, 4-methylphenyl, 4-ethylphenyl, 4-propylphenyl, 4-butylphenyl, 4-(2-bromoethyl)phenyl, 4-(3-bromopropyl)phenyl, 4-(4-bromobutyl)phenyl, 4-(trifluoromethyl)phenyl, 4-(4-methylphenyl)phenyl, 4-(4-bromomethylphenyl)phenyl, 4-(4-butylphenyl)phenyl, 4-(4-*tert*-butylphenyl)phenyl, 2-chlorophenyl, 4-chlorophenyl, 2,4-dichlorophenyl, 3,4-dichlorophenyl, 2,5-dichlorophenyl, 2,4-dimethylphenyl, 2,5-dimethylphenyl, 3,4-dimethylphenyl, 3,5-dimethylphenyl, 4-(4-chlorophenyl)phenyl, 4-(3,5-dichlorophenyl)phenyl, 4-(2,3-dichlorophenyl)phenyl, 4-(3,5-dimethylphenyl)phenyl, 4-(2,4,5-trichlorophenyl)phenyl, 4-(4-trifluoromethylphenyl)phenyl, 2-phenanthrenyl, 3-indolyl, 2-pyrrolyl, and 4-(benzyl)phenyl.


6. The compound of claim 5 wherein Ar is selected from the group consisting of 4-(2-bromoethyl)phenyl, 4-(3-bromopropyl)phenyl, 4-(2-azidoethyl)phenyl, 4-(3-azidopropyl)phenyl, 4-butylphenyl, 4-*t*-butylphenyl, 2-naphthalenyl, 3-indolyl, 4-biphenyl, 4'-chloro[1,1'-biphenyl]-4-yl, 3',5'-dichloro[1,1'-biphenyl]-4-yl, 2',3'-dichloro[1,1'-biphenyl]-4-yl, 4'-methyl[1,1'-biphenyl]-4-yl, 4'-trifluoromethyl[1,1'-biphenyl]-4-yl, 4'-bromomethyl[1,1'-biphenyl]-4-yl, 3',5'-dimethyl[1,1'-biphenyl]-4-yl, 4'-butyl[1,1'-biphenyl]-4-yl, 4'-*tert*-butyl[1,1'-biphenyl]-4-yl, 4-(phenylmethyl)phenyl, 9H-fluoren-2-yl, 9-anthracenyl, 2-phenanthrenyl, 9-phenanthrenyl.

7. The compound of claim 6 wherein Ar is 2-phenanthrenyl.

8. The compound of claim 1 wherein R is selected from aminoacetamide and guanidine.

9. The compound of claim 1 wherein X is CF₃, Ar is 2-phenanthrenyl, and R is selected from aminoacetamide and guanidine.


10. A compound of formula 2



II

X is selected from the group consisting of alkyl and haloalkyl;

R is selected from the group consisting of -CN, -CH₂CN, -CH₂CH₂CN, -CH₂CH₂CH₂CN –



or metabolites or pharmaceutically acceptable salts thereof.


11. The compound of claim 10 wherein X is C₁ to C₄ haloalkyl.

12. The compound of claim 11 wherein X is CF₃.

13. The compound of claim 10 wherein R is selected from aminoacetamide and guanidine.


or a pharmaceutically acceptable salt thereof.

14. A compound of formula III:



X is selected from the group consisting of alkyl and haloalkyl;

R is selected from the group consisting of -CN, -CH₂CN, -CH₂CH₂CN, -CH₂CH₂CH₂CN –



or metabolites or pharmaceutically acceptable salts thereof.


15. The compound of claim 14 wherein X is C₁ to C₄ haloalkyl.

16. The compound of claim 15 wherein X is CF₃.

17. The compound of claim 14 wherein R is selected from aminoacetamide and guanidine.

or a pharmaceutically acceptable salt thereof.


18. A compound of formula IV:



IV

or metabolites or pharmaceutically acceptable salts thereof.


19. A compound of formula V



V

or metabolites or pharmaceutically acceptable salts thereof.

20. A method of inducing apoptosis in unwanted rapidly proliferating cells, the method comprising the step of contacting a therapeutically effective amount of a compound of formula I




wherein

X is selected from the group consisting of alkyl and haloalkyl;

Ar is selected from the group consisting of phenyl, biphenyl, naphthyl, anthryl, phenanthryl, and fluorenyl;


R is selected from the group consisting of -CN, -CH₂CN, -CH₂CH₂CN, -CH₂CH₂CH₂CN -




or metabolites or pharmaceutically acceptable salts thereof, with the rapidly proliferating cells.

21. The method of claim 20, wherein the rapidly proliferating cells are cancer cells, wherein the cancer cells are selected from the group consisting of leukemia, non-small cell lung cancer, colon cancer, central nervous system cancer, melanoma, ovarian cancer, renal cancer, prostate cancer, and breast cancer.

22. The method of claim 20 wherein the compound is selected from compounds IV and V




IV



V

or a combination thereof, or metabolites or pharmaceutically acceptable salts thereof.

23. A method for treating, inhibiting, or delaying the onset of cancer, wherein the cancer is selected from the group consisting of leukemia, non-small cell lung cancer, colon cancer, central nervous system cancer, melanoma, ovarian cancer, renal cancer, prostate cancer, and breast cancer, in a subject in need of such treatment, the method comprising administering a therapeutically effective amount of a compound of formula I:




I

wherein

X is selected from the group consisting of alkyl and haloalkyl;



Ar is selected from the group consisting of phenyl, biphenyl, naphthyl, anthryl, phenanthryl, and fluorenyl;

R is selected from the group consisting of -CN, -CH₂CN, -CH₂CH₂CN, -CH₂CH₂CH₂CN -



or metabolites or pharmaceutically acceptable salts thereof, to the subject in need of such treatment.


24. The method of claim 23 wherein the compound is selected from compounds IV and V



or a combination thereof, or metabolites or pharmaceutically acceptable salts thereof.

25. The method of claim 23 wherein the subject is a human.

26. A method for preventing restenosis in a subject that has undergone an angioplasty or stent procedure, the method comprising administering a therapeutically effective amount of a compound of formula I:




wherein

X is selected from the group consisting of alkyl and haloalkyl;

Ar is selected from the group consisting of phenyl, biphenyl, naphthyl, anthryl, phenanthryl, and fluorenyl;

R is selected from the group consisting of -CN, -CH₂CN, -CH₂CH₂CN, -CH₂CH₂CH₂CN –



or metabolites or pharmaceutically acceptable salts thereof, to the subject in need of such treatment.