

where R1 and R2 are the same or different and are each independently

- 1) $-(C_6-C_{14})$ -aryl where aryl is unsubstituted or mono-, di- or trisubstituted independently by R11,

where R11 is

- a) F, Cl, I or Br,
 - b) $-(C_1-C_4)$ -alkyl,
 - c) -CN,
 - d) $-CF_3$,
 - e) $-OR^5$ in which R^5 is a hydrogen atom or $-(C_1-C_4)$ -alkyl,
 - f) $-N(R^5)-R^6$ in which R^5 and R^6 are each independently a hydrogen atom or $-(C_1-C_4)$ -alkyl,
 - g) $-C(O)-R^5$ in which R^5 is a hydrogen atom or $-(C_1-C_4)$ -alkyl, or
 - h) $-S(O)_x-R^5$ in which x is the integer zero, 1 or 2, and R^5 is a hydrogen atom or $-(C_1-C_4)$ -alkyl, or
- 2) a 4- to 15-membered Het ring where the Het ring is unsubstituted or mono-, di- or trisubstituted independently by $-(C_1-C_5)$ -alkyl, $-(C_1-C_5)$ -alkoxy, halogen, nitro, amino, trifluoromethyl, hydroxyl, hydroxy- $-(C_1-C_4)$ -alkyl, methylenedioxy, ethylenedioxy, formyl, acetyl, cyano, hydroxycarbonyl, aminocarbonyl or $-(C_1-C_4)$ -alkoxycarbonyl,

- R3 is 1) $-(C_6-C_{14})$ -aryl in which aryl is unsubstituted or mono-, di- or trisubstituted independently

by $-NO_2$, $-O-(C_1-C_4)$ -alkyl, F, Cl or bromine,

- 2) $-O-C(CH_3)_3$, or

- 3) $-O-CH(R7)$ -aryl in which aryl is unsubstituted or mono-, di- or trisubstituted independently

by $-NO_2$, $-O-CH_3$, F, Cl or bromine,

in which R7 is a hydrogen atom or $-(C_1-C_4)$ -alkyl,

- R4 is 1) a hydrogen atom,


- 2) $-(C_1-C_4)$ -alkyl or

- 3) $-CH(R8)$ -aryl

in which R8 is a hydrogen atom or $-(C_1-C_4)$ -alkyl.


1b

In a second aspect, there is provided a compound of the formula IV



where R1, R2, R3 and R4 are each as defined as in the compound of the formula II according to the first aspect.

In a third aspect, there is provided a process for obtaining the compound of the formula I



where R1 is

- 1) a hydrogen atom,
- 2) $-(C_1-C_4)$ -alkyl,
- 3) $-(C_6-C_{14})$ -aryl where aryl is unsubstituted or mono-, di- or trisubstituted independently by R11,

where R11 is

- a) F, Cl, I or Br,
- b) $-(C_1-C_4)$ -alkyl,
- c) -CN,
- d) -CF₃,
- e) -OR⁵ in which R⁵ is a hydrogen atom or $-(C_1-C_4)$ -alkyl,
- f) -N(R⁵)-R⁶ in which R⁵ and R⁶ are each independently a hydrogen atom or $-(C_1-C_4)$ -alkyl,

- g) $-C(O)-R^5$ in which R^5 is a hydrogen atom or $-(C_1-C_4)$ -alkyl, or
 h) $-S(O)_x-R^5$ in which x is the integer zero, 1 or 2, and R^5 is a hydrogen atom or $-(C_1-C_4)$ -alkyl,

5 4) $-CH(R7)$ -aryl in which aryl is unsubstituted or mono-, di- or trisubstituted independently
 by $-NO_2$, $-O-CH_3$, F, Cl or bromine, where R7 is a hydrogen atom or $-(C_1-C_4)$ -alkyl, or

10 5) a 4- to 15-membered Het ring where the Het ring is unsubstituted or mono-, di- or trisubstituted independently by
 $-(C_1-C_5)$ -alkyl, $-(C_1-C_5)$ -alkoxy, halogen, nitro, amino, trifluoromethyl, hydroxyl, hydroxy- $-(C_1-C_4)$ -alkyl, methylenedioxy, ethylenedioxy, formyl, acetyl, cyano, hydroxycarbonyl, aminocarbonyl or $-(C_1-C_4)$ -alkoxycarbonyl,

15 R2 is 1) $-(C_6-C_{14})$ -aryl where aryl is unsubstituted or mono-, di- or trisubstituted independently by R11,

2) $-CH(R7)$ -aryl in which aryl is unsubstituted or mono-, di- or trisubstituted independently
 by $-NO_2$, $-O-CH_3$, F, Cl or bromine, where R7 is a hydrogen atom or $-(C_1-C_4)$ -alkyl, or

20 3) a 4- to 15-membered Het ring where the Het ring is unsubstituted or mono-, di- or trisubstituted
 independently by $-(C_1-C_5)$ -alkyl, $-(C_1-C_5)$ -alkoxy, halogen, nitro, amino, trifluoromethyl, hydroxyl,
 25 hydroxy- $-(C_1-C_4)$ -alkyl, methylenedioxy, ethylenedioxy, formyl, acetyl, cyano, hydroxycarbonyl, aminocarbonyl
 or $-(C_1-C_4)$ -alkoxycarbonyl,

R3 is 1) a hydrogen atom,

2) $-(C_1-C_4)$ -alkyl,

30 3) $-(C_6-C_{14})$ -aryl in which aryl is unsubstituted or mono-, di- or trisubstituted independently
 by $-NO_2$, $-O-(C_1-C_4)$ -alkyl, F, Cl or bromine,

4) $-O-C(CH_3)_3$, or


35 5) $-O-CH(R7)$ -aryl in which aryl is unsubstituted or mono-, di- or trisubstituted independently
 by $-NO_2$, $-O-CH_3$, F, Cl or bromine,
 where R7 is a hydrogen atom or $-(C_1-C_4)$ -alkyl,

1d

- R4 is 1) a hydrogen atom,
2) $-(C_1-C_4)$ -alkyl or
3) $-CH(R_8)$ -aryl


in which R8 is a hydrogen atom or $-(C_1-C_4)$ -alkyl,

5 which comprises hydrogenating a compound of the formula II



10 in which R1, R2, R3 and R4 are each as defined in the formula I and
the compound of the formula II may be present in the E or in the Z
configuration on the double bond,
in the presence of hydrogen and a catalyst.


15 In a fourth aspect, there is provided a process for obtaining the compound of the
formula III



20 or salts thereof in which R1, R2 and R4 are each as defined in the
formula I according to the third aspect,
which comprises


- 25 a) hydrogenating the compound of the formula II

1e



in which R1, R2, R3 and R4 are each as defined in the formula I,


in the presence of hydrogen and a catalyst, and converting it to a compound of the formula I



and

- b) converting the resulting compound of the formula I to a compound of the formula III.

The disclosure herein relates to a process for obtaining the compound of the formula I




where R1 and R2 are the same or different and are each independently

- 1) a hydrogen atom,
 - 2) $-(C_1-C_4)$ -alkyl,
 - 5 3) $-(C_6-C_{14})$ -aryl where aryl is unsubstituted or mono-, di- or trisubstituted independently by R11, where R11 is
 - a) F, Cl, I or Br,
 - b) $-(C_1-C_4)$ -alkyl,
 - 10 c) $-CN$,
 - d) $-CF_3$,
 - e) $-OR^5$ in which R^5 is a hydrogen atom or $-(C_1-C_4)$ -alkyl,
 - f) $-N(R^5)-R^6$ in which R^5 and R^6 are each independently a hydrogen atom or $-(C_1-C_4)$ -alkyl,
 - 15 g) $-C(O)-R^5$ in which R^5 is a hydrogen atom or $-(C_1-C_4)$ -alkyl, or
 - h) $-S(O)_x-R^5$ in which x is the integer zero, 1 or 2, and R^5 is a hydrogen atom or $-(C_1-C_4)$ -alkyl,
 - 4) $-CH(R7)$ -aryl in which aryl is unsubstituted or mono-, di- or trisubstituted independently
20 by $-NO_2$, $-O-CH_3$, F, Cl or bromine, where R7 is a hydrogen atom or $-(C_1-C_4)$ -alkyl, or
 - 5) a 4- to 15-membered Het ring where the Het ring is unsubstituted or mono-, di- or trisubstituted independently by
25 $-(C_1-C_5)$ -alkyl, $-(C_1-C_5)$ -alkoxy, halogen, nitro, amino, trifluoromethyl, hydroxyl, hydroxy- (C_1-C_4) -alkyl, methylenedioxy, ethylenedioxy, formyl, acetyl, cyano, hydroxycarbonyl, aminocarbonyl or $-(C_1-C_4)$ -alkoxycarbonyl,
- R3 is
- 30 1) a hydrogen atom,
 - 2) $-(C_1-C_4)$ -alkyl,
 - 3) $-(C_6-C_{14})$ -aryl in which aryl is unsubstituted or mono-, di- or trisubstituted independently
by $-NO_2$, $-O-(C_1-C_4)$ -alkyl, F, Cl or bromine,
 - 4) $-O-C(CH_3)_3$, or
 - 35 5) $-O-CH(R7)$ -aryl in which aryl is unsubstituted or mono-, di- or trisubstituted independently
by $-NO_2$, $-O-CH_3$, F, Cl or bromine,
where R7 is a hydrogen atom or $-(C_1-C_4)$ -alkyl,

- R4 is
- 1) a hydrogen atom,
 - 2) $-(C_1-C_4)$ -alkyl or
 - 3) $-CH(R_8)$ -aryl
- in which R8 is a hydrogen atom or $-(C_1-C_4)$ -alkyl,

5 which comprises hydrogenating a compound of the formula II



10 in which R1, R2, R3 and R4 are each as defined in the formula I and the compound may be present in the E or in the Z configuration on the double bond,
in the presence of hydrogen and a catalyst.

15 The disclosure herein further relates to a process for obtaining the compound of the formula I


where R1 is phenyl or a hydrogen atom,

R2 is phenyl, pyridyl or thiazolyl, in which phenyl, pyridyl or thiazolyl is unsubstituted or substituted by fluorine or chlorine, and

R3 is phenyl or $-O-CH_2$ -phenyl and

20 R4 is methyl or ethyl.


The disclosure herein further relates to a process for obtaining the compound of the formula




25

or salts thereof in which R1, R2 and R4 are each as defined in the formula I,
which comprises

- a) hydrogenating the compound of the formula II



- 5 in which R1, R2, R3 and R4 are each as defined in the formula I,
in the presence of hydrogen and a catalyst, and converting it to a compound of the formula I



10

and

- b) converting the resulting compound of the formula I to a compound of the formula III.

15

Process step b) is performed, for example, according to the reaction conditions as described by T. Greene, P. Wuts in *Protective Groups in Organic Synthesis*, Wiley-Interscience, for the cleavage of amides or carbamates. Depending on the reaction conditions selected, here strongly basic reaction conditions in particular, the direct conversion of compounds of the formula I to compounds of the formula III leads to racemization of the chiral center formed by the asymmetric hydrogenation or to other undesired side reactions. This can be prevented when the compound of the formula I is converted to the compound IV with, for example, tert-butyl dicarbonate or another reagent for the introduction of tert-butoxycarbonyl protecting groups. The tert-butoxycarbonyl protecting group is introduced in a suitable solvent such as acetonitrile, tetrahydrofuran or toluene, preferably with the aid of an acylation catalyst such as N,N-dimethylaminopyridine (DMAP).

20

25


The reaction temperature is from 0°C to 120°C, preferably from 20°C to 40°C.

The reaction time is generally from 0.5 to 24 hours, depending on the composition of the mixture and the temperature range selected.

- 5 The resulting compound of the formula IV is then converted to a compound of the formula Ia under mild conditions, such as magnesium methoxide.


The conversion to the compounds of the formula III is typically done under reaction conditions known from the literature, as described by T. Greene, P. Wuts in *Protective Groups in Organic Synthesis*, Wiley-Interscience, for the cleavage of tert-butyloxycarbonyl (BOC) protecting groups.

10




The disclosure herein further relates to a process for obtaining the compound of the formula III,
 15 which comprises

- a) hydrogenating the compound of the formula II
 in which R1, R2, R3 and R4 are each as defined in the
 formula I
 20 in the presence of hydrogen and a catalyst and converting it to a compound of the formula I,
- b) reacting the resulting compound of the formula I with a tert-butyl dicarbonate and an acylation catalyst such as dimethylaminopyridine (DMAP) to give a compound of the
 25 formula IV



- in which R1, R2, R3 and R4 are each as defined in formula I,
 c) then converting the resulting compound of the formula IV to
 the compound of the formula Ia



- in which R1, R2, R3 and R4 are each as defined in formula I,
 and
 d) converting the resulting compound of the formula Ia to the
 compound of the formula III or salts thereof, in which R1, R2
 and R4 are each as defined in formula I.

The conversion of the compounds of the formula IV to the compound of the
 formula Ia is achieved, for example, by treatment with bases, such as
 lithium hydroxide, hydrazine or magnesium methoxide (literature: J. Org.
 Chem. **1997**, 62, 7054-7057). The tert-butoxycarbonyl group is detached
 to give compounds of the formula III under standard conditions, such as
 treatment with trifluoroacetic acid (TFA), hydrochloric acid or
 p-toluenesulfonic acid in suitable solvents.

The undesired enantiomer is depleted by crystallization of the compounds
 of the formula I or III from suitable solvents such as methanol, ethanol,
 1-propanol, 2-propanol, n-butanol, 2-butanol and esters thereof. In the case
 of compounds of the formula III, the crystallization is preferably performed
 in the form of their (acidic) salts such as hydrochloride, methanesulfonate

or p-toluenesulfonate. Under these conditions, optical purities of > 99% are achieved. Appropriately, the entire reaction sequence can be performed in a one-pot process without isolation of the compounds IV and Ia. The yields and optical purities achieved here correspond to the values mentioned
5 above.

The term "catalyst" refers to compounds as described, for example, by E. N. Jacobson, A. Pfaltz, H. Yamamoto in *Comprehensive Asymmetric Catalysis*, Springer-Verlag, 1999 or X. Zhang, *Chemical Reviews*, 2003,
10 103, 3029-3069 and the literature cited there, for example optically active rhodium, ruthenium or iridium complexes or mixtures thereof. The catalytically active complex is formed by reaction of a metal complex with an optically active phosphine. In the case of the above-described acylated 2,3-diaminopropionic acid derivatives, the Me-Duphos or Et-Duphos-
15 rhodium complexes exhibited very good enantioselectivities and conversions. It is also known that chiral β -amino acids can be prepared by using rhodium complexes of the BICP, t-Bu-BisP, BDPMI, Et-FerroTANE, MalPHOS and MonoPHOS type as catalysts.

20 The terms "-(C₁-C₄)-alkyl" or "-(C₁-C₅)-alkyl" are understood to mean hydrocarbon radicals whose carbon chain is straight-chain or branched and contains from 1 to 4 or from 1 to 5 carbon atoms, for example methyl, ethyl, propyl, isopropyl, butyl, isobutyl, tert-butyl or pentyl.

25 The terms "-CH(R₇)-" or "-CH(R₈)-" are understood to mean straight-chain or branched hydrocarbon radicals such as methylene, ethylene, isopropylene, isobutylene or pentylene. For example, in the case that R₇ is a hydrogen atom and aryl is phenyl, the "-CH(R₇)-aryl" radical is the benzyl radical.

30 The terms "-(C₆-C₁₄)-aryl" or "aryl" are understood to mean aromatic carbon radicals having from 6 to 14 carbon atoms in the ring. -(C₆-C₁₄)-Aryl radicals are, for example, phenyl, naphthyl, for example 1-naphthyl, 2-naphthyl, anthryl or fluorenyl. Naphthyl radicals and in particular phenyl
35 radicals are preferred aryl radicals.

The term "4- to 15-membered Het ring" is understood to mean ring systems having from 4 to 15 carbon atoms, which are present in one, two or three

ring systems bonded to one another and which contain one, two, three or four identical or different heteroatoms from the group of oxygen, nitrogen or sulfur. Examples of these ring systems are the acridinyl, azepinyl, azetidiny, aziridinyl, benzimidazalinyl, benzimidazolyl, benzofuranyl, benzothiofuranyl, benzothiophenyl, benzoxazolyl, benzothiazolyl, benzotriazolyl, benzotetrazolyl, benzisoxazolyl, benzisothiazolyl, carbazolyl, 4aH-carbazolyl, carbolinyl, quinazoliny, quinolinyl, 4H-quinoliziny, quinoxaliny, quinuclidiny, chromanyl, chromenyl, cinnoliny, decahydroquinoliny, dibenzofuranyl, dibenzothiophenyl, dihydrofuran[2,3-b]tetrahydrofuranyl, dihydrofuranyl, dioxolyl, dioxanyl, 2H, 6H-1,5,2-dithiaziny, furanyl, furazanyl, imidazolidiny, imidazoliny, imidazolyl, 1H-indazolyl, indoliny, indoliziny, indolyl, 3H-indolyl, isobenzofuranyl, isochromanyl, isoindazolyl, isoindoliny, isoindolyl, isoquinoliny (benzimidazolyl), isothiazolidiny, 2-isothiazoliny, isothiazolyl, isoxazolyl, isoxazolidiny, 2-isoxazoliny, morpholiny, naphthyridiny, octahydroisoquinoliny, oxadiazolyl, 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl, oxazolidiny, oxazolyl, oxazolidiny, oxothiolanyl, pyrimidiny, phenanthridiny, phenanthroliny, phenaziny, phenothiaziny, phenoxathiiny, phenoxaziny, phthalaziny, piperaziny, piperidiny, pteridiny, puriny, pyranyl, pyraziny, pyrazolidiny, pyrazoliny, pyrazolyl, pyridaziny, pyridooxazolyl, pyridoimidazolyl, pyridothiazolyl, pyridothiophenyl, pyridyl, pyrimidiny, pyrrolidiny, pyrroliny, 2H-pyrrolyl, pyrrolyl, tetrahydrofuranyl, tetrahydroisoquinoliny, tetrahydroquinoliny, tetrahydropyridiny, 6H-1,2,5-thiadiaziny, 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, 1,2,5-thiadiazolyl, 1,3,4-thiadiazolyl, thianthrenyl, thiazolyl, thienyl, thienothiazolyl, thienooxazolyl, thienoimidazolyl, thiomorpholiny, thiophenyl, triaziny, 1,2,3-triazolyl, 1,2,3-triazolyl, 1,2,4-triazolyl, 1,2,5-triazolyl, 1,3,4-triazolyl and xanthenyl radicals.

The asterisk on a carbon atom in the compound of the formula I or II means that the particular carbon atom is chiral and that the compound is present either as the R- or S-enantiomer.

The asymmetric hydrogenation of the compounds of the formula II is advantageously performed at a temperature of from 10°C to 200°C and a hydrogen pressure of from 1 bar to 200 bar. The molar catalyst-reactant ratio is advantageously from 1:100 to 1:10 000.


Suitable solvents for the asymmetric hydrogenation are, for example,

water, lower alcohols such as methanol, ethanol, propanol or isopropanol, aromatic hydrocarbons such as toluene, ketones such as acetone, halogenated hydrocarbons such as dichloromethane, carboxylic esters such as ethyl acetate, and ethers such as tetrahydrofuran.

5

The optically active 2,3-diaminopropionic acid derivatives of the formula I, III and IV as such, including in the form of their enantiomer mixtures and their salts, likewise form part of the subject matter of the present invention. Enantiomer mixtures should be understood here to mean in particular those in which one enantiomer is enriched compared to the other.

The compounds of the formula II are either known or can be prepared, for example, by reacting compounds of the formula IV in which R³ and R⁴ are each as defined above with an amine of the formula V in which R¹ and R² are each as defined above.



The reaction temperature is from 0°C to 120°C, preferably from 20°C to 60°C.


The reaction time is generally from 0.5 to 8 hours, depending on the composition of the mixture and the selected temperature range. The resulting compound of the formula II is then removed from the reaction mixture by aqueous workup and extraction with a suitable solvent, for example ethyl acetate or dichloromethane, or by crystallization.

A further aspect of the invention relates to novel compounds of the formula II where R1 and R2 are the same or different and are each independently

- 30 1) $-(C_6-C_{14})$ -aryl where aryl is unsubstituted or mono-, di- or trisubstituted independently by R11,
where R11 is
a) F, Cl, I or Br,
b) $-(C_1-C_4)$ -alkyl,

- 5
- c) -CN,
d) -CF₃,
e) -OR⁵ in which R⁵ is a hydrogen atom or -(C₁-C₄)-alkyl,
f) -N(R⁵)-R⁶ in which R⁵ and R⁶ are each independently a
hydrogen atom or -(C₁-C₄)-alkyl,
g) -C(O)-R⁵ in which R⁵ is a hydrogen atom or -(C₁-C₄)-
alkyl, or
h) -S(O)_x-R⁵ in which x is the integer zero, 1 or 2, and R⁵
is a hydrogen atom or -(C₁-C₄)-alkyl, or
- 10 2) a 4- to 15-membered Het ring where the Het ring is
unsubstituted or mono-, di- or trisubstituted independently by
-(C₁-C₅)-alkyl, -(C₁-C₅)-alkoxy, halogen, nitro, amino, trifluoro-
methyl, hydroxyl, hydroxy-(C₁-C₄)-alkyl, methylenedioxy,
ethylenedioxy, formyl, acetyl, cyano, hydroxycarbonyl, amino-
15 carbonyl or -(C₁-C₄)-alkoxycarbonyl,
- R3 is 1) -(C₆-C₁₄)-aryl in which aryl is unsubstituted or mono-,
di- or trisubstituted independently
by -NO₂, -O-(C₁-C₄)-alkyl, F, Cl or bromine,
2) -O-C(CH₃)₃, or
20 3) -O-CH(R7)-aryl in which aryl is unsubstituted or mono-,
di- or trisubstituted independently
by -NO₂, -O-CH₃, F, Cl or bromine,
in which R7 is a hydrogen atom or -(C₁-C₄)-alkyl,
- R4 is 1) a hydrogen atom,
25 2) -(C₁-C₄)-alkyl or
3) -CH(R8)-aryl
in which R8 is a hydrogen atom or -(C₁-C₄)-alkyl.

30 A further aspect of the invention relates to novel compounds of the formula
IV



where R1 and R2 are the same or different and are each independently

- 1) $-(C_6-C_{14})$ -aryl where aryl is unsubstituted or mono-, di- or trisubstituted independently by R11,

where R11 is

- a) F, Cl, I or Br,
- b) $-(C_1-C_4)$ -alkyl,
- c) $-CN$,
- d) $-CF_3$,
- e) $-OR^5$ in which R^5 is a hydrogen atom or $-(C_1-C_4)$ -alkyl,
- f) $-N(R^5)-R^6$ in which R^5 and R^6 are each independently a hydrogen atom or $-(C_1-C_4)$ -alkyl,
- g) $-C(O)-R^5$ in which R^5 is a hydrogen atom or $-(C_1-C_4)$ -alkyl, or
- h) $-S(O)_x-R^5$ in which x is the integer zero, 1 or 2, and R^5 is a hydrogen atom or $-(C_1-C_4)$ -alkyl, or

- 2) a 4- to 15-membered Het ring where the Het ring is unsubstituted or mono-, di- or trisubstituted independently by $-(C_1-C_5)$ -alkyl, $-(C_1-C_5)$ -alkoxy, halogen, nitro, amino, trifluoromethyl, hydroxyl, hydroxy- $-(C_1-C_4)$ -alkyl, methylenedioxy, ethylenedioxy, formyl, acetyl, cyano, hydroxycarbonyl, aminocarbonyl or $-(C_1-C_4)$ -alkoxycarbonyl,

R3 is

- 1) $-(C_6-C_{14})$ -aryl in which aryl is unsubstituted or mono-, di- or trisubstituted independently by $-NO_2$, $-O-(C_1-C_4)$ -alkyl, F, Cl or bromine,
- 2) $-O-C(CH_3)_3$, or
- 3) $-O-CH(R7)$ -aryl in which aryl is unsubstituted or mono-, di- or trisubstituted independently by $-NO_2$, $-O-CH_3$, F, Cl or bromine, in which R7 is a hydrogen atom or $-(C_1-C_4)$ -alkyl,

- R4 is
- 1) a hydrogen atom,
 - 2) $-(C_1-C_4)$ -alkyl or
 - 3) $-CH(R_8)$ -aryl


in which R8 is a hydrogen atom or $-(C_1-C_4)$ -alkyl.

5

The invention therefore further relates to a process for obtaining the novel compounds of the formula IV, which comprises

- a) hydrogenating the compound of the formula II,
in which R1, R2, R3 and R4 are each as defined in the novel compound of the formula II,
in the presence of hydrogen and a catalyst and converting it to a compound of the formula I, and
- b) reacting the resulting compound of the formula I with a tert-butyl dicarbonate and an acylation catalyst such as dimethylaminopyridine (DMAP) to give a compound of the formula IV

15



20

in which R1, R2, R3 and R4 are each as defined in the novel compound of the formula II.

The compounds of the formulae I, II, III and IV are suitable as intermediates for the preparation of I κ B kinase inhibitors (WO 01/30774 A1).

25

The invention is illustrated in detail hereinafter with reference to examples. End products are generally determined by ¹H NMR (400 MHz, in DMSO-D₆); in each case, the main peak or the two main peaks are reported. Temperatures are reported in degrees Celsius; RT means room temperature (22°C to 26°C). Abbreviations used are either explained or


30

correspond to the usual conventions.

Example 1

Preparation of methyl 2-benzoylamino-3-diphenylaminoacrylate

5



66 g (266 mmol) of methyl 2-benzoylamino-3-dimethylaminoacrylate and 50 g (295 mmol) of diphenylamine were dissolved at 40°C in 1300 ml of isopropanol. The solution was admixed with 60 ml (725 mmol) of concentrated hydrochloric acid within 5 minutes (min) and stirred for a further 10 min. 550 ml of solvent were evaporated off under reduced pressure, the suspension was cooled to 10°C and the crystallized product was filtered off.


15 Yield: 83.5 g (84% of theory)

¹H NMR: 3.62 (s, 3H), 6.95-7.10 (m, 6H), 7.20-7.30 (m, 8H), 7.32-7.40 (m, 1H), 7.61 (s, 1H), 8.70 (s, 1H)

Example 2

Preparation of methyl 2-benzyloxycarbonylamino-3-diphenylaminoacrylate

20



36 g (129 mmol) of methyl 2-benzyloxycarbonylamino-3-dimethylaminoacrylate and 24.12 g (142 mmol) of diphenylamine were dissolved in 630 ml of isopropanol at 40°C. Subsequently, the solution was admixed with 17.4 ml of concentrated hydrochloric acid within 5 min and stirred at 40°C for a further 30 min. The reaction solution was concentrated

25


to 300 ml and admixed slowly with 300 ml of water. The crystallized product was filtered off with suction and dried at 40°C under reduced pressure.

Yield: 30.5 g (59% of theory)

¹H NMR: 3.62 (s, 3H), 4.68 (s, 2H), 6.95-7.10 (m, 6H), 7.20-7.50 (m, 9H), 7.61 (s, 1H)

Example 3

Preparation of racemic methyl 2-benzoylamino-3-diphenylaminopropionate



With exclusion of oxygen, an autoclave was charged with 1 g (2.68 mmol) of methyl 2-benzoylamino-3-diphenylaminoacrylate and 40 mg (0.042 mmol) of tris(triphenylphosphine)rhodium(I) chloride. After purging with argon, 40 ml of oxygen-free methanol were added. The autoclave was sealed gas-tight and the solution was hydrogenated at RT for 20 hours (h). The autoclave was decompressed and purged with nitrogen. The solvent was evaporated off under reduced pressure and the residue was chromatographed through a column filled with silica gel 60 (eluent: 1:1 ethyl acetate/heptane). After the solvents had been evaporated off under reduced pressure, a white solid remained, which was utilized for the formulation of a method and as a system test for the determination of the enantiomeric purity by HPLC on chiral phase.

HPLC column: Chiralpak OD 4×250

Eluent: 45:2:1 hexane/EtOH/MeOH + 0.1% diethylamine

Temperature: 30°C

Reactant retention time: 13.2 minutes

S-Isomer retention time: 11.8 minutes

R-Isomer retention time: 14.2 minutes


Yield: 0.5 g (50% of theory)

¹H NMR: 3.62 (s, 3H), 4.15-4.35 (m, 2H), 4.75-4.90 (m, 1H), 6.90-7.05

(m, 6H), 7.20-7.30 (m, 4H), 7.40-7.48 (m, 2H), 7.50-7.60 (m, 1H), 7.70-7.78 (d, 2H), 8.85 (d, 1H)

Example 4

5 Preparation of methyl (S)-2-benzoylamino-3-diphenylaminopropionate



With exclusion exclusion of oxygen and moisture, an ampule was charged
10 with 100 mg of methyl 2-benzoylamino-3-diphenylaminoacrylate
(0.255 mmol) and 1.9 mg (0.0026 mmol, 0.01 equivalent) of [(S,S)-Et-
FerroTANE-Rh]BF₄. After purging with argon, 5 ml of oxygen-free methanol
were added. The ampule was sealed gas-tight and hydrogenated in an
autoclave at 20 bar of hydrogen pressure for 24 h. The autoclave was
15 decompressed and purged with nitrogen. The conversion of the
hydrogenation was determined by HPLC. The enantioselectivity was
determined by HPLC on chiral phase by the method described in
Example 4. The [(R,R)-Et-FerroTANE-Rh]BF₄ catalyst afforded the
corresponding R derivative in the same yield and enantiomeric purity.
20 ee: 87%

Example 5

Preparation of methyl (S)- and (R)-2-benzoylamino- 3-diphenylaminopropionate

25 Analogously to Example 4, methyl 2-benzoylamino-
3-diphenylaminoacrylate was hydrogenated with various catalysts and
solvents. In the case of the catalysts which had not been prepared
beforehand, the active catalyst was prepared in situ from the optically
30 active phosphine ligand and equimolar amounts of [Rh(cod)Cl]₂ as the
rhodium(I) salt. The results are compiled in Table 1 below.

Table 1:

RCS	Catalyst/ligand	Rhodium salt	Pressure [bar]	Solvent	Conversion [%]	ee [%]
1:100	[(R,R)-Et-FerroTANE-Rh]BF ₄ [268220-96-8]	no	20	Toluene	97	90
1:100	[(R,R)-Et-FerroTANE-Rh]BF ₄ [268220-96-8]	no	20	Methanol	95	R 87
1:100	[(R,R)-Et-FerroTANE-Rh]BF ₄ [268220-96-8]	no	20	Dichloromethane	97	85
1:100	[(S,S)-Et-FerroTANE-Rh]BF ₄ [268220-96-8]	no	20	Methanol	94	S 86
1:100	[(R,R)-Me-DUPHOS-Rh]CF ₃ SO ₃ [136705-77-6]	no	20	Toluene	NC	nd
1:100	[(R,R)-Me-DUPHOS-Rh]CF ₃ SO ₃ [136705-77-6]	no	20	Methanol	71	90
1:100	[(R,R)-Me-DUPHOS-Rh]CF ₃ SO ₃ [136705-77-6]	no	20	Dichloromethane	65	86
1:100	(R)-(S)-JOSIPHOS [155806-35-2]	yes	20	Toluene	68	36
1:100	(R)-(S)-JOSIPHOS [155806-35-2]	yes	20	Methanol	91	31
1:100	(R)-(S)-JOSIPHOS [155806-35-2]	yes	20	Dichloromethane	82	11
1:100	L-BPPM-E [61478-28-2]	yes	20	Toluene	21	68
1:100	L-BPPM-E	yes	20			

RCS	Catalyst/ligand	Rhodium salt	Pressure [bar]	Solvent	Conversion [%]	ee [%]
	[61478-28-2]			Methanol	80	35
1:100	L-BPPM-E [61478-28-2]	yes	20	Dichloromethane	5	45
1:100	(S)-BINAPHANE [544461-38-3]	yes	20	Toluene	37	6
1:100	(S)-BINAPHANE [544461-38-3]	yes	20	Methanol	31	48
1:100	(S)-BINAPHANE [544461-38-3]	yes	20	Dichloromethane	19	46
1:100	(R)-(-)-tert-Ferro [155830-69-6]	yes	20	Methanol	59	99
1:100	(R)-(-)-Cyclohexyl-Ferro [167416-28-6]	yes	20	Methanol	11	99
1:100	(R,R)-BDPP [96183-46-9]	yes	20	Methanol	NC	nd
1:100	(S,S)-CHIRAPHOS [64896-28-2]	yes	20	Methanol	NC	nd
1:100	(R,R)-DIOP [32305-98-9]	yes	20	Methanol	5	95
1:100	(R)-PROPHOS [67884-32-6]	yes	20	Methanol	NC	nd
1:100	(S,S)-NORPHOS [71042-55-2]	yes	20	Methanol	NC	nd
1:100	(R,R)-iPr-DUPHOS [136705-65-2]	yes	20	Methanol	30	99
1:100	[(R,R)-Et-BPE [136705-62-9]	yes	20	Methanol	62	99
1:100	[(R,R)-Me-BPE- Rh]CF ₃ SO ₃ [213343-69-2]	no	20	Methanol	55	96
1:100	(R)-Me-BOPHOZ [406680-93-1]	yes	20	Methanol	NC	nd
1:100	MonoPhos	yes	20			

RCS	Catalyst/ligand	Rhodium salt	Pressure [bar]	Solvent	Conversion [%]	ee [%]
	[157488-65-8]			Methanol	NC	nd
1:100	MonoPhos [157488-65-8]	yes	20	Dichloromethane	NC	nd
1:100	MonoPhos [490023-37-5]	yes	20	Methanol	NC	nd
1:100	MonoPhos [490023-37-5]	yes	20	Dichloromethane	NC	nd
1:100	MonoPhos [380230-02-4]	yes	20	Methanol	NC	nd
1:100	MonoPhos [380230-02-4]	yes	20	Dichloromethane	NC	nd

RCS = Molar ratio of catalyst to substrate

NC = No conversion

nd = Not determined

- 5 R or S in the "ee [%]" column means the particular R or S enantiomer

Example 6

- 10 With exclusion of oxygen, an autoclave was charged with the amounts of methyl 2-benzoylamino-3-diphenylaminoacrylate and [(R,R)-Me-DUPHOS-Rh]CF₃SO₃ specified in Table 1. After purging with argon, the amount of oxygen-free methanol specified below was added. The autoclave was sealed gas-tight and the solution was hydrogenated at 30 bar of hydrogen pressure at RT for 20 h. The autoclave was decompressed and purged with
- 15 nitrogen. The conversion of the hydrogenation was determined by HPLC. The enantioselectivity was determined by HPLC on chiral phase by the method described in Example 4. The result is shown in Table 2.

Table 2:

20

Catalyst/substrate ratio	Catalyst	Reactant [g]	Conversion [%]	ee [%]
1:1000	[(R,R)-Me-DUPHOS-Rh]CF ₃ SO ₃	26	25	87

Example 7


With exclusion of oxygen, an autoclave was charged with the amounts of methyl 2-benzoylamino-3-diphenylaminoacrylate and [(S,S)-Et-FerroTANE-Rh]BF₄ specified in Table 2. After purging with argon, the amount of oxygen-free methanol specified below was added. The autoclave was sealed gas-tight and the solution was hydrogenated at 30 bar of hydrogen pressure at RT for 20 h. The autoclave was decompressed and purged with nitrogen. The solution was filtered, admixed with the same amount of water at 40°C and stirred at RT for 2 h. The crystallized product is filtered off with suction and dried to constant weight under reduced pressure at 45°C. The conversion of the hydrogenation was determined by HPLC. The enantioselectivity was determined by HPLC on chiral phase by the method described in Example 4. The result is shown in Table 3.

Table 3:

Catalyst/substrate ratio	Catalyst	Reactant [g]	Conversion [%]	ee [%]	Yield [%]
1:1000	[(S,S)-Et-FerroTANE-Rh]BF ₄ [268220-96-8]	26	98	85	89
1:2500	[(S,S)-Et-FerroTANE-Rh]BF ₄ [268220-96-8]	26	99	84	88
1:5000	[(S,S)-Et-FerroTANE-Rh]BF ₄ [268220-96-8]	26	98	86	90
1:10 000	[(S,S)-Et-FerroTANE-Rh]BF ₄ [268220-96-8]	26	73	85	n.i.
1:5000	[(S,S)-Et-FerroTANE-Rh]BF ₄ [268220-96-8]	260	98	85	89

n.i. = Not isolated


Example 8



- 5 With exclusion of oxygen, an autoclave was charged with 1 g (2.68 mmol) of methyl 2-benzyloxycarbonylamino-3-diphenylaminoacrylate and 40 mg (0.042 mmol) of tris(triphenylphosphine)rhodium(I) chloride. After purging with argon, 40 ml of oxygen-free methanol were added. The autoclave was sealed gas-tight and the solution was hydrogenated at RT for 20 h. The autoclave was decompressed and purged with nitrogen. The solvent was evaporated off under reduced pressure and the residue was purified by means of a column filled with silica gel 60 (eluent: 1:1 ethyl acetate/heptane). After the solvents had been evaporated off under reduced pressure, a white solid remained, which was utilized for the formulation of a method and as a system test for the determination of the enantiomeric purity by HPLC on chiral phase.

	HPLC column:	Chiralpak OD 4×250
	Eluent:	50:2:1 hexane/EtOH/MeOH + 0.1% diethylamine
20	Temperature:	30°C
	Reactant retention time:	19.2 minutes
	S-Isomer retention time:	14.6 minutes
	R-Isomer retention time:	16.0 minutes
	Yield:	0.2 g (20% of theory)
25	¹ H NMR:	3.60 (s, 3H), 3.95-4.15 (m, 2H), 4.35-4.45 (m, 1H), 4.92-5.05 (m, 2H), 6.90-7.00 (m, 6H), 7.15-7.40 (m, 9H), 7.85-7.90 (d, 1H)

Example 9



5 Analogously to Example 5, methyl 2-benzyloxycarbonylamino-3-diphenylaminoacrylate was hydrogenated with various catalysts and solvents. In the case of the catalysts which had not been prepared beforehand, the active catalyst was prepared in situ from the optically active phosphine ligand and equimolar amounts of $[\text{Rh}(\text{cod})\text{Cl}]_2$ as the rhodium(I) salt. The results are compiled in Table 4 below.

10 Table 4:

RCS	Catalyst/ligand	Rhodium salt	Pressure [bar]	Solvent	Conversion [%]	ee [%]
1:100	[(R,R)-Et-FerroTANE-Rh] BF_4 [268220-96-8]	no	20	Toluene	15	43
1:100	[(R,R)-Et-FerroTANE-Rh] BF_4 [268220-96-8]	no	20	Methanol	2	nd
1:100	[(R,R)-Et-FerroTANE-Rh] BF_4 [268220-96-8]	no	20	Dichloromethane	4	nd
1:100	[(R,R)-Me-DUPHOS-Rh] CF_3SO_3 [136705-77-6]	no	20	Toluene	15	nd
1:100	[(R,R)-Me-DUPHOS-Rh] CF_3SO_3 [136705-77-6]	no	20	Methanol	25	61

RCS	Catalyst/ligand	Rhodium salt	Pressure [bar]	Solvent	Conversion [%]	ee [%]
1:100	[(R,R)-Me-DUPHOS-Rh]CF ₃ SO ₃ [136705-77-6]	no	20	Dichloromethane	29	46
1:100	(R)-(S)-JOSIPHOS [155806-35-2]	yes	20	Toluene	13	nd
1:100	(R)-(S)-JOSIPHOS [155806-35-2]	yes	20	Methanol	35	28
1:100	(R)-(S)-JOSIPHOS [155806-35-2]	yes	20	Dichloromethane	17	43
1:100	L-BPPM-E [61478-28-2]	yes	20	Toluene	5	nd
1:100	L-BPPM-E [61478-28-2]	yes	20	Methanol	7	nd
1:100	L-BPPM-E [61478-28-2]	yes	20	Dichloromethane	6	nd
1:100	(S)-BINAPHANE [544461-38-3]	yes	20	Toluene	< 5	nd
1:100	(S)-BINAPHANE [544461-38-3]	yes	20	Methanol	< 5	nd
1:100	(S)-BINAPHANE [544461-38-3]	yes	20	Dichloromethane	< 5	nd
1:100	[(R,R)-Me-BPE-Rh]CF ₃ SO ₃	no	20	Methanol	36	73
1:100	(R)-(-)-tert-Ferro [155830-69-6]	yes	20	Methanol	85	21
1:100	(R)-(S)-JOSIPHOS	yes	20	Methanol	99	24
1:100	[(R,R)-Et-FerroTANE-Rh]BF ₄	yes	20	Methanol	52	60
1:100	(R,R)-BDPP [96183-46-9]	yes	20	Methanol	NC	nd
1:100	(S,S)-CHIRAPHOS [64896-28-2]	yes	20	Methanol	NC	nd


RCS	Catalyst/ligand	Rhodium salt	Pressure [bar]	Solvent	Conversion [%]	ee [%]
1:100	(R,R)-DIOP [32305-98-9]	yes	20	Methanol	NC	nd
1:100	(R)-PROPHOS [67884-32-6]	yes	20	Methanol	NC	nd
1:100	(S,S)-NORPHOS [71042-55-2]	yes	20	Methanol	NC	nd
1:100	[(R,R)-Et-BPE-Rh]BF ₄	yes	20	Methanol	6	23
1:100	(R)-Me-BOPHOZ (Eastman)	no	20	Methanol	25	16

RCS = Ratio of catalyst to substrate;
nd = Not determined;

NC = No conversion
< = Less than

5 Example 10

Preparation of methyl (S)-2-(benzoyl-tert-butoxycarbonylamino)-3-diphenylaminopropionate



10

18.7 g of methyl (S)-2-(benzoyl-tert-butoxycarbonylamino)-3-diphenylaminopropionate, ee = 85%, 20.6 of di-tert-butyl dicarbonate and 1.2 g of N,N-dimethylaminopyridine were dissolved in 90 ml of acetonitrile and stirred at 40°C for 3 hours. The acetonitrile was evaporated off under reduced pressure and the remaining residue was taken up in 300 ml of diisopropyl ether and hot-filtered. The product crystallized out overnight as a colorless solid.

15

Yield: 23.7 g (88% of theory)

¹H NMR: 1.38 (s, 9H), 3.70 (s, 3H), 4.35-4.58 (m, 2H), 5.45-5.52 (m,

1H), 6.93-7.05 (m, 6H), 7.13-7.18 (m, 2H), 7.22-7.30 (m, 4H),
7.32-7.30 (m, 2H), 7.45-7.52 (m, 1H)

Example 11

- 5 Preparation of methyl (S)-2-(tert-butoxycarbonylamino)-3-diphenylaminopropionate



- 10 1.3 g of methyl (S)-2-(benzoyl-tert-butoxycarbonylamino)-3-diphenylaminopropionate were dissolved in 13 ml of methanol and admixed with 2.74 ml of a 1M solution of magnesium methoxide in methanol. The solution was stirred at RT overnight and concentrated under reduced pressure. The residue was taken up in ethyl acetate and washed with water. The ethyl acetate phase was dried over sodium sulfate, filtered and concentrated under reduced pressure. The residue was crystallized from a little diisopropyl ether/heptane.


Yield: 0.95 g (90% of theory)

- 1H NMR: 1.38 (s, 9H), 3.55 (s, 3H), 4.10-4.25 (m, 2H), 4.50-4.62 (m, 1H), 5.10-5.25 (m, 1H), 6.90-7.05 (m, 6H), 7.20-7.30 (m, 4H)

Example 12

Preparation of methyl (S)-2-amino-3-diphenylaminopropionate p-toluenesulfonate

25



18.5 g of methyl (S)-2-(tert-butoxycarbonylamino)-


3-diphenylaminopropionate (ee = 85%) were dissolved in 100 ml of dichloromethane and with and admixed with 50 ml of trifluoroacetic acid (TFA). The solution was heated under reflux for 30 minutes and then concentrated to a volume of 100 ml under reduced pressure. The solution was washed with water and admixed with 9 g of p-toluenesulfonic acid. 125 ml of n-butanol were added and the remaining dichloromethane was evaporated off. To crystallize the p-toluenesulfonic salt, the solution was cooled to RT and stirred overnight. The solid was filtered off with suction and dried to constant weight under reduced pressure.

Yield: 16.6 g (81% of theory, based on desired isomer)
1H NMR: 2.38 (s, 3H), 3.30 (s, 3H), 4.10-4.35 (m, 3H), 5.45-5.52 (m, 1H), 6.73-6.95 (m, 6H), 7.01-7.05 (m, 2H), 7.10-7.18 (m, 4H), 7.58-7.62 (m, 2H), 8.30-8.55 (s, broad, 3H, NH)
ee: 99%

15

Example 13

Preparation of methyl 2-benzoylamino-3-phenylaminoacrylate



20


10 g (39.5 mmol) of methyl 2-benzoylamino-3-dimethylaminoacrylate and 11.1 g (118 mmol) of aniline were dissolved at 40°C in 200 ml of isopropanol. The solution was admixed with 3.6 ml (43.5 mmol) of concentrated hydrochloric acid within 5 minutes (min) and stirred for a further 10 min. 200 ml of deionized water were added, the suspension was cooled to 10°C and the crystallized product was filtered off.

Yield: 11.5 g (92% of theory)
1H NMR: 3.62 (s, 3H); 6.90-7.00 (m, 1H); 7.19 (d, 2H); 7.25-7.30 (m, 2H); 7.48-7.61 (m, 3H); 7.93 (d, 1H); 8.02 (d, 2H); 8.90 (d, 1H); 9.15 (s, 1H)

30

Example 14

Preparation of methyl 2-benzoylamino-3-(4-fluorophenylamino)acrylate




- 5 10 g (39.5 mmol) of methyl 2-benzoylamino-3-dimethylaminoacrylate and 11.4 ml (118 mmol) of 4-fluoroaniline were dissolved in 200 ml of isopropanol at 40°C. The solution was admixed with 3.6 ml (43.5 mmol) of concentrated hydrochloric acid within 5 minutes (min) and stirred for a further 30 min. The suspension was cooled to 10°C and the crystallized product was filtered off.

Yield: 12.4 g (94% of theory)

¹H NMR: 3.62 (s, 3H); 7.05-7.24 (m, 4H); 7.48-7.52 (m, 3H); 7.88 (d, 1H); 8.00-8.04 (m, 2H); 8.90 (d, 1H); 9.15 (s, 1H)

15 Example 15

Preparation of methyl 2-benzoylamino-3-(pyridin-2-ylamino)acrylate



- 20 10 g (39.5 mmol) of methyl 2-benzoylamino-3-dimethylaminoacrylate and 11.3 g (118 mmol) of 2-aminopyridine were dissolved in 200 ml of isopropanol at 40°C. The solution was admixed with 3.96 ml (48 mmol) of concentrated hydrochloric acid within 5 minutes (min) and stirred for a further 30 min. The suspension was cooled to 10°C and the crystallized product was filtered off.

Yield: 7.3 g (60% of theory)


¹H NMR: 3.62 (s, 3H); 6.92-6.97 (m, 1H); 7.02 (d, 1H); 7.45-7.70 (m, 4H); 8.02 (d, 2H); 8.22-8.24 (m, 1H); 8.60 (d, 1H); 9.22 (s,

1H); 9.45 (d, 1H)

Example 16

Preparation of methyl 2-benzoylamino-3-(thiazol-2-ylamino)acrylate

5




10 g (39.5 mmol) of methyl 2-benzoylamino-3-dimethylaminoacrylate and 11.3 g (118 mmol) of 2-aminothiazole were dissolved at 40°C in 200 ml of isopropanol. The solution was admixed with 3.96 ml (48 mmol) of concentrated hydrochloric acid within 5 minutes (min) and stirred for a further 60 min. 75 ml of deionized water were added, the suspension was cooled overnight and the crystallized product was filtered off.

Yield: 8.6 g (70% of theory)

15 ¹H NMR: 3.62 (s, 3H); 7.08 (d, 1H); 7.32 (d, 1H); 7.45-7.60 (m, 3H); 8.02 (d, 2H); 8.22 (d, 1H); 9.28 (s, 1H); 10.45 (d, 1H)

Claims

1. A compound of the formula II



where R1 and R2 are the same or different and are each independently

- 1) $-(C_6-C_{14})$ -aryl where aryl is unsubstituted or mono-, di- or trisubstituted independently by R11, where R11 is

- a) F, Cl, I or Br,
- b) $-(C_1-C_4)$ -alkyl,
- c) -CN,
- d) $-CF_3$,
- e) $-OR^5$ in which R^5 is a hydrogen atom or $-(C_1-C_4)$ -alkyl,
- f) $-N(R^5)-R^6$ in which R^5 and R^6 are each independently a hydrogen atom or $-(C_1-C_4)$ -alkyl,
- g) $-C(O)-R^5$ in which R^5 is a hydrogen atom or $-(C_1-C_4)$ -alkyl, or
- h) $-S(O)_x-R^5$ in which x is the integer zero, 1 or 2, and R^5 is a hydrogen atom or $-(C_1-C_4)$ -alkyl, or

- 2) a 4- to 15-membered Het ring where the Het ring is unsubstituted or mono-, di- or trisubstituted independently by $-(C_1-C_5)$ -alkyl, $-(C_1-C_5)$ -alkoxy, halogen, nitro, amino, trifluoromethyl, hydroxyl, hydroxy- $-(C_1-C_4)$ -alkyl, methylenedioxy, ethylenedioxy, formyl, acetyl, cyano, hydroxycarbonyl, aminocarbonyl or $-(C_1-C_4)$ -alkoxycarbonyl,


- R3 is 1) $-(C_6-C_{14})$ -aryl in which aryl is unsubstituted or mono-, di- or trisubstituted independently by $-NO_2$, $-O-(C_1-C_4)$ -alkyl, F, Cl or bromine,
- 2) $-O-C(CH_3)_3$, or
 - 3) $-O-CH(R7)$ -aryl in which aryl is unsubstituted or mono-,

di- or trisubstituted independently
by $-\text{NO}_2$, $-\text{O}-\text{CH}_3$, F, Cl or bromine,
in which R7 is a hydrogen atom or $-(\text{C}_1-\text{C}_4)\text{-alkyl}$,

- R4 is 1) a hydrogen atom,
2) $-(\text{C}_1-\text{C}_4)\text{-alkyl}$ or
3) $-\text{CH}(\text{R8})\text{-aryl}$

in which R8 is a hydrogen atom or $-(\text{C}_1-\text{C}_4)\text{-alkyl}$.

2. A compound of the formula IV




where R1, R2, R3 and R4 are each as defined as in the compound
of the formula II as claimed in claim 1.

5

10

15

3. A process for obtaining the compound of the formula I



20

where R1 is

- 1) a hydrogen atom,
- 2) $-(C_1-C_4)$ -alkyl,
- 3) $-(C_6-C_{14})$ -aryl where aryl is unsubstituted or mono-, di- or trisubstituted independently by R11,

25

where R11 is

- a) F, Cl, I or Br,
- b) $-(C_1-C_4)$ -alkyl,
- c) $-CN$,
- d) $-CF_3$,
- e) $-OR^5$ in which R^5 is a hydrogen atom or $-(C_1-C_4)$ -alkyl,
- f) $-N(R^5)-R^6$ in which R^5 and R^6 are each independently a hydrogen atom or $-(C_1-C_4)$ -alkyl,


30

- g) $-C(O)-R^5$ in which R^5 is a hydrogen atom or $-(C_1-C_4)$ -alkyl, or
- h) $-S(O)_x-R^5$ in which x is the integer zero, 1 or 2, and R^5 is a hydrogen atom or $-(C_1-C_4)$ -alkyl,
- 5 4) $-CH(R7)$ -aryl in which aryl is unsubstituted or mono-, di- or trisubstituted independently by $-NO_2$, $-O-CH_3$, F, Cl or bromine, where R7 is a hydrogen atom or $-(C_1-C_4)$ -alkyl, or
- 10 5) a 4- to 15-membered Het ring where the Het ring is unsubstituted or mono-, di- or trisubstituted independently by $-(C_1-C_5)$ -alkyl, $-(C_1-C_5)$ -alkoxy, halogen, nitro, amino, trifluoromethyl, hydroxyl, hydroxy- $-(C_1-C_4)$ -alkyl, methylenedioxy, ethylenedioxy, formyl, acetyl, cyano, hydroxycarbonyl, aminocarbonyl or $-(C_1-C_4)$ -alkoxycarbonyl,
- 15 R2 is 1) $-(C_6-C_{14})$ -aryl where aryl is unsubstituted or mono-, di- or trisubstituted independently by R11,
- 2) $-CH(R7)$ -aryl in which aryl is unsubstituted or mono-, di- or trisubstituted independently by $-NO_2$, $-O-CH_3$, F, Cl or bromine, where R7 is a
- 20 hydrogen atom or $-(C_1-C_4)$ -alkyl, or
- 3) a 4- to 15-membered Het ring where the Het ring is unsubstituted or mono-, di- or trisubstituted independently by $-(C_1-C_5)$ -alkyl, $-(C_1-C_5)$ -alkoxy, halogen, nitro, amino, trifluoromethyl, hydroxyl,
- 25 hydroxy- $-(C_1-C_4)$ -alkyl, methylenedioxy, ethylenedioxy, formyl, acetyl, cyano, hydroxycarbonyl, aminocarbonyl or $-(C_1-C_4)$ -alkoxycarbonyl,
- R3 is 1) a hydrogen atom,
- 2) $-(C_1-C_4)$ -alkyl,
- 30 3) $-(C_6-C_{14})$ -aryl in which aryl is unsubstituted or mono-, di- or trisubstituted independently by $-NO_2$, $-O-(C_1-C_4)$ -alkyl, F, Cl or bromine,
- 4) $-O-C(CH_3)_3$, or
- 5) $-O-CH(R7)$ -aryl in which aryl is unsubstituted or mono-, di- or trisubstituted independently
- 35 by $-NO_2$, $-O-CH_3$, F, Cl or bromine, where R7 is a hydrogen atom or $-(C_1-C_4)$ -alkyl,

- R4 is 1) a hydrogen atom,
 2) $-(C_1-C_4)$ -alkyl or
 3) $-CH(R_8)$ -aryl

in which R8 is a hydrogen atom or $-(C_1-C_4)$ -alkyl,

which comprises hydrogenating a compound of the formula II




in which R1, R2, R3 and R4 are each as defined in the formula I and the compound of the formula II may be present in the E or in the Z configuration on the double bond, in the presence of hydrogen and a catalyst.

4. The process as claimed in claim 3, wherein the compound of the formula I is obtained where R1 is phenyl or a hydrogen atom, R2 is phenyl, pyridyl or thiazolyl, in which phenyl, pyridyl or thiazolyl is unsubstituted or substituted by fluorine or chlorine, and R3 is phenyl or $-O-CH_2$ -phenyl and R4 is methyl or ethyl.
5. The process as claimed in claim 3 or 4, wherein the catalyst used comprises rhodium complexes, ruthenium complexes or iridium complexes or mixtures thereof.
6. The process as claimed in one or more of claims 3 to 5, wherein the molar ratio of catalyst to the compound of the formula II is from 1:100 to 1:10 000.
7. The process as claimed in one or more of claims 3 to 6, wherein the hydrogenation is performed at a temperature of from $10^{\circ}C$ to $200^{\circ}C$ and a hydrogen pressure of from 1 bar to 200 bar.


8. The process as claimed in one or more of claims 3 to 7, wherein the solvent used comprises water, alcohols such as methanol, ethanol, propanol or isopropanol, aromatic hydrocarbons such as toluene, ketones such as acetone, halogenated hydrocarbons such as dichloromethane, carboxylic esters such as ethyl acetate, and ethers such as tetrahydrofuran, or mixtures thereof.

9. A process for obtaining the compound of the formula III




or salts thereof in which R1, R2 and R4 are each as defined in the formula I as claimed in claim 3, which comprises

- a) hydrogenating the compound of the formula II



in which R1, R2, R3 and R4 are each as defined in the formula I, in the presence of hydrogen and a catalyst, and converting it to a compound of the formula I



and

- b) converting the resulting compound of the formula I to a compound of the formula III.

5


10. The process for obtaining the compound of the formula III as claimed in claim 9, which comprises

10

- a) hydrogenating the compound of the formula II in which R1, R2, R3 and R4 are each as defined in the formula I in the presence of hydrogen and a catalyst and converting it to a compound of the formula I,

15

- b) reacting the resulting compound of the formula I with a tert-butyl dicarbonate and an acylation catalyst such as N,N-dimethylaminopyridine to give a compound of the formula IV




20

in which R1, R2, R3 and R4 are each as defined in formula I,

and

- c) then converting the resulting compound of the formula IV to the compound of the formula Ia



5


and

- d) converting the resulting compound of the formula Ia to the compound of the formula III or salts thereof, in which R1, R2 and R4 are each as defined in formula I.

10

5

11. A process for obtaining the compound of the formula II as claimed in claim 1, which comprises reacting the compound of the formula VI




10

in which R3 and R4 are each as defined in formula II as claimed in claim 1

with an amine of the formula V

15



in which R1 and R2 are each as defined in formula II as claimed in claim 1 to give the compound of the formula II.

12. A process for obtaining the compound of the formula IV as claimed in claim 2, which comprises

20


- a) hydrogenating the compound of the formula II as claimed in claim 1,

in which R1, R2, R3 and R4 are each as defined in claim 2

in the presence of hydrogen and a catalyst and converting it to a compound of the formula I, and

25

- b) reacting the resulting compound of the formula I with a tert-butyl dicarbonate and an acylation catalyst such as dimethylaminopyridine to give a compound of the formula IV



in which R1, R2, R3 and R4 are each as defined in claim 2.