

FIG. 1






FIG. 2



Manufacturing Method of
Chondrocyte Mass (Continued)

FIG. 3A




FIG. 3B





FIG. 4



4 / 3 8

Manufacturing Method of Graft Material

FIG. 5




FIG. 6




FIG. 7




FIG. 8




FIG. 9




FIG. 10






FIG. 11A

Cartilage tissue



After 6 days of culturing

FIG. 11B



After 21 days of culturing

FIG. 12




FIG. 13A




FIG. 13B




FIG. 13C



FIG. 13D



FIG. 14A




FIG. 14B




FIG. 14C




FIG. 14D




FIG. 15




FIG. 16





FIG. 17






FIG. 18




FIG. 19




FIG. 20

FIG. 21




FIG. 22




FIG. 23




FIG. 24




FIG. 25




FIG. 26





FIG. 27



25/38

FIG. 28A



FIG. 28B




FIG. 29





FIG. 30



27/38

FIG. 31





FIG. 32








FIG. 33






FIG. 34A




FIG. 34B




FIG. 34C

FIG. 35




FIG. 36A




FIG. 36B




FIG. 36C





FIG. 37



Culturing time

2 hours

18 hours

FIG. 38A




FIG. 38B




FIG. 38C




FIG. 38D




FIG. 39




FIG. 40




FIG. 41




FIG. 42

