

PREPARATIONS CONTAINING METHYLSULFONYLMETHANE AND METHOD
OF PURIFICATION

The present invention relates to compositions for conditioning the skin, nails, other tissue and/or body fluids of a human or other animal subject. More specifically, it relates to compositions which can be used to soften, smooth, lubricate and preserve the pliancy of human tissue, for reducing the brittleness of finger and toe nails, and as diluents for blood.

There presently exists a number of substances which are known to provide softening and other benefits to the skin when applied topically. The most popular of these is lanolin which is safe for topical applications and is used widely in cosmetic preparations.

To have a substantial effect, however, lanolin must be applied to the skin frequently and in large quantities. Even then, the softening effect on skin is less than has been obtained using lesser amounts of other skin softening substances. Compositions containing lanolin in sufficient quantity can leave the skin with a greasy feeling and may stain clothing if not thoroughly absorbed by the skin.

Due to the disadvantages of lanolin as a skin softener, substantial efforts have been made to find suitable alternative substances for skin conditioning. Unfortunately, most of the substances discovered have been synthetically derived and are not naturally occurring substances. Such substances are medically suspect and in some instances produce side-effects which make them unsuitable for use in cosmetics

or other preparations for treating human tissue or body fluids.

One exception has been urea which is a naturally occurring substance in animals. Urea has been used with reasonable success for softening the skin and for treating skin diseases. Typically, the urea is applied in an aqueous solution or in the form of a cream. Such urea preparations have had only limited utility, however, because the urea tends to spontaneously decompose. As a result, preservatives for the urea must be added if such a skin treating composition is to have a reasonable shelf life.

The urea decomposition problem has been explored by a number of researchers. For example, Gunner P.E. Swanbeck has discovered various additives for stabilizing urea in a skin treating composition. In his U.S. Patent No. 3,666,863, lactic acid is suggested as a preservative for skin treating compositions containing carbamide. And, in British Patent Specification No. 1,411,432, the same inventor describes a skin treating composition containing urea in combination with one or more salts selected from the group including sodium chloride, sodium thiosulfate and sodium dyhydrogen phosphate.

Even stabilized urea compositions of the type described in the Swanbeck references have their limitations. Most particularly, the effectiveness of urea as a skin treating agent is somewhat less than desired.

Another substance proposed for use in skin treating compositions is dimethyl sulfoxide (DMSO). As described in

U.S. Patent No. 3,549,770, DMSO has utility in relieving signs and symptoms of pain and tissue inflammation and for promoting tissue repair of a subject having a skin graft or suffering from burns or tissue damage. But,

5 DMSO frequently causes adverse skin reactions so that DMSO has previously been considered unsuitable for use in cosmetics and other non-prescription preparations.

In general, it has been difficult to find substances which can be used to treat the tissues and body fluids of
10 animal subjects without creating adverse side-effects. For example, after much research, dextran has become a widely accepted extender for blood plasma and is commonly used as a blood diluent. Despite the wide acceptance of dextran, this substance has a higher and more variable molecular
15 weight than most normal blood constituents; and many subjects who receive the substance react allergically. Other blood diluents have even greater drawbacks; so there is no universally accepted blood diluent or plasma expander.

It has now been discovered that compositions containing
20 methylsulfonylmethane (MSM) can be used effectively to soften skin, to dilute blood, and for a variety of other useful purposes. MSM compositions are stable and safe for administration to human or other animal subjects and therefore eliminate many problems of the prior art. In
25 particular, topically applied MSM compositions benefit the skin to a far greater degree than lanolin or urea. And, MSM can be combined with lanolin and/or urea to form skin preparations which are exceptionally effective.

MSM has proved to have varied and useful properties when applied to any animal tissue subject to undesired chemical bond formation including cross-linking. It has been observed to beautify the complexion, to enhance scalp and hair, and generally to help make the body of the user more flexible and comfortable. Various MSM formulations are also exceedingly useful as bland vehicles for pharmaceuticals.

Manicuring preparations can advantageously include MSM. Such preparations can increase nail toughness by reducing brittleness and can be used for softening cuticle for easy removal.

Depending on its intended use, a preparation can contain MSM in solution or in a dispersion. It may take the form of a cream, lotion, gel or paste for topical administration or a liquid, for administration by injection,

The presence of MSM tends to stabilize compositions containing urea by inhibiting spontaneous urea decomposition. Similarly, urea stabilizes MSM in compositions so that MSM and urea are exceptionally well suited for use together.

Furthermore, DMSO may be included with certain MSM compositions to enhance their effectiveness. The adverse skin reactions frequently associated with use of DMSO, do not occur when certain of these MSM-DMSO compositions are used.

An aqueous composition, with the proper concentration of MSM, is an excellent diluent for blood. In such a

composition MSM is nonallergenic, nonpyretic and has no interfering or undesirable pharmacological effect, even when administered in relatively large amounts for a blood diluent. MSM has a far lower molecular weight yet remains in the body longer than dextran.

MSM is a naturally occurring substance found in the tissues and body fluids of at least all higher animals. It can be isolated from mammalian milk, e.g. cow's milk. It is also a naturally occurring excretory product in higher animals, such as man, where it is found in the urine.

MSM is of exceedingly low toxicity to all forms of life, plant as well as animal. Except for its beneficial effect on animal tissue, it appears to be totally inert to the diverse chemical reactions involved in the processes of life.

MSM used in cosmetic preparations and other preparations administered to tissue and/or body fluids of a human subject should be substantially pure. Accordingly an improved process for purification of MSM has been developed.

It is an object of this invention to provide a composition suitable for application to the tissue of a human or other animal subject, which composition is safe and effective in softening, smoothing and/or comforting the skin or other tissue.

Another object is to provide such a composition that is stable over a long period of time so that it can be practically sold for use as a cosmetic or consumer skin care treatment.

A further object is to provide MSM compositions in a variety of forms and to provide a variety of methods of

treatment with MSM in accord with the conditions or body parts to be treated. A particular related object is to provide such compositions in a viscous liquid form so that they can be easily applied topically and in an injectable liquid form,

It is also an object to provide a method of treatment to enhance moisture retention of the skin, relieve discomfort resulting from skin irritation and to soften, smooth and lubricate the skin.

Yet another object is to provide methods for treating connective tissue of a human or animal subject to resist cross-linking of collagen and other protein in the tissue and reduce dehydration or other alteration of cementing substrates such a hyaluronic acid.

An object is to provide a stable, neutral vehicle for pharmaceuticals, which vehicle has no interfering or undesirable pharmacological activity.

An additional object is to provide an effective composition and method for reducing the brittleness of finger and toe nails.

Also an object is to provide a method and composition for the softening of cuticle tissue so that it can be removed by gentle rubbing.

A specific object is to provide a safe and universally acceptable blood diluent.

Another object is to provide a method for purification of MSM to a purity at which it can be included in cosmetics

or other preparations for administration to the tissue
or body fluids of animal subjects.

According to the present invention there is provided
an aqueous solution suitable for topical application

5

comprising methylsulfonylmethane and at least one
member selected from urea, dextrose and sodium chloride.
The methylsulfonylmethane is preferably present in an amount
of from 1 to 50 weight percent of the solution.

10

In a preferred embodiment, the methylsulfonylmethane
presence is controlled to approximate the oncotic pressure
of the animal recipient of said solution.

15

When urea is present in the aqueous solution same is
preferably present in an amount of from 1 to 20 weight percent
of the solution.

The invention also provides an aqueous solution, suitable
as a blood diluent for warm blooded animals, comprising
about 20 percent methylsulfonylmethane, about 5 percent
of dextrose and the remainder water.

20

Also in accordance with the present invention there is
provided a cosmetic composition in the form of liquid, lotion,
cream or gel comprising a skin softening amount of methyl-
sulfonylmethane. Urea may be included in such cosmetic
composition preferably in an amount of from at least a
promotional amount for said methylsulfonylmethane to 20 weight
percent of the cosmetic composition. In a preferred
arrangement, the methylsulfonylmethane and urea are present

25

in about the same amount of from 1 to 10 weight percent.

Also in accordance with the present invention there is provided a nail-polish composition suitable for application to human finger-nails, said composition
5 containing about 5 weight percent of methylsulfonylmethane.

Methylsulfonylmethane (MSM) is an active agent which can be used safely and effectively for treating human or other animal subjects.

10 Advantageously, MSM can be included in a cosmetic or other preparations applied to the skin. Such compositions when applied topically, can beautify the complexion, improve the condition of the scalp and hair and help to make the body of the user more flexible and comfortable. Liquid
15 cosmetic preparations can include MSM in aqueous or nonaqueous solutions or emulsions.

Compositions containing MSM may be in the form of an injectable liquid, For topical

application, the composition will preferably be a solution or emulsion containing thickeners suitable to form a cream, lotion or gel. MSM can be administered in a mixture, e.g. a mixture of MSM crystals and peanut oil,-

or in a dry compound of body powders.

Certain aqueous MSM compositions can be administered intravenously as nontoxic, non-allergenic blood diluents and modifiers of internal tissues.

A substantial advantage of MSM, regardless of how or why it is administered, is the noninteraction of MSM with the diverse chemical reactions involved in the process of life. Except for its inhibiting effect on cross-linking, MSM appears to be inert in the tissue and body fluids of living subjects.

Because of its inertness, MSM can be combined with non-irritating solvents and/or emulsifiers to provide excellent vehicles for pharmaceutical substances. Such vehicles deliver the pharmaceutical and simultaneously improve the condition of the subject's connective tissue. Inertness further makes MSM a superior substance for formulating aqueous blood diluent solutions as will be further described below.

To formulate MSM compositions, it is important to consider the physical properties of MSM. Commercially available MSM

is an odourless, white, crystalline substance with a melting point of $109^{\circ}\text{C}.$, and a boiling point of $238^{\circ}\text{C}.$ It tends to sublime with heating. MSM has a molecular weight of 94.13, specific gravity of about 1.45 and an index of a refraction of 1.42 at $20^{\circ}\text{C}.$

The solubility characteristics of MSM are broad and somewhat unusual. These characteristics must be considered in preparing and using MSM compositions according to the present invention. MSM is highly soluble in hot water, but poorly soluble near the freezing point of water. For example, it is possible to dissolve over 1000g. of MSM in 100g. of water at $88^{\circ}\text{C}.$, while only about 33.9g. of MSM will dissolve in 100g. of water at $26^{\circ}\text{C}.$ MSM is soluble in a wide range of organic solvents and itself solvates many organic and inorganic compounds.

Preparations containing MSM in aqueous solution should generally contain only about 20 weight percent MSM and preferably no more than about 15 weight percent. Solutions containing MSM in greater concentrations can be formulated by using other pharmaceutically acceptable solvents e.g. glycerine and/or ethanol. High concentration aqueous MSM compositions, suitable for certain specific purposes, can also be made by dissolving crystalline MSM in heated water immediately prior to application.

MSM also works well in emulsions such as those formed with a polymer of acrylic acid cross-linked with a poly-functional agent. One suitable polymer is Carbopol 934P(RM)

manufactured by B.F. Goodrich Chemical Company, Cleveland, Ohio. Other suitable emulsions can be formulated using modified cellulose polymers such as hydroxyethylcellulose.

When MSM is to be administered to human or other animal subjects it should be present in a composition which is pharmaceutically acceptable for the intended route of administration. In other words, the composition should not contain MSM in so great a concentration that it would be injurious; and the composition should not include unnecessarily toxic substances or substances which react with MSM to produce undesired substances.

a. MSM Administered to Connective Tissue

The reason why MSM softens or otherwise benefits animal skin, are not fully understood. It is believed that a normal part of the ageing process is the progressive cross-linking or bonding of protein such as collagen in animal connective tissue, especially skin. Such cross-links (possibly covalent, hydrophobic or hydrogen bonds and/or van der Waals associations) stiffen and harden the tissue. It appears that MSM acts to soften and preserve the pliancy of connective tissue by inhibiting bond structing of connective tissue, protein and other components in the tissue, and possibly by attacking existing cross-links. Thus, the benefits observed when MSM is used may result from reductions in dehydration of body substances such as hyaluronic acid and elastin, or from beneficial effects of MSM on fibrinogen.

The amount of MSM to be included in a particular composition may vary greatly depending upon the intended use of the composition. When treating the naturally soft and pliant skin of infants, a composition containing even minute quantities of MSM may be beneficial when applied to the skin. When treating the connective tissue of adult subjects, MSM should comprise at least about one weight percent of the composition administered. Disproportionately superior results are achieved using compositions containing at least about five weight percent MSM.

The maximum concentration of MSM in compositions for topical application is about 15 weight percent, preferably only about 10 weight percent. Greater amounts of MSM could, however, be used if the composition is heated and mixed prior to topical application.

The following examples are provided to illustrate the activity of MSM in softening, smoothing and otherwise inhibiting the ageing of skin and other animal tissue and to show the non-interactivity which makes MSM suitable for use in vehicles for pharmaceutical substances and in blood diluents. The examples are not to be taken as an exhaustive list of the possible methods for using MSM and the benefits resulting from its use. The examples do, however, give some indication of the broad scope of this invention, specifically the suitability of MSM for a variety of purposes.

Example 1

To determine the effectiveness of MSM in preserving the pliancy of animal connective tissue, young rabbits were sacrificed and bled. A portion of the vascular beds of these animals were removed and divided into three groups for testing. Those in the first group were infiltrated with a 10 weight percent solution of MSM in water. Those of the second group with an aqueous solution containing 10 weight percent each of MSM and urea. The third group was infiltrated with an isotonic saline solution.

After four hours, sections of the veins and arteries from the beds of each group were removed and immersed in a standard formaldehyde-based fixative hardener solution of the type used to preserve tissue for histological study.

After immersion for 20 minutes, the veins and arteries were removed from the hardening solution and examined. No hardening of the tissue was observed in the veins and arteries from the first two experimental groups. Vein and artery tissue from vascular beds of the third group displayed substantial hardening. From these results it appears that the MSM, with or without urea, prevented hardening of vein and artery tissue, apparently by blocking formaldehyde-induced cross-linking in connective tissue.

Example 2

A series of tests were conducted to determine the effect of MSM on a hydrolyzed collagen representative of the collagen which is found in the exterior layer of animal skin and which hardens with dehydration.

Five aqueous solutions of MSM were prepared, the solutions containing 2,4,6,8 and 10 weight percent of MSM, respectively. Each of the solutions was heated and combined with a sample of modified collagen at a weight ratio of nine to one. Normally, collagen combined with heated water will swell and form physical and chemical bonds to form a gel. However, when the mixtures of this experiment cooled to room temperature, no gelation had occurred in the mixtures containing 8 and 10 weight percent MSM solutions.

b. MSM and Urea Administered to Connective Tissue

A further discovery is that the addition of urea to an MSM composition enhances the effectiveness of the composition as a tissue treating agent, beyond expectations. Although urea is known to have some beneficial effects when applied to the skin, it is not believed that urea alone has any effect in inhibiting chemical bond linking of collagen.

As will best be understood from the following examples, compositions containing both MSM and urea have a much greater activity in inhibiting bond structuring of substances in connective tissue than compositions containing either one of the substances alone. It appears that the presence of urea accelerates tissue penetration of MSM. Furthermore, experimental results show increased benefits beyond those expected to result from accelerated MSM penetration alone.

In MSM compositions for application to the tissue of an animal subject, even small amounts of urea may be helpful in improving the overall softness of the skin.

The composition becomes especially effective, however, when urea is present in an amount effective to multiply the activity of the MSM. Such a composition will contain at least about one weight percent urea and preferably at least about 5 weight percent.

Example 3

The procedure of Example 2 was repeated twice. In the first repetition, 2, 4, 6, 8 and 10 weight percent aqueous urea solutions, instead of MSM solutions, were added to five collagen samples. When the collagen samples had cooled to room temperature, it was observed that urea had no effect in reducing collagen cross-linking in any of the mixtures.

In the second repetition, aqueous solutions containing equal weights of both MSM and urea were mixed with the collagen samples. A first solution contained 2 weight percent each MSM and urea, a second 4 weight percent each, a third 6 weight percent each, a fourth 8 weight percent each; and a fifth solution contained 10 weight percent of each.

When cooled to room temperature none of the collagen mixtures formed a gel except for the mixture containing first solution which had only 2 weight percent each of MSM and urea. This is in contrast with the procedure of Example 2 which showed no cross-linking inhibition using MSM solutions of less than 8 weight percent and the first repetition which showed no cross-linking inhibition by solutions containing urea alone.

5 Because cross-linking of collagen can so effectively be reduced by the application of MSM, and that the reduction can be increased by the addition of urea, such compositions can be administered to animal tissue to counter the cross-linking of collagen which is associated with ageing and various disease processes. It specifically appears that the softness and pliancy associated with youthful, healthy skin thus can be maintained and that unhealthy, abused, old skin at least partially can be restored.

10

Example 4

15

Sections of human cadaver skin were brought to a constant weight in a 50% relative humidity oven. The sections were divided into four groups. Each group of sections was immersed in a different aqueous solution as set forth in the following table:

20

<u>Group</u>	<u>Solution (aqueous)</u>	
1	MSM	10 wt. percent
2	Urea	10 wt. percent
3	MSM	10 wt. percent
	Urea	10 wt. percent
4	Isotonic saline	

25

In each case, the sections were immersed for two hours, blotter dried and then stabilized at 50% relative humidity.

The cadaver skin, which can be expected to behave similarly to the outer layers of the epidermis of a healthy human, was found to retain less than 15% moisture when treated with the isotonic saline (Group 4) solution and then stabilized. The skin treated with solution containing MSM alone (Group 2) was substantially more flexible and

retained 30-35% moisture. The highest moisture levels and greatest flexibility was observed in the sections which were treated with both MSM and urea (Group 3). The sections treated with the urea solution (Group 2) showed some improved moisture retention, but less than that of the Group 3 sections.

As this example illustrates, MSM may serve as an additive for embalming fluid to maintain tissue pliancy.

Example 5

A variety of test applications were made to determine the effectiveness of compositions containing MSM and urea in treating human subjects having skin with a thickened stratum corneum.

A first solution containing 10 grams MSM, 45 grams water and 45 grams ethanol was applied three times daily to a human subject having skin with a thickened stratum corneum. After applications three times daily for a period of one week, a substantial softening and smoothing of the skin was observed.

A second solution containing 10 grams urea, 45 grams water and 45 grams ethanol was applied to a human subject having skin with a thickened stratum corneum. After one week, small improvements in skin pliancy were observed; but there was no increase in softness or smoothness of the skin.

A third solution containing 5 grams MSM, 5 grams urea, 45 grams water and 45 grams ethanol was applied to several human subjects having skin with a thickened stratum corneum. After a one week period of application three times a day, the skin of these subjects was exceptionally soft and smooth in the areas of application. Even though the composition contained

only 5 weight percent MSM, the extent of the physiological effect resulting from the application of this combined solution was greater than the effect which resulted from application of the first two solutions containing 10 weight percent MSM and 10 weight percent urea, respectively.

Example 6

To determine the amount of MSM and urea needed to produce a desired effect, a series of tests on healthy skin of human subjects was performed. Eleven separate compositions were formulated containing different amounts of MSM and urea. As a solvent, each composition included a liquid comprising equal amounts, by volume, of water and ethanol.

Compositions were applied daily to separate subjects. And, the date was recorded for each subject when an increase in skin softness and smoothness was first detected. The results of this testing were compiled and appear in the following table:

	<u>Composition</u>		<u>Result</u>
	<u>Wt. % MSM</u>	<u>Wt. % Urea</u>	
20	1	1	Benefit seen in three weeks
	1	2	Benefit seen in three weeks
	2	2	Benefit seen in three weeks
	3	3	Benefit seen in two weeks
	4	4	Benefit seen in one week
25	5	5	Benefit seen in less than one week
	6	6	Benefit seen in less than one week
	7	7	Benefit seen in less than one week

8	8	Benefit seen in less than one week
9	9	Benefit seen in less than one week
10	10	Benefit seen in less than one week

As can be seen from the table, softness and smoothness were improved when solutions containing even low amounts of MSM and urea were applied over a period of time. Specifically, it was found that a composition produced desirable effects at concentrations of MSM and urea of as low as 1 weight percent each. Rapid improvement in skin softness and smoothness were observed when the solution contained 5 or more weight percent each of MSM and urea.

Example 7

Compositions containing MSM and urea improve the softness and pliability of skin even of persons suffering from adverse skin conditions. In one test, two human subjects suffering from "hide bound disease" or progressive systemic sclerosis were poorly responsive to standard treatments for softening the skin and making hands and feet more comfortable.

For treatment of the subjects, a preparation containing 20 weight percent MSM, 20 weight percent urea, 30 weight percent diamethyl sulfoxide and 30 weight percent water was prepared. A solution was formed by heating to about 40°C.

The subjects were treated by placing 15 milliliters of the solution in a plastic bag, placing a hand or foot to be treated in the bag. The hand or foot with plastic bag overwrap was then immersed in a heated water bath maintained at a temperature as warm as the subject would tolerate, taking

care not to dilute the solution.

5 The hands and feet were thus immersed for 30 minutes, 3 times daily, for a period of 2 weeks. The result was a reduction in discomfort and an increased skin softness and pliancy.

Dimethyl sulfoxide was used in solution to enhance penetration of MSM and urea into the effected tissue. No adverse side-effects resulted from administration of the DMSO.

Example 8

10 Properly formulated compositions containing MSM and urea can be applied to skin which is injured or inflamed without causing adverse reactions.

A gel containing MSM and urea was made according to the following formulation:

15	H ₂ O	780 grams
	urea	80 grams
	MSM	50 grams
	Carbowax 600(PEG-12)	50 grams
	Glycerin	20 grams
20	Carbopol 940 (carbomer-940)	4 grams
	Na ₂ CO ₃ (in 10 ml. of H ₂ O)	0.25 grams

25 This formulation was applied to the skin of subjects having a variety of adverse skin conditions including diaper rash, abrasions, wind burn, thermal burn and skin tears from berry bushes. In none of these applications did the subjects experience instantaneous discomfort of the type frequently encountered when medical preparations were applied. In a number of instances the preparation was soothing to the effected skin and provided a protective coating for the

affected area.

Example 9

It is found that MSM and urea can be used effectively in more complex compositions, such as the creams, lotions and gels which are most preferred for cosmetics preparations. One such formulation, a gel containing Carbopol 940 as a thickening agent, was made by combining the following ingredients:

	H ₂ O	796 grams
10	urea	80 grams
	MSM	50 grams
	Carbowax 600 (PEG-12)	50 grams
	Glycerin	20 grams
	Carbopol 940 (carbomer-940)	4 grams
15	2,2',2"-nitrilotriethanol	1.75 grams
	KOH	0.25 grams

Fifty human subjects applied the formulation to the skin of the right hand and foot for three weeks on a morning and night (twice daily) routine. The gel was applied in sufficient amount to form a white matt film over the right hand and foot after which the excess gel was washed off with water. The subjects were requested to abstain from using other cosmetic preparations during the test period.

After one week, the evaluators noted a substantial improvement in skin softness, smoothness and comfort. Seven of the subjects continued to apply the formulation to the right hand and foot for a second week. At the end of this

- 22 -

period, further improvements in softness, smoothness and comfort were observed.

Example 10

5 MSM, alone or with urea, can also be added to a variety of cosmetic preparations other than skin lotions, creams and gels. For instance, a shampoo was formulated with the following ingredients:

	H ₂ O	800 grams
	MSM	100 grams
10	Urea	100 grams
	Carbowax 400 (PEG-8)	50 grams
	Sodium lauryl sulfate	100 grams
	Ammonium lauryl sulfate	75 grams
	Cocamide MEA	5 grams

15 This shampoo formulation was tested both by subjects with healthy hair and scalp and by others with problem seborrheic dermatitis. Both groups found the shampoo to be effective, to leave the hair easily manageable and to soften the scalp. Those subjects having a dandruff problem
20 found there was a reduction in itching, scaling and scalp inflammation after only four to five washings.

Example 11

A hair styling gel including MSM and urea was prepared according to the following formulation:

25	H ₂ O (with added ammonia)	76 grams
	Polyvinylpyrrolidone (PVP) 45% aqueous solution	6.0 grams
	Carbopol 940 (carbomer-940)	0.5 grams
	MSM	5 grams

Urea	5 grams
Ethanol	2 grams
Carbowax 600 (PEG-12)	5 grams

When applied to the hair of human subjects, this composition proved to be more effective in hair management than a comparable commercial product. Several subjects, having persistent itching scalp problem, observed a reduction in the itching. In two subjects dandruff was reduced.

Example 12

Subjects used the gel described in Example 8 as a pre-shave preparation with good results. If applied before retiring, or even minutes before shaving (using a commercial shaving cream), subjects obtained a more comfortable smoother, and cleaner shave, using a blade razor, than if the commercial shaving cream was used alone. It also appeared that drag of the razor was reduced.

When the gel was used, leg hair was easily shaved without a commercial shaving cream overcoat. In all instances, use of the formulation left the skin feeling softer and smoother.

Example 13

A nail conditioner was formulated from the following ingredients:

H ₂ O	50 wt. percent
MSM	10 wt. percent
Urea	10 wt. percent
Glycerin	5 wt. percent
Dimethyl sulfoxide	20 wt. percent
Glyoxal (30% aqueous)	5 wt. percent

344

The formulation was applied with cotton pads to the nails of human subjects and allowed to remain for at least 15 minutes. At the end of that time, the nails were toughened, i.e. less brittle, and the cuticle was softened such that it could be removed by gentle rubbing.

It is not fully understood how the brittleness of the nail is reduced by application of the formulation. It appears, however, that the MSM may inhibit directional cross-linking in the nail. It may even break down naturally formed cross-links that are the cause of undesirably rigid structures responsible for nail brittleness.

The above composition is enhanced by the presence of 20 wt. percent dimethylsulfoxide (DMSO) which accelerates penetration of the MSM and urea into the nail. Subjects using this formulation had no reddening or irritation of the skin surrounding the nail. DMSO is an irritant to the epidermis of many people, so that the absence of adverse skin reactions was unexpected.

Example 14

Nail polishes containing MSM were formulated by preparing 25 weight percent MSM solutions in ethyl or amyl acetate. Such solutions were mixed with the commercial nail polishes of several well-known U.S. manufacturers in such a ratio that the resulting mixtures contained about 5 weight percent MSM.

Nails of subjects using the resulting polish mixtures became less brittle with repeated applications. Also, the reformulated polishes were more easily removed with

commercial nail polish removers than the original commercial polishes.

Example 15

Reductions in nail brittleness were also observed when commercial nail polish removers, reformulated with MSM, were used repeatedly. Commercial nail polish removers containing esters, ketones and alcohols were combined with MSM in such amounts that the resulting mixtures contained 5-10 weight percent MSM. These proved to be beneficial in reducing nail brittleness.

c. MSM Administered Orally

Example 16

To determine whether living animal subjects would react adversely to orally administered compositions of MSM, a 40 weight percent solution of MSM in distilled water was prepared. This solution was administered orally to laboratory rats at such a rate that each rat received 20 grams of MSM per kilogram of body weight per day.

After six weeks of administration, none of the animals had died or displayed unusual symptoms or behaviour.

d. MSM in Vehicle Compositions

Because MSM is substantially inert to the chemistry of the body, compositions containing this substance are excellent vehicles for pharmaceuticals. When applied, the MSM will produce the desirable results mentioned above. MSM does not react with common pharmaceutical substances. And, as discussed below, MSM can actually stabilize certain substances such as carbamide so that the shelf life of certain pharmaceutical preparations can be extended by the inclusion of MSM.

Example 17

The formulation of Example 9 was evaluated as a potential vehicle for agents known to have beneficial cosmetic, skin treating and/or insect repelling activities. A variety of compositions were made using the formulation of Example 9 as a vehicle.

In each instance, a 150 gram sample of the formulation described in Example 9 was combined, in a household, high sheer mixer, with 10 grams of an active agent and the resulting combination thoroughly mixed. Combined formulations were made in this manner with each of the following agents:

- | | |
|-----------------------------|-------------------------------|
| propyl palmitate | ethyl aminobenzoate |
| hydrated lanolin | (10% solution) |
| USP mineral oil | iodine (10% solution) |
| olive oil | nicotinic acid (10% solution) |
| peanut oil | fluorouracil (2% solution) |
| thiabendazole solution | vitamin A and vitamin D |
| (3% in DMSO) | concentrates |
| erythromycin phosphate | disodium edetate |
| (10% in aqueous DMSO) | junipar tar (10% solution) |
| testosterone (5% solution) | meta delphene (50% alcohol |
| triamcinolone (5% solution) | solution) |

All of the compositions formed were stable. When applied to the skin, each composition produced the desirable tissue softening effects described in Example 9 without reducing the effectiveness of the agent. From this sampling it is logical to conclude that formulation of Example 9 could be very beneficial as a vehicle for a wide variety of pharmaceutical and cosmetic substances.

e. MSM-Carbamide Stabilization

An unexpected discovery resulting from experimentation with the above preparations was that compositions containing both MSM and urea are more stable than similar compositions containing MSM or urea alone, apparently because MSM and

urea stabilize one another. MSM is thus useful as a stabilizer in almost any composition which contains the urea subject to spontaneous decomposition and vice versa.

Example 18

5 To demonstrate the ability of MSM to inhibit the spontaneous decomposition of urea, laboratory tests were conducted. In these tests, the gel described in Example 8 was compared with a control formula that was an identical gel except that it contained no MSM. Multiple samples
10 of each formula were placed in plastic coated metal tubes that were sealed and capped.

Tubes containing each formulation, were subjected to successive freeze-thaw cycles. The formulation without MSM began to break down to a liquid after two to three such
15 cycles; while no break down of the composition including MSM was observed after ten freeze-thaw cycles.

Other tubes were maintained for two weeks in a dry oven at a temperature of 75°C. At the end of this time, gas pressure was elevated in the tubes containing formula
20 without MSM such that the tubes were bloated and hissed upon opening. Gas in these tubes was determined, by odour, to contain ammonia which indicates that break down of the carbamide occurred. In tubes containing the formulation with MSM, no such gas pressure or ammonia odour was
25 observed. MSM thus stabilized urea in the composition of Example 8 under accelerated ageing conditions.

f. MSM in Blood Diluents

As discussed above, test applications of MSM have demonstrated only positive effects on human or other animal tissue. Its physical characteristics and inertness in normal bodily chemical reactions make MSM ideal for use in formulating diluents for body fluids such as blood. It is especially suitable as a plasma extender because subjects do not react allergically to MSM as they do when dextran is used.

An MSM composition for use as blood diluent should be formulated as a solution which will create substantially the same oncotic pressure as normal plasma of the blood to be diluted. In such compositions, which primarily contain water and MSM, the MSM should comprise between 0.1 and 50 weight percent of the composition. Use of MSM amounts at the low end of this range presumes the presence of other dissolved substances which, together with the MSM, cause the solution to have substantially the same oncotic pressure as the body fluid to be diluted.

Example 19

To determine the safety of intravenous administrations of MSM compositions for blood dilution and/or systemic treatment of collagen related diseases, MSM was dissolved in a standard 5 percent dextrose solution so that the resulting liquid composition contained 20 weight percent MSM. The composition was delivered intravenously to a dog weighing 9.5 kilograms at such a rate that the subject received 2 grams of MSM per kilogram of body weight per day.

39

The subject did not react adversely to the intravenous administrations and did not require physical restraint. After several weeks of this treatment there was no evidence of physical discomfort or injury to the subject.

5

g. Purification of MSM

A simple procedure may be used to obtain MSM of sufficient purity for use in compositions for treating the tissues of human or other animal subjects. MSM is combined with heated water to form a solution, the temperature of the water determining the amount of MSM which can be dissolved for purification. Activated charcoal is then added to the solution along with a substantially water insoluble base, such as aluminium or magnesium hydroxide, in an amount effective to neutralize the solution. The charcoal and base are then mechanically separated from the solution and the solution cooled to a temperature where purified MSM crystallizes. In most instances, crystals form when a solution is cooled to 10°C.; but enhanced crystallization occurs when the solution is cooled to below 5°C. The purified MSM crystals are then mechanically separated from the water. The remaining water still contains some dissolved MSM. Losses of MSM can thus be minimized by recycling the water for use as a solvent to purify additional MSM crystals according to this process.

10

15

20

25

Example 20

In one such purification procedure, 400 grams of unpurified MSM were dissolved in 800 grams of heated

water to form a solution. To the heated solution were added 10 grams of activated charcoal and four grams of magnesium hydroxide. The solution was then allowed to stand, with stirring, for 0.5 hours and then was filtered while still hot to remove the activated charcoal and magnesium hydroxide. The remaining solution, containing dissolved MSM, was chilled to below 10°C. to crystallize the MSM. MSM crystals were then separated by filtration and the resulting filtrate was heated and re-used to dissolve additional MSM to be purified by the same process.

MSM produced by this process was extremely pure and thus suitable for use in any appropriate composition to be applied to human tissue or body fluids.



CLAIMS

1. An aqueous solution suitable for topical application comprising methylsulfonylmethane and at least one member selected from urea, dextrose and sodium chloride.
2. An aqueous solution as claimed in claim 1, wherein said
5 methylsulfonylmethane is present in an amount of from 1 to 50 weight percent of said solution.
3. An aqueous solution as claimed in claim 1 or 2, wherein said urea is present in an amount of from 1 to 20 weight percent of said solution.
- 10 4. An aqueous solution as claimed in any one of claims 1 to 3 substantially as hereinbefore described and exemplified.
5. An aqueous solution, suitable as a blood diluent for warm blooded animals, comprising about 20% methylsulfonylmethane, about 5% dextrose, and the remainder water.
- 15 6. An aqueous solution as claimed in claim 5, wherein said methylsulfonylmethane presence is controlled to approximate the oncotic pressure of the animal recipient of said solution.
7. A cosmetic composition in the form of a liquid, lotion,
20 cream or gel comprising a skin softening amount of methylsulfonylmethane.
8. A cosmetic composition as claimed in claim 7, wherein urea is a component in an amount from at least a promotional amount for said methylsulfonylmethane to 20 weight percent
25 of said cosmetic composition.



9. A cosmetic composition as claimed in claim 8, wherein said methylsulfonylmethane and said urea are present in about the same amount of from 1 to 10 weight percent.
10. A cosmetic composition as claimed in claim 7, 8 or 9 substantially as hereinbefore described and exemplified.
11. A nail polish composition suitable for application to human finger nails, containing about 5 weight percent of methylsulfonylmethane.
12. A nail polish composition as claimed in claim 11, substantially as hereinbefore described and exemplified.
13. A cosmetic gel, suitable for application to the human skin, consisting of:

		<u>Approx. Weight %</u>
	Methylsulfonylmethane	5.
15	Urea	8.
	Glycerin	2.
	Polyethylene Glycol	5.
	Carbovinyl polymer	0.4
	Sodium carbonate	Trace
20	Water	Remainder

The text of the specification has been reproduced by photocopying the applicants original typescript. It may contain a few amendments which are difficult to read. The original typescript containing these amendments may be inspected on the premises of the Patent Office