CLAIMS

[Claim 1]

A leucine-rich α2 glycoprotein composition comprising leucine-rich α2 glycoprotein, which is a calibration sample solution for measuring the leucine-rich α2 glycoprotein and is filled in a preservation container,

wherein a pH of the composition is 7.0 to 9.3~~.~~,

wherein a concentration of a salt of the composition is 200 mM to 800 mM, and

wherein the salt is selected from the group consisting of sodium salts, potassium salts, and magnesium salts.

~~[Claim 2]~~

~~The leucine-rich α2 glycoprotein composition according to claim 1, which is a calibration sample solution for measuring the leucine-rich α2 glycoprotein.~~

~~[Claim 3]~~

~~The leucine-rich α2 glycoprotein composition according to claim 1 or 2, which is filled in a preservation container.~~

[Claim ~~4~~2]

The leucine-rich α2 glycoprotein composition according to ~~any one of claims 1 to 3~~claim 1, wherein the pH of the composition is 7.8 to 9.0.

~~[Claim 5]~~

~~The leucine-rich α2 glycoprotein composition according~~

~~to any one of claims 1 to 4, wherein a concentration of a salt of the composition is 200 mM to 800 mM.~~

[Claim ~~6~~3]

The leucine-rich α2 glycoprotein composition according to claim ~~5~~1 or 2, wherein the sodium salt is ~~a metal salt~~sodium chloride.

[Claim ~~7~~4]

A method for measuring leucine-rich α2 glycoprotein comprising:

using the leucine-rich α2 glycoprotein composition according to any one of claims 1 to ~~6~~3.

[Claim ~~8~~5]

A kit for measuring leucine-rich α2 glycoprotein comprising the leucine-rich α2 glycoprotein composition according to any one of claims 1 to ~~6~~3.

[Claim ~~9~~6]

A preservation stabilizing method for leucine-rich α2 glycoprotein comprising:

bringing leucine-rich α2 glycoprotein into contact with a solvent having a pH of 7.0 to 9.3~~.~~,

wherein a concentration of a salt in the solvent is 200 mM to 800 mM, and

wherein the salt is selected from the group consisting of sodium salts, potassium salts, and magnesium salts.

[Claim ~~10~~7]

The preservation stabilizing method for leucine-rich α2 glycoprotein according to claim ~~9~~6, wherein the pH of the solvent is 7.8 to 9.0.

~~[Claim 11]~~

~~The preservation stabilizing method for leucine-rich α2 glycoprotein according to claim 9 or 10, wherein a concentration of a salt in the solvent is 200 mM to 800 mM.~~

[Claim ~~12~~8]

The preservation stabilizing method for leucine-rich α2 glycoprotein according to ~~any one of claims 9 to 11~~claim 6 or 7, wherein the sodium salt is ~~a metal salt~~sodium chloride.