



166166

TITLE OF THE INVENTION**16-SUBSTITUTED-4-AZA-ANDROSTANE 5-ALPHA-REDUCTASE
ISOZYME 1 INHIBITORS****FIELD OF THE INVENTION**

The present invention provides novel compounds, novel compositions, methods of their use and methods of their manufacture, where such compounds are generally pharmacologically useful as agents in therapies whose mechanism of action rely on the selective inhibition of the isozyme 5 α -reductase 1.

BACKGROUND OF THE INVENTION

Certain undesirable physiological manifestations, such as acne vulgaris, seborrhea, female hirsutism, androgenetic alopecia which includes female and male pattern baldness, and benign prostatic hyperplasia, are the result of hyper-androgenic stimulation caused by an excessive accumulation of testosterone ("T") or similar androgenic hormones in the metabolic system. Early attempts to provide a chemotherapeutic agent to counter the undesirable results of hyperandrogenicity resulted in the discovery of several steroidal antiandrogens having undesirable hormonal activities of their own. The estrogens, for example, not only counteract the effect of the androgens but have a feminizing effect as well. Non-steroidal antiandrogens have also been developed, for example, 4'-nitro-3'-trifluoromethylisobutyranilide. See Neri, *et al.*, *Endocrinol.* 1972, 91 (2). However, these products, though devoid of hormonal effects, compete with all natural androgens for receptor sites, and hence have a tendency to feminize a male host or the male fetus of a female host and/or initiate feed-back effects which would cause hyperstimulation of the testes.

The principal mediator of androgenic activity in some target organs, e.g., the prostate, is 5 α -dihydrotestosterone ("DHT"), formed locally in the target organ by the action of testosterone-5 α -reductase (or simply 5 α -reductase). Inhibitors of 5 α -reductase will serve to prevent or lessen symptoms of hyperandrogenic stimulation in these organs. See



G00002

- 2 -


especially United States Patent Nos. 4,377,584, issued March 22, 1983, and 4,760,071, issued July 26, 1988, both assigned to Merck & Co., Inc. It is now known that a second 5 α -reductase isozyme exists, which interacts with skin tissues, especially in scalp tissues. See, e.g., G. Harris, *et al.*, *Proc. Natl. Acad. Sci. USA*, Vol. 89, pp. 10787-10791 (Nov. 1992). The isozyme that principally interacts in skin tissues is conventionally designated as 5 α -reductase 1 (or 5 α -reductase type 1), while the isozyme that principally interacts within the prostatic tissues is designated as 5 α -reductase 2 (or 5 α -reductase type 2).

In the treatment of androgenic sensitive disease conditions, e.g., benign prostatic hyperplasia (BPH) and/or the prevention and treatment of prostatic cancer, it would be desirable to have one drug entity which is active against both isozymes in the prostate to significantly inhibit all dihydrotestosterone production. It would also be desirable to have another drug entity which is highly selective for inhibiting the isozyme 5 α -reductase 1 associated with the scalp, for use in treating conditions of the skin and scalp, e.g., acne vulgaris, male pattern baldness and hirsutism in females. Additionally, a selective 5 α -reductase 1 inhibitor could be used in combination with a 5 α -reductase 2 inhibitor such as, e.g., finasteride (PROSCAR[®]), for therapy in the treatment of hyperandrogenic conditions such as BPH and/or the prevention and treatment of prostatic cancer, and for the treatment of skin and scalp-related disorders such as acne vulgaris, seborrhea, female hirsutism, and androgenic alopecia. Still further, the 5 α -reductase 1 inhibitors of this invention could be used in combination with a potassium channel opener such as minoxidil for the treatment of these skin and scalp-related disorders. Therefore it is an object of this invention to provide compounds that have sufficient activity in the inhibition of 5 α -reductase isozyme 1.

- 3 -

SUMMARY OF THE INVENTION


The novel compounds of this invention have the formula



and are selective 5 α -reductase 1 inhibitors. It is an object of this invention to provide compounds that alone or in combination with inhibitors of 5 α -reductase 2 are useful in the treatment of benign prostatic hyperplasia, prostatitis, and/or the prevention and treatment of prostatic cancer. It is an additional object of this invention to provide compounds that alone or in combination with inhibitors of 5 α -reductase 2 are useful in the treatment of acne vulgaris, female hirsutism, androgenic alopecia (also known as androgenetic alopecia and human pattern baldness), and insufficient plasma levels of high density lipoproteins. The compounds of the invention have utility in one or more of the aforementioned areas.

DETAILED DESCRIPTION OF THE INVENTION

The novel compounds of the present invention have the general structural Formula I:



or a pharmaceutically acceptable salt or ester thereof wherein:

- 4 -

the C1-C2 carbon-carbon bond may be a single bond, or a double bond as indicated by the dashed line;

R^1 is selected from the group consisting of hydrogen and C1-10 alkyl;

R^2 is selected from the group consisting of hydrogen and C1-10 alkyl;

one of R^3 and R^4 is selected from the group consisting of hydrogen and methyl, and the other is selected from the group consisting of:

- (a) amino;
- (b) cyano;
- (c) fluoro;
- (d) methyl;
- (e) OH;
- (f) $-C(O)NR_bR_c$, where R_b and R_c are independently H, C1-6 alkyl, aryl, or arylC1-6alkyl; wherein the alkyl moiety can be substituted with 1-3 of: halo; C1-4alkoxy; or trifluoromethyl; and the aryl moiety can be substituted with 1-3 of: halo; C1-4alkyl; C1-4 alkoxy; or trifluoromethyl;
- (g) C1-10 alkyl-X-;
- (h) C2-10 alkenyl-X-;

wherein the C1-10 alkyl in (g) and C2-10 alkenyl in (h) can be unsubstituted or substituted with one to three of:

- i) halo; hydroxy; cyano; nitro; mono-, di- or trihalomethyl; oxo; hydroxysulfonyl; carboxy;
- ii) hydroxyC1-6alkyl; C1-6alkyloxy; C1-6 alkylthio; C1-6alkylsulfonyl; C1-6 alkyloxycarbonyl; in which the C1-6 alkyl moiety can be further substituted with 1-3 of: halo; C1-4 alkoxy; or trifluoromethyl;
- iii) arylthio; aryl; aryloxy; arylsulfonyl; aryloxycarbonyl; in which the aryl moiety can be further substituted with 1-3 of: halo; C1-4 alkyl; C1-4 alkoxy; or trifluoromethyl;
- iv) $-C(O)NR_bR_c$; $-N(R_b)-C(O)-R_c$; $-NR_bR_c$; where R_b and R_c are defined above;

- 5 -

- (i) aryl-X-;
- (j) heteroaryl-X-, wherein heteroaryl is a 5, 6 or 7 membered heteroaromatic ring containing at least one member selected from the group consisting of: one ring oxygen atom, one ring sulfur atom, 1-4 ring nitrogen atoms, or combinations thereof; in which the heteroaromatic ring can also be fused with one benzo or heteroaromatic ring;
wherein the aryl in (i) and heteroaryl in (j) can be unsubstituted or substituted with one to three of:
 - v) halo; hydroxy; cyano; nitro; mono-, di- or trihalomethyl; mono-, di- or trihalomethoxy; C2-6 alkenyl; C3-6 cycloalkyl; formyl; hydrosulfonyl; carboxy; ureido;
 - vi) C1-6 alkyl; hydroxy C1-6 alkyl; C1-6 alkyloxy; C1-6 alkyloxy C1-6alkyl; C1-6 alkylcarbonyl; C1-6 alkylsulfonyl; C1-6 alkylthio; C1-6 alkylsulfinyl; C1-6 alkylsulfonamido; C1-6 alkylarylsulfonamido; C1-6 alkyloxy-carbonyl; C1-6 alkyloxycarbonyl C1-6alkyl; $R_bR_cN-C(O)-C1-6alkyl$; C1-6 alkanoylamino C1-6 alkyl; aroylamino C1-6 alkyl; wherein the C1-6 alkyl moiety can be substituted with 1-3 of: halo; C1-4alkoxy; or trifluoromethyl;
 - vii) aryl; aryloxy; arylcarbonyl; arylthio; arylsulfonyl; arylsulfinyl; arylsulfonamido; aryloxycarbonyl; wherein the aryl moiety can be substituted with 1-3 of: halo; C1-4alkyl; C1-4alkoxy; or trifluoromethyl;
 - viii) $-C(O)NR_bR_c$; $-O-C(O)-NR_bR_c$; $-N(R_b)-C(O)-R_c$; $-NR_bR_c$; $R_b-C(O)-N(R_c)-$; where R_b and R_c are defined in (f) above; and $-N(R_b)-C(O)-OR_g$, wherein R_g is C1-6alkyl or aryl, in which the alkyl moiety can be substituted with 1-3 of: halo; C1-4alkoxy; or trifluoromethyl, and the

- 6 -

aryl moiety can be substituted with 1-3 of: halo; C₁-4alkyl; C₁-4 alkoxy, or trifluoromethyl; -N(R_b)-C(O)NR_cR_d, wherein R_d is selected from H, C₁-6 alkyl, and aryl; in which said C₁-6alkyl and aryl can be substituted as described above in (f) for R_b and R_c;

- ix) a heterocyclic group, which is a 5, 6 or 7 membered ring, containing at least one member selected from the group consisting of: one ring oxygen atom, one ring sulfur atom, 1-4 ring nitrogen atoms, or combinations thereof; in which the heterocyclic ring can be aromatic, unsaturated, or saturated, wherein the heterocyclic ring can be fused with a benzo ring, and wherein said heterocyclic ring can be substituted with one to three substituents, as defined above for v), vi), vii) and viii), excluding ix) a heterocyclic group; and
- (k) R³ and R⁴ taken together can be carbonyl oxygen;
- (l) R³ and R⁴ taken together can be =CH-R_g, wherein R_g is defined in viii); and wherein:

X is selected from the group consisting of:

-O-; -S(O)_n-; -C(O)-; -CH(R_e)-; -C(O)-O-*; -C(O)-N(R_e)-*;
 -N(R_e)-C(O)-O-*; -O-C(O)-N(R_e)-*; -N(R_e)C(O)-N(R_e)-;
 -O-CH(R_e)-*; -N(R_e)-; wherein R_e is H, C₁-3 alkyl, aryl, aryl-C₁-3 alkyl, or unsubstituted or substituted heteroaryl, as defined above in (j);

wherein the asterisk (*) denotes the bond which is attached to the 16-position in Structure I; and n is zero, 1 or 2.

Combinations of substituents and/or variables are permissible only if such combinations result in stable compounds.

- 7 -

In one embodiment of the instant invention are compounds of Formula I wherein R^1 is hydrogen or methyl and R^2 is hydrogen or methyl.

A further embodiment of the instant invention are compounds of Formula I wherein:
one of R^3 and R^4 is selected from the group consisting of hydrogen and methyl, and the other is selected from the group consisting of:

- (b) cyano;
- (c) fluoro;
- (e) OH;
- (g) C₁-10 alkyl-X-; or C₁-10 alkyl-X-, where alkyl can be substituted with aryl, and wherein aryl in turn can be substituted with 1-2 of halo or C₁-6alkyl;
- (h) C₂-10 alkenyl-X-;
- (i) aryl-X-;
- (j) heteroaryl-X-, wherein heteroaryl is a 5 or 6 membered heteroaromatic ring containing 1-2 ring nitrogen atoms; wherein the aryl in (i) and heteroaryl in (j) can be unsubstituted or substituted with one to two of:
 - x) halo; cyano; nitro; trihalomethyl; trihalomethoxy; C₁-6 alkyl; aryl; C₁-6 alkylsulfonyl; C₁-6 alkyl-arylsulfonamino;
 - xi) -NR_bR_c; R_b-C(O)-N(R_c)-; wherein R_b and R_c are independently H, C₁-6 alkyl, aryl, or arylC₁-6alkyl; wherein the alkyl moiety can be substituted with 1-3 of: halo; C₁-4alkoxy; or trifluoromethyl; and the aryl moiety can be substituted with 1-3 of: halo; C₁-4alkyl; C₁-4 alkoxy; or trifluoromethyl;
 - xii) a heterocyclic group, which is a 5 membered aromatic ring, containing one ring nitrogen atom, or one ring oxygen and one ring nitrogen atom; and
- (k) wherein R^3 and R^4 taken together can be carbonyl oxygen; and wherein:

- 8 -

X is selected from the group consisting of:

-O-; -S(O)_n-; -CH(R_e)-; -C(O)-N(R_e)-*;

-O-C(O)-N(R_e)-*;

wherein R_e is H, C₁-3 alkyl, aryl, aryl C₁-3 alkyl;

wherein the asterisk (*) denotes the bond which is attached to the 16-position in Structure I; and n is zero or 2.

Novel compounds of the present invention exemplified by this embodiment include but are not limited to the following compounds:

4-aza-4,7β-dimethyl-5α-androstane-3,16-dione;

4-aza-4-methyl-5α-androstan-3,16-dione;

3-oxo-4-aza-4-methyl-16β-hydroxy-5α-androstane;

3-oxo-4-aza-4-methyl-16β-(benzylaminocarbonyloxy)-5α-androstane;

3-oxo-4-aza-4-methyl-16β-benzoylamino-5α-androstane;

3-oxo-4-aza-4-methyl-16β-methoxy-5α-androstane;

3-oxo-4-aza-4-methyl-16β-allyloxy-5α-androstane;

3-oxo-4-aza-4-methyl-16β-(n-propyloxy)-5α-androstane;

3-oxo-4-aza-4-methyl-16α-hydroxy-5α-androstane;

3-oxo-4-aza-4-methyl-16β-(phenoxy)-5α-androstane;

3-oxo-4-aza-7β-methyl-16β-(phenoxy)-5α-androst-1-ene;

3-oxo-4-aza-4-methyl-16α-methoxy-5α-androstane;

3-oxo-4-aza-4-methyl-16β-(4-chlorophenoxy)-5α-androstane;

3-oxo-4-aza-7β-methyl-16β-(4-chlorophenoxy)-5α-androst-1-ene;

3-oxo-4-aza-7β-methyl-16β-(4-chlorophenoxy)-5α-androstane;

3-oxo-4-aza-7β-methyl-16β-(3-chloro-4-methylphenoxy)-5α-androstane;

3-oxo-4-aza-7β-methyl-16β-(4-methylphenoxy)-5α-androstane;

3-oxo-4-aza-7β-methyl-16β-(4-methylphenoxy)-5α-androst-1-ene;

3-oxo-4-aza-7β-methyl-16β-[4-(1-pyrrolyl)phenoxy]-5α-androst-1-ene;

3-oxo-4-aza-4,7β-dimethyl-16β-hydroxy-5α-androstane;

3-oxo-4-aza-4,7β-dimethyl-16β-methoxy-5α-androstane;

3-oxo-4-aza-4,7β-dimethyl-16β-allyloxy-5α-androstane;

3-oxo-4-aza-4,7β-dimethyl-16β-(3,3-dimethylallyloxy)-5α-androstane;

3-oxo-4-aza-4,7β-dimethyl-16β-(n-propyloxy)-5α-androstane;

3-oxo-4-aza-4,7β-dimethyl-16β-(iso-pentyloxy)-5α-androstane;

3-oxo-4-aza-4,16α-dimethyl-16β-hydroxy-5α-androstane;

3-oxo-4-aza-4,7 β -dimethyl-16 β -ethyloxy-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -benzyloxy-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 α -hydroxy-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -methylthio-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(n-propylthio)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -fluoro-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -cyano-5 α -androstane;
3-oxo-4-aza-4-methyl-16 β -(1-hexyl)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(n-propyl)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -benzyl-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-chlorobenzyl)-5 α -androstane;
3-oxo-4-aza-4,16 α -dimethyl-16 β -methoxy-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-cyanophenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(3-cyanophenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-nitrophenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(1-naphthyloxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(3-chloro-4-methylphenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-methylphenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(tert-butyloxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(3-methyl-1-butyloxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 α -(n-propyloxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-trifluoromethylphenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-trifluoromethoxyphenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -ethylthio-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -ethylsulfonyl-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-methylsulfonylphenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -[4-(4-tolylsulfonylamino)phenoxy]-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(3-pyridyloxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -[(4-phenyl)phenoxy]-5 α -androstane;

3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-fluorophenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(2-pyrazinyloxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -[4-(5-oxazolyl)phenoxy]-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(2-pyrimidinyloxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -[4-(1-pyrryl)phenoxy]-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-aminophenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-acetylaminoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-benzoylaminoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-chlorophenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(phenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(2-chlorophenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(3-chlorophenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-chlorophenoxy)-5 α -androst-1-ene;
3-oxo-4-aza-4,7 β -dimethyl-16-(4-chlorobenzylidene)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16-benzylidene-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16-(4-methylbenzylidene)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16-(4-chlorobenzyl)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16-(4-methylbenzyl)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16-(3-pyridylmethyl)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 α -methanesulfonyl-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -thiophenoxy-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-chlorothiophenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-fluorothiophenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-methylthiophenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-methoxythiophenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -phenylsulfinyl-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -phenylsulfonyl-5 α -androstane;
3-oxo-4-aza-4,7 β ,16 α -trimethyl-16 β -(4-trifluoromethylphenoxy)-5 α -androstane,
3-oxo-4-aza-4,7 β ,16 α -trimethyl-16 β -hydroxy-5 α -androstane;
3-oxo-4-aza-4,7 β ,16 α -trimethyl-16 β -methoxy-5 α -androstane;

- 11 -

pharmaceutically acceptable salts thereof, and analogs of the above-described compounds wherein the C1-C2 carbon-carbon bond is a double bond, and/or R¹ is -H, and/or R² is -H or methyl, where appropriate.

In another embodiment of this invention are compounds of Formula I further limited to those wherein the C1-C2 carbon-carbon bond is a single bond, R¹ is methyl, R² is methyl, R³ is selected from unsubstituted or substituted aryloxy, and R⁴ is hydrogen.

Some non-limiting examples of compounds within this embodiment are:

3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-cyanophenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(3-cyanophenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-nitrophenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(1-naphthyloxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(3-chloro-4-methylphenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-methylphenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-trifluoromethylphenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-trifluoromethoxyphenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-methylsulfonylphenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -[4-(4-tolylsulfonylamino)phenoxy]-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -[(4-phenyl)phenoxy]-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-fluorophenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -[4-(5-oxazolyl)phenoxy]-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -[4-(1-pyrryl)phenoxy]-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-aminophenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-acetylaminophenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-benzoylaminophenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-chlorophenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(phenoxy)-5 α -androstane;

- 12 -

3-oxo-4-aza-4,7 β -dimethyl-16 β -(2-chlorophenoxy)-5 α -androstan-3-one;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(3-chlorophenoxy)-5 α -androstan-3-one;
and the pharmaceutically acceptable salts thereof.

A useful compound of the present invention is 3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-chlorophenoxy)-5 α -androstan-3-one, or a pharmaceutically acceptable salt thereof.

As used herein "alkyl" is intended to include both branched and straight-chain saturated aliphatic hydrocarbon groups having the specified number of carbon atoms, e.g., methyl (Me), ethyl (Et), propyl, butyl, pentyl, hexyl, heptyl, octyl, nonyl, decyl, iso-propyl (i-Pr), isobutyl (i-Bu), tert-butyl (t-Bu), sec-butyl (s-Bu), iso-pentyl, and the like. "Alkyloxy" (or "alkoxy") represents an alkyl group having the indicated number of carbon atoms attached through an oxygen bridge, e.g., methoxy, ethoxy, propyloxy, and the like. "Alkenyl" is intended to include hydrocarbon groups of either a straight or branched configuration with one or more carbon-carbon double bonds which may occur in any stable point along the chain, such as ethenyl, propenyl or allyl, butenyl, pentenyl, and the like. Included in this invention are all E, Z diastereomers.

The alkyl and alkenyl groups can be unsubstituted or substituted with one or more, and preferably 1-3, of:

- i) halo; hydroxy; cyano; nitro; mono-, di- or trihalomethyl; oxo; hydroxysulfonyl; carboxy;
- ii) hydroxyC₁₋₆alkyl; C₁₋₆alkyloxy; C₁₋₆ alkylthio; C₁₋₆alkylsulfonyl; C₁₋₆ alkyloxycarbonyl; in which the C₁₋₆ alkyl moiety can be further substituted with 1-3 of: halo; C₁₋₄ alkoxy; or trifluoromethyl;
- iii) arylthio; aryl; aryloxy; arylsulfonyl; aryloxycarbonyl; in which the aryl moiety can be further substituted with 1-3 of: halo; C₁₋₄ alkyl; C₁₋₄ alkoxy; or trifluoromethyl;
- iv) -C(O)NR_bR_c; -N(R_b)-C(O)-R_c; -NR_bR_c; where R_b and R_c are defined above; and halo is F, Cl, Br or I.

The term "oxo", as used herein, indicates an oxo radical which can occur in any stable point along the carbon chain resulting in a

- 13 -

formyl group, if at the end of the chain, or an acyl or aroyl group at other points along the carbon chain.

As used herein the term "aryl", i.e., C₆-10 aryl, is intended to mean phenyl or naphthyl, including 1-naphthyl or 2-naphthyl, either unsubstituted or substituted as described below.

The term "heteroaryl" as used herein, is intended to include a 5, 6 or 7 membered heteroaromatic radical containing at least one member selected from the group consisting of: one ring oxygen atom, one ring sulfur atom, 1-4 ring nitrogen atoms, or combinations thereof; in which the heteroaryl ring can also be fused with one benzo or heteroaromatic ring. This category includes the following either unsubstituted or substituted heteroaromatic rings (as described below): pyridyl, furyl, pyrrol, thienyl, isothiazolyl, imidazolyl, benzimidazolyl, tetrazolyl, pyrazinyl, pyrimidyl, quinolyl, quinazolinyl, isoquinolyl, benzofuryl, isobenzofuryl, benzothienyl, pyrazolyl, indolyl, isoindolyl, purinyl, carbazolyl, isoxazolyl, thiazolyl, isothiazolyl, oxazolyl, benzthiazolyl, and benzoxazolyl. The heteroaryl ring may be attached within structural Formula I by a heteroatom, e.g., N, or carbon atom in the ring, which results in the creation of a stable structure. The heteroaryl ring can also be fused to a benzo ring.

The one to three, and more usefully one to two substituents which can be on the C₆-10 aryl and heteroaryl groups named above are independently selected from:

v) halo; hydroxy; cyano; nitro; mono-, di- or trihalomethyl; mono-, di- or trihalomethoxy; C₂-6 alkenyl; C₃-6 cycloalkyl; formyl; hydrosulfonyl; carboxy; ureido;

vi) C₁-6 alkyl; hydroxy C₁-6 alkyl; C₁-6 alkyloxy; C₁-6 alkyloxy C₁-6 alkyl; C₁-6 alkylcarbonyl; C₁-6 alkylsulfonyl; C₁-6 alkylthio; C₁-6 alkylsulfinyl; C₁-6 alkylsulfonamido; C₁-6 alkylarylsulfonamido; C₁-6 alkyloxy-carbonyl; C₁-6 alkyloxycarbonyl C₁-6 alkyl; R_bR_cN-C(O)-C₁-6 alkyl; C₁-6 alkanoylamino C₁-6 alkyl; aroylamino C₁-6 alkyl; wherein the C₁-6 alkyl moiety can be substituted with 1-3 of: halo; C₁-4alkoxy; or trifluoromethyl;

- 14 -

vii) aryl; aryloxy; arylcarbonyl; arylthio; arylsulfonyl; arylsulfinyl; arylsulfonamido; aryloxy carbonyl; wherein the aryl moiety can be substituted with 1-3 of: halo; C₁-4alkyl; C₁-4alkoxy; or trifluoromethyl;

viii) -C(O)NR_bR_c; -O-C(O)-NR_bR_c; -N(R_b)-C(O)-R_c; -NR_bR_c; R_b-C(O)-N(R_c)-; where R_b and R_c are defined in (e) above; and -N(R_b)-C(O)-OR_c, wherein this instance R_c is C₁-6alkyl or aryl; -N(R_b)-C(O)NR_cR_d, wherein R_d is selected from H, C₁-6 alkyl, and aryl; in which said C₁-6alkyl and aryl can be substituted as described above in (e) for R_b and R_c;

ix) a heterocyclic group, which is a 5, 6 or 7 membered ring, containing at least one member selected from the group consisting of: one ring oxygen atom, one ring sulfur atom, 1-4 ring nitrogen atoms, or combinations thereof; in which the heterocyclic ring can be aromatic, unsaturated, or saturated, and wherein the heterocyclic ring can be fused with a benzo ring, and wherein said heterocyclic ring can be substituted with one to three substituents, as defined above for v), vi), vii) and viii), excluding ix) a heterocyclic group.

The fused heteroaromatic ring systems include: purine, imidazoimidazole, imidazothiazole, pyridopyrimidine, pyridopyridazine, pyrimidopyrimidine, imidazopyridazine, pyrrolopyridine, imidazopyridine, and the like.

The "heterocyclic" group includes the fully unsaturated heteroaryl rings described above and also their respective dihydro, tetrahydro and hexahydro derivatives resulting in partially unsaturated and fully saturated versions of the ring systems. Examples include: dihydroimidazolyl, dihydrooxazolyl, dihydropyridyl, tetrahydrofuryl, dihydropyrryl, tetrahydrothienyl, dihydroisothiazolyl, 1,2-dihydrobenzimidazolyl, 1,2-dihydrotetrazolyl, 1,2-dihydropyrazinyl, 1,2-dihydropyrimidyl, 1,2-dihydroquinolyl, 1,2,3,4-tetrahydroisoquinolyl, 1,2,3,4-tetrahydrobenzofuryl, 1,2,3,4-tetrahydroisobenzofuryl, 1,2,3,4-tetrahydrobenzothienyl, 1,2,3,4-tetrahydropyrazolyl, 1,2,3,4-tetrahydroindolyl, 1,2,3,4-tetrahydroisoindolyl, 1,2,3,4-tetrahydropuriny, 1,2,3,4-

- 15 -

tetrahydrocarbazolyl, 1,2,3,4-tetrahydroisoxazolyl, 1,2,3,4-tetrahydrothiazolyl, 1,2,3,4-tetrahydrooxazolyl, 1,2,3,4-tetrahydrobenzthiazolyl, and 1,2,3,4-tetrahydrobenzoxazolyl. and the like.

The heterocyclic group can be substituted in the same fashion as described above for heteroaryl.

Whenever the terms "alkyl", "alkenyl", "alkyloxy (or alkoxy)", "aryl" or "heteroaryl", or one of their prefix roots, appear in a name of a substituent in Formula I, (e.g., aralkoxyaryloxy) they shall have the same definitions as those described above for "alkyl", "alkenyl", "alkyloxy (or alkoxy)", "aryl" and "heteroaryl", respectively. Designated numbers of carbon atoms (e.g., C₁₋₁₀) shall refer independently to the number of carbon atoms in an alkyl or alkenyl moiety or to the alkyl or alkenyl portion of a larger substituent in which alkyl or alkenyl appears as its prefix root.

Also included within the scope of this invention are pharmaceutically acceptable salts of the compounds of Formula I, where a basic or acidic group is present on the structure. When an acidic substituent is present, e.g., -COOH, there can be formed the ammonium, sodium, potassium, calcium salt, and the like, for use as the dosage form. Where a basic group is present, i.e., amino or a basic heteroaryl radical such as, e.g., 4-pyridyl, an acidic salt, i.e., hydrochloride, hydrobromide, acetate, pamoate, and the like, can be used as the dosage form.

Also, in the case of the -COOH group being present, pharmaceutically acceptable esters can be employed, e.g., C₁₋₅ alkyl, pivaloyloxymethyl, and the like, and those esters known in the art for modifying solubility or hydrolysis characteristics for use as sustained release or prodrug formulations.

Representative salts include the following salts: acetate, lactobionate, benzenesulfonate, laurate, benzoate, malate, bicarbonate, maleate, bisulfate, mandelate, bitartrate, mesylate, borate, methylbromide, bromide, methylnitrate, calcium edetate, methylsulfate, camsylate, mucate, carbonate, napsylate, chloride, nitrate, clavulanate, N-methylglucamine, citrate, ammonium salt, dihydrochloride, oleate, edetate, oxalate, edisylate, pamoate (embonate), estolate, palmitate,

- 16 -

esylate, pantothenate, fumarate, phosphate/diphosphate, gluceptate, polygalacturonate, gluconate, salicylate, glutamate, stearate, glycollylarsanilate, sulfate, hexylresorcinate, subacetate, hydrabamine, succinate, hydrobromide, tannate, hydrochloride, tartrate, hydroxy-naphthoate, teoclate, iodide, tosylate, isothionate, triethiodide, lactate, and valerate.

In addition, some of the compounds of the instant invention may form solvates with water or common organic solvents. Such solvates are encompassed within the scope of this invention.

The term "therapeutically effective amount" shall mean that amount of a drug or pharmaceutical agent that will elicit the biological or medical response of a tissue, system, animal or human that is being sought by a researcher, veterinarian, medical doctor or other clinician.

The compounds of the present invention have chiral centers and may occur as racemates, racemic mixtures and as individual enantiomers or diastereomers, with all isomeric forms being included in the present invention as well as mixtures thereof. Furthermore, some of the crystalline forms for compounds of the present invention may exist as polymorphs and as such are intended to be included in the present invention.

Accordingly, the present invention has the objective of providing methods of treating the hyperandrogenic conditions of androgenic alopecia including male pattern baldness, acne vulgaris, seborrhea, and female hirsutism by oral, systemic, parenteral or topical administration of the novel compounds of Formula I either alone or in combination with a 5α -reductase 2 inhibitor, and/or further in combination with: a potassium channel opener, e.g., minoxidil; an anti-androgen, e.g., flutamide; a retinoid, e.g., tretinoin or isotretinoin; an alpha-1 receptor antagonist, e.g., terazocin.

The term "treating androgenic alopecia" is intended to include the arresting and/or reversing of androgenic alopecia, and the promotion of hair growth. The present invention has the further objective of providing methods of treating benign prostatic hyperplasia, prostatitis, and treating and/or preventing prostatic carcinoma by oral, systemic or

- 17 -

parenteral administration of the novel compounds of Formula I either alone or in combination with a 5 α -reductase 2 inhibitor.

The present invention also has the objective of providing suitable topical, oral, systemic and parenteral pharmaceutical formulations for use in the novel methods of treatment of the present invention. The compositions containing the present compounds as the active ingredient for use in the treatment of the above-noted hyperandrogenic conditions can be administered in a wide variety of therapeutic dosage forms in conventional vehicles for systemic administration. For example, the compounds can be administered in such oral dosage forms as tablets, capsules (each including timed release and sustained release formulations), pills, powders, granules, elixirs, tinctures, solutions, suspensions, syrups and emulsions, or by injection. Likewise, they may also be administered in intravenous (both bolus and infusion), intraperitoneal, subcutaneous, topical with or without occlusion, or intramuscular form, all using forms well known to those of ordinary skill in the pharmaceutical arts.

The pharmaceutical compositions included herein include those comprising a pharmaceutically acceptable carrier and a therapeutically effective amount of a compound of Formula I of this invention; which can further contain:

(1) a therapeutically effective amount of an inhibitor of 5 α -reductase 2, or a pharmaceutically acceptable salt thereof, e.g., finasteride, epristeride or turosteride;

(2) a potassium channel opener, or a pharmaceutically acceptable salt thereof, e.g., minoxidil;

(3) a therapeutically effective amount of a retinoid, or a pharmaceutically acceptable salt thereof, e.g., tretinoin, isotretinoin,

(4) a therapeutically effective amount of an anti-androgen, or a pharmaceutically acceptable salt thereof, e.g., flutamide, spironolactone, or casodex.

The daily dosage of the products may be varied over a range from 0.1 to 1,000 mg per adult human/per day. For oral administration, the compositions are preferably provided in the form of scored or

- 18 -

unscored tablets containing 0.1, 0.5, 1.0, 2.5, 5.0, 10.0, 15.0, 25.0, and 50.0 milligrams of the active ingredient for the symptomatic adjustment of the dosage to the patient to be treated. An effective amount of the drug is ordinarily supplied at a dosage level of from about 0.002 mg/kg to about 50 mg/kg of body weight per day. The range is more particularly from about 0.01 mg/kg to 7 mg/kg of body weight per day.

Advantageously, compounds of the present invention may be administered in a single daily dose, or the total daily dosage may be administered in divided doses of two, three or four times daily. Furthermore, compounds for the present invention can be administered in intranasal form via topical use of suitable intranasal vehicles, or via transdermal routes, using those forms of transdermal skin patches well known to those of ordinary skill in that art. To be administered in the form of a transdermal delivery system, the dosage administration will, of course, be continuous rather than intermittent throughout the dosage regimen.

For the treatment of androgenic alopecia including male pattern baldness, acne vulgaris, seborrhea, and female hirsutism, the compounds of the present invention may be administered in a pharmaceutical composition comprising the active compound in combination with a pharmaceutically acceptable carrier adapted for topical administration. Topical pharmaceutical compositions may be, e.g., in the form of a solution, cream, ointment, gel, lotion, shampoo or aerosol formulation adapted for application to the skin. These topical pharmaceutical compositions containing the compounds of the present invention ordinarily include about 0.001% to 15% by weight of the active compound in admixture with a pharmaceutically acceptable vehicle.

For the treatment of acne vulgaris, androgenic alopecia including male pattern baldness, seborrhea, female hirsutism, benign prostatic hyperplasia, prostatitis and the prevention and/or treatment of prostatic cancer, the compounds of the instant invention can be used alone or can be combined with a therapeutically effective amount of a 5 α -reductase 2 inhibitor, such as finasteride, in a single oral, systemic, or parenteral pharmaceutical dosage formulation. Also, for the skin and

- 19 -

scalp related disorders of acne vulgaris, androgenic alopecia including male pattern baldness, seborrhea, and female hirsutism, the compounds of the instant invention and a 5 α -reductase 2 inhibitor can be formulated for topical administration. Alternatively, a combined therapy can be employed wherein the compound of Formula I and the 5 α -reductase 2 inhibitor are administered in separate oral, systemic, parenteral or topical dosage formulations. For example, a compound of Formula I and e.g., finasteride can be administered in a single oral or topical dosage formulation, or each active agent can be administered in a separate dosage formulation, e.g., in separate oral dosage formulations, or an oral dosage formulation of finasteride in combination with a topical dosage formulation of a compound of Formula I. See, e.g., U.S. Patent No.'s 4,377,584 and 4,760,071 which describe dosages and formulations for 5 α -reductase inhibitors. Where the active agents are in separate dosage formulations, they can be administered concomitantly, or they each can be administered at separately staggered times.

Other 5 α -reductase type 2 inhibitors useful in the above-described method are also included within the scope of this invention. Examples which are non-limiting are: 17 β -(N-tert-butyl-carbamoyl)androsta-3,5-diene-3-carboxylic acid (epristeride, Smith Kline & Beecham, SKF 105657), which is described in WO9113550 and WO9319758; and 17 β -[N-isopropyl-N-(isopropylcarbamoyl)carbamoyl]-4-methyl-4-aza-5 α -androstane-3-one, (turosteride, Farmitalia, FCE 26073), which is described in USP 5,155,107, and derivatives thereof.

Furthermore, administration of a compound of the present invention in combination with a therapeutically effective amount of a potassium channel opener, such as minoxidil, cromakalim, pinacidil, a compound selected from the classes of S-triazine, thiane-1-oxide, benzopyran, and pyridinopyran derivatives or a pharmaceutically acceptable salt thereof, may be used for the treatment of androgenic alopecia including male pattern baldness. The active agents can be administered in a single topical dosage formulation, or each active agent can be administered in a separate dosage formulation, e.g., in separate topical dosage formulations, or an oral dosage formulation of a

- 20 -

compound of Formula I in combination with a topical dosage formulation of, e.g., minoxidil. See, e.g., U.S. Patent No.'s 4,596,812, 4,139,619 and WO 92/02225, published 20 February 1992, for dosages and formulations of calcium channel openers. Where the active agents are in separate dosage formulations, they can be administered concurrently, or they each can be administered at separately staggered times.

Furthermore, for the treatment of acne vulgaris and/or androgenic alopecia, a combined therapy can be used by administering a therapeutically effective amount of a compound of Formula I, alone or in combination with a 5α -reductase 2 inhibitor, or further in combination with a therapeutically effective amount of a retinoid, e.g., retinoic acid or an ester or amide derivative thereof, such as e.g., tretinoin (RETIN A) or isotretinoin (ACCUTANE, Roche, see USP 3,006,939; USP 3,746,730; USP 4,556,518).

Also, for the treatment of acne vulgaris, androgenic alopecia, seborrhea, female hirsutism, benign prostatic hyperplasia, prostatitis and the prevention and/or treatment of prostatic cancer, a combined therapy can be used by administering a therapeutically effective amount of a compound of Formula I with a therapeutically effective amount of an anti-androgen, such as, e.g., flutamide, spironolactone or casodex.

Also covered in this invention is a method of for treating benign prostatic hyperplasia comprising the step of administering to a mammal in need of such treatment a therapeutically effective amount of a compound of Formula I, or a therapeutically effective amount of a compound of Formula I in combination with an inhibitor of 5α -reductase 2. This method also includes the use of an alpha-1 receptor antagonist, e.g., terazosin (Abbott, see USP 4,026,894; 4,251,532).

Also covered is a method of inhibiting the biosynthetic conversion of testosterone to dihydrotestosterone in a mammal in need of such treatment comprising the step of administering to said mammal a therapeutically effective amount of a compound of Formula I , or a therapeutically effective amount of a compound of Formula I in combination with an inhibitor of 5α -reductase 2.

- 21 -

Also covered is a method of inhibiting 5 α -reductase or the isozymes thereof, in a mammal in need of such treatment, comprising the step of administering to said mammal a therapeutically effective amount of a compound of Formula I, or a therapeutically effective amount of a compound of Formula I in combination with an inhibitor of 5 α -reductase 2.

The dosage regimen utilizing the compounds of the present invention is selected in accordance with a variety of factors including type, species, age, weight, sex and medical condition of the patient; the severity of the condition to be treated; the route of administration; the renal and hepatic function of the patient; and the particular compound thereof employed. A physician or veterinarian of ordinary skill can readily determine and prescribe the effective amount of the drug required to prevent, counter or arrest the progress of the condition. Optimal precision in achieving concentration of drug within the range that yields efficacy without toxicity requires a regimen based on the kinetics of the drug's availability to target sites. This involves a consideration of the distribution, equilibrium, and elimination of a drug.

In the methods of the present invention, the compounds herein described in detail can form the active ingredient, and are typically administered in admixture with suitable pharmaceutical diluents, excipients or carriers (collectively referred to herein as "carrier" materials) suitably selected with respect to the intended form of administration, that is, oral tablets, capsules, elixirs, syrups and the like, and consistent with conventional pharmaceutical practices.

For instance, for oral administration in the form of a tablet or capsule, the active drug component can be combined with an oral, non-toxic pharmaceutically acceptable inert carrier such as ethanol, glycerol, water and the like. Moreover, when desired or necessary, suitable binders, lubricants, disintegrating agents and coloring agents can also be incorporated into the mixture. Suitable binders include, without limitation, starch, gelatin, natural sugars such as glucose or beta-lactose, corn sweeteners, natural and synthetic gums such as acacia, tragacanth or sodium alginate, carboxymethyl-cellulose, polyethylene glycol, waxes

- 22 -

and the like. Lubricants used in these dosage forms include, without limitation, sodium oleate, sodium stearate, magnesium stearate, sodium benzoate, sodium acetate, sodium chloride and the like. Disintegrators include, without limitation, starch, methyl cellulose, agar, bentonite, xanthan gum and the like.

The liquid forms in suitably flavored suspending or dispersing agents such as the synthetic and natural gums, for example, tragacanth, acacia, methyl-cellulose and the like. Other dispersing agents which may be employed include glycerin and the like. For parenteral administration, sterile suspensions and solutions are desired. Isotonic preparations which generally contain suitable preservatives are employed when intravenous administration is desired.

Topical preparations containing the active drug component can be admixed with a variety of carrier materials well known in the art, such as, e.g., alcohols, aloe vera gel, allantoin, glycerine, vitamin A and E oils, mineral oil, PPG2 myristyl propionate, and the like, to form, e.g., alcoholic solutions, topical cleansers, cleansing creams, skin gels, skin lotions, and shampoos in cream or gel formulations. See, e.g., EP 0 285 382.

The compounds of the present invention can also be administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles and multilamellar vesicles. Liposomes can be formed from a variety of phospholipids, such as cholesterol, stearylamine or phosphatidylcholines.

Compounds of the present invention may also be delivered by the use of monoclonal antibodies as individual carriers to which the compound molecules are coupled. The compounds of the present invention may also be coupled with soluble polymers as targetable drug carriers. Such polymers can include polyvinyl-pyrrolidone, pyran copolymer, polyhydroxypropylmethacrylamide-phenol, polyhydroxy-ethylaspartamidephenol, or polyethylene-oxidepolylysine substituted with palmitoyl residues. Furthermore, the compounds of the present invention may be coupled to a class of biodegradable polymers useful in achieving controlled release of a drug, for example, polylactic acid,

- 23 -

polyepsilon caprolactone, polyhydroxy butyric acid, polyorthoesters, polyacetals, polydihydropyrans, polycyanoacrylates and cross-linked or amphipathic block copolymers of hydrogels.

The compounds of the present invention can be prepared readily according to the following reaction Schemes and Examples or modifications thereof using readily available starting materials, reagents and conventional synthesis procedures. In these reactions, it is also possible to make use of variants which are themselves known to those of ordinary skill in this art, but are not mentioned in greater detail. Specific definitions of variables in the Schemes (e.g., R = CH₃) are illustrative only, and are not intended to limit the procedures described. Some abbreviations used herein are as follows: Ph is phenyl; Ac is an acyl group; t-Bu is tert-butyl; Et is ethyl; Me is methyl; i-Am is iso-amyl; EtOAc is ethyl acetate.

The inhibitors described in Scheme 1 can be prepared as follows. 4-Aza-4-methyl-5 α -androstane-3,17-dione (A) is first converted into the isomeric 3,16-dione **1** by the following sequence of reactions: (1) treatment of A with isoamyl nitrite in t-butanol in the presence of potassium t-butoxide to generate the intermediate 16-oximino-17-ketone; (2) reduction of the 17-keto group with hydrazine hydrate and potassium hydroxide in ethylene glycol at elevated temperatures to give 16-oxime **B**; and (3) cleavage of the 16-oximino group in **B** either by hydrolysis with aqueous acetic acid at elevated temperatures or with sodium bisulfite followed by treatment with aqueous hydrochloric acid to afford **1**. Reduction of the 16-ketone **1** to the 16 β -alcohol **2** is carried out with a suitable hydride-based reducing agent, such as sodium borohydride in methanol or lithium tri-sec-butylborohydride in tetrahydrofuran (THF). Alcohol **2** is converted into its alkyl ether derivatives **3** and **4**, by first generating the alkoxide anion with potassium hydride in N,N-dimethylformamide (DMF) or potassium hydroxide in dimethyl sulfoxide (DMSO) followed by addition of the appropriate alkyl bromide or iodide. The 16 β -(n-propyloxy) derivative **5** is obtained from the precursor 16 β -(allyloxy) derivative **4** by catalytic hydrogenation.

- 24 -

The inhibitors described in Scheme 2 can be prepared as follows. 16-Oxime **B** is converted into the 16 β -amine **C** by catalytic hydrogenation in the presence of a heterogeneous catalyst such as platinum oxide in aqueous acetic acid. Acylation of **C** is effected with the appropriate acid anhydride or acid chloride in the presence of an acid acceptor such as pyridine, triethylamine, and 4-dimethylaminopyridine (DMAP). In this fashion are obtained examples 6 and 7. Carbamates, such as **8** depicted in Scheme 3, are made by treatment of alcohol **2** with the appropriate isocyanate in the presence of an organic base, such as triethylamine, pyridine, and 4-dimethylaminopyridine.

The inhibitors described in Scheme 4 can be prepared as follows. The 16 β -alcohol **2** is converted into the 16 α -alcohol **9** by treatment with 4-nitro-benzoic acid in the presence of diethyl azadicarboxylate (DEAD) and triphenylphosphine to generate the intermediate 16 α -(p-nitrobenzoate) ester **D** followed by hydrolysis in aqueous base in an appropriate alcohol solvent. Alkylation of **9** is carried out in an analogous fashion as described above with alcohol **2** to yield the desired 16 α -alkyl ethers, such as the 16 α -methoxy derivative (example 10) shown in Scheme 4.

The 7 β -methyl inhibitors described in Scheme 5 are prepared in a similar manner as that described above for the examples in Scheme 1, but using instead as starting material, 4-aza-4,7 β -dimethyl-5 α -androstan-3,17-dione (**E**).

The 7 β -methyl inhibitors described in Scheme 6 are prepared as follows. Compound **20** is prepared by treatment of alcohol **12** with *t*-butyl trichloroacetimidate in the presence of an organic sulfonic acid, such as trifluoromethanesulfonic acid. The 16 β -aryloxy derivatives, such as Examples **21-24**, are obtained by first generating the alkoxide anion from alcohol **12** with potassium or sodium hydride in tetrahydrofuran or *N,N*-dimethylformamide or potassium hydroxide in dimethylsulfoxide and subsequent addition of the appropriately substituted fluorobenzene.

The 7 β -methyl inhibitors described in Scheme 7 are prepared in a similar manner as that described above for the examples in

- 25 -

Scheme 4, but using instead as starting material the 7 β -methyl-16 β -ol intermediate **12**. Inversion of configuration at the 16-position to form (F) is effected using a Mitsunobu-based transformation as shown in Scheme 7. O-Alkylation to generate 16 α -ethers, such as **26**, is performed as already described above.

The inhibitors described in Scheme 8 are prepared as follows. Addition of methylmagnesium bromide in tetrahydrofuran to either ketone **1** or **11** affords the corresponding 16 α -methyl-16 β -alcohol **27** or **28**. O-Alkylation or O-arylation is then carried out as described in the previous Schemes to afford the 16 α -methyl-16 β -ether derivatives, such as Examples **29** and **30**.

The inhibitors described in Scheme 9 are prepared by the following reaction routes. The 7 β -methyl-16 α -alcohol **25** is converted into the 16 β -thiol **H** by treatment with thiolacetic acid in the presence of diisopropyl azodicarboxylate (DIAD) and triphenylphosphine to give the intermediate 16 β -thioacetate **G**, which is then hydrolyzed under basic conditions to yield thiol **H**. Alkylation is effected by generating the mercaptide anion with sodium hydride or potassium hydride in tetrahydrofuran or N,N-dimethyl-formamide followed by addition of the appropriate alkyl halide. In this fashion are prepared Examples **31-33**. The corresponding sulfones, such as Example **34**, are obtained by treatment of the precursor thioethers **31-33** with an oxidizing agent, such as organic peracid or potassium peroxymonosulfate (OXONE), the latter in aqueous methanol.

The inhibitors described in Scheme 10 are prepared by the following synthetic pathways. The p-nitrophenoxy derivative **50** is reduced with Pd on carbon at room temperature in a H₂ atmosphere to yield the p-amino-phenoxy derivative **51**. The amine is then acylated with acetyl chloride in methylene chloride in the presence of pyridine to yield the p-acetylaminophenoxy derivative **52**, or likewise treated with benzoyl chloride to yield the corresponding p-benzoylamino analog **53**. Alternately, the amino compound **51** is treated with tosyl chloride to yield the p-tosylamino analog **54**.

- 26 -

The inhibitors described in Scheme 11 are prepared as follows. The N-2,4-dimethoxybenzyl protected 16-alcohol **55** is treated with p-fluorochlorobenzene and potassium hydride in dimethylformamide to yield the p-chlorophenoxy derivative **56**, which is then treated with trifluoroacetic acid in methylene chloride to remove the N-2,4-dimethoxybenzyl protecting group to yield **57**. This is treated with hydrogen gas and a palladium on carbon catalyst in methanol to dechlorinate the phenyl ring to yield the phenoxy derivative **58**. This compound is treated with methyl iodide and sodium hydride in dimethylformamide to methylate the ring nitrogen to yield **61**. Alternately, **58** is treated with DDQ and BSTFA in toluene to introduce a double bond at the 1-position to yield **59**. Utilizing the same reduction reaction scheme, the 1,2-dihydro androstane **57** yields the p-chloroandrost-1-ene **60**. This is then methylated at the 1-position by treatment with methyl iodide, sodium hydride in dimethylformamide to yield **62**.

The inhibitors described in Scheme 12 are prepared via similar reaction pathways as described in Scheme 11. The N-2,4-dimethoxybenzyl protected 16-alcohol **55** is treated with 4-methyl-3-chlorofluorobenzene and potassium hydride in dimethylformamide to yield the 4-methyl-3-chlorophenoxy derivative **63**, which is then treated with trifluoroacetic acid in methylene chloride to remove the N-2,4-dimethoxybenzyl protecting group to yield **64**. This is treated with hydrogen gas and a palladium on carbon catalyst in methanol to dechlorinate the phenyl ring to yield the p-methylphenoxy derivative **65**. This compound is treated with methyl iodide and sodium hydride in dimethylformamide to methylate the ring nitrogen to yield **67**. Alternately, **65** is treated with DDQ and BSTFA in toluene to introduce a double bond at the 1-position to yield **66**.


The inhibitors in Scheme 13 are prepared as follows. The starting 16-alcohol **25** is treated with methanesulfonic acid in pyridine containing DMAP to yield the mesylate **77**. This in turn is treated with an appropriate thiophenol in anhydrous THF containing sodium hydride to yield the thiophenoxy **78**, 4-chlorothiophenoxy **79**, 4-

- 27 -



fluorothiophenoxy 80, 4-methylthiophenoxy 81) and the 4-methoxythiophenoxy 82 derivatives. Treatment of the thiophenoxy 78 derivative with m-chloroperbenzoic acid in methylene chloride at 0°C for one hour yields the phenylsulfinyl derivative 83. Treatment of the phenylsulfinyl compound 83 under the same reaction conditions prolonged however for three hours, yields the phenylsulfonyl derivative 84.

The inhibitors for Scheme 14 are prepared as follows. The 16-ketone (11) is treated with an appropriate arylmethyl diethylphosphonate under Wittig conditions using sodium hydride in DMF at 80-100°C to yield the corresponding 4-chlorobenzylidene 71, benzylidene 72 and 4-methylbenzylidene 73 analogs. These are reduced in ethanol under a hydrogen atmosphere using a 5% rhodium on carbon catalyst to yield the corresponding 4-chlorobenzyl 74 and 4-methylbenzyl 75 derivatives. The 3-pyridyl-methyl 76 analog is made in the same two step manner.


- 28 -

SCHEME 1


- 29 -

SCHEME 2SCHEME 3


- 30 -

SCHEME 4


- 31 -

SCHEME 5


- 32 -

SCHEME 6


- 33 -

SCHEME 7


- 34 -

SCHEME 8


- 35 -

SCHEME 9


- 36 -

SCHEME 10


- 37 -


SCHEME 11

- 38 -

SCHEME 12

- 39 -

SCHEME 13

SCHEME 14

The following examples are provided to further illustrate details for the preparation of the compounds of the present invention. The examples are not intended to be limitations on the scope of the instant invention in any way, and they should not be so construed. Furthermore, the compounds described in the following examples are not to be construed as forming the only genus that is considered as the invention, and any combination of the compounds or their moieties may itself form a genus. Those skilled in the art will readily understand that known variations of the conditions and processes of the following preparative procedures can be used to prepare these compounds. All temperatures are in degrees Celsius unless noted otherwise.

- 41 -

The starting material 4-aza-4-methyl-5 α -androstan-3,17-dione (Compound A in Scheme 1 above) can be made according to the methods described in Rasmusson, *et al.*, *J. Med. Chem.*, 27, p. 1690-1701 (1984). The starting material 4-aza-4,7 β -dimethyl-5 α -androstan-3,17-dione can be synthesized according to the procedure described in Example 36, below.

EXAMPLE 1

4-Aza-4-methyl-5 α -androstan-3,16-dione

Step 1: 4-aza-4-methyl-5 α -androstan-3,17-dione-16-oxime

To 2-methyl-2-propanol (14 mL) in a round-bottom flask under a stream of nitrogen gas was added potassium *tert*-butoxide (740 mg, 6.59 mmol). After complete solution was achieved, 4-aza-4-methyl-5 α -androstan-3,17-dione (1.0 g, 3.30 mmol) was added and stirring was continued for 1 hour affording a gold-colored solution. To the reaction mixture was added dropwise with stirring isoamyl nitrite (0.884 mL, 6.58 mmol), and stirring was continued overnight at room temperature affording a deep-orange solution. The mixture was then diluted with an equal volume of water and acidified to pH ~2 with 2 N hydrochloric acid. Diethyl ether was added, and the solid that formed was filtered, washed with ether, and dried *in vacuo*, to yield the title compound.

Step 2: 4-aza-4-methyl-5 α -androstan-3-one-16-oxime

To a mixture of 4-aza-4-methyl-5 α -androstan-3,17-dione-16-oxime (596 mg, 1.79 mmol) in ethylene glycol (5 mL) were added 98% hydrazine (57 μ L, 1.74 mmol) and powdered potassium hydroxide (568 mg, 10.12 mmol). The mixture was heated for 16 h at 140°, cooled, and neutralized with 2N hydrochloric acid. The resulting solid was filtered, washed with water, and dried *in vacuo* to yield the title compound; mass spectrum: m/z 318(M).

- 42 -

Step 3: 4-aza-4-methyl-5 α -androstan-3,16-dione

A mixture of 4-aza-4-methyl-5 α -androstan-3-one-16-oxime (218 mg, 0.684 mmol) and sodium bisulfite (249 mg, 23.9 mmol) in 50% aqueous ethanol (10 mL) was heated for 3 h at reflux temperature. Dilute hydrochloric acid (0.5 N, 33 mL) and methylene chloride (50 mL) were added, and the mixture was vigorously agitated for several minutes. The organic layer was separated and washed with sodium hydrogencarbonate solution, saturated brine solution, dried (Na₂SO₄), and evaporated. The desired product was purified by flash silica gel chromatography using 15% acetone/methylene chloride as eluant to yield the title compound; FAB mass spectrum: *m/z* 304 (*M*+1). 400 MHz ¹H NMR (CDCl₃): δ 0.89 (s, 3H); 0.91 (s, 3H); 2.90 (s, 3H); and 3.05 (dd, 1H).

EXAMPLE 2**3-Oxo-4-aza-4-methyl-16 β -hydroxy-5 α -androstan-3-one**

A solution of 4-aza-4-methyl-5 α -androstan-3,16-dione (100 mg, 0.330 mmol) in methanol (2 mL) was cooled in an ice bath and treated with sodium borohydride (38 mg, 0.989 mmol) for 1 h. The reaction mixture was diluted with water and extracted with methylene chloride (2 x 20 mL). The combined organic extracts were washed with saturated brine solution, dried (Na₂SO₄), and evaporated. The desired product was purified by flash silica gel chromatography using 10% acetone/methylene chloride as eluant to yield the title compound; FAB mass spectrum: *m/z* 306 (*M*+1). 400 MHz ¹H NMR (CDCl₃): δ 0.88 (s, 3H); 0.95 (s, 3H); 2.90 (s, 3H); 3.00 (dd, 1H); and 4.39 (m, 1H).

- 43 -

EXAMPLE 33-Oxo-4-aza-4-methyl-16 β -methoxy-5 α -androstande

To a solution of 3-oxo-4-aza-4-methyl-16 β -hydroxy-5 α -androstande (35 mg, 0.115 mmol) in dimethyl sulfoxide (1.0 mL) was added powdered potassium hydroxide (32 mg, 0.575 mmol). After stirring for 15 min at room temperature under an nitrogen atmosphere, iodomethane (36 μ l, 0.575 mmol) was added and stirring was continued for a further 4 hours. The mixture was diluted with diethyl ether (30 mL), which was washed with water, saturated brine solution, dried (Na₂SO₄), and evaporated. The desired product was purified by flash silica gel chromatography using 10% acetone/methylene chloride as eluant to yield the title compound; mass spectrum: *m/z* 391 (M). 400 MHz ¹H NMR (CDCl₃): δ 0.88 (s, 6H); 2.90 (s, 3H); 3.00 (dd, 1H); 3.21 (s, 3H); and 3.83 (m, 1H).

EXAMPLE 43-Oxo-4-aza-4-methyl-16 β -allyloxy-5 α -androstande

This compound was prepared in a similar fashion as Example 3, but substituting allyl bromide in place of iodomethane to yield the title compound; mass spectrum: *m/z* 345 (M). 400 MHz ¹H NMR (CDCl₃): δ 0.88 (s, 3H); 0.90 (s, 3H); 2.90 (s, 3H); 3.00 (dd, 1H); 3.90 (m, 2H); 3.99 (m, 1H); 5.11-5.27 (m, 2H); and 5.83-5.93 (m, 1H).

EXAMPLE 53-Oxo-4-aza-4-methyl-16 β -(n-propyloxy)-5 α -androstande

A solution of 3-oxo-4-aza-4-methyl-16 β -allyloxy-5 α -androstande in ethyl acetate (0.85 mL) was hydrogenated at atmospheric pressure in the presence of platinum oxide (4 mg) for 30 min at room temperature. The catalyst was removed by filtration through a Millex-HV 0.45 μ m Filter Unit. Purification was achieved by flash silica gel

- 44 -

chromatography using 10% acetone/methylene chloride as eluant to yield the title compound; mass spectrum: m/z 348 (M+1).
400 MHz NMR ($CDCl_3$): δ 0.88 (s, 3H); 0.89 (s, 3H); 2.90 (s, 3H); 3.00 (dd, 1H); 3.28 (t, 2H); and 3.92 (m, 1H).

EXAMPLE 6

3-Oxo-4-aza-4-methyl-16 β -(acetamido)-5 α -androstane

Step 1: 3-Oxo-4-aza-4-methyl-16 β -(amino)-5 α -androstane

A solution of 4-aza-4-methyl-5 α -androstan-3-one-16-oxime (150 mg, 0.471 mmol) in ethanol (15 mL) - acetic acid (7 mL) was hydrogenated at atmospheric pressure in the presence of platinum oxide (50 mg) overnight at room temperature. The catalyst was removed by filtration through a Millex-HV 0.45 μ m Filter Unit, and the filtrate was evaporated. The residue was dissolved in methylene chloride (50 mL), and the solution was washed with saturated sodium hydrogencarbonate solution, saturated brine solution, dried (Na_2SO_4), and evaporated to afford the desired amine.

Step 2: 3-Oxo-4-aza-4-methyl-16 β -(acetamido)-5 α -androstane

The amine from Step 1 (56 mg, 0.184 mmol) was dissolved in methylene chloride (1.0 mL) and treated with pyridine (0.6 mL), 4-dimethylaminopyridine (5 mg), and acetic anhydride (0.3 mL) for 2 h at room temperature. The mixture was diluted with methylene chloride (50 mL), and the solution was washed with water, 1N hydrochloric acid, saturated sodium hydrogencarbonate solution, saturated brine solution, dried (Na_2SO_4), and evaporated. The product was purified by flash silica gel chromatography using 2% methanol/methylene chloride as eluant to yield the title compound; mass spectrum: m/z 346 (M).
400 MHz 1H NMR ($CDCl_3$): δ 0.82 (s, 3H); 0.87 (s, 3H); 1.93 (s, 3H); 2.90 (s, 3H); 3.00 (dd, 1H); 4.28 (m, 1H); and 5.54 (d, 1H).

- 45 -

EXAMPLE 7

3-Oxo-4-aza-4-methyl-16 β -(benzamido)-5 α -androstane

This compound was prepared in a similar fashion as Example 6, but substituting benzoyl chloride in place of acetic anhydride to yield the title compound; mass spectrum: m/z 408 (M). 400 MHz ^1H NMR (CDCl_3): δ 0.89 (s, 3H); 0.90 (s, 3H); 2.90 (s, 3H); 3.01 (dd, 1H); 4.48 (m, 1H); and 6.12 (d, 1H).

EXAMPLE 8

3-Oxo-4-aza-4-methyl-16 β -(benzylaminocarbonyloxy)-5 α -androstane

To a solution of 3-oxo-4-aza-4-methyl-16 β -hydroxy-5 α -androstane (40 mg, 0.131 mmol) in methylene chloride (2 mL) were added triethylamine (67 μL , 0.481 mmol), 4-dimethylaminopyridine (2 mg), and benzyl isocyanate (50 μL , 0.405 mmol). The reaction mixture was stirred for 48 h at room temperature, evaporated, and then subjected to flash silica gel chromatography using 15% acetone/methylene chloride as eluant to yield the title compound; FAB mass spectrum: m/z 439 (M+1).

400 MHz ^1H NMR (CDCl_3): δ 0.87 (s, 6H); 2.90 (s, 3H); 3.00 (dd, 1H); 4.33 (m, 2H); 4.90 (m, 1H) and 5.11 (m, 1H).

EXAMPLE 9

3-Oxo-4-aza-4-methyl-16 α -hydroxy-5 α -androstane

Step 1: 3-Oxo-4-aza-4-methyl-16 α -(4-nitrobenzoyloxy)-5 α -androstane

To a solution of 3-oxo-4-aza-4-methyl-16 β -hydroxy-5 α -androstane (34 mg, 0.0111 mmol) in dry benzene (1.5 mL) were added triphenylphosphine (35 mg, 0.134 mmol), 4-nitrobenzoic acid (22 mg, 0.134 mmol), and diethyl azodicarboxylate (21 μL , 0.134 mmol). The reaction mixture was heated for one hour at 80° (oil bath temperature)

- 46 -

under a nitrogen atmosphere. After removal of the benzene by evaporation under diminished pressure, the crude product mixture was subjected to flash silica gel chromatography using 2% methanol/methylene chloride as eluant to give desired product contaminated with some triphenylphosphine (97 mg) which was saponified as described in Step 2.

Step 2: 3-Oxo-4-aza-4-methyl-16 α -hydroxy-5 α -androstane

The crude product from Step 1 (97 mg) was suspended in ethanol (0.5 mL) and treated with 0.4 N sodium hydroxide (0.36 mL, 0.144 mmol). After stirring 90 min at room temperature, the reaction mixture was neutralized with several drops of glacial acetic acid, extracted with ethyl acetate (2 x 20 mL), washed with water (20 mL), saturated brine solution, dried (sodium sulfate), and evaporated. The product was obtained pure by flash silica gel chromatography using 20% acetone/methylene chloride as eluant; mass spectrum: m/z 305 M. 400 MHz ^1H NMR (CDCl_3): δ 0.70 (s, 3H); 0.85 (s, 3H); 2.90 (s, 3H); 3.02 (dd, 1H); and 4.47 (m, 1H).

EXAMPLE 10

3-Oxo-4-aza-4-methyl-16 α -methoxy-5 α -androstane

To a solution of 3-oxo-4-aza-4-methyl-16 α -hydroxy-5 α -androstane (20 mg, 0.065 mmol) in dimethyl sulfoxide (0.6 mL) was added powdered potassium hydroxide (18 mg, 0.325 mmol). After stirring for 15 min at room temperature under an nitrogen atmosphere, iodomethane (20 μl , 0.325 mmol) was added and stirring was continued overnight at room temperature. The mixture was diluted with diethyl ether (25 mL), which was washed with water (2 x 10 mL), dried (Na_2SO_4), and evaporated. The desired product was purified by flash silica gel chromatography using 10% acetone/methylene chloride as eluant to yield the title compound; mass spectrum: m/z 319 (M). 400 MHz ^1H NMR (CDCl_3): δ 0.70 (s, 3H); 0.87 (s, 3H); 2.90 (s, 3H); 3.01 (dd, 1H); 3.22 (s, 3H); and 3.92 (m, 1H).

EXAMPLE 114-Aza-4,7 β -dimethyl-5 α -androstan-3,16-dioneStep 1: 4-aza-4,7 β -dimethyl-5 α -androstan-3,17-dione-16-oxime

To 2-methyl-2-propanol (28 mL) in a round-bottom flask under a stream of nitrogen gas was added potassium *tert*-butoxide (1.35 g, 12.1 mmol). After complete solution was achieved, 4-aza-4,7 β -dimethyl-5 α -androstan-3,17-dione (1.92 g, 6.0 mmol) was added and stirring was continued for 1 hour affording a gold-colored solution. To the reaction mixture was added dropwise with stirring isoamyl nitrite (1.63 mL, 12.1 mmol), and stirring was continued overnight at room temperature affording a deep-orange solution. The mixture was then diluted with an equal volume of water, acidified to pH ~2 with 2 N hydrochloric acid, and extracted with diethyl ether (3 x 50 mL). The combined ether extracts were washed with saturated brine solution, dried (sodium sulfate), and evaporated. The crude product was subjected to flash silica gel chromatography using 5% methanol/methylene chloride as eluant to yield the title compound.

Step 2: 4-aza-4,7 β -dimethyl-5 α -androstan-3-one-16-oxime

To a mixture of 4-aza-4,7 β -dimethyl-5 α -androstan-3,17-dione-16-oxime (2.7 g, 7.79 mmol) in ethylene glycol (30 mL) were added 98% hydrazine (0.27 mL, 8.57 mmol) and powdered potassium hydroxide (2.62 g, 46.8 mmol). The mixture was heated for 3 h at 140°, cooled, diluted with water (100 mL), neutralized with concentrated hydrochloric acid to give a tan precipitate that was filtered and dried (1.7 g). Flash silica gel chromatography of this material using initially 2% methanol/methylene chloride and subsequently 5% methanol/methylene chloride as eluant gave pure product.

Step 3: 4-aza-4,7 β -dimethyl-5 α -androstan-3,16-dione

A mixture of 4-aza-4,7 β -dimethyl-5 α -androstan-3-one-16-oxime (0.55 g, 1.65 mmol) in 60% acetic acid (20 mL) was heated at

- 48 -

reflux temperature for 48 hours. The cooled mixture was diluted with water (25 mL) and extracted with methylene chloride (3 x 50 mL). The combined extracts were washed with saturated sodium hydrogen-carbonate solution, dried (sodium sulfate), and evaporated. Flash silica gel chromatography using 2% methanol/methylene chloride afforded pure product; mass spectrum: m/z 317 (M).

400 MHz ^1H NMR (CDCl_3): δ 0.88 (s, 3H); 0.89 (s, 3H); 1.00 (d, 3H); 2.90 (s, 3H); and 3.07 (dd, 1H).

EXAMPLE 12

3-Oxo-4-aza-4,7 β -dimethyl-16 β -hydroxy-5 α -androstane

A solution of 4-aza-4,7 β -dimethyl-5 α -androstan-3,16-dione (390 mg, 1.23 mmol) in methanol (8 mL) was cooled in an ice bath and treated with sodium borohydride (140 mg, 3.68 mmol) for 30 min. The reaction mixture was diluted with water and extracted with methylene chloride (3 x 50 mL). The combined organic extracts were washed with saturated brine solution, dried (Na_2SO_4), and evaporated. The desired product was purified by flash silica gel chromatography using initially 10% acetone/methylene chloride and subsequently 20% acetone/methylene chloride as eluant to yield the title compound; mass spectrum: m/z 391(M).

400 MHz ^1H NMR (CDCl_3): δ 0.83 (s, 3H); 0.96 (s, 3H); 1.03 (d, 3H); 2.90 (s, 3H); 3.00 (dd, 1H); and 4.36 (m, 1H).

EXAMPLE 13

3-Oxo-4-aza-4,7 β -dimethyl-16 β -methoxy-5 α -androstane

To a solution of 3-oxo-4-aza-4,7 β -dimethyl-16 β -hydroxy-5 α -androstane (20 mg, 0.063 mmol) in dimethyl sulfoxide (0.5 mL) was added powdered potassium hydroxide (18 mg, 0.313 mmol). After stirring for 15 min at room temperature under a nitrogen atmosphere, iodomethane (20 μl , 0.313 mmol) was added and stirring was continued overnight at room temperature. The mixture was diluted with diethyl

- 49 -

ether (25 mL), which was washed with water, saturated brine solution, dried (Na₂SO₄), and evaporated. The desired product was purified by flash silica gel chromatography using 1.5% methanol/methylene chloride as eluant to yield the title compound; mass spectrum: m/z 334 (M+1). 400 MHz ¹H NMR (CDCl₃): δ 0.83 (s, 3H); 0.89 (s, 3H); 1.03 (d, 3H); 2.90 (s, 3H); 3.00 (dd, 1H); 3.24 (s, 3H); and 3.80 (m, 1H).

EXAMPLE 14

3-Oxo-4-aza-4,7β-dimethyl-16β-ethyloxy-5α-androstane

This compound was prepared in a similar fashion as Example 13, but substituting iodoethane in place of iodomethane and potassium hydride in N,N-dimethylformamide in place of potassium hydroxide in dimethyl sulfoxide; mass spectrum: m/z 347 (M). 400 MHz ¹H NMR (CDCl₃): δ 0.83 (s, 3H); 0.90 (s, 3H); 1.03 (d, 3H); 1.18 (t, 3H); 2.90 (s, 3H); 3.00 (dd, 1H); 3.39 (m, 2H); and 4.40 (m, 1H).

EXAMPLE 15

3-Oxo-4-aza-4,7β-dimethyl-16β-allyloxy-5α-androstane

This compound was prepared in a similar fashion as Example 13, but substituting allyl bromide in place of iodomethane; mass spectrum: m/z 359 (M). 400 MHz ¹H NMR (CDCl₃): δ 0.83 (s, 3H); 0.91 (s, 3H); 1.04 (d, 3H); 2.90 (s, 3H); 3.00 (dd, 1H); 3.90 (m, 2H); 3.96 (m, 1H); 5.11-5.29 (m, 2H); and 5.85-5.93 (m, 1H).

EXAMPLE 16

3-Oxo-4-aza-4,7β-dimethyl-16β-benzyloxy-5α-androstane

This compound was prepared in a similar fashion as Example 14, but substituting benzyl bromide in place of iodoethane; mass spectrum: m/z 409 (M).

- 50 -

400 MHz ^1H NMR (CDCl_3): δ 0.85 (s, 3H); 0.95 (s, 3H); 1.04 (d, 3H); 2.90 (s, 3H); 3.00 (dd, 1H); 4.01 (m, 1H); 4.43 (q, 2H); and 7.31 (m, 5H).

EXAMPLE 17

3-Oxo-4-aza-4,7 β -dimethyl-16 β -(3,3-dimethylallyloxy)-5 α -androstane

This compound was prepared in a similar fashion as Example 13 but substituting 3,3-dimethylallyl bromide in place of iodomethane;

400 MHz ^1H NMR (CDCl_3): δ 0.82 (s, 3H); 0.90 (s, 3H); 1.02 (d, 3H); 1.67 (s, 3H); 1.71 (s, 3H); 2.90 (s, 3H); 3.00 (dd, 1H); 3.93 (m, 1H); and 5.31 (m, 1H).

EXAMPLE 18

3-Oxo-4-aza-4,7 β -dimethyl-16 β -(n-propyloxy)-5 α -androstane

A solution of 3-oxo-4-aza-4,7 β -dimethyl-16 β -allyloxy-5 α -androstane (13.0 mg, 0.036 mmol) in ethyl acetate (0.5 mL) was hydrogenated at atmospheric pressure in the presence of platinum oxide (4 mg) for 30 min at room temperature. The catalyst was removed by filtration through a Millex-HV 0.45 μm Filter Unit. Purification was achieved by flash silica gel chromatography using 1% methanol/methylene chloride as eluant to yield the title compound; mass spectrum: m/z 361 (M).

400 MHz ^1H NMR (CDCl_3): δ 0.82 (s, 3H); 0.89 (s, 3H); 0.89 (t, 3H); 1.05 (d, 3H); 2.90 (s, 3H); 3.00 (dd, 1H); 3.29 (t, 2H); and 3.89 (m, 1H).

EXAMPLE 19

3-Oxo-4-aza-4,7 β -dimethyl-16 β -(3-methyl-1-butyloxy)-5 α -androstane

A solution of 3-oxo-4-aza-4,7 β -dimethyl-16 β -(3,3-dimethylallyloxy)-5 α -androstane (12 mg) in ethyl acetate (0.5 mL) was hydrogenated at atmospheric pressure in the presence of 10% palladium-on-charcoal (3 mg) for 30 min at room temperature. The catalyst was removed by filtration through a Millex-HV 0.45 μm Filter Unit.

- 51 -

Purification was achieved by flash silica gel chromatography using 2% methanol/methylene chloride as eluant to yield the title compound; mass spectrum: m/z 389 M.

400 MHz ^1H NMR (CDCl_3): δ 0.82 (s, 3H); 0.88 (s, 3H); 1.03 (d, 3H); 2.90 (s, 3H); 3.00 (dd, 1H); 3.33 (m, 2H); and 3.88 (m, 1H).

EXAMPLE 20

3-Oxo-4-aza-4,7 β -dimethyl-16 β -(t-butoxy)-5 α -androstane

To a solution of 3-oxo-4-aza-4,7 β -dimethyl-16 β -hydroxy-5 α -androstane (20 mg, 0.063 mmol) in methylene chloride (0.5 mL) cooled in an ice-bath were added t-butyl trichloroacetimidate (23 μL , 0.126 mmol) and trifluoromethanesulfonic acid (0.56 μL , 0.0063 mmol). The reaction mixture was allowed to reach room temperature, and after one hour additional amounts of t-butyl trichloroacetimidate (23 μL) and trifluoromethanesulfonic acid (0.56 μL) were added. After one hour, a third addition of each reagent was made, and the reaction mixture was stirred for 5 h at room temperature. The mixture was diluted with diethyl ether (50 mL), washed with 1N aqueous sodium hydroxide (10 mL), 1N hydrochloric acid (10 mL), saturated sodium hydrogencarbonate solution, dried (sodium sulfate), and evaporated. The crude product was purified by flash silica gel chromatography using 10% acetone/methylene chloride as eluant to yield the title compound; mass spectrum: m/z 375 (M). 400 MHz ^1H NMR (CDCl_3): δ 0.82 (s, 3H); 0.90 (s, 3H); 1.03 (d, 3H); 1.11 (s, 9H); 2.90 (s, 3H); 3.00 (dd, 1H); and 4.00 (m, 1H).

EXAMPLE 21

3-Oxo-4-aza-4,7 β -dimethyl-16 β -(4-cyanophenoxy)-5 α -androstane

To a solution of 3-oxo-4-aza-4,7 β -dimethyl-16 β -hydroxy-5 α -androstane (20 mg, 0.063 mmol) in N,N-dimethylformamide (0.5 mL) was added powdered potassium hydride (35 weight%) (15 mg, 0.126 mmol). After stirring for 15 min at room temperature under an nitrogen atmosphere, 4-fluorobenzonitrile (38 mg, 0.315 mmol) was added and

- 52 -

stirring was continued for 2 hours at room temperature. The mixture was diluted with methylene chloride (25 mL) and quenched in ice-water. The aqueous layer was extracted with methylene chloride (3 x 25 mL) and the combined organic layers were washed with saturated brine solution, dried (sodium sulfate) and evaporated. The desired product was purified by flash silica gel chromatography using initially 1.5% methanol/methylene chloride and subsequently 2% methanol/methylene chloride as eluant to yield the title compound; mass spectrum: m/z 420 (M).

400 MHz ^1H NMR (CDCl_3): δ 0.86 (s, 3H); 0.92 (s, 3H); 1.04 (d, 3H); 2.90 (s, 3H); 3.02 (dd, 1H); 4.76 (m, 1H); 6.87 (m, 2H); and 7.53 (m, 2H).

EXAMPLE 22

3-Oxo-4-aza-4,7 β -dimethyl-16 β -(4-trifluoromethylphenoxy)-5 α -androstane

This compound was prepared in a similar fashion as Example 21, but substituting 4-fluorobenzotrifluoride in place of 4-fluorobenzonitrile; mass spectrum: m/z 463 (M).

400 MHz ^1H NMR (CDCl_3): δ 0.85 (s, 3H); 0.93 (s, 3H); 1.04 (d, 3H); 2.90 (s, 3H); 3.02 (dd, 1H); 4.76 (m, 1H); 6.88 (d, 2H); and 7.50 (d, 2H).

EXAMPLE 23

3-Oxo-4-aza-4,7 β -dimethyl-16 β -(4-chlorophenoxy)-5 α -androstane

This compound was prepared in a similar fashion as Example 21, but substituting 1-chloro-4-fluorobenzene in place of 4-fluorobenzonitrile; mass spectrum: m/z 430 (M+1).

400 MHz ^1H NMR (CDCl_3): δ 0.85 (s, 3H); 0.93 (s, 3H); 1.03 (d, 3H); 2.90 (s, 3H); 3.02 (dd, 1H); 5.28 (m, 1H); 6.74 (d, 2H); and 7.19 (d, 2H).

EXAMPLE 243-Oxo-4-aza-4,7 β -dimethyl-16 β -(4-fluorophenoxy)-5 α -androstane

This compound was prepared in a similar fashion as Example 21, but substituting 1,4-difluorobenzene in place of 4-fluorobenzonitrile; mass spectrum: m/z 414 ($M+1$).
400 MHz ^1H NMR (CDCl_3): δ 0.85 (s, 3H); 0.94 (s, 3H); 1.04 (d, 3H); 2.91 (s, 3H); 3.02 (dd, 1H); 4.65 (m, 1H); 6.75 (m, 2H); and 6.92 (m, 2H).

EXAMPLE 253-Oxo-4-aza-4,7 β -dimethyl-16 α -hydroxy-5 α -androstane

Step 1: 3-Oxo-4-aza-4,7 β -dimethyl-16 α -(4-nitrobenzoyloxy)-5 α -androstane

To a solution of 3-oxo-4-aza-4,7 β -dimethyl-16 β -hydroxy-5 α -androstane (178 mg, 0.560 mmol) in dry benzene (10 mL) were added triphenylphosphine (294 mg, 1.12 mmol), 4-nitrobenzoic acid (187 mg, 1.12 mmol), and diethyl azodicarboxylate (176 μL , 1.12 mmol). The reaction mixture was heated for one hour at 80° (oil bath temperature) under a nitrogen atmosphere. After removal of the benzene by evaporation under diminished pressure, the crude product mixture was subjected to flash silica gel chromatography using 2% methanol/methylene chloride as eluant to give desired product contaminated with some triphenylphosphine (404 mg) which was saponified as described in Step 2.

400 MHz ^1H NMR (CDCl_3): δ 0.80 (s, 3H); 0.88 (s, 3H); 1.03 (d, 3H); 2.90 (s, 3H); 3.05 (dd, 1H); and 5.48 (m, 1H).

Step 2: 3-Oxo-4-aza-4,7 β -dimethyl-16 α -hydroxy-5 α -androstane

The crude product from Step 1 (404 mg) was suspended in ethanol (5 mL) and treated with 0.4 N sodium hydroxide (1.82 mL, 0.728 mmol). After stirring 90 min at room temperature, the reaction mixture

- 54 -

was neutralized with several drops of glacial acetic acid, extracted with ethyl acetate (100 mL), washed with water (2 x 25 mL), saturated brine solution, dried (sodium sulfate), and evaporated. The product was obtained pure by flash silica gel chromatography using 20% acetone/methylene chloride as eluant; mass spectrum: m/z 319 (M). 400 MHz ^1H NMR (CDCl_3): δ 0.71 (s, 3H); 0.82 (s, 3H); 1.02 (d, 3H); 2.90 (s, 3H); 3.03 (dd, 1H); and 4.42 (m, 1H).

EXAMPLE 26

3-Oxo-4-aza-4,7 β -dimethyl-16 α -(n-propyloxy)-5 α -androstane

To a solution of 3-oxo-4-aza-4,7 β -dimethyl-16 α -hydroxy-5 α -androstane (20 mg, 0.063 mmol) in *N,N*-dimethylformamide (0.65 mL) was added potassium hydride (35 weight%) (15 mg, 0.126 mmol). After stirring for 15 min at room temperature under a nitrogen atmosphere, allyl bromide (27 μL , 0.315 mmol) was added and stirring was continued for 2 h. Additional amounts of potassium hydride (15 mg) and allyl bromide (27 μL) were added, and stirring was continued overnight. The mixture was diluted with diethyl ether (50 mL) and water (10 mL). The organic layer was washed with 1 N hydrochloric acid (10 mL), water (10 mL), saturated brine solution, dried (Na_2SO_4), and evaporated. The desired product was purified by flash silica gel chromatography using 2% methanol/methylene chloride as eluant. This material was hydrogenated in ethyl acetate (0.5 mL) in the presence of 10% palladium-on-charcoal for 2 hours. The catalyst was removed by filtration through a Millex-HV 0.45 μm Filter Unit. Purification was achieved by flash silica gel chromatography using 10% isopropanol/hexane as eluant to yield the title compound; mass spectrum: m/z 361 (M).

400 MHz ^1H NMR (CDCl_3): δ 0.76 (s, 3H); 0.82 (s, 3H); 0.90 (t, 3H); 1.02 (d, 3H); 2.90 (s, 3H); 3.02 (dd, 1H); 3.29 (t, 2H); and 3.98 (m, 1H).

- 55 -

EXAMPLE 273-Oxo-4-aza-4,16 α -dimethyl-16 β -hydroxy-5 α -androstane

To a solution of 4-aza-4-methyl-5 α -androstane-3,16-dione (50 mg, 0.165 mmol) cooled to -40° was added dropwise with stirring methylmagnesium bromide (3.0 M solution in diethyl ether) (275 μ L, 0.825 mmol). The reaction mixture was allowed to reach room temperature and stirred for 2 h under a nitrogen atmosphere. The reaction was quenched with saturated ammonium chloride solution (25 mL) and extracted with methylene chloride (2 x 50 mL). The combined organic extracts were washed with saturated brine solution, dried (sodium sulfate) and evaporated. The desired product was obtained pure by flash silica gel chromatography using 2% methanol/methylene chloride as eluant; mass spectrum: m/z 319 (M). 400 MHz 1 H NMR (CDCl₃): δ 0.88 (s, 3H); 0.98 (s, 3H); 1.31 (s, 3H); 2.90 (s, 3H); and 3.00 (dd, 1H).

EXAMPLE 283-Oxo-4-aza-4,7 β ,16 α -trimethyl-16 β -hydroxy-5 α -androstane

This compound was prepared in a similar fashion as Example 27, but substituting 4-aza-4,7 β -dimethyl-5 α -androstane-3,16-dione in place of 4-aza-4-methyl-5 α -androstane-3,16-dione as starting material; mass spectrum: m/z 333 (M). 400 MHz 1 H NMR (CDCl₃): δ 0.82 (s, 3H); 0.98 (s, 3H); 1.01 (d, 3H); 1.30 (s, 3H); 2.90 (s, 3H); and 3.00 (dd, 1H).

EXAMPLE 293-Oxo-4-aza-4,16 α -dimethyl-16 β -methoxy-5 α -androstane

To a solution of 3-oxo-4-aza-4,16 α -dimethyl-16 β -hydroxy-5 α -androstane (31 mg, 0.097 mmol) in N,N-dimethylformamide (0.5 mL) was added potassium hydride (35 weight%) (23 mg, 0.194 mmol). After stirring for 15 min at room temperature, iodomethane (32 μ L, 0.485

- 56 -

mmol) was added, and stirring was continued overnight at room temperature. The reaction mixture was diluted with diethyl ether, washed with 2N hydrochloric acid (10 mL), water (10 mL), saturated brine solution, dried (sodium sulfate), and evaporated. The desired product was obtained pure by flash silica gel chromatography using 2% methanol/methylene chloride as eluant; mass spectrum: m/z 333 (M). 400 MHz ^1H NMR (CDCl_3): δ 0.88 (s, 3H); 0.90 (s, 3H); 1.22 (s, 3H); 2.90 (s, 3H); 3.00 (dd, 1H); and 3.17 (s, 3H).

EXAMPLE 30

3-Oxo-4-aza-4,7 β ,16 α -trimethyl-16 β -methoxy-5 α -androstande

This compound was prepared in a similar fashion as Example 29, but substituting 3-oxo-4-aza-4,7 β ,16 α -trimethyl-16 β -hydroxy-5 α -androstande in place of 3-oxo-4-aza-4,16 α -dimethyl-16 β -hydroxy-5 α -androstande as starting material; mass spectrum: m/z 347 (M).

400 MHz ^1H NMR (CDCl_3): δ 0.82 (s, 3H); 0.90 (s, 3H); 1.02 (d, 3H); 1.22 (s, 3H); 2.90 (s, 3H); 3.00 (dd, 1H); and 3.18 (s, 3H).

EXAMPLE 31

3-Oxo-4-aza-4,7 β -dimethyl-16 β -methanethio-5 α -androstande

Step 1: 3-Oxo-4-aza-4,7 β -dimethyl-16 β -(acetylthio)-5 α -androstande

A 25-mL round-bottom flask was charged with dry tetrahydrofuran (4 mL) and triphenylphosphine (177 mg, 0.676 mmol) under a nitrogen atmosphere. The flask was cooled in an ice-bath and diisopropyl azodicarboxylate (133 μL , 0.676 mmol) was added, and the mixture was stirred for 30 min at 0°. To the reaction mixture was added a solution of 3-oxo-4-aza-4,7 β -dimethyl-16 α -hydroxy-5 α -androstande (108 mg, 0.338 mmol) and thiolacetic acid (49 μL , 0.676 mmol) in tetrahydrofuran (2.0 mL). The reaction mixture was stirred for 1 h at 0° and then an additional hour at room temperature. The mixture was

- 57 -

evaporated and subjected to flash chromatography on silica gel using 10% acetone/methylene chloride as eluant to give the desired product contaminated with some triphenylphosphine. The mixture was used without further purification in Step 2.

400 MHz ^1H NMR (CDCl_3): δ 0.80 (s, 3H); 0.82 (s, 3H); 1.00 (d, 3H); 2.28 (s, 3H); 2.90 (s, 3H); 3.00 (dd, 1H); and 3.80 (m, 1H).

Step 2: 3-Oxo-4-aza-4,7 β -dimethyl-16 β -(mercapto)-5 α -androstande

To a solution of product mixture from Step 1 (208 mg) in ethanol (4.0 mL) was added 0.4N sodium hydroxide (1.8 mL, 0.716 mmol) under a nitrogen atmosphere. The reaction mixture was stirred at room temperature for 1 h, neutralized with several drops of acetic acid, diluted with ethyl acetate (100 mL), washed with water (2 x 10 mL), saturated brine solution, dried (sodium sulfate), and evaporated. Pure 16-mercaptan was obtained by flash silica gel chromatography using 20% acetone/hexane as eluant.

400 MHz ^1H NMR (CDCl_3): δ 0.82 (s, 3H); 0.93 (s, 3H); 1.02 (d, 3H); 2.90 (s, 3H); 3.00 (dd, 1H); and 3.28 (m, 1H).

Step 3: 3-Oxo-4-aza-4,7 β -dimethyl-16 β -(methanethio)-5 α -androstande

To a solution of 3-oxo-4-aza-4,7 β -dimethyl-16 β -(mercapto)-5 α -androstande (18 mg, 0.054 mmol) in dry tetrahydrofuran (0.5 mL) was added sodium hydride (80% dispersion in mineral oil) (3.2 mg, 0.108 mmol) under a nitrogen atmosphere. After stirring 15 min at room temperature, iodomethane (17 μL , 0.270 mmol) was added, and stirring was continued for 3 h at room temperature. The reaction mixture was diluted with methylene chloride (50 mL), washed with water (10 mL), saturated brine solution, dried (sodium sulfate), and evaporated. Flash silica gel chromatography using 10% isopropanol/hexane as eluant afforded pure desired product; mass spectrum: m/z 349 (M).

400 MHz ^1H NMR (CDCl_3): δ 0.82 (s, 3H); 0.91 (s, 3H); 1.04 (d, 3H); 2.10 (s, 3H); 2.90 (s, 3H); 3.01 (dd, 1H); and 3.08 (m, 1H).

- 58 -

EXAMPLE 32**3-Oxo-4-aza-4,7 β -dimethyl-16 β -ethanethio-5 α -androstane**

This compound was prepared in a similar fashion as Example 31, but substituting iodoethane in place of iodomethane in Step 3; mass spectrum: m/z 363 (M).

400 MHz ^1H NMR (CDCl_3): δ 0.82 (s, 3H); 0.91 (s, 3H); 1.03 (d, 3H); 1.24 (t, 3H); 2.57 (q, 2H); 2.90 (s, 3H); 3.00 (dd, 1H); and 3.18 (m, 1H).

EXAMPLE 33**3-Oxo-4-aza-4,7 β -dimethyl-16 β -(1-propanethio)-5 α -androstane**

This compound was prepared in a similar fashion as Example 31, but substituting 1-iodopropane in place of iodomethane in Step 3; mass spectrum: m/z 377 (M).

400 MHz ^1H NMR (CDCl_3): δ 0.82 (s, 3H); 0.90 (s, 3H); 0.98 (t, 3H); 1.03 (d, 3H); 2.51 (t, 2H); 2.90 (s, 3H); 3.01 (dd, 1H); and 3.13 (m, 1H).

EXAMPLE 34**3-Oxo-4-aza-4,7 β -dimethyl-16 β -ethanesulfonyl-5 α -androstane**

To a solution of 3-oxo-4-aza-4,7 β -dimethyl-16 β -ethanethio-5 α -androstane (17 mg, 0.047 mmol) in methanol (1.0 mL) was added a solution of OXONE, monopersulfate compound (19 mg) in water (1 mL). After stirring 2 h at room temperature, an additional amount of OXONE (19 mg) in water (0.5 mL) was added, and stirring was continued for 10 min. The reaction mixture was diluted with water (25 mL) and extracted with methylene chloride (3 x 50 mL). The combined organic extracts were washed with saturated brine solution, dried (sodium sulfate), and evaporated. Flash silica gel chromatography using 2% methanol/methylene chloride as eluant afforded pure desired product; mass spectrum: m/z 395 (M).

400 MHz ^1H NMR (CDCl_3): δ 0.85 (s, 3H); 0.92 (s, 3H); 1.03 (d, 3H); 1.39 (t, 3H); 2.91 (s, 3H); 2.99 (q, 2H); 3.00 (dd, 1H); and 3.41 (m, 1H).

EXAMPLE 35**3-Oxo-4-aza-4,7 β -dimethyl-16 β -fluoro-5 α -androstane**

To a solution of 3-oxo-4-aza-4,7 β -dimethyl-16 α -hydroxy-5 α -androstane (18 mg, 0.056 mmol) in methylene chloride (0.5 mL) at room temperature diethylaminosulfur trifluoride (19 μ L, 0.144 mmol). After stirring one hour at room temperature, the reaction mixture was diluted with methylene chloride (25 mL), washed with water (25 mL), saturated sodium hydrogencarbonate solution (10 mL), saturated brine solution (10 mL), dried (sodium sulfate), and evaporated. The product was purified by flash silica gel chromatography using 10% acetone/methylene chloride as eluant to yield the title compound; mass spectrum: m/z 321 (M).

400 MHz ^1H NMR spectrum (CDCl_3): δ 0.87 (s, 3H); 0.92 (s, 3H); 1.04 (d, 3H); 2.90 (s, 3H); 3.01 (dd, 1H); and 5.12 (dm, 1H).

EXAMPLE 36

Preparation of 4-aza-4,7 β -dimethyl-5 α -androstan-3,17-dione
(Compound E in Scheme 5 above)

Step 1: Synthesis of 3-acetoxy-androst-5-en-17-ol

To a solution of 100 mg. (0.303 mmol) of 3-acetoxy-androst-5-en-17-one in 3 ml EtOH at -10°C , was added 22.9 mg (0.606 mmol) of sodium borohydride with stirring. After the reaction mixture was stirred for one and 1/2 hours, the mixture was diluted with 10 ml water, the ethanol solvent removed under vacuum, and the residue extracted with ethyl acetate. The organic layer was washed with aqueous Na_2CO_3 , brine, dried over sodium sulfate and concentrated to leave a residue of crude title compound. Proton NMR confirmed the assigned structure.

- 60 -

Step 2: **Synthesis of 3-acetoxy-androst-5-en-17-ol, 17-t-butyl-dimethyl-silyl ether**

To a solution of the androstan-17-ol, from the previous synthesis being 4.5 g (13.55 mmol) in 50 ml dimethylformamide at 23°C was added 2.76 g (40-65 mmol) imidazole followed by 3.063 g (20.32 mmol) of t-butyldimethylsilyl chloride. The reaction mixture was stirred and a solid began to precipitate. Twenty additional ml of DMF were added and the mixture further stirred overnight. The mixture was poured into 1 liter water, the solid filtered and washed with water. The solid was dissolved in ethylacetate, the organic layer washed with brine and dried over sodium sulfate, concentrated to yield the silyl protected 17-ol title compound. The proton NMR confirmed the assigned structure.

Step 3: **7-one-17 β -ol, 17-t-butyldimethylsilyl ether**

To a solution of the TBMS protected 17-ol from the previous synthesis, being 5.6 g (12.55 mmol) in 100 ml acetonitrile at 23°C was added 90% t-butyl hydrogen peroxide, 3.958 g (43.92 mol), and 138 mg chromium hexacarbonyl. After refluxing the mixture under nitrogen for 24 hours, the reaction mixture was poured into one liter water, solid was filtered, the residue washed with 500 ml water and the residue dissolved in 350 ml methylene chloride. The organic layer was washed with brine, dried over sodium sulfate and concentrated to yield crude material. Thin layer chromatography (3:1 hexane/ethyl acetate on silica gel) showed the presence of starting material. The solid was purified by column chromatography over silica gel by elution with 7% ethyl acetate/hexane to yield the title compound. Proton NMR confirmed the assigned structure.

Step 4: **Synthesis of 3,7-dihydroxy-7-methyl-androst-5-en-17 β -ol, 17-t-butyldimethylsilyl ether**

To a solution of the product from the previous synthesis, being 440 mg (0.956 mmol) in dry tetrahydrofuran at 0°C was added dropwise methyl magnesium chloride over 5-10 minutes. The reaction mixture was then allowed to stir at room temperature for 24 hours, then

- 61 -

poured into saturated aqueous ammonium chloride. The THF solvent was removed under vacuum and the aqueous phase extracted with ethyl acetate. The organic layer was washed with brine, dried, concentrated to yield crude product. Proton NMR confirmed the assigned structure of the title compound which was used in the next step without further purification.

Step 5: **Synthesis of 7-methyl-androst-4,6-dien-3-one-17 β -ol, 17-t-butyltrimethylsilyl ether**

The above Grignard product, 3.5 g (7.142 mmol) was dissolved in 50 ml toluene/50 ml cyclohexanone and 20 ml of solvent distilled off under vacuum. To this was added 4.54 g aluminum isopropoxide and the reaction mixture refluxed overnight for 15 hours. The mixture was cooled, diluted with ethyl acetate, washed with sodium potassium tartarate, brine, and the organic layer was concentrated under vacuum and the residue steam distilled. The residue was extracted with ethyl acetate, washed with brine, dried and purified by column chromatography on silica gel, eluting with 5% EtOAc/hexane to yield the title compound.

Step 6: **Synthesis of 7 β -methyl-androst-5-en-3-one-17 β -ol, t-butyltrimethylsilyl ether**

To a solution of 370 mg of the product of the previous synthesis, in 5.5 ml ammonia, 1 ml THF, 1 ml toluene, was added 50 mg of metallic lithium in small pieces. After stirring the blue solution for 2 hours, a solution of 1,2-dibromethane in 2 ml THF was added. After stirring the solution at -78°C for 10 minutes, 250 mg of ammonium chloride was added and the mixture stirred for 10 minutes. The excess ammonia was removed by evaporation under a nitrogen stream. The reaction mixture was diluted with brine, extracted with ethyl acetate. The organic layer was washed with brine, dried and concentrated to yield crude material which was used as such in the next synthesis.

Step 7: **Synthesis of 7 β -methyl-androst-4-en-3-on-17 β -ol, t-butyltrimethylsilyl ether**

To a solution of the product of the previous synthesis, being 432 mg in 4 ml THF was added 150 microliters DBU (1,8-diazabicyclo-[5.4.0] undec-7-ene under nitrogen with stirring. The mixture was refluxed for 1.5 hours, then cooled, diluted with NH₄Cl solution. The solvent THF was removed under vacuum and the residue extracted with ethyl acetate. The organic layer was washed with brine, dried and concentrated under reduced pressure to yield crude material. The titled product was purified by chromatography on silica gel using 10% EtOAc/hexane as eluant.

Step 8: **Synthesis of 17 β -(t-butyltrimethylsilyloxy)-7 β -methyl-5-oxo-A-nor-3,5-secoandrost-3-ene acid**

To a solution of 884 mg of the product of the previous synthesis in 15 ml t-butyl alcohol at 80°C was added 248 mg sodium carbonate in 1.5 ml water followed by a dropwise addition over 15-20 minutes of a mixture of 2.273 g sodium periodate with 16.8 mg potassium permanganate in 8 ml water. The reaction mixture was heated at 80°C for 2 hours, cooled, filtered, the residue washed with water, and then the extract L- concentrated under vacuum. The extract was acidified with aqueous HCl, extracted with ethyl acetate and the organic layer washed with aqueous NaHSO₃, brine, dried and concentrated to yield crude 9. The proton NMR confirmed the assigned structure.

Step 9: **Synthesis of 4,7 β -dimethyl-4-aza-androst-5-en-3-one-17 β -ol, t-butyltrimethylsilyl ether**

To a solution of the product of the previous synthesis, 840 mg in 5 ml ethylene glycol, was added 1.5 g sodium acetate and 737 mg methylamine hydrochloride. After stirring the reaction mixture 4 hours at 180°C, the mixture was cooled, diluted with water, extracted with ethyl acetate, dried and concentrated to afford crude title compound. Proton NMR confirmed the assigned structure.

Step 10: Synthesis of 4,7 β -dimethyl-4-aza-androst-5-en-3-one-17 β -ol

To a solution of 700 mg of the product of the previous example, in 20 ml of acetonitrile at 0°C, was added 500 microliters. aqueous HF. After stirring the reaction mixture for one hour, the HF was neutralized with aqueous sodium carbonate, diluted with water, acetonitrile removed under vacuum, and the residue extracted with ethyl acetate. The organic layer was dried, concentrated to give crude title compound which was further purified by preparative chromatography on silica gel using 3:1 chloroform/acetone.

Step 11: Synthesis of 4,7 β -dimethyl-4-aza-androstan-3-one-17 β -ol

To a solution of the product of the previous synthesis, being 350 mg in 10 ml acetic acid was added 100 mg platinum dioxide and the resulting mixture was evacuated and flushed with hydrogen. The reaction was shaken overnight at room temperature under 40 Psig hydrogen pressure. The solution was filtered concentrated. The residue was worked up with ethyl acetate, the organic layer was then concentrated under vacuum, diluted with ethyl acetate, washed with aqueous NaHCO₃, brine, dried, concentrated to yield the title compound. Mass Spec: 320 (M+1).

Step 12: Synthesis of 4-aza-4,7 β -dimethyl-5 α -androstan-3,17-dione

The product of the previous synthesis, 1.013 g (3.176 mmol) was placed with 6 ml methylene chloride into a dry flask. Powdered molecular 4Å sieves, 1.6 g, and 0.558 g (4.76 mmol) of N-methyl-morpholine-N-oxide (NMO) and then tetrapropylammonium perruthanate (TPAP), 55 mg (0.159 mmol) were added. The reaction was stirred for 2 hours, diluted with 150 ml ethyl acetate and filtered. The filtrate was evaporated to dryness to yield crude product which was recrystallized from EtOAc to yield pure product, mp 135-138°C.

Elemental Analysis Calc'd for C₂₀H₃₁NO₂, mw=317.48

Calc'd: C, 75.67; H, 9.84; N, 4.41.

Found: C, 75.16; H, 10.22; N, 4.13.

Mass Spec. 318 (M+1).

- 64 -

The following Examples (37 to 49) are prepared in a similar fashion as Example 21, but substituting appropriate 4-fluoro derivatives in place of 4-fluorobenzonitrile.

EXAMPLE 37

3-Oxo-4-aza-4,7 β -dimethyl-16 β -(4-methylsulfonylphenoxy)-5 α -androstane

Mass spectrum: m/z 474 (M+1).

400 MHz ^1H NMR (CDCl_3): δ 0.85 (s, 3H); 0.93 (s, 3H); 1.04 (d, 3H); 2.90 (s, 3H); 3.00 (s, 3H); 4.80 (m, 1H); 6.92 (d, 2H); 7.81 (d, 2H).

EXAMPLE 38

3-Oxo-4-aza-4,7 β -dimethyl-16 β -(3-pyridyloxy)-5 α -androstane

Mass spectrum: m/z 397 (M+1).

400 MHz ^1H NMR (CDCl_3): δ 0.85 (s, 3H); 0.94 (s, 3H); 1.04 (d, 3H); 2.91 (s, 3H); 3.02 (dd, 1H); 4.75 (m, 1H); 7.21 (m, 2H); 8.22 (m, 2H).

EXAMPLE 39

3-Oxo-4-aza-4,7 β -dimethyl-16 β -(4-phenylphenoxy)-5 α -androstane

Mass spectrum: m/z 472 (M+1).

400 MHz ^1H NMR (CDCl_3): δ 0.85 (s, 3H); 0.96 (s, 3H); 1.05 (d, 3H); 2.91 (s, 3H); 3.02 (dd, 1H); 4.76 (m, 1H); 6.9 (d, 2H); 7.26 (m, 1H); 7.43 (m, 2H); 7.52 (m, 4H).

EXAMPLE 40

3-Oxo-4-aza-4,7 β -dimethyl-16 β -(3-chlorophenoxy)-5 α -androstane

Mass spectrum: m/z 431 (M+1).

400 MHz ^1H NMR (CDCl_3): δ 0.85 (s, 3H); 0.93 (s, 3H); 1.05 (d, 3H); 2.90 (s, 3H); 4.68 (m, 1H); 6.71 (m, 1H); 6.80 (m, 1H); 6.88 (m, 1H); 7.13 (m, 1H).

- 65 -

EXAMPLE 413-Oxo-4-aza-4,7 β -dimethyl-16 β -(4-trifluoromethoxyphenoxy)-5 α -androstande

Mass spectrum: m/z 480(M+1).

400 MHz ^1H NMR (CDCl_3): δ 0.85 (s, 3H); 0.94 (s, 3H); 1.04 (d, 3H); 2.91 (s, 3H); 3.02 (dd, 1H); 4.69 (m, 1H); 6.78 (m, 2H); 7.09 (m, 2H).EXAMPLE 423-Oxo-4-aza-4,7 β -dimethyl-16 β -(2-chlorophenoxy)-5 α -androstande

Mass spectrum: m/z 431 (M+1).

400 MHz ^1H NMR (CDCl_3): δ 0.85 (s, 3H); 0.99 (s, 3H); 1.04 (d, 3H); 2.91 (s, 3H); 3.03 (dd, 1H); 4.80 (m, 1H); 6.81 (m, 2H); 7.24 (m, 1H); 7.32 (m, 2H).EXAMPLE 433-Oxo-4-aza-4,7 β -dimethyl-16 β -(2-pyrazinyloxy)-5 α -androstande

Mass spectrum: m/z 398 (M+1).

400 MHz ^1H NMR (CDCl_3): δ 0.85 (s, 3H); 0.95 (s, 3H); 1.04 (d, 3H); 2.90 (s, 3H); 3.02 (dd, 1H); 5.34 (m, 1H); 8.04 (d, 2H); 8.15 (1H).EXAMPLE 443-Oxo-4-aza-4,7 β -dimethyl-16 β -(2-pyrimidinyloxy)-5 α -androstande

Mass spectrum: m/z 398 (M+1).

400 MHz ^1H NMR (CDCl_3): δ 0.85 (s, 3H); 0.95 (s, 3H); 1.04 (d, 3H); 2.90 (s, 3H); 3.02 (dd, 1H); 5.35 (m, 1H); 6.89 (m, 1H); 8.15 (d, 2H);

- 66 -

EXAMPLE 453-Oxo-4-aza-4,7 β -dimethyl-16 β -[4-(1-pyrrol)-phenoxy]-5 α -androstande

Mass spectrum: m/z 461 (M+1).

400 MHz ^1H NMR (CDCl_3): δ 0.85 (s, 3H); 0.95 (s, 3H); 1.05 (d, 3H); 2.91 (s, 3H); 4.73 (m, 1H); 6.30 (m, 2H); 6.84 (m, 2H); 6.96 (m, 2H); 7.25 (m, 2H).EXAMPLE 463-Oxo- 4-aza-4,7 β -dimethyl-16 β -(3-cyanophenoxy)-5 α -androstande

Mass spectrum: m/z 420 (M).

400 MHz ^1H NMR (CDCl_3): δ 0.85 (s, 3H); 0.93 (s, 3H); 1.04 (d, 3H); 2.91 (s, 3H); 3.02 (dd, 1H); 4.71 (m, 1H); 7.05 (m, 2H); 7.22 (m, 1H); 7.32 (m, 1H).EXAMPLE 473-Oxo-4-aza-4,7 β -dimethyl-16 β -(1-naphthyloxy)-5 α -androstande

Mass spectrum: m/z 445 (M).

400 MHz ^1H NMR (CDCl_3): δ 0.86 (s, 3H); 1.03 (s, 3H); 1.07 (d, 3H); 2.92 (s, 3H); 3.02 (dd, 1H); 6.70 (d, 1H); 7.32 (m, 2H); 7.44 (m, 2H); 7.78 (m, 1H); 8.24 (1H).EXAMPLE 483-Oxo-4-aza-4,7 β -dimethyl-16 β -(3-chloro-4-methylphenoxy)-5 α -androstande

Mass spectrum: m/z 445(M+1).

400 MHz ^1H NMR (CDCl_3): δ 0.84 (s, 3H); 0.92 (s, 3H); 1.04 (d, 2H); 2.26 (s, 3H); 2.92 (s, 3H); 4.76 (m, 1H); 6.62 (m, 1H); 6.81 (m, 1H); 7.12 (d, 1H).

EXAMPLE 493-Oxo-4-aza-4,7 β -dimethyl-16 β -[4-(5-oxazolyl)phenoxy]-5 α -androstane

Mass spectrum: m/z 463(M+1).

400 MHz ^1H NMR (CDCl_3): δ 0.85 (s, 3H); 0.95 (s, 3H); 1.05 (d, 3H); 2.91 (s, 3H); 4.76 (m, 1H); 6.86 (d, 2H); 7.21 (s, 1H); 7.53 (d, 2H); 7.84 (s, 1H).

EXAMPLE 503-oxo-4-aza-4,7 β -dimethyl-16 β -(4-nitrophenoxy)-5 α -androstane

This compound was prepared in a similar fashion as Example 21, but substituting 1-fluoro-4-nitrobenzene in place of 4-fluorobenzonitrile; 400 MHz ^1H NMR (CDCl_3): δ 0.85 (s, 3H); 0.94 (s, 3H); 1.05 (d, 3H); 2.92 (s, 3H); 3.03 (dd, 1H); 4.81 (q, 1H); 6.87 (d, 2H); 8.17 (d, 2H).

EXAMPLE 513-oxo-4-aza-4,7 β -dimethyl-16 β -(4-aminophenoxy)-5 α -androstane

To a solution of 3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-aminophenoxy)-5 α -androstane (163 mg, 0.36 mmol) in ethylacetate (8 mL) and methanol (8 mL) was added 10% Pd on carbon (25 mg, 0.23 mmol). It was then stirred for four hours under a hydrogen atmosphere at room temperature. It was then filtered through celite and evaporated to afford 148 mg of the title compound. No purification was needed. Mass spectrum: m/z 411 (M+1). 400 MHz ^1H NMR (CDCl_3): δ 0.84 (s, 3H); 0.94 (s, 3H); 1.37 (d, 3H); 2.90 (s, 3H); 3.03 (dd, 1H); 4.64 (q, 1H); 6.70 (d, 2H); 6.78 (d, 2H).

- 68 -

EXAMPLE 523-oxo-4-aza-4,7 β -dimethyl-16 β -(4-acetylaminophenoxy)-5 α -androstande

To a solution of 3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-amino-phenoxy)-5 α -androstande (48 mg, 0.116 mmol) in methylene chloride (1 mL) and pyridine (0.037 mL, 0.46 mmol) was added acetic anhydride (0.022 mL, 0.23 mmol) and DMAP (5 mg, 0.04 mmol). The reaction was stirred overnight at room temperature under a nitrogen atmosphere. It was then diluted with methylene chloride (50 mL), washed with water (50 mL) and brine (50 mL). The organic phase was then dried over sodium sulfate and evaporated. The crude product was purified by preparative TLC (silica gel, 1000 microns) using 5% methanol/methylene chloride to give 51 mg of the title compound. Mass spectrum: m/z 453 (M+1). 400 MHz ^1H NMR (CDCl_3): δ 0.84 (s, 3H); 0.93 (s, 3H); 1.04 (d, 3H); 2.13 (s, 3H); 2.92 (s, 3H); 3.03 (dd, 1H); 4.68 (q, 1H); 6.76 (d, 2H); 7.11 (s, 1H); 7.33 (d, 2H).

EXAMPLE 533-oxo-4-aza-4,7 β -dimethyl-16 β -(4-benzoylaminophenoxy)-5 α -androstande

This compound was prepared in a similar fashion as Example 52, but substituting benzoyl chloride in place of acetic anhydride, triethylamine in place of pyridine and DMAP was not used; mass spectrum: m/z 515 (M+1). 400 MHz ^1H NMR (CDCl_3): δ 0.84 (s, 3H); 0.94 (s, 3H); 1.05 (d, 3H); 2.92 (s, 3H); 3.04 (dd, 1H); 4.73 (q, 1H); 6.82 (d, 2H); 7.49 (m, 5H); 7.72 (s, 1H); 7.84 (d, 2H).

EXAMPLE 543-oxo-4-aza-4,7 β -dimethyl-16 β -(4-methylsulfonamidophenoxy)-5 α -androstande

This compound was prepared in a similar fashion as Example 52, but substituting tosyl chloride in place of acetic anhydride;

- 69 -

mass spectrum: m/z 565 ($M+1$). 400 MHz ^1H NMR (CDCl_3): δ 0.84 (s, 3H); 0.93 (s, 3H); 1.03 (d, 3H); 2.37 (s, 3H); 2.92 (s, 3H); 3.02 (dd, 1H); 4.63 (q, 1H); 6.34 (s, 1H); 6.67 (d, 2H); 6.91 (d, 2H); 7.19 (d, 2H); 7.56 (d, 2H).

EXAMPLE 55

3-oxo-4-aza-4-(2,4-dimethoxybenzyl)-7 β -methyl-16 β -hydroxy-5 α -androstane

The compound 55 was prepared in a similar fashion as compound 12 described in the Scheme 5, except that the corresponding benzyl analog of (Compound E in Scheme 5) was made via similar synthesis of Example 36, in which 2,4-dimethoxy-benzylamine was used in place of methylamine.

EXAMPLE 56

3-oxo-4-aza-4-(2,4-dimethoxybenzyl)-7 β -methyl-16 β -(4-chlorophenoxy)-5 α -androstane

This compound was prepared in a similar fashion as Example 23, but substituting 3-oxo-4-aza-4-(2,4-dimethoxy benzyl)-7 β -methyl-16 β -hydroxy-5 α -androstane in place of 3-oxo-4-aza-4,7 β -dimethyl-16 β -hydroxy-5 α -androstane. No purification was done prior to the next reaction.

EXAMPLE 57

3-oxo-4-aza-7 β -methyl-16 β -(4-chlorophenoxy)-5 α -androstane

To a solution of 3-oxo-4-aza-4-(2,4-dimethoxy benzyl)-7 β -methyl-16 β -(4-chlorophenoxy)-5 α -androstane (130 mg, 0.23 mmol) in methylene chloride (1 mL) was added trifluoroacetic acid (1 mL). The reaction was stirred overnight at room temperature. Then the solvent was evaporated and the residue taken up in methylene chloride. The organic phase was washed with saturated sodium bicarbonate and brine. It was

- 70 -

then dried over sodium sulfate and evaporated. The crude compound was purified by preparative TLC (silica gel, 1000 microns) using 20% acetone/methylene chloride to yield the title compound. 400 MHz ^1H NMR (CDCl_3): δ 0.86 (s, 3H); 0.93 (s, 3H); 1.01 (d, 3H); 3.07 (dd, 1H); 4.67 (q, 1H); 5.49 (s, 1H); 6.73 (d, 2H); 7.18 (d, 2H).

EXAMPLE 58

3-oxo-4-aza-7 β -methyl-16 β -phenoxy-5 α -androstane

To a solution of 3-oxo-4-aza-7 β -methyl-16 β -(4-chlorophenoxy)-5 α -androstane in methanol was added 20% Pd on carbon. This solution was shaken under a hydrogen atmosphere at 48 psig for one day. It was then filtered through celite and evaporated. The crude compound was then purified by flash silica gel chromatography using 20% acetone/methylene chloride to elute the title compound. 400 MHz ^1H NMR (CDCl_3): δ 0.86 (s, 3H); 0.95 (s, 3H); 1.01 (d, 2H); 3.08 (dd, 1H); 4.71 (q, 1H); 5.48 (s, 1H); 6.81 (d, 2H); 6.89 (t, 1H); 7.24 (t, 2H).

EXAMPLE 59

3-oxo-4-aza-7 β -methyl-16 β -phenoxy-5 α -androst-1-ene

To a solution of 3-oxo-4-aza-7 β -methyl-16 β -phenoxy-5 α -androstane (145 mg, 0.35 mmol) in toluene (3 mL) was added DDQ (95 mg, 0.42 mmol), BSTFA (360 mg, 1.4 mmol) and triflic acid (4.04 mg, 0.027 mmol). This solution was stirred overnight at room temperature under a nitrogen atmosphere. Then methylacetoacetate (4.06 mg, 0.035 mmol) was added and the solution was stirred. After one hour, the reaction was refluxed overnight. It was then poured into water (75 mL) containing sodium carbonate (160 mg) and sodium bisulfite (120 mg). The aqueous phase was then extracted with methylene chloride (40 mL) (3x) and the organic phases were combined. The organic phase was washed with water (50 mL) and brine (50 mL). It was dried over sodium sulfate and evaporated. The crude compound was purified by flash silica

- 71 -

gel chromatography using 15% acetone/methylene chloride to elute the title compound. 400 MHz ^1H NMR (CDCl_3): δ 0.92 (s, 3H); 0.96 (s, 3H); 1.02 (d, 3H); 3.34 (dd, 1H); 4.72 (q, 1H); 5.31 (s, 1H); 5.80 (d, 1H); 6.80 (d, 1H); 6.82 (d, 2H); 6.89 (t, 1H); 7.24 (t, 2H).

EXAMPLE 60

3-oxo-4-aza-7 β -methyl-16 β -(4-chlorophenoxy)-5 α -androst-1-ene

This compound was prepared in a similar fashion as Example 59, but substituting 3-oxo-4-aza-7 β -methyl-16 β -(4-chlorophenoxy)-5 α -androstane in place of 3-oxo-4-aza-7 β -methyl-16 β -phenoxy-5 α -androstane. 400 MHz ^1H NMR (CDCl_3): δ 0.92 (s, 3H); 0.95 (s, 3H); 1.02 (d, 2H); 3.34 (dd, 1H); 4.67 (q, 1H); 5.27 (s, 1H); 5.80 (d, 1H); 6.73 (d, 2H); 6.78 (d, 1H); 7.18 (d, 2H).

EXAMPLE 61

3-oxo-4-aza-4,7 β -dimethyl-16 β -phenoxy-5 α -androstane

To a solution of 3-oxo-4-aza-7 β -methyl-16 β -phenoxy-5 α -androstane (60 mg, 0.16 mmol) in N,N-dimethylformamide (1 mL) was added sodium hydride (8 mg, 0.21 mmol), a 60% dispersion in mineral oil. After stirring for 30 min at room temperature under a nitrogen atmosphere, methyl iodide (40 mg, 0.28 mmol) was added. The reaction was stirred overnight. It was diluted with ethylacetate (50 mL) and washed with 1N hydrochloric acid (50 mL), water (50 mL) and brine (50 mL). The organic phase was dried over sodium sulfate and evaporated. The crude product was purified by flash silica gel chromatography using 10% acetone/methylene chloride to elute the title compound. 400 MHz ^1H NMR (CDCl_3): δ 0.85 (s, 3H); 0.95 (s, 3H); 1.05 (d, 3H); 2.91 (s, 3H); 3.02 (dd, 1H); 4.72 (q, 1H); 6.81 (d, 2H); 6.89 (t, 1H); 7.24 (t, 2H).

- 72 -

EXAMPLE 623-oxo-4-aza-4,7 β -dimethyl-16 β -(4-chlorophenoxy)-5 α -androst-1-ene

This compound was prepared in a similar fashion as Example 61, but substituting 3-oxo-4-aza-7 β -methyl-16 β -(4-chlorophenoxy)-5 α -androstane in place of 3-oxo-4-aza-7 β -methyl-16 β -phenoxy-5 α -androstane. 400 MHz ^1H NMR (CDCl_3): δ 0.87 (s, 3H); 0.95 (s, 3H); 1.07 (d, 2H); 2.93 (s, 1H); 3.34 (dd, 1H); 4.68 (q, 1H); 5.84 (d, 1H); 6.69 (d, 1H); 6.73 (d, 2H); 7.18 (d, 2H).

EXAMPLE 633-oxo-4-aza-4-(2,4-dimethoxybenzyl)-7 β -methyl-16 β -(3-chloro-4-methylphenoxy)-5 α -androstane

This compound was prepared in a similar fashion as Example 56, but substituting 2-chloro-4-fluorotoluene in place of 1-chloro-4-fluorobenzene. No purification was done prior to the next reaction.

EXAMPLE 643-oxo-4-aza-7 β -methyl-16 β -(3-chloro-4-methylphenoxy)-5 α -androstane

This compound was prepared in a similar fashion as Example 57, but substituting 3-oxo-4-aza-4-(2,4-dimethoxy benzyl)-7 β -methyl-16 β -(3-chloro-4-methylphenoxy)-5 α -androstane in place of 3-oxo-4-aza-4-(2,4-dimethoxy benzyl)-7 β -methyl-16 β -(4-chlorophenoxy)-5 α -androstane. 400 MHz ^1H NMR (CDCl_3): δ 0.86 (s, 3H); 0.93 (s, 3H); 1.01 (d, 3H); 2.26 (s, 3H); 3.08 (dd, 1H); 4.66 (q, 1H); 5.59 (s, 1H); 6.62 (m, 1H); 6.81 (d, 1H); 7.06 (d, 1H).

- 73 -

EXAMPLE 653-oxo-4-aza-7 β -methyl-16 β -(4-methylphenoxy)-5 α -androstane

This compound was prepared in a similar fashion as Example 58, but substituting 3-oxo-4-aza-7 β -methyl-16 β -(3-chloro-4-methyl-phenoxy)-5 α -androstane in place of 3-oxo-4-aza-7 β -methyl-16 β -(4-chloro-phenoxy)-5 α -androstane. 400 MHz ^1H NMR (CDCl_3): δ 0.86 (s, 3H); 0.94 (s, 3H); 1.03 (d, 3H); 2.25 (s, 3H); 4.69 (q, 1H); 6.71 (d, 2H); 7.03 (d, 2H).

EXAMPLE 663-oxo-4-aza-7 β -methyl-16 β -(4-methylphenoxy)-5 α -androst-1-ene

This compound was prepared in a similar fashion as Example 59, but substituting 3-oxo-4-aza-7 β -methyl-16 β -(4-methylphenoxy)-5 α -androstane in place of 3-oxo-4-aza-7 β -methyl-16 β -phenoxy-5 α -androstane. 400 MHz ^1H NMR (CDCl_3): δ 0.92 (s, 3H); 0.96 (s, 3H); 1.03 (d, 3H); 2.25 (s, 3H); 3.34 (dd, 1H); 4.68 (q, 1H); 5.35 (s, 1H); 5.81 (d, 1H); 6.71 (d, 2H); 6.79 (d, 1H); 7.03 (d, 2H).

EXAMPLE 673-oxo-4-aza-7 β -methyl-16 β -(4-methylphenoxy)-5 α -androstane

This compound was prepared in a similar fashion as Example 61, but substituting 3-oxo-4-aza-7 β -methyl-16 β -(4-methyl-phenoxy)-5 α -androstane in place of 3-oxo-4-aza-7 β -methyl-16 β -phenoxy-5 α -androstane. 400 MHz ^1H NMR (CDCl_3): δ 0.85 (s, 3H); 0.94 (s, 3H); 1.04 (d, 3H); 2.25 (s, 3H); 2.91 (s, 3H); 3.05 (dd, 1H); 4.69 (q, 1H); 6.71 (d, 2H); 7.04 (d, 2H).

- 74 -

EXAMPLE 68**3-Oxo-4-aza-4,7 β -dimethyl-16 β -fluoro-5 α -androstande**

This compound was prepared by treatment of intermediate (12) (Scheme 5) with diethylaminosulfur trifluoride in methylene chloride at room temperature followed by chromatography on silica gel; 64% yield; m/z 321 (M); 400 MHz ^1H NMR spectrum (CDCl_3): 0.87 (s, 3H); 0.92 (s, 3H); 1.04 (d, 3H); 2.90 (s, 3H), 5.12 (m, H).

EXAMPLE 69**3-Oxo-4-aza-4,7 β -dimethyl-16 β -cyano-5 α -androstande**

This compound is obtained by conversion of intermediate (25) (Scheme 7) to its methanesulfonate derivative by treatment with methanesulfonyl chloride or methanesulfonic anhydride in methylene chloride in the presence of an organic base, such as pyridine and triethylamine, and 4-dimethylaminopyridine (DMAP). Displacement of the methanesulfonate group is effected by heating in an appropriate solvent, such as N,N-dimethylformamide or dimethylsulfoxide, in the presence of sodium or potassium cyanide.

EXAMPLE 70**3-Oxo-4-aza-4-methyl-16 β -(1-hexyl)-5 α -androstande****Step 1: 3-Oxo-4-aza-4-methyl-16 β -(1-hexenyl)-5 α -androstande**

To a 50-mL round-bottom flask under nitrogen was added 1-hexyl-triphenylphosphonium bromide (141 mg, 0.33 mmol) followed by freshly distilled tetrahydrofuran (1 mL). The mixture was cooled to 0°C, and butyllithium (2.5M solution in hexanes, 132 mL, 0.33 mmol) affording a bright orange solution. The solution was stirred at 0°C for 10 min., and was charged with a solution of 4-aza-4-methyl-5 α -androstan-3,16-dione (50 mg, 0.165 mmol) in tetrahydrofuran (0.5 mL). The reaction mixture was allowed to reach room temperature and stirred

- 75 -

overnight. The mixture was then partitioned between water (10 mL) and ethyl acetate (20 mL), the organic layer separated, washed with 0.5N hydrochloric acid (2 x 10 mL), saturated brine solution, dried (Na₂SO₄), and evaporated. The title compound was purified by flash silica gel chromatography using 1% methanol/methylene chloride as eluant. This material (29.6 mg) was used without further purification in Step 2.

Step 2: 3-Oxo-4-aza-4-methyl-16 β -(1-hexyl)-5 α -androstane

A solution of the product obtained in Step 1 (22 mg) in ethyl acetate (0.5 mL) was hydrogenated in the presence of platinum oxide (5 mg) under a balloon atmosphere of hydrogen gas for 1 hour at room temperature. The catalyst was removed by filtration through a Millex-HV disposable filter, and the filtrate was evaporated. The title compound was purified by flash silica gel chromatography using 20% acetone/hexane as eluant; yield 5.2 mg. Mass spectrum: m/z 374 (M+1). 400 MHz ¹H NMR (CDCl₃): δ 2.42 (dd, 2H), 2.90 (s, 3H), and 3.00 (dd, 1H).

EXAMPLE 71

4-aza-4,7 β -dimethyl-16b-(4-chlorobenzylidene)-5 α -androstan-3-one

Following Reaction Scheme 14, solution of 4-aza-4,7 β -dimethyl-5 α -androstan-3,17-dione (11) (32 mg, 0.1 mmol), sodium hydride (5 mg, 1.02 eq), diethyl 4-chlorobenzylphosphonate (27 mg, 1.02 eq) and DMF (0.5mL) were heated to 80°C for 1 hour. The reaction was cooled, diluted with dichloromethane and washed with water (x2), brine, dried over anhydrous magnesium sulfate, filtered and concentrated. The desired product was purified by silica gel chromatography (hexanes:isopropanol 4:1) t1.5:1 mixture of E/Z isomers: m/z=389 ¹H NMR(500 MHz, CDCl₃):0.75 (s, 3H); 0.82 (s, 3H); 0.90 (d, 3H); 2.96 (s, 3H); 3.08 (dd, 1H); 6.34 (s, 0.4H); 6.41 (s, 0.6H); 7.18-7.38 (m, 5H).

- 76 -

EXAMPLE 724-aza-4,7 β -dimethyl-16-benzylidene-5 α -androstan-3-one

This example was prepared in a similar fashion as 4-aza-4,7 β -dimethyl-16-(4-chlorobenzylidene)-5 α -androstan-3-one (71) but substituting diethyl benzylphosphonate for diethyl 4-chlorobenzylphosphonate: m/z= 390

¹H NMR(500 MHz, CDCl₃): 0.75 (s, 3H); 0.88 (s, 3H); 1.05 (d, 3H); 2.94 (s, 3H); 3.08 (dd, 1H); 6.28 (s, 0.4H); 6.35 (s, 0.6H); 7.15-7.35(m, 5H).

EXAMPLE 734-aza-4,7 β -dimethyl-16-(4-methylbenzylidene)-5 α -androstan-3-one

This example was prepared in a similar fashion as 4-aza-4,7 β -dimethyl-16-(4-chlorobenzylidene)-5 α -androstan-3-one (71) but substituting diethyl 4-methylphosphonate for diethyl 4-chlorobenzylphosphonate: m/z= 404

¹H NMR (500 MHz, CDCl₃): 0.78 (s, 3H); 0.85 (s, 3H); 1.1 (d, 3H); 2.32 (s, 3H); 2.94 (s, 3H); 3.08 (dd, 1H); 6.30 (s, 0.4H); 6.38 (s, 0.6H); 7.10-7.24 (m, 5H).

EXAMPLE 744-aza-4,7 β -dimethyl-16-(4-chlorobenzyl)-5 α -androstan-3-one

To a solution of 4-aza-4,7 β -dimethyl-16-(4-chlorobenzylidene)-5 α -androstan-3-one (71) (33 mg) in ethanol (4 mL) was added 5% Rh/C and the black suspension stirred under a hydrogen balloon. After 2 hours the mixture was filtered to remove catalyst, concentrated and purified on silica gel (hexanes:acetone 3:1) to give the desired product as a 3:1 mixture of isomers: m/z 427

¹H NMR (500 MHz, CDCl₃): 0.84 (s, 3H); 0.86 (s, 3H); 1.02 (d, 3H); 2.92 (bs, 2.7H); 2.93 (bs, 1.3H); 2.98 (s, 3H); 3.02 (dd, 1H); 7.10 (d, 2H), 7.25 (d, 2H).

- 77 -

EXAMPLE 754-aza-4,7 β -dimethyl-16-(4-methylbenzyl)-5 α -androstan-3-one

This example was prepared similarly to the procedure used for 4-aza-4,7 β -dimethyl-16-(4-chlorobenzyl)-5 α -androstan-3-one (74):
m/z= 408

¹H NMR (500 MHz, CDCl₃): 0.86 (s, 6H); 1.04 (d, 3H); 2.33 (s, 3H); 2.95 (s, 2H); 2.96 (s, 1H); 3.05 (dd, 1H); 7.06-7.11 (m, 4H).

EXAMPLE 764-aza-4,7 β -dimethyl-16-(3-pyridylmethyl)-5 α -androstan-3-one

This example was prepared similarly to the procedure used for 4-aza-4,7 β -dimethyl-16-(4-chlorobenzyl)-5 α -androstan-3-one (74) except 3-pyridylmethyl-dimethylphosphonate was used:
m/z= 395

¹H NMR (500 MHz, CDCl₃): 0.89 (s, 3H); 0.88 (s, 3H); 1.03 (d, 3H); 2.93 (bs, 2H); 2.94 (bs, 1H); 2.98 (s, 3H); 3.04 (dd, 1H); 7.10 (d, 2H), 7.25 (d, 2H); 7.58 (s, 1H); 8.55 (s, 2H).

EXAMPLE 774-aza-4,7 β -dimethyl-16 α -methanesulfonyl-5 α -androstan-3-one

Following Reaction Scheme 13, to a solution of 4-aza-4,7 β -dimethyl-16 α -hydroxy-5 α -androstan-3-one (25) (65 mg, 0.2 mmol) in anhydrous dichloromethane was added a catalytic amount of DMAP followed with methanesulfonic anhydride (45 mg, 1.1 eq). After 15 min, the reaction was diluted with dichloromethane, washed with 1M HCl (x3), 1M sodium bicarbonate, water and brine, dried over anhydrous magnesium sulfate filtered and concentrated to yield the desired compound of sufficient purity: m/z=398

¹H NMR (500 MHz, CDCl₃): 0.78 (s, 3H); 0.85 (s, 3H); 1.02 (d, 3H); 2.95 (s, 3H); 3.1 (dd, 2H); 5.18 (m, 1H);

- 78 -

EXAMPLE 784-aza-4,7 β -dimethyl-16 β -thiophenoxy-5 α -androstan-3-one

To a solution of thiophenol (50 μ L, 2.5 eq) in anhydrous THF was added sodium hydride (20 mg, 2.6 eq). After stirring 20 min, a THF solution of 4-aza-4,7 β -dimethyl-16 α -methanesulfonyl-5 α -androstan-3-one (77) (65 mg, 0.2 mmol) was added and the mixture stirred 20 hours at ambient temperature. The reaction was quenched with 1M ammonium chloride and diluted with ethyl acetate, washed with water and brine, dried over anhydrous magnesium sulfate filtered and concentrated. The desired compound was purified by silica gel chromatography (hexanes:isopropanol 9:1): m/z =412
 ^1H NMR (500 MHz, CDCl_3): 0.86 (s, 3H); 0.96 (s, 3H); 1.06 (d, 3H); 2.94 (s, 3H); 3.06 (dd, 2H); 3.65 (m, 1H); 7.26-7.70 (m, 5H).

EXAMPLE 794-aza-4,7 β -dimethyl-16 β -(4-chlorothiophenoxy)-5 α -androstan-3-one

This compound was prepared in a similar fashion as 4-aza-4,7 β -dimethyl-16 β -thiophenoxy-5 α -androstan-3-one (78) but substituting 4-chlorothiophenol in place of thiophenol: m/z =446
 ^1H NMR (500 MHz, CDCl_3): 0.85 (s, 3H); 0.96 (s, 3H); 1.04 (d, 3H); 2.94 (s, 3H); 3.02 (dd, 2H); 3.61 (m, 1H); 7.22 (d, 2H); 7.32 (d, 2H).

EXAMPLE 804-aza-4,7 β -dimethyl-16 β -(4-fluorothiophenoxy)-5 α -androstan-3-one

This compound was prepared in a similar fashion as 4-aza-4,7 β -dimethyl-16 β -thiophenoxy-5 α -androstan-3-one (78) but substituting 4-fluorothiophenol in place of thiophenol: m/z =431
 ^1H NMR (500 MHz, CDCl_3): 0.85 (s, 3H); 0.96 (s, 3H); 1.05 (d, 3H); 2.92 (s, 3H); 3.03 (dd, 2H); 3.51 (m, 1H); 6.99 (d, 2H); 7.35 (d, 2H).

- 79 -

EXAMPLE 814-aza-4,7 β -dimethyl-16 β -(4-methylthiophenoxy)-5 α -androstan-3-one

This compound was prepared in a similar fashion as 4-aza-4,7 β -dimethyl-16 β -thiophenoxy-5 α -androstan-3-one (78) but substituting 4-methylthiophenol in place of thiophenol: m/z=426

¹H NMR(500 MHz, CDCl₃): 0.75 (s, 3H); 0.95 (s, 3H); 1.1 (d, 3H); 2.31 (s, 3H); 2.94 (s, 3H); 3.02 (dd, 2H); 3.59 (m, 1H); 7.09 (d, 2H); 7.22 (d, 2H).

EXAMPLE 824-aza-4,7 β -dimethyl-16 β -(4-methoxythiophenoxy)-5 α -androstan-3-one

This compound was prepared in a similar fashion as 4-aza-4,7 β -dimethyl-16 β -thiophenoxy-5 α -androstan-3-one (78) but substituting 4-methoxythiophenol in place of thiophenol: m/z= 443

¹H NMR (500 MHz, CDCl₃): 0.81 (s,3H); 0.93 (s,3H); 1.18 (d,3H); 2.93 (s,3H); 3.02 (dd,2H); 3.50 (m,1H); 3.81 (s,3H); 7.45 (d,2H); 7.67 (d,2H).

EXAMPLE 834-aza-4,7 β -dimethyl-16 β -phenylsulfinyl-5 α -androstan-3-one

To a solution of 4-aza-4,7 β -dimethyl-16 β -thiophenoxy-5 α -androstan-3-one (78) (20 mg, 0.05 mmol) in dichloromethane at 0°C was added mCPBA (11 mg, 1 eq) and the solution stirred 1 hour. The reaction was diluted with dichloromethane and washed with 1M sodium bicarbonate, water, brine and dried over anhydrous sodium sulfate. The desired compound was purified by silica gel chromatography to yield a 4.6:1 mixture of diastereomers: m/z=428

¹H NMR(500 MHz, CDCl₃): 0.83 (s, 3H); 0.92 (s, 3H); 1.01 (d, 3H); 2.92 (s, 3H); 3.01 (dd, 2H); 3.19 (m, 0.85H); 3.55 (m, 0.15H); 7.5-7.70 (m, 5H).

- 80 -

EXAMPLE 84**4-aza-4,7 β -dimethyl-16 β -phenylsulfonyl-5 α -androstan-3-one**

A solution of 4-aza-4,7 β -dimethyl-16 β -phenylsulfinyl-5 α -androstan-3-one (83) 912 mg, 0.03 mmol) in dichloromethane was treated with mCPBA (9 mg, 1.5 eq) for 3 hours. The reaction was diluted with dichloromethane and washed with 1M sodium bicarbonate, water, brine and dried over anhydrous sodium sulfate. The desired compound was purified by silica gel chromatography (hexanes:isopropanol 7:3):
m/z= 444

¹H NMR (500 MHz, CDCl₃): 0.85 (s, 3H); 0.91 (s, 3H); 1.0 (d, 3H); 2.95 (s, 3H); 3.05 (dd, 2H); 3.55 (m, 0.15H); 7.41 (t, 1H); 7.55 (t, 2H); 7.90 (d, 1H), 7.90 (d, 1H)

EXAMPLE 85**3-Oxo-4-aza-4,16 β -dimethyl-5 α -androstane**

This compound is made by converting the readily available 4-aza-4,16 β -dimethyl-androstan-3,17-dione to the 17-triflate. Reduction of the triflate through conventional methods yields the titled 16- β -methyl analog.

Mass spectrum: m/z 304 (M+1)

400MHz NMR (CDCl₃): δ 0.76 (s, 3H); 0.85 (s, 3H); 1.04 (d, 3H); 2.90 (s, 3H); 3.01 (dd, 1H).

BIOLOGICAL ASSAYS**Preparation of Human prostatic and scalp 5 α -reductases**

Samples of human tissue were pulverized using a freezer mill and homogenized in 40 mM potassium phosphate, pH 6.5, 5 mM magnesium sulfate, 25 mM potassium chloride, 1 mM phenylmethylsulfonyl fluoride, 1 mM dithiothreitol (DTT) containing 0.25 M sucrose using a Potter-Elvehjem homogenizer. A crude nuclear pellet was prepared by centrifugation of the homogenate at 1,500 x g for 15 min.

The crude nuclear pellet was washed two times and resuspended in two volumes of buffer. Glycerol was added to the resuspended pellet to a final concentration of 20%. The enzyme suspension was frozen in aliquots at -80°C. The prostatic and scalp reductases were stable for at least 4 months when stored under these conditions.

Cloned enzyme protocol:

For IC₅₀ determinations, the test 5 α -reductase 1 and 2 inhibitors were dissolved in ethanol and serially diluted to the appropriate concentration. The baculovirus-expressed recombinant type 1 5 α -reductase was preincubated with inhibitor (0.1-1,000 nM) in 40 mM sodium phosphate, pH 7.0, 500 μ M NADPH, 1mM DTT and 1 mg/ml BSA for 18 h at 4°C. The reaction was initiated by the addition of [7-³H]T (NEN, 20 Ci/mmol) and NADPH to a final concentration of 0.3 μ M and NADPH and incubated at 37°C for 90 min. Similarly, baculovirus-expressed type 2 5 α -reductase was preincubated with inhibitor (1-10,000 nM) in 40 mM sodium citrate, pH 5.5, 500 μ M NADPH, 1mM DTT and 1 mg/ml BSA for 18 h at 4°C. The reaction was initiated by the addition of [7-³H]T (NEN, 20 Ci/mmol) and NADPH to a final concentration of 0.3 μ M and 500 μ M, respectively. The conversion of T to DHT was monitored using a radioflow detector following separation by reverse phase HPLC (Whatman RACII C18 column, 1ml/min 0.1% TFA in water:methanol (42:58); retention times T, 6.3 min, DHT, 9.7 min).

5 α -reductase assay

The reaction mixture for the type 1 5 α -reductase contained 40 mM potassium phosphate, pH 6.5, 5 μ M [7-³H]-testosterone, 1 mM dithiothreitol and 500 μ M NADPH in a final volume of 100 μ l. The reaction mixture for the type 2 5 α -reductase contained 40 mM sodium citrate, pH 5.5, 0.3 μ M [7-³H]-testosterone, 1 mM dithiothreitol and 500 μ M NADPH in a final volume of 100 μ l. Typically, the assay was initiated by the addition of 50-100 μ g prostatic homogenate or 75-200 μ g scalp homogenate and incubated at 37°C. After 10-50 min the reaction

- 82 -

was quenched by extraction with 250 μ l of a mixture of 70% cyclohexane: 30% ethyl acetate containing 10 μ g each DHT and T. The aqueous and organic layers were separated by centrifugation at 14,000 rpm in an Eppendorf microfuge. The organic layer was subjected to normal phase HPLC (10 cm Whatman partisil 5 silica column equilibrated in 1 ml/min 70% cyclohexane: 30% ethyl acetate; retention times: DHT, 6.8-7.2 min; androstenediol, 7.6-8.0 min; T, 9.1-9.7 min). The HPLC system consisted of a Waters Model 680 Gradient System equipped with a Hitachi Model 655A autosampler, Applied Biosystems Model 757 variable UV detector, and a Radiomatic Model A120 radioactivity analyzer. The conversion of T to DHT was monitored using the radioactivity flow detector by mixing the HPLC effluent with one volume of Flo Scint 1 (Radiomatic). Under the conditions described, the production of DHT was linear for at least 25 min. The only steroids observed with the human prostate and scalp preparations were T, DHT and androstenediol.

Inhibition studies

Compounds were dissolved in 100% ethanol. IC₅₀ values represent the concentration of inhibitor required to decrease enzyme activity to 50% of the control. IC₅₀ values were determined using a 6 point titration where the concentration of the inhibitor was varied from 0.1 to 1000 nM.

Representative compounds of this invention were tested in the above described assay for 5 α -reductase type 1 and type 2 inhibition. For the inhibition of 5 α -reductase type 1, the compounds have IC₅₀ values lower than 600 nM, with the majority of compounds having IC₅₀ values ranging from about 0.3 nM to about 200 nM. For the inhibition of 5 α -reductase type 2, the same compounds have IC₅₀ values greater than about 155 nM, with the majority of compounds having IC₅₀ values greater than 1000 nM. Each compound tested has at least a 2-fold greater selectivity for inhibition of 5 α -reductase type 1 over type 2, with the majority of the compounds having a 10-fold or greater selectivity for inhibition of 5 α -reductase type 1 over type 2. These results demonstrate

the utility of the compounds of the instant invention for the treatment of androgenetic conditions.

A compound referred to herein as a 5 α -reductase 2 inhibitor is a compound that shows inhibition of the 5 α -reductase 2 isozyme in the above-described assay; likewise, a compound referred to herein as a 5 α -reductase 1 inhibitor is a compound that shows inhibition of the 5 α -reductase 1 isozyme in the above-described assay.

Human Dermal Papilla Cell Assay

The dermal papilla is a small group of cells at the base of each hair follicle, and it is presently thought that these cells are stem cells that form the basis for hair growth. These cells have been shown to have 5 alpha reductase activity, and it is therefore possible to test inhibitors of 5 alpha reductase in these cell culture systems.

Isolated and cultured dermal papilla cells are prepared according to the methods of Messenger, A.G., *The Culture of Dermal Papilla Cells From Human Hair Follicles*, Br. J. Dermatol. 110:685-689, 1984 and Itami, S. *et al.*, *5 α -Reductase Activity In Cultured Human Dermal Papilla Cells From Beard Compared With Reticular Dermal Fibroblasts*, J. Invest. Dermatol. 94:150-152, 1990. Beard dermal papilla cells and occipital scalp hair of two different individuals are used throughout the study. All experiments are performed at confluency after the fourth to sixth subculture. Confluent monolayers are rinsed twice with phosphate-buffered saline, scraped from dishes by rubber policemen, and collected into a centrifuge tube. The cell suspensions are centrifuged at 1,500 rpm for 10 min at 4°C. The pellets are resuspended in 20 mM Tris-HCl buffer, pH 7.5, at 4°C, containing 250 mM sucrose, 1 mM MgCl₂, and 2 mM CaCl₂, by vortexing and 10 passes through a 25-gauge needle. The crude homogenate is further homogenized by a teflon-glass homogenizer, and is used as the cell homogenate. For the study of subcellular localization of 5 α -reductase, the cell homogenate is centrifuged at 800 x g for 10 min to yield a crude nuclear pellet. The resultant supernatant is centrifuged at 10,000 x g for 15 min to produce a crude mitochondrial pellet. The supernatant is centrifuged at 100,000 x g

- 84 -

for 60 min to yield a microsomal pellet and cytosol. Each particulate fraction is washed twice and resuspended in the buffer.

A standard incubation mixture will consist of 50 nM [^3H]-testosterone, 1 mM NADPH, 100 mM sodium citrate, pH 5.5 or 100 mM Tris-HCl, pH 7.5, and 50 μl of the cell homogenate, in a final volume of 100 μl . Each tube contains 50-100 μg of cellular protein. Incubation is carried out at 37°C for 30 min. During this incubation, the reaction is proportional to the time. For the study of optimum pH, citrate buffer is used at pH 4.5-6.5, and the Tris HCl buffer at pH 7.0-9.0. The protein content is determined by the method of Lowry, *et al.*, *Protein Measurement With The Folin Phenol Reagent* J. Biol. Chem. 193:265-275, 1951.

After incubation, the reaction is stopped by adding 4 times volume of chloroform-methanol (2/1:V/V) containing 110 μg each of carrier steroids. The extracted steroids are analyzed by thin-layer chromatography as previously described by Gomez, *et al.*, *In Vitro Metabolism Of Testosterone-4- ^{14}C and Δ -androstene-3, 17-dione-4- ^{14}C In Human Skin*. Biochem. 7:24-32, 1968, and the purity of each steroid is determined by the recrystallization method. The activity of 5 α -reductase is expressed by the sum of dihydrotestosterone, androstanediol and androstanedione formed. [1,2- ^3H]-testosterone (55.2 Ci/mmol) is obtainable from New England Nuclear Corporation (Boston, MA) and unlabeled steroids can be purchased from Sigma Chemical Company (St. Louis, MO). Fetal calf serum is obtainable from Hazleton (Lenaxa, Kansas). All other chemicals are of reagent grade.

Fuzzy Rat Acne Model

Adult fuzzy rats are a variety of rat that has stunted hair growth, brown colored seborrhea covering their entire back skin and abnormally increased sebum production after puberty that has been demonstrated to be due to circulating androgens. 0.1, 0.05 and 0.025% solutions of a selected 5 α -reductase inhibitor of interest are prepared in a vehicle of propylene glycol, isopropanol, isopropyl myristate and water (50/30/2/18%), and is topically applied onto the backs of adult male

- 85 -

fuzzy rats, 0.2 ml per animal daily for 4 weeks. Controls receive the vehicle alone and 5 of them are castrated. After 2 weeks seborrhea will be dose-dependently depleted and after 4 weeks bromodeoxyuridine (BrdU, 200 mg/kg) is intraperitoneally injected 2 hours before sacrifice. The skin tissues are incubated with EDTA (20 mM) in phosphate buffer, 1.5 hours at 37°C. The pilo-sebaceous unit attached to the epidermis is striped from the dermis and fixed with formalin for immuno-staining of BrdU. DNA synthesis cells showing a BrdU-positive nucleus are located in the outer glandular border. The number of S-phase cells per lobe is determined with a micro-image apparatus. Using formalin fixed skin, frozen serial sections are stained with 1% osmium and the size of the lobes is measured. A positive inhibitor of skin 5 α -reductrase will induce suppression of sebum production by inhibiting the rate of glandular cell turnover, and showing reduced lobular size.

The following describes an example methodology that can be used for detection of hair growth.

MACROPHOTOGRAPHY AND GLOBAL PHOTOGRAPHY PROCEDURE FOR DETECTION OF HAIR GROWTH

A. Macrophotographic Procedure

Location: ID card

Haircount target area

Equipment: Film: Kodak-T-max 24 exposure each of same emulsion lot
number

Camera: Nikon N-6000

Lens: Nikkor 60 mm f2.8

Flashes: Nikon SB-21B Macroflash

Device: registration device

Photographic Procedure:

In these clinical photographs, the only variable allowed is the haircount. Film emulsion, lighting, framing, exposure, and reproduction ratios are held constant.

- 86 -

1. The haircount area on the patient is prepared as follows: A small (~1mm) dot tattoo is placed at the beginning of the study at the leading edge of the bald area directly interior to the center of the vertex bald spot, using a commercial tattooing machine or manually (needle and ink). An area approximately one square inch in size, centered at the tattoo at the leading edge of the balding area, is clipped short (~2mm). Cut hairs are removed from the area to be photographed, using tape. Compressed air and/or ethanol wipes may also be used to facilitate removal of cut hairs.
2. Magnification: Each lens supplied has a fixed reproduction ratio of 1:1.2.
Aperture: Every photograph is taken at f/22.
Film: T-Max 100 (24 exposure) is used.
3. Patient's haircount target area. Three exposures (-2/3, 0, and +2/3 f-stop).

A trained technician places a transparency over the photograph and, using a felt tip pen, places a black dot over each visible hair. The dot map transparency is then counted using image analysis with computer assistance.

Photographs are coded with a random number corresponding to study site, visit number and patient allocation number to insure blinding to time. At Month 6, baseline and Month 6 photographs are counted and data analyzed for interim analysis. At Month 12, baseline, Month 6 and Month 12 photographs are counted and data analyzed for the primary endpoint.

Methodology for detection of hair growth is also described in Olsen, E.A. and DeLong, E., *J. American Academy of Dermatology*, Vol. 23, p. 470 (1990).

- 87 -

B. Global Photographic Procedure

Locations: Color card/patient Id

Global photograph

Equipment: Film: Kodachrome KR-64 24 exposure each of same
emulsion lot number

Camera: Nikon N-6000

Lens: Nikkor 60 mm f2.8

Flashes: Nikon SB-23

Color card/patient Id

Photographic Procedure

In these clinical photographs, the only variable allowed is the global area's appearance. Anything extraneous to the area (clothing, furniture, walls, etc.) is eliminated from the fields to be photographed.


1. Patients will have global photographs taken prior to hair clipping with the head in a fixed position (determined by the supplied stereotactic device). Hair on the patient's head is positioned consistently so as to not obscure the bald area.
2. Magnification: Each lens supplied has a fixed reproduction ratio of 1:6.
Aperture: Every photograph will be taken at f/11.
Film: Kodachrome (24 exposure) is used.
3. Patient's global photographs. Three exposures at zero compensation.

While the invention has been described and illustrated with reference to certain particular embodiments thereof, those skilled in the art will appreciate that various changes, modifications and substitutions can be made therein without departing from the spirit and scope of the invention. For example, effective dosages other than the particular dosages as set forth herein above may be applicable as a consequence of variations in the responsiveness of the mammal being treated for any of the indications for the compounds of the invention indicated above. Likewise, the specific pharmacological responses observed may vary

- 88 -

according to and depending upon the particular active compound selected or whether there are present pharmaceutical carriers, as well as the type of formulation and mode of administration employed, and such expected variations or differences in the results are contemplated in accordance with the objects and practices of the present invention. It is intended, therefore, that the invention be defined by the scope of the claims which follow and that such claims be interpreted as broadly as is reasonable.

- 89 -

WHAT IS CLAIMED IS:**1. A compound of the Formula I**

or a pharmaceutically acceptable salt or ester thereof
wherein:

the C1-C2 carbon-carbon bond may be a single bond, or a double bond as indicated by the dashed line;

R^1 is selected from the group consisting of hydrogen and C1-10 alkyl;

R^2 is selected from the group consisting of hydrogen and C1-10 alkyl;

one of R^3 and R^4 is selected from the group consisting of hydrogen and methyl, and the other is selected from the group consisting of:

- (a) amino;
 - (b) cyano;
 - (c) fluoro,
 - (d) methyl;
 - (e) OH;
 - (f) $-C(O)NR_bR_c$, where R_b and R_c are independently H, C1-6 alkyl, aryl, or arylC1-6alkyl; wherein the alkyl moiety can be substituted with 1-3 of: halo; C1-4alkoxy; or trifluoromethyl; and the aryl moiety can be substituted with 1-3 of: halo; C1-4alkyl; C1-4 alkoxy; or trifluoromethyl;
 - (g) C1-10 alkyl-X-;
 - (h) C2-10 alkenyl-X-;
- wherein the C1-10 alkyl in (g) and C2-10 alkenyl in (h) can be unsubstituted or substituted with one to three of:

- 90 -

- i) halo; hydroxy; cyano; nitro; mono-, di- or trihalomethyl; oxo; hydroxysulfonyl; carboxy;
 - ii) hydroxyC₁₋₆alkyl; C₁₋₆alkyloxy; C₁₋₆alkylthio; C₁₋₆alkylsulfonyl; C₁₋₆alkyloxycarbonyl; in which the C₁₋₆ alkyl moiety can be further substituted with 1-3 of: halo; C₁₋₄ alkoxy; or trifluoromethyl;
 - iii) arylthio; aryl; aryloxy; arylsulfonyl; aryloxycarbonyl; in which the aryl moiety can be further substituted with 1-3 of: halo; C₁₋₄ alkyl; C₁₋₄ alkoxy; or trifluoromethyl;
 - iv) -C(O)NR_bR_c; -N(R_b)-C(O)-R_c; -NR_bR_c; where R_b and R_c are defined above;
- (i) aryl-X-;
- (j) heteroaryl-X-, wherein heteroaryl is a 5, 6 or 7 membered heteroaromatic ring containing at least one member selected from the group consisting of: one ring oxygen atom, one ring sulfur atom, 1-4 ring nitrogen atoms, or combinations thereof; in which the heteroaromatic ring can also be fused with one benzo or heteroaromatic ring;
- wherein the aryl in (i) and heteroaryl in (j) can be unsubstituted or substituted with one to three of:
- v) halo; hydroxy; cyano; nitro; mono-, di- or trihalomethyl; mono-, di- or trihalomethoxy; C₂₋₆ alkenyl; C₃₋₆ cycloalkyl; formyl; hydrosulfonyl; carboxy; ureido;
 - vi) C₁₋₆ alkyl; hydroxy C₁₋₆ alkyl; C₁₋₆ alkyloxy; C₁₋₆ alkyloxy C₁₋₆alkyl; C₁₋₆ alkylcarbonyl; C₁₋₆ alkylsulfonyl; C₁₋₆ alkylthio; C₁₋₆ alkylsulfinyl; C₁₋₆ alkylsulfonamido; C₁₋₆ alkylarylsulfonamido; C₁₋₆ alkyloxy-carbonyl; C₁₋₆ alkyloxycarbonyl C₁₋₆alkyl; R_bR_cN-C(O)-C₁₋₆alkyl; C₁₋₆ alkanoylamino C₁₋₆ alkyl; aroylamino C₁₋₆ alkyl; wherein the C₁₋₆

- 91 -

- alkyl moiety can be substituted with 1-3 of:
halo; C₁₋₄alkoxy; or trifluoromethyl;
- vii) aryl; aryloxy; arylcarbonyl; arylthio;
arylsulfonyl; arylsulfinyl; arylsulfonamido;
aryloxycarbonyl; wherein the aryl moiety can
be substituted with 1-3 of: halo; C₁₋₄alkyl; C₁₋₄
alkoxy; or trifluoromethyl;
- viii) -C(O)NR_bR_c; -O-C(O)-NR_bR_c; -N(R_b)-C(O)-
R_c; -NR_bR_c; R_b-C(O)-N(R_c)-; where R_b and
R_c are defined in (f) above; and -N(R_b)-C(O)-
OR_g, wherein R_g is C₁₋₆alkyl or aryl, in which
the alkyl moiety can be substituted with 1-3 of:
halo; C₁₋₄alkoxy; or trifluoromethyl, and the
aryl moiety can be substituted with 1-3 of: halo;
C₁₋₄alkyl; C₁₋₄ alkoxy, or trifluoromethyl;
-N(R_b)-C(O) NR_cR_d, wherein R_d is selected
from H, C₁₋₆ alkyl, and aryl; in which said C₁₋₆
alkyl and aryl can be substituted as described
above in (f) for R_b and R_c;
- ix) a heterocyclic group, which is a 5, 6 or 7
membered ring, containing at least one member
selected from the group consisting of: one ring
oxygen atom, one ring sulfur atom, 1-4 ring
nitrogen atoms, or combinations thereof; in
which the heterocyclic ring can be aromatic,
unsaturated, or saturated, wherein the
heterocyclic ring can be fused with a benzo
ring, and
wherein said heterocyclic ring can be
substituted with one to three substituents, as
defined above for v), vi), vii) and viii),
excluding ix) a heterocyclic group; and
- (k) R³ and R⁴ taken together can be carbonyl oxygen;

- 92 -

- (l) R^3 and R^4 taken together can be $=CH-R_g$, wherein R_g is defined in *viii*); and wherein:

X is selected from the group consisting of:

-O-; -S(O)_n-; -C(O)-; -CH(R_e)-; -C(O)-O-*; -C(O)-N(R_e)-*;
 -N(R_e)-C(O)-O-*; -O-C(O)-N(R_e)-*; -N(R_e)C(O)-N(R_e)-;
 -O-CH(R_e)-*; -N(R_e)-; wherein R_e is H, C₁₋₃ alkyl, aryl, aryl-
 C₁₋₃ alkyl, or unsubstituted or substituted heteroaryl, as defined
 above in (j);

wherein the asterisk (*) denotes the bond which is attached to
 the 16-position in Structure I; and n is zero, 1 or 2.

2. The compound of Claim 1 wherein R^1 is hydrogen or methyl and R^2 is hydrogen or methyl.

3. The compound of Claim 1 wherein heteroaryl is selected from unsubstituted or substituted: pyridyl; furyl; pyrrolyl; thienyl; isothiazolyl; imidazolyl; benzimidazolyl; tetrazolyl; pyrazinyl; pyrimidyl; quinolyl; isoquinolyl; quinazolyl; benzofuryl; isobenzofuryl; benzothienyl; pyrazolyl; indolyl; isoindolyl; purinyl; carbazolyl; isoxazolyl; thiazolyl; isothiazolyl; oxazolyl; benzthiazolyl; and benzoxazolyl.

4. The compound of Claim 3 wherein said heteroaryl can be substituted with one to two substituents.

5. The compound of Claim 1 wherein said heterocyclic group is selected from unsubstituted or substituted: pyridyl; furyl; pyrrolyl; thienyl; isothiazolyl; imidazolyl; benzimidazolyl; tetrazolyl; pyrazinyl; pyrimidinyl; quinolyl; isoquinolyl; benzofuryl; isobenzofuryl; benzothienyl; pyrazolyl; indolyl; isoindolyl; purinyl; carbazolyl; isoxazolyl; thiazolyl; oxazolyl; benzthiazolyl; benzoxazolyl; and dihydro, tetrahydro, hexahydro; and saturated derivatives thereof.

- 93 -

6. The compound of Claim 5 wherein said heterocyclic group can be substituted with one to two substituents.

7. The compound of Claim 2 wherein:
one of R^3 and R^4 is selected from the group consisting of hydrogen and methyl, and the other is selected from the group consisting of:

- (b) cyano;
- (c) fluoro;
- (e) OH;
- (g) C₁-10 alkyl-X-; or C₁-10 alkyl-X-, where alkyl can be substituted with aryl, and wherein aryl in turn can be substituted with 1-2 of halo or C₁-6alkyl;
- (h) C₂-10 alkenyl-X-;
- (i) aryl-X-;
- (j) heteroaryl-X-, wherein heteroaryl is a 5 or 6 membered heteroaromatic ring containing 1-2 ring nitrogen atoms; wherein the aryl in (i) and heteroaryl in (j) can be unsubstituted or substituted with one to two of:
 - x) halo; cyano; nitro; trihalomethyl; trihalomethoxy; C₁-6 alkyl; aryl; C₁-6 alkylsulfonyl; C₁-6 alkyl-arylsulfonamino;
 - xi) -NR_bR_c; R_b-C(O)-N(R_c)-; wherein R_b and R_c are independently H, C₁-6 alkyl, aryl, or arylC₁-6alkyl; wherein the alkyl moiety can be substituted with 1-3 of: halo; C₁-4alkoxy; or trifluoromethyl; and the aryl moiety can be substituted with 1-3 of: halo; C₁-4alkyl; C₁-4 alkoxy; or trifluoromethyl;
 - xii) a heterocyclic group, which is a 5 membered aromatic ring, containing one ring nitrogen atom, or one ring oxygen and one ring nitrogen atom; and
- (k) wherein R^3 and R^4 taken together can be carbonyl oxygen; and wherein:

X is selected from the group consisting of:

- 94 -

-O-; -S(O)_n-; -CH(R_e)-; -C(O)-N(R_e)-*;

-O-C(O)-N(R_e)-*;

wherein R_e is H, C₁-3 alkyl, aryl, aryl C₁-3 alkyl;

wherein the asterisk (*) denotes the bond which is attached to the 16-position in Structure I; and n is zero or 2.

8. The compound of Claim 1 selected from the group consisting of:

4-aza-4,7β-dimethyl-5α-androstane-3,16-dione;

4-aza-4-methyl-5α-androstan-3,16-dione;

3-oxo-4-aza-4-methyl-16β-hydroxy-5α-androstane;

3-oxo-4-aza-4-methyl-16β-(benzylaminocarbonyloxy)-5α-androstane;

3-oxo-4-aza-4-methyl-16β-benzoylamino-5α-androstane;

3-oxo-4-aza-4-methyl-16β-methoxy-5α-androstane;

3-oxo-4-aza-4-methyl-16β-allyloxy-5α-androstane;

3-oxo-4-aza-4-methyl-16β-(n-propyloxy)-5α-androstane;

3-oxo-4-aza-4-methyl-16α-hydroxy-5α-androstane;

3-oxo-4-aza-4-methyl-16β-(phenoxy)-5α-androstane;

3-oxo-4-aza-7β-methyl-16β-(phenoxy)-5α-androst-1-ene;

3-oxo-4-aza-4-methyl-16α-methoxy-5α-androstane;

3-oxo-4-aza-4-methyl-16β-(4-chlorophenoxy)-5α-androstane;

3-oxo-4-aza-7β-methyl-16β-(4-chlorophenoxy)-5α-androst-1-ene;

3-oxo-4-aza-7β-methyl-16β-(4-chlorophenoxy)-5α-androstane;

3-oxo-4-aza-7β-methyl-16β-(3-chloro-4-methylphenoxy)-5α-androstane;

3-oxo-4-aza-7β-methyl-16β-(4-methylphenoxy)-5α-androstane;

3-oxo-4-aza-7β-methyl-16β-(4-methylphenoxy)-5α-androst-1-ene;

3-oxo-4-aza-7β-methyl-16β-[4-(1-pyrrolyl)phenoxy]-5α-androst-1-ene;

3-oxo-4-aza-4,7β-dimethyl-16β-hydroxy-5α-androstane;

3-oxo-4-aza-4,7β-dimethyl-16β-methoxy-5α-androstane;

3-oxo-4-aza-4,7β-dimethyl-16β-allyloxy-5α-androstane;

3-oxo-4-aza-4,7β-dimethyl-16β-(3,3-dimethylallyloxy)-5α-androstane;

3-oxo-4-aza-4,7β-dimethyl-16β-(n-propyloxy)-5α-androstane;

3-oxo-4-aza-4,7β-dimethyl-16β-(iso-pentyloxy)-5α-androstane;

3-oxo-4-aza-4,16α-dimethyl-16β-hydroxy-5α-androstane;

3-oxo-4-aza-4,7 β -dimethyl-16 β -ethyloxy-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -benzyloxy-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 α -hydroxy-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -methylthio-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(n-propylthio)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -fluoro-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -cyano-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(1-hexyl)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(n-propyl)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -benzyl-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-chlorobenzyl)-5 α -androstane;
3-oxo-4-aza-4,16 α -dimethyl-16 β -methoxy-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-cyanophenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(3-cyanophenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-nitrophenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(1-naphthyloxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(3-chloro-4-methylphenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-methylphenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(tert-butyloxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(3-methyl-1-butyloxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 α -(n-propyloxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-trifluoromethylphenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-trifluoromethoxyphenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -ethylthio-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -ethylsulfonyl-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-methylsulfonylphenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -[4-(4-tolylsulfonylamino)phenoxy]-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(3-pyridyloxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -[(4-phenyl)phenoxy]-5 α -androstane;

3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-fluorophenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(2-pyrazinyloxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -[4-(5-oxazolyl)phenoxy]-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(2-pyrimidinyloxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -[4-(1-pyrryl)phenoxy]-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-aminophenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-acetylaminophenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-benzoylaminophenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-chlorophenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(phenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(2-chlorophenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(3-chlorophenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-chlorophenoxy)-5 α -androst-1-ene;
3-oxo-4-aza-4,7 β -dimethyl-16-(4-chlorobenzylidene)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16-benzylidene-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16-(4-methylbenzylidene)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16-(4-chlorobenzyl)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16-(4-methylbenzyl)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16-(3-pyridylmethyl)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 α -methanesulfonyl-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -thiophenoxy-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-chlorothiophenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-fluorothiophenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-methylthiophenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-methoxythiophenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -phenylsulfinyl-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -phenylsulfonyl-5 α -androstane;
3-oxo-4-aza-4,7 β ,16 α -trimethyl-16 β -(4-trifluoromethylphenoxy)-5 α -androstane,
3-oxo-4-aza-4,7 β ,16 α -trimethyl-16 β -hydroxy-5 α -androstane;
3-oxo-4-aza-4,7 β ,16 α -trimethyl-16 β -methoxy-5 α -androstane;
and pharmaceutically acceptable salts thereof.

- 97 -

9. The compound of Claim 7 wherein the C₁-C₂ carbon-carbon bond is a single bond, R¹ is methyl, R² is methyl, R³ is selected from unsubstituted or substituted aryloxy, and R⁴ is hydrogen.

10. The compound of Claim 1 selected from the group consisting of:

3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-cyanophenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(3-cyanophenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-nitrophenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(1-naphthyloxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(3-chloro-4-methylphenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-methylphenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-trifluoromethylphenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-trifluoromethoxyphenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-methylsulfonylphenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -[4-(4-tolylsulfonylamino)phenoxy]-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -[(4-phenyl)phenoxy]-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-fluorophenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -[4-(5-oxazolyl)phenoxy]-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -[4-(1-pyrryl)phenoxy]-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-aminophenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-acetylamino)phenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-benzoylamino)phenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-chlorophenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(phenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(2-chlorophenoxy)-5 α -androstane;
3-oxo-4-aza-4,7 β -dimethyl-16 β -(3-chlorophenoxy)-5 α -androstane;
and the pharmaceutically acceptable salts thereof.

- 98 -

11. The compound 3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-chlorophenoxy)-5 α -androstane, or a pharmaceutically acceptable salt thereof.

12. A use of a compound of Claim 1, or a compound of Claim 1 in combination with an inhibitor of 5 α -reductase 2, for the preparation of a medicament useful for inhibiting 5 α -reductase or the isozymes thereof.

13. A use of a compound of Claim 1, or a compound of Claim 1 in combination with an inhibitor of 5 α -reductase 2, for the preparation of a medicament useful for treating the conditions of acne vulgaris, androgenic alopecia, female hirsutism, benign prostatic hyperplasia, prostatitis, and the treatment and/or prevention of prostatic cancer.

14. The use of Claim 13 wherein the androgenic alopecia is male pattern baldness.

15. The use of Claim 13 wherein the inhibitor of 5 α -reductase 2 is finasteride, epristeride or turosteride.

16. The use of Claim 13, wherein said compound of Claim 1 is 3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-chlorophenoxy)-5 α -androstane, or a pharmaceutically acceptable salt thereof.

17. The use of a compound of Claim 1, or a compound of Claim 1 in combination with an inhibitor of 5 α -reductase 2 as claimed in Claim 13, further in combination with a potassium channel opener for the preparation of a medicament useful for treating the condition of androgenic alopecia.

- 99 -

18. The use of Claim 17 wherein the potassium channel opener is minoxidil, or a pharmaceutically acceptable salt thereof.

19. The use of a compound of Claim 1, or a compound of Claim 1 in combination with an inhibitor of 5 α -reductase 2 as claimed in Claim 13, further in combination with a retinoid for the preparation of a medicament useful for treating the condition of acne vulgaris.

20. The use of Claim 19 wherein the retinoid is tretinoin, isotretinoin, or a pharmaceutically acceptable salt thereof.

21. The use of a compound of Claim 1, or a compound of Claim 1 in combination with an inhibitor of 5 α -reductase 2 as claimed in Claim 13, further in combination with an anti-androgen for the preparation of a medicament useful for treating and/or preventing the condition of prostatic cancer.

22. The use of Claim 21 wherein the anti-androgen is selected from: flutamide, spironolactone, casodex, or a pharmaceutically acceptable salt thereof.

23. The use of a compound of Claim 1, or a compound of Claim 1 in combination with an inhibitor of 5 α -reductase 2 as claimed in Claim 13 for the preparation of a medicament useful for treating benign prostatic hyperplasia.

- 100 -

24. The use of Claim 23 wherein the inhibitor of 5 α -reductase 2 is finasteride, epristeride, turosteride, or a pharmaceutically acceptable salt thereof.

25. The use of a compound of Claim 1, or a compound of Claim 1 in combination with an inhibitor of 5 α -reductase 2 as claimed in Claim 23, in further combination with an alpha-1 receptor antagonist for the preparation of a medicament useful for treating benign prostatic hyperplasia.

26. The use of Claim 25 wherein the alpha-1 receptor antagonist is terazosin.

27. A use of a compound of Claim 1, or a compound of Claim 1 in combination with an inhibitor of 5 α -reductase 2 for the preparation of a medicament useful for arresting and reversing androgenic alopecia and promoting hair growth.

28. A use of a compound of Claim 1, or a compound of Claim 1 in combination with an inhibitor of 5 α -reductase 2 for the preparation of a medicament for inhibiting the biosynthetic conversion of testosterone to dihydrotestosterone.

29. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and a therapeutically effective amount of a compound of Claim 1.

- 101 -

30. The pharmaceutical composition of Claim 29 further comprising a therapeutically effective amount of an inhibitor of 5 α -reductase 2, or a pharmaceutically acceptable salt thereof.

31. The pharmaceutical composition of Claim 30 wherein said 5 α -reductase 2 inhibitor is finasteride, epristeride or turosteride.

32. The pharmaceutical composition of Claim 29 wherein said compound of Claim 1 is 3-oxo-4-aza-4,7 β -dimethyl-16 β -(4-chlorophenoxy)-5 α -androstane, or a pharmaceutically acceptable salt thereof.

33. The pharmaceutical composition of Claim 29 further comprising a therapeutically effective amount of a potassium channel opener, or a pharmaceutically acceptable salt thereof.

34. The pharmaceutical composition of Claim 33 wherein said potassium channel opener is minoxidil, or a pharmaceutically acceptable salt thereof.

35. The pharmaceutical composition of Claim 33 further comprising a pharmaceutically acceptable carrier adapted for topical application.

36. The pharmaceutical composition of Claim 29 further comprising a retinoid, or a pharmaceutically acceptable salt thereof.

37. The pharmaceutical composition of Claim 36 wherein said retinoid is tretinoin, isotretinoin, or a pharmaceutically acceptable salt thereof.

38. The pharmaceutical composition of Claim 29 further comprising an anti-androgen, or a pharmaceutically acceptable salt thereof.

- 102 -

39. The pharmaceutical composition of Claim 38 wherein said anti-androgen is flutamide, spironolactone, casodex, or a pharmaceutically acceptable salt thereof.

40. The compound 3-oxo-4-aza-7 β -methyl-16 β -(4-methylphenoxy)-5 α -androst-1-ene.

41. A use of the compound of Claim 40 for the preparation of a medicament useful for inhibiting 5 α -reductase or the isozymes thereof.

42. A use of the compound of Claim 40 in combination with an inhibitor of 5 α -reductase 2 for the preparation of a medicament useful for inhibiting 5 α -reductase or the isozymes thereof in a human in need thereof.

43. The use according to Claim 42 wherein the human is male and the inhibitor of 5 α -reductase 2 is finasteride.

44. A use of a compound of Claim 40 for the preparation of a medicament useful for treating acne vulgaris.

45. A use of a compound of Claim 40 in combination with an inhibitor of 5 α -reductase 2 for the preparation of a medicament useful for treating acne vulgaris in a human in need thereof.

46. The use according to Claim 45 wherein the human is male and the inhibitor of 5 α -reductase 2 is finasteride.

47. A use of a compound of Claim 40 for the preparation of a medicament useful for treating androgenic alopecia.

48. A use of a compound of Claim 40 in combination with an inhibitor of 5α -reductase 2 for the preparation of a medicament useful for treating androgenic alopecia in a human need thereof.

49. The use according to Claim 48 wherein the human is male and the inhibitor of 5α -reductase 2 is finasteride.

50. A use of a compound of Claim 40 for the preparation of a medicament useful for treating female hirsutism.

51. A use of a compound of Claim 40 in combination with finasteride for the preparation of a medicament useful for treating female hirsutism.

52. A use of a compound of Claim 40 for the preparation of a medicament useful for treating benign prostatic hyperplasia.

53. A use of a compound of Claim 40 in combination with an inhibitor of 5α -reductase 2 for the preparation of a medicament useful for treating benign prostatic hyperplasia in a human in need thereof.

54. The use according to Claim 53 wherein the human is male and the inhibitor of 5α -reductase 2 is finasteride.

55. A use of a compound of Claim 40 in combination with an alpha-1 receptor antagonist for the preparation of a medicament useful for treating benign prostatic hyperplasia.

-104-

56. A use of a compound of Claim 40 for the preparation of a medicament useful for preventing prostatic cancer in a man at risk for developing prostatic cancer.

57. A use of a compound of Claim 40 in combination with an inhibitor of 5α -reductase 2 for the preparation of a medicament useful for preventing prostatic cancer in a man at risk for developing prostatic cancer.

58. The use according to Claim 57 wherein the inhibitor of 5α -reductase 2 is finasteride.

59. A use of a compound of Claim 40 for the preparation of a medicament useful for treating prostatic cancer in a man in need thereof.

60. A use of a compound of Claim 40 in combination with an inhibitor of 5α -reductase 2 for the preparation of a medicament useful for treating prostatic cancer in a man in need thereof.

61. The use according to Claim 60 wherein the inhibitor of 5α -reductase 2 is finasteride.