

Novel carboxylic acid derivatives, their preparation and use

The present invention relates to novel carboxylic acid
 5 derivatives, to their preparation and to their use.


Endothelin is a peptide which is composed of 21 amino acids and is synthesized and released by the vascular endothelium. Endothelin exists in three isoforms, ET-1, ET-2 and ET-3. In the
 10 following text, "Endothelin" or "ET" signifies one or all isoforms of endothelin. Endothelin is a potent vasoconstrictor and has a potent effect on vessel tone. It is known that this vasoconstriction is caused by binding of endothelin to its receptor (Nature, 332, 1988, 411-415; FEBS Letters, 231, 1988,
 15 440-444 und Biochem. Biophys. Res. Commun., 154, 1988, 868-875).

Increased or abnormal release of endothelin causes persistent vasoconstriction in the peripheral, renal and cerebral blood vessels, which may lead to illnesses. It has been reported in the
 20 literature that elevated plasma levels of endothelin were found in patients with hypertension, acute myocardial infarct, pulmonary hypertension, Raynaud's syndrome, atherosclerosis and in the airways of asthmatics (Japan J. Hypertension, 12, (1989), 79, J. Vascular Med. Biology 2, (1990) 207, J. Am. Med. Associa-
 25 tion 264, (1990) 2868).

Accordingly, substances which specifically inhibit the binding of endothelin to the receptor ought also to antagonize the various
 30 abovementioned physiological effects of endothelin and therefore be valuable drugs.

We have found that certain carboxylic acid derivatives are good inhibitors of endothelin receptors.

35 The invention relates to carboxylic derivatives of the formula I



where R is tetrazole [sic], nitrile [sic], a group COOH or a radical which can be hydrolyzed to COOH, and the other
 45 substituents have the following meanings:

R² hydrogen, hydroxyl, NH₂, NH(C₁-C₄-alkyl), N(C₁-C₄-alkyl)₂,

2

halogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy or C₁-C₄-alkylthio, or CR² is linked to CR¹² as indicated below to form a 5- or 6-membered ring;

5 X nitrogen or CR¹² where R¹² is hydrogen or C₁₋₅-alkyl, or CR¹² forms together with CR² or CR³ a 5- or 6-membered alkylene or alkenylene ring which can be substituted by one or two C₁₋₄-alkyl groups and in which in each case one methylene group can be replaced by oxygen, sulfur, -NH or -NC₁₋₄-alkyl;

10

Y nitrogen or methine;

R³ hydrogen, hydroxyl, NH₂, NH(C₁-C₄-alkyl), N(C₁-C₄-alkyl)₂, halogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, -NH-O-C₁₋₄-alkyl, C₁-C₄-alkylthio; or CR³ is linked to CR¹² as indicated above to form a 5- or 6-membered ring;

15

R⁴ and R⁵ (which can be identical or different):

20

phenyl or naphthyl, each of which can be substituted by one or more of the following radicals: halogen, nitro, cyano, hydroxyl, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, phenoxy, C₁-C₄-alkylthio, amino, C₁-C₄-alkylamino or C₁-C₄-dialkylamino; or

25

phenyl or naphthyl, which are connected together in ortho positions by a direct linkage, a methylene, ethylene or ethenylene group, an oxygen or sulfur atom or an SO₂-, NH- or N-alkyl group, or C₃-C₇-cycloalkyl;

30

R⁶ hydrogen, C₁-C₈-alkyl, C₃-C₆-alkenyl, C₃-C₆-alkynyl or C₃-C₈-cycloalkyl, it being possible for each of these radicals to be substituted one or more times by: halogen, hydroxyl, mercapto, carboxyl, nitro, cyano, C₁-C₄-alkoxy, C₃-C₆-alkenyloxy, C₃-C₆-alkynyloxy, C₁-C₄-alkylthio, C₁-C₄-haloalkoxy, C₁-C₄-alkylcarbonyl, C₁-C₄-alkoxycarbonyl, C₃₋₈-alkylcarbonylalkyl, C₁-C₄-alkylamino, di-C₁-C₄-alkylamino, phenyl or phenoxy or phenyl, substituted one or more times, e.g. once to three times, by halogen, nitro, cyano, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy or C₁-C₄-alkylthio;

35

40

phenyl or naphthyl, each of which can be substituted by one or more of the following radicals: halogen, nitro, cyano, hydroxyl, amino, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, phenoxy, C₁-C₄-alkylthio, C₁-C₄-alkylamino,

45

C₁-C₄-dialkylamino, dioxomethylene [sic] or dioxoethylene [sic];

- 5 a five- or six-membered heteroaromatic system which contains one to three nitrogen atoms and/or one sulfur or oxygen atom and which can carry one to four halogen atoms and/or one or two of the following radicals: C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-alkylthio, phenyl, phenoxy or phenylcarbonyl, it being possible for the phenyl radicals in turn to carry one to five halogen atoms and/or one to three of the following radicals: C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy and/or C₁-C₄-alkylthio;
- 10
- 15 Z sulfur or oxygen.

The compounds, and the intermediates for preparing them, such as II and IV, may have one or more asymmetrically substituted carbon atoms. Such compounds may be in the form of the pure enantiomers or pure diastereomers or a mixture thereof. The use of an enantiomerically pure compound as active substance is preferred.


20

The invention furthermore relates to the use of the abovementioned carboxylic acid derivatives for producing drugs, in particular for producing endothelin receptor inhibitors.

25


Compounds of the general formula IV where Z is sulfur or oxygen can be prepared, also in enantiomerically pure form, as described in P 44 36 851.8.

30



40 Compounds according to the invention in which the substituents have the meanings stated under the general formula I can be prepared, for example, by reacting the carboxylic acid derivatives of the general formula IV in which the substituents have the stated meanings with compounds of the general formula V,

45



where R^{13} is halogen or $R^{14}-SO_2-$, where R^{14} can be C_1-C_4 -alkyl, C_1-C_4 -haloalkyl or phenyl. The reaction preferably takes place in an inert solvent or diluent with the addition of a suitable base, i.e. a base which deprotonates the intermediate IV, at a temperature in the range from room temperature to the boiling point of the solvent.

Examples of such solvents or diluents are aliphatic, alicyclic and aromatic hydrocarbons, each of which may be chlorinated, such as hexane, cyclohexane, petroleum ether, naphtha, benzene, toluene, xylene, methylene chloride, chloroform, carbon tetrachloride, ethyl chloride and trichloroethylene, ethers such as diisopropyl ether, dibutyl ether, methyl tert-butyl ether, propylene oxide, dioxane and tetrahydrofuran, nitriles, such as acetonitrile and propionitrile, acid amides such as dimethylformamide, dimethylacetamide and N-methylpyrrolidone, sulfoxides and sulfones, such as dimethyl sulfoxide and sulfolane.

Compounds of the formula V are known, and some of them can be bought, or they can be prepared in a conventional manner.

The base which can be used is an alkali metal or alkaline earth metal hydride such as sodium hydride, potassium hydride or calcium hydride, a carbonate such as an alkali metal carbonate, e.g. sodium or potassium carbonate, an alkali metal or alkaline earth metal hydroxide such as sodium or potassium hydroxide, an organometallic compound such as butyllithium, or an alkali metal amide such as lithium diisopropylamide.

Compounds of the formula I can also be prepared by starting from the corresponding carboxylic acids, i.e. compounds of the formula I where R is COOH, and first converting the latter in a conventional way into an activated form, such as an acid halide, an anhydride or imidazolidine, and then reacting the latter with an appropriate hydroxyl compound HOR^8 . This reaction can be carried out in conventional solvents and often requires the addition of a base, in which case the abovementioned are suitable. These two steps can also be simplified, for example, by allowing the

5


carboxylic acid to act on the hydroxyl compound in the presence of a dehydrating agent such as a carbodiimide.

- Moreover, compounds of the formula I can also be prepared by starting from salts of the corresponding carboxylic acids, i.e. from compounds of the formula I where R is a group COR^1 and R^1 is OM, where M can be an alkali metal cation or the equivalent of an alkaline earth metal cation. These salts can be reacted with many compounds of the formula $\text{R}^1\text{-A}$ where A is a conventional nucleofugic leaving group, for example halogen such as chlorine, bromine, iodine or unsubstituted or halogen-, alkyl- or haloalkyl-substituted aryl- or alkylsulfonyl, such as toluenesulfonyl and methylsulfonyl, or another equivalent leaving group. Compounds of the formula $\text{R}^1\text{-A}$ with a reactive substituent A are known or can easily be obtained with general expert knowledge. This reaction can be carried out in conventional solvents and advantageously takes place with the addition of a base, in which case the abovementioned are suitable.
- The radical R in formula I can be varied widely. R is, for example, a group



where R^1 has the following meanings:

- a) hydrogen;
- b) a succinylimidoxy [sic] group;
- c) a 5-membered heteroaromatic system linked via a nitrogen atom, such as pyrrolyl, pyrazolyl, imidazolyl and triazolyl, which may then carry one or two halogen atoms or one or two $\text{C}_1\text{-C}_4\text{-alkyl}$ or one or two $\text{C}_1\text{-C}_4\text{-alkoxy}$ groups.
- d) R^1 is furthermore a group



- where k can assume the values 0, 1 and 2, p can assume the values 1, 2, 3 and 4, and R^7 is

C₁-C₄-alkyl, C₃-C₇-cycloalkyl, C₃-C₆-alkenyl, C₃-C₆-alkynyl or unsubstituted or substituted phenyl which can be substituted by one or more, e.g. one to three, of the following radicals: halogen, nitro, cyano, C₁-C₄-alkyl, C₁-C₄-haloalkyl, hydroxyl, C₁-C₄-alkoxy, C₁-C₄-alkylthio, mercapto, amino, C₁-C₄-alkylamino, C₁-C₄-dialkylamino;

e) R¹ is furthermore a radical OR⁸ where R⁸ is:

hydrogen, the cation of an alkali metal such as lithium, sodium, potassium, or the cation of an alkaline earth metal such as calcium, magnesium and barium or a physiologically tolerated organic ammonium ion such as tertiary C₁-C₄-alkylammonium or the ammonium ion;

C₃-C₈-cycloalkyl such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and [sic] cycloheptyl, cyclooctyl,

C₁-C₈-alkyl, in particular C₁-C₄-alkyl such as methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, tert-butyl;

CH₂-phenyl, which may be substituted by one or more of the following radicals: halogen, nitro, cyano, C₁-C₄-alkyl, C₁-C₄-haloalkyl, hydroxyl, C₁-C₄-alkoxy, mercapto, C₁-C₄-alkylthio, amino, C₁-C₄-alkylamino, C₁-C₄-dialkylamino, a C₃-C₆-alkenyl or C₃-C₆-alkynyl group, it being possible for this group in turn to carry one to five halogen atoms;


R⁸ can furthermore be a phenyl radical which can carry one to five halogen atoms and/or one to three of the following radicals: nitro, cyano, C₁-C₄-alkyl, C₁-C₄-haloalkyl, hydroxyl, C₁-C₄-alkoxy, mercapto, C₁-C₄-alkylthio, amino, C₁-C₄-alkylamino, C₁-C₄-dialkylamino;

a 5-membered heteroaromatic system which is linked via a nitrogen atom and contains one to three nitrogen atoms and which can carry one or two halogen atoms and/or one or two of the following radicals: C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, phenyl, C₁-C₄-haloalkoxy and/or C₁-C₄-alkylthio. Particular mention may be made of 1-pyrazolyl,

3-methyl-1-pyrazolyl, 4-methyl-1-pyrazolyl, 3,5-dimethyl-1-pyrazolyl, 3-phenyl-1-pyrazolyl, 4-phenyl-1-pyrazolyl, 4-chloro-1-pyrazolyl, 4-bromo-1-pyrazolyl, 1-imidazolyl, 1-benzimidazolyl, 1,2,4-triazol-1-yl, 3-methyl-1,2,4-triazol-1-yl, 5-methyl-1,2,4-triazol-1-yl,

1-benzotriazolyl, 3,4-dichloro-1-imidazolyl;

f) R^1 is furthermore a radical



10 where R^9 is:

C_1 - C_4 -alkyl, C_3 - C_6 -alkenyl, C_3 - C_6 -alkynyl, C_3 - C_8 -cycloalkyl as mentioned above in particular, it being possible for these radicals to carry a C_1 - C_4 -alkoxy, C_1 - C_4 -alkylthio and/or a phenyl radical as mentioned above;

15

phenyl, unsubstituted or substituted, in particular as mentioned above;


20 g) R^1 is a radical



where R^9 has the abovementioned meaning;

h) R^1 can furthermore be

30



35

where R^{10} and R^{11} can be identical or different and have the following meanings:

hydrogen, C_1 - C_7 -alkyl, C_3 - C_7 -cycloalkyl [sic], C_3 - C_7 -alkenyl, C_3 - C_7 -alkynyl, benzyl, phenyl, unsubstituted or substituted, as described above,

40

or R^{10} and R^{11} together form a C_4 - C_7 -alkylene chain which is closed to form a ring, is unsubstituted or substituted, e.g. substituted by C_1 - C_4 -alkyl, and may contain a heteroatom, e.g.

45

oxygen, sulfur or nitrogen, such as $-(CH_2)_4-$, $-(CH_2)_5-$,

$-(CH_2)_6-$, $-(CH_2)_7-$, $-(CH_2)_2-O-(CH_2)_2-$, $-CH_2-S-(CH_2)_2-$,
 $-CH_2-NH-(CH_2)_2-$, $-(CH_2)_2-N-(CH_2)_2-$.

With a view to the biological effect, preferred carboxylic acid
 5 derivatives of the general formula I, both as pure enantiomers
 and pure diastereomers or as mixtures thereof, are those where
 the substituents have the following meanings:

10 R^2 hydrogen, hydroxyl, $N(C_1-C_4-alkyl)_2$, the $C_1-C_4-alkyl$,
 $C_1-C_4-haloalkyl$, $C_1-C_4-alkoxy$, $C_1-C_4-haloalkoxy$,
 $C_1-C_4-alkylthio$ groups and halogen atoms specifically
 mentioned for R^1 , in particular chlorine, methyl, methoxy,
 ethoxy, difluoromethoxy and trifluoromethoxy, or CR^2 is
 15 linked to CR^{12} as indicated below to form a 5- or 6-membered
 ring;

X nitrogen or or CR^{12} , where

20 R^{12} is hydrogen or alkyl, or CR^{12} forms together with CR^2 or CR^3 a
 5- to 6-membered alkylene or alkenylene ring in which, in
 each case, a methylene group can be replaced by oxygen or
 sulfur, such as

25 $-CH_2-CH_2-O-$, $-CH=CH-O-$, $-CH_2-CH_2-CH_2-O-$, $-CH=CH-CH_2O-$, in
 particular hydrogen, $-CH_2-CH_2-O-$, $-CH(CH_3)-CH(CH_3)-O-$,
 $-C(CH_3)=C(CH_3)-O-$, $-CH=C(CH_3)-O-$ or $-C(CH_3)=C(CH_3)-S$;

Y nitrogen or methine;

30 R^3 hydrogen, hydroxyl, $N(C_1-C_4-alkyl)_2$, $C_1-C_4-alkyl$,
 $C_1-C_4-haloalkyl$, $C_1-C_4-alkoxy$, $C_1-C_4-haloalkoxy$,
 $C_1-C_4-alkylthio$ groups and halogen atoms, in particular
 chlorine, methyl, methoxy, ethoxy, difluoromethoxy and
 trifluoromethoxy, or $C-R^3$ is linked to $C-R^{12}$ as mentioned
 35 above to form a 5- or 6-membered ring;

40 R^4 and R^5 , which may be identical or different, phenyl or
 naphthyl, each of which may be substituted by one or more,
 e.g. one to three, of the following radicals: halogen, nitro,
 cyano, hydroxyl, mercapto, amino, $C_1-C_4-alkyl$,
 $C_1-C_4-haloalkyl$, $C_1-C_4-alkoxy$, $C_1-C_4-haloalkoxy$,
 $C_1-C_4-alkylthio$, $C_1-C_4-alkylamino$, di- $C_1-C_4-alkylamino$,
 $C_1-C_4-alkylcarbonyl$, $C_1-C_4-alkoxycarbonyl$;

45

phenyl or naphthyl, which may be connected together in ortho positions by a direct linkage, a methylene, ethylene or ethenylene group, an oxygen or sulfur atom or an SO₂, NH or N-alkyl group, or C₃-C₇ cycloalkyl;

5

R⁶ C₁-C₈-alkyl, C₃-C₆-alkenyl, C₃-C₆-alkynyl or C₃-C₈-cycloalkyl as mentioned above in particular, it being possible for each of these radicals to be substituted one or more times by: halogen, hydroxyl, mercapto, carboxyl, nitro, cyano,

10

C₁-C₄-alkoxy, C₃-C₆-alkenyloxy, C₃-C₆-alkynyloxy, C₁-C₄-alkylthio, C₁-C₄-haloalkoxy, C₁-C₄-alkylcarbonyl, hydroxycarbonyl, C₁-C₄-alkoxycarbonyl, C₁-C₄-alkylamino, di-C₁-C₄-alkylamino or unsubstituted or substituted phenyl or phenoxy, as mentioned above in particular;

15

phenyl or naphthyl, which can be substituted by one or more of the following radicals: halogen, nitro, cyano, hydroxyl, amino, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy,

20

C₁-C₄-haloalkoxy, phenoxy, C₁-C₄-alkylthio, C₁-C₄-alkylamino or C₁-C₄-dialkylamino;

a five- or six-membered heteroaromatic system which contains one to three nitrogen atoms and/or one sulfur or oxygen atom and which can carry one to four halogen atoms and/or one or two of the following radicals: C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-alkylthio, phenyl, phenoxy or phenylcarbonyl, it being possible for the phenyl radicals in turn to carry one to five halogen atoms and/or

25

30

one to three of the following radicals: C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy and/or C₁-C₄-alkylthio, as mentioned for R⁴ in particular;

Z sulfur or oxygen.

35

Particularly preferred compounds of the formula I, both as pure enantiomers and pure diastereomers or as mixture thereof, are those in which the substituents have the following meanings:

40 R² C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-alkylthio, trifluoromethyl or is linked to R¹² as mentioned below to form a 5- or 6-membered ring;

X nitrogen or CR¹², where

45

10

- R^{12} is hydrogen or alkyl, or CR^{12} forms together with CR^2 or CR^3 a 4- to 5-membered alkylene or alkenylene ring, e.g. $-CH_2-CH_2-CH_2-$ or $-CH=CH-CH_2-$, in which, in each case, one methylene group can be replaced by oxygen or sulfur, such as
- 5 $-CH_2-CH_2-O-$, $-CH=CH-O-$,
- $-CH_2-CH_2-CH_2-O-$, $-CH=CH-CH_2O-$, in particular hydrogen, $-CH_2-CH_2-O-$, $-CH(CH_3)-CH(CH_3)-O-$, $-C(CH_3)=C(CH_3)-O-$, $-CH=C(CH_3)-O-$ or $-C(CH_3)=C(CH_3)-S$;
- 10 Y nitrogen or methine;
- R^3 the C_1-C_4 -alkyl, C_1-C_4 -alkoxy, C_1-C_4 -alkylthio groups mentioned for R^1 or $C-R^3$ is linked to $C-R^{12}$ as mentioned above
- 15 to form a 5- or 6-membered ring;
- R^4 and R^5 phenyl (identical or different), which may be substituted by one or more, e.g. one to three, of the following radicals: halogen, nitro, hydroxyl, C_1-C_4 -alkyl,
- 20 C_1-C_4 -alkoxy, C_1-C_4 -alkylthio or
- R^4 and R^5 are phenyl groups which are connected together in ortho positions by a direct linkage, a methylene, ethylene or ethenylene group, an oxygen or sulfur atom or an SO_2 , NH or
- 25 N-alkyl group; or
- R^4 and R^5 are C_3-C_7 -cycloalkyl;
- R^6 C_1-C_8 -alkyl, C_3-C_6 -alkenyl or C_3-C_8 -cycloalkyl, it being
- 30 possible for each of these radicals to be substituted one or more times by: halogen, mercapto, caboxyl [sic], hydroxyl, nitro, cyano,
- C_1-C_4 -alkoxy, C_3-C_6 -alkenyloxy, C_1-C_4 -alkylthio;
- 35 phenyl or naphthyl, which can be substituted by one or more of the following radicals: halogen, nitro, cyano, hydroxyl, amino, C_1-C_4 -alkyl, C_1-C_4 -haloalkyl, C_1-C_4 -alkoxy, C_1-C_4 -haloalkoxy, phenoxy, C_1-C_4 -alkylthio, C_1-C_4 -akylamino
- 40 [sic] or C_1-C_4 -dialkylamino;
- a five- or six-membered heteroaromatic system which contains a nitrogen atom and/or a sulfur or oxygen atom and which can carry one to four halogen atoms and/or one or two of the
- 45 following radicals: C_1-C_4 -alkyl, C_1-C_4 -haloalkyl, C_1-C_4 -alkoxy, C_1-C_4 -alkylthio, phenyl, phenoxy or phenylcarbonyl, it being possible for the phenyl radicals in turn to carry one to five

11

halogen atoms and/or one to three of the following radicals:
C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy and/or
C₁-C₄-alkylthio;

5 Z sulfur or oxygen.

Examples of preferred compounds are listed in the following
table.

10

15

20

25


30

35

40

45

Table I [sic]



No.	R ¹	R ⁴ , R ⁵	R ⁶	R ³	R ²	X	Y	Z
I-1	OH	phenyl	methyl	OMe	OMe	N	N	O
I-2	OH	phenyl	methyl	OMe	Me	N	N	O
I-3	OH	phenyl	methyl	Me	OMe	N	N	O
I-4	OH	phenyl	methyl	ethyl	Me	N	N	O
I-5	OH	phenyl	methyl	Me	ethyl	N	N	O
I-6	OH	phenyl	methyl	Me	Me	N	N	O
I-7	OH	phenyl	methyl	OMe	SMe	N	N	O
I-8	OH	phenyl	methyl	OMe	CF ₃	N	N	O
I-9	OH	phenyl	methyl	OMe	OMe	N	N	S
I-10	OH	phenyl	methyl	OMe	Me	N	N	S
I-11	OH	phenyl	methyl	Me	OMe	N	N	S
I-12	OH	phenyl	methyl	ethyl	Me	N	N	S

No.	R ¹	R ⁴ , R ⁵	R ⁶	R ³	R ²	X	Y	Z
I-13	OH	phenyl	ethyl	OMe	OMe	N	N	O
I-14	OH	phenyl	ethyl	OMe	Me	N	N	O
I-15	OH	phenyl	ethyl	Me	OMe	N	N	O
I-16	OH	phenyl	ethyl	ethyl	Me	N	N	O
I-17	OH	phenyl	ethyl	Me	ethyl	N	N	O
I-18	OH	phenyl	HO-CH ₂ -CH ₂ -	OMe	OMe	N	N	O
I-19	OH	phenyl	HO-CH ₂ -CH ₂ -	OMe	Me	N	N	O
I-20	OH	phenyl	HO-CH ₂ -CH ₂ -	Me	OMe	N	N	O
I-21	OH	phenyl	HO-CH ₂ -CH ₂ -	ethyl	Me	N	N	O
I-22	OH	phenyl	HO-CH ₂ -CH ₂ -	Me	ethyl	N	N	O
I-23	OH	phenyl	HO-CH ₂ -CH ₂ -	Me	Me	N	N	O
I-24	OH	phenyl	HO-CH ₂ -CH ₂ -	OMe	SMe	N	N	O
I-25	OH	phenyl	HO ₂ C-(CH ₂) ₂ -	OMe	OMe	N	N	O
I-26	OH	phenyl	HO ₂ C-(CH ₂) ₂ -	OMe	Me	N	N	O
I-27	OH	phenyl	HO ₂ C-(CH ₂) ₂ -	Me	OMe	N	N	O
I-28	OH	phenyl	HO ₂ C-(CH ₂) ₂ -	ethyl	Me	N	N	O
I-29	OH	phenyl	HO ₂ C-(CH ₂) ₂ -	Me	ethyl	N	N	O
I-30	OH	phenyl	HO ₂ C-(CH ₂) ₂ -	Me	Me	N	N	O
I-31	OH	phenyl	HO ₂ C-(CH ₂) ₂ -	OMe	SMe	N	N	O
I-32	OH	phenyl	HO ₂ C-CH ₂ -	OMe	OMe	N	N	O
I-33	OH	phenyl	HO ₂ C-CH ₂ -	OMe	Me	N	N	O
I-34	OH	phenyl	HO ₂ C-CH ₂ -	Me	OMe	N	N	O
I-35	OH	phenyl	HO ₂ C-CH ₂ -	ethyl	Me	N	N	O
I-36	OH	phenyl	HO ₂ C-CH ₂ -	Me	ethyl	N	N	O
I-37	OH	phenyl	HO ₂ C-CH ₂ -	Me	Me	N	N	O

No.	R ¹	R ⁴ , R ⁵	R ⁶	R ³	R ²	X	Y	Z
I-38	OH	phenyl	HO ₂ C-CH ₂ -	OMe	SMe	N	N	O
I-39	OH	phenyl	HO-CH ₂ -(CH-OH)-CH ₂ -	OMe	OMe	N	N	O
I-40	OH	phenyl	HO-CH ₂ -(CH-OH)-CH ₂ -	OMe	Me	N	N	O
I-41	OH	phenyl	HO-CH ₂ -(CH-OH)-CH ₂ -	Me	OMe	N	N	O
I-42	OH	phenyl	HO-CH ₂ -(CH-OH)-CH ₂ -	ethyl	Me	N	N	O
I-43	OH	phenyl	HO-CH ₂ -(CH-OH)-CH ₂ -	Me	ethyl	N	N	O
I-44	OH	phenyl	HO-CH ₂ -(CH-OH)-CH ₂ -	Me	Me	N	N	O
I-45	OH	phenyl	HO-CH ₂ -(CH-OH)-CH ₂ -	OMe	SMe	N	N	O
I-46	OH	phenyl	HO-CH ₂ -(CH-OH)-CH ₂ -	OMe	CF ₃	N	N	O
I-47	OH	phenyl	HO-CH ₂ -(CH-OH)-CH ₂ -	OMe	OMe	N	N	S
I-48	OH	phenyl	HO-CH ₂ -(CH-OH)-CH ₂ -	OMe	Me	N	N	S
I-49	OH	phenyl	HO-CH ₂ -(CH-OH)-CH ₂ -	Me	OMe	N	N	S
I-50	OH	phenyl	HO-CH ₂ -(CH-OH)-CH ₂ -	ethyl	Me	N	N	S
I-51	OH	phenyl	HO-CH ₂ -CH ₂ -	OMe	OMe	N	N	S
I-52	OH	phenyl	HO-CH ₂ -CH ₂ -	OMe	Me	N	N	S
I-53	OH	phenyl	HO-CH ₂ -CH ₂ -	Me	OMe	N	N	S
I-54	OH	phenyl	HO-CH ₂ -CH ₂ -	ethyl	Me	N	N	S
I-55	OH	phenyl	HO-(CH ₂) ₃ -	OMe	OMe	N	N	O
I-56	OH	phenyl	HO-(CH ₂) ₃ -	OMe	Me	N	N	O
I-57	OH	phenyl	HO-(CH ₂) ₃ -	Me	OMe	N	N	O
I-58	OH	phenyl	HO-(CH ₂) ₃ -	ethyl	Me	N	N	O
I-59	OH	phenyl	HO-(CH ₂) ₃ -	Me	ethyl	N	N	O
I-60	OH	phenyl	HO-(CH ₂) ₃ -	Me	Me	N	N	O
I-61	OH	phenyl	HO-(CH ₂) ₃ -	OMe	SMe	N	N	O
I-62	OH	phenyl	HO-(CH ₂) ₃ -	OMe	CF ₃	N	N	O

No.	R ¹	R ⁴ , R ⁵	R ⁶	R ³	R ²	X	Y	Z
I-63	OH	phenyl	HO-(CH ₂) ₃ -	OMe	OMe	N	N	S
I-64	OH	phenyl	HO-(CH ₂) ₃ -	OMe	Me	N	N	S
I-65	OH	phenyl	HO-(CH ₂) ₃ -	Me	OMe	N	N	S
I-66	OH	phenyl	HO-(CH ₂) ₃ -	ethyl	Me	N	N	S
I-67	OH	phenyl	n-propyl	OMe	OMe	N	N	O
I-68	OH	phenyl	n-propyl	OMe	Me	N	N	O
I-69	OH	phenyl	n-propyl	Me	OMe	N	N	O
I-70	OH	phenyl	n-propyl	ethyl	Me	N	N	O
I-71	OH	phenyl	n-propyl	Me	ethyl	N	N	O
I-72	OH	phenyl	n-propyl	Me	Me	N	N	O
I-73	OH	phenyl	n-propyl	OMe	SMe	N	N	O
I-74	OH	phenyl	n-propyl	OMe	CF ₃	N	N	O
I-75	OH	phenyl	n-propyl	OMe	OMe	N	N	S
I-76	OH	phenyl	n-propyl	OMe	Me	N	N	S
I-77	OH	phenyl	n-propyl	Me	OMe	N	N	S
I-78	OH	phenyl	n-propyl	ethyl	Me	N	N	S
I-79	OH	phenyl	iso-propyl	OMe	OMe	N	N	O
I-80	OH	phenyl	iso-propyl	OMe	Me	N	N	O
I-81	OH	phenyl	iso-propyl	Me	OMe	N	N	O
I-82	OH	phenyl	iso-propyl	ethyl	Me	N	N	O
I-83	OH	phenyl	iso-propyl	Me	ethyl	N	N	O
I-84	OH	phenyl	iso-propyl	Me	Me	N	N	O
I-85	OH	phenyl	iso-propyl	OMe	SMe	N	N	O
I-86	OH	phenyl	iso-propyl	OMe	CF ₃	N	N	O
I-87	OH	phenyl	iso-propyl	OMe	OMe	N	N	S

No.	R ¹	R ⁴ , R ⁵	R ⁶	R ³	R ²	X	Y	Z
I-88	OH	phenyl	iso-propyl	OMe	Me	N	N	S
I-89	OH	phenyl	iso-propyl	Me	OMe	N	N	S
I-90	OH	phenyl	iso-propyl	ethyl	Me	N	N	S
I-91	OH	phenyl	methyl	CF ₃	tert-Butyl	N	N	O
I-92	OH	phenyl	methyl	CF ₃	OMe	N	N	O
I-93	OH	phenyl	methyl	SMe	OMe	N	N	O
I-94	OH	phenyl	methyl	CF ₃	Me	N	N	O
I-95	OH	phenyl	methyl	Me	CF ₃	N	N	O
I-96	OH	phenyl	benzyl	OMe	Me	N	N	O
I-97	OH	phenyl	benzyl	Me	OMe	N	N	O
I-98	OH	phenyl	benzyl	ethyl	Me	N	N	O
I-99	OH	phenyl	benzyl	Me	ethyl	N	N	O
I-100	OH	phenyl	benzyl	Me	Me	N	N	O
I-101	OH	phenyl	benzyl	OMe	OMe	N	N	O
I-102	OH	phenyl	phenyl	OMe	Me	N	N	O
I-103	OH	phenyl	phenyl	Me	OMe	N	N	O
I-104	OH	phenyl	phenyl	ethyl	Me	N	N	O
I-105	OH	phenyl	phenyl	Me	ethyl	N	N	O
I-106	OH	phenyl	phenyl	Me	Me	N	N	O
I-107	OH	phenyl	phenyl	OMe	OMe	N	N	O
I-108	OH	phenyl	4-OMe-benzyl	OMe	Me	N	N	O
I-109	OH	phenyl	4-OMe-benzyl	Me	OMe	N	N	O
I-110	OH	phenyl	4-OMe-benzyl	ethyl	Me	N	N	O
I-111	OH	phenyl	4-OMe-benzyl	Me	ethyl	N	N	O
I-112	OH	phenyl	4-OMe-benzyl	Me	Me	N	N	O

No.	R ¹	R ⁴ , R ⁵	R ⁶	R ³	R ²	X	Y	Z
I-113	OH	phenyl	4-OMe-benzyl	OMe	OMe	N	N	O
I-114	OH	phenyl	4-Cl-benzyl	OMe	Me	N	N	O
I-115	OH	phenyl	4-Cl-benzyl	Me	OMe	N	N	O
I-116	OH	phenyl	4-Cl-benzyl	ethyl	Me	N	N	O
I-117	OII	phenyl	4-Cl-benzyl	Me	ethyl	N	N	O
I-118	OH	phenyl	4-Cl-benzyl	Me	Me	N	N	O
I-119	OH	phenyl	4-Cl-benzyl	OMe	OMe	N	N	O
I-120	OII	cyclohexyl	methyl	OMe	OMe	N	N	O
I-121	OH	cyclohexyl	methyl	OMe	Me	N	N	O
I-122	OII	cyclohexyl	methyl	Me	OMe	N	N	O
I-123	OII	cyclohexyl	methyl	ethyl	Me	N	N	O
I-124	OH	cyclohexyl	methyl	Me	ethyl	N	N	O
I-125	OH	cyclohexyl	methyl	Me	CF ₃	N	N	O
I-126	OH	cyclohexyl	ethyl	OMe	Me	N	N	O
I-127	OH	cyclohexyl	ethyl	Me	OMe	N	N	O
I-128	OH	cyclohexyl	ethyl	ethyl	Me	N	N	O
I-129	OH	cyclohexyl	ethyl	OMe	ethyl	N	N	O
I-130	OH	cyclohexyl	HO-CH ₂ -CH ₂ -	OMe	Me	N	N	O
I-131	OH	cyclohexyl	HO-CH ₂ -CH ₂ -	Me	OMe	N	N	O
I-132	OH	cyclohexyl	HO-CH ₂ -CH ₂ -	ethyl	Me	N	N	O
I-133	OH	cyclohexyl	HO-CH ₂ -CH ₂ -	Me	ethyl	N	N	O
I-134	OH	cyclohexyl	HO-CH ₂ -CH ₂ -	Me	Me	N	N	O
I-135	OH	4-fluorophenyl	methyl	OMe	OMe	N	N	O
I-136	OH	4-fluorophenyl	methyl	OMe	Me	N	N	O
I-137	OH	4-fluorophenyl	methyl	Me	OMe	N	N	O

No.	R ¹	R ⁴ , R ⁵	R ⁶	R ³	R ²	X	Y	Z
I-138	OH	4-fluorophenyl	methyl	ethyl	Me	N	N	O
I-139	OH	4-fluorophenyl	methyl	Me	ethyl	N	N	O
I-140	OH	4-fluorophenyl	methyl	Me	CF ₃	N	N	O
I-141	OH	4-fluorophenyl	ethyl	OMe	Me	N	N	O
I-142	OH	4-fluorophenyl	ethyl	Me	OMe	N	N	O
I-143	OH	4-fluorophenyl	ethyl	Me	Et	N	N	O
I-144	OH	4-fluorophenyl	ethyl	OMe	ethyl	N	N	O
I-145	OH	4-fluorophenyl	HO-CH ₂ -CH ₂ -	OMe	Me	N	N	O
I-146	OH	4-fluorophenyl	HO-CH ₂ -CH ₂ -	Me	OMe	N	N	O
I-147	OH	4-fluorophenyl	HO-CH ₂ -CH ₂ -	ethyl	Me	N	N	O
I-148	OH	4-fluorophenyl	HO-CH ₂ -CH ₂ -	Me	ethyl	N	N	O
I-149	OH	4-fluorophenyl	HO-CH ₂ -CH ₂ -	Me	Me	N	N	O
I-150	OH	4-fluorophenyl	HO-CH ₂ -CH ₂ -	OMe	OMe	N	N	O
I-151	OH	4-chlorophenyl	methyl	OMe	Me	N	N	O
I-152	OH	4-chlorophenyl	methyl	Me	OMe	N	N	O
I-153	OH	4-chlorophenyl	methyl	ethyl	Me	N	N	O
I-154	OH	4-chlorophenyl	methyl	Me	ethyl	N	N	O
I-155	OH	4-chlorophenyl	methyl	Me	CF ₃	N	N	O
I-156	OH	4-chlorophenyl	ethyl	OMe	Me	N	N	O
I-157	OH	4-chlorophenyl	ethyl	Me	OMe	N	N	O
I-158	OH	4-chlorophenyl	ethyl	Me	Et	N	N	O
I-159	OH	4-chlorophenyl	ethyl	OMe	ethyl	N	N	O
I-160	OH	4-chlorophenyl	HO-CH ₂ -CH ₂ -	OMe	Me	N	N	O
I-161	OH	4-chlorophenyl	HO-CH ₂ -CH ₂ -	Me	OMe	N	N	O
I-162	OH	4-chlorophenyl	HO-CH ₂ -CH ₂ -	ethyl	Me	N	N	O

No.	R ¹	R ⁴ , R ⁵	R ⁶	R ³	R ²	X	Y	Z
I-163	OH	4-chlorophenyl	HO-CH ₂ -CH ₂ -	Me	ethyl	N	N	O
I-164	OH	4-chlorophenyl	HO-CH ₂ -CH ₂ -	Me	Me	N	N	O
I-165	OH	4-methylphenyl	methyl	Me	Me	N	N	O
I-166	OH	phenyl	methyl	CH ₃	CH=CH-CH=CH-C		N	O
I-167	OH	phenyl	methyl	H	CH=CH-CH=CH-C		N	O
I-168	OH	phenyl	methyl	Me	Me	CH	N	O
I-169	OH	phenyl	methyl	Me	ethyl	CH	N	O
I-170	OH	phenyl	methyl	ethyl	Me	CH	N	O
I-171	OH	phenyl	methyl	OMe	Me	CH	N	O
I-172	OH	phenyl	methyl	Me	CH ₂ -CH ₂ -CH ₂ -C		N	O
I-173	OH	phenyl	methyl	ethyl	CH ₂ -CH ₂ -CH ₂ -C		N	O
I-174	OH	phenyl	methyl	CF ₃	Me	CH	N	O
I-175	OH	phenyl	methyl	CF ₃	ethyl	CH	N	O
I-176	OH	phenyl	ethyl	CH ₃	CH=CH-CH=CH-C		N	O
I-177	OH	phenyl	ethyl	H	CH=CH-CH=CH-C		N	O
I-178	OH	phenyl	ethyl	Me	Me	CH	N	O
I-179	OH	phenyl	ethyl	Me	ethyl	CH	N	O
I-180	OH	phenyl	ethyl	ethyl	Me	CH	N	O
I-181	OH	phenyl	ethyl	OMe	Me	CH	N	O
I-182	OH	phenyl	ethyl	Me	CH ₂ -CH ₂ -CH ₂ -C		N	O
I-183	OH	phenyl	ethyl	ethyl	CH ₂ -CH ₂ -CH ₂ -C		N	O
I-184	OH	phenyl	ethyl	CF ₃	Me	CH	N	O
I-185	OH	phenyl	ethyl	CF ₃	ethyl	CH	N	O
I-186	OH	phenyl	n-propyl	CH ₃	CH=CH-CH=CH-C		N	O
I-187	OH	phenyl	n-propyl	H	CH=CH-CH=CH-C		N	O

No.	R ¹	R ⁴ , R ⁵	R ⁶	R ³	R ²	X	Y	Z
I-188	OH	phenyl	n-propyl	Me	Me	CH	N	O
I-189	OH	phenyl	n-propyl	Me	ethyl	CH	N	O
I-190	OH	phenyl	n-propyl	ethyl	Me	CH	N	O
I-191	OH	phenyl	n-propyl	OMe	Me	CH	N	O
I-192	OH	phenyl	n-propyl	Me	CH ₂ -CH ₂ -CH ₂ -C		N	O
I-193	OH	phenyl	n-propyl	ethyl	CH ₂ -CH ₂ -CH ₂ -C		N	O
I-194	OH	phenyl	n-propyl	CF ₃	Me	CH	N	O
I-195	OH	phenyl	n-propyl	CF ₃	ethyl	CH	N	O
I-196	OH	phenyl	HO-CH ₂ -CH ₂ -	CH ₃	CH=CH-CH=CH-C		N	O
I-197	OH	phenyl	HO-CH ₂ -CH ₂ -	H	CH=CH-CH=CH-C		N	O
I-198	OH	phenyl	HO-CH ₂ -CH ₂ -	Me	Me	CH	N	O
I-199	OH	phenyl	HO-CH ₂ -CH ₂ -	Me	ethyl	CH	N	O
I-200	OH	phenyl	HO-CH ₂ -CH ₂ -	ethyl	Me	CH	N	O
I-201	OH	phenyl	HO-CH ₂ -CH ₂ -	OMe	Me	CH	N	O
I-202	OH	phenyl	HO-CH ₂ -CH ₂ -	Me	CH ₂ -CH ₂ -CH ₂ -C		N	O
I-203	OH	phenyl	HO-CH ₂ -CH ₂ -	ethyl	CH ₂ -CH ₂ -CH ₂ -C		N	O
I-204	OH	phenyl	HO-CH ₂ -CH ₂ -	CF ₃	Me	CH	N	O
I-205	OH	phenyl	HO-CH ₂ -CH ₂ -	CF ₃	ethyl	CH	N	O
I-206	OH	phenyl	HOOC-CH ₂ -CH ₂ -	CH ₃	CH=CH-CH=CH-C		N	O
I-207	OH	phenyl	HOOC-CH ₂ -CH ₂ -	H	CH=CH-CH=CH-C		N	O
I-208	OH	phenyl	HOOC-CH ₂ -CH ₂ -	Me	Me	CH	N	O
I-209	OH	phenyl	HOOC-CH ₂ -CH ₂ -	Me	ethyl	CH	N	O
I-210	OH	phenyl	HOOC-CH ₂ -CH ₂ -	ethyl	Me	CH	N	O
I-211	OH	phenyl	HOOC-CH ₂ -CH ₂ -	OMe	Me	CH	N	O
I-212	OH	phenyl	HOOC-CH ₂ -CH ₂ -	Me	CH ₂ -CH ₂ -CH ₂ -C		N	O

No.	R ¹	R ⁴ , R ⁵	R ⁶	R ³	R ²	X	Y	Z
I-213	OH	phenyl	HOOC-CH ₂ -CH ₂ -	ethyl		CH ₂ -CH ₂ -CH ₂ -C	N	O
I-214	OH	phenyl	HOOC-CH ₂ -CH ₂ -	CF ₃	Me	CH	N	O
I-215	OH	phenyl	HOOC-CH ₂ -CH ₂ -	CF ₃	ethyl	CH	N	O
I-216	OH	phenyl	HO-CH ₂ -(CH-OH)-CH ₂ -	Me	Me	CH	N	O
I-217	OH	phenyl	HO-CH ₂ -(CH-OH)-CH ₂ -	Me	ethyl	CH	N	O
I-218	OH	phenyl	HO-CH ₂ -(CH-OH)-CH ₂ -	ethyl	Me	CH	N	O
I-219	OH	phenyl	HO-CH ₂ -(CH-OH)-CH ₂ -	OMe	Me	CH	N	O
I-220	OH	phenyl	iso-propyl	Me	Me	CH	N	O
I-221	OH	phenyl	iso-propyl	Me	ethyl	CH	N	O
I-222	OH	phenyl	iso-propyl	ethyl	Me	CH	N	O
I-223	OH	phenyl	iso-propyl	OMe	Me	CH	N	O
I-224	OH	phenyl	methyl	Me	Me	CH	N	S
I-225	OH	phenyl	methyl	Me	ethyl	CH	N	S
I-226	OH	phenyl	methyl	ethyl	Me	CH	N	S
I-227	OH	phenyl	methyl	OMe	Me	CH	N	S
I-228	OH	4-fluorophenyl	methyl	Me	Me	CH	N	O
I-229	OH	4-fluorophenyl	methyl	Me	ethyl	CH	N	O
I-230	OH	4-fluorophenyl	methyl	ethyl	Me	CH	N	O
I-231	OH	4-fluorophenyl	methyl	OMe	Me	CH	N	O
I-232	OH	4-fluorophenyl	methyl	CF ₃	Me	CH	N	O
I-233	OH	4-fluorophenyl	methyl	CF ₃	ethyl	CH	N	O
I-234	OH	4-fluorophenyl	HO-CH ₂ -CH ₂ -	Me	Me	CH	N	O
I-235	OH	4-fluorophenyl	HO-CH ₂ -CH ₂ -	Me	ethyl	CH	N	O
I-236	OH	4-fluorophenyl	HO-CH ₂ -CH ₂ -	ethyl	Me	CH	N	O
I-237	OH	4-fluorophenyl	HO-CH ₂ -CH ₂ -	OMe	Me	CH	N	O

No.	R ¹	R ⁴ , R ⁵	R ⁶	R ³	R ²	X	Y	Z
I-238	OH	4-chlorophenyl	methyl	Me	Me	CH	N	O
I-239	OH	4-chlorophenyl	methyl	Me	ethyl	CH	N	O
I-240	OH	4-chlorophenyl	methyl	ethyl	Me	CH	N	O
I-241	OH	4-chlorophenyl	methyl	OMe	Me	CH	N	O
I-242	OH	4-chlorophenyl	methyl	CF ₃	Me	CH	N	O
I-243	OH	4-chlorophenyl	methyl	CF ₃	ethyl	CH	N	O
I-244	OH	4-chlorophenyl	HO-CH ₂ -CH ₂ -	Me	Me	CH	N	O
I-245	OH	4-chlorophenyl	HO-CH ₂ -CH ₂ -	Me	ethyl	CH	N	O
I-246	OH	4-chlorophenyl	HO-CH ₂ -CH ₂ -	ethyl	Me	CH	N	O
I-247	OH	4-chlorophenyl	HO-CH ₂ -CH ₂ -	OMe	Me	CH	N	O
I-248	OH	phenyl	ethyl	Me	Me	N	N	O
I-249	OH	phenyl	ethyl	OMe	SMe	N	N	O
I-250	OH	phenyl	ethyl	OMe	CF ₃	N	N	O
I-251	OH	phenyl	methyl	OMe	OMe	N	CH	O
I-252	OH	phenyl	methyl	OMe	Me	N	CH	O
I-253	OH	phenyl	methyl	Me	Me	N	CH	O
I-254	OH	phenyl	methyl	ethyl	Me	N	CH	O
I-255	OH	phenyl	methyl	ethyl	ethyl	N	CH	O
I-256	OH	phenyl	methyl	CF ₃	Me	N	CH	O
I-257	OH	phenyl	methyl	CF ₃	ethyl	N	CH	O
I-258	OH	phenyl	methyl	OMe	CF ₃	N	CH	O
I-259	OH	phenyl	methyl	OMe	OMe	N	CH	S
I-260	OH	phenyl	methyl	OMe	Me	N	CH	S
I-261	OH	phenyl	methyl	Me	Me	N	CH	S
I-262	OH	phenyl	methyl	ethyl	Me	N	CH	S

No.	R ¹	R ⁴ , R ⁵	R ⁶	R ³	R ²	X	Y	Z
I-263	OH	phenyl	ethyl	OMe	OMe	N	CH	O
I-264	OH	phenyl	ethyl	OMe	Me	N	CH	O
I-265	OH	phenyl	ethyl	Me	Me	N	CH	O
I-266	OH	phenyl	ethyl	ethyl	Me	N	CH	O
I-267	OH	phenyl	HO-CH ₂ -CH ₂ -	OMe	OMe	N	CH	O
I-268	OH	phenyl	HO-CH ₂ -CH ₂ -	OMe	Me	N	CH	O
I-269	OH	phenyl	HO-CH ₂ -CH ₂ -	Me	Me	N	CH	O
I-270	OH	phenyl	HO-CH ₂ -CH ₂ -	ethyl	Me	N	CH	O
I-271	OH	phenyl	HO-CH ₂ -(CH-OH)-CH ₂ -	OMe	OMe	N	CH	O
I-272	OH	phenyl	HO-CH ₂ -(CH-OH)-CH ₂ -	OMe	Me	N	CH	O
I-273	OH	phenyl	HO-CH ₂ -(CH-OH)-CH ₂ -	Me	Me	N	CH	O
I-274	OH	phenyl	HO-CH ₂ -(CH-OH)-CH ₂ -	ethyl	Me	N	CH	O
I-275	OH	phenyl	HOOC-CH ₂ -CH ₂ -	OMe	OMe	N	CH	O
I-276	OH	phenyl	HOOC-CH ₂ -CH ₂ -	OMe	Me	N	CH	O
I-277	OH	phenyl	HOOC-CH ₂ -CH ₂ -	Me	Me	N	CH	O
I-278	OH	phenyl	HOOC-CH ₂ -CH ₂ -	ethyl	Me	N	CH	O
I-279	OH	phenyl	propyl	OMe	OMe	N	CH	O
I-280	OH	phenyl	propyl	OMe	Me	N	CH	O
I-281	OH	phenyl	propyl	Me	Me	N	CH	O
I-282	OH	phenyl	propyl	ethyl	Me	N	CH	O
I-283	OH	phenyl	4-OMe-benzyl	OMe	OMe	N	CH	O
I-284	OH	phenyl	4-OMe-benzyl	OMe	Me	N	CH	O
I-285	OH	phenyl	4-OMe-benzyl	Me	Me	N	CH	O
I-286	OH	phenyl	4-OMe-benzyl	ethyl	Me	N	CH	O
I-287	OH	4-fluorophenyl	methyl	OMe	OMe	N	CH	O

No.	R ¹	R ⁴ , R ⁵	R ⁶	R ³	R ²	X	Y	Z
I-288	OH	4-fluorophenyl	methyl	OMe	Me	N	CH	O
I-289	OH	4-fluorophenyl	methyl	Me	Me	N	CH	O
I-290	OH	4-fluorophenyl	methyl	ethyl	Me	N	CH	O
I-291	OH	4-fluorophenyl	ethyl	OMe	OMe	N	CH	O
I-292	OH	4-fluorophenyl	ethyl	OMe	Me	N	CH	O
I-293	OH	4-fluorophenyl	ethyl	Me	Me	N	CH	O
I-294	OH	4-fluorophenyl	ethyl	ethyl	Me	N	CH	O
I-295	OH	4-fluorophenyl	HO-CH ₂ -CH ₂ -	OMe	OMe	N	CH	O
I-296	OH	4-fluorophenyl	HO-CH ₂ -CH ₂ -	OMe	Me	N	CH	O
I-297	OH	4-fluorophenyl	HO-CH ₂ -CH ₂ -	Me	Me	N	CH	O
I-298	OH	4-fluorophenyl	HO-CH ₂ -CH ₂ -	ethyl	Me	N	CH	O
I-299	OH	4-chlorophenyl	methyl	OMe	OMe	N	CH	O
I-300	OH	4-chlorophenyl	methyl	OMe	Me	N	CH	O
I-301	OH	4-chlorophenyl	methyl	Me	Me	N	CH	O
I-302	OH	4-chlorophenyl	methyl	ethyl	Me	N	CH	O
I-303	OH	4-chlorophenyl	ethyl	OMe	OMe	N	CH	O
I-304	OH	4-chlorophenyl	ethyl	OMe	Me	N	CH	O
I-305	OH	4-chlorophenyl	ethyl	Me	Me	N	CH	O
I-306	OH	4-chlorophenyl	ethyl	ethyl	Me	N	CH	O
I-307	OH	4-chlorophenyl	HO-CH ₂ -CH ₂ -	OMe	OMe	N	CH	O
I-308	OH	4-chlorophenyl	HO-CH ₂ -CH ₂ -	OMe	Me	N	CH	O
I-309	OH	4-chlorophenyl	HO-CH ₂ -CH ₂ -	Me	Me	N	CH	O
I-310	OH	4-chlorophenyl	HO-CH ₂ -CH ₂ -	ethyl	Me	N	CH	O

The compounds of the present invention provide a novel therapeutic potential for the treatment of hypertension, pulmonary hypertension, myocardial infarct, angina pectoris, acute kidney failure, renal insufficiency, cerebral vasospasms, cerebral ischemia, subarachnoidal hemorrhages, migraine, asthma, atherosclerosis, endotoxic shock, endotoxin-induced organ failure, intravascular coagulation, restenosis after angioplasty, benign prostate hyperplasia, or hypertension or kidney failure caused by ischemia or intoxication, and of cancers, especially prostate cancer and skin cancer. The invention further relates to the combination of compounds of the formula I with inhibitors of the renin-angiotensin system (RAS). RAS inhibitors are disclosed in, for example, EP 634 175.

The combinations according to the invention are suitable for treating disorders for which compounds of the formula I also shows efficacy on their own, especially for treating hypertension and chronic heart failure.

The good effect of the compounds can be shown in the following tests:

Receptor binding studies

Cloned human ET_A receptor-expressing CHO cells and guinea pig cerebellar membranes with $> 60\%$ ET_B compared with ET_A receptors were used for binding studies.

Membrane preparation

The ET_A receptor-expressing CHO cells were grown in F_{12} medium containing 10% fetal calf serum, 1% glutamine, 100 E/ml penicillin and 0.2% streptomycin (Gibco BRL, Gaithersburg, MD, USA). After 48 h, the cells were washed with PBS and incubated with 0.05% trypsin-containing PBS for 5 min. Neutralization was then carried out with F_{12} medium and the cells were collected by centrifugation at 300 x g. To lyse the cells, the pellet was briefly washed with lysis buffer (5 mM tris-HCl, pH 7.4 with 10% glycerol) and then incubated at a concentration of 10^7 cells/ml of lysis buffer at 4°C for 30 min. The membranes were centrifuged at 20,000 x g for 10 min, and the pellet was stored in liquid nitrogen.

Guinea pig cerebella were homogenized in a Potter-Elvehjem homogenizer and were obtained by differential centrifugation at 1,000 x g for 10 min and repeated centrifugation of the supernatant at 20,000 x g for 10 min.

5

Binding assays

For the ET_A and ET_B receptor binding assay, the membranes were suspended in incubation buffer (50 mM tris-HCl, pH 7.4 with 5 mM
10 of $MnCl_2$, 40 μ g/ml bacitracin and 0.2% BSA) at a concentration of 50 μ g of protein per assay mixture and incubated with 25 pM [^{125}I [sic]]- ET_1 (ET_A receptor assay) or 25 pM [^{125}I [sic]]- RZ_3 (ET_B receptor assay) in the presence and absence of test substance. The nonspecific binding was determined with 10^{-7} M ET_1 . After 30
15 min, filtration was carried out through GF/B glass fiber filters (Whatman, England) in a Skatron cell collector (Skatron, Lier, Norway) to separate free and bound radioligands, and the filters were washed with ice-cold tris-HCl buffer, pH 7.4 with 0.2% BSA. The radioactivity collected on the filters was quantified using a
20 Packard 2200 CA liquid scintillation counter.

Functional in vitro assay system to look for endothelin receptor (subtype A) antagonists

25 This assay system is a functional, cell-based assay for endothelin receptors. When certain cells are stimulated with endothelin 1 (ET_1) they show an increase in the intracellular calcium concentration. This increase can be measured in intact cells loaded with calcium-sensitive dyes.

30

1-Fibroblasts which had been isolated from rats and in which an endogenous endothelin receptor of the A subtype had been detected were loaded with the fluorescent dye Fura 2-am as follows: after trypsinization, the cells were resuspended in buffer A (120 mM
35 NaCl, 5 mM KCl, 1.5 mM $MgCl_2$, 1 mM $CaCl_2$, 25 mM HEPES, 10 mM glucose, pH 7.4) to a density of 2×10^6 /ml and incubated with Fura 2-am (2 μ M), Pluronic [sic] F-127 (0.04%) and DMSO (0.2%) at 37°C in the dark for 30 min. The cells were then washed twice with buffer A and resuspended at 2×10^6 /ml.

40

The fluorescence signal from 2×10^5 cells per ml with Ex/Em 380/510 was recorded continuously at 30°C. The test substances and, after an incubation time of 3 min, ET_1 were [lacuna] to the cells, the maximum change in the fluorescence was determined. The
45 response of the cells to ET_1 without previous addition of a test substance was used as control and was set equal to 100%.

Testing of ET antagonists in vivo

Male SD rats weighing 250-300 g were anesthetized with amobarbital, artificially ventilated, vagotomized and pithed. The
5 carotid artery and jugular vein were cathetized [sic].

In control animals, intravenous administration of 1 µg/kg ET1 led to a distinct rise in blood pressure which persisted for a lengthy period.

10

The test animals received an i.v. injection of the test compounds (1 ml/kg) 5 min before administration of ET1. To determine the ET-antagonistic properties, the rise in blood pressure in the test animals was compared with that in the control animals.

15

Endothelin-1 induced "sudden death" in mice

The principle of the test is the inhibition of the sudden heart death caused in mice by endothelin, which is probably induced by
20 constriction of the coronary vessels, by pretreatment with endothelin receptor antagonists. Intravenous injection of 10 nmol/kg endothelin in a volume of 5 ml/kg of bodyweight results in death of the animals within a few minutes.

25 The lethal endothelin-1 dose is checked in each case on a small group of animals. If the test substance is administered intravenously, the endothelin-1 injection which was lethal in the reference group usually takes place 5 min thereafter. With other modes of administration, the times before administration are
30 extended, where appropriate up to several hours.

The survival rate is recorded, and effective doses which protect 50% of the animals (ED 50) from endothelin-induced heart death for 24 h or longer are determined.

35

Function test on vessels for endothelin receptor antagonists

Segments of rabbit aorta are, after an initial tension of 2 g and a relaxation time of 1 h in Krebs-Henseleit solution at 37°C and
40 pH 7.3-7.4, first induced to contract with K⁺. After washing out, an endothelin dose-effect plot up to the maximum is constructed.

Potential endothelin antagonists are administered to other preparations of the same vessel 15 min before starting the
45 endothelin dose-effect plot. The effects of the endothelin are calculated as % of the K⁺-induced contraction. Effective

endothelin antagonists result in a shift to the right in the endothelin dose-effect plot.

- The compounds according to the invention can be administered
- 5 orally or parenterally (subcutaneously, intravenously, intramuscularly, intraperitoneally [sic]) in a conventional way. Administration can also take place with vapors or sprays through the nasopharyngeal space.
- 10 The dosage depends on the age, condition and weight of the patient and on the mode of administration. The daily dose of active substance is, as a rule, about 0.5 to 50 mg/kg of bodyweight on oral administration and about 0.1-10 mg/kg of bodyweight on parenteral administration.
- 15 The novel compounds can be used in conventional solid or liquid pharmaceutical forms, e.g. as uncoated or (film-)coated tablets, capsules, powders, granules, suppositories, solutions, ointments, creams or sprays. These are produced in a conventional way. The
- 20 active substances can for this purpose be processed with conventional pharmaceutical aids such as tablet binders, bulking agents, preservatives, tablet disintegrants, flow regulators, plasticizers, wetting agents, dispersants, emulsifiers, solvents, release-slowing agents, antioxidants and/or propellant gases (cf.
- 25 H. Sucker et al.: Pharmazeutische Technologie, Thieme-Verlag, Stuttgart, 1991). The forms obtained in this way normally contain from 0.1 to 90% by weight of active substance.

Synthesis examples

30

Example 1

Methyl 2-hydroxy-3-methoxy-3,3-diphenylpropionate

- 5 g (19.6 mmol) of methyl 3,3-diphenyl-2,3-epoxypropionate were
- 35 dissolved in 50 ml of absolute methanol, and, at 0°C, 0.1 ml of boron trifluoride etherate was added. The mixture was stirred at 0°C for 2 h and at room temperature for a further 12 h. The solvent was removed by distillation, the residue was taken up in ethyl acetate, and the solution was washed with sodium
- 40 bicarbonate solution and water and dried over magnesium sulfate. After removal of the solvent by distillation, 5.5 g (88%) of a pale yellow oil remained.

Example 2

- 45 Methyl 2-(2,6-dimethoxy-4-pyrimidyloxy)-3-methoxy-3,3-diphenylpropionate

1.15 g (4 mmol) of methyl 2-hydroxy-3-methoxy-3,3-diphenylpropionate were dissolved in 10 ml of dimethylformamide, and 276 mg (2 mmol) of potassium carbonate were added. Then 524 mg (3 mmol) of 2,6-dimethoxy-4-chloropyrimidine were added, and the mixture was stirred at 100°C for 6 hours. It was then cautiously hydrolyzed with 10 ml of water, the pH was adjusted to 5 with citric acid and, after extraction with ethyl acetate, the organic phase was washed with water then dried over magnesium sulfate, and the solvent was removed by distillation. The oily residue (1.9 g) was chromatographed on silica gel, resulting in 617 mg of a slightly impure oil.

Example 3

2-(2,6-Dimethoxy-4-pyrimidinyl-oxy)-3-methoxy-3,3-diphenylpropionic acid

550 mg (1.3 mmol) of methyl 2-(2,6-dimethoxy-4-pyrimidinyl-oxy)-3-methoxy-3,3-diphenylpropionate were dissolved in 5 ml of dioxane, 2.6 ml of 1N KOH solution were added, and the mixture was stirred at 100°C for 3 h. The solution was diluted with 300 ml of water and extracted with ethyl acetate to remove unreacted ester. The aqueous phase was then adjusted to pH 1-2 with dilute hydrochloric acid and was extracted with ethyl acetate. After drying over magnesium sulfate and removal of the solvent by distillation, the residual foam (425 mg) was purified by MPLC, resulting in 205 mg (40%) of product as foam.

Example 4

Methyl 2-(4-methyl-2-quinolinyloxy)-3-methoxy-3,3-diphenylpropionate

5.7 g (20 mmol) of Methyl 2-hydroxy-3-methoxy-3,3-diphenylpropionate were dissolved in 90 ml of dimethylformamide, and 0.98 g (22 mmol) of sodium hydride (55% in oil) was added. After stirring for 15 minutes, 3.9 g (22 mmol) of 2-chloro-4-methylquinoline were added. The dark red solution was stirred at room temperature overnight, then cautiously hydrolyzed with 20 ml of water and subsequently extracted with ethyl acetate. The organic phase was washed with water, and dried over magnesium sulfate, and the solvent was removed by distillation. The residue (8.1 g) was then chromatographed on silica gel, resulting in 2.6 g of impure product as oil.

Example 5

2-(4-Methyl-2-quinolinyloxy)-3-methoxy-3,3-diphenylpropionic acid

30

1.8 g (4.2 mmol) of methyl 2-(4-methyl-2-quinolinyloxy)-3-methoxy-3,3-diphenylpropionate were dissolved in 25 ml of dioxane, 8.4 ml of 1N KOH solution were added, and the mixture was stirred at 60°C for 20 hours. The solution was diluted with 300 ml of water and extracted with ethyl acetate to remove unreacted ester and impurities. The aqueous phase was then adjusted to pH 1-2 with dilute hydrochloric acid and was extracted with ethyl acetate. Drying over magnesium sulfate and removal of the solvent by distillation resulted in 1.0 g of a foam. This was dissolved in a little ethyl acetate and left to stand at room temperature overnight, during which 265 mg of the product separated out as white crystals. Repetition of this procedure with the mother liquor afforded a further 166 mg of the product.

15 Total yield: 431 mg $\approx$ 25%
Melting point: 131-133°C
MS: M^+ = 413

Example 6

20 Benzyl 2-(2-tert-butyl-6-trifluoromethyl-4-pyrimidinyloxy)-3-methoxy-3,3-diphenylpropionate

2.0 g (5.5 mmol) of benzyl 2-hydroxy-3-methoxy-3,3-diphenylpropionate (see P44 36 851.8) were dissolved in 20 ml of dimethylformamide, and 0.76 g (5.5 mmol) of potassium carbonate was added. Then 1.32 g (5.5 mmol) of 2-tert-butyl-6-trifluoromethyl-4-chloropyrimidine were added, and the mixture was stirred at room temperature overnight and then at 100°C for 5 hours. To complete the reaction, a further 0.5 g of the pyrimidine and 0.5 g of NaH were added, and the mixture was stirred at 60°C for a further 6 hours. Finally, it was poured into ice-water and extracted with ethyl acetate, the organic phase was washed with saturated NaCl solution and dried over magnesium sulfate, and the solvent was removed by distillation. The oily residue (3.5 g) crystallized in n-heptane/ethyl acetate (5%). The precipitate was filtered off with suction and dried.

Yield: 1.6 g $\approx$ 52%
MS: M^+ = 564

40

Example 7

2-(2-tert-Butyl-6-trifluoromethyl-4-pyrimidinyloxy)-3-methoxy-3,3-diphenylpropionic acid

45 2.0 g (5.5 mmol) of benzyl 2-(2-tert-butyl-6-trifluoromethyl-4-pyrimidinyloxy)-3-methoxy-3,3-diphenylpropionate were dissolved in 250 ml of methanol, 120 mg of palladium on active

31

carbon (5%) were added, and the mixture was stirred under hydrogen at room temperature for 2 hours. After hydrogen uptake ceased, the catalyst was removed by filtration through celite, and the solvent was stripped off in a rotary evaporator.

5

Yield: 1.0 g $\cong$ 98%

$^1\text{H-NMR}$ (360 MHz, in CDCl_3 , data in ppm):

1.35 (s, 9H); 3.3 (s, 3H); 6.3 (s, 1H); 6.8 (s, 1H);

7.15 - 7.45 (m, 10H)

10

The compounds listed in Table 1 can be prepared in a similar way.

Example 8

15 Receptor binding data were measured for the compounds listed below using the binding assay described above.

The results are shown in Table 2.

20 Table 2

Receptor binding data (K_i values)

25	Compound	ET_A [μM]	ET_B [μM]
	I-1	0.038	8
	I-91	3.3	> 10
	I-166	0.5	4
30	I-167	3	> 7


35

40


45

We claim:

1. A carboxylic acid derivative of the formula I




where R is a tetrazole [sic], nitrile [sic], or a group



where R^1 has the following meanings:

- a) hydrogen;
- b) a succinylimidoxy [sic] group;
- c) a 5-membered heteroaromatic system linked via a nitrogen atom, such as pyrrolyl, pyrazolyl, imidazolyl and triazolyl, which may carry one or two halogen atoms or one or two C₁-C₄-alkyl or one or two C₁-C₄-alkoxy groups.
- d) R¹ is furthermore a group



where k can assume the values 0, 1 and 2, p can assume the values 1, 2, 3 and 4, and R^7 is

C₁-C₄-alkyl, C₃-C₇-cycloalkyl, C₃-C₆-alkenyl, C₃-C₆-alkynyl or unsubstituted or substituted phenyl which can be substituted by one or more, e.g. one to three, of the following radicals:

halogen, nitro, cyano, C₁-C₄-alkyl, C₁-C₄-haloalkyl, hydroxyl, C₁-C₄-alkoxy, C₁-C₄-alkylthio, mercapto, amino, C₁-C₄-alkylamino, C₁-C₄-dialkylamino;

- e) R^1 is furthermore a radical OR^8 where R^8 is:



2

hydrogen, the cation of an alkali metal such as lithium, sodium, potassium, or the cation of an alkaline earth metal such as calcium, magnesium and barium or a physiologically tolerated organic ammonium ion such as tertiary C₁-C₄-alkylammonium or the ammonium ion;

C₃-C₈-cycloalkyl such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and [sic] cycloheptyl, cyclooctyl,

C₁-C₈-alkyl, in particular C₁-C₄-alkyl such as methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, tert-butyl;

CH₂-phenyl, which may be substituted by one or more of the following radicals: halogen, nitro, cyano, C₁-C₄-alkyl, C₁-C₄-haloalkyl, hydroxyl, C₁-C₄-alkoxy, mercapto,

C₁-C₄-alkylthio, amino, C₁-C₄-alkylamino, C₁-C₄-dialkylamino,

a C₃-C₆-alkenyl or C₃-C₆-alkynyl group, it being possible for this group in turn to carry one to five halogen atoms;


R⁸ can furthermore be a phenyl radical which can carry one to five halogen atoms and/or one to three of the following radicals: nitro, cyano, C₁-C₄-alkyl, C₁-C₄-haloalkyl, hydroxyl, C₁-C₄-alkoxy, mercapto, C₁-C₄-alkylthio, amino, C₁-C₄-alkylamino, C₁-C₄-dialkylamino;

a 5-membered heteroaromatic system which is linked via a nitrogen atom and contains one to three nitrogen atoms and which can carry one or two halogen atoms and/or one or two of the following radicals: C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, phenyl, C₁-C₄-haloalkoxy and/or C₁-C₄-alkylthio. Particular mention may be made of 1-pyrazolyl, 3-methyl-1-pyrazolyl, 4-methyl-1-pyrazolyl, 3,5-dimethyl-1-pyrazolyl, 3-phenyl-1-pyrazolyl, 4-phenyl-1-pyrazolyl, 4-chloro-1-pyrazolyl, 4-bromo-1-pyrazolyl, 1-imidazolyl, 1-benzimidazolyl, 1,2,4-triazol-1-yl, 3-methyl-1,2,4-triazol-1-yl, 5-methyl-1,2,4-triazol-1-yl, 1-benzotriazolyl, 3,4-dichloro-1-imidazolyl;

f) R¹ is furthermore a radical



3



5

where R^9 is:

10 $\text{C}_1\text{-C}_4\text{-alkyl}$, $\text{C}_3\text{-C}_6\text{-alkenyl}$, $\text{C}_3\text{-C}_6\text{-alkynyl}$, $\text{C}_3\text{-C}_8\text{-cycloalkyl}$ as mentioned above in particular, it being possible for these radicals to carry a $\text{C}_1\text{-C}_4\text{-alkoxy}$, $\text{C}_1\text{-C}_4\text{-alkylthio}$ and/or a phenyl radical as mentioned above;

15 phenyl, unsubstituted or substituted, in particular as mentioned above.

g) R^1 is a radical



20

where R^9 has the abovementioned meaning.

25

h) R^1 can furthermore be



30

where R^{10} and R^{11} can be identical or different and have the following meanings:

35

hydrogen, $\text{C}_1\text{-C}_7\text{-alkyl}$, $\text{C}_3\text{-C}_7\text{-cycloalkyl}$ [sic], $\text{C}_3\text{-C}_7\text{-alkenyl}$, $\text{C}_3\text{-C}_7\text{-alkynyl}$, benzyl, phenyl, unsubstituted or substituted, as described above, or R^{10} and R^{11} together form a $\text{C}_4\text{-C}_7\text{-alkylene}$ chain which is closed to form a ring, is unsubstituted or substituted, e.g. substituted by $\text{C}_1\text{-C}_4\text{-alkyl}$, and may contain a heteroatom, e.g. oxygen, sulfur or nitrogen, such as $\text{---(CH}_2\text{)}_4\text{---}$, $\text{---(CH}_2\text{)}_5\text{---}$, $\text{---(CH}_2\text{)}_6\text{---}$, $\text{---(CH}_2\text{)}_7\text{---}$, $\text{---(CH}_2\text{)}_2\text{---O---(CH}_2\text{)}_2\text{---}$, $\text{---CH}_2\text{---S---(CH}_2\text{)}_2\text{---}$, $\text{---CH}_2\text{---NH---(CH}_2\text{)}_2\text{---}$, $\text{---(CH}_2\text{)}_2\text{---N---(CH}_2\text{)}_2\text{---}$;

40

45



* C L M S *

the other substituents have the following meanings:

- 5 R^2 hydrogen, hydroxyl, NH_2 , $NH(C_1-C_4-alkyl)$, $N(C_1-C_4-alkyl)_2$,
halogen, $C_1-C_4-alkyl$, $C_1-C_4-haloalkyl$, $C_1-C_4-alkoxy$,
 $C_1-C_4-haloalkoxy$ or $C_1-C_4-alkylthio$, or CR^2 is linked to
 CR^{12} as indicated below to form a 5- or 6-membered ring;
- 10 X nitrogen or CR^{12} where R^{12} is hydrogen or $C_{1-5-alkyl}$, or
 CR^{12} forms together with CR^2 or CR^3 a 5- or 6-membered
alkylene or alkenylene ring which can be substituted by
one or two $C_{1-4-alkyl}$ groups and in which in each case
one methylene group can be replaced by oxygen, sulfur,
-NH or -NC_{1-4-alkyl};
- 15 Y nitrogen or methine;
- 20 R^3 hydrogen, hydroxyl, NH_2 , $NH(C_1-C_4-alkyl)$, $N(C_1-C_4-alkyl)_2$,
halogen, $C_1-C_4-alkyl$, $C_1-C_4-haloalkyl$, $C_1-C_4-alkoxy$,
 $C_1-C_4-haloalkoxy$, -NH-O- $C_{1-4-alkyl}$, $C_1-C_4-alkylthio$; or CR^3
is linked to CR^{12} as indicated above to form a 5- or
6-membered ring;
- 25 R^4 and R^5 (which can be identical or different):
phenyl or naphthyl, each of which can be substituted by
one or more of the following radicals: halogen, nitro,
cyano, hydroxyl, $C_1-C_4-alkyl$, $C_1-C_4-haloalkyl$,
 $C_1-C_4-alkoxy$, $C_1-C_4-haloalkoxy$, phenoxy, $C_1-C_4-alkylthio$,
amino, $C_1-C_4-alkylamino$ or $C_1-C_4-dialkylamino$; or
30 phenyl or naphthyl, which are connected together in ortho
positions by a direct linkage, a methylene, ethylene or
ethenylene group, an oxygen or sulfur atom or an SO_2- ,
NH- or N-alkyl group, or $C_3-C_7-cycloalkyl$;
- 35 R^6 hydrogen, $C_1-C_8-alkyl$, $C_3-C_6-alkenyl$, $C_3-C_6-alkynyl$ or
 $C_3-C_8-cycloalkyl$, it being possible for each of these
radicals to be substituted one or more times by: halogen,
hydroxyl, mercapto, carboxyl, nitro, cyano, $C_1-C_4-alkoxy$,
40 $C_3-C_6-alkenyloxy$, $C_3-C_6-alkynyloxy$, $C_1-C_4-alkylthio$,
 $C_1-C_4-haloalkoxy$, $C_1-C_4-alkylcarbonyl$,
 $C_1-C_4-alkoxycarbonyl$,
45 $C_{3-8-alkylcarbonylalkyl}$, $C_1-C_4-alkylamino$,



5

di-C₁-C₄-alkylamino, phenyl or phenoxy or phenyl, substituted one or more times, e.g. once to three times, by halogen, nitro, cyano, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy or C₁-C₄-alkylthio;

5

phenyl or naphthyl, each of which can be substituted by one or more of the following radicals: halogen, nitro, cyano, hydroxyl, amino, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, phenoxy, C₁-C₄-alkylthio, C₁-C₄-alkylamino, C₁-C₄-dialkylamino, dioxomethylene [sic] or dioxoethylene [sic];

10

a five- or six-membered heteroaromatic system which contains one to three nitrogen atoms and/or one sulfur or oxygen atom and which can carry one to four halogen atoms and/or one or two of the following radicals: C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-alkylthio, phenyl, phenoxy or phenylcarbonyl, it being possible for the phenyl radicals in turn to carry one to five halogen atoms and/or one to three of the following radicals: C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy and/or C₁-C₄-alkylthio;

15

20

Z sulfur or oxygen.

25

2. A carboxylic acid derivative as claimed in claim 1, wherein R is COOH.

30

3. A carboxylic acid derivative as claimed in any of the preceding claims, wherein at least one of the radicals R⁴ and R⁵ is phenyl.

35

4. A carboxylic acid derivative as claimed in claim 3, wherein R⁴ and R⁵ are both phenyl.

5. A carboxylic acid derivative as claimed in claim 3, wherein R⁶ is C₁-C₈-alkyl, unsubstituted or substituted by OH or C₁-C₄-alkoxy, and Z is O.

40 6. A carboxylic acid derivative as claimed in any of the preceding claims, wherein X is CH.

7. A carboxylic acid derivative as claimed in any of the preceding claims, wherein at least one of the radicals X, Y is nitrogen (N).

45



8. A carboxylic acid derivative as claimed in any of the preceding claims, wherein at least one of the radicals R^2 , R^3 is C_1 - C_4 -alkyl.

5 9. The use of compounds as claimed in any of claims 1 to 8 for producing medicines for treating hypertension, pulmonary hypertension, acute and chronic kidney failure, chronic heart failure, cerebral ischemia, restenosis after angioplasty, prostate cancer.

10

10. The use of a combination of a compound as claimed in any of claims 1 to 8 with an inhibitor of the renin-angiotensin system (RAS).

15

20

25

30

35

40

45

