


Abstract

**2-UREIDO-THIAZOLE DERIVATIVES, PROCESS FOR THEIR PREPARATION,
AND THEIR USE AS ANTITUMOR AGENTS**

Compounds which are 2-ureido-1, 3-thiazole derivatives of formula (I) wherein R is a halogen atom, a nitro group, an optionally substituted amino group or it is a group, optionally further substituted, selected from: i) straight or branched C 1 -C 6 alkyl; ii) C 3 -C 6 cycloalkyl; iii) aryl or arylalkyl with from 1 to 6 carbon atoms within the straight or branched alkyl chain; R 1 is an optionally substituted group selected from: i) straight or branched C 1 -C 6 alkyl; ii) 3 to 6 membered carbocycle or 5 to 7 membered heterocycle ring; iii) aryl or arylcarbonyl; iv) arylalkyl with from 1 to 6 carbon atoms within the straight or branched alkyl chain; R 2 is hydrogen, a straight or branched C 1 -C 4 alkyl or C 2 -C 4 alkenyl or alkynyl group; or, taken together with the nitrogen atom to which they are bonded, R 1 and R 2 form a substituted or unsubstituted group selected from: i) an optionally benzocondensed or bridged 5 to 7 membered heterocycle; or ii) a 9 to 11 membered spiro-heterocyclic compound; or a pharmaceutically acceptable salt thereof; are useful for treating cell proliferative disorders associated with an altered cell dependent kinase activity.



(I)



162162



G00002