

Description

This invention concerns the field of antimicrobial agents, and in particular novel compounds which have microbicidal activity.

Slager, U.S.-A-3,923,870, discloses iodopropargyl oxycarbamates which are N-substituted with alkyl, aryl, or alkylaryl groups as having fungicidal activity.

DE-A-2515091 (1975), Ciba Geigy AG, discloses microbicidal anilides which are acylanilides which have an amino acid moiety and an iodopropargyl fragment as shown, but the point of attachment of the iodopropargyl fragment is not directly on the amino acid portion of the molecule. These compounds are N-iodopropargyloxy acid amides.


Kokai Tokkyo Koho JP 54/125614 (1979), Shigaken Pharmaceutical Co., Ltd. discloses antimicrobial iodopropargyloxy acid amide derivatives useful for bactericides and agricultural fungicides.

U.S.-A-4,535,088-A, assigned to Shionogi and Co., shows 2-(3-iodo)propynyl-amino-thiazole derivatives as useful as antimicrobials, i.e. antibacterial and antifungal agents.

Peptides: Synthesis, Structure and Function; Proc. Am. Pept. Symposium, 7th, (1981), pp 101-4 describes the synthesis of C-iodopropargyl glycine. There is no mention of antibacterial or antifungal activity.

Many of the antimicrobials of the prior art have toxicity and/or environmental problems. It is an object of the present invention to provide novel antimicrobial compounds which have improved toxicity profiles and are not harmful to the environment.

These objects, and others which will become apparent from the following disclosure, are achieved by the present invention which comprises in one aspect compounds of the formula



wherein R¹ is H, lower (C₁-C₄)alkyl, alkyl aryl, CH₂OR, CH₂SR, or CH(CH₃)OR; and

R, R², and R³ are independently H, (C₁-C₄)alkyl, (C₆-C₁₀)aryl, (C₆-C₁₄)arylalkyl, or (C₆-C₁₄)alkaryl.

The compounds of the invention have been discovered to be unexpectedly effective antimicrobials. The compounds are designed to biodegrade down to non-toxic amino acid components.

Some representative compounds include the following:

TABLE OF COMPOUNDS (R³ = H)

Compound Number	R ¹	R ²
1	H	H
2	H	CH ₃
3	H	C ₈ H ₁₇
4	H	CH ₂ -C(C-1
5	CH ₃	H
6	CH ₃	CH ₃
7	CH ₃	C ₈ H ₁₇
8	CH ₂ Ph	H
9	CH ₂ Ph	CH ₃
10	CH ₂ Ph	C ₈ H ₁₇

The invention also includes within its scope compositions comprising a compound as defined above, and an agronomically acceptable carrier, a cosmetic agent, a cutting oil, a soap or synthetic detergent, a stabilizer, or a film forming material, preferably wherein said compound is present in an amount of from

0.0001 to 50, more preferably from 0.001 to 10 % by weight. A further aspect of the invention comprises the use of a compound as defined above as a microbicide, and a still further aspect is a method of preventing or inhibiting the growth of bacteria, fungi or algae in a locus susceptible or subject thereto, comprising incorporating into or onto the locus, in an amount effective to adversely affect said growth, a compound or composition as defined above.

The compounds of this invention can be prepared by converting the appropriate D- or L- α -amino acid ester hydrochloride to the corresponding 2-isocyanato alkyl ester. The resultant isocyanate compound is reacted with iodopropargyl alcohol to give the N-iodopropargyloxycarbonyl amino acid ester. The free N-iodopropargyloxycarbonyl amino acids are prepared via alkaline hydrolysis of the corresponding ester. Other N-iodopropargyloxycarbonyl amino acid esters are prepared by esterification of the free N-iodopropargyloxycarbonyl amino acid with an appropriate alcohol or alkylhalide utilizing standard known methods.

Iodopropargyl alcohol can be prepared by the procedure described in Bulletin of the Chemical Society of Japan, Ando, T., Shioi, S., Nakagawa, M., (1972)45, 2611.

The 2-isocyanato alkyl esters are obtained from the corresponding amino acid ester hydrochloride salts by adapting the procedure for preparing 3-isocyanatopropanoyl chloride as described in Organic Synthesis: Collective Volume VI, (1988),715.

As stated above, compositions comprising a compound according to formula I and either an agronomically acceptable carrier, a cosmetic agent, a cutting oil, a soap or synthetic detergent, a stabilizer, a film forming material, or the like have a wide range of utility for protecting against or controlling microorganisms from a wide variety of classes including fungus, bacteria, algae, viruses and yeasts. The preferred utilities of the compositions are to protect wood, paint, adhesive, glue, paper, textile, leather, plastics, cardboard, lubricants, cosmetics, food, caulking, feed and industrial cooling water from microorganisms.

The following lists specific industries and applications of the compounds or compositions:

<u>Industry</u>	<u>Application</u>
Adhesives, sealants	adhesives caulks sealants
Agriculture/food chain	adjuvant preservation agricultural active ingredient agricultural chemical preservative agricultural formulations preservation animal feed preservation dairy chemicals fertilizer preservation food preservation food processing chemicals grain preservation post-harvest produce protection sugar processing tobacco
Construction products	asphalt / concrete cement modifiers construction products roof mastics synthetic stucco wall mastics joint cement
Cosmetics and toiletries	cosmetics raw materials for cosmetics, toiletries toiletries
Disinfectants, antiseptics	antiseptic disinfectant

5	Emulsions, dispersions	aqueous dispersions dispersed pigments latex photographic emulsions pigment slurries polymer latices
10	Formulated consumer & industrial products	air fresheners fabric softeners hand cleaners polishes, floor, furniture, shoe sponges & towelettes spray strach waxes
15		
20	Industrial processing, misc	dry cleaning fluids preservation electrodeposition paint, baths, rinses. electrodeposition pre-treatment, post rinses industrial fluids preservation pasteurization baths process aid preservation
25		
30	Industrial water treatment	air washers cooling towers cooling water water cooling
35		
40	Laundry	household laundry products laundered goods laundry wash water pre-washers sanitizers-laundry removers, spot & stain
45		
	Leather, leather products	leather and hide leather and hide products
50	Lubricants, hydraulic aids	automotive lubricants and fluids

55

5		conveyor lubricants greases hydraulic fluids hydraulic oils lubricants
10	Medical devices	diagnostic enzymes diagnostic kits medical devices
15	Metalworking & related app's	cutting fluids metal cleaning metalworking fluids
20	Odor control (active ingredient)	air conditioning animal bedding cat litter chemical toilet prep'ns deodorizers humidifiers industrial deodorants sanitary formulations toilet bowls
25		
30	Paints and coatings	coating emulsions paints
35		
40	Paper and wood pulp, their products	absorbant materials of paper and wood pulp packaging materials of paper and wood pulp paper paper products paper treatment soap wrap wood pulp wood pulp products
45		
50	Paper mill	paper mill slimicides pulp and paper slurries
55		

Petroleum refining, fuels

aviation fuels (jet fuel, aviation
gas)
burner, diesel and turbine fuel
oils
coal slurries
diesel fuel additives
diesel fuels
fuels
gasoline
heating oils
hydrocarbons
kerosene
liquefied petroleum gas
petrochemical feedstocks
petroleum products, storage,
transportation and production
recycled petroleum products
residual fuel oils
turbine oils

Pharmaceutical

topical antifungal and
antibacterial

Photographic chemicals and process

photographic processing - wash water,
rinses
photoplate processing chemicals
(developers, stabilizers etc)

Printing

fountain solutions (printing)
ink components (pigments, resins,
solvents, etc)
inks

Sanitizers (active)

sanitizers
sanitizers-dairy
sanitizers-dental
sanitizers-fermentation
sanitizers-food preparation
sanitizers-food processing
sanitizers-medical
sanitizers-rendering
sanitizers-veterinary

Soaps, detergents, cleaners

5

10

15

cleaners
detergents, hand automatic laundry,
other
household cleaners
industrial cleaners
liquid soaps, hand, dish,
laundry
oil and grease remover
powdered soaps
raw materials for cleaning
products
soaps
surfactants

Textiles, textile products

25

30

35

40

45

50

55

bonded fabrics
burlap
canvas
canvas goods
carpet backing
carpets
clothing
coated fabrics
curtains
draperies
engineering textiles
fibers
geotextiles
goods made of textiles
knitted fabrics
nets
nonwoven fabrics
rope
rugs
textile accessories
textile products
textiles
upholstery
woven fabrics
yarn

Textile processing

dye fixatives
dyes
fiber lubricants
hand modifiers
sizes
textile processing fluids

Therapeutic (active or preservative)

animal health/veterinary
aquaculture
dental
human health
pharmaceutical /therapeutic

Water purification

charcoal beds
deionization resins
filters
membranes
reverse osmosis membranes
ultrafilters
water purification
water purification pipes, tubing

Wood applications

lazes (wood stains)
wood
wood products

Miscellaneous

alcohols
bedding incorporating water or
gels
ceramic
contact lens cases-leaching
electronic circuitry
electronics chemicals
enzymes-food production
enzymes-industrial
gel cushions
laboratory reagents
marine antifoulants
mildewcides
mining applications
natural rubber latex

oil field applications
 pipes
 plastics
 polymer systems
 polymers and resins (synthetic and
 natural)
 reagent preservation
 rubber
 rubber products
 skin remover
 solid protective/decorative
 films
 swimming pools
 waste treatment
 water beds

The amounts of the compound to be used depend on the application. The useful amounts for a particular application are similar to amounts used for other microbiocide compounds.


The compound can be used in combination with other microbiocides. The term "microbiocide" is considered equivalent to "antimicrobial" as used herein.

Suitable methods of application of compounds of formula I to control fungi, bacteria, algae, viruses, yeasts, and the like are in amounts and with carriers, etc., as well known in the art.

The following examples are presented to illustrate a few embodiments of the invention, but are not to be considered as limiting.

Example 1

Preparation of N-iodopropargyloxycarbonyl glycine methyl ester




Trichloromethylchloroformate (18.00 ml., 149.2 mmole) was added to a slurry of glycine methyl ester hydrochloride (12.56 g., 100.0 mmole) in anhydrous 1,4-dioxane (200 ml) and refluxed for 5 hours and then stirred at room temperature for 16 hours. The excess trichloromethylchloroformate and 1,4-dioxane were removed from the reaction mixture by distillation under slightly reduced pressure. The resultant residue was distilled to afford the desired intermediate 2-isocyanatoacetic acid methyl ester (10.02 g., 87.1% yield) as a clear colorless liquid. B.P. 67-68°C (15-20 mm Hg.); ¹H-NMR (200 MHz, CDCl₃) δ_{ppm} = 3.84, s, 3H (-OCH₃), 3.97, s, 2H (-N-CH₂-C).

The 2-isocyanatoacetic acid methyl ester (10.02 g. 87.06 mmole) was dissolved in anhydrous toluene (100 ml) and treated with an anhydrous toluene (50 ml) solution of iodopropargyl alcohol (15.84 g., 87.05 mmole), followed by 6 drops of di-n-butyl tin dilaurate. The reaction mixture was heated at 85°C for 17.5 hours. After cooling to room temperature, the heterogeneous reaction mixture was diluted with ethyl acetate. The solution was washed with 3 X 60 ml. of water, 3 X 60 ml. of 10% aqueous potassium hydrogen sulfate, 3 X 60 ml. of saturated aqueous sodium bicarbonate, 60 ml. of saturated aqueous sodium chloride, dried over magnesium sulfate, filtered and concentrated under reduced pressure to afford 25.50 g. (94.7% crude yield) as a yellow crystalline solid. Recrystallization from methylene chloride/hexane gave 12.97 g. of N-iodopropargyloxycarbonyl glycine methyl ester (Compound 2) as a white crystalline solid (m.p. 93.5-

95.0 °C). ¹H-NMR (200 MHz, CDCl₃) δ_{ppm} = 3.78, s, 3H (-OCH₃), 4.01, d, 2H (-CH₂-NH-), 4.86, s, 2H (-CH₂-), 5.33, broad s, 1H, (NH). The mother liquor of compound 2 was concentrated under reduced pressure and the residue chromatographed on 200 g. of silica gel and eluted with a 6:1 mixture of 25% methylene chloride in hexane and ethyl acetate to afford 8.43 g. of N-iodopropargyloxycarbonyl glycine methyl ester (compound 2) as a white crystalline solid. This second crop of material had a m.p. 88-92 °C. The combined total yield for this reaction was 82.8%.

Example 2


10 Preparation of N-iodopropargyloxycarbonyl Glycine



N-iodopropargyloxycarbonyl glycine methyl ester (Compound 2, 8.03 g., 27.03 mmole) was dissolved in 100 ml. of tetrahydrofuran, cooled to 0 °C with an ice bath and treated during 15 minutes with 30.5 ml. of 1M aqueous lithium hydroxide. The reaction mixture was stirred at 0 °C for 3/4 hr. or until the starting material was no longer present. The reaction progress was monitored by thin layer chromatography (silica gel, methylene chloride/acetic acid 90:10). After the reaction was complete, the aqueous solution was acidified (pH=2) with 10% aqueous potassium hydrogen sulfate (100 ml) and 3N aqueous hydrogen chloride (10 ml.). This mixture was extracted several times with diethyl ether. The ether layers were combined and washed with water, dried over magnesium sulfate, filtered and concentrated under reduced pressure to afford N-iodopropargyloxycarbonyl glycine (Compound 1) as a pale amber oil, which crystallized upon standing (6.83 g., 86.7% yield after drying in vacuo over P₂O₅). m.p. 108-110 °C. ¹H-NMR (200 MHz., DMSO-d₆) δ_{ppm} = 3.67., d, 2H (-NH-CH₂-), 4.76, s, 2H (C-CH₂-O), 7.66, t, 1H (NH).

Example 3


Preparation of N-iodopropargyloxycarbonyl Glycine N-Octyl Ester



N-iodopropargyloxycarbonyl glycine (compound 1, 6.50 g., 22.97 mmole) was dissolved in anhydrous acetonitrile (100 ml.), treated with 1,8-diazabicyclo[5.4.0]undec-7-ene (3.5 ml., 23.40 mmole) and stirred at room temperature for 5 minutes. n-Octylbromide (4.00 ml., 23.16 mmole) was added and the reaction mixture was stirred at room temperature for 17 hours, then heated at reflux temperature for 6 hours. The reaction mixture was concentrated under reduced pressure and the resultant residue was dissolved in ethyl acetate. The ethyl acetate solution was washed with water, 5% aqueous citric acid, saturated aqueous sodium bicarbonate, saturated aqueous sodium chloride, dried over magnesium chloride, filtered and concentrated under reduced pressure. The resultant residue was purified utilizing column chromatography (silica gel, 200 g.) and eluted with ethyl acetate/hexane (1:10) to afford N-iodopropargyloxycarbonyl glycine n-octyl ester (Compound 3) as a clear colorless liquid which crystallized on standing. m.p. 43-46 °C. Elemental Analysis: C₁₄H₂₂NO₄I: Calc'd. C: 42.55, H: 5.61, N: 3.54. Found: C: 43.73, H: 5.62, N: 3.54. (Ethyl acetate was present in 0.5 mole%). ¹H-NMR (200 MHz, CDCl₃) δ_{ppm} = 0.89, m, 3H (-CH₃), 1.30, m, 10H (-CH₂)₅-CH₃), 1.64, m, 2H (-CH₂-), 4.0, d, J=3.4 Hz., 2H (-N-CH₂-), 4.1-4.2, m, 2H (-CH₂-), 4.86, s, 2H (C-CH₂-O), 5.32, m, 1H (-NH-). IR (neat) 3460, 2920, 2850, 2200, 1740, 1560, 1480, 1400, 1360, 1200, 1060, 980 cm⁻¹.

Example 4


Preparation of N-Iodopropargyloxycarbonyl Glycine Iodopropargyl Ester



N-Iodopropargyloxycarbonyl glycine (compound 1, 4.96 g., 17.52 mmole) was dissolved in a mixture of methylene chloride and 1,4-dioxane (450 ml., 20 ml. respectively) and cooled to 0°C with an ice bath. Iodopropargyl alcohol (3.51 g., 19.29 mmole) was added, followed by 4-dimethylaminopyridine (0.22 g., 1.80 mmole) and N-(dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride (4.17 g., 21.75 mmole). The reaction mixture was stirred at 4°C for 17 hours. Upon warming to room temperature, the reaction mixture was diluted with methylene chloride, washed with water, 10% aqueous potassium hydrogen sulfate, saturated aqueous sodium bicarbonate, saturated aqueous sodium chloride, dried over magnesium sulfate, filtered and concentrated under reduced pressure to afford N-Iodopropargyloxycarbonyl glycine iodopropargyl ester (compound 4) as a white crystalline solid (8.26 g.). m.p. 86-88°C; ¹H-NMR (200 MHz, CDCl₃) δ_{ppm} = 4.05, d, J = 5.6 Hz., 2H (NH-CH₂-), 4.89, d, J = 10.2 Hz., 4H (C-CH₂-O), 5.32, m, 1H (-NH-).

Example 5

Preparation of N-Iodopropargyloxycarbonyl Alanine Methyl Ester




L-Alanine methyl ester hydrochloride (6.00 g., 42.99 mmole) was slurried in anhydrous 1,4-dioxane (125 ml.), treated with trichloromethylchloroformate (7.8 ml., 64.66 mmole) and heated at 55°C for 20.5 hours. The excess trichloromethylchloroformate and 1,4-dioxane were removed from the reaction mixture by distillation under slightly reduced pressure. The resultant residue was subjected to distillation to afford the desired intermediate 2-isocyanatopropionic acid methyl ester (4.64 g., 83.6% yield) as a clear colorless liquid. B.P. 58°C (15-20 mm Hg.). ¹H-NMR (200 MHz, CDCl₃) δ_{ppm} = 1.54, d, J = 8 Hz., 3H (-CH₃), 3.87, s, 3H (-OCH₃), 4.13, q, J = 8 Hz., 1H (-CH-CH₃), IR (neat) 3000, 2960, 2270, 1745, 1455, 1445, 1385, 1305, 1230, 1170 cm⁻¹.

An anhydrous toluene solution (30 ml.) of iodopropargyl alcohol (6.60 g., 36.27 mmole) was added to an anhydrous solution (40 ml.) of 2-isocyanatopropionic acid methyl ester (4.64 g., 35.94 mmole) and the reaction mixture was stirred at room temperature for 2.5 hours, then at 70°C for 17 hours. When cool, the reaction mixture was diluted with toluene and washed with water, 10% aqueous potassium hydrogen sulfate, water, saturated aqueous sodium chloride, dried over magnesium sulfate, filtered and concentrated under reduced pressure to form a yellow crystalline solid (9.55 g., 85.4% crude yield). The crude material was recrystallized from ethyl acetate/hexane to afford N-Iodopropargyloxycarbonyl alanine methyl ester (compound 6, 5.85g.) as a white crystalline solid. m.p. 100-102°C. ¹H-NMR (200 MHz, CDCl₃) δ_{ppm} = 1.43, d, J = 8 Hz., 3H (-CH₃), 3.77, s, 3H (-OCH₃), 4.38, m, 1H (-CH-CH₃), 4.85, d, J = 2.7 Hz., 2H (-C-CH₂-O), 5.39, m, 1H (-NH). IR (CHCl₃) 3450, 2970, 2210, 1735, 1515, 1469, 1350, 1080 cm⁻¹.

Example 6


Preparation of N-iodopropargyloxycarbonyl Alanine



N-iodopropargyloxycarbonyl alanine methyl ester (compound 6, 4.00g., 12.86 mmole) was dissolved in a tetrahydrofuran/water mixture (3:1, 100 ml.) and cooled to 0°C with an ice bath. Lithium hydroxide (0.35 g., 14.61 mmole) was added and the reaction mixture was stirred for 2 hours. The reaction mixture was filtered, concentrated under reduced pressure, redissolved in water and acidified with 10% aqueous potassium hydrogen sulfate. The aqueous mixture was extracted with methylene chloride several times. The combined methylene chloride layers were washed with water, dried over magnesium sulfate, filtered and concentrated under reduced pressure to afford N-iodopropargyloxycarbonyl alanine (compound 5, 2.72 g., 71.2% yield) as a white crystalline solid. m.p. 95-99°C; ¹H-NMR (200 MHz, DMSO-d₆) δ_{ppm} = 1.28, d, J = 8 Hz., 3H (-CH-CH₃), 4.00, m, 1H (-CH-CH₃), 4.75, s, 2H (-CH₂-O), 7.7, d, J = 8 Hz., 1H (-NH-), 12.57, s, 1H (-COOH). IR (KBr) 3500-2000, 2200, 1670-1715, 1530, 1450, 1335, 1250, 1075, 1050 cm⁻¹.

Example 7


Preparation of N-iodopropargyloxycarbonyl Alanine n-Octyl Ester



N-iodopropargyloxycarbonyl alanine (compound 5, 0.95 g., 3.20 mmole) was dissolved in anhydrous N,N-dimethylformamide (15 ml.) and treated with the following: n-octanol (0.80 ml., 3.81 mmole), N-(dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride (0.68 g., 4.07 mmole) and 4-dimethylaminopyridine (0.041 g., 0.336 mmole). The reaction mixture was stirred at room temperature for 10 minutes, then held at 4°C for 17 hours. The reaction mixture was diluted with ethyl acetate and washed with water, 10% aqueous potassium hydrogen sulfate, water, saturated aqueous sodium carbonate, water, saturated aqueous sodium chloride, dried over magnesium sulfate, filtered and concentrated under reduced pressure to afford an oil. The oil was chromatographed on silica gel (50 g.) and eluted with hexanes / ethyl acetate (9:1) to afford N-iodopropargyloxycarbonyl alanine n-octyl ester (compound 7, 0.97 g., 74% yield) as a clear colorless oil. ¹H-NMR (200 MHz, CDCl₃) δ_{ppm} = 0.93, m, 3H (-CH₂-CH₃), 1.34, m, 10H (-CH₂)₅-CH₃), 1.47, d, J = 8 Hz., 3H (-CH-CH₃), 1.7, m, 2H (-CH₂-), 4.18, t, J = 8 Hz., 2H (-O-CH₂-CH₂-), 4.4, m, 1H (-CH-CH₃), 4.87, s, 2H (-C-CH₂-O), 5.42, m, 1H (-NH-). IR (CHCl₃) 3440, 2940, 2870, 2210, 1745, 1505, 1455, 1340, 1070 cm⁻¹. IR (neat) 3350, 2930, 2860, 2200, 1740, 1515, 1445, 1070 cm⁻¹.

Example 8

Preparation of N-Iodopropargyloxycarbonyl Phenylalanine Methyl Ester




L-Phenylalanine methyl ester hydrochloride (6.00 g., 27.82 mmole) was slurried in anhydrous 1,4-dioxane (125 ml.), treated with trichloromethylchloroformate (5.05 ml., 41.86 mmole) and heated at 55 °C for 17.5 hours. The excess trichloromethylchloroformate and 1,4-dioxane were removed from the reaction mixture by distillation under slightly reduced pressure. The resultant residue was subjected to high vacuum distillation to afford the desired intermediate 2-isocyanato-3-phenylpropionic acid methyl ester (4.78 g., 84% yield) B.P. 83-85 °C (100 μ m Hg.) as a clear colorless liquid. ¹H-NMR (200 MHz, CDCl₃) δ_{ppm} = 3.12, dd, J = 8.5 Hz., 2H (-CH₂-Ph), 3.83, s, 3H (-COOCH₃), 4.28, m, 1H (-CH-CH₂-), 7.15-7.4, m, 5H (Ar-H).

An anhydrous toluene solution (50 ml.) of iodopropargyl alcohol (4.66 g., 25.61 mmole) was added to an anhydrous toluene solution (50 ml.) of 2-isocyanato-3-phenylpropionic acid methyl ester (4.78 g., 23.29 mmole) containing 5 drops of di-n-butyl tin dilaurate and the reaction mixture was stirred at room temperature for 17 hours. When cool, the reaction mixture was diluted with toluene and washed with water, 10% aqueous potassium hydrogen sulfate, water, saturated aqueous sodium chloride, dried over magnesium sulfate, filtered and concentrated under reduced pressure to form N-iodopropargyloxycarbonyl phenylalanine methyl ester (compound 9) as a white crystalline solid (8.87 g., 98.3% yield). m.p. 100-103 °C; ¹H-NMR (200 MHz, CDCl₃) δ_{ppm} = 3.13, m, 2H (-CH₂-Ph), 3.75, s, 3H (COOCH₃), 4.65, m, 1H (-CH-), 4.83, s, 2H (C-CH₂-O), 5.25, m, 1H (-NH-), 7.05-7.4, m, 5H (Ar-H).

Example 9


Preparation of N-Iodopropargyloxycarbonyl Phenylalanine



N-Iodopropargyloxycarbonyl phenylalanine methyl ester (compound 9, 7.05g., 18.21 mmole) was dissolved in a tetrahydrofuran/water mixture (3:1, 100 ml.) and cooled to 0 °C with an ice bath. Lithium hydroxide (0.483 g., 20.17 mmole) was added and the reaction mixture was stirred for 1.5 hours. An additional amount of lithium hydroxide (45 mg.) was added to the reaction mixture and stirring was continued for 0.5 hr. The reaction mixture was concentrated under reduced pressure to remove the tetrahydrofuran, redissolved in water and acidified with 10% aqueous potassium hydrogen sulfate. The resultant white precipitate was isolated by filtration. The solid was dissolved in diethyl ether and washed with water, saturated aqueous sodium chloride, dried over magnesium sulfate, filtered and concentrated under reduced pressure to afford N-iodopropargyloxycarbonyl phenylalanine (compound 8, 5.97 g., 87.8% yield) as a white crystalline solid. ¹H-NMR (200 MHz., DMSO-d₆) δ_{ppm} = 2.95, dd, J = 4.16 Hz., 2H (-CH₂-Ph), 4.15, m, 1H (-CH-CH₂-), 4.7, s, 2H (-C-CH₂-O), 7.3, m, 5H (Ar-H), 7.75, d, J = 8 Hz., 1H (-NH-), 12.8, m, 1H (COOH). IR (CHCl₃) 3600-2400, 3420, 2200, 1730, 1500, 1065 cm⁻¹.

Example 10

Preparation of N-Iodopropargyloxycarbonyl Phenylalanine n-Octyl Ester



N-iodopropargyloxycarbonyl phenylalanine (compound 8, 2.00 g., 5.36 mmole) was dissolved in anhydrous 1,4-dioxane (20 ml.) and treated with the following: n-octanol (0.95 ml., 6.03 mmole), N-(dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride (1.08 g., 6.47 mmole) and 4-dimethylaminopyridine (0.073 g., 0.598 mmole). N,N-Dimethylformamide (5.00 ml.) was added to the heterogeneous reaction mixture to effect a homogeneous solution. The reaction mixture was stirred at room temperature for 10 minutes, then held at 4°C for 17 hours. The reaction mixture was concentrated under reduced pressure to remove the 1,4-dioxane, diluted with methylene chloride and washed with water, 10% aqueous potassium hydrogen sulfate, water, saturated aqueous sodium carbonate, water, saturated aqueous sodium chloride, dried over magnesium sulfate, filtered and concentrated under reduced pressure to afford a viscous pale yellow oil which solidified on standing. The oil was chromatographed on silica gel (70 g.) and eluted with hexanes/ethyl acetate (9:1) to afford N-iodopropargyloxycarbonyl phenylalanine n-octyl ester (compound 10, 1.98 g., 76.2% yield) as a white crystalline solid. ¹H-NMR (200 MHz., CDCl₃) δ_{ppm} = 0.9, m, 3H (-CH₃), 1.3, m, 10H (-CH₂)₅-CH₃), 1.58, m, 2H (-CH₂-), 3.13, d, J = 5 Hz., 2H (-CH₂-Ph), 4.1, t, J = 5 Hz., 2H (-O-CH₂-CH₂-), 4.63, m, 1H (-CH-CH₂-), 4.72, s, 2H (C-CH₂-O), 5.27, d, J = 8 Hz., 1H (-NH-), 7.1-7.35, m, 5H (Ar-H). IR (CHCl₃) 3440, 2940, 2870, 1730, 1505, 1355, 1120 cm⁻¹.

Example 11 - Biological Activity

A. Biocidal Activity:

Biocidal evaluations (bactericidal, algicidal, and fungicidal) were carried out.

A minimum inhibitory concentration (MIC) value is obtained using a broth, two-fold serial dilution test performed as follows: A stock solution or dispersion of the test compound, typically at a concentration of 1%, is made in a 5:3:2 solvent solution of acetone, methanol, and water. A volume of the stock solution is dispensed into culture media to give an initial starting test concentration of 500 ppm compound.

When the test is ready to be done, each vessel in the dilution series, except the first vessel, contains an equal volume of compound free broth. The first vessel contains twice the volume of broth with the starting concentration of test compound. One half of the broth from the first vessel is transferred to the second vessel. After being mixed, one half the resulting volume is removed from the second vessel and transferred to the third vessel. The entire cycle is repeated sufficiently to give a series of concentrations amounting to 500, 250, 125, 63, 31, 16, 8, and 4 ppm (or 100, 50, 25, 12.5, 6.2, 3.1, 1.6, and 0.8), respectively.

Each vessel is then inoculated with a cell suspension of the appropriate test organism. Bacteria are grown in broth, fungi on agar slants for a time and at a temperature appropriate to the species being tested, and algae are a mixture of green algae and blue-green bacteria grown in a nutrient media. At the end of the growth period, in the case of bacteria, the broth is vortexed to disperse the cells.

In the case of fungi, the spores are harvested by pipetting water onto the slant and dislodging the spores with a sterile loop. The cell/spore suspension is standardized by controlling incubation time, temperature, and the volume of the diluent. The suspension is then used to inoculate the vessels containing the broth compound.

The algae culture contains green algae and blue-green bacteria, and is obtained from a cooling tower in Spring House, Pennsylvania. The algae culture is grown in Allen's medium on a rotary shaker under fluorescent room lighting. This culture is further diluted with Allen's medium and then added to the test vessel.

The vessels are then incubated at the appropriate temperature. After the incubation, the vessels are examined for growth/no growth. The minimum inhibitory concentration (MIC) is defined as the lowest concentration of compound that results in complete inhibition of growth of the test organism.

The organisms tested to demonstrate biocidal activity include:

BACTERIA:

- 5 Pseudomonas fluorescens (PSFL), gram negative
Pseudomonas aeruginosa (PSAE), gram negative
Escherichia coli (ECOL), gram negative
Staphylococcus aureus (SAUR), gram positive

10 **FUNGI:**

- Aspergillus niger (ANIG)
Aureobasidium pullulans (APUL)

15 The results of the minimum inhibitory concentration (MIC) tests of compounds of this invention are shown in Table I against the microorganisms shown in Table II.

TABLE 1

BIOCIDES SECONDARY MIC TEST DATA IN PPM

20 These compounds were tested in M9G (minimal, salts media with glucose)


25	Compound #	(G-) PSFL	(G-) PSAE	(G-) ECOL	(G+) SAUR	(F) ANIG	(F) APUL	ALGAE
30	1	250	>500	250	250	>500	>500	
	2	16	32	32	16	<4	<4	
	3	8	>500	>500	8	<4	<4	
	4	32	125	125	32	<0.12	<0.12	12.5
35	5	125	>500	125	250	125	32	
	6	<4	32	<4	16	<4	<4	
	7	8	125	<4	63	<4	<63	
	8	63	250	<4	125	250	250	
40	9	125	>500	>500	125	<4	<4	
	10	32	63	<4	32	125	125	

TABLE II
MICROORGANISMS USED IN THE BIOCIDES TESTS

	<u>Name</u>		<u>GRAM ATCC No.</u>	<u>Abbreviation used</u>
	BACTERIA			
10	1. <i>Pseudomonas aeruginosa</i>	(-)	15442	PSAE
	2. <i>Staphylococcus aureus</i>	(+)	6538	SAUR
	3. <i>Escherichia coli</i>	(-)	11229	ECOL
	4. <i>Pseudomonas fluorescens</i>	(-)	948	PSFL
15	FUNGI			
	5. <i>Aspergillus niger</i>		6275	ANIG
20	6. <i>Aureobasidium pullulans</i>		9348	APUL

Claims

- 25 1. Compound of the formula



- 35 wherein R¹ is H, lower (C₁-C₄)alkyl, alkyl aryl, CH₂OR, CH₂SR, or CH(CH₃)OR; and
 R, R², and R³ are each independently H, (C₁-C₄)alkyl, (C₆-C₁₀)aryl, (C₆-C₁₄)arylalkyl, or (C₆-C₁₄)-alkaryl.
- 40 2. Compound according to claim 1 which is
N-iodopropargyloxycarbonyl glycine methyl ester;
N-iodopropargyloxycarbonyl glycine; N-iodopropargyloxycarbonyl glycine n-octyl ester; N-iodopropargyloxycarbonyl glycine iodopropargyl ester;
N-iodopropargyloxycarbonyl alanine methyl ester;
N-iodopropargyloxycarbonyl alanine; N-iodopropargyloxycarbonyl alanine n-octyl ester; N-iodopropargyloxycarbonyl phenylalanine methyl ester;
 45 N-iodopropargyloxycarbonyl phenylalanine; or N-iodopropargyloxycarbonyl phenylalanine n-octyl ester.
- 50 3. Composition comprising a compound according to claim 1 or 2, and an agronomically acceptable carrier, a cosmetic agent, a cutting oil, a soap or synthetic detergent, a stabilizer, or a film forming material, preferably wherein said compound is present in an amount of from 0.0001 to 50, more preferably from 0.001 to 10 % by weight.
4. Use of compound as defined in claim 1 or 2 as a microbicide.
- 55 5. Method of preventing or inhibiting the growth of bacteria, fungi or algae in a locus susceptible or subject thereto, comprising incorporating into or onto the locus, in an amount effective to adversely affect said growth, a compound or composition according to any of claims 1 to 3.




TABLE II
MICROORGANISMS USED IN THE BIOCIDES TESTS

	<u>Name</u>	<u>GRAM</u>	<u>ATCC No.</u>	<u>Abbreviation used</u>
BACTERIA				
10	1. Pseudomonas aeruginosa	(-)	15442	PSAE
	2. Staphylococcus aureus	(+)	6538	SAUR
	3. Escherichia coli	(-)	11229	ECOL
	4. Pseudomonas fluorescens	(-)	948	PSFL
15	FUNGI			
	5. Aspergillus niger		6275	ANIG
20	6. Aureobasidium pullulans		9348	APUL

Claims

- 25 1. Compound of the formula




- 35 wherein R¹ is H, lower (C₁-C₄)alkyl, alkyl aryl, CH₂OR, CH₂SR, or CH(CH₃)OR; and
R, R², and R³ are each independently H, (C₁-C₄)alkyl, (C₆-C₁₀)aryl, (C₆-C₁₄)arylalkyl, or (C₆-C₁₄)-alkaryl.
- 40 2. Compound according to claim 1 which is
N-iodopropargyloxycarbonyl glycine methyl ester;
N-iodopropargyloxycarbonyl glycine; N-iodopropargyloxycarbonyl glycine n-octyl ester; N-iodopropargyloxycarbonyl glycine iodopropargyl ester;
N-iodopropargyloxycarbonyl alanine methyl ester;
N-iodopropargyloxycarbonyl alanine; N-iodopropargyloxycarbonyl alanine n-octyl ester; N-iodopropargyloxycarbonyl phenylalanine methyl ester;
45 N-iodopropargyloxycarbonyl phenylalanine; or N-iodopropargyloxycarbonyl phenylalanine n-octyl ester.
- 50 3. Composition comprising a compound according to claim 1 or 2, and an agronomically acceptable carrier, a cosmetic agent, a cutting oil, a soap or synthetic detergent, a stabilizer, or a film forming material, preferably wherein said compound is present in an amount of from 0.0001 to 50, more preferably from 0.001 to 10 % by weight.
4. Use of compound as defined in claim 1 or 2 as a microbicide.
- 55 5. Method of preventing or inhibiting the growth of bacteria, fungi or algae in a locus susceptible or subject thereto, comprising incorporating into or onto the locus, in an amount effective to adversely affect said growth, a compound or composition according to any of claims 1 to 3.



6. Method according to claim 5, wherein said locus is wood, paint, adhesive, glue, paper, textile, leather, plastics, cardboard, lubricants, cosmetics, food, caulking, feed or industrial cooling water.

Patentansprüche

1. Verbindung der Formel



in der R¹ H, niederes (C₁ -C₄)-Alkyl, Alkylaryl, CH₂ OR, CH₂ SR oder CH(CH₃)OR ist und R² und R³ jeweils unabhängig H, (C₁ -C₄)-Alkyl, (C₆ -C₁₀)-Aryl, (C₆ -C₁₄)-Arylalkyl oder (C₆ -C₁₄)-Alkaryl sind.

2. Verbindung nach Anspruch 1,

dadurch gekennzeichnet,

daß sie N-Jodpropargyloxycarbonylglycinmethylester,

N-Jodpropargyloxycarbonylglycin,

N-Jodpropargyloxycarbonylglycin-n-octylester,

N-Jodpropargyloxycarbonylglycinjodpropargylester,

N-Jodpropargyloxycarbonylalaninmethylester,

N-Jodpropargyloxycarbonylalanin,

N-Jodpropargyloxycarbonylalanin-n-octylester,

N-Jodpropargyloxycarbonylphenylalaninmethylester,

N-Jodpropargyloxycarbonylphenylalanin oder

N-Jodpropargyloxycarbonylphenylalanin-n-octylester ist.

3. Zusammensetzung, enthaltend eine Verbindung nach Anspruch 1 oder 2 und einen landwirtschaftlich verträglichen Träger, ein Kosmetikum, ein Schneidöl, eine Seife oder synthetisches Waschmittel, einen Stabilisator oder ein filmbildendes Material, wobei vorzugsweise die Verbindung in einer Menge von 0,0001 bis 50, besonders bevorzugt von 0,001 bis 10 Gew.-% vorhanden ist.

4. Verwendung einer Verbindung nach Anspruch 1 oder 2 als Mikrobiozid.

5. Verfahren zum Verhindern oder Hemmen des Wachstums von Bakterien, Pilzen oder Algen an einem Ort, der anfällig dafür oder damit verunreinigt ist, durch Aufbringen auf den Ort oder in den Ort einer wirksamen Menge einer Verbindung oder Zusammensetzung nach einem der Ansprüche 1 bis 3, um das Wachstum gegenteilig zu beeinflussen.

6. Verfahren nach Anspruch 5,


dadurch gekennzeichnet,

daß der Ort Holz, Lack, Klebstoff, Leim, Papier, Textilien, Leder, Kunststoffe, Karton, Schmiermittel, Kosmetika, Nahrungsmittel, Dichtungsmassen, Frischwasser oder industrielles Kühlwasser ist.



Revendications

1. Composé de formule



où R^1 est H, un alkyle inférieur (C_1 à C_4), un alkyl aryle, CH_2OR , CH_2SR , ou $\text{CH}(\text{CH}_3)\text{OR}$; et R , R^2 et R^3 sont chacun indépendamment H, un alkyle (C_1 à C_4), un aryle (C_6 à C_{10}), un arylalkyle (C_6 à C_{14}), ou un alkaryle (C_6 à C_{14}).

2. Composé selon la revendication 1 qui est le N-iodopropargyloxycarbonyl glycine méthyl ester; la N-iodopropargyloxycarbonyl glycine; le N-iodopropargyloxycarbonyl glycine n-octyl ester; le N-iodopropargyloxycarbonyl glycine iodopropargyl ester; le N-iodopropargyloxycarbonyl alanine méthyl ester; la N-iodopropargyloxycarbonyl alanine; le N-iodopropargyloxycarbonyl alanine n-octyl ester; le N-iodopropargyloxycarbonyl phénylalanine méthyl ester; la N-iodopropargyloxycarbonyl phénylalanine; ou le N-iodopropargyloxycarbonyl phénylalanine n-octyl ester.
3. Composition comprenant un composé selon la revendication 1 ou 2, et un support agronomiquement acceptable, un agent cosmétique, une huile de coupe, un savon ou un détergent synthétique, un stabilisant, ou un matériau filmogène, de préférence dans laquelle ledit composé est présent en une quantité allant de 0,0001 à 50, avec une plus grande préférence de 0,001 à 10% en poids.
4. Utilisation du composé tel que défini dans la revendication 1 ou 2 comme microbicide.
5. Procédé pour empêcher ou inhiber le développement de bactéries, de champignons ou d'algues dans un site susceptible de ou sujet à ce développement, comprenant l'incorporation dans ou sur le site, en une quantité efficace pour affecter défavorablement ledit développement, d'un composé ou d'une composition selon l'une quelconque des revendications 1 à 3.
6. Procédé selon la revendication 5, dans lequel ledit site est du bois, une peinture, un adhésif, une colle, du papier, du textile, du cuir, du plastique, du carton, des lubrifiants, des cosmétiques, des aliments, du calfatage, de l'eau d'alimentation ou de l'eau de refroidissement industrielle.