

(43) International Publication Date
11 September 2009 (11.09.2009)(10) International Publication Number
WO 2009/109765 A1

PCT

(51) International Patent Classification:
C07F 5/06 (2006.01) **C07D 317/36** (2006.01)(21) International Application Number:
PCT/GB2009/000624(22) International Filing Date:
6 March 2009 (06.03.2009)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0804345.7 7 March 2008 (07.03.2008) GB
0821092.4 18 November 2008 (18.11.2008) GB(71) Applicant (for all designated States except US): **UNIVERSITY OF NEWCASTLE UPON TYNE** [GB/GB];
6 Kensington Terrace, Newcastle Upon Tyne, Tyne and Wear NE1 7RU (GB).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **NORTH, Michael** [GB/GB]; School of Natural Sciences, Bedson Building, University of Newcastle Upon Tyne, Newcastle Upon Tyne Tyne and Wear NE1 7RU (GB).(74) Agents: **WATSON, Robert** et al.; Mewburn Ellis LLP, 33 Gutter Lane, London, Greater London EC2V 8AS (GB).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).


Declarations under Rule 4.17:

— of inventorship (Rule 4.17(iv))

Published:

— with international search report (Art. 21(3))

(54) Title: SYNTHESIS OF CYCLIC CARBONATES



(57) **Abstract:** A dimeric aluminium(salen) catalyst of formula I: herein: Y-Q is CR_{C1}=N or CR_{C1}R_{C2}-NR_{N1}, where R_{C1}, R_{C2} and R_{N1} are independently selected from H, halo, optionally substituted C₁₋₂₀ alkyl, optionally substituted C₅₋₂₀ aryl, ether and nitro; each of the substituents R¹, R², R³, R⁴, R⁵, R⁶, R⁷, R⁸, R⁹, R¹⁰, R¹¹, R¹², R¹³, R¹⁴, R¹⁵, and R¹⁶, is independently selected from H, halo, optionally substituted C₁₋₂₀ alkyl, optionally substituted C₅₋₂₀ aryl, optionally substituted C₃₋₂₀ heterocyclyl, ether and nitro; X₁ and X₂ are independently either (i) a C₂₋₅ alkylene chain, which is optionally substituted by one or more groups selected from C₁₋₄ alkyl and C₅₋₇ aryl, or a C₁₋₃ bisoxyalkylene chain, which is optionally substituted by one or more groups selected from C₁₋₄ alkyl and C₅₋₇ aryl or (ii) represent a divalent group selected from C₅₋₇ arylene, C₅₋₇ cyclic alkylene and C₃₋₇ heterocyclylene, which may be optionally substituted; (i) (a) at least one of R¹, R², R³, R⁴, R⁵, R⁶, R⁷, R⁸, R⁹, R¹⁰, R¹¹, R¹², R¹³, R¹⁴, R¹⁵, and R¹⁶ is selected from L-A, where L is a single bond or a C₁₋₁₀ alkylene group and A is an ammonium group paired with a counterion selected from Cl, Br and I; and/or (b) at least one of X¹ and X² is a divalent C₃₋₇ heterocyclene group, containing a ring atom which is a quaternary nitrogen atom paired with a counterion selected from Cl, Br and I; and/or (c) at least one of X¹ and X² is a C₂₋₅ alkylene chain or a C₁₋₃ bisoxyalkylene chain, substituted by a group -Q-L-A, where Q is either -C(=O)-O-, -C(=O)-NH-, or a single bond; and/or (ii) (a) one of R¹, R², R³, R⁴, R⁵, R⁶, R⁷, R⁸, R⁹, R¹⁰, R¹¹, R¹², R¹³, R¹⁴, R¹⁵, and R¹⁶ is L-A', where L is as defined above and A' is an ammonium linking group bound to a solid support and paired with a counterion selected from Cl, Br and I; or (b) one of X¹ and X² is a divalent C₃₋₇ heterocyclene group, containing a ring atom which is a quaternary nitrogen forming part of an ammonium linking group bound to a solid support and paired with a counterion selected from Cl, Br and I; or (c) one of X¹ and X² is a C₂₋₅ alkylene chain or a C₁₋₃ bisoxyalkylene chain, substituted by a group -Q-L-A'.