

METHODS AND REAGENTS FOR TREATMENT AND DIAGNOSIS OF AGE-RELATED MACULAR DEGENERATION

[0001] This application claims benefit of U.S. Provisional Application Nos. 60/650,078 (filed February 14, 2005), 60/717,861 (filed September 16, 2005), 60/715,503 (filed September 9, 2005), and 60/735,697 (filed November 9, 2005), the entire contents of which are incorporated herein by reference.

STATEMENT AS TO RIGHTS TO INVENTIONS MADE UNDER FEDERALLY SPONSORED RESEARCH AND DEVELOPMENT

[0002] Work described in this application has been supported, in part, by NIH Eye Institute grant EY11515. The U.S. Government may have certain rights in the invention.

BACKGROUND OF THE INVENTION

[0003] Age-related macular degeneration (AMD) is the leading cause of irreversible vision loss in the developed world (for reviews see Zarbin, 1998, 2004; Klein et al., 2004; Ambati et al., 2003; de Jong, 2004; van Leeuwen et al., 2003) affecting approximately 15% of individuals over the age of 60. An estimated 600 million individuals are in this age demographic. The prevalence of AMD increases with age; mild, or early forms occur in nearly 30%, and advanced forms in about 7%, of the population that is 75 years and older (Klein et al., 1992; Vingerling et al., 1995a, 1995b). Clinically, AMD is characterized by a progressive loss of central vision attributable to degenerative changes that occur in the macula, a specialized region of the neural retina and underlying tissues. In the most severe, or exudative, form of the disease neovascular fronds derived from the choroidal vasculature breach Bruch's membrane and the retinal pigment epithelium (RPE) typically leading to detachment and subsequent degeneration of the retina.

[0004] AMD, a late-onset complex disorder, appears to be caused and/or modulated by a combination of genetic and environmental factors (Seddon and Chen, 2004; Tuo et al., 2004; Klein and Francis, 2003). Familial aggregation studies have estimated the genetic component to be primarily involved in as much as 25% of the disorder (Klaver et al., 1998a). According to the prevailing hypothesis, the majority of AMD cases is not a collection of multiple single-gene disorders, but instead represents a quantitative phenotype, an expression of interaction

of multiple susceptibility loci. The number of loci involved, the attributable risk conferred, and the interactions between various loci remain obscure.

[0005] Linkage and candidate gene screening analyses have provided limited insight into the genetics of AMD. Reliable association of one gene with increased risk, *ABCA4* (Allikmets et al., 1997) and one gene with decreased risk, *ApoE4* (Klaver et al., 1998b, Souied et al., 1998) for AMD have been reported. In addition, several groups have reported results of genome-wide linkage analyses (reviewed in Tuo et al., 2004; Weeks et al., 2004). Linkage of one family with AMD phenotype to a specific chromosomal region, 1q25-q31 (*ARMD1*) has been documented (Klein et al., 1998). *HEMICENTIN-1* has been suggested to be the causal gene (Schultz et al., 2003) although its role has not been reliably confirmed. The identification of overlapping loci on chromosome 1q in several studies (Weeks et al., 2001; Iyengar et al., 2003; Weeks et al., 2004) suggests that this locus may harbor AMD-associated gene(s).

[0006] Recent studies of drusen, the hallmark ocular lesions associated with the onset of AMD, have implicated a role for inflammation and other immune-mediated processes, in particular complement activation, in the etiology of early and late forms of AMD (Hageman et al., 1999, 2001; Mullins et al., 2000, 2001; Russell et al., 2000; Anderson et al., 2002, 2004; Johnson et al., 2000, 2001; Crabb et al., 2002; Ambati et al., 2003; Penfold et al., 2001; Espinosa-Heidman et al., 2003). These studies have revealed the terminal pathway complement components (C5, C6, C7, C8 and C9) and activation-specific complement protein fragments of the terminal pathway (C3b, iC3b, C3dg and C5b-9) as well as various complement pathway regulators and inhibitors (including Factor H, Factor I, Factor D, CD55 and CD59) within drusen, along Bruch's membrane (an extracellular layer comprised of elastin and collagen that separates the RPE and the choroid) and within RPE cells overlying drusen (Johnson et al., 2000, 2001; Mullins et al. 2000, 2001; Crabb et al., 2002). Many of these drusen-associated molecules are circulating plasma proteins previously thought to be synthesized primarily by the liver. Interestingly, many also appear to be synthesized locally by RPE and/or choroidal cells.

[0007] Activation of the complement system plays a key role in normal host defense and in the response to injury (Kinoshita, 1991). Inappropriate activation and/or control of this system, often caused by mutations in specific complement-associated genes, can contribute to autoimmune sequelae and local tissue destruction (Holers, 2003; Liszewski and Atkinson, 1991; Morgan and Walport, 1991; Shen and Meri, 2003), as has been shown in

atherosclerosis (Torzewski et al., 1997; Niculescu et al., 1999), Alzheimer's disease (Akiyama et al., 2000) and glomerulonephritis (Schwartz et al., 2001).

[0008] Membranoproliferative glomerulonephritis type 2 (MPGN II) is a rare disease that is associated with uncontrolled systemic activation of the alternative pathway of the complement cascade. The disease is characterized by the deposition of abnormal electron-dense material comprised of C3 and C3c, proteins involved in the alternative pathway of complement, within the renal glomerular basement membrane, which eventually leads to renal failure. Interestingly, many patients with MPGNII develop macular drusen, RPE detachments and choroidal neovascular membranes that are clinically and compositionally indistinguishable from those that form in AMD, although they are often detected in the second decade of life (Mullins et al., 2001; O'Brien et al., 1993; Huang et al., 2003; Colville et al., 2003; Duvall-Young et al., 1989a, 1989b; Raines et al., 1989; Leys et al., 1990; McAvoy and Silvestri, 2004; Bennett et al., 1989; Orth and Ritz, 1998; Habib et al., 1975).

[0009] In most patients with MPGNII, the inability to regulate the complement cascade is mediated by an autoantibody directed against C3bBb. Other MPGN II patients, however, harbor mutations in Factor H (Ault et al., 1997; Dragon-Durey et al., 2004) a major inhibitor of the alternative complement pathway. A point mutation in Factor H (H166R) causes MPGNII in the Yorkshire pig (Jansen et al., 1998) and Factor H deficient mice develop severe glomerulonephritis (Pickering et al., 2002). Moreover, affected individuals within some extended families with MPGNIII, a related disorder, show linkage to chromosome 1q31-32 (Neary et al., 2002) a region that overlaps a locus that has been identified in genome-wide linkage studies for AMD (see above). This particular locus contains a number of complement pathway-associated genes. One group of these genes, referred to as the regulators of complement activation (RCA) gene cluster, contains the genes that encode Factor H, five Factor H-related genes (*CFHR1*, *CFHR2*, *CFHR3*, *CFHR4* and *CFHR5*), and the beta subunit of coagulation factor XIII. A second cluster of complement pathway-associated genes, including C4BPA, C4BPB, C4BPAL2, DAF (CD55) CR1, CR2, CR1L and MCP (CD46) lies immediately adjacent to the 1q25-31 locus.

BRIEF SUMMARY OF THE INVENTION

[0010] The disclosure relates to polymorphisms and haplotypes in the complement Factor H gene that are associated with development of age-related macular degeneration (AMD) and membranoproliferative glomerulonephritis type 2 (MPGNII). The disclosure also relates to

polymorphisms and haplotypes in the complement Factor H-related 5 (CFHR5) genes that are associated with development of AMD and MPGNII. The disclosure provides methods of diagnosing, monitoring, and treating these and other diseases.

[0010a] In a first aspect, there is provided a method of screening for susceptibility to developing age-related macular degeneration (AMD) in a subject, comprising screening for the presence or absence of a haplotype in the Complement Factor H (CFH) gene, wherein the haplotype comprises

- (a) rs1061170 with a nucleotide encoding histidine at position 402; or
- (b) rs800292 with a nucleotide encoding isoleucine at position 62.

[0010b] In a second aspect, there is provided a method of screening for susceptibility to age-related macular degeneration (AMD) in a subject, comprising screening for a haplotype, wherein the haplotype is:

- a) a risk haplotype in the CFH gene that is more frequently present in individuals diagnosed with AMD compared to healthy individuals, or
- b) a protective haplotype in the CFH gene that is more frequently present in healthy individuals compared to individuals diagnosed with AMD or
- c) a neutral haplotype in the CFH gene; and

wherein said screening comprises detecting the presence or absence of a variant sequence at any one or more of rs1061170, rs800292, rs1061147, intron 2 (IVS2 or insTT); rs2274700; rs1061170, exon 10A or exon 22.

[0010c] In a third aspect, there is provided a device or kit for screening susceptibility to AMD in a subject, comprising oligonucleotides that distinguish alleles at one, or more than one, CFH gene polymorphic site listed in Table A.

[0010d] In a fourth aspect, there is provided a device or kit for screening for susceptibility to AMD in a subject, comprising oligonucleotides that distinguish alleles at one, or more than one, of rs1061170, rs800292, rs1061147, intron 2 (IVS2 or insTT); rs2274700; rs1061170, rs9427661, rs9427662, rs12097550, exon 10A, and exon 22.

[0011] The disclosure provides a diagnostic method for determining a subject's propensity to develop age-related macular degeneration (AMD), comprising detecting the presence or absence of a variation or variations at polymorphic site or polymorphic sites of the Factor H gene. In one embodiment, the disclosure provides methods of diagnosing an increased susceptibility to developing AMD involving detecting the presence or absence of a polymorphism in the Factor H gene of an individual. The methods may include obtaining the DNA from an individual and analyzing the DNA from the individual to determine whether the DNA contains a polymorphism in the Factor H gene. Certain polymorphisms indicate the individual has an increased susceptibility to developing AMD relative to a control population. Certain polymorphisms indicate the individual has a reduced likelihood of developing AMD. Certain polymorphisms indicate the individual has neither an increased nor a reduced likelihood of developing AMD.

[0012] In one embodiment, a method of diagnosing a propensity to develop age-related macular degeneration (AMD) in a subject involves obtaining a sample of DNA from the subject and detecting in the DNA of the patient the presence or absence of a polymorphism associated with development of AMD, the presence of the polymorphism being an indication that the subject has an increased propensity to develop AMD and the absence of the polymorphism being an indication that the subject has a reduced propensity to develop AMD.

[0013] The disclosure also provides methods of diagnosing susceptibility to developing AMD involving determining an individual's Factor H haplotype. The methods include obtaining the DNA from an individual and analyzing the DNA of the individual to determine their Factor H haplotype. Certain haplotypes (risk haplotypes) indicate the individual has an increased susceptibility to develop AMD. Certain haplotypes (protective haplotypes) indicate the individual has a decreased susceptibility to develop AMD. Certain haplotypes (neutral haplotypes) indicate the individual has neither an increased nor a reduced likelihood of developing AMD.

[0014] In a related embodiment the presence or absence of a variation at a polymorphic site of the Factor H gene is determined by analysis of a gene product, such as an RNA or a Factor H protein (e.g., protein isoform) encoded by the gene. Expression of a variant protein is an indication of a variation in the Factor H gene and can indicate an increased or reduced

propensity to develop AMD. Proteins can be detected using immunoassays and other methods.

[0015] The disclosure further provides methods of diagnosing susceptibility to developing AMD or other diseases by detecting a variant Factor H polypeptide in a biological sample of an individual. In one embodiment, an antibody-based assay is used to diagnose AMD or other diseases in an individual by contacting a biological sample, e.g., a serum sample, of the individual with the antibody and detecting the presence or absence of the variant Factor H polypeptide. In an embodiment, the antibody specifically interacts with an epitope specific to a variant Factor H polypeptide (i.e., not found in the wild-type Factor H polypeptide). In an embodiment, a separation-based assay (e.g., PAGE) is used to diagnose AMD or other diseases in an individual by detecting the presence or absence of the variant Factor H polypeptide in a biological sample, e.g., a serum sample, of the individual.

[0016] The disclosure provides methods of treating an individual with AMD (e.g., an individual in whom a polymorphism or haplotype indicative of elevated risk of developing symptomatic AMD is detected) or other disease involving a variant Factor H gene by modulating the type and/or amount of systemic and/or ocular levels of Factor H. The Factor H polypeptide may be a wild-type Factor H polypeptide or a variant Factor H polypeptide. The Factor H polypeptide may be a Factor H polypeptide with a sequence encoded by neutral or protective alleles rather than alleles associated with a risk haplotype. In one embodiment, the method includes administering to the individual a Factor H polypeptide in an amount effective to reduce a symptom of the disease. In one embodiment, the method includes administering to an individual a Factor H polypeptide in an amount effective to reduce the propensity to develop symptoms of the disease and delay development or progression of the disease. In one embodiment, the method includes administering blood that contains Factor H. In one embodiment, the methods include administering a nucleic acid (e.g., transgene) including a nucleotide sequence encoding a Factor H polypeptide. In one embodiment, the methods include administering cells that express a Factor H polypeptide.

[0017] The disclosure also provides methods of treating an individual with AMD (e.g., an individual in whom a polymorphism or haplotype indicative of elevated risk of developing symptomatic AMD is detected) or other disease involving a variant Factor H gene. In one embodiment, the method includes administering to the patient an agent that decreases the amount of a variant Factor H or expression of a gene encoding Factor H in an

amount effective to reduce a symptom of the disease in the patient. In a related embodiment a therapeutic amount of an inhibitor (e.g., inactivator) of the variant Factor H polypeptide in the individual is administered.

[0018] In one embodiment an inhibitory nucleic acid (e.g., an RNA complementary to at least a portion of the nucleotide sequence of the variant Factor H polypeptide) in the individual is administered. In one embodiment, purified anti-sense RNA complementary to RNA encoding a variant Factor H polypeptide is administered.

[0019] In another embodiment a therapeutic amount of an anti-CFH antibody sufficient to partially inactivate the variant Factor H polypeptide in the individual is administered.

[0020] In another embodiment, the individual is treated to remove deleterious forms of Factor H from blood (e.g., by plasmaphoresis, antibody-directed plasmaphoresis, or complexing with a Factor H binding moiety, e.g., heparin).

[0021] The disclosure provides purified DNA encoding a variant Factor H polypeptide, purified RNA encoding a variant Factor H polypeptide, purified anti-sense RNA complementary to the RNA encoding a variant Factor H polypeptide, and purified variant Factor H polypeptide. The disclosure also provides nucleic acids for expressing wild-type or variant Factor H polypeptides or biologically active fragments of Factor H.

[0022] The disclosure provides gene therapy vectors comprising nucleic acid encoding the Factor H polypeptide. The vector may include a promoter that drives expression of the Factor H gene in multiple cell types. Alternatively, the vector may include a promoter that drives expression of the Factor H gene only in specific cell types, for example, in cells of the retina or in cells of the kidney. Pharmaceutical compositions are disclosed containing a gene therapy vector encoding a Factor H protein and a pharmaceutically acceptable excipient, where the composition is free of pathogens and suitable for administration to a human patient. In one embodiment the encoded Factor H polypeptide is a protective variant.

[0023] The disclosure provides a composition containing recombinant or purified Factor H polypeptide, where the polypeptide is a protective variant.

[0024] The disclosure also provides a pharmaceutical composition containing recombinant or purified Factor H polypeptide and a pharmaceutically acceptable excipient, where the composition is free of pathogens and suitable for administration to a human patient. In one embodiment the encoded Factor H polypeptide has the wild-type sequence. In one embodiment the encoded Factor H polypeptide is a protective variant.

[0025] The disclosure provides antibodies that specifically interact with a variant Factor H polypeptide but not with a wild-type Factor H polypeptide. These antibodies may be polyclonal or monoclonal and may be obtained by subtractive techniques. These antibodies may be sufficient to inactivate a variant Factor H polypeptide.

The disclosure also provides pharmaceutical compositions containing an anti-Factor H antibody and a pharmaceutically acceptable excipient, where the composition is free of pathogens and suitable for administration to a human patient.

[0026] The disclosure provides methods for identifying variant Factor H proteins associated with increased or reduced risk of developing AMD. In one embodiment, the disclosure provides a method of identifying a protective Factor H protein by (a) identifying an individual as having a protective haplotype and (b) determining the amino acid sequence(s) of Factor H encoded in the genome of the individual, where a protective Factor H protein is encoded by an allele having a protective haplotype. In one embodiment, the disclosure provides a method of identifying a neutral Factor H protein by (a) identifying an individual as having a neutral haplotype and (b) determining the amino acid sequence(s) of Factor H encoded in the genome of the individual, where a neutral Factor H protein is encoded by an allele having a neutral haplotype. In a related embodiment, the disclosure provides a method of identifying a variant form of Factor H associated with decreased risk of developing AMD comprising (a) identifying an individual as having a haplotype or diplotype associated with a decreased risk of developing AMD; (b) obtaining genomic DNA or RNA from the individual; and (c) determining the amino acid sequence(s) of the Factor H encoded in the individual's genome, where a protective Factor H protein is encoded by an allele having a haplotype associated with a decreased risk of developing AMD. In an embodiment, the protective or neutral Factor H proteins do not have the amino acid sequence of the wild-type Factor H polypeptide.

[0027] In a related method, a form of Factor H associated with increased risk of developing AMD is identified by (a) identifying an individual as having a risk haplotype and (b) determining the amino acid sequence(s) of Factor H encoded in the genome of the individual, where a risk Factor H protein is encoded by an allele having a risk haplotype. In a related embodiment, the disclosure provides a method of identifying a variant form of Factor H associated with increased risk of developing AMD comprising (a) identifying an individual as having a haplotype or diplotype associated with an increased risk of developing AMD; (b) obtaining genomic DNA or RNA from the individual; and (c) determining the amino acid

sequence(s) of the Factor H encoded in the individual's genome, where a risk Factor H protein is encoded by an allele having a haplotype associated with an increased risk of developing AMD. In an embodiment, the risk Factor H proteins do not have the amino acid sequence of the wild-type Factor H polypeptide.

[0028] The disclosure provides methods of diagnosing a propensity or susceptibility to develop AMD or other diseases by detecting the ratio of full-length Factor H to truncated Factor H in a biological sample of a patient. In one embodiment, a method of diagnosing a propensity or susceptibility to develop AMD in a subject involves obtaining a sample of RNA from the subject and detecting in the RNA of the patient the ratio of expression of exon 10 (i.e., full-length Factor H) to exon 10A (i.e., truncated Factor H), the increase in ratio being an indication that the subject has an increased propensity or susceptibility to develop AMD and the decrease in ratio being an indication that the subject has a reduced propensity or susceptibility to develop AMD. In one embodiment, a method of diagnosing a propensity or susceptibility to develop AMD in a subject involves obtaining a sample of protein from the subject and detecting in the protein of the patient the ratio of expression of full-length Factor H to truncated Factor H, the increase in ratio being an indication that the subject has an increased propensity or susceptibility to develop AMD and the decrease in ratio being an indication that the subject has a reduced propensity or susceptibility to develop AMD.

[0029] The disclosure provides cells containing recombinant or purified nucleic acid encoding a Factor H protein or fragment thereof, e.g., a nucleic acid derived from the Factor H gene. The cells may be bacterial or yeast, or any other cell useful for research and drug development. Thus, the disclosure provides an isolated host cell or cell line expressing a recombinant variant human Factor H. In an embodiment, the variant is a risk variant and has a histidine at amino acid position 402. In an embodiment, the variant is a protective variant and has isoleucine at amino acid position 62. In an embodiment, the variant is a neutral variant. In an embodiment, the risk, protective or neutral variant Factor H proteins do not have the amino acid sequence of the wild-type Factor H polypeptide.

[0030] The disclosure provides transgenic non-human animals whose somatic and germ cells contain a transgene encoding a human variant Factor H polypeptide. Transgenic animals of the disclosure are used as models for AMD and for screening for agents useful in treating AMD. The animal may be a mouse, a worm, or any other animal useful for research and drug development (such as recombinant production of Factor H). In an

embodiment, the Factor H is a variant human Factor H, wherein said variant has isoleucine amino acid 62 or has histidine at amino acid 402.

[0031] The disclosure provides methods of screening for polymorphic sites linked to polymorphic sites in the Factor H gene described in TABLES 1A, 1B and 1C. These methods involve identifying a polymorphic site in a gene that is linked to a polymorphic site in the Factor H gene, wherein the polymorphic form of the polymorphic site in the Factor H gene is associated AMD, and determining haplotypes in a population of individuals to indicate whether the linked polymorphic site has a polymorphic form in equilibrium disequilibrium with the polymorphic form of the Factor H gene that associates with the AMD phenotype.

[0032] The disclosure provides diagnostic, therapeutic and screening methods for MPGNII, carried out as described above for AMD.

[0033] The disclosure provides a diagnostic method for determining a subject's propensity to develop AMD or MPGNII, comprising detecting the presence or absence of a variation or variations at polymorphic site or polymorphic sites of the CFHR5 gene. In one embodiment, the disclosure provides methods of diagnosing an increased susceptibility to developing AMD or MPGNII involving detecting the presence or absence of a polymorphism in the CFHR5 gene of an individual. The methods may include obtaining the DNA from an individual and analyzing the DNA from the individual to determine whether the DNA contains a polymorphism in the CFHR5 gene. Certain polymorphisms indicate the individual has an increased susceptibility to developing AMD or MPGNII. Certain polymorphisms indicate the individual has a reduced likelihood of developing AMD or MPGNII. Certain polymorphisms indicate the individual has neither an increased nor a reduced likelihood of developing AMD or MPGNII.

[0034] In one embodiment, a method of diagnosing a propensity to develop AMD or MPGNII in a subject involves obtaining a sample of DNA from the subject and detecting in the DNA of the patient the presence or absence of a polymorphism associated with development of AMD or MPGNII, the presence of the polymorphism being an indication that the subject has an increased propensity to develop AMD or MPGNII and the absence of the polymorphism being an indication that the subject has a reduced propensity to develop AMD or MPGNII.

[0035] In a related embodiment the presence or absence of a variation at a polymorphic site of the CFHR5 gene is determined by analysis of a gene product, such as an RNA or a CFHR5

protein (e.g., protein isoform) encoded by the gene. Expression of a variant protein is an indication of a variation in the CFHR5 gene and can indicate an increased or reduced propensity to develop AMD or MPGNII. Proteins can be detected using immunoassays and other methods.

[0036] The disclosure provides methods of diagnosing susceptibility to developing AMD or MPGNII involving determining an individual's CFHR5 haplotype. The methods include obtaining the DNA from an individual and analyzing the DNA of the individual to determine their CFHR5 haplotype. Certain haplotypes (risk haplotypes) indicate the individual has an increased susceptibility to develop AMD or MPGNII relative to a control population. Certain haplotypes (protective haplotypes) indicate the individual has an decreased susceptibility to develop AMD or MPGNII. Certain haplotypes (neutral haplotypes) indicate the individual has neither an increased nor a reduced likelihood of developing AMD or MPGNII.

[0037] The disclosure provides methods of diagnosing susceptibility to developing AMD or MPGNII or other diseases by detecting a variant CFHR5 polypeptide in a biological sample of an individual. In one embodiment, an antibody-based assay is used to diagnose AMD or MPGNII or other diseases in an individual by contacting a biological sample, e.g., a serum sample, of the individual with the antibody and detecting the presence or absence of the variant CFHR5 polypeptide. In an embodiment, the antibody specifically interacts with an epitope specific to a variant CFHR5 polypeptide (i.e., not found in the wild-type CFHR5 polypeptide). In an embodiment, a separation-based assay (e.g., PAGE) is used to diagnose MPGNII or other diseases in an individual by detecting the presence or absence of the variant CFHR5 polypeptide in a biological sample, e.g., a serum sample, of the individual. Various types of immunoassay formats can be used to assay CFH or CFHR5 polypeptide or protein in a sample. These include sandwich ELISA, radioimmunoassay, fluoroimmunoassay, immunohistochemistry assay, dot-blot, dip-stick and Western Blot.

[0038] The disclosure provides methods of treating an individual with or at risk for AMD or MPGNII (e.g., an individual in whom a polymorphism or haplotype indicative of elevated risk of developing symptomatic AMD or MPGNII is detected) or other disease involving a variant CFHR5 gene by modulating the type and/or amount of systemic and/or renal levels of CFHR5. The CFHR5 polypeptide may be a CFHR5 polypeptide encoded by neutral or protective alleles rather than alleles associated with a risk haplotype. In one embodiment, the method includes administering to the individual a CFHR5 polypeptide in an

amount effective to reduce a symptom of the disease. In one embodiment, the method includes administering to an individual a CFHR5 polypeptide in an amount effective to reduce the propensity to develop symptoms of the disease and delay development or progression of the disease. In one embodiment, the method includes administering blood, which contains CFHR5. In one embodiment, the methods include administering a nucleic acid (e.g., transgene) including a nucleotide sequence encoding a CFHR5 polypeptide.

[0039] The disclosure provides methods of treating an individual with AMD or MPGNII (e.g., an individual in whom a polymorphism or haplotype indicative of elevated risk of developing symptomatic AMD or MPGNII is detected) or other disease involving a variant CFHR5 gene. In one embodiment, the method includes administering to the patient an agent that decreases the amount of a variant CFHR5 or expression of a gene encoding CFHR5 in an amount effective to reduce a symptom of the disease in the patient. The CFHR5 polypeptide may be a wild-type CFHR5 polypeptide or a variant CFHR5 polypeptide.

[0040] In one embodiment an inhibitory nucleic acid (e.g., an RNA complementary to at least a portion of the nucleotide sequence of the variant CFHR5 polypeptide) in the individual is administered. In one embodiment, purified anti-sense RNA complementary to RNA encoding a variant CFHR5 polypeptide is administered.

[0041] In another embodiment a therapeutic amount of an anti-CFHR5 antibody sufficient to partially inactivate the variant CFHR5 polypeptide in the individual is administered.

[0042] In a related embodiment a therapeutic amount of an inhibitor (e.g., inactivator) of the variant CFHR5 polypeptide in the individual is administered.

[0043] In another embodiment, the individual is treated to remove deleterious forms of CFHR5 from blood (e.g., by plasmaphoresis, antibody-directed plasmaphoresis, or complexing with a CFHR5 binding moiety, e.g., heparin).

[0044] The disclosure provides purified DNA encoding a variant CFHR5 polypeptide, purified RNA encoding a variant CFHR5 polypeptide, purified anti-sense RNA complementary to the RNA encoding a variant CFHR5 polypeptide, and purified variant CFHR5 polypeptide. The disclosure also provides nucleic acids for expressing wild-type or variant CFHR5 polypeptides or biologically active fragments of CFHR5.

[0045] The disclosure provides gene therapy vectors comprising nucleic acid encoding the CFHR5 polypeptide. The vector may include a promoter that drives expression of the CFHR5 gene in multiple cell types. Alternatively, the vector may include a promoter

that drives expression of the CFHR5 gene only in specific cell types, for example, in cells of the retina or cells of the kidney (e.g., endothelial cells, mesangial cells, podocytes).

Pharmaceutical compositions are disclosed containing a gene therapy vector encoding a CFHR5 protein and a pharmaceutically acceptable excipient, where the composition is free of pathogens and suitable for administration to a human patient. In one embodiment the encoded CFHR5 polypeptide is a protective variant.

[0046] The disclosure provides a composition containing recombinant or purified CFHR5 polypeptide, where the polypeptide is a protective variant.

[0047] The disclosure provides a pharmaceutical composition containing recombinant or purified CFHR5 polypeptide and a pharmaceutically acceptable excipient, where the composition is free of pathogens and suitable for administration to a human patient. In one embodiment the encoded CFHR5 polypeptide has the wild-type sequence. In one embodiment the encoded CFHR5 polypeptide is a protective variant.

[0048] The disclosure provides antibodies that specifically interact with a variant CFHR5 polypeptide but not with a wild-type CFHR5 polypeptide. These antibodies may be polyclonal or monoclonal and may be obtained by subtractive techniques. These antibodies may be sufficient to inactivate a variant CFHR5 polypeptide. The disclosure also provides pharmaceutical compositions containing an anti-CFHR5 antibody and a pharmaceutically acceptable excipient, where the composition is free of pathogens and suitable for administration to a human patient.

[0049] The disclosure provides methods for identifying variant CFHR5 proteins associated with increased or reduced risk of developing AMD or MPGNII. In one embodiment, the disclosure provides a method of identifying a protective CFHR5 protein by (a) identifying an individual as having a protective haplotype and (b) determining the amino acid sequence(s) of CFHR5 encoded in the genome of the individual, where a protective CFHR5 protein is encoded by an allele having a protective haplotype. In one embodiment, the disclosure provides a method of identifying a neutral CFHR5 protein by (a) identifying an individual as having a neutral haplotype and (b) determining the amino acid sequence(s) of CFHR5 encoded in the genome of the individual, where a neutral CFHR5 protein is encoded by an allele having a neutral haplotype. In a related embodiment, the disclosure provides a method of identifying a variant form of CFHR5 associated with decreased risk of developing AMD or MPGNII comprising (a) identifying an individual as having a haplotype or diplotype associated with a decreased risk of developing AMD or MPGNII; (b) obtaining genomic

DNA, or RNA, from the individual; and (c) determining the amino acid sequence(s) of the CFHR5 encoded in the individual's genome, where a protective CFHR5 protein is encoded by an allele having a haplotype associated with a decreased risk of developing AMD or MPGNII. In an embodiment, the protective or neutral CFHR5 proteins do not have the amino acid sequence of the wild-type CFHR5 polypeptide.

[0050] In a related method, a form of CFHR5 associated with increased risk of developing AMD or MPGNII is identified by (a) identifying an individual as having a risk haplotype and (b) determining the amino acid sequence(s) of CFHR5 encoded in the genome of the individual, where a risk CFHR5 protein is encoded by an allele having a risk haplotype. In a related embodiment, the disclosure provides a method of identifying a variant form of CFHR5 associated with increased risk of developing AMD or MPGNII comprising (a) identifying an individual as having a haplotype or diplotype associated with an increased risk of developing AMD or MPGNII; (b) obtaining genomic DNA or RNA from the individual; and (c) determining the amino acid sequence(s) of the CFHR5 encoded in the individual's genome, where a risk CFHR5 protein is encoded by an allele having a haplotype associated with an increased risk of developing AMD or MPGNII. In an embodiment, the risk CFHR5 proteins do not have the amino acid sequence of the wild-type CFHR5 polypeptide.

[0051] The disclosure provides cells containing recombinant or purified nucleic acid derived from the CFHR5 gene. The cells may be bacterial or yeast, or any other cell useful for research and drug development. Thus, the disclosure provides an isolated host cell or cell line expressing a recombinant variant human CFHR5. In an embodiment, the CFHR5 variant is a risk variant and has a serine at amino acid position 46. In an embodiment, the CFHR5 variant is a neutral variant. In an embodiment, the risk, protective or neutral variant CFHR5 proteins does not have the amino acid sequence of the wild-type CFHR5 polypeptide.

[0052] The disclosure provides transgenic non-human animals whose somatic and germ cells contain a transgene encoding a human variant CFHR5 polypeptide. Transgenic animals of the disclosure are used as models for AMD or MPGNII and for screening for agents useful in treating AMD or MPGNII. The animal may be a mouse, a worm, or any other animal useful for research and drug development (such as recombinant production of CFHR5). In an embodiment, the CFHR5 is a variant human CFHR5, wherein said CFHR5 variant has serine at amino acid 46.

[0053] The disclosure provides methods of screening for polymorphic sites linked to polymorphic sites in the CFHR5 gene described in TABLE 14 or TABLE 15.

These methods involve identifying a polymorphic site in a gene that is linked to a polymorphic site in the CFHR5 gene, wherein the polymorphic form of the polymorphic site in the CFHR5 gene is associated with AMD or MPGNII, and determining haplotypes in a population of individuals to indicate whether the linked polymorphic site has a polymorphic form in equilibrium disequilibrium with the polymorphic form of the CFHR5 gene that associates with the AMD or MPGNII phenotype.

[0054] The disclosure provides kits for analysis of a Factor H haplotype. The kits may be used for diagnosis of AMD in a patient. The kits may include one or more Factor H Factor H allele-specific oligonucleotides (e.g., allele-specific primers or probes), or antibodies that specifically recognize the Factor H polypeptide. The Factor H allele-specific oligonucleotides may include sequences derived from the coding (exons) or non-coding (promoter, 5' untranslated, introns or 3' untranslated) region of the Factor H gene. The Factor H-specific antibodies may recognize the normal or wild-type H polypeptide or variant Factor H polypeptides in which one or more non-synonymous single nucleotide polymorphisms (SNPs) are present in the Factor H coding region. The kits may be used to diagnose AMD, as well as other diseases associated with SNPs in the Factor H gene, such as MPGNII. The kits may include instead, or in addition, one or more Factor H-Related 5 (CFHR5) allele-specific oligonucleotides (e.g., primers and probes), or antibodies that specifically recognize the CFHR5 polypeptide. The CFHR5 allele-specific primers and Factor H-related 5 allele-specific oligonucleotides may include sequences derived from the coding (exons) or non-coding (promoter, 5' untranslated, introns or 3' untranslated) region of the Factor H-related 5 gene. The Factor H-related 5-specific antibodies may recognize the normal or wild-type H polypeptide or variant Factor H-related 5 polypeptides in which one or more non-synonymous single nucleotide polymorphisms (SNPs) are present in the Factor H-related 5 coding region.

[0055] In one embodiment the kit contains probes or primers that distinguish alleles at a polymorphic site listed in TABLE 1A, TABLE 1B and/or TABLE 1C. In an embodiment the probes are primers for nucleic acid amplification of a region spanning a Factor H gene polymorphic site listed in TABLE 1A, TABLE 1B and/or TABLE 1C. In an embodiment the kit has probes or primers that distinguish alleles at more than one polymorphic site listed in TABLE 1A, TABLE 1B and/or TABLE 1C. In an embodiment the kit has probes or primers that distinguish alleles at more than one polymorphic site, where the polymorphic site includes: (a) rs529825; (b) rs800292; (c) rs3766404; (d) rs1061147; (e) rs1061170; (f)

rs203674; (g) at least one of rs529825 and rs800292; (h) at least one of rs1061147, rs1061170 and rs203674; (i) at least one of rs529825 and rs800292; and rs3766404; and at least one of rs1061147, rs1061170 and rs203674; or (j) at least rs529825, rs800292, rs3766404, rs1061170 and rs203674.

[0056] In a related embodiment the kit has probes or primers that distinguish alleles at more than one polymorphic site, where the polymorphic site includes: (a) rs529825; (b) rs800292; (c) intron 2 (IVS2 or insTT) (d) rs3766404; (e) rs1061147; (f) rs1061170; (g) exon 10A; (h) rs203674; (i) rs375046; (j) rs529825 and rs800292; (k) at least two or three of rs1061147, rs1061170 and rs203674; (l) at least one of rs529825 and rs800292; and intron 2; and rs3766404; and at least one of rs1061147, rs1061170 and rs203674; and exon 10A; and, rs375046; (m) at least rs529825; rs800292; intron 2; rs3766404; rs1061170; exon 10A; rs203674; and rs375046; (n) at least two, or at least three, or at least four of rs529825, rs800292, intron 2; rs3766404, rs1061170, exon 10A, rs203674, and rs375046; (o) exon 22 (1210); or (p) exon 22 (1210) in combination with any aforementioned variation or set of variations (a-o). In an embodiment the kit has probes or primers that distinguish alleles at one or both of rs460897 and rs460184. In an embodiment the kit has probes or primers that distinguish alleles at more than one polymorphic site, where the polymorphic sites are selected from: (a) rs3753394; (b) rs529825; (c) rs800292; (d) intron 2 (IVS2 or insTT); (e) rs3766404; (f) rs1061147; (g) rs1061170; (h) rs2274700; (i) rs203674; (j) rs3753396; and (k) rs1065489.

[0057] In one embodiment the kit contains, instead of, or in addition to, the probes described above, probes, primers, antibodies and the like that distinguish polymorphic sites in the CFHR5 gene. The disclosure provides kits for the diagnosis of AMD or MPGNII in a patient based on variation in the CFHR5 gene. The kits may include one or more CFHR5-specific probes or CFHR5 allele-specific oligonucleotides, or antibodies that specifically recognize the CFHR5 polypeptide. The CFHR5-specific primers and CFHR5 allele-specific oligonucleotides may include sequences derived from the coding (exons) or non-coding (promoter, 5' untranslated, introns or 3' untranslated) region of the CFHR5 gene. The CFHR5-specific antibodies may recognize the normal or wild-type CFHR5 polypeptide or variant CFHR5 polypeptides in which one or more non-synonymous single nucleotide polymorphisms (SNPs) are present in the CFHR5 coding region. The kits may be used to diagnose AMD or MPGNII, as well as other diseases associated with SNPs in the CFHR5 gene.

[0058] In one embodiment the kit contains probes or primers that distinguish alleles at a polymorphic site listed in TABLE 14 or TABLE 15. In an embodiment the probes are primers for nucleic acid amplification of a region spanning a CFHR5 gene polymorphic site listed in TABLE 14 or TABLE 15. In an embodiment the kit has probes or primers that distinguish alleles at more than one polymorphic site listed in TABLE 14 or TABLE 15. In an embodiment the kit comprises probes or primers that distinguish alleles one, two or all of the following polymorphic sites: rs9427661 (-249T>C); rs9427662 (-20T>C); and rs12097550 (P46S).

[0059] In one embodiment the kit contains probes or primers that distinguish alleles at a polymorphic site in the CFH gene and at a polymorphic site in a CFHR gene, such as CFHR5.

[0060] The disclosure provides devices for determining a subject's haplotype. The devices are useful for, for example, the diagnosis of AMD or other diseases in a patient. In one embodiment the device contains probes or primers that distinguish alleles at a polymorphic site listed in TABLE 1A, 1B and/or 1C. In an embodiment the probes are primers for nucleic acid amplification of a region spanning a Factor H gene polymorphic site listed in TABLE 1A, 1B and/or 1C. In an embodiment the device has probes or primers that distinguish alleles at more than one polymorphic site listed in TABLE 1A, 1B and/or 1C. In an embodiment the device has probes or primers that distinguish alleles at more than one polymorphic site, where the polymorphic site includes (a) rs529825; (b) rs800292; (c) rs3766404; (d) rs1061147; (e) rs1061170; (f) rs203674; (g) at least one of rs529825 and rs800292; (h) at least one of rs1061147, rs1061170 and rs203674; (i) at least one of rs529825 and rs800292; and rs3766404; and at least one of rs1061147, rs1061170 and rs203674; or (j) at least rs529825, rs800292, rs3766404, rs1061170 and rs203674.

[0061] The kits described above and their contents may also be used to identity a propensity to develop MPGNII or to determine a Factor H haplotype for any purpose.

[0062] In a related embodiment the device has probes or primers that distinguish alleles at more than one polymorphic site, where the polymorphic site includes: (a) rs529825; (b) rs800292; (c) intron 2 (IVS2 or insTT) (d) rs3766404; (e) rs1061147; (f) rs1061170; (g) exon 10A; (h) rs203674; (i) rs375046; (j) rs529825 and rs800292; (k) at least two or three of rs1061147, rs1061170 and rs203674; (l) at least one of rs529825 and rs800292; and intron 2; and rs3766404; and at least one of rs1061147, rs1061170 and rs203674; and exon 10A; and rs375046; (m) at least rs529825; rs800292; intron 2; rs3766404; rs1061170; exon 10A;

rs203674; and rs375046; (n) at least two, or at least three, or at least four of rs529825; rs800292; intron 2; rs3766404; rs1061170; exon 10A; rs203674; and rs375046; (o) exon 22 (1210); or (p) exon 22 (1210) in combination with any aforementioned variation or set of variations (a-o). In an embodiment the device has probes or primers that distinguish alleles at one or both of rs460897 and rs460184. In an embodiment the device has probes or primers that distinguish alleles at more than one polymorphic site, where the polymorphic sites are selected from: (a) rs3753394; (b) rs529825; (c) rs800292; (d) intron 2 (IVS2 or insTT); (e) rs3766404; (f) rs1061147; (g) rs1061170; (h) rs2274700; (i) rs203674; (j) rs3753396; and (k) rs1065489. In an embodiment the device has probes or primers that distinguish alleles at more than one polymorphic site, where the polymorphic sites are selected from: (a) rs3753394; (b) rs529825; (c) rs800292; (d) intron 2 (IVS2 or insTT); (e) rs3766404; (f) rs1061147; (g) rs1061170; (h) rs2274700; (i) rs203674; (j) rs3753396; and (k) rs1065489.

[0063] The disclosure provides devices for the diagnosis of AMD or MPGNII in a patient. In one embodiment the device contains probes or primers that distinguish alleles at a polymorphic site listed in TABLE 14 or TABLE 15. In an embodiment the probes are primers for nucleic acid amplification of a region spanning a CFHR5 gene polymorphic site listed in TABLE 14 or TABLE 15. In an embodiment the device has probes or primers that distinguish alleles at more than one polymorphic site listed in TABLE 14 or TABLE 15. Devices of the disclosure may contain probes or primers that distinguish between both Factor H and CFHR5 variants, including any combination of the sites described above and elsewhere in this disclosure.

[0064] The devices described above and their contents may also be used to identify a propensity to develop MPGNII or to determine a Factor H haplotype for any purpose.

[0065] In one embodiment the device contains, instead of, or in addition to, the probes or primers described above, probes, primers, antibodies and the like that distinguish polymorphic sites in the CFHR5 gene. The disclosure provides devices for the diagnosis of AMD or MPGNII in a patient based on variation in the CFHR5 gene. The devices may include one or more CFHR5-specific probes or CFHR5 allele-specific oligonucleotides, or antibodies that specifically recognize the CFHR5 polypeptide. The CFHR5-specific primers and CFHR5 allele-specific oligonucleotides may include sequences derived from the coding (exons) or non-coding (promoter, 5' untranslated, introns or 3' untranslated) region of the CFHR5 gene. The CFHR5-specific antibodies may recognize the normal or wild-type CFHR5 polypeptide or variant CFHR5 polypeptides in which one or

more non-synonymous single nucleotide polymorphisms (SNPs) are present in the CFHR5 coding region. The devices may be used to diagnose AMD or MPGNII, as well as other diseases associated with SNPs in the CFHR5 gene.

[0066] In one embodiment the device contains probes or primers that distinguish alleles at a polymorphic site listed in TABLE 14 or TABLE 15. In an embodiment the probes are primers for nucleic acid amplification of a region spanning a CFHR5 gene polymorphic site listed in TABLE 14 or TABLE 15. In an embodiment the device has probes or primers that distinguish alleles at more than one polymorphic site listed in TABLE 14 or TABLE 15. In an embodiment the kit comprises probes or primers that distinguish alleles one, two or all of the following polymorphic sites: rs9427661 (-249T>C); rs9427662 (-20T>C); and rs12097550 (P46S).

[0067] In one embodiment the device contains probes or primers that distinguish alleles at a polymorphic site in the CFH gene and at a polymorphic site in a CFHR gene, such as CFHR5.

[0068] Additional embodiments of the invention will be apparent upon reading the entire disclosure.

BRIEF DESCRIPTION OF THE FIGURES

[0069] FIGURES 1A-1L show the immunolocalization of Factor H (Figs. 1A-1H) and the terminal complement complex (C5b-9) (Figs. 1I-1L) in the human retinal pigmented epithelium. Abbreviations: (RPE)-choroid (Chor) complex; Bruch's membrane (BM); Retina (Ret); Drusen (Dr).

[0070] FIGURE 2 shows RT-PCR analysis of Factor H gene expression (CFH and the truncated form HFL1) using RNA extracted from the human eye.

[0071] FIGURE 3 is a diagram of the human Factor H gene showing the approximate locations of 12 SNPs used in the analysis, the 22 exons of the Factor H gene, the 20 short consensus repeats (SCRs), the binding sites for pathogens and other substrates, and the linkage disequilibrium (LD) blocks. The diagram, showing all 22 exons of CFH (but not introns) is not drawn to scale.

[0072] FIGURE 4 is a haplotype network diagram of human Factor H gene SNPs showing the relationship between the risk (filled-in circles), protective (lined circles), neutral (open circles) and ancestral (indicated) haplotypes and the relative frequency of the haplotypes, as indicated by the sizes and positions of the circles.

[0073] FIGURE 5 shows an association analysis of human Factor H gene haplotypes and diplotypes. Eight informative SNPs were analyzed for pairwise linkage disequilibrium in AMD cases and controls. The nucleotide on the coding strand at the indicated polymorphic sites is shown, except for IVS1, where the nucleotide on the non-coding strand is shown.

[0074] FIGURE 6 shows the 3926 base nucleotide sequence of the reference form of human Factor H cDNA (GenBank accession number Y00716 [SEQ ID NO:1]). The ATG initiation codon begins at nucleotide position 74 and the TAG termination codon ends at nucleotide position 3769.

[0075] FIGURE 7 shows the polypeptide sequence encoded by SEQ ID NO:1 (GenBank accession number Y00716 [SEQ ID NO:2]). The 1231 amino acid Factor H polypeptide includes an 18 amino acid N-terminal signal peptide.

[0076] FIGURE 8 shows the 1658 base nucleotide sequence of the reference form of HFL1, the truncated form of the human Factor H (GenBank accession number X07523 [SEQ ID NO:3]). The ATG initiation codon begins at nucleotide position 74 and the TGA termination codon ends at nucleotide position 1423.

[0077] FIGURE 9 shows the polypeptide sequence of the reference form of HFL1 encoded by SEQ ID NO:3 (GenBank accession number X07523 [SEQ ID NO:4]). The 449 amino acid HFL1 polypeptide includes an 18 amino acid N-terminal signal peptide.

[0078] FIGURE 10 shows the polypeptide sequence of an exemplary protective variant of human Factor H [SEQ ID NO:5]. This protective variant Factor H polypeptide has a isoleucine at amino acid position 62 and a tyrosine at amino acid position 402 (indicated in bold).

[0079] FIGURE 11 shows the polypeptide sequence of an exemplary protective variant of HFL1, the truncated form of human Factor H (SEQ ID NO:6). This protective variant truncated Factor H polypeptide has a isoleucine at amino acid position 62 and tyrosine at amino acid position 402 (indicated in bold).

[0080] FIGURE 12 shows marked glomerular hypercellularity with dense intramembranous deposits that cause capillary wall thickening in a patient with MPGNII, as viewed by (A) light microscopy and (B) electron microscopy. The deposits can form a segmental, discontinuous or diffuse pattern in the lamina densa of the glomerular basement membrane (GBM). By light microscopy, they are eosinophilic and refractile, stain brightly with periodic acid-Schiff and are highly osmophilic, which explains their electron-dense appearance (A). Even by electron microscopy the deposits lack substructure and appear as

very dark homogeneous smudges (B). The exact composition of dense deposits remains unknown (bar, 5 μ m).

[0081] FIGURE 13 is a diagram showing the activation and regulation of the alternative pathway of the complement cascade, which is systematically activated at a high level in patients with AMD and MPGNII. The alternative pathway of the complement cascade is systematically activated at a high level in patients with MPGN II/DDD. Normally, continuous low-level activation of C3 occurs by a process of spontaneous hydrolysis known as tick-over. C3 hydrolysis is associated with a large conformational protein change shown at the top of the diagram. The conformational change makes C3(H₂O) similar to C3b, a C3 cleavage product. The initial convertase, C3(H₂O)Bb, activates C3 to form C3b. Although C3b has a fleeting half-life, if it binds to IgG, cells or basement membranes, it is protected from immediate inactivation. (C3b)₂-IgG complexes form in the fluid phase and bind properdin (P), which facilitates factor B binding and the generation of C3bBb, the convertase of the alternative pathway, shown here as a Bb(C3b)₂-IgG-properdin complex. The amplification loop is depicted by the arrows. C3NeF prolongs the half-life of C3 convertase and is shown in the inset. One mechanism to degrade C3 convertase is through its interaction with complement Factor H (CFH), shown at the bottom right as fH. Deficiency of and mutations in Factor H are associated with MPGN II/DDD.

[0082] FIGURE 14 is a diagram showing the organization of the regulators-of-complement-activation (RCA) gene cluster on chromosome 1q32 and the arrangement of approximately 60-amino acid domains known as short consensus repeats (SCRs) in complement Factor H (CFH), Factor H-Like 1 (CFHL1) and Factor H-Related 1, 2, 3, 4 and 5 (CFHR1, CFHR2, CFHR3, CFHR4 and CFHR5). CFH has 20 SCRs. The interacting partners with some of these SCRs has been determined and is shown on the top right (CRP, C reactive protein; Hep, heparin). Complement factor H-like 1 (CFHL1) is a splice isoform of CFH, while complement factor H-related proteins 1-5 (CFHR1-5) are each encoded by a unique gene (CFHR1-5). The SCRs of CFHR1-5 are similar to some of the SCRs in CFH, as denoted by the numbers in the ovals. For example, CFHR5 has 9 SCRs, with the first two being similar to SCRs 6 and 7 of Factor H and therefore having CRP and heparin binding properties. SCRs5-7 of CFHR5 have the numbers 12-14 within the corresponding ovals because these SCRs are similar to SCRs 12-14 of Factor H and have C3b and heparin binding properties.

[0083] FIGURE 15 shows a linkage disequilibrium plot indicating that A307A and Y402H are in linkage disequilibrium in Factor H and -249T>C and -20T>C are in linkage disequilibrium in *CFHR5*.

[0084] FIGURE 16 shows the 2821 base nucleotide sequence of the reference form of human *CFHR5* (GenBank accession number AF295327 [SEQ ID NO:7]. The ATG initiation codon begins at nucleotide position 94 and the TGA termination codon ends at nucleotide position 1803.

[0085] FIGURE 17 shows the polypeptide sequence encoded by SEQ ID NO:7 (GenBank accession number AAK15619 [SEQ ID NO:8]. The 569 amino acid *CFHR5* polypeptide includes an 18 amino acid N-terminal signal peptide.

[0086] FIGURE 18 shows genomic duplications in the genes for CFH and the Factor H-related proteins. Exons are indicated as vertical lines. Regions labeled with the same letter (e.g., A, A', and A'') have substantially identical sequences.

DETAILED DESCRIPTION OF THE INVENTION

I. INTRODUCTION

[0087] The invention provides a collection of polymorphisms and haplotypes comprised of multiple variations in the Factor H gene, and in Factor H-related genes such as Factor H-Related 5 gene. These polymorphisms and haplotypes are associated with age related macular degeneration (AMD) and other Factor H-related conditions. Certain of these polymorphisms and haplotypes result in variant Factor H polypeptides. Detection of these and other polymorphisms and sets of polymorphisms (e.g., haplotypes) is useful in designing and performing diagnostic assays for AMD. Polymorphisms and sets of polymorphisms can be detected by analysis of nucleic acids, by analysis of polypeptides encoded by Factor H coding sequences (including polypeptides encoded by splice variants), or by other means known in the art. Analysis of such polymorphisms and haplotypes is also useful in designing prophylactic and therapeutic regimes for AMD.

[0088] Factor H is a multifunctional protein that functions as a key regulator of the complement system. See Zipfel, 2001, "Factor H and disease: a complement regulator affects vital body functions" *Semin Thromb Hemost.* 27:191-9. The Factor H protein activities include: (1) binding to C-reactive protein (CRP), (2) binding to C3b, (3) binding to heparin, (4) binding to sialic acid; (5) binding to endothelial cell surfaces, (6) binding to cellular

integrin receptors (7) binding to pathogens, including microbes (see FIGURE 3), and (8) C3b co-factor activity. The Factor H gene, known as HFI, CFH and HF, is located on human chromosome 1, at position 1q32. The 1q32 particular locus contains a number of complement pathway-associated genes. One group of these genes, referred to as the regulators of complement activation (RCA) gene cluster, contains the genes that encode Factor H, five Factor H-related genes (FHR-1, FHR-2, FHR-3, FHR-4 and FHR-5 or *CFHR1*, *CFHR2*, *CFHR3*, *CFHR4* and *CFHR5*, respectively), and the gene encoding the beta subunit of coagulation factor XIII. The Factor H and Factor H related genes is composed almost entirely of short consensus repeats (SCRs). Factor H and FHL1 are composed of SCRs 1-20 and 1-7, respectively. FHR-1, FHR-2, FHR-3, FHR-4 and FHR-5 are composed of 5, 4, 5, 5 and 8 SCRs, respectively (see FIGURE 14). The order of genes, from centromere to telomere is FH/FHL1, FHR-3, FHR-1, FHR-4, FHR-2 and FHR-5.

Factor H Gene

[0089] The reference form of human Factor H cDNA (SEQ ID NO:1) (see Ripoche et al., 1988, *Biochem J* 249:593-602) and genomic sequences have been determined. The Factor H cDNA encodes a polypeptide 1231 amino acids in length (SEQ ID NO:2) having an apparent molecular weight of 155 kDa. There is an alternatively spliced form of Factor H is known as FHL-1 (and also has been referred to as HFL1 or CFHT). FHL-1 (SEQ ID NO:3) corresponds essentially to exons 1 through 9 of Factor H (see Ripoche et al., 1988, *Biochem J* 249:593-602). The FHL1 cDNA encodes a polypeptide 449 amino acids in length (SEQ ID NO:4) having an apparent molecular weight of 45-50 kDa. The first 445 amino acids of FH1 and FHL1 are identical, with FHL1 having a unique C-terminal 4 amino acids (exon 10A). The alternative exon 10A is located in the intron between exon 9 and exon 10. cDNA and amino acid sequence data for human Factor H and FHL1 are found in the EMBL/GenBank Data Libraries under accession numbers Y00716 and X07523, respectively. The 3926 base nucleotide sequence of the reference form of human Factor H cDNA (GenBank accession number Y00716 [SEQ ID NO:1]) is shown in FIGURE 6, and the polypeptide sequence encoded by SEQ ID NO:1 (GenBank accession number Y00716 [SEQ ID NO:2]) is shown in FIGURE 7. The 1658 base nucleotide sequence of the reference form of HFL1, the truncated form of the human Factor H (GenBank accession number X07523 [SEQ ID NO:3]) is shown in FIGURE 8, and the polypeptide sequence encoded by SEQ ID NO:3 (GenBank accession number X07523 [SEQ ID NO:4]) is shown in FIGURE 9. The Factor H gene

sequence (150626 bases in length) is found under GenBank accession number AL049744. The Factor H promoter is located 5' to the coding region of the Factor H gene.

FHR-1 Gene

[0090] The FHR-1 gene is also known as *CHFR1*, *CFHL1*, *CFHL*, *FHR1* and *HFL1*. The reference form of human HFR-1 cDNA (see Estaller et al., 1991, *J. Immunol.* 146:3190-3196) and genomic sequences have been determined. The FHR-1 cDNA encodes a polypeptide 330 amino acids in length having an predicted molecular weight of 39 kDa. cDNA and amino acid sequence data for human FHR-1 are found in the EMBL/GenBank Data Libraries under accession number M65292. The FHR-1 gene sequence is found under GenBank accession number AL049741.

FHR-2 Gene

[0091] The FHR-2 gene is also known as *CHFR2*, *CFHL2*, *FHR2* and *HFL3*. The reference form of human HFR-2 cDNA (see Strausberg et al., *Proc. Natl. Acad. Sci USA* 99:16899-16903) and genomic sequences have been determined. The FHR-2 cDNA encodes a polypeptide 270 amino acids in length having a predicted molecular weight of 31 kDa. cDNA and amino acid sequence data for human FHR-2 are found in the EMBL/GenBank Data Libraries under accession number BC022283. The FHR-2 gene sequence is found under GenBank accession number AL139418.

FHR-3 Gene

[0092] The FHR-3 gene is also known as *CFHR3*, *CFHL3*, *FHR3* and *HLF4*. The reference form of human HFR-3 cDNA (see Strausberg et al., *Proc. Natl. Acad. Sci USA* 99:16899-16903) and genomic sequences have been determined. The FHR-3 cDNA encodes a polypeptide 330 amino acids in length having a predicted molecular weight of 38 kDa. cDNA and amino acid sequence data for human FHR-3 are found in the EMBL/GenBank Data Libraries under accession number BC058009. The FHR-3 gene sequence is found under GenBank accession number AL049741.

FHR-4 Gene

[0093] The FHR-4 gene is also known as *CFHR4*, *CFHL4* and *FHR4*. The reference form of human HFR-4 cDNA (see Skerka et al., 1991, *J. Biol. Chem.* 272:5627-5634) and genomic

sequences have been determined. The FHR-4 cDNA encodes a polypeptide 331 amino acids in length having a predicted molecular weight of 38 kDa. cDNA and amino acid sequence data for human FHR-4 are found in the EMBL/GenBank Data Libraries under accession number X98337. The FHR-4 gene sequence is found under GenBank accession numbers AF190816 (5' end), AL139418 (3' end) and BX248415.

FHR-5 Gene

[0094] The FHR-5 gene is also known as *CFHR5*, *CFHL5* and *FHR5*. The reference form of human *CFHR5* cDNA (SEQ ID NO:83) (see McRae et al., 2001, *J. Biol.Chem.* 276:6747-6754) and genomic sequences have been determined. The *CFHR5* cDNA encodes a polypeptide 569 amino acids in length (SEQ ID NO:8) having an apparent molecular weight of 65 kDa. cDNA and amino acid sequence data for human *CFHR5* are found in the EMBL/GenBank Data Libraries under accession number AF295327. The 2821 base nucleotide sequence of the reference form of human *CFHR5* (GenBank accession number AF295327 [SEQ ID NO:7]) is shown in FIGURE 16, and the polypeptide sequence encoded by SEQ ID NO:7 (GenBank accession number AAK15619 [SEQ ID NO:8]) is shown in FIGURE 17. The *CFHR5* gene sequence is found under GenBank accession numbers AL139418 (5' end) and AL353809 (3' end). The FHR-5 promoter is located 5' to the coding region of the *CFHR5* gene.

II. DEFINITIONS

[0095] The following definitions are provided to aid in understanding the invention. Unless otherwise defined, all terms of art, notations and other scientific or medical terms or terminology used herein are intended to have the meanings commonly understood by those of skill in the arts of medicine and molecular biology. In some cases, terms with commonly understood meanings are defined herein for clarity and/or for ready reference, and the inclusion of such definitions herein should not be assumed to represent a substantial difference over what is generally understood in the art.

[0096] A "nucleic acid", "polynucleotide" or "oligonucleotide" is a polymeric form of nucleotides of any length, may be DNA or RNA, and may be single- or double-stranded. Nucleic acids may include promoters or other regulatory sequences. Oligonucleotides are usually prepared by synthetic means. Nucleic acids include segments of DNA, or their complements spanning or flanking any one of the polymorphic sites shown in TABLE 1A,

TABLE 1B and/or TABLE 1C or otherwise known in the Factor H gene. The segments are usually between 5 and 100 contiguous bases, and often range from a lower limit of 5, 10, 12, 15, 20, or 25 nucleotides to an upper limit of 10, 15, 20, 25, 30, 50 or 100 nucleotides (where the upper limit is greater than the lower limit). Nucleic acids between 5-10, 5-20, 10-20, 12-30, 15-30, 10-50, 20-50 or 20-100 bases are common. The polymorphic site can occur within any position of the segment. A reference to the sequence of one strand of a double-stranded nucleic acid defines the complementary sequence and except where otherwise clear from context, a reference to one strand of a nucleic acid also refers to its complement. For certain applications, nucleic acid (e.g., RNA) molecules may be modified to increase intracellular stability and half-life. Possible modifications include, but are not limited to, the use of phosphorothioate or 2'-O-methyl rather than phosphodiesterase linkages within the backbone of the molecule. Modified nucleic acids include peptide nucleic acids (PNAs) and nucleic acids with nontraditional bases such as inosine, queosine and wybutosine and acetyl-, methyl-, thio- and similarly modified forms of adenine, cytidine, guanine, thymine, and uridine which are not as easily recognized by endogenous endonucleases.

[0097] "Hybridization probes" are nucleic acids capable of binding in a base-specific manner to a complementary strand of nucleic acid. Such probes include nucleic acids and peptide nucleic acids (Nielsen et al., 1991). Hybridization may be performed under stringent conditions which are known in the art. For example, see, e.g., Berger and Kimmel (1987) *Methods In Enzymology*, Vol. 152: *Guide To Molecular Cloning Techniques*, San Diego: Academic Press, Inc.; Sambrook et al. (1989) *Molecular Cloning: A Laboratory Manual*, 2nd Ed., Vols. 1-3, Cold Spring Harbor Laboratory; Sambrook (2001) 3rd Edition; Rychlik, W. and Rhoads, R.E., 1989, *Nucl. Acids Res.* 17, 8543; Mueller, P.R. et al. (1993) In: *Current Protocols in Molecular Biology* 15.5, Greene Publishing Associates, Inc. and John Wiley and Sons, New York; and Anderson and Young, *Quantitative Filter Hybridization in Nucleic Acid Hybridization* (1985)). As used herein, the term "probe" includes primers. Probes and primers are sometimes referred to as "oligonucleotides."

[0098] The term "primer" refers to a single-stranded oligonucleotide capable of acting as a point of initiation of template-directed DNA synthesis under appropriate conditions, in an appropriate buffer and at a suitable temperature. The appropriate length of a primer depends on the intended use of the primer but typically ranges from 15 to 30 nucleotides. A primer sequence need not be exactly complementary to a template but must be sufficiently complementary to hybridize with a template. The term "primer site" refers to the area of the

target DNA to which a primer hybridizes. The term "primer pair" means a set of primers including a 5' upstream primer, which hybridizes to the 5' end of the DNA sequence to be amplified and a 3' downstream primer, which hybridizes to the complement of the 3' end of the sequence to be amplified.

[0099] Exemplary hybridization conditions for short probes and primers is about 5 to 12 degrees C below the calculated T_m . Formulas for calculating T_m are known and include: $T_m = 4^{\circ}\text{C} \times (\text{number of G's and C's in the primer}) + 2^{\circ}\text{C} \times (\text{number of A's and T's in the primer})$ for oligos <14 bases and assumes a reaction is carried out in the presence of 50mM monovalent cations. For longer oligos, the following formula can be used: $T_m = 64.9^{\circ}\text{C} + 41^{\circ}\text{C} \times (\text{number of G's and C's in the primer} - 16.4)/N$, where N is the length of the primer. Another commonly used formula takes into account the salt concentration of the reaction (Rychlik, supra, Sambrook, supra, Mueller, supra.): $T_m = 81.5^{\circ}\text{C} + 16.6^{\circ}\text{C} \times (\log_{10}[\text{Na}^+] + [\text{K}^+]) + 0.41^{\circ}\text{C} \times (\% \text{GC}) - 675/N$, where N is the number of nucleotides in the oligo. The aforementioned formulae provide a starting point for certain applications; however, the design of particular probes and primers may take into account additional or different factors. Methods for design of probes and primers for use in the methods of the invention are well known in the art.

[0100] The terms "risk," "protective," and "neutral" are used to describe variations, SNPS, haplotypes, diplotypes, and proteins in a population encoded by genes characterized by such patterns of variations. A risk haplotype is an allelic form of a gene, herein Factor H or a Factor H-related gene, comprising at least one variant polymorphism, and preferably a set of variant polymorphisms, associated with increased risk for developing AMD. The term "variant" when used in reference to a Factor H or Factor H-related gene, refers to a nucleotide sequence in which the sequence differs from the sequence most prevalent in a population, herein humans of European-American descent. The variant polymorphisms can be in the coding or non-coding portions of the gene. An example of a risk Factor H haplotype is the allele of the Factor H gene encoding histidine at amino acid 402 and/or cysteine at amino acid 1210. The risk haplotype can be naturally occurring or can be synthesized by recombinant techniques. A protective haplotype is an allelic form of a gene, herein Factor H or a Factor H-related gene, comprising at least one variant polymorphism, and preferably a set of variant polymorphisms, associated with decreased risk of developing AMD. For example, one protective Factor H haplotype has an allele of the Factor H gene encoding isoleucine at amino acid 62. The protective haplotype can be naturally occurring or synthesized by recombinant

techniques. A neutral haplotype is an allelic form of a gene, herein Factor H or a Factor H-related gene, that does not contain a variant polymorphism associated in a population or ethnic group with either increased or decreased risk of developing AMD. It will be clear from the following discussion that a protein encoded in a "neutral" haplotype may be protective when administered to a patient in need of treatment or prophylaxis for AMD or other conditions. That is, both "neutral" and "protective" forms of CFH or CFHR5 can provide therapeutic benefit when administered to, for example, a subject with AMD or risk for developing AMD, and thus can "protect" the subject from disease.

[0101] The term "wild-type" refers to a nucleic acid or polypeptide in which the sequence is a form prevalent in a population, herein humans of European-American descent (approximately 40% prevalence; see FIGURE 5). For purposes of this disclosure, a "wild-type" Factor H protein has the sequence of SEQ ID NO:2 (FIGURE 7), except that the amino acid at position 402 is tyrosine (Y; [SEQ ID NO:337]). For purposes of this disclosure, a Factor H gene encoding a wild-type Factor H protein has the sequence of SEQ ID NO:1 (FIGURE 6), except that the codon beginning at base 1277, corresponding to the amino acid at position 402 encodes tyrosine (TAT [SEQ ID NO:336]).

[0102] The term "variant" when used in reference to a Factor H or Factor H-related polypeptide, refers to a polypeptide in which the sequence differs from the normal or wild-type sequence at a position that changes the amino acid sequence of the encoded polypeptide. For example, some variations or substitutions in the nucleotide sequence of Factor H gene alter a codon so that a different amino acid is encoded (for example and not for limitation, having an alternative allele at one or more of I62V, Y402H, D936E) resulting in a variant polypeptide. Variant polypeptides can be associated with risk (e.g., having histidine at position 402), associated with protection (e.g., having isoleucine at position 62), or can be encoded by a neutral haplotype (e.g., having aspartic acid at position 936). Variant CFHR5 polypeptides can be associated with risk (e.g., having serine at position 46), associated with protection, or can be neutral.

[0103] The term "reference" when referring to a Factor H polypeptide means a polypeptide in which the amino acid sequence is identical to the sequence described by Ripoché et al., 1988, *Biochem J.* 249:593-602) for full-length (FH1, SEQ ID NO:2) or truncated (FHL1, SEQ ID NO:4) human Factor H. The term "reference" when referring to a CFHR5 polypeptide means a polypeptide in which the amino acid sequence is identical to the sequence described by McRae et al., 2001, *J. Biol.Chem.* 276:6747-6754) for full-length

human CFHR5 (SEQ ID NO:8). The first identified allelic form is arbitrarily designated the reference form or allele; other allelic forms are designated as alternative or variant alleles.

Wild-type and variant forms may have substantial sequence identity with the reference form (e.g., the wild-type or variant form may be identical to the reference form at at least 90% of the amino acid positions of the wild-type or variant, sometimes at least 95% of the positions and sometimes at least 98% or 99% of the positions). A variant may differ from a reference form in certain regions of the protein due to a frameshift mutation or splice variation.

[0104] The term "polymorphism" refers to the occurrence of two or more genetically determined alternative sequences or alleles in a population. A "polymorphic site" is the locus at which sequence divergence occurs. Polymorphic sites have at least two alleles. A diallelic polymorphism has two alleles. A triallelic polymorphism has three alleles. Diploid organisms may be homozygous or heterozygous for allelic forms. A polymorphic site may be as small as one base pair. Examples of polymorphic sites include: restriction fragment length polymorphisms (RFLPs); variable number of tandem repeats (VNTRs); hypervariable regions; minisatellites; dinucleotide repeats; trinucleotide repeats; tetranucleotide repeats; and simple sequence repeats. As used herein, reference to a "polymorphism" can encompass a set of polymorphisms (i.e., a haplotype).

[0105] A "single nucleotide polymorphism (SNP)" occurs at a polymorphic site occupied by a single nucleotide, which is the site of variation between allelic sequences. The site is usually preceded by and followed by highly conserved sequences of the allele. A SNP usually arises due to substitution of one nucleotide for another at the polymorphic site. Replacement of one purine by another purine or one pyrimidine by another pyrimidine is called a transition. Replacement of a purine by a pyrimidine or vice versa is called a transversion. A synonymous SNP refers to a substitution of one nucleotide for another in the coding region that does not change the amino acid sequence of the encoded polypeptide. A non-synonymous SNP refers to a substitution of one nucleotide for another in the coding region that changes the amino acid sequence of the encoded polypeptide. A SNP may also arise from a deletion or an insertion of a nucleotide or nucleotides relative to a reference allele.

[0106] A "set" of polymorphisms means more than one polymorphism, e.g., at least 2, at least 3, at least 4, at least 5, at least 6, or more than 6 of the polymorphisms shown in TABLE 1A, TABLE 1B and/or TABLE 1C or otherwise known in the Factor H gene or other gene.

[0107] The term "haplotype" refers to the designation of a set of polymorphisms or alleles of polymorphic sites within a gene of an individual. For example, a "112" Factor H haplotype refers to the Factor H gene comprising allele 1 at each of the first two polymorphic sites and allele 2 at the third polymorphic site. A "diplotype" is a haplotype pair.

[0108] An "isolated" nucleic acid means a nucleic acid species that is the predominant species present in a composition. Isolated means the nucleic acid is separated from at least one compound with which it is associated in nature. A purified nucleic acid comprises (on a molar basis) at least about 50, 80 or 90 percent of all macromolecular species present.

[0109] Two amino acid sequences are considered to have "substantial identity" when they are at least about 80% identical, preferably at least about 90% identical, more preferably at least about 95%, at least about 98% identical or at least about 99% identical. Percentage sequence identity is typically calculated by determining the optimal alignment between two sequences and comparing the two sequences. Optimal alignment of sequences may be conducted by inspection, or using the local homology algorithm of Smith and Waterman, 1981, *Adv. Appl. Math.* 2: 482, using the homology alignment algorithm of Needleman and Wunsch, 1970, *J. Mol. Biol.* 48: 443, using the search for similarity method of Pearson and Lipman, 1988, *Proc. Natl. Acad. Sci. U.S.A.* 85: 2444, by computerized implementations of these algorithms (e.g., in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, Wis.) using default parameters for amino acid comparisons (e.g., for gap-scoring, etc.). It is sometimes desirable to describe sequence identity between two sequences in reference to a particular length or region (e.g., two sequences may be described as having at least 95% identity over a length of at least 500 basepairs). Usually the length will be at least about 50, 100, 200, 300, 400, 500, 600, 700, 800, 900 or 1000 amino acids, or the full length of the reference protein. Two amino acid sequences can also be considered to have substantial identity if they differ by 1, 2, or 3 residues, or by from 2-20 residues, 2-10 residues, 3-20 residues, or 3-10 residues.

[0110] "Linkage" describes the tendency of genes, alleles, loci or genetic markers to be inherited together as a result of their location on the same chromosome. Linkage can be measured by percent recombination between the two genes, alleles, loci or genetic markers. Typically, loci occurring within a 50 centimorgan (cM) distance of each other are linked. Linked markers may occur within the same gene or gene cluster. "Linkage disequilibrium" or "allelic association" means the preferential association of a particular allele or genetic marker with a specific allele or genetic marker at a nearby chromosomal location more frequently

than expected by chance for any particular allele frequency in the population. A marker in linkage disequilibrium can be particularly useful in detecting susceptibility to disease, even if the marker itself does not cause the disease.

[0111] The terms "diagnose" and "diagnosis" refer to the ability to determine whether an individual has the propensity to develop disease (including with or without signs or symptoms). Diagnosis of propensity to develop disease can also be called "screening" and, as used herein, the terms diagnosis and screening are used interchangeably. It will be appreciated that having an increased or decreased propensity to developing a condition refers to the likelihood of developing the condition relative to individuals in the population without the condition.

III. TABLES

[0112] Certain tables referred to herein are provided following the Examples, *infra*. The following descriptions are provided to assist the reader:

[0113] TABLES 1A-1C show human Factor H gene polymorphisms and their association with age-related macular degeneration. (1A) The dbSNP no., location, sequences of the coding (top, 5' to 3' direction) and non-coding (bottom) strands spanning the polymorphisms, amino acid changes, allele frequencies for the control and AMD cases, and χ^2 and P-values for 1 SNPs in the human Factor H gene are shown. (1B) The dbSNP no., interrogated sequences, corresponding nucleotide in the chimp Factor H gene, location, amino acid changes, and sets of primers and probes for 11 SNPs in the human Factor H gene are shown. (1C) The location, sequences spanning the polymorphisms, amino acid position and amino acid change, if any, for 14 SNPs in the human Factor H gene that are not found in the dbSNP database are shown.

[0114] TABLE 2 shows a haplotype analysis of eight SNPs in the human Factor H gene in a cohort of AMD cases and controls.

[0115] TABLE 3 shows a haplotype analysis of six SNPs in the human Factor H gene and their association with AMD.

[0116] TABLE 4 shows the association of 11 human Factor H gene SNPs with age-related macular degeneration.

[0117] TABLE 5 shows the primers used for SSCP, DHPLC and DNA sequencing analysis for human Factor H.

[0118] TABLE 6 shows genotyping data of AMD patients and controls.

- [0119] TABLE 7 shows the frequency of an at-risk haplotype in various ethnic groups.
- [0120] TABLE 8 shows several Factor H diplotypes. Common risk and protective diplotypes are indicated.
- [0121] TABLE 9 shows the sequences of primers used to amplify the Factor H coding sequence.
- [0122] TABLE 10 shows the sequences of primers used to amplify the CFHR5 coding sequence.
- [0123] TABLE 11 shows an analysis of Factor H SNPs in 22 MPGNII patients.
- [0124] TABLE 12 shows a comparison of Factor H SNP frequencies in 22 MPGNII patients and AMD-negative, ethnically matched controls.
- [0125] TABLE 13 lists Factor H SNPs associated with MPGNII and their related SCR.
- [0126] TABLE 14 shows an analysis of *CFHR5* SNPs in 22 MPGNII patients.
- [0127] TABLE 15 shows a comparison of *CFHR5* SNP frequencies in 22 MPGNII patients and AMD-negative, ethnically matched controls.
- [0128] TABLE 16 shows exemplary allele-specific probes (16A) and primers (16B) useful for detecting polymorphisms in the Factor H gene.

IV. COMPLEMENT FACTOR H POLYMORPHISMS

- [0129] In one aspect, the invention provides new diagnostic, treatment and drug screening methods related to the discovery that polymorphic sites in the Complement Factor H (HF1) gene are associated with susceptibility to and development of AMD.
- [0130] Factor H polymorphisms associated with AMD were identified as described in Example 1, by examining the coding and adjacent intronic regions of Factor H (including exon 10A, which is transcribed for the Factor H isoform FHL1) for variants using SSCP analysis, DHPLC analysis, and direct sequencing, according to standard protocols. Remaining polymorphisms were typed by the 5' nuclease (Taqman, ABI) methodology. Taqman genotyping and association analysis were performed as described (Gold et al., 2004). Primers for SSCP and DNA sequencing analyses were designed to amplify each exon and its adjacent intronic regions using MacVector software. PCR-derived amplicons were screened for sequence variation by SSCP and DHPLC according to standard protocols. All changes detected by SSCP and DHPLC were confirmed by bidirectional sequencing according to standard protocols. Statistical analyses were performed using chi-square (χ^2) and Fisher's exact tests (P values).

[0131] Two independent groups of AMD cases and age-matched controls were used. All participating individuals were of European-American descent, over the age of 60 and enrolled under IRB-approved protocols following informed consent. These groups were comprised of 352 unrelated patients with clinically documented AMD (mean age 79.5 ± 7.8 years) and 113 unrelated, control patients (mean age 78.4 ± 7.4 years; matched by age and ethnicity) from the University of Iowa, and 550 unrelated patients with clinically documented AMD (mean age 71.32 ± 8.9 years) and 275 unrelated, matched by age and ethnicity, controls (mean age 68.84 ± 8.6 years) from Columbia University. Patients were examined by indirect ophthalmoscopy and slit-lamp microscopy by retina fellowship-trained ophthalmologists.

[0132] Fundus photographs were graded according to a standardized, international classification system (Bird et al., 1995). Control patients were selected and included if they did not exhibit any distinguishing signs of macular disease or have a known family history of AMD. The AMD patients were subdivided into phenotypic categories: early AMD (ARM), geographic atrophy (GA), and exudative (CNV) AMD, based upon the classification of their most severe eye at the time of their entry into the study. The University of Iowa ARM and GA cases were further subdivided into distinct phenotypes (RPE changes alone, >10 macular hard drusen, macular soft drusen, BB (cuticular) drusen, PED, "Cherokee" atrophy, peninsular geographic atrophy and pattern geographic atrophy). The earliest documentable phenotype for all cases was also recorded and employed in the analyses.

[0133] As shown in TABLE 1A, a highly significant association of polymorphic sites in the Factor H gene with AMD was found in an examination of two independent cohorts that together included approximately 900 AMD cases and 400 matched controls. Sixteen (16) polymorphisms in the Factor H gene are listed in TABLES 1A-1B. Of these twelve (12) are found in the SNP database (dbSNP) which may be found in the National Center for Biotechnology Information (NCBI). The dbSNP is a collection of SNPs in the human Factor H gene which are dispersed among the 22 coding exons of the Factor H gene and among the promoter; the 5' untranslated region, the introns, and the 3' untranslated region of the Factor H gene. Listed below are the accession numbers for 379 SNPs in the human Factor H gene that are found in the dbSNP database. These SNPs can be used in carrying out methods of the invention.

TABLE A

rs17575212	rs11582939	rs7551203	rs5014736	rs2019724	rs534479	rs395963
rs17573867	rs11580821	rs7546015	rs5014735	rs1984894	rs534399	rs395544
rs16840522	rs11579439	rs7540032	rs5014734	rs1928433	rs529825	rs395129

rs16840465	rs11539862	rs7539005	rs5014733	rs1928432	rs528298	rs393955
rs16840462	rs11398897	rs7537967	rs5003626	rs1887973	rs521605	rs386258
rs16840422	rs11390840	rs7535653	rs5003625	rs1831282	rs520992	rs385892
rs16840419	rs11340441	rs7535263	rs5003624	rs1831281	rs519839	rs385543
rs16840410	rs11339120	rs7529589	rs5002880	rs1831280r	rs518957	rs383191
rs16840401	rs11318544	rs7526622	rs5002876	rs1410997	rs551397	rs405306
rs16840397	rs11285593	rs7524776	rs5002875	rs1410996	rs544889	rs403846
rs16840394	rs10922109	rs7522681	rs5002874	rs1329429	rs543879	rs402991
rs16840381	rs10922108	rs7519439	rs4658046	rs1329428	rs536564	rs402056
rs16840379	rs10922107	rs7514261	rs4657826	rs1329427	rs536539	rs399469
rs12756364	rs10922106	rs7513157	rs4350148	rs1329424	rs515299	rs398248
rs12746361	rs10922105	rs7415913	rs4044888	rs1329423	rs514756	rs381974
rs12740961	rs10922104	rs7413999	rs4044884	rs1329422	rs514591	rs380390
rs12726401	rs10922103	rs7413137	rs4044882	rs1329421	rs513699	rs380060
rs12566629	rs10922102	rs6695321	rs3834020	rs1299282	rs512900	rs379489
rs12565418	rs10922101	rs6691749	rs3766405	rs1292487	rs508505	rs375046
rs12406047	rs10922100	rs6690982	rs3766404	rs1292477	rs499807	rs374896
rs12405238	rs10922099	rs6689826	rs3766403	rs1292476	rs495968	rs374231
rs12402808	rs10922098	rs6689009	rs3753397	rs1292475	rs495222	rs371647
rs12238983	rs10922097	rs6688272	rs3753396	rs1292474	rs493367	rs368465
rs12144939	rs10922096	rs6685249	rs3753395	rs1292473	rs491480	rs364947
rs12136675	rs10922095	rs6682138	rs3753394	rs1292472	rs490864	rs203688
rs12134975	rs10922094	rs6680396	rs3043115	rs1292471	rs488738	rs203687
rs12134598	rs10922093	rs6677604	rs3043113	rs1292466	rs487114	rs203686
rs12127759	rs10922092	rs6677460	rs3043112	rs1156679	rs482934	rs203685
rs12124794	rs10801561	rs6677089	rs3043111	rs1156678	rs480266	rs203684
rs12116702	rs10801560	rs6675088	rs2878649	rs1089031	rs466287	rs203683r
rs12096637	rs10801559	rs6674960	rs2878648	rs1065489	rs464798	s203682
rs12085209	rs10801558	rs6673106	rs2878647	rs1061171	rs463726	rs203681
rs12081550	rs10801557	rs6664877	rs2860102	rs1061170	rs460897	rs203680
rs12069060	rs10801556	rs6664705	rs2746965	rs1061147	rs460787	rs203679
rs12047565	rs10801555	rs6660100	rs2336225	rs1061111	rs460184	rs203678
rs12047106	rs10801554	rs6428357	rs2336224	rs1060821	rs459598	rs203677
rs12047103	rs10801553	rs6428356	rs2336223	rs1048663	rs454652	rs203676
rs12045503	rs10754200	rs5779848	rs2336222	rs1040597	rs436337	rs203675
rs12042805	rs10754199	rs5779847	rs2336221	rs800295	rs435628	rs203674
rs12041668	rs10737680	rs5779846	rs2300430	rs800293	rs434536	rs203673
rs12040718	rs10737679	rs5779845	rs2300429	rs800292	rs430173	rs203672
rs12039905	rs10733086	rs5779844	rs2284664	rs800291	rs428060	rs203671
rs12038674	rs10688557	rs5022901	rs2284663	rs800290	rs424535	rs203670
rs12038333	rs10685027	rs5022900	rs2274700	rs800280	rs422851	rs203669
rs12033127	rs10664537	rs5022899	rs2268343	rs800271	rs422404	rs70621
rs12032372	rs10616982	rs5022898	rs2173383	rs800269	rs420922	rs70620
rs12030500	rs10545544	rs5022897	rs2143912	rs766001	rs420921	rs15809
rs12029785	rs10540668	rs5016801	rs2104714	rs765774	rs419137	rs14473
rs12025861	rs10536523	rs5014740	rs2064456	rs742855	rs414539	rs3645
rs11809183	rs10489456	rs5014739	rs2020130	rs731557	rs412852	
rs11801630	rs10465603	rs5014738	rs2019727	rs572515	rs410232	
rs11799956	rs10465586	rs5014737	rs1803696	rs570618	rs409953	
rs11799595	rs9970784	rs5002879	rs1587325	rs569219	rs409319	

rs11799380	rs9970075	rs5002878	rs1576340	rs564657	rs409308	
rs11584505	rs9427909	rs5002877	rs1474792	rs559350	rs407361	

[0134] Two frequent non-synonymous variants, I62V in exon 2 and Y420H in exon 9, and a less frequent variant, R1210C in exon 22, exhibited the most significant association with AMD.

[0135] Three additional polymorphisms in TABLES 1A-1B are not found in the SNP database: a polymorphism in the promoter (promoter 1 in TABLE 1A); a polymorphism in intron 2 in which two T nucleotides are inserted; and a polymorphism in Exon 10A.

[0136] The first column in TABLE 1A lists the dbSNP number for polymorphisms in the Factor H gene. For example, rs800292 is the dbSNP designation for a polymorphism in the Factor H gene. A description of this polymorphism, as well as the other Factor H gene polymorphisms in dbSNP, is available at <http://www.ncbi.nlm.nih.gov> (<http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=snp&cmd=search&term=>). The second column lists the location of the polymorphism. For example, the rs800292 polymorphism is located in exon 2 of the Factor H gene. Polymorphisms not identified by a database number can be referred to by location (e.g., "intron 2"). The third column lists the nucleic acid sequence of the coding (top, 5' to 3' direction) and non-coding (bottom) strands of DNA spanning the polymorphisms. For example, the rs800292 polymorphism, G or A as indicated in the brackets for the coding strand, is flanked by the 20 nucleotides shown 5' and 3' to the polymorphism. "N" in the sequence spanning the Exon 10A polymorphism indicates the insertion of a single nucleotide, either A, C, G or T, in the variant allele. The fourth column lists the SEQ ID NO: for the sequences. The fifth column lists the amino acid change, if any, associated with the polymorphism. For example, the rs800292 polymorphism results in a change in the amino acid sequence from valine (V) to isoleucine (I) at position 62 of the Factor H polypeptide. The sixth column lists the allele frequency of the polymorphism in a control population. The numbers 1 and 2 refer to the alleles that correspond to the first and second nucleotide, respectively, at the polymorphic site in the third column. For example, for the rs800292 polymorphism, G is present in 78% and A is present in 22% of the alleles sequenced from the control population. The seventh column lists the allele frequency of the polymorphism in an AMD population. For example, for the rs800292 polymorphism, G is present in 91% and A is present in 9% of the alleles sequenced from an AMD population. The eighth column lists the chi-square and Fisher's exact tests (χ^2 and P values, respectively) for the comparison between the allele frequencies of the polymorphism in the control and

AMD populations. For example, for the rs800292 polymorphism, the χ^2 value is 16.19 and the P value is 5.74×10^{-5} , indicating that the G allele is associated with AMD.

[0137] The first column in TABLE 1B parts (1), (2) and (3) lists the dbSNP number for polymorphisms in the Factor H gene. For part (1), the second column lists the nucleic acid sequence spanning the polymorphisms (interrogated sequence). For the rs529825 (intron 1), rs800292 (exon 2), and rs203674 (intron 10) polymorphisms, the sequences of the non-coding strand of the human Factor H gene are shown. The third column lists the SEQ ID NOs: for the sequences. The fourth column lists the allele present in the chimp Factor H gene. The fifth column lists the location of the SNP. The sixth column lists the amino acid change, if any, associated with the polymorphism. For part (2), the second and fourth columns list the forward and reverse primers or AOD numbers for amplifying the polymorphisms. The third and fifth columns list the SEQ ID NOs: for the primers. For part (3), the second and fourth columns list probes used for detecting the polymorphisms. The third and fifth columns list the SEQ ID NOs: for the probes.

[0138] It should be understood that additional polymorphic sites in the Factor H gene, which are not listed in TABLES 1A-1B may be associated with AMD. Exemplary polymorphic sites in the Factor H gene are listed, for example and not limitation, above. TABLE 1C lists an additional 14 polymorphic sites in the Factor H gene, which are not found in the dbSNP database, that may be associated with AMD or other diseases. The first column lists the location of the SNP. The second column lists the nucleic acid sequence spanning the polymorphisms. "notG" in the sequence spanning the Exon 5 polymorphism indicates the presence of an A, C or T nucleotide in the variant allele. "not C" in the sequences spanning the Exon 6 polymorphisms indicates the presence of an A, G or T nucleotide in the variant allele. "N" in the sequences spanning the Exon 21 polymorphism indicates the insertion of a single nucleotide, either A, C, G or T, in the variant allele. The third column lists the amino acid change, if any associated with the polymorphism. The fourth column lists the SEQ ID NOs: for the sequences. These SNPs can also be used in carrying out methods of the invention. Moreover, it will be appreciated that these CFH polymorphisms are useful for linkage and association studies, genotyping clinical populations, correlation of genotype information to phenotype information, loss of heterozygosity analysis, and identification of the source of a cell sample.

[0139] TABLE 2 shows a haplotype analysis of eight SNPs in the human Factor H gene in AMD cases and controls. The at-risk haplotypes are shown in stippled boxes, with the

haplotype determining SNPs (Y402H and IVS10) shown in denser stippling. The protective haplotypes are shown in diagonal-lined boxes, with the haplotype determining SNPs (IVS1, I62V and IVS6) shown in denser diagonal lines. The first column lists the allele of the polymorphism in the promoter (Prom). The second column lists the allele of the non-coding strand of the polymorphism in intron 1 (IVS1). The third column lists the allele of the non-coding strand of the polymorphism in exon 2 (I62V). The fourth column lists the allele of the polymorphism in intron 6 (IVS6). The fifth column lists the allele of the polymorphism in Exon 9 (Y402H). The sixth column lists the allele of the non-coding strand of the polymorphism in intron 10 (IVS10). The seventh column lists the allele of the polymorphism in Exon 13 (Q672Q). The eighth column lists the allele of the polymorphism in Exon 18 (D936E). The dbSNP designations for these eight SNPs are listed in TABLES 1A-1B. The ninth column lists the Odds Ratio (OR) for the haplotype. The tenth column lists the P value for the at-risk and two protective haplotypes. The eleventh and twelfth columns list the frequencies of the haplotype in AMD cases and controls.

[0140] TABLE 3 shows a haplotype analysis of six Factor H polymorphisms with AMD. The first column lists certain alleles of the polymorphism in the promoter (rs3753394). The second column lists the allele of the polymorphism in intron 1 (rs529825). The third column lists the allele of the polymorphism in intron 6 (rs3766404). The fourth column lists the polymorphism in intron 10 (rs203674). The fifth column lists the allele of the polymorphism in exon 13 (rs3753396). The sixth column lists the allele of the polymorphism in exon 18 (rs1065489). The numbers 1 and 2 in columns 1 to 6 refer to the alleles that correspond to the first and second nucleotide, respectively, at each of the polymorphic sites (see TABLE 1A). Thus, columns 1 to 6 list the alleles of polymorphisms from 5' to 3' in the Factor H gene. The seventh column lists the Factor H haplotype based on the polymorphisms listed in columns 1 to 6. The eighth column lists the frequency of the indicated Factor H haplotype in a control population. The ninth column lists the frequency of the indicated Factor H haplotype in the AMD population. As shown in TABLE 3, the haplotype analysis suggests that multiple variants contribute to the association and may confer either elevated or reduced risk of AMD.

[0141] TABLE 8 shows a diplotype analysis of seven Factor H polymorphisms. The first column indicates whether the diplotype is associated with increased (risk diplotype) or decreased (protective diplotype) risk of developing AMD. Common risk and protective diplotypes are indicated. The second column lists the alleles of the polymorphism in exon 2

(I62V). The third column lists the alleles of the polymorphism in intron 2 (IVS2-18). The fourth column lists the alleles of the polymorphism in exon 9 (Y402H). The fifth column lists the alleles of the polymorphism in exon 18 (D936E). The sixth column lists the alleles of the polymorphism in intron 20 (IVS20).

Risk-Associated ("Risk") Polymorphisms and Haplotypes

[0142] Sites comprising polymorphisms associated with increased risk for AMD are shown in TABLE 1A and TABLE 2. Polymorphisms particularly associated with increased risk include a variant allele at: rs1061170 (402H; exon 9); rs203674 (intron 10) and the polymorphism at residue 1210 (R1210C; exon 22).

[0143] Certain haplotypes associated with increased risk for AMD are shown in TABLES 2 and 6 and FIGURE 5. As shown in TABLE 2 and FIGURE 5, one common at-risk haplotype is the H1 haplotype, which includes the variant allele at position 402 (encoding histidine) and the variant allele at IVS10 (intron 10, rs203674) and is found in 49% of AMD cases, but only in 26% of controls. Homozygotes for the risk diplotype (H1/H1) are significantly at risk. Other at-risk haplotypes and diplotypes are shown in TABLES 2 and 8. Similar data are presented in TABLE 3, which shows an at-risk haplotype (111211) found in 48% of AMD cases, but only in 28% of controls.

[0144] Notably, seventy percent of MPGN II (membranoproliferative glomerulonephritis type II) patients harbor this at-risk haplotype (see TABLE 7), indicating that propensity to develop MPGNII can be detected and treated as described herein for AMD.

[0145] Significant associations of these polymorphic sites were also found with various AMD subtypes, as disclosed in Example 1.

[0146] The non-synonymous polymorphism at amino acid position 1210 in exon 22 of the Factor H gene is strongly associated with AMD (see TABLE 1A). The variant allele, which encodes a cysteine instead of an arginine, is found in the heterozygous state in 5% of AMD cases, and no controls in a cohort comprised of 919 individuals ascertained at the University of Iowa. No 1210C homozygotes have been identified to date. The presence of cysteine at amino acid position 1210 of Factor H, therefore, provides a strong indication that the individual has AMD or is likely to develop AMD. Remarkably, 1210C is indicative of propensity to develop AMD or other complement mediated conditions even when detected on allele that is otherwise protective (e.g., Y402). Variation at CFH position 1210 (R1210C) is known to cause atypical hemolytic uremic syndrome (aHUS), a complement related disease

with renal manifestations. By extension, other CFH variations or mutations known to cause aHUS may be associated with an increased risk for developing AMD. The most common established aHUS-causing variations include, but are not limited to, T956M, Q1076E, D1119G, W1183L, T1184R, L1189R, L1189F, S1191W, S1191L, V1197A, and R1215G (Esparza-Gordillo et al 2005; Perez-Caballero et al 2001; Richards et al 2001; Sanchez-Corral et al 2002); additional aHUS-causing mutations are described in Saunders (Saunders et al 2006). In one aspect of the present invention, a biological sample from a subject (e.g., protein or nucleic acid) is assayed for the presence of one or more aHUS-associated variations or mutations, the presence of which is indicative of a propensity to develop AMD.

[0147] It will be appreciated that additional polymorphic sites in the Factor H gene, which are not listed in TABLES 1A-1C, may further refine this haplotype analysis. A haplotype analysis using non-synonymous polymorphisms in the Factor H gene is useful to identify variant Factor H polypeptides. Other haplotypes associated with risk may encode a protein with the same sequence as a protein encoded by a neutral or protective haplotype, but contain an allele in a promoter or intron, for example, that changes the level or site of Factor H expression. It will also be appreciated that a polymorphism in the Factor H gene, or in a Factor H-related gene, may be linked to a variation in a neighboring gene. The variation in the neighboring gene may result in a change in expression or form of an encoded protein and have detrimental or protective effects in the carrier.

Protective Polymorphisms and Haplotypes

[0148] Unexpectedly, protective polymorphisms and haplotypes were also discovered. For example, as shown in TABLE 2 and FIGURE 5, the protective H2 haplotype, including a variant allele in IVS6 (intron 6, rs3766404) occurs in 12% of controls, but only in 6% of AMD cases. The protective H4 haplotype includes the variant allele in IVS1 (intron 1, rs529825) and the variant allele (I62) (exon 2, rs800292) and occurs in 18% of controls, but only in 12% of AMD cases. Similar data is presented in TABLE 3, where the haplotype 121111 occurs in 21% of controls, but only in 13% of AMD cases and the haplotype 112111 occurs in 13% of controls, but only in 6% of AMD cases. As shown in FIGURE 5, homozygotes with a protective haplotype are significantly protected.

[0149] In some cases the protein encoded by a gene characterized by a protective haplotype has a sequence different from risk haplotype proteins (e.g., due to the presence of a nonsynonymous SNP). For example, a protective form of Factor H protein generally does not

have histidine at position 402. In some embodiments a protective form has isoleucine at position 62. Additional protective forms can be identified by (1) identifying an individual or individuals with a protective haplotype and (2) determining the sequence(s) of Factor H cDNA or protein from the individuals. Other protective forms are identified as described below in Section VIII.

Neutral Polymorphisms and Haplotypes

[0150] Certain haplotypes are associated in a population with neither increased risk nor decreased risk of developing AMD and are referred to as "neutral." Examples of neutral haplotypes identified in a Caucasian population are shown in FIGURE 5 (H3 and H5). Additional or different neutral haplotypes may be identified in racially/ethnically different populations. Proteins encoded by a gene characterized by a neutral haplotype are "neutral" Factor H proteins. As explained supra, "neutral" Factor H proteins could provide therapeutic benefit when administered to patients having a risk haplotype or diagnosed with AMD. For example, exemplary proteins encoded by genes characterized by a neutral haplotype include proteins not having histidine at position 402 and/or not having isoleucine at position 62. A protein not having histidine at position 402 may have tyrosine at that position, or may have an amino acid other than histidine or tyrosine. A protein not having isoleucine at position 62 may have valine at that position, or may have an amino acid other than valine or isoleucine. A neutral form of Factor H protein generally does not have cysteine at position 1210.

V. FACTOR H RELATED 5 (CFHR5) GENE POLYMORPHISMS

[0151] In one aspect, the invention provides new diagnostic, treatment and drug screening methods related to the discovery that polymorphic sites in the Factor H and CFHR5 genes are associated with susceptibility to and development of MPGNII.

[0152] Factor H and CFHR5 polymorphisms associated with MPGNII were identified as described in Example 2, by examining the coding and adjacent intronic regions of Factor H or CFHR5 for variants using PCR amplification, followed by agarose gel electrophoresis and bi-directional sequencing according to standard protocols to verify PCR products. Novel and reported SNPs were typed in the control population by denaturing high performance liquid chromatography (DHPLC). Primers used to amplify the Factor H and *CFHR5* coding sequences are shown in TABLES 9 and 10, respectively.

[0153] The test group consisted of patients with biopsy-proven MPGNII were ascertained in nephrology divisions and enrolled in this study under IRB-approved guidelines. The control group consisted of ethnically-matched, but not age-matched, unrelated persons in whom AMD had been excluded by ophthalmologic examination.

[0154] As shown in TABLES 11 and 12, a significant association of polymorphic sites in the Factor H gene with MPGNII was found in an examination of 22 MPGNII cases and 131 ethnically-matched controls. Eleven (11) polymorphisms in the Factor H gene are listed in TABLES 11 and 12. Of these, six (6) are found in the SNP database (dbSNP) which may be found in the National Center for Biotechnology Information (NCBI). The dbSNP is a collection of SNPs in the human genome. The SNPs in the Factor H gene are dispersed among the 22 coding exons of the Factor H gene and among the promoter, the 5' untranslated region, the introns, and the 3' untranslated region of the Factor H gene. The accession numbers for 379 SNPs in the human Factor H gene that are found in the dbSNP database are listed above. These SNPs can be used in carrying out methods of the invention.

[0155] Five additional polymorphisms in TABLES 11 and 12 are not found in the SNP database: a polymorphism in intron 2 in which two T nucleotides are inserted (IVS2 - 18insTT); a polymorphism in intron 7 (IVS7 -53G>T); a polymorphism in intron 15 (IVS15 - 30C>A); a polymorphism in intron 18 (IVS18 -89T>C; and a polymorphism in Exon 20 (N1050Y). These polymorphisms are useful in the methods of the invention. Moreover, it will be appreciated that these CFHR5 polymorphisms are useful for linkage and association studies, genotyping clinical populations, correlation of genotype information to phenotype information, loss of heterozygosity analysis, and identification of the source of a cell sample.

[0156] The first row of TABLE 11 lists the exon or intron position of the SNP in the Factor H gene. For exon SNPs, the amino acid position and change, if any, is listed. For example the exon 2 SNP is at position 62 of the Factor H polypeptide and a change from valine (V) to isoleucine (I). For intron SNPs, the nature the SNP is indicated. For example, the intron 2 SNP is an insertion of two nucleotides, TT. The second row of TABLE 11 lists dbSNP number, if any, for the polymorphism. For example, rs800292 is the dbSNP designation for a polymorphism in exon 2 in the Factor H gene. A description of this polymorphism, as well as the other Factor H (CFH) gene polymorphisms in dbSNP, is available at <http://www.ncbi.nlm.nih.gov> (<http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=snp&cmd=search&term=>). The third to fifth rows of TABLE 11 lists number of times a particular diplotype is present among the 22 MPGNII patients. For example, for

the exon 2 SNP, GG is present in 20 patients, GA is present in 2 patient, and AA is present in no patients, with MPGNII. The sixth and seventh rows of TABLE 11 list the frequency that a particular haplotype is present among the 22 MPGNII patients. For example, for the exon 2 SNP, G is present in 95%, and A is present in 5%, of the alleles of the 22 MPGNII patients. The eighth row lists the nucleotide of the common haplotype in the Factor H gene for the 22 MPGNII patients. For example, G is the more frequent nucleotide in the exon 2 SNP, and 9 T nucleotides is more frequently observed than 11 T nucleotides in the intron 2 SNP, in the Factor H gene for the 22 MPGNII patients. The remaining rows list the diplotype for the 11 SNPs in the Factor H gene for each of the 22 MPGNII patients.

[0157] It should be understood that additional polymorphic sites in the Factor H gene, which are not listed in TABLE 11 may be associated with MPGNII. Exemplary polymorphic sites in the Factor H gene are listed, for example and not limitation, above.

[0158] TABLE 12 shows a comparison of the SNP frequencies in patients with MPGNII *versus* AMD-negative, ethnically-matched control individuals. The first column of TABLE 12 lists the SNP in the Factor H gene. The second and third columns of TABLE 12 list the frequencies that a particular haplotype is present among the 22 MPGNII patients. The fourth and fifth columns of TABLE 12 list the frequencies that a particular haplotype is present among the 131 control individuals. The sixth column of TABLE 12 lists the P-value calculated for each data set.

[0159] As shown in TABLES 11 and 12, two frequent non-synonymous variants, I62V in exon 2 and Y420H in exon 9, a synonymous variant, A307A in exon 10, and a polymorphism in intron 2 exhibited significant association with MPGNII.

[0160] As shown in TABLES 14 and 15, a significant association of polymorphic sites in the CFHR5 gene with MPGNII was found in an examination of 22 MPGNII cases and 103 ethnically-matched controls. Five (5) polymorphisms in the CFHR5 gene are listed in TABLES 14 and 15; these are found in dbSNP in the NCBI. The SNPs in the CFHR5 gene are dispersed among the 10 coding exons of the CFHR5 gene and among the promoter, the 5' untranslated region, the introns, and the 3' untranslated region of the CFHR5 gene. Listed below are the accession numbers for 82 SNPs in the human CFHR5 gene that are found in the dbSNP database. These SNPs can be used in carrying out methods of the invention.

TABLE B

rs16840956	rs12116643	rs10922151	rs9427662	rs7535993	rs6694672	rs1332664
rs16840946	rs12097879rs	rs10801584	rs9427661	rs7532441	rs6692162	rs1325926
rs16840943	12097550	rs10801583	rs9427660	rs7532068	rs6657256	rs1170883
rs12755054	rs12092294	rs10622350	rs9427659	rs7528757	rs6657171	rs1170882
rs12750576	rs12091602	rs10614978	rs7555407	rs7527910	rs5779855	rs1170881
rs12745733	rs12064805	rs10613146	rs7555391	rs7522952	rs3748557	rs1170880
rs12736097	rs12049041	rs10588279	rs7554757	rs7522197	rs2151137	rs1170879
rs12736087	rs12039272	rs9727516	rs7550970	rs7419075	rs2151136	rs1170878
rs12735776	rs11583363	rs9427942	rs7550735	rs7366339	rs1855116	rs928440
rs12731848	rs11306823	rs9427941	rs7550650	rs6702632	rs1759016	rs928439
rs12731209	rs10922153	rs9427664	rs7547265	rs6702340	rs1750311	
rs12142971	rs10922152	rs9427663	rs7537588	rs6674853	rs1412636	

[0161] The first row of TABLE 14 lists the exon, promoter or intron position of the SNP in the CFHR5 gene. For exon SNPs, the amino acid position and change, if any, is listed. For example, the exon 2 SNP is at position 46 of the CFHR5 polypeptide and a change from proline (P) to serine (S). For promoter and intron SNPs, the nature of the SNP is indicated. For example, the promoter SNP at position -249 replaces T with C. The second row of TABLE 14 lists dbSNP number, if any, for the polymorphism. For example, rs9427661 is the dbSNP designation for a polymorphism in the promoter region of the CFHR5 gene. A description of this polymorphism, as well as the other CFHR5 gene polymorphisms in dbSNP, is available at <http://www.ncbi.nlm.nih.gov> (<http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=snp&cmd=search&term=>). The third to fifth rows of TABLE 14 lists number of times a particular diplotype is present among the 22 MPGNII patients. For example, for the exon 2 SNP, CC is present in 19 patients, CT is present in 3 patients, and TT is present in no patients, with MPGNII. The sixth and seventh rows of TABLE 14 list the frequency that a particular haplotype is present among the 22 MPGNII patients. For example, for the exon 2 SNP, C (encoding proline) is present in 93%, and T (encoding serine) is present in 7%, of the alleles of the 22 MPGNII patients. The eighth row lists the nucleotide of the common haplotype in the CFHR5 gene for the 22 MPGNII patients. For example, C is the more frequent nucleotide in the exon 2 SNP in the CFHR5 gene for the 22 MPGNII patients. The remaining rows list the diplotype for the 5 SNPs in the CFHR5 gene for each of the 22 MPGNII patients.

[0162] It should be understood that additional polymorphic sites in the CFHR5 gene, which are not listed in TABLE 14 may be associated with MPGNII. Exemplary polymorphic sites in the CFHR5 gene are listed, for example and not limitation, above.

[0163] TABLE 15 shows a comparison of the SNP frequencies in patients with MPGNII versus AMD-negative, ethnically-matched control individuals. The first column of TABLE 15 lists the SNP in the CFHR5 gene. The second and third columns of TABLE 15 list the frequencies that a particular haplotype is present among the 22 MPGNII patients. The fourth and fifth columns of TABLE 15 list the frequencies that a particular haplotype is present among the 103 control individuals. The sixth column of TABLE 15 lists the P-value calculated for each data set.

[0164] As shown in TABLES 14 and 15, one non-synonymous variant, P46S in exon 2, and two promoter polymorphisms, -249T>C and -2-T>C, exhibited significant association with MPGNII.

Risk-Associated ("Risk") Polymorphisms and Haplotypes Identified In MPGNII Patients

[0165] Sites comprising polymorphisms in Factor H and CFHR5 associated with increased risk for MPGNII are shown in TABLES 11 and 12 and TABLES 14 and 15, respectively. Polymorphisms particularly associated with increased risk in Factor H and CFHR5 include a variant allele at rs1061170 (Y420H in exon 9) and rs12097550 (P46S in exon 2), respectively.

[0166] Certain haplotypes associated with increased risk for MPGNII are shown in TABLES 12 and 15. As shown in TABLE 12, one at-risk haplotype in the Factor H gene includes the variant allele (encoding histidine) at position 402 and is found in 64% of MPGNII cases, but only in 33% of controls. As shown in TABLE 15, one at-risk haplotype in the CFHR5 gene includes the variant allele (encoding serine) at position 46 and is found in 7% of MPGNII cases, but only in <1% of controls.

[0167] It will be appreciated that additional polymorphic sites in the Factor H and *CFHR5* genes, which are not listed in TABLES 11-12 and 14-15, may further refine these haplotype analyses. A haplotype analysis using non-synonymous polymorphisms in the Factor H or CFHR5 gene is useful to identify variant Factor H or CFHR5 polypeptides. Other haplotypes associated with risk may encode a protein with the same sequence as a protein encoded by a neutral or protective haplotype, but contain an allele in a promoter or intron, for example, that changes the level or site of Factor H or CFHR5 expression.

Protective Polymorphisms and Haplotypes

[0168] Unexpectedly, protective polymorphisms and haplotypes were also discovered. For example, as shown in TABLE 12, the haplotype with the variant allele in exon 2 (rs800292, I62V) occurs in 23% of controls, but only in <3% of MPGNII cases and the haplotype with the variant allele in IVS2 (intron 2, -18insTT) occurs in 26% of controls, but only in <3% of MPGNII cases. The haplotype with the variant allele in exon 10 (rs2274700, A473A) occurs at higher frequency in controls than in MPGNII cases.

[0169] In some cases the protein encoded by a gene characterized by a protective haplotype has a sequence different from risk haplotype proteins. For example, a protective form of Factor H protein generally does not have histidine at position 402. In some embodiments a protective form has isoleucine at position 62. Additional protective forms can be identified by (1) identifying an individual or individuals with a protective haplotype and (2) determining the sequence(s) of Factor H cDNA or protein from the individuals. Some protective forms are less than full-length. Protective forms of CFHR5 protein may be similarly identified.

Neutral Polymorphisms and Haplotypes

[0170] Certain haplotypes are associated with neither increased risk or decreased risk of developing MPGNII and are referred to as "neutral." Proteins encoded by a gene characterized by a neutral haplotype are "neutral" Factor H or CFHR5 proteins. For example, exemplary proteins encoded by genes characterized by a neutral haplotype include Factor H proteins not having histidine at position 402 or isoleucine at position 62, and CFHR5 proteins not having serine at position 46.

Significance of Polymorphisms in MPGNII Patients

[0171] As shown in Example 2, it has been discovered that the same CFH polymorphisms associated with propensity to develop AMD are also associated with development of membranoproliferative glomerulonephritis type 2 (MPGN II). Indeed, the risk haplotypes originally found in AMD patients (Y402 H and IVS10) are also found in 70% of patients tested having membranoproliferative glomerulonephritis type 2 (MPGN II), indicating that the diagnostic methods of the invention are useful to detect this condition. In addition, variations and haplotypes in the CFHR5 gene were strongly associated with increased risk of

having MPGNII. One conclusion that emerges from these data is that MPGNII and AMD are alternative manifestations of the same genetic lesion. Notably, patients with MPGNII develop drusen that are clinically and compositionally indistinguishable from drusen that form in AMD. The single feature that distinguishes these two fundus phenotypes is age of onset - drusen in MPGNII develop early, often in the second decade of life, while drusen in AMD develop later in life. We conclude that polymorphisms in the Factor H gene and CFHR5 gene identified in either population (AMD or MPGNII) are predictive of susceptibility to both diseases. There are likely other factors that contribute to MPGNII and account for the early manifestation. Because AMD is very common and MPGNII is rare, the haplotype analysis of both CFH and CFHR5 genes and other methods described herein will be useful for screening and treatment of patients with AMD, or with an increased likelihood of developing AMD.

Loss of Function

[0172] Loss of the normal or wild-type function of Factor H or CFHR5 may be associated with AMD. Non-synonymous polymorphisms in the Factor H gene, such as those shown in TABLES 1A, 1B, 1C, 11, 14 and 15, showing the strongest correlation with AMD and resulting in a variant Factor H polypeptide or variant CFHR5 polypeptide, are likely to have a causative role in AMD. Such a role can be confirmed by producing a transgenic non-human animal expressing human Factor H or CFHR5 bearing such a non-synonymous polymorphism(s) and determining whether the animal develops AMD. Polymorphisms in Factor H or CFHR5 coding regions that introduce stop codons may cause AMD by reducing or eliminating functional Factor H or CFHR5 protein. Stop codons may also cause production of a truncated Factor H or CFHR5 peptide with aberrant activities relative to the full-length protein. Polymorphisms in regulatory regions, such as promoters and introns, may cause AMD by decreasing Factor H or CFHR5 gene expression. Polymorphisms in introns (e.g., intron 2 of CFH) may also cause AMD by altering gene splicing patterns resulting in an altered Factor H or CFHR5 protein. CFH RNA or proteins can be assayed to detect changes in expression of splice variants, where said changes are indicative of a propensity to develop AMD. Alternative splice patterns have been reported for the Factor H gene itself.

[0173] The effect of polymorphisms in the Factor H gene or CFHR5 gene on AMD can be determined by several means. Alterations in expression levels of a variant Factor H or CFHR5 polypeptide can be determined by measuring protein levels in samples from groups

of individuals having or not having AMD or various subtypes of AMD. Alterations in biological activity of variant Factor H or CFHR5 polypeptides can be detected by assaying for in vitro activities of Factor H or CFHR5, for example, binding to C3b or to heparin, in samples from the above groups of individuals.

VI. POLYMORPHISMS AT SITES OF GENOMIC DUPLICATION

[0174] As illustrated in FIGURE 18, the genes for CFH and the factor H related (CFHR) 1-5 genes have regions of shared, highly conserved, sequence which likely arose from genomic duplications. Certain SNPs and variations found in CFH or CFHR5, such as those described herein, are also expected in the corresponding sequences of CFHR1, CFHR2, CFHR3, and CFHR4. For example, sequences corresponding to *CFH* exon 22 are found in CFH, CFHR1 and CFHR2, and it is possible that polymorphisms identified in exon 22 of CFH (e.g., R1210C) are also found in CFHR1 and/or CFHR2 and these variants might be linked to propensity to development of AMD, MPGNII, and other complement related conditions. Homologous blocks of sequence flanking the polymorphic sites identified in CFH and CFHR5 can be identified by alignment of the cDNA or genomic sequences in those regions. The conserved sequences flanking the polymorphic site usually comprise at least 10 bp (on either side of the polymorphic site) and more often at least 20 bp, or at least 50 bp, or at least 100 bp with at least 95% identity at the nucleotide level, and sometimes with at least 98% identity, at least 99% identity, or even 100% identity). Identity can be determined by inspection or using well know algorithms (Smith and Waterman, 1981 or Needleman and Wunsch, 1970, both *supra*). The invention therefore provides methods of determining a subject's propensity to develop age-related macular degeneration (AMD) or other conditions by detecting the presence or absence of a variation at a polymorphic site of a Factor H-related gene that corresponds to a homologous polymorphic site in the CFH or CFHR5 gene.

[0175] Sequences for CFH and the factor H related genes are known in the art (see sequences and accession numbers provided elsewhere herein). Also see Rodriquez de Cordoba, S., et al, 2004, *Mol Immunol* 41:355-67; Zipfel et al, 1999, *Immunopharmacology* 42:53-60; Zipfel et al., Factor H family proteins: on complement, microbes and human diseases, *Biochem Soc Trans.* 2002 Nov;30(Pt 6):971-8; Diaz-Guillen MA, et al., A radiation hybrid map of complement factor H and factor H-related genes, *Immunogenetics*, 1999 Jun;49(6):549-52; Skerka C, et al., A novel short consensus repeat-containing molecule is related to human complement factor H, *J Biol Chem.* 1993 Feb 5;268(4):2904-8; Skerka C, et al., The human

factor H-related gene 2 (FHR2): structure and linkage to the coagulation factor XIIIb gene, *Immunogenetics*, 1995;42(4):268-74; Male DA, et al., Complement factor H: sequence analysis of 221 kb of human genomic DNA containing the entire fh, fHR-1 and fHR-3 genes, *Mol Immunol*, 2000 Jan-Feb;37(1-2):41-52; Hellwage J, et al., Biochemical and functional characterization of the factor-H-related protein 4 (FHR-4), *Immunopharmacology*, 1997 Dec;38(1-2):149-57; Skerka C, et al., The human factor H-related protein 4 (FHR-4). A novel short consensus repeat-containing protein is associated with human triglyceride-rich lipoproteins, *J Biol Chem*, 1997 Feb 28;272(9):5627-34; Hellwage J, et al., Functional properties of complement factor H-related proteins FHR-3 and FHR-4: binding to the C3d region of C3b and differential regulation by heparin, *FEBS Lett*, 1999 Dec 3;462(3):345-52; Jozsi M, et al., FHR-4A: a new factor H-related protein is encoded by the human FHR-4 gene, *Eur J Hum Genet*, 2005 Mar;13(3):321-9; McRae JL, et al., Location and structure of the human FHR-5 gene, *Genetica*, 2002 Mar;114(2):157-61; McRae JL, et al., Human factor H-related protein 5 has cofactor activity, inhibits C3 convertase activity, binds heparin and C-reactive protein, and associates with lipoprotein, *J Immunol*, 2005 May 15;174(10):6250-6; Murphy B, et al., Factor H-related protein-5: a novel component of human glomerular immune deposits, *Am J Kidney Dis*, 2002 Jan;39(1):24-7.

VII. DETECTION AND ANALYSIS OF FACTOR H POLYMORPHISMS ASSOCIATED WITH AMD

[0176] The discovery that polymorphic sites and haplotypes in the Factor H gene and CFHR5 gene are associated with AMD (and MPGNII) has a number of specific applications, including screening individuals to ascertain risk of developing AMD and identification of new and optimal therapeutic approaches for individuals afflicted with, or at increased risk of developing, AMD. Without intending to be limited to a specific mechanism, polymorphisms in the Factor H gene may contribute to the phenotype of an individual in different ways. Polymorphisms that occur within the protein coding region of Factor H may contribute to phenotype by affecting the protein structure and/or function. Polymorphisms that occur in the non-coding regions of Factor H may exert phenotypic effects indirectly via their influence on replication, transcription and/or translation. Certain polymorphisms in the Factor H gene may predispose an individual to a distinct mutation that is causally related to a particular AMD phenotype. Alternatively, as noted above, a polymorphism in the CFH gene, or in a CFHR5, may be linked to a variation in a neighboring gene (including but not limited to

CFHR-1, 2, 3, or 4). The variation in the neighboring gene may result in a change in expression or form of an encoded protein and have detrimental or protective effects in the carrier.

A. Preparation of Samples for Analysis

[0177] Polymorphisms are detected in a target nucleic acid isolated from an individual being assessed. Typically genomic DNA is analyzed. For assay of genomic DNA, virtually any biological sample containing genomic DNA or RNA, *e.g.*, nucleated cells, is suitable. For example, in the experiments described in Example 1, genomic DNA was obtained from peripheral blood leukocytes collected from case and control subjects (QIAamp DNA Blood Maxi kit, Qiagen, Valencia, CA). Other suitable samples include saliva, cheek scrapings, biopsies of retina, kidney or liver or other organs or tissues; skin biopsies; amniotic fluid or CVS samples; and the like. Alternatively RNA or cDNA can be assayed. Alternatively, as discussed below, the assay can detect variant Factor H proteins. Methods for purification or partial purification of nucleic acids or proteins from patient samples for use in diagnostic or other assays are well known.

B. Detection of Polymorphisms in Target Nucleic Acids

[0178] The identity of bases occupying the polymorphic sites in the Factor H gene and the Factor H-Related 5 gene shown in TABLES 1A, 1B, 1C, 11, 14 and 15, as well as others in the dbSNP collection that are located in or adjacent to the Factor H or CFHR5 genes (see lists above), can be determined in an individual, *e.g.*, in a patient being analyzed, using any of several methods known in the art. Examples include: use of allele-specific probes; use of allele-specific primers; direct sequence analysis; denaturing gradient gel electrophoresis (DGGE) analysis; single-strand conformation polymorphism (SSCP) analysis; and denaturing high performance liquid chromatography (DHPLC) analysis. Other well known methods to detect polymorphisms in DNA include use of: Molecular Beacons technology (see, *e.g.*, Piatek et al., 1998; *Nat. Biotechnol.* 16:359-63; Tyagi, and Kramer, 1996, *Nat. Biotechnology* 14:303-308; and Tyagi, et al., 1998, *Nat. Biotechnol.* 16:49-53), Invader technology (see, *e.g.*, Neri et al., 2000, *Advances in Nucleic Acid and Protein Analysis* 3826:117-125 and U.S. Patent No. 6,706,471), nucleic acid sequence based amplification (Nasba) (Compton, 1991), Scorpion technology (Thelwell et al., 2000, *Nuc. Acids Res.* 28:3752-3761 and Solinas et al., 2001, "Duplex Scorpion primers in SNP analysis and FRET applications" *Nuc. Acids Res.*

29:20.), restriction fragment length polymorphism (RFLP) analysis, and the like. Additional methods will be apparent to the one of skill.

[0179] The design and use of allele-specific probes for analyzing polymorphisms are described by *e.g.*, Saiki et al., 1986; Dattagupta, EP 235,726, Saiki, WO 89/11548. Briefly, allele-specific probes are designed to hybridize to a segment of target DNA from one individual but not to the corresponding segment from another individual, if the two segments represent different polymorphic forms. Hybridization conditions are chosen that are sufficiently stringent so that a given probe essentially hybridizes to only one of two alleles. Typically, allele-specific probes are designed to hybridize to a segment of target DNA such that the polymorphic site aligns with a central position of the probe.

[0180] Exemplary allele-specific probes for analyzing Factor H polymorphisms are shown in TABLE 16A. Using the polymorphism dbSNP No. rs1061170 as an illustration, examples of allele-specific probes include: 5'-TTTCTTCCATAATTTTG-3' [SEQ ID NO:234] (reference allele probe) and 5'-TTTCTTCCATGATTTTG-3' [SEQ ID NO:235] (variant allele probe); and 5'-TAATCAAAATTATGGAA-3' [SEQ ID NO:232] (reference allele probe) and 5'-TAATCAAAATCATGGAA -3' [SEQ ID NO:233] (variant allele probe). In this example, the first set of allele-specific probes hybridize to the non-coding strand of the Factor H gene spanning the exon 9 polymorphism. The second set of allele-specific probes hybridize to the coding strand of the Factor H spanning the exon 9 polymorphism. These probes are 17 bases in length. The optimum lengths of allele-specific probes can be readily determined using methods known in the art.

[0181] Allele-specific probes are often used in pairs, one member of a pair designed to hybridize to the reference allele of a target sequence and the other member designed to hybridize to the variant allele. Several pairs of probes can be immobilized on the same support for simultaneous analysis of multiple polymorphisms within the same target gene sequence.

[0182] The design and use of allele-specific primers for analyzing polymorphisms are described by, *e.g.*, WO 93/22456 and Gibbs, 1989. Briefly, allele-specific primers are designed to hybridize to a site on target DNA overlapping a polymorphism and to prime DNA amplification according to standard PCR protocols only when the primer exhibits perfect complementarity to the particular allelic form. A single-base mismatch prevents DNA amplification and no detectable PCR product is formed. The method works best when

the polymorphic site is at the extreme 3'-end of the primer, because this position is most destabilizing to elongation from the primer.

[0183] Exemplary allele-specific primers for analyzing Factor H polymorphisms are shown in TABLE 16B. Using the polymorphism dbSNP No. rs1061170 as an illustration, examples of allele-specific primers include: 5'-CAAAC TTTCTTCCATA-3' [SEQ ID NO:294] (reference allele primer) and 5'-CAAAC TTTCTTCCATG-3' [SEQ ID NO:295] (variant allele primer); and 5'-GGATATAATCAA AATT-3' [SEQ ID NO:292] (reference allele primer) and 5'-GGATATAATCAA AATC-3' [SEQ ID NO:293] (variant allele primer). In this example, the first set of allele-specific primers hybridize to the non-coding strand of the Factor H gene directly adjacent to the polymorphism in exon 9, with the last nucleotide complementary to the reference or variant polymorphic allele as indicated. These primers are used in standard PCR protocols in conjunction with another common primer that hybridizes to the coding strand of the Factor H gene at a specified location downstream from the polymorphism. The second set of allele-specific primers hybridize to the coding strand of the Factor H gene directly adjacent to the polymorphic site in exon 9, with the last nucleotide complementary to the reference or variant polymorphic allele as indicated. These primers are used in standard PCR protocols in conjunction with another common primer that hybridizes to the non-coding strand of the Factor H gene at a specified location upstream from the polymorphism. The common primers are chosen such that the resulting PCR products can vary from about 100 to about 300 bases in length, or about 150 to about 250 bases in length, although smaller (about 50 to about 100 bases in length) or larger (about 300 to about 500 bases in length) PCR products are possible. The length of the primers can vary from about 10 to 30 bases in length, or about 15 to 25 bases in length. The sequences of the common primers can be determined by inspection of the Factor H genomic sequence, which is found under GenBank accession number AL049744.

[0184] Many of the methods for detecting polymorphisms involve amplifying DNA or RNA from target samples (e.g., amplifying the segments of the Factor H gene of an individual using Factor H-specific primers) and analyzing the amplified gene. This can be accomplished by standard polymerase chain reaction (PCR & RT-PCR) protocols or other methods known in the art. The amplifying may result in the generation of Factor H allele-specific oligonucleotides, which span the single nucleotide polymorphic sites in the Factor H gene. The Factor H-specific primer sequences and Factor H allele-specific oligonucleotides

may be derived from the coding (exons) or non-coding (promoter, 5' untranslated, introns or 3' untranslated) regions of the Factor H gene.

[0185] Amplification products generated using PCR can be analyzed by the use of denaturing gradient gel electrophoresis (DGGE). Different alleles can be identified based on sequence-dependent melting properties and electrophoretic migration in solution. See Erlich, ed., PCR Technology, Principles and Applications for DNA Amplification, Chapter 7 (W.H. Freeman and Co, New York, 1992).

[0186] Alleles of target sequences can be differentiated using single-strand conformation polymorphism (SSCP) analysis. Different alleles can be identified based on sequence- and structure-dependent electrophoretic migration of single stranded PCR products (Orita et al., 1989). Amplified PCR products can be generated according to standard protocols, and heated or otherwise denatured to form single stranded products, which may refold or form secondary structures that are partially dependent on base sequence.

[0187] Alleles of target sequences can be differentiated using denaturing high performance liquid chromatography (DHPLC) analysis. Different alleles can be identified based on base differences by alteration in chromatographic migration of single stranded PCR products (Frueh and Noyer-Weidner, 2003). Amplified PCR products can be generated according to standard protocols, and heated or otherwise denatured to form single stranded products, which may refold or form secondary structures that are partially dependent on the base sequence.

[0188] Direct sequence analysis of polymorphisms can be accomplished using DNA sequencing procedures that are well-known in the art. See Sambrook et al., Molecular Cloning, A Laboratory Manual (2nd Ed., CSHP, New York 1989) and Zyskind et al., Recombinant DNA Laboratory Manual (Acad. Press, 1988).

[0189] A wide variety of other methods are known in the art for detecting polymorphisms in a biological sample. See, e.g., Ullman et al. "Methods for single nucleotide polymorphism detection" U.S. Pat. No. 6,632,606; Shi, 2002, "Technologies for individual genotyping: detection of genetic polymorphisms in drug targets and disease genes" *Am J Pharmacogenomics* 2:197-205; Kwok et al., 2003, "Detection of single nucleotide polymorphisms" *Curr Issues Biol.* 5:43-60).

[0190] It will be apparent to the skilled practitioner guided by this disclosure that various polymorphisms and haplotypes can be detected to assess the propensity of an individual to develop a Factor H related condition. The following examples and combinations, and others

provided herein, are provided for illustration and not limitation. In one aspect of the invention, the allele of the patient at one of more of the following polymorphic sites in the Factor H gene is determined: rs529825; rs800292; rs3766404; rs1061147; rs1061170; and rs203674. In one embodiment the allele of the patient at rs529825 is determined. In one embodiment the allele of the patient at rs800292 is determined. In one embodiment the allele of the patient at rs3766404 is determined. In one embodiment the allele of the patient at rs1061147 is determined. In one embodiment the allele of the patient at rs1061170 is determined. In one embodiment the allele of the patient at rs203674 is determined. In one embodiment at least one of rs529825 and rs800292 is determined. In one embodiment at least one of rs1061147, rs1061170 and rs203674 is determined. In one embodiment at least one of rs529825 and rs800292 is determined, rs3766404 is determined, and at least one of rs1061147, rs1061170 and rs203674 is determined. In one embodiment alleles at rs529825, rs800292, rs3766404, rs1061170 and rs203674 are determined. The aforementioned polymorphisms and combinations of polymorphisms are provided herein for illustration and are not intended to limit the invention in any way. That is, other polymorphisms and haplotypes useful in practicing the invention will be apparent from this disclosure.

[0191] In a related aspect of the invention, the allele of the patient at one of more of the following polymorphic sites in the Factor H gene is determined: rs529825; rs800292; intron 2 (IVS2 or insTT); rs3766404; rs1061147; rs1061170; exon 10A; rs203674; rs375046; and exon 22 (1210). In one embodiment the allele of the patient at rs529825 is determined. In one embodiment the allele of the patient at rs800292 is determined. In one embodiment the allele of the patient at intron 2 is determined. In one embodiment the allele of the patient at rs3766404 is determined. In one embodiment the allele of the patient at rs1061147 is determined. In one embodiment the allele of the patient at rs1061170 is determined. In one embodiment the allele of the patient at exon 10A is determined. In one embodiment the allele of the patient at rs203674 is determined. In one embodiment the allele of the patient at rs375046 is determined. In one embodiment the allele of the patient at exon 22 (1210) is determined. In one embodiment at least one of rs529825 and rs800292 is determined; intron 2 is determined; rs3766404 is determined; at least one of rs1061147, rs1061170 and rs203674 is determined; exon 10A is determined; rs375046 is determined; and exon 22 (1210) is determined. In one embodiment alleles at rs529825, rs800292, intron 2; rs3766404, rs1061170, exon 10A, rs203674, rs375046, and exon 22 (1210) are determined. In one embodiment one, two, three, four five, or more than five of the following polymorphic sites

in the Factor H gene is determined: rs529825; rs800292; intron 2 (IVS2 or insTT); rs3766404; rs1061147; rs1061170; rs2274700; exon 10A; rs203674; rs375046; and exon 22 (1210). The aforementioned polymorphisms and combinations of polymorphisms are provided for illustration and are not intended to limit the invention in any way.

[0192] As discussed above, the non-synonymous polymorphism at amino acid position 1210 in exon 22 of the Factor H gene is strongly associated with AMD, and the presence of cysteine at amino acid position 1210 of Factor H, therefore, provides a strong indication that the individual has AMD or is likely to develop AMD. Remarkably, 1210C is indicative of propensity to develop AMD or other complement mediated conditions even when detected on allele that is otherwise protective (e.g., Y402). Thus, the allele of the patient at exon 22 (1210) is highly informative with respect to risk of developing AMD or other Factor H-associated diseases.

[0193] In a related aspect of the invention, the allele of an individual at one of more of the following polymorphic sites in the *CFHR5* gene is determined: rs9427661 (-249T>C); rs9427662 (-20T>C); and rs12097550 (P46S). In one embodiment the allele of the patient at rs9427661 is determined. In one embodiment the allele of the patient at rs9427662 is determined. In one embodiment the allele of the patient at rs12097550 is determined. In one embodiment at least one of rs9427661 and rs9427662 is determined. In one embodiment at least one of rs9427661 and rs9427662 is determined, and rs12097550 is determined. In one embodiment rs9427661, rs9427662 and rs12097550 is determined. The aforementioned polymorphisms and combinations of polymorphisms are provided for illustration and are not intended to limit the invention in any way. That is, other polymorphisms and haplotypes useful in practicing the invention will be apparent from this disclosure.

C. Detection of Protein Variants

[0194] In one embodiment of the invention, a protein assay is carried out to characterize polymorphisms in a subject's CFH or CFHR5 genes. Methods that can be adapted for detection of variant CFH, HFL1 and CFHR5 are well known. These methods include analytical biochemical methods such as electrophoresis (including capillary electrophoresis and two-dimensional electrophoresis), chromatographic methods such as high performance liquid chromatography (HPLC), thin layer chromatography (TLC), hyperdiffusion chromatography, mass spectrometry, and various immunological methods such as fluid or gel precipitin reactions, immunodiffusion (single or double), immunoelectrophoresis,

radioimmunoassay (RIA), enzyme-linked immunosorbent assays (ELISAs), immunofluorescent assays, western blotting and others.

[0195] For example, a number of well established immunological binding assay formats suitable for the practice of the invention are known (see, e.g., Harlow, E.; Lane, D. Antibodies: a laboratory manual. Cold Spring Harbor, N.Y: Cold Spring Harbor Laboratory; 1988; and Ausubel et al., (2004) Current Protocols in Molecular Biology, John Wiley & Sons, New York NY. The assay may be, for example, competitive or non-competitive. Typically, immunological binding assays (or immunoassays) utilize a "capture agent" to specifically bind to and, often, immobilize the analyte. In one embodiment, the capture agent is a moiety that specifically binds to a variant CFH or CFHR5 polypeptide or subsequence. The bound protein may be detected using, for example, a detectably labeled anti-CFH/CFHR5 antibody. In one embodiment, at least one of the antibodies is specific for the variant form (e.g., does not bind to the wild-type CFH or CFHR5 polypeptide. In one embodiment, the variant polypeptide is detected using an immunoblot (Western blot) format.

D. Patient Screening/Diagnosis of AMD

[0196] Polymorphisms in the Factor H gene, such as those shown in TABLE 1A, TABLE 1B, TABLE 1C or identified as described herein, which correlate with AMD or with particular subtypes of AMD, are useful in diagnosing AMD or specific subtypes of AMD, or susceptibility thereto. Polymorphisms in the CFHR5 gene, such as those shown in TABLES 14 and 15 or identified as described herein, which correlate with AMD or with particular subtypes of AMD, are useful in diagnosing AMD or specific subtypes of AMD, or susceptibility thereto. These polymorphisms are also useful for screening for MPGNII and other Factor H-associated diseases.

[0197] Individuals identified as at high risk for developing AMD can take steps to reduce risk, including frequent ophthalmological examinations and treatments described below, known in the art, or developed in the future.

[0198] As described in Example 1, an at-risk CFH haplotype in combination with a triggering event (e.g., infection) appears to be sufficient for disease manifestation. Patients identified as at-risk for AMD can receive aggressive therapy (e.g., using antibiotics, anti-inflammatory agents, treatment with protective forms of CFH/CFHR5, or treatment with other modulators of CFH activity) at early signs of infection.

[0199] Combined detection of several such polymorphic forms (i.e., the presence or absence of polymorphisms at specified sites), for example, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or all of the polymorphisms in the Factor H gene listed in TABLE 1A, TABLE 1B and/or TABLE 1C, alone or in combination with additional Factor H gene polymorphisms not included in TABLES 1A-1C, may increase the probability of an accurate diagnosis. Similarly, combined detection of several polymorphic forms in the CFHR5 gene, for example, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or all of the polymorphisms in the CFHR5 gene listed in TABLES 14 and 15, alone or in combination with additional CFHR5 gene polymorphisms not included in TABLES 14 and 15, may increase the probability of an accurate diagnosis. In one embodiment, screening involves determining the presence or absence of at least one polymorphism in the Factor H gene and at least one polymorphism in the CFHR5 gene. In one embodiment, screening involves determining the presence or absence of at least 2, 3, or 4 polymorphisms in the Factor H gene in combination with and at least 2, 3, or 4 polymorphisms in the CFHR5 gene.

[0200] The polymorphisms in the Factor H and CFHR5 genes are useful in diagnosing AMD or specific subtypes of AMD, or susceptibility thereto, in family members of patients with AMD, as well as in the general population.

[0201] In diagnostic methods, analysis of Factor H polymorphisms and/or CFHR5 polymorphisms can be combined with analysis of polymorphisms in other genes associated with AMD, detection of protein markers of AMD (see, e.g., Hageman et al., patent publications US20030017501; US20020102581; WO0184149; and WO0106262), assessment of other risk factors of AMD (such as family history), with ophthalmological examination, and with other assays and procedures.

E) Identification of Patients for Drug Therapy

[0202] Polymorphisms in the Factor H gene and CFHR5 gene are also useful for identifying suitable patients for conducting clinical trials for drug candidates for AMD. Such trials are performed on treated or control populations having similar or identical polymorphic profiles at a defined collection of polymorphic sites in the Factor H gene and/or CFHR5 gene, or having similar or identical Factor H haplotypes and/or CFHR5 haplotypes. The use of genetically matched populations eliminates or reduces variation in treatment outcome due to genetic factors, leading to a more accurate assessment of the efficacy of a potential drug.

F) Screening Donor Tissue for Transplantation

[0203] Transplantation of organs (e.g., liver) and tissues (e.g., blood, hepatocytes) is increasingly common. It is desirable, in carrying out such transplantation, to avoid introducing into the recipient a deleterious form of Factor H or a Factor H-Related Protein and thereby increasing the recipient's risk of developing AMD. Thus, in one aspect of the invention, a donor tissue is tested to detect the presence or absence of a variation at a polymorphic site of a Factor H or CFHR5 gene to identify host tissues carrying risk haplotypes or other deleterious sequences. In addition or alternatively, organs and tissues can be tested for the expression of forms of Factor H or CFHR proteins, for example by using immunological assays as described herein. In one embodiment the transplanted tissue is blood or plasma (i.e., given in a blood transfusion or plasma replacement). Routine screening of donated blood to avoid administration of a protein associated with risk (e.g., the 1210C form of CFH) may avoid compromising the recipient.

G) Phenotypic Categories

[0204] Susceptibility to specific subtypes of AMD can be identified based on the association with particular haplotypes. Thus, the screening can be used to determine suitable therapies for groups of patients with different genetic subtypes of AMD.

[0205] The methods may be used for the diagnosis of AMD, which may be subdivided into phenotypic categories (for example, early AMD (ARM) geographic atrophy (GA) and exudative AMD (CNV)). The ARM and GA phenotypes may be further subdivided into distinct phenotypes (for example, RPE changes alone, >10 macular hard drusen, macular soft drusen, BB (cuticular) drusen, pigment epithelial detachment (PED), "Cherokee" atrophy, peninsular geographic atrophy and pattern geographic atrophy). For descriptions of these phenotypes see, e.g., Bird et al., 1995, *Surv Ophthalmol* 39, 367-74; and Klaver et al., 2001, *Invest Ophthalmol Vis Sci* 42, 2237-41.

H) Other Diseases

[0206] Polymorphisms in the Factor H and CFHR5 gene, such as those shown in TABLES 1A, 1B, 1C, 11, 14 and 15, can also be tested for association with other diseases, (for example, Alzheimer's disease, multiple sclerosis, lupus, and asthma) and conditions (for example, burn injuries, transplantation, and stroke) which involve dysregulation of the alternative complement pathway, that have known but hitherto unmapped genetic

components. Without being limited to any particular mechanism of action, it is suggested herein that expression of variant Factor H and/or CFHR5 polypeptides is associated with dysregulation of the alternative complement pathway. The variant forms of Factor H and/or CHFR5 may have a causal effect on diseases involving a defect in the alternative complement pathway, or the presence of variant forms of Factor H and/or CFHR5 may indicate that another gene involved in the alternative complement pathway has a causal effect.

[0207] Polymorphisms in the Factor H gene may also be useful in mapping and treating diseases that map to chromosome 1q, in particular at or near 1q32 where the Factor H gene is located. This particular locus contains a number of complement pathway-associated genes. One group of these genes, referred to as the regulators of complement activation (RCA) gene cluster, contains the genes that encode Factor H, five Factor H-related genes and the beta subunit of coagulation factor XIII. A second cluster of complement-associated genes, including C4BPA, C4BPB, C4BPAL2, DAF (CD55) CR1, CR2, CR1L and MCP (CD46) lies immediately adjacent to the 1q25-31 locus.

VIII. PREVENTION AND TREATMENT OF AMD

[0208] A patient with a Factor H polymorphism can be treated by administering to a patient an antagonist of the variant Factor H polypeptide and/or variant CHFR5 polypeptide. An antagonist may include a therapeutic amount of an RNA complementary to the nucleotide sequence of a variant Factor H polypeptide and/or variant CHFR5 polypeptide or an antibody that specifically interacts with and neutralizes the activity of a variant Factor H polypeptide and/or variant CHFR5 polypeptide. Alternatively, AMD associated with the Factor H polymorphism and/or CFHR5 polymorphism can be treated by administering to a patient a form of Factor H and/or CHFR5 not associated with increased risk, such as the normal or wild-type Factor H protein and/or normal or wild-type CHFR5 polypeptide. In one method of the invention, a protective variant form of Factor H and/or protective variant form of CHFR5 is administered to a patient.

[0209] Therapeutic and prophylactic approaches in subjects identified as being at high risk for AMD include, but are not limited to, (1) increasing the amount or expression of neutral or protective forms of Factor H and/or neutral or protective forms of CHFR5; (2) decreasing the amount or expression of risk-associated forms of Factor H and/or risk-associated form of CHFR5; and (3) reducing activation of the complement alternative pathway. Examples of such therapeutic and prophylactic approaches include: (1) administration of neutral or

protective forms of Factor H protein or therapeutically active fragments and/or neutral or protective forms of CFHR5 or therapeutically active fragments; (2) other wise increasing expression or activity of neutral and protective forms of Factor H; (3) interfering with expression of variant Factor H and/or variant CFHR5 proteins encoded by individuals with a risk haplotype by (e.g., by administration of antisense RNA); (4) reducing the amount of activity of a detrimental variant form.

[0210] Therapeutic agents (e.g., agents that increase or decrease levels of wild-type or variant Factor H or modulate its activity and/or agents that increase or decrease levels of wild-type or variant CFHR5 or modulate its activity) can be administered systemically (e.g., by i.v. injection or infusion) or locally (e.g., to the vicinity of the ocular RPE for treatment of AMD). Methods for administration of agents to the eye are well known in the medical arts and can be used to administer AMD therapeutics described herein. Exemplary methods include intraocular injection (e.g., retrobulbar, subretinal, intravitreal and intrachoroidal), iontophoresis, eye drops, and intraocular implantation (e.g., intravitreal, sub-Tenons and sub-conjunctival). For examples, anti-VEGF antibody has been introduced into cynomolgus monkeys by intravitreal injection (see, e.g., Gaudreault et al., 2005, "Preclinical pharmacokinetics of Ranibizumab (rhuFabV2) after a single intravitreal administration" *Invest Ophthalmol Vis Sci.* 46:726-33), and bioactive VEGF and bFGF have been expressed in the eye via intravitreal implantation of sustained release pellets (Wong et al., 2001, "Intravitreal VEGF and bFGF produce florid retinal neovascularization and hemorrhage in the rabbit" *Curr Eye Res.* 22:140-7). Importantly, it has been discovered that Factor H is synthesized locally by the retinal pigment epithelium (see Example 1), indicating that local administration of agents has therapeutic benefit.

A. Administration of Therapeutic Factor H Polypeptides

[0211] Administration of neutral or protective forms of Factor H polypeptides and/or neutral or protective forms of CFHR5 polypeptides to subjects at risk for developing AMD (and/or with early stage disease) can be used to ameliorate the progression of the disease.

[0212] In one approach, recombinant Factor H polypeptide is administered to the patient. In one embodiment, the recombinant Factor H is encoded by a neutral haplotype sequence, which may be full-length (CFH/HF1), truncated (FHL1), or alternatively spliced form, or a biologically active fragment thereof. In another embodiment the recombinant Factor H has the sequence of a protective allele, either full-length or truncated form, or a protective

biologically active fragment thereof. Methods for production of therapeutic recombinant proteins are well known and include methods described hereinbelow. The therapeutic polypeptide can be administered systemically (e.g., intravenously or by infusion) or locally (e.g., directly to an organ or tissue, such as the eye or the liver).

[0213] Some protective forms of Factor H and the CHFL1 protein are less than full-length. For example, fragments of neutral or protective forms of Factor H may be administered for treatment or prevention of AMD or MPGNII. In a particular embodiment, polypeptides encoded by CFH splice variants expressed in individuals with a protective phenotype are administered. These proteins can be identified by screening expression of CFH-related RNA in individuals homozygous for a protective or neutral haplotype.

[0214] In particular embodiments, the protective protein has a sequence corresponding to one or more exons of the CFH gene sequence. For example, the protective protein may have the sequence of full-length or truncated CFH protein, except that the amino acid residues encoded by 1, 2, 3 or more exons (which may or may not be contiguous) are deleted.

[0215] In one embodiment a protective Factor H protein of the invention has an amino acid sequence substantially identical to SEQ ID NO:2, with the proviso that the residue at position 402 is not histidine and the residue at position 1210 is not cysteine. In one embodiment the residue at position 62 is not valine. Preferably, the residue at position 62 is isoleucine. Preferably the residue at position 62 is isoleucine, the residue at position 402 is tyrosine and the residue at position 1210 is arginine. Preferably the protective Factor H protein has 95% amino acid identity to SEQ ID NO:2 or a fragment thereof; sometimes at least 95% amino acid identity, sometimes at least 98% amino acid identity, and sometimes at least 99% identity to the reference Factor H polypeptide of SEQ ID NO:2. The polypeptide sequence of an exemplary protective variant of human Factor H [SEQ ID NO:5] is shown in FIGURE 10. This protective variant Factor H polypeptide has an isoleucine at amino acid position 62 and a tyrosine at amino acid position 402 (indicated in bold). The polypeptide sequence of an exemplary protective variant of HFL1, the truncated form of human Factor H (SEQ ID NO:6) is shown in FIGURE 11. This protective variant truncated Factor H polypeptide has a isoleucine at amino acid position 62 and tyrosine at amino acid position 402 (indicated in bold).

[0216] In one embodiment a protective Factor H protein of the invention has an amino acid sequence substantially identical to SEQ ID NO:4 (FHL1). In one embodiment the residue at position 62 is not valine. Preferably, the residue at position 62 is isoleucine. Preferably the

protective Factor H protein has 95% amino acid identity to SEQ ID NO:4 or a fragment thereof; sometimes at least 95% amino acid identity, sometimes at least 98% amino acid identity, and sometimes at least 99% identity to the reference Factor H polypeptide of SEQ ID NO:4.

[0217] In some embodiments the protective Factor H protein has one or more activities of the reference Factor H polypeptide. In one embodiment the activity is binding to heparin. In one embodiment the activity is binding to CRP. In one embodiment the activity is binding to C3b. In one embodiment the activity is binding to endothelial cell surfaces. In one embodiment the activity is C3b co-factor activity. In one embodiment, the protective Factor H protein has activity that is higher with respect to its normal function than the protein of SEQ ID NO:2. In one embodiment, the protective Factor H protein has activity with respect to its normal function that is higher than the protein of SEQ ID NO:4.

[0218] Assays for Factor H activities are well known and described in the scientific literature. For illustration and not limitation, examples of assays will be described briefly.

Binding of Protective Proteins (CFH Variants) to C3b or CRP.

[0219] Interactions between C3b and CFH proteins can be analyzed by surface resonance using a Biacore 3000 system (Biacore AB, Uppsala, Sweden), as described previously (Manuelian et al., 2003, Mutations in factor H reduce binding affinity to C3b and heparin and surface attachment to endothelial cells in hemolytic uremic syndrome. *J Clin Invest* 111, 1181-90). In brief, C3b (CalBiochem, Inc), are coupled using standard amine-coupling to flow cells of a sensor chip (Carboxylated Dextran Chip CM5, Biacore AB, Uppsala, Sweden). Two cells are activated and C3b (50 µg/ml, dialyzed against 10 mM acetate buffer, pH 5.0) is injected into one flow cell until a level of coupling corresponding to 4000 resonance units is reached. Unreacted groups are inactivated using ethanolamine-HCl. The other cell is prepared as a reference cell by injecting the coupling buffer without C3b. Before each binding assay, flow cells will be washed thoroughly by two injections of 2 M NaCl in 10 mM acetate buffer, pH 4.6 and running buffer (PBS, pH 7.4). The Factor H protein is injected into the flow cell coupled with C3b or into the control cell at a flow rate of 5 µl/min at 25°C. Binding of Factor H to C3b is quantified by measuring resonance units over time, as described in Manuelian et al., 2003, *supra*.

[0220] Interactions between CRP and CHF proteins can be analyzed by surface resonance in an identical manner by substituting CRP for C3b in flow cells of a sensor chip

Binding to Endothelial Cell Surface

[0221] Binding of CHF proteins to endothelial cell surfaces is assayed by immunofluorescence staining of HUVECs and FACS analysis. HUVEC cells are kept in serum free DMEM (BioWhittaker) for 24 hrs prior to the assay. Cells are detached from the surface with DPBS/EDTA and washed twice with DPBS; 5×10^5 cells will be transferred into plastic tubes and unspecific binding sites will be blocked with 1 % BSA/DPBS for 15 min prior to incubation with purified allele variants of factor H (5 μ g). Controls are performed in the absence of the factor H isoform. Following binding of factor H, cells are thoroughly washed with DPBS. Polyclonal goat anti-human FH antiserum is used as a primary antibody (CalBiochem) (diluted 1:100), incubating cells at 4°C for 15 minutes. Alexa-fluor 488-conjugated goat antiserum diluted 1:100 in blocking buffer is used as the secondary antibody. Cells are examined by flow cytometry (FACScalibur, Becton-Dickinson Immunocytometry, Mountain View, California, USA). Typically, 10,000 events are counted.

Cofactor Activity in Fluid Phase

[0222] For the fluid phase cofactor assay, C3b biotin (100 ng/reaction), Factor I (200 ng/reaction) and 100 ng of purified factor H are used in a total volume of 30 μ l. Samples taken before and after addition of Factor I are separated by SDS-PAGE under reducing conditions and analyzed by Western blotting, detecting and quantitating C3b degradation products by Streptavidin-POD-conjugation (1:10000). C3b (40 μ g) (CalBiochem) is biotinylated using the Biotin Labeling Kit (Roche Diagnostics, Mannheim, Germany), according to the manufacturer's instructions. In brief, 30 μ g of C3b (CalBiochem) is labeled with D-biotinyl-epsilon-aminocaproic acid-N-hydroxysuccinimide ester for 2 hours at 25°C. Excess biotin is removed by gel filtration using a PBS equilibrated PD10 column (Amersham Biosciences). Also see Sanchez-Corral et al., 2002, *Am J. Hum. Genet.* 71:1285-95.

Heparin Binding Assay

[0223] Binding of purified CFH proteins (CFH402Y and CFH402H) to heparin is analyzed using heparin affinity chromatography in a high-performance liquid chromatograph (HPLC) system. 10 μ g of CFH protein is diluted in 1/2xPBS and applied to a heparin-Sepharose affinity column (HiTrap, Amersham Biosciences) at a flow rate of 0.5 ml/min. The column is

extensively washed with 1/2xPBS, and the bound CFH protein eluted using a linear salt gradient ranging from 75 to 500 mM NaCl, in a total volume of 10 ml and at a flow rate of 0.5 ml/min. Eluted fractions are assayed by SDS-PAGE and Western blot analysis. Elution of isoforms in different fractions is indicative that specific amino acid variations in the CFH protein can modulate binding of the protein to heparin. Also see, e.g., Pangburn et al., 1991, Localization of the heparin-binding site on complement Factor H, *J Biol Chem.* 266:16847-53.

CFHR5 Administration

[0224] In another approach, recombinant CFHR5 polypeptide is administered to the patient. In one embodiment, the recombinant CFHR5 has a neutral-type sequence, or a biologically active fragment thereof. In another embodiment the recombinant CFHR5 has the sequence of a protective allele, or a protective biologically active fragment thereof. Methods for production of therapeutic recombinant proteins are well known and include methods described hereinbelow. The therapeutic polypeptide can be administered systemically (e.g., intravenously or by infusion) or locally (e.g., directly to an organ or tissue, such as the eye or the liver).

Therapeutic Compositions Containing CFH or CFHR5 Polypeptides

[0225] The invention provides therapeutic preparations of Factor H polypeptides, which may be wild-type or variants (e.g., neutral or protective variants), and may be full length forms, truncated forms, or biologically active fragments of the variant Factor H polypeptides. As described herein, protective Factor H proteins (and genes encoding them) can be identified by identifying an individual as having a protective haplotype and determining the amino acid sequence(s) of Factor H encoded in the genome of the individual, where a protective Factor H protein is encoded by an allele having a protective haplotype. Biologically active fragments may include any portion of the full-length Factor H polypeptide which confers a biological function on the variant protein. In some cases, a protective haplotype will be associated with expression of a less-than full length form of Factor H (i.e., in addition to FHL-1) due, for example, to the presense of a premature stop codon in the gene.

[0226] Therapeutically active fragments can also be identified by testing the effect of the protein on expression of AMD biomarkers. Exemplary AMD biomarkers include

complement pathway components (for example, Factor I, Factor H, C1r, C3, C3a), C reactive protein, haptoglobin, apolipoprotein E, immunoglobulin heavy or light chain(s), alpha 1 antitrypsin, alpha 2 macroglobulin, transthyretin, creatinine, and others described in copending provisional application No. 60/715,503 entitled "Biomarkers Associated With Age-Related Macular Degeneration."

[0227] The invention provides therapeutic preparations of CFHR5 polypeptides, which may be wild-type or variants (e.g., neutral or protective variants), and may be full length forms or biologically active fragments of the variant CFHR5 polypeptides. As described herein, protective CFHR5 proteins (and genes encoding them) can be identified by identifying an individual as having a protective haplotype and determining the amino acid sequence(s) of CFHR5 encoded in the genome of the individual, where a protective CFHR5 protein is encoded by an allele having a protective haplotype. Biologically active fragments may include any portion of the full-length CFHR5 polypeptide which confers a biological function on the variant protein. Therapeutically active fragments can also be identified by testing the effect of the protein on expression of AMD biomarkers as described above for Factor H.

[0228] Some forms of Factor H and CFHR5 can be isolated from the blood of genotyped donors, from cultured or transformed RPE cells derived from genotyped ocular donors, or from cell lines (e.g., glial or hepatic) that express endogenous Factor H. Alternatively, therapeutic proteins can be recombinantly produced (e.g., in cultured bacterial or eukaryotic cells) and purified using methods well known in the art and described herein. As noted above, some forms of Factor H and CFHR5 have been recombinantly expressed for research purposes. However, such research preparations are not suitable for therapeutic use. The present invention provides recombinant polypeptides suitable for administration to patients including polypeptides that are produced and tested in compliance with the Good Manufacturing Practice (GMP) requirements. For example, recombinant polypeptides subject to FDA approval must be tested for potency and identity, be sterile, be free of extraneous material, and all ingredients in a product (i.e., preservatives, diluents, adjuvants, and the like) must meet standards of purity, quality, and not be deleterious to the patient.

[0229] The invention provides a composition comprising a Factor H polypeptide or CFHR5 polypeptide, and a pharmaceutically acceptable excipient or carrier. The term "pharmaceutically acceptable excipient or carrier" refers to a medium that is used to prepare a desired dosage form of a compound. A pharmaceutically acceptable excipient or carrier can

include one or more solvents, diluents, or other liquid vehicles; dispersion or suspension aids; surface active agents; isotonic agents; thickening or emulsifying agents; preservatives; solid binders; lubricants; and the like. Remington's Pharmaceutical Sciences, Fifteenth Edition, E.W. Martin (Mack Publishing Co., Easton, PA, 1975) and Handbook of Pharmaceutical Excipients, Third Edition, A.H. Kibbe ed. (American Pharmaceutical Assoc. 2000), disclose various carriers used in formulating pharmaceutical compositions and known techniques for the preparation thereof. In one embodiment, the pharmaceutically acceptable excipient is not deleterious to a mammal (e.g., human patient) if administered to the eye (e.g., by intraocular injection). For intraocular administration, for example and not limitation, the therapeutic agent can be administered in a Balanced Salt Solution (BSS) or Balanced Salt Solution Plus (BSS Plus) (Alcon Laboratories, Fort Worth, Texas, USA). In a related aspect, the invention provides a sterile container, e.g. vial, containing a therapeutically acceptable Factor H protein, optionally a lyophilized preparation. Therapeutic Factor H proteins or CFHR5 polypeptides can be made recombinantly, as described above. Alternatively, Factor H protein or CFHR5 polypeptide can be isolated from cultured RPE cells (e.g., primary cultures) or other cells that express Factor H or CFHR5 endogenously.

[0230] The amount of neutral or protective forms of Factor H or truncated Factor H, or biologically active fragments thereof, or neutral or protective forms of CFHR5, or biologically active fragments thereof, to be administered to an individual can be determined. The normal plasma concentration of Factor H varies between 116 and 562 micrograms/ml and the half-life of Factor H in plasma is about 6 1/2 days (for a recent review, see Esparza-Gordillo et al., 2004 "Genetic and environmental factors influencing the human factor H plasma levels" *Immunogenetics* 56:77-82). In one embodiment, exogenous Factor H can be administered to an individual in an amount sufficient to achieve a level similar to the plasma concentration of Factor H in a healthy individual, i.e., an amount sufficient to achieve a plasma level of from 50 to 600 mg/ml, such as from 100 to 560 mg/ml. The amount of Factor H to be administered to an individual (e.g., a 160 pound subject) can be, for example and not for limitation, from 10 milligrams to 5000 milligrams per dose, from 50 milligrams to 2000 milligrams per dose, from 100 milligrams to 1500 milligrams per dose, from 200 milligrams to 1000 milligrams per dose, or from 250 milligrams to 750 milligrams per dose. The frequency with which Factor H can be administered to an individual can be, for example and not for limitation, twice per day, once per day, twice per week, once per week, once every two weeks, once per month, once every two months, once every six months, or once

per year. The amount and frequency of administration of Factor H to an individual can be readily determined by a physician by monitoring the course of treatment.

B) Gene Therapy Methods

[0231] In another approach, Factor H protein or CFHR5 polypeptide is administered by in vivo expression of protein encoded by exogenous polynucleotide (i.e., via gene therapy). In one example, gene therapy involves introducing into a cell a vector that expresses Factor H polypeptide or biologically active fragment or CFHR5 polypeptide or biologically active fragment.

[0232] Vectors can be viral or nonviral. A number of vectors derived from animal viruses are available, including those derived from adenovirus, adeno-associated virus, retroviruses, pox viruses, alpha viruses, rhabdoviruses, and papillomaviruses. Usually the viruses have been attenuated to no longer replicate (see, e.g., Kay et al. 2001, *Nature Medicine* 7:33-40).

[0233] The nucleic acid encoding the Factor H polypeptide or CFHR5 polypeptide is typically linked to regulatory elements, such as a promoters and an enhancers, which drive transcription of the DNA in the target cells of an individual. The promoter may drive expression of the Factor H gene or CFHR5 gene in all cell types. Alternatively, the promoter may drive expression of the Factor H gene or CFHR5 gene only in specific cell types, for example, in cells of the retina or the kidney. The regulatory elements, operably linked to the nucleic acid encoding the Factor H polypeptide or CFHR5 polypeptide, are often cloned into a vector.

[0234] As will be understood by those of skill in the art, gene therapy vectors contain the necessary elements for the transcription and translation of the inserted coding sequence (and may include, for example, a promoter, an enhancer, other regulatory elements). Promoters can be constitutive or inducible. Promoters can be selected to target preferential gene expression in a target tissue, such as the RPE (for recent reviews see Sutanto et al., 2005, "Development and evaluation of the specificity of a cathepsin D proximal promoter in the eye" *Curr Eye Res.* 30:53-61; Zhang et al., 2004, "Concurrent enhancement of transcriptional activity and specificity of a retinal pigment epithelial cell-preferential promoter" *Mol Vis.* 10:208-14; Esumi et al., 2004, "Analysis of the VMD2 promoter and implication of E-box binding factors in its regulation" *J Biol Chem* 279:19064-73; Camacho-Hubner et al., 2000, "The Fugu rubripes tyrosinase gene promoter targets transgene expression to pigment cells in the mouse" *Genesis.* 28:99-105; and references therein).

[0235] Suitable viral vectors include DNA virus vectors (such as adenoviral vectors, adeno-associated virus vectors, lentivirus vectors, and vaccinia virus vectors), and RNA virus vectors (such as retroviral vectors). In one embodiment, an adeno-associated viral (AAV) vector is used. For recent reviews see Auricchio et al., 2005, "Adeno-associated viral vectors for retinal gene transfer and treatment of retinal diseases" *Curr Gene Ther.* 5:339-48; Martin et al., 2004, Gene therapy for optic nerve disease, *Eye* 18:1049-55; Ali, 2004, "Prospects for gene therapy" *Novartis Found Symp.* 255:165-72; Hennig et al., 2004, "AAV-mediated intravitreal gene therapy reduces lysosomal storage in the retinal pigmented epithelium and improves retinal function in adult MPS VII mice" *Mol Ther.* 10:106-16; Smith et al., 2003, "AAV-Mediated gene transfer slows photoreceptor loss in the RCS rat model of retinitis pigmentosa" *Mol Ther.* 8:188-95; Broderick et al., 2005, "Local administration of an adeno-associated viral vector expressing IL-10 reduces monocyte infiltration and subsequent photoreceptor damage during experimental autoimmune uveitis" *Mol Ther.* 12:369-73; Cheng et al., 2005, "Efficient gene transfer to retinal pigment epithelium cells with long-term expression. *Retina* 25:193-201; Rex et al., "Adenovirus-mediated delivery of catalase to retinal pigment epithelial cells protects neighboring photoreceptors from photo-oxidative stress. *Hum Gene Ther.* 15:960-7; and references cited therein).

[0236] Gene therapy vectors must be produced in compliance with the Good Manufacturing Practice (GMP) requirements rendering the product suitable for administration to patients. The present invention provides gene therapy vectors suitable for administration to patients including gene therapy vectors that are produced and tested in compliance with the GMP requirements. Gene therapy vectors subject to FDA approval must be tested for potency and identity, be sterile, be free of extraneous material, and all ingredients in a product (i.e., preservatives, diluents, adjuvants, and the like) must meet standards of purity, quality, and not be deleterious to the patient. For example, the nucleic acid preparation is demonstrated to be mycoplasma-free. See, e.g, Islam et al., 1997, An academic centre for gene therapy research and clinical grade manufacturing capability, *Ann Med* 29, 579-583.

[0237] Methods for administering gene therapy vectors are known. In one embodiment, Factor H or CFHR5 expression vectors are introduced systemically (e.g., intravenously or by infusion). In one embodiment, Factor H or CFHR5 expression vectors are introduced locally (i.e., directly to a particular tissue or organ, e.g., liver). In one preferred embodiment, Factor H or CFHR5 expression vectors are introduced directly into the eye (e.g., by ocular

injection). For recent reviews see, e.g., Dinculescu et al., 2005, "Adeno-associated virus-vectored gene therapy for retinal disease" *Hum Gene Ther.* 16:649-63; Rex et al., 2004, "Adenovirus-mediated delivery of catalase to retinal pigment epithelial cells protects neighboring photoreceptors from photo-oxidative stress" *Hum Gene Ther.* 15:960-7; Bennett, 2004, "Gene therapy for Leber congenital amaurosis" *Novartis Found Symp.* 255:195-202; Hauswirth et al., "Range of retinal diseases potentially treatable by AAV-vectored gene therapy" *Novartis Found Symp.* 255:179-188, and references cited therein).

[0238] Thus in one aspect, the invention provides a preparation comprising a gene therapy vector encoding a Factor H protein or CFHR5 polypeptide, optionally a viral vector, where the gene therapy vector is suitable for administration to a human subject and in an excipient suitable for administration to a human subject (e.g., produced using GLP techniques). Optionally the gene therapy vector comprising a promoter that is expressed preferentially or specifically in retinal pigmented epithelium cells.

[0239] Nonviral methods for introduction of Factor H or CFHR5 sequences, such as encapsulation in biodegradable polymers (e.g., polylactic acid (PLA); polyglycolic acid (PGA); and co-polymers (PLGA) can also be used (for recent reviews see, e.g., Bejjani et al., 2005, "Nanoparticles for gene delivery to retinal pigment epithelial cells" *Mol Vis.* 11:124-32; Mannermaa et al., 2005, "Long-lasting secretion of transgene product from differentiated and filter-grown retinal pigment epithelial cells after nonviral gene transfer" *Curr Eye Res.* 2005 30:345-53; and references cited therein). Alternatively, the nucleic acid encoding a Factor H polypeptide or CFHR5 polypeptide may be packaged into liposomes, or the nucleic acid can be delivered to an individual without packaging without using a vector.

C) DNA Repair

[0240] In another approach, subjects at risk for developing AMD (and/or with early stage disease) can have a risk form of Factor H or CFHR5 replaced by a neutral or protective form of Factor H or CFHR5 by DNA repair. In one embodiment, triplex forming oligonucleotides designed to specifically bind to polymorphic sites in the Factor H or CFHR5 gene associated with a risk haplotype can be administered to an individual by viral or nonviral methods. Triplex-forming oligonucleotides bind to the major groove of duplex DNA in a sequence-specific manner and provoke DNA repair, resulting in the targeted modification of the genome (for a recent review see Kuan et al., 2004, "Targeted gene modification using triplex-forming oligonucleotides" *Methods Mol Biol.* 262:173-94). A triplex-forming

oligonucleotide that binds to a sequence spanning a polymorphism associated with a risk haplotype provokes DNA repair, resulting in the modification of the sequence from a risk allele to a neutral or protective allele and can ameliorate the development or progression of disease.

D) Introduction of Cells, Tissues, or Organs Expressing a Neutral or Protective Form of Factor H Protein or CFHR5 Polypeptide

[0241] In another approach, cells expressing neutral or protective forms of Factor H or Factor H-Related proteins (e.g., CFHR5) are administered to a patient. In an embodiment the recipient is heterozygous or, more often, homozygous for a risk haplotype. For example, hepatocyte transplantation has been used as an alternative to whole-organ transplantation to support many forms of hepatic insufficiency (see, e.g., Ohashi et al., Hepatocyte transplantation: clinical and experimental application, *J Mol Med.* 2001 79:617-30). According to this method, hepatocytes or other CFH or CFHR5-expressing cells are administered (e.g., infused) to a patient in need of treatment. These cells migrate to the liver or other organ, and produce the therapeutic protein. Also see, e.g., Alexandrova et al., 2005, "Large-scale isolation of human hepatocytes for therapeutic application" *Cell Transplant.* 14(10):845-53; Cheong et al., 2004, "Attempted treatment of factor H deficiency by liver transplantation" *Pediatr Nephrol.* 19:454-8; Ohashi et al., 2001, "Hepatocyte transplantation: clinical and experimental application" *J Mol Med.* 79:617-30; Serralta et al., 2005, "Influence of preservation solution on the isolation and culture of human hepatocytes from liver grafts" *Cell Transplant.* 14(10):837-43; Yokoyama et al., 2006, "In vivo engineering of metabolically active hepatic tissues in a neovascularized subcutaneous cavity" *Am. J. Transplant.* 6(1):50-9; Dhawan et al., 2005, "Hepatocyte transplantation for metabolic disorders, experience at King's College hospital and review of literature." *Acta Gastroenterol. Belg.* 68(4):457-60; Bruns et al., 2005, "Injectable liver: a novel approach using fibrin gel as a matrix for culture and intrahepatic transplantation of hepatocytes" *Tissue Eng.* 11(11-12):1718-26. Other cell types that may be used include, for illustration and not limitation, kidney and pancreatic cells. In one embodiment, the administered cells are engineered to express a recombinant form of the protein.

[0242] In another, related approach, therapeutic organ transplantation is used. Most of the body's systemic Factor H is produced by the liver, making transplation of liver tissue the

preferred method. See, Gerber et al., 2003, "Successful (?) therapy of hemolytic-uremic syndrome with factor H abnormality" *Pediatr Nephrol.* 18:952-5.

[0243] In another approach, a protective form of CFH protein is delivered to the back of the eye by injection into the eye (e.g. intravitreal) or via encapsulated cells. Neurotech's Encapsulated Cell Technology (ECT), as an example, is a unique technology that allows for the sustained, longterm delivery of therapeutic factors to the back of the eye. See (<http://www.neurotech.fr>). ECT implants consist of cells that have been genetically modified to produce a specific therapeutic protein that are encapsulated in a semi-permeable hollow fiber membrane. The cells continuously produce the therapeutic protein that diffuses out of the implant and into the eye (Bush et al 2004). CNTF delivered to the human eye by ECT devices was recently shown to be completely successful and associated with minimal complications in 10 patients enrolled in a Phase I clinical trial (Sieving et al 2005). Also see Song et al., 2003; Tao 2002., and Hammang et al., U.S. Pat. No. 6,649,184. In one embodiment of the present invention, a protective form of Factor H (including the so-called neutral form) is expressed in cells and administered in an encapsulated form. In one embodiment, the cells used are the NTC-201 human RPE line (ATCC # CRL-2302) available from the American Type Culture Collection P.O. Box 1549, Manassas, VA 20108.

E) Therapy to Decrease Levels of Risk Variant of Factor H or CFHR5

[0244] Loss of the normal or protective function of Factor H or CFHR5 may be associated with AMD. Non-synonymous polymorphisms in the Factor H and CFHR5 genes, such as those shown in TABLES 1A, 1B, 1C, 11, 14 and 15, showing the strongest correlation with AMD and resulting in a variant Factor H polypeptide or CFHR5 polypeptide, are likely to have a causative role in AMD. For example, the variant Factor H or CFHR5 may act as a so-called "dominant-negative" mutant interfering with normal Factor H or CFHR5 function.

[0245] Any method of reducing levels of the risk forms of Factor H or CFHR5 in the eye or systemically may be used for treatment including, for example, inhibiting transcription of a Factor H or CFHR5 gene, inhibiting translation of Factor H or CFHR5 RNA, increasing the amount or activity of a neutral or protective form of Factor H or truncated Factor H, or biologically active fragment thereof, increasing the amount or activity of a neutral or protective form of CFHR5 polypeptide, or biologically active fragment thereof, or decreasing the amount or activity of Factor H protein or CFHR5 polypeptides (e.g., by plasmaphoresis, antibody-directed plasmaphoresis, or complexing with a Factor H or CFHR5 binding moiety,

e.g., heparin or variant specific antibody). In some embodiments levels of Factor H or CFHR5 are preferentially reduced in the eye (e.g., RPE) relative to other tissues. For illustration and not limitation, several methods are briefly described below.

[0246] In one approach, a subject identified as being at risk for AMD is treated by administration of heparin. Heparin and heparin derivatives (including heparinoids) may have promising therapeutic properties for the treatment various complement-associated diseases, including MPGNII (Floege et al., 1993; Girardi, 2005; Diamond and Karnovsky, 1986; Striker, 1999; Rops et al., 2004). In view of the association between AMD and MPGNII disclosed herein, heparin and heparin derivatives (including heparinoids) may be efficacious for the treatment of AMD. In a clinical trial of patients with chronic proliferative glomerulonephritis receiving daily subcutaneous injections of heparin for over one year, Cade and colleagues reported improved creatinine clearance and a regression of glomerular hypercellularity (Cade et al., 1971). Both heparin and low molecular weight heparin (Enoxaparin) have been shown to prevent the progression of antiphospholipid antibody syndrome in mice by blocking the alternative and classical pathways of the complement cascade (Girardi et al., 2004). The anti-complement activity of heparin includes blockade of the formation of C3bBb, the amplification convertase by the alternative pathway; fluid phase heparin prevents the generation of C3bBb by inhibiting the interaction of C3b with factor B and factor D (Weiler et al., 1976).

F) Administration of Inhibitory Nucleic Acids

[0247] *Antisense nucleic acids* -Antisense nucleic acids, such as purified anti-sense RNA complementary to the RNA encoding a variant Factor H polypeptide can be used to inhibit expression of a Factor H gene associated with a risk haplotype. For recent reviews see, e.g., Gomes et al., 2005, "Intraocular delivery of oligonucleotides" *Curr Pharm Biotechnol.* 6:7-15; and Henry et al., 2004, "Setting sights on the treatment of ocular angiogenesis using antisense oligonucleotides" *Trends Pharmacol Sci* 25:523-7; and references cited therein.

[0248] *RNA Interference*- Double stranded RNA (dsRNA) inhibition methods can also be used to inhibit expression of HF1. The RNA used in such methods is designed such that at least a region of the dsRNA is substantially identical to a region of the HF1 gene; in some instances, the region is 100% identical to the HF1 gene. For use in mammals, the dsRNA is typically about 19-30 nucleotides in length (i.e., short interfering RNAs are used (siRNA or RNAi)), and most often about 21 nucleotides in length. Methods and compositions useful for

performing dsRNAi and siRNA are discussed, for example, in PCT Publications WO 98/53083; WO 99/32619; WO 99/53050; WO 00/44914; WO 01/36646; WO 01/75164; WO 02/44321; and U.S. Patent No. 6,107,094. siRNA can be synthesized in vitro and administered to a patient. Alternatively, RNAi strategies can be successfully combined with vector-based approaches to achieve synthesis in transfected cells of small RNAs from a DNA template (see, e.g., Sui et al., 2002, "A DNA vector-based RNAi technology to suppress gene expression in mammalian cells" *Proc Natl Acad Sci USA* 99:5515-20; and Kasahara and Aoki, 2005, "Gene silencing using adenoviral RNAi vector in vascular smooth muscle cells and cardiomyocytes" *Methods Mol Med* 112:155-72; and references cited therein).

[0249] *Ribozymes* - Ribozymes are enzymatic RNA molecules capable of catalyzing the specific cleavage of RNA. The mechanism of ribozyme action involves sequence-specific hybridization of the ribozyme molecule to complementary target RNA, followed by endonucleolytic cleavage. Within the scope of the invention are engineered hammerhead motif ribozyme molecules that can specifically and efficiently catalyze endonucleolytic cleavage of the sequence encoding human Factor H. Specific ribozyme cleavage sites within any potential RNA target are initially identified by scanning the target molecule for ribozyme cleavage sites which include the sequences such as, GUA, GUU and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides corresponding to the region of the target gene containing the cleavage site may be evaluated for secondary structural features which may render the oligonucleotide inoperable. The suitability of candidate targets may also be evaluated by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays. Properties of ribozymes are well known in the art; for a general description see patents by Cech (US6180399; US5869254; US6025167; US5854038; US5591610; US5667969; US5354855; US5093246; US5180818; US5116742; US5037746; and US4987071). Ribozymes and other inhibitory nucleic acids can be designed to preferentially inhibit expression of a gene having a sequence associated with a risk haplotype. Thus, a ribozyme that recognizes the sequence spanning the polymorphism and cleaving adjacent to GUA recognizes the risk form but not the neutral or protective form, allowing selective cleavage (Dawson et al., 2000, "Hammerhead ribozymes selectively suppress mutant type I collagen mRNA in osteogenesis imperfecta fibroblasts" *Nucleic Acids Res.* 28:4013-20; Blalock et al., 2004 "Hammerhead ribozyme targeting connective tissue growth factor mRNA blocks transforming growth factor-beta mediated cell proliferation" *Exp Eye Res.* 78:1127-36).

[0250] *Triplex-Forming Oligonucleotides* - Triplex-forming oligonucleotides bind to the major groove of duplex DNA in a sequence-specific manner and provoke DNA repair, resulting in the targeted modification of the genome (for a recent review see Kuan et al., 2004, "Targeted gene modification using triplex-forming oligonucleotides" *Methods Mol Biol.* 262:173-94). Oligonucleotides can be designed to specifically bind to polymorphic sites in the Factor H gene associated with a risk haplotype. A triplex-forming oligonucleotide that binds to a sequence spanning a polymorphism associated with a risk haplotype provokes DNA repair, resulting in the modification of the sequence from a risk allele to a neutral or protective allele.

[0251] Similar antisense nucleic acid, RNA interference, ribozyme and triplex-forming plionucleotide methodologies as described above may be used to reduce levels of risk forms of CFHR5 in the eye or systemically for treatment of AMD.

[0252] It will be understood that inhibitory nucleic acids can be administered as a pharmaceutical composition or using gene therapy methods.

G) Antibody Therapy

[0253] In one aspect, an anti-HF1 antibody that specifically interacts with and neutralizes the activity of a variant Factor H polypeptide is administered to an individual with or at risk for AMD. In one embodiment, the antibody recognizes both wild-type and variant Factor H protein. In one embodiment, the antibody recognizes the variant but not the wild-type Factor H protein. In another aspect, an anti-CFHR5 antibody that specifically interacts with and neutralizes the activity of a variant CFHR5 polypeptide is administered to an individual with or at risk for AMD. In one embodiment, the antibody recognizes both wild-type and variant CFHR5 protein. In one embodiment, the antibody recognizes the variant but not the wild-type CFHR5 protein. The antibody can be administered systemically or locally (see, e.g., Gaudreault et al., 2005, "Preclinical pharmacokinetics of Ranibizumab (rhuFabV2) after a single intravitreal administration" *Invest Ophthalmol Vis Sci.* 46:726-33). Methods for making anti-HF1 and anti-CFHR5 antibodies are known in the art, and include methods described below. In a related aspect, an agent that preferentially interacts with and reduces the activity of a variant Factor H polypeptide and/or CFHR5 polypeptide is administered to an individual with or at risk for AMD.

H) Modulators of the Alternative Pathway

[0254] In one aspect the invention provides methods for treating AMD by administering an agent (e.g., native protein, recombinant protein, antibody, or small molecule) directed at modulating the alternative pathway (AP) of the complement cascade, either locally in the eye or at the systemic level. In one embodiment, the treatment comprises administering an agent that modulates the AP directly. In one embodiment, the treatment comprises administering an agent that modulates the triggering of the AP (e.g., microbes). In one embodiment, the treatment comprises administering an agent that modulates pathways downstream from the AP. Exemplary agents that modulate the AP are known in the art and include, but are not limited to, DFP, PR226, BCX-1470, FUT-175, sMCP, PS-oligo, Compstatin, Fucan, and GCRF (see, e.g., Makrides, 1998, "Therapeutic inhibition of the complement system" *Pharmacol Rev.* 50:59-87; Holland et al., 2004, "Synthetic small molecule complement inhibitors" *Curr Opin Investig Drugs* 5:1163-73; Holers et al., 2004, "The alternative pathway of complement in disease: opportunities for therapeutic targeting" *Mol Immunol.* 41:147-52). AP modulators can be administered systemically or by intraocular injection or other methods known for delivery of compounds to the eye.

I) Drug Screening/Antagonists of Risk Variant Factor H or Variant CFHR5

[0255] The invention provides a method of screening for an agent effective for treatment of AMD by contacting a variant protein, host cell or transgenic animal expressing a Factor H or CFHR5 variant, and monitoring binding, expression, processing or activity of the variant. In an embodiment, the Factor H variant has valine at amino acid 62 and/or has histidine at amino acid 402 and/or has cysteine at amino acid 1210. In an embodiment, the CFHR5 variant has a serine at amino acid 46.

[0256] Antagonists of variant Factor H polypeptides (e.g., variants associated with risk haplotypes) can be used to treat AMD. Antagonists may suppress expression of variant Factor H, suppress activity, or reduce RNA or protein stability. Antagonists can be obtained by producing and screening large combinatorial libraries, which can be produced for many types of compounds in a step-wise and high throughput fashion. Such compounds include peptides, polypeptides, beta-turn mimetics, polysaccharides, phospholipids, hormones, prostaglandins, steroids, aromatic compounds, heterocyclic compounds, benzodiazepines, oligomeric N-substituted glycines and oligocarbamates, and the like. Large combinatorial libraries of the compounds can be constructed by methods known in the art. See e.g., WO

95/12608; WO 93/06121; WO 94/08051; WO 95/35503; WO 95/30642 and WO 91/18980. Libraries of compounds are initially screened for specific binding to the variant Factor H polypeptide. Compounds with *in vitro* binding activity can also be assayed for their ability to interfere with a biological activity of the variant Factor H polypeptide, for example, binding to C3b or to heparin. Antagonist activity can be assayed in either a cell-based system or in a transgenic animal model in which exogenous variant Factor H polypeptide is expressed.

[0257] Antagonists of variant CFHR5 polypeptides (e.g., variants associated with risk haplotypes) can be used treat AMD and can be obtained as described above for variant Factor H antagonists.

J) Patient Specific Therapy

[0258] Customized therapies can be devised for groups of patients with different genetic subtypes of AMD, based upon the presence of certain polymorphisms in the Factor H gene or CFHR5 gene having causative roles in AMD and having elucidated the effect of these polymorphisms on the expression level and/or biological activity of variant Factor H polypeptides or CFHR5 polypeptides. For example, if a polymorphism in Factor H or CFHR5 causes AMD in an animal model by increasing the expression level and/or biological activity of a variant Factor H polypeptide or CFHR5 polypeptide, AMD associated with the Factor H or CFHR5 polymorphism can be treated by administering to a patient an antagonist of the variant Factor H polypeptide or variant CFHR5 polypeptide.

K) Assessing Therapeutic Efficacy Using AMD Biomarkers

[0259] As noted above, therapeutic efficacy of particular fragments of CFH or CFHR proteins can also be determined by testing the effect of the protein on expression of AMD biomarkers. Exemplary AMD biomarkers include those described hereinabove. These AMD-associated proteins (biomarkers) are present in individuals with AMD at different (elevated or reduced) levels compared to healthy individuals. The invention provides methods of assessing the efficacy of treatment of AMD and monitoring the progression of AMD by determining a level of a biomarker(s) in an individual with AMD being treated for the disease and comparing the level of the biomarker(s) to an earlier determined level or a reference level of the biomarker. As described in copending provisional application No. 60/715,503, the level of the biomarker(s) can be determined by any suitable method, such as conventional techniques known in the art, including, for example and not for limitation, separation-based methods (e.g., gel electrophoresis), immunoassay methods (e.g., antibody-

based detection) and function-based methods (e.g., enzymatic or binding activity). In one embodiment, a method of assessing the efficacy of treatment of AMD in a individual involves obtaining a sample from the individual and determining the level of the biomarker(s) by separating proteins by 2-dimensional difference gel electrophoresis (DIGE).

VIII. FACTOR H AND CFHR5 NUCLEIC ACIDS

A) Primers and Probes

[0260] The invention provides nucleic acids adjacent to or spanning the polymorphic sites. The nucleic acids can be used as probes or primers (including Invader, Molecular Beacon and other fluorescence resonance energy transfer (FRET) type probes) for detecting Factor H polymorphisms. In one embodiment, the probes or primers recognize the insertion in intron 2 but do not recognize the wild-type sequence. Exemplary nucleic acids comprise sequences that span at least one of the polymorphisms listed in TABLES 1A, 1B, 1C, 11, 14 and 15 in which the polymorphic position is occupied by an alternative base for that position. The base for that position, which is found more frequently in the control population, is denoted the normal or wild-type sequence, whereas the alternative base for that position, which is found less frequently in the control population, is denoted the variant sequence. The nucleic acids also comprise sequences that span other polymorphisms known in the Factor H and CFHR5 genes, such as polymorphisms identified in Tables A and B above.

B) Expression Vectors and Recombinant Production of Factor H and CFHR5

Polypeptides.

[0261] The invention provides vectors comprising nucleic acid encoding the Factor H polypeptide. The Factor H polypeptide may be wild-type or a variant (e.g., a protective variant) and may be a full-length form (e.g., HF1) or a truncated form. The nucleic acid may be DNA or RNA and may be single-stranded or double-stranded.

[0262] Some nucleic acids encode full-length, variant forms of Factor H polypeptides. The variant Factor H polypeptide may differ from normal or wild-type Factor H at an amino acid encoded by a codon including one of any non-synonymous polymorphic position known in the Factor H gene. In one embodiment, the variant Factor H polypeptides differ from normal or wild-type Factor H polypeptides at an amino acid encoded by a codon including one of the non-synonymous polymorphic positions shown in TABLE 1A, TABLE 1B and/or TABLE 1C, that position being occupied by the amino acid shown in TABLE 1A, TABLE 1B and/or

TABLE 1C. It is understood that variant Factor H genes may be generated that encode variant Factor H polypeptides that have alternate amino acids at multiple polymorphic sites in the Factor H gene.

[0263] The invention provides vectors comprising nucleic acid encoding the CFHR5 polypeptide. The CFHR5 polypeptide may be wild-type or a variant (e.g., a protective variant). The nucleic acid may be DNA or RNA and may be single-stranded or double-stranded.

[0264] Some nucleic acids encode full-length, variant forms of CFHR5 polypeptides. The variant CFHR5 polypeptide may differ from normal or wild-type CFHR5 at an amino acid encoded by a codon including one of any non-synonymous polymorphic position known in the CFHR5 gene. In one embodiment, the variant CFHR5 polypeptides differ from normal or wild-type CFHR5 polypeptides at an amino acid encoded by a codon including one of the non-synonymous polymorphic positions shown in TABLES 14 and 15, that position being occupied by the amino acid shown in TABLES 14 and 15. It is understood that variant CFHR5 genes may be generated that encode variant CFHR5 polypeptides that have alternate amino acids at multiple polymorphic sites in the CFHR5 gene.

[0265] Expression vectors for production of recombinant proteins and peptides are well known (see Ausubel *et al.*, 2004, Current Protocols In Molecular Biology, Greene Publishing and Wiley-Interscience, New York). Such expression vectors include the nucleic acid sequence encoding the Factor H polypeptide linked to regulatory elements, such a promoter, which drive transcription of the DNA and are adapted for expression in prokaryotic (e.g., *E. coli*) and eukaryotic (e.g., yeast, insect or mammalian cells) hosts. A variant Factor H or CFHR5 polypeptide can be expressed in an expression vector in which a variant Factor H or CFHR5 gene is operably linked to a promoter. Usually, the promoter is a eukaryotic promoter for expression in a mammalian cell. Usually, transcription regulatory sequences comprise a heterologous promoter and optionally an enhancer, which is recognized by the host cell. Commercially available expression vectors can be used. Expression vectors can include host-recognized replication systems, amplifiable genes, selectable markers, host sequences useful for insertion into the host genome, and the like.

[0266] Suitable host cells include bacteria such as *E. coli*, yeast, filamentous fungi, insect cells, and mammalian cells, which are typically immortalized, including mouse, hamster, human, and monkey cell lines, and derivatives thereof. Host cells may be able to process the variant Factor H or CFHR5 gene product to produce an appropriately processed, mature

polypeptide. Such processing may include glycosylation, ubiquitination, disulfide bond formation, and the like.

[0267] Expression constructs containing a variant Factor H or CFHR5 gene are introduced into a host cell, depending upon the particular construction and the target host. Appropriate methods and host cells, both procaryotic and eukaryotic, are well-known in the art.

Recombinant full-length human Factor H has been expressed for research purposes in *Sf9* insect cells (see Sharma and Pangburn, 1994, Biologically active recombinant human complement factor H: synthesis and secretion by the baculovirus system, *Gene* 143:301-2). Recombinant fragments of human Factor H have been expressed for research purposes in a variety of cell types (see, e.g., Cheng et al., 2005, "Complement factor H as a marker for detection of bladder cancer" *Clin Chem.* 5:856-63; Vaziri-Sani et al., 2005, "Factor H binds to washed human platelets" *J Thromb Haemost.* 3:154-62; Gordon et al., 1995, "Identification of complement regulatory domains in human factor H" *J Immunol.* 155:348-56).

Recombinant full-length human *CFHR5* has been expressed for research purposes in *Sf9* insect cells (see McRae et al., 2001, Human Factor H-related Protein 5 (FHR-5), *J. Biol. Chem.* 276:6747-6754).

[0268] A variant Factor H or CFHR5 polypeptide may be isolated by conventional means of protein biochemistry and purification to obtain a substantially pure product. For general methods see Jacoby, *Methods in Enzymology* Volume 104, Academic Press, New York (1984); Scopes, *Protein Purification, Principles and Practice*, 2nd Edition, Springer-Verlag, New York (1987); and Deutscher (ed) *Guide to Protein Purification, Methods in Enzymology*, Vol. 182 (1990). Secreted proteins, like Factor H or CFHR5, can be isolated from the medium in which the host cell is cultured. If the variant Factor H or CFHR5 polypeptide is not secreted, it can be isolated from a cell lysate.

[0269] In one embodiment the vector is an expression vector for production of a variant Factor H protein having a sequence having non-wildtype sequence at one or more of the polymorphic sites shown in TABLES 1A, 1B and/or 1C.

[0270] In one embodiment the vector is an expression vector for production of a variant Factor H protein having a sequence of a protective variant of Factor H.

[0271] In one embodiment the vector is an expression vector for production of a variant CFHR5 protein having a sequence having non-wildtype sequence at one or more of the polymorphic sites shown in TABLES 14 and 15.

[0272] In one embodiment the vector is an expression vector for production of a variant CFHR5 protein having a sequence of a protective variant of Factor H.

C) Gene Therapy Vectors

[0273] Methods for expression of Factor H polypeptides or CFHR5 polypeptides for gene therapy are known and are described in Section IV(A) above.

XI. ANTIBODIES

[0274] The invention provides Factor H-specific antibodies that may recognize the normal or wild-type Factor H polypeptide or a variant Factor H polypeptide in which one or more non-synonymous single nucleotide polymorphisms (SNPs) are present in the Factor H coding region. In one embodiment, the invention provides antibodies that specifically recognize variant Factor H polypeptides or fragments thereof, but not Factor H polypeptides not having a variation at the polymorphic site.

[0275] The invention also provides CFHR5-specific antibodies that may recognize the normal or wild-type CFHR5 polypeptide or a variant CFHR5 polypeptide in which one or more non-synonymous single nucleotide polymorphisms (SNPs) are present in the CFHR5 coding region. In one embodiment, the invention provides antibodies that specifically recognize variant CFHR5 polypeptides or fragments thereof, but not CFHR5 polypeptides not having a variation at the polymorphic site.

[0276] The antibodies can be polyclonal or monoclonal, and are made according to standard protocols. Antibodies can be made by injecting a suitable animal with a variant Factor H or variant CFHR5 polypeptide, or fragment thereof, or synthetic peptide fragments thereof. Monoclonal antibodies are screened according to standard protocols (Koehler and Milstein 1975, *Nature* 256:495; Dower et al., WO 91/17271 and McCafferty et al., WO 92/01047; and Vaughan et al., 1996, *Nature Biotechnology*, 14: 309; and references provided below). In one embodiment, monoclonal antibodies are assayed for specific immunoreactivity with the variant Factor H or CFHR5 polypeptide, but not the corresponding wild-type Factor H or CFHR5 polypeptide, respectively. Methods to identify antibodies that specifically bind to a variant polypeptide, but not to the corresponding wild-type polypeptide, are well-known in the art. For methods, including antibody screening and subtraction methods; see Harlow & Lane, *Antibodies, A Laboratory Manual*, Cold Spring Harbor Press, New York (1988); *Current Protocols in Immunology* (J.E. Coligan et al., eds., 1999,

including supplements through 2005); Goding, *Monoclonal Antibodies, Principles and Practice* (2d ed.) Academic Press, New York (1986); Burioni et al., 1998, "A new subtraction technique for molecular cloning of rare antiviral antibody specificities from phage display libraries" *Res Virol.* 149(5):327-30; Ames et al., 1994, Isolation of neutralizing anti-C5a monoclonal antibodies from a filamentous phage monovalent Fab display library. *J Immunol.* 152(9):4572-81; Shinohara et al., 2002, Isolation of monoclonal antibodies recognizing rare and dominant epitopes in plant vascular cell walls by phage display subtraction. *J Immunol Methods* 264(1-2):187-94. Immunization or screening can be directed against a full-length variant protein or, alternatively (and often more conveniently), against a peptide or polypeptide fragment comprising an epitope known to differ between the variant and wild-type forms. Particular variants include the Y402H or I62V variants of CFH and HFL1, the R1210C variant of CFH, the P46S variant of CFHR5, and truncated forms of CFH. In one embodiment the HF1 is measured. As discussed above, in one embodiment the ratio of HFL1 and CHF is measured. Monoclonal antibodies specific for variant Factor H or CFHR5 polypeptides (i.e., which do not bind wild-type proteins, or bind at a lower affinity) are useful in diagnostic assays for detection of the variant forms of Factor H or CFHR5, or as an active ingredient in a pharmaceutical composition.

[0277] The present invention provides recombinant polypeptides suitable for administration to patients including antibodies that are produced and tested in compliance with the Good Manufacturing Practice (GMP) requirements. For example, recombinant antibodies subject to FDA approval must be tested for potency and identity, be sterile, be free of extraneous material, and all ingredients in a product (i.e., preservatives, diluents, adjuvants, and the like) must meet standards of purity, quality, and not be deleterious to the patient.

[0278] The invention provides a composition comprising an antibody that specifically recognizes a Factor H or CFHR5 polypeptide (e.g., a normal or wild-type Factor H polypeptide or a variant Factor H polypeptide, or a normal or wild-type CFHR5 polypeptide or a variant CFHR5 polypeptide) and a pharmaceutically acceptable excipient or carrier.

[0279] In a related aspect, the invention provides a sterile container, e.g. vial, containing a therapeutically acceptable Factor H-specific or CFHR5-specific antibody. In one embodiment it is a lyophilized preparation.

[0280] In a related aspect, the invention provides pharmaceutical preparations of human or humanized anti-Factor H or anti-CFHR5 antibodies for administration to patients.

Humanized antibodies have variable region framework residues substantially from a human

antibody (termed an acceptor antibody) and complementarity determining regions substantially from a mouse-antibody, (referred to as the donor immunoglobulin). See, Peterson, 2005, Advances in monoclonal antibody technology: genetic engineering of mice, cells, and immunoglobulins, *ILAR J.* 46:314-9, Kashmiri et al., 2005, SDR grafting - a new approach to antibody humanization, *Methods* 356:25-34, Queen et al., *Proc. Natl. Acad. Sci. USA* 86:10029-10033 (1989), WO 90/07861, U.S. Pat. No. 5,693,762, U.S. Pat. No. 5,693,761, U.S. Pat. No. 5,585,089, U.S. Pat. No. 5,530,101, and Winter, U.S. Pat. No. 5,225,539. The constant region(s), if present, are also substantially or entirely from a human immunoglobulin. The human variable domains are usually chosen from human antibodies whose framework sequences exhibit a high degree of sequence identity with the murine variable region domains from which the CDRs were derived. The heavy and light chain variable region framework residues can be derived from the same or different human antibody sequences. The human antibody sequences can be the sequences of naturally occurring human antibodies or can be consensus sequences of several human antibodies. See Carter et al., WO 92/22653. Certain amino acids from the human variable region framework residues are selected for substitution based on their possible influence on CDR conformation and/or binding to antigen. Investigation of such possible influences is by modeling, examination of the characteristics of the amino acids at particular locations, or empirical observation of the effects of substitution or mutagenesis of particular amino acids.

[0281] For example, when an amino acid differs between a murine variable region framework residue and a selected human variable region framework residue, the human framework amino acid should usually be substituted by the equivalent framework amino acid from the mouse antibody when it is reasonably expected that the amino acid: (1) noncovalently binds antigen directly, (2) is adjacent to a CDR region, (3) otherwise interacts with a CDR region (e.g. is within about 6 Å of a CDR region), or (4) participates in the VL-VH interface.

[0282] Other candidates for substitution are acceptor human framework amino acids that are unusual for a human immunoglobulin at that position. These amino acids can be substituted with amino acids from the equivalent position of the mouse donor antibody or from the equivalent positions of more typical human immunoglobulins. Other candidates for substitution are acceptor human framework amino acids that are unusual for a human immunoglobulin at that position. The variable region frameworks of humanized

immunoglobulins usually show at least 85% sequence identity to a human variable region framework sequence or consensus of such sequences.

IX. IDENTIFICATION OF RISK, PROTECTIVE, AND NEUTRAL VARIATIONS AND HAPLOTYPES

[0283] The invention provides methods of screening for polymorphic sites linked to polymorphic sites in the Factor H gene and/or CFHR5 gene described in TABLES 1A, 1B, 1C, 11, 14 and 15. These methods involve identifying a polymorphic site in a gene that is linked to a polymorphic site in the Factor H gene or CFHR5 gene, wherein the polymorphic form of the polymorphic site in the Factor H gene or CFHR5 gene is associated AMD (e.g., increased or decreased risk), and determining haplotypes in a population of individuals to indicate whether the linked polymorphic site has a polymorphic form in equilibrium or disequilibrium with the polymorphic form of the Factor H gene or CFHR5 gene that correlates with the AMD phenotype.

[0284] Polymorphisms in the Factor H gene or CFHR5 gene, such as those shown in TABLES 1A, 1B, 1C, 11, 14 and 15, can be used to establish physical linkage between a genetic locus associated with a trait of interest and polymorphic markers that are not associated with the trait, but are in physical proximity with the genetic locus responsible for the trait and co-segregate with it. Mapping a genetic locus associated with a trait of interest facilitates cloning the gene(s) responsible for the trait following procedures that are well-known in the art.

[0285] Polymorphisms in the Factor H gene or CFHR5 gene, such as those shown in TABLES 1A, 1B, 1C, 11, 14 and 15, can be used in familial linkage studies to determine which polymorphisms co-segregate with a phenotypic trait, to determine individuals who require therapy, and to determine the effects of therapy.

[0286] Linkage is analyzed by calculation of a LOD (log of the odds) score, which is the $\log_{10}$ of the ratio of the likelihood of obtaining observed segregation data for a marker and a genetic locus when the two are located at a recombination fraction θ , versus the situation in which the two are not linked (segregating independently). See Thompson & Thompson, Genetics in Medicine (5th ed, W.B. Saunders Company, Philadelphia, 1991) and Strachan, "Mapping the human genome" in The Human Genome (BIOS Scientific Publishers Ltd, Oxford) Chapter 4). A LOD score of 3 indicates a 1000 to 1 odds against an apparent observed linkage being a coincidence. A LOD score of +3 or greater is considered definitive

evidence that two loci are linked, whereas LOD score of -2 or less is considered definitive evidence against linkage.

X. TRANSGENIC NON-HUMAN ANIMALS

[0287] The invention provides transgenic non-human animals capable of expressing human variant Factor H or CFHR5 polypeptides. Transgenic non-human animals may have one or both of alleles of the endogenous Factor H or CFHR5 gene inactivated. Expression of an exogenous variant Factor H or CFHR5 gene is usually achieved by operably linking the gene to a promoter and optionally an enhancer, and then microinjecting the construct into a zygote following standard protocols. See Hogan et al., "Manipulating the Mouse Embryo, A Laboratory Manual," Cold Spring Harbor Laboratory. The endogenous Factor H or CFHR5 genes can be inactivated by methods known in the art (Capecchi, 1989). Factor H deficient mice are available for the introduction of exogenous human variant Factor H genes.

Transgenic animals expressing human or non-human variant Factor H or CFHR5 polypeptides provide useful drug screening systems and as models of AMD and other complement related diseases. Transgenic animals may also be used for production of recombinant CFH and CFHR5 proteins of the invention (see, e.g. U.S. Pat. Nos. 6,066,725; 6,013,857; 5,994,616; and 5,959,171; Lillico et al., 2005; Houdebine, 2000).

XI. KITS

[0288] The invention provides reagents, devices and kits detecting Factor H or CFHR5 polymorphisms and haplotypes. Although particularly suited for screening for risk of developing AMD and/or for identifying appropriate therapy for preventing or ameliorating AMD in a subject, it will be understood that in certain embodiments these reagents, devices and kits can be used for analysis of Factor H and CFHR5 polymorphisms and haplotypes for any purpose, including but not limited to determining risk of developing MPGNII or any other complement associated condition.

[0289] A number of assay systems are known in the art, and it is within the skill of the art to arrive at means to determine the presence of variations associated with AMD. The kit reagents, such as multiple primers, multiple probes, combinations of primers, or combinations of probes, may be contained in separate containers prior to their use for diagnosis or screening. In an embodiment, the kit contains a first container containing a probe, primer, or primer pair for a first CFH or CFHR5 allele described herein, and a second

container containing a probe, primer, or primer pair for a second CFH or CFHR5 allele described herein.

[0290] In one embodiment, the invention provides kits comprising at least one Factor H or CFHR5 allele-specific oligonucleotide that hybridizes to a specific polymorphism in the Factor H or CFHR5 gene. The kits may contain one or more pairs of Factor H or CFHR5 allele-specific oligonucleotides hybridizing to different forms of a polymorphism. The Factor H or CFHR5 allele-specific oligonucleotides may include sequences derived from the coding (exons) or non-coding (promoter, 5' untranslated, introns or 3' untranslated) region of the Factor H or CFHR5 gene. The Factor H or CFHR5 allele-specific oligonucleotides may be provided immobilized on a substrate. The substrate may comprise Factor H or CFHR5 allele-specific oligonucleotide probes for detecting at least 2, 3, 4, 5, more than 5, (e.g., at least 6, 7, or 8) or all of the polymorphisms shown in TABLES 1A, 1B, 1C, 11, 14 and 15 and/or other polymorphisms in the Factor H or CFHR5 gene (e.g., including polymorphisms listed above that are found in the SNP database). In one embodiment the kit is used to diagnose AMD. In a related embodiment, the kit is used to screen for another disease associated with variation in the Factor H or CFHR5 gene.

[0291] The kit may include at least one Factor H- or CFHR5- specific primer that hybridizes spanning or adjacent to a specific polymorphism in the Factor H or CFHR5 gene. The Factor H- or CFHR5- specific primers may include sequences derived from the coding (exons) or non-coding (promoter, 5' untranslated, introns or 3' untranslated) region of the Factor H or CFHR5 gene. Often, the kits contain one or more pairs of Factor H- or CFHR5- specific primers that hybridize to opposite strands of nucleic acid adjacent to a specific polymorphism in the Factor H or CFHR5 gene. In the presence of appropriate buffers and enzymes, the Factor H- or CFHR5- specific primer pairs are useful in amplifying specific polymorphisms in the Factor H or CFHR5 gene.

[0292] It will be apparent to the skilled practitioner guided by this disclosure that various polymorphisms and haplotypes can be detected to assess the propensity of an individual to develop a Factor H related condition. The following examples and combinations are provided for illustration and not limitation. In some cases, the assay identifies the allele at at least one, at least two, at least three, at least four, at least five or at least six polymorphic sites in the Factor H or CFHR5 gene. In some cases, the assay identifies the allele at 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or all of the polymorphisms in the Factor H or CFHR5 gene listed in TABLES 1A, 1B, 1C, 11, 14 and 15. In one embodiment, the sites are selected from: rs529825;

rs800292; rs3766404; rs1061147; rs1061170; rs203674; and optionally including exon 22 (R120C). In one embodiment, the sites are selected from rs529825; rs800292; intron 2 (IVS2 or insTT); rs3766404; rs1061147; rs1061170; exon 10A; rs203674; rs375046; and optionally including exon 22 (R120C). In one embodiment, the sites are selected from: rs3753394; rs529825; rs800292; intron 2 (IVS2 or insTT); rs3766404; rs1061147; rs1061170; rs2274700; rs203674; rs3753396; rs1065489; and optionally including exon 22 (R120C). In one embodiment, the sites are selected from: rs800292 (I62V); IVS 2 (-18insTT); rs1061170 (Y402H); and rs2274700 (A473A). In one embodiment, the sites are selected from: rs9427661 (-249T>C); rs9427662 (-20T>C); and rs12097550 (P46S). In a preferred embodiment, a diagnostic/screening assay of the invention identifies the allele at at least two polymorphic sites in the Factor H or *CFHR5* gene. In a preferred embodiment, a diagnostic/screening assay of the invention identifies the allele at at least three polymorphic sites in the Factor H or *CFHR5* gene. In a preferred embodiment, a diagnostic/screening assay of the invention identifies the allele at at least four polymorphic sites in the Factor H or *CFHR5* gene.

[0293] In some cases, the kit includes primers or probes ("oligonucleotides") to identify the allele at 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or all of the polymorphisms in the Factor H or *CFHR5* gene listed in TABLES 1A, 1B, 1C, 11, 14 and 15. In one embodiment the kit includes primers or probes to determine the allele at at least one of the following polymorphic sites: rs529825; rs800292; rs3766404; rs1061147; rs1061170; rs203674; and optionally including exon 22 (R120C). In one embodiment the kit includes primers or probes to determine the allele at at least one of the following polymorphic sites: rs529825; rs800292; intron 2 (IVS2 or insTT); rs3766404; rs1061147; rs1061170; exon 10A; rs203674; rs375046; and optionally including exon 22 (R120C). In one embodiment the kit includes primers or probes to determine the allele at at least one of the following polymorphic sites: rs3753394; rs529825; rs800292; intron 2 (IVS2 or insTT); rs3766404; rs1061147; rs1061170; rs2274700; rs203674; rs3753396; rs1065489; and optionally including exon 22 (R120C). In one embodiment, the sites are selected from: rs800292 (I62V); IVS 2 (-18insTT); rs1061170 (Y402H); and rs2274700 (A473A). In one embodiment, the sites are selected from: rs9427661 (-249T>C); rs9427662 (-20T>C); and rs12097550 (P46S).

[0294] The kit can include primers or probes to determine the allele at two of the above sites, at least three, at least four, at least five or at least six. In one embodiment the primers or probes distinguish alleles at rs529825. In one embodiment the primers or probes distinguish

alleles at rs800292. In one embodiment the primers or probes distinguish alleles at rs3766404. In one embodiment the primers or probes distinguish alleles at rs1061147. In one embodiment the primers or probes distinguish alleles at rs1061170. In one embodiment the primers or probes distinguish alleles at rs203674. In one embodiment; the primers or probes distinguish alleles at exon 22 (R1210C). In one embodiment the primers or probes distinguish alleles at rs529825 and rs800292. In one embodiment the primers or probes distinguish alleles at two or three of rs1061147, rs1061170 and rs203674. In one embodiment the primers or probes distinguish alleles at at least one of rs529825 and rs800292; and rs3766404; and at least one of rs1061147, rs1061170 and rs203674. In one embodiment the primers or probes distinguish alleles at rs529825, rs800292, rs3766404, rs1061170 and rs203674. In one embodiment, the primers or probes distinguish alleles at exon 22 (R1210C) and at: (a) rs529825; rs800292; rs3766404; rs1061147; rs1061170; rs203674; rs529825 rs 800292; (b) at two or three of rs1061147, rs1061170 and rs203674; at rs529825 and rs800292, rs3766404, and two or three of rs1061147, rs1061170 and rs203674; or at rs529825, rs800292, rs3766404, rs1061170 and rs203674. In one embodiment the primers or probes distinguish alleles at at least one of rs529825 and rs800292; and rs3766404; and at least one of rs1061147, rs1061170 and rs203674. In one embodiment the primers or probes distinguish alleles at rs529825, rs800292, rs3766404, rs1061170 and rs203674.

[0295] The kit can include primers or probes to determine the allele at two of the above sites, or at least three of the above sites. In one embodiment, the primers or probes distinguish alleles at rs800292. In one embodiment, the primers or probes distinguish alleles at rs1061170. In one embodiment, the primers or probes distinguish alleles at exon 22 (R1210C). In one embodiment, the primers or probes distinguish alleles at exon 22 (R1210C) and rs800292 and/or exon 22 and rs1061170 and exon 22. In one embodiment, the primers or probes distinguish alleles at rs800292, rs1061170 and exon 22 (R1210C).

[0296] The kit can include primers or probes to determine the allele at two of the above sites, at least three, at least four, at least five or at least six. In one embodiment the primers or probes distinguish alleles at rs529825. In one embodiment the primers or probes distinguish alleles at rs800292. In one embodiment the primers or probes distinguish alleles at intron 2 (IVS2 or insTT). In one embodiment the primers or probes distinguish alleles at rs3766404. In one embodiment the primers or probes distinguish alleles at rs1061147. In one embodiment the primers or probes distinguish alleles at rs1061170. In one embodiment the

primers or probes distinguish alleles at rs2274700. In one embodiment the primers or probes distinguish alleles at exon 10A. In one embodiment the primers or probes distinguish alleles at rs203674. In one embodiment the primers or probes distinguish alleles at rs375046. In one embodiment the primers or probes distinguish alleles at exon 22 (R1210C). In one embodiment the primers or probes distinguish alleles at rs529825 and rs800292. In one embodiment the primers or probes distinguish alleles at two or three of rs1061147, rs1061170 and rs203674. In one embodiment the primers or probes distinguish alleles at rs529825 and rs800292, at intron 2, at rs3766404, at two or three of rs1061147, rs1061170 and rs203674, at rs2274700, at exon 10A, and at rs375046. In one embodiment the primers or probes distinguish alleles at rs529825, rs800292, intron 2 (IVS2 or insTT), rs3766404, rs1061170, rs2274700, exon 10A, rs203674, and rs375046. In one embodiment, the primers or probes distinguish alleles at exon 22 (R1210C) and at any one or more of rs529825; rs800292; intron 2 (IVS2 or insTT); rs3766404; rs1061147; rs1061170; rs2274700, exon 10A; rs203674; rs375046; rs529825 and rs 800292. In one embodiment, the primers or probes distinguish alleles at exon 22 (R1210C) and at: (a) any two or three of rs1061147, rs1061170 and rs203674; (b) at rs529825 and rs800292, intron 2 (IVS2 or insTT), rs3766404, two or three of rs1061147, rs1061170 and rs203674, rs2274700, exon 10A, and rs375046; or at rs529825, rs800292, intron 2 (IVS2 or insTT), rs3766404, rs1061170, rs2274700, exon 10A, rs203674, and rs375046.

[0297] In one embodiment the kit contains probes or primers for detecting at least one variation in the Factor H gene as well as probes or primers for detecting at least one variation in the CHFR-5 gene. In this embodiment, the kit optionally contains probes or primers for detecting more than one variation in either or both of the Factor H and CHFR-5 genes, such as two, three or more than three variations.

[0298] A number of assay formats are known for determining haplotypes and can be adapted to the present invention. See, e.g., Görgens et al., 2004, "One-Step Analysis of Ten Functional Haplotype Combinations of the Basic RET Promoter with a LightCycler Assay" *Clinical Chemistry* 50:1693-1695; Dawson, 1989, "Carrier identification of cystic fibrosis by recombinant DNA techniques." *Mayo Clin Proc* 64:325-34; Lee et al., 2005, "Gene SNPs and mutations in clinical genetic testing: haplotype-based testing and analysis." *Mutat Res.* 573:195-204.

[0299] In one embodiment, the primers or probes distinguish alleles at (a) any one or more of rs529825; rs800292; rs3766404; rs1061147; rs1061170; and rs203674; (b) any one of more

of intron 2 (IVS2 or insTT); rs2274700; exon 10A; and rs375046; (c) one or both of rs529825 and rs800292; (d) one or more of rs1061147, rs1061170 and rs203674; (e) at least one of rs529825 and rs800292; and rs3766404; and at least one of rs1061147, rs1061170 and rs203674; (f) at least rs529825, rs800292, rs3766404, rs1061170, and rs203674; (g) exon 22 (R1210C); (h) exon 22 (R1210C) and any of (a)-(g); or (i) any one or more of rs529825; rs800292; rs3766404; rs1061147; rs1061170; rs203674; intron 2 (IVS2 or insTT); rs2274700; exon 10A; rs375046; and exon 22 (R1210C) and any one or more of rs9427661, rs9427662 and rs12097550. In one embodiment, the kits include oligonucleotide sufficient to distinguish any combination of alleles at the sites listed below in the context of devices.

[0300] The kits may include antibodies that specifically recognize the Factor H or CFHR5 polypeptide. The Factor H- or CFHR5-specific antibodies may recognize the normal, functional Factor H or CFHR5 polypeptide or variant Factor H or CFHR5 polypeptides in which one or more non-synonymous single nucleotide polymorphisms (SNPs) are present in the Factor H or CFHR5 coding region.

XII. DIAGNOSTIC DEVICES

[0301] Devices and reagents useful for diagnostic, prognostic, drug screening, and other methods are provided. In one aspect, a device comprising immobilized primer(s) or probe(s) specific for one or more Factor H and/or CFHR5 polymorphic sites is provided. In an embodiment the Factor H and/or CFHR5 polymorphic sites are those described herein as associated with AMD.

[0302] In one aspect, a device comprising immobilized primer(s) or probe(s) specific for one or more Factor H and/or CFHR5 gene products (polynucleotides or proteins) is provided. The primers or probes can bind polynucleotides (e.g., based on hybridization to specific polymorphic sites) or polypeptides (e.g., based on specific binding to a variant polypeptide).

[0303] In one embodiment, an array format is used in which a plurality (at least 2, usually at least 3 or more) of different primers or probes are immobilized. The term "array" is used in its usual sense and means that each of a plurality of primers or probes, usually immobilized on a substrate, has a defined location (address) e.g., on the substrate. The number of primers or probes on the array can vary depending on the nature and use of the device. For example, a dipstick format array can have as few as 2 distinct primers or probes, although usually more than 2 primers or probes, and often many more, will be present. One method for attaching the nucleic acids to a surface is by making high-density oligonucleotide arrays (see, Fodor et

al., 1991, *Science* 251:767-73; Lockhart et al., 1996, *Nature Biotech* 14:1675; and U.S. Pat. Nos. 5,578,832; 5,556,752; and 5,510,270). It is also contemplated that, in some embodiments, a device comprising a single immobilized probe can be used.

[0304] In one embodiment, an array format is used in which a plurality (at least 2, usually at least 3 or more) of different primers or probes are immobilized. The term "array" is used in its usual sense and means that each of a plurality of primers or probes, usually immobilized on a substrate, has a defined location (address) e.g., on the substrate. The number of primers or probes on the array can vary depending on the nature and use of the device.

[0305] In one embodiment, the immobilized probe is an antibody or other Factor H or *CFHR5* binding moiety.

[0306] It will be apparent to the skilled practitioner guided by this disclosure that various polymorphisms and haplotypes can be detected to assess the propensity of an individual to develop a Factor H related condition. The following examples and combinations are provided for illustration and not limitation. In some cases, the array includes primers or probes to identify the allele at 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or all of the polymorphisms in the Factor H or *CFHR5* gene listed in TABLES 1A, 1B, 1C, 11, 14 and 15. In one embodiment the array includes primers or probes to determine the allele at at least one of the following polymorphic sites: rs529825; rs800292; rs3766404; rs1061147; rs1061170; rs203674; and optionally including exon 22 (R1210C). In one embodiment the array includes primers or probes to determine the allele at at least one of the following polymorphic sites: rs529825; rs800292; intron 2 (IVS2 or insTT); rs3766404; rs1061147; rs1061170; exon 10A; rs203674; rs375046; and optionally including exon 22 (R1210C). In an embodiment the array includes primers or probes to determine the allele at at least one of the following polymorphic sites: (a) rs3753394; (b) rs529825; (c) rs800292; (d) intron 2 (IVS2 or insTT); (e) rs3766404; (f) rs1061147; (g) rs1061170; (h) rs2274700; (i) rs203674; (j) rs3753396; (j) rs1065489; and optionally including exon 22 (R1210C). In one embodiment, the array includes primers or probes to determine the allele at at least one of the following polymorphic sites: rs800292 (I62V); IVS 2 (-18insTT); rs1061170 (Y402H); and rs2274700 (A473A). In one embodiment, the array includes primers or probes to determine the allele at at least one of the following polymorphic sites: rs9427661 (-249T>C); rs9427662 (-20T>C); and rs12097550 (P46S).

[0307] The array can include primers or probes to determine the allele at two of the above sites, at least three, at least four, at least five or at least six. In one embodiment the primers or

probes distinguish alleles at rs529825. In one embodiment the primers or probes distinguish alleles at rs800292. In one embodiment the primers or probes distinguish alleles at rs3766404. In one embodiment the primers or probes distinguish alleles at rs1061147. In one embodiment the primers or probes distinguish alleles at rs1061170. In one embodiment the primers or probes distinguish alleles at rs203674. In one embodiment the primers or probes distinguish alleles at exon 22 (R1210C). In one embodiment the primers or probes distinguish alleles at rs529825 and rs800292. In one embodiment the primers or probes distinguish alleles at two or three of rs1061147, rs1061170 and rs203674. In one embodiment the primers or probes distinguish alleles at rs529825 and rs800292, at rs3766404, two or three of rs1061147, rs1061170 and rs203674. In one embodiment the primers or probes distinguish alleles at rs529825, rs800292, rs3766404, rs1061170 and rs203674. In one embodiment, the primers or probes distinguish alleles at exon 22 (R1210C) and at rs529825; at rs800292; at rs3766404; at rs1061147; at rs1061170; at rs203674; at rs529825 and rs 800292; at two or three of rs1061147, rs1061170 and rs203674; at rs529825 and rs800292, rs3766404, and two or three of rs1061147, rs1061170 and rs203674; or at rs529825, rs800292, rs3766404, rs1061170 and rs203674. In one embodiment, the primers or probes distinguish alleles at (a) any one or more of rs529825; rs800292; rs3766404; rs1061147; rs1061170; and rs203674; (b) any one or more of intron 2 (IVS2 or insTT); rs2274700; exon 10A; and rs375046; (c) one or both of rs529825 and rs800292; (d) one or more of rs1061147, rs1061170 and rs203674; (e) at least one of rs529825 and rs800292; and rs3766404; and at least one of rs1061147, rs1061170 and rs203674; (f) at least rs529825, rs800292, rs3766404, rs1061170, and rs203674; (g) exon 22 (R1210C); (h) exon 22 (R1210C) and any of (a)-(g); or (i) any one or more of rs529825; rs800292; rs3766404; rs1061147; rs1061170; rs203674; intron 2 (IVS2 or insTT); rs2274700; exon 10A; rs375046; and exon 22 (R1210C) and any one or more of rs9427661, rs9427662 and rs12097550.

[0308] The array can include primers or probes to determine the allele at two of the above sites, at least three, at least four, at least five or at least six. In one embodiment the primers or probes distinguish alleles at rs529825. In one embodiment the primers or probes distinguish alleles at rs800292. In one embodiment the primers or probes distinguish alleles at intron 2 (IVS2 or insTT). In one embodiment the primers or probes distinguish alleles at rs3766404. In one embodiment the primers or probes distinguish alleles at rs1061147. In one embodiment the primers or probes distinguish alleles at rs1061170. In one embodiment the primers or probes distinguish alleles at exon 10A. In one embodiment the primers or probes

distinguish alleles at rs2274700. In one embodiment the primers or probes distinguish alleles at rs203674. In one embodiment the primers or probes distinguish alleles at rs375046. In one embodiment the primers or probes distinguish alleles at exon 22 (R1210C). In one embodiment the primers or probes distinguish alleles at rs529825 and rs800292. In one embodiment the primers or probes distinguish alleles at two or three of rs1061147, rs1061170 and rs203674. In one embodiment the primers or probes distinguish alleles at of rs529825 and rs800292, at intron 2, at rs3766404, at two or three of rs1061147, rs1061170 and rs203674, at exon 10A, at rs2274700, and at rs375046. In one embodiment the primers or probes distinguish alleles at rs529825, rs800292, intron 2 (IVS2 or insTT), rs3766404, rs1061170, exon 10A, rs2274700, rs203674, and rs375046. In one embodiment, the primers or probes distinguish alleles at exon 22 (R1210C) and at either at rs529825; at rs800292; at intron 2 (IVS2 or insTT); at rs3766404; at rs1061147; at rs1061170; at rs2274700, at exon 10A; at rs203674; at rs375046; at rs529825 and rs 800292; at two or three of rs1061147, rs1061170 and rs203674; at rs529825 and rs800292, intron 2 (IVS2 or insTT), rs3766404, two or three of rs1061147, rs1061170 and rs203674, rs2274700, exon 10A, and rs375046; or at rs529825, rs800292, intron 2 (IVS2 or insTT), rs3766404, rs1061170, rs2274700, exon 10A, rs203674, and rs375046. In one embodiment, the device distinguishes any combination of alleles at the sites listed above in the context of kits.

[0309] In one embodiment, the substrate comprises fewer than about 1000 distinct primers or probes, often fewer than about 100 distinct primers or probes, fewer than about 50 distinct primers or probes, or fewer than about 10 distinct primers or probes. As used in this context, a primer is "distinct" from a second primer if the two primers do not specifically bind the same polynucleotide (i.e., such as cDNA primers for different genes). As used in this context, a probe is "distinct" from a second probe if the two probes do not specifically bind the same polypeptide or polynucleotide (i.e., such as cDNA probes for different genes). Primers or probes may also be described as distinct if they recognize different alleles of the same gene (i.e., CFH or CFHR5). Thus, in one embodiment diagnostic devices of the invention detect alleles of CFH only, CFHR5 only, CFH and CFHR5 only, or CFH, CFHR5 and up to 20, preferably up to 10, or preferably up to 5 genes other than CFH and/or CFHR5. That is, the device is particularly suited to screening for AMD and related complement-associated diseases. In one embodiment, the device comprises primers or probes that recognize CFH and/or one or more of CFHR1-5 only. In a related embodiment, the device

contains primers and probes for up to 20, preferably up to 10, or preferably up to 5 other genes than CFH or CFHR1-5.

[0310] In one embodiment, the immobilized primer(s) is/are an allele-specific primer(s) that can distinguish between alleles at a polymorphic site in the Factor H or *CHRF5* gene. Exemplary allele-specific primers to identify alleles at polymorphic sites in the Factor H gene are shown in TABLE 16A. The immobilized allele-specific primers hybridize preferentially to nucleic acids, either RNA or DNA, that have sequences complementary to the primers. The hybridization may be detected by various methods, including single-base extension with fluorescence detection, the oligonucleotide ligation assay, and the like (see Shi, M.M., 2001, "Enabling large-scale pharmacogenetic studies by high-throughput mutation detection and genotyping technologies" *Clin. Chem.* 47(2):164-172). Microarray-based devices to detect polymorphic sites are commercially available, including Affymetrix (Santa Clara, CA), Protogene (Menlo Park, CA), Genometrix (The Woodland, TX), Motorola BioChip Systems (Northbrook, IL), and Perlegen Sciences (Mountain View, CA).

[0311] In one embodiment, the immobilized probe(s) is/are an allele-specific probe(s) that can distinguish between alleles at a polymorphic site in the Factor H or CFHR5 gene. Exemplary allele specific probes to identify alleles at polymorphic sites in the Factor H gene are shown in TABLE 16B. The immobilized allele-specific probes hybridize preferentially to nucleic acids, either RNA or DNA, that have sequences complementary to the probes. The hybridization may be detected by various methods, including fluorescence of hybridized nucleic acids (see Shi, M.M., 2001, "Enabling large-scale pharmacogenetic studies by high-throughput mutation detection and genotyping technologies." *Clin. Chem.* 47(2):164-172). Microarray-based devices to detect polymorphic sites are commercially available, including Affymetrix (Santa Clara, CA), Protogene (Menlo Park, CA), Genometrix (The Woodland, TX), Motorola BioChip Systems (Northbrook, IL), and Perlegen Sciences (Mountain View, CA).

[0312] In certain embodiments probes or primers specific for particular SNPs and variations are excluded from a kit or a device of the invention. For example, in some embodiments, a kit or device does not include one or more of the following SNPs can be excluded: (i) rs529825; (ii) rs900292; (iii) intron 2 (IVS2 or insTT); (iv) rs3766404; (v) rs1061147; (vi) rs1061170; (vii) rs2274700; (viii) exon 10A; (ix) rs203674; (x) rs375046; (xi) rs3753396; (xii) rs1065489; or (xiii) exon 22 (R1210C).

XIII. EXAMPLES

EXAMPLE 1

Common Haplotype in the Factor H Gene (*HF1/CFH*) Predisposes Individuals to Age-related Macular Degeneration

[0313] Age-related macular degeneration (AMD) is the most frequent cause of irreversible blindness in the elderly in developed countries, affecting more than 50 million individuals worldwide. Our previous studies implicated activation of the alternative complement pathway in the formation of ocular drusen, the hallmark lesion of AMD. We have also shown that macular drusen in AMD patients are indistinguishable from those that form at an early age in individuals with membranoproliferative glomerulonephritis type 2 (MPGNII), a disease characterized by uncontrolled activation of the alternative pathway of the complement cascade. Here we show that Factor H protein (HF1), the major inhibitor of the alternative complement pathway, accumulates within drusen, and is synthesized locally by the retinal pigment epithelium. Previous linkage analyses identified chromosome 1q25-32, which harbors the Factor H gene (*HF1/CFH*), as a major AMD susceptibility locus. We analyzed *HF1* for genetic variation in two independent cohorts comprised of approximately 900 AMD cases and 400 matched controls. We find a highly significant association of 8 common *HF1* SNPs with AMD in these cohorts; two common missense variants exhibit highly significant associations (I62V; $\chi^2=36.1$, $p=3.2 \times 10^{-7}$ and Y402H; $\chi^2=54.4$, $p=1.6 \times 10^{-13}$). Haplotype analysis suggests that multiple *HF1* variants confer either an elevated or a reduced risk of AMD. One common at-risk haplotype is present at a frequency of 49% in AMD cases and 26% in controls (OR=2.67, 95%CI [1.80-2.85]). Homozygotes for this haplotype account for 22.1% of cases and 5.1% of controls (OR=5.26, 95%CI [2.84-9.76]). Several protective haplotypes are also identified (OR=0.44-0.55). Further strengthening these data is the finding of the risk haplotype in 70% of MPGNII patients. We propose that genetically pre-determined variation in regulators of the complement system, when combined with triggering events such as infection, underlie a major proportion of AMD in the human population.

Introduction

[0314] Age-related macular degeneration (AMD) is the leading cause of irreversible vision loss in the developed world (Klein et al., 2004; van Leeuwen et al., 2003), affecting 15% of individuals over the age of 60 or an estimated 600 million individuals. AMD is characterized

by a progressive loss of central vision attributable to degenerative and neovascular changes which occur at the interface between the neural retina and the underlying choroid. At this location lie the photoreceptors, the adjacent retinal pigmented epithelium (RPE), a basement membrane complex known as Bruch's membrane (BM), and a network of choroidal capillaries.

[0315] The prevailing view is that AMD is a complex disorder attributable to the interaction of multiple genetic and environmental risk factors (Klein et al., 2003; Tuo et al., 2004). Familial aggregation studies indicate that a genetic component can be identified in up to 25% of the cases (Klaver et al., 1998). As such, AMD appears to be a product of the interaction between multiple susceptibility loci rather than a collection of single-gene disorders. The number of loci involved, the attributable risk conferred, and the interactions between various loci remain obscure.

[0316] Linkage analyses and candidate gene screening have provided limited insight into the genetics of AMD. Reliable associations of one gene with increased risk, *ABCA4* (Allikmets et al., 1997; Allikmets, 2000), and one gene with decreased risk, *ApoE4* (Klaver et al., 1998; Souied et al., 1998), have been reported. Several groups have reported results from genome-wide linkage analyses (Tuo et al., 2004; Weeks et al., 2001). To date, linkage of one AMD phenotype (ARMD1; MIM 603075) to a specific chromosomal region, 1q25-q31, has been documented (Klein et al., 1998). HEMICENTIN-1, also known as *Fib16*, has been tentatively identified as the causal gene (Schultz et al., 2003), although it does not account for a significant disease load (Abecasis et al., 2004; Hayashi et al., 2004). The identification of overlapping loci on chromosome 1q by several groups (Weeks et al., 2001; Iyengar et al., 2003) indicates that this locus is likely to harbor a major AMD-associated gene.

[0317] In AMD, as well as many other diseases such as Alzheimer's disease (Akiyama et al., 2000), atherosclerosis (Torzewski et al., 1997), and glomerulonephritis (Schwartz et al., 2001), characteristic lesions and deposits contribute to the pathogenesis and progression of the disease. Although the molecular pathogenesis of these diseases may be diverse, the deposits contain many shared molecular constituents that are attributable, in part, to local inflammation and activation of the complement cascade, a key element of host defense in the innate immune system. Drusen are the hallmark deposits associated with early AMD, and recent studies have implicated local inflammation and activation of the complement cascade in their formation as well (Hageman et al., 1999; Espinosa-Heidmann et al., 2003). Drusen contain a variety of complement activators, inhibitors, activation-specific complement

fragments, and terminal pathway components including the membrane attack complex (MAC), the lytic complex formed as a consequence of complement activation. The MAC is potentially lethal to host cells and tissues as well as foreign pathogens.

[0318] Many individuals with membranoproliferative glomerulonephritis type II (MPGNII), a rare kidney disease characterized by uncontrolled systemic activation of the alternative activation pathway of complement, also develop ocular drusen in the macula that are indistinguishable in composition and appearance from those in AMD (Mullins et al., 2001; O'Brien et al., 1993; McAvoy et al., 2004). Furthermore, one patient diagnosed with MPGNII harbors a mutation in *HF1* (*HF1*), a major inhibitor of the alternative pathway of complement activation (Zipfel, personal communication). Additionally, individuals in a few extended families with MPGNIII, a related disorder, show linkage to a region of chromosome mapped in 1q31-32 (Neary et al., 2002) that overlaps the locus identified in genome-wide linkage studies for AMD. Collectively, these findings provided the impetus for examining whether *HF1* is involved in the development of AMD and MPGNII.

[0319] In this investigation, we determined the frequencies of *HF1* sequence variants in AMD and MPGNII patients and matched controls, and analyzed their association with disease phenotype. We also examined *HF1* transcription and the distribution of *HF1* protein in the macular RPE-choroid complex from normal and AMD donors.

Methods

[0320] *Patients, Phenotyping and DNA* - Two independent groups of AMD cases and age-matched controls were used for this study. All participating individuals were of European-American descent, over the age of 60, and enrolled under IRB approved protocols following informed consent. These groups were comprised of 404 unrelated patients with clinically documented AMD (mean age 79.5 ± 7.8) and 131 unrelated, control individuals (mean age 78.4 ± 7.4 ; matched by age and ethnicity) from the University of Iowa, and 550 unrelated patients with clinically documented AMD (mean age 71.32 ± 8.9 years), and 275 unrelated, matched by age and ethnicity, controls (mean age 68.84 ± 8.6 years) from the Columbia University. Patients were examined by indirect ophthalmoscopy and slit-lamp microscopy by retina fellowship-trained ophthalmologists.

[0321] Dr. Caroline Klaver, and later individuals trained by Dr. Klaver, graded fundus photographs at both institutions according to a standardized, international classification system (Bird et al., 1995). Control patients were selected and included if they did not exhibit

any distinguishing signs of macular disease or have a known family history of AMD. The AMD patients were subdivided into phenotypic categories -- early AMD (eAMD), geographic atrophy (GA) and exudative (CNV) AMD -- based on the classification of their most severe eye at the time of their entry into the study. The University of Iowa eAMD and GA cases were further subdivided into distinct phenotypes (RPE changes alone, >10 macular hard drusen, macular soft drusen, BB (cuticular) drusen, PED, "Cherokee" atrophy, peninsular geographic atrophy and pattern geographic atrophy). The earliest documented phenotype for all cases was also recorded and employed in the analyses.

[0322] Genomic DNA was generated from peripheral blood leukocytes collected from case and control subjects using QIAamp DNA Blood Maxi kits (Qiagen, Valencia, CA).

[0323] Rapanui - Following an informed consent process approved by the Unidad de Bioetica, Ministerio de Salud (Santiago, Chile), 447 (66% female; 34% male) Easter Island inhabitants were provided a complete eye examination that included a dilated funduscopy examination. Medical, family and ophthalmic histories were taken and records and assistance from local physicians and community leaders were used to classify the ethnicity of the subjects. 49% of those patients examined were pure Rapanui, 9% were admixed (mixture of Rapanui and European, Chilean, Mapuchi and/or recent Polynesian), and 42% were continental (largely Chilean European). Peripheral venous blood and sera were collected from 201 of the older individuals; 114 of these individuals were pure Rapanui (108 were >50 years old; 89 were >60 years old). DNA from 60 of the pure Rapanui inhabitants and 13 of the Chilean residents over the age of 65 was used in this study.

[0324] Human Donor Eyes - Human donor eyes were obtained from the Iowa Lions Eye Bank (Iowa City, IA), the Oregon Lions Eye Bank (Portland, OR) and the Central Florida Lions Eye and Tissue Bank (Tampa, FL) within five hours of death. Gross pathologic features of these eyes, as well as fundus photographs and angiograms, when available, were read and classified by retinal specialists. Fundi were graded according to a modified version of the International AMD grading system (Bird et al., 1995) by at least two individuals.

[0325] Total RNA was prepared from retina, RPE/choroid, and RPE cells derived from eyes using an RNeasy Mini Kit (Qiagen, Valencia, CA). Genomic DNA was sheared using a QiaShredder (Qiagen, Valencia, CA) and residual genomic DNA digested with DNase (Promega). RNA integrity was assessed using an Agilent BioAnalyzer.

[0326] DNA derived from 38 unrelated donors with clinically documented AMD (mean age 81.5 ± 8.6) and 19 unrelated, control donors (mean age 80.5 ± 8.8 ; matched by age and

ethnicity) were employed for SSCP analyses and to assess potential genotype-phenotype correlations.

[0327] Immunohistochemistry – Wedges of posterior poles, including the ora serrata and macula were fixed and processed as described previously (Hageman et al., 1999). Some posterior poles were embedded directly in OCT without prior fixation. Tissues were sectioned to a thickness of 6-8µm on a cryostat and immunolabeling was performed as described previously (Hageman et al., 1999). Adjacent sections were incubated with secondary antibody alone, to serve as controls. Some immunolabeled specimens were prepared and viewed by confocal laser scanning microscopy, as described previously (Anderson et al., 2002).

[0328] Polymerase Chain Reaction (PCR) - First strand cDNA was synthesized from total RNA using Superscript reverse transcriptase (Gibco BRL) and random hexamers. PCR reactions were carried out using the following primer sets: FH1 (exon 8 to exon 10) 5'-GAACATTTTGAGACTCCGTC-3' [SEQ ID NO:324] and 5'-ACCATCCATCTTTCCCAC-3' [SEQ ID NO:325]; FH1 (exon 9 to exon 10) 5'-TCCTGGCTACGCTCTTC-3' [SEQ ID NO:326] and 5'-ACCATCCATCTTTCCCAC-3' [SEQ ID NO:325]; HFL1 (exon 8 to exon 10) 5'-TCCGTCAGGAAGTTACTGG-3' [SEQ ID NO:327] and 5'-AGTCACCATACTCAGGACCC-3' [SEQ ID NO:328]; HFL1 (exon 9 to exon 10), 5'-GGCTACGCTCTTCCAAAAG-3' [SEQ ID NO:329] and 5'-AGTCACCATACTCAGGACCC-3' [SEQ ID NO:330]. PCR primers (IDT, Coralville, IA) were designed using MacVector software (San Diego, CA). Reaction parameters were one cycle at 94°C for 3 minutes, 40 cycles at 94°C for 45 sec, 51.4°C (FH1)/55°C (HFL1) for 1 min, 72°C for 1 min, and one cycle at 72°C for 3 min. The PCR products were run on 2% agarose gels and recorded using a Gel Doc 2000™ Documentation System accompanied by Quantity One® software (Bio-Rad, Hercules, CA).

[0329] Microarray Analyses: DNA microarray analyses were performed using total RNA extracted from native human RPE or the RPE-Choroid complex (RNeasy minikit, Qiagen, Valencia, CA) collected within <5 hours of death. Three different platforms were used: an 18,380 non-redundant DNA microarray (Incyte Pharmaceuticals; St. Louis, MO); the Affymetrix gene chip system; and a Whole Human Genome or Human 1A V2 oligo-microarray (Agilent Inc., Palo Alto, CA). The individual protocols followed each of the manufacturer's instructions. For the Incyte analyses, cDNA derived from 6mm punches of macular and mid-peripheral regions was labeled with 33-P in a random-primed reaction,

purified and hybridized to the Nylon-based arrays containing 18,380 non-redundant cDNAs. The membranes were phosphoimaged, signals were normalized and data were analyzed using the Genome Discovery Software package. For the Affymetrix analyses, RPE and RPE/choroid (from 6-8mm macular and peripheral punches) cRNA was hybridized directly to Affymetrix GeneChips (HG-U133A) using standardized protocols. These procedures were conducted in the University of Iowa DNA core facility, which is equipped with a fluidics station and a GeneArray scanner. The Agilent data was obtained from punches of the macula and mid-periphery. CY3 and CY5 labeled amplified cRNA derived from macula and peripheral RPE/choroid was generated using an Agilent Low Input RNA Amplification Kit using the macular and peripheral RNA from the *same* donor. The Agilent array data were collected from 3 normal young donors, 3 AMD donors, and 3 age-matched non-AMD controls using a VersArray Scanner; data were quantified using the VersArray Analyzer Software (BioRad). The median net intensity of each spot was calculated using global background subtraction and the data was normalized using the local regression method.

[0330] Mutation Screening and Analysis - Coding and adjacent intronic regions of *HF1* (including exon 10A that is transcribed to generate the truncated FHL1 isoform) were examined for variants using single-strand conformation polymorphism (SSCP) analyses, denaturing high performance liquid chromatography (DHPLC) and direct sequencing. The remaining SNPs were typed by the 5' nuclease (Taqman, ABI) methodology. Taqman genotyping and association analyses were performed as described (Gold et al., 2004). Primers for SSCP, DHPLC and DNA sequencing analyses (TABLE 5) were designed to amplify each exon and its adjacent intronic regions using MacVector software (San Diego, CA). PCR-derived amplicons were screened for sequence variation by SSCP and DHPLC, as described previously (Allikmets et al., 1997; Hayashi et al., 2004). All changes detected by SSCP and DHPLC were confirmed by bidirectional sequencing according to standard protocols. Statistical analyses were performed using chi-square and Fisher's exact tests.

Results

Factor H at the RPE-Choroid Interface

[0331] The distribution of Factor H protein in the RPE/choroid complex from the macula and extramacula was assessed in eyes obtained from six donors with a history of early AMD and three donors of similar age without AMD or drusen (FIGURES 1A-1L). In donors with AMD, intense HF1 immunoreactivity (IR) is present in drusen, beneath the RPE (*i.e.* the sub-

RPE space), and around the choroidal capillaries (FIGURES 1A-1D, 1E, 1G). In the absence of the primary antibody, labeling in the RPE/choroid is absent (FIGURES 1F). All of the Factor H antibodies labeled drusen to some degree, in a homogeneous fashion (FIGURES 1C and 1E). One antibody also labeled substructural elements within drusen (FIGURES 1A and 1B). Such structures are also labeled using antibodies to the activated complement component C3b/iC3b that binds HF1 (Anderson et al., 2004; Johnson et al., 2001). Factor H immunoreactivity is more robust in donors with AMD compared to age-matched controls; it is also more pronounced in the maculas of AMD donors than in the periphery (FIGURES 1G and 1H). The anti-HF1 pattern in the macula (FIGURE 1G) is highly similar to the anti-C5b-9 pattern (FIGURES 1I and 1K); in both cases, labeling typically includes the choroidal capillaries. Extramacular locations show much less anti-C5b-9 immunoreactivity (FIGURE 1J). Little or no C5b-9 immunoreactivity in the RPE-choroid is observed in donors under the age of 50 and without AMD (FIGURE 1L).

Description of Figure 1

[0332] (A-B) High magnification confocal immunofluorescence images from an 84 year old male donor diagnosed with atrophic AMD. Anti-HF1 (Advanced Research Technologies) labeling of substructural elements (white arrows) in drusen and the sub-RPE space is imaged on the Cy2/fluorescein channel. The sub-RPE space is the extracellular compartment between the basal RPE surface and the inner collagenous layer of Bruch's membrane. Such elements also display immunoreactivity (IR) using monoclonal antibodies directed against C3 fragments (iC3b, C3d, C3dg) that bind covalently to complement activating surfaces (Johnson et al, 2001; 2003). The intense anti-Factor H labeling in the lumens of choroidal capillaries (asterisks) from this donor most likely reflects the high circulating levels of HF1 in the bloodstream. Autofluorescent lipofuscin granules in the RPE cytoplasm are labeled on the Cy3/Texas Red channel. Magnification bars. A) 5 μ m; B) 3 μ m.

[0333] (C-D) Confocal immunofluorescence localization of HF1 in drusen and the sub-RPE space in an 83 year old male with AMD using a different HF1 polyclonal antibody (Quidel) (Cy2/fluorescein channel; green). C) In this donor eye, the drusen (Dr) labeling pattern is homogeneous. D) Low magnification image of the RPE-choroid. Anti-HF1 IR is present throughout the choroid and in the sub-RPE space (arrows), the anatomical compartment where drusen and other deposits associated with aging and AMD form. Lipofuscin autofluorescence (Cy3 channel; red). Magnification bars. C) 10 μ m; D) 20 μ m.

[0334] (E-F) Immunohistochemical localization of HF1 in drusen. E) Anti-HF1 monoclonal antibody (Quidel) labeling, signified by the purple alkaline phosphatase reaction product, is apparent in drusen, along Bruch's membrane, and on the choroidal capillary walls (arrows). F) Control section from the same eye. In the absence of the primary antibody, no labeling is present. Brown pigmentation in the RPE cytoplasm and choroid is melanin.

Magnification bars = 10 μ m.

[0335] (G-H) Immunolocalization of HF1 in the macula. G) Extensive labeling is present along BM, the choroidal capillary walls, and the intercapillary pillars (arrows) in a donor with AMD. H) Control section from the macula of a donor without AMD; much less labeling is apparent in same structures. Magnification bars = 20 μ m.

[0336] (I-J) Immunohistochemical localization of the complement membrane attack complex (C5b-9) in the RPE-choroid underlying the macula (FIGURE 1I) and extramacula (FIGURE 1J) from the same AMD donor eye. In the macula, intense anti-C5b-9 labeling is associated with drusen, Bruch's membrane, and the choroidal capillary endothelium. Anti-C5b-9 labeling outside the macula is restricted to a narrow zone in the vicinity of Bruch's membrane. Brown pigment in the RPE cytoplasm and choroid represents melanin pigmentation. Magnification bars = 20 μ m.

[0337] (K-L) Immunohistochemical location of C5b-9 in the macula from a donor with AMD (FIGURE 1K) and from a second donor without AMD (FIGURE 1L). Brown pigmentation in the RPE cytoplasm and choroid represents melanin. The anti-C5b-9 labeling is associated primarily with the choroidal capillary walls (black arrows) and the intercapillary pillars (white arrows). Labeling is much more intense in the AMD eye. Note the strong similarity to the anti-HF1 labeling pattern in the macula from the same donor, as shown in Figure G. Magnification bars. K = 15 μ m; L = 20 μ m.

The Retinal Pigment Epithelium is a Local Source of Factor H

[0338] Expression of *HF1* and *FHL1* in the RPE, RPE/choroid and retina was assessed by RT-PCR and DNA microarray analysis. Appropriately sized PCR products for both gene products are present in freshly isolated RPE and the RPE/choroid complex, but not neural retina, in human eyes derived from donors with and without AMD (FIGURE 2). Primers were chosen within exons 8, 9, 10A (the exon employed to generate the truncated isoform *FHL1*) and 10 of the *HF1* coding sequence. The PCR reactions were performed with cDNA prepared from RNA extracted from human neurosensory retina (lanes 2), RPE and choroid

(lanes 3), and freshly isolated RPE cells (lanes 4) derived from a donor with a clinically documented history of AMD. Genomic DNA was employed as a template for amplification (lanes 5); no template was added to the mixtures depicted in lanes 6. Lanes 1 contains the 100 bp ladder. Amplimers spanning from exon 8 to exon 10 (left panel), and from exon 9 to exon 10 (right panel), of *HF1* and from exon 8 to exon 10A (left panel), and exon 8 to exon 10A (right panel), of *FHL1* were of the expected sizes (376, 210, 424 and 248 bp, respectively). Transcripts for FHRs 1-5 are not detected in RPE or RPE/choroid, but FHRs 1-4 are detected in neural retina by RT-PCR (data not shown).

[0339] Gene expression array data derived from three platforms confirm that *HF1* and *FHL1* transcripts, but few if any of the *HF1*-related protein transcripts (*FHR1* being the possible exception), are expressed locally by RPE and choroid cells. Data derived from Incyte arrays probed with RPE/choroid cDNA derived from nine donors with AMD and three age-matched controls show elevated levels that average 2-3 times that of *HF1* mRNA in the donors with AMD. There is also a trend toward slightly higher levels in the macula regions as compared to the extramacular regions, although the difference is not statistically significant. The data generated from the examination of isolated RPE and adjacent RPE/choroid preparations from two donors with AMD and two age-matched control donors using Affymetrix arrays confirm the presence of *HF1* transcripts in these tissues and shows that a significant proportion of the *HF1* message is present within the RPE layer (data not shown).

Variants in HF1 are Associated with AMD and MPGN II

[0340] To test whether allele variants of *HF1* gene are associated with AMD, all 22 coding exons and 50-100bp flanking intronic sequences were screened, in a cohort of 404 AMD patients and 131 matched controls at the University of Iowa. A total of 26 sequence variants are detected; 17 SNPs in the coding region (cSNPs), including 5 synonymous and 12 non-synonymous substitutions, and 9 intronic SNPs (some of the variants shown in FIGURE 3). FIGURE 3 shows the approximate locations of 11 SNPs used in the analysis, the 20 short consensus repeats (SCRs), and the linkage disequilibrium (LD) blocks, and the approximate binding sites for pathogens and other substrates are depicted below the diagram based on previously published data (Zipfel et al., 2002; Rodriguez de Cordoba et al., 2004). cSNPs included previously described common non-synonymous variants, such as I62V in exon 2, Y402H in exon 9, and D936E in exon 18 (FIGURE 3). An example of a common intronic

SNP with a potentially functional effect is the IVS2-18insTT variant. Five rare (<0.5%) variants are also detected (data not shown) in both AMD patients and controls, excluding the possibility of the disease phenotype being caused by rare *HFI* alleles (i.e., disease-causing mutations). Detailed genotyping data were obtained for 6 SNPs in some or all of the 404 patients and 131 controls (TABLES 4 and 6A-6C) and association analyses were performed using a case-control study design. Highly significant associations are found with several individual variants, including I62V ($\chi^2=15.0$, $p=1.1 \times 10^{-4}$) and Y402H ($\chi^2=49.4$, $p=2.1 \times 10^{-12}$). The strongest association with AMD in this cohort is observed with a synonymous A473A variant in exon 10, resulting in an odds ratio (OR) of 3.42 (95% confidence interval (CI) [2.27-5.15]).

[0341] These results were confirmed in an independent cohort of AMD patients (n=550) and matched controls (n=275) obtained at Columbia University, New York (TABLE 4). The same two non-synonymous SNPs are also highly associated with AMD in this second cohort (I62V; $\chi^2=36.1$, $p=3.2 \times 10^{-7}$ and Y402H; $\chi^2=54.4$, $p=1.6 \times 10^{-13}$). In addition, several other intronic SNPs were selected based on frequency and the availability of commercial assays (for a total of 11 SNPs). The strongest association in this cohort is observed with SNP rs203674 in intron 10 ($\chi^2=66.1$, $p=4.29 \times 10^{-16}$). This variant shows an OR of 2.44 with AMD (95% CI=1.97-3.03). Although the OR is modest, the variant is very common; 30.5% of the cases are homozygous for allele B, but only 12.9% of the controls. The Q672Q and D936E alleles in exon 13 and 18 show no statistically significant association, suggesting that variation in the N-terminal half of *HFI*, which includes domains involved in pathogen and substrate molecule recognition (FIGURE 3, also see below), are associated with AMD. The two sets of data are strikingly similar, in that the genotyped SNPs are not only associated with AMD in a highly significant fashion, but the frequencies and extent of association are very similar in the two cohorts (TABLES 4 and 6).

[0342] The association is highly significant when the entire AMD patient cohorts are compared to controls (TABLE 4). When the major sub-phenotypes of AMD, such as the early AMD (eAMD, characterized by macular drusen and/or pigmentary abnormalities), CNV (neovascular membranes and/or disciform scars), and GA (geographic atrophy) are analyzed separately, the association is especially prominent in cases of eAMD and CNV. The GA group shows some deviation from the general trend in some cases, especially with the haplotype defined by exon 13 (Q672Q) and 18 (D936E) alleles (data not shown). While

this deviation may be significant in terms of varying etiology, it did not reach statistical significance, most likely due to relatively small numbers of patients with GA.

[0343] Linkage disequilibrium (LD) analysis showed extensive LD across the entire *HF1* gene (TABLE 2 and FIGURE 3). Three SNPs in the exon 2-3 region are in virtually complete LD as are the A307A and Y402H variants in exons 7 and 9, and the Q672Q and D936E variants in exon 13 and 18 (TABLE 6 and FIGURE 5). Haplotype estimation in cases and controls identified the most frequent at-risk haplotype in 49% of cases versus only in 26% of controls (OR=2.93 95%CI [2.29-3.74]). Homozygotes for this haplotype are present in 22.1% of the Columbia cases and 5.1% of the controls (OR=5.26, 95% CI [2.84-9.76]). Two common protective haplotypes are found in 30% of controls and 18% of cases (OR=0.476 95%CI [0.349-0.650] and OR=0.472, 95%CI [0.320-0.698]). These haplotypes differ only in the exon 2-3 locus SNPs and the intron 10 SNP. As shown in FIGURE 4 and TABLE 2, these protective haplotypes are closely related to each other and are both at least five steps away from the risk haplotype. Interestingly, the 3 SNPs, (promoter -257C>T, A473A, and D936E) previously shown to be associated with HUS are all on one relatively common haplotype (12%) that is neutral for AMD risk (see the discussion below). For each SNP we identified the base present in the consensus chimpanzee genome. The haplotype generated represents the likely ancestral human haplotype and is closely related to the protective haplotypes (data not shown).

[0344] SNPs IVS2-18insTT and Y402H were also genotyped in 20 unrelated MPGNII patients, 52 Rapanui natives and small cohorts of Hispanic Americans, African Americans and European Americans (TABLE 7). The frequency of the at-risk haplotype was estimated in samples from different populations from genotypes of the Y402H variant and/or the IVS10 locus. These include Rapanui natives over the age of 65 (AMD is extremely rare, and most likely absent, in this Easter Island population), controls (>65 years of age) from Columbia University, Hispanics general population, controls (>65 years of age) from the University of Iowa, African Americans general population, AMD cases from Columbia University, European Americans general population, AMD cases from the University of Iowa and individuals with MPGNII. N=number of individuals. In the MPGNII cohort, the frequency of the risk haplotype is approximately 70%. In addition, the risk haplotype appears to be lower in frequency in Hispanic Americans and African Americans (35-45%), groups with lower incidences of AMD. However, the number of samples typed in these populations is small. The Rapa Nui population on Easter Island has a remarkably low level of AMD. From

the analysis of 52 AMD-free Rapanui natives over the age of 65, we estimate the frequency of the risk haplotype to be only 19%.

Discussion

Factor H Polymorphisms and AMD

[0345] The data presented here links a major proportion of AMD cases in two independent cohorts to specific polymorphisms in the complement regulatory gene, Factor H (*HF1/CFH*) (Zipfel, 2001; Rodriguez de Cordoba et al., 2004). Haplotype analysis shows the most frequent at-risk haplotype to be present in almost half of individuals with AMD, compared to approximately 25% of controls. The frequencies and extent of SNP associations are very similar in the two cohorts and, genotyped SNPs show highly significant association with AMD in each. The associations are especially prominent in cases of early AMD or choroidal neovascularization, less so for geographic atrophy. The magnitude of the observed association of specific *HF1* haplotypes with AMD is striking when compared to those genetic abnormalities previously linked to AMD.

[0346] Further support for the conclusion that specific *HF1* haplotypes confer increased risk for an AMD disease phenotype was obtained by genotyping of SNPs IVS2-18insTT and Y402H in 20 unrelated patients with MPGNII, a renal disease associated with *HF1* mutations in which patients develop early onset macular drusen, and in 52 Rapanui natives, a race with a remarkably low incidence of AMD, if any. Approximately 70% of MPGN II patients and 19% of Rapanui were found to harbor the *HF1* at-risk haplotype in this study. Analysis of larger sample sets will be required to confirm these results, but the data do suggest that the *HF1* association with AMD may not be restricted to European-derived populations.

Protective haplotypes we also identified, further implicating *HF1* function in the pathogenetic mechanisms underlying AMD.

Functional Implications of Factor H Polymorphisms

[0347] Factor H deficiencies in humans are associated with MPGN II and atypical hemolytic uremic syndrome (aHUS) (Zipfel et al., 2001). *HF1* deficiency arises from mutations leading to protein truncations or amino acid substitutions that result in protein retention in the endoplasmic reticulum. Reduced levels of plasma *HF1* ensue, leading to uncontrolled activation of the alternative complement pathway with concomitant consumption of C3 and other complement components. The *HF1* mutations that lead to

aHUS, in contrast, are typically missense mutations that limit the complement-inhibitory functions of FH1 at the cell surface. Recent studies have revealed an association between three common SNPs and aHUS in individuals both with and without *FHI* mutations (Caprioli et al., 2003). Furthermore, insults such as infection have been documented to trigger the manifestation of aHUS.

[0348] Most of the genotyped SNPs of *HFI* are located within important functional domains of the encoded protein (FIGURE 3), which consists of 20 short consensus repeats (SCR) of 60 amino acids. The SCRs contain binding sites for C3b (SCR1-4, SCR12-14 and SCR19-20), heparin, sialic acid (SCR7, SCR13 and SCR19-20) and C-reactive protein (CRP) (SCR7). Therefore, SNPs located within the functional domains, although common, presumably affect protein function through variability in expression levels, binding efficiency, and other molecular properties. For example, the exon 2 I62V variant is located in SCR2, which is included in the first C3b binding site, and the exon 9 Y402H variant is within SCR7 domain, which binds both heparin and CRP. Intronic SNPs, such as the IVS2-18insTT variant, can affect splicing. For example, the analysis of the effect of the TT insertion (on the <https://splice.cmh.edu/server>) suggested a creation of a new cryptic splice acceptor 6 bp upstream of the natural acceptor site (data not shown). It is also possible that some of the studied SNPs affect the expression of *HFI* isoforms. For example, I62V is present in a predicted exon splice enhancer (Wang et al., 2004) (data not shown).

[0349] The functional consequences of common SNPs may be modest, since they are involved in late-onset phenotypes and not subjected to (rigorous) evolutionary constraint. *HFI* variants with more substantial effects, i.e., disease-causing mutations, are implicated in early-onset, severe (recessive) diseases, such as HFI deficiency and aHUS (Zipfel et al., 2002; Rodriguez de Cordoba et al., 2004; Caprioli et al., 2003; Zipfel, 2001). Of interest is the fact that true disease-causing mutations have been identified in only about 25% to 35% of HUS patients after complete screening of the *HFI* gene (Caprioli et al., 2003). At the same time, a disease-associated haplotype defined by variants -257C>T (promoter), A473A (exon 13) and D936E (exon 18) has been identified in HUS patients, predominantly in those persons with no disease-causing mutations (Caprioli et al., 2003). Moreover, the same study identified several families where affected probands had inherited a mutated allele from one parent and a susceptibility allele, from another. By contrast, healthy siblings of affected probands, carriers of the disease-causing mutation, had inherited a protective allele. In affected individuals from these families a disease-triggering effect, specified as bacterial or

viral infection in >60% of cases, was identified in >80% of all cases and in >90% of cases with no apparent disease-associated mutation (Caprioli et al., 2003).

[0350] Together, these data strongly suggest that an at-risk *HF1* haplotype in combination with an infectious triggering event is sufficient for disease manifestation. Interestingly, the at-risk *HF1* haplotype in HUS (mainly C-terminal) does not overlap with that in AMD and/or MPGNII (TABLE 2), suggesting different triggers in HUS as opposed to MPGNII and AMD. Observation of early-onset drusen in MPGNII and those of the same composition in AMD, but not in HUS, support a different etiology for these diseases.

[0351] Disease-causing mutations in *HF1* are rare in MPGN II and have not been reported in AMD, nor did we find them after extensive screening in this study. However, we observed the same at-risk haplotype at a frequency of 70% of patients with MPGN II and approximately 50% in patients with AMD. These data are consistent with an at-risk *HF1* haplotype that, if triggered by an infectious agent, substantially increase one's susceptibility to disease. The combined effect of these factors determines the severity the resulting phenotype, ranging from AMD to MPGNII.

[0352] Evolutionary analysis of the *HF1* haplotype indicates that the risk haplotype has evolved significantly from the ancestral haplotype found in the chimpanzee. FIGURE 4 depicts a haplotype network diagram of *HF1* SNPs which shows the relationship between the haplotypes and the size of the circles is proportional to the frequency of the haplotype. The large filled-in circle represents the major risk haplotype, the vertically-lined circles are the two significant protective haplotypes, and the large open circle is the haplotype that contains the three SNPs associated with atypical hemolytic uremic syndrome (HUS), which is neutral for AMD risk. The putative ancestral haplotype is also indicated. It is possible that different forms of the *HF1* gene emerged in response to pathogens that activate the alternative complement pathway. Weakly acting *HF1* haplotypes could provide reduced complement inhibition and stronger protection against bacterial infection. However, such weak alleles could have the consequence of predisposing individuals to the consequences of complement system dysfunction. It is interesting that the AMD risk haplotype is principally different in the 5' end of the gene that produces both the full length HF1 and the FHL1 protein. In contrast the HUS mutations cluster in the 3' portion of the gene that is found only in HF1. Therefore, it will be important to determine the role of these two forms of the protein in disease.

Biological Model of Factor H Dysfunction in AMD

[0353] One of the primary functions of the complement system is to provide defense against such infectious agents. It mediates the opsonization and lysis of microorganisms, removal of foreign particles, recruitment of inflammatory cells, regulation of antibody production, and elimination of immune complexes (Morgan et al., 1991; Kinoshita, 1991). Activation of the system triggers a sequential, amplifying, proteolytic cascade that gives rise to modifications of activating surfaces, to the release of soluble anaphylatoxins that stimulate inflammatory cells, and ultimately, to formation of the membrane attack complex (MAC), a macromolecular complex that promotes cell lysis through the formation of transmembrane pores. Uncontrolled activation of complement can lead to bystander damage to host cells and tissues. As a result HF1, as well as other circulating and membrane-associated proteins, have evolved to modulate the system (Morgan, 1999).

[0354] A spectrum of complement components have been identified either within drusen (and/or the RPE cells that flank or overlie them), along Bruch's membrane, and/or on the choroidal endothelial cell membrane (Hageman et al., 2001; Mullins et al., 2000; Mullins et al., 2001; Anderson et al., 2002; Johnson, et al., 2000; Johnson et al., 2001; Crabb et al., 2002; Johnson et al., 2002; Mullins et al., 1997). These include terminal pathway complement components, activation-specific fragments of the terminal pathway, as well as various complement modulators. There is evidence that cell-mediated events may also contribute to this process (Penfold et al., 2001; Seddon et al., 2004; Miller et al., 2004).

[0355] We now show that HF1 is also a constituent of drusen in human donors with a prior history of AMD. Secondly, we show that HF1 co-localizes with its ligand C3b in amyloid-containing substructural elements within drusen, further implicating these structures as candidate complement activators (Anderson et al., 2004; Johnson et al., 2002). We also demonstrate that HF1 and the MAC, as shown by C5b-9 immunoreactivity, co-distribute at the RPE-choroid interface and that these deposits are more robust in eyes from donors with prior histories of AMD. Finally, HF1 and C5b-9 immunoreactivities are more intense in the macula compared to more peripheral locations from the same eye. All of these findings are consistent with the fact that AMD pathology is manifested primarily in the macula and with the conclusion that complement activation at the level of Bruch's membrane is a key element in the process of drusen formation as well as a contributing factor in the pathogenesis of AMD Hageman et al., 2001; Anderson et al., 2002).

[0356] The distribution of HFI at the RPE-choroid interface is strikingly similar to that of C5b-9, implying that significant amounts of the MAC are generated and deposited at the RPE-choroid interface. This suggests that the protein associated with the at-risk *HFI* haplotype(s) may undergo a reduction in its normal ability to attenuate complement activation. Thus, the *HFI* variants associated with AMD may put RPE and choroidal cells at sustained risk for alternative pathway-mediated complement attack, drusen formation and the disruption of Bruch's membrane integrity that is associated with late-stage neovascular AMD. Since Bruch's membrane is significantly thinner in the macula than elsewhere (Chong et al., 2005), it may be more susceptible to subsequent neovascular invasion. Because Bruch's membrane is significantly thinner in the macula than in the periphery, it may be more likely to become degraded to an extent that it is susceptible to neovascular invasion.

[0357] In summary, the results of this investigation provide strong evidence that common haplotypes in *HFI* predispose individuals to AMD. We propose that alterations in genes that regulate the alternative pathway of the complement cascade, in combination with events that activate the system, underlie a major proportion of AMD in the human population.

EXAMPLE 2

Variations in the Complement Regulatory Genes Factor H (*CFH*) and Factor H Related 5 (*CFHR5*) are Associated with Membranoproliferative Glomerulonephritis Type II (Dense Deposit Disease)

Introduction

[0358] The membranoproliferative glomerulonephritides are diseases of diverse and often obscure etiology that account for 4% and 7% of primary renal causes of nephrotic syndrome in children and adults, respectively (Orth et al., 1998). Based on renal immunopathology and ultrastructural studies, three subtypes are recognized. Membranoproliferative glomerulonephritis (MPGN) types I and III are variants of immune complex-mediated disease; MPGNII, in contrast, has no known association with immune complexes (Appel et al., 2005).

[0359] MPGNII accounts for less than 20% of cases of MPGN in children and only a fractional percentage of cases in adults (Orth et al., 1998; Habib et al., 1975; Habib et al., 1987). Both sexes are affected equally, with the diagnosis usually made in children between the ages of 5-15 years who present with non-specific findings like hematuria, proteinuria,

acute nephritic syndrome or nephrotic syndrome (Appel et al., 2005). More than 80% of patients with MPGNII are also positive for serum C3 nephritic factor (C3NeF), an autoantibody directed against C3bBb, the convertase of the alternative pathway of the complement cascade (Schwartz et al., 2001). C3NeF is found in up to one-half of persons with MPGN types I and III and also in healthy individuals, making the electron microscopic demonstration of dense deposits in the glomerular basement membrane (GBM) necessary for a definitive diagnosis of MPGN II (Appel et al., 2005). This morphological hallmark is so characteristic of MPGN II that the disease is more accurately referred to as Dense Deposit Disease (MPGNII/DDD) (FIGURE 12).

[0360] Spontaneous remissions of MPGNII/DDD are uncommon (Habib et al., 1975; Habib et al., 1987; Cameron et al., 1983; Barbiano di Belgiojoso et al., 1977). The more common outcome is chronic deterioration of renal function leading to end-stage renal disease (ESRD) in about half of patients within 10 years of diagnosis (Barbiano di Belgiojoso et al., 1977; Swainson et al., 1983). In some patients, rapid fluctuations in proteinuria occur with episodes of acute renal deterioration in the absence of obvious triggering events; in other patients, the disease remains stable for years despite persistent proteinuria.

[0361] C3NeF persists throughout the disease course in more than 50% of patients with MPGNII/DDD (Schwartz et al., 2001). Its presence is typically associated with evidence of complement activation, such as a reduction in CH50, decrease in C3, increase in C3dg/C3d and persistently high levels of activation of the alternative pathway of the complement cascade. C3, the most abundant complement protein in serum (~1.2 mg/ml), normally undergoes low levels of continuous autoactivation by hydrolysis of its thioester in a process known as tick-over. C3 hydrolysis induces a large conformational protein change, making C3(H₂O) similar to C3b, a cleavage product of C3. C3(H₂O) associates with factor B to form C3(H₂O)Bb, which cleaves C3 to C3b in an amplification loop that consumes C3 and produces C3bBb² (FIGURE 13).

[0362] In MPGNII/DDD, C3NeF binds to C3bBb (or to the assembled convertase) to prolong the half-life of this enzyme, resulting in persistent C3 consumption that overwhelms the normal regulatory mechanisms to control levels of C3bBb and complement activation (Appel et al., 2005). Normal control involves at least seven proteins, four of which are present in serum (complement Factor H (CFH), complement factor H-like protein 1 (CFHL1), complement factor I (CFI) and C4 binding protein (C4BP)) and three of which are cell membrane-associated (membrane co-factor protein (MCP, CD46), decay accelerating

factor (DAF, CD55) and complement receptor 1 (CR1, CD35)) (Appel et al., 2005; Meri et al., 1994; Pascual et al., 1994).

[0363] Of particular relevance to MPGNII/DDD is Factor H, one of 7 proteins in the Factor H family. In pigs and mice, its deficiency is associated with the development of renal disease that is similar at the light and electron microscopic level to MPGNII/DDD, and in humans its deficiency as well as mutations in the Factor H gene have been reported in patients with MPGNII/DDD (Meri et al., 1994; Dragen-Durey et al., 2004; Zipfel et al., 2005) (FIGURE 14).

[0364] The other 6 members of the Factor H family include FHL1, which is a splice isoform of Factor H, and five CFH-related proteins encoded by distinct genes (*CFHR1-5*). There is little known about the latter five proteins, although they do show varying degrees of structural similarity to Factor H (Appel et al., 2005). Most interesting in this group with respect to MPGNII/DDD is *CFHR5*, because it shows the highest similarity to Factor H and has been demonstrated in renal biopsies of patients with other types of glomerulonephritis (Appel et al., 2005; Murphy et al., 2002). *In vitro* studies have also shown that V is present on surfaces exposed to complement attack, suggesting a possible role in the complement cascade (Murphy et al., 2002).

[0365] A possible relationship between Factor H/ *CFHR5* and MPGNII/DDD is further strengthened by the observation that patients with MPGNII/DDD develop an ocular phenotype called drusen. Drusen result from the deposition of abnormal extracellular deposits in the retina within the ocular Bruch's membrane beneath the retinal pigment epithelium. The drusen of MPGNII/DDD are clinically and compositionally indistinguishable from drusen that form in age-related macular degeneration (AMD) (Mullins et al., 2001; Anderson et al., 2002), which is the most common form of visual impairment in the elderly (Klein et al., 2004; van Leeuwen et al., 2003). The single feature that distinguishes these two types of drusen is age of onset - drusen in MPGNII/DDD develop early, often in the second decade of life, while drusen in AMD are found in the elderly.

[0366] Four recent studies have implicated specific allele variants of Factor with AMD, suggesting that subtle differences in Factor H-mediated regulation of the alternative pathway of complement may play a role in a substantial proportion of AMD cases (Hageman et al., 2005; Edwards et al., 2005; Haines et al., 2005; Klein et al., 2005). One of these studies also showed that MPGNII/DDD and AMD patients segregate several of the same Factor H risk

alleles (Hageman et al., 2005). In this study, we sought to refine the association of allele variations of Factor H and *CFHR5* with MPGNII/DDD.

Materials and Methods

[0367] *Patients and Controls.* Patients with biopsy-proven MPGNII/DDD were ascertained in nephrology divisions and enrolled in this study under IRB-approved guidelines. The control group was ascertained from ethnically-matched but not age-matched unrelated persons in whom AMD had been excluded by ophthalmologic examination.

[0368] *Mutation Screening and Analysis.* Coding and adjacent intronic regions of Factor H and *CFHR5* were PCR amplified for 35 cycles of 30 seconds each at 94°C denaturing, 61°C annealing and 70°C extension. The sequences of primers used to amplify the Factor H and *CFHR5* coding sequences are shown in TABLES 10 and 11, respectively. Product generation was verified by agarose gel electrophoresis and amplicons were then bi-directionally sequenced in patients with MPGNII/DDD. All novel and reported SNPs identified through data mining (Ensemble database, dbSNP, Applied Biosystems) were typed in the control population by denaturing high performance liquid chromatography (DHPLC) (Tables 9 and 10). In brief, DHPLC analysis of each amplicon was performed at three different temperatures. Amplicons were analyzed using Wavemaker software to estimate optimal temperature, run time and acetonitrile gradient. Predicted temperatures were bracketed by +/- 2°C to optimize sensitivity and maximize the likelihood that novel mutations would be detected (Prasad et al., 2004).

[0369] *Haplotype Analysis.* Construction of block structures with distribution of haplotypes was completed using Haploview, a publicly available software program developed at the Whitehead Institute (<http://www.broad.mit.edu/mpg/haploview/>) (see Barrett et al.). Two datasets, one consisting of each control's sex and genotype, and the other describing marker information including SNP identification and chromosomal location, were assimilated in Excel files, which were up-loaded into the Haploview program. The output consisted of linkage disequilibrium (LD) plots and the corresponding population frequencies with crossover percentages.

[0370] *Statistical Analysis.* The chi-square test of independence was used to detect differences in SNP frequencies between patients with MPGNII/DDD and controls. P-values ≤ 0.05 were considered significant. The LD plots for Factor H and *CFHR5* were created using the control population.

Results

[0371] *Patients and Controls.* Twenty-two patients with biopsy-proven MPGNII/DDD and 131 persons without AMD participated in this study. Mean age of the control group was 78.4 years, reflecting our ascertainment criterion to exclude AMD.

[0372] *Factor H, CFHR5 and MPGN II/DDD.* Allele frequencies of four of seven Factor H SNPs genotyped in the MPGNII/DDD patient group and the control population showed a significant association with the MPGNII/DDD disease phenotype at $p < 0.05$. These SNPs included exon 2 I62V, IVS 2-18insTT, exon 9 Y402H and exon 10 A473A. Allele frequencies for exon 7 A307A, exon 13 Q672Q and exon 18 D936E were not significantly different between groups (TABLES 11 to 13).

[0373] Five *CFHR5* SNPs were genotyped in the MPGNII/DDD patient group and control population, including one non-synonymous SNP (exon 2 P46S), two promoter SNPs (-249T>C, -20 T>C) and two intronic SNPs (IVS1+75T>A, IVS2+58C>T). Allele frequencies of three SNPs – exon 2 P46S, -249T>C and -20 T>C – were significantly different between groups at $p < 0.05$ (TABLES 14 and 15).

[0374] *Haploblocks.* Haplotype blocks showed that A307A and Y402H are in linkage disequilibrium in Factor H while -249T>C and -20T>C are in linkage disequilibrium in *CFHR5* (FIGURE 15).

Discussion

[0375] The alternative pathway of complement represents an elegant system to protect humans from pathogens. Its central component, C3, circulates at a high concentration in plasma and is distributed throughout body fluids (Walport, 2001). Its activation creates a toxic local environment that damages foreign surfaces and results in the elimination of microbes. To prevent unrestricted complement activation, host cells and tissue surfaces down-regulate the amplification loop using a combination of surface-attached and membrane-bound regulators of complement. Some host cells express a single membrane-bound regulator of complement in high copy number, while other cells express several membrane-bound regulators and also attach soluble fluid-phase regulators. A few tissues lack membrane-bound regulators and depend exclusively on the attachment of soluble regulators (Appel et al., 2005).

[0376] In the kidney, endothelial and mesangial cells express two membrane-bound regulators of complement, MCP and DAF (van den Dobbela et al., 1994; Timmerman et al., 1996). Podocytes express four: MCP, DAF, CR1 and CD59. Both mesangial cells and podocytes also secrete the soluble regulator, Factor H, which is up-regulated in membranous nephropathy in response to complement activation and inflammation (Angaku et al., 1998; Bao et al., 2002). Factor H acts in an autocrine fashion by binding directly to the secreting mesangial cells and podocytes.

[0377] The GBM, in contrast, is unique. It lacks endogenous membrane-bound regulators to protect it from complement-mediated injury, however its highly negatively charged surface binds and absorbs Factor H (Zipfel et al., 2005). The dependency of the GBM on Factor H for local complement control is consistent with the finding of pathologic mutations in Factor H in a few persons with MPGNII/DDD (Ault et al., 1997; Dragen-Durey et al., 2004).

[0378] Our data identifying several allele variants of Factor H and *CFHR5* associated with MPGNII/DDD is consistent with the hypothesis that complement control plays a role in the pathogenesis of this disease. A comparison of our data with reported AMD data adds additional support, as the allele frequency for each of the identified at-risk SNP variants we observed in Factor H is higher in the MPGNII/DDD patient cohort than in the AMD patient cohort, and strong evidence implicates Factor H in AMD (Hageman et al., 2005; Edwards et al., 2005; Haines et al., 2005; Klein et al., 2005). Although it is not known whether the amino acid changes in exons 2 and 9 of Factor H impact function, these changes are found in domains that interact with C3b and heparin, and differences in C3b/C3d and heparin binding have been demonstrated with several amino acid changes in Factor H that are associated with another renal disease, atypical hemolytic uremic syndrome (Manuelian et al., 2003) (TABLES 12 and 13).

[0379] With the exception of Factor H, the function of other members of the Factor H - related family is largely unknown and their expression patterns have not been explored, however studies of *CFHR5* have shown that it has properties similar to Factor H, including heparin, CRP and C3b binding (Murphy et al., 2002) (FIGURE 14). This similarity suggests that like Factor H, *CFHR5* plays a role in MPGNII/DDD. Consistent with this is our finding of *CFHR5* expression in renal biopsies from two patients with MPGNII/DDD (data not shown).

[0380] Our genotyping data show that some allele variants of *CFHR5* preferentially associate with the MPGNII/DDD disease phenotype. Included are two SNPs in the promoter

region of *CFHR5* which could affect transcription, one by removing a binding site for C/EBP β and the other by adding a GATA-1 binding site. The other significant association changes a proline to serine in exon 2. Since exons 1 and 2 of *CFHR5* encode a domain homologous to short consensus repeat 6 (SCR6) of Factor H, which is integral to heparin and CRP binding, this change could affect complement activation and control.

EXAMPLE 3

PRODUCTION OF PROTECTIVE FORM OF FACTOR H PROTEIN

[0381] An exemplary protective form of human complement factor H (CFH) was prepared based on haplotype H2 (FIGURE 5). Briefly, RNA was isolated from ocular tissues (RPE/choroid complexes) of four donors. The RNA was amplified by reverse transcription-polymerase chain reaction using the following primers:

5'-GAAGATTGCAATGAACTTCCTCCAAG-3' [SEQ ID NO:331]

5'-AAGTTCTGAATAAAGGTGTGC-3' [SEQ ID NO:332].

RT-PCR reactions were performed using Superscript III One-Step High Fidelity with Platinum Taq, as described by the manufacturer (Invitrogen, Carlsbad, CA). The appropriate sized products (3,769 bp) were excised from agarose gels and isolated using spin columns.

[0382] The PCR products were cloned using the TOPO-TA cloning system, as recommended by the manufacturer (Invitrogen) in the vector pCR2.1-TOPO. The complete genetic sequences of the clones derived from each of the four patients were determined by direct sequencing. The DNA derived from one patient (patient #498-01) had the fewest number of nucleotide polymorphisms relative to that of an exemplary protective reference sequence (H2), although this DNA encoded the risk sequence at amino acid position 402 (histidine) and encoded valine at amino acid position 62. To prepare a gene encoding a protective form of CFH we changed the bases encoding amino acid 62 such that it coded for isoleucine and those at position 402 such that they encoded a tyrosine, using the QuikChange Mutagenesis system (Stratagene, La Jolla, CA), resulting in SEQ ID NO:335. The amino acid at position 1210 of this protein is arginine. The oligonucleotides employed were as follows (plus the appropriate antisense version):

62: 5'-TATAGATCTCTTGGAATATAATAATGGTATGCAGG-3' [SEQ ID NO:333]

402: 5'-ATGGATATAATCAAAATTATGGAAGAAAGTTTGTAC-3' [SEQ ID NO:334]

The fidelity of the introduced mutations were confirmed by direct sequencing of the entire gene. The resulting protective gene was cloned into the eukaryotic expression vector

pcDNA3.1 (Invitrogen) under control of the cytomegalovirus promoter. This expression vector was transfected into the human lung carcinoma cell line A549 (ATCC, Manassas, VA) using the Exgen 500 transfection reagent (Fermentas, Hanover, MD). Subsequent to transfection, cells were grown in serum-free media (Hybridoma-SFM, Invitrogen).

[0383] Supernatants were collected 48 hours after transfection and subjected to Western blot analyses. The presence of the appropriate-sized product (approximately 150kDa) was confirmed using monoclonal and polyclonal antibodies directed against human CFH (Quidel, San Diego, CA).

[0384] Patient #498-01 (62I, 402Y) CFH Gene [SEQ ID NO:335]

```
AGTTAGCTGGTAAATGTCCTCTTAAAAGATCCAAAAAATGAGACTTCTAG
CAAAGATTATTTGCCTTATGTTATGGGCTATTTGTGTAGCAGAAGATTGC
AATGAACTTCCTCCAAGAAGAAATACAGAAATTCTGACAGGTTCTGGTC
TGACCAAACATATCCAGAAGGCACCCAGGCTATCTATAAATGCCGCCCTG
GATATAGATCTCTTGGAAATATAATAATGGTATGCAGGAAGGGAGAATGG
GTTGCTCTTAATCCATTAAGGAAATGTCAGAAAAGGCCCTGTGGACATCC
TGGAGATACTCCTTTTGGTACTTTTACCCTTACAGGAGGAAATGTGTTTG
AATATGGTGTAAAAGCTGTGTATACATGTAATGAGGGGTATCAATTGCTA
GGTGAGATTAAATTACCGTGAATGTGACACAGATGGATGGACCAATGATAT
TCCTATATGTGAAGTTGTGAAGTGTTTACCAGTGACAGCACCAGAGAATG
GAAAAATTGTCAGTAGTGCAATGGAACCAGATCGGGAATACCATTTTGGG
CAAGCAGTACGGTTTGTATGTAACCTCAGGCTACAAGATTGAAGGAGATGA
AGAAATGCATTGTTTCAGACGATGGTTTTTGGAGTAAAGAGAAAACCAAAGT
GTGTGGAAATTTTCATGCAATCCCCAGATGTTATAAATGGATCTCCTATA
TCTCAGAAGATTATTTATAAGGAGAATGAACGATTTCAATATAAATGTAA
CATGGGTTATGAATACAGTGAAAGAGGAGATGCTGTATGCACTGAATCTG
GATGGCGTCCGTTGCCTTCATGTGAAGAAAAATCATGTGATAATCCTTAT
ATTCCAAATGGTGACTACTCACCTTTAAGGATTAAACACAGAACTGGAGA
TGAAATCACGTACCAGTGTAGAAATGGTTTTTATCCTGCAACCCGGGGAA
ATACAGCAAAATGCACAAGTACTGGCTGGATACCTGCTCCGAGATGTACC
TTGAAACCTTGTGATTATCCAGACATTAAACATGGAGGTCTATATCATGA
GAATATGCGTAGACCATACTTTCCAGTAGCTGTAGGAAAATATTACTCCT
ATTACTGTGATGAACATTTTGGAGACTCCGTCAGGAAGTTACTGGGATCAC
ATTCATTGCACACAAGATGGATGGTCGCCAGCAGTACCATGCCTCAGAAA
ATGTTATTTTCCTTATTTGGAAAATGGATATAATCAAAATTATGGAAGAA
AGTTTGTACAGGGTAAATCTATAGACGTTGCCTGCCATCCTGGCTACGCT
CTTCCAAAAGCGCAGACCACAGTTACATGTATGGAGAATGGCTGGTCTCC
TACTCCCAGATGCATCCGTGTCAAAACATGTTCCAAATCAAGTATAGATA
TTGAGAATGGGTTTATTTCTGAATCTCAGTATACATATGCCTTAAAAGAA
AAAGCGAAATATCAATGCAAACCTAGGATATGTAACAGCAGATGGTGAAAC
ATCAGGATCAATTACATCTGGGAAAGATGGATGGTCAGCTCAACCCACGT
GCATTAAATCTTGTGATATCCAGTATTTTATGAATGCCAGAACTAAAAAT
GACTTCACATGGTTTAAAGCTGAATGACACATTGGACTATGAATGCCATGA
TGGTTATGAAAGCAATACTGGAAGCACCCTGGTTCCATAGTGTGTGGTT
ACAAATGGTTGGTCTGATTTACCCATATGTTATGAAAGAGAATGCGAAGTT
```

CCTAAAATAGATGTACACTTAGTTCCTGATCGCAAGAAAGACCAGTATAA
AGTTGGAGAGGTGTTGAAATTCTCCTGCAAACCAGGATTTACAATAGTTG
GACCTAATTCCGTTCAAGTGCTACCACTTTGGATTGTCTCCTGACCTCCCA
ATATGTAAAGAGCAAGTACAATCATGTGGTCCACCTCCTGAACTCCTCAA
TGGGAATGTTAAGGAAAAAACGAAAGAAGAATATGGACACAGTGAAAGTGG
TGGAATATTATTGCAATCCTAGATTTCTAATGAAGGGACCTAATAAAATT
CAATGTGTTGATGGAGAGTGGACAACCTTTACCAGTGTGTATTGTGGAGGA
GAGTACCTGTGGAGATATACCTGAACTTGAACATGGCTGGGCCCAGCTTT
CTTCCCCTCCTTATTACTATGGAGATTCAGTGGAATTC AATTGCTCAGAA
TCATTTACAATGATTGGACACAGATCAATTACGTGTATT CATGGAGTATG
GACCCAACCTCCCCAGTGTGTGGCAATAGATAAACTTAAGAAGTGCAAAT
CATCAAATTTAATTATACTTGAGGAACATTTAAAAACAAGAAGGAATTC
GATCATAATTCTAACATAAGGTACAGATGTAGAGGAAAAAGAAGGATGGAT
ACACACAGTCTGCATAAATGGAAGATGGGATCCAGAAGTGAACCTGCTCAA
TGGCACAAATACAATTATGCCCACCTCCACCTCAGATTCCCAATTCTCAC
AATATGACAACCACACTGAATTATCGGGATGGAGAAAAAGTATCTGTTCT
TTGCCAAGAAAATTATCTAATTCAGGAAGGAGAAGAAATTACATGCAAAG
ATGGAAGATGGCAGTCAATACCACTCTGTGTTGAAAAAATTCATGTTCA
CAACCACCTCAGATAGAACACGGAACCATTAATTCATCCAGGTCTTCACA
AGAAAGTTATGCACATGGGACTAAATTGAGTTATACTTGTGAGGGTGGTT
TCAGGATATCTGAAGAAAATGAAACAACATGCTACATGGGAAAATGGAGT
TCTCCACCTCAGTGTGAAGGCCTTCTTGTAAATCTCCACCTGAGATTTCT
TCATGGTGTGTAGCTCACATGTCAGACAGTTATCAGTATGGAGAAGAAG
TTACGTACAAATGTTTTGAAGGTTTTGGAATTGATGGGCCTGCAATTGCA
AAATGCTTAGGAGAAAAATGGTCTCACCCTCCATCATGCATAAAACAGA
TTGTCTCAGTTTACCTAGCTTTGAAAATGCCATACCCATGGGAGAGAAGA
AGGATGTGTATAAGGCGGGTGAGCAAGTGACTTACACTTGTGCAACATAT
TACAAAATGGATGGAGCCAGTAATGTAACATGCATTAATAGCAGATGGAC
AGGAAGGCCAACATGCAGAGACACCTCCTGTGTGAATCCGCCCACAGTAC
AAAATGCTTATATAGTGTGCGAGACAGATGAGTAAATATCCATCTGGTGAG
AGAGTACGTTATCAATGTAGGAGCCCTTATGAAATGTTTGGGGATGAAGA
AGTGATGTGTTTAAATGGAAACTGGACGGAACCACCTCAATGCAAAGATT
CTACAGGAAAATGTGGGCCCCCTCCACCTATTGACAATGGGGACATTACT
TCATTCCCGTTGTCAGTATATGCTCCAGCTTCATCAGTTGAGTATCAATG
CCAGAACTTGTATCAACTTGAGGGTAACAAGCGAATAACATGTAGAAATG
GACAATGGTCAGAACCACCAAATGCTTACATCCGTGTGTAATATCCCGA
GAAATTATGGAAAATTATAACATAGCATTAAGGTGGACAGCCAAACAGAA
GCTTTATTCGAGAACAGGTGAATCAGTTGAATTTGTGTGTAAACGGGGAT
ATCGTCTTTCATCACGTTCTCACACATTGCGAACAACATGTTGGGATGGG
AAACTGGAGTATCCAACCTGTGCAAAAAGATAGAATCAATCATAAAGTGC
ACACCTTTATTCAGAACTT

TABLE 1A

dbSNP No.	Location	Sequence Spanning Polymorphism	SEQ ID No.	AA Change	Allele Freq. CTL	Allele Freq. AMD	Chi2 & P Value
	Promoter 1	GGGTTTCTGGGATGATTA[A/G]ATGTTCTAGTGTTCACCTT CCCCAAGACCCACATTA[T/C]TACAGTACCAAACTGGA	9		1-0.944:2-0.056	1-0.96:2-0.04	
rs3753394	Promoter 4	TTATGAATCCAGAGGATAT[C/T]ACCACTCTGATTTGCACA AAATCTTAGCTCTCTCTATA[G/A]TGGTCGACACTAAACGGT	10		1-0.31:2-0.69	1-0.25:2-0.75	6.485:0.039
rs529825	Intron 1	AGTCCAGTTTACACACATAC[G/A]ATACACATACCATTCGCA TCAGGTCAAATGTCTCTG[C/T]TATCTGATGGGTACGGTT	11		1-0.74:2-0.26	1-0.84:2-0.16	28.07:2.18E-06
rs800292	Exon 2	GATATAGTCTCTTGGAAAT[G/A]TAATAATGSTATGAGGAG CTATATCTAGAGACCTTA[C/T]ATTATTACCATACGCTTC	12	I62V	1-0.78:2-0.22	1-0.91:2-0.09	16.19:5.74E-05
	Intron 2	TAATTCATACCTTTTCTTTT[-/TT]CGTTTAGAAGCCCTGTG ATTAGTATTCAGAAAATAA[-/AA]GCAAAATCTTCGGGAC	13		1-0.77:2-0.23	1-0.89:2-0.11	22.19:2.47E-06
rs3766404	Intron 6	AAAGGATACATTTAGGACT[C/T]ATTGTTGAGTTAGTCTCACA TTTCTTATGTATATCCCTGA[G/A]TAACCTCAATCAGCTGT	14		1-0.83:2-0.17	1-0.91:2-0.09	23.82:6.71E-06
rs1061147	Exon 7	CAACCCGGGGAATACAGC[A/C]AAATGCACATGACTGCTG GTGGGCCCCCTTATGCTG[T/G]TTTACGTGTCATGACCGAC	15	A307A	1-0.34:2-0.66	1-0.59:2-0.41	50.39:1.26E-12
rs1061170	Exon 9	AAATGGATATATCAAAAT[T/C]ATGGAGAGAACTTTGTACAG TTTTACCTATATTAAGTTTA[A/G]TACCTCTTTCACATCTC	16	Y402H	1-0.66:2-0.34	1-0.46:2-0.54	55.20:1.03E-12
rs2274700	Exon 10	TATCCCTTAAGAGAAAGG[C/A]AAATATCATCTCAGCTAGG ATACGGAATTTCTTTTTCG[C/T]TTTATGATGACTGTTGATCC	17	A473A	1-0.54:2-0.46	1-0.80:2-0.20	36.48:1.58E-09
	Exon 10A	CAGCTTCAGTGTCAAGA[-/N]TGACAAGGGCCATGACACCC GTGCACTCACCTAGTTCT[-/N]ACTGTTCCCGGTACCTGG	18		1-1.00:2-0.00	1-0.933:2-0.067	
rs203674	Intron 10	ACGCTACCTATTATAGTA[C/T]ATCTTATCAATAAAGCTTTT TGCCATGATTAATTAATCAT[C/A]TAGATTAGTTATTTGAAA	19		1-0.66:2-0.34	1-0.44:2-0.56	66.97:2.86E-15
rs203674	Intron 10*	AAAGCTTATGATAGAT[A/C]TACTAATAAATAGGTACCGT	63		1-0.66:2-0.34	1-0.44:2-0.56	66.97:2.86E-15
rs3753396	Exon 13	AAAGGCTTAATAAATCA[A/G]TGTGTTGATGGAGATGGAC TTCCCTGATTAATTTAAGT[T/C]ACACAACCTCTCTACCTG	20	Q672Q	1-0.84:2-0.16	1-0.86:2-0.14	0.308:0.579
rs375046	Intron 15	TTTTTTTATTTTATATTA[A/C]AATTAATTAATTTTAAAT AAAAAATAAATAATATTT[G/T]TAATTAATTAATTAATTA	21		1-0.87:2-0.13	1-0.85:2-0.14	
rs1065489	Exon 18	CCTTGGAAATCTCCACCTGA[G/T]ATTCTCATGCTGTGAGC GGACATTTAGAGGAGGACT[C/A]TAAAGAGTACCAACATCG	22	D936E	1-0.87:2-0.13	1-0.86:2-0.15	0.155:0.694
	Exon 22	GGGATATCGTCTTTTCA[A/C]TCTCTCACACATTCGACACA CCCTATAGCAGAAAGTAGT[G/A]CAGAGTGTGTAACCTGT	23	R1210C	1-1.00:2-0.00	1-0.95:2-0.05	

(*)Shows the non-coding strand of the genomic sequence.

TABLE 1B

SNP name	Interrogated Sequence	SEQ ID NO:	Chimp	Location	AA
rs3753394	AAATCCAGAGGATATCTACGAGCTGCTGATTT	24	C	Promoter	
rs529825	AATGGGTAAGTCTATCTGTACTGTGTAAGCTT	25	T	Intron 1	
rs800292	TGCATACCATTATTAICTATTTCCAGAGATCT	26	T	Exon 2	I 62V
Intron 2 InsTT	ACATACTAATTCATAAGCTTTTCTTTTTCGTTTAG	27		Intron 2	
rs3766404	AATACATTAGGACTTCATTTGAAGTTAGTGT	28	C	Intron 6	
rs1061147	CCGGGGAATACAGCCATAAATGCCACAAGTACT	29	A	Exon 7	A307A
rs1061170	GGATATAATCAAAATTCATGGAAGAAAGTTTG	30	T	Exon 9	Y402H
rs2274700	CITAAAAGAAAAGCGCAATAATATCAATGCAAA	31	G	Exon 10	A473A
rs203674	CTTTATTGATTAGATATGCTACTAATAATAGGT	32	A	Intron 10	
rs3753396	ACCTAATAAATTCATGATGTTGATGGAGAG	33	A	Exon 13	Q672Q
rs1065489	TAAATCTCCACCTGAGTATTTCTCATGGTGT	34	G	Exon 18	D936E

(2)

SNP name	Forward Primer or AOD number	SEQ ID NO:	Reverse Primer	SEQ ID NO:
rs3753394	C 2530387 10			
rs529825	C 2250476 10			
rs800292	C 2530382 10			
Intron 2 InsTT	ACTTGTTCCTCCACTCTCTAC	35	CCCTCTTTTCGTATGGACTAC	36
rs3766404	C 11890065 10			
rs1061147	TGAATCAGTACCGAGTGAGAAATGG	37	CAGGTATCCAGCCAGTACTTGT	38
rs1061170	CTTTATTATCATGTATGGTCTTAGGAAATGTTATTT	39	GGCAGGCAACGCTATAGATTACC	40
rs2274700	TCACCATCTGCTGTACATATCTCTAGT	41	TGGGTTTATTTCTGAATCTCAGTATACATATGC	42
rs203674	C 2530311 10			
rs3753396	C 2530296 10			
rs1065489	C 2530274 10			

(3)

SNP name	VIC Probe	SEQ ID NO:	FAM Probe	SEQ ID NO:
rs1061147	AATACAGCAAAATGC	43	ATACAGCCAAATGC	46
rs1061170	TTTCTTCCATGATTTTG	44	TTCTTCCATATTTTG	47
rs2274700	AAGAAAAAGCGGAATAT	45	AAGAAAAAGCAAAATAT	48

TABLE 1C

Location	Sequence Spanning Polymorphism	AA Position	SEQ ID NO:
Exon 2	CCAGGCTATCTATAAATGCC [G/A] CCCTGGATATAGATCTCTTG	R53H	49
Exon 3	TTGGTACTTTTACCCCTTACA [G/T] GAGGAAATGTGTTGAATAT	G100R	50
Exon 5	ACGATGGTTTTTGGAGTAAA [G/notG] AGAAACCAAAAGTGTGGGT	201	51
Exon 6	TTATTTATAAGGAGAAATGAA [C/notC] GATTCAATATAAATGTAAC	R232X	52
Exon 6	CACTGAATCTGGATGGCGTC [C/notC] GTTGCCTTCATGTGAAG (end Exon 6)	258	53
Exon 8	AAGATGGATGGTGCCAGCA [G/C] [-/C] TACCATGCCTCA (end Exon 8)	V383L	54
Exon 16	ACAATTATGCCACCTCCAC [C/G] TCAGATTCCTCAATTCACACA	815	55
Exon 17	CAACCACCTCAGATAGAACA [C/T] GGAACCATTAATTCATCCAG	H878H	56
Exon 17	GTCTTCACAAGAAAGTTATG [C/T] ACATGGGACTAAATTGAGTT	A892V	57
Exon 18	CACATGTCAGACAGTTATCA [G/T] TATGGAGAAGAAAGTTACGTA	Q950H	58
Exon 18	TCAGTATGGAGAAGAAGTTA [C/T] GTACAAATGTTTGAAGGTT	S956L	59
Intron 18	(begin IVS18) GTATGG [G/T] GCATTGAATTTTATTATATG		60
Exon 20	(begin Exon 20) ACACCTCCTGTGTG [A/T] ATCCGCCACACAGTACAAAAT	N1050Y	61
Exon 21	CTTGTATCAACTTGAGGGTA [-/N] A [-/N] CAAGCGAATAACATGTAGAAA	1147	62

TABLE 2

Haplotype Analysis of HF1 SNPs in AMD Cases and Controls

Prom	IVS1	162V	IVS6	Y402H	IVS10	Q672Q	D936E	OR est	P	Cases	Con
C	C	C	T	C	G	A	G	2.67	<0.000001	0.49	0.29
C	G	G	T	T	A	A	G	0.49	<0.000011	0.06	0.12
C	T	T	T	T	A	A	G	0.63	0.00035	0.12	0.18
G	C	C	T	C	C	A	A	2.68		0.48	0.26
G	C	G	T	T	A	A	A	0.50		0.12	0.12
G	T	T	T	T	A	A	A	0.62		0.12	0.18
C	C	C	T	C	C	A	A	2.65		0.49	0.26
C	G	G	T	T	A	A	A	0.50		0.06	0.12
C	T	T	T	T	A	A	A	0.6		0.12	0.19
G	C	C	T	C	C	A	A	2.56		0.49	0.27
G	C	G	T	T	A	A	A	0.49		0.06	0.12
G	T	T	T	T	A	A	A	0.6		0.12	0.19
C	C	C	T	C	C	A	A	2.33		0.51	0.31
C	G	G	T	T	A	A	A	0.51		0.06	0.12
C	T	T	T	T	A	A	A	0.56		0.13	0.21
C	C	C	T	C	C	A	G	2.54		0.51	0.29
C	G	G	T	T	A	A	G	0.48		0.07	0.14
T	T	T	T	T	A	A	G	0.58		0.12	0.19
C	C	C	T	C	C	A	A	2.55		0.51	0.29
C	G	G	T	T	A	A	A	0.48		0.07	0.14
T	T	T	T	T	A	A	A	0.58		0.12	0.19
C	C	C	T	C	C	A	A	2.48		0.51	0.30
C	G	G	T	T	A	A	A	0.47		0.07	0.14
T	T	T	T	T	A	A	A	0.56		0.12	0.20
C	C	C	T	C	C	A	A	2.47		0.51	0.30
C	G	G	T	T	A	A	A	0.48		0.07	0.14
T	T	T	T	T	A	A	A	0.55		0.12	0.20
C	C	C	T	C	C	A	A	2.12		0.76	0.60
C	G	G	T	T	A	A	A	0.51		0.08	0.14
T	T	T	T	T	A	A	A	0.53		0.13	0.21
C	C	C	T	C	C	A	G	2.49		0.52	0.30
C	G	G	T	T	A	A	G	0.45		0.29	0.47
T	T	T	T	T	A	A	G	0.58		0.12	0.19

TABLE 3

rs3753394	rs529825	rs3766404	rs203674	rs3753396	rs1065489	Haplotype	Freq. CTL	Freq. AMD
Promoter	Intron 1	Intron 6	Intron 10	Exon 13	Exon 18			
1	1	1	2	1	1	11211	0.28436	0.478059
1	2	1	1	1	1	12111	0.210856	0.131313
2	1	1	1	2	2	21122	0.149247	0.126697
1	1	2	1	1	1	11211	0.129893	0.061917
2	1	1	1	1	1	21111	0.094861	0.07125
2	1	1	2	1	1	21121	0.046438	0.06834
1	2	2	1	1	1	12111	0.026677	0.014859
2	1	2	1	1	1	21211	0.012731	0.01473
1	1	1	1	1	1	11111	0.012686	0.007305
1	2	1	2	1	1	12121	0.012684	0.004723
1	2	1	1	2	2	12122	0.009249	0.000945
1	1	1	1	2	2	11122	0.007487	0.008705
2	1	2	1	2	1	21212	0.002049	
1	2	2	2	1	1	12211	0.00078	0.000488
2	1	2	1	2	2	21212	0.000001	
2	1	1	2	1	2	21121		0.002175
2	1	1	1	1	2	21112		0.001869
2	2	1	1	1	1	22111		0.00169
2	1	2	2	1	1	21211		0.001061
1	2	1	2	2	2	12122		0.00096
1	1	2	1	1	2	11212		0.00095
2	1	1	1	2	1	21121		0.00095
2	1	2	2	1	2	21212		0.00069
2	2	1	2	1	1	22121		0.000322
1	2	2	1	1	2	12212		0.000001

TABLE 4

HFI SNP Association with Age-related Macular Degeneration

	Promoter rs3753394	IVS1 rs529825	Exon 2 162V	IVS2 insTT	IVS6 rs3766404	Exon 7/9 A307AY402H	Exon 10 A473A	IVS10 rs203674	Exon 13 Q672Q	Exon 18 D936E
Iowa										
# Controls	126	266	261	273	271	272	264	264	265	264
# Cases	329	547	546	549	546	549	542	545	545	536
X2	8.61	25.4	36.12	28.4	23.04	54.4	66.1	66.1	2.05	0.53
P	0.00334	4.66X10 ⁻⁰⁷	3.24X10 ⁻⁰⁷	9.87X10 ⁻⁰³	1.59X10 ⁻⁰⁶	1.64X10 ⁻¹³	1.60X10 ⁻¹¹	4.29X10 ⁻¹⁶	0.15	0.46
OR	0.70	1.92	1.95	2.042	2.105	2.25	2.10	2.44	1.24	1.12
95%CI	0.56-0.89	1.49-2.48	1.51-2.52	1.57-2.63	1.56-2.85	1.79-2.75	1.69-2.61	1.97-3.03	0.937-1.65	0.846-1.49
Columbia										
# Controls	126	266	261	273	271	272	264	264	265	264
# Cases	329	547	546	549	546	549	542	545	545	536
X2	8.61	25.4	36.12	28.4	23.04	54.4	66.1	66.1	2.05	0.53
P	0.00334	4.66X10 ⁻⁰⁷	3.24X10 ⁻⁰⁷	9.87X10 ⁻⁰³	1.59X10 ⁻⁰⁶	1.64X10 ⁻¹³	1.60X10 ⁻¹¹	4.29X10 ⁻¹⁶	0.15	0.46
OR	0.70	1.92	1.95	2.042	2.105	2.25	2.10	2.44	1.24	1.12
95%CI	0.56-0.89	1.49-2.48	1.51-2.52	1.57-2.63	1.56-2.85	1.79-2.75	1.69-2.61	1.97-3.03	0.937-1.65	0.846-1.49

The frequency of allele 1 and allele 2 from each SNP was compared between cases and controls and the Yates Chi squared (X2) and P values were calculated along with the Odds Ratio (OR) and 95% confidence interval (95% CI). The actual counts of each genotype are given in Tables 6A-6C.

TABLE 5

SSCP, DHPLC and Sequencing Primers

Exon	Region	Forward Primer (5'-3')	SEQ ID NO:	Reverse Primer (5'-3')	SEQ ID NO:
1	5'upstream-int1	GCAAAAGTTTCTGATAGGC	64	AATCTTACCTTCTGCTACAC	65
2	int1-int2	TTAGATAGACCTGTGACTG	66	TCAGGCATAATTGCTAC	67
3	int2-int3	ACTTGTTCCCCCACTC	68	CCTCTTTTCGTATGGACTAC	69
	int2-ex3	TTGTTCCCCCACTCCTAC	70	ACACATTTCTCCTGTAAAGG	71
	ex3-int3	CCCTGTGGACATCCTGG	72	AACCTCTTTTCGTATGGACTAC	73
4	int3-ex4	ATGCTGTTTCATTITCC	74	CCATCCATCTGTGTCAC	75
	ex4-int4	ATTACCGTGAATGTGAC	76	TTGTATGAGAAAAAAGAAC	77
5	int4-int5	TCCAATCTTATCCTGAGG	78	TCITACCCACACACTTTG	79
6	int5-ex6	GTCTGGTCCACAGTCC	80	GCATACAGCATCTCCTC	81
	ex6-int6	GCACTGAATCTGGATG	82	ATGAACCTTGAACACAG	83
7	int6-ex7	CGGATACTTATTTCTGC	84	CGTGATTTTCATCTCCAG	85
	ex7-int7	AGAACTGGAGATGAAATC	86	TGAATGGAACCTTACAGG	87
8	int7-ex8	GTGAAACCTTGTGATTATC	88	TCCCAGTAACTTCCTG	89
	ex8-int8	CTGTGATGAACATTTTGAG	90	TGCTCTCCTTTCTTCG	91
9	int8-int9	CATTGTTATGGTCTTAGG	92	ACATGCTAGGATTCAGAG	93
10	int9-ex9	CTTTTCTTATTCCTTCCC	94	TCACCATCTGCTGTTAC	95
	ex9-int10	TGTAACAGCAGATGGTG	96	CCCACAAAAAGACTAAAG	97
11	int10-ex11	GGGAAATACTCAGATTG	98	ATGGCATTTCATAGTCC	99
	ex11-int11	CCAGAACTAAAAATGACTTC	100	GGTAAATCAGACCAACC	101
	ex11-int11	ATAGTGTGTGGTTACAATG	102	GTTTATGTCAAATCAGGAG	103
12	int11-int12	CAAGAAAGAGAATGCGAAC	104	AGATTACAGGCAATGGG	105
13	int12-ex13	TTGATTGTTTAGGATGC	106	TTGAGGAGTTCAGGAGGTGG	107
	ex13-int13	CTGAACTCCTCAATGG	108	ATTACCAATACACACTGG	109
14	int13-int14	TTACATAGTGGAGGAGAG	110	TGGAAATGTTGAGGC	111
15	int14-int15	AGTTGGTTTGATTCCTATC	112	TTGAGCAGTTCACCTCTG	113
16	int15-ex16	TTATGCCACCTCCAC	114	ATACACTACTGACCAACAC	115
	ex16-int16	GTCTATGAGAATACAAGCC	116	GAATCTGAGGTGGAGG	117
17	int16-ex17	CCCTTTGATTTTCATTTC	118	AGAACTCCATTTTCCC	119
	ex17-int17	CACAACCACCTCAGATAG	120	GCCTAACCTTCACACTG	121
18	int17-ex18	GTCATAGTAGCTCCTGTATTG	122	ACGTAACTTCTCTCCATAC	123
	ex18-int18	CTTCCTTGTAATCTCCAC	124	CAATGCACCATACTTATGC	125
19	int18-ex19	TAAAGATTGCGGAAC	126	GGCTCCATCCATTTTG	127
	ex19-int19	TTACAAAATGGATGGAG	128	AAGTGCTGGGATTACAGGCG	129
20	int19-ex20	CTACTCAAAATGAACACTAGG	130	TTTAAACCCTGCTATACTCC	131
	ex20-int20	TAAATGGAAACTGGACG	132	ACCCTATTACTTGTGTTCTG	133
21	int20-int21	GTGTTTGCGTTTGCC	134	GAGATTTTCCAGCCAC	135
22	int21-ex22	TCTCACACATGCGAAC	136	ACCGTTAGTTTCCAGG	137
	ex22-3'downstream	GGTTTGGATAGTGTTTTGAG	138	ATGTTGTTCCGAATGTG	139

TABLE 6

Cohort 1	Promoter	Promoter	rs3753394	Intron 1	IVS1	rs523825	Exon 2	162V	rs800292	Intron 2	IVS2	InsTT
Sum												
freq allele 1							GG	CON	AMD	SS	CON	AMD
freq allele 2							GA	18	34	SL	38	73
AMD association							AA	6	4	LL	10	7
Chi square							G	68	228	S	126	390
Count Allele 1							A	0.78	0.91	L	0.77	0.89
Count Allele 2								0.22	0.09		0.23	0.11
Yates χ^2 /P value												
OR/95%CI												
Cohort 2												
Sum							Allele 1	X2	P		X2	P
freq allele 1							Allele 2	16.19	5.73703E-05		22.19	2.4667E-06
freq allele 2								2.79			2.38	
Count allele 1								106	414		194	693
Count allele 2								30	42		58	87
Yates χ^2 /P value								15	0.000107511		22.21	2.44398E-06
OR/95%CI								2.79	1.67-4.65		2.38	1.65-3.44
Cohort 2												
Sum												
freq allele 1												
freq allele 2												
Count allele 1												
Count allele 2												
Yates χ^2 /P value												
OR/95%CI												

TABLE 6 (continued 1)

Cohort 1	Intron 6	IVS6	rs376644	Exon 7/9	A307A/ Y402H	rs1061147/ rs1061170	Exon 7	A307A	rs1061147	Exon 9	Y402H	rs1061170
Sum				AA/CC	CON	AMD		CON	AMD	CC	CON	AMD
freq allele 1				AC/CT	16	146	AA	16	146	CT	16	146
freq allele 2				CC/TT	56	183	AC	56	183	TT	56	183
AMD association					59	75	CC	59	75		59	75
				A/C	131	404		131	404	C	131	404
				C/T	0.34	0.59	A	0.34	0.59	T	0.34	0.59
					0.66	0.41	C	0.66	0.41		0.66	0.41
Chi-square					X2	P		X2	P		X2	P
Count Allele 1					50.4	1.2593E-12		50.4	1.2593E-12		50.4	1.2593E-12
Count Allele 2					2.82			2.82			2.82	
Yates χ^2 /P value					88	475		88	475		88	475
OR/95%CI					174	333		174	333		174	333
					49.4	2.08746E-12		49.4	2.08746E-12		49.4	2.08746E-12
					2.82	2.11-3.78		2.82	2.11-3.78		2.82	2.11-3.78
Cohort 2												
Sum												
freq allele 1												
freq allele 2												
Count allele 1												
Count allele 2												
Yates χ^2 /P value												
OR/95%CI												

TABLE 6 (continued 2)

TABLE 7

Frequency of the At-risk Allele in Various Ethnic Groups with or without AMD

Risk Haplotype	Rapanui	Columbia Controls	Hispanic	Iowa Controls	African American	Columbia Cases	European American	Iowa Cases	MPGN II
N	52	272	24	131	49	549	56	404	20
	0.20	0.35	0.35	0.36	0.47	0.55	0.57	0.60	0.69

The frequency of the at-risk haplotype was estimated in samples from different populations from genotypes of the Y402H variant and/or the IVS10 locus. These include Rapanui natives over the age of 65 (AMD is extremely rare, and most likely absent, in this Easter Island population), controls (>65 years of age) from Columbia University, Hispanics general population, controls (>65 years of age) from the University of Iowa, African Americans general population, AMD cases from Columbia University, European Americans general population, AMD cases from the University of Iowa and individuals with MPGNII. N=number of individuals.

TABLE 8
Factor H Diplotypes

	I62V	IVS2-18	Y402H	D936E	IVS20
Risk	GG	SS	CC	GG	TT
Protective	AA	LL	TT	GG	CC
Protective	AA	LL	CT	GG	CC
	AA	LL	TT	GG	CC
	AA	LL	TT	GT	TT
	AA	LL	TT	GT	CC
	AA	LS	CT	GG	CC
	AA	SS	CC	GG	TT
	GA	LS	CT	GG	TT
	GA	LS	CT	GG	CT
	GA	LS	CT	GG	CC
	GA	LS	CT	GG	CC
	GA	LS	CT	GT	CT
	GA	LS	TT	GG	CT
	GA	LS	TT	GG	CC
	GA	LS	TT	GG	TT
	GA	LS	TT	GT	CC
	GA	LS	TT	TT	CC
Risk	GA	SS	CT	GG	CC
	GG	SS	CT	GT	CC
	GG	SS	TT	GT	TT
	GG	SS	TT	TT	CT
	GG	LL	TT	GG	TT
	GG	SS	CC	GG	TT
	GG	SS	CT	GG	CT
	GG	SS	CT	GG	CC
	GG	SS	CT	GG	CC
	GG	SS	CT	GT	TT
	GG	SS	CT	GT	CT
	GG	SS	CT	GT	CC
	GG	SS	CT	GT	CC
	GG	SS	CT	GT	CC
	GG	SS	CT	GT	CC
	GG	SS	CT	TT	CT
	GG	SS	TT	GG	TT
	GG	SS	TT	GG	CT
	GG	SS	TT	GT	TT
	GG	SS	TT	GT	CT
	GG	SS	TT	GT	GT
	GG	SS	TT	GT	CC
	GG	SS	TT	TT	CT
	GG	SS	TT	TT	CC
	GG	SS	CC	GG	TT
	GG	SS	CT	GG	TT
	GG	SS	CT	GT	CC

G, A, T, C refer to nucleotides at the indicated polymorphisms. S, L refer to the short and long (insertion of 2 T nucleotides) alleles of the intron 2 polymorphism.

TABLE 9

Primers Used to Amplify the Factor H Coding Sequence

Exon	Forward	SEQ ID NO:	Reverse	SEQ ID NO:
1	TGGGAAGTCAGTGAGAATTG	140	GCTAATGATGCTTTTCACAGGA	141
2	CCTGTGACTGTCTAGGCATTTT	142	TATGCCTGAATTATATCACTATTGCC	143
3	GCTTTGCTATGTTTAATTTTCCTT	144	AACATGATGGAAATAATTAAATCTGG	145
4	TGCATATGCTGTTCAATTTT	146	GTCTTACATTAAAATATCTTAAAGTCTC	147
5	TTTCCTCCAATCTTATCCTGAG	148	CGTTCATTCTAAGGAATATCAGCA	149
6	CCTGATGGAAACAACATTTCTG	150	AACAGGGCCAGAAAAGTTCA	151
7	TGTTCAATTTAATGCCATTTTG	152	AGTTTTCGAAGTTGCCGAAA	153
8	CCTAGAAACCCTAATGGAATGTG	154	TGTTCAAGCAAAGTGACCAAA	155
9	TGAGCAAATTTATGTTTCTCATTT	156	ATGTCACCTTGTTTACCAATGG	157
10	TGAATGCTTATGGTTATCCAGGT	158	AAAACCTGCAGGAACAAAGC	159
11	TCTTAGAATGGGAAATACTCAGATTG	160	TGGTTTTTCAGAAATTCATTTTCA	161
12	ATGTAAAATTAACCTTGCCAATGA	162	TTGCTGAAATAAGAATTAGAAGTTTG	163
13	TGAATAAAAGAAGAAAATCTTTCCA	164	ATCTAAAACACATACATCATGTTTCA	165
14	AAAACACATACATCATGTTTTCACAA	166	GATATGCCTCAACATTTCCAGTC	167
15	GTTGGTTTGATTCTATCATTTG	168	TTGGAAAAGTAATAGGTATGTGTGTC	169
16	CTATGAGAATACAAGCCAAAAGTTC	170	TCTCTGTGCTTCGTGTAAACAA	171
17	AACCCTTTGATTTTCATTCTTCA	172	TCAAAGTGAGGGGAATAATTGA	173
18	AATTTATGAGTTAGTGAAACCTGAAT	174	TCTTCATTCAAAGTGTAAGTGGTACC	175
19	ACAAAATGGCTAATATATTTCTCAAG	176	TAATGTGTGGGCCAGCC	177
20	CAAAATGAACACTAGGTGGAACC	178	ATTTGGGGGAGTATAGCAGG	179
21	CTGTGTTTGCGTTTGCCTTA	180	TTACGTGGCTGGAAAAATC	181
22	TTGAAAACCTGAAAGTCTATGAAGA	182	TCAATCATAAAGTGCACACCTTT	183

TABLE 10

Primers Used to Amplify the *CFHR5* Coding Sequence

Exon	Forward	SEQ ID NO:	Reverse	SEQ ID NO:
1	CAGTCCCATTTCTGATTGTTCCA	184	GCTGAGGATAATTTGAAGGGG	185
2	GTGATTCATCGATGTAGCTCTTT	186	AATGACCAGAGGAGCCTGGAA	187
3	TGATGTCAGTTTTCAAAGTTTCC	188	ACCACTCTCTCAGTTTGTCTAATTAT	189
4	CACATTAAATTTGTTTCTGCAATGA	190	AGAAGTGATGAAACAAGAATTTGA	191
5	CCATTTAAGCATTATTTATGGTTTC	192	AAACAGGACAGTTACTATTACTTTGCA	193
6	AAATATTTTCAGAGTAAGCACTCATT	194	TTTATCATTTTGATTGGGATTGT	195
7	TGCAGATATTTATTGACATAATTGTT	196	GTTGATCTTGTTGCTTCTTACAAGA	197
8	CCATTTTCTGAAACACTACCC	198	TCTGTTGCACTGTACCCCAA	199
9	AATTATTTGAATTTCCAGACACCTT	200	TTTTGGACTAATTTTCATAGAATAACCC	201
10	CTFAAATGCAATTTCACTATTCTATGA	202	TAGCCATTATGTAGCC	203

TABLE 11

CFH SNPs in 22 Patients Segregating with MPGNII
(Allele Frequencies (f1 and f2) and Number of Patients by Genotype are Shown)

	EX2	EX7	EX9	EX10	EX13	EX18	EX20	EX21
	rs300292	rs105147	rs1051170	rs2274700	rs3753396	rs105489	rs105489	rs105489
	GG 20 GA 2 AA 0	CC 3 CA 10 AA 9	CC 8 CT 10 TT 3	GG 16 GA 4 AA 0	AA 13 AG 9 GG 0	GG 13 GT 9 TT 0	TT 19 TC 3 CC 0	AA 21 AT 1 TT 0
f1	.95 (T)9	.64A	.36Y	.30G	.80A	.80G	.93T	.98A
f2	.05 (T)11	.36C	.64H	.10A	.20G	.20T	.07C	.02T
8	At-Risk Haplotype	A	C	G	A	G	T	A
1	MPGN2-1	AC	G,T	G,G	G,A	G,T	T,T	A,A
2	MPGN2-2	C,C	T,T	G,G	G,A	G,T	T,T	A,A
3	MPGN2-3	AC	C,T	G,G	G,A	G,T	T,T	A,A
4	MPGN2-4	AC	C,T	G,G	G,A	G,T	T,T	A,A
5	MPGN2-5	AC	C,T	G,G	G,A	G,T	T,T	A,A
6	MPGN2-6	AC	C,T	G,G	G,A	G,T	T,T	A,A
7	MPGN2-7	AA	C,C	G,G	G,A	G,T	T,T	A,A
8	MPGN2-8	AA	C,C	G,G	G,A	G,T	T,T	A,A
9	MPGN2-9	AA	C,C	G,G	G,A	G,T	T,T	A,A
10	MPGN2-10	AA	C,C	G,G	G,A	G,T	T,T	A,A
11	MPGN2-11	AA	C,C	G,G	G,A	G,T	T,T	A,A
12	MPGN2-12	AA	C,C	G,G	G,A	G,T	T,T	A,A
13	MPGN2-13	AA	C,C	G,G	G,A	G,T	T,T	A,A
14	MPGN2-14	AA	C,C	G,G	G,A	G,T	T,T	A,A
15	MPGN2-15	AA	C,C	G,G	G,A	G,T	T,T	A,A
16	MPGN2-16	AA	C,C	G,G	G,A	G,T	T,T	A,A
17	MPGN2-17	AA	C,C	G,G	G,A	G,T	T,T	A,A
18	MPGN2-18	AA	C,C	G,G	G,A	G,T	T,T	A,A
19	MPGN2-19	AA	C,C	G,G	G,A	G,T	T,T	A,A
20	MPGN2-20	AA	C,C	G,G	G,A	G,T	T,T	A,A
21	MPGN2-21	AA	C,C	G,G	G,A	G,T	T,T	A,A
22	MPGN2-22	AA	C,C	G,G	G,A	G,T	T,T	A,A
23	MPGN2-23	AA	C,C	G,G	G,A	G,T	T,T	A,A
24	MPGN2-24	AA	C,C	G,G	G,A	G,T	T,T	A,A
25	MPGN2-25	AA	C,C	G,G	G,A	G,T	T,T	A,A
26	MPGN2-26	AA	C,C	G,G	G,A	G,T	T,T	A,A
27	MPGN2-27	AA	C,C	G,G	G,A	G,T	T,T	A,A
28	MPGN2-28	AA	C,C	G,G	G,A	G,T	T,T	A,A
29	MPGN2-29	AA	C,C	G,G	G,A	G,T	T,T	A,A
30	MPGN2-30	AA	C,C	G,G	G,A	G,T	T,T	A,A

TABLE 12

Comparison of Factor H SNP Frequencies in 22 MPGNII Patients Versus Controls
(Allele Frequencies Given as f1 and f2)

SNP	f1 MPGNII	f2 MPGNII	f1 Controls	f2 Controls	P-value
Exon 2 I62V	42 (G)	2 (A)	202 (G)	60 (A)	0.0051
IVS2 -18insTT	42 (short)	2 (long)	194 (short)	68 (long)	0.0018
Exon 7 A307A	16 (C)	28 (A)	88 (A)	174 (C)	0.72
Exon 9 Y402H	28 (H)	16 (Y)	88 (H)	174 (Y)	0.00014
Exon 10 A473A	40 (G)	4 (A)	74 (G)	62 (A)	0.000013
Exon 13 Q672Q	35 (A)	9 (G)	217 (A)	41 (G)	0.45
Exon 18 D936E	35 (D)	9 (E)	115 (D)	19 (E)	0.32

TABLE 13

**Coding SNPs Associated with MPGNII and the Related Short Consensus Repeat (SCR) of
Factor H**

SNP	SCR	Function of SCR
Exon 2 I62V	1	Interaction with C3b
Exon 9 Y402H	7	Heparin binding Interaction with C reactive protein
Exon 10 A473A	8	Interaction with C reactive protein

TABLE 14

CFHR5 SNPs in 22 Patients Segregating with MPGNII
(Allele Frequencies (f1 and f2) and Number of Patients by Genotype are Shown)

	Promoter rs9427661	Promoter rs9427662	IVS1	75T>A rs3748557	Exon2 rs12097550	IVS2	58C>T rs12097550
1.	TT	TT	TT	16	CC	CC	16
2	TC	TC	TA	5	CT	CT	5
3	CC	CC	AA	1	TT	TT	1
4							.84C
5				.84T	.93P		.16T
6				.16A	.07S		
7							
8	At-Risk Haplotype	T	T	T	C	C	C
MPGN2-02	T,T	T,T	T,T	T,T	C,C	C,C	C,C
MPGN2-03	T,T	T,T	T,T	T,T	C,C	C,C	C,C
MPGN2-07	T,T	T,T	T,T	T,T	C,C	C,C	C,C
MPGN2-09	T,T	T,T	A,T	A,T	C,C	C,T	C,T
MPGN2-10	T,T	T,T	T,T	T,T	C,C	C,C	C,C
MPGN2-11	T,T	T,T	A,T	A,T	C,C	C,T	C,T
MPGN2-12	T,T	T,T	T,T	T,T	C,C	C,C	C,C
MPGN2-13	T,T	T,T	A,T	A,T	C,T	C,T	T,T
MPGN2-14	T,T	T,T	A,A	A,A	C,C	C,C	C,C
MPGN2-15	T,T	T,T	T,T	T,T	C,T	C,T	C,C
MPGN2-16	C,T	C,T	A,T	A,T	C,C	C,C	C,T
MPGN2-17	T,T	T,T	T,T	T,T	C,C	C,C	C,C
MPGN2-18	T,T	T,T	A,T	A,T	C,C	C,T	C,C
MPGN2-19	T,T	T,T	T,T	T,T	C,T	C,C	C,C
MPGN2-20	T,T	T,T	T,T	T,T	C,C	C,C	C,C
MPGN2-21	T,T	T,T	T,T	T,T	C,C	C,C	C,C
MPGN2-22	T,T	T,T	T,T	T,T	C,C	C,C	C,C
MPGN2-23	T,T	T,T	T,T	T,T	C,C	C,C	C,C
MPGN2-24	T,T	T,T	T,T	T,T	C,C	C,C	C,C
MPGN2-27-2	T,T	T,T	T,T	T,T	C,C	C,C	C,C
MPGN2-29	T,T	T,T	T,T	T,T	C,C	C,C	C,C
MPGN2-30	T,T	T,T	T,T	T,T	C,C	C,C	C,C

TABLE 15

Comparison of *CFHR5* SNP Frequencies in 22 MPGNII Patients Versus Controls
(Allele Frequencies Given as f1 and f2)

SNP	f1 MPGN II	f2 MPGN II	f1 Controls	f2 Controls	P-value
Promoter -249T>C	43 (T)	1 (C)	178 (G)	28(A)	0.033
Promoter -20T>C	43 (T)	1 (C)	178 (G)	28(A)	0.033
IVS1 +75T>A	37(T)	7 (A)	161 (A)	41 (C)	0.38
Exon 2 P46S	41 (P)	3(S)	205 (P)	1 (S)	0.00023
IVS2 +58C>T	37 (C)	7 (T)	158 (C)	28 (T)	0.28

TABLE 16A

SNP Name	Location	Probes			SEQ ID NO:	Variant Allele	SEQ ID NO:
		Reference Allele	Reference Allele	Reference Allele			
	Promoter 1	5'-TCTGGGATGTAATAATG-3'	5'-TCTGGGATGTAATAATG-3'	5'-TCTGGGATGTAATAATG-3'	204	5'-TCTGGGATGTAATAATG-3'	205
		5'-GAACATTATTACATCCC-3'	5'-GAACATTATTACATCCC-3'	5'-GAACATTATTACATCCC-3'	206	5'-GAACATTATTACATCCC-3'	207
rs3753394	Promoter 4	5'-CAGAGGATATCACCAGC-3'	5'-CAGAGGATATCACCAGC-3'	5'-CAGAGGATATCACCAGC-3'	208	5'-CAGAGGATATCACCAGC-3'	209
		5'-AGCAGCTGGTGATATCC-3'	5'-AGCAGCTGGTGATATCC-3'	5'-AGCAGCTGGTGATATCC-3'	210	5'-AGCAGCTGGTGATATCC-3'	211
rs29825	Intron 1	5'-TACACAGTACGATAGAC-3'	5'-TACACAGTACGATAGAC-3'	5'-TACACAGTACGATAGAC-3'	212	5'-TACACAGTACGATAGAC-3'	213
		5'-TAAAGTCTATCGTACTGT-3'	5'-TAAAGTCTATCGTACTGT-3'	5'-TAAAGTCTATCGTACTGT-3'	214	5'-TAAAGTCTATCGTACTGT-3'	215
rs800292	Exon 2	5'-TCTTGGGAAATGTAATAA-3'	5'-TCTTGGGAAATGTAATAA-3'	5'-TCTTGGGAAATGTAATAA-3'	216	5'-TCTTGGGAAATGTAATAA-3'	217
		5'-ACCATTTATACATTTC-3'	5'-ACCATTTATACATTTC-3'	5'-ACCATTTATATATTC-3'	218	5'-ACCATTTATATATTC-3'	219
	Intron 2	5'-TTTTTTTTTTCGTTTAG-3'	5'-TTTTTTTTTTCGTTTAG-3'	5'-TTTTTTTTTTCGTTT-3'	220	5'-TTTTTTTTTTCGTTT-3'	221
		5'-CTTTCIAAAACGAAAAA-3'	5'-CTTTCIAAAACGAAAAA-3'	5'-TCTAAAAACGAAAAA-3'	222	5'-TCTAAAAACGAAAAA-3'	223
rs3766404	Intron 6	5'-TTTAGGACTCATTTGAA-3'	5'-TTTAGGACTCATTTGAA-3'	5'-TTTAGGACTTATTGAA-3'	224	5'-TTTAGGACTTATTGAA-3'	225
		5'-TAACTTCAAATGAGTCC-3'	5'-TAACTTCAAATGAGTCC-3'	5'-TAACTTCAAATAAGTCC-3'	226	5'-TAACTTCAAATAAGTCC-3'	227
rs1061147	Exon 7	5'-GAAATACAGCAAAATGC-3'	5'-GAAATACAGCAAAATGC-3'	5'-GAAATACAGCCAAATGC-3'	228	5'-GAAATACAGCCAAATGC-3'	229
		5'-ACTTGTGTCATTTTGTG-3'	5'-ACTTGTGTCATTTTGTG-3'	5'-ACTTGTGTCATTTGGCTG-3'	230	5'-ACTTGTGTCATTTGGCTG-3'	231
rs1061170	Exon 9	5'-TAATCAAAATATGGAA-3'	5'-TAATCAAAATATGGAA-3'	5'-TAATCAAAATCATGGAA-3'	232	5'-TAATCAAAATCATGGAA-3'	233
		5'-TTTCTTCCATAATTTTG-3'	5'-TTTCTTCCATAATTTTG-3'	5'-TTTCTTCCATGATTTTG-3'	234	5'-TTTCTTCCATGATTTTG-3'	235
rs2274700	Exon 10	5'-AAGAAAAAGCGAAATAT-3'	5'-AAGAAAAAGCGAAATAT-3'	5'-AAGAAAAAGCAAAATAT-3'	236	5'-AAGAAAAAGCAAAATAT-3'	237
		5'-TTGATATTTCGCTTTT-3'	5'-TTGATATTTCGCTTTT-3'	5'-TTGATATTTCGCTTTT-3'	238	5'-TTGATATTTCGCTTTT-3'	239
	Exon 10A	5'-GGATCAAAAGATGACAA-3'	5'-GGATCAAAAGATGACAA-3'	5'-GGATCAAAAGATGACAA-3'	240	5'-GGATCAAAAGATGACAA-3'	241
		5'-GCCCTTGTCA[-]TCTTTG-3'	5'-GCCCTTGTCA[-]TCTTTG-3'	5'-GCCCTTGTCA[N]TCTTTG-3'	242	5'-GCCCTTGTCA[N]TCTTTG-3'	243
rs203674	Intron 10	5'-TTTATTAGTAGATCTAA-3'	5'-TTTATTAGTAGATCTAA-3'	5'-TTTATTAGTATATCTAA-3'	244	5'-TTTATTAGTATATCTAA-3'	245
		5'-TTGATTAGATCTACTAA-3'	5'-TTGATTAGATCTACTAA-3'	5'-TTGATTAGATATACTAA-3'	246	5'-TTGATTAGATATACTAA-3'	247
rs3753396	Exon 13	5'-ATAAAATTCAATGTGT-3'	5'-ATAAAATTCAATGTGT-3'	5'-ATAAAATTCAATGTGT-3'	248	5'-ATAAAATTCAATGTGT-3'	249
		5'-CCATCAACACATGGAAT-3'	5'-CCATCAACACATGGAAT-3'	5'-CCATCAACACACTGAAT-3'	250	5'-CCATCAACACACTGAAT-3'	251
rs375046	Intron 15	5'-TTTATATACATTAAAT-3'	5'-TTTATATACATTAAAT-3'	5'-TTTATATATAAAATTAAT-3'	252	5'-TTTATATATAAAATTAAT-3'	253
		5'-TATAAATTAATGTTATAA-3'	5'-TATAAATTAATGTTATAA-3'	5'-TATAAATTAATTTTATAA-3'	254	5'-TATAAATTAATTTTATAA-3'	255

SNP Name	Location	Probes			
		Reference Allele	SEQ ID NO:	Variant Allele	SEQ ID NO:
rs1065489	Exon 18	5'-CTCCACCTGAGATTCT-3'	256	5'-CTCCACCTGATATTCT-3'	257
		5'-CATGAGAGAAATCTCAGGT-3'	258	5'-CATGAGAGAAATATCAGGT-3'	259
	Exon 22	5'-TCTTTTCATCAGTTC-3'	260	5'-TCTTTTCATCATGTTC-3'	261
		5'-GTGTGAGAACGTTGATGA-3'	262	5'-GTGTGAGAACATGATGA-3'	263

TABLE 16A (continued)

TABLE 16B

SNP Name	Location	Primers			SEQ ID NO:
		Reference Allele	SEQ ID NO:	Variant Allele	
rs3753394	Promoter 1	5'-TTTCTGGGATGTAATA-3' (forward)	264	5'-TTTCTGGGATGTAATG-3' (forward)	265
	Promoter 4	5'-CAAAAACACTGAACATT-3' (reverse)	266	5'-CAAAAACACTGAACATC-3' (reverse)	267
		5'-AAATCCAGAGGATATC-3' (forward)	268	5'-AAATCCAGAGGATA TT-3' (forward)	269
rs529825	Intron 1	5'-AAATCAGCAGCTGGTG-3' (reverse)	270	5'-AAATCAGCAGCTGGTA-3' (reverse)	271
		5'-AAGTTTACACAGTACG-3' (forward)	272	5'-AAGTTTACACAGTACA-3' (forward)	273
	rs800292	Exon 2	5'-AATGGGTAAAGTCTATC-3' (reverse)	274	5'-AATGGGTAAAGTCTATT-3' (reverse)
5'-AGATCTCTTGGAAATG-3' (forward)			276	5'-AGATCTCTTGGAAAATA-3' (forward)	277
Intron 2		5'-TGCATACCAATTATTAC-3' (reverse)	278	5'-TGCATACCAATTATTAT-3' (reverse)	279
	5'-TCATAAAGCTTTTTT-3' (forward)	280	5'-ATAACTTTTTTTTTT-3' (forward)	281	
	5'-GGGCCCTTCTTAAACG-3' (reverse)	282	5'-GCCCTTCTTAAAAACGAA-3' (reverse)	283	
rs3766404	Intron 6	5'-AATACATTTAGGACTC-3' (forward)	284	5'-AATACATTTAGGACTT-3' (forward)	285
		5'-ACACTAACTTCAAAATG-3' (reverse)	286	5'-ACACTAACTTCAAAATA-3' (reverse)	287
	rs1061147	Exon 7	5'-CCGGGGAATACAGCA-3' (forward)	288	5'-CCGGGGGAAATACAGCC-3' (forward)
5'-AGTACTTGTGCAATTT-3' (reverse)			290	5'-AGTACTTGTGCATTG-3' (reverse)	291
rs1061170		Exon 9	5'-GGATATAATCAAAATT-3' (forward)	292	5'-GGATATAATCAAAATC-3' (forward)
	5'-CAAACTTCTTCCATA-3' (reverse)		294	5'-CAAAACTTCTTCCATG-3' (reverse)	295
	rs2274700	Exon 10	5'-CTTAAAGAAAAAGCG-3' (forward)	296	5'-CTTAAAGAAAAAGCA-3' (forward)
5'-TTTGCATTGATATTTC-3' (reverse)			298	5'-TTTGCATTGATATTTT-3' (reverse)	299
Exon 10A		5'-TGAGTGGATCAAGA[-]-3' (forward)	300	5'-TGAGTGGATCAAGA[IN]-3' (forward)	301
	5'-CATTTGGCCCTTGCA[-]-3' (reverse)	302	5'-CATTTGGCCCTTGCA[IN]-3' (reverse)	303	
	5'-ACCTATTATTAGTAG-3' (forward)	304	5'-ACCTATTATTAGTAT-3' (forward)	305	
rs203674	Intron 10	5'-CITTATTGATTAGATC-3' (reverse)	306	5'-CITTATTGATTAGATA-3' (reverse)	307
		5'-ACCTAATAAAATTCAA-3' (forward)	308	5'-ACCTAATAAAATTACAG-3' (forward)	309
	rs3753396	Exon 13	5'-CTCTCCATCAACACAT-3' (reverse)	310	5'-CTCTCCATCAACACAC-3' (reverse)
5'-TATTTTTTATTATAAC-3' (forward)			312	5'-TATTTTTTATTATAAA-3' (forward)	313
rs375046		Intron 15	5'-AAAAATATAATTAATG-3' (reverse)	314	5'-AAAAATATAATTAATT-3' (reverse)

SNP Name	Location	Primers			
		Reference Allele	SEQ ID NO:	Variant Allele	SEQ ID NO:
rs1065489	Exon 18	5'-TAAATCTCCACCTGAG-3' (forward)	316	5'-TAAATCTCCACCTGAT-3' (forward)	317
		5'-AACACCATGAGAAATC-3' (reverse)	318	5'-AACACCATGAGAAATA-3' (reverse)	319
		5'-TATCGTCTTTCATCAC-3' (forward)	320	5'-TATCGTCTTTCATCAT-3' (forward)	321
	Exon 22	5'-GCAATGTGTGAGAACG-3' (reverse)	322	5'-GCAATGTGTGAGAACG-3' (reverse)	323

TABLE 16B (continued)

XV. REFERENCES

[0385] Full citations are provided below for references cited by author and date above:

[0386] Abecasis et al. "Age-related macular degeneration: a high-resolution genome scan for susceptibility loci in a population enriched for late-stage disease." *American Journal of Human Genetics* 74, 482-94 (2004).

[0387] Akiyama et al. "Inflammation and Alzheimer's disease." *Neurobiol. Aging* 2000; 21:383-421.

[0388] Allikmets et al. "Mutation of the Stargardt disease gene (ABCR) in age-related macular degeneration." *Science* 1997; 277:1805-1807.

[0389] Allikmets. "Further evidence for an association of ABCR alleles with age-related macular degeneration. The International ABCR Screening Consortium." *Am J Hum Genet* 67, 487-91 (2000).

[0390] Ambati et al. "Age-related macular degeneration: etiology, pathogenesis, and therapeutic strategies." *Surv Ophthalmol* 2003; 48(3):257-293.

[0391] Anderson et al. "A role for local inflammation in the formation of drusen in the aging eye." *Am J Ophthalmol* 2002, 134:411-431.

[0392] Anderson et al. "Characterization of β -amyloid assemblies in drusen: the deposits associated with aging and age-related macular degeneration." *Exp. Eye Res.* 2004; 78:243-256.

[0393] Angaku-. "Complement regulatory proteins in glomerular diseases." *Kidney Int* 1998; 54:1419-1428.

[0394] Appel et al. "Membranoproliferative glomerulonephritis type II (Dense Deposit Disease): an update." *J Am Soc Nephrol* 2005; 16:1392-1403.

[0395] Ault et al. "Human factor H deficiency. Mutations in framework cysteine residues and block in H protein secretion and intracellular catabolism." *J Biol Chem* 1997; 272:25168-25175.

[0396] Bao et al. "Decay-accelerating factor expression in the rat kidney is restricted to the apical surface of podocytes." *Kidney Int* 2002;62:2010-2021.

[0397] Barbiano di Belgiojoso et al. "The prognostic value of some clinical and histological parameters in membranoproliferative glomerulonephritis." *Nephron* 1977;19:250-258.

[0398] Barrett et al. "Haploview: analysis and visualization of LD and haplotype maps." *Bioinformatics* 2004;21:263-5.

- [0399] Bennett et al. "Mesangiocapillary glomerulonephritis type 2 (dense deposit disease): Clinical features of progressive disease." *Am J Kidney Dis* 1989; 13:469-476.
- [0400] Bird et al. "An international classification and grading system for age-related maculopathy and age-related macular degeneration. The International ARM Epidemiological Study Group." *Surv Ophthalmol* 1995, 39: 367-374.
- [0401] Bush RA, Lei B, Tao W, Raz D, Chan CC, Cox TA, Santos-Muffley M, Sieving PA. 2004. Encapsulated cell-based intraocular delivery of ciliary neurotrophic factor in normal rabbit: dose-dependent effects on ERG and retinal histology. *Invest Ophthalmol Vis Sci*. 45:2420-30.
- [0402] Cade JR, DeQuesada AM, Shires DL, Levin DM, Hackett RL, Spooner GR, Schlein EM, Pickering MJ, Holcomb A. 1971. The Effect of Long Term High Dose Heparin Treatment on the Course of Chronic Proliferative Glomerulonephritis. *Nephron*. 8:67-80.
- [0403] Cameron et al. "Idiopathic mesangiocapillary glomerulonephritis. Comparison of types I and II in children and adults and long-term prognosis." *Am J Med* 1983;74:175-192.
- [0404] Capecchi. *Science* 1989; 244:1288-1292.
- [0405] Caprioli et al. "Complement factor H mutations and gene polymorphisms in haemolytic uraemic syndrome: the C-257T, the A2089G and the G2881T polymorphisms are strongly associated with the disease." *Hum Mol Genet* 12, 3385-95 (2003).
- [0406] Chong et al. "Decreased thickness and integrity of the macular elastic layer of Bruch's membrane correspond to the distribution of lesions associated with age-related macular degeneration" *Am J Pathol* 166, 241-51 (2005).
- [0407] Colville et al. "Visual impairment caused by retinal abnormalities in mesangiocapillary (membranoproliferative) glomerulonephritis type II ("dense deposit disease")." *Am J Kidney Dis* 2003; 42:E2-5.
- [0408] Compton. *Nature* 1991; 350:91-91.
- [0409] Cousins et al. "Monocyte activation in patients with age-related macular degeneration: a biomarker of risk for choroidal neovascularization?" *Arch Ophthalmol* 122, 1013-8 (2004).
- [0410] Crabb et al. "Drusen proteome analysis: An approach to the etiology of age-related macular degeneration." *Proc. Natl. Acad. Sci. USA*. 2002; 99:14682-14687.
- [0411] de Jong. "Risk profiles for ageing macular disease." *Ophthalmologia* 2004; 218 Suppl 1:5-16.

- [0412] Diamond JR, Karnovsky MJ. 1986. Nonanticoagulant Protective Effect of Heparin in Chronic Aminonucleoside Nephrosis. *Renal Physiol. Basel* 9:366-374.
- [0413] Dragon-Durey et al. "Heterozygous and homozygous factor H deficiencies associated with hemolytic uremic syndrome or membranoproliferative glomerulonephritis: report and genetic analysis of 16 cases." *J Am Soc Nephrol* 2004; 15:787-795.
- [0414] Droz et al. "Évolution a long terme des glomérulonéphrites membranoproliférative de l'adulte: remissionspontanée durable chez 13 malades avec étude de biopsies rénales itératives dans 5 cas." *Nephrologie* 1982;3:6-11.
- [0415] Duvall-Young et al. "Fundus changes in (type II) mesangiocapillary glomerulonephritis stimulating drusen: a histopathological report." *Br J Ophthalmol* 1989a; 73(4):297-302.
- [0416] Duvall-Young et al. "Fundus changes in mesangiocapillary glomerulonephritis type II: clinical and fluorescein angiographic findings." *Br J Ophthalmol* 1989b; 73(11):900-906.
- [0417] Edwards et al. "Complement factor H Polymorphism and age-related macular degeneration." *Science* 2005;308:421-424.
- [0418] Esparza-Gordillo J, Goicoechea de Jorge E, Buil A, Carreras Berges L, Lopez-Trascasa M, Sanchez-Corral P, Rodriguez de Cordoba S. 2005. Predisposition to atypical hemolytic uremic syndrome involves the concurrence of different susceptibility alleles in the regulators of complement activation gene cluster in 1q32. *Human Mol. Genetics* 14:703-712.
- [0419] Espinosa-Heidman et al. "Macrophage depletion diminishes lesion size and severity in experimental choroidal neovascularization." *Invest. Ophthalmol. Vis. Sci.* 2003; 44:3586-3592.
- [0420] Estaller et al. "Cloning of the 1.4-kb mRNA species of human complement factor H reveals a novel member of the short consensus repeat family related to the carboxy terminal of the classical 150-kDa molecule." *J Immunol* 1991; 146(9):3190-3196.
- [0421] Floege J, Eng E, Young BA, Couser WG, Johnson RJ. 1993. Heparin suppresses mesangial cell proliferation and matrix expansion in experimental mesangioproliferative glomerulonephritis. *Kidney International* 43:369-380.
- [0422] Frueh et al. *Clin Chem Lab Med* 2003; 41(4):452-461.
- [0423] Gibbs. *Nucl Acids Res* 1989; 17:2427-2448.
- [0424] Girardi G, Redecha P, Salmon JE. 2004. Heparin prevents antiphospholipid antibody-induced fetal loss by inhibiting complement activation. *Nature Medicine* 10:1222-1226.

- [0425] Girardi G. 2005. Heparin treatment in pregnancy loss: Potential therapeutic benefits beyond anticoagulation. *J. Reproduc. Immunol.* 66:45-51.
- [0426] Gold et al. "Estrogen receptor genotypes and haplotypes associated with breast cancer risk." *Cancer Res* 2004; 64:8891-8900.
- [0427] Habib et al. "Dense deposit disease. A variant of membranoproliferative glomerulonephritis." *Kidney Int* 1975; 7:204-15.
- [0428] Habib et al. "Glomerular lesions in the transplanted kidney in children." *Am J Kidney Diseases* 1987;10:198-207.
- [0429] Hageman et al. "An integrated hypothesis that considers drusen as biomarkers of immune-mediated processes at the RPE-Bruch's membrane interface in aging and age-related macular degeneration." *Prog Retin Eye Res* 2001; 20: 705-732.
- [0430] Hageman et al. "Common haplotype in the complement regulatory gene, factor H (*HFI/CFH*), predisposes individuals to age-related macular degeneration." *Proc Nat Acad Sci* 2005; 102: 7227-32.
- [0431] Hageman et al. "Vitronectin is a constituent of ocular drusen and the vitronectin gene is expressed in human retinal pigmented epithelial cells." *FASEB Journal* 1999; 13:477-484.
- [0432] Haines et al. "Complement factor H variant increases the risk of age-related macular degeneration." *Science* 2005;308:419-421.
- [0433] Hayashi et al. "Evaluation of the ARMD1 locus on 1q25-31 in patients with age-related maculopathy: genetic variation in laminin genes and in exon 104 of HEMICENTIN-1." *Ophthalmic Genetics* 25, 111-9 (2004).
- [0434] Houdebine, 2000, Transgenic animal bioreactors, *Transgenic Res.* 9:305-20
- [0435] Holers. "The complement system as a therapeutic target in autoimmunity." *Clin Immunol* 2003; 107:140.
- [0436] Holz, et al. "Pathogenesis of lesions in late age-related macular disease." *Am J Ophthalmol* 2004;137:504-510.
- [0437] Huang et al. "Peripheral drusen in membranoproliferative glomerulonephritis." *Retina* 2003; 23(3):429-431.
- [0438] Iyengar et al. "Model Free Linkage Analysis in Extended Families Confirms a Susceptibility Locus for Age Related Macular Degeneration (ARMD) on 1q31 [ARVO Abstract]." *Invest Ophthalmol Vis Sci* 2003; 44:2113.

- [0439] Jansen et al. "In situ complement activation in porcine membranoproliferative glomerulonephritis type II." *Kidney Int* 1998; 53(2):331-349.
- [0440] Johnson et al. "A potential role for immune complex pathogenesis in drusen formation." *Exp Eye Res* 2000; 70:441-449.
- [0441] Johnson et al. "Complement activation and inflammatory processes in drusen formation and age-related macular degeneration." *Exp. Eye Res.* 2001; 73:887-896.
- [0442] Johnson et al. "The Alzheimer's A beta -peptide is deposited at sites of complement activation in pathologic deposits associated with aging and age-related macular degeneration." *Proc Natl Acad Sci USA* 99, 11830-5 (2002).
- [0443] Kinoshita. "Biology of complement: the overture." *Immunol.Today* 1991; 12:291.
- [0444] Klaver et al. "Genetic risk of age-related maculopathy. Population-based familial aggregation study." *Arch Ophthalmol* 1998a; 116:1646-1651.
- [0445] Klaver et al. Genetic association of apolipoprotein E with age-related macular degeneration. *Am J Hum Genet* 1998b; 63:200-206.
- [0446] Klein et al. "Age-related macular degeneration. Clinical features in a large family and linkage to chromosome 1q." *Arch Ophthalmol* 1998; 116:1082-1088.
- [0447] Klein et al. "Complement factor H polymorphism in age-related macular degeneration." *Science* 2005; 308:385-389.
- [0448] Klein et al. "Genetics of age-related macular degeneration." *Ophthalmol Clin North Am* 2003; 16(4):575-582.
- [0449] Klein et al. "Prevalence of age-related maculopathy." *Ophthalmol* 1992; 99(6):933-943.
- [0450] Klein et al. "The epidemiology of age-related macular degeneration." *Am J Ophthalmol* 2004; 137(3):504-510.
- [0451] Leys et al. "Subretinal neovascular membranes associated with chronic membranoproliferative glomerulonephritis type II." *Graefe's Arch Clin Exper Ophthalmol* 1990; 228:499-504.
- [0452] Lillico et al., 2005, Transgenic chickens as bioreactors for protein-based drugs. *Drug Discov Today*. 10:191-6
- [0453] Liszewski et al. "The role of complement in autoimmunity." *Immunol Ser* 1991; 54:13.

- [0454] Manuelian et al. "Mutations in factor H reduce binding affinity to C3b and heparin and surface attachment to endothelial cells in hemolytic uremic syndrome." *J Clin Invest* 2003;111:1181-1190.
- [0455] McAvoy et al. "Retinal changes associated with type 2 glomerulonephritis." *Eye* 2005 19:985-9
- [0456] McEnery. "Membranoproliferative glomerulonephritis: The Cincinnati experience cumulative renal survival from 1957 to 1989." *J Pediatr* 1990;116:S109-S114.
- [0457] McRae et al. "Human factor H-related protein 5 (FHR-5). A new complement-associated protein." *J Biol Chem* 2001; 276 (9):6747-6754.
- [0458] Meri et al. "Regulation of alternative pathway complement activation by glycosaminoglycans: specificity of the polyanion binding site on factor H." *Biochem Biophys Res Commun* 1994;198:52-59.
- [0459] Miller et al. "The association of prior cytomegalovirus infection with neovascular age-related macular degeneration." *Am J Ophthalmol* 138, 323-8 (2004).
- [0460] Morgan et al. "Complement deficiency and disease." *Immunol Today* 1991; 12:301.
- [0461] Morgan. "Regulation of the complement membrane attack pathway." *Crit Rev Immunol* 19, 173-98 (1999).
- [0462] Mullins et al. "Characterization of drusen-associated glycoconjugates." *Ophthalmology* 104, 288-94 (1997).
- [0463] Mullins et al. "Drusen associated with aging and age-related macular degeneration contain molecular constituents common to extracellular deposits associated with atherosclerosis, elastosis, amyloidosis and dense deposit disease." *FASEB J.* 2000; 14:835-846.
- [0464] Mullins et al. "Structure and composition of drusen associated with glomerulonephritis: Implications for the role of complement activation in drusen biogenesis." *Eye* 2001; 15:390-395.
- [0465] Murphy et al. "Factor H-related protein-5: a novel component of human glomerular immune deposits." *Am J Kid Dis* 2002;39:24-27.
- [0466] Neary et al. "Linkage of a gene causing familial membranoproliferative glomerulonephritis type III to chromosome 1." *J Am Soc Nephrol.* 2002; 13(8):2052-2057.
- [0467] Neri et al. *Adv. Nucl Acid Prot Analysis* 2000; 3826:117-125.
- [0468] Niculescu et al. "Complement activation and atherosclerosis." *Mol. Immunol.* 1999; 36:949-955.

- [0469] Nielsen et al. *Science* 1991; 254:1497-1500.
- [0470] O'Brien et al. "Electrophysiology of type II mesangiocapillary glomerulonephritis with associated fundus abnormalities." *Br J Ophthalmol* 1993; 77:778-80.
- [0471] Orita et al. *Proc Natl Acad Sci* 1989; 86:2766-2770.
- [0472] Orth et al. The nephrotic syndrome. *New Engl J Med* 1998; 338:1202-1211.
- [0473] Pascual et al. "Identification of membrane-bound CR1 (CD35) in human urine: evidence for its release by glomerular podocytes." *J Exp Med* 1994;79:889-899.
- [0474] Penfold et al. "Immunological and aetiological aspects of macular degeneration." *Progress in Retinal and Eye Research* 2001; 20:385-414.
- [0475] Perez-Caballero D, Gonzalez-Rubio C, Gallardo ME, Vera M, Lopez-Trascasa M, Rodriguez de Cordoba S, Sanchez-Corral P. 2001. Clustering of Missense Mutations in the C-Terminal Region of Factor H in Atypical Hemolytic Uremic Syndrome. *Am. J. Hum. Genet.* 68:478-484.
- [0476] Piatek et al. *Nat Biotechnol* 1998; 16:359-363.
- [0477] Pickering et al. "Uncontrolled C3 activation causes membranoproliferative glomerulonephritis in mice deficient in complement factor H." *Nat Genet* 2002; 31:424-428.
- [0478] Prasad et al. "Pendred syndrome and DFNB4 - Mutation screening of *SLC26A4* by denaturing high-performance liquid chromatography and the identification of seven novel mutations." *Am J Med Genet* 2004;124A:1-9.
- [0479] Raines et al. "Fundus changes in mesangiocapillary glomerulonephritis type II: vitreous fluorophotometry." *Br J Ophthalmol* 1989; 73:907-910.
- [0480] Richards A, Buddles MR, Donne RL, Kaplan BS, Kirk E, Venning MC, Tielemans CL, Goodship JA, Goodship THJ. 2001. Factor H Mutations in Hemolytic Uremic Syndrome Cluster in Exons 18-20, a Domain Important for Host Cell Recognition. *Am. J. Hum. Genet.* 68:485-490.
- [0481] Ripoché et al. "The complete amino acid sequence of human complement factor H." *Biochem J* 1988, 249:593-602.
- [0482] Rodriguez de Cordoba et al. "The human complement factor H: functional roles, genetic variations and disease associations." *Mol Immunol* 41, 355-67 (2004).

- [0483] Rops Angelique L.W.M.M., Van Der Vlag J, Lensen Joost F.M., Wijnhoven Tessa J.M., Van Den Heuvel Lambert P.W.J., van Kuppevelt TH, Berden Jo H.M. 2004. Heparan sulfate proteoglycans in glomerular inflammation. *Kidney International* 65:768-785.
- [0484] Russell et al. "Location, substructure and composition of basal laminar drusen compared with drusen associated with aging and age-related macular degeneration." *Am. J. Ophthalmol.* 2000; 129:205-214.
- [0485] Saiki et al. *Nature* 1986; 324:163-166.
- [0486] Sanchez-Corral P, Perez-Caballero D, Huarte O, Simckes AM, Goicoechea E, Lopez-Trascasa M, Rodriguez de Cordoba S. 2002. Structural and Functional Characterization of Factor H Mutations Associated with Atypical Hemolytic Uremic Syndrome. *Am. J. Genet.* 71:1285-1295.
- [0487] Saunders RE, Goodship THJ, Zipfel PF, Perkins SJ. An Interactive Web Database of Factor H-Associated Hemolytic Uremic Syndrome Mutations: Insights Into the Structural Consequences of Disease-Associated Mutations. *Human Mutation* 2006. 27:21-30.
- [0488] Schultz et al. "Analysis of the ARMD1 locus: evidence that a mutation in HEMICENTIN-1 is associated with age-related macular degeneration in a large family." *Hum Mol Genet* 2003; 12(24):3315-3323.
- [0489] Schwertz et al. "Complement analysis in children with idiopathic membranoproliferative glomerulonephritis: A long-term follow-up." *Pediatr Allergy Immunol* 2001; 12:166-172.
- [0490] Seddon et al. "Association between C-reactive protein and age-related macular degeneration." *Jama* 291, 704-10 (2004).
- [0491] Seddon et al. "The epidemiology of age-related macular degeneration." *Ophthalmol Clin* 2004; 44:17-39.
- [0492] Sharma et al. "Biologically active recombinant human complement factor H: synthesis and secretion by the baculovirus system." *Gene* 143:301-302.
- [0493] Sharma et al. "Identification of three physically and functionally distinct binding sites in human complement factor H by deletion mutagenesis." *Proc Natl Acad Sci USA* 1996, 93:10996-11001.
- [0494] Shen et al. "Ying and Yang: complement activation and regulation of Alzheimer's disease." *Prog Neurobiol* 2003; 70(6):463-472.

- [0495] Sieving, P.A., R.C. Caruso, W. Tao, D.J.S. Thompson, K.R. Fullmer, H. Rodriguez Coleman and R.A. Bush. 2005. Phase I study of ciliary neurotrophic factor (CNF) delivered by intravitreal implant of encapsulated cell technology (ECT) device in patients with retinitis pigmentosa.
- [0496] Skerka et al. "The human factor H-related protein 4 (FHR-4). A novel short consensus repeat-containing protein is associated with human triglyceride-rich lipoproteins." *J Biol Chem* 1997; 272(9):5627-5634.
- [0497] Song Y, Zhao L, Tao W, Laties AM, Luo Z, and Wen R. Photoreceptor protection by cardiotrophin-1 in transgenic rats with the rhodopsin mutation s334ter. *IOVS*, 44(9):4069-75. 2003.
- [0498] Souied et al. "The epsilon4 allele of the apolipoprotein E gene as a potential protective factor for exudative age-related macular degeneration." *Am J Ophthalmol*. 1998; 125:353-359.
- [0499] Strausberg et al. "Generation and initial analysis of more than 15,000 full-length human and mouse cDNA sequences." *Proc Natl Acad Sci USA* 2002; 99(26):16899-16903.
- [0500] Striker GE. 1999. Therapeutic uses of heparinoids in renal disease patients. *Nephrol. Dial. Transplant*. 14:540-543.
- [0501] Swainson et al. "Mesangiocapillary glomerulonephritis: A long-term study of 40 cases." *J Pathol* 1983;141:449-468.
- [0502] Tao W, Wen R, Goddard MB, Sherman S, O'Rourke PJ, Stabila PF, Bell WJ, Dean BJ, Kauper KA, Budz VA, Tsiaras WG, Acland GM, Pearce-Kelling S, Laties AM, and Aguirre GD Encapsulated Cell-Based Delivery of CNTF Reduces Photoreceptor Degeneration in Animal Models of Retinitis Pigmentosa. *IOVS*, Vol. 43 (10) 3292-3298. 2002.
- [0503] Thelwell et al. *Nucleic Acids Res* 2000; 28:3752-3761.
- [0504] Timmerman et al. "Differential expression of complement components in human fetal and adult kidneys." *Kidney Int* 1996;49:730-740.
- [0505] Torzierski et al. "Processes in atherogenesis complement activation." *Atherosclerosis*. 1997; 132:131-138.
- [0506] Tuo et al. "Genetic factors in age-related macular degeneration." *Prog Retin Eye Res* 2004; 23(2):229-249.
- [0507] van den Dobbelsteen et al. "Regulation of C3 and factor H synthesis of human glomerular mesangial cells by IL-1 and interferon-gamma." *Clin Exp Immunol* 1994;95:173-180.

- [0508] Van Leeuwen et al. "Epidemiology of age-related macular degeneration." *Eur J Epidemiol* 2003; 18(9):845-854.
- [0509] Vingerling et al. "Epidemiology of age-related maculopathy." *Epidemiol Rev.* 1995;17(2):347-360.
- [0510] Vingerling et al. "The prevalence of age-related maculopathy in the Rotterdam Study." *Ophthalmol* 1995 Feb;102(2):205-210.
- [0511] Walport. "Complement. First of two parts." *N Engl J Med* 2001;344:1058-1066.
- [0512] Wang et al. "Systematic identification and analysis of exonic splicing silencers." *Cell* 119, 831-45 (2004).
- [0513] Weeks et al. "Age-related maculopathy: a genomewide scan with continued evidence of susceptibility loci within the 1q31, 10q26, and 17q25 regions." *Am J Hum Genet* 2004; 75:174.
- [0514] Weeks et al. "Age-related maculopathy: an expanded genome-wide scan with evidence of susceptibility loci within the 1q31 and 17q25 regions." *Am J Ophthalmol* 2001; 132:682-692.
- [0515] Weiler JM, Daha MR, Austen KF, Fearon DT. 1976. Control of the amplification convertase of complement by the plasma protein β 1H. *Proc. Natl. Acad. Sci. USA* 73:3268-3272.
- [0516] Zarbin. "Age Related Macular Degeneration: a review of pathogenesis." *Eur J Ophthalmol* 1998, 8:199-206.
- [0517] Zarbin. "Current concepts in the pathogenesis of age-related macular degeneration." *Arch Ophthalmol* 2004; 122(4):598-614.
- [0518] Zipfel et al. "Complement factor H and hemolytic uremic syndrome." *Int Immunopharmacol* 1, 461-8 (2001).
- [0519] Zipfel et al. "Factor H family proteins: on complement, microbes and human diseases." *Biochem Soc Trans* 30, 971-8 (2002).
- [0520] Zipfel et al. "The role of complement in membranoproliferative glomerulonephritis." In *Complement and Kidney Disease* 2005
- [0521] Zipfel. "Complement factor H: physiology and pathophysiology." *Semin Thromb Hemost* 27, 191-9 (2001).
- [0522] Zipfel. "Hemolytic uremic syndrome: how do factor H mutants mediate endothelial damage?" *Trends Immunol* 22, 345-8 (2001).

[0523] Although the present invention has been described in detail with reference to specific embodiments, those of skill in the art will recognize that modifications and improvements are within the scope and spirit of the invention, as set forth in the claims which follow. All publications and patent documents cited herein are incorporated herein by reference as if each such publication or document was specifically and individually indicated to be incorporated herein by reference. Citation of publications and patent documents (patents, published patent applications, and unpublished patent applications) is not intended as an admission that any such document is pertinent prior art, nor does it constitute any admission as to the contents or date of the same. The invention having now been described by way of written description, those of skill in the art will recognize that the invention can be practiced in a variety of embodiments and that the foregoing description is for purposes of illustration and not limitation of the following claims.

CLAIMS

1. A method of screening for susceptibility to developing age-related macular degeneration (AMD) in a subject, comprising screening for the presence or absence of a haplotype in the Complement Factor H (CFH) gene, wherein the haplotype comprises
 - (a) rs1061170 with a nucleotide encoding histidine at position 402; or
 - (b) rs800292 with a nucleotide encoding isoleucine at position 62.
2. The method of claim 1 comprising detecting in a biological sample from the subject the presence of a variant CFH polypeptide comprising histidine at position 402.
3. The method of claim 2 wherein the biological sample is a serum sample.
4. The method of claim 1 comprising detecting variation at a polymorphic site in a Factor H gene in the genome of the subject, wherein the variation is selected from a single nucleotide polymorphism at rs800292, rs1061170, or rs1065489.
5. The method of claim 4 wherein the detecting comprises (i) combining a nucleic acid sample from the subject with one or more polynucleotide probes capable of hybridizing selectively to a CFH gene allele and (ii) detecting the presence or absence of hybridization.
6. The method of claim 5 wherein the probes are oligonucleotides capable of priming polynucleotide synthesis in an amplification reaction.
7. The method of claim 1, comprising detecting a variant CFH allele in a polynucleotide from the subject.
8. The method of claim 1 comprising detecting a variant form of CFH protein in a sample from the subject.
9. The method of claim 7, wherein the polymorphic site is a non-synonymous single nucleotide polymorphism in the CFH gene selected from rs800292 encoding valine at position 62, rs1061170 encoding histidine at position 402, and a polymorphism in exon 22 encoding cysteine at position 1210.
10. The method of claim 7, wherein the polymorphic site is a single nucleotide polymorphism in the CFH gene selected from rs1061147, intron 2 (IVS2 or insTT); rs2274700;

rs1061170, and exon 10A.

11. The method of claim 8, wherein the detecting step comprises
 - (i) combining a nucleic acid sample from the subject with one or more polynucleotide probes capable of hybridizing selectively to a CFH gene allele; and
 - (ii) detecting the presence or absence of hybridization.
12. The method of claim 8 wherein the variant CFH protein comprises valine at position 62, histidine at position 402, or cysteine at position 1210.
13. The method of claim 11, wherein the probes are oligonucleotides capable of priming polynucleotide synthesis in an amplification reaction.
14. The method of claim 8 or 12, wherein the sample is blood or serum, and said detecting comprise combining the sample and an antibody specific for a variant form of the protein and detecting binding between the antibody and the variant form of the protein, if present.
15. The method of any one of claims 7-14, wherein the presence of the variation is an indication that the patient has an increased propensity to develop AMD.
16. The method of any one of claims 7-15, wherein the presence of the variation is an indication that the patient has a decreased propensity to develop AMD.
17. A method of screening for susceptibility to age-related macular degeneration (AMD) in a subject, comprising screening for a haplotype, wherein the haplotype is:
 - a) a risk haplotype in the CFH gene that is more frequently present in individuals diagnosed with AMD compared to healthy individuals, or
 - b) a protective haplotype in the CFH gene that is more frequently present in healthy individuals compared to individuals diagnosed with AMD or
 - c) a neutral haplotype in the CFH gene; andwherein said screening comprises detecting the presence or absence of a variant sequence at any one or more of rs1061170, rs800292, rs1061147, intron 2 (IVS2 or insTT); rs2274700; rs1061170, exon 10A or exon 22.
18. The method of claim 17, wherein the haplotype is in the gene.

19. The method of claim 18, wherein the haplotype is a risk haplotype in the CFHR5 gene that comprises a variant sequence at least one of rs9427661, rs9427662 and rs12097550.

20. A device or kit for screening susceptibility to AMD in a subject, comprising oligonucleotides that distinguish alleles at one, or more than one, CFH gene polymorphic site listed in Table A.

21. The device or kit of claim 20, wherein the oligonucleotides are primers for nucleic acid amplification of a region spanning a CFH gene polymorphic site listed in Table A, or probes for nucleic acid hybridization to a region spanning a CFH gene polymorphic site.

22. A device or kit for screening for susceptibility to AMD in a subject, comprising oligonucleotides that distinguish alleles at one, or more than one, of rs1061170, rs800292, rs1061147, intron 2 (IVS2 or insTT); rs2274700; rs1061170, rs9427661, rs9427662, rs12097550, exon 10A, and exon 22.

23. The method of claim 1 wherein the step of screening for the presence or absence of a haplotype consists of screening for the presence or absence of one variant polymorphism and said one variant polymorphism is a single nucleotide polymorphism (SNP) at rs800292, rs1061170, or rs1065489.

24. The method of claim 23 wherein the detecting comprises (i) combining a nucleic acid sample from the subject with one or more polynucleotide probes capable of hybridizing selectively to a CFH gene allele and (ii) detecting the presence or absence of hybridization.

25. The method of claim 24 wherein the probes are oligonucleotides capable of priming polynucleotide synthesis in an amplification reaction.

26. The method of any of claims 1 to 25, further comprising screening for specific subtypes of AMD.

27. The method of claim 26, wherein the subtype is early AMD, geographic atrophy, or exudative AMD (CNV).

SEQUENCE LISTING

<110> University of Iowa Research Foundation
Hageman, Gregory S.

<120> Methods and Reagents for the Treatment of Age-Related Macular
Degeneration

<130> 020618-001220PC

<140> WO 00/000,000

<141> 2006-02-14

<150> US 60/650,078

<151> 2005-02-14

<150> US 60/717,861

<151> 2005-09-16

<150> US 60/715,503

<151> 2005-09-09

<150> US 60/735,697

<151> 2005-11-09

<160> 337

<170> PatentIn version 3.3

<210> 1

<211> 3926

<212> DNA

<213> Homo sapiens

<400> 1

```

aattcttggga agaggagaac tggacgttgt gaacagagtt agctggtaaa tgtcctctta      60
aaagatccaa aaaatgagac ttctagcaaa gattatttgc cttatgttat gggctatttg      120
tgtagcagaa gattgcaatg aacttcotcc aagaagaaat acagaaaatc tgacagggttc      180
ctgggtctgac caaacatatc cagaaggcac ccaggctatc tataaatgcc gccctggata      240
tagatctott ggaaatgtaa taatggtatg caggaaggga gaatgggttg ctcttaatcc      300
attaaggaaa tgtcagaaaa ggccctgtgg acatcctgga gatactcctt ttggtacttt      360
tacccttaca ggaggaaatg tgtttgaata tgggtgtaaaa gctgtgtata catgtaatga      420
ggggtatcaa ttgctaggtg agattaatta ccgtgaatgt gacacagatg gatggaccaa      480
tgatattcct atatgtgaag ttgtgaagtg tttaccagtg acagcaccag agaatggaaa      540
aattgtcagt agtgcaatgg aaccagatcg ggaataccat tttggacaag cagtacggtt      600
tgtatgtaac tcaggctaca agattgaagg agatgaagaa atgcattggt cagacgatgg      660
tttttgagat aaagagaaac caaagtgtgt ggaaatttca tgcaaatccc cagatgttat      720
aaatggatct cctatatctc agaagattat ttataaggag aatgaacgat ttcaatataa      780
atgtaacatg ggttatgaat acagtgaaag aggagatgct gtatgcactg aatctggatg      840

```

gcgtccgttg ccttcattgtg aagaaaaatc atgtgataat ccttatattc caaatgggtga 900
 ctactcacct ttaaggatta aacacagAAC tggagatgaa atcacgtacc agtgtagaaa 960
 tggtttttat cctgcaaccc ggggaaatac agccaaatgc acaagtactg gctggatacc 1020
 tgctccgaga tgtacottga aacotttgtga ttatccagac attaaacatg gaggtotata 1080
 tcatgagaat atgcgtagac catactttcc agtagctgta ggaaaatatt actcctatta 1140
 ctgtgatgaa ctttttgaga ctccgtcagg aagttactgg gatcacattc attgcacaca 1200

 agatggatgg tgcacagcag taccatgcct cagaaaatgt ttttttctt atttgaaaa 1260
 tggatataat caaatcatg gaagaaagt tgtacagggt aaatctatag acgttgccctg 1320
 ccatcctggc tacgtcttc caaaagcgca gaccacagtt acatgtatgg agaatggctg 1380
 gtctcctact ccagatgca tccgtgtcaa aacatgttcc aaatcaagta tagatattga 1440
 gaatgggttt atttctgaat ctcagtatac atatgcctta aaagaaaaag cgaaatatca 1500
 atgcaaaacta ggatatgtaa cagcagatgg tgaacatca ggatcaatta gatgtgggaa 1560
 agatggatgg tcagctcaac ccacgtgcat taaatcttgt gatatcccag ttttatgaa 1620
 tgccagaact aaaaatgact tcacatgggt taagctgaat gacacattgg actatgaatg 1680
 coatgatgg tatgaaagca atactggaag caccactgggt tccatagtgt gtggttacaa 1740
 tggttgggtct gatttaccca tatgttatga aagagaatgc gaacttccta aaatagatgt 1800
 acacttagtt cctgatcgca agaaagacca gtataaagt ggagaggtgt tgaaattctc 1860
 ctgcaaacca ggatttacia tagttggacc taattccgtt cagtgtctacc actttggatt 1920
 gtctcctgac ctcccaatat gttaaagagca agtacaatca tgtgggtccac ctctgaact 1980
 cctcaatggg aatgttaagg aaaaaacgaa agaagaatat ggacacagtg aagtgggtga 2040
 atattattgc aatcctagat ttctaataa gggacctaataaaaattcaat gtgtgatgg 2100
 agagtggaca actttaccag tgtgtattgt ggaggagagt acctgtggag atatacctga 2160
 acttgaacat ggctgggccc agctttcttc cctccttat tactatggag attcagtga 2220
 attcaattgc tcagaatcat ttacaatgat tggacacaga tcaattacgt gtattcatgg 2280
 agtatggacc caacttcccc agtgtgtggc aatagataaa cttagaagt gcaaatcatc 2340
 aaatttaatt atacttgagg aacattttaa aaacaagaag gaattcgatc ataattctaa 2400
 cataaggtag agatgtagag gaaaagaagg atggatacac acagtctgca taaatggaag 2460
 atgggatcca gaagtgaact gctcaatggc acaaatataa ttatgccac ctccacotca 2520
 gattcccaat tctcacaata tgacaaccac actgaattat cgggatggag aaaaagtatc 2580
 tgttctttgc caagaaaatt atctaattca ggaaggagaa gaaattacat gcaaagatgg 2640

aagatggcag tcaataccac totgtgttga aaaaattcca tgttcacaac cacctcagat 2700
agaacacgga accattaatt catccaggto ttcacaagaa agttatgcac atgggactaa 2760
attgagttat acttgtgagg gtgggtttcag gatattctgaa gaaaatgaaa caacatgcta 2820
catgggaaaaa tggagttctc cacctcagtg tgaaggcctt ccttgtaaatt ctccacctga 2880
gattttotcat ggtgtttag ctccacatgtc agacagttat cagtatggag aagaagttac 2940
gtacaaatgt tttgaagggt ttggaattga tgggcctgca attgcaaaat gcttaggaga 3000
aaaatggtct caccctccat catgcataaa aacagattgt ctccagtttac cttagctttga 3060
aaatgccata cccatgggag agaagaagga tgtgtataag gcgggtgagc aagtgactta 3120
cacttgtgca acatattaca aaatggatgg agccagtaat gtaacatgca ttaatagcag 3180
atggacagga aggccaaacat gcagagacac ctctgtgtg aatccgccca cagtacaaaa 3240
tgcttatata gtgtcgagac agatgagtaa atatccatct ggtgagagag tacgttatca 3300
atgtaggagc cttatgaaa tgtttgggga tgaagaagtg atgtgtttaa atggaaactg 3360
gacggaacca cctcaatgca aagattctac aggaaaatgt gggccccctc cacctattga 3420
caatggggac attacttcat tcccgttgtc agtatatgct ccagcttcat cagttgagta 3480
ccaatgccag aacttgtatc aacttgaggg taacaagcga ataacatgta gaaatggaca 3540
atggtcagaa ccacaaaaat gttacatcc gtgtgtaata tcccgagaaa ttatggaaaa 3600
ttataacata gcattaaggt ggacagccaa acagaagctt tattcgagaa caggtgaatc 3660
agttgaatth gtgtgtaaac ggggatatcg tctttcatca cgtttctaca cattgcgaac 3720
aacatgttgg gatgggaaac tggagtatcc aacttgtgca aaaagataga atcaatcata 3780
aagtgcacac ctttattcag aactttagta ttaaatcagt tctcaatttc attttttatg 3840
tattgtttta ctctttttta ttcatacgtg aaattttgga ttaattttgtg aaaatgtaat 3900
tataagctga gaccggtggc tctctt 3926

<210> 2

<211> 1231

<212> PRT

<213> Homo sapiens

<400> 2

Met Arg Leu Leu Ala Lys Ile Ile Cys Leu Met Leu Trp Ala Ile Cys
1 5 10 15

Val Ala Glu Asp Cys Asn Glu Leu Pro Pro Arg Arg Asn Thr Glu Ile
20 25 30

Leu Thr Gly Ser Trp Ser Asp Gln Thr Tyr Pro Glu Gly Thr Gln Ala
 35 40 45
 Ile Tyr Lys Cys Arg Pro Gly Tyr Arg Ser Leu Gly Asn Val Ile Met
 50 55 60
 Val Cys Arg Lys Gly Glu Trp Val Ala Leu Asn Pro Leu Arg Lys Cys
 65 70 75 80
 Gln Lys Arg Pro Cys Gly His Pro Gly Asp Thr Pro Phe Gly Thr Phe
 85 90 95
 Thr Leu Thr Gly Gly Asn Val Phe Glu Tyr Gly Val Lys Ala Val Tyr
 100 105 110
 Thr Cys Asn Glu Gly Tyr Gln Leu Leu Gly Glu Ile Asn Tyr Arg Glu
 115 120 125
 Cys Asp Thr Asp Gly Trp Thr Asn Asp Ile Pro Ile Cys Glu Val Val
 130 135 140
 Lys Cys Leu Pro Val Thr Ala Pro Glu Asn Gly Lys Ile Val Ser Ser
 145 150 155 160
 Ala Met Glu Pro Asp Arg Glu Tyr His Phe Gly Gln Ala Val Arg Phe
 165 170 175
 Val Cys Asn Ser Gly Tyr Lys Ile Glu Gly Asp Glu Glu Met His Cys
 180 185 190
 Ser Asp Asp Gly Phe Trp Ser Lys Glu Lys Pro Lys Cys Val Glu Ile
 195 200 205
 Ser Cys Lys Ser Pro Asp Val Ile Asn Gly Ser Pro Ile Ser Gln Lys
 210 215 220
 Ile Ile Tyr Lys Glu Asn Glu Arg Phe Gln Tyr Lys Cys Asn Met Gly
 225 230 235 240
 Tyr Glu Tyr Ser Glu Arg Gly Asp Ala Val Cys Thr Glu Ser Gly Trp
 245 250 255
 Arg Pro Leu Pro Ser Cys Glu Glu Lys Ser Cys Asp Asn Pro Tyr Ile
 260 265 270

Pro Asn Gly Asp Tyr Ser Pro Leu Arg Ile Lys His Arg Thr Gly Asp
 275 280 285

Glu Ile Thr Tyr Gln Cys Arg Asn Gly Phe Tyr Pro Ala Thr Arg Gly
 290 295 300

Asn Thr Ala Lys Cys Thr Ser Thr Gly Trp Ile Pro Ala Pro Arg Cys
 305 310 315 320

Thr Leu Lys Pro Cys Asp Tyr Pro Asp Ile Lys His Gly Gly Leu Tyr
 325 330 335

His Glu Asn Met Arg Arg Pro Tyr Phe Pro Val Ala Val Gly Lys Tyr
 340 345 350

Tyr Ser Tyr Tyr Cys Asp Glu His Phe Glu Thr Pro Ser Gly Ser Tyr
 355 360 365

Trp Asp His Ile His Cys Thr Gln Asp Gly Trp Ser Pro Ala Val Pro
 370 375 380

Cys Leu Arg Lys Cys Tyr Phe Pro Tyr Leu Glu Asn Gly Tyr Asn Gln
 385 390 395 400

Asn His Gly Arg Lys Phe Val Gln Gly Lys Ser Ile Asp Val Ala Cys
 405 410 415

His Pro Gly Tyr Ala Leu Pro Lys Ala Gln Thr Thr Val Thr Cys Met
 420 425 430

Glu Asn Gly Trp Ser Pro Thr Pro Arg Cys Ile Arg Val Lys Thr Cys
 435 440 445

Ser Lys Ser Ser Ile Asp Ile Glu Asn Gly Phe Ile Ser Glu Ser Gln
 450 455 460

Tyr Thr Tyr Ala Leu Lys Glu Lys Ala Lys Tyr Gln Cys Lys Leu Gly
 465 470 475 480

Tyr Val Thr Ala Asp Gly Glu Thr Ser Gly Ser Ile Arg Cys Gly Lys
 485 490 495

Asp Gly Trp Ser Ala Gln Pro Thr Cys Ile Lys Ser Cys Asp Ile Pro
 500 505 510

Val Phe Met Asn Ala Arg Thr Lys Asn Asp Phe Thr Trp Phe Lys Leu

515	520	525
Asn Asp Thr Leu Asp Tyr Glu Cys His Asp Gly Tyr Glu Ser Asn Thr		
530	535	540
Gly Ser Thr Thr Gly Ser Ile Val Cys Gly Tyr Asn Gly Trp Ser Asp		
545	550	555
Leu Pro Ile Cys Tyr Glu Arg Glu Cys Glu Leu Pro Lys Ile Asp Val		
	565	570
His Leu Val Pro Asp Arg Lys Lys Asp Gln Tyr Lys Val Gly Glu Val		
	580	585
Leu Lys Phe Ser Cys Lys Pro Gly Phe Thr Ile Val Gly Pro Asn Ser		
	595	600
Val Gln Cys Tyr His Phe Gly Leu Ser Pro Asp Leu Pro Ile Cys Lys		
	610	615
Glu Gln Val Gln Ser Cys Gly Pro Pro Pro Glu Leu Leu Asn Gly Asn		
	625	630
Val Lys Glu Lys Thr Lys Glu Glu Tyr Gly His Ser Glu Val Val Glu		
	645	650
Tyr Tyr Cys Asn Pro Arg Phe Leu Met Lys Gly Pro Asn Lys Ile Gln		
	660	665
Cys Val Asp Gly Glu Trp Thr Thr Leu Pro Val Cys Ile Val Glu Glu		
	675	680
Ser Thr Cys Gly Asp Ile Pro Glu Leu Glu His Gly Trp Ala Gln Leu		
	690	700
Ser Ser Pro Pro Tyr Tyr Tyr Gly Asp Ser Val Glu Phe Asn Cys Ser		
	705	710
Glu Ser Phe Thr Met Ile Gly His Arg Ser Ile Thr Cys Ile His Gly		
	725	730
Val Trp Thr Gln Leu Pro Gln Cys Val Ala Ile Asp Lys Leu Lys Lys		
	740	745
Cys Lys Ser Ser Asn Leu Ile Ile Leu Glu Glu His Leu Lys Asn Lys		

755	760	765
Lys Glu Phe Asp His Asn Ser Asn Ile Arg Tyr Arg Cys Arg Gly Lys		
770	775	780
Glu Gly Trp Ile His Thr Val Cys Ile Asn Gly Arg Trp Asp Pro Glu		
785	790	795 800
Val Asn Cys Ser Met Ala Gln Ile Gln Leu Cys Pro Pro Pro Pro Gln		
	805	810 815
Ile Pro Asn Ser His Asn Met Thr Thr Thr Leu Asn Tyr Arg Asp Gly		
	820	825 830
Glu Lys Val Ser Val Leu Cys Gln Glu Asn Tyr Leu Ile Gln Glu Gly		
	835	840 845
Glu Glu Ile Thr Cys Lys Asp Gly Arg Trp Gln Ser Ile Pro Leu Cys		
	850	855 860
Val Glu Lys Ile Pro Cys Ser Gln Pro Pro Gln Ile Glu His Gly Thr		
	865	870 875 880
Ile Asn Ser Ser Arg Ser Ser Gln Glu Ser Tyr Ala His Gly Thr Lys		
	885	890 895
Leu Ser Tyr Thr Cys Glu Gly Gly Phe Arg Ile Ser Glu Glu Asn Glu		
	900	905 910
Thr Thr Cys Tyr Met Gly Lys Trp Ser Ser Pro Pro Gln Cys Glu Gly		
	915	920 925
Leu Pro Cys Lys Ser Pro Pro Glu Ile Ser His Gly Val Val Ala His		
	930	935 940
Met Ser Asp Ser Tyr Gln Tyr Gly Glu Glu Val Thr Tyr Lys Cys Phe		
	945	950 955 960
Glu Gly Phe Gly Ile Asp Gly Pro Ala Ile Ala Lys Cys Leu Gly Glu		
	965	970 975
Lys Trp Ser His Pro Pro Ser Cys Ile Lys Thr Asp Cys Leu Ser Leu		
	980	985 990
Pro Ser Phe Glu Asn Ala Ile Pro Met Gly Glu Lys Lys Asp Val Tyr		
	995	1000 1005

Lys Ala Gly Glu Gln Val Thr Tyr Thr Cys Ala Thr Tyr Tyr Lys
 1010 1015 1020
 Met Asp Gly Ala Ser Asn Val Thr Cys Ile Asn Ser Arg Trp Thr
 1025 1030 1035
 Gly Arg Pro Thr Cys Arg Asp Thr Ser Cys Val Asn Pro Pro Thr
 1040 1045 1050
 Val Gln Asn Ala Tyr Ile Val Ser Arg Gln Met Ser Lys Tyr Pro
 1055 1060 1065
 Ser Gly Glu Arg Val Arg Tyr Gln Cys Arg Ser Pro Tyr Glu Met
 1070 1075 1080
 Phe Gly Asp Glu Glu Val Met Cys Leu Asn Gly Asn Trp Thr Glu
 1085 1090 1095
 Pro Pro Gln Cys Lys Asp Ser Thr Gly Lys Cys Gly Pro Pro Pro
 1100 1105 1110
 Pro Ile Asp Asn Gly Asp Ile Thr Ser Phe Pro Leu Ser Val Tyr
 1115 1120 1125
 Ala Pro Ala Ser Ser Val Glu Tyr Gln Cys Gln Asn Leu Tyr Gln
 1130 1135 1140
 Leu Glu Gly Asn Lys Arg Ile Thr Cys Arg Asn Gly Gln Trp Ser
 1145 1150 1155
 Glu Pro Pro Lys Cys Leu His Pro Cys Val Ile Ser Arg Glu Ile
 1160 1165 1170
 Met Glu Asn Tyr Asn Ile Ala Leu Arg Trp Thr Ala Lys Gln Lys
 1175 1180 1185
 Leu Tyr Ser Arg Thr Gly Glu Ser Val Glu Phe Val Cys Lys Arg
 1190 1195 1200
 Gly Tyr Arg Leu Ser Ser Arg Ser His Thr Leu Arg Thr Thr Cys
 1205 1210 1215
 Trp Asp Gly Lys Leu Glu Tyr Pro Thr Cys Ala Lys Arg
 1220 1225 1230

<210> 3
 <211> 1658
 <212> DNA
 <213> Homo sapiens

<400> 3
 aattcttgga agaggagaac tggacgttgt gaacagagtt agctggtaaa tgcctcttta 60
 aaagatccaa aaaatgagac ttctagcaaa gattatattgc cttatgttat gggctatttg 120
 tgtagcagaa gattgcaatg aacttcctcc aagaagaaat acagaaattc tgacaggttc 180
 ctggtctgac caaacatata cagaaggcac ccaggctata tataaatgcc gccctggata 240
 tagatctctt ggaaatgtaa taatggtatg caggaaggga gaatgggttg ctcttaatcc 300
 attaaggaaa tgcagaaaaa ggccctgttg acatcctgga gatactcctt ttggtacttt 360
 tacccttaca ggaggaaatg tgtttgaata tgggtgtaaaa gctgtgtata catgtaatga 420
 ggggtatcaa ttgctagggtg agattaatta ccgtgaatgt gacacagatg gatggaccaa 480
 tgatattcct atatgtgaag ttgtgaagtg tttaccagtg acagcaccag agaatggaaa 540
 aattgtcagt agtgcaatgg aaccagatcg ggaataccat tttggacaag cagtacgggt 600
 tgtatgtaac tcaggctaca agattgaagg agatgaagaa atgcattgtt cagacgatgg 660
 tttttggagt aaagagaaac caaagtgtgt ggaaatttca tgcaaatccc cagatgttat 720
 aaatggatct cctatatctc agaagattat ttataaggag aatgaacgat ttcaatataa 780
 atgtaacatg ggttatgaat acagtgaag aggagatgct gtatgcactg aatctggatg 840
 gcgtccgttg ccttcagtgt aagaaaaatc atgtgataat ccttatattc caaatggtga 900
 ctactcacct ttaaggatta aacacagaac tggagatgaa atcacgtacc agtgtagaaa 960
 tggtttttat cctgcaaccc ggggaaatac agccaaatgc acaagtactg gctggatacc 1020
 tgctccgaga tgtaccttga aaccttgtga ttatccagac attaaacatg gaggtctata 1080
 tcatgagaat atgcgtagac catactttcc agtagctgta ggaaaatatt actcctatta 1140
 ctgtgatgaa cattttgaga ctccgtcagg aagtactgg gatcacattc attgcacaca 1200
 agatggatgg togccagcag taccatgcct cagaaaatgt tattttcctt atttgaaaaa 1260
 tggatataat caaaattatg gaagaaagtt tgtacagggt aaatctatag acgttgccctg 1320
 ccacctctgc tacgctottc caaaagcgca gaccacagtt acatgtatgg agaatggctg 1380
 gtctcctact ccagatgca tccgtgtcag ctttaccctc tgaacttctg atcgaaggtc 1440
 atccctctcc agcttgagtg gatcaaagat gacaagggcc aatggaacca agtttgagtc 1500
 ttgccaggtc aatacttggg tcttgagtat ggtgactagt atctgttttg ttatgtgtgt 1560
 attattccag ccagaatggg aaatgctaata tcagctcctc caggcagcca atggggctgg 1620

tggttttgag attattaaac ttttctggat cctctacg

1658

<210> 4

<211> 449

<212> PRT

<213> Homo sapiens

<400> 4

Met Arg Leu Leu Ala Lys Ile Ile Cys Leu Met Leu Trp Ala Ile Cys
1 5 10 15

Val Ala Glu Asp Cys Asn Glu Leu Pro Pro Arg Arg Asn Thr Glu Ile
20 25 30

Leu Thr Gly Ser Trp Ser Asp Gln Thr Tyr Pro Glu Gly Thr Gln Ala
35 40 45

Ile Tyr Lys Cys Arg Pro Gly Tyr Arg Ser Leu Gly Asn Val Ile Met
50 55 60

Val Cys Arg Lys Gly Glu Trp Val Ala Leu Asn Pro Leu Arg Lys Cys
65 70 75 80

Gln Lys Arg Pro Cys Gly His Pro Gly Asp Thr Pro Phe Gly Thr Phe
85 90 95

Thr Leu Thr Gly Gly Asn Val Phe Glu Tyr Gly Val Lys Ala Val Tyr
100 105 110

Thr Cys Asn Glu Gly Tyr Gln Leu Leu Gly Glu Ile Asn Tyr Arg Glu
115 120 125

Cys Asp Thr Asp Gly Trp Thr Asn Asp Ile Pro Ile Cys Glu Val Val
130 135 140

Lys Cys Leu Pro Val Thr Ala Pro Glu Asn Gly Lys Ile Val Ser Ser
145 150 155 160

Ala Met Glu Pro Asp Arg Glu Tyr His Phe Gly Gln Ala Val Arg Phe
165 170 175

Val Cys Asn Ser Gly Tyr Lys Ile Glu Gly Asp Glu Glu Met His Cys
180 185 190

Ser Asp Asp Gly Phe Trp Ser Lys Glu Lys Pro Lys Cys Val Glu Ile
195 200 205

Ser Cys Lys Ser Pro Asp Val Ile Asn Gly Ser Pro Ile Ser Gln Lys
 210 215 220

Ile Ile Tyr Lys Glu Asn Glu Arg Phe Gln Tyr Lys Cys Asn Met Gly
 225 230 235 240

Tyr Glu Tyr Ser Glu Arg Gly Asp Ala Val Cys Thr Glu Ser Gly Trp
 245 250 255

Arg Pro Leu Pro Ser Cys Glu Glu Lys Ser Cys Asp Asn Pro Tyr Ile
 260 265 270

Pro Asn Gly Asp Tyr Ser Pro Leu Arg Ile Lys His Arg Thr Gly Asp
 275 280 285

Glu Ile Thr Tyr Gln Cys Arg Asn Gly Phe Tyr Pro Ala Thr Arg Gly
 290 295 300

Asn Thr Ala Lys Cys Thr Ser Thr Gly Trp Ile Pro Ala Pro Arg Cys
 305 310 315 320

Thr Leu Lys Pro Cys Asp Tyr Pro Asp Ile Lys His Gly Gly Leu Tyr
 325 330 335

His Glu Asn Met Arg Arg Pro Tyr Phe Pro Val Ala Val Gly Lys Tyr
 340 345 350

Tyr Ser Tyr Tyr Cys Asp Glu His Phe Glu Thr Pro Ser Gly Ser Tyr
 355 360 365

Trp Asp His Ile His Cys Thr Gln Asp Gly Trp Ser Pro Ala Val Pro
 370 375 380

Cys Leu Arg Lys Cys Tyr Phe Pro Tyr Leu Glu Asn Gly Tyr Asn Gln
 385 390 395 400

Asn Tyr Gly Arg Lys Phe Val Gln Gly Lys Ser Ile Asp Val Ala Cys
 405 410 415

His Pro Gly Tyr Ala Leu Pro Lys Ala Gln Thr Thr Val Thr Cys Met
 420 425 430

Glu Asn Gly Trp Ser Pro Thr Pro Arg Cys Ile Arg Val Ser Phe Thr
 435 440 445

Leu

<210> 5
 <211> 1231
 <212> PRT
 <213> Artificial

<220>
 <223> Synthetic Factor H variant

<400> 5

Met Arg Leu Leu Ala Lys Ile Ile Cys Leu Met Leu Trp Ala Ile Cys
 1 5 10 15

Val Ala Glu Asp Cys Asn Glu Leu Pro Pro Arg Arg Asn Thr Glu Ile
 20 25 30

Leu Thr Gly Ser Trp Ser Asp Gln Thr Tyr Pro Glu Gly Thr Gln Ala
 35 40 45

Ile Tyr Lys Cys Arg Pro Gly Tyr Arg Ser Leu Gly Asn Ile Ile Met
 50 55 60

Val Cys Arg Lys Gly Glu Trp Val Ala Leu Asn Pro Leu Arg Lys Cys
 65 70 75 80

Gln Lys Arg Pro Cys Gly His Pro Gly Asp Thr Pro Phe Gly Thr Phe
 85 90 95

Thr Leu Thr Gly Gly Asn Val Phe Glu Tyr Gly Val Lys Ala Val Tyr
 100 105 110

Thr Cys Asn Glu Gly Tyr Gln Leu Leu Gly Glu Ile Asn Tyr Arg Glu
 115 120 125

Cys Asp Thr Asp Gly Trp Thr Asn Asp Ile Pro Ile Cys Glu Val Val
 130 135 140

Lys Cys Leu Pro Val Thr Ala Pro Glu Asn Gly Lys Ile Val Ser Ser
 145 150 155 160

Ala Met Glu Pro Asp Arg Glu Tyr His Phe Gly Gln Ala Val Arg Phe
 165 170 175

Val Cys Asn Ser Gly Tyr Lys Ile Glu Gly Asp Glu Glu Met His Cys
 180 185 190

Ser Asp Asp Gly Phe Trp Ser Lys Glu Lys Pro Lys Cys Val Glu Ile
195 200 205

Ser Cys Lys Ser Pro Asp Val Ile Asn Gly Ser Pro Ile Ser Gln Lys
210 215 220

Ile Ile Tyr Lys Glu Asn Glu Arg Phe Gln Tyr Lys Cys Asn Met Gly
225 230 235 240

Tyr Glu Tyr Ser Glu Arg Gly Asp Ala Val Cys Thr Glu Ser Gly Trp
245 250 255

Arg Pro Leu Pro Ser Cys Glu Glu Lys Ser Cys Asp Asn Pro Tyr Ile
260 265 270

Pro Asn Gly Asp Tyr Ser Pro Leu Arg Ile Lys His Arg Thr Gly Asp
275 280 285

Glu Ile Thr Tyr Gln Cys Arg Asn Gly Phe Tyr Pro Ala Thr Arg Gly
290 295 300

Asn Thr Ala Lys Cys Thr Ser Thr Gly Trp Ile Pro Ala Pro Arg Cys
305 310 315 320

Thr Leu Lys Pro Cys Asp Tyr Pro Asp Ile Lys His Gly Gly Leu Tyr
325 330 335

His Glu Asn Met Arg Arg Pro Tyr Phe Pro Val Ala Val Gly Lys Tyr
340 345 350

Tyr Ser Tyr Tyr Cys Asp Glu His Phe Glu Thr Pro Ser Gly Ser Tyr
355 360 365

Trp Asp His Ile His Cys Thr Gln Asp Gly Trp Ser Pro Ala Val Pro
370 375 380

Cys Leu Arg Lys Cys Tyr Phe Pro Tyr Leu Glu Asn Gly Tyr Asn Gln
385 390 395 400

Asn Tyr Gly Arg Lys Phe Val Gln Gly Lys Ser Ile Asp Val Ala Cys
405 410 415

His Pro Gly Tyr Ala Leu Pro Lys Ala Gln Thr Thr Val Thr Cys Met
420 425 430

Glu Asn Gly Trp Ser Pro Thr Pro Arg Cys Ile Arg Val Lys Thr Cys
 435 440 445

Ser Lys Ser Ser Ile Asp Ile Glu Asn Gly Phe Ile Ser Glu Ser Gln
 450 455 460

Tyr Thr Tyr Ala Leu Lys Glu Lys Ala Lys Tyr Gln Cys Lys Leu Gly
 465 470 475 480

Tyr Val Thr Ala Asp Gly Glu Thr Ser Gly Ser Ile Arg Cys Gly Lys
 485 490 495

Asp Gly Trp Ser Ala Gln Pro Thr Cys Ile Lys Ser Cys Asp Ile Pro
 500 505 510

Val Phe Met Asn Ala Arg Thr Lys Asn Asp Phe Thr Trp Phe Lys Leu
 515 520 525

Asn Asp Thr Leu Asp Tyr Glu Cys His Asp Gly Tyr Glu Ser Asn Thr
 530 535 540

Gly Ser Thr Thr Gly Ser Ile Val Cys Gly Tyr Asn Gly Trp Ser Asp
 545 550 555 560

Leu Pro Ile Cys Tyr Glu Arg Glu Cys Glu Leu Pro Lys Ile Asp Val
 565 570 575

His Leu Val Pro Asp Arg Lys Lys Asp Gln Tyr Lys Val Gly Glu Val
 580 585 590

Leu Lys Phe Ser Cys Lys Pro Gly Phe Thr Ile Val Gly Pro Asn Ser
 595 600 605

Val Gln Cys Tyr His Phe Gly Leu Ser Pro Asp Leu Pro Ile Cys Lys
 610 615 620

Glu Gln Val Gln Ser Cys Gly Pro Pro Pro Glu Leu Leu Asn Gly Asn
 625 630 635 640

Val Lys Glu Lys Thr Lys Glu Glu Tyr Gly His Ser Glu Val Val Glu
 645 650 655

Tyr Tyr Cys Asn Pro Arg Phe Leu Met Lys Gly Pro Asn Lys Ile Gln
 660 665 670

Cys Val Asp Gly Glu Trp Thr Thr Leu Pro Val Cys Ile Val Glu Glu
675 680 685

Ser Thr Cys Gly Asp Ile Pro Glu Leu Glu His Gly Trp Ala Gln Leu
690 695 700

Ser Ser Pro Pro Tyr Tyr Tyr Gly Asp Ser Val Glu Phe Asn Cys Ser
705 710 715 720

Glu Ser Phe Thr Met Ile Gly His Arg Ser Ile Thr Cys Ile His Gly
725 730 735

Val Trp Thr Gln Leu Pro Gln Cys Val Ala Ile Asp Lys Leu Lys Lys
740 745 750

Cys Lys Ser Ser Asn Leu Ile Ile Leu Glu Glu His Leu Lys Asn Lys
755 760 765

Lys Glu Phe Asp His Asn Ser Asn Ile Arg Tyr Arg Cys Arg Gly Lys
770 775 780

Glu Gly Trp Ile His Thr Val Cys Ile Asn Gly Arg Trp Asp Pro Glu
785 790 795 800

Val Asn Cys Ser Met Ala Gln Ile Gln Leu Cys Pro Pro Pro Pro Gln
805 810 815

Ile Pro Asn Ser His Asn Met Thr Thr Thr Leu Asn Tyr Arg Asp Gly
820 825 830

Glu Lys Val Ser Val Leu Cys Gln Glu Asn Tyr Leu Ile Gln Glu Gly
835 840 845

Glu Glu Ile Thr Cys Lys Asp Gly Arg Trp Gln Ser Ile Pro Leu Cys
850 855 860

Val Glu Lys Ile Pro Cys Ser Gln Pro Pro Gln Ile Glu His Gly Thr
865 870 875 880

Ile Asn Ser Ser Arg Ser Ser Gln Glu Ser Tyr Ala His Gly Thr Lys
885 890 895

Leu Ser Tyr Thr Cys Glu Gly Gly Phe Arg Ile Ser Glu Glu Asn Glu
900 905 910

Thr Thr Cys Tyr Met Gly Lys Trp Ser Ser Pro Pro Gln Cys Glu Gly

915	920	925
Leu Pro Cys Lys Ser Pro Pro Glu Ile Ser His Gly Val Val Ala His		
930	935	940
Met Ser Asp Ser Tyr Gln Tyr Gly Glu Glu Val Thr Tyr Lys Cys Phe		
945	950	955 960
Glu Gly Phe Gly Ile Asp Gly Pro Ala Ile Ala Lys Cys Leu Gly Glu		
	965	970 975
Lys Trp Ser His Pro Pro Ser Cys Ile Lys Thr Asp Cys Leu Ser Leu		
	980	985 990
Pro Ser Phe Glu Asn Ala Ile Pro Met Gly Glu Lys Lys Asp Val Tyr		
	995	1000 1005
Lys Ala Gly Glu Gln Val Thr Tyr Thr Cys Ala Thr Tyr Tyr Lys		
	1010	1015 1020
Met Asp Gly Ala Ser Asn Val Thr Cys Ile Asn Ser Arg Trp Thr		
	1025	1030 1035
Gly Arg Pro Thr Cys Arg Asp Thr Ser Cys Val Asn Pro Pro Thr		
	1040	1045 1050
Val Gln Asn Ala Tyr Ile Val Ser Arg Gln Met Ser Lys Tyr Pro		
	1055	1060 1065
Ser Gly Glu Arg Val Arg Tyr Gln Cys Arg Ser Pro Tyr Glu Met		
	1070	1075 1080
Phe Gly Asp Glu Glu Val Met Cys Leu Asn Gly Asn Trp Thr Glu		
	1085	1090 1095
Pro Pro Gln Cys Lys Asp Ser Thr Gly Lys Cys Gly Pro Pro Pro		
	1100	1105 1110
Pro Ile Asp Asn Gly Asp Ile Thr Ser Phe Pro Leu Ser Val Tyr		
	1115	1120 1125
Ala Pro Ala Ser Ser Val Glu Tyr Gln Cys Gln Asn Leu Tyr Gln		
	1130	1135 1140
Leu Glu Gly Asn Lys Arg Ile Thr Cys Arg Asn Gly Gln Trp Ser		
	1145	1150 1155

Glu Pro Pro Lys Cys Leu His Pro Cys Val Ile Ser Arg Glu Ile
1160 1165 1170

Met Glu Asn Tyr Asn Ile Ala Leu Arg Trp Thr Ala Lys Gln Lys
1175 1180 1185

Leu Tyr Ser Arg Thr Gly Glu Ser Val Glu Phe Val Cys Lys Arg
1190 1195 1200

Gly Tyr Arg Leu Ser Ser Arg Ser His Thr Leu Arg Thr Thr Cys
1205 1210 1215

Trp Asp Gly Lys Leu Glu Tyr Pro Thr Cys Ala Lys Arg
1220 1225 1230

<210> 6
<211> 449
<212> PRT
<213> Artificial

<220>
<223> Synthetic Truncated Factor H variant

<400> 6

Met Arg Leu Leu Ala Lys Ile Ile Cys Leu Met Leu Trp Ala Ile Cys
1 5 10 15

Val Ala Glu Asp Cys Asn Glu Leu Pro Pro Arg Arg Asn Thr Glu Ile
20 25 30

Leu Thr Gly Ser Trp Ser Asp Gln Thr Tyr Pro Glu Gly Thr Gln Ala
35 40 45

Ile Tyr Lys Cys Arg Pro Gly Tyr Arg Ser Leu Gly Asn Ile Ile Met
50 55 60

Val Cys Arg Lys Gly Glu Trp Val Ala Leu Asn Pro Leu Arg Lys Cys
65 70 75 80

Gln Lys Arg Pro Cys Gly His Pro Gly Asp Thr Pro Phe Gly Thr Phe
85 90 95

Thr Leu Thr Gly Gly Asn Val Phe Glu Tyr Gly Val Lys Ala Val Tyr
100 105 110

Thr Cys Asn Glu Gly Tyr Gln Leu Leu Gly Glu Ile Asn Tyr Arg Glu

115		120		125
Cys Asp Thr Asp Gly Trp Thr Asn Asp Ile Pro Ile Cys Glu Val Val				
130		135		140
Lys Cys Leu Pro Val Thr Ala Pro Glu Asn Gly Lys Ile Val Ser Ser				
145		150		155
Ala Met Glu Pro Asp Arg Glu Tyr His Phe Gly Gln Ala Val Arg Phe				
	165		170	175
Val Cys Asn Ser Gly Tyr Lys Ile Glu Gly Asp Glu Glu Met His Cys				
	180		185	190
Ser Asp Asp Gly Phe Trp Ser Lys Glu Lys Pro Lys Cys Val Glu Ile				
	195		200	205
Ser Cys Lys Ser Pro Asp Val Ile Asn Gly Ser Pro Ile Ser Gln Lys				
	210		215	220
Ile Ile Tyr Lys Glu Asn Glu Arg Phe Gln Tyr Lys Cys Asn Met Gly				
225		230		235
Tyr Glu Tyr Ser Glu Arg Gly Asp Ala Val Cys Thr Glu Ser Gly Trp				
	245		250	255
Arg Pro Leu Pro Ser Cys Glu Glu Lys Ser Cys Asp Asn Pro Tyr Ile				
	260		265	270
Pro Asn Gly Asp Tyr Ser Pro Leu Arg Ile Lys His Arg Thr Gly Asp				
	275		280	285
Glu Ile Thr Tyr Gln Cys Arg Asn Gly Phe Tyr Pro Ala Thr Arg Gly				
	290		295	300
Asn Thr Ala Lys Cys Thr Ser Thr Gly Trp Ile Pro Ala Pro Arg Cys				
305		310		315
Thr Leu Lys Pro Cys Asp Tyr Pro Asp Ile Lys His Gly Gly Leu Tyr				
	325		330	335
His Glu Asn Met Arg Arg Pro Tyr Phe Pro Val Ala Val Gly Lys Tyr				
	340		345	350
Tyr Ser Tyr Tyr Cys Asp Glu His Phe Glu Thr Pro Ser Gly Ser Tyr				
	355		360	365

Trp Asp His Ile His Cys Thr Gln Asp Gly Trp Ser Pro Ala Val Pro
370 375 380

Cys Leu Arg Lys Cys Tyr Phe Pro Tyr Leu Glu Asn Gly Tyr Asn Gln
385 390 395 400

Asn Tyr Gly Arg Lys Phe Val Gln Gly Lys Ser Ile Asp Val Ala Cys
405 410 415

His Pro Gly Tyr Ala Leu Pro Lys Ala Gln Thr Thr Val Thr Cys Met
420 425 430

Glu Asn Gly Trp Ser Pro Thr Pro Arg Cys Ile Arg Val Ser Phe Thr
435 440 445

Leu

<210> 7
<211> 2823
<212> DNA
<213> Homo sapiens

<400> 7
ggcaggtgct tgttactgtt aatgaaagca gatttaaagc aacaccacca tcaactggagt 60
atTTTTtagtt atatacgatt gagactacca agcatgtttgc tcttattcag tgtaatccta 120
atctcatggg tatccactgt tgggggagaa ggaacacttt gtgattttcc aaaaatacac 180
catggatttc tgtatgatga agaagattat aacccttttt cccaagttcc tacaggggaa 240
gttttctatt actcctgtga atataatttt gtgtctcctt caaaatcctt ttggactcgc 300
ataacatgca cagaagaagg atggtcacca acaccgaagt gtctcagaat gtgttccttt 360
ccttttgtga aaaatgggtca ttctgaatct tcaggactaa tacatctgga aggtgatact 420
gtacaaatta ttgcaacac aggatacagc cttcaaaaaca atgagaaaaa ctttctgtgt 480
gtagaacggg gctgggtccac tcctcccata tgcagcttca ctaaaggaga atgtcatgtt 540
ccaatttttag aagccaatgt agatgctcag ccaaaaaaag aaagctacaa agttggagac 600
gtgttgaaat tctcctgcag aaaaaatctt ataagagttg gatcagactc agttcaatgt 660
taccaatttg ggtggtcacc taactttcca acatgcaaag gacaagtacg atcatgtggt 720
ccacctctc aactctccaa tgggtgaagtt aaggagataa gaaaagagga atatggacac 780
aatgaagtag tggaatatga ttgcaatcct aattttataa taaacgggcc taagaaaata 840

caatgtgtgg atggagaatg gacaacttta cccacttgtg ttgaacaagt gaaaacatgt 900
 ggatacatac ctgaactcga gtacggttat gttcagccgt ctgtccctcc ctatcaacat 960
 ggagtttcag tcgaggtgaa ttgcagaaat gaatatgcaa tgattggaaa taacatgatt 1020
 acctgtatta atggaatatg gacagagctt cctatgtgtg ttgcaacaca ccaacttaag 1080
 aggtgcaaaa tagcaggagt taatataaaa acattactca agctatctgg gaaagaattt 1140
 aatcataatt ctagaatacg ttacagatgt tcagacatct tcagatacag gcactcagtc 1200
 tgtataaacg ggaaatggaa tcctgaagta gactgcacag aaaaaaggga acaattctgc 1260
 ccacgccac ctcagatacc taatgctcag aatatgacaa ccacagtga ttatcaggat 1320
 ggagaaaaag tagctgttct ctgtaaagaa aactatctac ttccagaagc aaaagaaatt 1380
 gtatgtaaag atggacgatg gcaatcatta ccacgctgtg ttgagtctac tgcataattgt 1440
 gggccccctc catctattaa caatggagat accacctcat tcccattatc agtatatcct 1500
 ccagggtcaa cagtgcagta ccgttgccag tccttctata aactccaggg ctctgtaact 1560
 gtaacatgca gaaataaaca gtggtcagaa ccaccaagat gcctagatcc atgtgtggta 1620
 tctgaagaaa acatgaacaa aaataacata cagttaaaat ggagaaacga tggaaaactc 1680
 tatgcaaaaa caggggatgc tgttgaattc cagtgtaaat tcccacataa agcgatgata 1740

 tcatcaccac catttcgagc aatctgtcag gaagggaaat ttgaatatcc tatatgtgaa 1800
 tgaagcaagc ataattttcc tgaatatatt ctccaacat ccactctacg taaaagtagc 1860
 cattatgtag ccaattctgt agttacttct tttattcttt caggtgttgt ttaactcagt 1920
 tttatttaga actctggatt tttagagctt tagaaatttg taagctgaga gaacaatggt 1980
 tcacttaata ggaggggtgc ttagtcata ttacattggt ataacagagt atcacagact 2040
 ggataacttc taaccaatag tttatttggt tcataaatct aaaagctgag aagtccaaga 2100
 tgggtggggt gcctctggtg agggctctct cgaagcatca taatatgctg gaaggcatca 2160
 caacatggtg gaagggatca cgtggcaaaa gagcatgtac atgggagtga gagaaaaaga 2220
 gagagagaga cagagtggcg ggggccgggg aggagcgcaa actcatcctt tataaagaca 2280
 ccactcctga gataacaatc caatcccatg ataatgacat taatccattc aagaagatag 2340
 agctctcgtg acttaatcac cttctaaaga tctcacctga caacactggt gcattggcag 2400
 ttaagtttcc acgtaaactt tcggggacac attcaaacca caggagaaac tcaaattggt 2460
 cctgggcaaa tcacaacatg gggaatttta ttcataaatg tccacagaaa cagtaaatgt 2520
 tctcgottca gaacttaatt catctaatec ctctgtttg tctcaaatta taggataact 2580
 ttgaaacttt ctgaattaac gttatttaaa aggaaatgta gatgttattt tagtotctat 2640

cttcagggtta ttatcactta aaaacctgcg aaagctgtca acttttgtgg ttgtagcaag 2700
 tattaataaaa tatttataaaa tcctctaatag taagtctagc tacctatcca atactaaata 2760
 ccccttaaag tattaatatgc actatctgct gtaaacggaa aaaaaaaaaa aaaaaaaaaa 2820
 aaa 2823

<210> 8
 <211> 569
 <212> PRT
 <213> Homo sapiens

<400> 8

Met Leu Leu Leu Phe Ser Val Ile Leu Ile Ser Trp Val Ser Thr Val
 1 5 10 15

Gly Gly Glu Gly Thr Leu Cys Asp Phe Pro Lys Ile His His Gly Phe
 20 25 30

Leu Tyr Asp Glu Glu Asp Tyr Asn Pro Phe Ser Gln Val Pro Thr Gly
 35 40 45

Glu Val Phe Tyr Tyr Ser Cys Glu Tyr Asn Phe Val Ser Pro Ser Lys
 50 55 60

Ser Phe Trp Thr Arg Ile Thr Cys Thr Glu Glu Gly Trp Ser Pro Thr
 65 70 75 80

Pro Lys Cys Leu Arg Met Cys Ser Phe Pro Phe Val Lys Asn Gly His
 85 90 95

Ser Glu Ser Ser Gly Leu Ile His Leu Glu Gly Asp Thr Val Gln Ile
 100 105 110

Ile Cys Asn Thr Gly Tyr Ser Leu Gln Asn Asn Glu Lys Asn Ile Ser
 115 120 125

Cys Val Glu Arg Gly Trp Ser Thr Pro Pro Ile Cys Ser Phe Thr Lys
 130 135 140

Gly Glu Cys His Val Pro Ile Leu Glu Ala Asn Val Asp Ala Gln Pro
 145 150 155 160

Lys Lys Glu Ser Tyr Lys Val Gly Asp Val Leu Lys Phe Ser Cys Arg
 165 170 175

Lys Asn Leu Ile Arg Val Gly Ser Asp Ser Val Gln Cys Tyr Gln Phe

180	185	190
Gly Trp Ser Pro Asn Phe Pro Thr Cys Lys Gly Gln Val Arg Ser Cys 195 200 205		
Gly Pro Pro Pro Gln Leu Ser Asn Gly Glu Val Lys Glu Ile Arg Lys 210 215 220		
Glu Glu Tyr Gly His Asn Glu Val Val Glu Tyr Asp Cys Asn Pro Asn 225 230 235 240		
Phe Ile Ile Asn Gly Pro Lys Lys Ile Gln Cys Val Asp Gly Glu Trp 245 250 255		
Thr Thr Leu Pro Thr Cys Val Glu Gln Val Lys Thr Cys Gly Tyr Ile 260 265 270		
Pro Glu Leu Glu Tyr Gly Tyr Val Gln Pro Ser Val Pro Pro Tyr Gln 275 280 285		
His Gly Val Ser Val Glu Val Asn Cys Arg Asn Glu Tyr Ala Met Ile 290 295 300		
Gly Asn Asn Met Ile Thr Cys Ile Asn Gly Ile Trp Thr Glu Leu Pro 305 310 315 320		
Met Cys Val Ala Thr His Gln Leu Lys Arg Cys Lys Ile Ala Gly Val 325 330 335		
Asn Ile Lys Thr Leu Leu Lys Leu Ser Gly Lys Glu Phe Asn His Asn 340 345 350		
Ser Arg Ile Arg Tyr Arg Cys Ser Asp Ile Phe Arg Tyr Arg His Ser 355 360 365		
Val Cys Ile Asn Gly Lys Trp Asn Pro Glu Val Asp Cys Thr Glu Lys 370 375 380		
Arg Glu Gln Phe Cys Pro Pro Pro Pro Gln Ile Pro Asn Ala Gln Asn 385 390 395 400		
Met Thr Thr Thr Val Asn Tyr Gln Asp Gly Glu Lys Val Ala Val Leu 405 410 415		
Cys Lys Glu Asn Tyr Leu Leu Pro Glu Ala Lys Glu Ile Val Cys Lys 420 425 430		

Asp Gly Arg Trp Gln Ser Leu Pro Arg Cys Val Glu Ser Thr Ala Tyr
 435 440 445

Cys Gly Pro Pro Pro Ser Ile Asn Asn Gly Asp Thr Thr Ser Phe Pro
 450 455 460

Leu Ser Val Tyr Pro Pro Gly Ser Thr Val Thr Tyr Arg Cys Gln Ser
 465 470 475 480

Phe Tyr Lys Leu Gln Gly Ser Val Thr Val Thr Cys Arg Asn Lys Gln
 485 490 495

Trp Ser Glu Pro Pro Arg Cys Leu Asp Pro Cys Val Val Ser Glu Glu
 500 505 510

Asn Met Asn Lys Asn Asn Ile Gln Leu Lys Trp Arg Asn Asp Gly Lys
 515 520 525

Leu Tyr Ala Lys Thr Gly Asp Ala Val Glu Phe Gln Cys Lys Phe Pro
 530 535 540

His Lys Ala Met Ile Ser Ser Pro Pro Phe Arg Ala Ile Cys Gln Glu
 545 550 555 560

Gly Lys Phe Glu Tyr Pro Ile Cys Glu
 565

<210> 9
 <211> 41
 <212> DNA
 <213> Artificial

<220>
 <223> Factor H gene polymorphism

<400> 9
 ggggttttct gggatgtaat ratgttcagt gttttgacct t

41

<210> 10
 <211> 41
 <212> DNA
 <213> Artificial

<220>
 <223> Polymorphism

<400> 10
 ttatgaaatc cagaggatat yaccagctgc tgatttgacac a

41

<210> 11
<211> 41
<212> DNA
<213> Artificial

<220>
<223> Polymorphism

<400> 11
agtccaagtt tacacagtao ratagactta cccattgcca a

41

<210> 12
<211> 41
<212> DNA
<213> Artificial

<220>
<223> Factor H gene polymorphism

<400> 12
gatatagatc tottggaat rtaataatgg tatgcaggaa g

41

<210> 13
<211> 42
<212> DNA
<213> Artificial

<220>
<223> Factor H gene polymorphism

<220>
<221> misc_feature
<222> (21)..(22)
<223> Residues 21-22 may be absent

<400> 13
taattcataa cttttttttt ttogtttttag aaaggccctg tg

42

<210> 14
<211> 41
<212> DNA
<213> Artificial

<220>
<223> Factor H gene polymorphism

<400> 14
aaaggaatac atttaggact yatttgaagt tagtgtcaac a

41

<210> 15
<211> 40
<212> DNA
<213> Artificial

<220>

<223> Factor H gene polymorphism

<400> 15

caacccgggg aaatacagcm aaatgcacaa gtactggctg

40

<210> 16

<211> 41

<212> DNA

<213> Artificial

<220>

<223> Factor H gene polymorphism

<400> 16

aaaatggata taatcaaaat yatggaagaa agtttgtaca g

41

<210> 17

<211> 41

<212> DNA

<213> Artificial

<220>

<223> Factor H gene polymorphism

<400> 17

tatgoccttaa aagaaaaagc raaatatcaa tgcaaactag g

41

<210> 18

<211> 41

<212> DNA

<213> Artificial

<220>

<223> Factor H gene polymorphism

<220>

<221> misc_feature

<222> (21)..(21)

<223> Residue 21 is optionally absent

<400> 18

cagcttgagt ggatcaaaga ntgacaaggc ccaatggaac c

41

<210> 19

<211> 41

<212> DNA

<213> Artificial

<220>

<223> Factor H gene polymorphism

<400> 19

acggtaccta tttattagta katctaataca ataaagotit t

41

<210> 20
<211> 41
<212> DNA
<213> Artificial

<220>
<223> Factor H gene polymorphism

<400> 20
aagggaccta ataaaaattca rtgtgttgat ggagagtgga c 41

<210> 21
<211> 41
<212> DNA
<213> Artificial

<220>
<223> Factor H gene polymorphism

<400> 21
ttttttatatt tttattataa mattaattat atttttaata t 41

<210> 22
<211> 41
<212> DNA
<213> Artificial

<220>
<223> Factor H gene polymorphism

<400> 22
ccttgtaaat ctccacctga katttctcat ggtgttgtag c 41

<210> 23
<211> 41
<212> DNA
<213> Artificial

<220>
<223> Factor H gene polymorphism

<400> 23
ggggatatcg tttttcatca ygttctcaca cattgcgaac a 41

<210> 24
<211> 31
<212> DNA
<213> Artificial

<220>
<223> Factor H gene polymorphism

<400> 24
aaatccagag gatatyacca gctgctgatt t 31

<210> 25

<211> 31
<212> DNA
<213> Artificial

<220>

<223> Factor H gene polymorphism

<400> 25
aatgggtaag tctatygtag tgtgtaaact t

31

<210> 26
<211> 31
<212> DNA
<213> Artificial

<220>

<223> Factor H gene polymorphism

<400> 26
tgcataccat tattayattt ccaagagatc t

31

<210> 27
<211> 36
<212> DNA
<213> Artificial

<220>

<223> Factor H gene polymorphism

<220>
<221> misc_feature
<222> (18)..(19)
<223> Residues 18-19 may be optionally absent

<400> 27
acatactaatt tcataacttt ttttttttcg ttttag

36

<210> 28
<211> 31
<212> DNA
<213> Artificial

<220>

<223> Factor H gene polymorphism

<400> 28
aatacattta ggactyattt gaagttagtg t

31

<210> 29
<211> 31
<212> DNA
<213> Artificial

<220>

<223> Factor H gene polymorphism

<400> 29
ccgggggaaat acagcmaa at gcacaagtac t 31

<210> 30
<211> 31
<212> DNA
<213> Artificial

<220>
<223> Factor H gene polymorphism

<400> 30
ggatataatc aaaatyatgg aagaaagttt g 31

<210> 31

<211> 31
<212> DNA
<213> Artificial

<220>
<223> Factor H gene polymorphism

<400> 31
cttaaaagaa aaagcraaat atcaatgcaa a 31

<210> 32
<211> 31
<212> DNA
<213> Artificial

<220>
<223> Factor H gene polymorphism

<400> 32
ctttattgat tagatmtact aataaatagg t 31

<210> 33
<211> 31
<212> DNA
<213> Artificial

<220>
<223> Factor H gene polymorphism

<400> 33
acctaataaa attcartgtg ttgatggaga g 31

<210> 34
<211> 31
<212> DNA
<213> Artificial

<220>
<223> Factor H gene polymorphism

<400> 34
taaattctcca cctgakatatt ctcattggtgt t 31

<210> 35
<211> 20
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 35
acttgttccc ccactcctac 20

<210> 36
<211> 20
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 36
cctcttttctg tatggactac 20

<210> 37
<211> 27
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 37
tgaaatcacg taccagtgtg gaaatgg 27

<210> 38
<211> 22
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 38
caggtatcca gccagtactt gt 22

<210> 39
<211> 45
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 39

ctttatttat ttatcattgt tatggtcott aggaaaatgt tttt

45

<210> 40
<211> 25
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 40
ggcaggcaac gtctatagat ttacc

25

<210> 41
<211> 27
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 41
tcaccatctg ctgttacata tcttagt

27

<210> 42
<211> 33
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 42
tgggtttatt tctgaatctc agtatacata tgc

33

<210> 43
<211> 15
<212> DNA
<213> Artificial

<220>
<223> Probe

<400> 43
aatacagcaa aatgc

15

<210> 44
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Probe

<400> 44
tttcttccat gattttg

17

<210> 45
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Probe

<400> 45
aagaaaaagc gaaatat

17

<210> 46
<211> 14
<212> DNA
<213> Artificial

<220>
<223> Probe

<400> 46
atacagccaa atgc

14

<210> 47
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Probe

<400> 47
ttcttccata attttg

16

<210> 48
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Probe

<400> 48
aagaaaaagc aaaatat

17

<210> 49
<211> 41
<212> DNA
<213> Artificial

<220>
<223> Polymorphism

<400> 49
ccaggctatc tataaatgcc ccctggata tagatctctt g

41

<210> 50
<211> 41
<212> DNA
<213> Artificial

<220>
<223> Polymorphism

<400> 50
ttggtacttt taccottaca kgaggaaatg tgtttgaata t

41

<210> 51
<211> 41
<212> DNA
<213> Artificial

<220>
<223> Polymorphism

<220>
<221> misc_feature
<222> (21)..(21)
<223> Residue 21 is G or H

<400> 51
acgatggttt ttggagtaaa nagaaaccaa agtgtgtggg t

41

<210> 52
<211> 41
<212> DNA
<213> Artificial

<220>
<223> Polymorphism

<220>
<221> misc_feature
<222> (21)..(21)
<223> Residue is C or D

<400> 52
ttatttataa ggagaatgaa ngatttcaat ataaatgtaa c

41

<210> 53
<211> 38
<212> DNA
<213> Artificial

<220>
<223> Polymorphism

<220>
<221> misc_feature
<222> (21)..(21)

<223> Residue is C or D

<400> 53

cactgaatct ggatggcgtc ngttgccttc atgtgaag

38

<210> 54

<211> 33

<212> DNA

<213> Artificial

<220>

<223> Polymorphism

<220>

<221> misc feature

<222> (21)..(21)

<223> Residue 21 is optionally absent

<400> 54

aagatggatg gtcgccagca staccatgcc tca

33

<210> 55

<211> 41

<212> DNA

<213> Artificial

<220>

<223> Polymorphism

<400> 55

acaattatgc ccacotccac stcagattcc caattctcac a

41

<210> 56

<211> 41

<212> DNA

<213> Artificial

<220>

<223> Polymorphism

<400> 56

caaccacctc agatagaaca yggaaccatt aattcatcca g

41

<210> 57

<211> 41

<212> DNA

<213> Artificial

<220>

<223> Polymorphism

<400> 57

gtcttcacaa gaaagttatg yacatgggac taaattgagt t

41

<210> 58

<211> 41
<212> DNA
<213> Artificial

<220>
<223> Polymorphism

<400> 58
cacatgtcag acagttatca ktatggagaa gaagttacgt a

41

<210> 59
<211> 41
<212> DNA
<213> Artificial

<220>
<223> Polymorphism

<400> 59
tcagtatgga gaagaagtta ygtacaaatg ttttgaaggt t

41

<210> 60
<211> 27
<212> DNA
<213> Artificial

<220>
<223> Polymorphism

<400> 60
gtatggkgca ttgaatttta ttatatg

27

<210> 61
<211> 35
<212> DNA
<213> Artificial

<220>
<223> Polymorphism

<400> 61
acacctcctg tgtgwatccg cccacagtac aaaat

35

<210> 62
<211> 44
<212> DNA
<213> Artificial

<220>
<223> Polymorphism

<220>
<221> misc_feature
<222> (21)..(21)
<223> Residue 21 may be optionally absent

<220>
<221> misc_feature
<222> (23)..(23)
<223> Residue 23 may be optionally absent

<400> 62
cttgatatcaa cttgagggtg nancaagcga ataacatgta gaaa 44

<210> 63
<211> 41
<212> DNA
<213> Artificial

<220>
<223> Polymorphism

<400> 63
aaaagcttta ttgattagat mtactaataa ataggtagcg t 41

<210> 64
<211> 19
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 64
gcaaaaagttt ctgataggc 19

<210> 65
<211> 20

<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 65
aatcttacct tctgctacac 20

<210> 66
<211> 19
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 66
ttagatagac ctgtgactg 19

<210> 67
<211> 17
<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 67

tcaggcataa ttgctac

17

<210> 68

<211> 16

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 68

acttggtccc ccactc

16

<210> 69

<211> 20

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 69

cctottttcg tatggactac

20

<210> 70

<211> 18

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 70

ttggtccccc actcctac

18

<210> 71

<211> 20

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 71

acacatttcc tcctgtaagg

20

<210> 72

<211> 17

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 72

ccctgtggac atcctgg

17

<210> 73

<211> 22

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 73

aacctctttt cgtatggact ac

22

<210> 74

<211> 16

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 74

atgctgttca ttttcc

16

<210> 75

<211> 17

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 75

ccatccatct gtgtcac

17

<210> 76

<211> 17

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 76

attaccgtga atgtgac

17

<210> 77

<211> 19

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 77

ttgtatgaga aaaaaaac

19

<210> 78

<211> 18

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 78

tccaatctta tcttgagg

18

<210> 79

<211> 18

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 79

tcttaccac acaatttg

18

<210> 80

<211> 16

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 80

gtcctggtea cagtec

16

<210> 81

<211> 17

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 81

gcatacagca tctcctc

17

<210> 82

<211> 16

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 82

gcactgaatc tggatg

16

<210> 83

<211> 17

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 83

atgaaccttg aacacag

17

<210> 84

<211> 17

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 84

cggatactta tttctgc

17

<210> 85

<211> 17

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 85

cgtgatttca tctccag

17

<210> 86

<211> 18

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 86

agaactggag atgaaatc

18

<210> 87

<211> 17

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 87
tgaatggaac ttacagg

17

<210> 88
<211> 19
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 88
gtgaaacctt gtgattatc

19

<210> 89
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 89
tcccagtaac ttcctg

16

<210> 90
<211> 19
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 90
ctgtgatgaa cattttgag

19

<210> 91
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 91
tgctctcctt tcttcg

16

<210> 92
<211> 19
<212> DNA
<213> Artificial

<220>

<223> Primer

<400> 92
cattggttatg gtccttagg

19

<210> 93
<211> 19
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 93
acatgctagg atttcagag

19

<210> 94
<211> 20
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 94
ctttttctta ttctcttccc

20

<210> 95
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 95
tcaccatctg ctgttac

17

<210> 96
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 96
tgtaacagca gatgggtg

17

<210> 97
<211> 18

<212> DNA
<213> Artificial

<220>

<223> Primer

<400> 97

cccacaaaaa gactaaag

18

<210> 98

<211> 17

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 98

gggaaatact cagattg

17

<210> 99

<211> 16

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 99

atggcattca tagtcc

16

<210> 100

<211> 20

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 100

ccagaactaa aaatgacttc

20

<210> 101

<211> 17

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 101

ggtaaatacag accaacc

17

<210> 102

<211> 19

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 102
atagtgtgtg gttacaatg 19

<210> 103
<211> 19
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 103
gtttatgtca aatcaggag 19

<210> 104
<211> 19
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 104
caagaaagag aatgcgaac 19

<210> 105
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 105
agattacagg caatggg 17

<210> 106
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 106
ttgattgttt aggatgc 17

<210> 107
<211> 20
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 107
ttgaggagtt caggaggtgg 20

<210> 108
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 108
ctgaactcct caatgg 16

<210> 109
<211> 18
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 109
attaccaata cacactgg 18

<210> 110
<211> 18
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 110
ttacatagtg gaggagag 18

<210> 111
<211> 15
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 111
tggaaatgtt gaggc 15

<210> 112
<211> 19
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 112

agttggtttg attcctatc

19

<210> 113
<211> 18
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 113
ttgagcagtt cacttctg

18

<210> 114
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 114
ttatgccac ctccac

16

<210> 115
<211> 19
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 115
atacactact gaccaacac

19

<210> 116
<211> 19
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 116
gtctatgaga atacaagcc

19

<210> 117
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 117
gaatctgagg tggagg

16

<210> 118
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 118
ccctttgatt ttcattc

17

<210> 119
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 119
agaactccat tttccc

16

<210> 120
<211> 18
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 120
cacaaccacc tcgatag

18

<210> 121
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 121
gcctaacctt cacactg

17

<210> 122
<211> 21
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 122
gtcatagtag ctccgtgtatt g

21

<210> 123
<211> 20
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 123
acgtaacttc ttctccatac

20

<210> 124
<211> 19
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 124
cttccttgta aatctccac

19

<210> 125
<211> 19
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 125
caatgcacca tacttatgc

19

<210> 126
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 126
taaagatttg cggaac

16

<210> 127
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 127
ggctccatcc attttg

16

<210> 128
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 128
ttacaaaatg gatggag

17

<210> 129
<211> 20
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 129
aagtgctggg attacaggcg

20

<210> 130
<211> 21
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 130
ctactcaaaa tgaacactag g

21

<210> 131
<211> 19
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 131
tttaaccctg ctatactcc

19

<210> 132
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 132
taaagtggaaa ctggacg

17

<210> 133

<211> 20
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 133
accctattac ttgtgttctg

20

<210> 134
<211> 15
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 134
gtgttttgcgt ttgcc

15

<210> 135
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 135
gagatttttcc cagccac

17

<210> 136
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 136
tctcacacat tgcgaac

17

<210> 137
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 137
accgtaggtt ttccagg

17

<210> 138

<211> 20
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 138
ggtttgata gtgttttgag

20

<210> 139
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 139
atgttgttcg caatgtg

17

<210> 140
<211> 20
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 140
tgggagtga gtagaattg

20

<210> 141
<211> 22
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 141
gctaattgatg cttttcacag ga

22

<210> 142
<211> 22
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 142
cctgtgactg tctaggcatt tt

22

<210> 143

<211> 26
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 143
tatgcctgaa ttatatcact attgcc

26

<210> 144
<211> 24
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 144
gctttgotat gtttaatttt cctt

24

<210> 145
<211> 27
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 145
aactatgatg gaaataatta aatctgg

27

<210> 146
<211> 20
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 146
tgcatatgct gttcattttc

20

<210> 147
<211> 28
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 147
gtcttacatt aaaatatctt aaagtctc

28

<210> 148
<211> 22

<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 148
tttcctccaa tcttatcctg ag

22

<210> 149
<211> 24
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 149
cgttcattct aaggaatata agca

24

<210> 150
<211> 22
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 150
cctgatggaa acaacatttc tg

22

<210> 151
<211> 20
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 151
aacagggcca gaaaagttca

20

<210> 152
<211> 22
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 152
tgttcatttt aatgccattt tg

22

<210> 153
<211> 20
<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 153

agtttttcgaa gttgccgaaa

20

<210> 154

<211> 23

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 154

octagaaacc ctaatggaat gtg

23

<210> 155

<211> 21

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 155

tggtcaagca aagtgaccaa a

21

<210> 156

<211> 24

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 156

tgagcaaatt tatgtttctc attt

24

<210> 157

<211> 23

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 157

atgtcacctt gttttaccaa tgg

23

<210> 158

<211> 23

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 158

tgaatgctta tggttatcca ggt

23

<210> 159

<211> 20

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 159

aaaacctgca ggaacaaagc

20

<210> 160

<211> 26

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 160

tccttagaatg ggaaatactc agattg

26

<210> 161

<211> 24

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 161

tggtttttca gaaattcatt ttca

24

<210> 162

<211> 24

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 162

atgtaaaatt aactttggca atga

24

<210> 163

<211> 26

<212> DNA

<213> Artificial

<220>
<223> Primer

<400> 163
ttgctgaaat aagaattaga actttg 26

<210> 164
<211> 25
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 164
tgaataaaag aagaaaatct ttcca 25

<210> 165
<211> 27
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 165
atctaaaaca catacatcat gttttca 27

<210> 166
<211> 26
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 166
aaaacacata catcatgttt tcacaa 26

<210> 167
<211> 23
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 167
gatatgctc aacattttcca gtc 23

<210> 168
<211> 23
<212> DNA
<213> Artificial

<220>

<223> Primer

<400> 168

gttggtttga ttctatcat ttg

23

<210> 169

<211> 26

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 169

ttggaaaagt aataggtatg tgtgtc

26

<210> 170

<211> 25

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 170

ctatgagaat acaagccaaa agttc

25

<210> 171

<211> 23

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 171

tctctgtgc ttctgtgaaa caa

23

<210> 172

<211> 23

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 172

aacctttga tttcattct tca

23

<210> 173

<211> 22

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 173

tcaaagtgag gggaataatt ga

22

<210> 174

<211> 26

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 174

aatttatgag ttagtgaaac ctgaat

26

<210> 175

<211> 26

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 175

tcttcattca aagtgttaagt ggtacc

26

<210> 176

<211> 27

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 176

acaaaatggc taatatatatt tctcaag

27

<210> 177

<211> 18

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 177

taatgtgtgg gccagcc

18

<210> 178

<211> 23

<212> DNA

<213> Artificial

<220>
<223> Primer

<400> 178
caaaatgaac actaggtgga acc

23

<210> 179
<211> 21
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 179
atthttggggg agtatagcag g

21

<210> 180
<211> 20
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 180
ctgtgtttgc gtttgcctta

20

<210> 181
<211> 20
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 181
ttcacgtggc tggaaaaatc

20

<210> 182
<211> 25
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 182
ttgaaaacct gaaagtctat gaaga

25

<210> 183
<211> 23
<212> DNA
<213> Artificial

<220>

<223> Primer

<400> 183

tcaatcataa agtgcacacc ttt

23

<210> 184

<211> 23

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 184

cagtccatt tctgattgtt cca

23

<210> 185

<211> 21

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 185

gctgaggata atttgaagg g

21

<210> 186

<211> 23

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 186

gtgattcatc gatgtagctc ttt

23

<210> 187

<211> 21

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 187

aatgaccaga ggagcctgga a

21

<210> 188

<211> 24

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 188

tgatgtcagt tttcaaagtt ttcc

24

<210> 189

<211> 26

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 189

accactctct cagttttgct aattat

26

<210> 190

<211> 25

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 190

cacattaaat ttgtttctgc aatga

25

<210> 191

<211> 24

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 191

agaagtgatg aaacaagaat ttga

24

<210> 192

<211> 25

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 192

ccattttaagc attatttatg gtttc

25

<210> 193

<211> 27

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 193

aaacaggaca gttactatta ctttgca

27

<210> 194

<211> 27

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 194

aaatatatttc agagtaagca ctcattt

27

<210> 195

<211> 23

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 195

tttatcattt tgattgggat tgt

23

<210> 196

<211> 27

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 196

tcgagatatt ttattgacat aattggt

27

<210> 197

<211> 26

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 197

gttgatcttg ttgcttcttt acaaga

26

<210> 198

<211> 22

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 198

ccatttttcct gaaacactac cc

22

<210> 199

<211> 25

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 199

aattatttga atttccagac acctt

25

<210> 200

<211> 25

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 200

aattatttga atttccagac acctt

25

<210> 201

<211> 27

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 201

ttttggacta atttcataga ataaccc

27

<210> 202

<211> 27

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 202

cttaaatgca atttcactat tctatga

27

<210> 203

<211> 16

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 203
tagccattat gtagcc

16

<210> 204
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Probe

<400> 204
tctgggatgt aataatg

17

<210> 205
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Probe

<400> 205
tctgggatgt aatgatg

17

<210> 206
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Probe

<400> 206
gaacattatt acatccc

17

<210> 207
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Probe

<400> 207
gaacatcatt acatccc

17

<210> 208
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Probe

<400> 208
cagaggatat caccagc 17

<210> 209
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Probe

<400> 209
cagaggatat taccagc 17

<210> 210
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Probe

<400> 210
agcagctggt gatatcc 17

<210> 211
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Probe

<400> 211
agcagctggt aatatcc 17

<210> 212
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Probe

<400> 212
tacacagtac gatagac 17

<210> 213
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Probe

<400> 213
tacacagtac aatagac

17

<210> 214
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Probe

<400> 214
taagtctatc gtactgt

17

<210> 215
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Probe

<400> 215
taagtctatt gtactgt

17

<210> 216
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Probe

<400> 216
tcttggaat gtaataa

17

<210> 217
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Probe

<400> 217
tcttggaat ataataa

17

<210> 218
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Probe

<400> 218

accattatta catttcc

17

<210> 219
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Probe

<400> 219
accattatta tatttcc

17

<210> 220
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Probe

<400> 220
tttttttttct gtttttag

17

<210> 221
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Probe

<400> 221
ttttttttttt tcgtttt

17

<210> 222
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Probe

<400> 222
ctttctaaaa cgaaaaa

17

<210> 223
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Probe

<400> 223
ttctaaaacg aaaaaaa

17

<210> 224
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Probe

<400> 224
tttaggactc atttgaa

17

<210> 225
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Probe

<400> 225
tttaggactt atttgaa

17

<210> 226
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Probe

<400> 226
taacttcaaa tgagtcc

17

<210> 227
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Probe

<400> 227
taacttcaaa taagtcc

17

<210> 228
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Probe

<400> 228
gaaatacagc aaaatgc

17

<210> 229
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Probe

<400> 229
gaaatacagc caaatgc

17

<210> 230
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Probe

<400> 230
acttgtgcat ttgtctg

17

<210> 231
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Probe

<400> 231
acttgtgcat ttggctg

17

<210> 232
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Probe

<400> 232
taatcaaaat tatggaa

17

<210> 233
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Probe

<400> 233
taatcaaaat catggaa

17

<210> 234
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Probe

<400> 234
tttcttccat aatcttg

17

<210> 235
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Probe

<400> 235
tttcttccat gatcttg

17

<210> 236
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Probe

<400> 236
aagaaaaagc gaaatat

17

<210> 237
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Probe

<400> 237
aagaaaaagc aaaatat

17

<210> 238
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Probe

<400> 238
ttgatatttc gcttttt

17

<210> 239
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Probe

<400> 239
ttgatatttt gcttttt

17

<210> 240
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Probe

<400> 240
ggatcaaaga tgacaa

16

<210> 241
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Probe

<220>
<221> misc_feature
<222> (11)..(11)
<223> n is a, c, g, or t

<400> 241
ggatcaaaga ntgacaa

17

<210> 242
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Probe

<400> 242
gcccttggtca tctttg

16

<210> 243
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Probe

<220>
<221> misc_feature
<222> (11)..(11)
<223> n is a, c, g, or t

<400> 243
gcccttgatca ntctttg

17

<210> 244
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Probe

<400> 244
tttattagta gatctaa

17

<210> 245
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Probe

<400> 245
tttattagta tatctaa

17

<210> 246
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Probe

<400> 246
ttgattagat ctactaa

17

<210> 247
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Probe

<400> 247
ttgattagat atactaa

17

<210> 248
<211> 17

<212> DNA
<213> Artificial

<220>
<223> Probe

<400> 248
ataaaaattca atgtggtt

17

<210> 249
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Probe

<400> 249
ataaaaattca gtgtggtt

17

<210> 250
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Probe

<400> 250
ccatcaacac attgaat

17

<210> 251
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Probe

<400> 251
ccatcaacac actgaat

17

<210> 252
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Probe

<400> 252
tttattataa cattaat

17

<210> 253
<211> 17
<212> DNA

<213> Artificial

<220>

<223> Probe

<400> 253

tttattataa aattaat

17

<210> 254

<211> 17

<212> DNA

<213> Artificial

<220>

<223> Probe

<400> 254

tataattaat gttataa

17

<210> 255

<211> 17

<212> DNA

<213> Artificial

<220>

<223> Probe

<400> 255

tataattaat tttataa

17

<210> 256

<211> 17

<212> DNA

<213> Artificial

<220>

<223> Probe

<400> 256

ctccacctga gatttct

17

<210> 257

<211> 17

<212> DNA

<213> Artificial

<220>

<223> Probe

<400> 257

ctccacctga tattttct

17

<210> 258

<211> 17

<212> DNA

<213> Artificial

<220>
<223> Probe

<400> 258
catgagaaat ctcaggt

17

<210> 259
<211> 17
<212> DNA

<213> Artificial

<220>
<223> Probe

<400> 259
catgagaaat atcaggt

17

<210> 260
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Probe

<400> 260
tctttcatca cgttctc

17

<210> 261
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Probe

<400> 261
tctttcatca tgttctc

17

<210> 262
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Probe

<400> 262
gtgtgagaac gtgatga

17

<210> 263
<211> 17
<212> DNA
<213> Artificial

<220>
<223> Probe

<400> 263
gtgtgagaac atgatga

17

<210> 264
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 264
tttctgggat gtaata

16

<210> 265
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 265
tttctgggat gtaatg

16

<210> 266
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 266
caaaacactg aacatt

16

<210> 267
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 267
caaaacactg aacatc

16

<210> 268
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 268
aaatccagag gatatc

16

<210> 269
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 269
aaatccagag gatatt

16

<210> 270
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 270
aaatcagcag ctggtg

16

<210> 271
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 271
aaatcagcag ctggta

16

<210> 272
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 272
aagtttacac agtacg

16

<210> 273
<211> 16
<212> DNA
<213> Artificial

<220>

<223> Primer

<400> 273
aagtttacac agtaca

16

<210> 274
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 274
aatgggtaag tctatc

16

<210> 275
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 275
aatgggtaag tctatt

16

<210> 276
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 276
agatctcttg gaaatg

16

<210> 277
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 277
agatctcttg gaaata

16

<210> 278
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 278
tgcataccat tattac

16

<210> 279
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 279
tgcataccat tattat

16

<210> 280
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 280
tcataacttt tttttt

16

<210> 281
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 281
ataacttttt tttttt

16

<210> 282
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 282
gggcctttct aaaacg

16

<210> 283
<211> 16
<212> DNA
<213> Artificial

<220>

<223> Primer

<400> 283
gccttttctaa aacgaa

16

<210> 284
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 284
aatacattta ggactc

16

<210> 285
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 285
aatacattta ggactt

16

<210> 286
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 286
acactaactt caaatg

16

<210> 287
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 287
acactaactt caaata

16

<210> 288
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 288
ccggggaaat acagca 16

<210> 289
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 289
ccggggaaat acagcc 16

<210> 290
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 290
agtacttgtg catttt 16

<210> 291
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 291
agtacttgtg catttg 16

<210> 292
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 292
ggatataatc aaaatt 16

<210> 293
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 293
ggatataatc aaaatc

16

<210> 294
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 294
caaactttct tccata

16

<210> 295
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 295
caaactttct tccatg

16

<210> 296
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 296
cttaaaagaa aaagcg

16

<210> 297
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 297
cttaaaagaa aaagca

16

<210> 298
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 298

tttgcattga tatttc

16

<210> 299
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 299
tttgcattga tatttt

16

<210> 300
<211> 15
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 300
tgagtggatc aaaga

15

<210> 301
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<220>
<221> misc_feature
<222> (16)..(16)
<223> n is a, c, g, or t

<400> 301
tgagtggatc aaagan

16

<210> 302
<211> 15
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 302
cattggccct tgtca

15

<210> 303
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<220>
<221> misc_feature
<222> (16)..(16)
<223> n is a, c, g, or t

<400> 303
cattggccct tgtcan 16

<210> 304
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 304
acctatttat tagtag 16

<210> 305
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 305
acctatttat tagtat 16

<210> 306
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 306
ctttattgat tagatc 16

<210> 307
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 307
ctttattgat tagata 16

<210> 308
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 308
acctaataaa attcaa

16

<210> 309
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 309
acctaataaa attcag

16

<210> 310
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 310
ctctccatca acacat

16

<210> 311
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 311
ctctccatca acacac

16

<210> 312
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 312
tatttttttat tataac

16

<210> 313
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 313
tatttttttat tataaa

16

<210> 314
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 314
aaaaatataa ttaatg

16

<210> 315
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 315
aaaaatataa ttaatt

16

<210> 316
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 316
taaattctcca cctgag

16

<210> 317
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 317
taaattctcca cctgat

16

<210> 318
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 318
aacacccatga gaaatc

16

<210> 319
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 319
aacacccatga gaaata

16

<210> 320
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 320
tatcgtcttt catcac

16

<210> 321
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 321
tatcgtcttt catcat

16

<210> 322
<211> 16
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 322
gcaatgtgtg agaacg

16

<210> 323

<211> 16

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 323

gcaatgtgtg agaacg

16

<210> 324

<211> 20

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 324

gaacattttg agactcgcgc

20

<210> 325

<211> 18

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 325

accatccatc tttcccac

18

<210> 326

<211> 17

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 326

tcctggctac gctcttc

17

<210> 327

<211> 19

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 327

tcogtcagga agttactgg

19

<210> 328
<211> 20
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 328
agtcaccata ctcaggaccc

20

<210> 329
<211> 19
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 329
ggctacgctc ttccaaaag

19

<210> 330
<211> 20
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 330
agtcaccata ctcaggaccc

20

<210> 331
<211> 26
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 331
gaagattgca atgaacttcc tccaag

26

<210> 332
<211> 21
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 332
aagttctgaa taaaggtgtg c

21

<210> 333

<211> 36
 <212> DNA
 <213> Artificial

<220>
 <223> Primer

<400> 333
 tatagatctc ttggaaatat aataatggta tgcagg 36

<210> 334
 <211> 36
 <212> DNA
 <213> Artificial

<220>
 <223> Primer

<400> 334
 atggatataa tcaaaattat ggaagaaagt ttgtac 36

<210> 335
 <211> 3769
 <212> DNA
 <213> Artificial

<220>
 <223> Synthetic human CFH variant

<400> 335
 agttagctgg taaatgtcct cttaaaagat ccaaaaaatg agacttctag caaagattat 60
 ttgccttatg ttatgggcta tttgtgtagc agaagattgc aatgaacttc ctccaagaag 120
 aaatacagaa attctgacag gttcctggtc tgaccaaaaca tatccagaag gcacccaggc 180
 tatctataaa tgccgccttg gatatagatc tcttggaaat ataataatgg tatgcaggaa 240
 gggagaatgg gttgctctta atccattaag gaaatgtcag aaaaggccct gtggacatcc 300
 tggagatact ccttttggta cttttaccct tacaggagga aatgtgtttg aatatggtgt 360
 aaaagctgtg tatacatgta atgaggggta tcaattgcta ggtgagatta attaccgtga 420
 atgtgacaca gatggatgga ccaatgatat tcttatatgt gaagttgtga agtgtttacc 480
 agtgacagca ccagagaatg gaaaaattgt cagtagtgca atggaaccag atcggaata 540
 ccattttgga caagcagtac ggtttgtatg taactcaggc tacaagattg aaggagatga 600
 agaaatgcat tgttcagacg atggtttttg gagtaaagag aaaccaaagt gtgtggaaat 660
 ttcattgcaaa tcccagatg ttataaatgg atctcctata tctcagaaga ttattttataa 720
 ggagaatgaa cgatttcaat ataaatgtaa catgggttat gaatacagtg aaagaggaga 780
 tgctgtatgc actgaatctg gatggcgtcc gttgccttca tgtgaagaaa aatcatgtga 840
 taatccttat attccaaatg gtgactactc acctttaagg attaaacaca gaactggaga 900

tgaaatcacg taccagtgtg gaaatgggtt ttatcctgca acccggggaa atacagcaaa 960
 atgcacaagt actggctgga tacctgctcc gagatgtacc ttgaaacctt gtgattatcc 1020
 agacattaaa catggagggtc tatatcatga gaatatgcgt agaccatact ttccagtagc 1080
 tgtaggaaaa tattactcct attactgtga tgaacatttt gagactccgt caggaagtta 1140
 ctgggatcac attcattgca cacaagatgg atggctcgcca gcagtaccat gcctcagaaa 1200
 atgttatttt ccttatgttg aaaatggata taatcaaaat tatggaagaa agtttgtaca 1260
 gggtaaatct atagacgttg cctgccatcc tggctacgct cttccaaaag cgcagaccac 1320
 agttacatgt atggagaatg gctgggtctcc tactccocaga tgcattccgtg tcaaaacatg 1380
 ttccaaatca agtatagata ttgagaatgg gtttatctct gaattctcagt atacatatgc 1440
 cttaaaagaa aaagcgaaat atcaatgcaa actaggatat gtaacagcag atggtgaaac 1500
 atcaggatca attacatgtg ggaaagatgg atggctcagct caaccacgt gcattaaatc 1560
 ttgtgatatc ccagtattta tgaatgccag aactaaaaat gacttcacat ggtttaagct 1620
 gaatgacaca ttggaactatg aatgccatga tgggttatgaa agcaatactg gaagcaccac 1680
 tggttccata gtgtgtgggtt acaatgggtg gtctgattta cccatatgtt atgaaagaga 1740
 atgcgaactt cctaaaatag atgtacactt agttcctgat cgcaagaaag accagtataa 1800
 agttggagag gtgttgaaat tctcctgcaa accaggattt acaatagttg gacctaatc 1860
 cgttcagtgc taccactttg gattgtctcc tgacctcca atatgtaaag agcaagtaca 1920
 atcatgtggt ccacctcctg aactcctcaa tgggaatgtt aaggaaaaaa cgaaagaaga 1980
 atatggacac agtgaagtgg tggaatatta ttgcaatcct agatttctaa tgaagggacc 2040
 taataaaatt caatgtgttg atggagagtg gacaacttta ccagtgtgta ttgtggagga 2100
 gagtacctgt ggagatatac ctgaacttga acatggctgg gccagcttt cttccctcc 2160
 ttattactat ggagattcag tggaattcaa ttgctcagaa tcatttaca tgattggaca 2220
 cagatcaatt acgtgtattc atggagtatg gacccaactt cccagtgtg tggcaataga 2280
 taaacttaag aagtgc aaat catcaaat aattatactt gaggaacatt taaaaaaca 2340
 gaaggaattc gatcataatt ctaacataag gtacagatgt agaggaaaag aaggatggat 2400
 acacacagtc tgcataaatg gaagatggga tccagaagtg aactgctcaa tggcacaat 2460
 acaattatgc ccacctccac ctccagattcc caattctcac aatatgacaa ccacactgaa 2520
 ttatcgggat ggagaaaaag tatctgttct ttgccaagaa aattatctaa ttcaggaagg 2580
 agaagaaatt acatgcaaag atggaagatg gcagtcaata ccactctgtg ttgaaaaaat 2640
 tccatgttca caaccacctc agatagaaca cggaaccatt aattcatcca ggtcttcaca 2700

agaaagttat gcacatggga ctaaattgag ttatacttgt gaggggtggtt tcaggatata 2760
 tgaagaaaat gaaacaacat gctacatggg aaaatggagt tctccacctc agtgtgaagg 2820
 ccttccttgt aaatctccac ctgagatttc tcatggtgtt gtagctcaca tgtcagacag 2880
 ttatcagtat ggagaagaag ttacgtacaa atgttttgaa ggttttggaa ttgatgggoc 2940
 tgcaattgca aaatgcttag gagaaaaatg gtctcaccct ccatcatgca taaaaacaga 3000
 ttgtctcagt ttacctagct ttgaaaatgc catacccatg ggagagaaga aggatgtgta 3060
 taaggcgggt gagcaagtga cttacacttg tgcaacatat taaaaatgg atggagocag 3120
 taatgtaaca tgcattaata gcagatggac aggaaggcca acatgcagag acacctcctg 3180
 tgtgaatccg cccacagtac aaaatgctta tatagtgtcg agacagatga gtaaatatcc 3240
 atctggtgag agagtacgtt atcaatgtag gagcccttat gaaatgtttg gggatgaaga 3300
 agtgatgtgt ttaaatggaa actggacgga accacctcaa tgcaaagatt ctacaggaaa 3360
 atgtggggcc cctccacctt ttgacaatgg ggacattact tcattcccggt tgtcagtata 3420
 tgctccagct tcatcagttg agtatcaatg ccagaacttg tatcaacttg agggtaacaa 3480
 gcgaataaca tgtagaaatg gacaatgggc agaaccacca aaatgcttac atccgtgtgt 3540
 aatatcccg gaaattatgg aaaattataa catagcatta aggtggacag ccaaacagaa 3600
 gctttattcg agaacaggtg aatcagttga atttgtgtgt aaacggggat atcgtctttc 3660
 atcacgttct cacacattgc gaacaacatg ttgggatggg aaactggagt atccaacttg 3720
 tgcaaaaaga tagaatcaat cataaagtgc acacctttat tcagaactt 3769

<210> 336
 <211> 3926
 <212> DNA
 <213> Homo sapiens

<400> 336
 aattcttggg agaggagaac tggacgttgt gaacagagtt agctgggtaaa tgcctcttta 60
 aaagatccaa aaaatgagac ttctagcaaa gattatttgc cttatgttat gggctatttg 120
 tgtagcagaa gattgcaatg aacttcctcc aagaagaaat acagaaattc tgacaggttc 180
 ctggtctgac caaacatata cagaaggcac ccaggctata tataaatgcc gccctggata 240
 tagatctctt ggaaatgtaa taatggtatg caggaaggga gaatgggttg ctcttaatcc 300
 attaaggaaa tgtcagaaaa ggccctgtgg acatcctgga gatactcctt ttggtacttt 360
 tacccttaca ggaggaaatg tgtttgaata tgggtgtaaaa gctgtgtata catgtaatga 420
 ggggtatcaa ttgctaggtg agattaatta ccgtgaatgt gacacagatg gatggaccaa 480
 tgatattcct atatgtgaag ttgtgaagtg tttaccagtg acagcaccag agaatggaaa 540

aattgtcagt agtgcaatgg aaccagatcg ggaataccat tttggacaag cagtacgggtt 600
tgtatgtaac tcaggctaca agattgaagg agatgaagaa atgcattgtt cagacgatgg 660
tttttgaggt aaagagaaac caaagtgtgt ggaaatttca tgcaaatccc cagatgttat 720
aaatggatct cctatatctc agaagattat ttataaggag aatgaacgat ttcaatataa 780
atgtaacatg ggttatgaat acagtgaag aggagatgct gtatgcactg aatctggatg 840
gcgtccgttg ccttcattgt aagaaaaatc atgtgataat ccttatattc caaatgggtga 900
ctactcacct ttaaggatta aacacagaac tggagatgaa atcacgtacc agtgtagaaa 960
tggtttttat cctgcaaccc ggggaaatac agccaaatgc acaagtactg gctggatacc 1020
tgctccgaga tgtacottga aacctgtgta ttatccagac attaaacatg gaggtctata 1080
tcatgagaat atgcgtagac catactttcc agtagctgta ggaaaatatt actcctatta 1140
ctgtgatgaa cattttgaga ctccgtcagg aagttactgg gatcacattc attgcacaca 1200
agatggatgg tcgccagcag taccatgcct cagaaaatgt tattttcctt atttggaaaa 1260
tggatataat caaaattatg gaagaaagtt tgtacagggg aaatctatag acgttgctg 1320
ccatcctggc tacgctcttc caaaagcgca gaccacagtt acatgtatgg agaatggctg 1380
gtctcctact ccagatgca tccgtgtcaa aacatgttcc aaatcaagta tagatattga 1440
gaatgggttt atttctgaat ctcagtatac atatgcctta aaagaaaaag cgaaatatca 1500
atgcaaaacta ggatattgaa cagcagatgg tgaaacatca ggatcaatta gatgtgggaa 1560
agatggatgg tcagctcaac ccacgtgoat taaatcttgt gatatoccag tatttatgaa 1620
tgccagaact aaaaatgact tcacatgggt taagctgaat gacacattgg actatgaatg 1680
ccatgatggg tatgaaagca atactggaag caccactggg tccatagtgt gtgggttacia 1740
tggttgggtc gatttaccoc tatgttatga aagagaatgc gaacttcta aaatagatgt 1800
acacttagtt cctgatcgca agaaagacca gtataaagtt ggagaggtgt tgaaattctc 1860
ctgcaaacca ggatttacia tagttggacc taattccgtt cagtgtacc actttggatt 1920
gtctcctgac ctcccaatat gtaaagagca agtacaatca tgtgggtccac ctctgaact 1980
cctcaatggg aatgttaagg aaaaaacgaa agaagaatat ggacacagtg aagtgggtga 2040
atattattgc aatcctagat ttctaataa gggacctaat aaaattcaat gtgttgatgg 2100
agagtggaca actttaccag tgtgtattgt ggaggagagt acctgtggag atatacctga 2160
acttgaacat ggetgggccc agctttcttc ccctccttat tactatggag attcagtggg 2220
attcaattgc tcagaatcat ttacaatgat tggacacaga tcaattacgt gtattcatgg 2280
agtatggacc caacttcccc agtgtgtggc aatagataaa ottaagaagt gcaaatcatc 2340
aaatttaatt atacttgagg aacattttaa aaacaagaag gaattcgatc ataattctaa 2400

cataaggtac agatgtagag gaaaagaagg atggatacac acagtctgca taaatggaag 2460
atgggatcca gaagtgaact gctcaatggc acaaatacaa ttatgcccac ctccacctca 2520
gattcccaat tctcacaata tgacaaccac actgaattat cgggatggag aaaaagtatc 2580
tggtctttgc caagaaaatt atctaattca ggaaggagaa gaaattacat gcaaagatgg 2640
aagatggcag tcaataccac tctgtgttga aaaaattcca tggtcacaac cacctcagat 2700
agaacacgga accattaatt catccaggtc ttcacaagaa agttatgcac atgggactaa 2760
attgagttat acttgtagg gtggtttcag gatatctgaa gaaaatgaaa caacatgcta 2820
catgggaaaa tggagtctc cacctcagtg tgaaggcctt ccttgtaa atctccacctga 2880
gattttotcat ggtgtttag ctccatgtc agacagttat cagtatggag aagaagttac 2940
gtacaaatgt tttgaagggt ttggaattga tgggcctgca attgcaaaat gcttaggaga 3000
aaaatggtct caccctccat catgcataaa aacagattgt ctccagtttac ctagctttga 3060
aaatgccata cccatgggag agaagaagga tgtgtataag gcgggtgagc aagtgaacta 3120
cacttggtgca acatattaca aaatggatgg agccagtaat gtaacatgca ttaatagcag 3180
atggacagga aggccaaacat gcagagacac ctctgtgtg aatccgccca cagtacaaaa 3240
tgottatata gtgtcgagac agatgagtaa atatccatct ggtgagagag tacgttatca 3300
atgtaggagc ccttatgaaa tgtttgggga tgaagaagtg atgtgtttta atggaaactg 3360
gacggaacca cctcaatgca aagattctac aggaaaatgt gggcccccctc cacctattga 3420
caatggggac attacttcat tccggttgtc agtatatgct ccagcttcat cagttgagta 3480
ccaatgccag aacttgatc aacttgagg taacaagcga ataacatgta gaaatggaca 3540
atggtcagaa ccacaaaaat gttacatcc gtgtgtaata tccogagaaa ttatggaaaa 3600
ttataacata gcattaaggt ggacagccaa acagaagctt ttttcgagaa caggtgaatc 3660
agttgaatth gtgtgtaaac ggggatatcg tctttcatca cgttctcaca cattgcgaac 3720
aacatgttgg gatgggaaac tggagtatcc aacttggtgca aaaagataga atcaatcata 3780
aagtgcacac ctttattcag aactttagta ttaaatcagt tctcaatttc attttttatg 3840
tattgtttta ctctttttta ttcatacgta aaattttgga ttaattttgtg aaaatgtaat 3900
tataagctga gaccggtggc tctctt 3926

<210> 337
<211> 1231
<212> PRT
<213> Homo sapiens

<400> 337

Met Arg Leu Leu Ala Lys Ile Ile Cys Leu Met Leu Trp Ala Ile Cys
 1 5 10 15
 Val Ala Glu Asp Cys Asn Glu Leu Pro Pro Arg Arg Asn Thr Glu Ile
 20 25 30
 Leu Thr Gly Ser Trp Ser Asp Gln Thr Tyr Pro Glu Gly Thr Gln Ala
 35 40 45
 Ile Tyr Lys Cys Arg Pro Gly Tyr Arg Ser Leu Gly Asn Val Ile Met
 50 55 60
 Val Cys Arg Lys Gly Glu Trp Val Ala Leu Asn Pro Leu Arg Lys Cys
 65 70 75 80
 Gln Lys Arg Pro Cys Gly His Pro Gly Asp Thr Pro Phe Gly Thr Phe
 85 90 95
 Thr Leu Thr Gly Gly Asn Val Phe Glu Tyr Gly Val Lys Ala Val Tyr
 100 105 110
 Thr Cys Asn Glu Gly Tyr Gln Leu Leu Gly Glu Ile Asn Tyr Arg Glu
 115 120 125
 Cys Asp Thr Asp Gly Trp Thr Asn Asp Ile Pro Ile Cys Glu Val Val
 130 135 140
 Lys Cys Leu Pro Val Thr Ala Pro Glu Asn Gly Lys Ile Val Ser Ser
 145 150 155 160
 Ala Met Glu Pro Asp Arg Glu Tyr His Phe Gly Gln Ala Val Arg Phe
 165 170 175
 Val Cys Asn Ser Gly Tyr Lys Ile Glu Gly Asp Glu Glu Met His Cys
 180 185 190
 Ser Asp Asp Gly Phe Trp Ser Lys Glu Lys Pro Lys Cys Val Glu Ile
 195 200 205
 Ser Cys Lys Ser Pro Asp Val Ile Asn Gly Ser Pro Ile Ser Gln Lys
 210 215 220
 Ile Ile Tyr Lys Glu Asn Glu Arg Phe Gln Tyr Lys Cys Asn Met Gly
 225 230 235 240
 Tyr Glu Tyr Ser Glu Arg Gly Asp Ala Val Cys Thr Glu Ser Gly Trp

245 250 255
 Arg Pro Leu Pro Ser Cys Glu Glu Lys Ser Cys Asp Asn Pro Tyr Ile
 260 265 270
 Pro Asn Gly Asp Tyr Ser Pro Leu Arg Ile Lys His Arg Thr Gly Asp
 275 280 285
 Glu Ile Thr Tyr Gln Cys Arg Asn Gly Phe Tyr Pro Ala Thr Arg Gly
 290 295 300
 Asn Thr Ala Lys Cys Thr Ser Thr Gly Trp Ile Pro Ala Pro Arg Cys
 305 310 315 320
 Thr Leu Lys Pro Cys Asp Tyr Pro Asp Ile Lys His Gly Gly Leu Tyr
 325 330 335
 His Glu Asn Met Arg Arg Pro Tyr Phe Pro Val Ala Val Gly Lys Tyr
 340 345 350
 Tyr Ser Tyr Tyr Cys Asp Glu His Phe Glu Thr Pro Ser Gly Ser Tyr
 355 360 365
 Trp Asp His Ile His Cys Thr Gln Asp Gly Trp Ser Pro Ala Val Pro
 370 375 380
 Cys Leu Arg Lys Cys Tyr Phe Pro Tyr Leu Glu Asn Gly Tyr Asn Gln
 385 390 395 400
 Asn Tyr Gly Arg Lys Phe Val Gln Gly Lys Ser Ile Asp Val Ala Cys
 405 410 415
 His Pro Gly Tyr Ala Leu Pro Lys Ala Gln Thr Thr Val Thr Cys Met
 420 425 430
 Glu Asn Gly Trp Ser Pro Thr Pro Arg Cys Ile Arg Val Lys Thr Cys
 435 440 445
 Ser Lys Ser Ser Ile Asp Ile Glu Asn Gly Phe Ile Ser Glu Ser Gln
 450 455 460
 Tyr Thr Tyr Ala Leu Lys Glu Lys Ala Lys Tyr Gln Cys Lys Leu Gly
 465 470 475 480
 Tyr Val Thr Ala Asp Gly Glu Thr Ser Gly Ser Ile Arg Cys Gly Lys
 485 490 495

Asp Gly Trp Ser Ala Gln Pro Thr Cys Ile Lys Ser Cys Asp Ile Pro
500 505 510

Val Phe Met Asn Ala Arg Thr Lys Asn Asp Phe Thr Trp Phe Lys Leu
515 520 525

Asn Asp Thr Leu Asp Tyr Glu Cys His Asp Gly Tyr Glu Ser Asn Thr
530 535 540

Gly Ser Thr Thr Gly Ser Ile Val Cys Gly Tyr Asn Gly Trp Ser Asp
545 550 555 560

Leu Pro Ile Cys Tyr Glu Arg Glu Cys Glu Leu Pro Lys Ile Asp Val
565 570 575

His Leu Val Pro Asp Arg Lys Lys Asp Gln Tyr Lys Val Gly Glu Val
580 585 590

Leu Lys Phe Ser Cys Lys Pro Gly Phe Thr Ile Val Gly Pro Asn Ser
595 600 605

Val Gln Cys Tyr His Phe Gly Leu Ser Pro Asp Leu Pro Ile Cys Lys
610 615 620

Glu Gln Val Gln Ser Cys Gly Pro Pro Pro Glu Leu Leu Asn Gly Asn
625 630 635 640

Val Lys Glu Lys Thr Lys Glu Glu Tyr Gly His Ser Glu Val Val Glu
645 650 655

Tyr Tyr Cys Asn Pro Arg Phe Leu Met Lys Gly Pro Asn Lys Ile Gln
660 665 670

Cys Val Asp Gly Glu Trp Thr Thr Leu Pro Val Cys Ile Val Glu Glu
675 680 685

Ser Thr Cys Gly Asp Ile Pro Glu Leu Glu His Gly Trp Ala Gln Leu
690 695 700

Ser Ser Pro Pro Tyr Tyr Tyr Gly Asp Ser Val Glu Phe Asn Cys Ser
705 710 715 720

Glu Ser Phe Thr Met Ile Gly His Arg Ser Ile Thr Cys Ile His Gly
725 730 735

Val Trp Thr Gln Leu Pro Gln Cys Val Ala Ile Asp Lys Leu Lys Lys
 740 745 750
 Cys Lys Ser Ser Asn Leu Ile Ile Leu Glu Glu His Leu Lys Asn Lys
 755 760 765
 Lys Glu Phe Asp His Asn Ser Asn Ile Arg Tyr Arg Cys Arg Gly Lys
 770 775 780
 Glu Gly Trp Ile His Thr Val Cys Ile Asn Gly Arg Trp Asp Pro Glu
 785 790 795 800
 Val Asn Cys Ser Met Ala Gln Ile Gln Leu Cys Pro Pro Pro Pro Gln
 805 810 815
 Ile Pro Asn Ser His Asn Met Thr Thr Thr Leu Asn Tyr Arg Asp Gly
 820 825 830
 Glu Lys Val Ser Val Leu Cys Gln Glu Asn Tyr Leu Ile Gln Glu Gly
 835 840 845
 Glu Glu Ile Thr Cys Lys Asp Gly Arg Trp Gln Ser Ile Pro Leu Cys
 850 855 860
 Val Glu Lys Ile Pro Cys Ser Gln Pro Pro Gln Ile Glu His Gly Thr
 865 870 875 880
 Ile Asn Ser Ser Arg Ser Ser Gln Glu Ser Tyr Ala His Gly Thr Lys
 885 890 895
 Leu Ser Tyr Thr Cys Glu Gly Gly Phe Arg Ile Ser Glu Glu Asn Glu
 900 905 910
 Thr Thr Cys Tyr Met Gly Lys Trp Ser Ser Pro Pro Gln Cys Glu Gly
 915 920 925
 Leu Pro Cys Lys Ser Pro Pro Glu Ile Ser His Gly Val Val Ala His
 930 935 940
 Met Ser Asp Ser Tyr Gln Tyr Gly Glu Glu Val Thr Tyr Lys Cys Phe
 945 950 955 960
 Glu Gly Phe Gly Ile Asp Gly Pro Ala Ile Ala Lys Cys Leu Gly Glu
 965 970 975

Lys Trp Ser His Pro Pro Ser Cys Ile Lys Thr Asp Cys Leu Ser Leu
 980 985 990

Pro Ser Phe Glu Asn Ala Ile Pro Met Gly Glu Lys Lys Asp Val Tyr
 995 1000 1005

Lys Ala Gly Glu Gln Val Thr Tyr Thr Cys Ala Thr Tyr Tyr Lys
 1010 1015 1020

Met Asp Gly Ala Ser Asn Val Thr Cys Ile Asn Ser Arg Trp Thr
 1025 1030 1035

Gly Arg Pro Thr Cys Arg Asp Thr Ser Cys Val Asn Pro Pro Thr
 1040 1045 1050

Val Gln Asn Ala Tyr Ile Val Ser Arg Gln Met Ser Lys Tyr Pro
 1055 1060 1065

Ser Gly Glu Arg Val Arg Tyr Gln Cys Arg Ser Pro Tyr Glu Met
 1070 1075 1080

Phe Gly Asp Glu Glu Val Met Cys Leu Asn Gly Asn Trp Thr Glu
 1085 1090 1095

Pro Pro Gln Cys Lys Asp Ser Thr Gly Lys Cys Gly Pro Pro Pro
 1100 1105 1110

Pro Ile Asp Asn Gly Asp Ile Thr Ser Phe Pro Leu Ser Val Tyr
 1115 1120 1125

Ala Pro Ala Ser Ser Val Glu Tyr Gln Cys Gln Asn Leu Tyr Gln
 1130 1135 1140

Leu Glu Gly Asn Lys Arg Ile Thr Cys Arg Asn Gly Gln Trp Ser
 1145 1150 1155

Glu Pro Pro Lys Cys Leu His Pro Cys Val Ile Ser Arg Glu Ile
 1160 1165 1170

Met Glu Asn Tyr Asn Ile Ala Leu Arg Trp Thr Ala Lys Gln Lys
 1175 1180 1185

Leu Tyr Ser Arg Thr Gly Glu Ser Val Glu Phe Val Cys Lys Arg
 1190 1195 1200

Gly Tyr Arg Leu Ser Ser Arg Ser His Thr Leu Arg Thr Thr Cys

1205

1210

1215

Trp	Asp	Gly	Lys	Leu	Glu	Tyr	Pro	Thr	Cys	Ala	Lys	Arg
1220						1225					1230	