

**TOCOTRIENOLS AND TOCOTRIENOL-LIKE
COMPOUNDS AND METHODS FOR THEIR USE**

TECHNICAL FIELD OF THE INVENTION

The present invention relates to novel
5 tocotrienols and tocotrienol-like compounds displaying
biological activity. The tocotrienols and tocotrienol-
like compounds of this invention may be conveniently
obtained from biological sources or by chemical
synthesis and may be used in pharmaceutical
10 compositions, foodstuffs and dietary supplements. This
invention also relates to the use of tocotrienols,
tocotrienol-like compounds, and mixtures thereof, as
hypocholesterolemic, antithrombotic, antioxidizing,
antiatharogenic, antiinflammatory and immunoregulatory
15 agents, or as agents useful to decrease lipoprotein (a)
concentration in the blood or to increase feed
conversion efficiency.

BACKGROUND OF THE INVENTION

Plant constituents have been proven useful in
20 the prevention and treatment of a wide variety of
diseases and conditions. For example, barley has been
shown to be particularly effective in lowering lipid
levels in animal models (A.A. Qureshi et al.,
"Suppression Of Cholesterogenesis By Plant
25 Constituents", Lipids, 20, pp. 817-24 (1985)). More
specifically, α -tocotrienol, a chromanol isolated from

barley extract, has been identified as a therapeutic agent for hypercholesterolemia (A.A. Qureshi et al., "The Structure Of An Inhibitor Of Cholesterol Biosynthesis Isolated From Barley", J. Biol. Chem., 5 261, pp. 10544-50 (1986)). In addition, tocotrienol, γ -tocotrienol and δ -tocotrienol have also been shown to reduce hypercholesterolemia in mammals (European patent application 421,419).

Hypercholesterolemia involves high serum 10 cholesterol levels and is a causative agent of diseases including arteriosclerosis, atherosclerosis, cardiovascular disease and xanthomatosis. In addition, high serum cholesterol levels are seen in patients suffering from diseases including diabetes mellitus, 15 familial hypercholesterolemia, acute intermittent prothyria, anorexia nervosa, nephrotic syndrome, primary cirrhosis and various liver disorders, such as hepatitis and obstructive jaundice. Improvement of lipoprotein profiles and a decrease in total serum and 20 low density lipoprotein cholesterol have been shown to retard the progression of such diseases, as well as to induce regression of clinically significant lesions in hypercholesterolemic patients.

Although the relationship between 25 hypercholesterolemia and its many associated diseases, most notably cardiovascular disease, has been extensively studied, no clear answer to this world-wide problem has yet been found. As a result, coronary artery disease remains the leading cause of death in 30 the United States and other developed countries. Coronary artery disease is the result of complex interactions between a large number of different processes, including lipoprotein metabolism, aggregation of blood platelets, blood coagulation and 35 fibrinolysis. Accordingly, the cardiovascular risk

profile of a given patient is dependent on these interactions.

In addition to lowering cholesterol levels, the cardiovascular risk profile of a patient may also be reduced by decreasing the levels of other factors in the serum and the blood. For example, reduction of thromboxane A_2 generation (measured by the levels of thromboxane B_2 , a stable metabolite of thromboxane A_2) and platelet factor 4 levels in the serum lessens the risk of cardiovascular disease because of decreased thrombogenic activity.

Thromboxane A_2 and platelet factor 4 levels are also associated with other biological activities. For example, when reduction of these factors is accompanied by a reduction in macrophage cell count, lower tumor necrosis factor (TNF) levels and lower arachidonic acid levels in bodily tissues, reduced levels of prostaglandins, leukotrienes and interleukins are implicated. Reduction of these factors, therefore, leads to a decrease in the inflammation accompanying a wide variety of diseases. In addition, since prostaglandins inhibit glucose-induced insulin release and increase glucagon secretion, an increased insulin to glucagon ratio may also result from the reduction in prostaglandins. Such an increase is useful in improving glucose intolerance in diabetes mellitus and restoration of acute glucose-induced insulin response in non-insulin-dependent diabetes mellitus.

It has been noted that there is a low incidence of cardiovascular disease in populations consuming large amounts of cereal grains. Soluble and insoluble fibers have, in the past, been viewed as the agents responsible for cholesterol reduction in such populations (see D. Kritchevsky et al., "Fiber, Hypercholesterolemia and Atherosclerosis", Lipids, 13,

pp. 366-69 (1978)). Recently, the hypocholesterolemic effects of cereal grains have been attributed to natural components of the grains -- tocotrienols ("T₃") and structurally similar compounds, such as tocopherols ("T"). Tocotrienols and tocopherols occur naturally in small quantities in a wide variety of plant sources, such as rice bran, palm oil and barley (A.A. Qureshi et al., "Lowering of Serum Cholesterol in Hypercholesterolemic Humans by Tocotrienols (Palmvitea)", Am. J. Clin. Nutr., 53, pp. 1021S-6S (1991)).

As a class, the tocopherols, including d- α -tocopherol (vitamin E), have been extensively studied. As a result of these studies, certain biological activities have been attributed to the tocopherols. Such activities include platelet aggregation and antioxidant functions (see, for example, E. Niki et al., "Inhibition of Oxidation of Biomembranes By Tocopherol", Annals of the New York Academy of Sciences, 570, pp. 23-31 (1989) and K. Fukuzawa et al., "Increased Platelet-Activating Factor (PAF) Synthesis in Polymorphonuclear Leukocytes of Vitamin E-Deficient Rats", Annals of the New York Academy of Sciences, 570, pp. 449-453 (1989)). Although the exact structure-function relationship is not known, several experiments have highlighted the importance of the phytyl side chain in the biological activity of tocopherols (see W.A. Skinner et al., "Antioxidant Properties of α -Tocopherol Derivatives and Relationships of Antioxidant Activity to Biological Activity", Lipids, 5(2), pp. 184-186 (1969) and A.T. Diplock, "Relationship of Tocopherol Structure to Biological Activity, Tissue Uptake, and Prostaglandin Biosynthesis", Annals of the New York Academy of Sciences, 570, pp. 73-84 (1989)).

In contrast to the tocopherols, interest in the tocotrienols has been limited, as those compounds were not typically considered to be biologically useful. Recently, however, studies have indicated that tocotrienols may be biologically active. For example, United States patent 4,603,142 identifies d- α -tocotrienol, isolated from barley extracts, as an inhibitor of cholesterol biosynthesis. See also A.A. Qureshi et al. (1986), *supra*. Various human and animal studies have confirmed the impact of pure tocotrienols, isolated from barley, oats and palm oil, on cholesterol biosynthesis, specifically LDL-cholesterol (A.A. Qureshi et al., "Dietary Tocotrienols Reduce Concentrations of Plasma Cholesterol, Apolipoprotein B, Thromboxane B₂ and Platelet Factor 4 In Pigs With Inherited Hyperlipidemias", Am. J. Clin. Nutr., pp. 1042S-46S (1991); A.A. Qureshi et al., "Lowering Of Serum Cholesterol In Hypercholesterolemic Humans By Tocotrienols (Palmvite)", Am. J. Clin. Nutr., 53, pp. 1021S-26S (1991); D.T.S. Tan et al., "The Effect Of Palm Oil Vitamin E Concentrate On The Serum And Lipoprotein Lipids In Humans", Am. J. Clin. Nutr., 53, pp. 1027S-30S (1991)). In addition, tocotrienol, γ - and δ -tocotrienol have been indicated for use in the treatment of hypercholesterolemia, hyperlipidemia and thromboembolic disorders (European patent application 421,419).

The five known naturally occurring tocotrienols have been designated tocotrienol, α -, β -, γ - and δ -tocotrienol. Those compounds exhibit varying degrees of hypercholesterolemic activity and have also been used as antithrombotic agents and antioxidants. α -T₃, for example, displays antioxidant activity against lipid peroxidation in rat liver microsomal membranes and against oxidative damage of cytochrome P-

450 (E. Serbinova, Free Radical Biology and Medicine,
in press (1991)). Despite these activities, the known
tocotrienols have not found wide-spread therapeutic
use.

5 Accordingly, the need still exists for
compounds which, as single agents, can safely and
effectively act as hypercholesterolemic,
antithrombotic, antioxidizing, antiatherogenic,
antiinflammatory and immunoregulatory agents.

10 SUMMARY OF THE INVENTION

 The present invention solves the problems
referred to above by providing novel tocotrienols,
tocotrienol-like compounds, and mixtures thereof, that
are useful as hypocholesterolemic, antithrombotic,
15 antioxidizing, antiatherogenic, antiinflammatory and
immunoregulatory agents. These novel tocotrienols and
tocotrienol-like compounds are also useful in
decreasing lipoprotein (a) ("Lp(a)") concentration in
the blood, increasing feed conversion efficiency, and
20 in the treatment or prevention of conditions such as
fever, edema, diabetes mellitus, cancer, signs of
aging, pain, rheumatoid diseases, septic shock, chronic
fatigue syndrome and functio laesa.

 The novel tocotrienols and tocotrienol-like
25 compounds of the present invention are useful in
inhibiting the synthesis of HMG-CoA reductase, reducing
lipogenesis and increasing the HDL/LDL cholesterol
ratio, as well as reducing total serum cholesterol, low
density lipoprotein-cholesterol, Lp(a), apolipoprotein
30 B, thromboxane A₂, platelet factor 4, triglycerides and
glucose. Because of such individual activities, these
compounds are useful in treating diseases, for example,
coronary artery disease, which result from the
interaction of processes such as lipoprotein

metabolism, aggregation of blood platelets, blood coagulation and fibrinolysis. Accordingly, the compounds of this invention are useful to improve the overall cardiovascular profile of patients.

5 This invention also provides novel uses for the known tocotrienols as antiinflammatory and immunoregulatory agents, to decrease Lp(a) concentration in the blood, to increase feed conversion efficiency, and for the treatment or prevention of
10 conditions such as fever, edema, diabetes mellitus, pain, septic shock, chronic fatigue syndrome and functio laesa.

Advantageously, the tocotrienol and tocotrienol-like compounds reduce serum TNF and IL-1
15 levels. These activities render the compounds useful in reducing active and chronic inflammation, such as that associated with rheumatoid disease. Also, such activities render the compounds useful in treating, preventing or lessening the severity of
20 immunoregulatory diseases, such as autoimmune diseases, and in preventing or treating pain, septic shock, chronic fatigue syndrome, functio laesa and oxidative conditions.

The effects of tocotrienol and tocotrienol-
25 like compounds on lipid metabolism also influence the regulation of antibody production. Thus, these compounds are useful in modulating immune function through adjustments in fatty acid levels.

Tocotrienol and tocotrienol-like compounds
30 also decrease glucose levels by mediating the levels of insulin and glucagon. By increasing the insulin to glucagon ratio in the blood, these compounds are useful in increasing glucose intolerance in diabetes mellitus and restoring acute glucose-induced insulin response in
35 non-insulin dependent diabetes mellitus patients.

The novel tocotrienols and tocotrienol-like compounds of this invention are characterized by specific structural characteristics and specific biological activity or, alternatively, by specific high performance liquid chromatography ("HPLC") elution profiles and specific biological activity. More particularly, the compounds of this invention may be characterized by three structural features: (1) a hydrogen donor group (or a group which can be hydrolyzed to a hydrogen donor group) attached to an aromatic ring system, (2) an atom having at least one lone pair of electrons, said electrons being in conjugation with the aromatic system and (3) a side chain comprising one or more isoprenoid or isoprenoid-like units attached to a position adjacent to that atom. This invention also encompasses the hydrolysis and oxidation products obtained from such compounds. In addition, the compounds of this invention having the above-mentioned structural characteristics are also characterized by the ability to inhibit the activity of β -hydroxy- β -methyl glutaryl coenzyme A (HMG-CoA) reductase. Furthermore, these compounds are effective in the treatment or prevention of one or more of the following diseases or conditions: hypercholesterolemic diseases, thrombotic diseases, oxidative conditions, inflammation or immunoregulatory diseases or, alternatively, in increasing feed conversion efficiency.

According to the alternate embodiment, the novel tocotrienols and tocotrienol-like compounds of this invention may be characterized by an elution time of at least 22 minutes under the following HPLC conditions: μ -Porasil column (Waters column, 10 μ , 4 mm x 30 cm) using an isocratic system of hexane and isopropanol (99.75%:0.25%, v/v) at a flow rate of 1.3

ml/min. The particular compound is detected at an
excitation wavelength of 295 nm and an emission
wavelength of 330 nm (fluorescence detector) and UV
absorption at 295 nm. In addition to the above-defined
5 elution profile, the novel tocotrienols of this
invention are also characterized by the ability to
inhibit the activity of HMG-CoA reductase.

According to this invention, tocotrienols and
tocotrienol-like compounds may be conveniently obtained
10 from biological sources. Alternatively, they may be
synthesized using conventional chemical methodologies.
In a preferred embodiment of this invention, the
tocotrienols and tocotrienol-like compounds are
isolated and purified from stabilized biological
15 sources. This invention also encompasses the
techniques used to purify such compounds.

In a preferred embodiment of this invention,
the tocotrienols, tocotrienol-like compounds and
mixtures thereof, may be administered to an animal or a
20 human as a pharmaceutical composition, a foodstuff or a
dietary supplement, to treat or prevent
hypercholesterolemic diseases, thrombotic diseases,
oxidative or atherogenic conditions, inflammation or
immunoregulatory diseases. Advantageously, these
25 compositions, foodstuffs and dietary supplements may
also be used to increase feed conversion efficiency,
decrease Lp(a) concentration in the blood and in the
treatment or prevention of fever, edema, diabetes
mellitus, cancer, signs of aging, pain, rheumatoid
30 diseases, septic shock, chronic fatigue syndrome and
functio laesa.



THIS SHEET HAS BEEN
REPLACED

- 10 -

BRIEF DESCRIPTION OF THE DRAWINGS

Figures 1-4 depict the effects of different tocotrienols from rice bran oil on fatty acid concentrations in various tissues of hyper-
5 cholesterolemic swine.

Figure 5 depicts the effects of GT301 (γ -T₃) on superoxide production in human peripheral blood neutrophils.

DETAILED DESCRIPTION OF THE INVENTION

10 In order that the invention herein described may be more fully understood, the following detailed description is set forth. In the description, the following terms are employed:

Biological source - Any natural or
15 recombinant plant source, natural or transgenic animal source, microbial source (e.g., bacteria), fungi, yeast, algae, higher plant source, or derivative thereof, which contains one or more tocopherol, tocotrienol or tocotrienol-like compounds.

20 Stabilization - A process effective to increase the recoverable amounts of tocopherols, tocotrienols and tocotrienol-like compounds in a biological source by one or a combination of:
(1) inactivating enzymes which are capable of degrading
25 tocotrienols and tocotrienol-like compounds in that biological source, (2) breaking bonds or otherwise interfering with the interactions -- such as hydrogen bonds, covalent bonds, ionic bonds, Van der Waals forces, London forces and hydrophobic or hydrophilic
30 interactions which bind the desired products to proteins, sugars, lipids, carbohydrates, glycoproteins, other biological molecules (i.e., amino acids or nucleotides) or other membrane components, or combinations thereof, in the biological source -- which

retain tocotrienols and tocotrienol-like compounds in that biological source, thus facilitating the release of those desired compounds, or (3) increasing the solubility of the tocotrienols and tocotrienol-like compounds of that biological source beyond that prior to stabilization or beyond the level of solubility of the tocotrienols of a corresponding non-stabilized biological source. As a result of stabilization, tocotrienols and tocotrienol-like compounds in a biological source may be recovered in higher yields than those realized from a corresponding non-stabilized biological source.

Tocol - A mixture of one or more compounds selected from tocopherols (T), tocotrienols (T_3), and tocotrienol-like (T_3 -like) compounds.







Tocotrienol-like - Any biologically active compound which is contained in or derived from a biological source and (1) which is released, or whose release is facilitated, upon stabilizing that source or (2) whose recoverable amount in that source is increased by stabilizing that source. Such tocotrienol-like compounds include any biologically active compound displaying the biological activity of a tocotrienol which inhibits the activity of HMG-CoA reductase as measured by an in vitro HMG-CoA reductase assay, such as that described in D.J. Shapiro et al., "Microassay for β -hydroxy- β -methylglutaryl-CoA Reductase in Rat Liver and in L-cell Fibroblasts", Biochim. Biophys. Acta, 370, p. 369 (1974).

Tocotrienol-like compounds include, but are not limited to, any electron transfer ring compounds, anti-oxidant type compounds, redox compounds and compounds similar to or containing the three structural features that characterize the tocotrienols of this invention. Specific examples of T_3 -like compounds are ubiquinones,

plastoquinones, isoquinones, phylloquinones, benzoquinones, flavanols, flavanoids, coumarins, unsaturated terpenoids and unsaturated isoprenoids. The term "T₃-like compound" also encompasses analogues, 5 homologs, isomers and derivatives of such compounds, such as prenylated derivatives or pyrolytic derivatives.

Preferred tocotrienol-like compounds have the following structures:

10



and salts, oxidation products and hydrolysis products thereof wherein:

R_1 and R_3 is each independently selected from the group consisting of H, halogen, OH, OCH_3 and C_1-C_8 branched or unbranched alkyls;


R_2 is selected from the group consisting of OH, NHR_8 , CO_2Y , $C(R_8)_2CO_2H$, and C_1-C_8 branched or unbranched alkyls substituted by OH, NHR_8 , CO_2H or $C(R_8)_2CO_2H$;

R_4 is selected from the group consisting of O, NH, $CH-R_9$, $C=O$ and $CH-OH$;

R_5 is selected from the group consisting of CH_2 , $CHOH$, O, S and NH;

R_6 is selected from the group consisting of H and C_1-C_8 branched or unbranched alkyls;

Each R_7 independently represents



wherein R_{10} is selected from the group consisting of H, NH_2 and C_1-C_8 branched or unbranched alkyls;

R_8 and R_9 is each independently selected from the group consisting of H and C_1-C_8 branched or unbranched alkyls;

Each R_{11} is independently selected from the group consisting of H, C_1-C_8 branched or unbranched alkyls, CH_2OH , CO_2H or OH;

Y is H or C_1-C_{18} branched or unbranched alkyls, preferably C_1-C_4 alkyls;

Z is H, halogen, OH, CH₂OH, CH₃, OCH₃ or COCH₃;

n is an integer from 0-4; and

m is an integer from 1-30, more preferably
5 1-20 or most preferably 3-10.

TRF - A tocotrienol-rich fraction obtained by the stabilization and extraction of a biological source. It typically contains varying amounts of the five known tocotrienols, the known tocopherols and the
10 novel tocotrienols and tocotrienol-like compounds of this invention. Most commonly, the TRF will be composed of about 50% to about 90% tocotrienols and tocotrienol-like compounds. The TRF, may be used for any of the uses described herein for either the known
15 tocotrienols or the novel tocotrienols and tocotrienol-like compounds of this invention.

TRF Standard - A tocotrienol-rich fraction (TRF) obtained from palm oil (A.A. Qureshi et al. (1991), supra). The TRF Standard contains varying
20 amounts of α -, γ - and δ -tocotrienol and α -tocopherol but essentially none of the novel tocotrienol and tocotrienol-like compounds of this invention.

Enhanced - The state of a stabilized biological source, wherein the recoverable amount of
25 the novel tocotrienols and tocotrienol-like compounds of this invention is increased beyond that recoverable from the biological source prior to stabilization.

Foodstuff - Substances that can be used or prepared for use as food for an animal or a human.
30 Foodstuffs include substances that may be used in the preparation of food (such as frying oils) or food additives. For example, foodstuffs include animals used for human consumption or any product therefrom, such as, for example, eggs. Such animals themselves
35 may have ingested or been treated with one or more

tocotrienol or tocotrienol-like compound, or foodstuffs containing them. As discussed herein, after ingesting or being otherwise administered tocotrienols or tocotrienol-like compounds, such animals retain the
5 advantageous hypocholesterolemic, antithrombotic, antioxidizing, antiatherogenic, antiinflammatory and immunoregulatory properties of the tocotrienols and tocotrienol-like compounds.

HPLC elution time or profile - Except where
10 otherwise noted, the term "HPLC elution time or profile" refers to the time necessary for a given compound to elute from the HPLC column or the characteristic profile of a mixture of compounds (20 μ l) under the following conditions: μ -Porasil
15 column (Waters column, 10 μ , 4 mm x 30 cm) using an isocratic system of hexane and isopropanol (99.75%:0.25%, v/v) at a flow rate of 1.3 ml/min. The compound is detected at an excitation wavelength of 295 nm and an emission wavelength of 330 nm (fluorescence
20 detector) and UV absorption at 295 nm.

The individual tocotrienols and tocotrienol-like compounds of this invention constitute a novel class of biologically active compounds. Several of these compounds occur naturally in small quantities in
25 biological sources. Therefore, in a preferred embodiment of this invention, tocotrienols and tocotrienol-like compounds are recovered from biological materials which have been stabilized and extracted according to the processes described in
30 copending International patent application PCT/US91/03626, filed on May 23, 1991. The stabilization and recovery process disclosed therein enhances or increases the yields of tocotrienols and tocotrienol-like compounds recovered from biological
35 sources.

The tocotrienol and tocotrienol-like compounds may be recovered from any biological materials including, but not limited to, oats, wheat, rye, barley, soybean, wheat germ, wheat bran, corn, rice (including whole kernel, husk or hull, endosperm and germ), cottonseed, milkweed, flax, sesame, rice bran, parboiled brown rice, brown rice flour, olives, vegetable oil distillant, fruit concentrate evaporate, barley bran, palm oil, wheat germ oil, rice bran oil, barley oil, coconut oil, cottonseed oil, soybean oil, other cereal grains and other cereal grain oils, plant tissues, flowers, bushes (such as juniper), trees (such as pine and rubber), fruits (such as melons, berries, tomatoes and citrus fruits), vegetables, grasses (such as alfalfa), fungi (such as mushrooms), leaves, seeds (such as sesame, millet and pine), such as sesame seeds and pine seeds, stems, bark, roots, nuts (such as cashews and almonds) and legumes (such as peanuts), or portions thereof. We have noted that the tocotrienols and tocotrienol-like compounds decompose in the biological source over time. Therefore, we prefer to use freshly harvested biological sources. Most preferably, the biological source is freshly harvested, stabilized rice bran.

Alternatively, the tocotrienols and tocotrienol-like compounds may be obtained from a non-stabilized source or synthesized according to known chemical methodology. Typical synthetic routes are described in J.W. Scott et al., "Synthesis of (2R, 4'R, 8'R)- α -Tocopherol and (2R, 3'E, 7'E)- α -Tocotrienol", Helv. Chem. Acta, 59, pp. 290-306 (1976) and P. Schudel et al., "Die Synthese von rac. all-trans δ - und ϵ -Tocopherol", Helv. Chem. Acta, 46, pp. 2517-2526 (1963). An alternative synthetic scheme is proposed in European patent application 421,419.

The tocotrienols and tocotrienol-like compounds of this invention may be altered by known chemical means to produce various derivatives or analogues. Such derivatives or analogues may be more easily isolated in pure form, or more resistant to degradation, or possess other desired characteristics. Such derivatives and analogues are also envisioned by this invention. Known chemical means to alter the compounds of this invention include, but are not limited to, heating in the presence or absence of air or in an inert or reactive gas environment (or in a mixture of inert and active gases) and pyrolysis.


This invention encompasses the d- or l-isomer and the d, l- racemic mixture of each tocotrienol and tocotrienol-like compound. However, the naturally occurring d-isomer is preferred. This invention also includes mixtures of at least one novel tocotrienol or tocotrienol-like compound of this invention with one or more of the known tocotrienols.

The tocotrienols and tocotrienol-like compounds of this invention may be characterized either by specific structural characteristics or alternatively, by a specific elution profile and specific biological activity. In the former case, the compounds of this invention share three structural features: (1) a hydrogen donor group (or a group which can be hydrolyzed to a hydrogen donor group) attached to an aromatic ring system, (2) an atom having at least one lone pair of electrons, said electrons being in conjugation with the aromatic system and (3) a side chain comprising one or more isoprenoid or isoprenoid-like units attached to a position adjacent to that atom. This invention also encompasses the hydrolysis and oxidation products obtained from such compounds. These compounds are effective in increasing feed

conversion efficiency, decreasing Lp(a) concentration in the blood or, alternatively, in the treatment or prevention of one or more of the following diseases or conditions: immunoregulatory disease, inflammation, fever, edema, diabetes mellitus, cancer, signs of aging, pain, rheumatoid diseases, septic shock, chronic fatigue syndrome and functio laesa.

According to an alternate embodiment, the novel tocotrienols and tocotrienol-like compounds of this invention may be characterized by an elution time and specific biological activity. These compounds elute after at least 22 minutes under the following high performance liquid chromatography (HPLC) conditions: Porasil column (Waters column, 10 μ , 4 mm x 30 cm) using an isocratic system of hexane and isopropanol (99.75%:0.25%, v/v) at a flow rate of 1.3 ml/min. The compound is detected at an excitation wavelength of 295 nm and an emission wavelength of 330 nm (fluorescence detector) and UV absorption at 295 nm. In addition to the above-defined elution profile, the novel tocotrienols and tocotrienol-like compounds of this invention are also characterized by the ability to inhibit the activity of β -hydroxy- β -methyl glutaryl coenzyme A reductase ("HMG-CoA reductase").


The preferred class of compounds of this invention is of formula I:




- 19 -

above. As herein described above, the compounds of formula I do not include dasmethyl-tocotrienol (3,4-dihydro-2-methyl-2-(4,8,12-trimethyltrideca-3'(E),7'(E),11'-trienyl)-2H-1-benzopyran-6-ol).*

- 5 Typical hydrolysis products of the compounds of formula I which are also envisioned by this invention include, but are not limited to, compounds of formula II:



This Sheet Has Been Replaced




- 10 wherein R₁-R₁₀, Z, n and m are as defined as above. Preferred compounds are the compounds of formula I, wherein R₁, R₃ and Z are each independently H, halogen, OH, OCH₃ or CH₃; R is OH, OCH₃ or NH₂; R₄ is C=O, CH₂ or NH; R₅ is O, S, CH₂ or NH; R₆ is H or CH₃; R₁₀ and
15 R₁₁ are each independently H or CH₃; n is 0 or 1; and m is 3-10.

- The most preferred compounds are the compounds of formula I, wherein R₁, R₃, Z, R₆ and R₉ are each hydrogen, R₂ is OH, R₄ is CH₂, R₅ is CH₂ or O, R₆
20 is H or CH₃, R₁₀ is H or CH₃, n is 1 and m is 3.

- The novel compound, 3,4-dihydro-2-(4,8,12-trimethyltrideca-3'(E),7'(E),11' trienyl)-2H-1-benzopyran-6-ol ("P₂₅"), is representative of the class of compounds according to formula I. The structure of
25 this compound is shown below:

* The compounds of formula I also exclude α-, β-, γ- and δ-tocotrienol.



Tocotrienols and tocotrienol-like compounds of this invention having the above-defined structural features are also characterized by biological activity.

- 5 They are useful in increasing feed conversion efficiency, decreasing Lp(a) concentration in the blood or in the treatment or prevention of one or more of the following diseases or conditions: immunoregulatory disease, inflammation, fever, edema, diabetes mellitus,
- 10 cancer, signs of aging, pain, rheumatoid diseases, septic shock, chronic fatigue syndrome and functio laesa. They may also reduce the levels of TNF, IL-1 and IL-1 stimulatory products including possibly IL-2, IL-6, IL-8, GMCSF, prostaglandins and gamma interferon,
- 15 while increasing antibody titers in response to foreign proteins. By virtue of these activities, these compounds may result in a decrease in the release of superoxide and other cytotoxins produced by neutrophils, mast cells, basophils, monocytes and
- 20 macrophages, eosinophils, platelets, lymphocytes and polymorphonuclear leukocytes, endothelial tissue and other immunoregulatory tissues. These compounds may also effect an increase in antibody titers.

- HMG-CoA reductase catalyzes the rate-
- 25 limiting step of cholesterol biosynthesis. Therefore, a reduction in its activity decreases the total amount of cholesterol in the serum of animals and humans alone, or in combination with a low fat, low cholesterol diet. The effects are most noticeable in
- 30 hypercholesterolemic individuals with poor dietary

regimens. Typically, the tocotrienols and tocotrienol-like compounds of this invention reduce the activity of HMG-CoA reductase by at least about 15% in the in vitro assay described in Shapiro (supra), at a concentration
5 of 10-20 µg/ml. More preferably, however, the HMG-CoA reductase activity is reduced by at least about 20% and most preferably by about 25%.

While not wishing to be bound by theory, we believe that some of the antiinflammatory,
10 antioxidizing and immunoregulatory properties of the tocotrienol and tocotrienol-like compounds is a result of an inhibition in the production of free arachidonic acid.

We believe that the inhibition in the
15 production of arachidonic acid is caused by either the inhibition of phospholipase A₂ or alternatively, by increasing the amount of corticosterone level in the blood. Phospholipase A₂ cleaves at C-2 of phosphate head groups, resulting in the release of free
20 arachidonic acid. Arachidonic acid can then be converted into a variety of biologically important molecules, such as the prostaglandins and thromboxanes (via the cyclooxygenase pathway) and the leukotrienes (via the lipoxygenase pathway). We believe that the
25 hydrogen donor group of the tocotrienol and tocotrienol-like compounds bonds with the phosphate or phospholipid groups of biological membranes. In conjunction with the aromatic system, it also plays a role in radical scavenging. The isoprenoid tail may
30 act to stabilize the membrane -- possibly by forming weak bonds with the lipid side chains. In addition, by optionally replacing the tocotrienol methyl with other substituents (such as, for example, hydrogen) at position 2 on the chromanol ring, we are able to
35 maximize the stabilization of the membrane. We believe

that the aromatic system may also aid in tightening the surface of the membrane, thus increasing the steric hindrance for phospholipase A₂ and phospholipase C and decreasing its permeability. The aromatic system may
5 also serve electron-transport and antioxidizing functions. Such a hypothesis would contribute to the explanation of the broad spectrum of biological functions that are effected by the administration of the tocotrienols and tocotrienol-like compounds
10 according to this invention.

This invention also includes several novel uses for known tocotrienols, as well as tocotrienols and tocotrienol compounds as described herein. Examples include their use as antiinflammatory,
15 antiatherogenic and immunoregulatory agents, for the treatment of fever, edema, diabetes mellitus, cancer, signs of aging, pain, septic shock, chronic fatigue syndrome and functio laesa. In addition, tocotrienols and tocotrienol-like compounds are useful for
20 increasing feed conversion efficiency -- resulting in conversion of a greater percentage of food into protein rather than fat -- and decreasing Lp(a) concentration in the blood. Known tocotrienols exhibit much of the same biological activity attributed to the novel
25 tocotrienols and tocotrienol-like compounds of this invention. For example, known tocotrienols are capable of reducing total serum and LDL-cholesterol, apolipoprotein B, thromboxane A₂, platelet factor 4, triglycerides and glucose and inhibiting the activity
30 of HMG-CoA reductase. They also reduce the levels of TNF and IL-1, and possibly IL-2 and gamma interferon, while increasing antibody titers in response to foreign proteins. In combination, these factors may produce a variety of advantageous and novel biological results,
35 including a decrease in the release of superoxide and

other cytotoxins produced by immunoregulatory cells and an increase in antibody titers.

Tocotrienols and tocotrienol-like compounds according to this invention are preferably recovered
5 from stabilized biological sources. Alternatively, tocotrienols and tocotrienol-like compounds may be recovered from biological sources subjected to conventional food processing or preparation techniques. The stabilization process, however, greatly increases
10 the recovered yields of these compounds. Without wishing to be bound by theory, we believe that a significant amount of tocotrienols and tocotrienol-like compounds are, in nature, bound to proteins or linked to phosphate or phospholipid groups in the
15 membranes of biological sources. The significant increase in recovered amounts of these compounds realized by stabilization may be attributed to their release from protein or cleavage of phosphate or pyrophosphate moieties attached to the hydroxy group of
20 the benzene ring in the tocotrienol or the hydrogen donor group of the tocotrienol-like compound. The application of heat and, optionally, pressure advantageously releases the compounds of this invention so that they may be recovered in good yield.
25 Preferably, a combination of heat and pressure are used. Depending on the method of recovery used, the compounds of this invention may be extracted together with other components of the TRF.

We have found that microwaving is
30 particularly effective to release membrane-bound T_3 and T_3 -like compounds. We have also found that the following protocol maximizes the desired stabilization results: An amount (typically from 1 g to 1 kg) of a ground biological source (preferably rice or rice bran)
35 is placed in a pyrex dish (typically, 10 cm - 20 cm in diameter and 1.5 cm - 15 cm in height). The dish can

be covered or uncovered and optionally, a second dish containing a ground biological source can be stacked above the first. We prefer using one covered dish at a time. The biological source is then heated in a
5 microwave oven at the maximum level (preferably 600-1500 watts) for 1-5 minute intervals, optionally in an ambient, vacuum or nitrogen blanket environment. Pressures of greater than 1 atmosphere may also optionally be used. After each heating interval, the
10 sample should be stirred to homogeneity. The heating and stirring process is repeated 1-10 times. We prefer repeating the process 3-5 time for a 0.5 kg sample. The TRF may then be extracted, as described herein.

The TRF may be used directly, or it may be
15 further separated in to its component compounds. It may be desired to isolate the novel tocotrienols and tocotrienol-like compounds from the TRF mixture.

The TRF typically contains varying degrees of each of the known tocotrienols, plus additional Tocol
20 products. Theoretically, each novel tocotrienol and tocotrienol-like compound should be separable from the known tocopherols and tocotrienols in the TRF using standard silica gel HPLC methodology. However, in the case of rice bran, a complex mixture results. This
25 resultant mixture contained 3,4-dihydro-2-methyl-2-(4,8,12-trimethyltrideca-3'(E),7'(E),11'-trienyl)-2H-1-benzopyran-6-ol), a known tocotrienol that eluted by HPLC after about 21 minutes under the specified conditions. For that reason, we refer to this compound
30 as "P₂₁". In addition, the mixture contained 3,4-dihydro-2-(4,8,12-trimethyltrideca-3'(E), 7'(E),11'-trienyl)-2H-1-benzopyran-6-ol), or "P₂₅", a novel tocotrienol that eluted by HPLC after about 25 minutes. And the mixture contained an apparent sterol which had
35 a UV absorption maximum at 315 nm that eluted by HPLC

after about 20 minutes ("P₂₀"). These three compounds cannot be separated using standard silica gel HPLC techniques.

Accordingly, the present invention includes
5 novel techniques to purify tocotrienols and tocotrienol-like compounds from contaminants that cannot be removed using a silica column. These techniques are especially well suited for separating tocopherols and tocotrienol-like compounds from sterol
10 and other waxy contaminants. In one process, the TRF is dissolved in an appropriate solvent, preferably hexane, and then bound to an amine or cyano column (1 ml). We prefer to use a Bond Elute amine or cyano column. The tocotrienols and tocotrienol-like
15 compounds of this invention may then be selectively eluted from the column by an appropriate solvent system. Preferably, this solvent system is a gradient of isopropanol in hexane. Using a small concentration of the polar solvent in the non-polar solvent
20 (preferably about 0.5% isopropanol in hexane) known tocotrienols and tocopherols may be eluted, while the novel tocotrienols and tocotrienol-like compounds of this invention are retained on the amine column. Then, by increasing the concentration of the more polar
25 solvent (preferably to about 3% isopropanol in hexane), the tocotrienols and tocotrienol-like compounds of this invention may be selectively eluted, while any sterol and waxy contaminants remain bound to the amine column. Finally, these contaminants may be eluted using a high
30 polarity solvent system, such as 6-10% isopropanol in hexane. Using this procedure, the desirable tocotrienols and tocotrienol-like compounds may be effectively separated.

An alternative technique also increases the
35 recovered yield of tocotrienols and tocotrienol-like

compounds from a biological source. In this alternative procedure, a biological source, stabilized or unstabilized, is first extracted with methanol. This step removes many of the unbound contaminants.

5 Then, heat and, optionally, pressure, are applied to the biological source to release the tocotrienols and tocotrienol-like compounds. Examples of heat sources include, but are not limited to radiant heat sources (i.e., visible light or radioactive materials),

10 convectional heat sources, microwaves, radio-frequency, friction or shearing. The preferred heat source is microwave heat. Alternatively, freeze/thaw methods or mechanical grinding might also be employed before, during or after heat and/or pressure treatment.

15 Alternatively, the tocotrienols and tocotrienol-like compounds may be released by the use of caustic agents, for example, acids, such as hydrochloric or sulfuric acid. According to an alternate embodiment, sonication or detergent treatment may be employed prior to,

20 concurrently with, or following the heat, pressure or caustic agent treatment. Experiments using the techniques described herein that employ various combinations of heat, pressure and reaction conditions will readily indicate the preferred conditions for a

25 given biological source.

In purified form, one of the preferred compounds of the present invention, P_{25} , exhibits greater biological activity as a cholesterol-lowering agent than any of the known tocotrienols: including

30 P_{21} (known also as "tocotrienol"), α - T_3 , β - T_3 , γ - T_3 or δ - T_3 . We believe that the double bonds on the isoprenoid side chain comprise the active portion of these compounds because their presence decreases the effect of London forces between the lipid side chain in

35 biological membranes. We also believe that the number

and the position of the alkyl substituents on the aromatic ring system and the isoprenoid side chain make a difference in the biological activity -- the fewer the number of alkyl substituents, the greater the activity. We believe that reducing the steric hindrance caused by the methyl groups allows P_{25} to penetrate more deeply into the membrane. Accordingly, the membrane may become more highly organized and thus, less permeable. Under this hypothesis, the compounds of this invention which substitute hydrogen for methyl at the 2-position of the chromanol ring demonstrate increased biological activity. However, alterations at that and other positions are also envisioned, as several different mechanisms which result in increased biological activity may be involved.

Because of their ability to lower total serum cholesterol and low density lipoprotein-cholesterol and increase the HDL-/LDL-cholesterol ratio, the novel tocotrienols and tocotrienol-like compounds of this invention may be used in the prevention and treatment of diseases associated with high levels of cholesterol.

Advantageously, the compounds of this invention do not substantially alter the serum levels of other blood components that contribute to the biodegradation of cholesterol. For example, the compounds of this invention do not substantially reduce the activity of cholesterol 7 α -hydroxylase -- the enzyme that is responsible for degradation of cholesterol into bile acids.

Examples of diseases associated with high levels of cholesterol that may be treated by the compounds of this invention include, but are not limited to, atherosclerosis, thrombosis, coronary artery disease and other forms of cardiovascular disease. In addition, the ability of the compounds of

this invention to lower serum glucose levels may increase the insulin production in Type 2 diabetics.

The compounds of this invention may also be used to alter the serum or plasma levels of several
5 other blood constituents. For example, these compounds lower the plasma levels of thromboxane A₂ and platelet factor 4. In addition, they may serve passively as simple antioxidants or actively by decreasing the release of superoxides by neutrophils and other
10 cytotoxins or cytokines, mast cells, macrophages, endothelial tissue and other immunoregulatory tissues. Antioxidation is accomplished in at least two ways. First, by reducing arachidonic acid metabolites, the neutrophils reduce the levels of superoxide production.
15 Second, these compounds scavenge radicals which are already present. Accordingly, they exert a protective effect on the endothelium, lipoproteins, smooth muscle cells and platelets. In addition, these compounds may also serve as antioxidants to prolong the shelf lives
20 of products prone to oxidation, such as food products. Advantageously, the tocotrienols and tocotrienol-like compounds of this invention may also prevent oxidative degradation of food products which results in the formation of carcinogenic compounds in such food
25 products.

The thromboxanes (whose plasma levels are decreased using the compounds of this invention) also induce platelet aggregation and vasoconstriction. Therefore, the tocotrienols and tocotrienol-like
30 compounds of this invention may be used to reduce blood clotting in a wide variety of applications. For example, these compounds may be used to treat or prevent diseases, such as thrombotic diseases, cardiovascular diseases, hypertension, pulmonary
35 diseases and renal diseases. Specifically, the

compounds of this invention may be used to prevent or reverse blood clots and lesions which may cause diseases such as myocardial infarction, stroke, pulmonary embolism, deep vein thrombosis, peripheral
5 arterial occlusion and other blood system thromboses.

Tocotrienols and tocotrienol-like compounds are also capable of acting as antiatherogenic agents by inhibiting or reversing the oxidation of LDL and by protecting vascular tissue in general from oxidative
10 damages. The oxidated form of LDL ("OX-LDL") is a major component in the formation of atheroma. Such formation commonly results in a narrowing of the arteries by atherosclerosis plaques. While not wishing to be bound by theory, we believe that by reducing serum LDL
15 levels, tocotrienols and tocotrienol-like compounds enhance the rate of metabolic LDL turnover and therefore, decrease the exposure of LDL to oxidative agents.

Tocotrienols and tocotrienol-like compounds
20 also decrease the concentration of lipoprotein (a) in the blood. It has been well established that elevated concentrations of Lp(a) are correlated with early onset and progression of atherosclerosis, premature myocardial ischemia and rheumatoid arthritis. In fact,
25 Lp(a) concentration is a more accurate indicator of coronary heart disease than LDL concentration. Lowering Lp(a) concentrations in individuals having high Lp(a) levels (above about 20 mg/dl) drastically reduces the probability of atherosclerosis and coronary
30 heart disease. Significantly, Lp(a) levels, unlike LDL concentration, are not affected by low-fat diets or commercial available hypocholesterolemic agents.

Tocotrienols and tocotrienol-like compounds also exhibit diuretic activity -- they are antagonistic
35 to vasopressin and angiotensin II. Accordingly, these

compounds are useful in the treatment and management, for example, of hypertension.

According to another embodiment of this invention, tocotrienol or tocotrienol-like compounds
5 may be administered pre-operatively to a patient in order to prevent septic shock. Advantageously, tocotrienols and tocotrienol-like compounds may be used in the treatment of extracorporeal blood. As used
10 herein, the term "extracorporeal blood" includes blood removed in line from a patient subjected to extracorporeal treatment, and returned to the patient in processes such as dialysis, or blood filtration or bypass timing surgery. And the term includes blood
15 products which are stored extracorporeally for eventual administration to a patient. Such products include whole blood, platelet concentrates and any other blood fraction in which inhibition of platelet aggregation is desired.

According to another embodiment of this
20 invention, tocotrienols and tocotrienol-like compounds may be formulated in compositions and methods for coating the surface of invasive devices, to lower the risk of platelet aggregation -- for example, the
25 surfaces of devices such as, vascular grafts, stents, catheters and artificial valves. Such devices may be coated with the tocotrienols and tocotrienol-like compounds using conventional methodologies including physical adsorption and chemical cross-linking.

Furthermore, we have found that the
30 tocotrienols and tocotrienol-like compounds of this invention, the known tocotrienols, and mixtures thereof also reduce the levels of tumor necrosis factor in response to lipopolysaccharide stimulation, lower
35 arachidonic acid in the tissues and reduce oxygen metabolites in the blood of animals and humans. These

results point to an overall reduction in prostaglandins and leukotrienes, both of which are synthesized from arachidonic acid, and a possible reduction in interleukin-1. Accordingly, the compounds of this invention may be employed for a variety of uses. For example, they may be used to prevent endothelial injury, such as ischemic and reperfused myocardium and ulcers. In addition, the inhibition of tumor necrosis factor biosynthesis would also be accompanied by a decrease in inflammation -- i.e., through inhibiting the respiratory bursts of neutrophils or through free radical scavenging. Therefore, the compounds of this invention and the known tocotrienols are also useful as antiinflammatory agents for the prevention and treatment of a wide variety of diseases and conditions involving minor, acute and chronic inflammation. These include, but are not limited to, fever, rheumatoid diseases, pain, functio laesa, hypertension and edema.

In addition to their role in inflammatory response, prostaglandins have also been shown to inhibit glucose-induced insulin release, increase glucose concentration and stimulate glucagon secretion. Consequently, use of the compounds of this invention typically leads to an increased insulin to glucagon ratio. Therefore, the novel tocotrienols and tocotrienol-like compounds of this invention, the known tocotrienols, and mixtures thereof, may be used to improve glucose intolerance in diabetes mellitus. They may also be used to restore acute glucose-induced insulin response in non-insulin-dependent diabetes mellitus.

In addition to the above-stated uses, the tocotrienols and tocotrienol-like compounds of this invention, the known tocotrienols, and mixtures thereof, may also be used to enhance the immune

response in animals and humans. These compounds typically reduce the amount of fatty acids in biological tissues. Since fatty acid levels effect the immune system, the compounds of this invention may
5 serve as immunoregulators. They may, for example, be used to increase antibody titers to foreign proteins.

In addition, the reductions in fatty acid, cholesterol, triglyceride and glucose levels effected by the compounds of this invention are obtained without
10 attendant substantial weight loss. The result is an increased feed to protein conversion ratio. Therefore, the novel tocotrienols and tocotrienol-like compounds of this invention, the known tocotrienols, and mixtures thereof, are useful in increasing feed conversion
15 efficiency.

Hypercholesterolemic diseases and conditions that may be treated using the compositions and mixtures described herein include, but are not limited to, arteriosclerosis, atherosclerosis, xanthomatosis,
20 hyperlipoproteinemias, and familial hypercholesterolemia.

Thrombotic diseases and conditions that may be treated using such compositions include, but are not limited to, pulmonary disease, in general (such as
25 reduced specific conductance, reduced dynamic compliance and constriction (contraction of smooth muscle), excess pulmonary fluids (such as pulmonary lymph, foam, or bronchoalveolar lavage), adult respiratory distress syndrome, astatis and rhinitic
30 disease (such as pulmonary and systemic hypertension, pulmonary edema, fluid accumulation (neutrophil infiltration) and pulmonary vascular permeability), pulmonary vasoconstriction (associated, for example, with endotoxemia, gram-negative organisms, anaphylaxis,
35 hemorrhagic shock or allergy to ragweed), cardiac

ischemia, microembolic and/or frank occlusion, reocclusion following transluminal angioplasty, myocardial infarction, cardiopulmonary bypass associated dysfunction, vasoconstriction (pulmonary and peripheral), organ dysfunction, platelet consumption and/or activation (and subsequent decreased function, aggregation and decreased numbers), mitral valve pathology associated with acute perioperative pulmonary hypertension, chronic obstructive arterial disease caused by arteriosclerosis, Maurice Raynaud's syndrome -- vasoconstriction, renal artery stenosis, myocardial infarction, stroke, pulmonary embolism, deep vein thrombosis, peripheral arterial occlusion and other blood system thromboses.

Antioxidizing uses include, but are not limited to, the treatment and prevention of endothelial injury, such as ischemic and reperfused myocardium. Because of their antioxidantizing activity, the tocotrienols and tocotrienol-like compounds of this invention may also be used in treating and preventing cancerous conditions (by, for example, preventing cancer-causing mutations in the genetic material of an animal or a human).

Antiatherogenic diseases and conditions that may be treated using such compositions include, but are not limited to, arteriosclerosis, atherosclerosis, myocardial infarction, ischemia (i.e., myocardial ischemia, brain ischemia and renal ischemia) and strokes.

Inflammatory diseases and conditions that may be treated using such compositions include, but are not limited to, essential hypertension, hypertension of congestive heart failure, renal dysfunction caused by reduced myocardial output, endotoxemia, chronic liver disease or hypertension, pulmonary inflammation in

asthma, bronchitic, pneumonia or acute lung injury, rheumatic diseases (such as rheumatoid arthritis and systemic lupus erythematosus), inflammatory bowel disease (such as ulcerative colitis), irritable bowel
5 disease (such as villous adenoma), gastrointestinal disorders caused by excess acids, pepsin or bile salts, Zollinger-Ellison syndrome, skin diseases or trauma (such as burns or acid or caustic injury), gout, Bartter's syndrome, fever, rheumatoid diseases, pain,
10 functio laesa, hypertension and edema.

Immunoregulatory diseases and diseases that may be treated using the compositions of this invention include, but are not limited to, chronic fatigue syndrome, graft rejections, autoimmune diseases, such
15 as AIDS, and other viral diseases that weaken the immune system.

The compounds and mixtures described herein are useful in pharmaceutical compositions, foodstuffs and dietary supplements. Advantageously, these
20 products are hypocholesterolemic, antithrombotic, antioxidizing, antiatherogenic, antiinflammatory and immunoregulatory agents.

Pharmaceutical compositions may take the form of tablets, capsules, emulsions, suspensions and
25 powders for oral administration, sterile solutions or emulsions for parenteral administration, sterile solutions for intravenous administration and gels, lotions and cremes for topical application. The pharmaceutical compositions may be administered to
30 humans and animals in a safe and pharmaceutically effective amount to elicit any of the desired results indicated for the compounds and mixtures described herein.

This invention also relates to prodrug forms
35 of tocotrienol and tocotrienol-like compounds which,

upon administration to a patient, undergo biotransformation into active form.

The pharmaceutical compositions of this invention typically comprise a pharmaceutically effective amount of a tocotrienol or tocotrienol-like compound of this invention, or a mixture thereof, and a pharmaceutically acceptable carrier. Such carriers may be solid or liquid, such as, for example, cornstarch, lactose, sucrose, olive oil or sesame oil. If a solid carrier is used, the dosage forms may be tablets, capsules or lozenges. Liquid dosage forms include soft gelatin capsules, syrup or liquid suspension.

Therapeutic and prophylactic methods of this invention comprise the step of treating patients in a pharmaceutically acceptable manner with the compositions and mixtures described herein. As used in this application, the term "pharmaceutically effective amount" or "cholesterol-lowering amount" refers to an amount effective to lower blood levels of LDL-cholesterol and total serum cholesterol, while increasing the ratio of HDL-cholesterol to LDL-cholesterol in the blood. Alternatively, the term "pharmaceutically effective amount" refers to an amount effective to decrease blood levels of LDL-cholesterol and total serum cholesterol associated with hypercholesterolemia, an amount effective for lipogenesis, an amount effective to inhibit platelet aggregation, an amount effective to decrease the release of superoxides by human peripheral blood neutrophils, an amount effective to reduce the level of tumor necrosis factor or interleukin-1, an amount effective to reduce the level of arachadonic acid, an amount effective to increase antibody titers in the blood, an amount effective for antithrombotic uses, an amount effective to treat, prevent or delay the onset

of any one of the following diseases or conditions including inflammatory diseases, immunoregulatory diseases, fever, edema, diabetes mellitus, cancer, signs of aging, pain, septic shock, chronic fatigue syndrome and functio laesa; or an amount effective to decrease the concentration of Lp(a) in the blood or to increase feed conversion efficiency.

The pharmaceutical compositions of this invention may be employed in a conventional manner for the treatment and prevention of any of the aforementioned diseases and conditions. Such methods of treatment and prophylaxis and their dosage levels and requirements are well-recognized in the art and may be chosen by those of ordinary skill in the art from the available methods and techniques. Dosage ranges may be from about 1 to about 1000 mg/day. However, lower or higher dosages may be employed. Specific dosage and treatment regimens will depend upon factors such as the patient's health status, the severity and course of the patient's disease or disposition thereto and the judgment of the treating physician.

Tocotrienols and tocotrienol-like compounds and mixtures thereof may also be used in combination with conventional therapeutics used in the treatment or prophylaxis of any of the aforementioned diseases. Such combination therapies advantageously utilize lower dosages of those conventional therapeutics, thus avoiding possible toxicity incurred when those agents are used as monotherapies. For example, tocotrienols or tocotrienol-like compounds may be used in combination with bile acid sequestrants, such as Cholestyramine and Colestipol; fibric acid derivatives, such as, Clofibrate, Gemfibrozil, Bezafibrate, Fenofibrate, and Ciprofibrate; HMGCR inhibitors, such as Lovastatin, Mavastatin, Pravastatin, Simvastatin and

SRI-62320; Probucol; Nicotinic Acid; its derivatives and conjugates, such as, 6-OH-Nicotinic Acid, Nicotinaric Acid, Nicotinamide, Nicotinamide-N-oxide, 6-OH-Nicotinamide, NAD, N-Methyl-2-pyridine-8-
5 carboxamide, N-Methyl-Nicotinamide, N-Ribosyl-2-Pyridone-8-Carboxamide, N-Methyl-4-pyridone-5-carboxamide, Bradillian, Niceritrol, Sorbinic acid and Hexanoic; Neomycin and d-Thyroxine.

In foodstuffs, tocotrienols and tocotrienol-
10 like compounds, and mixtures thereof, may be used with any biologically acceptable carrier to provide safe and effective means of lowering blood levels of LDL-cholesterol and total serum cholesterol, while increasing the ratio of HDL-cholesterol to LDL-
15 cholesterol in the blood. In addition, the foodstuffs may be used to inhibit platelet aggregation, to decrease the release of superoxides by human peripheral blood neutrophils, to reduce the levels of tumor necrosis factor and interleukin-1, to increase antibody
20 titers in the blood, or to treat, prevent or delay the onset of one or more of the following diseases or conditions: immunoregulatory disease, inflammation, fever, edema, diabetes mellitus, cancer, signs of aging, pain, rheumatoid diseases, septic shock, chronic
25 fatigue syndrome and functio laesa. Alternatively, they may be used to decrease Lp(a) concentrations in the blood or to increase feed conversion efficiency.

Foodstuffs containing the tocotrienols and tocotrienol-like compounds according to this invention,
30 and mixtures thereof, may be combined with any other foodstuff. For example, oils containing the compounds of this invention may be used as cooking oil, frying oil, or salad oil and may be used in any oil-based food, such as margarine, mayonnaise or peanut butter.
35 Grain flour fortified with the compounds of this

invention may be used in foodstuffs, such as baked goods, cereals, pastas and soups. Oils containing tocotrienols and tocotrienol-like compounds can be emulsified and used in a variety of water-based foodstuffs, such as drinks, including drink mixes. Advantageously, such foodstuffs may be included in low fat, low cholesterol or otherwise restricted dietary regimens.

The pharmaceutical compositions and foodstuffs of this invention may be administered to humans and animals such as, for example, livestock and poultry. Once an animal has ingested or otherwise been administered a tocotrienol or tocotrienol-like compound, or a mixture thereof, it advantageously retains the hypercholesterolemic, antithrombotic, antioxidizing, antiinflammatory, antiatherogenic, immunoregulatory and other advantageous biological activities of the administered compounds. Therefore, such an animal, or any product derived therefrom, such as, for example, milk, may be consumed by a human or another animal to derive the benefits of the tocotrienols and tocotrienol-like compounds. For example, a chicken which ingests a grain or meal fortified with the compounds of this invention may later be eaten by a human to derive the cholesterol-reducing benefits.

In addition, the administration of tocotrienols or tocotrienol-like compounds to animals results in an increase in feed conversion efficiency. In higher fat content animals, such as cattle, swine, sheep and lamb, tocotrienols or tocotrienol-like compounds advantageously lead to faster growth, lower cholesterol content, and higher percentage lean meat. When administered to fowl, tocotrienols and tocotrienol-like compounds result in production of eggs

characterized by reduced cholesterol content of the yolk and higher protein content of the egg white.

In order that this invention be more fully understood, the following examples are set forth.

- 5 These examples are for purposes of illustration only, and are not to be construed as limiting the scope of the invention in any way.

In these examples, mass spectra data was acquired using Model MS-902 manufactured by and
10 obtainable from Associate Electrical Industries, Ltd. (Department of Chemistry, University of Wisconsin, Madison). Samples were introduced directly into the ion source on a glass probe at 100°C. The potential of the ionizing electrical beam was 70 e.v. NMR in CDCl_3
15 were obtained using NMR-500 by Bruker, FDR at room temperature. IR was done as thin film using a Perkin Elmer instrument. Centrifugation was done in an Eppendorf Centrifuge (model 1240). Microwaving was done in a Whirlpool household oven (Model No.
20 MT6901XW-0 operating at 800 watts. HPLC was done using a Waters pump 6000A or a Gilson pump 307, a Shimadzu fluorescence detector RF-535 and integrator CR3A, a Gilson's autoinjector, Perkin Elmer 50A UV detector or a Shimadzu SPD-10 AV UV-VIS spectrum photometric
25 detector and a Kipp and Zonen BD41 recorder. Unless otherwise indicated, the following HPLC conditions were employed: 20 μl injections, Porasil column (Waters column, 10 μ , 4 mm x 30 cm) using an isocratic system of hexane and isopropanol (99.75%:0.25%, v/v) at a flow
30 rate of 1.3 ml/min. Detection was done at an excitation wavelength of 295 nm and an emission wavelength of 330 nm (fluorescence detector) and UV absorption at 295 nm.

All assays conducted on chicken or swine serum were done following the protocols described in
35 A.A. Qureshi et al., "Lowering Of Serum Cholesterol In

Hypercholesterolemic Humans By Tocotrienols"

(Palmvitee), Am. J. Clin. Nutr., 53, pp. 1021S-26S

(1991). All enzymatic assays were done following the protocols described in A.A. Qureshi et al., "Effects of

5 Cereals and Culture Filtrate of *Trichoderma viride* on Lipid Metabolism of Swine", Lipids, 17, pp. 924 (1982).

Plasma insulin, glucagon and TNF levels were measured using radioimmunoassay kits from Ventrex Laboratories, Inc. (Portland, Maine), ICN Biomedicals, Inc. (Costa

10 Mesa, California) and Genzyme Corp. (Cambridge, Massachusetts), respectively.

Example 1

The following protocol may be employed for carrying out the stabilization of a biological source,
15 using rice bran as the sample.

Rough rice (paddy rice) from a farm is dried in a commercial-type continuous flow, non-mixing, heated air dryer. Drying is carried out to lower the moisture content of the rice from a level of between
20 about 18 and 22 percent to a level between about 10 and 13 percent. The dried rice is then cleaned by removing dust, stones, seeds and sticks by aspiration in a commercial rice cleaning machine, followed by gravity separation in a stoner and particle size separation in
25 a disk grader and a drum separator. The husks are then removed using a rubber roller sheller. Paddy (husks or hulls) are removed using a paddy separator. The raw bran is then removed in a friction mill to yield polished rice. The raw bran is then pneumatically
30 conveyed either to the extruder or to storage until stabilization.

To carry out stabilization, the raw bran is pneumatically conveyed to a filter/sifter to remove residual broken rice. After sifting, the raw bran is

pneumatically conveyed to a mixing/tempering hopper tank. The raw bran is conveyed from the discharge of the mixing tank to the extruder inlet valve of a clamped barrel single screw extruder by a metered screw
5 conveyer feeder. The operating conditions of the extruder are maintained during stabilization in the following range:

	flow rate:	900-2000 lbs/hr
	pressure:	800-2000 PSI
10	temperature:	135°-210°C
	time:	5-90 seconds.

The dry heat stabilized raw bran is then fed directly into the feed hopper of an expander cooker. Alternatively, the feed to the cooker may be raw bran
15 that has been cooled with dry ice. Within the extruder, the bran is conveyed by a discontinuous worm shaft toward the discharge plate at a rate of around 341 lbs/hr. Water and steam are added through injection ports in the barrel of the extruder at a rate
20 of around 38 lbs/hr to completely mix the material and to raise the moisture level. The ambient temperature is about 30°C. Flow of the material is controlled by a discharge die plate at a rate of about 341 lbs/hr. The feed material moisture level is maintained at about
25 11.4% and the temperature is held between about 90° and 135°C for between about 15 and 90 seconds. The moisture level in the collet produced is about 14% at a discharge temperature of 125°C.

As the bran is extruded through the die
30 plate, the sudden decrease in pressure causes the liquid water to vaporize. During cooking, enzymes are denatured and some constituents of the bran are gelatinized into a fluid paste which binds the particles together. A compact pellet is formed.
35 Vaporization of water causes breakage within the cells

ideally suited for solvent migration percolation. The introduction of steam and water during the process raises the moisture content of the bran to about 22-25 percent. The extruder discharge is then sent
5 downstream at a rate of around 341 lbs/hr. to a dryer/cooler. Moisture flow to the extruder is maintained at about 96 lbs/hr and the temperature is kept in the range of 82°C to 130°C. The discharge from the dryer/cooler is maintained at a rate of about 341
10 lbs/hr and at a moisture level of about 8%. Stabilized rice bran is the result. These conditions also allow for storage of the stabilized bran.

The stabilized bran is immersed in hexane in a ratio by weight of about two to one. Typically,
15 about 10-100 g of material can be extracted using this protocol. The hexane is generally heated to about 60°C using a steam table incorporated into an explosion proof vented hood, but other solvents and other temperatures may also be employed. The hexane/oil
20 miscella is removed from the bran by filtration. About 5-6 washings are necessary to bring the oil content of the bran to less than one percent. The defatted bran and the hexane/oil miscella are both desolventized under gentle heating with steam.

25 If 100-500 lbs. of stabilized bran is to be extracted, it is more practical to use the following protocol. The stabilized bran is fed into a counter-current extractor at a flow rate of about 111 lbs/hour. Fresh hexane is introduced at a rate of around 312
30 lbs/hr. The fresh solvent (hexane) temperature is maintained at about 50°C, while the extractor temperature is maintained at around 52°C. Residence time in the extractor is typically around 45 minutes. The product is a defatted bran with an oil content of
35 less than one percent. The hexane/oil miscella exiting

the discharge of the extractor is filtered through a plate and frame filter press. The filtered miscella is then pumped to a steam heated still where the hexane is evaporated and collected by a condenser for reuse.

5 Following extraction and desolventization, the crude rice bran oil is typically degummed, dewaxed, bleached and physically refined using steam distillation. Degumming is carried out by a two stage addition under agitation of 2% water by weight and then
10 0.15% phosphoric acid (85% reagent grade) by weight. The temperature is held at about 82°C to 88°C for 10 minutes and the sludge containing the gums is then removed via ultrafugation. (See, e.g., United States Patent No. 4,049,686). The degummed bran is cooled to
15 about 5°C to 8°C and held for 24 hours. The dewaxed oils form a layer above the waxes which can be decanted using a vacuum pump. Bleaching is carried out according to the official American Oil Chemist's Society method 6c 8a-52. Physical refining is carried
20 out in a glass deodorizer at about 250°C and around 3 mm Hg for about 2 hours.

The following specific protocols are referred to in the subsequent examples. Stabilization protocols follow the general method set forth above, with the
25 precise conditions defined as follows:

Protocol I- Dry Heat Stabilization

Extruder: Wenger Model X-25

Standard Screw/Barrel Setup:

	<u>Barrel #</u>	<u>Standard Port</u>	<u>Screw #</u>	<u>Standard Port</u>
30	5	28714-9	5	28320-1
	4	28318-1	4	8326-9
	3	28372-9	3	28326-1

2	28318-1	2	28326-5
1	28350-1	1	28387-9

Standard Die Setup:

	<u>Die/Spacer</u>	<u>Measurement</u>	<u>Standard Port</u>
5	Spacer	0.375	28340-11
	Back Plate	0.625	28361-51
	Intermediate Plate	0.218	28316-723
	Front Plate	0.235	28389-507

10 Operating Conditions:

Feed Rate: 1000 lbs/hr
Temperature: 170°C at exit die
Pressure: 975-1025 psi
Moisture Feed: 12%
15 Moisture Discharge: 9.6%
Residence Time: 15 seconds
Run Duration: 8 hours
Sample Size: 50 lbs.

Protocol II-Dry Heat Followed by Wet Heat Stabilization

20 Dry Heat Stage: Protocol I

Wet Heat Stage:

Extruder: Anderson 4 inch

Screw/Barrel Configuration: Standard Cut Flight

Die Setup:

25 Diameter: 0.1875 inches
Land: 0.75 inches

Operating Conditions:

30 Feed Rate: 378 lbs/hr
Shaft Speed: 279 rpm
Steam injection: 36 lbs/hr (32 psi at #8 hole)
Mechanical Pressure: 750 psi (est.)
Moisture Feed: 11.4%

Discharge Moisture: 15%
Discharge Rate: 450 lbs/hr
Discharge Temp.: 121°C.

Protocol III-Drying/Cooling Procedure

5 The wet heat stabilized product of protocol
II (15% moisture) was discharged onto aluminum trays
and placed in a tray oven at 101.1°C until the moisture
content was 8-10% (approximately 1.5 hours). The trays
were then placed on tray racks and allowed to cool at
10 ambient temperature (approximately 20°C).

Protocol IV-Oil Extraction (Laboratory Method)

Oil to hexane ratio: 1 : 4
of washings: 3
Extraction temperature: 40°C

15 The hexane was removed from the extract by
mild heating (40°C) under a mild vacuum.

Protocol V-Oil Extraction (Pilot Plant Method)

Oil to hexane ratio: 1 : 4
of washings: 6
20 Extraction temperature: 60°C
Amount Extracted: 20 lbs
Yield: 16 lbs defatted bran
4 lbs crude oil

The hexane was removed from the extract by
25 heating to 115.5°C.

Protocol VI-Oil Extraction (Cold Extraction)

Oil to hexane ratio: 1 : 4
of washings: 6
Extraction temperature: 20°C (ambient)
30 Amount Extracted: 20 lbs
Yield: 16.4 lbs defatted bran
3.6 lbs crude oil

The hexane was removed from the extract by
heating to 115.5°C.

Protocol VII-Dewaxing

20 lbs of crude oil were reffridgerated for 24 hours at 4°F (-15.6°C). The supernatant, which contained the dewaxed oil, was decanted from the solidified waxes. The waxes were then centrifuged to remove entrained oil yielding waxes (0.59 lbs) and dewaxed oil (19.407 lbs).

Example 2 -- Purification of P₂₅

The rice bran was stabilized according to protocol I. 1.0 g of the stabilized rice bran was ground into a fine powder and extracted with 8 ml of methanol in a disposable screw capped tube using a shaker for ~30 minutes. The suspension was then centrifuged at 2000 rpm for 10 min. to recover the methanol layer, which was evaporated under vacuum at 40°C. The remaining dried residue was then extracted with 4 ml hexane by shaking on a shaker for 3 min. followed by centrifugation at 2000 rpm for 5 min. The resulting supernatant was transferred into an injecting vial, which was capped and centrifuged again for 2 min. Evaporation yielded the TRF.

Ten mg of the TRF, which contained novel tocotrienol P₂₅, was dissolved in 0.5 ml of hexane and bound to a Bond Elute Amine column (1.0 ml) that was first equilibrated with 2 ml hexane. The column was then washed with 1 ml of hexane, followed by 1 ml of 3% isopropanol in hexane. This step removed most of the known Tocol products. Then 1.0 ml of 5% isopropanol in hexane was run through the column. The recovered solution was evaporated, yielding 1.85 mg of P₂₅ and 0.44 g of tocotrienol (P₂₁). Finally, 1.0 ml of 6% isopropanol in hexane followed by 1.0 ml 10% isopropanol in hexane was run through the column. The

resultant fractions were analyzed by HPLC. No trace of the P₂₀ impurity was visible on the HPLC trace.

Example 3

Amine column purified compounds from
5 Example 2 were further purified by HPLC to separate
tocotrienol (P₂₁) and P₂₅. The peak fractions from
several runs were collected and analyzed by HPLC. The
purified samples each showed a single symmetrical peak
as observed by UV and fluorescence. The following
10 analytical data was obtained for tocotrienol (P₂₁) and
P₂₅:

Tocotrienol P₂₁:

MS(EI): m/e 382⁺ (molecular ion)
15 163⁺
123⁺

NMR (CDCl₃):

δ 1.26 (s, H), 1.50-1.84 (m, 4H), 1.58
(s, 9H), 1.66 (s, 3H), 1.90-2.15 (m, 10H), 2.69 (t, J=
6.7 Hz, 2H), 4.23 (s, 1H), 5.07 (m, 3H), 6.51-6.57 (m,
20 2H), 6.63 (d, J= 9.5 Hz, 1H).

IR (film):

(cm⁻¹) 3400, 2990, 2950, 2876, 1500,
1458, 1240, 750.


The MS data correspond to a molecular formula
25 of C₂₆H₃₈O₂. The pattern is characteristic of the
fragmentation observed with α-, β-, and γ-tocotrienols
as reported by Rao et al., "Identification and
Estimation of Tocopherols and Tocotrienols in Vegetable
Oils Using Gas Chromatography-Mass Spectrometry", J.
30 Agr. Food Chem., 20(2), pp.240-245 (1972). The peak at
m/e 163⁺ indicates loss of the side chain (C₁₆H₂₇)⁺
giving rise to the ion C₁₀H₁₁O₂ (163.0759). The peak
m/e 123⁺ originates from the cleavage of the side chain

by the breakdown of the chroman structure with hydrogen rearrangement and loss of a methyl acetylene fragment ($\text{CH}_3\text{-C}\equiv\text{CH}_2$).

The NMR spectra is identical to that of T_3 , except that the methyl group on the benzene ring at δ 2.2 normally present in T_3 is missing in the spectrum of P_{21} .

The IR indicates the presence of a phenolic OH group at 3700 cm^{-1} , a benzene ring at 1650 cm^{-1} and a chroman ring at 1680 cm^{-1} .

Based on these data, the structure of P_{21} was identified as desmethyl-tocotrienol (3,4-dihydro-2-methyl-2-(4,8,12-trimethyltrideca-3'(E),7'(E),11'-trienyl)-2H-1-benzopyran-6-ol):



P_{25i}

MS(EI):

20

m/e	386 ⁺	(molecular ion)
	175 ⁺	
	149 ⁺	
	123 ⁺	

NMR (CDCl_3):

25

δ 1.58 (s, 3H), 1.59 (s, 3H), 1.62 (s, 3H), 1.67 (s, 3H), 1.73 (m, 2H), 1.90-2.08 (m, 10H), 2.19 (m, 2H), 2.63-2.85 (m, 2H), 3.89 (m, 1H), 4.30 (br. s, 1H), 5.07 (m, 2H), 5.14 (t, $J=7.9\text{ Hz}$, 1H), 6.51 (d, $J=2.9\text{ Hz}$, 1H), 6.56 (d of d, $J=2.9, 8.6\text{ Hz}$, 1H), 6.66 E(d, $J=8.6\text{ Hz}$, 1H).

30 IR (film):


(cm^{-1}) 3388, 2924, 1494, 1450, 1352, 1280, 1218, 1080.

The MS data correspond to a molecular formula of $\text{C}_{25}\text{H}_{36}\text{O}_2$. The pattern is characteristic of the fragmentation observed for γ -tocotrienol, except that the methyl on the chroman ring adjacent to oxygen is missing.

The NMR spectra is similar to that of desmethyl- T_3 .

10 The IR indicates the presence of a phenolic OH group at 3388 cm^{-1} , a benzene ring at 1494 cm^{-1} and a chroman ring at 1352 cm^{-1} .

Based on these data, the structure of P_{25} was identified as didesmethyl-tocotrienol (3,4-dihydro-2-(4,8,12-trimethyltrideca-3'(E),7'(E),11'-trienyl)-2H-15 1-benzopyran-6-ol):



Example 4 -- Isolation and Purification of P_{10} and P_{18}

The rice bran was stabilized according to protocol I. 1.0 g of the stabilized rice bran was ground into a fine power and microwaved for 3 minutes (in 1 minute intervals) to release any bound T_3 and T_3 -like compounds. The microwaved rice bran was then extracted with 7 ml of methanol in a disposable screw capped tube using a shaker for 30 minutes. The suspension was then centrifuged at 2000 rpm for about 5 minutes. The supernatant was removed and the methanol was evaporated under vacuum at 40°C . The resulting

residue contained a mixture of free and previously bound T_3 and T_3 -like compounds.

The residue was dissolved in 250 μ l (200 mg) of hexane and applied to a 1 ml Bond Elute amine column that was first equilibrated with 2 ml of hexane. The column was then washed with 1 ml of hexane. Then 0.5 ml aliquots of the following solvents were eluted through the column successively: 0.5% isopropanol in hexane, 1.0% isopropanol in hexane, 1.5% isopropanol in hexane, 2.0% isopropanol in hexane, 2.5% isopropanol in hexane and 3% isopropanol in hexane. Finally, the column was washed with 0.5 ml of 10% isopropanol in hexane. The resultant fractions were analyzed and the fractions containing P_{16} and P_{18} were combined (only the fractions collected using 1.5%-3% isopropanol in hexane contained P_{16} and P_{18} in significant amounts).

In order to separate P_{16} and P_{18} , HPLC conditions identical to those described in this application were used, except that the concentration of isopropanol was increased from 0.25% to 0.5% in hexane. Under those conditions, P_{16} eluted at 13 minutes and P_{18} eluted at 15 minutes. The peak fractions were collected and analyzed by HPLC. These purified samples each showed a single symmetrical peak as observed by UV and fluorescence. The following analytical data was obtained for P_{16} and P_{18} :


P_{16} :

MS(EI): m/e 380+ (molecular ion)
161+

molecular formula: $C_{27}H_{40}O$
molecular weight (actual): 380.3084
(calculated): 380.3079

The fragmentation pattern was similar to that of P_{21} and P_{25} . The peak at m/e 161+ indicates loss of the $C_{16}H_{27}$ side chain giving rise to $C_{11}H_{13}O$. Based on

these data, the following structure for P₁₈ was identified:




5

MS (EI): m/e 424+ (molecular ion)
205+

molecular formula: C₂₈H₄₀O₃
molecular weight (actual): 424.3010
(calculated: 424.2977)

10 The fragmentation pattern was again similar to that of P₂₁ and P₂₅. Based upon these data, the following structure for P₁₈ was identified:



Example 5

The effects of different rice brans on the serum levels of total cholesterol, HDL-cholesterol, 15 LDL-cholesterol, apolipoprotein (a)₁ and apolipoprotein B was determined in chickens.

The following protocols were performed to yield the samples displayed in Table I:

entry 5: protocol I

20 entry 6: protocol I, followed by protocol IV.

All other samples were obtained from commercially available sources.

25 Chickens were chosen as our initial model because they synthesize cholesterol in the liver using a mechanism similar to that of humans (unlike rats and mice that synthesize cholesterol in the intestines).

Each group of chickens (6-week old female white leghorn) was administered either a chick mash control diet or a control diet containing a 20% equivalent of a test rice bran. The amount of feed consumed by all groups was comparable to the control group and the feeding period was 4 weeks. The birds were fasted for a period of 14 hours prior to sacrifice (at 0800 hours).

The chicken mash diet contained the following ingredients:

	<u>Ingredients</u>	<u>Weight (g)</u>
	Corn (8.8% protein)	615.0
	Soybean Meal	335.0
	Corn Oil	10.0
15	Calcium Carbonate	10.0
	Dicalcium Phosphate	20.0
	Iodized Salt	5.0
	Mineral Mixture ^a	2.5
	Vitamin Mixture ^b	2.5

^a Mineral mixture contained per kg feed: zinc sulfate · H₂O, 110 mg; manganese sulfate 5H₂O, 70 mg; ferric citrate · H₂O, 500.0 mg; copper sulfate · 5H₂O, 16.0 mg; sodium selenite, 0.2 mg; DL-methionine, 2.5 g; choline chloride (50%), 1.5 g; ethoxyquin (1,2-dihydro-6-ethoxy-2,2,4-trimethylquinoline), 125 mg; and thiamine-HCl, 1.8 mg.

^b Vitamin mixture contained per kg feed: vitamin A, 1,500 units; vitamin D₃, 400 units; vitamin E, 10 units; riboflavin, 3.6 mg; calcium pantothenate, 10.0 mg; niacin, 25.0 mg; pyridoxine-HCl, 3.0 mg; folacin, 0.55 mg; biotin, 0.15 mg; vitamin B₁₂, 0.01 mg; and vitamin K₁, 0.55 mg.

The results are displayed below in Table I. Percentages of increases or decreases are shown in parentheses.

TABLE I

5	Nutritional state ¹	Concentration in Serum (mg/100/ml)				
		Total Chol.	HDL-Chol.	LDL-Chol.	Apo-A ₁	Apo-B
10	1) Control Group Chick Mash (CTL)	165.9±6.5 ^A (100.0)	104.2±4.65 ^A (100.0)	47.9±4.64 ^A (100.0)	132.9±3.88 ^A (100.0)	31.4±1.68 ^A (100.0)
	2) Cellulose (20% Equ.)	155.4±3.5 ^B (93.7)	96.0±4.99 ^A (92.1)	42.6±4.12 ^B (88.9)	126.8±3.81 ^A (95.4)	30.4±1.31 ^A (96.8)
	3) California Rice Bran (20% Equ.)	151.2±3.7 ^B (91.1)	100.4±3.42 ^A (96.4)	39.0±4.22 ^{A,B} (81.4)	125.8±4.78 ^A (94.7)	28.8±1.54 ^A (91.7)
	4) Louisiana Rice Bran (20% Equ.)	142.0±4.3 ^C (85.6)	101.7±6.41 ^A (97.6)	34.6±3.09 ^B (72.2)	132.0±3.84 ^A (99.3)	27.6±1.41 ^B (87.9)
15	5) Louisiana Rice Bran stabilized (20% Equ.)	128.5±4.0 ^D (77.5)	98.1±7.25 ^A (94.1)	31.4±2.72 ^B (65.6)	133.2±3.32 ^A (100.2)	25.7±2.25 ^B (81.8)
	6) Louisiana Rice Bran Defatted (20% Equ.)	148.5±6.1 ^D (89.5)	100.2±4.69 ^A (96.2)	42.4±3.01 ^A (88.5)	125.7±5.56 ^A (94.6)	28.7±1.31 ^A (91.4)
20	7) Commercial Rice Bran (20% Equ.)	150.0±4.6 ^B (90.4)	98.9±6.89 ^A (94.9)	40.6±3.88 ^A (84.8)	125.1±5.56 ^A (94.1)	29.0±1.26 ^A (92.4)

¹ Feeding period was 4 weeks; time of killing was 0800 hours. The birds were fasted for 14 hours prior to killing. Data expressed as means ± SD; n = 6 chickens per group.

² Percentages of increases or decreases are in parentheses.

25 A-D Values not sharing a common superscript letter are different at P < 0.01.

The diet which included stabilized rice bran demonstrated superior cholesterol reducing activity when compared with the other diets. It advantageously lowered the LDL-cholesterol level by almost 35%, while
30 lowering the HDL-cholesterol level by only 5.9%, as compared to the control diet. In addition, it increased the ratio of apolipoprotein (a)₁ to apolipoprotein B by 21%. This ratio is used as an indicator for assessment of risk for coronary heart
35 disease (see Naito et al., "The Clinical Significance

Of Apolipoprotein Measurements," J. Clin. Immunoassay,
9(1), pp. 11-20 (1986)).

Example 6

We next determined the effects of different
5 rice brans on the hepatic enzymatic activities of HMG-
CoA reductase, cholesterol 7 α -hydroxylase and fatty
acid synthetase, and the serum levels of triglycerides
and glucose in chickens fed with them. The feeding
conditions were identical to those described in
10 Example 5. The samples were prepared as described in
Example 5.

The results are displayed below in Table II.
Percentages of increases and decreases are shown in
parentheses.

TABLE II

Hepatic Enzymatic Activities						
Nutritional State ¹		HMG-CoA Reductase (pmoles/mg/min) ³	Cholesterol 7 α -Hydroxylase (nmoles/mg/min) ⁴	Fatty Acid Synthetase (nmoles/mg/min) ⁵	Serum Triglycerides (mg/100 ml)	Serum Glucose (mg/100 ml)
5	1) Control Group Chick Mash (CTL)	405.3 $\pm$ 15.5 ^A (100.0) ²	0.808 $\pm$ 0.062 ^A (100.0) ²	69.0 $\pm$ 3.40 ^A (100.0) ²	75.7 $\pm$ 3.34 ^A (100.0) ²	109.8 $\pm$ 3.50 ^A (100.0) ²
	2) Cellulose (20% Equ.)	417.0 $\pm$ 14.6 ^A (102.9)	0.882 $\pm$ 0.031 ^A (109.2)	58.7 $\pm$ 2.25 ^B (85.1)	69.4 $\pm$ 2.93 ^B (91.7)	100.9 $\pm$ 1.89 ^B (91.8)
10	3) California Rice Bran (20% Equ.)	340.2 $\pm$ 17.6 ^B (83.9)	0.851 $\pm$ 0.093 ^A (105.3)	55.1 $\pm$ 4.39 ^B (79.9)	65.4 $\pm$ 2.84 ^{B,C} (86.4)	96.1 $\pm$ 0.94 ^{A,B} (87.5)
	4) Louisiana Rice Bran (20% Equ.)	330.5 $\pm$ 12.8 ^B (81.5)	0.935 $\pm$ 0.044 ^B (115.7)	68.7 $\pm$ 3.88 ^A (99.6)	63.2 $\pm$ 1.61 ^C (83.5)	93.1 $\pm$ 2.20 ^C (84.8)
15	5) Louisiana Rice Bran Stabilized (20% Equ.)	311.7 $\pm$ 14.6 ^B (76.9)	0.939 $\pm$ 0.039 ^B (116.2)	65.6 $\pm$ 3.69 ^A (95.1)	63.9 $\pm$ 0.92 ^C (84.5)	90.4 $\pm$ 1.38 ^C (82.3)
	6) Louisiana Rice Bran Defatted (20% Equ.)	388.7 $\pm$ 14.8 ^C (95.9)	0.852 $\pm$ 0.028 ^A (105.4)	70.4 $\pm$ 2.12 ^A (102.0)	72.3 $\pm$ 1.67 ^A (95.6)	99.3 $\pm$ 1.63 ^A (90.4)
20	7) Commercial Rice Bran (20% Equ.)	371.2 $\pm$ 15.5 ^C (91.6)	0.850 $\pm$ 0.034 ^A (105.2)	64.7 $\pm$ 3.66 ^A (93.8)	70.9 $\pm$ 1.61 ^A (93.7)	101.2 $\pm$ 1.21 ^A (92.1)
¹ Feeding period was four weeks; time of killing was 0800 hours. The birds were fasted for 14 hours prior to killing. Data expressed as means $\pm$ SD; n = 6 chickens per group. ² Percentages of increases or decreases are in parentheses. ³ p-moles of mevalonic acid synthesized per minute per mg of microsomal protein. ⁴ n-moles of [¹⁴ C]-cholesterol into [¹⁴ C] 7 α -hydroxycholesterol per minute per mg of microsomal protein. ⁵ n-moles of NADPH oxidized per minute per mg of cytosolic protein.						

A-G Values not sharing a common superscript letter are different at P < 0.01.

As shown in Table II, the diet which contained stabilized rice bran produced a marked reduction (over 23% as compared to the control diet) in the activity of HMG-CoA reductase, the rate-limiting enzyme of cholesterol biosynthesis. However, the activity of the enzyme responsible for the degradation of cholesterol, cholesterol 7 α -hydroxylase, was not substantially affected. Furthermore, this diet reduced the serum levels of triglycerides (fats) and glucose by 15.5% and 17.7%, respectively.

Example 7

We next compared the effects of rice bran oils extracted from various sources on serum lipid parameter of chickens fed with diets supplemented with
5 those oils. Each group of 6 chickens (6-week old female white leghorn) was fed the chicken mash diet described in Example 5 for 14 days. Following this period, the chickens were fed a diet consisting of the chicken mash diet containing a 5% supplement of various
10 oils. The control diet included a supplement of 5% corn oil. After 4 weeks, the birds were fasted for 36 hours and then refed for 48 hours prior to sacrifice (at 0800 hours). The amount of feed consumed by all groups was comparable to the control group. The
15 following protocols were performed to yield the samples displayed in Table III:

- entry 2: protocol I, followed by protocol IV
- entry 3: protocol III, followed by protocol IV
- entry 4: protocol III, followed by protocol IV
- 20 entry 5: protocol IV
- entry 6: protocol IV

All other samples were obtained from commercially available sources.

The results are displayed below in Table III.
25 Percentages of increases or decreases are shown in parentheses.

TABLE III

		SERUM CHOLESTEROL (mg/100/ml)				
Nutritional State ¹		Total Chol.	HDL-Chol.	LDL-Chol.	Triglycerides	Glucose
5	1) Chick Diet + 5.0% Corn Oil (CDC)	185.1±5.1 ^A (100.0) ²	110.3±4.95 ^A (110.0) ²	61.9±4.49 ^A (100.0) ²	90.2±2.17 ^A (100.0) ²	124.6±2.30 ^A (100.0) ²
	2) Chick Diet + 5.0% Louisiana Rice Bran Oil (LRBO)	129.7±4.1 ^B (70.7)	99.8±3.57 ^A (90.5)	27.9±2.60 ^B (45.1)	84.5±2.97 ^B (93.7)	117.3±5.92 ^{A,B} (94.1)
10	3) Chick Diet + 5.0% Hot Extracted LRBO	135.3±4.5 ^B (73.1)	102.6±4.93 ^A (93.0)	30.0±2.89 ^B (48.5)	81.3±3.94 ^{B,C} (90.1)	112.4±4.65 ^B (90.2)
	4) Chick Diet + 5.0% Cold Extracted LRBO	121.6±3.5 ^C (65.7)	98.4±4.99 ^A (89.2)	26.1±2.89 ^B (42.2)	77.2±4.81 ^C (85.6)	113.2 ±4.40 ^B (90.9)
15	5) Chick Diet + 5.0% Louisiana Crude Rice Bran Oil Unstabilized	154.5±3.4 ^D (83.5)	108.6±3.25 ^A (98.5)	44.7±3.25 ^C (73.4)	86.6±4.64 ^{A,B} (96.0)	116.5±6.21 ^B (93.5)
	6) Chick Diet + 5.0% Commercial Rice Bran Oil	163.3±5.3 ^E (88.2)	102.4±4.10 ^A (92.8)	55.8±3.84 ^D (90.1)	88.5±4.33 ^A (98.1)	116.2±4.78 ^B (93.3)
20	¹ Feeding period was 4 weeks; time of killing was 0800 hours. The birds were fasted for 16 hours prior to killing. Data expressed as means ± SD; n = 6 chickens per group. ² Percentage of increases or decreases are in parentheses. A-E Values not sharing a common superscript letter are different at P < 0.01.					

The diet supplemented with 5% cold extracted Louisiana rice bran oil demonstrated superior cholesterol-lowering ability over the other diets. Impressive decreases in total serum and LDL-cholesterol (34.3% and 57.8%, respectively) were recorded, while the level of HDL-cholesterol decreased only slightly (10.8%). Decreases in triglycerides and glucose levels were also observed.

Example 8

This example demonstrates that waxes (sterols) were not responsible for the cholesterol-lowering properties of Louisiana rice bran and its oil. Each group of 6 chickens (6-week old female white leghorn) was administered the chick mash diet described

in Example 5, supplemented with one of three different amounts of waxes (sterols), 50 ppm of TRF (prepared as in Example 2) or 5% equivalent of the methanol insoluble fraction Louisiana rice bran oil. The amount
5 of feed consumed by all groups was comparable to the control group and the feeding period was 4 weeks. The birds were fasted for a period of 14 hours prior to sacrifice (at 0800 hours). The following protocols were performed on the samples displayed in Table IV:
10 waxes (entries 2-4): Protocol VII.

TRF (entry 5) was obtained using the method described in Example 2. The methanol insoluble fraction (entry 6) was obtained by freezing Louisiana rice bran oil (prepared using protocols I and IV), then
15 centrifuging. The resultant supernatant was extracted 3 times with twice the volume of methanol.

All other samples were obtained from commercially available sources.

The results are displayed below in Table IV.
20 Percentages of increases or decreases are shown in parentheses.

TABLE IV

Nutritional State ¹	Serum Cholesterol (mg/100/ml)			HMG-CoA Reductase (pmoles/mg/min) ²	Cholesterol 7 α -Hydroxylase (nmoles/mg/min) ³
	Total Chol.	NDL-Chol.	LDL-Chol.		
5 1) Chick Diet + 5.0% Corn Oil (COCO)	185.1 $\pm$ 4.12 ^A (100.0) ⁴	110.3 $\pm$ 4.95 ^A (100.0) ⁴	61.9 $\pm$ 1.69 ^A (100.0) ⁴	344.3 $\pm$ 1.69 ^A (100.0) ⁴	0.855 $\pm$ 0.084 ^A (100.0) ⁴
2) Chick Diet + 5.0% Corn Oil + Waxes; 50 ppm	184.7 $\pm$ 6.50 ^A (99.8)	109.6 $\pm$ 2.83 ^A (99.4)	61.7 $\pm$ 1.71 ^A (99.7)	339.3 $\pm$ 19.7 ^A (98.5)	0.837 $\pm$ 0.081 ^A (97.9)
10 3) Chick Diet + 5.0% Corn Oil + Waxes; 5,000 ppm	173.8 $\pm$ 7.31 ^A (93.9)	106.2 $\pm$ 4.69 ^A (96.3)	58.1 $\pm$ 1.77 ^A (93.9)	317.1 $\pm$ 14.4 ^{A,B} (92.1)	0.846 $\pm$ 0.072 ^A (98.9)
4) Chick Diet + 5.0% Corn Oil + Waxes; 10,000 ppm	165.9 $\pm$ 4.90 ^B (89.6)	108.5 $\pm$ 4.68 ^A (98.4)	57.9 $\pm$ 1.48 ^A (93.5)	304.3 $\pm$ 14.4 ^B (88.4)	0.902 $\pm$ 0.080 ^A (105.5)
15 5) Chick Diet + 5.0% Corn Oil + Tocotrienol-Rich- Fraction; 50 ppm	134.8 $\pm$ 3.82 ^C (72.8)	104.3 $\pm$ 3.99 ^A (94.6)	25.9 $\pm$ 1.02 ^B (41.8)	276.0 $\pm$ 17.6 ^C (80.2)	1.068 $\pm$ 0.047 ^B (124.9)
20 6) Chick Diet + 5.0% Methanol Insoluble Fraction	180.2 $\pm$ 6.01 ^A (97.4)	104.0 $\pm$ 4.37 ^A (94.3)	62.6 $\pm$ 1.68 ^A (101.1)	321.1 $\pm$ 19.3 ^B (93.3)	.852 $\pm$ 0.090 ^A (99.6)

¹ Feeding period was 4 weeks; time of killing was 0800 hours. The birds were fasted for 14 hours prior to killing. Data expressed as means $\pm$ SD; n = 6 chickens per group.

25 ² p-moles of mevalonic acid synthesized per minute per mg of microsomal protein.

³ n-moles of [¹⁴C] 7 α -hydroxycholesterol per minute per mg of microsomal protein.

⁴ Percentages of increases or decreases are in parentheses.

A-D Values not sharing a common superscript letter are different at P < 0.01.

The diets supplemented with sterols were not effective in lowering total serum or LDL-cholesterol levels. The maximum decrease in total cholesterol recorded with the sterol supplemented diet was 10.6% and that for LDL-cholesterol was only 6.5%. Only a marginal decrease was observed in hepatic enzymatic activity of HMG-CoA reductase (11.6%). Conversely, chickens fed the diet supplemented with the methanol soluble fraction of the TRF recorded a substantial decrease in the levels of total serum cholesterol (27.2%), LDL-cholesterol (58.2%) and hepatic enzymatic activity of HMG-CoA reductase (19.8%). Therefore, the

methanol soluble fraction of the TRF (containing the tocotrienols according to this invention), rather than sterols, was responsible for the impressive cholesterol-reducing properties of rice bran oil.

5

Example 9

This study was conducted to determine if sterols were responsible for the reduction in serum levels of various enzymes and blood constituents observed with rice bran. The feeding conditions
10 described in Example 8 were used. The samples were prepared as described in Example 8.

The results are displayed below in Table V. Percentages of increases or decreases are shown in parentheses.

TABLE V

	Nutritional State ¹	Fatty Acid Synthetase ²	Triglycerides	Glucose	Thromboxane B ₂	Platelet Factor 4
		(pmoles/mg/min)	(mg/100 ml)	(mg/100 ml)	(mg/100 ml)	(ng/ml)
5	1) Chick Diet + 5.0% Corn Oil (CDCO)	61.4±2.6 ^A (100.0)	90.2±1.1 ^A (100.0)	124.6±2.3 ^A (100.0)	16.7±1.6 ^A (100.0)	7.2±0.4 ^A (100.0)
	2) Chick Diet + 5.0% Corn Oil + Waxes; 50 ppm	62.5±1.9 ^A (102.0)	91.5±1.6 ^A (101.4)	126.7±2.1 ^A (101.7)	15.8±1.2 ^A (94.6)	7.5±0.4 ^A (104.2)
10	3) Chick Diet + 5.0% Corn Oil + Waxes; 5,000 ppm	63.6±2.8 ^B (103.8)	95.2±1.0 ^A (105.5)	123.9±1.5 ^A (99.4)	16.4±1.6 ^A (98.2)	7.4±0.3 ^A (102.8)
15	4) Chick Diet + 5.0% Corn Oil + Waxes; 10,000 ppm	60.4±1.9 ^A (98.5)	96.1±1.2 ^A (106.5)	124.3±1.1 ^A (99.8)	16.8±1.6 ^A (100.6)	7.4±0.8 ^A (102.8)
	5) Chick Diet + 5.0% Corn Oil + Tocotrienol-Rich- Fraction; 50 ppm	68.5±2.1 ^B (111.7)	73.2±1.6 ^B (81.2)	86.4±1.5 ^B (69.3)	12.4±1.4 ^B (74.3)	5.7±0.6 ^B (79.2)
20	6) Chick Diet + 5.0% Methanol Insoluble Fraction	63.8±1.2 ^B (107.3)	89.5±1.2 ^A (99.2)	111.1±2.4 ^C (89.4)	15.4±1.6 ^A (92.2)	7.5±0.3 ^A (104.2)

25 ¹ Feeding period was 4 weeks; time of killing was 0800 hours. The birds were fasted for 14 hours prior to killing. Data expressed as means ± SD; n = 6 chickens per group.

² n-moles of NADPH oxidized per minute per mg of cytosolic protein.

³ Percentages of increases or decreases are in parentheses.

A-C Values not sharing a common superscript letter are different at P < 0.01.

30 These results show that the sterols are not
responsible for the reduction in the levels of
triglycerides, glucose, thromboxane B₂ or platelet
factor 4 levels observed with rice bran and its oil.
In fact, the sterols either increased or had no
35 significant effect on the level of each of those
factors. Conversely, the chickens fed with the diet
supplemented with the methanol soluble fraction of the
TRF of Louisiana rice bran induced significant
decreases in all of these factors.

Example 10

This study measured the effects of three tocotrienols isolated from rice bran on cholesterol-related enzyme activity in hepatocytes isolated from
5 livers of 8-week old female chickens. The chickens were fed the chick mash diet for 8 weeks. They were then fasted for 40 hours and finally refed for 48 hours before sacrifice. The hepatocytes were then prepared following standard methods.

10 Each compound was isolated from TRF using the following protocol: A bond Elute amine column was equilibrated with 2 ml hexane. 10 mg of TRF (dissolved in 0.5 ml hexane) was bound to the amine column. The column was washed with 1 ml of hexane, then with 1 ml
15 of 3% isopropanol in hexane. P-21 and P-25 were eluted off the column with 5% isopropanol in hexane. Then using 10% isopropanol in hexane, the impurity P-20 was eluted. α -tocotrienol was also tested.

The results are displayed below in Table VI.
20 Percentages of increases or decreases are shown in parentheses.

TABLE VI

Concentration in ¹ (µg/ml)		HMG-CoA Reductase pmoles/min/mg	Fatty Acid Synthase nmoles/min/mg	Cholesterol 7α- hydroxylase pmoles/min/mg
5	A) α-Tocotrienol			
	1. 0.0	26.8 (100.0)	17.2 (100.0)	2.31 (100.0)
	2. 10.0	21.4 (79.9)	15.2 (88.4)	2.12 (91.8)
	3. 20.0	17.3 (64.6)	14.3 (83.2)	2.27 (98.7)
	4. 40.0	16.7 (62.3)	12.4 (72.1)	2.34 (101.3)
10	B) P-21 (Tocotrienol)			
	1. 0.0	26.8 (100.0)	17.2 (100.0)	2.31 (100.0)
	2. 10.0	20.4 (76.1)	14.3 (83.1)	2.41 (104.3)
	3. 20.0	16.7 (62.3)	12.7 (73.8)	3.40 (103.9)
	4. 40.0	14.2 (52.9)	11.8 (68.6)	2.39 (103.5)
15	5. 80.0	14.3 (53.4)	10.2 (59.3)	1.37 (102.6)
	C) P-25			
	1. 0.0	26.8 (100.0)	17.2 (100.0)	2.31 (100.0)
	2. 10.0	19.0 (70.9)	16.2 (94.2)	2.33 (100.9)
	3. 20.0	15.3 (57.1)	14.3 (83.1)	2.37 (102.61)
20	4. 40.0	12.4 (46.3)	12.7 (73.8)	2.44 (105.6)
	5. 80.0	11.1 (41.4)	12.1 (70.3)	2.37 (102.6)
	D) P-20			
	1. 0.0	28.78(100.0)		
	2. 25.0	29.21(101.5)		
25	3. 55.0	28.87(100.3)		
	4. 100.0	28.21 (98.0)		

¹ Feeding period was 8 weeks; the birds were fasted for 40 hrs. and refed for 48 hrs. The hepatocytes were prepared at 10 PM from two livers by standard methods.

P₂₅ demonstrated very effective dose dependant inhibition of HMG-CoA reductase with maximum reductions in activity of 46.6% and 58.6%, respectively. In fact, P₂₅ exerted substantially better effects than the known hypercholesterolemic agent, α-tocotrienol. α-tocotrienol showed a maximum reduction of only 37.7%. P₂₀, the sterol contaminant, showed no significant effect on the activity of HMG-CoA reductase.

Example 11

We next measured the effects of various tocotrienols and tocotrienol-like compounds isolated from rice bran on the activity of HMG-CoA reductase.

5 20.0 g of stabilized rice bran was extracted with 200.0 ml of methanol to remove various UV absorbing impurities and tocotrienols (6.7 mg total). This step was repeated four times. The remaining residue was dried under vacuum in a desiccator and then heated at

10 180°C for 2 hrs under 30 psi pressure. Then the dried residue was extracted again with 200.0 ml methanol (3.4 mg). The various peaks were purified by HPLC except 100 µl was injected instead of 20µl. Twenty runs were done to obtain enough material for this

15 study. The peaks which eluted at 32, 36, 45 and 54 minutes were tested in chicken hepatocytes as in Example 10.

The results of the HMG-CoA reductase assay are displayed below in Table VII.

20

TABLE VII

	Concentration ¹ (µg/ml)	HMG-CoA Reductase (nmoles/min/mg of microsome)			
		P-32	P-36	P-45	P-54
	1) 0.00	28.78	28.78	28.78	28.78
	2) 10.00	24.67	23.78	22.61	25.46
25	3) 20.00	21.31	20.12	20.33	19.67
	4) 30.00	18.36	17.24	19.67	20.11
	5) 40.00	16.59	17.34	18.21	18.19
	6) 50.00	16.61	15.21	18.40	18.20

30

¹ Feeding period was 8 weeks; the birds were fasted for 40 hrs. and refed for 48 hrs. The hepatocytes were prepared at 10 PM from two livers by standard methods.

Each of these compounds demonstrated an ability to inhibit the activity of HMG-CoA reductase.

Most notably, P₃₆ showed a 47% maximum reduction of activity.

Example 12

This example measured the effects of the TRF
5 and its components on the hepatic enzymatic activity of
HMG-CoA reductase and cholesterol 7 α -hydroxylase. Each
group of 12 chickens (6-week old female white leghorn)
was administered either a control diet or a control
diet supplemented with the TRF, a component of the TRF
10 a commercial cholesterol inhibitor (either Lovastatin
or Geraniol) or a combination of the commercial
inhibitors. The amount of feed consumed ranged from
11.19-11.60 g per chicken and the feeding period was 4
weeks. The birds were fasted for a period of 36 hours
15 and refed for 48 hours prior to sacrifice (at
0800 hours).

The TRF (entry 2) was prepared as described
in Example 10. Geraniol was obtained from Sigma
Chemical Co. and Lovastatin was obtained under the
20 brand name LOVOCOR.

The chicken diet for this study contained the
following ingredients:

	<u>Ingredients</u>	<u>Percentage</u>
	Corn (9.3% protein)	61.5
25	Soybean Meal (44.0% protein)	30.0
	Meat scrap (50.0% protein)	5.0
	Calcium Carbonate	0.5
30	Dicalcium Phosphate	1.0
	Alfalfa (17% protein)	1.0
	Mineral Mixture ^a	0.5

Vitamin Mixture^b

0.5

^a Mineral mixture contained per kg feed: zinc sulfate, 50 mg; sodium chloride, 2.0 mg; and manganese dioxide, 50.0 mg.

- 5 ^b Vitamin mixture contained per kg feed: vitamin A, 2,000 units; vitamin D₃, 200 units; vitamin E, 10 units, riboflavin, 3.6 mg; vitamin B₁₂, 0.01 mg; and vitamin K₁, 0.50 mg.

The results are displayed below in

- 10 Table VIII. Percentages of increases or decreases are in parentheses.

TABLE VIII

Nutritional State ¹		HMG-CoA Reductase ² pmole/min./mg	Cholesterol 7 α -Hydroxylase ³ nmole/min./mg
5	1) Control Diet (CD)	513.26 $\pm$ 15.07 ^A (100.00) ⁴	10.22 $\pm$ 0.25 ^A (100.00) ⁴
	2) CD + TRF-RBO; 50 ppm	441.14 $\pm$ 7.28 ^B (85.95) ⁴	10.80 $\pm$ 0.18 ^A (105.00) ⁴
10	3) CD + α - T ₃ ; 50 ppm	459.02 $\pm$ 1505 ^B (89.43) ⁴	10.69 $\pm$ 0.20 ^A (104.60) ⁴
	4) CD + γ - T ₃ ; 50 ppm	401.99 $\pm$ 5.84 ^C (78.32) ⁴	10.97 $\pm$ 0.17 ^A (107.34) ⁴
	5) CD + δ - T ₃ ; 50 ppm	389.22 $\pm$ 8.54 ^C (75.83) ⁴	10.98 $\pm$ 0.16 ^A (107.44) ⁴
15	6) CD + P-21-T ₃ ; 50 ppm	373.78 $\pm$ 7.99 ^D (72.83) ⁴	11.49 $\pm$ 0.20 ^B (112.43) ⁴
	7) CD + P-25-T ₃ ; 50 ppm	366.10 $\pm$ 6.66 ^D (71.33) ⁴	11.39 $\pm$ 0.26 ^B (111.45) ⁴
20	8) CD + Geraniol; 100 ppm	449.89 $\pm$ 12.82 ^B (87.65) ⁴	10.28 $\pm$ 0.27 ^A (100.59) ⁴
	9) CD + Lovastatin; 100 ppm	527.11 $\pm$ 13.93 ^A (102.69) ⁴	11.34 $\pm$ 0.31 ^B (110.96) ⁴
	10) CD + Geraniol+Lovastatin 50 ppm + 50 ppm	442.48 $\pm$ 1.64 ^B (86.21) ⁴	11.24 $\pm$ 0.17 ^B (109.98) ⁴

25 ¹ Feeding period was 4 weeks. Time of sacrificing was 0800 hr. The birds were fasted for 36 hrs. and refed 48 hrs. at the end of feeding period (28 days), and then sacrificed. Data expressed as means $\pm$ SD; n = 12 birds per group.

² pmoles of HMG-CoA formed per minute per mg of microsomal protein.

30 ³ nmoles of [¹⁴C]-cholesterol into [¹⁴C] 7 α -hydroxycholesterol per minute per mg of microsomal protein.

⁴ Percentages of increases are in parentheses.

A-D Values not sharing a common superscript letter are different at P < 0.01.

P₂₅ was the best inhibitor of HMG-CoA reductase. The enzymatic activity of HMG-CoA reductase was reduced by about 28% with P₂₅, while twice the amount of the commercial cholesterol-lowering drugs provided a maximum reduction of only about 13%. These data indicate that the fused oxygen heterocycle of the tocotrienols increases the biological activity as compared with a single ring system, such as Geraniol.

We believe that a three ring system might provide additional activity. Alternatively, adding substituents such as carbonyls, hydroxyls and alkyls may also enhance the biological activity of
5 tocotrienol-like compounds.

Example 13

We next studied the effects of the TRF and its components on total serum cholesterol levels, the HDL-cholesterol/total cholesterol ratio and the HDL-
10 /LDL-cholesterol ratio. Feeding conditions were identical to those in Example 12.

The samples were prepared as described in Example 12.

The results are displayed below in Table IX.
15 Percentages of increases or decreases are in parentheses.

TABLE IX

NUTRITIONAL STATE ¹	Concentration in serum (mg/dl)				
	Total Cholesterol	HDL-Cholesterol	LDL-Cholesterol	HDL-Cholesterol/ Total Cholesterol	HDL-Cholesterol/ LDL-Cholesterol
5 1) Control Diet (CD)	142.77±1.916 ^A (100.00) ²	83.66±0.85 ^A (100.00) ²	55.06±0.89 ^A (100.00) ²	0.59 (100.00) ²	1.52 (100.00) ²
2) CD + TRF + RBO 50 ppm	125.10±1.50 ^B (87.62)	78.71±1.06 ^B (94.08)	43.47±0.78 ^B (78.95)	0.63 (102.40)	1.81 (119.20)
10 3) CD + α - T ₃ 30 ppm	131.71±1.08 ^C (92.25)	80.47±0.67 ^B (96.19)	47.97±0.56 ^C (87.12)	0.61 (104.30)	1.68 (110.40)
4) CD + γ - T ₃ 50 ppm	116.86±1.47 ^D (81.85)	78.39±0.96 ^B (93.70)	37.52±0.81 ^D (68.14)	0.67 (114.50)	2.09 (137.50)
5) CD + δ - T ₃ 50 ppm	119.40±1.41 ^D (83.63)	77.87±0.97 ^B (93.08)	36.14±0.97 ^D (65.64)	0.65 (111.30)	2.15 (141.80)
15 6) CD + P-21 - T ₃ 50 ppm	92.91±1.27 ^E (65.08)	68.88±1.52 ^C (82.33)	22.16±0.57 ^E (40.25)	0.74 (126.50)	3.11 (204.60)
7) CD + P-25 - T ₃ 50 ppm	95.51±1.11 ^E (66.90)	70.12±1.25 ^C (83.82)	23.00±1.27 ^E (41.77)	0.73 (125.30)	3.05 (200.60)
20 8) CD + Geraniol 100 ppm	122.53±1.37 ^D (85.82)	82.28±1.06 ^A (98.35)	41.07±1.13 ^F (96.30)	0.65 (110.10)	2.00 (131.50)
9) CD + Lovastatin 100 ppm	128.65±1.02 ^C (90.11)	81.36±0.73 ^A (97.28)	44.12±1.26 ^B (86.51)	0.63 (108.00)	1.84 (121.40)
10) CD + Geraniol + Lovastatin; 50 ppm + 50 ppm	119.92±0.91 ^D (84.00)	80.93±0.62 ^B (96.74)	41.31±1.13 ^B (89.29)	0.67 (115.20)	1.96 (128.90)

¹ Feeding period was 4 weeks. Time of drawing the blood was 0800 hr. Data expressed as means ± SD; n = 12 birds per group.

² Percentages of increases or decreases are in parentheses.

A-F Values not sharing common superscript letter are different at P < 0.01.

- 30 P₂₁ and P₂₅ decreased the levels of serum cholesterol to a substantially greater degree than any other compound tested, including Lovastatin and Geraniol. P₂₅ caused about a 33% reduction. In comparison, Geraniol effected a 14% reduction and
- 35 Lovastatin caused only a 10% reduction. P₂₁ and P₂₅ were also clearly superior than the other compounds in increasing the HDL-/LDL-cholesterol ratio. Both of these compounds induced over a 100% increase in the HDL-/LDL-cholesterol ratio, while Lovastatin and
- 40 Geraniol increased the ratio by a maximum of 31.5%.

Example 14

We next measured the effects of TRF and its components on serum levels of apolipoprotein (a)₁, apolipoprotein B, triglycerides and glucose and the
5 plasma levels of thromboxane B₂ and platelet factor 4. Feeding conditions were identical to Example 12.

The samples were prepared as described in Example 12.

The results are displayed below in Table X.
10 Percentages of increases or decreases are in parentheses.

TABLE X

5	Nutritional State ¹	Concentration in Serum (mg/100 ml)				Thromboxane B ₂ (pg/ml)	Platelet Factor 4 (ng/ml)
		APO A ₁	APO B	Triglycerides	Glucose		
	1) Control diet (CD)	140.40 +1.7 ^A (100.00) ²	33.10 ±0.28 ^A (100.00) ²	74.03 ±1.26 ^A (100.00) ²	224.30 ±5.30 ^A (100.00) ²	25.84 ±0.84 ^A (100.00) ²	13.09 ±0.49 ^A (100.00) ²
10	2) CD + TRF-RBO; 50 ppm	140.20 ±2.8 ^A (99.86)	28.90 ±0.28 ^B (87.31)	64.73 ±0.93 ^B (87.44)	212.50 ±0.98 ^B (94.74)	20.55 ±0.95 ^B (79.53)	9.67 ±0.43 ^B (73.87)
	3) CD + α - T ₃ ; 50 ppm	139.60 ±3.2 ^A (99.43)	29.50 ±0.16 ^B (88.52)	68.04 ±1.32 ^C (91.91)	216.90 ±1.62 ^B (96.70)	21.63 ±0.85 ^B (83.71)	8.78 ±0.72 ^C (67.07)
15	4) CD + γ - T ₃ ; 50 ppm	139.80 ±1.5 ^A (99.37)	29.53 ±3.47 ^B (89.21)	61.07 ±1.37 ^D (82.49)	208.70 ±1.41 ^C (93.05)	19.22 ±0.78 ^{B,C} (74.38)	8.36 ±0.79 ^C (63.87)
20	5) CD + δ - T ₃ ; 50 ppm	139.30 ±1.4 ^A (99.22)	27.30 ±0.28 ^B (82.48)	60.50 ±2.50 ^D (81.72)	205.30 ±1.83 ^C (91.53)	18.68 ±0.99 ^C (72.17)	8.23 ±0.77 ^C (62.87)
	6) CD + P-21-T ₃ ; 50 ppm	138.40 ±4.4 ^A (98.79)	25.40 ±0.60 ^C (76.74)	56.22 ±1.11 ^E (75.94)	198.80 ±2.10 ^D (88.63)	16.74 ±1.62 ^C (64.78)	7.76 ±1.67 ^C (59.28)
25	7) CD + P-23-T ₃ ; 50 ppm	138.70 ±2.7 ^A (98.78)	25.20 ±0.40 ^C (76.13)	54.65 ±1.42 ^E (73.82)	196.10 ±1.74 ^D (87.43)	16.42 ±1.36 ^C (63.54)	7.27 ±0.74 ^C (55.54)
	8) CD + Geraniol; 100 ppm	138.40 ±1.7 ^A (98.58)	29.20 ±0.48 ^B (88.22)	71.29 ±1.19 ^F (96.30)	227.60 ±2.03 ^A (101.47)	25.46 ±1.52 ^A (98.53)	12.38 ±1.26 ^B (94.57)
30	9) CD + Lovastatin; 100 ppm	139.90 ±1.4 ^D (99.29)	25.80 ±0.89 ^C (77.95)	64.04 ±1.03 ^B (86.31)	220.10 ±1.32 ^D (98.13)	24.89 ±0.88 ^A (96.32)	12.24 ±1.49 ^B (93.51)
35	10) CD + Geraniol + Lovastatin; 50 ppm + 50 ppm	139.60 ±4.8 ^A (99.43)	25.95 ±0.34 ^C (78.40)	66.10 ±0.99 ^B (89.29)	221.70 ±1.89 ^D (98.84)	24.95 ±0.97 ^A (96.56)	12.19 ±1.15 ^B (93.12)

¹ Feeding period was 4 weeks. Time of drawing the blood was 0800 hr. Data expressed as means ± SD; n = 12 birds per group.

² Percentages of increases or decreases are in parentheses.

A-F Values not sharing a common superscript letter are different at P < 0.01.

40 P₂₁ and P₂₅ were the most potent cholesterol inhibitors. They effected the highest Apo A₁ to Apo B ratio and resulted in the maximum decrease in serum levels of triglycerides and glucose. In addition, they also reduced plasma levels of thromboxane B₂ and

platelet factor 4 to a greater extent than any other compound tested.

Example 15

We next measured the effects of various
5 tocotrienols from rice bran on HMG-CoA reductase and
cholesterol 7 α -hydroxylase in hypercholesterolemic
swine. Since pigs are very similar to humans in their
cholesterol metabolism, they provide a useful model
from which to study hypercholesterolemia and related
10 diseases. The pigs used in this study carried Lpd⁵ and
Lpu¹ mutant alleles for apolipoprotein B and U.
Because of this genetic defect, these pigs demonstrate
spontaneously elevated LDL-cholesterol levels and
hypercholesterolemia. The serum cholesterol
15 concentration in these swine is typically above 300-
500 mg/dl, as compared to 120-160 mg/dl in normal adult
swine. As a result, the pigs develop complicated
atherosclerotic plaques that closely resemble advanced
atherosclerotic lesions found in humans.

20 Each group of three 5-month old swine was
administered either a control diet or a control diet
supplemented with 50 ppm of TRF or 50 ppm of an
individual tocotrienol. The samples were prepared as
described in Example 11. After a 12 hour fast, serum
25 and plasma samples were taken at 0 hrs, 21 days and
42 days from the start of the feeding period. After
taking serum and plasma samples from the pigs at the
end of the 42 day period, all of the swine were fasted
for a total of 40 hours, followed by a 48 hour
30 refeeding period. One swine from each group was
sacrificed at that time and the liver, intestine, lung,
heart, loin muscle, adipose tissue and ham muscle were
removed.

The swine diet contained the following ingredients:

	<u>Ingredients</u>	<u>Percentage</u>
	Corn (9.3% protein)	78.37
5	Soybean Meal (44.0% protein)	15.42
	Lard	3.00
	Calcium Carbonate	0.95
	Dicalcium Phosphate	0.96
10	Mineral Mixture ^a	0.30
	Vitamin Mixture ^b	1.00

^a Mineral mixture contained per kg feed: zinc sulfate H_2O , 110 mg; manganese sulfate $5H_2O$, 70 mg; ferric citrate H_2O , 500.0 mg; copper sulfate $5H_2O$, 16.0 mg; 15 sodium selenite, 0.2 mg; DL-methionine, 2.5 g; choline chloride (50%), 1.5 g; ethoxyquin (1,2-dihydro-6-ethoxy-2,2,4-trimethyl-quinoline), 125 mg; and thiamine-HCl, 1.8 mg.

^b Vitamin mixture contained per kg feed: vitamin A, 20 1,500 units; vitamin D_3 , 400 units; vitamin E, 10 units; riboflavin, 3.6 mg; calcium pantothenate, 10.0 mg; niacin, 25.0 mg; pyridoxine-HCl, 3.0 mg; folacin, 0.55 mg; biotin, 0.15 mg; vitamin B_{12} , 0.01 mg; and vitamin K_1 , 0.55 mg.

25 The gain in body weight in all groups was comparable to the control, although feed conversion efficiency increased 7% with P_{21} and 10% with P_{25} .

The results of this study are displayed in Table XI. Percentages of increases and decreases are 30 shown in parentheses.

TABLE XI

5	NUTRITIONAL STATE ¹	HMG-CoA REDUCTASE ² pg/min/mg of microsomes Experimental Diets Feeding Period			Percentage of Control Activity
		0-time	21 days	42 days	
	Hypercholesterolemic Swine	355.49 ±8.44 (100.00)	374.14 ±13.33 (105.23)	391.38 ±15.75 (110.09)	100.00 ^A
10	2) Control Diet (CD)				
	3) CD+TRF RS-oil; 50 ppm	348.92 ±10.82 (100.00)	300.72 ±10.53 (86.19)	285.13 ±15.58 (81.72)	74.23 ^A
15	4) CD+ γ -T ₃ ; 50 ppm	347.99 ±16.29 (100.00)	300.72 ±10.53 (86.19)	280.88 ±12.81 (80.71)	73.31 ^B
	5) CD+P-21-T ₃ ; 50 ppm	340.36 ±23.55 (100.00)	304.23 ±23.13 (89.38)	275.75 ±12.76 (81.02)	73.59 ^B
20	6) CD+P-25-T ₃ ; 50 ppm	350.86 ±23.79 (100.00)	307.27 ±23.36 (87.58)	269.43 ±13.23 (76.79)	69.75 ^B

1 Corn-soybean meal control diet or experimental diets were fed to 5-month-old swine for 42 days. Then all groups were fasted for 40 hrs. and refed for 48 days. The time of sacrificing was 9:00 a.m. All tissues were kept on ice. TRF = Tocotrienol-Rich-Fraction.

2 Data expressed as means ± SD; n = 6 samples of 3 swine per group;

A-B Values not sharing a common superscript letter are different at P < 0.01.

5	NUTRITIONAL STATE ¹	CHOLESTEROL 7 α -HYDROXYLASE ² pg/min/mg of microsomes Experimental Diets Feeding Period			Percentage of Control Activity
		0-Time	21 days	42 days	
	Hypercholesterolemic Swine	4.88 ± 0.75 (100.00) ^A	5.00 ± 0.60 (102.44) ^B	5.07 ± 0.61 (103.89) ^B	100.00 ^A
	2) Control Diet (CD)				
10	3) CD+TRF RS-oil; 50 ppm	4.58 ± 0.10 (100.00)	4.96 ± 0.13 (108.29)	5.02 ± 0.11 (109.61)	105.50 ^A
	4) CD+ γ -T ₃ ; 50 ppm	4.49 ± 0.13 (100.00)	4.53 ± 0.12 (100.89)	4.58 ± 0.19 (102.00)	98.18 ^A
15	5) CD+Desmethyl-T ₃ (P-21); 50 ppm	4.53 ± 0.10 (100.00)	4.50 ± 0.18 (99.34)	4.63 ± 0.56 (102.21)	98.38 ^A
20	6) CD+Didesmethyl-T ₃ (P-25); 50 ppm	4.57 ± 0.19 (100.00)	4.62 ± 0.19 (101.09)	4.68 ± 0.18 (102.41)	98.58 ^A
25	¹ Corn-soybean meal control diet or experimental diets were fed to 5-month-old swine for 42 days. Then all groups were fasted for 40 hrs. and refed for 48 days. The time of sacrificing was 9:00 a.m. All tissues were kept on ice. TRF = Tocotrienol-Rich-Fraction.				
	² Data expressed as means $\pm$ SD; n = 6 samples of 3 swine per group;				
	³ Percentages of increases or decreases in respect to baseline value.				
	A-B Values not sharing a common superscript letter are different at P < 0.01.				

The hepatic enzymatic activity of the HMG-CoA reductase decreased approximately 30% with all the tocotrienols tested. In addition, there was no significant decrease in the activity of cholesterol 7 α -hydroxylase. These results were all time dependent and point to an overall reduction in cholesterologenesis, with no increase in cholesterol biosynthesis.

Example 16

Further tests were conducted on the swine described in Example 16. The experimental conditions were the same, except that the two remaining swine in each group were transferred to the unsupplemented

control diet following the 42 day feeding period. After an additional 70 days, serum and plasma samples were collected from these swine after a 12 hour fast for testing.

- 5 The samples were prepared as described in Example 12.

The results of these studies are displayed below in Tables XII-XV. Percentages of increases and decreases are shown in parentheses.

10

TABLE XII

NUTRITIONAL STATE ¹	TOTAL CHOLESTEROL CONCENTRATION IN SERUM (mg/100ml) ²					
	Experimental Diets Feeding Period			Percentage of Control Activity	Control Diet Feeding Period	Percentage of Control Activity
	0-Time	21 days	42 days		70 days	
15 Normolipemic Swine						
1) Control Diet (CD)	116.53 ±5.64 (100.00) ³	131.83 ±6.64 (113.13) ³	142.50 ±2.75 (122.29) ³	--	147.20 ±4.45 (126.32) ³	--
20 Hypercholesterolemic Swine						
2) Control Diet, (CD)	228.63 ±21.89 (100.00)	251.07 ±16.82 (110.02)	356.87 ±8.69 (156.79)	100.00 ^A	374.36 ±13.50 (164.60)	100.00 ^A
25 3) CD+TRF RD-oil; 50 ppm	206.43 ±18.45 (100.00)	222.00 ±15.75 (107.39)	243.03 ±12.28 (117.72)	75.00 ^B	285.77 ±14.00 (139.16)	84.60 ^B
4) CD+T ₃ ; 50 ppm	209.73 ±8.50 (100.00)	225.20 ±8.36 (107.39)	233.93 ±5.35 (111.76)	71.00 ^B	271.70 ±6.84 (129.66)	78.80 ^B
30 5) CD+P-21-T ₃ ; 50 ppm	220.50 ±16.54 (100.00)	236.40 ±18.17 (107.20)	238.17 ±14.29 (108.20)	68.90 ^B	276.23 ±15.93 (125.39)	76.20 ^B
6) CD+P-25-T ₃ ; 50 ppm	216.20 ±12.50 (100.00)	229.90 ±13.97 (106.33)	237.64 ±12.15 (107.60)	68.60 ^B	268.90 ±11.41 (124.45)	75.60 ^B

- 35 ¹ Feeding period was 6 weeks. Time of drawing the blood was 0.009 hr. after fasted for 12 hrs.; TRF = Tocotrienol-Rich-Fraction.

² Data expressed as means ± SD; n = 6 samples of 3 swine per group.

³ Percentages of increases or decreases in respect to baseline value.

A-B Values not sharing a common superscript letter are different at P < 0.01.

TABLE XIII

		HDL-CHOLESTEROL CONCENTRATION IN SERUM (mg/100ml) ²					
		Experimental Diets			Percentage	Control Diet	Percentage
		Feeding Period			of Control	Feeding Period	of Control
5	NUTRITIONAL STATE ¹	D-Time	21 days	42 days	Activity	70 days	Activity
	Normolipemic Swine						
	1) Control Diet (CD)	20.76 ±0.67 (100.00) ³	23.63 ±0.62 (113.82) ³	28.41 ±1.47 (136.85) ³	--	29.78 ±1.43 (143.45) ³	--
10	Hypercholesterolemic Swine						
	2) Control Diet (CD)	24.04 ±1.53 (100.00)	25.82 ±0.86 (107.40)	28.27 ±0.69 (117.59)	100.00 ^A	31.42 ±1.67 (130.69)	100.00 ^A
15	3) CD+TRF RD-oil; 50 ppm	23.36 ±1.35 (100.00)	25.14 ±1.49 (107.62)	26.50 ±1.84 (113.44)	96.47 ^A	30.19 ±1.21 (129.24)	98.89 ^A
	4) CD+YT ₃ ; 50 ppm	21.86 ±1.67 (100.00)	25.27 ±1.62 (115.59)	28.55 ±1.62 (130.60)	111.06 ^B	30.47 ±1.62 (139.39)	106.66 ^A
20	5) CD+P-21-T ₃ ; 50 ppm	22.95 ±1.73 (100.00)	26.23 ±0.89 (114.29)	30.74 ±1.13 (133.94)	113.90 ^B	31.15 ±1.73 (135.72)	103.80 ^A
25	6) CD+P-25-T ₃ ; 50 ppm	21.86 ±0.99 (100.00)	25.68 ±1.67 (117.47)	31.81 ±1.75 (145.52)	123.75 ^B	32.65 ±1.09 (149.36)	114.29 ^A

¹ Feeding period was 6 weeks. Time of drawing the blood was 0.009 hr. after fasting for 12 hrs.

² Data expressed as means ± SD; n = 6 samples of 3 swine per group; TRF = Tocotrienol-Rich-Fraction.

³ Percentages of increases or decreases in respect to baseline value.

A-B Values not sharing a common superscript letter are different at P < 0.01.

TABLE XIV

		LDL-CHOLESTEROL CONCENTRATION IN SERUM (mg/100ml) ²					
5	NUTRITIONAL STATE ¹	Experimental Diets Feeding Period			Percentage of Control Activity	Control Diet Feeding Period	Percentage of Control Activity
		0-Time	21 days	42 days		70 days	
	Normolipemic Swine						
	1) Control Diet (CD)	92.84 ±1.01 (100.00) ³	107.31 ±1.22 (115.58) ³	110.88 ±9.66 (119.43) ³	..	115.70 ±11.69 (124.62) ³	--
10	Hypercholesterolemic Swine	192.84 ±11.89 (100.00)	216.39 ±13.46 (112.21)	319.39 ±6.02 (164.07)	100.00 ^A	274.11 ±13.55 (142.14)	100.00 ^A
15	3) CD+TRF RD-oil; 50 ppm	177.28 ±10.97 (100.00)	187.15 ±13.49 (105.57)	209.09 ±11.86 (117.94)	71.88 ^B	267.08 ±7.7 (150.65)	105.99 ^A
	4) CD+T ₃ ; 50 ppm	178.37 ±6.32 (100.00)	185.12 ±6.63 (103.78)	193.80 ±2.70 (108.65)	66.22 ^C	224.66 ±3.71 (125.95)	88.61 ^B
20	5) CD+P-21-T ₃ ; 50 ppm	182.09 ±14.03 (100.00)	194.17 ±11.74 (106.63)	198.76 ±9.57 (109.15)	66.53 ^C	227.82 ±11.33 (125.11)	88.02 ^B
	6) CD+P-25-T ₃ ; 50 ppm	182.37 ±11.92 (100.00)	191.05 ±8.78 (104.76)	198.01 ±6.04 (107.48)	65.91 ^C	233.75 ±7.49 (128.17)	90.17 ^B
25							

¹ Feeding period was 6 weeks. Time of drawing the blood was 0.009 hr.

² Data expressed as means ± SD;; n = 6 samples of 3 swine per group; TRF = Tocotrienol-Rich-Fraction.

³ Percentages of increases or decreases in respect to baseline value.

30 A-C Values not sharing a common superscript letter are different at P < 0.01.

TABLE XV

5	NUTRITIONAL STATE ¹	HDL:LDL CHOLESTEROL CONCENTRATION IN SERUM (mg/100ml) ²					
		Experimental Diet ³			Percentage of Control Activity	Control Diet	
		0-Time	21 days	42 days		Feeding Period	Percentage of Control Activity
	Normolipemic Swine						
	1) Control Diet (CD)	1:4.47 (100.00) ³	1:4.54 (101.57) ³	1:3.90 (87.25) ³	--	1:3.89 (87.02) ³	--
10	Hypercholesterolemic Swine						
	2) Control Diet (CD)	1:8.02 (100.00)	1:8.38 (104.49)	1:11.19 (139.53)	100.00 ^A	1:8.72	100.00 ^A
	3) CD+TRF RD-oil; 50 ppm	1:7.59 (100.00)	1:7.44 (98.02)	1:7.89 (103.95)	70.51 ^B	1:8.85	101.41 ^A
15	4) CD+Y ₂ ; 50 ppm	1:8.16 (100.00)	1:7.33 (89.83)	1:6.79 (82.84)	60.38 ^C	1:7.37	84.52 ^B
	5) CD+P-21-T ₂ ; 50 ppm	1:7.93 (100.00)	1:7.40 (93.32)	1:6.47 (81.59)	57.82 ^C	1:7.31	83.83 ^B
	6) CD+P-25-T ₂ ; 50 ppm	1:8.34 (100.00)	1:7.44 (89.20)	1:6.22 (74.58)	55.59 ^D	1:7.16	82.06 ^B

20 ¹ Feeding period was 6 weeks. Time of drawing the blood was 0.009 hr. TRF = Tocotrienol-Rich-Fraction.

² Data expressed as means $\pm$ SD; n = 6 samples of 3 swine per group.

³ Percentages of increases or decreases in respect to baseline value.

A-D Values not sharing a common superscript letter are different at P < 0.01.

25 All tocotrienols tested caused the levels of
total serum cholesterol and LDL-cholesterol to decrease
by about 25% and 10%, respectively. In addition, the
HDL-/LDL-cholesterol ratio was reduced by about 20%
with all three tocotrienols tested. These reductions
30 were all time dependent. Moreover, the reductions
observed after 42 days persisted over the 10 weeks
following termination of tocotrienol feeding. This
result suggests that the tocotrienols remain in the
blood stream over extended periods of time. While not
35 wishing to be bound by theory, we believe that the
tocotrienols might be transported in association with
the LDL- and HDL- lipoproteins. Alternatively, the
tocotrienols may also bind to other proteins.

Example 17

This example was conducted to measure the serum levels of apolipoprotein (a)₁, apolipoprotein B, triglycerides, glucose, insulin and glucagon and the plasma levels of thromboxane B₂ and platelet factor 4 in the swine described in Example 15.

The samples were prepared as described in Example 12.

The results of these studies are displayed in Tables XVI and XVII. Percentages of increases and decreases are shown in parentheses.

TABLE XVI

NUTRITIONAL STATE ¹	Concentration in Serum (mg/100 ml)		
	Apo A ₁	Apo B	Triglycerides
15 Hypercholesterolemic Swine			
1) Control Diet (CD)	25.69±1.19 (100.00) ^A	147.77±1.50 (100.00) ^A	76.54±1.22 (100.00) ^A
2) CD+TRF RB-oil; 50 ppm	26.26±1.07 (102.22) ^A	118.39±1.50 (80.12) ^A	66.67±1.39 (86.15) ^B
3) CD+γ-T ₃ ; 50 ppm	26.22±1.11 (102.06) ^A	114.29±1.79 (77.34) ^C	61.78±1.09 (80.69) ^C
20 4) CD+P-21-T ₃ ; 50 ppm	26.72±1.14 (104.01) ^A	111.73±1.80 (75.64) ^C	64.44±1.29 (84.17) ^B
5) CD+P-25-T ₃ ; 50 ppm	26.73±1.22 (104.05) ^A	106.63±1.72 (72.16) ^D	62.00±1.19 (80.98) ^C

¹ Feeding period was 6 weeks. Time of drawing the blood was 0.009 hr.

² Data expressed as means ± SD; n = 6 samples of 3 swine per group; TRF = Tocotrienol-Rich-Fraction.

³ Percentages of increases or decreases in respect to baseline value.

A-D Values not sharing a common superscript letter are different at P < 0.01.

TABLE XVII

NUTRITIONAL STATE ¹		Glucose (mg/100 ml)	Insulin (μ G/ML)	Glucagon (pg/ml)	Thromboxane B ₂ (pg/100 ml)	Platelet Factor 4 (ng/100 ml)
5	Hypercholesterolemic Swine	99.99	10.50	330.41	75.93	24.12
	1) Control Diet (CD)	$\pm 1.37^A$ (100.00)	$\pm 0.12^A$ (100.00)	$\pm 14.08^A$ (100.00)	$\pm 1.45^A$ (100.00)	$\pm 1.75^A$ (100.00)
10	2) CD+TRF RB-oil; 50 ppm	74.43 $\pm 1.03^B$ (77.56)	25.10 $\pm 1.08^B$ (112.68)	295.36 $\pm 10.45^B$ (89.39)	64.35 ± 1.288 (95.23)	20.32 $\pm 1.70^B$ (84.30)
	3) CD+ γ -T ₃ ; 50 ppm	74.40 $\pm 1.12^B$ (77.51)	21.21 $\pm 0.89^B$ (103.44)	285.41 $\pm 13.88^B$ (86.38)	62.54 $\pm 1.39^B$ (92.25)	20.65 $\pm 1.15^B$ (85.28)
15	4) CD+P-21-T ₃ ; 50 ppm	72.63 $\pm 1.58^B$ (75.66)	21.43 $\pm 0.87^B$ (104.63)	278.36 $\pm 16.42^B$ (84.25)	60.48 $\pm 1.46^B$ (87.66)	19.85 $\pm 1.27^{B,C}$ (82.59)
	5) CD+P-25-T ₃ ; 50 ppm	71.89 $\pm 1.89^B$ (74.89)	20.89 $\pm 1.79^B$ (101.90)	276.41 $\pm 12.32^B$ (83.05)	57.03 $\pm 1.95^{B,C}$ (83.02)	19.15 $\pm 1.45^{B,C}$ (79.36)

¹ Corn-soybean meal control diet or experimental diets were fed to 5-month-old swine for 42 days. Then all groups were fasted for 40 hrs. and refed for 48 days. The time of sacrificing was 9:00 a.m. All tissues were kept on ice. TRF = Tocotrienol-Rich-Fraction.

² Data expressed as means $\pm$ SD; n = 6 samples of 3 swine per group;

³ Percentages of increases or decreases in respect to baseline value.

A-G Values not sharing a common superscript letter are different at P < 0.01.

These results show that all the tocotrienols tested reduced levels of apolipoprotein B, triglycerides, glucose, glucagon and platelet factor 4 by about 30%, 20%, 25%, 17% and 21%, respectively. The tocotrienols did not substantially affect the levels of apolipoprotein (a), and insulin.

Example 18

We next determined the effect of tocotrienols on the total cholesterol, triglyceride and glucose concentrations in various tissues of hypercholesterolemic swine.

The samples were prepared as described in Example 12.

The results of these studies are displayed in Tables XVIII - XX. Percentages of increases and decreases are shown in parentheses.

TABLE XVIII

		TOTAL CHOLESTEROL CONCENTRATION (mg/100gm) ²					
		Liver	Intestine	Lung	Heart	Loan Muscle	Adipose Tissue Ham Muscle
NUTRITIONAL STATE ¹							
Normolipemic Swine							
10	1) Control Diet (CD)	151.59 ±6.22	175.10 ±6.61	76.45 ±7.19	60.50 ±3.49	97.89 ±4.14	92.86 ±3.51 86.09 ±2.37
	Hypercholesterolemic Swine	201.59 ±5.68 ^A	215.49 ±5.78 ^A	101.74 ±4.34 ^A	64.51 ±9.61 ^A	119.62 ±3.89 ^A	109.40 ±4.00 ^A 100.00 ±3.73 ^A
	2) Control Diet (CD)	(100.00)	(100.00)	(100.00)	(100.00)	(100.00)	(100.00)(100.00)
15	3) CD+TRF RB-oil; 50 ppm	172.22 ±4.19 ^B (85.43)	198.63 ±3.75 ^B (92.18)	91.12 ±2.28 ^B (89.56)	58.78 ±1.87 ^B (91.12)	112.69 ±2.63 ^B (94.21)	102.25 ±1.94 ^A (93.46) 93.05 ±3.44 ^B (93.05)
	4) CD+T ₃ ; 50 ppm	149.20 ±3.83 ^C (74.01)	187.06 ±4.27 ^C (86.81)	83.79 ±3.24 ^C (82.36)	56.68 ±1.91 ^B (87.86)	103.48 ±2.39 ^C (86.51)	94.55 ±3.03 ^C (86.43) 85.15 ±2.24 ^C (85.15)
20	5) CD+P-21-T ₃ 50 ppm	136.70 ±4.55 ^D (67.81)	180.59 ±3.03 ^{C,D} (83.80)	75.68 ±4.14 ^D (74.39)	53.63 ±2.01 ^C (83.13)	100.58 ±1.71 ^C (84.09)	89.85 ±2.17 ^B (82.13) 79.51 ±3.75 ^C (79.51)
25	6) CD+P-21-T ₃ ; 50 ppm	125.21 ±4.30 ^E (62.11)	176.67 ±5.41 ^E (81.98)	68.5 ±5.32 ^D (66.98)	51.53 ±2.20 ^C (82.79)	99.04 ±3.52 ^C (81.85)	89.54 ±1.98 ^B (81.85) 69.17 ±3.88 ^D (69.17)

¹ Corn-soybean meal control diet or experimental diets were fed to 5-month-old swine for 42 days. Then all groups were fasted for 40 hrs. and refed for 48 days. The time of sacrificing was 9:00 a.m. All tissues were kept on ice. TRF = Tocotrienol-Rich-Fraction.

30 ² Data expressed as means ± SD; n = 6 samples of 3 swine per group;

A-E Values not sharing a common superscript letter are different at P < 0.01.

Percentages of control activity are given in parentheses.

TABLE XIX

TOTAL TRIGLYCERIDES CONCENTRATION (mg/100gm)²

NUTRITIONAL STATE ¹		Liver	Intestine	Lung	Heart	Loan Muscle	Adipose Tissue	Ham Muscle
5	Normalipenic Swine							
	1) Control Diet (CD)	388.68 ±5.80	300.24 ±5.63	163.53 ±3.57	130.51 ±3.59	207.55 ±5.50	294.43 ±4.99	187.50 ±3.68
10	Hypercholesterolemic Swine							
	2) Control Diet (CD)	429.36 ±5.82 ^A (100.00)	357.63 ±4.48 ^A (100.00)	171.96 ±4.04 ^A (100.00)	144.22 ±3.29 ^A (100.00)	219.68 ±5.19 ^A (100.00)	435.68 ±3.79 ^A (100.00)	185.83 ±4.05 ^A (100.00)
	3) CD+TRF RB-oil; 50 ppm	409.80 ±6.48 ^B (95.44)	355.93 ±3.07 ^A (99.52)	166.66 ±2.49 ^A (96.92)	140.71 ±3.35 ^A (97.57)	212.27 ±7.09 ^B (96.63)	421.42 ±3.43 ^B (96.73)	184.99 ±2.40 ^A (99.55)
15	4) CD+γ-T ₃ ; 50 ppm	397.51 ±3.88 ^C (92.58)	351.53 ±2.80 ^{A,B} (98.24)	167.18 ±2.86 ^A (97.22)	142.72 ±1.28 ^A (98.96)	201.65 ±2.01 ^C (91.79)	391.58 ±7.59 ^C (89.88)	181.99 ±1.67 ^A (97.93)
	5) CD+P-21-T ₃ ; 50 ppm	386.25 ±5.10 ^D (89.96)	347.75 ±4.74 ^B (97.24)	165.47 ±2.26 ^A (96.22)	142.38 ±1.73 ^A (98.72)	194.73 ±1.72 ^D (88.65)	282.82 ±6.86 ^C (64.91)	182.16 ±1.28 ^A (98.03)
20	6) CD+P-25-T ₃ ; 50 ppm	384.00 ±2.98 ^D (89.44)	345.37 ±3.97 ^B (96.57)	163.42 ±2.46 ^A (95.03)	140.51 ±0.99 ^A (97.43)	189.18 ±1.68 ^E (86.12)	259.62 ±7.47 ^D (59.59)	182.98 ±1.76 ^A (98.44)

¹ Corn-soybean meal control diet or experimental diets were fed to 5-month-old swine for 42 days. Then all groups were fasted for 40 hrs. and refed for 48 days. The time of sacrificing was 9:00 a.m. All tissues were kept on ice. TRF = Tocotrienol-Rich-Fraction.

² Data expressed as means ± SD; n = 4 analysis of the samples per group;

A-D Values not sharing a common superscript letter are different at P < 0.01.

30 Percentages of control activity are given in parentheses.

TABLE XX
TOTAL GLUCOSE CONCENTRATION (mg/100gm)²

NUTRITIONAL STATE ¹		Liver	Intestine	Lung	Heart	Loft Muscle	Adipose Tissue	Ham Muscle
5	Normolipemic Swine							
	1) Control Diet (CD)	361.05 ±7.79	66.86 ±5.81	61.35 ±4.45	42.86 ±1.91	65.79 ±2.57	21.85 ±5.01	53.82 ±2.60
	Hypercholesterolemic Swine							
10	2) Control Diet (CD)	405.23 ±7.68 ^A (100.00)	71.97 ±2.97 ^A (100.00)	66.93 ±4.40 ^A (100.00)	49.25 ±2.91 ^A (100.00)	71.24 ±1.26 ^A (100.00)	18.34 ±2.60 ^A (100.00)	53.82 ±2.60 ^A (100.00)
	3) CD+TRF RB-oil; 50 ppm	395.54 ±3.82 ^A (97.61)	64.02 ±2.80 ^B (88.95)	66.16 ±3.82 ^A (98.85)	46.80 ±2.50 ^A (95.03)	64.48 ±2.70 ^B (90.51)	17.04 ±1.35 ^{A,B} (92.01)	52.10 ±1.90 ^A (96.80)
15	4) CD+γ-T ₃ ; 50 ppm	383.68 ±3.37 ^B (94.53)	62.50 ±2.36 ^{B,C} (86.84)	63.08 ±3.38 ^{A,B} (94.25)	46.81 ±3.21 ^A (95.05)	60.30 ±1.95 ^B (84.64)	15.93 ±1.77 ^{A,B} (86.86)	52.10 ±1.69 ^A (96.80)
	5) CD+P-21-T ₃ ; 50 ppm	374.22 ±3.19 ^C (92.35)	62.50 ±2.80 ^{B,C} (86.84)	61.92 ±2.63 ^B (92.51)	44.17 ±3.55 ^A (89.69)	55.45 ±1.88 ^C (77.84)	16.67 ±1.96 ^{A,B} (90.89)	54.01 ±1.75 ^A (100.35)
20	6) CD+P-25-T ₃ ; 50 ppm	362.99 ±4.31 ^D (89.58)	56.06 ±4.06 ^C (77.89)	60.96 ±5.41 ^B (91.08)	45.49 ±2.78 ^A (92.37)	53.01 ±1.30 ^C (74.41)	14.96 ±1.26 ^B (81.57)	53.05 ±2.61 ^A (98.57)

25 ¹ Corn-soybean meal control diet or experimental diets were fed to 5-month-old swine for 42 days. Then all groups were fasted for 40 hrs. and refed for 48 days. The time of sacrificing was 9:00 a.m. All tissues were kept on ice. TRF = Tocotrienol-Rich-Fraction.

² Data expressed as means ± SD; n = 4 analysis of the sample per group;

³ Percentages of control activity are given in parentheses.

30 A-D values not sharing a common superscript letter are different at P < 0.01.

Decreases in total serum cholesterol were found in the each of the tissues tested. Most notably, total cholesterol concentration was reduced by 37% and 30% with P₂₅ in the liver and ham muscle, respectively.

35 Total triglyceride and glucose concentrations were less affected.

Example 19

We next measured the effects of different tocotrienols of rice bran oil on fatty acid

compositions in various tissues of hypercholesterolemic swine.

The levels of fatty acids, measured as their esters, were estimated using the method described in K. Hirai et al., "Effects of Dietary Fats and Phytosterol on Serum Fatty Acid Composition and Lipoprotein Cholesterol in Rats", J. Nutr. Sci. Vitaminol., 30, pp. 101-112 (1984), except that 1 g of each tissue was mixed with 2 ml of saline solution, the homogenized 30 sec by polytron. Following homogenization, the sample was extracted with 8 ml of hexane by shaking for 20 min., then centrifuging for 10 min. at 2000 rpm. The hexane layer was dried at 40°C and the residue treated with 0.5 ml diazomethane to yield the fatty acid esters.

The identification of each fatty acid ester was established by comparison against a standard mixture of fatty acid esters (obtained from Sigma Chemical Co.).

The results, as displayed in Figures 1-4, demonstrate that the tocotrienols tested were able to lower the total fatty acid concentration in every tissue tested. The most impressive declines were seen in the heart (76%), the loin (81%) and the ham muscle (66%). The greatest reductions were seen in levels of arachidonic acid. This decrease may be due to an increase in the corticosterone levels. Such an increase would inhibit the activity of phospholipase A₂, which would in turn decrease the arachidonic acid metabolites. The overall fatty acid reduction is also important because it points towards an immunoregulatory role of these tocotrienols.

Example 20

We next determined the effect of tocotrienols on immune function. The model for this study was 6-week old female white leghorn chickens. Each group of 12 chickens was fed an unsupplemented chicken diet or a control diet supplemented with 50 ppm of a tocotrienol or 100 ppm of a commercially available hypercholesterolemic agent (Geraniol or Lovastatin). The samples were prepared as described in Example 11.

10 The chickens were fed over a period of 4 weeks, then the blood was drawn. The antibody response was determined by injecting the chickens i.p. with 0.5 ml of a 5% suspension of sheep red blood cells (RBCs) in PBS (v/v) which were previously washed three times.

15 Total antibody titers to the RBCs were determined using 50 µg of serum and the microhemagglutination technique described in Witlin, "Detection of Antibodies by Microtitration Techniques", Mycopath. Mycol. Appl., 33, pp. 241-257 (1967). Relative IgG levels were

20 determined by using 2-mercaptoethanol preincubation for 30 min. The relative IgM levels were calculated by subtracting the IgG antibody levels from total antibody levels. All antibody levels are expressed as the log base 2 of the highest titer that was needed to

25 hemagglutinate an equal volume of 0.5% RBC suspension. The results of this example are displayed below in Table XXI.

TABLE XXI
Antibody titers in Serum (Log 2)

5	Nutritional State ¹	Total Ab titers	IgG	IgM
	1) Control Group (CD)	6.30±2.99 ^A	1.40±0.55 ^A	3.67±3.09 ^A
	2) CD + TRF-RBO 50 ppm	8.75±1.66 ^A	1.50±0.54 ^A	7.75±1.55 ^A
	3) CD + α - T ₃ 50 ppm	7.10±1.51 ^A	1.00±0.00 ^A	6.60±1.24 ^A
	4) CD + γ - T ₃ 50 ppm	6.60±1.51 ^A	1.28±0.49 ^A	5.83±1.53 ^A
10	5) CD + δ - T ₃ 50 ppm	8.67±1.78 ^A	1.00±0.00 ^A	8.25±1.96 ^A
	6) CD + P-21-T ₃ 50 ppm	8.67±1.92 ^A	1.50±1.18 ^A	7.17±2.04 ^A
	7) CD + P-25-T ₃ 50 ppm	10.30±1.16 ^B	2.43±1.40 ^A	8.92±2.61 ^A
	8) CD + Geraniol	5.60±2.11 ^A	1.00±0.00 ^A	5.42±2.02 ^A
	9) Lovastatin 100 ppm	6.33±1.44 ^A	1.00±0.00 ^A	5.90±1.68 ^A
15	10) Geraniol+Lovastatin 50 ppm + 50 ppm	7.17±2.04 ^A	1.00±0.00 ^A	6.83±1.99 ^A

¹ Feeding period was 4 weeks. Time of drawing the blood was 0800 hr. Data expressed as means ± SD; n = 12 birds per group.

A-B values not sharing a common superscript letter are different at P < 0.01.

20 Every tocotrienol, most notably F₂₅, increased the total antibody titer and the IgM titer as compared to the control. Slight increases were also observed in the IgG levels.

Example 21

25 We determined the effect of a known tocotrienol, γ-T₃, on the release of superoxide in human peripheral blood neutrophils. Neutrophils are an extracellular source of oxygen free radicals. Once activated, these neutrophils can attach to endothelial
30 tissue where they release a potent toxin, superoxide. Superoxide amplifies the inflammatory response and impairs local circulation of the blood.

The neutrophils we tested were isolated by density centrifugation on Ficoll-Hypaque gradients

using conventional methods (see E. Serbinova et al.,
"Free Radical Recycling and Intramembrane Mobility in
the Antioxidant Properties of α -Tocopherol and α -
Tocotrienol", Free Rad. Bio. and Med., 10, pp. 263-75
5 (1991)). The neutrophils were then placed in a 96-
well plate. γ -T₃ and phorbol myrstate acetate were
added to the wells at the same time. The secretion of
superoxide anion was measured as the superoxide
dismutase-inhibitable reduction of ferricytochrome C.
10 The results of this study are displayed in
Figure 5.

The amount of superoxide released was reduced
from 19.7 nmole (for 5×10^5 cells/hour) in the control
to 8.0 and 0 nmole at γ -T₃ concentrations of 10^{-6} and
15 10^{-5} , respectively.

While we have described a number of
embodiments of this invention, it is apparent that our
basic constructions may be altered to provide other
embodiments which utilize the processes and products of
20 this invention. Therefore, it will be appreciated that
the scope of this invention is to be defined by the
appended claims, rather than by the specific
embodiments which have been presented by way of
example.



THIS SHEET HAS BEEN
REPLACED

CLAIMS


We claim:

1. The use of a tocotrienol or tocotrienol-like compound for increasing feed conversion efficiency.

2. The use of a tocotrienol or tocotrienol-like compound for decreasing lipoprotein (a) concentration in the blood.

3. The use of a tocotrienol or tocotrienol-like compound in the treatment or prevention of a disease or condition selected from the group consisting of immunoregulatory diseases, inflammation, fever, edema, diabetes mellitus, pain, rheumatoid diseases, atherogenic diseases, septic shock, chronic fatigue syndrome and functio laesa.

4. A tocotrienol or tocotrienol-like compound having the formula:



and salts, oxidation products and hydrolysis products thereof wherein:

R_1 and R_3 is each independently selected from the group consisting of H, halogen, OH, OCH_3 and $\text{C}_1\text{-C}_8$ branched or unbranched alkyls;

R_2 is selected from the group consisting of OH, NHR_8 , CO_2Y , $\text{C(R}_8)_2\text{CO}_2\text{H}$, and $\text{C}_1\text{-C}_8$ branched or

This Sheet Has Been Replaced





THIS SHEET HAS BEEN
REPLACED


unbranched alkyls substituted by OH, NHR_8 , CO_2H or $\text{C}(\text{R}_8)_2\text{CO}_2\text{H}$;

R_4 is selected from the group consisting of O, NH, CH-R_9 , C=O and CH-OH ;

R_5 is selected from the group consisting of CH_2 , CHOH , O, S and NH;

R_6 is selected from the group consisting of H and $\text{C}_1\text{-C}_6$ branched or unbranched alkyls;

Each R_7 independently represents



wherein R_{10} is selected from the group consisting of H, NH_2 and $\text{C}_1\text{-C}_6$ branched or unbranched alkyls;

R_8 and R_9 is each independently selected from the group consisting of H and $\text{C}_1\text{-C}_6$ branched or unbranched alkyls;

Each R_{11} is independently selected from the group consisting of H, $\text{C}_1\text{-C}_6$ branched or unbranched alkyls, CH_2OH , CO_2H or OH;

H or $\text{C}_1\text{-C}_{18}$ branched or unbranched alkyls;

Z is halogen, OH, CH_2OH , CH_3 , OCH_3 or COCH_3 ;

n is an integer from 0-4; and

m is an integer from 1-30.

5. The tocotrienol or tocotrienol-like compound according to claim 14, wherein R_1 , R_3 and Z are each independently H, halogen, OH, OCH_3 or CH_3 ; R is OH, OCH_3 or NH_2 ; R_4 is C=O , CH_2 or NH; R_5 is O, S,

This Sheet Has Been Replaced






THIS SHEET HAS BEEN
REPLACED

- 91 -


CH_2 or NH ; R_9 is H or CH_3 ; R_{10} and R_{11} are each independently H or CH_3 ; n is 0 or 1; and m is 3-10.

6. A tocotrienol having the formula 3,4-dihydro-2-(4,8,12-trimethyltrideca-3'(E),7'(E),11'trienyl)-2H-1-benzopyran-6-ol).

7. A tocotrienol having the formula:



8. The tocotrienol having the formula:



9. The use of the tocotrienol or tocotrienol-like compound of any one of claims 4-8, or mixtures thereof, as a hypocholesterolemic agent, an antithrombotic agent, an antioxidizing agent, an antiatherogenic agent, an antiinflammatory agent, or an immunoregulatory agent.

10. The use of the tocotrienol or tocotrienol-like compound of any one of claims 5-8 or mixtures thereof to increase feed conversion efficiency.

11. The use of the tocotrienol or tocotrienol-like compound of any one of claims 4-8 or

This Sheet Has Been Replaced





- 92 -

THIS SHEET HAS BEEN
REPLACED

mixtures thereof to decrease lipoprotein (a) concentration in the blood.

12. The use of the tocotrienol or tocotrienol-like compound of any one of claims 4-8 or mixtures thereof to prevent oxidative degradation of food products.

13. The use according to claim 9, for the treatment or prevention of a disease or condition selected from the group consisting of: atherosclerosis, hypercholesterolemia, thromboses, endothelial injury, immunoregulatory diseases, atherogenic diseases, inflammation, fever, edema, diabetes mellitus, cancer, signs of aging, pain, rheumatoid inflammation, septic shock, chronic fatigue syndrome and functio laesa.

14. The use of TRF for increasing feed conversion efficiency.

15. The use of TRF for decreasing lipoprotein (a) concentration in the blood.

16. The use of TRF in the treatment or prevention of diseases or conditions selected from the group consisting of: immunoregulatory diseases, inflammation, fever, edema, cancer, signs of aging, diabetes mellitus, pain, rheumatoid inflammation, septic shock, chronic fatigue syndrome and functio laesa.

17. A tocotrienol or tocotrienol-like compound having the following characteristics:

This Sheet Has Been Replaced





THIS SHEET HAS BEEN
REPLACED

- 93 -

(a) an HPLC elution time of greater than about 22 minutes under the following conditions: 20 μ l sample, μ -Porasil column (Waters column, 10 μ , 4 mm x 30 cm) using an isocratic system of hexane and isopropanol (99.75%:0.25%, v/v) at a flow rate of 1.3 ml/min.; and

(b) an ability to inhibit the synthesis of β -hydroxy- β -methyl glutaryl coenzyme A reductase.

18. The tocotrienol or tocotrienol-like compound according to claim 17, wherein said HPLC elution time is about 32, 36, 45 or 54 minutes.

19. The tocotrienol or tocotrienol-like compound according to any one of claims 4-8, 17 or 18, obtained from a biological source.

20. The tocotrienol or tocotrienol-like compound according to claim 19, wherein the biological source is stabilized.

21. The tocotrienol or tocotrienol-like compound according to claim 19 or 20, wherein the biological source is a plant source.

22. The tocotrienol or tocotrienol-like compound according to claim 21, wherein the plant source is selected from the group consisting of oats, oat bran, oat bran oil, barley, barley bran, barley bran oil, rice, rice bran and rice bran oil.

23. The tocotrienol or tocotrienol-like compound according to claim 22, wherein the plant source is rice bran or rice bran oil.

This Sheet Has Been Replaced





THIS SHEET HAS BEEN
REPLACED

24. The tocotrienol or tocotrienol-like compound according to any one of claims 4-8, 17 or 18, obtained by chemical synthesis.

25. A foodstuff comprising the tocotrienol or tocotrienol-like compound according to any one of claims 4-8, 17 or 18 or mixtures thereof.

26. The foodstuff of claim 54, further comprising TRF, P₂₁, α -, β -, γ - or δ -tocotrienol.

27. A dietary supplement comprising the tocotrienol or tocotrienol-like compound according to any one of claims 4-8, 17 or 18, or mixtures thereof.

28. The dietary supplement according to claim 27, further comprising TRF, tocotrienol, α -, β -, γ - or δ -tocotrienol.

29. A pharmaceutical composition (for treating or preventing hypercholesterolemic-related diseases, thrombotic diseases, atherogenic diseases, inflammation, immunodeficiencies, fever, edema, diabetes mellitus, cancer, signs of aging, pain, septic shock, chronic fatigue syndrome or functio laesa, or to decrease lipoprotein (a) concentration in the blood or to increase feed conversion efficiency) comprising the tocotrienol or tocotrienol-like compound according to any one of claims 4-8, 17 or 18 or mixtures thereof, and a pharmaceutical carrier.

30. The pharmaceutical composition according to claim 29, (further comprising TRF, P₂₁, α -, β -, γ - or δ -tocotrienol)

This Sheet Has Been Replaced





- 95 -

THIS SHEET HAS BEEN
REPLACED

31. A pharmaceutical composition for treating or preventing inflammation, immunoregulatory diseases, fever, edema, diabetes mellitus, cancer, signs of aging, pain, rheumatoid diseases, septic shock, chronic fatigue syndrome or functio laesa, to decrease lipoprotein (a) concentration in the blood or to increase feed conversion efficiency comprising TRF, P₂₁, α -, β -, δ - or γ -tocotrienol or a combination thereof.

32. A method for treating or preventing hypercholesterolemic-related diseases, thrombotic diseases, atherogenic diseases, inflammation, immunodeficiencies, fever, edema, diabetes mellitus, cancer, signs of aging, pain, septic shock, chronic fatigue syndrome or functio laesa, to decrease lipoprotein (a) concentration in the blood or to increase feed conversion efficiency in a patient comprising the step of administering to said patient a pharmaceutically effective amount of the pharmaceutical composition according to claim 29 or 30.

33. A method for treating or preventing inflammation, immunoregulatory diseases, fever, edema, diabetes mellitus, cancer, signs of aging, pain, septic shock, chronic fatigue syndrome and functio laesa, to decrease lipoprotein (a) concentration in the blood or to increase feed conversion efficiency in a patient comprising the step of administering to said patient a pharmaceutically effective amount of the pharmaceutical composition according to claim 31.

34. A method for decreasing the total serum cholesterol and LDL-cholesterol in a human or animal comprising the step of administering to said human or

This Sheet Has Been Replaced





animal a cholesterol-lowering effective amount of a foodstuff according to claim 25 or 26.

35. A method for decreasing the total serum cholesterol and LDL-cholesterol in a human or animal comprising the step of administering to said human or animal a cholesterol-lowering effective amount of a dietary supplement according to claim 27 or 28.

36. A method for increasing the HDL/LDL cholesterol ratio in a human or animal comprising the step of administering to said human or animal a cholesterol-lowering effective amount of a foodstuff according to claim 25 or 26.

37. A method for decreasing the HDL/LDL cholesterol ratio in a human or animal comprising the step of administering to said human or animal a cholesterol-lowering effective amount of a dietary supplement according to claim 27 or 28.)

38. A method to enhance the yield of tocotrienols and tocotrienol-like compounds from the tocotrienol rich fraction of a biological source, comprising the steps of:

- (a) dissolving the tocotrienol rich fraction from a biological source in a solvent,
- (b) contacting the dissolved tocotrienol rich fraction with an amine column, and
- (c) eluting the tocotrienols with a solvent that does not substantially elute impurities.

39. A method to enhance the yield of tocotrienols and tocotrienol-like compounds from a biological source, comprising the steps of: _____





THIS SHEET HAS BEEN
REPLACED

- (a) extracting the biological source with an organic solvent,
- (b) heating the extracted source, and
- (c) extracting the heated biological source.

40. The method according to claim 39, wherein the heating of step (b) is accomplished with microwave heat.

41. The method according to claim 38 or 39, wherein the biological source is a plant source.

42. The method according to claim 41, wherein the plant source is selected from the group consisting of oats, peanuts, pine nuts, oat bran, oat bran oil, barley, barley bran, barley bran oil, rice, rice bran and rice bran oil.

43. The method according to claim 42, wherein the plant source is rice bran or rice bran oil.


44. The method according to claim 38, wherein the solvent of step (a) is hexane.

45. The method according to claim 38, wherein the amine column of step (b) is a Bond Elute amine or cyano column.

46. The method according to claim 38, wherein the solvent of step (c) is a gradient system of 0.5% - 3% isopropanol in hexane.

47. The use of a stabilized biological source for decreasing lipoprotein (a) concentration in

This Sheet has Been Replaced





THIS SHEET HAS BEEN
REPLACED

- 98 -

the blood, increasing feed conversion efficiency or in the treatment or prevention of one or more of the following diseases or conditions: immunoregulatory diseases, inflammation, fever, edema, diabetes mellitus, cancer, signs of aging, pain, rheumatoid diseases, septic shock, chronic fatigue syndrome and *functio laesa*.

48. The use according to claim 47, wherein the stabilized biological source is stabilized rice bran.

This Sheet Has Been Replaced



ABSTRACT

The present invention relates to novel
tocotrienols and tocotrienol-like compounds displaying
biological activity. The tocotrienols and tocotrienol-
5 like compounds of this invention may be conveniently
obtained from biological sources or by chemical
synthesis and may be used in pharmaceutical
compositions, foodstuffs and dietary supplements. This
invention also relates to the use of tocotrienols,
10 tocotrienol-like compounds, and mixtures thereof, as
hypocholesterolemic, antithrombotic, antioxidizing,
antiatherogenic, antiinflammatory and immunoregulatory
agents, or as agents useful to decrease lipoprotein (a)
concentration in the blood or to increase feed
15 conversion efficiency.

This Sheet Has Been Replaced



FIG. 1A

	ME- Myristic Acid 14:0	ME- Palmitic Acid 16:0	ME- Palmitoleic Acid 16:1
LIVER			
Normolipemic Swine			
I) Control Diet (CD)	31.23	134.46	3.16
Hypercholesterolemic Swine			
II) Control Diet (CD)	32.58	271.61	20.39
III) CD & TRF RB oil; 50 ppm	20.22	138.52	17.83
IV) CD & Tocotrienol; 50 ppm	14.17	112.15	5.79
V) CD & P-21- Tocotrienol; 50 ppm	17.05	155.07	6.36
VI) CD & P-25- Tocotrienol; 50 ppm (P-25)	20.30	132.04	12.12
INTESTINE			
Normolipemic Swine			
I) Control Diet (CD)	34.67	142.66	---
Hypercholesterolemic Swine			
II) Control Diet (CD)	16.25	139.01	0.79
III) CD & TRF RB oil; 50 ppm	13.79	93.59	---
IV) CD & δ -Tocotrienol; 50 ppm	17.88	85.28	---
V) CD & P-21- Tocotrienol; 50 ppm	13.36	85.72	---
VI) CD & P-25- Tocotrienol; 50 ppm	16.84	106.87	---



2 / 13

FIG. 1B

	ME- Palmitelaic Acid 16:1	ME- Stearic Acid 18:0	ME- Oleac Acid 18:1	ME- Linoleac Acid 18:2
LIVER				
Normolipemic Swine				
I) Control Diet (CD)	0.49	183.54	3.49	185.89
Hypercholesterolemic Swine				
II) Control Diet (CD)	17.87	94.45	189.89	5.12
III) CD & TRF RB oil; 50 ppm	---	140.69	150.07	---
IV) CD & Tocotrienol; 50 ppm	17.21	141.89	157.99	1.94
V) CD & P-21- Tocotrienol; 50 ppm	19.59	183.63	198.40	---
VI) CD & P-25- Tocotrienol; 50 ppm (P-25)	18.59	116.89	161.05	---
INTESTINE				
Normolipemic Swine				
I) Control Diet (CD)	14.62	177.46	141.49	---
Hypercholesterolemic Swine				
II) Control Diet (CD)	12.03	150.68	194.12	2.74
III) CD & TRF RB oil; 50 ppm	3.99	127.93	156.46	---
IV) CD & δ -Tocotrienol; 50 ppm	13.41	124.04	72.37	---
V) CD & P-21- Tocotrienol; 50 ppm	10.97	107.39	151.61	2.29
VI) CD & P-25- Tocotrienol; 50 ppm	17.89	115.53	71.65	0.73



3/13

FIG. 1C

	ME- Linolenic Acid 18:3	ME- Arachidonic Acid 20:4	Total Fatty Acids
LIVER			
Normolipemic Swine			
I) Control Diet (CD)	0.68	0.4	543.36
Hypercholesterolemic Swine			
II) Control Diet (CD)	1.06	10.9	634.06
III) CD & TRF RB oil; 50 ppm	---	3.2	470.53
IV) CD & Tocotrienol; 50 ppm	1.94	1.50	454.58
V) CD & P-21- Tocotrienol; 50 ppm	5.08	1.02	603.25
VI) CD & P-25- Tocotrienol; 50 ppm (P-25)	2.08	1.08	482.15
INTESTINE			
Normolipemic Swine			
I) Control Diet (CD)	2.86	---	513.76
Hypercholesterolemic Swine			
II) Control Diet (CD)	---	1.28	516.90
III) CD & TRF RB oil; 50 ppm	---	1.02	396.78
IV) CD & δ -Tocotrienol; 50 ppm	---	0.96	313.94
V) CD & P-21- Tocotrienol; 50 ppm	---	0.78	372.12
VI) CD & P-25- Tocotrienol; 50 ppm	---	0.77	330.28

This Sheet Has Been Replaced



4 / 13

FIG. 2A

	ME- Myristic Acid 14:0	ME- Palmitic Acid 16:0	ME- Palmitoleic Acid 16:1
LUNG			
Normolipemic Swine			
I) Control Diet (CD)	15.12	87.75	---
Hypercholesterolemic Swine			
II) Control Diet (CD)	---	122.15	---
III) CD & TRF RB oil; 50 ppm	21.49	115.94	---
IV) CD & Tocotrienol; 50 ppm	---	75.46	---
V) CD & P-21- Tocotrienol; 50 ppm	---	68.19	---
VI) CD & Didesmethyl- Tocotrienol; 50 ppm (P-25)	---	76.89	---
HEART			
Normolipemic Swine			
I) Control Diet (CD)	---	42.63	---
Hypercholesterolemic Swine			
II) Control Diet (CD)	4.86	27.52	---
III) CD & TRF RB oil; 50 ppm	---	44.65	---
IV) CD & Tocotrienol; 50 ppm	---	42.23	---
V) CD & P-21- Tocotrienol; 50 ppm	---	4.19	---
VI) CD & P-25- Tocotrienol 50 ppm	---	3.44	---

This Sheet Has Been Replaced.



5/13

FIG. 2B

	ME- Palmitelaic Acid 16:1	ME- Stearic Acid 18:0	ME- Oleac Acid 18:1
LUNG			
Normolipemic Swine			
I) Control Diet (CD)	12.02	137.93	64.69
Hypercholesterolemic Swine			
II) Control Diet (CD)	---	158.81	128.29
III) CD & TRF RB oil; 50 ppm	9.68	118.94	54.61
IV) CD & Tocotrienol; 50 ppm	---	160.12	58.71
V) CD & P-21- Tocotrienol; 50 ppm	---	133.04	50.76
VI) CD & Didesmethyl- Tocotrienol; 50 ppm (P-25)	---	46.86	80.19
HEART			
Normolipemic Swine			
I) Control Diet (CD)	---	69.35	59.35
Hypercholesterolemic Swine			
II) Control Diet (CD)	11.87	49.18	47.43
III) CD & TRF RB oil; 50 ppm	---	99.32	53.42
IV) CD & Tocotrienol; 50 ppm	---	84.30	64.16
V) CD & P-21- Tocotrienol; 50 ppm	6.13	21.56	18.53
VI) CD & P-25- Tocotrienol 50 ppm	4.73	12.38	12.92

This Sheet Has Been Replaced





THIS SHEET HAS BEEN
REPLACED

6/13

FIG. 2C

	ME- Linoleic Acid 18:2	ME- Linolenic Acid 18:3	ME- Arachidonic Acid 20:4	Total Fatty Acids
LUNG				
Normolipemic Swine				
I) Control Diet (CD)	---	---	---	317.51
Hypercholesterolemic Swine				
II) Control Diet (CD)	---	---	---	409.25
III) CD & TRF RB oil; 50 ppm	---	---	---	321.36
IV) CD & Tocotrienol; 50 ppm	3.32	---	---	297.61
V) CD & P-21- Tocotrienol; 50 ppm	---	---	---	251.99
VI) CD & Didesmethyl- Tocotrienol; 50 ppm (P-25)	3.39	3.32	---	310.65
HEART				
Normolipemic Swine				
I) Control Diet (CD)	---	---	---	171.33
Hypercholesterolemic Swine				
II) Control Diet (CD)	---	---	---	140.86
III) CD & TRF RB oil; 50 ppm	---	---	---	197.39
IV) CD & Tocotrienol; 50 ppm	---	---	---	190.69
V) CD & P-21- Tocotrienol; 50 ppm	---	---	---	50.41
VI) CD & P-25- Tocotrienol 50 ppm	---	---	---	33.47

THIS SHEET HAS BEEN
REPLACED

This Sheet Has Been Replaced



7/13

FIG. 3A

	ME- Myristic Acid 14:0	ME- Palmitic Acid 16:0	ME- Palmitoleic Acid 16:1
LOIN MUSCLE			
Normolipemic Swine			
1) Control Diet (CD)	23.51	125.97	---
Hypercholesterolemic Swine			

8/13

FIG. 3B

	ME- Palmitelaic Acid 16:1	ME- Stearic Acid 18:0	ME- Oleic Acid 18:1
LOIN MUSCLE			
Normolipemic Swine			
I) Control Diet (CD)	5.28	359.83	185.80
Hypercholesterolemic Swine			
II) Control Diet (CD)	---	83.61	58.72
III) CD & TRF RB oil; 50 ppm	---	72.08	18.13
IV) CD & δ -Tocotrienol; 50 ppm	---	48.67	15.90
V) CD & P-21- Tocotrienol; 50 ppm	---	38.58	11.41
VI) CD & P-25- Tocotrienol; 50 ppm	1.37	12.39	5.87
ADIPOSE TISSUE			
Normolipemic Swine			
I) Control Diet (CD)	---	322.76	319.23
Hypercholesterolemic Swine			
II) Control Diet (CD)	21.87	160.42	209.76
III) CD & TRF RB oil; 50 ppm	7.21	192.17	231.54
IV) CD & δ -Tocotrienol; 50 ppm	9.95	231.54	184.89
V) CD & P-21- Tocotrienol; 50 ppm	8.22	146.77	16.80
VI) CD & P-25 Tocotrienol; 50 ppm	7.23	133.40	15.00

This Sheet Has Been Replaced





9/13

FIG. 3C

	ME- Linoleic Acid 18:2	ME- Linolenic Acid 18:3	ME- Arachidonic Acid 20:4	Total Fatty Acids
LOIN MUSCLE				
Normolipemic Swine				
I) Control Diet (CD)	---	---	---	700.39
Hypercholesterolemic Swine				
II) Control Diet (CD)	---	---	---	167.50
III) CD & TRF RB oil; 50 ppm	---	---	---	109.54
IV) CD & δ -Tocotrienol; 50 ppm	---	---	---	76.67
V) CD & P-21- Tocotrienol; 50 ppm	---	---	---	56.06
VI) CD & P-25- Tocotrienol; 50 ppm	---	---	---	31.20
ADIPOSE TISSUE				
Normolipemic Swine				
I) Control Diet (CD)	---	---	3.4	852.94
Hypercholesterolemic Swine				
II) Control Diet (CD)	---	---	7.8	728.54
III) CD & TRF RB oil; 50 ppm.	---	---	2.2	585.98
IV) CD & δ -Tocotrienol; 50 ppm	---	---	2.0	355.91
V) CD & P-21- Tocotrienol; 50 ppm	---	---	2.0	248.85
VI) CD & P-25 Tocotrienol; 50 ppm	---	---	1.9	230.35

10/13

FIG. 4 A

	ME- Myristic Acid 14:0	ME- Palmitic Acid 16:0	ME- Palmitoleic Acid 16:1
HAM MUSCLE			
Normolipemic Swine			
I) Control Diet (CD)	---	141.38	---
Hypercholesterolemic Swine			
II) Control Diet (CD)	3.41	24.15	---
III) CD & TRF RB oil; 50 ppm	---	38.92	---
IV) CD & δ -Tocotrienol; 50 ppm	10.62	45.04	---
V) CD & P-21- Tocotrienol; 50 ppm	5.35	25.21	---
VI) CD & P-25 Tocotrienol; 50 ppm	---	7.91	---

This Sheet Has Been Replaced




11/13

FIG. 4B

	ME- Palmitelaic Acid 16:1	ME- Stearic Acid 18:0	ME- Oleac Acid 18:1
HAM MUSCLE			
Normolipemic Swine			
I) Control Diet (CD)	---	---	59.94
Hypercholesterolemic Swine			
II) Control Diet (CD)	---	149.91	21.59
III) CD & TRF RB oil; 50 ppm	---	80.04	---
IV) CD & δ -Tocotrienol; 50 ppm	---	130.08	---
V) CD & P-21- Tocotrienol; 50 ppm	---	---	---
VI) CD & P-25 Tocotrienol; 50 ppm	---	60.04	---

This Sheet Has Been Replaced



12/13

FIG.4C

	ME- Linoleic Acid 18:2	ME- Linolenic Acid 18:3	ME- Arachidonic Acid 20:4	Total Fatty Acids
HAM MUSCLE				
Normolipemic Swine				
I) Control Diet (CD)	---	---	---	201.32
Hypercholesterolemic Swine				
II) Control Diet (CD)	---	---	---	199.06
III) CD & TRF RB oil; 50 ppm	---	---	---	118.96
IV) CD & δ -Tocotrienol; 50 ppm	---	---	---	158.74
V) CD & P-21- Tocotrienol; 50 ppm	---	---	---	30.56
VI) CD & P-25 Tocotrienol; 50 ppm	---	---	---	67.75

This Sheet Has Been Replaced



FIG. 5A ^{13/13}





FIG. 5B






23 OCT 1996

CLAIMS

1. The use of a tocotrienol or tocotrienol-like compound for the preparation of a pharmaceutical composition useful in increasing feed conversion efficiency.
2. The use of a tocotrienol or tocotrienol-like compound for the preparation of a pharmaceutical composition useful in decreasing lipoprotein (a) concentration in the blood.
3. The use of a tocotrienol or tocotrienol-like compound for the preparation of a pharmaceutical composition useful in the treatment or prevention of a disease or condition selected from the group consisting of immunoregulatory diseases, inflammation, fever, edema, diabetes mellitus, pain, rheumatoid diseases, atherogenic diseases, septic shock, chronic fatigue syndrome and functio laesa.
4. A tocotrienol or tocotrienol-like compound having the formula:



and salts, oxidation products and hydrolysis products thereof wherein:

R_1 and R_3 is each independently selected from the group consisting of H, halogen, OH, OCH_3 and $\text{C}_1\text{-C}_6$ branched or unbranched alkyls;

R_2 is selected from the group consisting of OH, NHR_8 , CO_2Y , $\text{C}(\text{R}_8)_2\text{CO}_2\text{H}$, and $\text{C}_1\text{-C}_6$ branched or



23 OCT 1996

- 90 -


unbranched alkyls substituted by OH, NHR_8 , CO_2H or $\text{C(R}_8)_2\text{CO}_2\text{H}$;

R_4 is selected from the group consisting of O, NH, CH-R_9 , C=O and CH-OH ;

R_5 is selected from the group consisting of CH_2 , CHOH , O, S and NH;

R_6 is selected from the group consisting of H and $\text{C}_1\text{-C}_6$ branched or unbranched alkyls;

Each R_7 independently represents



wherein R_{10} is selected from the group consisting of H, NH_2 and $\text{C}_1\text{-C}_6$ branched or unbranched alkyls;

R_8 and R_9 is each independently selected from the group consisting of H and $\text{C}_1\text{-C}_6$ branched or unbranched alkyls;

Each R_{11} is independently selected from the group consisting of H, $\text{C}_1\text{-C}_6$ branched or unbranched alkyls, CH_2OH , CO_2H or OH;

H or $\text{C}_1\text{-C}_{18}$ branched or unbranched alkyls;

Z is halogen, OH, CH_2OH , CH_3 , OCH_3 or COCH_3 ;

n is an integer from 0-4; and

m is an integer from 1-30.

5. The tocotrienol or tocotrienol-like compound according to claim 4, wherein R_1 , R_3 and Z are each independently H, halogen, OH, OCH_3 or CH_3 ; R is OH, OCH_3 or NH_2 ; R_4 is C=O , CH_2 or NH; R_5 is O, S,




23 OCT 1996


CH_2 or NH ; R_6 is H or CH_3 ; R_{10} and R_{11} are each independently H or CH_3 ; n is 0 or 1; and m is 3-10.

6. A tocotrienol having the formula 3,4-dihydro-2-(4,8,12-trimethyltrideca-3'(E),7'(E),11'trienyl)-2H-1-benzopyran-6-ol).

7. A tocotrienol having the formula:



8. The tocotrienol having the formula:



9. The use of the tocotrienol or tocotrienol-like compound of any one of claims 4-8, or mixtures thereof, for the preparation of a pharmaceutical composition useful as a hypocholesterolemic agent, an antithrombotic agent, an antioxidizing agent, an antiatherogenic agent, an antiinflammatory agent, or an immunoregulatory agent.

10. The use of the tocotrienol or tocotrienol-like compound of any one of claims 5-8 or mixtures thereof for the preparation of a pharmaceutical composition useful in increasing feed conversion efficiency.

11. The use of the tocotrienol or tocotrienol-like compound of any one of claims 4-8 or



23 OCT 1996

mixtures thereof for the preparation of a pharmaceutical composition useful in decreasing lipoprotein (a) concentration in the blood.

12. The use of the tocotrienol or tocotrienol-like compound of any one of claims 4-8 or mixtures thereof to prevent oxidative degradation of food products.

13. The use according to claim 9, wherein the pharmaceutical composition prepared is useful for the treatment or prevention of a disease or condition selected from the group consisting of:

atherosclerosis, hypercholesterolemia, thromboses, endothelial injury, immunoregulatory diseases, atherogenic diseases, inflammation, fever, edema, diabetes mellitus, cancer, signs of aging, pain, rheumatoid inflammation, septic shock, chronic fatigue syndrome and functio laesa.

14. The use of TRF for the preparation of a pharmaceutical composition useful in increasing feed conversion efficiency.

15. The use of TRF for the preparation of a pharmaceutical composition useful in decreasing lipoprotein (a) concentration in the blood.

16. The use of TRF for the preparation of a pharmaceutical composition useful for the treatment or prevention of diseases or conditions selected from the group consisting of: immunoregulatory diseases, inflammation, fever, edema, cancer, signs of aging, diabetes mellitus, pain, rheumatoid inflammation, septic shock, chronic fatigue syndrome and functio laesa.

17. A tocotrienol or tocotrienol-like compound having the following characteristics:



- 93 -

REPLACEMENT SHEET

23 OCT 1996

(a) an HPLC elution time of greater than about 22 minutes under the following conditions: 20 μ l sample, μ -Porasil column (Waters column, 10 μ , 4 mm x 30 cm) using an isocratic system of hexane and isopropanol (99.75%:0.25%, v/v) at a flow rate of 1.3 ml/min.; and

(b) an ability to inhibit the synthesis of β -hydroxy- β -methyl glutaryl coenzyme A reductase.

18. The tocotrienol or tocotrienol-like compound according to claim 17, wherein said HPLC elution time is about 32, 36, 45 or 54 minutes.

19. The tocotrienol or tocotrienol-like compound according to any one of claims 4-8, 17 or 18, obtained from a biological source.

20. The tocotrienol or tocotrienol-like compound according to claim 19, wherein the biological source is stabilized.

21. The tocotrienol or tocotrienol-like compound according to claim 19 or 20, wherein the biological source is a plant source.

22. The tocotrienol or tocotrienol-like compound according to claim 21, wherein the plant source is selected from the group consisting of oats, oat bran, oat bran oil, barley, barley bran, barley bran oil, rice, rice bran and rice bran oil.

23. The tocotrienol or tocotrienol-like compound according to claim 22, wherein the plant source is rice bran or rice bran oil.

23 OCT 1996

24. The tocotrienol or tocotrienol-like compound according to any one of claims 4-8, 17 or 18, obtained by chemical synthesis.

25. A foodstuff comprising the tocotrienol or tocotrienol-like compound according to any one of claims 4-8, 17 or 18 or mixtures thereof.

26. The foodstuff of claim 54, further comprising TRF, P_{21} , α -, β -, γ - or δ -tocotrienol.

27. A dietary supplement comprising the tocotrienol or tocotrienol-like compound according to any one of claims 4-8, 17 or 18, or mixtures thereof.

28. The dietary supplement according to claim 27, further comprising TRF, tocotrienol, α -, β -, γ - or δ -tocotrienol.

29. A pharmaceutical composition comprising the tocotrienol or tocotrienol-like compound according to any one of claims 4 to 8, 17 or 18 or mixtures thereof and a pharmaceutical carrier.

30. A pharmaceutical composition according to claim 29 for treating or preventing hypercholesterolemic-related diseases, thrombotic diseases, atherogenic diseases, inflammation, immunodeficiencies, fever, edema, diabetes mellitus, cancer, signs of aging, pain, septic shock, chronic fatigue syndrome or functio laesa, or to decrease lipoprotein (a) concentration in the blood or to increase feed conversion efficiency.

31. The pharmaceutical composition according to claim 29 or 30 further comprising TRF, P_{21} , α -, β -, γ - or δ -tocotrienol.





23 OCT 1996

32. A pharmaceutical composition for treating or preventing inflammation, immunoregulatory diseases, fever, edema, diabetes mellitus, cancer, signs of aging, pain, rheumatoid diseases, septic shock, chronic fatigue syndrome or functio laesa, to decrease lipoprotein (a) concentration in the blood or to increase feed conversion efficiency comprising TRF, P₂₁, α -, β -, δ - or γ -tocotrienol or a combination thereof.

33. The foodstuff according to claim 25 or 26 for use in decreasing the total serum cholesterol and LDL-cholesterol in a human or animal.

34. The dietary supplement according to claim 27 or 28 for use in decreasing the total serum cholesterol and LDL-cholesterol in a human or animal.

35. The foodstuff according to claim 25 or 26 for use in increasing the HDL/LDL cholesterol ratio in a human or animal.

36. The dietary supplement according to claim 27 or 28 for use in decreasing the HDL/LDL cholesterol ratio in a human or animal.

37. A method to enhance the yield of tocotrienols and tocotrienol-like compounds from the tocotrienol rich fraction of a biological source, comprising the steps of:

- (a) dissolving the tocotrienol rich fraction from a biological source in a solvent,
- (b) contacting the dissolved tocotrienol rich fraction with an amine column, and
- (c) eluting the tocotrienols with a solvent that does not substantially elute impurities.



23 OCT 1996

38. A method to enhance the yield of tocotrienols and tocotrienol-like compounds from a biological source, comprising the steps of:

- (a) extracting the biological source with an organic solvent,
- (b) heating the extracted source, and
- (c) extracting the heated biological source.

39. The method according to claim 38, wherein the heating of step (b) is accomplished with microwave heat.

40. The method according to claim 37 or 38, wherein the biological source is a plant source.

41. The method according to claim 40, wherein the plant source is selected from the group consisting of oats, peanuts, pine nuts, oat bran, oat bran-oil, barley, barley bran, barley bran oil, rice, rice bran and rice bran oil.

42. The method according to claim 41 wherein the plant source is rice bran or rice bran oil.

43. The method according to claim 37 wherein the solvent of step (a) is hexane.

44. The method according to claim 37 wherein the amine column of step (b) is a Bond Elute amine or cyano column.

45. The method according to claim 37 wherein the solvent of step (c) is a gradient system of 0.5% - 3% isopropanol in hexane.

46. The use of a stabilized biological source for decreasing lipoprotein (a) concentration in



- 97 -

REPLACEMENT SHEET

23 OCT 1996

the blood, increasing feed conversion efficiency or in the treatment or prevention of one or more of the following diseases or conditions: immunoregulatory diseases, inflammation, fever, edema, diabetes mellitus, cancer, signs of aging, pain, rheumatoid diseases, septic shock, chronic fatigue syndrome and functio laesa.

47. The use according to claim 46, wherein the stabilized biological source is stabilized rice bran.

THIS SHEET HAS BEEN
REPLACED

- 98 -

REPLACEMENT SHEET

23 OCT 1996

ABSTRACT

The present invention relates to novel
tocotrienols and tocotrienol-like compounds displaying
biological activity. The tocotrienols and tocotrienol-
5 like compounds of this invention may be conveniently
obtained from biological sources or by chemical
synthesis and may be used in pharmaceutical
compositions, foodstuffs and dietary supplements. This
invention also relates to the use of tocotrienols,
10 tocotrienol-like compounds, and mixtures thereof, as
hypocholesterolemic, antithrombotic, antioxidizing,
antiatherogenic, antiinflammatory and immunoregulatory
agents, or as agents useful to decrease lipoprotein (a)
concentration in the blood or to increase feed
15 conversion efficiency.

This Sheet Has Been Replaced



23 OCT 1996

FIG. 1A

	ME- Myristic Acid 14:0	ME- Palmitic Acid 16:0	ME- Palmitoleic Acid 16:1
LIVER			
Normolipemic Swine			
I) Control Diet (CD)	31.23	134.46	3.16
Hypercholesterolemic Swine			
II) Control Diet (CD)	32.58	271.61	20.39
III) CD & TRF RB oil; 50 ppm	20.22	138.52	17.83
IV) CD & -Tocotrienol; 50 ppm	14.17	112.15	5.79
V) CD & P-21- Tocotrienol; 50 ppm	17.05	155.07	6.36
VI) CD & P-25- Tocotrienol; 50 ppm (P-25)	20.30	132.04	12.12
INTESTINE			
Normolipemic Swine			
I) Control Diet (CD)	34.67	142.66	---
Hypercholesterolemic Swine			
II) Control Diet (CD)	16.25	139.01	0.79
III) CD & TRF RB oil; 50 ppm	13.79	93.59	---
IV) CD & δ -Tocotrienol; 50 ppm	17.88	85.28	---
V) CD & P-21- Tocotrienol; 50 ppm	13.36	85.72	---
VI) CD & P-25- Tocotrienol; 50 ppm	16.84	106.87	---

23 OCT 1996

FIG. 1B

	ME- Palmitelaic Acid 16:1	ME- Stearic Acid 18:0	ME- Oleac Acid 18:1	ME- Linoleac Acid 18:2
LIVER				
Normolipemic Swine				
I) Control Diet (CD)	0.49	183.54	3.49	185.89
Hypercholesterolemic Swine				
II) Control Diet (CD)	17.87	94.45	189.89	5.12
III) CD & TRF RB oil; 50 ppm	---	140.69	150.07	---
IV) CD & Tocotrienol; 50 ppm	17.21	141.89	157.99	1.94
V) CD & P-21- Tocotrienol; 50 ppm	19.59	183.63	198.40	---
VI) CD & P-25- Tocotrienol; 50 ppm (P-25)	18.59	116.89	161.05	---
INTESTINE				
Normolipemic Swine				
I) Control Diet (CD)	14.62	177.46	141.49	---
Hypercholesterolemic Swine				
II) Control Diet (CD)	12.03	150.68	194.12	2.74
III) CD & TRF RB oil; 50 ppm	3.99	127.93	156.46	---
IV) CD & δ -Tocotrienol; 50 ppm	13.41	124.04	72.37	---
V) CD & P-21- Tocotrienol; 50 ppm	10.97	107.39	151.61	2.29
VI) CD & P-25- Tocotrienol; 50 ppm	17.89	115.53	71.65	0.73

23 OCT 1996

FIG. 1C

	ME- Linolenic Acid 18:3	ME- Arachidonic Acid 20:4	Total Fatty Acids
LIVER			
Normolipemic Swine			
I) Control Diet (CD)	0.68	0.4	543.36
Hypercholesterolemic Swine			
II) Control Diet (CD)	1.06	10.9	634.06
III) CD & TRF RB oil; 50 ppm	---	3.2	470.53
IV) CD & Tocotrienol; 50 ppm	1.94	1.50	454.58
V) CD & P-21- Tocotrienol; 50 ppm	5.08	1.02	603.25
VI) CD & P-25- Tocotrienol; 50 ppm (P-25)	2.08	1.08	482.15
INTESTINE			
Normolipemic Swine			
I) Control Diet (CD)	2.86	---	513.76
Hypercholesterolemic Swine			
II) Control Diet (CD)	---	1.28	516.90
III) CD & TRF RB oil; 50 ppm	---	1.02	396.78
IV) CD & δ -Tocotrienol; 50 ppm	---	0.96	313.94
V) CD & P-21- Tocotrienol; 50 ppm	---	0.78	372.12
VI) CD & P-25- Tocotrienol; 50 ppm	---	0.77	330.28

23 OCT 1996

FIG. 2A

	ME- Myristic Acid 14:0	ME- Palmitic Acid 16:0	ME- Palmitoleic Acid 16:1
LUNG			
Normolipemic Swine			
I) Control Diet (CD)	15.12	87.75	---
Hypercholesterolemic Swine			
II) Control Diet (CD)	---	122.15	---
III) CD & TRF RB oil; 50 ppm	21.49	115.94	---
IV) CD & Tocotrienol; 50 ppm	---	75.46	---
V) CD & P-21- Tocotrienol; 50 ppm	---	68.19	---
VI) CD & Didesmethyl- Tocotrienol; 50 ppm (P-25)	---	76.89	---
HEART			
Normolipemic Swine			
I) Control Diet (CD)	---	42.63	---
Hypercholesterolemic Swine			
II) Control Diet (CD)	4.86	27.52	---
III) CD & TRF RB oil; 50 ppm	---	44.65	---
IV) CD & Tocotrienol; 50 ppm	---	42.23	---
V) CD & P-21- Tocotrienol; 50 ppm	---	4.19	---
VI) CD & P-25- Tocotrienol 50 ppm	---	3.44	---

23 OCT 1996


FIG. 2B

	ME- Palmitelaic Acid 16:1	ME- Stearic Acid 18:0	ME- Oleac Acid 18:1
LUNG			
Normolipemic Swine			
I) Control Diet (CD)	12.02	137.93	64.69
Hypercholesterolemic Swine			
II) Control Diet (CD)	---	158.81	128.29
III) CD & TRF RB oil; 50 ppm	9.68	118.94	54.61
IV) CD & -Tocotrienol; 50 ppm	---	160.12	58.71
V) CD & P-21- Tocotrienol; 50 ppm	---	133.04	50.76
VI) CD & Didesmethyl- Tocotrienol; 50 ppm (P-25)	---	46.86	80.19
HEART			
Normolipemic Swine			
I) Control Diet (CD)	---	69.35	59.35
Hypercholesterolemic Swine			
II) Control Diet (CD)	11.87	49.18	47.43
III) CD & TRF RB oil; 50 ppm	---	99.32	53.42
IV) CD & -Tocotrienol; 50 ppm	---	84.30	64.16
V) CD & P-21- Tocotrienol; 50 ppm	6.13	21.56	18.53
VI) CD & P-25- Tocotrienol 50 ppm	4.73	12.38	12.92

THIS SHEET HAS BEEN
REPLACED

6/13

This Sheet Has Been Replaced



REPLACEMENT SHEET

23 OCT 1996

FIG. 2C

	ME- Linoleic Acid 18:2	ME- Linolenic Acid 18:3	ME- Arachidonic Acid 20:4	Total Fatty Acids
LUNG				
Normolipemic Swine				
I) Control Diet (CD)	---	---	---	317.51
Hypercholesterolemic Swine				
II) Control Diet (CD)	---	---	---	409.25
III) CD & TRF RB oil; 50 ppm	---	---	---	321.36
IV) CD & Tocotrienol; 50 ppm	3.32	---	---	297.61
V) CD & P-21- Tocotrienol; 50 ppm	---	---	---	251.99
VI) CD & Didesmethyl- Tocotrienol; 50 ppm (P-25)	3.39	3.32	---	310.65
HEART				
Normolipemic Swine				
I) Control Diet (CD)	---	---	---	171.33
Hypercholesterolemic Swine				
II) Control Diet (CD)	---	---	---	140.86
III) CD & TRF RB oil; 50 ppm	---	---	---	197.39
IV) CD & Tocotrienol; 50 ppm	---	---	---	190.69
V) CD & P-21- Tocotrienol; 50 ppm	---	---	---	50.41

23 OCT 1976

FIG. 3A

	ME- Myristic Acid 14:0	ME- Palmitic Acid 16:0	ME- Palmitoleic Acid 16:1
LOIN MUSCLE			
Normolipemic Swine			
I) Control Diet (CD)	23.51	125.97	---
Hypercholesterolemic Swine			
II) Control Diet (CD)	3.31	21.86	---
III) CD & TRF RB oil; 50 ppm	---	19.33	---
IV) CD & δ -Tocotrienol; 50 ppm	0.62	11.48	---
V) CD & P-21- Tocotrienol; 50 ppm	---	6.07	---
VI) CD & P-25- Tocotrienol; 50 ppm	3.08	8.49	---
ADIPOSE TISSUE			
Normolipemic Swine			
I) Control Diet (CD)	11.13	196.40	---
Hypercholesterolemic Swine			
II) Control Diet (CD)	45.07	283.62	---
III) CD & TRF RB oil; 50 ppm	---	152.85	---
IV) CD & δ -Tocotrienol; 50 ppm	---	9.26	---
V) CD & P-21- Tocotrienol; 50 ppm	---	75.06	---
VI) CD & P-25 Tocotrienol; 50 ppm	14.34	58.45	---

23 OCT 1996

FIG. 3B

	ME- Palmitelaic Acid 16:1	ME- Stearic Acid 18:0	ME- Oleac Acid 18:1
LOIN MUSCLE			
Normolipemic Swine			
I) Control Diet (CD)	5.28	359.83	185.80
Hypercholesterolemic Swine			
II) Control Diet (CD)	---	83.61	58.72
III) CD & TRF RB oil; 50 ppm	---	72.08	18.13
IV) CD & δ -Tocotrienol; 50 ppm	---	48.67	15.90
V) CD & P-21- Tocotrienol; 50 ppm	---	38.58	11.41
VI) CD & P-25- Tocotrienol; 50 ppm	1.37	12.39	5.87
ADIPOSE TISSUE			
Normolipemic Swine			
I) Control Diet (CD)	---	322.76	319.23
Hypercholesterolemic Swine			
II) Control Diet (CD)	21.87	160.42	209.76
III) CD & TRF RB oil; 50 ppm	7.21	192.17	231.54
IV) CD & δ -Tocotrienol; 50 ppm	9.95	231.54	184.89
V) CD & P-21- Tocotrienol; 50 ppm	8.22	146.77	16.80
VI) CD & P-25 Tocotrienol; 50 ppm	7.23	133.40	15.00

9/13

REPLACEMENT SHEET

23 OCT 1996

FIG. 3C

	ME- Linoleic Acid 18:2	ME- Linolenic Acid 18:3	ME- Arachidonic Acid 20:4	Total Fatty Acids
LOIN MUSCLE				
Normolipemic Swine				
I) Control Diet (CD)	---	---	---	700.39
Hypercholesterolemic Swine				
II) Control Diet (CD)	---	---	---	167.50
III) CD & TRF RB oil; 50 ppm	---	---	---	109.54
IV) CD & δ -Tocotrienol; 50 ppm	---	---	---	76.67
V) CD & P-21- Tocotrienol; 50 ppm	---	---	---	56.06
VI) CD & P-25- Tocotrienol; 50 ppm	---	---	---	31.20
ADIPOSE TISSUE				
Normolipemic Swine				
I) Control Diet (CD)	---	---	3.4	852.94
Hypercholesterolemic Swine				
II) Control Diet (CD)	---	---	7.8	728.54
III) CD & TRF RB oil; 50 ppm	---	---	2.2	585.98
IV) CD & δ -Tocotrienol; 50 ppm	---	---	2.0	355.91
V) CD & P-21- Tocotrienol; 50 ppm	---	---	2.0	248.85
VI) CD & P-25 Tocotrienol; 50 ppm	---	---	1.9	230.35

23 OCT 1996

FIG. 4 A

	ME- Myristic Acid 14:0	ME- Palmitic Acid 16:0	ME- Palmitoleic Acid 16:1
HAM MUSCLE			
Normolipemic Swine			
I) Control Diet (CD)	---	141.38	---
Hypercholesterolemic Swine			
II) Control Diet (CD)	3.41	24.15	---
III) CD & TRF RB oil; 50 ppm	---	38.92	---
IV) CD & δ -Tocotrienol; 50 ppm	10.62	45.04	---
V) CD & P-21- Tocotrienol; 50 ppm	5.35	25.21	---
VI) CD & P-25 Tocotrienol; 50 ppm	---	7.91	---

23 OCT 1996

FIG. 4B

	ME- Palmitelaic Acid 16:1	ME- Stearic Acid 18:0	ME- Oleac Acid 18:1
HAM MUSCLE			
Normolipemic Swine			
I) Control Diet (CD)	---	---	59.94
Hypercholesterolemic Swine			
II) Control Diet (CD)	---	149.91	21.59
III) CD & TRF RB oil; 50 ppm	---	80.04	---
IV) CD & δ -Tocotrienol; 50 ppm	---	130.08	---
V) CD & P-21- Tocotrienol; 50 ppm	---	---	---
VI) CD & P-25 Tocotrienol; 50 ppm	---	60.04	---

23 OCT 1996

FIG.4C

	ME- Linoleic Acid 18:2	ME- Linolenic Acid 18:3	ME- Arachidonic Acid 20:4	Total Fatty Acids
HAM MUSCLE				
Normolipemic Swine				
I) Control Diet (CD)	---	---	---	201.32
Hypercholesterolemic Swine				
II) Control Diet (CD)	---	---	---	199.06
III) CD & TRF RB oil; 50 ppm	---	---	---	118.96
IV) CD & δ -Tocotrienol; 50 ppm	---	---	---	158.74
V) CD & P-21- Tocotrienol; 50 ppm	---	---	---	30.56
VI) CD & P-25 Tocotrienol; 50 ppm	---	---	---	67.75

THIS SHEET HAS BEEN
REPLACED

13/13

REPLACEMENT SHEET

23 OCT 1996

FIG. 5A





FIG. 5B



14 MAR 1997

- 10 -

BRIEF DESCRIPTION OF THE DRAWINGS

Figures 1-4 depict the effects of different tocotrienols from rice bran oil on fatty acid concentrations in various tissues of hyper-
5 cholesterolemic swine.

Figure 5 depicts the effects of GT301 (γ -T₃) on superoxide production in human peripheral blood neutrophils.

10 In order that the invention herein described may be more fully understood, the following detailed description is set forth. In the description, the following terms are employed:

Biological source - Any natural or
15 recombinant plant source, natural or transgenic animal source, microbial source (e.g., bacteria), fungi, yeast, algae, higher plant source, or derivative thereof, which contains one or more tocopherol, tocotrienol or tocotrienol-like compounds.


20 Stabilization - A process effective to increase the recoverable amounts of tocopherols, tocotrienols and tocotrienol-like compounds in a biological source by one or a combination of:
(1) inactivating enzymes which are capable of degrading
25 tocotrienols and tocotrienol-like compounds in that biological source, (2) breaking bonds or otherwise interfering with the interactions -- such as hydrogen bonds, covalent bonds, ionic bonds, Van der Waals forces, London forces and hydrophobic or hydrophilic
30 interactions which bind the desired products to proteins, sugars, lipids, carbohydrates, glycoproteins, other biological molecules (i.e., amino acids or nucleotides) or other membrane components, or combinations thereof, in the biological source -- which

14 MAR 1997

salts, oxidation products and hydrolysis products thereof, wherein R_1 - R_{10} , Z , n and m are as defined as above.

As herein described above, the compounds of formula I do not include desmethyl-tocotrienol (3,4-dihydro-2-methyl-2-(4,8,12-trimethyltrideca-3'(E),7'(E),11'-trienyl)-2H-1-benzopyran-6-ol). The compounds of formula I also exclude α -, β -, γ - and δ -tocotrienol.

Typical hydrolysis products of the compounds of formula I which are also envisioned by this invention include, but are not limited to, compounds of formula II:



wherein R_1 - R_{10} , Z , n and m are as defined as above. Preferred compounds are the compounds of formula I, wherein R_1 , R_3 and Z are each independently H, halogen, OH, OCH_3 or CH_3 ; R_4 is $C=O$, CH_2 or NH ; R_5 is O, S, CH_2 or NH ; R_6 is H or CH_3 ; R_{10} and R_{11} are each independently H or CH_3 ; n is 0 or 1; and m is 3-10.

The most preferred compounds are the compounds of formula I, wherein R_1 , R_3 , Z , R_8 and R_9 are each hydrogen, R_2 is OH, R_4 is CH_2 , R_5 is CH_2 or O, R_6 is H or CH_3 , R_{10} is H or CH_3 , n is 1 and m is 3.

The novel compound, 3,4-dihydro-2-(4,8,12-trimethyltrideca-3'(E),7'(E),11' trienyl)-2H-1-benzopyran-6-ol ("P₂₅"), is representative of the class of compounds according to formula I. The structure of this compound is shown below:

14 MAR 1997

6

FIG. 2C

	ME- Linoleic Acid 18:2	ME- Linolenic Acid 18:3	ME- Arachidonic Acid 20:4	Total Fatty Acids
LUNG				
Normolipemic Swine				
I) Control Diet (CD)	---	---	---	317.51
Hypercholesterolemic Swine				
II) Control Diet (CD)	---	---	---	409.25
III) CD & TRF RB oil; 50 ppm	---	---	---	321.36
IV) CD & Tocotrienol; 50 ppm	3.32	---	---	297.61
V) CD & P-21- Tocotrienol; 50 ppm	---	---	---	251.99
VI) CD & Didesmethyl- Tocotrienol; 50 ppm (P-25)	3.39	3.32	---	310.65
HEART				
Normolipemic Swine				
I) Control Diet (CD)	---	---	---	171.33
Hypercholesterolemic Swine				
II) Control Diet (CD)	---	---	---	140.86
III) CD & TRF RB oil; 50 ppm	---	---	---	197.39
IV) CD & Tocotrienol; 50 ppm	---	---	---	190.69
V) CD & P-21- Tocotrienol; 50 ppm	---	---	---	50.41
VI) CD & P-25- Tocotrienol 50 ppm	---	---	---	33.47

14 MAR 1997

13

FIG. 5A





FIG. 5B



REPLACEMENT SHEET

01 JUL 1997

- 98 -

**TOCOTRIENOLS AND TOCOTRIENOL-LIKE COMPOUNDS AND
METHODS FOR THEIR USE****ABSTRACT**

The present invention relates to novel tocotrienols and tocotrienol-like compounds displaying biological activity. The tocotrienols and tocotrienol-like compounds of this invention may be conveniently obtained from biological sources or by chemical synthesis and may be used in pharmaceutical compositions, foodstuffs and dietary supplements. This invention also relates to the use of tocotrienols, tocotrienol-like compounds, and mixtures thereof, as hypocholesterolemic, antithrombotic, antioxidizing, antiatherogenic, antiinflammatory and immunoregulatory agents, or as agents useful to decrease lipoprotein (a) concentration in the blood or to increase feed conversion efficiency.

FIGURE 5A

1-1801 04