

Description

Field of the Invention

This invention relates to a chemical compound, namely [hexakis(pentenitrilo)nickel II]bis[μ-(cyano)bis-(triphenylborane)(I)]. When the pentenenitrile is 3-pentenitrile the compound has the formula $[\text{Ni}(\text{CH}_3\text{CH}=\text{CHCH}_2\text{CN})_6][\phi_3\text{BCNB}\phi_3]_2$, where ϕ is the phenyl radical. This compound can be prepared by the reaction of known nickel compounds with pentenenitrile. The compound is useful as a promoter in the preparation of adiponitrile by the hydrocyanation of 3 or 4-pentenitrile.

Background of the Invention

Cone U.S. Patent 4,394,321 discloses the compound $\text{Ni}[\text{NC}(\text{CH}_2)_4\text{CN}]_2[\text{NCB}\phi_3]_2$, hereinafter referred to as NCBC. This compound may be reacted in pentenenitrile to produce the compound of this invention.


Various organo boron compounds have been employed in the prior art to promote the hydrocyanation of 3,4-pentenitrile to adiponitrile -- see, for example, Shook U.S. Patent 4,082,811 where triarylborane is so employed, and the triarylborane recovered from the catalyst residue. The compound of this invention may be used in a manner like that taught in Shook to produce adiponitrile. Novel compounds of the formula $\text{Ni}[(\text{NH}_3)_4][\text{NC}(\text{B}(\text{C}_6\text{H}_4\text{R}))_3]_2$ where R is H, halogen, C_1 to C_6 alkyl or C_6 to C_{10} aryl, prepared by reacting a nickel compound with the cyanide adduct of a triarylborane, are disclosed in GB-A-2 047 680

Summary of the Invention

This invention is the chemical compound: [hexakis(pentenitrilo)nickel II]bis[μ-(cyano)bis-(triphenylborane)(I)]; the method of preparation of the compound; and the use of the compound in the hydrocyanation of 3 or 4-pentenitrile to form adiponitrile.

The compound of the invention consists of a cation $\text{Ni}(\text{CH}_3\text{CH}=\text{CHCH}_2\text{CN})_6^{+2}$ and two anions $[\phi_3\text{BCNB}\phi_3]^{-1}$. The cation has perfect octahedral symmetry. The anion has a threefold axis passing through the two boron atoms and the carbon atoms attached to the nitrogen atom.

The compound of the invention may be prepared by heating NCBC in pentenenitrile to a temperature above about 50° C. The reaction steps are believed to be as follows:



The compound of the invention may also be prepared by reacting triphenylborane with solutions of pentenenitrile containing $\text{Ni}(\text{CH}_3\text{CH}=\text{CHCH}_2\text{CN})_6(\text{NCB}\phi_3)_2$. Example 3 below shows the preparation by this process. The triphenylborane employed in this process can be that contained in the catalyst residue that results from the hydrocyanation of 3,4-pentenitrile using triphenylborane as a promoter. The reaction proceeds even though the triphenylborane exists in the residue in a state that is no longer active as a promoter.

The compound of the invention can also be prepared by the reaction of triphenylborane, sodium cyanide, pentenenitrile, and nickel dichloride dimethoxyethane. Example 4 below illustrates this method of preparation.

The process for preparing the compound of the invention proceeds satisfactorily at atmospheric pressure and at temperatures between 50 and 200° C. The ingredients that react to form the compound may be present in approximately stoichiometric amounts; however, the reaction will proceed with different ratios of the ingredients, for example, with the pentenenitrile being present in a large excess.

The compound of the invention has the I.R. spectra shown in the Figure 1, and the single crystal X-ray analysis shown in Figs. 2 and 3, Figs. 2 depicting the cation, and Fig. 3 depicting the anion.

Detailed Description of the InventionExample 15 Preparation from NCBC

In a glass reaction vessel that contains a magnetic stirrer, 4.25 g of NCBC was mixed with sufficient 3-pentenitrile to give 50 ml of slurry. The reactor was placed in a heat bath at 125°C, and the reaction allowed to proceed for 5 minutes after the temperature reached 120°C. The reactor was removed from the
 10 heat bath and placed in an ice bath to quench the reaction. The contents were then cooled to about 25°C. The contents were filtered to remove the black nickel-rich solid product, and a clear dark green product solution was isolated. The solution was placed in a refrigerator overnight, which caused solids to precipitate. Infrared analysis of the solids showed them to be the compound of the invention.

15 Example 2Preparation of Large Crystals of Product for Analysis

In a glass reaction vessel that contains a magnetic stirrer, 2.125 g of NCBC was mixed with sufficient 3-pentenitrile to give 50 ml of slurry. The reactor was placed in a heat bath at 125°C, and the reaction allowed to proceed for 5 minutes after the temperature reaches 120°C. The reactor was removed from the
 20 heat bath and placed in an ice bath to quench the reaction. The contents were cooled to 25°C. The contents were filtered to remove the black nickel-rich solids, and a dark green product solution was obtained. After standing for two weeks, large (~ 1 mm) cubic crystals formed which gave an infrared
 25 spectrum typical of $\text{Ni}(\text{CH}_3\text{CH}=\text{CHCH}_2\text{CN})_2(\phi_3\text{BNCB}\phi_3)_2$.

These crystals were analysed by X-ray diffraction to establish chemical structure, as shown in Figs. 2 and 3.

Example 330 Preparation from $\text{Ni}(\text{CH}_3\text{-CH}=\text{CH-CH}_2\text{-CN})_2(\text{NCB}\phi_3)_2$ and $\phi_3\text{B}$

A solution containing 1 g of $\text{Ni}(\text{CH}_3\text{-CH}=\text{CH-CH}_2\text{-CN})_2(\text{NCB}\phi_3)_2$ dissolved in 9 g of 3-pentenitrile was prepared. A second solution was prepared which contained 0.65 g of $\phi_3\text{B}$ dissolved in 9.35 g of 3-pentenitrile. One gram of each solution was mixed together, which caused the immediate precipitation of
 35 a light purple solid. The solid was washed with cyclohexane and dried. An infrared scan of the solid showed it to be the compound of the invention.

Example 440 Direct Synthesis from $\phi_3\text{B}$, NaCN, and $\text{NiCl}_2 \cdot \text{DME}$

A reaction was carried out by reacting 9.7 g of $\phi_3\text{B}$, 1.0 g NaCN, and 50 g of 3-pentenitrile. This mixture was heated to 60°C for 1 hr with stirring. The mixture was then filtered, and 2.2 g of $\text{NiCl}_2 \cdot \text{DME}$
 45 (i.e., nickel dichloro dimethoxyethane) was added to the filtrate. The resulting mixture was heated to 100°C and immediately filtered. The filtrate was allowed to cool to room temperature, which caused solid product to precipitate. Infrared analysis of the solid product showed it to be the product of the invention.

Example 550 Hydrocyanation Performance

A catalyst solution was prepared by dissolving 4.40 g of $\text{Ni}[\text{para-tolyl phosphite}]_4$ and 3.75 cc of paratolyl phosphite in 92 cc of 3-pentenitrile under a nitrogen atmosphere. To a 10 cc aliquot of this
 55 catalyst solution was added 0.240 g of $[\text{hexakis(pentenitrilo)nickel II}] \text{bis}[\mu\text{-(cyano)bis(triphenylborane)}(\text{I})]$, which corresponds to 0.5 moles per mole of $\text{Ni}[\text{para-tolyl phosphite}]_4$. The mixture was heated in a thermostated oil bath to the desired reaction temperature. Hydrogen cyanide was added to the reaction mixture by passing nitrogen gas at 5 cc/min through liquid hydrogen cyanide at 0°C, and then to the

reaction vessel just above the liquid level. The reactions were run until adiponitrile production ceased. Reactions were run at 40, 51, and 80 °C, and the results are given in the following table.

Temp (°C)	3-pentenitrile conversion (%)	Yields (%)			
		Adiponitrile	Methylglutaronitrile	Ethylsuccinonitrile	2-pentene nitrile
40	31	92.8	3.1	0.3	3.9
51	35	90.3	3.3	0.3	6.0
80	29	75.5	3.8	0.4	20.3

In the above table, yield % of adiponitrile is defined by the equation:

$$\text{Yield \%} = \frac{\text{adiponitrile}}{\text{adiponitrile} + \text{methylglutaronitrile} + \text{ethylsuccinonitrile} + \text{2-pentenitrile}} \times 100$$

As a comparative, hydrocyanations were performed using 1 mole of $\phi_3\text{B}$ per mole of $\text{Ni}[\text{para-tolyl phosphite}]_4$, all other conditions being the same. The results for $\phi_3\text{B}$ promoter are as follows:

Temp (°C)	3-pentenitrile conversion (%)	Yields (%)			
		Adiponitrile	Methylglutaronitrile	Ethylsuccinonitrile	2-pentene nitrile
40	29	93.4	2.9	0.3	3.4
51	29	90.3	3.3	0.3	6.1
80	30	76.7	3.8	0.4	19.1

Claims

- [Hexakis(pentenitrilo)nickel (II)]bis[μ -(cyano)bis(triphenylborane)(I)].
- A process for the preparation of [hexakis(pentenitrilo)nickel (II)]bis[μ -(cyano)bis(triphenylborane)(I)] which comprises reacting a mixture containing pentenenitrile and a member of the class consisting of $\text{Ni}(\text{NC}(\text{CH}_2)_4\text{CN})_2(\text{NCB}\phi_3)_2$ and triphenylborane.
- The process of claim 2 in which the member of the class is triphenylborane, and the mixture also contains $\text{Ni}(\text{CH}_3\text{CH}=\text{CHCH}_2\text{CN})_4(\text{NCB}\phi_3)_2$.
- The process of claim 2 in which the member of the class is triphenylborane, and the mixture also contains sodium cyanide and nickel dichloro dimethoxyethane.
- Compound according to Claim 1 in which said pentenenitrile radical is 3-pentenitrile.
- Use of a compound according to Claim 1 or Claim 5 as a promoter in the preparation of adiponitrile by the hydrocyanation of 3- or 4-pentenitrile.

Patentansprüche

- [Hexakis(pentennitrilo)nickel (II)]bis[μ -(cyano)bis(triphenylboran)(I)].
- Verfahren zur Herstellung von [Hexakis(pentennitrilo)nickel(II)]bis[μ -(cyano)bis(triphenylboran)(I)], bei dem ein Gemisch, enthaltend Pentennitril und ein Glied der Klasse, bestehend aus $\text{Ni}(\text{NC}(\text{CH}_2)_4\text{CN})_2(\text{NCB}\phi_3)_2$ und Triphenylboran, zur Reaktion gebracht wird.
- Verfahren nach Anspruch 2, bei dem das Glied der Klasse Triphenylboran ist, und das Gemisch auch $\text{Ni}(\text{CH}_3\text{CH}=\text{CHCH}_2\text{CN})_4(\text{NCB}\phi_3)_2$ enthält.



EP 0 248 643 B1

reaction vessel just above the liquid level. The reactions were run until adiponitrile production ceased. Reactions were run at 40, 51, and 80 ° C, and the results are given in the following table.

Temp (° C)	3-pentenitrile conversion (%)	Yields (%)			
		Adiponitrile	Methylglutaronitrile	Ethylsuccinonitrile	2-pentene nitrile
40	31	92.8	3.1	0.3	3.9
51	35	90.3	3.3	0.3	6.0
80	29	75.5	3.8	0.4	20.3

In the above table, yield % of adiponitrile is defined by the equation:

$$\text{Yield \%} = \text{adiponitrile} \times 100 / (\text{adiponitrile} + \text{methylglutaronitrile} + \text{ethylsuccinonitrile} + \text{2-pentenitrile})$$

As a comparative, hydrocyanations were performed using 1 mole of $\phi_3\text{B}$ per mole of $\text{Ni}[\text{para-tolyl phosphite}]_4$, all other conditions being the same. The results for $\phi_3\text{B}$ promoter are as follows:

Temp (° C)	3-pentenitrile conversion (%)	Yields (%)			
		Adiponitrile	Methylglutaronitrile	Ethylsuccinonitrile	2-pentene nitrile
40	29	93.4	2.9	0.3	3.4
51	29	90.3	3.3	0.3	6.1
80	30	76.7	3.8	0.4	19.1

Claims

- [Hexakis(pentenitrilo)nickel (II)]bis[μ -(cyano)bis(triphenylborane)(I)].
- A process for the preparation of [hexakis(pentenitrilo)nickel (II)]bis[μ -(cyano)bis(triphenylborane)(I)] which comprises reacting a mixture containing pentenenitrile and a member of the class consisting of $\text{Ni}(\text{NC}(\text{CH}_2)_4\text{CN})_2(\text{NCB}\phi_3)_2$ and triphenylborane.
- The process of claim 2 in which the member of the class is triphenylborane, and the mixture also contains $\text{Ni}(\text{CH}_3\text{CH}=\text{CHCH}_2\text{CN})_4(\text{NCB}\phi_3)_2$.
- The process of claim 2 in which the member of the class is triphenylborane, and the mixture also contains sodium cyanide and nickel dichloro dimethoxyethane.
- Compound according to Claim 1 in which said pentenenitrile radical is 3-pentenitrile.
- Use of a compound according to Claim 1 or Claim 5 as a promoter in the preparation of adiponitrile by the hydrocyanation of 3- or 4-pentenitrile.

Patentansprüche

- [Hexakis(pentennitrilo)nickel (II)]bis[μ -(cyano)bis(triphenylboran)(I)].
- Verfahren zur Herstellung von [Hexakis(pentennitrilo)nickel(II)]bis[μ -(cyano)bis(triphenylboran)(I)], bei dem ein Gemisch, enthaltend Pentennitril und ein Glied der Klasse, bestehend aus $\text{Ni}(\text{NC}(\text{CH}_2)_4\text{CN})_2(\text{NCB}\phi_3)_2$ und Triphenylboran, zur Reaktion gebracht wird.
- Verfahren nach Anspruch 2, bei dem das Glied der Klasse Triphenylboran ist, und das Gemisch auch $\text{Ni}(\text{CH}_3\text{CH}=\text{CHCH}_2\text{CN})_4(\text{NCB}\phi_3)_2$ enthält.



4. Verfahren nach Anspruch 2, bei dem das Glied der Klasse Triphenylboran ist und das Gemisch auch Natriumcyanid und Nickeldichlorodimethoxyethan enthält.

5. Verbindung nach Anspruch 1, bei dem das Pentennitrilradikal 3-Pentennitril ist.

6. Verwendung einer Verbindung nach Anspruch 1 oder Anspruch 5 als Promotor bei der Herstellung von Adiponitril durch die Hydrocyanierung von 3- oder 4-Pentennitril.

Revendications

1. L'[hexakis(pentanenitrilo)nickel(II)]bis[μ -(cyano) bis(triphénylborane) (I)].

2. Un procédé de préparation de l'[hexakis(pentanenitrilo)nickel(II)]bis[μ -(cyano)bis(triphénylborane) (I)] qui consiste à faire réagir un mélange contenant du pentènitrile et un membre de la classe constituée de $\text{Ni}(\text{NC}(\text{CH}_2)_4\text{CN})_2(\text{NCB}\Phi_3)_2$ et du triphénylborane.

3. Le procédé de la revendication 2 dans lequel le membre de la classe est le triphénylborane, et le mélange contient aussi du $\text{Ni}(\text{CH}_3\text{CH}=\text{CHCH}_2\text{CN})_4(\text{NCB}\Phi_3)_2$.

4. Le procédé de la revendication 2 dans lequel le membre de la classe est le triphénylborane, et le mélange contient aussi du cyanure de sodium et du nickel dichloro diméthoxyéthane.

5. Composé selon la revendication 1 dans lequel ledit radical de pentènitrile est le radical 3-pentènenitrile.

6. Utilisation d'un composé selon la revendication 1 ou 5 en tant que promoteur dans la préparation d'adiponitrile par hydrocyanation de 3- ou 4-pentènenitrile.