

PROCESSES FOR THE PREPARATION OF 3-(4-(2,4-DIFLUOROBENZYLOXY)-3-BROMO-6-METHYL-2-OXOPYRIDIN-1(2H)-YL)-N,4-DIMETHYLBENZAMIDE

5 BACKGROUND

3-(4-(2,4-Difluorobenzoyloxy)-3-bromo-6-methyl-2-oxopyridin-1(2H)-yl)-N,4-dimethylbenzamide is known to be useful as therapeutic agent for treating many pathological conditions, including the treatment or prevention of inflammatory and respiratory diseases. The efficacy of 3-(4-(2,4-difluorobenzoyloxy)-3-bromo-6-methyl-2-oxopyridin-1(2H)-yl)-N,4-dimethylbenzamide is believed to be related to its ability to inhibit p38 kinase.

It is believed that p38 α kinase can cause or contribute to the effects of, for example, inflammation generally; arthritis; neuroinflammation; pain; fever; pulmonary disorders; cardiovascular diseases; cardiomyopathy; stroke; ischemia; reperfusion injury; renal reperfusion injury; brain edema; neurotrauma and brain trauma; neurodegenerative disorders; central nervous system disorders; liver disease and nephritis; gastrointestinal conditions; ulcerative diseases; ophthalmic diseases; ophthalmological conditions; glaucoma; acute injury to the eye tissue and ocular traumas; diabetes; diabetic nephropathy; skin-related conditions; viral and bacterial infections; myalgias due to infection; influenza; endotoxic shock; toxic shock syndrome; autoimmune disease; bone resorption diseases; multiple sclerosis; disorders of the female reproductive system; pathological (but non-malignant) conditions, such as hemangiomas, angiofibroma of the nasopharynx, and avascular necrosis of bone; benign and malignant tumors/neoplasia including cancer; leukemia; lymphoma; systemic lupus erythematosus (SLE); angiogenesis including neoplasia; and metastasis.

WO03/068,230A1, published on August 21, 2003, describes 3-(4-(2,4-difluorobenzoyloxy)-3-bromo-6-methyl-2-oxopyridin-1(2H)-yl)-N,4-dimethylbenzamide, its preparation and use in the treatment of inflammation.


3-(4-(2,4-Difluorobenzoyloxy)-3-bromo-6-methyl-2-oxopyridin-1(2H)-yl)-N,4-dimethylbenzamide and closely related analogs can be made by the following described racemic route (Route 1).

In efforts to make the discovery route more efficient and a scalable process, several modifications have been introduced in Route 1. These not only include improved reaction conditions and workup procedures but also new reactions. The sequence of steps 2, 3, and 4 of the discovery route- alkylation, hydrolysis and bromination- was changed to bromination, alkylation and hydrolysis for a variety of reasons. These include avoiding tedious workup at each step, such as multiple extractions and concentrations and to obtain higher yields. Route 1 avoids the use of costly materials where possible.


SUMMARY

The present invention provides an improved process for the preparation of (-)-3-(4-(2,4-difluorobenzyloxy)-3-bromo-6-methyl-2-oxopyridin-1(2H)-yl)-N,4-dimethylbenzamide, or a pharmaceutically acceptable salt thereof comprising the steps:

a) contacting a compound of formula V:




with a brominating reagent in the presence of at least one solvent to produce a compound of Formula IV:




IV

b) contacting a compound of Formula IV with an hydrolase in the presence of a buffer solution to produce a compound of Formula X:



X

5 c) contacting a compound of Formula X with an activating reagent in the presence of at least one solvent and then contacting the resulting mixture with methylamine to produce a compound of Formula IX:



IX

; and

d) contacting a compound of Formula IX with a 2,4-difluorobenzylhalide in the presence of a base and at least one solvent;

wherein

R¹ is a C₁-C₆ alkyl or aryl.

The present invention also provides novel intermediates selected from:

- 5 methyl 3-(3-bromo-4-hydroxy-6-methyl-2-oxopyridin-1(2H)-yl-4-methylbenzoate;
(-)-3-(3-bromo-4-hydroxy-6-methyl-2-oxopyridin-1(2H)-yl-4-methylbenzoic acid;
and
(-)-3-(3-bromo-4-hydroxy-6-methyl-2-oxopyridin-1(2H)-yl-N,4-
dimethylbenzamide.

10

DETAILED DESCRIPTION

The compound and salts of this invention can be prepared from materials generally available in the art.

- The term "alkyl" means a straight or branched chain hydrocarbon radical
15 having from 1 to 8 carbon atoms and includes, for example, methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, n-hexyl, n-heptyl, n-octyl, and the like.

- The term "aryl" means an aromatic hydrocarbon ring system containing at least one aromatic ring. The aromatic ring may optionally be fused or otherwise
20 attached to other aromatic hydrocarbon rings or non-aromatic hydrocarbon rings. Examples of aryl groups include, for example, phenyl, naphthyl, 1,2,3,4-tetrahydronaphthyl, indanyl, and biphenyl.

The term "Group IA metal" is also known as an alkali metal and includes, for example, lithium, sodium and potassium.

- 25 The term "Group IIA metal" is also known as an alkaline earth metal and includes, for example, calcium, barium, strontium, magnesium.

The term "pharmaceutically-acceptable" is used adjectivally in this specification to mean that the modified noun is appropriate for use as a pharmaceutical product or as a part of a pharmaceutical product.

- 30 With reference to the use of the words "comprise" or "comprises" or "comprising" in this patent (including the claims) those words are used on the basis and clear understanding that they are to be interpreted inclusively, rather

than exclusively, and that Applicants intend each of those words to be so interpreted in construing this patent, including the claims below.

The following are definitions for various abbreviations used herein:

MHz is Megahertz.

5 Hz is Hertz.

J is Coupling Constant.

m/z is mass to charge ratio.

°C is degrees Celsius.

g is gram.

10 mg is milligrams.

mmole is millimoles.

mL is milliliters.

μL is microliters.

M is molar.

15 HPLC is high performance liquid chromatography.

"DMAC" is N,N-dimethylacetamide.

"DMAP" is dimethylaminopyridine.

"DMF" is dimethylformamide.

"DMI" is 1,3-dimethylimidazolidinone.

20 "DMPU" is 1,3-dimethyl-3,4,5,6-tetrahydro-2(1H)-pyrimidinone.

"DMSO" is dimethylsulfoxide.

"ee" is enantiomer excess.

"HCl" is hydrochloric acid.

"MTBE" is methyl tert-butyl ether.

25 "NMP" is 1-methyl-2-pyrrolidinone.

"TEA" is triethylamine.

"TFA" is trifluoroacetic acid.

"THF" is tetrahydrofuran.


30 GENERIC SYNTHESIS

This invention is directed to processes for the preparation of (-)-3-(4-(2,4-difluorobenzyloxy)-3-bromo-6-methyl-2-oxopyridin-1(2H)-yl)-N,4-dimethylbenzamide, or a pharmaceutically acceptable salt thereof.

The compound described in the invention is an atropisomer, i.e., a chiral rotational isomer. The need for chiral column chromatography has been eliminated by utilizing a selective enzymatic hydrolysis approach.

5 Preparation of a Compound of Formula V – Method 1

A process for the preparation of a compound of formula V comprises contacting a compound of formula VI with a compound of formula VII in the presence of at least one solvent.



- 10 wherein R^1 is a C_1 - C_6 alkyl or aryl; R^2 is H, halo or a C_1 - C_6 alkyl; and R^3 is halo or a C_1 - C_6 alkyl.

Typically a compound of formula VI is contacted with a compound of formula VII in the presence of at least one solvent and a base.

- 15 The process is typically carried out at a temperature from about 30°C to about 300°C. In one embodiment the temperature is from about 45°C to about 150°C. In another embodiment the temperature is from about 60°C to about 100°C.


- 20 The process is typically carried out for a time period from about 4 hours to about 60 hours. In one embodiment the time period is from about 2 hours to about 40 hours. In another embodiment the time period is from about 15 hours to about 25 hours.

- 25 The solvent is a polar solvent or a nonpolar solvent. Examples of useful polar solvents include ethylene glycol and trifluoroethanol. Examples of useful nonpolar solvents include dichlorobenzene, xylenes and diphenyl ether. In one embodiment the solvent is dichlorobenzene, xylenes, diphenyl ether, ethylene glycol, or trifluoroethanol. In another embodiment the solvent is trifluoroethanol or ethylene glycol. In another embodiment the solvent is ethylene glycol.

In one embodiment the base is an inorganic base or organic base. In one embodiment the base is a carbonate, bicarbonate or alkoxide of a Group IA or IIA metal, for example, potassium carbonate, potassium t-butoxide, or sodium bicarbonate. In one embodiment the base is, for example, a hindered tertiary amine such as N,N-diisopropylethylamine, triethylamine (TEA) or dimethylaminopyridine (DMAP). In one embodiment the base is potassium carbonate.

Preparation of a Compound of Formula V- Method 2

Another process for the preparation of a compound of formula V comprises contacting a compound of formula VII with a compound of formula VIII.



wherein R^1 is a C_1 - C_6 alkyl or aryl; R^2 is H, halo or a C_1 - C_6 alkyl; and R^3 is halo or a C_1 - C_6 alkyl.

Typically a compound of formula VII is contacted with a compound of formula VIII in the presence of at least one solvent and an acid.

The process is typically carried out at a temperature from about 0°C to about 250°C . In one embodiment the temperature is from about 5°C to about 100°C . In another embodiment the temperature is from about 15°C to about 60°C .

The process is typically carried out for a time period from about 0.01 hour to about 10 hours. In one embodiment the time period is from about 0.5 hour to about 7 hours. In another embodiment the time period is from about 1 hour to about 7 hours.


The solvent is a polar solvent or a nonpolar solvent. Useful examples of polar solvents include acetic acid, lower alkyl carboxylic acid and dimethylformamide. Useful examples of nonpolar solvents include dioxane,

tetrahydrofuran, methyl tert-butyl ether, diethyl ether, toluene and dichloromethane. In one embodiment the solvent is dioxane, tetrahydrofuran, methyl tert-butyl ether, diethyl ether, acetic acid or lower alkyl carboxylic acid, dichloromethane, or dimethylformamide. In one embodiment the solvent is
5 dioxane, tetrahydrofuran, acetic acid or dichloromethane. In one embodiment the solvent is dioxane.

In one embodiment the acid is an organic acid. In another embodiment the organic acid is a lower alkyl or a substituted lower alkyl carboxylic acid, or a sulfonic acid such as p-toluenesulfonic acid, or methanesulfonic acid. In one
10 embodiment the sulfonic acid is methanesulfonic acid.

A Compound of Formula VIII Synthesis

A process for preparing a compound of formula VIII comprises contacting Meldrum's acid with diketene in the presence of a base and at least one solvent.



The following procedure is similar to the one reported in the literature at Kang, J; Kim, Y; Park, M; Lee, C.; Kim, W. Synthetic Communications (1984), 14(3), 265-9.

The process is typically carried out at a temperature from about 0°C to
20 about 100°C. In one embodiment the temperature is from about 5°C to about 50°C. In another embodiment the temperature is from about 20°C to about 25°C.


The process is typically carried out for a time period from about 0.01 hour to about 25 hours. In one embodiment the time period is from about 0.5 hour to
25 about 10 hours. In another embodiment the time period is from about 1 hour to about 5 hours.

In one embodiment the base is an organic or inorganic base. Examples of useful inorganic bases include Group IA or IIA carbonates, bicarbonates or alkoxides, such as potassium carbonate, potassium t-butoxide, and sodium
30 bicarbonate. Examples of useful organic bases include triethylamine (TEA),

dimethylaminopyridine (DMAP), and N,N-diisopropylethylamine. In one embodiment the base is potassium carbonate, TEA or DMAP. In another embodiment the base is TEA.

5 Halogenation

A process for preparing a compound of formula IV comprises contacting a compound of formula V with a halogenating reagent.



wherein R¹ is a C₁-C₆ alkyl or aryl; R² is H, halo or a C₁-C₆ alkyl; and R³ is halo or a C₁-C₆ alkyl.

Typically a compound of formula V is contacted with a halogenating reagent in the presence of at least one solvent.

The process is typically carried out at a temperature from about -40°C to about 50°C. In one embodiment the temperature is from about -15°C to about 20°C. In another embodiment the temperature is from about -10°C to about 10°C.

The process is typically carried out for a time period from about 0.5 hour to about 10 hours. In one embodiment the time period is from about 1.5 hour to about 5 hours. In one embodiment the time period is from about 1 hour to about 3 hours.

In one embodiment the solvent is a polar solvent. Useful examples include acetonitrile, acetic acid, or an acetic acid containing a co-solvent such as water or lower alkyl alcohols. In one embodiment the solvent is, for example, acetonitrile, acetic acid, or an acetic acid containing a co-solvent such as water or lower alkyl alcohols. In one embodiment the solvent is an acetic acid/water solution. In another embodiment the solvent is acetonitrile.


The halogenating reagent can be neat or a solution. In one embodiment the halogenating reagent is a brominating or a chlorinating reagent. In one

embodiment the brominating or chlorinating reagent is, for example, phenyl triethylammonium tribromide, 1,3-dibromo-5,5-dimethylhydantoin, N-bromosuccinimide, pyridinium bromide, perbromide, bromine, dibromotriphenylphosphorane, bromine chloride, N-bromohydantoin, N-bromocaprolactam, N-chlorosuccinimide, sodium hypochlorite, chlorine, sulfuryl chloride, cupric bromide, phosphorus pentachloride, or t-butyl hypochlorite. In one embodiment the brominating or chlorinating reagent is 1,3-dibromo-5,5-dimethylhydantoin, bromine, bromine chloride or chlorine. In another embodiment the brominating reagent is bromine. In another embodiment the brominating reagent is 1,3-dibromo-5,5-dimethylhydantoin.

The halogenation may be conducted in the presence of an acid. In one embodiment the acid is an organic or inorganic acid. Examples of useful acids include tetrafluoroboric acid, hydrobromic acid, hydrochloric acid, hydrofluoric acid, sulfuric acid and phosphoric acid. In one embodiment the acid is tetrafluoroboric acid.

A process for the preparation of an enantiomerically enriched mixture or exclusively a single enantiomer of a compound of formula I is generally presented in Scheme 2.

Scheme 2



wherein X^1, X^2, X^3, X^4, X^5 and X^6 are independently H, halo or a C_1 - C_6 alkyl; R^1 is a C_1 - C_6 alkyl or aryl; R^2 is H, halo or a C_1 - C_6 alkyl; R^3 is halo or a C_1 - C_6 alkyl; and R^4 is H or a C_1 - C_6 alkyl; R^5 is H, a C_1 - C_6 alkyl or aryl; or R^4, R^5 , and the nitrogen to which they are attached form a morpholinyl, thiomorpholinyl,


5 piperidinyl, pyrrolidinyl, or piperazinyl ring which is optionally substituted with 1 or 2 groups that are independently C_1 - C_4 alkyl, C_1 - C_4 alkoxy, hydroxy, hydroxy C_1 - C_4 alkyl, C_1 - C_4 dihydroxyalkyl, or halogen.

Compounds of Formula V and IV can be prepared as previously described.

10

Protease Reaction

A process for preparing a compound of formula X comprises contacting a compound of formula IV with a hydrolase.



15 wherein R^1 is a C_1 - C_6 alkyl or aryl, R^2 is H, halo or a C_1 - C_6 alkyl and R^3 is halo or a C_1 - C_6 alkyl.

Typically a compound of formula IV is contacted with hydrolase in the presence of a buffer solution.

The process is typically carried out at a temperature from about 5°C to about 80°C. In one embodiment the temperature is from about 10°C to about 60°C. In another embodiment the temperature is from about 20°C to about 40°C.

The process is typically carried out for a time period from about 5 hours to about 100 hours. In one embodiment the time period is from about 25 hours to about 75 hours. In another embodiment the time period is from about 30 hours to about 60 hours.

The process is typically carried out at a pH of about 6 to about 12. In one embodiment the pH is from about 8 to about 11. In another embodiment the pH is from about 9 to about 10. In another embodiment the pH is about 9.

In one embodiment the buffer solution is an inorganic buffer solution. In another embodiment the buffer solution is, for example, potassium phosphate buffer solution or an inorganic bicarbonate buffer. In one embodiment the buffer solution is dibasic potassium phosphate buffer solution. Typically the molarity of the buffer solution is about 1 M.

In one embodiment the appropriate enzymatic solution is a hydrolase such as lipase or protease. In one embodiment the hydrolase produces the (-) enantiomer of a compound of Formula X in excess. In one embodiment particularly useful hydrolases are those that produce the (-) enantiomer in at least 80% enantiomeric excess. In another embodiment particularly useful hydrolases are those that produce the (-) enantiomer in at least 85% enantiomeric excess. In another embodiment particularly useful hydrolases are those that produce the (-) enantiomer in at least 90% enantiomeric excess. In another embodiment particularly useful hydrolases are those that produce the (-) enantiomer in at least 95% enantiomeric excess. In another embodiment particularly useful hydrolases are those that produce the (-) enantiomer in at least 99% enantiomeric excess. In one embodiment the hydrolase is Bacillus sp. protease solution (Savinase®, Novozyme, Bagsvaerd, Denmark).

In one embodiment the hydrolase produces a compound of Formula X as a racemic mixture. In another embodiment the hydrolase produces the (+) enantiomer of a compound of Formula X in excess.

In one embodiment the (-) enantiomer of a compound of formula X is preferred. The (+) enantiomer of a compound of formula IV can be isolated and removed. The (+) enantiomer can be recycled by thermal isomerization to the racemic mixture in a typical aprotic organic solvent at a temperature in a range from about 100°C to about 300°C.

Typically the (+) enantiomer of a compound of formula IV is contacted with a high boiling aprotic solvent, halogenated or substituted aromatic or aliphatic solvent.


The high boiling aprotic solvent, halogenated or substituted aromatic or aliphatic solvent can be chlorobenzene, anisole, dioxane, NMP, N,N-dimethylformamide, or dimethylsulfoxide.

The process is typically carried out at a temperature from about 0°C to about 200°C. In one embodiment the temperature is from about 75°C to about 150°C. In another embodiment the temperature is from about 100°C to about 140°C

The process is typically carried out for a time period from about 30 hours to about 200 hours. In one embodiment the time period is from about 20 hours to about 100 hours. In another embodiment the time period is from about 50 hours to about 75 hours.

Amidation

A process for preparing a compound of formula IX comprises reacting a compound of formula X with an appropriate activating agent, and then contacting the resulting mixture with an appropriate primary or secondary amine to produce a compound of formula IX.



wherein X^1 is H, halo or a C_1 - C_6 alkyl; R^2 is H, halo or a C_1 - C_6 alkyl; R^3 is halo or a C_1 - C_6 alkyl; and R^4 is H or a C_1 - C_6 alkyl; R^5 is H, a C_1 - C_6 alkyl or aryl; or R^4 , R^5 , and the nitrogen to which they are attached form a morpholinyl, thiomorpholinyl, piperidinyl, pyrrolidinyl, or piperazinyl ring which is optionally substituted with 1 or 2 groups that are independently C_1 - C_4 alkyl, C_1 - C_4 alkoxy, hydroxy, hydroxy C_1 - C_4 alkyl, C_1 - C_4 dihydroxyalkyl, or halogen.

In one embodiment the compounds of Formulae X and IX are the (-) enantiomers.

Typically a compound of formula X is contacted with an activating reagent

in the presence of at least one solvent and then the resulting mixture is contacted with an amine.

The process is typically carried out at a temperature from about 0°C to about 100°C. In one embodiment the temperature is from about 10°C to about 50°C. In one embodiment the temperature is from about 15°C to about 30°C.

The process is typically carried out for a time period from about 0.1 hour to about 10 hours. In one embodiment the time period is from about 0.5 hour to about 5 hours. In another embodiment the time period is from about 1 hour to about 3 hours.

In one embodiment the activating reagent is carbonyldiimidazole, dicyclohexylcarbodiimide, thionyl chloride, oxalyl chloride, phosphorus oxychloride, phosphorus trichloride, phosphorus pentachloride, phenyl triethylammonium tribromide, pyridinium bromide, perbromide, bromine, dibromotriphenylphosphorane, bromine chloride, N-bromohydantoin or N-bromocaprolactam. In one embodiment the activating reagent is carbonyldiimidazole, dicyclohexylcarbodiimide, thionyl chloride, oxalyl chloride, phosphorus oxychloride, phosphorus trichloride or phosphorus pentachloride. In one embodiment the reagent is oxalyl chloride. In another embodiment the activating reagent is carbonyldiimidazole.

The solvent is a polar solvent or a nonpolar solvent. Useful examples of polar solvents include dimethylformamide, acetic acid and lower alkyl carboxylic acid. Examples of useful nonpolar solvents include dichloroethane, tetrahydrofuran, dioxane, methyl tert-butyl ether, diethylether and toluene. In one embodiment the solvent is, for example, dimethylformamide, dichloroethane, tetrahydrofuran, dioxane, methyl tert-butyl ether, or toluene. In one embodiment the solvent is dimethylformamide, dichloroethane, tetrahydrofuran or dioxane. In another embodiment the solvent is dichloroethane. In one embodiment the solvent is dimethylformamide.


In one embodiment the amine is HNR^4R^5 wherein R^4 is H or a $\text{C}_1\text{-C}_6$ alkyl; R^5 is H, a $\text{C}_1\text{-C}_6$ alkyl or aryl; or R^4 , R^5 , and the nitrogen to which they are attached form a morpholinyl, thiomorpholinyl, piperidinyl, pyrrolidinyl, or piperazinyl ring which is optionally substituted with 1 or 2 groups that are independently $\text{C}_1\text{-C}_4$ alkyl, $\text{C}_1\text{-C}_4$ alkoxy, hydroxy, hydroxy $\text{C}_1\text{-C}_4$ alkyl, $\text{C}_1\text{-C}_4$

dihydroxyalkyl, or halogen. In one embodiment R^4 is H. In one embodiment R^5 is a C_1 - C_6 alkyl. In one embodiment, the amine is methylamine. Typically the amine is combined with a solvent prior to addition. In one embodiment the solvent is methanol, ethanol, tetrahydrofuran or water. In another embodiment the solvent is methanol, tetrahydrofuran or water. In one embodiment the solvent is tetrahydrofuran. In one embodiment the amine is a solution of methylamine in tetrahydrofuran.

Optionally an acid can be used to work up the reaction. In one embodiment the acid is an organic or inorganic acid. Examples of useful acids include acetic acid, citric acid, HCl and sulfuric acid. In one embodiment the acid is acetic acid, citric acid, HCl, or sulfuric acid. In one embodiment the acid is HCl, acetic acid or sulfuric acid. In one embodiment the acid is HCl.

Benzylation

A process for the preparation of a compound of formula I comprises reacting a compound of formula IX with a substituted benzylhalide.



wherein X^1 , X^2 , X^3 , X^4 , X^5 and X^6 are independently H, halo or a C_1 - C_6 alkyl; R^2 is H, halo or a C_1 - C_6 alkyl; R^3 is halo or a C_1 - C_6 alkyl; and R^4 is H or a C_1 - C_6 alkyl; R^5 is H, a C_1 - C_6 alkyl or aryl; or R^4 , R^5 , and the nitrogen to which they are attached form a morpholinyl, thiomorpholinyl, piperidinyl, pyrrolidinyl, or piperazinyl ring which is optionally substituted with 1 or 2 groups that are independently C_1 - C_4 alkyl, C_1 - C_4 alkoxy, hydroxy, hydroxy C_1 - C_4 alkyl, C_1 - C_4 dihydroxyalkyl, or halogen.

In one embodiment the compounds of Formulae IX and I are the (-) enantiomers.

Typically a compound of formula IX is contacted with a substituted


benzylhalide in the presence of a base and at least one solvent.

The process is typically carried out at a temperature from about 0°C to about 200°C. In one embodiment the temperature is from about 25°C to about 100°C. In another embodiment the temperature is from about 50°C to about

5 75°C.

The process is typically carried out for a time period from about 0.5 hour to about 20 hours. In one embodiment the time period is from about 1 hour to about 10 hours. In another embodiment the time period is from about 2 hours to about 4 hours.

10 In one embodiment the substituted benzylhalide is:



wherein halo is chloride, bromide or iodide; and X^2 , X^3 , X^4 , X^5 and X^6 are independently H, halo, or a C_1 - C_6 alkyl. In one embodiment the halo-substituted benzylhalide is, for example, 2,4-difluorobenzylbromide, 2,3-

15 difluorobenzylbromide, 2,5-difluorobenzylbromide, 2,4-difluorobenzylchloride, 2,3-difluorobenzylchloride or 2,5-difluorobenzylchloride. In another embodiment the halo-substituted benzylchloride is 2,4-difluorobenzylchloride. In another embodiment the halo substituted benzyl halide is 2,4-difluorobenzylbromide.

In one embodiment the base is an organic or inorganic base. Examples of
20 useful inorganic bases include Group IA or IIA carbonates, bicarbonates or alkoxides, such as potassium carbonate, potassium t-butoxide, and sodium bicarbonate. Examples of useful organic bases include triethylamine (TEA) or dimethylaminopyridine (DMAP); and a hindered amine such as N,N-diisopropylethylamine. In one embodiment the base is potassium carbonate,
25 potassium t-butoxide or sodium bicarbonate. In another embodiment the base is potassium carbonate.

In one embodiment the solvent is a polar aprotic solvent. In one embodiment the polar aprotic solvent is, for example, N,N-dimethylformamide (N,N-dimethylformamide), N,N-dimethylacetamide (DMAC) or N-

methylpyrrolidinone (NMP), 1,3-dimethylimidazolidinone (DMI), 1,3-dimethyl-3,4,5,6-tetrahydro-2(1H)-pyrimidinone (DMPU). In another embodiment the polar aprotic solvent is N,N-dimethylformamide. In another embodiment the solvent is N-methylpyrrolidinone.

5 In one embodiment the product is purified by trituration or precipitation using an appropriate solvent and/or co-solvent system. In one embodiment the solvent is methanol, 1-butanol, ethanol, ethyl acetate or 2-propanol. In one embodiment the solvent is methanol, ethanol or ethyl acetate. In one embodiment the solvent is methanol.

10

The present invention provides novel intermediates. In one embodiment the compound is selected from methyl 3-(3-bromo-4-hydroxy-6-methyl-2-oxopyridin-1(2H)-yl-4-methylbenzoate; (-)-3-(3-bromo-4-hydroxy-6-methyl-2-oxopyridin-1(2H)-yl-4-methylbenzoic acid;

15 and

(-)-3-(3-bromo-4-hydroxy-6-methyl-2-oxopyridin-1(2H)-yl-N,4-dimethylbenzamide.

The compounds of this invention may be used in the form of salts derived from inorganic or organic acids. Depending on the particular compound, a salt of
20 the compound may be advantageous due to one or more of the salt's physical properties, such as enhanced pharmaceutical stability in differing temperatures and humidities, or a desirable solubility in water or oil. In some instances, a salt of a compound also may be used as an aid in the isolation, purification, and/or resolution of the compound.


25 Non-toxic pharmaceutically acceptable salts include, but are not limited to salts of inorganic acids such as hydrochloric, sulfuric, phosphoric, diphosphoric, hydrobromic, and nitric or salts of organic acids such as formic, citric, malic, maleic, fumaric, tartaric, succinic, acetic, lactic, methanesulfonic, p-toluenesulfonic, 2-hydroxyethylsulfonic, salicylic and stearic. Similarly,
30 pharmaceutically acceptable cations include, but are not limited to sodium, potassium, calcium, aluminum, lithium and ammonium. Those skilled in the art will recognize a wide variety of non-toxic pharmaceutically acceptable addition salts.

Atropisomers are chiral rotational isomers. These compounds can be, for example, racemates, chiral non-racemic or diastereomers. In these situations, the single enantiomers, i.e., optically active forms, can be obtained by asymmetric synthesis or by resolution of the racemates. Resolution of the

- 5 racemates can be accomplished, for example, by conventional methods such as crystallization in the presence of a resolving agent; chromatography, using, for example a chiral HPLC column; thermal or kinetic resolution; or derivatizing the racemic mixture with a resolving reagent to generate diastereomers, separating the diastereomers via chromatography or selective crystallization, and removing
10 the resolving agent to generate the original compound in enantiomerically enriched form. Any of the above procedures can be repeated or combined to increase the enantiomeric purity of a compound.

The following illustration generically shows a compound (Z) that can exist as atropisomers as well as its two possible atropisomers (A) and (B). This

- 15 illustration also shows each of atropisomers (A) and (B) in a Fischer projection. In this illustration, R^3 and X^1 carry the same definitions as set forth for Formula I, $R_{p'}$ is a substituent within the definition of R^2 , R_p is a substituent within the definition of $CONR^4R^5$, and D represents the substituted benzyloxy.



When enantiomerically enriched, one enantiomer is present in greater amounts than the other and the extent of enrichment can be defined by an expression of enantiomer excess ("ee"), which is defined as $100(2x-1)$ wherein x is the mole fraction of the dominant enantiomer in the enantiomer mixture (e.g., an ee of 20% corresponds to a 60:40 ratio of enantiomers).

Detailed Examples


The following examples are merely illustrative.

10


Reference Example 1

Schemes 7 and 8 depict the overall syntheses that are described in Reference Example 1.

Scheme 7




Scheme 8



The racemic process to prepare (-)-3-[3-bromo-4-[(2,4-difluorobenzyl)oxy]-6-methyl-2-oxopyridin-1(2H)-yl]-N,4-dimethylbenzamide (**1**) is described as follows:


Synthesis of methyl 3-(4-hydroxy-6-methyl-2-oxopyridin-1(2H)-yl)-4-methylbenzoate (**5**) Using Pyrone **6**:




4-hydroxy-6-methylpyranone (**6**) (1.18kg, 9.34 moles), methyl-3-amino-4-methyl benzoate (**7**) (1.0kg, 6.21 moles), catalytic K_2CO_3 (102 gm, 0.74 moles) and 2L of trifluoroethanol were mixed and heated to 80-87 °C, under nitrogen for

22 hours. After the completion of reaction, the mixture was cooled to 65 °C and 11.3 L ethyl acetate was added and the solution gradually cooled to 5 -10°C. A product was collected, washed with 2L ethyl acetate to give about 55% isolated yield of methyl 3-(4-hydroxy-6-methyl-2-oxopyridin-1(2H)-yl)-4-methylbenzoate.

Alternatively methyl 3-(4-hydroxy-6-methyl-2-oxopyridin-1(2H)-yl)-4-methylbenzoate (**5**) can be prepared using 5-(1-hydroxy-3-oxobutylidene)-2,2-dimethyl-1,3-dioxane-4,6-dione (**8**) as follows:




A mixture of methyl 3-amino-4-methylbenzoate (**7**) (2.0 g, 12.1 mmoles) and 5-(1-hydroxy-3-oxobutylidene)-2,2-dimethyl-1,3-dioxane-4,6-dione (**8**) (3.86 g, 16.94 mmol, 1.4 eq.) in dioxane (20 mL) was heated at reflux for 5 minutes and cooled to 50 °C. The mixture was then treated with 1.16 g of methanesulfonic acid and heated at reflux for five minutes. Liquid chromatography indicated the reaction (cyclization) was essentially complete. The mixture was poured in 60 mL of crushed ice-water and stirred for 1.5 hours. The precipitate was filtered, washed with water and air dried to give 2.47 g (75% yield) of the product.




The above procedure is a modification of the procedure described in the literature at Kang, J; Kim, Y; Park, M; Lee, C.;Kim, W. Synthetic Communications (1984), 14(3), 265-9. In essence Meldrum's acid was reacted with diketene in dichloromethane in the presence of TEA at 20-25 °C for 2 hours to give 80% of 5-(1-hydroxy-3-oxobutylidene)-2,2-dimethyl-1,3-dioxane-4,6-dione (**8**).

Synthesis of methyl 3-(3-bromo-4-hydroxy-6-methyl-2-oxopyridin-1(2H)-yl)-4-methylbenzoate (4)



A suspension of methyl 3-(4-hydroxy-6-methyl-2-oxopyridin-1(2H)-yl)-4-methylbenzoate (**5**) (1.6 kg, 5.85 moles) in 6.4L acetic acid/1.6L water and cooled to 15 °C. A solution of 973g bromine in 1.6L acetic acid was prepared and added slowly to the reaction via dropping funnel. During the addition of bromine, the mixture became homogeneous followed by the formation of a white precipitate. After the addition was completed, the mixture was stirred for another 15 minutes. The mixture was diluted with 16L water; and the product was filtered and washed with 12L water followed by 9.6L cold acetonitrile (0-5 °C). The solid was dried to yield 1.85 kg (90%) product. ¹H NMR (300 MHz, DMSO-d₆) δ 1.80 (s, 3H), 2.04 (s, 3H), 3.86 (s, 3H), 6.15 (s, 1H), 7.59 (d, *J* = 8.1 Hz), 7.73 (s, 1H), 7.98 (d, *J* = 8.1, 1H), 11.6 (bs, 1H). Anal. Calcd. for C₁₅H₁₄BrNO₄: C, 51.16; H, 4.01; N, 3.98. Found: C, 50.83; H, 4.02; N, 4.00.

Synthesis of methyl 3-(4-(2,4-difluorobenzyloxy)-3-bromo-6-methyl-2-oxopyridin-1(2H)-yl)-4-methylbenzoate (3)




Methyl 3-(3-bromo-4-hydroxy-6-methyl-2-oxopyridin-1(2H)-yl)-4-methylbenzoate (**4**) (1.9kg, 5.4 moles), N,N-dimethylformamide (4.8L) and

powdered K_2CO_3 (1.1kg, 8.1 moles) were mixed together and heated to 55 °C under nitrogen. 2,4-difluorobenzyl chloride (964g, 5.9 moles) was added at a rate such that the temperature was maintained below 65°C. After the addition, the mixture was heated at 65 °C for 3.5 hours. After 3 hours the reaction was

complete and the reaction mixture was cooled to 20-25 °C with the addition of 19L water. The solid was filtered and washed with 15L water. The crude product was then purified by trituration in refluxing methanol (4L). The mixture was cooled and the solid filtered to yield 2.32 kg (90%) product. 1H NMR (300 MHz, $CDCl_3$) δ 1.90 (s, 3H), 2.11 (s, 3H), 3.87 (s, 3H), 5.25 (s, 2H), 6.11 (s, 1H), 6.88 (dt, J = 4.8, 1.8 Hz, 1H), 6.96 (t, J = 4.8 Hz, 1H), 7.42 (d, J = 6 Hz, 1H), 7.59 (q, J = 4.8 Hz, 1H), 7.74 (s, 1H), 8.02 (d, J = 6 Hz, 1H). Anal. Calcd. for $C_{22}H_{18}BrF_2NO_4$: C, 55.25; H, 3.79; N, 2.93. Found: C, 55.34; H, 3.83; N, 3.14.

Preparation of 3-(4-(2,4-difluorobenzoyloxy)-3-bromo-6-methyl-2-oxopyridin-


1(2H)-yl)-4-methylbenzoic acid (**2**)



Methyl 3-(4-(2,4-difluorobenzoyloxy)-3-bromo-6-methyl-2-oxopyridin-1(2H)-yl)-4-methylbenzoate (**3**) (3.4kg, 7.1 moles), 2.5M NaOH (3.1L, 7.8 moles), CH_3CN (12L) and 8.6L water were mixed together and heated to 60 °C, under nitrogen. Once the mixture became homogeneous the mixture was stirred for an additional hour. The reaction mixture was cooled to 20°C and then treated with 865 ml concentrated HCl at a rate such that the temperature was maintained below 25°C. After the HCl addition was completed the mixture was stirred for another hour. The product was filtered and washed with 12L CH_3CN , dried to give 2.87 kg (87%) adduct. 1H NMR (300 MHz, CD_3OD) δ 7.87 (dd, J = 7.8, 1.6 Hz, 1H), 7.82 (d, J = 1.8 Hz, 1H), 7.69 (q, J = 8.1 Hz, 1H), 7.57 (d, J = 8.1 Hz, 1H), 7.09 (dt, J = 2.2, 8.6 Hz, 1H), 6.7 (s, 1H), 5.4 (s, 2H), 2.14 (s, 3H), 2.02 (s,

3H) (d, $J = 2.4$ Hz, 1H), 3.94 (s, 3H), 2.15 (s, 3H), 1.91 (s, 3H). ES-HRMS m/z 464.0275 ($M+H$ calcd for $C_{21}H_{17}BrF_2NO_4$ requires 464.0304). Anal. Calcd. for $C_{21}H_{16}BrF_2NO_4$: C, 54.33; H, 3.47; N, 3.02. Found: C, 54.40; H, 3.42; N, 3.17

5 Preparation of (-)-3-(4-(2,4-difluorobenzoyloxy)-3-bromo-6-methyl-2-oxopyridin-1(2H)-yl)-N,4-dimethylbenzamide (1)



3-(4-(2,4-difluorobenzoyloxy)-3-bromo-6-methyl-2-oxopyridin-1(2H)-yl)-4-methylbenzoic acid (**2**)(1.4kg, 3.0 moles) was suspended in 7L tetrahydrofuran with 28 ml N,N-dimethylformamide and chilled to 5°C. Oxalyl chloride (368 ml, 4.22 moles) was added to the reactor at a rate such that the temperature was maintained below 10°C and the gas evolution was under control. After the oxalyl chloride had been added the reaction mixture was allowed to warm to room temperature. The reaction was completed when the mixture became homogeneous. The reaction mixture was drained into separate container and set aside. 40% methylamine (4.4L, 50.3 moles) and 2.6L water were added to the reactor and chilled to below 5°C. The reaction mixture was then added to the chilled methylamine solution at a rate such that the temperature was maintained below 15°C. Once the addition was completed the mixture was allowed to warm to room temperature and stirred an additional 30 minutes. 12.6L water was added to the mixture and the stirring was continued for another hour. The product was filtered and washed with 10L water to give 1.25 kg (90%). 1H NMR (300 MHz, $DMSO-d_6$) δ 7.95 (bs, 1H), 7.88 (dd, $J = 1.2, 5.7$ Hz, 1H), 7.68 (q, $J = 5.1$ Hz, 1H), 7.64 (d, $J = 1.2$ Hz, 1H), 7.46 (d, $J = 6.0$ Hz, 1H), 7.36(bs, 1H), 7.29 (dt, $J = 1.8, 7.8$ Hz, 1H), 7.16 (dt, $J = 1.8, 76.0$ Hz, 1H), 6.70 (s, 1H), 6.18 (s, 1H), 5.33 (s, 2H), 3.32 (s, 3H), 1.97 (s, 3H), 1.88 (s, 3H). Anal.

Cald. for $C_{21}H_{17}BrF_2N_2O_3 + 0.66 \text{ EtOH} + 0.1 \text{ H}_2\text{O}$: C, 54.11; H, 4.30; N, 5.65.


Found: C, 54.03; H, 4.59; N, 5.79

The racemic product 3-(4-(2,4-difluorobenzoyloxy)-3-bromo-6-methyl-2-oxypyridin-1(2H)-yl)-N,4-dimethylbenzamide (**1**) produced was then chromatographed using a chiral stationary phase such as Chiralcel OJ™ or Chiralpak® AD™ (Daicel Chemical Industries, Japan) with methanol or ethanol as the mobile phase to isolate the corresponding (-)-3-(4-(2,4-difluorobenzoyloxy)-3-bromo-6-methyl-2-oxypyridin-1(2H)-yl)-N,4-dimethylbenzamide (**1**).

Example 2

Scheme 9 depicts the overall synthesis that is described in Example 2.


Scheme 9



Following the development of a process for the racemic 3-(4-(2,4-difluorobenzoyloxy)-3-bromo-6-methyl-2-oxypyridin-1(2H)-yl)-N,4-dimethylbenzamide, a process for the direct preparation of (-)-3-(4-(2,4-difluorobenzoyloxy)-3-bromo-6-methyl-2-oxypyridin-1(2H)-yl)-N,4-

dimethylbenzamide has been developed and this is described as follows:


Step 1



- 5 4-hydroxy-6-methylpyranone (**6**) (6.71 kg, 53.2 moles), methyl-3-amino-4-methyl benzoate (**7**) (4.50kg, 27.2 moles), catalytic K_2CO_3 (0.54 kg, 3.91 moles), acetonitrile (6.14 L) and ethylene glycol (6.14 L) were mixed under nitrogen and the acetonitrile was removed by vacuum distillation. The reaction was stirred for 18-24 hours at 65°C. Following the reaction the mixture was cooled to 25°C. A
- 10 1:1 acetonitrile/water solution (6.14 L each) was added to the mixture. The mixture was stirred for 90 minutes and cooled to 0°C. A product was collected, washed with a 1:1 acetonitrile/water solution (3.4 L each) and dried to give 60% isolated yield of methyl 3-(4-hydroxy-6-methyl-2-oxopyridin-1(2H)-yl)-4-methylbenzoate (**5**).

15


Step 2



- A solution of 1,3-dibromo-5,5-dimethylhydantoin (1.375 kg, 4.81 moles) in acetonitrile (11.5 L) was added to a solution of methyl 3-(4-hydroxy-6-methyl-2-oxopyridin-1(2H)-yl)-4-methylbenzoate (**5**) (2.50 kg, 9.15 mol) in acetonitrile
- 20 (4.8L). The solution was added at a rate that maintained the temperature less than -10 °C for 90 to 180 minutes. A portion of the acetonitrile (approximately 6 L) was removed after completion of the reaction via distillation. The product was precipitated by the addition of water (7.9 L) and filtered to yield 2.90kg (90%) of


methyl 3-(3-bromo-4-hydroxy-6-methyl-2-oxopyridin-1(2H)-yl)-4-methylbenzoate
(4)

Step 3




(+/-)Methyl 3-(3-bromo-4-hydroxy-6-methyl-2-oxopyridin-1(2H)-yl)-4-methylbenzoate (**4**) (1kg, 2.84 moles) was mixed with 1M dibasic potassium phosphate buffer solution (dibasic potassium phosphate (3.05 kg, 17.5 moles) and water (16.4 L)) and warmed to 25°C. The pH of the solution was adjusted to 9.1 with 10% NaOH solution (about 4.2 L) followed by the addition of Savinase® enzyme (Novozyme, Bagsvaerd, Denmark) and warmed to 30°C. After stirring for about 40-45 hours, the pH of the solution was adjusted to 6.0 using 6N HCl (2.3 L) and stirred for 30 minutes. The resulting precipitate was the unreacted enantiomer of the ester methyl (+)3-(3-bromo-4-hydroxy-6-methyl-2-oxopyridin-1(2H)-yl)-4-methylbenzoate (423gm), which was isolated by filtration and washed with water. The aqueous filtrate was washed with 5.0 L methylene chloride. It was then further acidified to pH 3.6 with 2.1 L of 6N HCl to precipitate and isolate the optically enriched acid (-)3-(3-bromo-4-hydroxy-6-methyl-2-oxopyridin-1(2H)-yl)-4-methylbenzoic acid (**10**) by filtration. 1.3 kg (80% of theory) of product was obtained after drying.

Step 4



(-)-3-(3-bromo-4-hydroxy-6-methyl-2-oxopyridin-1(2H)-yl)-4-methylbenzoic acid (**10**) (1.50 kg, 4.44 moles), 1,1'-carbonyldiimidazole (1.08 kg, 6.66 moles) and N,N-dimethylformamide (3 mL) were mixed together at ambient temperature. Once the activation reaction was deemed complete, 2 M methylamine solution in tetrahydrofuran (5.5 L, 11.1 moles) was added at a rate that maintained the temperature below 30°C. The mixture became homogeneous as the reaction reached completion. A solution of 1 N hydrochloric acid solution (15.4 kg) was added to reach pH 3 or less. The acidification precipitated the product. The product was filtered, washed with water and vacuum dried at 60°C overnight to give (-)-3-(3-bromo-4-hydroxy-6-methyl-2-oxopyridin-1(2H)-yl)-N,4-dimethylbenzamide (**9**) (1.56 kg, 4.44 moles, 97% purity).

15 Step 5



3-(3-bromo-4-hydroxy-6-methyl-2-oxopyridin-1(2H)-yl)-N,4-dimethylbenzamide (**9**) (1 kg, 2.85 moles), potassium carbonate (0.43 kg, 3.11 moles), 1-methyl-2-pyrrolidinone (4.0 L) and 2,4-difluorobenzyl bromide (0.71 kg, 3.42 moles) were mixed together and heated at 30°C for 2 hours. The reaction mixture was diluted with water (12.5 L) over a period of 30 to 60 minutes and then stirred for 30 to 60 minutes. The product was filtered and washed with water. (-)-3-(4-(2,4-difluorobenzyloxy)-3-bromo-6-methyl-2-oxopyridin-1(2H)-yl)-N,4-dimethylbenzamide (**1**) (1.36 kg, 95% purity, greater than 99% ee) was


obtained.

The product was then purified. (-)-3-(4-(2,4-difluorobenzyloxy)-3-bromo-6-methyl-2-oxopyridin-1(2H)-yl)-N,4-dimethylbenzamide (**1**) (4.0 kg, 8.4 moles) and methanol (13 L) were mixed together and brought to 60°C and stirred for 1
5 hour. The product was filtered and washed with ambient temperature methanol (3.0 L) to give (-)-3-(4-(2,4-difluorobenzyloxy)-3-bromo-6-methyl-2-oxopyridin-1(2H)-yl)-N,4-dimethylbenzamide (**1**) (2.9 kg, 6.1 moles, 98.4% purity).

Claims


1. A process for the preparation of (-)-3-(4-(2,4-difluorobenzyloxy)-3-bromo-6-methyl-2-oxopyridin-1(2H)-yl)-N,4-dimethylbenzamide, or a pharmaceutically acceptable salt thereof comprising the steps:

a) contacting a compound of formula V:




V

with a brominating reagent in the presence of at least one solvent to produce a compound of Formula IV:



IV


b) contacting a compound of Formula IV with an hydrolase in the presence of a buffer solution to produce a compound of Formula X:



X

;

c) contacting a compound of Formula X with an activating reagent in the presence of at least one solvent and then contacting the resulting mixture with methylamine to produce a compound of Formula IX:



IX


; and

d) contacting a compound of Formula IX with a 2,4-difluorobenzylhalide in the presence of a base and at least one solvent;


wherein

R¹ is a C₁-C₆ alkyl or aryl.


2. A process of Claim 1 wherein the preparation of a compound of Formula V:



comprising contacting a compound of formula VI:



with a compound of formula VII:



in the presence of at least one solvent and a base, wherein R¹ is a C₁-C₆ alkyl or aryl.

3. The process of Claim 1 wherein (-)-3-(4-(2,4-difluorobenzyloxy)-3-bromo-6-methyl-2-oxopyridin-1(2H)-yl)-N,4-dimethylbenzamide is present in enantiomeric excess of 80%.

4. The process of Claim 1 wherein the halogenating reagent, in step a, is 1,3-dibromo-5,5-dimethylhydantoin.

5. The process of Claim 1 wherein the solvent, in step a, is acetonitrile, acetic acid, or an acetic acid containing a co-solvent such as water, lower alkyl alcohols, or dioxane.
6. The process of Claim 1 wherein the hydrolase, in step b, is a *Bacillus* sp. protease.
7. The process of Claim 1 wherein the buffer solution, in step b, is a dibasic potassium phosphate buffer solution or an inorganic bicarbonate buffer solution.
8. The process of Claim 1 wherein the activating reagent, in step c, is 1,1'-carbonyldiimidazole.
9. The process of Claim 1 wherein the solvent, in step c, is dimethylformamide, dichloroethane, tetrahydrofuran, dioxane, methyl tert-butyl ether, or toluene.
10. The process of Claim 1 wherein the solvent, in step d, is N-methylpyrrolidinone.
11. The process of Claim 1 wherein the base, in step d, is potassium carbonate.
12. A compound selected from:
methyl 3-(3-bromo-4-hydroxy-6-methyl-2-oxopyridin-1(2H)-yl-4-methylbenzoate;
(-)-3-(3-bromo-4-hydroxy-6-methyl-2-oxopyridin-1(2H)-yl-4-methylbenzoic acid; and
(-)-3-(3-bromo-4-hydroxy-6-methyl-2-oxopyridin-1(2H)-yl-N,4-dimethylbenzamide.