

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



20000257-6
pos copy

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07C 213/10, 217/18	A1	(11) International Publication Number: WO 99/05088 (43) International Publication Date: 4 February 1999 (04.02.99)
(21) International Application Number: PCT/GB98/02171 (22) International Filing Date: 21 July 1998 (21.07.98) (30) Priority Data: 9715479.3 23 July 1997 (23.07.97) GB (71) Applicants (for all designated States except US): UNIVERSITY OF BRADFORD [GB/GB]; Bradford, West Yorkshire BD7 1DP (GB). GENERIC BIOLOGICALS LIMITED [GB/GB]; 8 Centre One, Old Sarum Park, Lysander Way, Salisbury SP4 6BU (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): DOUBLE, John [GB/GB]; University of Bradford, Bradford, West Yorkshire BD7 1DP (GB). MAITLAND, Derek [GB/GB]; University of Bradford, Bradford, West Yorkshire BD7 1DP (GB). POPA, Ioana [RO/GB]; University of Bradford, Bradford, West Yorkshire BD7 1DP (GB). (74) Agents: BRIERLEY, Anthony, Paul et al.; Appleyard Lees, 15 Clare Road, Halifax HX1 2HY (GB).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: TAMOXIFEN AND ANALOGUES THEREOF (57) Abstract A method of preparing a first geometric isomer of tamoxifen or an analogue comprises allowing the first isomer to crystallise in hexanol and the product thereof to crystallise in methanol and then derivatising the product of the methanol crystallisation to prepare said first isomer in very high purity.		