

Spec. as filed
in POT9604759-2
1103 copyTHIS SHEET HAS BEEN
REPLACEDUREASE-BASED VACCINE AGAINST HELICOBACTER
INFECTION

5 The present invention relates to the prevention and treatment of gastric infection in mammals, including humans. More particularly, the present invention relates to a vaccine suitable for use in the prevention and treatment of *Helicobacter* infection in mammals, including humans, and to a method of treatment of humans suffering from gastric infection, its

10 consequences such as chronic gastritis or peptic ulcer, and prevention of gastric cancer.

BACKGROUND

15 *Helicobacter* infection of human gastric epithelium cause gastritis, are a major factor in the development of peptic ulcers and gastric lymphoma, and may be a risk factor for the development of gastric cancer [1-3]. The most frequent infection agent is *Helicobacter pylori*, followed at a much lower frequency by *Helicobacter heilmannii*. *H. pylori* is a slender S-

20 shaped gram negative microorganism, which is routinely recovered from gastric biopsies of adults and children with histologic evidence of gastritis or peptic ulceration. Evidence for a causal relationship between *H. pylori* and gastroduodenal disease comes from studies in human volunteers, patients with ulcers and gastric cancer, gnotobiotic pigs, and germ-free

25 rodents. Regarding etiology, Koch's postulates were satisfied by creating histologically confirmed gastritis in previously uninfected individuals following consumption of viable microorganisms [4-11], and by treatment to eradicate *H. pylori*, with resolution of the gastritis and, in patients with peptic ulcer disease, a decrease in the recurrence rate [12].

30 In spite of *in vitro* susceptibility to many antimicrobial agents, *in vitro* eradication of established *H. pylori* infections with antimicrobial agents is often difficult to achieve [13]. The microorganism is found within

This Sheet Has Been F



REPLACED SHEETS		SG
☒	Description : paged 1 to paged 33	
☒	Claim(s) : paged 33 to paged 33	
☐	Drawing(s) : paged to paged	
☒	Abstract : paged 37 to paged 37	
DATE OF RECEIPT		

the mucous coat overlying the gastric epithelium and in gastric pits. These are locations which do not appear to allow for adequate antimicrobial levels to be achieved even when antibiotics are given orally at high doses. At the present time, most authorities recommend a "triple therapy", namely a bismuth salt in combination with drugs such as tetracycline and metronidazole for 2-4 weeks. However, the effectiveness of this or other chemotherapeutic regimens remains suboptimal. Furthermore, this treatment may produce serious adverse drug reactions.

At the present time, little is known regarding the role of the mucosal immune system in the stomach. The distribution of immunoglobulin (Ig) producing cells in the normal gastric antrum indicates that IgA plasma cells make up 80% of the total plasma cell population. In addition, the number of plasma IgA cells present in the gastric antrum is comparable to other mucous membranes [14, 15]. A number of studies in humans [16] and in animal models [8, 10] have demonstrated specific IgG and IgA responses in serum and in gastric secretions in response to *Helicobacter* infection. However, the observation that *H. pylori* infection persists as a chronic infection for years, despite inducing a local and systematic immune response, is not encouraging the development of immunization strategies.

Lee et al reported the ability to infect germ-free rodents with *Helicobacter felis*, a bacterium closely related to *H. pylori*, and reproducible document histologic gastritis [9, 10]. Since then, this bacterium-host pairing has been accepted as a good model to study *Helicobacter*-mediated gastritis and its initiating factors [17]. Czinn et al have shown that repetitive oral immunization with a crude lysate of *H. pylori* plus cholera toxin adjuvant induces a vigorous gastrointestinal IgA anti-*H. pylori* response in mice and ferrets [13]. In addition, Chen et al and Czinn et al have recently reported that oral immunization with a crude lysate of *H. felis* induced protection against *H. felis* infection in mice [21, 22]. The exact nature of the antigen(s) responsible for the induction of this protection, however, had not been determined, and no information suggested that the protective antigen(s) of *H. felis* that induced protection against this pathogen would

This Sheet Has Been Replaced



induce a cross-reactive protection extending to another *Helicobacter* species.

We have demonstrated for the first time that *H. pylori* and *H. felis* sonicates and showing that some of these antibodies, directed against *H. pylori*, would crossreact with *H. felis* and vice versa [24, 25]. The basis for these cross-reactivities were unknown.

Based on the homology existing between the different known urease amino acid sequences, it has been proposed that urease could be used as a vaccine against *H. pylori* [26]. Nevertheless, cross-reactivity is not the rule. Guo and Liu have shown years ago that ureases of *Proteus mirabilis*, *Proteus vulgaris* and *Providencia rettgeri* show cross-reactivity to each other, while ureases of jack bean and *Morganella morganii* are immunologically distinct from the three former ureases [23]. Even if an antigenic cross-reactivity of *H. pylori* urease with other *Helicobacter* ureases was a reasonable postulate, no data existed demonstrating that this was really the case until we showed that some *H. felis* monoclonal antibodies crossreacted with *H. pylori* urease [25]. J. Pappo has further demonstrated that mice which have been infected by *H. felis* produce antibodies which crossreact with *H. pylori* urease but not jack bean urease (J. Pappo, unpublished data, 1993).

The use of *H. pylori* urease, or of related ureases, as a vaccine against *H. pylori* infection has previously been proposed by A. Labigne in EPO 367,644 [28]. However, that application contains no evidence of vaccination of any mammal against any *Helicobacter* infection with urease.

Moreover, while sequence homology with other bacterial ureases might support the use of urease as a vaccine candidate against *H. pylori* infection, the current knowledge of human *H. pylori* infection would certainly not. First, despite the fact that infected individuals often mount a strong antibody response to urease, the anti-urease immune response does not result in clearance or control of the infection. Secondly, *H. pylori* is able to transport urease out of the cell and to shed it from its surface [19, 20]. Thus, urease may not represent an appropriate target for the

This Sheet Has Been Replaced



development of a protective mucosal immune response. Indeed, mucosal immune protection is thought to be mainly mediated by secretory IgA, the agglutinating activity of which would be impaired when the recognized antigen can be shed by the target pathogen and thereby serve as a decoy for the protective antibody. Thirdly, urease appears to be toxic for epithelial cells in culture, and has been suspected to play a role in mucous degradation and in peptic ulceration *in vivo*. Thus, its use as antigen may be toxic.

Nevertheless, we reasoned that this antigen could be a potentially efficient vaccine if:

- first, we would deliver it orally at a sufficiently high dose to elicit a stronger immune response than the naturally occurring one
- second, the amount of antibodies produced would be high enough to bind all the urease, shed or not shed
- third, we would use subunits of urease or a molecular species that was non-toxic.

In summary, there remains a need for effective treatment and prevention of *H. pylori*-induced gastric infection in humans. Recent data suggested the possibility to generate a vaccine against this infection, but have not provided a clear identification of defined antigen(s), common to all strains of *H. pylori*, that could be incorporated into a safe and effective vaccine.

In this invention, we have identified the urease antigen of *H. pylori* as a candidate vaccine and demonstrated its efficiency in an animal model. These results were unexpected in the light of the natural history *Helicobacter* infections.

SUMMARY OF THE INVENTION

We have discovered that immunity can be induced in mammals susceptible to gastrointestinal *Helicobacter* infection by exploiting urease epitopes displayed on or about the surface of *Helicobacter* organisms and using them as a vaccine target. The immunity can be induced by

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

5

immunization with nature urease, but can also be induced with recombinant urease subunit, produced as an enzymatically inactive, therefore non-toxic form. The invention provides a method of inducing immunity to *Helicobacter* infection by administering to a mucosal surface of a mammal a polyaminoacid preparation, i.e. a mixture of peptides and/or proteins, together with an appropriate adjuvant. This polyaminoacid preparation presents a plurality of epitopes characteristic of and exhibited by a urease enzyme endogenous to the infecting *Helicobacter* organism. The administration of the polyaminoacid preparation may be performed by the oral route.

The active ingredient of the preparation may comprise natural or biosynthetic epitopes and may take various forms. A non exhaustive list of possible preparations includes purified, naturally occurring or recombinantly produced urease preparations of bacterial or other origin, digests of urease, fusion proteins comprising urease epitopes, truncated forms of urease enzyme, or peptides homologous with the aminoacid sequence of urease. Since development of immunity depends on induction of humoral and/or cellular immune responses which bind to the infecting *Helicobacter* organism, preferred preparations are those which most closely duplicate the epitopes of the urease endogenous to the infecting organism. For example, preparations displaying the epitopes of urease of *H. pylori* are preferred for administration in humans susceptible to *H. pylori*. However, in accordance with an important aspect of the invention, it has been discovered that urease from other species may be used. For example, we have shown that *H. felis* infection in mice can be prevented by administration of urease from *H. pylori*.

According to one aspect of the invention, there is provided a method of eliciting in a mammalian host a protective immune response to *Helicobacter* infection wherein an immunologically effective amount of a urease antigen capable of eliciting such a protective immune response, preferably *H. pylori* urease or *H. pylori* urease B subunit, is administered to a mucosal surface of the host.

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

6

According to another aspect of the present invention, there is provided a vaccine composition suitable for prevention of *Helicobacter* infection, comprising an effective amount of a urease antigen, preferably *H. pylori* urease or *H. pylori* urease B subunit, capable of eliciting in a host a protective immune response to *Helicobacter* infection, in association with a pharmaceutically acceptable carrier or diluent.

According to a further aspect of the present invention, there is provided a method of imparting to a mammalian host passive protection to *Helicobacter* infection, comprising administering to a mucosal surface of the host an immunologically effective amount of a urease specific antibody produced in a host immunized with a urease, preferably *H. pylori* urease or *H. pylori* urease B subunit, capable of eliciting a protective immune response to *Helicobacter* infection.

BRIEF DESCRIPTION OF THE DRAWINGS

The invention will now be further described with reference to the accompanying drawings, in which Figures 1 through 6 are graphical representations of the results set forth in Tables 1 through 6.

DETAILED DESCRIPTION OF THE INVENTION

The present inventors have discovered that oral administration to mice of polyaminoacid preparations exhibiting the epitopes of *H. pylori* urease gives rise to a protective immunological response against *H. felis* in mice, an animal model of generally-accepted value for the study of the immune response to *Helicobacter* infection [9]. The effect of the protective immune response is that immunized animals, when challenged with pathogen, have a greatly reduced incidence of infection, in comparison to non-immunized animals. Furthermore, the inventors have discovered that oral immunization in mice using *H. pylori* urease B subunit, produced as an enzymatically-inactive recombinant protein, gives rise to a protective immunological response in mice against *H. felis*. The effect of the

This Sheet Has Been Replaced



protective immune response is that immunized animals, when challenged with pathogen, have also a greatly reduced incidence of infection, in comparison to non-immunized animals which do become infected.

Thus, in a first aspect, the present invention provides a method of
5 eliciting in a mammalian host a protective immune response to *Helicobacter* infection. The method comprises the step of administering to a mucosal surface of the mammal, including humans, an immunologically effective amount of a urease antigen, preferably *H. pylori* urease, capable of eliciting such a protective immune response.

10 In a second aspect, the present invention provides a method of eliciting in a mammalian host a protective immune response to *Helicobacter* infection. The method comprises the step of administering to a mucosal surface of the animal, including humans, an immunologically effective amount of recombinant, enzymatically inactive urease B subunit an antigen,
15 preferably recombinant *H. pylori* urease B subunit, capable of eliciting such a protective immune response.

The invention also includes within its scope the treatment or prophylaxis of mammals, including humans, for *Helicobacter* infection, wherein an immunologically effective amount of a urease, or its subunits,
20 capable of eliciting a protective immune response to *Helicobacter* infection, is administered to a mucosal surface of a patient. Preferably, the urease is *H. pylori* urease or *H. pylori* urease B subunit, and the urease may be administered either alone or linked to a hydroxylated calcium phosphate, for example hydroxyapatite as a carrier particle. Moreover, it is preferred to
25 administer the *H. pylori* urease in association with a mucosal adjuvant, the B subunit of cholera toxin, muramyl dipeptide or other such adjuvants.

While not being bound by any theory, the present inventors believe that administration of the urease antigen, or B subunit thereof, to a mucosal surface stimulates the common mucosal immune system and perhaps local
30 sites in the gastric mucosal including an immune response, including the appearance of *H. pylori* specific IgA antibodies in the gastric secretions, which prevent *Helicobacter* infection. Since it is a routine matter to conduct

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

8

pre-clinical trials of a candidate vaccines for human use in animal models, it is believed that the methodology of the present invention is effective in humans, especially in the prevention and treatment of peptic ulcers, gastritis, gastric malignancies and other conditions arising as a result of the presence of *H. pylori* and/or *H. heilmannii*.

A - Bacterial cultures and urease purification

The strain of *H. pylori* used in the study originates from a patient with a duodenal ulcer, and has been subcultured on BHI agarose plates to homogeneity. *H. pylori* is cultured in a suitable medium, typically, BHI (Brain-Heart Infusion) medium, containing 0.25% yeast extract and 10% fetal calf serum and supplemented with 0.4% *Campylobacter* selective complement (Skirrow supplement; Oxoid 69). The bacteria are incubated overnight under microaerophilic conditions at 37°C in bottles that are then sealed and shaken at 37°C for 2 to 3 days to produce a liquid culture. A culture may also be prepared in agarose plates consisting of BHI with 0.25% of yeast extract and 5% of sheep blood under microaerophilic conditions at 37°C for 3 days. The quantity of bacteria is determined by optical density of the BHI solution at 660 nm, with one optical density unit corresponding to 10⁸ bacteria. Cultures on agarose plates are first resuspended in 154mm NaCl.

One currently preferred source of polyaminoacid displaying urease epitopes is purified urease, e.g., *H. pylori* urease obtained by following the general method of Dunn et al. J. Biol. Chem. 265, 9464-9469, modified as described below. Following culturing, the *H. pylori* is harvested in water, spun vortexed and spun again to produce a supernatant. The solution containing the urease activity of *H. pylori* (assessed by rapid urease test, see below) is then chromatographed on a CL-6B sizing column and the fractions which present a strong urease activity are pooled and dialyzed overnight and again chromatographed on an anion exchanger gel. The fractions are eluted in increasing NaCl buffer and the collected fractions with a strong urease activity are individually submitted to a SDS gel followed by Coomassie staining. Two distinct bands corresponding to a

This Sheet Has Been Replaced



molecular weight of about 63 and about 29 kDa are identified as urease.

The fractions containing urease are pooled to give purified *H. pylori* urease having a purity in the region of 95% to 99%.

B - Oral immunization with urease purified from *H. pylori*

5 While it is preferred to employ purified *H. pylori* urease obtained as described as the antigenic material, it will be understood that it is also possible to use, as the antigenic material, any urease or subunit of urease, either naturally occurring or obtained by recombinant DNA techniques, as well as digested fragment thereof, fusion proteins comprising the fragments or the
10 whole urease, truncated urease constructs, or other peptide or protein preparations exhibiting urease epitopes which are capable of eliciting a protective immune response to *Helicobacter* infection (see below). Thus, it is possible to employ a urease having a substantial homology with respect to *H. pylori* urease and which is effective in raising a cross-protective
15 immune response to *Helicobacter*. An example of such a urease is jack bean urease, which possesses about 70% homology with *H. pylori* urease. The invention is therefore not limited to the use of intact urease, and covers the use of any polypeptide preparation which displays urease epitopes and is effective to generate a protective immunological response in a host to
20 *Helicobacter* infection. Typically, a urease having a homology of 70-95% homology, for example, 80-90% homology, with respect to *H. pylori* urease, may be employed as the urease antigen in the invention.

A non-limiting list of sources of potentially useful urease preparations includes endogenous urease enzymes of the different
25 *Helicobacter* species, urease from other bacteria such as *Klebsiella pneumoniae* or *Proteus mirabilis*, and, by analogy, any other urease with the condition that these ureases share cross-reactive epitopes with *H. pylori* urease. The urease genes of all the organisms mentioned above represent a potential tool for expressing recombinant urease products as a whole protein
30 or as a part thereof.

A non-limiting list of potentially useful urease preparations includes peptides generated from purified urease (the sources are mentioned above).

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

10

using physical and/or chemical cleavage procedures (i.e. CnBr) and/or proteolytic cleavage (using proteases e.g. V8-protease, trypsin or others); or peptides synthesized chemically and sharing consecutive epitopes with urease.

- 5 Other sources of potentially useful epitopes include epitopes identified by their crossreactivity with urease, as the result of screening with anti-urease antibodies. These peptides can be naturally occurring peptides or peptides resulting from chemical synthesis. Furthermore such peptides can result from the expression of recombinant random
10 oligonucleotide.

Another source of potentially useful epitopes includes epitopes similar to urease as a result of the generation of anti-idiotypic antibodies to urease. Such anti-idiotypic antibodies, generated in any immunocompetent host, are obtained by immunization of this host with anti-urease antibodies,
15 with the goal of generating antibodies directed against anti-urease antibodies, which share structural homologies with urease.

The discussion herein focuses on the use of urease naturally produced by *H. pylori* (section B). However, it will be appreciated that the urease or subunits or constructs thereof mentioned above, capable of
20 eliciting the desired protective immune response, may be produced by recombinant DNA techniques well known in the art. The efficacy of particular preparations may be determined by routine administration using animal models, oral administration of the candidate vaccine, and challenge with pathogen using a protocol substantially similar or identical to the
25 procedure described below.

Table 1 and 2 below and Figures 1-5 describe the results obtained when mice were orally immunized with purified *H. pylori* urease. In this first experiment, administration of the *H. pylori* antigen was carried out by orally administering to the mice *H. pylori* urease purified as described in
30 section A, and coupled to hydroxyapatite crystals, used as a carrier to enhance M cell binding and uptake. Cholera toxin (Sigma) was given as a mucosal adjuvant. In this experiment, groups of female SPF BALB/c six-

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

11

week old mice were each orally immunized with 30 ug of purified *H. pylori* urease coupled to 1 mg of hydroxyapatite plus 10 ug of cholera toxin adjuvant at day 0, 7, 14 and 21. The mice were then challenged twice with 10^5 *H. felis*, at day 28 and 30. For comparison purposes, similar female SPF

- 5 BALB/c six-week old mice were orally immunized with whole *H. pylori* lysate (sonicate) and 10 ug cholera toxin at day 0, 7, 14 and 21. The mice were challenged at day 28 and 30 with *H. felis*. The *H. pylori* sonicate was prepared by collecting *H. pylori* from cell cultures, pelleting by centrifugation and resuspending the pellet in 0.9% sodium chloride followed
10 by sonication.

As a control, female SPF BALB/c six-week old mice were orally sham-immunized with 10 ug of cholera toxin and 1 mg of hydroxyapatite at day 0, 7, 14 and 21. All mice were housed, immunized, and challenged in parallel. All mice subject to the study were sacrificed on day 35.

- 15 C- Oral immunization with recombinant urease subunits of *H. pylori*

Genes encoding the structural A and B subunits of *H. pylori* urease were obtained by polymerase chain reaction (PCR) cloning according to standard procedures, based on previously published sequences [29]. These genes were inserted in a vector (named pEV40) designed for high
20 expression and easy purification of foreign genes in *E. coli*. Briefly, the foreign gene is inserted downstream of a thermo-repressible promoter, and in frame of a sequence encoding for a repeat of six histidines. An ampR gene is present on this vector for selection of transformants. Under the appropriate temperature conditions, the recombinant protein obtained is
25 supplemented by six histidines at the N-terminal, which allow for a one-step affinity purification on a nickel column. Both *H. pylori* recombinant urease A and B subunits were expressed separately in *E. coli*, and purified on nickel column to .95% purity.

While it is preferred to employ recombinant *H. pylori* urease
30 obtained as described above as the antigenic material, it will be understood that it is also possible to use, as the antigenic material, any urease or subunit of urease obtained by recombinant techniques (e.g. fusion protein)

This Sheet Has Been Replaced



12

expressing antigenic sites of urease, which is capable of eliciting a protective immune response to *Helicobacter* infection. Thus, it is possible to employ in a construct a urease gene having a substantial homology with respect to *H. pylori* urease and which is effective in raising a cross-
5 protective immune response to *Helicobacter*. Examples of such a urease is jack bean urease, which possesses about 70% homology with *H. pylori* urease, or *H. felis* urease, which possesses about 88% homology with *H. pylori* urease. The invention is therefore not limited to the use of *H. pylori* urease genes and their gene products, and covers the use of any
10 recombinant urease, or the subunits thereof, which is effective to generate a protective immunological response in a host to *Helicobacter* infection. Typically, a recombinant urease having a homology of 70-95% homology, for example, 80-90% homology with respect to *H. pylori* urease, may be employed as the recombinant urease antigen in the invention.

15 The discussion herein focuses on the use of recombinant *H. pylori* urease A and B subunits produced by *E. coli* (section C). However, it will be appreciated that recombinant urease or subunits or constructs thereof mentioned above, capable of eliciting the desired protective immune response, may be produced using other recombinant DNA techniques and
20 other eukaryotic or prokaryotic expression vectors well known in the art.

Table 3, 4 and 5 below and Figure 6 describe the results obtained when mice were orally immunized with recombinant *H. pylori* urease subunits produced in *E. coli*. In this experiment, administration of the *H. pylori* antigen was carried out by orally administering to the mice
25 recombinant *H. pylori* urease A or B subunits produced in *E. coli* and purified as described above, and coupled to hydroxyapatite crystals, used as a carrier to enhance M cell binding and uptake. Cholera toxin (Sigma) was given as a mucosal adjuvant. In this experiment, groups of female SPF BALB/c six-week old mice were each orally immunized with 30 ug of
30 recombinant *H. pylori* urease A and B subunit, coupled to 1 mg of hydroxyapatite plus 10 ug of cholera toxin adjuvant at day 0, 8, 14 and 21. The mice were then challenged twice with 10^8 *H. felis*, at day 32, 34 and

This Sheet Has Been Replaced



13

36. For comparison purposes, similar female SPF BALB/c six-week old mice were orally immunized with 30 ug of recombinant *H. pylori* urease B subunit coupled to hydroxyapatite plus 10 ug cholera toxin at day 0, 8, 14 and 21. The mice were challenged three times, at day 32, 34 and 36, with *H. felis*. As a control, female SPF BALB/c six-week old mice were each orally sham-immunized with 10 ug of cholera toxin and 1 mg of hydroxyapatite at day 0, 8, 14 and 21. The mice were then challenged at day 32, 34 and 36 with *H. felis*. All mice subject to the study were immunized and challenged in parallel. Animals were sacrificed on day 48 (12 days after challenge) or 10 weeks after challenge.

D- Analysis of gastric biopsies, blood, and intestinal secretions

Biopsies were taken from the stomach and blood was obtained from the heart. The intestine were removed and washed with 1mM PMSF (Boeringer) in PBS buffer to obtain intestinal secretions for ELISA analysis.

To evaluate protection against *H. felis* colonization, gastric biopsies from each animal were screened for the presence of *H. felis* by assessing rapid urease activity by the Jatrox HP test (Rohm Pharma), according to the supplier's directions. Briefly, gastric biopsies are immersed in 0.5 ml supplier's mixture of urea and phenol red, a pH indicator. Urease activity generates ammonia and bicarbonate from urea, and is followed by the colorimetric change of the solution towards a higher absorbance at 550 nm. Urease activity was quantified by spectrophotometric analysis.

Gastric biopsies of each animal included in the experiment described in section B were also cultured on BHI agarose plates, supplemented as above, for the detection of *H. felis*. After incubation for 3 to 10 days in microaerophilic conditions, the presence of *H. felis* was confirmed by Gram staining and determination of urease activity. As a very significant correlation was obtained for the detection of *H. felis* cultures during the first set of experiments (see Table 3), only gastric biopsies urease tests were performed for the detection of *H. felis* in the experiment described in the experiment described in section C. Detection of *H. felis* was confirmed by

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

14

microscopy by two independent investigators, using two different stains (acridine orange and cresyl violet).

Blood samples were allowed to clot for 3 hours at RT, and sera harvested and frozen at -20°C until analyzed. Intestinal secretions were spun for 5 min. at 4°C to remove debris, and kept frozen at -20°C. Serum and intestinal samples of each animal were analyzed by ELISA for evaluation of anti-*Helicobacter* activity, according to standard procedures. Briefly, 96-well plates were coated with a sonicate of *H. pylori*, followed by saturation with 5% fat-free milk. Samples were serially diluted from 1:1 to 1:1000 and incubated overnight at 4°C on ELISA plates. Biotinylated anti-mouse IgG (serum) and anti-mouse IgA, followed by streptavidin-Horseradish peroxidase was used for the determination of the antibody levels.

The results of *H. felis* challenges following immunizations with purified *H. pylori* urease are set out in Tables 1-3 and Figures 1-4 and the results of *H. felis* challenges following immunizations with recombinant *H. pylori* urease A and B subunits are set out in Tables 4-6 and Figures 5 and 6.

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

15

TABLE 1

mouse		urease	culture	immunoglobulins			
number	immunization	test	Gram	Serum		intestinal secretion	
		12h		Ig	IgG	Ig	IgA
1	Urease+HF	+	<i>H. felis</i>	27	0	25	258
2	Urease+HF	0	0	264	279	221	246
3	Urease+HF	0	0	84	44	318	354
4	Urease+HF	+	<i>H. felis</i>	81	42	12	5
5	Urease+HF	0	0	98	137	126	234
6	Urease+HF	+	0	988	2093	31	22
7	Urease+HF	0	0	98	0	96	34
8	Urease+HF	0	0	247	1010	214	128
9	Urease+HF	0	0	N.D.	N.D.	48	23
10	Urease+HF	0	0	50	0	124	99
11	Urease	0	0	318	205	44	53
12	Urease	0	0	14	0	86	87

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

16

13	Urease	0	0	0	0	0	0
14	Urease	0	0	0	0	43	61
15	Urease	0	0	58	0	110	127
16	Urease	0	0	140	63	21	97
17	Urease	0	0	84	240	114	260
18	Urease	0	0	N.D.	N.D.	93	148
19	Urease	0	0	45	0	135	216
20	Urease	0	0	261	197	161	261
21	CT+HF	0	0	0	0	0	2
22	CT+HF	+	<i>H felis</i>	83	0	310	303
23	CT+HF	+	<i>H felis</i>	90	0	N.D.	N.D.
24	CT+HF	+	<i>H felis</i>	31	0	150	192
25	CT+HF	+	<i>H felis</i>	197	250	250	440
26	CT+HF	+	<i>H felis</i>	105	135	214	138
27	CT+HF	+	<i>H felis</i>	140	47	109	55
28	CT+HF	+	0	0	0	16	15
29	CT+HF	+	<i>H felis</i>	0	0	0	0
30	CT+HF	+	<i>H felis</i>	N.D.	N.D.	N.D.	N.D.
31	HP sonicate+HF	+	<i>H felis</i>	0	0	76	103
32	HP sonicate+HF	+	<i>H felis</i>	77	0	11	33
33	HP sonicate+HF	+	<i>H felis</i>	549	748	57	36
34	HP sonicate+HF	0	0	660	153	180	286
35	HP sonicate+HF	+	<i>H felis</i>	730	192	0	5
36	HP sonicate+HF	+	<i>H felis</i>	32	0	5	64
37	HP sonicate+HF	0	0	400	400	312	1149
38	HP sonicate+HF	+	<i>H felis</i>	1007	1350	149	26
39	HP sonicate+HF	0	0	220	188	133	122
40	HP sonicate	0	0	873	1016	352	514
41	HP sonicate	0	0	727	899	126	191
42	HP sonicate	0	0	109	68	44	83
43	HP sonicate	0	0	147	949	167	97
44	HP sonicate	0	0	845	1094	246	64
45	HP sonicate	0	0	1217	1198	210	157
46	HP sonicate	0	0	81	0	256	216
47	HP sonicate	0	0	329	210	241	276
48	HP sonicate	0	0	1049	737	197	211

This Sheet Has Been Replaced



In Table 1, which refers to the experiment described in section B, "h" means hours, "Ig" means immunoglobulin, "ND" means "not determined", "urease + HF" means that the mice were immunized with urease (coupled to hydroxyapatite, with cholera toxin) and then challenged with *H. felis*, "urease" means that the mice were immunized with urease (coupled to hydroxyapatite, with cholera toxin) and not challenged, "CT+HF" means that the mice were sham-immunized with cholera toxin and challenged with *H. felis*, "HP sonicate + HF" means that the mice were immunized with *H. pylori* sonicate with cholera toxin and challenged by *H. felis*, and "HP sonicate" means that the mice were immunized with *H. pylori* sonicate with cholera toxin and not challenged. In Table 1, the numbers for the antibody results are given as a measure of absorbance at 595 nm multiplied by 1000. The background measured in absence of the antibodies, was subtracted.

The results of experiment described in section B obtained on the basis of the gastric biopsies urease tests and on Gram staining of *H. felis* cultures are set out in Table 2. Infection was defined by mice with one or more markers of colonization by *H. felis*, including urease test or Gram staining of cultures.

TABLE 2

Immunization	Challenge	% infected	% protected
Urease	<i>H. felis</i>	3/10 (30%)	7/10 (70%) *
Sonicate	<i>H. felis</i>	6/9 (66%)	3/9 (33%) **
CT	<i>H. felis</i>	9/10 (90%)	1/10 (10%)

* p=0.0198 (two tailed Fisher exact test) compared to CT control

** p=0.303 (two tailed Fisher exact test) compared to CT control

It will be seen from the results set out in Tables 1 and 2 that statistically significant protection against *H. felis* challenge is obtained with oral immunization using *H. pylori* urease as compared to that obtained



using either *H. pylori* sonicate or cholera toxin. Referring to Table 2, it will be seen that from a total of 10 immunized animals, only 3 became infected, as compared to 6 of the animals immunized with *H. pylori* sonicate and 9 of the animals immunized with cholera toxin. Table 2 shows that 70% of the animals were protected from challenge by *H. felis* as compared to 33% of the animals immunized with *H. pylori* sonicate and 10 of the animals immunized with cholera toxin and then subjected to *H. felis* challenge. In other words, 90% of the control mice exposed to *H. felis* became infected by that pathogen whereas, in contrast, in mice immunized with *H. pylori* urease 28 days before exposure to *H. felis*, the infection rate was only 30%. This represents a significant reduction in infection ($p=0.0198$ in the Fisher exact test, as compared to the control mice). When the mice were orally immunized with *H. pylori* sonicate, the infection rate was 67% (not significant versus the control). The protection obtained using *H. pylori* urease is unexpected and could not have been predicted on the basis of the results observed using *H. pylori* sonicate.

Referring to Figures 1-4, Figure 1 represents graphically the results of tests for antibodies in serum (IgG) and intestinal secretion (IgA) in mice not protected after immunization with urease. These are mice numbers 1, 4 and 6 appearing in Table 1, and constitute Group A. Figure 2 shows the antibody response of mice that were protected after immunization with urease (Group B), i.e. mice 2, 3, 5 and 7-10.

Figures 3 and 4 relate to the results obtained with mice numbers 31-39. Figure 3 (Group C) depicts antibody responses of mice not protected after immunization with *H. pylori* sonicate (mice numbers 31, 32, 33, 35, 36 and 38) and Figure 4 (Group D) depicts the antibody responses of mice protected after immunization with *H. pylori* sonicate (mice numbers 34, 37 and 39). It is of interest to note with respect to Figures 3 and 4 that the IgA antibody responses (but not IgG) are higher in the mice exhibiting protection than in the mice that are not protected, suggesting a correlation between protection and IgA response. Serum IgG responses did not exhibit

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

19

a correlation. Mucosal IgA but not serum IgG antibodies are known to play a role in protection against bacterial infections of the gut [3].

The results of the correlation between the detection of *H. felis* in gastric biopsies by urease tests and by cultures are set out in Table 3.

TABLE 3

	Urease Test +	Urease Test-	Total
<i>H. felis</i> culture +	16	0	16
<i>H. felis</i> culture -	2	30	32
Total	18	30	48

Two-tailed Fisher's Exact Test: $p < 0.0001$

Table 3 shows that a very significant correlation exists between the results of urease tests performed on gastric biopsies and the identification of *H. felis* infection than urease tests, urease tests were preferred for the diagnosis of *H. felis* infection in mice in the next experiments, due to its better sensitivity. This approach allowed duplication of urease tests with larger fragments of the stomach of each mouse, and a further increase in the sensitivity of the urease test. Furthermore, the use of the method with the highest sensitivity prevent an overestimation the protection obtained by the vaccine preparation to be tested. When positive culture is used as the standard for infection, the protection induced after urease immunization during the experiment depicted in section B is as significant as with the combined use of urease test and culture ($p=0.021$ versus $p=0.019$).

The results of the experiments described in section C (recombinant urease subunits), obtained on the basis of the gastric biopsies urease tests, are set out in Table 4, 5 and 6 and depicted in Figure 6.

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

20

TABLE 4

Immunization	mice n°	Urease test	Infection
CT	20	0.49	+
	21	0.31	+
	22	0.62	+
	23	0.67	+
	24	0.55	+
	50	0.50	+
	51	0.37	+
	52	0.29	+
	53	0.79	+
	54	0.32	+
ureA + HAP + CT	40	0.67	+
	41	0.48	+
	42	0.42	+
	43	0.65	+
	44	0.56	+
	45	0.52	+
	46	0.33	+
	47	0.63	+
	48	0.22	+
	49	0.37	+
ureB + HAP + CT	25	0.15	-
	26	0.07	-
	27	0.03	-
	28	0.64	+
	29	0.13	-
	30	0.02	-
	31	0.66	+
	32	0.00	-
ureA + HAP + CT	68	0.00	-
	69	0.07	-
	70	0.42	+
	71	0.00	-
ureB + HAP + CT	72	0.00	-
	73	0.37	+
	74	0.00	-
	75	0.37	+
	76	0.00	-
	77	0.00	-
	78	0.00	-
	79	0.39	+
	80	0.00	-
	81	0.37	+
	82	0.00	-

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

This Sheet Has Been Replaced



21

In Table 4, "CT" means cholera toxin; "UreA" means recombinant *H. pylori* urease A subunit; "UreB" mean recombinant *H. pylori* urease B subunit; and "HAP" means hydroxyapatite crystals. Mice 20 to 54 were sacrificed 12 days post challenge and mice 68 to 82 10 weeks (106 days) post challenge. The results of the urease test performed from biopsies of the stomach of each animal are expressed as OD values at 550 nm. The positive and negative signs depicts the final status of infection of each animal, according to the positivity or negativity of the urease test for detection of *H. felis*. Positivity: OD550 values >0.2

10

TABLE 5: Protection as measured 12 days post challenge

Immunization	Challenge	% infected	% protected
Urease A subunit	<i>H. felis</i>	10/10 (%)	0/10 (0%)
Urease B subunit	<i>H. felis</i>	3/10 (30%)	7/10 (70%) *
CT	<i>H. felis</i>	10/10 (100%)	0/10 (0%)

* p=0.0031 (two tailed Fisher exact test) compared to CT control

TABLE 6: Protection as measured 10 weeks post challenge

Immunization	Challenge	% infected	% protected
Urease A subunit	<i>H. felis</i>	1/5 (20%)	4/5 (80%) *
Urease B subunit	<i>H. felis</i>	4/10 (40%)	6/10 (60%) **

* p=0.003 (two tailed Fisher exact test) compared to CT control

** p=0.01 (two tailed Fisher exact test) compared to CT control

25

It will be seen from the results set out in Tables 4, 5 and 6 that statistically significant protection against *H. felis* challenge is obtained with oral immunization using recombinant *H. pylori* urease B subunit as compared to that obtained using either recombinant *H. pylori* urease A subunit or cholera toxin. Referring to Table 4, it will be seen that, 12 days post challenge, from a total of 10 immunized animals, only 3 were found infected in the urease B subunit group, as compared to all 10 animals

immunized with *H. pylori* A subunit of urease and 10 out of 10 of the animals immunized with cholera toxin. Table 4 shows that 70% of the animals were protected from challenge by *H. felis* as compared to 0% of the animals immunized with *H. pylori* urease A subunit and 0% of the
5 animals immunized with cholera toxin and then subjected to *H. felis* challenge. In other words, 100% of the control mice challenged with *H. felis* became infected whereas, in contrast, in mice immunized with recombinant *H. pylori* urease B subunit the infection rate was only 30%. This represents a significant reduction in infection ($p=0.0031$, Fisher exact
10 test) as compared to the control mice.

The fact that the protection observed with *H. pylori* urease is entirely conferred by immunization with the B subunit of urease, and that the A subunit has no such effect, was not expected on the basis of our experiment with purified urease. This definition of the roles of the 2
15 structural subunits of urease in the development of a protective immune response is therefore novel. The protection obtained using recombinant urease B subunit, which is enzymatically inactive also teaches that non-toxic forms of urease can be used as oral vaccine against *Helicobacter* infection. Furthermore these results strongly suggest that recognition of the
20 active site is not required for protection, as an inactive urease B subunit is very unlikely to induce antibodies that will recognize and inhibit the catalytic site of native urease.

Referring to Table 6, it will be seen that, when mice are sacrificed 10 weeks post infection, 60% (6 mice out of 10) of the animals immunized
25 with urease B subunit and 80% (4 mice out of 5) of the animals immunized with *H. pylori* urease B subunit were protected against *H. felis* infection. The fact that protection obtained through immunization with urease B subunit lasts over time and that immunization with urease A induces a protection which is displaced compared to the one induced by urease B
30 subunit could not be expected from our experiment with purified urease or with other experiment performed earlier. The fact that urease B subunit immunization confers protection definitely proves that recognition of the

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

This Sheet Has Been Replaced



23

active site is not required for protection. Figure 6 summarizes results obtained after oral immunization with recombinant urease A and B subunits (described in Tables 5 and 6).

The present invention also provides vaccine compositions suitable for the prevention of *Helicobacter* infection. The compositions comprise an effective amount of a urease antigen, preferably *H. pylori* urease or recombinant *H. pylori* urease subunits, capable of eliciting in a host a protective immune response to *Helicobacter* infection, in association with a pharmaceutically acceptable carrier or diluent.

The vaccines of the invention are administered in amounts readily determined by persons of ordinary skill in this art. Thus, for adults a suitable dosage will be in the range of 10 ug to 100 milligrams, for example 50 ug to 50 mg. Similar dosage ranges will be applicable for children. Carrier systems in humans may include enteric release capsules protecting the antigen from the acidic environment of the stomach, and including urease antigen in an insoluble form as fusion proteins. The vaccine can be administered as a primary prophylactic agent in adults or in children, as a secondary prevention, after successful eradication of *H. pylori* in an infected host, or as a therapeutic agent in the aim to induce an immune response in the host susceptible to contribute to the eradication of *H. pylori*.

As noted above, a suitable mucosal adjuvant is cholera toxin. Others which may be used as muramyl dipeptide or its derivatives, non-toxic derivatives of cholera toxin, including its B subunit, and/or conjugates or genetically engineered fusions of the urease antigen plus cholera toxin or its B subunit. Other suitable delivery methods include biodegradable microcapsules or immune stimulating complexes (ISCOM'S) or liposomes, genetically engineered attenuated live vectors such as viruses or bacteria, and recombinant (chimeric) virus-like particles, e.g. bluetongue. The amount of mucosal adjuvant employed depends on the type of mucosal adjuvant used. For example, when the mucosal adjuvant is cholera toxin, it is suitably used in an amount of 5 ug to 50 ug, for example 10 ug to 35 ug. When used in the form of microcapsules, the amount used will depend on

THIS SHEET HAS BEEN
REPLACED

This Sheet Has Been Replaced



24

the amount employed in the matrix of the microcapsules to achieve the desired dosage. The determination of this amount is within the skill of a person of ordinary skill in this art. Suitable carriers for the vaccines of the invention are enteric coated capsules and polyactide-glycolide microspheres.

- 5 Suitable diluents are 0.2N NaHCO_3 and/or saline.

Particulate hydroxylated calcium phosphate (HCP) is especially useful as a carrier for the *H. pylori* urease to be applied to mucosal surfaces. It is believed that the *H. pylori* urease-hydroxylated calcium phosphate conjugate is transported across epithelium where it raises a poly
10 Ig immune response. Preferably, the hydroxylated calcium phosphate is in the form of microparticles suitable for the transport across the epithelium, particularly by cells specialized for this purpose (M cells). A preferred form of hydroxylated calcium phosphate is hydroxyapatite, a commercially available crystalline hydroxylated calcium phosphate $\text{Ca}_{10}(\text{PO}_4)_6(\text{OH})_2$.

- 15 Commercially available hydroxyapatite generally consists of slab-like crystals that are chemically and physically analogous to inorganic hydroxyapatite in normal bone tissue. Ingestion of hydroxyapatite should therefore be safe, as evidenced by the existence of nutritional calcium/phosphorous supplements derived from ground bone, which are
20 designed to be ingested. Commercially-high resolution hydroxyapatite (from CalBioChecm) consists of crystals varying widely in size. Crystals over 1 μm in length are unlikely to be taken up by M cells. Therefore, for use in the invention, commercial hydroxyapatite crystals are broken into small, relatively uniform crystalline fragments such as by sonication. Preferably, a
25 substantial proportion of the hydroxyapatite is present as fragments of about 0.01-0.0 μm . Fragmentation may be measured either by electron microscopy or light scattering, using standard techniques.

- Preferred modes of administration of the *H. pylori* urease antigen are orally, nasally, rectally or ocularly. Oral administration can provide delivery
30 to other G.I. mucosa including the intestinal mucosa.

The vaccines of the present invention may be administered to a mucosal surface in the form of an aerosol, suspension, capsule and/or

THIS SHEET HAS BEEN
REPLACED

25

suppository. The method of administration will be readily apparent to a person of ordinary skill in this art.

The present invention further includes the passive immunization of mammals, including humans, against *Helicobacter* infection. This is
5 achieved by administering to a mucosal surface of the patient an effective amount of a urease specific antibody. Preferably, an effective amount of a *H. pylori* urease specific IgA monoclonal antibody.

Since the urease of *H. pylori* is shown to represent the antigen involved in inducing protective immunity, a further aspect of the invention
10 is the use of *H. pylori* urease as a diagnostic reagent to measure the immune response of persons who have received a vaccine based on urease or to determine whether an individual is immune or susceptible (and thus in need of vaccination). The present invention also includes the use of urease and urease-specific antibodies, to construct assays and kits for diagnosis of
15 *Helicobacter* immunity, assessment of *Helicobacter* susceptibility, and definition of immune responses to vaccines.

EXAMPLES

The invention will now be further described by reference to the
20 following non-limiting examples.

a) The Bacterial Strains

H. felis was provided by J. Fo (division of Cooperative Medicine, Mass. Institute of Technology, Boston, USA). *H. pylori* was isolated from patients with ulcer disease (CHUV, Lausanne, Switzerland).

25 b) Bacterial Cultures

Liquid Culture - Bacteria were cultured on BHI (Brain-Heart Infusion, BioMerieux) liquid medium containing 0.25% of yeast extract (Difco) and 10% of fetal calf serum (Intech) supplemented with 0.4% of Campylobacter selected complement (Oxoid). The bacteria were incubated
30 overnight under microphilic conditions at 37°C and then shaken at 37°C for 2 to 3 days.

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

This Sheet Has Been Replaced



26

Culture on Agarose plates - The bacteria were cultured on agar plate consisting of BHI 0.25% of yeast extract and 5% of sheep blood under microphilic conditions at 37°C for 3 days.

Quantification - The quantity of bacteria was determined by the optical density of the BHI solution at 660 nm (1 optical density unit corresponding to 10⁸ bacterial).

c) Preparation of sonicates

H. pylori was collected from 31 blood agar plates in 0.15 M NaCl and spun 5 minutes at 1400g at 4°C. The pellet was resuspended in 3 ml of NaCl and sonicated for 4 minutes. The amount of proteins was evaluated by a Bradford assay (BioRad Kit according to supplier).

d) Coupling of immunogen to Hydroxyapatite

Immunogen (urease or subunit thereof) was incubated for 1 hour at 4°C with hydroxyapatite. 1.0 mg of hydroxyapatite was used for 30 ug of immunogen per mouse. At the end of the incubation, 10 ug of cholera toxin was added in a final volume of 200ul PBS.

EXAMPLE 1

a) Extraction

H. pylori from 30 blood agar plates was harvested in 0.15 M NaCl on ice. The solution was spun 5 minutes at 1400 g at 4°C. The pellet was resuspended in 20 ml of H₂O and vortexed for 45 seconds (maximum speed). The extract was then spun 20 minutes at 6700 g at 4°C. The supernatant was recovered and the quantity of protein was evaluated (see "Quantification" above) and precipitated with 70% of ammonium sulfate.

b) Purification of urease

The solution was chromatographed on a Sepharose CL-6B column (Pharmacia) with PBS (phosphate buffered saline) as mobile phase. The collected fractions which presented a strong urease activity were pooled and dialyzed overnight at 4°C against 3 liters of PEB (20 mM phosphate buffer, pH 7) and then chromatographed on a Q Sepharose fast flow (Pharmacia) with PEB as mobile phase. The fractions were eluted by 0 to 500 mM NaCl

27

gradient. Ten of the collected fractions with a strong urease activity were individually subjected to an SDA gel followed by a Coomassie staining. The 6 fractions presenting 2 distinct bands corresponding to MW-63 and -28 KDa were pooled and were considered as the purified urease.

5

EXAMPLE 2 (see also section B)

Mice employed in the immunization studies were lightly anesthetized with ether prior to intragastric immunization. And then, sonicate preparation or purified urease, hydroxyapatite and cholera toxin
10 was suspended in PBS and 200ul were delivered to the stomach of the respective mice using a polyethylene tubing attached to a hypodermic syringe. This procedure will be referred to as oral immunization.

Three oral immunization protocols were evaluated. These are described below.

15 Protocol B1 - Vaccination with purified urease

Female BALB/c six-week old mice (20) were orally immunized with 30 ug of purified of *H. pylori* urease and 1 mg of hydroxyapatite and 10 ug of cholera toxin at day 0, 7, 14 and 21. Ten mice were challenged at day 28 and 30 with 5×10^7 and 10^8 *H. felis* from liquid culture.

20 Protocol B2 - Vaccination with Helicobacter sonicates

Female BALB/c six-week old mice (20) were orally immunized with 2 mg of *H. pylori* sonicate solution at day 0, 7, 14 and 21. Ten mice were challenged at day 28 and 30 with 5×10^7 and 10^8 *H. felis*.

Protocol B3 - Control

25 Female BALB/c six-week old mice (20) were orally immunized with 1 hydrozapatite and 10ug of cholera toxin at day 0, 7, 14 and 21. The mice were challenged at day 28 and 30 with 5×10^7 and 10^8 *H. felis*.

At day 35 all mice were sacrificed and biopsies from the stomach were taken as well as intestinal secretions and blood.

30

Protection and evaluation

This Sheet Has Been Replaced



This Sheet Has Been Replaced

THIS SHEET HAS BEEN
REPLACED

28

To evaluate protection, biopsies were screened for the urease activity by the Jetrox HP test (Rohm Pharma) according to the instructions of the supplier. The urease is quantified by a spectrophotometric measurement at 550 nm. The biopsies were also cultured in the presence of *H. felis* and was estimated by Gram staining. Gastric antral biopsies were homogenized and diluted (1:10 and 1:1000) in 0.15 M NaCl and plated onto blood agar plates and incubated under microaerophilic conditions at 37°C for 4 to 10 days.

10 ELISA

Intestinal secretions and blood were analyzed by ELISA for the evaluation of antibody titer. The ELISA was carried out as follows. Polystyrene plates (96 wells) were coated with $\mu\text{g}/\text{well}$ of purified urease at 37°C for 2 hrs. Non specific binding sites were blocked with 5% powdered milk in PBS 0.1% Tween at 37°C for 30 minutes. The plates were washed once with PBS 0.1% Tween. Blood samples were test at dilution 1:100 and intestinal secretions 1:1. 100 μl of each sample were added to the antigen coated plates. After 2 hrs of incubation, plates were washed 3 times with PBS 0.1% Tween. Anti-mouse biotinylated whole antibody from goat and anti-mouse IgA, IgG and IgM biotinylated (Amersham) were added (100 μl) at dilution 1:500 except for IgA (1:250) and incubated at 37°C for 1 hr. The plates were washed 3 times with PBS 0.1% Tween and 100 μl of 1:1000 dilution of streptavidin Horseradish peroxidase in PBS 0.1% Tween were added and incubated at 37°C for 30 minutes. The plates were washed 3 times and 50 μl of 1:50 dilution of o-phenyl-diamine in citrate buffer pH 5.0 with 1 $\mu\text{l}/\text{ml}$ of 30% H_2O_2 were added and incubated at room temperature for 20 minutes. The absorbance at 495 nm was measured in each well.

EXAMPLE 3 (see also section C)

Mice employed in the immunization studies were slightly anesthetized with ether prior to intragastric immunization. Then, 30 μg recombinant *H. pylori* urease A and B subunit produced in *E. coli* bound hydroxyapatite and supplemented with cholera toxin was suspended in PBS

29

and 200ul were delivered to the stomach of the respective mice using a polyethylene tubing attached to a hypodermic syringe. This procedure will be referred to as oral immunization.

Three oral immunization protocols were evaluated. These are described below.

Protocol C1 - Vaccination with recombinant urease A subunit

Female BALB/c six-week old mice (10) were orally immunized with 30 ug of purified recombinant *H. pylori* urease A subunit and 1 mg of hydroxyapatite and 10 ug of cholera toxin at day 0, 8, 14 and 21. Ten mice were challenged at day 32, 34 and 36 with 10^8 *H. felis* from liquid culture.

Protocol C2 - Vaccination with recombinant urease B subunit

Female BALB/c six-week old mice (10) were orally immunized with 30 ug of recombinant *H. pylori* urease B subunit and 1 mg of hydroxyapatite and 10 ug of cholera toxin at day 0, 8, 14 and 21. Ten mice were challenged at day 32, 34 and 36 with 10^8 *H. felis* from liquid culture.

Protocol C3 - Control

Female BALB/c six-week old mice (10) were orally immunized with 1 mg hydroxyapatite and 10 ug of cholera toxin at day 0, 8, 14 and 21.

The mice were challenged at day 32, 34 and 36 with 10^8 *H. felis*.

At day 42, or at day 106, mice were sacrificed and multiple biopsies from the stomach were taken.

25 Protection and Evaluation

To evaluate protection, biopsies of the corpus and antrum of the stomach were screened for urease activity by the Jatrox IIP test (Rohm Pharma) according to the instructions of the supplier. The urease is quantified by a spectrophotometric measurement at 550nm. The total of corpus and antrum OD values were added to obtain a final OD value for each mouse.

This Sheet Has Been Replaced



References

1. Blaser, M.J. "Gastric Campylobacter-like organisms, gastritis and peptic ulcer disease" *Gastroenterology* 1987, 93, 371-383.
- 5 2. Graham, D.Y. "Campylobacter *pylori* and peptic ulcer disease" *Gastroenterology* 1989, 96, 615-625.
3. Parsonnet, J. et al "Helicobacter *pylori* infection in intestinal and diffuse-type gastric adenocarcinomas" *J. Natl. Cancer Inst.* 1991, 93, 640-643.
- 10 4. Marshall, B.J. et al "Attempt to fulfill Koch's postulate for *pyloric* Campylobacter" *Med. J. Aust.* 1985, 142, 436-439.
5. Morris, A. et al "Ingestion of Campylobacter *pyloridis* causes gastritis and raised fasting gastric pH" *Am. J. Gastroenterology*, 1987, 82, 192-199.
- 15 6. Engstrand, L. et al "Inoculation of barrier-born pigs with *Helicobacter pylori*: a useful animal model for gastritis type B" *Infect. Immun.* 1990, 58, 1763-1768.
7. Fox, J.G. et al "Gastric colonization by campylobacter *pylori* subsp. *mustelae* in ferrets" *Infect. Immun.* 1988, 56, 2994-2996.
- 20 8. Fox, J.G. et al "Helicobacter *mustelae*-associated gastritis in ferrets: an animal model of *Helicobacter pylori* gastritis in humans" *Gastroenterology* 1990, 99, 352-361.
9. Lee, A. et al "A small animal model of human *Helicobacter pylori* active chronic gastritis" *Gastroenterology* 1990, 99, 1315-1323.
- 25 10. Fox, J.G. et al "Helicobacter *felis* gastritis in gnotobiotic rats: an animal model of *Helicobacter pylori* gastritis" *Infect. Immun.* 1991, 59, 785-791.
11. Eaton, K.A. et al "Campylobacter *pylori* virulence factors in gnotobiotic piglets" *Infect. Immun.* 1989, 57, 1119-1125.
- 30 12. Peterson, W.L. "Helicobacter *pylori* and peptic ulcer disease" *N. Engl. J. Med.* 1991, 324, 1043-1048.

This Sheet Has Been Replaced



13. Czinn, S.J. and Nedrud, J.G. "Oral immunization against *Helicobacter pylori*" *Infect. Immun.* 1991, 2359-2363.
14. Brandtzaeg, P. "Role of H chain and secretory component in receptor-mediated glandular and hepatic transport of immunoglobulins in man" *Scand. J. Immunol.* 1985, 22, 111-146.
15. Brandtzaeg, P. "Production and secretion of immunoglobulins in the gastrointestinal tract" *Ann. Allergy* 1987, 59, 21-39.
16. Wyatt, J. I. "Local immune response to gastric campylobacter in non-ulcer dyspepsia" *J. Clin. Path.* 1986, 39, 863-870.
- 10 17. Lee, A. et al "Pathogenicity of *Helicobacter pylori*: A perspective" *Infect. Immun.* 1993, 61, 1601-1610.
18. Pallen, M.J. and Clayton, C.L. "Vaccination against *Helicobacter pylori* urease, letter" *Lancet*, 1990, 336, 186.
19. Evans, D.J. et al "Urease-associated heat shock protein of
15 *Helicobacter pylori* *Infect. Immun.* 1992, 60, 2125-2127.
20. Ferrero, R.L. and Lee, A. "The importance of urease in acid protection for the gastric-colonizing bacteria *Helicobacter pylori* and *Helicobacter felis* sp. nov." *Microb. Ecol. Health Dis.* 1991, 4, 121-134.
21. Chen, et al "Immunization against gastric *Helicobacter* infection in
20 a mouse/*Helicobacter felis* model, letter" *Lancet*, 1992, 339, 1120-1121.
22. Czinn, S. et al "Oral immunization protects germ-free mice against infection from *Helicobacter felis*". *Proceedings of the DDW, American Gastroenterological Association.* May 10-13, 1992, 1321, A-331.
23. Guo, M. and Liu, P.V. "Serological specificities of ureases of
25 *Proteus* species" *J. Gen. Microbiol.* 1965, 136, 1995-2000.
24. Michetti, P. et al "Specificity of mucosal IgAa response in Balb/c mice following *H. felis* or *H. pylori* challenges" *Proceedings of the DDW, American Gastroenterology Association.* May 10-13, 1992, 1001, A-251.
25. Davin, C. et al. "*H. pylori* urease elicits protection against *H. felis*
30 infection in mice" *Proceedings of the DDW, American Gastroenterology Association,* May 16-19, 1993, 1213, A-304.

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

32

26. Pallen, M.J. and Clayton, C.L. "Vaccination against *Helicobacter pylori* urease". *Lancet* 1990, 336, 186-7.
27. Pimentel, J.L. and Cook, M.E. "Improved growth in the progeny of hens immunized with jackbean urease" *Poultry Sci.* 1988, 64, 434-439.
28. Labigne, A. "Sequences of nucleotides coding for a protein having an urease activity". EPO patent application #EPO 0 367 644 A1, 1989. Intl publication # WO90/04030, 1990.
29. Clayton, C.L. et al. "Nucleotide sequence of two genes from *Helicobacter pylori* encoding for urease subunits". *Nucleic Acid Res.* 1990, 18, 362.
30. McGhee, J.R. and Kyono, H. "New perspectives in vaccine development: mucosal immunity to infections". *Infect Agents Dis.* 1993, 2, 55-73.

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

33

WHAT IS CLAIMED IS:

- 1 1. A method of eliciting in a mammalian host a protective immune
2 response to *Helicobacter* infection, said method comprising the step of:
3 administering to a mucosal surface of said mammal an
4 immunogenically effective amount of a polyaminoacid preparation
5 presenting a sufficient number of epitopes exhibited by a urease endogenous
6 to said *Helicobacter* organism to elicit a protective immune response to
7 infection by said organism.
8
- 1 2. A method according to claim 1, wherein said preparation comprises an
2 intact urease purified from an organism.
- 1 3. A method according to claim 1, wherein said preparation comprises
2 peptides homologous with enzymatically inactive portions of the aminoacid
3 sequences of a urease.
- 1 4. A method according to claim 1, wherein said preparation comprises
2 peptides non-homologous with the aminoacid sequences of a urease and
3 displaying epitopes cross-reacting of a urease.
- 1 5. A method according to claim 1, wherein said preparation comprises *H.*
2 *pylori* urease.
- 1 6. A method according to claim 1, wherein said preparation comprises at
2 least subunits of a urease, with or without enzymatic activity.
- 1 7. A method according to claim 1, wherein said preparation comprises
2 anti-idiotypic antibodies to a urease.
- 1 8. A method according to claim 1, wherein said preparation comprises
2 peptides immunologically cross-reacting with urease.

This Sheet Has Been Replaced




THIS SHEET HAS BEEN
REPLACED

34



- 1 9. A method according to claim 3 and 9, wherein said peptides are
2 obtained by chemical synthesis.
- 1 10. A method according to claim 1, wherein said preparation comprises
2 urease antigens producing using DNA recombinant techniques.
- 1 11. A method according to claim 1, wherein said preparation comprises
2 subgenic fragments of urease produced with recombinant techniques.
- 1 12. A method according to claim 1, wherein said preparation comprises
2 urease subgenic fragments produced as genetically fused proteins.
- 1 13. A method according to claim 12, wherein said fused proteins comprise
2 cholera toxin subunits.
- 1 14. A method according to claim 1, wherein said preparation is
2 administered in association with a mucosal adjuvant.
- 1 15. A method according to claim 14, wherein said mucosal adjuvant is
2 cholera toxin.
- 1 16. A method according to claim 1, wherein said mammalian host is
2 human.
- 1 17. A method according to claim 1, wherein said preparation is
2 administered in association with a hydroxylated calcium phosphate.
- 1 18. A method according to claim 17, wherein said hydroxylated calcium
2 phosphate is hydroxyapatite.

This Sheet Has Been Replaced





36

27. A method according to claim 27, comprising the additional step of administering a *Helicobacter* vaccine to said mammal prior to said administration.

This Sheet Has Been Replaced



THIS HAS BEEN
REPLACED

37

ABSTRACT OF THE DISCLOSURE

Method of eliciting in a mammalian host a protective immune
s response to *Helicobacter* infection by administering to the host an
immunogenically effective amount of a *Helicobacter* urease or urease
subunits as antigen. Vaccine compositions are also provided.

This Sheet Has Been Replaced

REPLACED SHEETS 3G

9604759-2
1805

☒ Description : paged ... to paged ...
☒ Claim(s) : paged ... to paged ...
☒ Drawing(s) : paged ... to paged ...
☒ Abstract : paged ... to paged ...

THIS SHEET HAS BEEN REPLACED

Urease vaccine against *Helicobacter pylori*, Michetti et al.

DATE OF RECEIPT

9604759-2 303

REPLACEMENT SHEET 3 0 5 9

13 FEB 1997

☒ Description : paged ... to paged ...
☒ Claim(s) : paged ... to paged ...
☒ Drawing(s) : paged ... to paged ...
☒ Abstract : paged ... to paged ...

DATE OF RECEIPT 13/2/97

UREASE-BASED VACCINE AGAINST *HELICOBACTER* INFECTION

The present invention relates to the prevention and treatment of gastric infection in mammals, including humans. More particularly, the present invention relates to a vaccine suitable for use in the prevention and treatment of *Helicobacter* infection in mammals, including humans, and to a method of treatment of humans suffering from gastric infection, its consequences such as chronic gastritis or peptic ulcer, and prevention of gastric cancer.

This Sheet Has Been Replaced

BACKGROUND OF THE INVENTION

Helicobacter infections of human gastric epithelium cause gastritis, are a major factor in the development of peptic ulcers and gastric lymphoma, and may be a risk factor for the development of gastric cancer [1-3]. The most frequent infection agent is *Helicobacter pylori*, followed at a much lower frequency by *Helicobacter heilmannii*. *H. pylori* is a slender S-shaped gram negative microorganism, which is routinely recovered from gastric biopsies of adults and children with histologic evidence of gastritis or peptic ulceration. Evidence for a causal relationship between *H. pylori* and gastroduodenal disease comes from studies in human volunteers, patients with ulcers and gastric cancer, gnotobiotic pigs, and germ-free rodents. Regarding etiology, Koch's postulates were satisfied by creating histologically confirmed gastritis in previously uninfected individuals following consumption of viable microorganisms [4-11], and by treatment to eradicate *H. pylori*, with resolution of the gastritis and, in patients with peptic ulcer disease, a decrease in the recurrence rate [12].

In spite of *in vitro* susceptibility to many antimicrobial agents, *in vivo* eradication of established *H. pylori* infections with antimicrobial agents is often difficult to achieve [13]. The microorganism is found within the mucous coat overlying the gastric epithelium and in gastric pits. These are locations which do not appear to allow for adequate antimicrobial levels to be achieved even when antibiotics are given orally at high doses. At the present time, most authorities recommend a "triple therapy", namely a bismuth salt in combination with drugs such as tetracycline and metronidazole for 2-4 weeks. However, the effectiveness of this or other chemotherapeutic regimens remains suboptimal. Furthermore, this treatment may produce serious adverse drug reactions.

At the present time little is known regarding the role of the mucosal immune systems in the stomach. The distribution of immunoglobulin (Ig) producing cells in the normal gastric antrum indicates that IgA plasma cells make up 80% of the total plasma cell population. In addition, the number of plasma IgA cells present in the gastric antrum is comparable to other mucous membranes [14, 15]. A number of studies in humans [16] and in animal models [8, 10] have demonstrated specific IgG and IgA responses in serum and in gastric secretions in response to *Helicobacter* infection. However, the observation that *H. pylori* infection persists as a chronic infection

13 FEB 1997

Urease vaccine against H. pylori, Michetti et al

for years, despite inducing a local and systemic immune response, is not encouraging the development of immunization strategies.

Lee et al have reported the ability to infect germ-free rodents with Helicobacter felis, a bacterium closely related to H. pylori, and reproducibly document histologic gastritis [9, 10]. Since then, this bacterium-host pairing has been accepted as a good model to study Helicobacter-mediated gastritis and its initiating factors [17]. Czinn et al have shown that repetitive oral immunization with a crude lysate of H. pylori plus cholera toxin adjuvant induces a vigorous gastrointestinal IgA anti-H. pylori response in mice and ferrets [13]. In addition, Chen et al and Czinn et al have recently reported that oral immunization with a crude lysate of H. felis induced protection against H. felis infection in mice [21, 22]. The exact nature of the antigen(s) responsible for the induction of this protection, however, had not been determined, and no information suggested that the protective antigen(s) of H. felis that induced protection against this pathogen would induce a cross-reactive protection extending to another Helicobacter species.

We have demonstrated for the first time that H. pylori and H. felis share antigenic determinants by obtaining monoclonal antibodies after oral immunization of mice with either H. pylori or H. felis sonicates and showing that some of these antibodies, directed against H. pylori, would crossreact with H. felis and vice versa [24;25]. The basis for these cross-reactivities were unknown.

Based on the homology existing between the different known urease amino acid sequences, it has been proposed that urease could be used as a vaccine against H. pylori [26]. Nevertheless, despite the homology among the different ureases sequences, cross-reactivity is not the rule. Guo and Liu have shown years ago that ureases of Proteus mirabilis, Proteus vulgaris and Providencia rettgeri show cross-reactivity to each other, while ureases of jack bean and Morganella morganii are immunologically distinct from the three former ureases [23]. Even if an antigenic cross-reactivity of H. pylori urease with other Helicobacter ureases was a reasonable postulate, no data existed demonstrating that this was really the case until we showed that some H. felis monoclonal antibodies crossreacted with H. pylori urease [25]. J. Pappo has further demonstrated that mice which have been infected by H. felis produce antibodies which crossreact with H. pylori urease but not jack bean urease (J. Pappo, unpublished data, 1993).

The use of H. pylori urease, or of related ureases, as a vaccine against H. pylori infection has previously been proposed by A. Labigne in EPO 367,644 [28]. However, that application contains no evidence of vaccination of any mammal against any Helicobacter infection with urease.

Moreover, while sequence homology with other bacterial ureases might support the use of urease as a vaccine candidate against H. pylori infection, the current knowledge of human H. pylori infection would certainly not. First, despite the fact that infected individuals often mount a strong antibody response to urease, the anti-urease immune response does not result in clearance or control of the



13 FEB 1997

-- 4

Urease vaccine against H. pylori, Michetti et al

infection. Secondly, H. pylori is able to transport urease out of the cell and to shed it from its surface [19, 20]. Thus, urease may not represent an appropriate target for the development of a protective mucosal immune response. Indeed, mucosal immune protection is thought to be mainly mediated by secretory IgA, the agglutinating activity of which would be impaired when the recognized antigen can be shed by the target pathogen and thereby serve as a decoy for the protective antibody. Thirdly, urease appears to be toxic for epithelial cells in culture, and has been suspected to play a role in mucous degradation and in peptic ulceration in vivo. Thus, its use as antigen may be toxic.

Nevertheless, we reasoned that this antigen could be a potentially efficient vaccine if:

- first, we would deliver it orally at a sufficiently high dose to elicit a stronger immune response than the naturally occurring one
- second, the amount of antibodies produced would be high enough to bind all the urease, shed or not shed
- third, we would use subunits of urease or a molecular species that was non toxic.

In summary, there remains a need for effective treatment and prevention of H. pylori-induced gastric infection in humans. Recent data suggested the possibility to generate a vaccine against this infection, but have not provided a clear identification of defined antigen(s), common to all strains of H. pylori, that could be incorporated into a safe and effective vaccine.

In this invention, we have identified the urease antigen of H. pylori as a candidate vaccine and demonstrated its efficacy in an animal model. These results were unexpected in the light of the natural history Helicobacter infections.

SUMMARY OF THE INVENTION

We have discovered that immunity can be induced in mammals susceptible to gastrointestinal Helicobacter infection by exploiting urease epitopes displayed on or about the surface of Helicobacter organisms and using them as a vaccine target. The immunity can be induced by immunization with native urease, but can also be induced with recombinant urease subunit, produced as an enzymatically inactive, therefore non-toxic form. The invention provides a method of inducing immunity to Helicobacter infection by administering to a mucosal surface of a mammal a polyaminoacid preparation, i.e. a mixture of peptides and/or proteins, together with an appropriate adjuvant. This polyaminoacid preparation presents a plurality of epitopes characteristic of and exhibited by a urease enzyme endogenous to the infecting Helicobacter organism. The administration of the polyaminoacid preparation may be performed by the oral route.

The active ingredient of the preparation may comprise natural or biosynthetic epitopes and may take various forms. A non exhaustive list of possible preparations includes purified, naturally occurring

13 FEB 1997

Urease vaccine against H. pylori, Michetti et al

- 5

or recombinantly produced urease preparations of bacterial or other origin, digests of urease, fusion proteins comprising urease epitopes, truncated forms of urease enzyme, or peptides homologous with the amino acid sequence of urease. Since development of immunity depends on induction of humoral and/or cellular immune responses which bind to the infecting Helicobacter organism, preferred preparations are those which most closely duplicate the epitopes of the urease endogenous to the infecting organism. For example, preparations displaying the epitopes of urease of H. pylori are preferred for administration in humans susceptible to H. pylori, and preparations displaying the epitopes of urease of H. heilmanii are preferred for the administration in humans susceptible to H. heilmanii. However, in accordance with an important aspect of the invention, it has been discovered that urease from other species may be used. For example, we have shown that H. felis infection in mice can be prevented by administration of urease from H. pylori.

According to one aspect of the invention, there is provided a method of eliciting in a mammalian host a protective immune response to Helicobacter infection wherein an immunologically effective amount of a urease antigen capable of eliciting such a protective immune response, preferably H. pylori urease or H. pylori urease B subunit, is administered to a mucosal surface of the host.

According to another aspect of the present invention, there is provided a vaccine composition suitable for prevention of Helicobacter infection, comprising an effective amount of a urease antigen, preferably H. pylori urease or H. pylori urease B subunit, capable of eliciting in a host a protective immune response to Helicobacter infection, in association with a pharmaceutically acceptable carrier or diluent.

According to a further aspect of the present invention, there is provided a method of imparting to a mammalian host passive protection to Helicobacter infection, comprising administering to a mucosal surface of the host an immunologically effective amount of a urease specific antibody produced in a host immunized with a urease, preferably H. pylori urease or H. pylori urease B subunit, capable of eliciting a protective immune response to Helicobacter infection.

This Sheet Has Been Replaced





13 FEB 1997

-- 6

Urease vaccine against H. pylori, Michotti et al**BRIEF DESCRIPTION OF THE DRAWINGS**

The invention will now be further described with reference to the accompanying drawings, in which Figures 1 through 6 are graphical representations of the results set forth in Tables 1 through 6.

DETAILED DESCRIPTION OF THE INVENTION

The present inventors have discovered that oral administration to mice of polyaminoacid preparations exhibiting the epitopes of H. pylori urease gives rise to a protective immunological response against H. felis in mice, an animal model of generally-accepted value for the study of the immune response to Helicobacter infection [9]. The effect of the protective immune response is that immunized animals, when challenged with pathogen, have a greatly reduced incidence of infection, in comparison to non-immunized animals. Furthermore, the inventors have discovered that oral immunization in mice using H. pylori urease B subunit, produced as an enzymatically-inactive recombinant protein, gives rise to a protective immunological response in mice against H. felis. The effect of the protective immune response is that immunized animals, when challenged with pathogen, have also a greatly reduced incidence of infection, in comparison to non-immunized animals which do become infected.

Thus, in a first aspect, the present invention provides a method of eliciting in a mammalian host a protective immune response to Helicobacter infection. The method comprises the step of administering to a mucosal surface of the mammal, including humans, an immunologically effective amount of a urease antigen, preferably H. pylori urease, capable of eliciting such a protective immune response.

In a second aspect, the present invention provides a method of eliciting in a mammalian host a protective immune response to Helicobacter infection. The method comprises the step of administering to a mucosal surface of the mammal, including humans, an immunologically effective amount of recombinant, enzymatically inactive, urease B subunit as antigen, preferably recombinant H. pylori urease B subunit, capable of eliciting such a protective immune response.

The invention also includes within its scope the treatment or prophylaxis of mammals, including humans, for Helicobacter infection, wherein an immunologically effective amount of a urease, or its subunits, capable of eliciting a protective immune response to Helicobacter infection, is administered to a mucosal surface of a patient. Preferably, the urease is H. pylori urease or H. pylori urease B subunit, and the urease may be administered either alone or linked to a hydroxylated calcium phosphate, for example hydroxyapatite as a carrier particle. Moreover, it is preferred to administer the H. pylori urease in association with a mucosal adjuvant, the B subunit of cholera toxin, muramyl dipeptide or other such adjuvants.



13 FEB 1997

Urease vaccine against H. pylori, Michetti et al

While not being bound by any theory, the present inventors believe that administration of the urease antigen, or B subunit thereof, to a mucosal surface stimulates the common mucosal immune system and perhaps local sites in the gastric mucosa inducing an immune response, including the appearance of H. pylori specific IgA antibodies in the gastric secretions, which prevent Helicobacter infection. Since it is a routine matter to conduct pre-clinical trials of candidate vaccines for human use in animal models, it is believed that the methodology of the present invention is effective in humans, especially in the prevention and treatment of peptic ulcers, gastritis, gastric malignancies and other conditions arising as a result of the presence of H. pylori and/or H. heilmanni.


A - Bacterial cultures and urease purification

The strain of H. pylori used in the study originates from a patient with a duodenal ulcer, and has been subcultured on BHI agarose plates to homogeneity. H. pylori is cultured in a suitable medium, typically, BHI (Brain-Heart Infusion) medium, containing 0.25% yeast extract and 10% fetal calf serum and supplemented with 0.4% Campylobacter selective complement (Skirrow supplement; Oxoid 69). The bacteria are incubated overnight under microaerophilic conditions at 37°C in bottles that are then sealed and shaken at 37°C for 2 to 3 days to produce a liquid culture. A culture may also be prepared in agarose plates consisting of BHI with 0.25% of yeast extract and 5% of sheep blood under microaerophilic conditions at 37°C for 3 days. The quantity of bacteria is determined by optical density of the BHI solution at 660 nm, with one optical density unit corresponding to 10^8 bacteria. Cultures on agarose plates are first resuspended in 154mM NaCl.

One currently preferred source of polyaminoacid displaying urease epitopes is purified urease, e.g., H. pylori urease obtained by following the general method of Dunn et al. J. Biol. Chem. 265, 9464-9469, modified as described below. Following culturing, the H. pylori is harvested in water, spun vortexed and spun again to produce a supernatant. The solution containing the urease activity of H. pylori (assessed by rapid urease test, see below) is then chromatographed on a CL-6B sizing column and the fractions which present a strong urease activity are pooled and dialysed overnight and again chromatographed on an anion exchanger gel. The fractions are eluted in increasing NaCl buffer and the collected fractions with a strong urease activity are individually submitted to a SDS gel followed by Coomassie staining. Two distinct bands corresponding to a molecular weight of about 63 and about 28 kDa are identified as urease. The fractions containing urease are pooled to give purified H. pylori urease having a purity in the region of 95% to 99%.

B - Oral immunization with urease purified from H. pylori

While it is preferred to employ purified H. pylori urease obtained as described above as the antigenic material, it will be understood that it is also possible to use, as the antigenic material, any urease or subunit of urease, either naturally occurring or obtained by recombinant DNA techniques, as well as digested fragment thereof, fusion proteins comprising the fragments or the whole urease,



13 FEB 1997

Urease vaccine against H. pylori, Michetti et al

truncated urease constructs, or other peptide or protein preparations exhibiting urease epitopes which are capable of eliciting a protective immune response to Helicobacter infection (See below). Thus, it is possible to employ a urease having a substantial homology with respect to H. pylori urease and which is effective in raising a cross-protective immune response to Helicobacter. An example of such a urease is jack bean urease, which possesses about 70% homology with H. pylori urease. The invention is therefore not limited to the use of intact urease, and covers the use of any polyaminoacid preparation which displays urease epitopes and is effective to generate a protective immunological response in a host to Helicobacter infection. Typically, a urease having a homology of 70-95% homology, for example, 80-90% homology, with respect to H. pylori urease, may be employed as the urease antigen in the invention.

A non-limiting list of sources of potentially useful urease preparations includes endogenous urease enzymes of the different Helicobacter species, urease from other bacteria such as Klebsiella pneumoniae or Proteus mirabilis, and, by analogy, any other urease with the condition that these ureases share cross-reactive epitopes with H. pylori urease. The urease genes of all the organisms mentioned above represent a potential tool for expressing recombinant urease products as a whole protein or as a part thereof.

A non-limiting list of potentially useful urease preparations includes peptides generated from purified urease (the sources are mentioned above), using physical and/or chemical cleavage procedures (i.e. CnBr) and/or proteolytic cleavage (using proteases e.g. V8-protease, trypsin or others); or peptides synthesised chemically and sharing crossreactive epitopes with urease.

Other sources of potentially useful epitopes include epitopes identified by their crossreactivity with urease, as the result of screening with anti-urease antibodies. These peptides can be naturally occurring peptides or peptides resulting from chemical synthesis. Furthermore such peptides can result from the expression of recombinant random oligonucleotides.

Another source of potentially useful epitopes includes epitopes similar to urease as a result of the generation of anti-idiotypic antibodies to urease. Such anti-idiotypic antibodies, generated in any immunocompetent host, are obtained by immunization of this host with anti-urease antibodies, with the goal of generating antibodies directed against anti-urease antibodies, which share structural homologies with urease.

The discussion herein focuses on the use of urease naturally produced by H. pylori (section B). However, it will be appreciated that the urease or subunits or constructs thereof mentioned above, capable of eliciting the desired protective immune response, may be produced by recombinant DNA techniques well known in the art. The efficacy of particular preparations may be determined by routine administration using animal models, oral administration of the candidate vaccine, and challenge with pathogen using a protocol substantially similar or identical to the procedure described below.

13 FEB 1997

- 9

Urease vaccine against H. pylori, Michetti et al.

Tables 1 and 2 below and Figures 1-5 describe the results obtained when mice were orally immunized with purified H. pylori urease. In this first experiment, administration of the H. pylori antigen was carried out by orally administering to the mice H. pylori urease purified as described in section A, and coupled to hydroxyapatite crystals, used as a carrier to enhance M cell binding and uptake. Cholera toxin (Sigma) was given as a mucosal adjuvant. In this experiment, groups of female SPF BALB/c six-week old mice were each orally immunized with 30 ug of purified H. pylori urease coupled to 1 mg of hydroxyapatite plus 10 ug of cholera toxin adjuvant at day 0, 7, 14 and 21. The mice were then challenged twice with 10^8 H. felis, at day 28 and 30. For comparison purposes, similar female SPF BALB/c six-week old mice were orally immunized with whole H. pylori lysate (sonicate) and 10 ug cholera toxin at day 0, 7, 14 and 21. The mice were challenged at day 28 and 30 with H. felis. The H. pylori sonicate was prepared by collecting H. pylori from cell cultures, pelleting by centrifugation and resuspending the pellet in 0.9% sodium chloride followed by sonication.

As a control, female SPF BALB/c six-week old mice were orally sham-immunized with 10 ug of cholera toxin and 1 mg of hydroxyapatite at day 0, 7, 14 and 21. The mice were then challenged at day 28 and 30 with H. felis. All mice were housed, immunized, and challenged in parallel. All mice subject to the study were sacrificed on day 35.

C - Oral immunization with recombinant urease subunits of H. pylori

Genes encoding the structural A and B subunits of H. pylori urease were obtained by polymerase chain reaction (PCR) cloning according to standard procedures, based on previously published sequences [29]. These genes were inserted in a vector (named pEV40) designed for high expression and easy purification of foreign genes in E. coli. Briefly, the foreign gene is inserted downstream of a thermo-repressible promoter, and in frame of a sequence encoding for a repeat of six histidines. An ampR gene is present on this vector for selection of transformants. Under the appropriate temperature conditions, the recombinant protein obtained is supplemented by six histidines at the N-terminal, which allow for a one-step affinity purification on a nickel column. Both H. pylori recombinant urease A and B subunits were expressed separately in E. coli, and purified on nickel column to >95% purity.

While it is preferred to employ recombinant H. pylori urease obtained as described above as the antigenic material, it will be understood that it is also possible to use, as the antigenic material, any urease or subunit of urease obtained by recombinant techniques (e.g. fusion protein) expressing antigenic sites of urease, which is capable of eliciting a protective immune response to Helicobacter infection. Thus, it is possible to employ in a construct a urease gene having a substantial homology with respect to H. pylori urease and which is effective in raising a cross-protective immune response to Helicobacter. Examples of such a urease is jack bean urease, which possesses about 70% homology with H. pylori urease, or H. felis urease, which possesses about 88% homology with H. pylori urease. The invention is therefore not

13 FEB 1997

- 10

Urease vaccine against H. pylori, Michetti et al

limited to the use of H. pylori urease genes and their gene products, and covers the use of any recombinant urease, or the subunits thereof, which is effective to generate a protective immunological response in a host to Helicobacter infection. Typically, a recombinant urease having a homology of 70-95% homology, for example, 80-90% homology, with respect to H. pylori urease, may be employed as the recombinant urease antigen in the invention.

The discussion herein focuses on the use of recombinant H. pylori urease A and B subunits produced by E. coli (section C). However, it will be appreciated that recombinant urease or subunits or constructs thereof mentioned above, capable of eliciting the desired protective immune response, may be produced using other recombinant DNA techniques and other eukaryotic or prokaryotic expression vectors well known in the art.

Tables 3, 4 and 5 below and Figure 6 describe the results obtained when mice were orally immunized with recombinant H. pylori urease subunits produced in E. coli. In this experiment, administration of the H. pylori antigen was carried out by orally administering to the mice recombinant H. pylori urease A or B subunits produced in E. coli and purified as described above, and coupled to hydroxyapatite crystals, used as a carrier to enhance M cell binding and uptake. Cholera toxin (Sigma) was given as a mucosal adjuvant. In this experiment, groups of female SPF BALB/c six-week old mice were each orally immunized with 30 ug of recombinant H. pylori urease A subunit, coupled to 1 mg of hydroxyapatite plus 10 ug of cholera toxin adjuvant at day 0, 8, 14 and 21. The mice were then challenged twice with 10^8 H. felis, at day 32, 34 and 36. For comparison purposes, similar female SPF BALB/c six-week old mice were orally immunized with 30 ug of recombinant H. pylori urease B subunit coupled to hydroxyapatite plus 10 ug cholera toxin at day 0, 8, 14 and 21. The mice were challenged three times, at day 32, 34 and 36, with H. felis. As a control, female SPF BALB/c six-week old mice were each orally sham-immunized with 10 ug of cholera toxin and 1 mg of hydroxyapatite at day 0, 8, 14 and 21. The mice were then challenged at day 32, 34 and 36 with H. felis. All mice subject to the study were immunized and challenged in parallel. Animals were sacrificed on day 48 (12 days after challenge) or 10 weeks after challenge.

D - Analysis of gastric biopsies, blood, and intestinal secretions

Biopsies were taken from the stomach and blood was obtained from the heart. The intestines were removed and washed with 1mM PMSF (Boehringer) in PBS buffer to obtain intestinal secretions for ELISA analysis.

To evaluate protection against H. felis colonization, gastric biopsies from each animal were screened for the presence of H. felis by assessing rapid urease activity by the Jatrox HP test (Rohm Pharma), according to the supplier's directions. Briefly, gastric biopsies are immersed in 0.5 ml supplier's mixture of urea and phenol red, a pH indicator. Urease activity generates ammonia and bicarbonate from urea, and is followed by the colorimetric change of



13 FEB 1997

- 11

Urease vaccine against *H. pylori*, Michetti et al

the solution towards a higher absorbance at 550 nm. Urease activity was quantified by spectrophotometric analysis.


Gastric biopsies of each animal included in the experiment described in section B were also cultured on BHI agarose plates, supplemented as above, for the detection of *H. felis*. After incubation for 3 to 10 days in microaerophilic conditions, the presence of *H. felis* was confirmed by Gram staining and determination of urease activity. As a very significant correlation was obtained for the detection of *H. felis* in gastric biopsies between urease tests and *H. felis* cultures during the first set of experiments (see Table 3), only gastric biopsies urease tests were performed for the detection of *H. felis* in the experiment described in section C. Detection of *H. felis* was confirmed by microscopy by two independent investigators, using two different stains (acridine orange and cresyl violet).

Blood samples were allowed to clot for 3 hours at RT, and sera harvested and frozen at -20°C until analyzed. Intestinal secretions were spun for 5 min. at 4°C to remove debris, and kept frozen at -20°C. Serum and intestinal samples of each animal were analyzed by ELISA for evaluation of anti-*Helicobacter* activity, according to standard procedures. Briefly, 96-well plates were coated with a sonicate of *H. pylori*, followed by saturation with 5% fat-free milk. Samples were serially diluted from 1:1 to 1:1000 and incubated overnight at 4°C on ELISA plates. Biotinylated anti-mouse IgG (serum) and anti-mouse IgA, followed by streptavidin-Horseradish peroxidase was used for the determination of the antibody levels.

The results of *H. felis* challenges following immunizations with purified *H. pylori* urease are set out in Tables 1-3 and Figures 1-4 and the results of *H. felis* challenges following immunizations with recombinant *H. pylori* urease A and B subunits are set out in Tables 4-6 and Figures 5 and 6.

TABLE 1

mouse		urease	culture	Immunoglobulins			
number	Immunization	test	Gram	Serum		Intestinal secretion	
		12h		Ig	IgG	Ig	IgA
1	Urease+HF	+	<i>H felis</i>	27	0	25	258
2	Urease+HF	0	0	264	273	221	246
3	Urease+HF	0	0	84	44	318	354
4	Urease+HF	+	<i>H felis</i>	81	42	12	5
5	Urease+HF	0	0	98	137	126	234
6	Urease+HF	+	0	968	2093	31	22
7	Urease+HF	0	0	98	0	96	34
8	Urease+HF	0	0	247	1010	214	128
9	Urease+HF	0	0	N.D.	N.D.	48	23
10	Urease+HF	0	0	50	0	124	99
11	Urease	0	0	319	205	44	53
12	Urease	0	0	14	0	86	87



13 FEB 1997

- 12

Urease vaccine against *H. pylori*, Michetti et al

13	Urease	0	0	0	0	0	0
14	Urease	0	0	0	0	43	61
15	Urease	0	0	58	0	110	127
16	Urease	0	0	140	63	21	37
17	Urease	0	0	84	240	114	280
18	Urease	0	0	N.D.	N.D.	93	148
19	Urease	0	0	45	0	135	216
20	Urease	0	0	261	197	161	261
21	CT+HF	0	0	0	0	0	2
22	CT+HF	+	<i>H felis</i>	63	0	310	303
23	CT+HF	+	<i>H felis</i>	90	0	N.D.	N.D.
24	CT+HF	+	<i>H felis</i>	31	0	150	192
25	CT+HF	+	<i>H felis</i>	197	250	250	440
26	CT+HF	+	<i>H felis</i>	105	135	214	138
27	CT+HF	+	<i>H felis</i>	140	47	109	55
28	CT+HF	+	0	0	0	16	15
29	CT+HF	+	<i>H felis</i>	0	0	0	0
30	CT+HF	+	<i>H felis</i>	N.D.	N.D.	N.D.	N.D.
31	HP sonicate+HF	+	<i>H felis</i>	0	0	76	103
32	HP sonicate+HF	+	<i>H felis</i>	77	0	11	33
33	HP sonicate+HF	+	<i>H felis</i>	549	748	57	36
34	HP sonicate+HF	0	0	660	153	180	286
35	HP sonicate+HF	+	<i>H felis</i>	730	192	0	5
36	HP sonicate+HF	+	<i>H felis</i>	32	0	5	64
37	HP sonicate+HF	0	0	400	400	312	1149
38	HP sonicate+HF	+	<i>H felis</i>	1007	1360	149	26
39	HP sonicate+HF	0	0	220	186	133	122
40	HP sonicate	0	0	873	1016	352	514
41	HP sonicate	0	0	727	899	126	191
42	HP sonicate	0	0	109	68	44	83
43	HP sonicate	0	0	147	949	167	97
44	HP sonicate	0	0	845	1094	246	64
45	HP sonicate	0	0	1217	1198	210	157
46	HP sonicate	0	0	81	0	256	218
47	HP sonicate	0	0	329	210	241	276
48	HP sonicate	0	0	1049	737	197	211

In Table 1, which refers to the experiment described in section B, "h" means hour, "Ig" means immunoglobulin, "ND" means "not determined", "urease + HF" means that the mice were immunized with urease (coupled to hydroxyapatite, with cholera toxin) and then challenged with *H. felis*, "urease" means that the mice were immunized with urease (coupled to hydroxyapatite, with cholera toxin) and not challenged, "CT+HF" means that the mice were sham-immunized with cholera toxin and challenged with *H. felis*, "HP sonicate + HF" means that the mice were immunized with *H. pylori* sonicate with cholera toxin and challenged by *H. felis*, and "HP sonicate" means that the mice were immunized with *H. pylori* sonicate



13 FEB 1997

- 13

Urease vaccine against *H. pylori*, Michetti et al.

with cholera toxin and not challenged. In Table 1, the numbers for the antibody results are given as a measure of absorbance at 595 nm multiplied by 1000. The background measured in absence of the antibodies, was subtracted.

The results of experiment described in section B obtained on the basis of the gastric biopsies urease tests and on Gram staining of *H. felis* cultures are set out in Table 2. Infection was defined by mice with one or more markers of colonization by *H. felis*, including urease test or Gram staining of cultures.

TABLE 2

<u>Immunization</u>	<u>Challenge</u>	<u>% infected</u>	<u>% protected</u>
Urease	<i>H. felis</i>	3/10 (30%)	7/10 (70%) *
Sonicate	<i>H. felis</i>	6/9 (66%)	3/9 (33%) **
CT	<i>H. felis</i>	9/10 (90%)	1/10 (10%)

* $p=0.0198$ (two tailed Fisher exact test) compared to CT control

** $p=0.303$ (two tailed Fisher exact test) compared to CT control

It will be seen from the results set out in Tables 1 and 2 that statistically significant protection against *H. felis* challenge is obtained with oral immunization using *H. pylori* urease as compared to that obtained using either *H. pylori* sonicate or cholera toxin. Referring to Table 2, it will be seen that from a total of 10 immunized animals, only 3 became infected, as compared to 6 of the animals immunized with *H. pylori* sonicate and 9 of the animals immunized with cholera toxin. Table 2 shows that 70% of the animals were protected from challenge by *H. felis* as compared to 33% of the animals immunized with *H. pylori* sonicate and 10% of the animals immunized with cholera toxin and then subjected to *H. felis* challenge. In other words, 90% of the control mice exposed to *H. felis* became infected by that pathogen whereas, in contrast, in mice immunized with *H. pylori* urease 28 days before exposure to *H. felis*, the infection rate was only 30%. This represents a significant reduction in infection ($p=0.0198$ in the Fisher exact test, as compared to the control mice). When the mice were orally immunized with *H. pylori* sonicate, the infection rate was 67% (not significant versus the control). The protection obtained using *H. pylori* urease is unexpected and could not have been predicted on the basis of the results observed using *H. pylori* sonicate.

Referring to Figures 1-4, Figure 1 represents graphically the results of tests for antibodies in serum (IgG) and intestinal secretion (IgA) in mice not protected after immunization with urease. These are mice numbers 1, 4 and 6 appearing in Table 1, and constitute Group A. Figure 2 shows the antibody responses of mice that were protected after immunization with urease (Group B), i.e. mice 2, 3, 5 and 7-10.



13 FEB 1997

Urease vaccine against *H. pylori*, Michetti et al.

Figures 3 and 4 relate to the results obtained with mice numbers 31-39. Figure 3 (Group C) depicts antibody responses of mice not protected after immunization with *H. pylori* somicate (mice numbers 31, 32, 33, 35, 36 and 38) and Figure 4 (Group D) depicts the antibody responses of mice protected after immunization with *H. pylori* sonicate (mice numbers 34, 37 and 39). It is of interest to note with respect to Figures 3 and 4 that the IgA antibody responses (but not IgG) are higher in the mice exhibiting protection than in the mice that are not protected, suggesting a correlation between protection and IgA response. Serum IgG responses did not exhibit a correlation. Mucosal IgA but not serum IgG antibodies are known to play a role in protection against bacterial infections of the gut [30].

The results of the correlation between the detection of *H. felis* in gastric biopsies by urease tests and by cultures are set out in Table 3.

TABLE 3

	Urease Test +	Urease Test -	Total
<i>H. felis</i> culture +	16	0	16
<i>H. felis</i> culture -	2	30	32
Total	18	30	48

Two-tailed Fisher's Exact Test : $p < 0.00001$

Table 3 shows that a very significant correlation exists between the results of urease tests performed on gastric biopsies and the identification of *H. felis* by cultures. As cultures detected less often *H. felis* infection than urease tests, urease tests were preferred for the diagnosis of *H. felis* infection in mice in the next experiments, due to its better sensitivity. This approach allowed the duplication of urease tests with larger fragments of the stomach of each mouse, and a further increase in the sensitivity of the urease test. Furthermore, the use of the method with the highest sensitivity prevent an overestimation the protection obtained by the vaccine preparation to be tested. When positive culture is used as the standard for infection, the protection induced after urease immunization during the experiment depicted in section B is as significant as with the combined use of urease test and culture ($p=0.021$ versus $p=0.019$).

The results of the experiments described in section C (recombinant urease subunits), obtained on the basis of the gastric biopsies urease tests, are set out in Table 4, 5 and 6 and depicted in Figure 6.

TABLE 4

immunization	mice n°	Urease test	Infection
CT	20	0.49	+
	21	0.31	+
	22	0.62	+

THIS SHEET HAS BEEN
REPLACED

REPLACEMENT SHEET

13 FEB 1997

Urease vaccine against *H. pylori*, Michetti et al

This Sheet Has Been Replaced



	23	0.67	+
	24	0.55	+
	50	0.50	+
	51	0.37	+
	52	0.29	+
	53	0.79	+
	54	0.32	+
ureA + HAP + CT	40	0.67	+
	41	0.48	+
	42	0.42	+
	43	0.65	+
	44	0.56	+
	45	0.52	+
	46	0.33	+
	47	0.63	+
	48	0.22	+
	49	0.37	+
ureB + HAP + CT	25	0.15	-
	26	0.07	-
	27	0.03	-
	28	0.64	+
	29	0.13	-
	30	0.02	-
	31	0.66	+
	32	0.00	-
ureA + HAP + CT	68	0.00	-
	69	0.07	-
	70	0.42	+
	71	0.00	-
	72	0.00	-
ureB + HAP + CT	73	0.37	+
	74	0.00	-
	75	0.37	+
	76	0.00	-
	77	0.00	-
	78	0.00	-
	79	0.39	+
	80	0.00	-
	81	0.37	+
	82	0.00	-

In Table 4, "CT" means cholera toxin; "UreA" means recombinant *H. pylori* urease A subunit; "UreB" means recombinant *H. pylori* urease B subunit; and "HAP" means hydroxyapatite crystals. Mice 20 to 54 were sacrificed 12 days post challenge and mice 68 to 82 10 weeks (106 days) post challenge. The results of the urease test performed from biopsies of the stomach of each animal are expressed as OD values at 550 nm. The positive and negative signs depicts the final status of infection of each animal, according to the positivity or negativity of the urease test for detection of *H. felis*. Positivity: OD550 values >0.2



13 FEB 1997

16

Urease vaccine against *H. pylori*, Michetti et al

TABLE 5: Protection as measured 12 days post challenge

<u>Immunization</u>	<u>Challenge</u>	<u>% infected</u>	<u>% protected</u>
Urease A subunit	<i>H. felis</i>	10/10 (%)	0/10 (0%)
Urease B subunit	<i>H. felis</i>	3/10 (30%)	7/10 (70%) *
CT	<i>H. felis</i>	10/10 (100%)	0/10 (0%)

* p=0.0031 (two tailed Fisher exact test) compared to CT control

TABLE 6: Protection as measured 10 weeks post challenge


<u>Immunization</u>	<u>Challenge</u>	<u>% infected</u>	<u>% protected</u>
Urease A subunit	<i>H. felis</i>	1/5 (20%)	4/5 (80%) *
Urease B subunit	<i>H. felis</i>	4/10 (40%)	6/10 (60%) **

* p=0.003 (two tailed Fisher exact test) compared to CT control

** p=0.01 (two tailed Fisher exact test) compared to CT control

It will be seen from the results set out in Tables 4, 5 and 6 that statistically significant protection against *H. felis* challenge is obtained with oral immunization using recombinant *H. pylori* urease B subunit as compared to that obtained using either recombinant *H. pylori* urease A subunit or cholera toxin. Referring to Table 4, it will be seen that, 12 days post challenge, from a total of 10 immunized animals, only 3 were found infected in the urease B subunit group, as compared to all 10 animals immunized with *H. pylori* A subunit of urease and 10 out of 10 of the animals immunized with cholera toxin. Table 4 shows that 70% of the animals were protected from challenge by *H. felis* as compared to 0% of the animals immunized with *H. pylori* urease A subunit and 0% of the animals immunized with cholera toxin and then subjected to *H. felis* challenge. In other words, 100% of the control mice challenged with *H. felis* became infected whereas, in contrast, in mice immunized with recombinant *H. pylori* urease B subunit the infection rate was only 30%. This represents a significant reduction in infection (p=0.0031, Fisher exact test) as compared to the control mice.

The fact that the protection observed with *H. pylori* urease is entirely conferred by immunization with the B subunit of urease, and that the A subunit has no such effect, was not expected on the basis of our experiment with purified urease. This definition of the roles of the 2 structural subunits of urease in the development of a protective immune response is therefore novel. The protection obtained using recombinant urease B subunit, which is enzymatically inactive also teaches that non toxic forms of urease can be used as oral vaccine against *Helicobacter* infection. Furthermore these results strongly suggest that recognition of the active site is not required for protection, as an inactive urease B subunit is very



13 FEB 1997

- 17

Urease vaccine against H. pylori, Michetti et al

unlikely to induce antibodies that will recognize and inhibit the catalytic site of native urease.

Referring to Table 6, it will be seen that, when mice are sacrificed 10 weeks post infection, 60% (6 mice out of 10) of the animals immunized with urease B subunit and 80% (4 mice out of 5) of the animals immunized with H. pylori urease A subunit were protected against H. felis infection. The fact that protection obtained through immunization with urease B subunit lasts over time and that immunization with urease A induces a protection which is delayed compared to the one induced by urease B subunit could not be expected from our experiment with purified urease or with other experiment performed earlier. The fact that urease A subunit immunization confers protection definitively proves that recognition of the active site is not required for protection. Figure 6 summarizes results obtained after oral immunization with recombinant urease A and B subunits (described in Table 5 and 6)

The present invention also provides vaccine compositions suitable for the prevention of Helicobacter infection. The compositions comprise an effective amount of a urease antigen, preferably H. pylori urease or recombinant H. pylori urease subunits, capable of eliciting in a host a protective immune response to Helicobacter infection, in association with a pharmaceutically acceptable carrier or diluent.

The vaccines of the invention are administered in amounts readily determined by persons of ordinary skill in this art. Thus, for adults a suitable dosage will be in the range of 10ug to 100 milligrams, for example 50ug to 50mg. Similar dosage ranges will be applicable for children. Carrier systems in humans may include enteric release capsules protecting the antigen from the acidic environment of the stomach, and including urease antigen in a insoluble form as fusion proteins. The vaccine can be administered as a primary prophylactic agent in adults or in children, as a secondary prevention, after successful eradication of H. pylori in an infected host, or as a therapeutic agent in the aim to induce an immune response in the host susceptible to contribute to the eradication of H. pylori.

As noted above, a suitable mucosal adjuvant is cholera toxin. Others which may be used are muramyl dipeptide or its derivatives, non-toxic derivatives of cholera toxin, including its B subunit, and/or conjugates or genetically engineered fusions of the urease antigen plus cholera toxin or its B subunit. Other suitable delivery methods include biodegradable microcapsules or immune stimulating complexes (ISCOM'S) or liposomes, genetically engineered attenuated live vectors such as viruses or bacteria, and recombinant (chimeric) virus-like particles, e.g. bluetongue. The amount of mucosal adjuvant employed depends on the type of mucosal adjuvant used. For example, when the mucosal adjuvant is cholera toxin, it is suitably used in an amount of 5 ug to 50 ug, for example 10ug to 35ug. When used in the form of microcapsules, the amount used will depend on the amount employed in the matrix of the microcapsules to achieve the desired dosage. The determination of this amount is within the skill of a person of ordinary skill in this art.



13 FEB 1997

- 18

Urease vaccine against H. pylori, Michetti et al

Suitable carriers for the vaccines of the invention are enteric coated capsules and polylactide-glycolide microspheres. Suitable diluents are 0.2N NaHCO_3 and/or saline.

Particulate hydroxylated calcium phosphate (HCP) is especially useful as a carrier for the H. pylori urease to be applied to mucosal surfaces. It is believed that the H. pylori urease-hydroxylated calcium phosphate conjugate is transported across epithelium where it raises a poly Ig immune response. Preferably, the hydroxylated calcium phosphate is in the form of microparticles suitable for transport across the epithelium, particularly by cells specialized for this purpose (M cells). A preferred form of hydroxylated calcium phosphate is hydroxyapatite, a commercially available crystalline hydroxylated calcium phosphate $\text{Ca}_{10}(\text{PO}_4)_6(\text{OH})_2$.

Commercially available hydroxyapatite generally consists of slab-like crystals that are chemically and physically analogous to inorganic hydroxyapatite in normal bone tissue. Ingestion of hydroxyapatite should therefore be safe, as evidenced by the existence of nutritional calcium/phosphorus supplements derived from ground bone, which are designed to be ingested. Commercially-high resolution hydroxyapatite (from CalBioChecm) consists of crystals varying widely in size. Crystals over 1um in length are unlikely to be taken up by M cells. Therefore, for use in the invention, commercial hydroxyapatite crystals are broken into small, relatively uniform crystalline fragments such as by sonication. Preferably, a substantial proportion of the hydroxyapatite is present as fragments of about 0.01-1.0um. Fragmentation may be measured either by electron microscopy or light scattering, using standard techniques.

Preferred modes of administration of the H. pylori urease antigen are orally, nasally, rectally or ocularly. Oral administration can provide delivery to other G.I. mucosa including the intestinal mucosa.

The vaccines of the present invention may be administered to a mucosal surface in the form of an aerosol, suspension, capsule and/or suppository. The method of administration will be readily apparent to a person of ordinary skill in this art.

The present invention further includes the passive immunization of mammals, including humans, against Helicobacter infection. This is achieved by administering to a mucosal surface of the patient an effective amount of a urease specific antibody. Preferably, an effective amount of a H. pylori urease specific IgA monoclonal antibody.

Since the urease of H. pylori is shown to represent the antigen involved in inducing protective immunity, a further aspect of the invention is the use of H. pylori urease as a diagnostic reagent to measure the immune response of persons who have received a vaccine based on urease or to determine whether an individual is immune or susceptible (and thus in need of vaccination). The present invention also includes the use of urease and urease-specific antibodies to construct assays and kits for diagnosis of Helicobacter immunity,



13 FEB 1997

- 19

Urease vaccine against H. pylori, Michetti et al.

assessment of Helicobacter susceptibility, and definition of immune responses to vaccines.

EXAMPLES

The invention will now be further described by reference to the following non-limiting examples.

a) The Bacterial Strains

H. felis was provided by J. Fox (division of Cooperative Medicine, Mass. Institute of Technology, Boston, USA). H. pylori was isolated from patients with ulcer disease (CHUV, Lausanne, Switzerland).

b) Bacterial Cultures

Liquid Culture - Bacteria were cultured on BHI (Brain-Heart Infusion, BioMerieux) liquid medium containing 0.25% of yeast extract (Difco) and 10% of fetal calf serum (Inotech) supplemented with 0.4% of Campylobacter selective complement (Oxoid). The bacteria were incubated overnight under microaerophilic conditions at 37°C and then shaken at 37°C for 2 to 3 days.

Culture on Agarose plates - The bacteria were cultured on agar plate consisting on BHI with 0.25% of yeast extract and 5% of sheep blood under microaerophilic conditions at 37°C for 3 days.

Quantification - The quantity of bacteria was determined by the optical density of the BHI solution at 660 nm (1 optical density unit corresponding to 10^8 bacteria).

c) Preparation of sonicates

H. pylori was collected from 31 blood agar plates in 0.15 M NaCl and spun 5 minutes at 1400g at 4°C. The pellet was resuspended in 3 ml of NaCl and sonicated for 4 minutes. The amount of proteins was evaluated by a Bradford assay (BioRad Kit according to supplier).

d) Coupling of immunogen to Hydroxyapatite

Immunogen (urease or subunit thereof) was incubated for 1 hour at 4°C with hydroxyapatite. 1.0 mg of hydroxyapatite was used for 30ug of immunogen per mouse. At the end of the incubation, 10 µg of cholera toxin was added in a final volume of 200ul PBS.

EXAMPLE 1

a) Extraction

H. pylori from 30 blood agar plates was harvested in 0.15 M NaCl on ice. The solution was spun 5 minutes at 1400 g at 4°C. The pellet was resuspended in 20 ml of H₂O and vortexed for 45 seconds (maximum speed). The extract was then spun 20 minutes at 6700 g at 4°C. The supernatant was recovered and the quantity of protein was evaluated (see "Quantification" above) and precipitated with 70% of ammonium sulfate.

13 FEB 1997

- 20

Urease vaccine against *H. pylori*, Michetti et al**b) Purification of urease**

The solution was chromatographed on a Sepharose CL-6B column (Pharmacia) with PBS (phosphate buffered saline) as mobile phase. The 22 collected fractions which presented a strong urease activity were pooled and dialyzed overnight at 4°C against 3 liters of PEB (20 mM phosphate buffer, pH 7) and then chromatographed on a Q Sepharose fast flow (Pharmacia) with PEB as mobile phase. The fractions were eluted by 0 to 500 mM NaCl gradient. Ten of the collected fractions with a strong urease activity were individually subjected to an SDS gel followed by a Coomassie staining. The 6 fractions presenting 2 distinct bands corresponding to MW-63 and -28 KDa were pooled and were considered as the purified urease.

EXAMPLE 2 (see also section B)

Mice employed in the immunization studies were lightly anesthetized with ether prior to intragastric immunization. And then, sonicate preparation or purified urease, hydroxyapatite and cholera toxin was suspended in PBS and 200ul were delivered to the stomach of the respective mice using a polyethylene tubing attached to a hypodermic syringe. This procedure will be referred to as oral immunization.

Three oral immunization protocols were evaluated. These are described below.

Protocol B1 - Vaccination with purified urease

Female BALB/c six-week old mice (20) were orally immunized with 30ug of purified of *H. pylori* urease and 1 mg of hydroxyapatite and 10ug of cholera toxin at day 0, 7, 14 and 21. Ten mice were challenged at day 28 and 30 with 5×10^7 and 10^8 *H. felis* from liquid culture.

Protocol B2 - Vaccination with Helicobacter sonicates

Female BALB/c six-week old mice (20) were orally immunized with 2 mg of *H. pylori* sonicate solution at day 0, 7, 14 and 21. Ten mice were challenged at day 28 and 30 with 5×10^7 and 10^8 *H. felis*.

Protocol B3 - Control

Female BALB/c six-week old mice were orally immunized with 1 mg hydroxyapatite and 10ug of cholera toxin at day 0, 7, 14 and 21. The mice were challenged at day 28 and 30 with 5×10^7 and 10^8 *H. felis*.

At day 35 all mice were sacrificed and biopsies from the stomach were taken as well as intestinal secretions and blood.

Protection and evaluation

This Sheet Has Been Replaced





13 FEB 1997

21

Urease vaccine against *H. pylori*, Michetti et al.

To evaluate protection, biopsies were screened for the urease activity by the Jetrox HP test (Rohm Pharma) according to the instructions of the supplier. The urease is quantified by a spectrophotometric measurement at 550nm. The biopsies were also cultured in the presence of *H. felis* and was estimated by Gram staining. Gastric antral biopsies were homogenized and diluted (1:10 and 1:1000) in 0.15 M NaCl and plated onto blood agar plates and incubated under microaerophilic conditions at 37°C for 4 to 10 days.

ELISA

Intestinal secretions and blood were analyzed by ELISA for the evaluation of antibody titer. The ELISA was carried out as follows. Polystyrene plates (96 wells) were coated with 1 μ g/well of purified urease at 37°C for 2 hrs. Non specific binding sites were blocked with 5% powdered milk in PBS 0.1% Tween at 37°C for 30 minutes. The plates were washed once with PBS 0.1% Tween. Blood samples were tested at dilution 1:1000 and intestinal secretions 1:1. 100 μ l of each sample were added to the antigen coated plates. After 2 hrs of incubation, plates were washed 3 times with PBS 0.1% Tween. Anti-mouse biotinylated whole antibody from goat and anti-mouse IgA, IgG and IgM biotinylated (Amersham) were added (100 μ l) at dilution 1:500 except for IgA (1:250) and incubated at 37°C for 1 hr. The plates were washed 3 times with PBS 0.1% Tween and 100 μ l of 1:1000 dilution of streptavidin Horseradish peroxidase in PBS 0.1% Tween were added and incubated at 37°C for 30 minutes. The plates were washed 3 times and 50 μ l of 1:50 dilution of o-phenyl-diamine in citrate buffer pH 5.0 with 1 μ l/ml of 30% H₂O₂ were added and incubated at room temperature for 20 minutes. The absorbance at 495 nm was measured in each well.

EXAMPLE 3 (see also section C)

Mice employed in the immunization studies were lightly anesthetized with ether prior to intragastric immunization. Then, 30 μ g recombinant *H. pylori* urease A or B subunit produced in *E. coli* bound hydroxyapatite and supplemented with cholera toxin was suspended in PBS and 200 μ l were delivered to the stomach of the respective mice using a polyethylene tubing attached to a hypodermic syringe. This procedure will be referred to as oral immunization.

Three oral immunization protocols were evaluated. These are described below.

Protocol C1 - Vaccination with recombinant urease A subunit

Female BALB/c six-week old mice (10) were orally immunized with 30 μ g of purified recombinant *H. pylori* urease A subunit and 1 mg of hydroxyapatite and 10 μ g of cholera toxin at day 0, 8, 14 and 21. Ten mice were challenged at day 32, 34 and 36 with 10⁸ *H. felis* from liquid culture.

Protocol C2 - Vaccination with recombinant urease B subunit

13 FEB 1997

22

Urease vaccine against *H. pylori*, Michétti et al

Female BALB/c six-week old mice (10) were orally immunized with 30 ug of recombinant *H. pylori* urease B subunit and 1 mg of hydroxyapatite and 10ug of cholera toxin at day 0, 8, 14 and 21. Ten mice were challenged at day 32, 34 and 36 with 10^8 *H. felis* from liquid culture.

Protocol C3 - Control

Female BALB/c six-week old mice were orally immunized with 1 mg of hydroxyapatite and 10ug of cholera toxin at day 0, 8, 14 and 21. The mice were challenged at day 32, 34 and 36 with 10^8 *H. felis*.

At day 42, or at day 106, mice were sacrificed and multiple biopsies from the stomach were taken.

Protection and evaluation

To evaluate protection, biopsies of the corpus and antrum of the stomach were screened for urease activity by the Jatrox HP test (Rohm Pharma) according to the instructions of the supplier. The urease is quantified by a spectrophotometric measurement at 550nm. The total of corpus and antrum OD values were added to obtain a final OD value for each mouse.

This Sheet Has Been Replaced



References

1. Blaser, M.J. "Gastric Campylobacter-like organisms, gastritis and peptic ulcer disease" Gastroenterology 1987, 93, 371-383.
2. Graham, D.Y. "Campylobacter *pylori* and peptic ulcer disease" Gastroenterology 1989, 96, 615-625.
3. Parsonnet, J. et al "Helicobacter *pylori* infection in intestinal and diffuse-type gastric adenocarcinomas" J. Natl. Cancer Inst. 1991, 93, 640-643.
4. Marshall, B.J. et al "Attempt to fulfill Koch's postulate for pyloric Campylobacter" Med. J. Aust. 1985, 142, 436-439.
5. Morris, A. et al "Ingestion of Campylobacter *pyloridis* causes gastritis and raised fasting gastric pH" Am. J. Gastroenterol. 1987, 82, 192-199.
6. Engstrand, L. et al "Inoculation of barrier-born pigs with Helicobacter *pylori*: a useful animal model for gastritis type B" Infect. Immun. 1990, 58, 1763-1768.
7. Fox, J.G. et al "Gastric colonization by campylobacter *pylori* subsp. mustelae in ferrets" Infect. Immun. 1988, 56, 2994-2996.

13 FEB 1997
- 23Urease vaccine against H. pylori, Michetti et al . . .

- Fox, J.G. et al "Helicobacter mustelae-associated gastritis in ferrets: an animal model of Helicobacter pylori gastritis in humans" Gastroenterology 1990, 99, 352-361.
1. Lee, A. et al "A small animal model of human Helicobacter pylori active chronic gastritis" Gastroenterology 1990, 99, 1315-1323.
 10. Fox, J.G. et al "Helicobacter felis gastritis in gnotobiotic rats: an animal model of Helicobacter pylori gastritis" Infect. Immun. 1991, 59, 785-791.
 11. Eaton, K.A. et al "Campylobacter pylori virulence factors in gnotobiotic piglets" Infect. Immun. 1989, 57, 1119-1125.
 12. Peterson, W.L. "Helicobacter pylori and peptic ulcer disease" N. Engl. J. Med. 1991, 324, 1043-1048.
 13. Czinn, S.J. and Nedrud, J.G. "Oral immunization against Helicobacter pylori" Infect. Immun. 1991, 59, 2359-2363.
 14. Brandtzaeg, P. "Role of H chain and secretory component in receptor-mediated glandular and hepatic transport of immunoglobulins in man" Scand. J. Immunol. 1985, 22, 111-146.
 15. Brandtzaeg, P. et al "Production and secretion of immunoglobulins in the gastrointestinal tract" Ann. Allergy 1987, 59, 21-39.
 16. Wyatt, J.I. et al "Local immune response to gastritic campylobacter in non-ulcer dyspepsia" J. Clin. Path. 1986, 39, 863-870.
 17. Lee, A. et al. "Pathogenicity of Helicobacter pylori: A Perspective" Infect. Immun. 1993, 61, 1601-1610.
 18. Pallen, M.J. and Clayton, C.L. "Vaccination against Helicobacter pylori urease, letter" Lancet, 1990, 336, 186.
 19. Evans, D.J. et al. "Urease-associated heat shock protein of Helicobacter pylori" Infect. Immun. 1992, 60, 2125-2127.
 20. Ferrero, R.L. and Lee, A. "The importance of urease in acid protection for the gastric-colonising bacteria Helicobacter pylori and Helicobacter felis sp. nov." Microb. Ecol. Health Dis. 1991, 4, 121-134.
 21. Chen, et al. "Immunization against gastric Helicobacter infection in a mouse/Helicobacter felis model, letter" Lancet, 1992, 339, 1120-1121.
 22. Czinn, S. et al. "Oral immunization protects germ-free mice against infection from Helicobacter felis". Proceedings of the DDW, American Gastroenterological Association. May 10-13, 1992, 1321, A-331.
 23. Guo, M. and Liu, P.V. "Serological specificities of ureases of Proteus species". J. Gen. Microbiol. 1965, 136, 1995-2000.

13 FEB 1997
24Urease vaccine against H. pylori, Michetti et al

24. Michetti, P. et al. " Specificity of mucosal IgA response in Balb/C mice following H. felis or H. pylori challenges. .Proceedings of the DDW, American Gastroenterological Association. May 10-13, 1992, 1001, A-251.
25. Davin, C. et al. "H. pylori urease elicits protection against H. felis infection in mice". Proceedings of the DDW, American Gastroenterological Association. May 16-19, 1993, 1213, A-304.
26. Pallen, M.J. and Clayton, C.L. "Vaccination against Helicobacter pylori urease". Lancet 1990, 336, 186-7.
27. Pimentel, J.L. and Cook, M.E. "Improved growth in the progeny of hens immunised with jackbean urease". Poultry Sci. 1988, 64, 434-439.
28. Labigne, A. "Sequences of nucleotides coding for a protein having an urease activity". EPO patent application # EP O 367 644 A1, 1989. International publication # WO 90/04030, 1990.
29. Clayton, C.L. et al. S. "Nucleotide sequence of two genes from Helicobacter pylori encoding for urease subunits". Nucleic Acid Res. 1990, 18, 362.
30. McGhee, J.R and Kyono, H. "New perspectives in vaccine development: mucosal immunity to infections". Infect Agents Dis. 1993, 2, 55-73.

This Sheet Has Been Replaced





THIS SHEET HAS BEEN
REPLACED

REPLACEMENT

13 FEB 199

Urease vaccine against H. pylori, Michetti et al

WE CLAIM:

1. A method of eliciting in a mammalian host a protective immune response to Helicobacter infection, said method comprising the step of:

administering to a mucosal surface of said mammal an immunogenically effective amount of a polyaminoacid preparation presenting a sufficient number of epitopes exhibited by a urease endogenous to said Helicobacter organism to elicit a protective immune response to infection by said organism.
2. A method according to claim 1, wherein said preparation comprises an intact urease purified from an organism.
3. A method according to claim 1, wherein said preparation comprises peptides homologous with enzymatically inactive portions of the aminoacid sequences of a urease.
4. A method according to claim 1, wherein said preparation comprises peptides non-homologous with the aminoacid sequences of a urease and displaying epitopes cross-reacting with a urease.
5. A method according to claim 1, wherein said preparation comprises H. pylori urease.
6. A method according to claim 1, wherein said preparation comprises at least subunits of a urease, with or without enzymatic activity.
7. A method according to claim 1, wherein said preparation comprises anti-idiotypic antibodies to a urease.
8. A method according to claim 1, wherein said preparation comprises peptides immunologically cross-reacting with urease.
9. A method according to claim 3 and 9, wherein said peptides are obtained by chemical synthesis.
10. A method according to claim 1, wherein said preparation comprises urease antigens produced using DNA recombinant techniques.
11. A method according to claim 1, wherein said preparation comprises subgenic fragments of urease produced with recombinant techniques.
12. A method according to claim 1, wherein said preparation comprises urease subgenic fragments produced as genetically fused proteins.
13. A method according to claim 13, wherein said fused proteins comprise cholera toxin subunits.
14. A method according to claim 1, wherein said preparation is administered in association with a mucosal adjuvant.
15. A method according to claim 15, wherein said mucosal adjuvant is cholera toxin.

This Sheet Has Been Replaced





THIS SHEET HAS BEEN
REPLACED

REPLACEMENT SHEET

13 FEB 1997
26

Urease vaccine against H. pylori, Michetti et al.

16. A method according to claim 1, wherein said mammalian host is human.
17. A method according to claim 1, wherein said preparation is administered in association with a hydroxylated calcium phosphate.
18. A method according to claim 18, wherein said hydroxylated calcium phosphate is hydroxyapatite.
19. A method according to claim 19, wherein said hydroxyapatite is in the form of particles suitable for transport across epithelium.
20. A method according to claim 1, wherein said urease is administered orally, nasally, rectally or ocularly.
21. A vaccine for inducing a protective immune response to Helicobacter infection in a mammal, the vaccine comprising a polyaminoacid preparation presenting epitopes exhibited by a urease endogenous to said Helicobacter organism disposed in a pharmaceutically acceptable carrier or diluent.
22. The vaccine of claim 22 further comprising a mucosal adjuvant.
23. A method of imparting to a mammal passive protection to Helicobacter infection, the method comprising administering to a mucosal surface of said mammal an immunologically effective amount of a urease-specific IgA antibody produced in a host by immunization with a urease which elicits a protective immune response to Helicobacter.
24. A method according to claim 24, wherein said antibody is a Helicobacter pylori urease specific IgA antibody.
25. A method according to claim 24, wherein said mammal is human.
26. A method for assessing the endogenous immune response of a mammal infected by Helicobacter organism, the method comprising determining in a sample from the gastrointestinal tract of said mammal the presence of antibody reactive with epitopes exhibited by urease endogenous to said Helicobacter organism.
27. A method according to claim 27, comprising the additional step of administering a Helicobacter vaccine to said mammal prior to said determination.

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

REPLACEMENT SHEET

13 FEB 1997


27

Urease vaccine against H. pylori, Michetti et al. . . .

ABSTRACT OF THE DISCLOSURE

Method of eliciting in a mammalian host a protective immune response to Helicobacter infection by administering to the host an immunogenically effective amount of a Helicobacter urease or urease subunits as antigen. Vaccine compositions are also provided.

This Sheet Has Been Replaced



13 FEB 1997
28

Urease vaccine against *H. pylori*, Michetti et al.

FIGURE 1

GROUP A: MICE NOT PROTECTED AFTER IMMUNIZATION WITH
UREASE





FIGURE 2

GROUP B: MICE PROTECTED AFTER IMMUNIZATION WITH
UREASE



13 FEB 1997

Urease vaccine against *H. pylori*, Michetti et al

FIGURE 3

GROUP C: MICE NOT PROTECTED AFTER IMMUNIZATION WITH H.p.
SONICATE






FIGURE 4

GROUP D: MICE PROTECTED AFTER IMMUNIZATION WITH
H.p SONICATE




13 FEB 1997

39

Urease vaccine against *H. pylori*, Michetti et al

Figure 5




immunization :	<i>H. pylori</i> sonicate	<i>H. pylori</i> urease	control
----------------	------------------------------	----------------------------	---------

protected :	3/9	7/10	1/10
-------------	-----	------	------

* Two-tailed Fisher Exact Test

This Sheet Has Been Replaced







13 FEB 1997

- 31

Urease vaccine against *H. pylori*, Michetti et al

Figure 6





THIS SHEET HAS BEEN
REPLACED

9604759-2
1805 copy

REPLACEMENT SHEET

17 DEC 1997

25

1. Use of a polypeptide presenting a sufficient number of epitopes of a *Helicobacter* antigen in the preparation of a mucosal vaccine for inducing an immune response to *Helicobacter* in a mammal.
2. The use of claim 1, wherein said mammal is at risk of developing, but does not have, *Helicobacter* infection.
3. The use of claim 1, wherein said mammal is infected by *Helicobacter*.
4. The use of claim 1, wherein said polypeptide comprises a *Helicobacter* urease.
5. The use of claim 1, wherein said *Helicobacter* antigen is a *Helicobacter pylori* antigen.
6. The use of claim 1, wherein said *Helicobacter* antigen is a *Helicobacter* urease.
7. The use of claim 6, wherein said polypeptide comprises a subunit of said *Helicobacter* urease.
8. The use of claim 7, wherein said subunit is the urease A subunit.
9. The use of claim 7, wherein said subunit is the urease B subunit.

This Sheet Has Been Replaced



REPLACED SHEETS SG

- ☐ Description : paged to paged
- ☒ Claim(s) : paged 25... to paged 28...
- ☐ Drawing(s) : paged to paged
- ☐ Abstract : paged to paged

DATE OF RECEIPT

REPLACEMENT SHEETS SG

- ☐ Description : paged to paged
- ☒ Claim(s) : paged 25... to paged 28...
- ☐ Drawing(s) : paged to paged
- ☐ Abstract : paged to paged

DATE OF RECEIPT 17/12/97.....



THIS SHEET HAS BEEN
REPLACED

REPLACEMENT SHEET

17 DEC 1997

26

10. The use of claim 1, wherein said polypeptide comprises an anti-idiotypic antibody to a *Helicobacter urease*.

11. The use of claim 1, wherein said polypeptide is recombinant.

12. The use of claim 1, wherein said polypeptide is produced as a fusion protein.

13. The use of claim 12, wherein said fusion protein comprises a cholera toxin subunit.

14. The use of claim 1, wherein said mucosal vaccine is administered to said mammal with a mucosal adjuvant.

15. The use of claim 14, wherein said mucosal adjuvant comprises a cholera toxin.

16. The use of claim 1, wherein said mammal is human.

17. The use of claim 1, wherein said mucosal vaccine is administered to said mammal with a hydroxylated calcium phosphate.

18. The use of claim 17, wherein said hydroxylated calcium phosphate is hydroxyapatite.

This Sheet Has Been Replaced





THIS SHEET HAS BEEN
REPLACED

REPLACEMENT SHEET

17 DEC 1997

27

19. The use of claim 18, wherein said hydroxyapatite is in the form of particles suitable for transport across epithelium.

20. The use of claim 1, wherein said mucosal vaccine is administered to an oral, intranasal, rectal, or ocular surface of said mammal.

21. A vaccine for inducing a mucosal immune response to *Helicobacter* in a mammal, said vaccine comprising a polypeptide presenting a sufficient number of epitopes of a *Helicobacter* antigen to induce said immune response, in a pharmaceutically acceptable carrier or diluent.

22. The vaccine of claim 21, further comprising a mucosal adjuvant.


23. The vaccine of claim 21, wherein said polypeptide comprises a *Helicobacter* urease or a subunit of a *Helicobacter* urease.

24. The vaccine of claim 21, wherein said *Helicobacter* antigen is a *Helicobacter pylori* antigen.

25. The vaccine of claim 21, wherein said *Helicobacter* antigen is a *Helicobacter* urease.

26. Use of an IgA antibody specific for a *Helicobacter* antigen in the preparation of a medicament for imparting passive

This Sheet Has Been Replaced



THIS SHEET HAS
REPLACED

REPLACEMENT SHEET

17 DEC 1997



28

protection to *Helicobacter* infection in a mammal by mucosal administration.

27. The use of claim 26, wherein said *Helicobacter* antigen is a *Helicobacter pylori* antigen.

28. The use of claim 26, wherein said *Helicobacter* antigen is a *Helicobacter* urease.

29. The use of claim 26, wherein said mammal is human.

30. A method for assessing the immune response of a mammal infected by a *Helicobacter* organism, said method comprising detecting the presence of antibodies reactive with a polypeptide of a *Helicobacter* in a sample from the gastrointestinal tract of said mammal.

31. The method of claim 30, wherein said polypeptide of a *Helicobacter* is a *Helicobacter* urease.

32. The method of claim 30, further comprising administering a polypeptide of a *Helicobacter* to a mucosal surface of said mammal as a vaccine antigen prior to said detecting.

This Sheet Has Been Replaced



REPLACEMENT SHEET

17 MAR 1999

9604759-2
EPDS copy

9604759-2 EPDS

REPLACEMENT SHEETS SG
☒ Description : paged .1. to paged .31.
☒ Claim(s) : paged .12. to paged .42.
☒ Drawing(s) : paged .43. to paged .46.
☒ Abstract : paged .47. to paged .50.

DATE OF RECEIPT 11/3/99

1

UREASE-BASED VACCINE AGAINST HELICOBACTER INFECTION

The present invention relates to the prevention and treatment of gastric infection in mammals, including humans. More particularly, the present invention relates to a vaccine suitable for use in the prevention and treatment of Helicobacter infection in mammals, including humans, and to a method of treatment of humans suffering from gastric infection, its consequences such as chronic gastritis or peptic ulcer, and prevention of gastric cancer.

BACKGROUND OF THE INVENTION

Helicobacter infections of human gastric epithelium cause gastritis, are a major factor in the development of peptic ulcers and gastric lymphoma, and may be a risk factor for the development of gastric cancer [1-3]. The most frequent infection agent is Helicobacter pylori, followed at a much lower frequency by Helicobacter heilmanii. H. pylori is a slender S-shaped gram negative microorganism, which is routinely recovered from gastric biopsies of adults and children with histologic evidence of gastritis or peptic ulceration. Evidence for a causal relationship between H. pylori and gastroduodenal disease comes from studies in human volunteers, patients with ulcers and gastric cancer, gnotobiotic pigs, and germ-free rodents. Regarding etiology, Koch's postulates were satisfied by creating histologically confirmed gastritis in previously uninfected individuals following consumption of viable microorganisms [4-11], and by treatment to eradicate H. pylori, with resolution of the gastritis and, in patients with peptic ulcer disease, a decrease in the recurrence rate [12].

In spite of in vitro susceptibility to many antimicrobial agents, in vivo eradication of established H. pylori infections with antimicrobial agents is often difficult to achieve

17 MAR 1999

[13]. The microorganism is found within the mucous coat overlying the gastric epithelium and in gastric pits. These are locations which do not appear to allow for adequate antimicrobial levels to be achieved even when antibiotics are given orally at high doses. At the present time, most authorities recommend a "triple therapy", namely a bismuth salt in combination with drugs such as tetracycline and metronidazole for 2-4 weeks. However, the effectiveness of this or other chemotherapeutic regimens remains suboptimal. Furthermore, this treatment may produce serious adverse drug reactions.

At the present time little is known regarding the role of the mucosal immune systems in the stomach. The distribution of immunoglobulin (Ig) producing cells in the normal gastric antrum indicates that IgA plasma cells make up 80% of the total plasma cell population. In addition, the number of plasma IgA cells present in the gastric antrum is comparable to other mucous membranes [14, 15]. A number of studies in humans [16] and in animal models [8, 10] have demonstrated specific IgG and IgA responses in serum and in gastric secretions in response to Helicobacter infection. However, the observation that H. pylori infection persist as a chronic infection for years, despite inducing a local and systemic immune response, is not encouraging the development of immunization strategies.

Lee et al have reported the ability to infect germ-free rodents with Helicobacter felis, a bacterium closely related to H. pylori, and reproducibly document histologic gastritis [9, 10]. Since then, this bacterium-host pairing has been accepted as a good model to study Helicobacter-mediated gastritis and its initiating factors [17]. Czinn et al have shown that repetitive oral immunization with a crude lysate of H. pylori plus cholera toxin adjuvant induces a vigorous gastrointestinal IgA anti-H. pylori response in mice and ferrets [13]. In addition, Chen et al and Czinn et al have recently reported that oral

17 MAR 1999

immunization with a crude lysate of H. felis induced protection against H. felis infection in mice [21, 22]. The exact nature of the antigen(s) responsible for the induction of this protection, however, had not been determined, and no information suggested that the protective antigen(s) of H. felis that induced protection against this pathogen would induce a cross-reactive protection extending to another Helicobacter species.

We have demonstrated for the first time that H. pylori and H. felis share antigenic determinants by obtaining monoclonal antibodies after oral immunization of mice with either H. pylori or H. felis sonicates and showing that some of these antibodies, directed against H. pylori, would crossreact with H. felis and vice versa [24; 25]. The basis for these cross-reactivities were unknown.

Based on the homology existing between the different known urease amino acid sequences, it has been proposed that urease could be used as a vaccine against H. pylori [26]. Nevertheless, despite the homology among the different ureases sequences, cross-reactivity is not the rule. Guo and Liu have shown years ago that ureases of Proteus mirabilis, Proteus vulgaris and Providencia rettgeri show cross-reactivity to each other, while ureases of jack bean and Morganella morganii are immunologically distinct from the three former ureases [23]. Even if an antigenic cross-reactivity of H. pylori urease with other Helicobacter ureases was a reasonable postulate, no data existed demonstrating that this was really the case until we showed that some H. felis monoclonal antibodies crossreacted with H. pylori urease [25]. J. Pappo has further demonstrated that mice which have been infected by H. felis produce antibodies which crossreact with H. pylori urease but not jack bean urease (J. Pappo, unpublished data, 1993).

17 MAR 1999

The use of H. pylori urease, or of related ureases, as a vaccine against H. pylori infection has previously been proposed by A. Labigne in EPO 367,644 [28]. However, that application contains no evidence of vaccination of any mammal against any Helicobacter infection with urease.

Moreover, while sequence homology with other bacterial ureases might support the use of urease as a vaccine candidate against H. pylori infection, the current knowledge of human H. pylori infection would certainly not. First, despite the fact that infected individuals often mount a strong antibody response to urease, the anti-urease immune response does not result in clearance or control of the infection. Secondly, H. pylori is able to transport urease out of the cell and to shed it from its surface [19, 20]. Thus, urease may not represent an appropriate target for the development of a protective mucosal immune response. Indeed, mucosal immune protection is thought to be mainly mediated by secretory IgA, the agglutinating activity of which would be impaired when the recognized antigen can be shed by the target pathogen and thereby serve as a decoy for the protective antibody. Thirdly, urease appears to be toxic for epithelial cells in culture, and has been suspected to play a role in mucous degradation and in peptic ulceration in vivo. Thus, its use as antigen may be toxic.

Nevertheless, we reasoned that this antigen could be a potentially efficient vaccine if:

- first, we would deliver it orally at a sufficiently high dose to elicit a stronger immune response than the naturally occurring one
- second, the amount of antibodies produced would be high enough to bind all the urease, shed or not shed

17 MAR 1999

- third, we would use subunits of urease or a molecular species that was non toxic.

In summary, there remains a need for effective treatment and prevention of H. pylori-induced gastric infection in humans. Recent data suggested the possibility to generate a vaccine against this infection, but have not provided a clear identification of defined antigen(s), common to all strains of H. pylori, that could be incorporated into a safe and effective vaccine.

In this invention, we have identified the urease antigen of H. pylori as a candidate vaccine and demonstrated its efficacy in an animal model. These results were unexpected in the light of the natural history Helicobacter infections.

SUMMARY OF THE INVENTION

We have discovered that immunity can be induced in mammals susceptible to gastrointestinal Helicobacter infection by exploiting urease epitopes displayed on or about the surface of Helicobacter organisms and using them as a vaccine target. The immunity can be induced by immunization with native urease, but can also be induced with recombinant urease subunit, produced as an enzymatically inactive, therefore non-toxic form. The invention provides a method of inducing immunity to Helicobacter infection by administering to a mucosal surface of a mammal a polyaminoacid preparation, i.e. a mixture of peptides and/or proteins, together with an appropriate adjuvant. This polyaminoacid preparation presents a plurality of epitopes characteristic of and exhibited by a urease enzyme endogenous to the infecting Helicobacter organism. The administration of the polyaminoacid preparation may be performed by the oral route.

17 MAR 1999

The active ingredient of the preparation may comprise natural or biosynthetic epitopes and may take various forms. A non exhaustive list of possible preparations includes purified, naturally occurring or recombinantly produced urease preparations of bacterial or other origin, digests of urease, fusion proteins comprising urease epitopes, truncated forms of urease enzyme, or peptides homologous with the aminoacid sequence of urease. Since development of immunity depends on induction of humoral and/or cellular immune responses which bind to the infecting Helicobacter organism, preferred preparations are those which most closely duplicate the epitopes of the urease endogenous to the infecting organism. For example, preparations displaying the epitopes of urease of H. pylori are preferred for administration in humans susceptible to H. pylori, and preparations displaying the epitopes of urease of H. heilmannii are preferred for the administration in humans susceptible to H. heilmanni. However, in accordance with an important aspect of the invention, it has been discovered that urease from other species may be used. For example, we have shown that H. felis infection in mice can be prevented by administration of urease from H. pylori.

According to one aspect of the invention, there is provided a method of eliciting in a mammalian host a protective immune response to Helicobacter infection wherein an immunologically effective amount of a urease antigen capable of eliciting such a protective immune response, preferably H. pylori urease or H. pylori urease B subunit, is administered to a mucosal surface of the host.

According to another aspect of the present invention, there is provided a vaccine composition suitable for prevention of Helicobacter infection, comprising an effective amount of a urease antigen, preferably H. pylori urease or H. pylori urease B subunit, capable of eliciting in a host a protective immune response to Helicobacter infection, in association with a pharmaceutically acceptable carrier or diluent.

17 MAR 1999

According to a further aspect of the present invention, there is provided a method of imparting to a mammalian host passive protection to Helicobacter infection, comprising administering to a mucosal surface of the host an immunologically effective amount of a urease specific antibody produced in a host immunized with a urease, preferably H. pylori urease or H. pylori urease B subunit, capable of eliciting a protective immune response to Helicobacter infection.

BRIEF DESCRIPTION OF THE DRAWINGS

The invention will now be further described with reference to the accompanying drawings, in which Figures 1 through 6 are graphical representations of the results set forth in Tables 1 through 6.

DETAILED DESCRIPTION OF THE INVENTION

The present inventors have discovered that oral administration to mice of polyaminoacid preparations exhibiting the epitopes of H. pylori urease gives rise to a protective immunological response against H. felis in mice, an animal model of generally-accepted value for the study of the immune response to Helicobacter infection [9]. The effect of the protective immune response is that immunized animals, when challenged with pathogen, have a greatly reduced incidence of infection, in comparison to non-immunized animals. Furthermore, the inventors have discovered that oral immunization in mice using H. pylori urease B subunit, produced as an enzymatically-inactive recombinant protein, gives rises to a protective immunological response in mice against H. felis. The effect of the protective immune response is that immunized animals, when challenged with pathogen, have also a greatly reduced

17 MAR 1999

incidence of infection, in comparison to non-immunized animals which do become infected.

Thus, in a first aspect, the present invention provides a method of eliciting in a mammalian host a protective immune response to Helicobacter infection. The method comprises the step of administering to a mucosal surface of the mammal, including humans, an immunologically effective amount of a urease antigen, preferably H. pylori urease, capable of eliciting such a protective immune response.

In a second aspect, the present invention provides a method of eliciting in a mammalian host a protective immune response to Helicobacter infection. The method comprises the step of administering to a mucosal surface of the mammal, including humans, an immunologically effective amount of recombinant, enzymatically inactive, urease B subunit as antigen, preferably recombinant H. pylori urease B subunit, capable of eliciting such a protective immune response.

The invention also includes within its scope the treatment or prophylaxis of mammals, including humans, for Helicobacter infection, wherein an immunologically effective amount of a urease, or its subunits, capable of eliciting a protective immune response to Helicobacter infection, is administered to a mucosal surface of a patient. Preferably, the urease is H. pylori urease or H. pylori urease B subunit, and the urease may be administered either alone or linked to a hydroxylated calcium phosphate, for example hydroxyapatite as a carrier particle. Moreover, it is preferred to administer the H. pylori urease in association with a mucosal adjuvant, the B subunit of cholera toxin, muramyl dipeptide or other such adjuvants.

While not being bound by any theory, the present inventors believe that administration

17 MAR 1999

of the urease antigen, or B subunit thereof, to a mucosal surface stimulates the common mucosal immune system and perhaps local sites in the gastric mucosa inducing an immune response, including the appearance of H. pylori specific IgA antibodies in the gastric secretions, which prevent Helicobacter infection. Since it is a routine matter to conduct pre-clinical trials of candidate vaccines for human use in animal models, it is believed that the methodology of the present invention is effective in humans, especially in the prevention and treatment of peptic ulcers, gastritis, gastric malignancies and other conditions arising as a result of the presence of H. pylori and/or H. heilmanii.

A - Bacterial cultures and urease purification

The strain of H. pylori used in the study originates from a patient with a duodenal ulcer, and has been subcultured on BHI agarose plates to homogeneity. H. pylori is cultured in a suitable medium, typically, BHI (Brain-Heart Infusion) medium, containing 0.25% yeast extract and 10% fetal calf serum and supplemented with 0.4% Campylobacter selective complement (Skirrow supplement; Oxoid 69). The bacteria are incubated overnight under microaerophilic conditions at 37°C in bottles that are then sealed and shaken at 37°C for 2 to 3 days to produce a liquid culture. A culture may also be prepared in agarose plates consisting of BHI with 0.25% of yeast extract and 5% of sheep blood under microaerophilic conditions at 37°C for 3 days. The quantity of bacteria is determined by optical density of the BHI solution at 660 nm, with one optical density unit corresponding to 10^8 bacteria. Cultures on agarose plates are first resuspended in 154mM NaCl.

One currently preferred source of polyaminoacid displaying urease epitopes is purified urease, e.g., H. pylori urease obtained by following the general method of Dunn et al.

17 MAR 1999

J. Biol. Chem. 265, 9464-9469, modified as described below. Following culturing, the H. pylori is harvested in water, spun vortexed and spun again to produce a supernatant. The solution containing the urease activity of H. pylori (assessed by rapid urease test, see below) is then chromatographed on a CL-6B sizing column and the fractions which present a strong urease activity are pooled and dialysed overnight and again chromatographed on an anion exchanger gel. The fractions are eluted in increasing NaCl buffer and the collected fractions with a strong urease activity are individually submitted to a SDS gel followed by Coomassie staining. Two distinct bands corresponding to a molecular weight of about 63 and about 28 kDa are identified as urease. The fractions containing urease are pooled to give purified H. pylori urease having a purity in the region of 95% to 99%.

B - Oral immunization with urease purified from H. pylori

While it is preferred to employ purified H. pylori urease obtained as described above as the antigenic material, it will be understood that it is also possible to use, as the antigenic material, any urease or subunit of urease, either naturally occurring or obtained by recombinant DNA techniques, as well as digested fragment thereof, fusion proteins comprising the fragments or the whole urease, truncated urease constructs, or other peptide or protein preparations exhibiting urease epitopes which are capable of eliciting a protective immune response to Helicobacter infection (See below). Thus, it is possible to employ a urease having a substantial homology with respect to H. pylori urease and which is effective in raising a cross-protective immune response to Helicobacter. An example of such a urease is jack bean urease, which possesses about 70% homology with H. pylori urease. The invention is therefore not limited to the use of intact urease, and covers the use of any polyaminoacid preparation which displays urease epitopes and is effective to generate a protective immunological response in a

17 MAR 1999

host to Helicobacter infection. Typically, a urease having a homology of 70-95% homology, for example, 80-90% homology, with respect to H. pylori urease, may be employed as the urease antigen in the invention.

A non-limiting list of sources of potentially useful urease preparations includes endogenous urease enzymes of the different Helicobacter species, urease from other bacteria such as Klebsiella pneumoniae or Proteus mirabilis, and, by analogy, any other urease with the condition that these ureases share cross-reactive epitopes with H. pylori urease. The urease genes of all the organisms mentioned above represent a potential tool for expressing recombinant urease products as a whole protein or as a part thereof.

A non-limiting list of potentially useful urease preparations includes peptides generated from purified urease (the sources are mentioned above), using physical and/or chemical cleavage procedures (i.e. CnBr) and/or proteolytic cleavage (using proteases e.g. V8-protease, trypsin or others); or peptides synthesised chemically and sharing crossreactive epitopes with urease.

Other sources of potentially useful epitopes include epitopes identified by their crossreactivity with urease, as a result of screening with anti-urease antibodies. These peptides can be naturally occurring peptides or peptides resulting from chemical synthesis. Furthermore such peptides can result from the expression of recombinant random oligonucleotides.

Another source of potentially useful epitopes includes epitopes similar to urease as a result of the generation of anti-idiotypic antibodies to urease. Such anti-idiotypic antibodies, generated in any immunocompetent host, are obtained by immunization of

17 MAR 1999

this host with anti-urease antibodies, with the goal of generating antibodies directed against anti-urease antibodies, which share structural homologies with urease.

The discussion herein focuses on the use of urease naturally produced by H. pylori (section B). However, it will be appreciated that the urease or subunits or constructs thereof mentioned above, capable of eliciting the desired protective immune response, may be produced by recombinant DNA techniques well known in the art. The efficacy of particular preparations may be determined by routine administration using animal models, oral administration of the candidate vaccine, and challenge with pathogen using a protocol substantially similar or identical to the procedure described below.

Tables 1 and 2 below and Figures 1-5 describe the results obtained when mice were orally immunized with purified H. pylori urease. In this first experiment, administration of the H. pylori antigen was carried out by orally administering to the mice H. pylori urease purified as described in section A, and coupled to hydroxyapatite crystals, used as a carrier to enhance M cell binding and uptake. Cholera toxin (Sigma) was given as a mucosal adjuvant. In this experiment, groups of female SPF BALB/c six-week old mice were each orally immunized with 30 ug of purified H. pylori urease coupled to 1 mg of hydroxyapatite plus 10 ug of cholera toxin adjuvant at day 0, 7, 14 and 21. The mice were then challenged twice with 10^8 H. felis, at day 28 and 30. For comparison purposes, similar female SPF BALB/c six-week old mice were orally immunized with whole H. pylori lysate (sonicate) and 10 ug cholera toxin at day 0, 7, 14 and 21. The mice were challenged at day 28 and 30 with H. felis. The H. pylori sonicate was prepared by collecting H. pylori from cell cultures, pelleting by centrifugation and resuspending the pellet in 0.9% sodium chloride followed by sonication.

17 MAR 1999

As a control, female SPF BALB/c six-week old mice were orally sham-immunized with 10 µg of cholera toxin and 1 mg of hydroxyapatite at day 0, 7, 14 and 21. The mice were then challenged at day 28 and 30 with H. felis. All mice were housed, immunized, and challenged in parallel. All mice subject to the study were sacrificed on day 35.

C - Oral immunization with recombinant urease subunits of H. pylori

Genes encoding the structural A and B subunits of H. pylori urease were obtained by polymerase chain reaction (PCR) cloning according to standard procedures, based on previously published sequences [29]. These genes were inserted in a vector (named pEV40) designed for high expression and easy purification of foreign genes in E. coli. Briefly, the foreign gene is inserted downstream of a thermo-repressible promoter, and in frame of a sequence encoding for a repeat of six histidines. An ampR gene is present on this vector for selection of transformants. Under the appropriate temperature conditions, the recombinant protein obtained is supplemented by six histidines at the N-terminal, which allow for a one-step affinity purification on a nickel column. Both H. pylori recombinant urease A and B subunits were expressed separately in E. coli, and purified on nickel column to >95% purity.

While it is preferred to employ recombinant H. pylori urease obtained as described above as the antigenic material, it will be understood that it is also possible to use, as the antigenic material, any urease or subunit of urease obtained by recombinant techniques (e.g. fusion protein) expressing antigenic sites of urease, which is capable of eliciting a protective immune response to Helicobacter infection. Thus, it is possible to employ in a construct a urease gene having a substantial homology with respect to H. pylori urease and which is effective in raising a cross-protective immune response to

17 MAR 1999

Helicobacter. Examples of such a urease is jack bean urease, which possesses about 70% homology with H. pylori urease, or H. felis urease, which possesses about 88% homology with H. pylori urease. The invention is therefore not limited to the use of H. pylori urease genes and their gene products, and covers the use of any recombinant urease, or the subunits thereof, which is effective to generate a protective immunological response in a host to Helicobacter infection. Typically, a recombinant urease having a homology of 70-95% homology, for example, 80-90% homology, with respect to H. pylori urease, may be employed as the recombinant urease antigen in the invention.

The discussion herein focuses on the use of recombinant H. pylori urease A and B subunits produced by E. coli (section C). However, it will be appreciated that recombinant urease or subunits or constructs thereof mentioned above, capable of eliciting the desired protective immune response, may be produced using other recombinant DNA techniques and other eukaryotic or prokaryotic expression vectors well known in the art.

Tables 3, 4 and 5 below and Figure 6 describe the results obtained when mice were orally immunized with recombinant H. pylori urease subunits produced in E. coli. In this experiment, administration of the H. pylori antigen was carried out by orally administering to the mice recombinant H. pylori urease A or B subunits produced in E. coli and purified as described above, and coupled to hydroxyapatite crystals, used as a carrier to enhance M cell binding and uptake. Cholera toxin (Sigma) was given as a mucosal adjuvant. In this experiment, groups of female SPF BALB/c six-week old mice were each orally immunized with 30 ug of recombinant H. pylori urease A subunit, coupled to 1 mg of hydroxyapatite plus 10 ug of cholera toxin adjuvant at day 0, 8, 14 and 21. The mice were then challenged twice with 10^8 H. felis, at day 32, 34

17 MAR 1999

and 36. For comparison purposes, similar female SPF BALB/c six-week old mice were orally immunized with 30 ug of recombinant H. pylori urease B subunit coupled to hydroxyapatite plus 10 ug cholera toxin at day 0, 8, 14 and 21. The mice were challenged three times, at day 32, 34 and 36, with H. felis. As a control, female SPF BALB/c six-week old mice were each orally sham-immunized with 10 ug of cholera toxin and 1 mg of hydroxyapatite at day 0, 8, 14 and 21. The mice were then challenged at day 32, 34 and 36 with H. felis. All mice subject to the study were immunized and challenged in parallel. Animals were sacrificed on day 48 (12 days after challenge) or 10 weeks after challenge.

D - Analysis of gastric biopsies, blood, and intestinal secretions

Biopsies were taken from the stomach and blood was obtained from the heart. The intestines were removed and washed with 1mM PMSF (Boehringer) in PBS buffer to obtain intestinal secretions for ELISA analysis.

To evaluate protection against H. felis colonization, gastric biopsies from each animal were screened for the presence of H. felis by assessing rapid urease activity by the Jatrox HP test (Rohm Pharma), according to the supplier's directions. Briefly, gastric biopsies are immersed in 0.5 ml supplier's mixture of urea and phenol red, a pH indicator. Urease activity generates ammonia and bicarbonate from urea, and is followed by the colorimetric change of the solution towards a higher absorbance at 550 nm. Urease activity was quantified by spectrophotometric analysis.

Gastric biopsies of each animal included in the experiment described in section B were also cultured on BHI agarose plates, supplemented as above, for the detection of H. felis. After incubation for 3 to 10 days in microaerophilic conditions, the presence of

17 MAR 1999

H. felis was confirmed by Gram staining and determination of urease activity. As a very significant correlation was obtained for the detection of H. felis in gastric biopsies between urease tests and H. felis cultures during the first set of experiments (see Table 3), only gastric biopsies urease tests were performed for the detection of H. felis in the experiment described in section C. Detection of H. felis was confirmed by microscopy by two independent investigators, using two different stains (acridine orange and cresyl violet).

Blood samples were allowed to clot for 3 hours at RT, and sera harvested and frozen at -20°C until analyzed. Intestinal secretions were spun for 5 min. at 4°C to remove debris, and kept frozen at -20°C. Serum and intestinal samples of each animal were analyzed by ELISA for evaluation of anti-Helicobacter activity, according to standard procedures. Briefly, 96-well plates were coated with a sonicate of H. pylori, followed by saturation with 5% fat-free milk. Samples were serially diluted from 1:1 to 1:1000 and incubated overnight at 4°C on ELISA plates. Biotinylated anti-mouse IgG (serum) and anti-mouse IgA, followed by streptavidin-Horseradish peroxidase was used for the determination of the antibody levels.

The results of H. felis challenges following immunizations with purified H. pylori urease are set out in Tables 1-3 and Figures 1-4 and the results of H. felis challenges following immunizations with recombinant H. pylori urease A and B subunits are set out in Tables 4-6 and Figures 5 and 6.

17 MAR 1999

TABLE 1

mouse		urease	Culture	Immunoglobulins			
number	Immunization	test	Gram	Serum		Intestinal secretion	
		12th		Ig	Ig G	Ig	IgA
1	Urease + HF	+	<i>H felis</i>	27	0	25	258
2	Urease + HF	0	0	264	273	221	246
3	Urease + HF	0	0	84	44	318	354
4	Urease + HF	+	<i>H felis</i>	81	42	12	5
5	Urease + HF	0	0	98	137	126	234
6	Urease + HF	+	0	968	2093	31	22
7	Urease + HF	0	0	98	0	96	34
8	Urease + HF	0	0	247	1010	214	128
9	Urease + HF	0	0	N.D.	N.D.	48	23
10	Urease + HF	0	0	50	0	124	99
11	Urease	0	0	319	205	44	53
12	Urease	0	0	14	0	86	87
13	Urease	0	0	0	0	0	0
14	Urease	0	0	0	0	43	61
15	Urease	0	0	58	0	110	127
16	Urease	0	0	140	63	21	37
17	Urease	0	0	84	240	114	280
18	Urease	0	0	N.D.	N.D.	93	148
19	Urease	0	0	45	0	135	216

17 MAR 1999

20	Urease	0	0	261	197	161	261
21	CT+HF	0	0	0	0	0	2
22	CT+HF	+	<i>H felis</i>	63	0	310	303
23	CT+HF	+	<i>H felis</i>	90	0	N.D.	N.D.
24	CT+HF	+	<i>H felis</i>	31	0	150	192
25	CT+HF	+	<i>H felis</i>	197	250	250	440
26	CT+HF	+	<i>H felis</i>	105	135	214	138
27	CT+HF	+	<i>H felis</i>	140	47	109	55
28	CT+HF	+	0	0	0	16	15
29	CT+HF	+	<i>H felis</i>	0	0	0	0
30	CT+HF	+	<i>H felis</i>	N.D.	N.D.	N.D.	N.D.
31	HP sonicate+HF	+	<i>H felis</i>	0	0	76	103
32	HP sonicate+HF	+	<i>H felis</i>	77	0	11	33
33	HP sonicate+HF	+	<i>H felis</i>	549	748	57	36
34	HP sonicate+HF	0	0	660	153	180	286
35	HP sonicate+HF	+	<i>H felis</i>	730	192	0	5
36	HP sonicate+HF	+	<i>H felis</i>	32	0	5	64
37	HP sonicate+HF	0	0	400	400	312	1149
38	HP sonicate+HF	+	<i>H felis</i>	1007	1360	149	26
39	HP sonicate+HF	0	0	220	186	133	122
40	HP sonicate	0	0	873	1016	352	514
41	HP sonicate	0	0	727	899	126	191
42	HP sonicate	0	0	109	68	44	83

17 MAR 1999

43	HP sonicate	0	0	147	949	167	97
44	HP sonicate	0	0	845	1094	246	64
45	HP sonicate	0	0	1217	1198	210	157
46	HP sonicate	0	0	81	0	256	218
47	HP sonicate	0	0	329	210	241	276
48	HP sonicate	0	0	1049	737	197	211

In Table 1, which refers to the experiment described in section B, "h" means hour, "Ig" means immunoglobulin, "ND" means "not determined", "urease + HF" means that the mice were immunized with urease (coupled to hydroxyapatite, with cholera toxin) and then challenged with H. felis, "urease" means that the mice were immunized with urease (coupled to hydroxyapatite, with cholera toxin) and not challenged, "CT+HF" means that the mice were sham-immunized with cholera toxin and challenged H. felis, "HP sonicate +HF" means that the mice were immunized with H. pylori sonicate with cholera toxin and challenged by H. felis, and "HP sonicate" means that the mice were immunized with H. pylori sonicate with cholera toxin and not challenged. In Table 1, the numbers for the antibody results are given as a measure of absorbance at 595 nm multiplied by 1000. The background measured in absence of the antibodies, was subtracted.

The results of experiment described in section B obtained on the basis of the gastric biopsies urease tests and on Gram staining of H. felis cultures are set out in Table 2. Infection was defined by mice with one or more markers of colonization by H. felis, including urease test or Gram staining of cultures.

17 MAR 1999

TABLE 2

<u>Immunization</u>	<u>Challenge</u>	<u>% infected</u>	<u>% protected</u>
Urease	<u>H. felis</u>	3/10 (30%)	7/10 (70%) *
Sonicate	<u>H. felis</u>	6/9 (66%)	3/9 (33%) **
CT	<u>H. felis</u>	9/10 (90%)	1/10 (10%)

*p=0.0198 (two tailed Fisher exact test) compared to CT control

** p=0.303 (two tailed Fisher exact test) compared to CT control

It will be seen from the results set out in Tables 1 and 2 that statistically significant protection against H. felis challenge is obtained with oral immunization using H. pylori urease as compared to that obtained using either H. pylori sonicate or cholera toxin. Referring to Table 2, it will be seen that from a total of 10 immunized animals, only 3 became infected, as compared to 6 of the animals immunized with H. pylori sonicate and 9 of the animals immunized with cholera toxin. Table 2 shows that 70% of the animals were protected from challenge by H. felis as compared to 33% of the animals immunized with H. pylori sonicate and 10% of the animals immunized with cholera toxin and then subjected to H. felis challenge. In other words, 90% of the control mice exposed to H. felis became infected by that pathogen whereas, in contrast, in mice immunized with H. pylori urease 28 days before exposure to H. felis, the infection rate was only 30%. This represents a significant reduction in infection (p=0.0198 in the Fisher exact test, as compared to the control mice). When the mice were orally immunized with H. pylori sonicate, the infection rate was 67% (not significant versus the control). The protection obtained using H. pylori urease is unexpected and could not have been predicted on the basis of results observed using H. pylori sonicate.

17 MAR 1999

Referring to Figures 1-4, Figure 1 represents graphically the results of tests for antibodies in serum (IgG) and intestinal secretion (IgA) in mice not protected after immunization with urease. These are mice numbers 1, 4 and 6 appearing in Table 1, and constitute Group A. Figure 2 shows the antibody responses of mice that were protected after immunization with urease (Group B), i.e. mice 2, 3, 5 and 7-10.

Figures 3 and 4 relate to the results obtained with mice numbers 31-39. Figure 3 (Group C) depicts antibody responses of mice not protected after immunization with H. pylori sonicate (mice numbers 31, 32, 33, 35, 36 and 38) and Figure 4 (Group D) depicts the antibody responses of mice protected after immunization with H. pylori sonicate (mice numbers 34, 37 and 39). It is of interest to note with respect to Figures 3 and 4 that the IgA antibody responses (but not IgG) are higher in the mice exhibiting protection than in the mice that are not protected, suggesting a correlation between protection and IgA response. Serum IgG responses did not exhibit a correlation. Mucosal IgA but not serum IgG antibodies are known to play a role in protection against bacterial infections of the gut [30].

The results of the correlation between the detection of H. felis in gastric biopsies by urease tests and by cultures are set out in Table 3.

TABLE 3

	Urease Test +	Urease Test-	Total
<u>H. felis</u> culture +	16	0	16
<u>H. felis</u> culture -	2	30	32
Total	18	30	48

Two-tailed Fisher's Exact Test: $p > 0.00001$

17 MAR 1999

Table 3 shows that a very significant correlation exists between the results of urease tests performed on gastric biopsies and the identification of H. felis by cultures. As cultures detected less often H. felis infection than urease tests, urease tests were preferred for the diagnosis of H. felis infection in mice in the next experiments, due to its better sensitivity. This approach allowed the duplication of urease tests with larger fragments of the stomach of each mouse, and a further increase in the sensitivity of the urease test. Furthermore, the use of the method with the highest sensitivity prevented an overestimation of the protection obtained by the vaccine preparation to be tested. When positive culture is used as the standard for infection, the protection induced after urease immunization during the experiment depicted in section B is as significant as with the combined use of urease test and culture ($p=0.021$ versus $p=0.019$).

The results of the experiments described in section C (recombinant urease subunits), obtained on the basis of the gastric biopsies urease test, are set out in Table 4, 5 and 6 and depicted in Figure 6.

TABLE 4

immunization	mice n°	Urease Test	infection
CT	20	0.49	+
	21	0.31	+
	22	0.62	+
	23	0.67	+
	24	0.55	+
	50	0.50	+
	51	0.37	+
	52	0.29	+

17 MAR 1999

	53	0.79	+
	54	0.32	+
ureA + HAP + CT	40	0.67	+
	41	0.48	+
	42	0.42	+
	43	0.65	+
	44	0.56	+
	45	0.52	+
	46	0.33	+
	47	0.63	+
	48	0.22	+
	49	0.37	+
ureB + HAP + CT	25	0.15	-
	26	0.07	-
	27	0.03	-
	28	0.64	+
	29	0.13	-
	30	0.02	-
	31	0.66	+
	32	0.00	-
ureA + HAP + CT	68	0.00	-
	69	0.07	-
	70	0.42	+
	71	0.00	-
	72	0.00	-

17 MAR 1999

ureB + HAP + CT	73	0.37	+
	74	0.00	-
	75	0.37	+
	76	0.00	-
	77	0.00	-
	78	0.00	-
	79	0.39	+
	80	0.00	-
	81	0.37	+
	82	0.00	-

In Table 4, "CT" means cholera toxin; "UreA" means recombinant *H. pylori* urease A subunit; "UreB" means recombinant *H. pylori* urease B subunit; and "HAP" means hydroxyapatite crystals. Mice 20 to 54 were sacrificed 12 days post challenge and mice 68 to 82 10 weeks (106 days) post challenge. The results of the urease test performed from biopsies of the stomach of each animal are expressed as OD values at 550 nm. The positive and negative signs depicts the final status of infection of each animal, according to the positivity or negativity of the urease test for detection of *H. felis*. Positivity: OD₅₅₀ values >0.2

TABLE 5: Protection as measured 12 days post challenge

<u>Immunization</u>	<u>Challenge</u>	<u>% infected</u>	<u>% protected</u>
Urease A subunit	<u>H. felis</u>	10/10 (%)	0/10 (0%)
Urease B subunit	<u>H. felis</u>	3/10 (30%)	7/10 (70%) *

17 MAR 1999

CT	<u>H. felis</u>	10/10 (100%)	0/10 (0%)
----	-----------------	--------------	-----------

* p=0.0031 (two tailed Fisher exact test) compared to CT control

TABLE 6: Protection as measured 10 weeks post challenge

<u>Immunization</u>	<u>Challenge</u>	<u>% infected</u>	<u>% protected</u>
Urease A subunit	<u>H. felis</u>	1/5 (20%)	4/5 (80%) *
Urease B subunit	<u>H. felis</u>	4/10 (40%)	6/10 (60%) **

* p=0.003 (two tailed Fisher exact test) compared to CT control

** p=0.01 (two tailed Fisher exact test) compared to CT control

It will be seen from the results set out in Tables 4, 5 and 6 that statistically significant protection against H. felis challenge is obtained with oral immunization using recombinant H. pylori urease B subunit as compared to that obtained using either recombinant H. pylori urease A subunit or cholera toxin. Referring to Table 4, it will be seen that, 12 days post challenge, from a total of 10 immunized animals, only 3 were found infected in the urease B subunit group, as compared to all 10 animals immunized with H. pylori A subunit of urease and 10 out of 10 of the animals immunized with cholera toxin. Table 4 shows that 70% of the animals were protected from challenge by H. felis as compared to 0% of the animals immunized with H. pylori urease A subunit and 0% of the animals immunized with cholera toxin and then subjected to H. felis challenge. In other words, 100% of the control mice challenged with H. felis became infected whereas, in contrast, in mice immunized with

17 MAR 1999

recombinant H. pylori urease B subunit the infection rate was only 30%. This represents a significant reduction in infection ($p=0.0031$, Fisher exact test) as compared to the control mice.

The fact that the protection observed with H. pylori urease is entirely conferred by immunization with the B subunit of urease, and that the A subunit has no such effect, was not expected on the basis of our experiment with purified urease. This definition of the roles of the 2 structural subunits of urease in the development of a protective immune response is therefore novel. The protection obtained using recombinant urease B subunit, which is enzymatically inactive also teaches that non toxic forms of urease can be used as oral vaccine against Helicobacter infection. Furthermore these results strongly suggest that recognition of the active site is not required for protection, as an inactive urease B subunit is very unlikely to induce antibodies that will recognize and inhibit the catalytic site of native urease.

Referring to Table 6, it will be seen that, when mice are sacrificed 10 weeks post infection, 60% (6 mice out of 10) of the animals immunized with urease B subunit and 80% (4 mice out of 5) of the animals immunized with H. pylori urease A subunit were protected against H. felis infection. The fact that protection obtained through immunization with urease B subunit lasts over time and that immunization with urease A induces a protection which is delayed compared to the one induced by urease B subunit could not be expected from our experiment with purified urease or with other experiment performed earlier. The fact that urease A subunit immunization confers protection definitively proves that recognition of the active site is not required for protection. Figure 6 summarizes results obtained after oral immunization with recombinant urease A and B subunits (described in Table 5 and 6)

17 MAR 1999

The present invention also provides vaccine compositions suitable for the prevention of Helicobacter infection. The compositions comprise an effective amount of a urease antigen, preferable H. pylori urease or recombinant H. pylori urease subunits, capable of eliciting in a host a protective immune response to Helicobacter infection, in association with a pharmaceutically acceptable carrier or diluent.

The vaccines of the invention are administered in amounts readily determined by persons of ordinary skill in this art. Thus, for adults a suitable dosage will be in the range of 10ug to 100 milligrams, for example 50ug to 50mg. Similar dosage ranges will be applicable for children. Carrier systems in humans may include enteric release capsules protecting the antigen from the acidic environment of the stomach, and including urease antigen in a insoluble form as fusion proteins. The vaccine can be administered as a primary prophylactic agent in adults or in children, as a secondary prevention, after successful eradication of H. pylori in an infected host, or as a therapeutic agent in the aim to induce an immune response in the host susceptible to contribute to the eradication of H. pylori.

As noted above, a suitable mucosal adjuvant is cholera toxin. Others which may be used are muramyl dipeptide or its derivatives, non-toxic derivatives of cholera toxin, including its B subunit and/or conjugates or genetically engineered fusions of the urease antigen plus cholera toxin or its B subunit. Other suitable delivery methods include biodegradable microcapsules or immune stimulating complexes (ISCOM'S) or liposomes, genetically engineered attenuated live vectors such as viruses or bacteria, and recombinant (chimeric) virus-like particles, e.g. bluetongue. The amount of mucosal adjuvant employed depends on the type of mucosal adjuvant used. For example, when the mucosal adjuvant is cholera toxin, it is suitably used in an amount of 5 ug to 50 ug, for example 10ug to 35ug. When used in the form of microcapsules,

17 MAR 1999

the amount used will depend on the amount employed in the matrix of the microcapsules to achieve the desired dosage. The determination of this amount is within the skill of a person of ordinary skill in this art.

Suitable carriers for the vaccines of the invention are enteric coated capsules and polylactide-glycolide microspheres. Suitable diluents are 0.2N NaHCO₃ and/or saline.

Particulate hydroxylated calcium phosphate (HCP) is especially useful as a carrier for the H. pylori urease to be applied to mucosal surfaces. It is believed that the H. pylori urease-hydroxylated calcium phosphate conjugate is transported across epithelium where it raises a poly Ig immune response. Preferably, the hydroxylated calcium phosphate is in the form of microparticles suitable for transport across the epithelium, particularly by cells specialized for this purpose (M cells). A preferred form of hydroxylated calcium phosphate is hydroxyapatite, a commercially available crystalline hydroxylated calcium phosphate Ca₁₀(PO₄)₆(OH)₂.

Commercially available hydroxyapatite generally consists of slab-like crystals that are chemically and physically analogous to inorganic hydroxyapatite in normal bone tissue. Ingestion of hydroxyapatite should therefore be safe, as evidenced by the existence of nutritional calcium/phosphorus supplements derived from ground bone, which are designed to be ingested. Commercially-high resolution hydroxyapatite (from CalBioChecm) consists of crystals varying widely in size. Crystals over 1um in length are unlikely to be taken up by M cells. Therefore, for use in the invention, commercial hydroxyapatite crystals are broken into small, relatively uniform crystalline fragments such as by sonication. Preferably, a substantial proportion of the hydroxyapatite is present as fragments of about 0.01-1.0um. Fragmentation may be measured either by electron microscopy or light scattering, using standard techniques.

17 MAR 1999

Preferred modes of administration of the H. pylori urease antigen urease antigen are orally, nasally, rectally or ocularly. Oral administration can provide delivery to other G.I. mucosa including the intestinal mucosa.

The vaccines of the present invention may be administered to a mucosal surface in the form of an aerosol, suspension, capsule and/or suppository. The method of administration will be readily apparent to a person of ordinary skill in this art.

The present invention further includes the passive immunization of mammals, including humans, against Helicobacter infection. This is achieved by administering to a mucosal surface of the patient an effective amount of a urease specific antibody. Preferably, an effective amount of a H. pylori urease specific IgA monoclonal antibody.

Since the urease of H. pylori is shown to represent the antigen involved in inducing protective immunity, a further aspect of the invention is the use of H. pylori urease as a diagnostic reagent to measure the immune response of persons who have received a vaccine based on urease or to determine whether an individual is immune or susceptible (and thus in need of vaccination). The present invention also includes the use of urease and urease-specific antibodies to construct assays and kits for diagnosis of Helicobacter immunity, assessment of Helicobacter susceptibility, and definition of immune responses to vaccines.

EXAMPLES

The invention will now be further described by reference to the following non-limiting

17 MAR 1999

examples.

a) The Bacterial Strains

H. felis was provided by J. Fox (division of Cooperative Medicine, Mass. Institute of Technology, Boston, USA). H. pylori was isolated from patients with ulcer disease (CHUV, Lausanne, Switzerland).

b) Bacterial Cultures

Liquid Culture - Bacteria were cultured on BHI (Brain-Heart Infusion, BioMerieux) liquid medium containing 0.25% of yeast extract (Difco) and 10% of fetal calf serum (Inotech) supplemented with 0.4% of Campylobacter selective complement (Oxoid). The bacteria were incubated overnight under microaerophilic conditions at 37°C and then shaken at 37°C for 2 to 3 days.

Culture on Agarose plates - The bacteria were cultured on agar plate consisting on BHI with 0.25% of yeast extract and 5% of sheep blood under microaerophilic conditions at 37°C for 3 days.

Quantification - The quantity of bacteria was determined by the optical density of the BHI solution at 660 nm (1 optical density unit corresponding to 10^8 bacteria).

c) Preparation of sonicates

H. pylori was collected from 31 blood agar plates in 0.15 M NaCl and spun 5 minutes at 1400g at 4°C. The pellet was resuspended in 3 ml of NaCl and sonicated for 4

17 MAR 1999

minutes. The amount of proteins was evaluated by a Bradford assay (BioRad Kit according to supplier).

d) Coupling of immunogen to Hydroxyapatite

Immunogen (urease or subunit thereof) was incubated for 1 hour at 4°C with hydroxyapatite. 1.0 mg of hydroxyapatite was used for 30ug of immunogen per mouse. At the end of the incubation, 10 ug of cholera toxin was added in a final volume of 200ul PBS.

EXAMPLE 1

a) Extraction

H. pylori from 30 blood agar plates was harvested in 0.15 M NaCl on ice. The solution was spun 5 minutes at 1400 g at 4°C. The pellet was resuspended in 20 ml of H₂O and vortexed for 45 seconds (maximum speed). The extract was then spun 20 minutes at 6700 g at 4°C. The supernatant was recovered and the quantity of protein was evaluated (see "Quantification" above) and precipitated with 70% of ammonium sulfate.

b) Purification of urease

The solution was chromatographed on a Sepharose CL-6B column (Pharmacia) with PBS (phosphate buffered saline) as mobile phase. The 22 collected fractions which presented a strong urease activity were pooled and dialyzed overnight at 4°C against 3

17 MAR 1999

liters of PEB (20 mM phosphate buffer, pH 7) and then chromatographed on a Q Sepharose fast flow (Pharmacia) with PEB as mobile phase. The fractions were eluted by 0 to 500 mM NaCl gradient. Ten of the collected fractions with a strong urease activity were individually subjected to an SDS gel followed by a Coomassie staining. The 6 fractions presenting 2 distinct bands corresponding to MW-63 and -28 KDa were pooled and were considered as the purified urease.

EXAMPLE 2 (see also section B)

Mice employed in the immunization studies were lightly anesthetized with ether prior to intragastric immunization. And then, sonicate preparation or purified urease, hydroxyapatite and cholera toxin was suspended in PBS and 200ul were delivered to the stomach of the respective mice using a polyethylene tubing attached to a hypodermic syringe. This procedure will be referred to as oral immunization.

Three oral immunization protocols were evaluated. These are described below.

Protocol B1 - Vaccination with purified urease

Female BALB/c six-week old mice (20) were orally immunized with 30 ug of purified H. pylori of purified urease and 1 mg of hydroxyapatite and 10 ug of cholera toxin at day 0, 7, 14 and 21. The mice were challenged at day 28 and 30 with 5×10^7 and 10^8 H. felis from liquid culture.

Protocol B2 - Vaccination with purified urease

Female BALB/c six-week old mice (20) were orally immunized with 2 mg of H. pylori

17 MAR 1999

sonicate solution at day 0, 7, 14 and 21. Ten mice were challenged at day 28 and 30 with 5×10^7 and 10^8 H. felis.

Protocol B3 - Control

Female BALB/c six-week old mice were orally immunized with 1 mg hydroxyapatite and 10ug of cholera toxin at day 0, 7, 14 and 21. The mice were challenged at day 28 and 30 with 5×10^7 and 10^8 H. felis.

At day 35 all mice were sacrificed and biopsies from the stomach were taken as well as intestinal secretions and blood.

Protection and evaluation

To evaluate protection, biopsies were screened for the urease activity by the Jetrox HP test (Rohm Pharma) according to the instructions of the supplier. The urease is quantified by a spectrophotometric measurement at 550nm. The biopsies were also cultured in the presence of H. felis and was estimated by Gram staining. Gastric antral biopsies were homogenized and diluted (1:10 and 1:1000) in 0.15 M NaCl and plated onto blood agar plates and incubated under microaerophilic conditions at 37°C for 4 to 10 days.

ELISA

Intestinal secretions and blood were analyzed by ELISA for the evaluation of antibody titer. The ELISA was carried out as follows. Polystyrene plates (96 wells) were coated with 1 ug/well of purified urease at 37°C for 2 hrs. Non specific binding sites were

blocked with 5% powdered milk in PBS 0.1% Tween at 37°C for 30 minutes. The plates were washed once with PBS 0.1% Tween. Blood samples were tested at dilution 1:1000 and intestinal secretions 1:1. 100ul of each sample were added to the antigen coated plates. After 2 hrs of incubation, plates were washed 3 times with PBS 0.1% Tween. Anti-mouse biotinylated whole antibody from goat and anti-mouse IgA, IgG and IgM biotinylated (Amersham) were added (100ul) at dilution 1:500 except for IgA (1:250) and incubated at 37°C for 1 hr. The plates were washed 3 times with PBS 0.1% Tween and 100ul of 1:1000 dilution of streptavidin Horseradish peroxidase in PBS 0.1% Tween were added and incubated at 37°C for 30 minutes. The plates were washed 3 times and 50 ul of 1:50 dilution of o-phenyl-diamine in citrate buffer pH 5.0 with 1ul/ml of 30% H₂O₂ were added and incubated at room temperature for 20 minutes. The absorbance at 495 nm was measured in each well.

EXAMPLE 3 (see also section C)

Mice employed in the immunization studies were lightly anesthetized with ether prior to intragastric immunization. Then, 30 ug recombinant H. pylori urease A or B subunit produced in E. coli bound hydroxyapatite and supplemented with cholera toxin was suspended in PBS and 200ul were delivered to the stomach of the respective mice using a polyethylene tubing attached to a hypodermic syringe. This procedure will be referred to as oral immunization.

Three oral immunization protocols were evaluated. These are described below.

Protocol C1 - Vaccination with recombinant urease A subunit

Female BALB/c six-week old mice (10) were orally immunized with 30ug of purified

17 MAR 1999

recombinant H. pylori urease A subunit and 1 mg of hydroxyapatite and 10ug of cholera toxin at day 0, 8, 14 and 21. Ten mice were challenged at day 32, 34 and 36 with 10^8 H. felis from liquid culture.

Protocol C2 - Vaccination with recombinant urease B subunit

Female BALB/c six-week old mice (10) were orally immunized with 30 ug of recombinant H. pylori urease B subunit and 1 mg of hydroxyapatite and 10ug of cholera toxin at day 0, 8, 14 and 21. Ten mice were challenged at day 32, 34 and 36 with 10^8 H. felis from liquid culture.

At day 42, or at day 106, mice were sacrificed and multiple biopsies from the stomach were taken.

Protection and evaluation

To evaluate protection, biopsies of the corpus and antrum of the stomach were screened for urease activity by the Jatrox HP test (Rohm Pharma) according to the instructions of the supplier. The urease is quantified by a spectrophotometric measurement at 550nm. The total of corpus and antrum OD values were added to obtain a final OD value for each mouse.

References

1. Blaser, M.J. "Gastric Campylobacter-like organisms, gastritis and peptic ulcer disease" Gastroenterology 1987, 93, 371-383.

17 MAR 1999

2. Graham, D.Y. "Campylobacter pylori and peptic ulcer disease" Gastroenterology 1989, 96, 615-625.
3. Parsonnet, J. et al "Helicobacter pylori infection in intestinal and diffuse-type gastric adenocarcinomas" J. Natl. Cancer Inst. 1991, 93, 640-643.
4. Marshall, B.J. et al "Attempt to fulfill Koch's postulate for pyloric Campylobacter" Med. J. Aust. 1985, 142, 436-439.
5. Morris, A. et al "Ingestion of Campylobacter pyloridis causes gastritis and raised fasting gastric pH" Am. J. Gastroenterol. 1987, 82, 192-199.
6. Engstrand, L. et al "Inoculation of barrier-born pigs with Helicobacter pylori: a useful animal model for gastritis type B" Infect. Immun. 1990, 53, 1763-1768.
7. Fox, J.G. et al "Gastric colonization by campylobacter pylori subsp. mustelae in ferrets" Infect. Immun. 1988, 56, 2994-2996.
8. Fox, J. G. et al "Helicobacter mustelae-associated gastritis in ferrets: an animal model of Helicobacter pylori gastritis in humans" Gastroenterology 1990, 99, 352-361.
9. Lee, A. et al "A small animal model of human Helicobacter pylori active chronic gastritis" Gastroenterology 1990, 99, 1315-1323.
10. Fox, J. G. et al "Helicobacter felis gastritis in gnotobiotic rats: an animal model of Helicobacter pylori gastritis" Infect. Immun. 1991, 59, 785-791.

17 MAR 1999

11. Eaton, K.A. et al "Campylobacter pylori virulence factors in gnotobiotic piglets" Infect. Immun. 1989, 57, 1119-1125.

12. Peterson, W.L. "Helicobacter pylori and peptic ulcer disease" N. Engl. J. Med. 1991, 324, 1043-1048.

13. Czinn, S.J. and Nedrud, J.G. "Oral immunization against Helicobacter pylori" Infect. Immun. 1991, 59, 2359-2363.

14. Brandtzaeg, P. "Role of H chain and secretory component in receptor-mediated glandular and hepatic transport of immunoglobulins in man" Scand. J. Immunol. 1985, 22, 111-146.

15. Brandtzaeg, P. et al "Production and secretion of immunoglobulins in the gastrointestinal tract" Ann. Allergy 1987, 59, 21-39.

16. Wyatt, J.I. et al "Local immune response to gastritic campylobacter in non-ulcer dyspepsia" J. Clin. Path. 1986, 39, 863-870.

17. Lee, A. et al. "Pathogenicity of Helicobacter pylori: A Perspective" Infect. Immun. 1993, 61, 1601-1610.

18. Pallen, M.J. and Clayton, C.L. "Vaccination against Helicobacter pylori urease, letter" Lancet, 1990, 336, 186.

19. Evans, D.J. et al. "Urease-associated heat shock protein of Helicobacter pylori" Infect. Immun. 1992, 60, 2125-2127.

17 MAR 1999

20. Ferrero, R.L and Lee, A. "The importance of urease in acid protection for the gastric-colonising bacteria Helicobacter pylori and Helicobacter felis sp. nov." Microb. Ecol. Health Dis. 1991, 4, 121-134.
21. Chen, et al. "Immunization against gastric Helicobacter infection in a mouse/Helicobacter felis model, letter" Lancet, 1992, 339, 1120-1121.
22. Czinn, S. et al. "Oral immunization protects germ-free mice against infection from Helicobacter felis" Proceedings of the DDW, American Gastroenterological Association. May 10-13, 1992, 1321, A-331.
23. Guo, M. and Liu, P.V. "Serological specificities of ureases of Proteus species". J. Gen. Microbiol, 1965, 136, 1995-2000.
24. Michetti, P. et al. "Specificity of mucosal IgA response in Balb/C mice following H. felis or H. pylori challenges. Proceedings of the DDW, American Gastroenterological Association. May 10-13, 1992, 1001, A-251.
25. Davin, C. et al. "H. pylori urease elicits protection against H. felis infection in mice". Proceedings of the DDW, American Gastroenterological Association. May 16-19, 1993, 1213, A-304.
26. Pallen, M.J. and Clayton, C.L. "Vaccination against Helicobacter pylori urease". Lancet 1990, 336, 186-7.
27. Pimentel, J.L. and Cook, M.E. "Improved growth in the progeny of hens

17 MAR 1999

immunised with jackbean urease". Poultry Sci. 1988, 64, 434-439.

28. Labigne, A. "Sequences of nucleotides coding for a protein having an urease activity". EPO patent application # EP O 367 644 A1, 1989. International publication # WO 90/04030, 1990.

29. Clayton, C.L. et al. S. "Nucleotide sequence of two genes from *Helicobacter pylori* encoding for urease subunits". Nucleic Acid Res. 1990, 18, 362.

30. McGhee, J.R and Kyono, H. "New perspectives in vaccine development: mucosal immunity to infections". Infect Agents Dis. 1993, 2, 55-73.



17 MAR 1999

Claims

1. Use of a polypeptide presenting a sufficient number of epitopes of a *Helicobacter* antigen in the preparation of a mucosal vaccine for inducing an immune response to *Helicobacter* in a mammal.
2. The use of claim 1, wherein said mammal is at risk of developing, but does not have, *Helicobacter* infection.
3. The use of claim 1, wherein said mammal is infected by *Helicobacter*.
4. The use of claim 1, wherein said polypeptide comprises a *Helicobacter* urease.
5. The use of claim 1, wherein said *Helicobacter* antigen is a *Helicobacter pylori* antigen.
6. The use of claim 1, wherein said *Helicobacter* antigen is a *Helicobacter* urease.
7. The use of claim 6, wherein said polypeptide comprises a subunit of said *Helicobacter* urease.
8. The use of claim 7, wherein said subunit is the urease A subunit.
9. The use of claim 7, wherein said subunit is the urease B subunit.



17 MAR 1999

41

10. The use of claim 1, wherein said polypeptide comprises an anti-idiotypic antibody to a *Helicobacter urease*.
11. The use of claim 1, wherein said polypeptide is recombinant.
12. The use of claim 1, wherein said polypeptide is produced as a fusion protein.
13. The use of claim 12, wherein said fusion protein comprises a cholera toxin subunit.
14. The use of claim 1, wherein said mucosal vaccine is administered to said mammal with a mucosal adjuvant.
15. The use of claim 14, wherein said mucosal adjuvant comprises a cholera toxin.
16. The use of claim 1, wherein said mammal is human.
17. The use of claim 1, wherein said mucosal vaccine is administered to said mammal with a hydroxylated calcium phosphate.
18. The use of claim 17, wherein said hydroxylated calcium phosphate is hydroxyapatite.



17 MAR 1999

19. The use of claim 18, wherein said hydroxyapatite is in the form of particles suitable for transport across epithelium.

20. The use of claim 1, wherein said mucosal vaccine is administered to an oral, intranasal, rectal, or ocular surface of said mammal.

21. A vaccine for inducing a mucosal immune response to *Helicobacter* in a mammal, said vaccine comprising a polypeptide presenting a sufficient number of epitopes of a *Helicobacter* antigen to induce said immune response, in a pharmaceutically acceptable carrier or diluent.

22. The vaccine of claim 21, further comprising a mucosal adjuvant.

23. The vaccine of claim 21, wherein said polypeptide comprises a *Helicobacter* urease or a subunit of a *Helicobacter* urease.

24. The vaccine of claim 21, wherein said *Helicobacter* antigen is a *Helicobacter pylori* antigen.

25. The vaccine of claim 21, wherein said *Helicobacter* antigen is a *Helicobacter* urease.

26. Use of an IgA antibody specific for a *Helicobacter* antigen in the preparation of a medicament for imparting passive



17 MAR 1999

43

protection to *Helicobacter* infection in a mammal by mucosal administration.

27. The use of claim 26, wherein said *Helicobacter* antigen is a *Helicobacter pylori* antigen.

28. The use of claim 26, wherein said *Helicobacter* antigen is a *Helicobacter urease*.

29. The use of claim 26, wherein said mammal is human.

30. A method for assessing the immune response of a mammal infected by a *Helicobacter* organism, said method comprising detecting the presence of antibodies reactive with a polypeptide of a *Helicobacter* in a sample from the gastrointestinal tract of said mammal.

31. The method of claim 30, wherein said polypeptide of a *Helicobacter* is a *Helicobacter urease*.

32. The method of claim 30, further comprising administering a polypeptide of a *Helicobacter* to a mucosal surface of said mammal as a vaccine antigen prior to said detecting.

17 MAR 1999

ABSTRACT**Urease-Based Vaccine against Helicobacter Infection**

Method of eliciting in a mammalian host a protective immune response to Helicobacter infection by administering to the host an immunogenically effective amount of a Helicobacter urease or urease subunits as antigen. Vaccine compositions are also provided.

No Figure

17 MAR 1999

1/4

FIGURE 1

GROUP A: MICE NOT PROTECTED AFTER IMMUNIZATION WITH
UREASE





FIGURE 2

GROUP B: MICE PROTECTED AFTER IMMUNIZATION WITH
UREASE



17 MAR 1999

2/4

FIGURE 3

GROUP C: MICE NOT PROTECTED AFTER IMMUNIZATION WITH H.p.
SONICATE





FIGURE 4


GROUP D: MICE PROTECTED AFTER IMMUNIZATION WITH
H.p SONICATE



17 MAR 1999

3/4

FIGURE 5



immunization :	<i>H. pylori</i> sonicate	<i>H. pylori</i> urease	control
----------------	------------------------------	----------------------------	---------

protected :	3/9	7/10	1/10
-------------	-----	------	------

* Two-tailed Fisher Exact Test

17 MAR 1999

4/4

FIGURE 6

