




FIG. 1




FIG. 2

Murine soluble Lag-3, extracellular, without signal peptide. Accession Number:

UniProtKB-Q61790

SGPGKELPVVWAQEGAPVHLPCLSKSPNLDPNFLRRGGVWQHQPDSGQPTPIPALDLHQ
 GMPSPRQPAAGRYTVLSVAPGGLRSGRQPLHHPVQLEERGLQRGDFSLWLRPALRTDAGE
 YHATVRLFNRLSCSLRLRVGQASMIASPSGVKLKLSDWVLLNCSFSRPDRPVSVHWFQGG
 NRVFVYNSPRHFLAETFLLLPQVSFLDSGTWGCVLTYRDGFNVSIITYNLKVLGLEPVAPL
 TVYAAEGSRVELPCHLPFGVGTFSLLIAKWTFFGGGPELPVAGKSGNFTLHLEAVGLAQA
 GTYTCSIHLQGGQLNATVTILAVITVTPKSFGLPGSRGKLLCEVTPASGKERFVWRPLNNL
 SRSCPGPVLEIQEARLLAERWQCQLYEGQRLLGATVYAAESSGAHSARRISGDLKGGHL

Human LAG-3 proteins. Accession number: NP_002277, 525 amino acids full length

mweaqflql flqlplwvapv kplqpgaevp vvwaqegapa qlpcseptipl qdlslrrag
 vtqhqpdsq ppaaapghpl apgphpaaps swgprprryt vlevpggqlr sgrlplqprv
 qldergrgrg dfelwlrpar radageyraa vhlrdralec rlrirlqgas mtasppqslr
 asdwilnce ferpdrrpasv hwfrnrgqgr vpvresphhh laesflflpq vspmdsgpwg
 ciltyrdgfn veimynltvl glepptpltv yagagervgl porlpagvgt rslftakwtp
 pgggpdllvt qdngdftlrl edveqagagt ytchihlqeq qlnatvtlai itvtpksfgs
 pgslgkllce vtpvsgqerf vwsaldtpsq rsfsgpwlea qeaqlleqpw qcglyqgerl
 lgaavyftel sspgaqrgr apgalpaghl llflilgvl llllvtgafg fhlwrrqwrp
 rrfsaleqgi hppqaqskie eleqepesep epepepep epeql

FIG. 4

Human LAG-3 proteins. Accession number: XXX, truncated soluble LAG-3




Table 2. Multivariable regression analysis of independent predictors for plasma LAG-3 (A) and HDL-C (B): MESA.**A. Multivariable regression analysis of independent predictors for plasma LAG-3: MESA.**

Covariates	N	Beta	SE	P value
	5137			
Rs10846744:				
GG:		183.98	91.1	0.04
GC:		-261.8	80.1	0.001
Race:				0.0005
White		-641.37	408.5	
Chinese		491.2	1061.6	
African American		862.7	450.2	
Age		-19.81	6.7	0.003
Lipid meds		153.22	70.5	0.03
Smoking		-10.95	2.72	<0.0001

Covariates included race, PCs of ancestry, age, study sites, sex, HgbA_{1c}, BMI, lipid medications, lipids (TC, LDL-C, HDL-C), smoking pack years, systolic BP, and diastolic BP. LAG3 was log-transformed.

B. Multivariable regression analysis of independent predictors for HDL-C: MESA.

Covariates	N	Beta	SE	P value
	4953			
LAG3		-0.770	0.28	0.007
Age		0.056	0.02	0.006
Triglycerides		-0.062	0.002	<0.0001
Sex: Men		5.67	0.19	<0.0001
BMI		-0.623	0.035	<0.0001
HgbA _{1c}		-0.450	0.181	0.01
Alcohol		3.28	0.370	<0.0001
Systolic BP		0.027	0.01	0.03

Covariates included race, PCs of ancestry, age, study sites, sex, HgbA_{1c}, BMI, lipid medications, lipids (triglycerides), smoking pack years, current alcohol use, systolic BP, and diastolic BP. LAG3 was log-transformed.

FIG. 5

Reference (GG) expressing cell: Cell line

Isotype Control LAG-3




FIG. 6

Panel 2







Risk (CC) expressing cell**Isotype Control****LAG-3****Untreated****Vehicle (1 h)****Cocktail (1 h)**

FIG. 6(Cont.)

Panel 2 Risk (CC) expressing cell

Isotype Control LAG-3




FIG. 6(Cont.)

Reference (GG) expressing cell: Cell line
Isotype Control **LAG-3**




FIG. 6(Cont.)




FIG. 7




FIG. 8




FIG. 9




FIG. 10




FIG. 11

Table 3. Association of plasma LAG3 with CHD in MESA participants.

Variable	Odds Ratio	95% CI
**LAG3 (log scale)	1.15	1.02-1.28
*Age	2.41	1.81-3.21
*Sex	2.10	1.50-2.94
*Systolic blood pressure	1.60	1.25-2.06
**LDL cholesterol	1.79	1.11-2.87
††Total cholesterol	0.60	0.37-0.97
*Lipid medication	1.60	1.18-2.17
ΔΔHgb _{A1c}	1.09	1.01-1.14
*Smoking pack-years	1.07	1.00-1.14

Table 4. Association of plasma LAG3 with CHD in MESA participants with HDL-C ≥ 60 mg/dL.

Variable	Odds Ratio	95% CI
*LAG3 (log scale)	1.45	1.12-1.85
**Age	2.79	1.35-5.78
*Sex	3.80	1.68-8.56
††Diastolic blood pressure	0.49	0.26-0.92

Table 5. Plasma LAG3 as a CHD risk predictor compared with Framingham risk score.

	Beta	SE	P-value
Model 1: (N=5468)			
Framingham risk score (log scale)	0.872	0.093	< 0.0001
Model 2: (N=5468)			
Framingham risk score (log scale)	0.875	0.094	< 0.0001
LAG3 (log scale)	-0.065	0.031	0.039
Model 3: (N=5468)			
Framingham risk score (log scale)	0.891	0.095	< 0.0001
Model 4: (N=5468)			
Framingham risk score (log scale)	0.893	0.095	< 0.0001
LAG3 (log scale)	-0.064	0.031	0.044

Table 6. Association of plasma LAG3 with inflammatory markers in MESA participants.

Outcomes	N	Beta	SE	P-value
IL-2	2386	-0.002	0.004	0.624
IL-6	5414	-0.006	0.044	0.203
IL-10	2350	0.035	0.007	< 0.0001
CD40 ligand	851	-0.0004	0.010	0.966
sTNFαR	2396	-0.001	0.003	0.824
hs-CRP	5533	-0.013	0.007	0.087

FIG. 12

CD4⁺ Lag3⁺ T cell Activation in *Ldlr*^{-/-} Following High Fat Feeding

- *LDLr*^{-/-} mice fed high-fat diet (HFD) for 10 weeks
- CD4⁺ in spleen investigated for Lag-3 expression




FIG. 13

Bone Marrow Transfer of Lag-3^{-/-} Cells Significantly Associated with Aortic Root Activated CD4⁺ T Cells






FIG. 14

Lag-3^{-/-} Deficiency Significantly Associated with Impaired Tregs

(exTregs)

Significant Association of Para-Aortic Tregs and Lesion Size






FIG. 15

% Tregs (Foxp3⁺CD25⁺ of CD4⁺)


Recombinant human monomeric LAG3




Wild-type mouse B splenocytes (approximately 1 million) were incubated with varying concentrations of recombinant human LAG3 (µg protein/ml) and then stained with anti-human LAG3 (upper panels, 5 µl per condition) or control mouse IgG1 antibodies and then analyzed by flow cytometry.

FIG. 16

Recombinant human monomeric LAG3





Anti-mouse IgG1 APC



Wild-type mouse B splenocytes (approximately 1 million) were incubated with varying concentrations of recombinant human LAG3 (µg protein/ml) and then stained with anti-human LAG3 (upper panels, 5 µl per condition) or control mouse IgG1 antibodies and then analyzed by flow cytometry.

FIG. 16(cont.)

Anti-human LAG3 (5 μ l Ab/condition)



Anti-human LAG3 (2 μ l Ab/condition)





FIG. 17

Human LAG-3 Exon 3 Transcription in Lymphocytes under
Basal and Post-stimulation Conditions




FIG. 18