

DESCRIPTION

Novel Glycopeptide and Peptide Having a Kokumi Taste Imparting Function, and Method of Imparting the Kokumi Taste to Foods

Technical Field of the Invention

The present invention relates to glycopeptides and peptides which have a taste improving function, particularly a kokumi taste imparting function. The present invention more specifically relates to such glycopeptides and peptides per se, a method for imparting the kokumi taste to foods or seasonings with the use of the glycopeptide(s) and/or the peptide(s), a method for improving foods in kokumi taste with the use of a seasoning containing the glycopeptide(s) and/or the peptide(s), and foods or seasonings to which the kokumi taste has been imparted by such a kokumi taste improving method.

Background Art

The kokumi taste refers to a taste which cannot be expressed by the five basic tastes (sweet taste, salt taste, sour taste, bitter taste, and umami taste), and which is obtainable by enhancing not only the basic tastes but also tastes around, or peripheral to, the basic tastes, such as thickness, spread, continuity, unity, and the like.

Heretofore, some methods for imparting a kokumi taste have been

reported. Among them, there are known a method in which glutathione is added (see Japanese Patent No. 1464928), a method in which a heated product of gelatin and tropomyosin is added (see Japanese Patent Application Laid-open (Kokai) No. Hei 10-276709), a method in which a sulfone group-containing compound is added (see Japanese Patent Application Laid-open (Kokai) No. Hei 8-289760), and the like.

Further, as a kokumi taste imparting method other than the methods described above, an attempt has been made to impart a kokumi taste by increasing the ratio occupied by peptides in a seasoning (see "Food Processing and Ingredients", Vol 31, No. 12, pp. 17-20 (1996)). This is an attempt to increase the amount of peptides in the whole of a food to impart a kokumi taste to the whole of the food or foods containing the peptides. However, such an attempt failed to provide such a desired effect, because, for example, the titer was low and peptides with bitterness imparted a bitter taste. Further, as described in Japanese Patent Application Laid-open (Kokai) No. 2002-335904, an attempt has been made to provide a seasoning for imparting a kokumi taste obtained by binding sugar and unspecified peptide according to the Maillard reaction. However, the effect of such seasoning was not so strong, and the seasoning often imparts a negative effect when added to foods, such as impartment of a bitter taste or coloring due to the browned product.

Disclosure of the Invention


[Problems to be Solved by the Invention]

In view of the background of the background art described above, it is an object of the present invention to provide a food material which can be used for wider purposes and has a stronger effect of improving the taste, and which can impart a kokumi taste, that is, enhance the basic tastes and impart thickness, spread, continuity, unity, and the like, associated with the enhancement of the basic tastes.

[Means for Solving the Problems]

A first aspect of the present invention provides a method of imparting a kokumi taste to a food or a seasoning which method comprises adding to said food or said seasoning, a glycopeptide or peptide having the chemical structure represented by the below-mentioned general formula (a) in such that said glycopeptide and/or peptide shall be contained in an amount of 1 ppb to 1,000 ppm (w/w), in said food or seasoning.

General formula (a)




wherein Asn and His represent an asparagines residue and a histidine residue, respectively, X represents a hydrogen atom or a sugar chain, and A and B each independently represent a

hydrogen atom or an optical amino acid residue.

A second aspect of the present invention provides a method of imparting a kokumi taste to a food which comprises adding the seasoning as set forth in the first aspect in such that it shall be contained in an amount of 0.01-10% (w/w), in said food.

A third aspect of the present invention provides a glycopeptide or peptide having the chemical structure represented by the following general formula (b):





wherein Val, Asn, His And Thr represent a valine residue, an asparagines residue, a histidine residue, and a threonine residue, respectively, X represents a hydrogen atom or a sugar chain.

A fourth aspect of the present invention provides a method of imparting a kokumi taste to a food or a seasoning which method comprises adding to said food or said seasoning, said glycopeptide and/or peptide as set forth in the third aspect in such that said glycopeptide and/or peptide shall be contained in an amount of 1 ppb to 1,000 ppm (w/w), in said food or seasoning.

A fifth aspect of the present invention provides a method of imparting a kokumi taste to a food which comprises adding the seasoning as set forth in the fourth aspect in such that

it shall be contained in an amount of 0.01-10% (w/w), in said food.

A sixth aspect of the present invention provides a glycopeptide or peptide having the chemical structure represented by the following Sequence formula (I), (II), (III), (IV), (V) or (VI):



wherein Val, Asn, His, and Thr have the same meanings as in the general formula (b) given above, and GlcNAc represents an N-acetyl-glucosamine residue, Fuc represents a fucose residue, Man represents a mannose residue, and Xyl represents a xylose residue.

A seventh aspect of the present invention provides a

method of imparting a kokumi taste to a food or a seasoning which method comprises adding to said food or said seasoning, at least one of the glycopeptides and peptides having the chemical structure represented by the sequence formula (I), (II), (III), (IV), (V) or (VI) in such that said glycopeptide(s) and/or peptide(s) shall be contained in an amount of 1 ppb to 1,000 ppm (w/w), in said food or seasoning.

An eighth aspect of the present invention provides a method of imparting a kokumi taste to a food which comprises adding the seasoning as set forth in the seventh aspect in such that it shall be contained in an amount of 0.01-10% (w/w), in said food.

A ninth aspect of the present invention provides the food or seasoning to which a kokumi taste has been imparted by any one of the methods as set forth above.

The present inventors have made extensive and intensive studies to achieve the above object, and as a result have found glycopeptides and peptides which have an excellent function of imparting a kokumi taste to foods (including seasonings). Hereinafter, this will be described in detail.

In their attempt to complete the present invention, a compound having a kokumi taste imparting function has been to be isolated from a seasoning which has a very strong function of imparting a kokumi taste when added to foods and drinks, in searching compound(s) having a taste improving effect such as impartment of a kokumi taste. Further, since there is a possibility that the amounts of compound(s) to be isolated are

very small, judgment according to organoleptic evaluation is also employed.


First, in selecting a starting raw material, the present

inventors have selected, as such starting raw material, a seasoning material which had a very strong function of imparting a kokumi taste when added even in a small amount to foods or drinks, and tried to isolate the target compound(s) from such a starting raw material. This is because, in conventional experiments for the purpose of isolating a kokumi taste-imparting compound, the function often becomes vague as fractionation progresses so that fractionation cannot be continued until the compound has been isolated. For these reasons, a seasoning material obtained by decomposing a material containing wheat protein with the use of a Koji -mold, was selected as the starting raw material as being suitable for achieving the object of the present invention. Since this seasoning material exhibits a kokumi taste imparting function when added in an amount of as small as 0.01% to foods or drinks, it is one of the most suitable starting raw materials for achieving the object of the present invention.

As the results of their extensive and intensive investigation, the present inventors have found that a compound having a kokumi taste imparting function in the raw material described above is also contained in a large amount in the fraction for a molecular weight of 1,000 or higher, and that not only peptide which has been conventionally said to be contained therein but also polysaccharide is contained in a large amount.

Therefore, the present inventors have further fractionated the fraction for a molecular weight of 1,000 or

higher, of the raw material described above by various procedures to separate-obtain compounds having a kokumi taste imparting function, followed by subjecting the compound to analysis and organoleptic evaluation. As a result, glycopeptides represented by the following sequence formula (I) or (II) were isolated as a compound having a kokumi taste imparting function. There is a report on only the sugar chain moieties of both of the glycopeptides (see J.Chromatogr., 434: pp. 51-60 (1988)). However, there has been no report that the sugar chain moieties have a flavoring function. Further, peptides which are common to both of the glycopeptides and are represented by the following sequence formula (III), (IV), (V) or (VI) are novel peptides, and there has been no report on the glycopeptide represented by the sequence formula (I) or (II) which is obtained by bonding the peptide to the sugar moiety. These glycopeptides are completely novel glycopeptides.



Sequence formula (III)

Val—Asn—His—Thr

Sequence formula (IV)

Val—Asn—His

Sequence formula (V)

Asn—His—Thr

Sequence formula (VI)

Asn—His

In the formulas, Val, Asn, His, and Thr are the same as those in the general formula (b) given later, and GlcNAc is an N-acetyl-glucosamine residue, Fuc is a fucose residue, Man is a mannose residue, and Xyl is a xylose residue.

It has been found, when these glycopeptides and peptides were subjected to organoleptic evaluation, that all the six were substantially tasteless in the form of an aqueous solution, but they exhibited the function of imparting a kokumi taste to the food when added even in a very small amount thereto. These findings have led the present invention to completion.

Accordingly, as has been described above, the present invention relates to glycopeptides and peptides which have a taste improving function, particularly a kokumi taste imparting function, specifically to such glycopeptides and peptides per se, a method for imparting the kokumi taste to foods or seasonings with the use of the glycopeptide(s) and/or the peptide(s), a method for improving foods in kokumi taste with the use of a seasoning containing the glycopeptide(s) and/or the peptide(s), and foods or seasonings to which the kokumi taste has been imparted by such a kokumi taste improving method.

In this connection, these glycopeptides and peptides are

collectively represented by the following general formula (a), and out of these, those represented by the following general formula (b) are particularly excellent in taste improving function.

General formula (a)

A—Asn—His—B



General formula (b)

Val—Asn—His—Thr



Best Mode for Carrying out the Invention

Hereinafter, the present invention will be described in detail.

First, the present invention relates to glycopeptides and peptides which are represented by the following general formula (a), and specific examples thereof include those represented by the sequence formula (I), (II), (III), (IV), (V) or (VI). As described above, these glycopeptides and peptides have a feature in that they have no taste to be specially mentioned in the form of an aqueous solution, but have a taste improving function such as impartment of a kokumi taste when added to food even at a concentration of 1 ppb with respect to the food.

General formula (a)

A—Asn—His—B



wherein Asn and His represent an asparagine residue and

a histidine residue, respectively, and X represents a hydrogen atom or a sugar chain, and A and B each represent independently a hydrogen atom or an optional amino acid residue.

In the present invention, the glycopeptide is one in which one or more sugars are bonded to the asparagine which is one of the amino acids constituting the peptide moiety, and the kind of sugar constituting the sugar chain is not particularly limited. The peptide per se of the present invention represented by the sequence formula (III), (IV), (V) or (VI) has a taste improving effect such as impartment of a kokumi taste, and the like, but the peptide has a feature in that the kokumi taste imparting function thereof is enhanced when the peptide is bonded to a sugar to form a glycopeptide.

The glycopeptides and peptides having the kokumi taste imparting function, of the present invention, should have at least a dipeptide structure of Asn-His in their structure, and a tripeptide and a tetrapeptide are in this order increased in taste enhancing capability.

Although the glycopeptides and the peptides of the present invention can be synthesized, they can be usually obtained from a hydrolyzate which has been obtained by enzymatically hydrolyzing a raw material containing protein.

Such a raw material containing protein is not particularly limited, and any protein and a raw material containing any protein can be used, as long as the hydrolyzate thereof contains the glycopeptide(s) or the peptide(s) represented by the general formula (a) or (b), or any one of

Sequence formulas (I) to (IV). For example, any protein such as vegetable protein, animal protein, or protein derived from yeast can be used. Examples of the vegetable protein include seed proteins such as wheat protein, soybean protein, corn protein, and the like, and the like. Among them, wheat protein is particularly suitable in terms of the abundance ratio of amino acids.

As an enzyme to be used for the hydrolysis of the protein according to the present invention, a commercially available enzyme preparation as well as a naturally-derived enzyme which has been metabolized using a microorganism or the like can be used as long as it can enzymatically decompose a protein as the raw material or the starting raw material containing the protein. These enzymes may be used alone or in combination of two or more thereof.

As for the pH and the reaction temperature used at the time when any protein or a starting raw material containing the protein is subjected to an enzymatic hydrolysis treatment, the optimum conditions for the enzyme to be used or conditions close thereto may be appropriately employed. The pH can be adjusted by adding an acid or an alkali which is acceptable for foods and drinks.

The treatment time that the protein or the starting raw material containing the protein should be subjected to the enzymatic hydrolysis varies depending on the conditions relating to the decomposition, such as the kind of enzyme to be used for the hydrolysis of the protein, the amount of the

enzyme to be used, the temperature, the pH, and the like. It is, however, preferably 20 to 100 hours. If the treatment time is much longer than necessary, there is a case that decomposition and browning unnecessarily occur, which may have an adverse influence on the quality of the hydrolyzate. Those skilled in the art can easily determine the preferred conditions for the hydrolysis treatment of a protein to obtain the glycopeptide or the peptide represented by any one of Sequence formulas (I) to (VI) by sampling the hydrolyzate from time to time during the hydrolysis treatment and then carrying out analysis.

The seasoning containing the glycopeptide(s) and/or the peptide(s) of the present invention can be obtained by collecting the liquid part of the raw material containing the enzymatically hydrolyzed protein by the use of ordinary procedures such as filtration, centrifugal separation or the like. Such a seasoning may be subjected to a decolorizing treatment using activated carbon, an ultrafilter, or the like, a separation and purification treatment using various chromatographic techniques or membrane separation using a permeable membrane, or the like, and a concentration treatment such as membrane concentration or vacuum concentration, or the like. The thus-treated seasoning may be used as a purified seasoning which has been subjected to decolorization, purification, concentration, and the like. In addition, it is possible to obtain a powdery seasoning having excellent storage stability without having been added with table salt or the like,

by spray-drying, freeze-vacuum drying or the like, such a seasoning containing the glycopeptide and/or the peptide of the present invention.

Further, in a case where it is necessary to obtain a glycopeptide and/or a peptide per se of the present invention, it can be isolated and purified by treating such above-mentioned hydrolyzate of a protein or a seasoning containing a glycopeptide and/or a peptide of the present invention, using known separation and purification procedures such as ultrafiltration, reverse osmosis, dialysis, normal phase HPLC, reversed phase HPLC, ion-exchange chromatography, gel filtration chromatography, affinity chromatography, and the like.

The seasoning containing a glycopeptide and/or a peptide of the present invention can impart not only a kokumi taste but also the tightness of aftertaste, ripe feeling, and the like, to foods and drinks when added thereto, and has the effect of improving the taste of the whole of a food or a drink without particularly adding a flavor-enhancing seasoning (an umami seasoning).

There are a variety of foods such as soups, processed foods and the like, in respect of which the taste improving effect of a glycopeptide and/or a peptide of the present invention, such as impartment of a kokumi taste, can be provided, and in particular, the effect can be conspicuously provided in respect of fermented seasonings or fermented foods. That is, the glycopeptide and/or the peptide of the present invention

exhibits conspicuously the effect on fermented seasonings or fermented foods such as soy sauce, soybean paste, cheese, and the like, and foods and drinks made therefrom, and can impart taste improving effects such as impartment of a kokumi taste e.g., thickness, spread, unity, and the like, as well as improvement of the tightness of aftertaste, to these foods.

The state of the glycopeptide and/or the peptide of the present invention at the time when they are added to foods is not limited, and they may be in a dry powder state, a paste state, a liquid state or the like. Further, since the glycopeptide and/or the peptide exhibits a kokumi taste imparting function as long as they are added to foods or drinks in such that the concentration thereof therein would be in the range of 1 ppb to 1,000 ppm by weight, preferably 1 ppb to 100 ppm by weight, upon eating, the glycopeptide and/or the peptide can, at any time, be added to foods or seasonings to realize the kokumi taste imparting effect. For example, the glycopeptide and/or the peptide can be added to a raw material before production or during production, or to a finished product after completion, or to a finished product at just before upon eating, during eating, or the like.

Examples

Hereinafter, the present invention will be described with reference to examples. However, the technical scope of the present invention is not limited to these examples.

<Example 1: Fractionation of Glycopeptide, Synthesis of
Peptide, and Kokumi taste Imparting Function>

500 g of wheat gluten "SWP-5A" (manufactured by Amylum) was added to, and sufficiently dispersed in, 2 L of water, and the resulting mixture was sterilized by heating at 120°C for 20 minutes, whereby a dispersion liquid of wheat gluten was prepared. On the other hand, 30 g of soybean protein "ESUSAN PROTEIN F" (manufactured by Ajinomoto Oil Mills, Inc.) was added to, and dispersed in, 2 L of water, and the resulting mixture was sterilized by heating at 120°C for 20 minutes, whereby a dispersion liquid of defatted soybean was prepared. To this dispersion liquid of defatted soybean, an *Aspergillus oryzae* pre-cultured in a culture medium was added in such an amount that it would be present in an amount of 1% (v/v), and the mixture was subjected to culturing in a fermenter jar at 30°C for 36 hours. 0.6 L of the *Aspergillus oryzae*-cultured broth was added to 2 L of the dispersion liquid of wheat gluten, and the resulting mixture was subjected to hydrolysis reaction in a fermenter jar at 36°C for 72 hours while being agitated with aeration. The thus obtained hydrolyzate was subjected to solid-liquid separation using a Nutsche funnel to obtain a filtrate. To the filtrate was then added 60 g of activated carbon. The resulting mixture was heated at 60°C for 10 minutes to decolorize the filtrate. The activated carbon was removed from the obtained decolorized liquid with a Nutsche funnel, and the filtrate was dried with a lyophilizer to obtain powdery, enzymatically

decomposed wheat gluten seasoning.

The thus obtained powdery, enzymatically-decomposed wheat gluten seasoning was dissolved in water, and the resulting aqueous solution was fractionated using an ultrafilter membrane "Prep/Scale-TFF PLAC 1K" (manufactured by MILLIPORE Corporation). The obtained fraction of a molecular weight of 1,000 or higher was further fractionated with the use of a gel filtration "Superdex 75 pg26/60" (manufactured by Amersham Biosciences) to obtain a fraction in respect of which an absorption of 280 nm was confirmed with a UV detector. This fraction was fractionated into Nos. 1 to 12 fractions by reversed phase HPLC (Capcellpack C-18 UG, manufactured by Shiseido Co., Ltd.). For the fractionation, two solvents A and B were used. Solvent A was a 0.05% by volume trifluoroacetic acid (TFA) aqueous solution, and Solvent B was a 0.05% by volume TFA 50% by volume acetonitrile aqueous solution. The column was equilibrated with Solvent A. After injecting a sample, the proportion of Solvent B was increased linearly to 100% in 50 min.

2 g of a commercially available chicken consommé soup (manufactured by Ajinomoto Co., Inc.) was added to 100 mL of water to prepare a chicken consommé soup, and then each of the fractions was added to the chicken consommé soup. The resulting mixtures were subjected to organoleptic evaluation, to confirm that there were two fractions which had a particularly strong effect of improving taste. These two fractions were each again fractionated by reversed phase HPLC ("Carbon-500" manufactured

by Tosoh Corporation). The following conditions were used to obtain two peaks at 44.8 minutes and 46.6 minutes. For the fractionation, two solvents A and B were used. Solvent A was a 0.05% by volume trifluoroacetic acid (TFA) aqueous solution, and Solvent B was a 0.05% by volume TFA 50% by volume acetonitrile aqueous solution. The column was equilibrated with Solvent A. After injecting a sample, the proportion of Solvent B was increased linearly to 100% in 167 min. The fractionated two peaks were analyzed with an LC-MS/MS (liquid chromatography-tandem mass spectrometry), to confirm that the former peak had fragments, that is, $[M+H]^+ = m/z$ 1640, 1494, 1478, 1362, 1332, 1316, 1200, 1184, 1038, 876, 819, 673, and 470, and that the latter peak had fragments, that is, $[M+H]^+ = m/z$ 1478, 1346, 1332, 1316, 1200, 1184, 1038, 876, 819, 673, and 470. Further, the former and latter peaks were analyzed by some known methods of analysis such as NMR, a peptide sequencer, an enzymatic decomposition treatment, and the like (R.Takahashi, "Biochemical Experiment Methods 23: Method for Studying Glycoprotein Sugar Chain", published by Gakkai Shuppan Center, 1989), to confirm that the peaks were due to the glycopeptides having the structures represented by Sequence formulas (I) and (II), respectively.

In these two glycopeptides, common peptide moieties (Sequence formulas (III), (IV), (V) and (VI)) existed. Therefore, in order to confirm the taste improving function of the peptides, those peptides represented by Sequence formulas (III), (IV), (V) and (VI) were obtained by synthesis using a

"433A Peptide Synthesizer" (manufactured by Applied Biosystems), followed by purifying by reversed phase HPLC ("Carbon-500", manufactured by Tosoh Corporation). For the fractionation, two solvents A and B were used. Solvent A was a 0.05% by volume trifluoroacetic acid (TFA) aqueous solution, and Solvent B was a 0.05% by volume TFA 50% by volume acetonitrile aqueous solution. The column was equilibrated with Solvent A. After injecting a sample, the proportion of Solvent B was increased linearly to 100% in 167 min.

In order to confirm the kokumi taste imparting function and the threshold value of the glycopeptides and the peptide represented by Sequence formulas (I) to (III) (hereinafter, the glycopeptides and the peptide represented by Ssequence formulas (I) to (III) may also be referred to as Glycopeptide I, Glycopeptide II, and Peptide III, respectively), Glycopeptide I and Glycopeptide II were each added to a chicken consommé soup in such that the concentration thereof would be 0.1 ppb to 1.0 ppm by weight upon eating, and Peptide III was added to the chicken consommé soup in such that the concentration thereof would be 0.1 ppb to 100 ppm by weight upon eating. As a control, a non-added chicken consommé soup was prepared. For each of the three glycopeptides and peptide, organoleptic evaluation was carried out by 16 taste panelists according to a paired comparison test (the panelists should choose one with a stronger kokumi taste). The results will be shown in the following Table 1. In the above organoleptic evaluation, there was a significant difference in the effect caused by adding

Glycopeptide I or II or Peptide III in the case where the concentration thereof was 1 ppb by weight or more upon eating. Therefore, the lower limit of the threshold value was set to 1 ppb by weight.

Table 1: Evaluation of chicken consommé soup (n=16)

Amount added	Comparative organoleptic evaluation test A		Comparative organoleptic evaluation test B		Comparative organoleptic evaluation test C	
	Non-added	Glyco-peptide I	Non-added	Glyco-peptide II	Non-added	Peptide III
0.1 ppb	4	12	5	11	7	9
1.0 ppb	2	14**	2	14**	3	13*
10 ppb	0	16***	0	16***	2	14**
1.0 ppm	0	16***	0	16***	0	16***
100 ppm	-	-	-	-	0	16***

(***: Significance level of 0.1%; **: Significance level of 1%; and *: Significance level of 5%.)

<Example 2: Taste Improving Function of Glycopeptides and Peptide>

In order to verify a taste improving function for solutions having five basic tastes, Glycopeptide I, Glycopeptide II, and Peptide III were respectively added in an amount of 100 ppb by weight to a 0.45% by weight sodium glutamate aqueous solution, a 0.05% by weight sodium inosinate aqueous solution, a 0.06% by weight sodium dl-tartarate aqueous solution, a 2.4% by weight saccharose aqueous solution, a 0.9% by weight sodium chloride aqueous solution, and a 0.1% by weight caffeine aqueous solution. Each resulting solution was subjected to organoleptic evaluation using non-added aqueous

solutions as the controls in the same manner as in the organoleptic evaluation of Example 1. As a result, it was found that all the three glycopeptides and peptide had the effect of improving sweet taste and sour taste, and that the two glycopeptides enhanced bitter taste.

<Example 3: Taste Improving Function for "Mentsuyu">

In order to verify a taste improving function for "Mentsuyu" (a soup for Japanese noodle) in the same manner as in the organoleptic evaluation of Example 1, the three glycopeptides and peptide were each added in an amount of 1 ppb by weight to Mentsuyu prepared according to the formulation shown in the following Table 2. Each resulting solution was subjected to organoleptic evaluation by 16 taste panelists according to a paired comparison test, using non-added Mentsuyu as the control. The results are shown in the following Table 3.

Table 2: Formulation of "Mentsuyu"

Materials	Composition (wt. %)
Soy sauce	15.0
Sugar	6.0
Soup stock of dried bonito (manufactured by Ajinomoto Co., Inc. under the product name of "Ichibandashi-katsuodashi")	4.0
Table Salt	1.5
Water	73.5
Total	100.0

Soy sauce, sugar, soup stock and table salt were mixed with, and dissolved in, water, and the resulting mixture was heated at 85°C for 1 hour.

Table 3: Evaluation of "Mentsuyu" (n=16)

	Comparative organoleptic evaluation test A		Comparative organoleptic evaluation test B		Comparative organoleptic evaluation test C	
	Non-added	Glyco-peptide I	Non-added	Glyco-peptide II	Non-added	Peptide III
Stronger "kokumi" taste	1	15***	1	15***	3	13*
Comments	Lack of unity	Taste became thick.	Lack of unity	Taste was spread.	Lack of unity	Taste was spread.

(***: Significance level of 0.1%; **: Significance level of 1%; and *: Significance level of 5%.)

<Example 4: Taste Improving Function for Cheese Sauce>

In order to verify a taste improving function for cheese sauce in the same manner as in the organoleptic evaluation of Example 1, the three glycopeptides and peptide were each added in an amount of 1 ppm by weight to a cheese sauce prepared according to the formulation shown in the following Table 4. Each of the sauces added with one of the three glycopeptides and peptide was subjected to organoleptic evaluation by 14 taste panelists according to a paired comparison test, using non-added cheese sauce as the control. The results are shown in the following Table 5.

Table 4: Formulation of
cheese sause

Materials	Composition (wt. %)
Cheese	50.0
Potato starch	2.0
Wheat flour	0.5
Water	47.5
Total	100.0

Heat-stirred until creamy, and then used.

Table 5: Evaluation of cheese sauce (n=14)

	Comparative organoleptic evaluation test A		Comparative organoleptic evaluation test B		Comparative organoleptic evaluation test C	
	Non-added	Glyco-peptide I	Non-added	Glyco-peptide II	Non-added	Peptide III
Stronger "kokumi" taste	0	14***	2	12*	2	12*
Comments	-	Aftertaste lasted.	-	Taste was expanded.	-	Taste was expanded.

(***: Significance level of 0.1%; **: Significance level of 1%; and *: Significance level of 5%.)

<Example 5: Taste Improving Function for Beef Extract>

In order to verify a taste improving function for a commercially available beef extract (manufactured by Bordon) in the same manner as in the organoleptic evaluation of Example 1, the three glycopeptides and peptide were each added in an amount of 10 ppb by weight to a beef extract solution formulated according to the mixing ratio shown in the following Table 6. Each resulting solution was subjected to organoleptic evaluation by 16 taste panelists according to a paired comparison test, using the non-added beef extract solution as

the control. The results are shown in the following Table 7.

Table 6: Formulation of beef extract solution

Materials	Composition (wt. %)
Beef extract (manufactured by Bordon)	2.0
Table salt	0.2
Hot water	97.8
Total	100.0

Stir-dissolved, and then used.

Table 7: Evaluation of beef extract solution (n=16)

	Comparative organoleptic evaluation test A		Comparative organoleptic evaluation test B		Comparative organoleptic evaluation test C	
	Non-added	Glyco- peptide I	Non-added	Glyco- peptide II	Non-added	Peptide III
Stronger "kokumi" taste	1	15***	3	13*	3	13*
Comments	Lack of thickness	Tightened in aftertaste	Lack of thickness	Taste was spread.	Lack of thickness	Taste was spread.

(***: Significance level of 0.1%; **: Significance level of 1%; and *: Significance level of 5%.)

<Example 6: Taste Improving Function for "Mentsuyu">

In order to verify the taste improving function of a liquid comprising enzymatically decomposed wheat gluten, that is, a seasoning containing the three glycopeptides and peptide, for Mentsuyu in the same manner as in the organoleptic evaluation of Example 1, the seasoning comprising enzymatically decomposed wheat gluten (that is, the decolorized liquid of Example 1 from which the activated carbon had been removed) was added in an amount of 0.1% by volume to Mentsuyu prepared according to the formulation shown in Table 2. Organoleptic evaluation was

carried out by 20 taste panelists according to a paired comparison test, using the non-added Mentsuyu as the control. The results are shown in the following Table 8.

Table 8: Evaluation of "mentsuyu" (n=20)

	Comparative organoleptic evaluation test	
	Non-added	Enzymatically decomposed wheat gluten liquid, added.
Stronger "kokumi" taste	0	20***
Comments	Lack of unity.	Became thick, Unity provided.

(***: Significance level of 0.1%;
 **: Significance level of 1%; and
 *: Significance level of 5%.)

<Example 7: Taste Improving Function for Cheese Sauce>

In order to verify the taste improving function of the liquid comprising enzymatically decomposed wheat gluten, that is, a seasoning containing the three glycopeptides and peptide, for cheese sauce in the same manner as in the organoleptic evaluation of Example 1, the seasoning comprising enzymatically decomposed wheat gluten (that is, the powdered seasoning of Example 1 formed from the decolorized liquid from which the activated carbon had been removed) was added in an amount of 0.05% by weight to a cheese sauce prepared according to the formulation shown in Table 4. Organoleptic evaluation was carried out by 20 taste panelists according to a paired comparison test, using the non-added cheese sauce as the control. The results are shown in the following Table 9.

Table 9: Evaluation of cheese sauce (n=20)

	Comparative organoleptic evaluation test	
	Non-added	Enzymatically decomposed wheat gluten liquid, added.
Stronger "kokumi" taste	0	20***
Comments	-	Taste was expanded. Titer was strengthened.

(***: Significance level of 0.1%;

**: Significance level of 1%; and

*: Significance level of 5%.)

<Example 8: Taste Improving Function for Beef Extract>

In order to verify the taste improving function of a liquid comprising enzymatically decomposed wheat gluten, that is, a seasoning containing the three glycopeptides and peptide, for a commercially available beef extract in the same manner as in the organoleptic evaluation of Example 1, the seasoning comprising enzymatically decomposed wheat gluten (that is, the powdered seasoning of Example 1 formed from the decolorized liquid from which the activated carbon had been removed) was added in an amount of 0.01% by weight to a beef extract solution formulated according to the mixing ratio shown in Table 6. Organoleptic evaluation was carried out by 16 taste panelists according to a paired comparison test, using the non-added beef extract solution as a control. The results are shown in the following Table 10.

Table 10: Evaluation of beef extract solution (n=16)

	Comparative organoleptic evaluation test	
	Non-added	Enzymatically decomposed wheat gluten liquid, added
Stronger "kokumi" taste	2	14**
Comments	Thickness-less	Taste was spread. Continuity was provided.

(***: Significance level of 0.1%;

**: Significance level of 1%; and

*: Significance level of 5%.)

<Example 9: Taste Improving Function for Tomato Soup>

In order to verify the taste improving function of a liquid comprising enzymatically decomposed wheat gluten, that is, a seasoning containing the three glycopeptides and peptide, for a tomato soup in the same manner as in the organoleptic evaluation of Example 1, the seasoning comprising enzymatically decomposed wheat gluten (that is, the powdered seasoning of Example 1 formed from the decolorized liquid from which the activated carbon had been removed) was added in an amount of 0.05% by weight to a tomato soup prepared according to the formulation shown in the following Table 11. Organoleptic evaluation was carried out by 16 taste panelists according to a paired comparison test, using the non-added tomato soup as the control. The results are shown in the following Table 12.

Table 11: Formulation of
Tomato soup

Materials	Composition (wt. %)
Tomato juice	60.00
Sugar	1.80
Skimmed milk	1.70
Potato starch	1.60
Table salt	0.60
MSG	0.30
Yeast extract	0.20
Onion powder	0.17
Garlic powder	0.01
Paprika powder	0.01
Carrot powder	0.01
water	33.60
Total	100.00

Dissolved and heat-stirred, and then used.

Table 12: Evaluation of Tomato soup (n=16)

	Comparative organoleptic evaluation test	
	Non-added	Enzymatically decomposed wheat gluten liquid, added
Stronger "kokumi" taste	1	15***
Comments	-	Unity and ripe feeling were increased.

(***: Significance level of 0.1%;

**: Significance level of 1%; and

*: Significance level of 5%.)

<Example 10: Taste Improving Function for Soybean Paste>

In order to verify the taste improving function of a seasoning comprising enzymatically decomposed wheat gluten,

that is, a seasoning containing the three glycopeptides and peptide, for soybean paste in the same manner as in the organoleptic evaluation of Example 1, the seasoning comprising enzymatically decomposed wheat gluten (that is, a powdered seasoning of Example 1 formed from the decolorized liquid from which the activated carbon had been removed) was added in an amount of 0.05% by weight to a soybean paste-flavored soup for Chinese noodle prepared according to the formulation shown in the following Table 13. Organoleptic evaluation was carried out by 13 taste panelists according to a paired comparison test, using the non-added soybean paste-flavored soup for Chinese noodle as the control. The results are shown in the following Table 14.

Table 13: Formulation of hot aqueous
soybean paste solution

Materials	Composition (wt. %)
Soybean paste	20.00
Soy sauce	1.50
Sugar	3.00
Table salt	0.80
MSG	2.20
Nucleic acid (IN)	0.04
Chicken extract	0.90
Pork extract	1.30
Garlic paste	0.89
Ginger paste	0.20
White pepper	0.05
Capsicum powder	0.05
Lard	4.40
Vegetable oil (sesame)	1.30
Hot water	63.36
Total	100.0

Used after stir-dissolved.

Table 14: Evaluation of hot aqueous soybean
paste solution (n=13)

	Comparative organoleptic evaluation test	
	Non-added	Enzymatically decomposed wheat gluten liquid, added
Stronger "kokumi" taste	0	13***
Comments	-	Unity and ripe feeling were provided.

(***: Significance level of 0.1%;

**: Significance level of 1%; and

*: Significance level of 5%.)

<Example 11: Taste Improving Function for Beef Extract>

In order to verify a taste improving function for a commercially available beef extract (manufactured by Bordon) in the same manner as in the organoleptic evaluation of Example 1, the glycopeptides and peptides listed in the following Table 15, were each added in an amount of 10 ppb, 1 ppm or 100 ppm by weight to beef extract solutions formulated according to the mixing ratio shown in Table 6 above. Each resulting solution was subjected to organoleptic evaluation by 14 taste panelists according to a paired comparison test, using the non-added beef extract solution as the control. The results are also shown in the following Table 15.

Incidentally, in the Table 15 indicating the results, Marks Δ , O and OO represent "Equal or inferior in the kokumi taste to the control", "Stronger in the kokumi taste than the control", and "Remarkably stronger in the kokumi taste than the control", respectively.

Table 15: Evaluation of beef extract

Samples	10 ppb	1 ppm	100 ppm
Glycopeptide I	OO	OO	OO
Peptide III (Val-Asn-His-Thr)	OO	OO	OO
Val-Asn-His	O	OO	OO
Asn-His-Thr	O	OO	OO
Val-Asn	Δ	Δ	Δ
Asn-His	O	O	OO
Val-Gln	Δ	Δ	Δ
His-Ser	Δ	Δ	Δ

Industrial Applicability

According to the present invention, it is possible to provide a glycopeptide and/or a peptide which have the function of imparting a kokumi taste to foods when added thereto.


Further, according to the present invention, it is possible to provide a seasoning containing the glycopeptide and/or the peptide, which has a strong effect of improving taste such as the function of imparting the kokumi taste to foods.

Furthermore, according to the present invention, it is possible to impart a kokumi taste to a food and improve the taste of foods by directly adding such a glycopeptide and/or peptide to foods or by adding a seasoning which contains such a glycopeptide and/or peptide to a food.

CLAIMS

1. A method of imparting a kokumi taste to a food or a seasoning which method comprises adding to said food or said seasoning, a glycopeptide or peptide having the chemical structure represented by the below-mentioned general formula (a) in such that said glycopeptide and/or peptide shall be contained in an amount of 1 ppb to 1,000 ppm (w/w), in said food or seasoning.


General formula (a)



wherein Asn and His represent an asparagine residue and a histidine residue, respectively, X represents a hydrogen atom or a sugar chain, and A and B each independently represent a hydrogen atom or an optional amino acid residue.

2. A method of imparting a kokumi taste to a food which comprises adding the seasoning as set forth in Claim 1 in such that it shall be contained in an amount of 0.01-10% (w/w), in said food.

3. A glycopeptide or peptide having the chemical structure represented by the following general formula (b):








wherein Val, Asn, His and Thr represent a valine residue, an asparagine residue, a histidine residue, and a threonine residue, respectively, X represents a hydrogen atom or a sugar chain.

4. A method of imparting a kokumi taste to a food or a seasoning which method comprises adding to said food or said seasoning, said glycopeptide and/or peptide as set forth in Claim 3 in such that said glycopeptide and/or peptide shall be contained in an amount of 1 ppb to 1,000 ppm (w/w), in said food or seasoning.

5. A method of imparting a kokumi taste to a food which comprises adding the seasoning as set forth in Claim 4 in such that it shall be contained in an amount of 0.01-10% (w/w), in said food.

6. A glycopeptide or peptide having the chemical structure represented by the following Sequence formula (I),

(II), (III), (IV), (V) or (VI):



wherein Val, Asn, His, and Thr have the same meanings as in the general formula (b) given above, and GlcNAc represents an N-acetyl-glucosamine residue, Fuc represents a fucose residue, Man represents a mannose residue, and Xyl represents a xylose residue.

7. A method of imparting a kokumi taste to a food or a seasoning which method comprises adding to said food or said seasoning, at least one of the glycopeptides and peptides having the chemical structure represented by the sequence formula (I), (II), (III), (IV), (V) or (VI) in such that said glycopeptide(s)

and/or peptide(s) shall be contained in an amount of 1 ppb to 1,000 ppm (w/w), in said food or seasoning.

8. A method of imparting a kokumi taste to a food which comprises adding the seasoning as set forth in Claim 7 in such that it shall be contained in an amount of 0.01-10% (w/w), in said food.

9. The food or seasoning to which a kokumi taste has been imparted by any one of the methods as set forth in Claims 1-2, 4-5 and 7-8.