

DESCRIPTION

AN OXIME DERIVATIVE AND PREPARATIONS THEREOF

5 TECHNICAL FIELD

The present invention relates to a novel oxime derivative having an excellent glucokinase activation effect, which is useful as a medicine.

BACKGROUND ART

10 Glucokinase (GK) is one of four hexokinases found in mammalian animals. The hexokinases catalyze a conversion of glucose into glucose-6-phosphate which is the first step of glucose metabolism. GK is localized mainly in hepatic parenchymal cells and pancreatic β cells, and plays an important role in whole body glucose homeostasis as a
15 rate-controlling enzyme for glucose metabolism in these cells. The hepatic and pancreatic forms of the enzyme are different in N-terminal 15 amino-acid sequence depending on the difference of each splicing, but are functionally indistinguishable.

Three hexokinases except GK are saturated in enzymatic activity
20 at a glucose concentration below 1 mM, but K_m of GK is 8 mM, which is within a physiological range of blood-glucose levels. Therefore, GK-mediated intracellular glucose metabolism is activated as the concentration of blood-glucose increases from normal level (5 mM) to postprandial level (10 to 15 mM).

25 A hypothesis that GK functions as a glucose sensor of pancreatic β cells and hepatocyte has been proposed (nonpatent document 1).

Thereafter, it has been clarified that GK actually plays a definitely important role in whole body glucose homeostasis according to the results of GK genetically-modified animal studies. GK KO mice die soon
30 after birth (nonpatent document 2), while both normal and diabetic

mice overexpressing GK showed lower glucose level than wild type animals (nonpatent document 3).

In maturity-onset diabetes of the young type II (MODY-2), which is one of the genetically determined diabetes, loss of function mutations in the GK has been found and it is thought that the low GK activity in MODY-2 results in hyperglycemia (nonpatent document 4). On the other hand, families having a GK mutation with increased enzymatic activity have been found and these people show hypoglycemia (nonpatent document 5). Accordingly, GK is believed to be a glucose sensor and to play an important role in maintenance of glucose homeostasis in human as well. It is expected that a GK activating compound has an insulinotropic action in β cells, an enhancing effect of glucose uptake in liver and inhibitory effect of hepatic output since such a compound activates a GK sensor system, and hence, it is believed that such a compound is useful for treating, for example, Type 2 diabetes.

Recently, it has been shown that a pancreatic β cell type glucokinase is distributed locally in feeding center (Ventromedial hypothalamus, VMH) in rat brain. About 20% of nerve cells in VMH are referred to as glucose responsive neurons and it has been thought from the past that they play important roles in controlling of body weights. An intracerebral administration of glucose in rat decreases food intake, but on the contrary, rat becomes overeating by an intracerebral administration of a glucose analog glucosamine, which cause the suppression of glucose metabolism. In electrophysiological experiments, glucose responsive neurons in VMH are stimulated when glucose increases from 5 to 20 mM, and the activity is blocked by glucosamine or the like (nonpatent document: Diabetes. 1999 Sep; 48(9): 1763-72). It is thought that a glucose sensor mechanism of VHM is similar to that of pancreatic β cells. Therefore, a GK activating

substance has a possibility of ameliorating obesity which is one of the major problems in Type 2 diabetes as well as correcting hyperglycemia.

Accordingly, a compound having a GK activation effect is useful as a treating and/or preventing agent of diabetes, or chronic
5 complication of diabetes such as retinopathy, nephropathy, neuropathy, ischemic heart disease or arteriosclerosis, or even obesity.

A compound having a GK activation effect includes, for example, pyridinecarboxylic acid derivatives (patent document 1), 2-pyridine-carboxamide derivatives (patent document 2), heteroarylcarbamoyl-
10 benzene derivatives (patent document 3), heteroaryl derivatives (patent document 4), substituted arylcyclopropylacetamide derivatives (patent document 5), 5-substituted pyrazine or pyridine derivatives (patent document 6), substituted (thiazol-2-yl)amide or sulfonamide derivatives (patent document 7), substituted phenylacetamide derivatives (patent
15 document 8) or amide derivatives (patent document 9).

A method for preparing a 5-substituted 2-aminothiazole, which is an intermediate for the oxime derivative of the present invention, has been described in patent documents 10 and 11, wherein 5-fluoro-2-aminothiazole hydrochloride is prepared by treating 5-bromo-2-
20 trifluoroacetyl aminothiazole derived from 5-bromo-2-aminothiazole hydrochloride with n-butyllithium, followed by treating the resultant with N-fluorobenzenesulfonylimide (patent document 10, Preparation 61; patent document 11, Preparation 21). It is also described in patent document 12 that 5-formyl-2-aminothiazole hydrobromide is prepared
25 by a reaction of bromomalonaldehyde with thiourea. However, the methods disclosed in patent document 10 and patent document 11 give the product in low yield and are not advantageous as an industrial method. Additionally, the method disclosed in patent document 12 gives 2-aminothiazole as a by-product which is difficult to remove, and
30 hence it is difficult to obtain the desired compound in a high purity.

Besides, said method can not be applied to preparations of wide range of 5-substituted 2-fluoro aminothiazoles other than 5-formyl-2-aminothiazole.

Compounds having an oxime structure therein have been described in patent documents 13 to 16 and nonpatent documents 6 to 8.

[patent document 1] WO05/044801

[patent document 2] WO04/081001

[patent document 3] WO04/076420

10 [patent document 4] WO04/063194

[patent document 5] WO04/063179

[patent document 6] WO04/052869

[patent document 7] WO04/050645

[patent document 8] WO03/095438

15 [patent document 9] WO03/055482

[patent document 10] WO04/072031

[patent document 11] WO04/072066

[patent document 12] US4,225,719

[patent document 13] WO05/023761

20 [patent document 14] WO01/012189

[patent document 15] WO00/026202

[patent document 16] WO96/023763

[nonpatent document 1] American Journal Physiology, volume 247 (3Pt2) 1984, p527-536

25 [nonpatent document 2] Cell, volume 83, 1995, p69-78

[nonpatent document 3] Proceedings of the National Academy of Sciences of the U.S.A., volume 93, 1996, p7225-7230

[nonpatent document 4] Nature Genetics, volume 356, 1992, p721-722

30 [nonpatent document 5] New England Journal of Medicine,

volume 338, 1998, p226-230

[nonpatent document 6] Bulletin des Societes Chimiques Belges (1994), 103(5-6), 213-18

[nonpatent document 7] Bulletin of the Chemical Society of Japan (1993), 66(8), 2335-8

[nonpatent document 8] Pharmazie (1988), 43(8), 535-6

DISCLOSURE OF INVENTION


The present invention provides a novel glucokinase activator, which is for the prophylaxis and/or treatment of diseases involving glucokinase, such as diabetes, complication associated with diabetes, or obesity.

The present invention also provides a novel compound having an excellent glucokinase activation effect which is useful as an active ingredient of a medicine.

According to extensive studies for problems to be solved by the present inventions, it has been found that an oxime derivative of the following formula has an excellent glucokinase activation effect, and the present invention has been completed.

The present invention includes the following embodiments.

(1) An oxime derivative of the general formula [I]:



wherein Ring A is aryl or heteroaryl;

Q is cycloalkyl, heterocycle, alkyl or alkenyl;

Ring T is heteroaryl or heterocycle;

R¹ is hydrogen atom, halogen atom, cycloalkylsulfonyl, alkylsulfonyl, alkylsulfinyl, alkylthio, or substituted or unsubstituted tetrazolyl;

5 R² is hydrogen atom, halogen atom, cycloalkylsulfonyl, substituted or unsubstituted alkylsulfonyl, substituted or unsubstituted alkylthio, nitro, substituted or unsubstituted amino, substituted or unsubstituted aminosulfonyl, substituted or unsubstituted heterocyclyl-thio, substituted or unsubstituted
10 heterocyclyl-sulfinyl, substituted or unsubstituted heterocyclyl-sulfonyl, substituted or unsubstituted heteroarylsulfonyl, alkenyloxy, substituted or unsubstituted alkoxy, substituted or unsubstituted alkylsulfinyl, substituted or unsubstituted heteroaryl, or substituted or unsubstituted heteroarylthio;

15 R³ and R⁴ are independently hydrogen atom, alkoxy, substituted or unsubstituted heterocycle, substituted or unsubstituted heteroaryl, alkoxyalkoxy, substituted or unsubstituted cycloalkyl, cyano, substituted or unsubstituted aryl, substituted or unsubstituted carbamoyl, hydroxy, alkanoyl, alkylthio, alkoxycarbonyl, substituted or
20 unsubstituted aryloxy, halogen atom, oxo, or substituted or unsubstituted arylcarbonyloxy;

R⁵ is hydrogen atom, formyl, halogen atom, oxo, substituted or unsubstituted alkoxy, substituted or unsubstituted aminosulfonyl, substituted or unsubstituted alkylthio, cyano, substituted or
25 unsubstituted heterocyclyl-sulfonyl, nitro, substituted or unsubstituted cycloalkyl, alkoxycarbonyl, alkenyl, alkylsulfonyl, substituted or unsubstituted carbamoyl, substituted or unsubstituted heteroarylthio, substituted or unsubstituted amino, carboxyl, substituted or unsubstituted heteroaryl, substituted or unsubstituted alkynyl,
30 substituted or unsubstituted heterocyclyl-carbonyl, substituted or

unsubstituted heterocyclyl-oxy, substituted or unsubstituted heterocycle, substituted or unsubstituted heterocyclyl-thio, substituted or unsubstituted cycloalkyloxy, alkanoyl, or substituted or unsubstituted alkyl;

5 R⁶ is hydrogen atom, substituted or unsubstituted alkyl, halogen atom, or carboxyl;

or a pharmaceutically acceptable salt thereof.

10 (2) The oxime derivative of (1) wherein Ring A is aryl or heteroaryl, provided that Ring A is not thiazolyl or thiadiazolyl, or a pharmaceutically acceptable salt thereof.

(3) The oxime derivative of (1) wherein Ring A is aryl, or a pharmaceutically acceptable salt thereof.

(4) The oxime derivative of (1) wherein Ring A is phenyl or pyridyl, or a pharmaceutically acceptable salt thereof.


15 (5) The oxime derivative of any one of (1) to (4) wherein Q is cycloalkyl, heterocycle or alkyl, or a pharmaceutically acceptable salt thereof.

(6) The oxime derivative of any one of (1) to (4) wherein Q is cycloalkyl or heterocycle, or a pharmaceutically acceptable salt thereof.

20 (7) The oxime derivative of any one of (1) to (4) wherein Q is heterocycle, or a pharmaceutically acceptable salt thereof.

(8) The oxime derivative of any one of (1) to (4) wherein Q is tetrahydrofuryl group, or a pharmaceutically acceptable salt thereof.


25 (9) The oxime derivative of any one of (1) to (8) wherein Ring T is heteroaryl or heterocycle of



or a pharmaceutically acceptable salt thereof.

(10) The oxime derivative of any one of (1) to (8) wherein Ring T

is heteroaryl of



or a pharmaceutically acceptable salt thereof.

(11) The oxime derivative of any one of (1) to (8) wherein Ring T is thiazolyl, thiazolopyridinyl, pyridyl, pyrazinyl, benzothiazolyl, quinolyl, thiadiazolyl, pyrazolyl, thiazolopyrazinyl, thiazolopyrimidinyl, cyclohexanothiazolyl or dihydrothiazolopyridinyl, or a pharmaceutically acceptable salt thereof.

(12) The oxime derivative of any one of (1) to (8) wherein Ring T is thiazolyl, thiazolopyridinyl, pyridyl, pyrazinyl, benzothiazolyl, thiadiazolyl, thiazolopyrazinyl, thiazolopyrimidinyl, cyclohexanothiazolyl or dihydrothiazolopyridinyl, or a pharmaceutically acceptable salt thereof.

(13) The oxime derivative of any one of (1) to (8) wherein Ring T is thiazolyl, thiazolopyridinyl, pyrazinyl, thiadiazolyl, thiazolopyrazinyl or thiazolopyrimidinyl, or a pharmaceutically acceptable salt thereof.

(14) The oxime derivative of any one of (1) to (8) wherein Ring T is thiazolyl or thiazolopyridinyl, or a pharmaceutically acceptable salt thereof.

(15) The oxime derivative of any one of (1) to (14) wherein R¹ is hydrogen atom or halogen atom, or a pharmaceutically acceptable salt thereof.

(16) The oxime derivative of any one of (1) to (14) wherein R¹ is hydrogen atom, or a pharmaceutically acceptable salt thereof.

(17) The oxime derivative of any one of (1) to (16) wherein R² is cycloalkylsulfonyl, substituted or unsubstituted alkylsulfonyl, substituted or unsubstituted alkylthio, nitro, substituted or unsubstituted amino, substituted or unsubstituted aminosulfonyl,

substituted or unsubstituted heterocyclyl-thio, substituted or unsubstituted heterocyclyl-sulfonyl, substituted or unsubstituted alkylsulfinyl, substituted or unsubstituted heteroarylsulfonyl, substituted or unsubstituted heteroarylthio, or substituted or unsubstituted heteroaryl, or a pharmaceutically acceptable salt thereof.

(18) The oxime derivative of any one of (1) to (16) wherein R^2 is cycloalkylsulfonyl, substituted or unsubstituted alkylsulfonyl, substituted or unsubstituted aminosulfonyl, or substituted or unsubstituted heterocyclyl-sulfonyl, or a pharmaceutically acceptable salt thereof.

(19) The oxime derivative of any one of (1) to (16) wherein R^2 is cycloalkylsulfonyl, substituted or unsubstituted alkylsulfonyl, or substituted or unsubstituted aminosulfonyl, or a pharmaceutically acceptable salt thereof.

(20) The oxime derivative of any one of (1) to (16) wherein R^2 is cycloalkylsulfonyl, substituted or unsubstituted aminosulfonyl, substituted or unsubstituted heterocyclyl-sulfonyl, or substituted or unsubstituted heteroarylsulfonyl, or a pharmaceutically acceptable salt thereof.

(21) The oxime derivative of any one of (1) to (16) wherein R^2 is cycloalkylsulfonyl, or a pharmaceutically acceptable salt thereof.

(22) The oxime derivative of any one of (1) to (20) wherein the substituent of the "substituted aminosulfonyl" in R^2 is substituted or unsubstituted alkyl, cycloalkyl, substituted or unsubstituted heterocycle, or alkoxy, or a pharmaceutically acceptable salt thereof.

(23) The oxime derivative of any one of (1) to (19) wherein the substituent of the "substituted alkylsulfonyl" in R^2 is alkoxy, or a pharmaceutically acceptable salt thereof.

(24) The oxime derivative of any one of (1) to (23) wherein R^3 and R^4 are independently hydrogen atom, alkoxy, substituted or

unsubstituted heterocycle, substituted or unsubstituted heteroaryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted aryl, substituted or unsubstituted carbamoyl, hydroxy, alkanoyl, alkylthio, substituted or unsubstituted aryloxy, halogen atom, oxo, or substituted or unsubstituted arylcarbonyloxy, or a pharmaceutically acceptable salt thereof.

(25) The oxime derivative of any one of (1) to (23) wherein R^3 and R^4 are independently hydrogen atom, alkoxy, substituted or unsubstituted heteroaryl, substituted or unsubstituted cycloalkyl, or hydroxy, or a pharmaceutically acceptable salt thereof.

(26) The oxime derivative of any one of (1) to (4) and (9) to (23) wherein the group of $-Q(R^3)(R^4)$ is cycloalkyl substituted with one or two groups selected from alkoxy and hydroxy, heterocycle, or alkyl substituted with 1 to 2 groups selected from hydroxy and substituted or unsubstituted heteroaryl, or a pharmaceutically acceptable salt thereof.

(27) The oxime derivative of any one of (1) to (26) wherein when Q is cycloalkyl, alkyl or alkenyl, then R^3 and R^4 are not any combination of two groups independently selected from hydrogen, alkoxy, cyano, substituted or unsubstituted aryl, hydroxy, alkylthio, alkoxycarbonyl, or halogen atom, or a pharmaceutically acceptable salt thereof.

(28) The oxime derivative of any one of (1) to (4) and (9) to (23) wherein Q is heterocycle and both of R^3 and R^4 are hydrogen atom, or a pharmaceutically acceptable salt thereof.

(29) The oxime derivative of any one of (1) to (28) wherein R^5 is hydrogen atom, formyl, halogen atom, oxo, substituted or unsubstituted alkoxy, substituted or unsubstituted aminosulfonyl, substituted or unsubstituted alkylthio, cyano, substituted or unsubstituted heterocyclyl-sulfonyl, nitro, substituted or unsubstituted cycloalkyl, alkoxycarbonyl, alkenyl, alkanoyl, substituted or unsubstituted carbamoyl, substituted or unsubstituted heteroarylthio,

substituted or unsubstituted amino, substituted or unsubstituted heteroaryl, substituted or unsubstituted alkynyl, substituted or unsubstituted heterocyclyl-oxy, or substituted or unsubstituted alkyl, or a pharmaceutically acceptable salt thereof.

5 (30) The oxime derivative of any one of (1) to (28) wherein R⁵ is hydrogen atom, halogen atom, substituted or unsubstituted alkoxy, substituted or unsubstituted alkylthio, cyano, substituted or unsubstituted cycloalkyl, alkanoyl, substituted or unsubstituted carbamoyl, substituted or unsubstituted amino, substituted or
10 unsubstituted heterocyclyl-oxy, or substituted or unsubstituted alkyl, or a pharmaceutically acceptable salt thereof.

(31) The oxime derivative of any one of (1) to (28) wherein R⁵ is halogen atom, substituted or unsubstituted alkoxy, substituted or unsubstituted alkylthio, substituted or unsubstituted amino,
15 substituted or unsubstituted heterocyclyl-oxy, or substituted or unsubstituted alkyl, or a pharmaceutically acceptable salt thereof.

(32) The oxime derivative of any one of (1) to (28) wherein R⁵ is substituted or unsubstituted alkoxy, substituted or unsubstituted amino, substituted or unsubstituted heterocyclyl-oxy, or substituted or
20 unsubstituted alkyl, or a pharmaceutically acceptable salt thereof.

(33) The oxime derivative of any one of (1) to (32) wherein the substituent of the "substituted alkyl" in R⁵ is substituted or unsubstituted heterocycle, substituted or unsubstituted amino, substituted or unsubstituted alkoxy, substituted or unsubstituted
25 carbamoyl, hydroxy, trialkylsilyloxy, alkylthio, alkylsulfonyl, substituted or unsubstituted heterocyclyl-oxy, heteroaryl, substituted or unsubstituted hydroxyimino, halogen atom, carboxyl, alkoxycarbonyl, or alkanoyloxy, or a pharmaceutically acceptable salt thereof.

(34) The oxime derivative of any one of (1) to (33) wherein R⁶ is
30 hydrogen atom, or substituted or unsubstituted alkyl, or a

pharmaceutically acceptable salt thereof.

(35) The oxime derivative of any one of (1) to (33) wherein R⁶ is hydrogen atom, or a pharmaceutically acceptable salt thereof.

(36) A pharmaceutical composition comprising a compound of any one of (1) to (35) or a pharmaceutically acceptable salt thereof as an active ingredient.

(37) A method for preventing or treating diabetes, or complication associated with diabetes including retinopathy, nephropathy, neuropathy, ischemic heart disease or arteriosclerosis, or obesity, which comprises administering an effective dose of a compound of any one of (1) to (35) or a pharmaceutically acceptable salt thereof.

(38) Use of a compound of any one of (1) to (35) or a pharmaceutically acceptable salt thereof in the manufacture of a medicament for treating or preventing diabetes, or complication associated with diabetes including retinopathy, nephropathy, neuropathy, ischemic heart disease or arteriosclerosis, or obesity.

The substituents on the group of each symbol in the compound [I] mean as defined below.

In R¹-R⁶ of the compound [I], substituents in "substituted amino", "substituted aminosulfonyl", "substituted aminoalkyl", "substituted aminoalkanoyl", "substituted carbamoyl", "substituted carbamoylalkyl", "substituted alkyl", "substituted alkylthio", "substituted alkylsulfinyl", "substituted alkylsulfonyl", "substituted alkoxy", "substituted alkanoyl", "substituted alkynyl", "substituted cycloalkyl", "substituted cycloalkyloxy", "substituted cycloalkylcarbonyl", "substituted cycloalkylsulfonyl", "substituted aryl", "substituted aryloxy", "substituted arylcarbonyl", "substituted arylcarbonyloxy", "substituted arylsulfonyl", "substituted arylalkylcarbonyl", "substituted heteroaryl", "substituted heteroarylthio", "substituted heteroarylsulfonyl", "substituted heteroarylalkyl", "substituted heterocycle", "substituted

heterocycl-yl-oxy", "substituted heterocycl-yl-carbonyl", "substituted heterocycl-yl-thio", "substituted heterocycl-yl-sulfinyl", "substituted heterocycl-yl-sulfonyl", "substituted hydroxyimino", and "substituted phenyl", "substituted pyridyl", "substituted thiazolopyridinyl", "substituted pyrazinyl", "substituted pyrazolyl", "substituted imidazolyl", "substituted thiazolyl", "substituted benzothiazolyl", "substituted quinolyl", "substituted thiadiazolyl", "substituted pyrazolyl", "substituted thiazolopyrazinyl", "substituted thiazolopyrimidinyl", "substituted cyclohexanothiazolyl", "substituted dihydrothiazolopyridinyl", "substituted triazolyl", "substituted pyrimidinyl", "substituted pyrrolidinyl", "substituted tetrahydrofuryl", "substituted thiacyclohexyl", "substituted cyclopentyl", "substituted piperazinyl", "substituted piperazinylsulfonyl", "substituted homopiperazinyl", "substituted piperidinyl", "substituted morpholinyl", "substituted thiomorpholinyl", "substituted perhydrodiazepinyl", and "substituted tetrazolyl" include those specifically indicated in EXAMPLES. Such substituents include (1) alkyl being optionally substituted with hydroxy, alkoxy, amino, mono- or di-alkylamino, carbamoyl, tetrahydrofuryl or pyridyl, (2) cycloalkyl, (3) hydroxy, (4) alkoxy, (5) cyano, (6) halogen atom, (7) mono- or di-alkylamino, (8) amino being optionally substituted with alkanoyl, alkoxyalkanoyl or alkoxycarbonyl, (9) pyridyl, (10) carboxyl, (11) formyl, (12) alkanoyl being optionally substituted with mono- or di-alkylamino, hydroxy, alkoxy or alkanoyloxy, (13) cycloalkylcarbonyl, (14) alkoxycarbonyl, (15) oxo, (16) alkylsulfonyl, or the like. The R¹-R⁶ groups may have the same or different 1 to 3 substituents selected from the above groups.

Additionally, each substituent is explained depending on each symbol (A, Q, T, R¹-R⁶) of the compound [I]. The groups of those symbols may have the same or different 1 to 3 substituents selected from the groups as defined below.

A preferable substituent of substituted tetrazolyl in R¹ includes alkyl.

A preferable substituent of substituted alkylsulfonyl in R² includes alkoxycarbonyl, alkoxy, cycloalkyl (preferably, cyclopropyl),
5 hydroxy, substituted or unsubstituted amino (substituent(s): 1 or 2 groups selected from alkyl, alkanoyl), substituted or unsubstituted heteroaryl (preferably, imidazolyl, triazolyl) (substituent(s): alkyl), alkylsulfonyl, cyano, substituted or unsubstituted heterocycle (preferably, tetrahydrofuryl, tetrahydropyranyl, dihydro-3H-isoindolyl)
10 (substituent(s): oxo, dioxo). More preferable one among them is alkoxy, cycloalkyl (preferably, cyclopropyl), hydroxy, particularly preferable one is alkoxy.

A preferable substituent of substituted alkylthio in R² includes alkoxy, cycloalkyl, alkoxycarbonyl, hydroxy, cyano, alkylthio,
15 substituted or unsubstituted heterocycle (preferably, tetrahydrofuryl, tetrahydropyranyl, dihydro-3H-isoindolyl) (substituent(s): oxo, dioxo), heteroaryl (preferably, pyridyl). More preferable one among them is alkoxy, cycloalkyl, alkoxycarbonyl, hydroxy, cyano, alkylthio, heteroaryl (preferably, pyridyl).

20 A preferable substituent of substituted amino in R² includes heteroarylcarbonyl (preferably, pyridylcarbonyl), heteroarylalkanoyl (thienylalkanoyl), cycloalkylcarbonyl, cycloalkylsulfonyl, alkoxy-carbonylcarbonyl, heteroarylsulfonyl, alkylsulfonyl. More preferable one among them is alkoxycarbonylcarbonyl, alkylsulfonyl.

25 A preferable substituent of the substituted alkyl which is the substituent of substituted aminosulfonyl in R² includes amino being optionally substituted with mono- or di-alkyl; carbamoyl being optionally substituted with mono- or di-alkyl; hydroxy; alkoxy; heteroaryl being optionally substituted with alkyl; cycloalkyl;
30 alkoxycarbonyl; hydroxyalkoxy; heterocycle being optionally substituted

with alkyl; halogen; alkylthio. More preferable one among them is amino being optionally substituted with mono- or di-alkyl; carbamoyl being optionally substituted with mono- or di-alkyl; hydroxy; alkoxy; cycloalkyl; alkoxycarbonyl; heterocycle being optionally substituted with alkyl; halogen atom; particularly hydroxy, alkoxy.

A preferable substituent of the substituted heterocycle which is a substituent of substituted aminosulfonyl in R² includes alkyl.

A preferable substituent of substituted heterocyclyl-thio in R² includes hydroxy; alkyl; oxo; alkanoyl; hydroxyalkyl; carbamoyl being optionally substituted with mono- or di-alkyl; heteroaryl; aminosulfonyl being optionally substituted with mono- or di-alkyl; amino being optionally substituted with mono- or di-alkyl; alkylsulfonyl; alkoxy; alkoxyalkyl. More preferable one among them is hydroxy; alkyl; carbamoyl being optionally substituted with mono- or di-alkyl; oxo; alkoxy; alkoxyalkyl; particularly alkyl.

A preferable substituent of substituted heterocyclyl-sulfinyl in R² includes hydroxy; alkyl; oxo; alkanoyl; hydroxyalkyl; carbamoyl being optionally substituted with mono- or di-alkyl; heteroaryl; aminosulfonyl being optionally substituted with mono- or di-alkyl; amino being optionally substituted with mono- or di-alkyl; alkylsulfonyl; alkoxy; alkoxyalkyl. More preferable one among them is hydroxy; alkyl; carbamoyl being optionally substituted with mono- or di-alkyl; oxo; alkoxy; alkoxyalkyl; particularly alkyl.

A preferable substituent of substituted heterocyclyl-sulfonyl in R² includes hydroxy; alkyl; oxo; alkanoyl; hydroxyalkyl; carbamoyl being optionally substituted with mono- or di-alkyl; heteroaryl; aminosulfonyl being optionally substituted with mono- or di-alkyl; amino being optionally substituted with mono- or di-alkyl; alkylsulfonyl; alkoxy; alkoxyalkyl. More preferable one among them is hydroxy; alkyl; carbamoyl being optionally substituted with mono- or di-alkyl; oxo;

alkoxy; alkoxyalkyl.

A preferable substituent of substituted heteroarylsulfonyl in R² includes alkyl.

5 A preferable substituent of substituted alkoxy in R² includes cycloalkyl.

10 A preferable substituent of substituted alkylsulfinyl in R² includes alkoxycarbonyl, alkoxy, alkoxyalkyl, cycloalkyl (preferably, cyclopropyl), hydroxy, substituted or unsubstituted amino (substituent(s): 1 or 2 groups selected from alkyl, alkanoyl), substituted or unsubstituted heteroaryl (preferably; imidazolyl, triazolyl) (substituent(s): alkyl), alkylsulfonyl, cyano, substituted or unsubstituted heterocycle (preferably, tetrahydrofuryl, tetrahydropyranyl, dihydro-3H-isoindolyl) (substituent(s): oxo, dioxo). More preferable one among them is alkoxy, cycloalkyl (preferably, cyclopropyl), hydroxy, particularly hydroxy.

15 A preferable substituent of substituted heteroaryl in R² includes alkyl.

A preferable substituent of substituted heterocycle in R³ and R⁴ includes alkoxycarbonyl, oxo, alkyl, alkanoyl.

20 A preferable substituent of substituted heteroaryl in R³ and R⁴ includes alkyl; amino being optionally substituted with mono- or di-alkyl. More preferable one among them is alkyl.

A preferable substituent of substituted cycloalkyl in R³ and R⁴ includes benzoyloxy, oxo, hydroxy, alkanoyl. More preferable one among them is oxo, hydroxy.

25 A preferable substituent of substituted aryl in R³ and R⁴ includes alkyl, cyano, halogen atom, alkoxy.

A preferable substituent of substituted carbamoyl in R³ and R⁴ includes alkyl.

30 A preferable substituent of substituted aryloxy in R³ and R⁴ includes alkyl, cyano, halogen atom, alkoxy.

A preferable substituent of substituted arylcarbonyloxy in R³ and R⁴ includes alkyl, cyano, halogen atom, alkoxy.

5 A preferable substituent of substituted alkoxy in R⁵ includes substituted or unsubstituted amino (substituent(s): 1 or 2 groups selected from alkyl, alkoxycarbonyl); alkoxycarbonyl; carbamoyl being optionally substituted with mono- or di-alkyl; carboxyl; hydroxy; substituted or unsubstituted heterocycle (substituent(s): oxo); trialkylsilyloxy; alkoxy. More preferable one is amino being optionally substituted with mono- or di-alkyl; carbamoyl being optionally substituted with mono- or di-alkyl; hydroxy; particularly amino being optionally substituted with mono- or di-alkyl; hydroxy.

A preferable substituent of substituted aminosulfonyl in R⁵ includes alkyl. Therefore, said substituent is mono-alkyl or di-alkyl, preferably di-alkyl.

15 A preferable substituent of substituted alkylthio in R⁵ includes amino being optionally substituted with mono- or di-alkyl; alkoxycarbonylamino; halogen atom; hydroxy; carboxyl; carbamoyl being optionally substituted with mono- or di-alkyl; alkoxycarbonyl. More preferable one among them is amino being optionally substituted with mono- or di-alkyl; alkoxycarbonylamino; hydroxy; carbamoyl being optionally substituted with mono- or di-alkyl; particularly dialkylcarbamoyl.

A preferable substituent of substituted heterocyclyl-sulfonyl in R⁵ includes alkyl.

25 A preferable substituent of substituted cycloalkyl in R⁵ includes amino being optionally substituted with mono- or di-alkyl.

A preferable substituent of substituted cycloalkyloxy in R⁵ includes amino being optionally substituted with mono- or di-alkyl.

30 A preferable substituent of substituted carbamoyl in R⁵ includes substituted or unsubstituted alkyl (substituent(s): 1 or 2 groups

selected from hydroxy; cycloalkyl; heterocycle; amino being optionally substituted with mono- or di-alkyl; heteroaryl), cycloalkyl, heteroaryl. More preferable one among them is substituted or unsubstituted alkyl (substituent(s): 1 or 2 groups selected from hydroxy, heterocycle, dialkylamino, heteroaryl), cycloalkyl.

A preferable substituent of substituted heteroarylthio in R⁵ includes alkyl.

A preferable substituent of substituted amino in R⁵ includes alkyl, substituted or unsubstituted aminoalkyl (substituent(s): 1 or 2 groups selected from alkyl, alkanoyl), alkanoyl, hydroxyalkyl, alkoxyalkyl. More preferable one among them is alkyl, and hence, mono-alkyl or di-alkyl, particularly di-alkyl.

A preferable substituent of substituted heteroaryl in R⁵ includes alkyl.

A preferable substituent of substituted alkynyl in R⁵ includes hydroxy, amino being optionally substituted with mono- or di-alkyl. More preferable one among them is hydroxy, dialkylamino.

A preferable substituent of substituted heterocyclyl-carbonyl in R⁵ includes hydroxy, alkyl, oxo, hydroxyalkyl, alkanoyl. More preferable one among them is hydroxy, alkyl, hydroxyalkyl.

A preferable substituent of substituted heterocyclyl-oxy in R⁵ includes hydroxy, alkyl, oxo, hydroxyalkyl, alkanoyl. More preferable one among them is alkyl, oxo.

A preferable substituent of substituted heterocycle in R⁵ includes hydroxy, alkyl, oxo, hydroxyalkyl, alkanoyl. More preferable one among them is oxo.

A preferable substituent of substituted heterocyclyl-thio in R⁵ includes hydroxy, alkyl, oxo, hydroxyalkyl, alkanoyl. More preferable one is alkyl, alkanoyl.

A preferable substituent of substituted alkyl in R⁵ includes

substituted or unsubstituted heterocycle, substituted or unsubstituted amino, substituted or unsubstituted alkoxy, substituted or unsubstituted carbamoyl, hydroxy, trialkylsilyloxy, substituted or unsubstituted alkylthio, substituted or unsubstituted heterocyclyl-oxy, heteroaryl, substituted or unsubstituted hydroxyimino, halogen atom, more preferably substituted or unsubstituted heterocycle, substituted or unsubstituted amino, substituted or unsubstituted alkoxy, hydroxy, substituted or unsubstituted alkylthio, substituted or unsubstituted heterocyclyl-oxy, substituted or unsubstituted hydroxyimino, halogen atom, further preferably substituted or unsubstituted heterocycle, substituted or unsubstituted alkoxy, substituted or unsubstituted heterocyclyl-oxy, particularly substituted or unsubstituted heterocycle, substituted or unsubstituted alkoxy, further particularly substituted or unsubstituted heterocycle.

A preferable substituent of substituted heterocycle which is the substituent of substituted alkyl in R⁵ includes alkyl; oxo; alkoxyalkanoyl; alkanoyl; alkoxy; alkanoylamino; cycloalkyl-carbonylamino; tri(halogeno)alkanoylamino; formylamino; alkoxy-carbonylamino; hydroxy; cycloalkylcarbonyl; tri(halogeno)alkyl; alkoxy carbonyl; formyl; amino being optionally substituted with mono- or di-alkyl; aminosulfonyl being optionally substituted with mono- or di-alkyl; alkylsulfonyl; heteroaryl; alkoxy carbonylalkyl; alkanoyloxyalkanoyl; alkoxy carbonylcarbonyl; aminoalkanoyl being optionally substituted with mono- or di-alkyl; substituted or unsubstituted carbamoyl (substituent(s): 1 or 2 groups selected from alkyl, alkoxy); hydroxyalkanoyl; di(halogeno)alkanoyl; substituted or unsubstituted heterocyclyl-carbonyl (substituent(s): oxo); substituted or unsubstituted hydroxyimino (substituent(s): alkoxy carbonyl); carboxyl; hydroxyalkoxy; alkoxyalkoxy; halogen atom; alkanoyloxy. More preferable one among them is alkyl; oxo; alkoxyalkanoyl; alkanoyl;

alkoxy; alkanoylamino; cycloalkylcarbonylamino;
tri(halogeno)alkanoylamino; formylamino; alkoxycarbonylamino;
cycloalkylcarbonyl; tri(halogeno)alkyl; alkoxycarbonyl; formyl; amino
being optionally substituted with mono- or di-alkyl; aminosulfonyl being
5 optionally substituted with mono- or di-alkyl; alkylsulfonyl; heteroaryl;
alkoxycarbonylalkyl; alkanoyloxyalkanoyl; alkoxycarbonylcarbonyl;
aminoalkanoyl being optionally substituted with mono- or di-alkyl;
carbamoyl being optionally substituted with mono- or di-alkyl;
hydroxyalkanoyl; di(halogeno)alkanoyl; substituted or unsubstituted
10 heterocyclyl-carbonyl (substituent(s): oxo); substituted or unsubstituted
hydroxyimino (substituent(s): alkoxycarbonyl); more preferably alkyl;
oxo; alkoxyalkanoyl; alkanoyl; formyl; amino being optionally
substituted with mono- or di-alkyl; alkylsulfonyl; alkanoyloxyalkanoyl;
aminoalkanoyl being optionally substituted with mono- or di-alkyl;
15 hydroxyalkanoyl; more preferably alkyl, alkanoyl, formyl,
hydroxyalkanoyl, particularly alkyl, alkanoyl.

A preferable substituent of the substituted amino which is the
substituent of substituted alkyl in R⁵ includes alkyl; carbamoylalkyl
being optionally substituted with mono- or di-alkyl; substituted or
20 unsubstituted aminoalkyl (substituent(s): 1 or 2 groups selected from
alkyl, alkanoyl); alkoxyalkyl; hydroxyalkyl; alkoxyalkanoyl; heteroaryl;
heteroarylalkyl. More preferable one among them is alkyl;
carbamoylalkyl being optionally substituted with mono- or di-alkyl;
aminoalkyl being optionally substituted with mono- or di-alkyl;
25 alkoxyalkyl; heteroaryl; particularly alkyl.

A preferable substituent of the substituted alkoxy which is the
substituent of substituted alkyl in R⁵ includes hydroxy, alkoxy.

A preferable substituent of the substituted carbamoyl which is
the substituent of substituted alkyl in R⁵ includes alkyl, alkoxy.

30 A preferable substituent of the substituted heterocyclyl-oxy which


is the substituent of substituted alkyl in R⁵ includes alkanoyl, alkyl, formyl, cycloalkylcarbonyl, alkoxyalkanoyl, alkylsulfonyl. More preferable one among them is alkanoyl, alkyl, particularly alkanoyl.

A preferable substituent of the substituted hydroxyimino which is the substituent of substituted alkyl in R⁵ includes alkoxycarbonyl.

Among the compounds [I] of the present invention, an example of preferable compounds are those in which R⁵ is substituted or unsubstituted alkyl.

Among the compounds [I], other preferable compounds are those of formula [I-A] as shown below, and the present invention includes also the following embodiments:

(1) An oxime derivative of the general formula [I-A]:



wherein Ring A is aryl or heteroaryl;

Q is cycloalkyl, heterocycle, alkyl or alkenyl;

Ring T is heteroaryl or heterocycle;

R¹ and R² are independently hydrogen atom, halogen atom, cycloalkylsulfonyl, alkylsulfonyl, alkylsulfinyl, alkylthio, or substituted or unsubstituted tetrazolyl;

R³ and R⁴ are independently hydrogen atom, hydroxy, oxo, halogen atom, cyano, alkylthio, alkoxy, alkanoyl, alkoxyalkoxy, alkoxycarbonyl, substituted or unsubstituted carbamoyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted heterocycle, substituted or unsubstituted

cycloalkyl, or substituted or unsubstituted aryloxy;

R⁵ is hydrogen atom, halogen atom, cyano, nitro, tetrazolyl, oxo, cycloalkyl, alkenyl, alkylthio, alkylsulfonyl, alkoxy, formyl, alkanoyl, alkoxycarbonyl, substituted or unsubstituted carbamoyl, substituted or unsubstituted aminosulfonyl, substituted or unsubstituted heterocyclyl-carbonyl, substituted or unsubstituted heterocyclyl-sulfonyl, or substituted or unsubstituted alkyl; or a pharmaceutically acceptable salt thereof.

(2) The oxime derivative of (1) wherein Ring A is phenyl or pyridyl, or a pharmaceutically acceptable salt thereof.

(3) The oxime derivative of (1) wherein Ring A is phenyl, or a pharmaceutically acceptable salt thereof.

(4) The oxime derivative of any one of (1) to (3) wherein Q is cycloalkyl, tetrahydrofuryl, alkyl or alkenyl, or a pharmaceutically acceptable salt thereof.

(5) The oxime derivative of any one of (1) to (3) wherein Q is tetrahydrofuryl, or a pharmaceutically acceptable salt thereof.

(6) The oxime derivative of any one of (1) to (3) wherein Q is (3R)-3-tetrahydrofuryl, or a pharmaceutically acceptable salt thereof.

(7) The oxime derivative of any one of (1) to (6) wherein Ring T is thiazolyl, pyrazinyl, thiadiazolyl, thiazolopyridinyl, benzothiazolyl, cyclohexanothiazolyl or dihydrothiazolopyridinyl, or a pharmaceutically acceptable salt thereof.

(8) The oxime derivative of any one of (1) to (6) wherein Ring T is thiazolyl, or a pharmaceutically acceptable salt thereof.

(9) The oxime derivative of any one of (1) to (6) wherein Ring T is 2-thiazolyl, or a pharmaceutically acceptable salt thereof.

(10) The oxime derivative of any one of (1) to (9) wherein one of R¹ and R² is hydrogen atom and the other is cycloalkylsulfonyl, alkylsulfonyl, alkylsulfinyl, alkylthio, or substituted or unsubstituted

tetrazolyl, or a pharmaceutically acceptable salt thereof.

(11) The oxime derivative of any one of (1) to (9) wherein one of R^1 and R^2 is hydrogen atom and the other is cycloalkylsulfonyl, or a pharmaceutically acceptable salt thereof.

5 (12) The oxime derivative of (11) wherein Ring A is phenyl and the cycloalkylsulfonyl is substituted to the 4-position of the phenyl, or a pharmaceutically acceptable salt thereof.

(13) The oxime derivative of any one of (1) to (12) wherein R^3 and R^4 are independently hydrogen atom, hydroxy, alkoxy, substituted or
10 unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted heterocycle, or substituted or unsubstituted cycloalkyl, or a pharmaceutically acceptable salt thereof.

(14) The oxime derivative of any one of (1) to (12) wherein R^3 and R^4 are independently hydrogen atom, hydroxy, alkoxy, substituted or
15 unsubstituted phenyl, substituted or unsubstituted pyrazolyl, substituted or unsubstituted imidazolyl, substituted or unsubstituted thiazolyl, substituted or unsubstituted triazolyl, substituted or unsubstituted pyridyl, substituted or unsubstituted pyrimidinyl, substituted or unsubstituted pyrrolidinyl, substituted or unsubstituted
20 thiacyclohexyl, or substituted or unsubstituted cyclopentyl, or a pharmaceutically acceptable salt thereof.

(15) The oxime derivative of any one of (1) to (12) wherein both of R^3 and R^4 are hydrogen atom, or a pharmaceutically acceptable salt thereof.

25 (16) The oxime derivative of any one of (1) to (15) wherein R^5 is hydrogen atom, halogen atom, cyano, oxo, alkenyl, alkylthio, formyl, alkanoyl, substituted or unsubstituted carbamoyl, substituted or unsubstituted aminosulfonyl, substituted or unsubstituted heterocyclyl-sulfonyl, or substituted or unsubstituted alkyl, or a
30 pharmaceutically acceptable salt thereof.

(17) The oxime derivative of any one of (1) to (16) wherein a substituent of the substituted alkyl in R⁵ is 1 to 3 substituents selected from substituted or unsubstituted amino, substituted or unsubstituted hydroxyimino, hydroxy, alkoxy, halogen atom, carboxyl, alkoxycarbonyl, substituted or unsubstituted carbamoyl, alkanoyloxy, and substituted or unsubstituted heterocycle, or a pharmaceutically acceptable salt thereof.

(18) The oxime derivative of any one of (1) to (15) wherein R⁵ is hydrogen atom, halogen atom, cyano, oxo, alkenyl, alkylthio, formyl, alkanoyl, substituted or unsubstituted carbamoyl, substituted or unsubstituted aminosulfonyl, substituted or unsubstituted piperazinylsulfonyl, or alkyl, or alkyl substituted with 1 to 3 groups selected from substituted or unsubstituted amino, substituted or unsubstituted hydroxyimino, hydroxy, alkoxy, halogen atom, alkoxycarbonyl, substituted or unsubstituted piperazinyl, substituted or unsubstituted homopiperazinyl, substituted or unsubstituted piperidinyl, substituted or unsubstituted morpholinyl and substituted or unsubstituted thiomorpholinyl, or a pharmaceutically acceptable salt thereof.

(19) The oxime derivative of any one of (1) to (15) wherein R⁵ is fluorine atom, or alkyl substituted with 1 to 3 groups selected from substituted or unsubstituted piperazinyl, substituted or unsubstituted morpholinyl and substituted or unsubstituted thiomorpholinyl, or a pharmaceutically acceptable salt thereof.

(20) The oxime derivative of any one of (1) to (15) wherein R⁵ is fluorine atom, or alkyl substituted with piperazinyl being optionally substituted with 1 to 3 substituents selected from alkyl, oxo, alkanoyl and alkoxyalkanoyl, or a pharmaceutically acceptable salt thereof.

(21) The oxime derivative of any one of (1) to (15) wherein R⁵ is fluorine atom, or piperazinylmethyl being optionally substituted with

alkyl or oxo on the carbon or being optionally substituted with alkyl, alkanoyl or alkoxyalkanoyl on the nitrogen, or a pharmaceutically acceptable salt thereof.

(22) A medicine comprising as an active ingredient the oxime derivative of any one of (1) to (21) or a pharmaceutically acceptable salt thereof.

(23) A glucokinase activating agent comprising as an active ingredient the oxime derivative of any one of (1) to (21) or a pharmaceutically acceptable salt thereof as an active ingredient.

The substituents on the group of each symbol of the compound [I-A] means as defined below.

A substituent on the substituted alkyl in R⁵ includes substituted or unsubstituted amino, substituted or unsubstituted hydroxyimino, hydroxy, alkoxy, halogen atom, carboxyl, alkoxycarbonyl, substituted or unsubstituted carbamoyl, alkanoyloxy, substituted or unsubstituted heterocycle, preferably substituted or unsubstituted amino, substituted or unsubstituted hydroxyimino, hydroxy, alkoxy, halogen atom, alkoxycarbonyl, substituted or unsubstituted piperazinyl, substituted or unsubstituted homopiperazinyl, substituted or unsubstituted piperidinyl, substituted or unsubstituted morpholinyl, substituted or unsubstituted thiomorpholinyl, or the like. The alkyl is substituted with the same or different 1 to 3 substituents selected from the above groups.

In the compound [I-A], substituents in "substituted aryl", "substituted aryloxy", "substituted heteroaryl", "substituted heterocycle", "substituted heterocyclyl-carbonyl", "substituted heterocyclyl-sulfonyl", "substituted cycloalkyl", "substituted phenyl", "substituted pyrazolyl", "substituted imidazolyl", "substituted thiazolyl", "substituted triazolyl", "substituted pyridyl", "substituted pyrimidinyl", "substituted pyrrolidinyl", "substituted thiacyclohexyl", "substituted cyclopentyl",

"substituted piperazinyl", "substituted piperazinylsulfonyl", "substituted homopiperazinyl", "substituted piperidinyl", "substituted morpholinyl", "substituted thiomorpholinyl", "substituted tetrazolyl", "substituted carbamoyl", "substituted aminosulfonyl", "substituted amino" or "substituted hydroxyimino" are the same or different 1 to 3 substituents selected from those groups which include (1) alkyl being optionally substituted with hydroxy, alkoxy, amino, mono- or di-alkylamino, carbamoyl, tetrahydrofuryl or pyridyl, (2) cycloalkyl, (3) hydroxy, (4) alkoxy, (5) cyano, (6) halogen, (7) mono- or di-alkylamino, (8) amino being optionally substituted with alkanoyl, alkoxyalkanoyl or alkoxycarbonyl, (9) pyridyl, (10) carboxyl, (11) formyl, (12) alkanoyl being optionally substituted with mono- or di-alkylamino, hydroxy, alkoxy or alkanoyloxy, (13) cycloalkylcarbonyl, (14) alkoxycarbonyl, (15) oxo, (16) alkylsulfonyl, or the like.

In the compound [I-A], a preferable substituent in substituted tetrazolyl on R¹ and R² includes alkyl or the like.

A preferable substituent in substituted carbamoyl in R³ and R⁴ includes alkyl or the like, and it may be the same or different 1 to 2 groups.

A preferable substituent in substituted aryl, substituted aryloxy and substituted phenyl on R³ and R⁴ includes cyano, halogen atom, alkoxy, alkyl, mono- or di-alkylamino or the like, particularly cyano or halogen atom. The substituent may be the same or different 1 to 3 groups selected from these groups.

A preferable substituent in substituted heteroaryl, substituted pyrazolyl, substituted imidazolyl, substituted thiazolyl, substituted triazolyl, substituted pyridyl and substituted pyrimidinyl on R³ and R⁴ includes alkyl, mono- or di-alkylamino or the like, particularly alkyl. The substituent may be the same or different 1 to 2 groups selected from these groups.

A preferable substituent in substituted heterocycle, substituted pyrrolidinyl and substituted thiacyclohexyl on R³ and R⁴ includes oxo, alkoxy carbonyl, alkyl, alkanoyl or the like, particularly oxo or alkyl. The substituent may be the same or different 1 to 2 groups selected from these groups.

A preferable substituent in substituted cycloalkyl and substituted cyclopentyl in R³ and R⁴ includes oxo, hydroxy or the like, particularly hydroxy.

A preferable substituent in substituted carbamoyl in R⁵ includes alkoxy, alkyl, cycloalkyl, hydroxyalkyl, dialkylaminoalkyl, cycloalkyl, tetrahydrofurylalkyl, pyridylalkyl, alkoxy, pyridyl or the like, particularly hydroxyalkyl, dialkylaminoalkyl, pyridylalkyl, pyridyl or the like. The substituent may be the same or different 1 to 2 groups selected from these groups.

A preferable substituent in substituted aminosulfonyl in R⁵ includes alkyl or the like, and the substituent may be the same or different 1 to 2 groups selected from these groups.

A preferable substituent in substituted heterocycle, substituted heterocyclyl-carbonyl, substituted heterocyclyl-sulfonyl, substituted piperazinyl, substituted piperazinylsulfonyl, substituted homopiperazinyl, substituted piperidinyl, substituted morpholinyl and substituted thiomorpholinyl in R⁵ includes alkoxy carbonylamino, hydroxy, hydroxyalkyl, alkanoylamino, alkoxyalkanoylamino, oxo, alkyl, formyl, alkanoyl, hydroxyalkanoyl, cycloalkylcarbonyl, carboxyl, alkoxy carbonyl, alkoxyalkanoyl, alkanoyloxyalkanoyl, mono- or di-alkylaminoalkanoyl, alkylsulfonyl or the like, particularly oxo, alkyl, formyl, alkanoyl, hydroxyalkanoyl, cycloalkylcarbonyl, alkoxy carbonyl, alkoxyalkanoyl, alkanoyloxyalkanoyl or alkylsulfonyl. The substituent may be the same or different 1 to 3 groups selected from these groups.

A preferable substituent in substituted amino in R⁵ includes alkyl,

alkoxyalkyl, pyridyl, pyridylalkyl, dialkylaminoalkyl, carbamoylalkyl or the like. The substituent may be the same or different 1 to 2 groups selected from these groups.

5 A preferable substituent in substituted hydroxyimino in R⁵ includes alkoxycarbonyl or the like.

In the compound [I-A], the aryl in Ring A includes preferably phenyl.

The heteroaryl in Ring A includes preferably thienyl or pyridyl, particularly pyridyl.

10 The heterocycle in Q includes, for example, 5 to 6-membered monocyclic heterocycle, specifically tetrahydrofuryl, pyrrolidinyl, tetrahydropyranyl, thiacyclohexyl, piperidinyl or the like, particularly tetrahydrofuryl.

15 The heteroaryl in Ring T includes, for example, 5 to 9-membered monocyclic or bicyclic aromatic ring, specifically thiazolyl, thiadiazolyl, pyridyl, pyrazinyl, benzothiazolyl, thiazolopyridinyl or the like. The heterocycle in Ring T includes, for example, 9-membered bicyclic aromatic ring, specifically cyclohexanothiazolyl, dihydrothiazolopyridinyl or the like.

20 The aryl in R³ and R⁴ includes preferably phenyl.

The heteroaryl in R³ and R⁴ includes, for example, 5 to 6-membered monocyclic aromatic ring, specifically pyrazolyl, imidazolyl, isoxazolyl, thiazolyl, triazolyl, pyridyl, pyrimidinyl or the like.

25 The heterocycle in R³ and R⁴ includes, for example, 5 to 6-membered monocyclic heterocycle, specifically pyrrolidinyl, tetrahydrofuryl, dioxolanyl, piperidinyl, thiacyclohexyl or the like.

The cycloalkyl in R³ and R⁴ includes preferably 3 to 6-membered cycloalkyl, specifically cyclopropyl or cyclopentyl.

30 The heterocycle in R⁵ includes, for example, 4 to 6-membered monocyclic heterocycle, specifically azetidiny, pyrrolidinyl, piperidinyl,

piperazinyl, morpholinyl, thiomorpholinyl or the like.

Among the compound [I-A], a preferable compound includes a compound wherein Ring A is phenyl, Q is 3-tetrahydrofuryl, Ring T is 2-thiazolyl, one of R¹ and R² is hydrogen atom, the other is
5 cyclopropylsulfonyl, both of R³ and R⁴ are hydrogen atom, R⁵ is piperazinyl substituted alkyl being optionally substituted with 1 to 3 substituents selected from alkyl, oxo, alkanoyl and alkoxyalkanoyl.

Other preferable compound among the compounds [I] of the present invention includes a compound described in any of all
10 EXAMPLES.

The following terms used herein mean as defined below.

A "halogen atom" includes fluorine atom, chlorine atom, bromine atom or iodine atom, preferably fluorine atom or chlorine atom.

An "alkyl", which includes "alkyl" moiety in a group bound with
15 other groups such as "alkylthio" or "hydroxyalkyl" (the same for other groups defined hereinafter), includes, for example, straight- or branched-chain alkyl of C₁₋₆, preferably C₁₋₄, specifically methyl, ethyl, propyl, isopropyl, isobutyl, tert-butyl, pentyl, hexyl or the like.

An "alkenyl" includes, for example, straight- or branched-chain
20 alkenyl of C₂₋₆, preferably C₂₋₄, specifically vinyl, propenyl, isopropenyl, butenyl, pentenyl, hexenyl or the like.

An "alkynyl" includes, for example, straight- or branched-chain alkynyl of C₂₋₆, preferably C₂₋₄, specifically acetylenyl, propynyl, butynyl, pentynyl, hexynyl or the like.

25 An "alkoxy" includes, for example, straight- or branched-chain alkoxy of C₁₋₆, preferably C₁₋₄, specifically methoxy, ethoxy, propoxy, isopropoxy, butoxy, tert-butoxy, pentyloxy, hexyloxy or the like.

An "alkanoyl" includes, for example, straight- or branched-chain alkanoyl of C₂₋₇, preferably C₂₋₅, specifically acetyl, propionyl, butyryl,
30 pentanoyl or the like.

A "cycloalkyl" includes, for example, cycloalkyl of C₃₋₈, preferably C₃₋₆, specifically cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl or cycloheptyl.

5 An "aryl" includes 6 to 14-membered, preferably 6 to 10-membered monocyclic, bicyclic or tricyclic aromatic hydrocarbon, specifically phenyl, naphthyl, phenanthryl, anthryl or the like, preferably phenyl in particular.

10 A "heteroaryl" includes 4 to 10-membered, preferably 5 to 9-membered, monocyclic or bicyclic aromatic hydrocarbon wherein 1 to 3 carbon atoms are substituted with heteroatoms independently selected from oxygen atom, sulfur atom and nitrogen atom, specifically thienyl, thiazolyl, pyrazolyl, imidazolyl, isoxazolyl, triazolyl, thiadiazolyl, pyridyl, pyrimidinyl, pyrazinyl, quinolyl, benzothiazolyl, thiazolopyridinyl, thiazolopyrazinyl, thiazolopyrimidinyl or the like.

15 A "heterocycle" includes 4 to 10-membered, preferably 4 to 9-membered, monocyclic or bicyclic non-aromatic hydrocarbon wherein 1 to 3 carbon atoms are substituted with heteroatoms independently selected from oxygen atom, sulfur atom and nitrogen atom, specifically oxetanyl, azetidiny, pyrrolidinyl, tetrahydrofuryl, dioxolanyl, 20 piperidinyl, piperazinyl, homopiperazinyl, tetrahydropyranyl, thiacyclohexyl, morpholinyl, thiomorpholinyl, cyclohexanothiazolyl, dihydrothiazolopyridinyl, tetrahydrothiazolopyridinyl or the like.

Alternatively, illustrative embodiments of "halogen atom", "alkyl", "alkenyl", "alkynyl", "alkoxy", "alkanoyl", "cycloalkyl", "aryl", "heteroaryl", 25 "heterocycle" include those specifically indicated in EXAMPLES.

Additionally, each term is explained depending on each symbol (A, Q, T, R¹-R⁶) of the compound [I].

A preferable "aryl" in Ring A includes phenyl.

30 A preferable "heteroaryl" in Ring A includes thienyl, pyridyl, particularly pyridyl.

A preferable "cycloalkyl" in Q includes, for example, 5 to 6-membered monocyclic cycloalkyl, specifically cyclopentyl, cyclohexyl or the like, particularly cyclopentyl.

A preferable "heterocycle" in Q includes, for example, 4 to 6-membered monocyclic heterocycle optionally having 1 to 3 heteroatoms independently selected from oxygen atom, sulfur atom and nitrogen atom, specifically oxetanyl, tetrahydrofuryl, pyrrolidinyl, piperidinyl, tetrahydropyranyl, tetrahydrothiopyranyl or the like, particularly tetrahydrofuryl.

A "heteroaryl" in Ring T includes, for example, 5 to 9-membered monocyclic, bicyclic heteroaryl optionally having 1 to 3 heteroatoms independently selected from oxygen atom, sulfur atom and nitrogen atom, specifically thiazolyl, pyrazolyl, thiadiazolyl, pyridyl, pyrazinyl, benzothiazolyl, thiazolopyridinyl, thiazolopyrazinyl, thiazolopyrimidinyl, quinolyl or the like. A preferable one among them is thiazolyl, thiadiazolyl, pyridyl, pyrazinyl, benzothiazolyl, thiazolopyridinyl, thiazolopyrazinyl, thiazolopyrimidinyl, more preferably thiazolyl, thiadiazolyl, pyrazinyl, thiazolopyridinyl, thiazolopyrazinyl, particularly thiazolyl, thiazolopyridinyl, further particularly thiazolyl.

A "heterocycle" in Ring T includes, for example, 5 to 9-membered monocyclic, bicyclic heterocycle optionally having 1 to 3 heteroatoms independently selected from oxygen atom, sulfur atom and nitrogen atom, preferably 9-membered bicyclic heterocycle, specifically cyclohexanethiazolyl, dihydrothiazolopyridinyl or the like.

A "cycloalkyl" of cycloalkylsulfonyl in R² includes, for example, 3 to 4-membered cycloalkyl, specifically cyclopropyl, cyclobutyl or the like, preferably cyclopropyl in particular.

A "heterocycle" which is the substituent of the substituted or unsubstituted alkyl which is the substituent of substituted or unsubstituted aminosulfonyl in R² includes, for example, 5 to 9-

membered monocyclic, bicyclic heterocycle optionally having 1 to 3 heteroatoms independently selected from oxygen atom, sulfur atom and nitrogen atom, preferably 5-membered monocyclic heterocycle. Particularly, tetrahydrofuryl is preferable.

5 A "heterocycle" of substituted or unsubstituted heterocycl-
sulfonyl in R² includes, for example, 5 to 9-membered monocyclic,
bicyclic heterocycle optionally having 1 to 3 heteroatoms independently
selected from oxygen atom, sulfur atom and nitrogen atom, preferably 5
to 7-membered monocyclic heterocycle, specifically azetidiny, 10
pyrrolidiny, tetrahydrofuryl, piperidiny, piperaziny, morpholiny,
thiomorpholiny, oxazepidiny, perhydrodiazepiny.

A "heteroaryl" of substituted or unsubstituted heteroaryl in R³
and R⁴ includes, for example, 5 to 9-membered monocyclic, bicyclic
heteroaryl optionally having 1 to 3 heteroatoms independently selected
15 from oxygen atom, sulfur atom and nitrogen atom, preferably 5 to 6-
membered monocyclic heteroaryl optionally having 1 to 3 nitrogen
atoms, specifically pyrazolyl, imidazolyl, thiazolyl, triazolyl, pyridyl,
pyrimidinyl or the like, particularly pyrimidinyl.

A "cycloalkyl" of substituted or unsubstituted cycloalkyl in R³ and
20 R⁴ includes preferably 3 to 6-membered monocyclic cycloalkyl,
specifically cyclopropyl, cyclopentyl.

A "heterocycle" of substituted or unsubstituted heterocycle which
is the substituent of substituted or unsubstituted alkyl in R⁵ includes,
for example, 5 to 9-membered monocyclic, bicyclic heterocycle
25 optionally having 1 to 3 heteroatoms independently selected from
oxygen atom, sulfur atom and nitrogen atom, preferably 4 to 6-
membered monocyclic heterocycle optionally having 1 to 3 nitrogen
atoms, specifically azetidiny, pyrrolidiny, piperidiny, piperaziny,
morpholiny, thiomorpholiny, perhydrodiazepiny,
30 octahydropyrrolo[1,2-a]piperaziny or the like. More preferable one

among them is piperazinyl, morpholinyl, particularly piperazinyl.

A "heterocycle" of the substituted or unsubstituted heterocycl-
oxy which is the substituent of substituted alkyl in R⁵ includes, for
example, 5 to 9-membered monocyclic, bicyclic heterocycle optionally
5 having 1 to 3 heteroatoms independently selected from oxygen atom,
sulfur atom and nitrogen atom, preferably 4 to 6-membered monocyclic
heterocycle optionally having 1 to 3 nitrogen atoms, specifically
azetidiny, pyrrolidinyl, piperidinyl, piperazinyl, morpholinyl,
thiomorpholinyl, perhydrodiazepinyl, octahydropyrrolo[1,2-a]piperazinyl
10 or the like. More preferable one among them is piperidinyl.

The compound [I] of the present invention includes a mixture of
stereoisomers, or each stereoisomer with pure or substantively pure
forms. For example, the compound [I] can exist in enantiomer or
diastereomer or a mixture thereof when the compound of the present
15 invention has one or more asymmetric centers in any of carbon atoms.
The compound of the present invention includes its isomers or a
mixture thereof. Also, in case that the compound [I] of the present
invention contains double bonds, geometric isomers (cis isomer, trans
isomer) may exist and in case that the compound [I] of the present
20 invention contains unsaturated bonds such as carbonyl, tautomers may
exist, but the compound of the present invention includes all these
isomers or a mixture thereof.

A pharmaceutically acceptable salt of the compound [I] includes,
for example, an inorganic acid salt such as hydrochloride, sulfate,
25 phosphate or hydrobromide, or an organic acid salt such as acetate,
fumarate, oxalate, citrate, methanesulfonate, benzenesulfonate, tosylate
or maleate. Also, in case of having a substituent such as carboxyl, said
salt includes a salt with a base such as, for example, alkali metal salt
such as sodium salt or potassium salt, or alkali earth metal salt such as
30 calcium salt.

The pharmaceutically acceptable salt of the compound [I] of the present invention includes also an intramolecular salt, and the compounds [I] and their salts may be in the form of a solvate thereof such as a hydrate.

5 The compound [I] of the present invention or a pharmaceutically acceptable salt thereof can be formulated to a pharmaceutical composition comprising a therapeutically effective amount of the compound and a pharmaceutically acceptable carrier. The pharmaceutically acceptable carrier can include diluents, binding
10 agents (syrup, gum acacia, gelatin, sorbit, tragacanth or polyvinylpyrrolidone), excipients (lactose, sucrose, cornstarch, potassium phosphate, sorbit or glycine), lubricants (magnesium stearate, talc, polyethylene glycol or silica), disintegrants (Irish potato starch) and wetting agents (sodium lauryl sulfate), or the like.


15 The compound [I] of the present invention or a pharmaceutically acceptable salt thereof can be administered orally or parenterally and used in an appropriate pharmaceutical formulation. The appropriate pharmaceutical formulation for oral administration includes, for example, solid formulations such as tablet, granule, capsule or powder,
20 or in the form of a solution, a suspension or an emulsion. The appropriate pharmaceutical formulation for parenteral administration includes a suppository, an injectable solution or an intravenous fluid preparation using distilled water for injection, saline or glucose aqueous solution, or an inhaler, or the like.

25 The compound [I] of the present invention or a pharmaceutically acceptable salt thereof, or a pharmaceutical formulation thereof can be combined with other one or more medicines selected from antidiabetic and antihyperglycemic agents. In this case, the concept of the term "combine" includes administering with these other medicines
30 simultaneously or separately with optional interval as well as

administering as one pharmaceutical formulation formulated together with these other medicines. These other medicines include sulfonylurea (for example, glyburide, glimepiride, glipiride, glipizide, chlorpropamide, gliclazide, glisoxepide, acetohexamide, glibonuride, tolbutamide, 5 tolazamide, carbutamide, gliquidone, glihexamid, phenbutamide, tolcyclamide or the like), biguanide (for example, metformin, phenformin, buformin or the like), glucagon antagonist (for example, peptidic or nonpeptidic glucagon antagonist), glucosidase inhibitor (for example, acarbose, miglitol or the like), insulin sensitizer (for example, 10 troglitazone, rosiglitazone, pioglitazone or the like), antiobesity agent (for example, sibutramine, orlistat or the like) or the like.

The dose of the compound [I] of the present invention or a pharmaceutically acceptable salt thereof depends on methods of administration, ages, body weights or conditions of patients, but 15 usually about 0.01 to about 100 mg/kg per day, preferably about 0.1 to about 10 mg/kg.

The compound [I] of the present invention can be prepared according to the following methods.



(In the above scheme, Z¹ is halogen atom, hydroxy or alkoxy, Z² is hydrogen atom or alkyl, Z³ is hydroxy, halogen atom or arylsulfonyloxy, alkylsulfonyloxy, Z⁴ is halogen atom, dialkoxyboryl, dihydroxyboryl or trialkylstannyl, lithio, Z⁵ is hydrogen atom, halogen atom, dialkoxyboryl, dihydroxyboryl or trialkylstannyl, lithio, and the other symbols have the same meanings as mentioned above.)

(1) The reaction of preparing the compound [VI] (Z² is alkyl) from the compound [VII] (Z⁵ is hydrogen atom) and the compound [VIII] (Z¹ is halogen atom, Z² is alkyl) can be carried out under so-called Friedel-Crafts reaction condition. For example, the reaction can be carried out in an appropriate solvent (chloroform, methylene chloride, nitromethane

or the like) in the presence of an appropriate acid (aluminum chloride or the like).

The compound [VI] (Z^2 is alkyl) can be also prepared by reacting the compound [VII] (Z^5 is dialkoxyboryl, dihydroxyboryl or trialkylstannyl) with the compound [VIII] (Z^1 is halogen atom, Z^2 is alkyl) in an appropriate solvent (THF, methylene chloride, dioxane, water, DMF, toluene, 1,2-dimethoxyethane or the like, or a mixture thereof) using a metal catalyst (for example, dichlorobis(triphenylphosphine)palladium, tetrakis-(triphenylphosphine)palladium, tris(dibenzylideneacetone)dipalladium, dichloro[1,1'-bis(diphenylphosphino)ferrocene]palladium or the like), and the process preferably proceeds at -78°C to 200°C .

Additionally, the compound [VI] (Z^2 is alkyl) can be also prepared by reacting the compound [VII] (Z^5 is lithio) with the compound [VIII] (Z^1 is alkoxy, Z^2 is alkyl) in an appropriate solvent (THF, dioxane, DMF, toluene, 1,2-dimethoxyethane or the like, or a mixture thereof), and the process can be preferably carried out at -78°C to 200°C . Also, in case of using the compound [VII] (Z^5 is halogen atom), the compound [VI] (Z^2 is alkyl) can be prepared by converting the compound [VII] (Z^5 is halogen atom) into the compound [VII] (Z^5 is lithio) with an appropriate alkyl lithium (n-butyllithium, sec-butyllithium, t-butyllithium or the like) in an appropriate solvent (THF, diethyl ether, toluene, 1,2-dimethoxyethane or the like, or a mixture thereof) to convert, followed by reacting with the compound [VIII] (Z^1 is alkoxy, Z^2 is alkyl) in the similar manner as the above-mentioned.

(2) The reaction of the compound [VI] (Z^2 is alkyl) with hydroxylamine or a salt thereof with an appropriate acid (hydrochloride, sulfate or the like) can be carried out in any conventional manner converting ketone into hydroxyimino. For example, the reaction can be carried out in an appropriate solvent (alcoholic solvent such as methanol, ethanol, or

THF, dioxane, water or the like, or a mixed solvent thereof) in the presence or absence of a base. The base used in the reaction includes pyridine, picoline, lutidine, N,N-dimethylaniline, triethylamine or the like. An oxime generated in a cis-isomer or a mixture of cis- and trans-
5 isomers can be converted into the desired trans-isomer by treating with acid (trifluoroacetic acid, acetic acid, hydrochloric acid, sulfuric acid, phosphoric acid or the like). Also an oxime generated in the reactions described below can also be converted into the desired trans-isomer by treating in the similar manner as the above.

10 (3) The reaction of the compound [IV] (Z^2 is alkyl) with the compound [V] wherein Z^3 is hydroxy can be carried out by using, in the presence of triphenylphosphine, an activating agent (diethyl azodicarboxylate, diisopropyl azodicarboxylate or the like), or, in absence of triphenylphosphine, cyanomethyl tri-n-butyl phosphorane or the like in
15 an appropriate solvent (THF, methylene chloride or the like) (so-called Mitsunobu reaction). Also, the reaction with the compound [V] wherein Z^3 is halogen atom, arylsulfonyloxy or alkylsulfonyloxy can be carried out in an appropriate solvent (acetone, ethanol, THF, dimethyl sulfoxide, DMF, dioxane, N,N-dimethylacetamide, N-methylpyrrolidone
20 or the like, or a mixed solvent thereof) in the presence of a base such as potassium carbonate, potassium tert-butoxide, sodium hydride, cesium carbonate or the like. A product resulted in this way can be converted into the compound [II] (Z^2 is hydrogen atom) in any conventional manner hydrolyzing alkoxycarbonyl to carboxyl, for example, by treating
25 with lithium hydroxide, sodium hydroxide, potassium carbonate or the like in an appropriate solvent (alcoholic solvent such as methanol, ethanol, or THF, dioxane, water or the like, or a mixed solvent thereof) to hydrolyze Z^2 group.

30 (4) The reaction of the compound [II] (Z^2 is hydrogen atom) with the compound [III] can be carried out in an appropriate solvent in the

presence or absence of a condensing agent by using any conventional method for amide formation usually used in peptide synthesis or the like. As the condensing agent, any of N-ethyl-N'-(3-diethylaminopropyl)-carbodiimide, N,N'-dicyclohexylcarbodiimide, 1-methyl-2-bromopyridinium iodide, N,N'-carbonyldiimidazole, diphenyl phosphoryl azide, benzotriazol-1-yloxytris(dimethylamino)phosphonium hexafluorophosphate, 4-(4,6-dimethoxy-1,3,5-triazin-2-yl)-4-methylmorpholinium chloride, fluoro-N,N,N',N'-tetramethylformamidinium hexafluorophosphate or the like can be preferably used. As the solvent, any of a single solvent or a mixed solvent of water, methanol, isopropanol, ethanol, methylene chloride, THF, dioxane, DMF, dimethylacetamide, chloroform or the like can be preferably used. The reaction preferably proceeds at -78°C to 100°C, more preferably at -25°C to 25°C. The proceed of the reaction can be accelerated by adding an inorganic base such as potassium carbonate, sodium carbonate, sodium bicarbonate or an organic base such as triethylamine, diisopropylethylamine, N-methylmorpholine, pyridine, N,N-dimethylaminopyridine, picoline, lutidine or the like as a base, and N-hydroxysuccinimide, 3-hydroxy-3,4-dihydro-4-oxo-1,2,3-benzotriazole, N,N-dimethylaminopyridine or N-hydroxybenzotriazole or the like as an additive.

The reaction from the compound [II] (Z^2 is hydrogen atom) to the compound [I] can be carried out by converting the compound [II] (Z^2 is hydrogen atom) into a reactive intermediate such as acid chloride or a mixed acid anhydride, followed by reacting with the compound [III]. The conversion into acid chloride can be preferably carried out by using thionyl chloride, oxalyl chloride, phosphorus oxychloride, phosphorus pentachloride, or triphenylphosphine in the presence of carbon tetrachloride, or the like, and the conversion into a mixed acid anhydride can be carried out by using diphenyl phosphoryl chloride, diethyl phosphorocyanidate, methanesulfonyl chloride, ethyl

chloroformate, isobutyl chloroformate or the like in the presence of a base such as triethylamine. As the solvent, any of a single solvent or a mixed solvent of methylene chloride, chloroform, THF, DMF or the like can be preferably used. The reaction preferably proceeds at -78°C to 100°C , more preferably -25°C to 25°C . The reaction of an acid chloride or a mixed acid anhydride resulted in this way with the compound [III] proceeds in the presence of a base such as pyridine, triethylamine, N,N-dimethylaminopyridine, diisopropylethylamine or the like preferably at -78°C to 100°C , more preferably -25°C to 25°C and as the solvent, any of a single solvent or a mixed solvent of methylene chloride, chloroform, THF, DMF or the like can be preferably used.

(5) The reaction of the compound [VI] (Z^2 is hydrogen atom or alkyl) with the compound [III] can be carried out in case that Z^2 is hydrogen atom in the similar manner as the reaction of the above (4), or in case that Z^2 is alkyl via the compound [VI'] and the compound [X'] below.


The conversion from the compound [VI] ($Z^2 = \text{alkyl}$) to the compound [VI'] (Z^2 is hydrogen atom) can be carried out in any conventional manner reducing ketone to alcohol, for example, by treating with a reducing agent such as zinc borohydride, sodium

triacetoxyborohydride, sodium borohydride or the like in an appropriate solvent (water, methanol, ethanol, chloroform, methylene chloride or the like, or a mixed solvent thereof), followed by hydrolysis in any conventional manner hydrolyzing alkoxycarbonyl to carboxyl, for example, by treating with lithium hydroxide, sodium hydroxide or the like in an appropriate solvent (methanol, ethanol, THF, dioxane, water or the like, or a mixed solvent thereof) to hydrolyze Z^2 group.

The reaction of the compound [VI'] (Z^2 is hydrogen atom) with the compound [III] can be carried out in the similar manner as the reaction of the above (4).

The conversion from the compound [X'] to the compound [X] can

be carried out in any conventional manner oxidizing alcohol to ketone, for example, by dimethylsulfoxide oxidation using an activating agent such as oxalyl chloride (Swern oxidation), or by using an oxidizing agent (activated manganese dioxide, sulfur trioxide-pyridine complex, 1-hydroxy-1,2-benziodoxol-3(1H)-one 1-oxide, 1,1,1-triacetoxy-1,1-dihydro-1,2-benziodoxol-3(1H)-one, pyridinium chlorochromate, pyridinium dichromate or the like) in the presence or absence of a base (triethylamine or the like) in an appropriate solvent (dimethylsulfoxide, chloroform, methylene chloride or the like).



(6) The reaction from the compound [X] to the compound [IX] can be carried out in the similar manner as the reaction of the above (2).

(7) The reaction from the compound [IX] and the compound [V] to the compound [I] can be carried out in the similar manner as the reaction of the above (3).

(8) The reaction of the compound [VI] with the compound [XI] can be carried out in the similar manner as the reaction of the above (2) by using O-substituted hydroxylamine or a salt thereof (alkyloxyamine, cycloalkyloxyamine, heterocyclyl-oxyamine, benzyloxyamine or the like) as an alternative to hydroxylamine in the reaction of (2).

(9) The reaction of the compound [X] with the compound [XI] can be carried out in the similar manner as the reaction of the above (8).

(10) The reaction of the compound [XI] with the compound [VIII] (Z^1 is halogen atom or hydroxy, Z^2 is alkyl) can be carried out in any conventional manner of amide formation usually used in peptide synthesis or the like, for example in the similar manner as the reaction of the above (4).


(11) The reaction from the compound [XIII] (Z^2 is alkyl) to the compound [XII] (Z^4 is hydrogen atom) can be carried out by using any conventional method for converting amide into haloimino, preferably the manner of reference: WO9520569, for example, by using the compound [XIII] (Z^2 is alkyl) with a halogenating agent (phosphorus oxychloride, phosphorus pentachloride or the like) in an appropriate solvent (acetonitrile, chloroform, methylene chloride, THF or the like, or a mixed solvent thereof). The reaction can be also carried out by using carbon tetrachloride, carbon tetrabromide, N-bromosuccinimide, N-chlorosuccinimide, iodine or the like in the presence of triphenylphosphine.

(12) The reaction from the compound [XII] (Z^2 is alkyl) and the compound [VII], for example in case that Z^5 is dihydroxyboryl, to the compound [II] (Z^2 is alkyl) can be carried out by using a metal catalyst (for example, dichlorobis(triphenylphosphine)palladium, tetrakis(triphenylphosphine)palladium, tris(dibenzylideneacetone)-dipalladium, dichloro[1,1'-bis(diphenylphosphino)ferrocene]palladium or the like) in an appropriate solvent (for example, dioxane, toluene, THF, 1,2-dimethoxyethane, methanol, ethanol, DMF, N-methylpyrrolidone or the like, or a mixed solvent thereof) in the presence of a base (sodium carbonate, potassium carbonate, triethylamine, diisopropylethylamine or the like). The reaction preferably proceeds under inert gas such as argon at room temperature to 200°C or in the exposure of microwave.

The compound [I] can be converted further in the following methods.


(A) The compound containing sulfinyl (SO) or sulfonyl (SO₂) on R¹-R⁶ among the objective compound [I] of the present invention can be prepared by oxidation using any conventional method for converting the corresponding sulfide compound into a sulfinyl or sulfonyl compound. For example, the oxidation can be carried out by treating with an oxidizing agent in an appropriate solvent (methylene chloride, chloroform, THF, methanol, water, or the like or a mixed solvent thereof). As the oxidizing agent, peracids such as hydrogen peroxide, m-chloroperbenzoic acid, peracetic acid or the like as well as oxone™ ("a mixture of potassium peroxybisulfate, dipotassium sulfate and potassium bisulfate" manufactured by DuPont) can be preferably used, and the reaction can be preferably carried out at -78°C to 100°C.

(B) The compound having a group of the formula:



wherein R¹¹ and R¹² are substituents of the substituted amino group described herein or R¹¹ and R¹² form together with N atom of said amino group a heterocycle having 1 to 3 heteroatoms independently selected from oxygen atom, sulfur atom and nitrogen atom wherein the heterocycle may be substituted,

on R¹-R⁶ among the objective compound [I] can be also prepared by so-called "reductive amination", by reacting the compound wherein the corresponding site is formyl with a substituted or unsubstituted amine of the formula:



wherein the symbols have the same meanings as mentioned above (hereinafter, this compound is referred to as "a substituted or unsubstituted amine", and the group after removing of hydrogen atom from the substituted or unsubstituted amine is referred to as "a

substituted or unsubstituted amino"),
under reductive condition. The reaction can be carried out in any
conventional manner of reductive amination. For example, the reaction
can be preferably carried out by using a reducing agent (sodium
5 borohydride, sodium triacetoxyborohydride, sodium cyanoborohydride
or the like) in an appropriate solvent (methanol, methylene chloride,
chloroform or the like) at -78°C to 100°C .

(C) Among the objective compound [I], the compound wherein
nitrogen atom on $\text{R}^1\text{-R}^6$ is substituted with a substituted or
10 unsubstituted alkanoyl such as alkanoyl, cycloalkylcarbonyl,
alkoxyalkanoyl, alkanoyloxyalkanoyl or the like, which is simply
referred to as substituted or unsubstituted alkanoyl hereinafter, can be
also prepared by alkanoylation of the compound wherein the
corresponding N atom is unsubstituted (for example, the compound
15 wherein R^5 is piperazinylmethyl, piperazinylcarbonyl or piperazinyl-
sulfonyl or the like). The alkanoylation can be carried out by using any
conventional method of amide formation usually used in peptide
synthesis or the like. For example, the alkanoylation can be preferably
carried out by using acid chloride, acid anhydride or ester in an
20 appropriate solvent (methylene chloride, THF, DMF, N,N-
dimethylacetamide, chloroform or a mixed solvent thereof) in the
presence or absence of a base (triethylamine, pyridine or the like) at -78°C to 100°C . The reaction can be also carried out, for example, in an
appropriate solvent in the presence or absence of a condensing agent.
25 As the condensing agent, any of N-ethyl-N'-(3-diethylaminopropyl)-
carbodiimide, N,N'-dicyclohexylcarbodiimide, 1-methyl-2-bromo-
pyridinium iodide, N,N'-carbonyldiimidazole, diphenylphosphoryl azide,
benzotriazol-1-yloxytris(dimethylamino)phosphonium hexafluoro-
phosphate, 4-(4,6-dimethoxy[1.3.5]triazin-2-yl)-4-methylmorpholinium
30 chloride, fluoro-N,N,N',N'-tetramethylformamidinium hexafluoro-

phosphate or the like can be preferably used. As the solvent, any of a single solvent or a mixed solvent of water, methanol, isopropanol, ethanol, methylene chloride, THF, DMF, N,N-dimethylacetamide, chloroform or the like can be preferably used. The reaction preferably
5 proceeds at -78°C to 100°C , more preferably -25°C to 25°C . The proceed of the reaction can be promoted by adding potassium carbonate, sodium carbonate, sodium bicarbonate or triethylamine, diisopropylethylamine, N-methylmorpholine, pyridine, N,N-dimethylaminopyridine, picoline, lutidine or the like as a base, and N-
10 hydroxysuccinimide or 3-hydroxy-3,4-dihydro-4-oxo-1,2,3-benzotriazole, N,N-dimethylaminopyridine, N-hydroxybenzotriazole or the like as an additive.

(D) The compound having substituted or unsubstituted amino-carbonyl, i.e. substituted or unsubstituted carbamoyl, on $\text{R}^1\text{-R}^6$ among
15 the objective compound [I] can be prepared by reacting the compound wherein the corresponding site is carboxy with a substituted or unsubstituted amine. The reaction can be carried out in the similar manner as the reaction of the above (C).

(E) The compound wherein R^5 or R^6 has substituted or unsubstituted
20 alkoxymethyl, or substituted or unsubstituted heteroarylmethyl among the objective compound [I] can be also prepared by converting the compound wherein the corresponding site is hydroxymethyl into alkanoylmethyl, preferably acetyloxymethyl, in any conventional esterification manner, followed by condensing substituted or
25 unsubstituted alkanol, cycloalkanol, alkylthio or a heterocycle compound having hydroxyl group, or substituted or unsubstituted heteroaryl compound having hydrogen atom on nitrogen atom, for example pyrazole or the like. The condensing reaction can be preferably carried out as neat or in an appropriate solvent (THF, dioxane,
30 methylene chloride, chloroform, toluene, benzene or the like) in the

presence or absence of an acid (p-toluenesulfonic acid, hydrochloric acid, sulfuric acid, trifluoroacetic acid or the like) at -78°C to 200°C , more preferably 25°C to 100°C .

(F) The compound having hydroxymethyl on $\text{R}^1\text{-R}^6$ among the objective compound [I] can be prepared by reducing the compound wherein the corresponding site is formyl in any conventional manner by reducing formyl to alcohol. For example, the reaction can be preferably carried out by using a reducing agent (sodium borohydride, sodium triacetoxyborohydride, diborane, diisobutylaluminum hydride, lithium aluminum hydride or the like) in an appropriate solvent (methanol, ethanol, methylene chloride, chloroform, dioxane, THF or the like) at -78°C to 100°C .

(G) The compound having carboxyl on $\text{R}^1\text{-R}^6$ among the objective compound [I] can be prepared by oxidizing the compound wherein the corresponding site is formyl in any conventional manner by oxidizing formyl to carboxyl. The oxidation can be preferably carried out by, for example, using an oxidizing agent (sodium chlorite, potassium permanganate, pyridinium dichromate or the like) in an appropriate solvent (DMF, dimethylsulfoxide, acetone, tert-butanol, water, methylene chloride, chloroform or the like) at -78°C to 100°C .

(H) The compound having alkoxycarbonyl on $\text{R}^1\text{-R}^6$ among the objective compound [I] can be also prepared by esterifying the compound wherein the corresponding site is carboxyl in any conventional manner by esterifying carboxyl to alkoxycarbonyl. The esterification can be preferably carried out by, for example, using an acid (sulfuric acid, hydrochloric acid, p-toluenesulfonic acid) in an appropriate solvent (methanol, ethanol, isopropanol, tert-butanol or the like) at -78°C to 200°C , more preferably 0°C to 100°C .

Additionally, the esterification can be also carried out by converting a carboxyl compound to a reactive intermediate such as an

acid halide with a halogenating agent (oxalyl chloride, thionyl chloride or the like) in an appropriate solvent (methylene chloride, chloroform, THF, dioxane or the like), followed by using alkanol (methanol, ethanol, isopropanol or the like) at -78°C to 200°C .

- 5 (I) The compound having carboxyl on $\text{R}^1\text{-R}^6$ among the objective compound [I] can be also prepared by hydrolyzing the compound wherein the corresponding site is alkoxycarbonyl in any conventional manner of ester hydrolysis. The hydrolysis can be preferably carried out by using a base (sodium hydroxide, potassium hydroxide,
10 potassium carbonate, lithium hydroxide or the like) in an appropriate solvent (alcoholic solvent such as methanol, ethanol, or dioxane, THF, water or the like, or a mixed solvent thereof) at -78°C to 200°C , more preferably 0°C to 100°C .

15 Additionally, the hydrolysis can be also preferably carried out by using an acid (sulfuric acid, hydrochloric acid or the like) in an appropriate solvent (THF, dioxane, acetic acid, water or the like, or a mixed solvent thereof) at -78°C to 200°C .

- (J) The compound having formyl on $\text{R}^1\text{-R}^6$ among the objective compound [I] can be also prepared from the compound wherein the
20 corresponding site is carboxyl in any conventional manner by reducing carboxyl to aldehyde. The reaction can be preferably carried out by using a halogenating agent (oxalyl chloride, thionyl chloride or the like) in an appropriate solvent (methylene chloride, chloroform, THF or the like, or a mixed solvent thereof) to synthesize acid halide, followed by
25 reducing the acid halide with a metal catalyst (palladium carbon, platinum dioxide or the like) under hydrogen at -78°C to 200°C .

- (K) The compound having hydroxymethyl on $\text{R}^1\text{-R}^6$ among the objective compound [I] can be prepared by using any conventional method of reduction of ester or carboxylic acid to alcohol. For example,
30 the reaction can be preferably carried out by treating the corresponding

carboxyl or alkoxycarbonyl with a reducing agent (sodium borohydride, diborane, lithium aluminum hydride, diisobutylaluminum hydride or the like) in an appropriate solvent (methylene chloride, chloroform, THF or the like) at -78°C to 200°C .

5 (L) The compound having carboxyl on $\text{R}^1\text{-R}^6$ among the objective compound [I] can be prepared by using the conventional method of oxidation of primary alcohol to carboxylic acid. For example, the reaction can be preferably carried out by using the compound wherein the corresponding site is hydroxymethyl with an oxidizing agent
10 (chromium trioxide, pyridinium dichromate or the like) in an appropriate solvent (methylene chloride, acetone, chloroform, DMF or the like) at, for example, 0°C to 100°C .

(M) The compound having amino on $\text{R}^1\text{-R}^6$ among the objective compound [I] can be carried out by using any conventional method of
15 reduction of nitro to amine. For example, the reaction can be carried out by treating the compound wherein the corresponding site is nitro with a metal catalyst (palladium carbon, platinum dioxide or the like) in an appropriate solvent (methanol, ethanol, DMF, THF, dioxane or the like) under hydrogen at -78°C to 200°C .

20 Additionally, the process can be also preferably carried out by using a reducing agent (stannous chloride, iron, zinc or the like) in an appropriate solvent (alcoholic solvent such as methanol, ethanol, or methylene chloride, chloroform, THF, dioxane, acetic acid, water or the like, or a mixed solvent thereof) at -78°C to 200°C , more preferably 0°C
25 to 100°C .

(N) The compound having halogenosulfonyl on $\text{R}^1\text{-R}^6$ among the objective compound [I] can be prepared by reacting the compound wherein the corresponding site is amino under so-called Sandmayer reaction condition to halogenosulfonylate via a diazonium salt. The
30 formation of a diazonium salt can be preferably carried out by, for

example, using an oxidizing agent (sodium nitrite, isoamyl nitrite, tert-butyl nitrite or the like) in an appropriate solvent (water, methylene chloride, chloroform, THF or the like, or a mixed solvent thereof) in the presence or absence of an appropriate acid (hydrochloric acid, sulfuric acid or the like) and/or an additive (cupric chloride or the like) at -78°C to 200°C. The following halogenosulfonylation can be carried out by adding a sulfonylating agent (sulfur dioxide, sodium bisulfite or the like) to the resulting reaction solution at -78°C to 200°C.

(O) The compound having substituted or unsubstituted aminosulfonyl on R¹-R⁶ among the objective compound [I] can be also prepared by reacting the compound wherein the corresponding site is halogenosulfonyl with a substituted or unsubstituted amine. The reaction can be preferably carried out in an appropriate solvent (methylene chloride, chloroform, THF, dioxane, water or the like) in the presence or absence of a base (pyridine, triethylamine, sodium hydroxide, sodium carbonate or the like) at -78°C to 200°C.

(P) The compound having alkylthio, cycloalkylthio, heterocyclyl-thio on R¹-R⁶ among the objective compound [I] can be also prepared by, for example, converting the compound wherein the corresponding site is methylsulfinyl into thiol in the same manner as described in a literature (Young R. N., et al., Tetrahedron Lett., 1984, 25(17), 1753.), followed by reacting with an alkylating agent (haloalkyl, halocycloalkyl, haloheterocycle compound, alkyl mesylate, cycloalkyl mesylate, heterocyclyl mesylate, alkyl tosylate, cycloalkyl tosylate, heterocyclyl tosylate or the like) in the presence or absence of a base (sodium hydride, cesium carbonate, potassium carbonate, potassium tert-butoxide, triethylamine, diazabicycloundecene or the like).

(Q) The compound having substituted or unsubstituted alkanoyl-amino on R¹-R⁶ among the objective compound [I] can be also prepared by alkanoylating the compound wherein the corresponding site is

amino. The alkanoylation can be carried out in a similar manner as in the reaction of the above (C). Also, the alkanoylation can be carried out in the compound wherein the corresponding site is secondary amine as well as primary amine.

- 5 (R) The compound having substituted sulfonylamino such as alkylsulfonylamino, heteroarylsulfonylamino, heterocycl-
sulfonylamino or the like on R^1 - R^6 among the objective compound [I]
can be also prepared by sulfonylating the compound wherein the
corresponding site is amino. The sulfonylation can be carried out in an
10 appropriate solvent (water, THF, methylene chloride, chloroform or the
like) in the presence or absence of a base (triethylamine,
diisopropylethylamine, pyridine or the like) at -78°C to 200°C . Also, the
sulfonylation can be carried out in the compound wherein the
corresponding site is secondary amine as well as primary amine.
- 15 (S) The compound having secondary alcohol on R^1 - R^6 among the
objective compound [I] can be prepared by using any conventional
method for converting ketone into secondary alcohol. For example, the
reaction can be carried out by using the compound having the
corresponding oxo in the similar manner as the reaction of the above
20 (K).
- (T) The compound having oxo on R^1 - R^6 among the objective
compound [I] can be prepared by using any conventional method for
converting secondary alcohol into ketone. For example, the reaction
can be carried out by dimethylsulfoxide oxidation with an activating
25 agent such as oxalyl chloride in an appropriate solvent
(dimethylsulfoxide, chloroform, methylene chloride or the like) (Swern
oxidation), or by using an oxidizing agent (activated manganese dioxide,
sulfur trioxide-pyridine complex, 1-hydroxy-1,2-benziodoxol-3(1H)-one-
1-oxide, 1,1,1-triacetoxy-1,1-dihydro-1,2-benziodoxol-3(1H)-one,
30 pyridinium chlorochromate, pyridinium dichromate or the like) in the

presence or absence of a base (triethylamine or the like).

(U) The compound having secondary alcohol on R^1 - R^6 among the objective compound [I] can be prepared by using any conventional method for converting the compound having formyl into secondary alcohol. For example, the reaction can be preferably carried out by using the corresponding formyl and a metal reagent (alkylmagnesium halide, alkyllithium, dialkylzinc or the like) in an appropriate solvent (THF, toluene, diethyl ether or the like) at -78°C to 100°C .

(V) The compound having hydroxyamidino on R^1 - R^6 among the objective compound [I] can be prepared by using any conventional method for converting cyano group into hydroxyamidino group. For example, the reaction can be preferably carried out by reacting the compound having the corresponding cyano with hydroxylamine (or a salt with an appropriate acid thereof) in the presence or absence of a base (sodium carbonate, potassium carbonate, sodium hydroxide, potassium hydroxide, potassium tert-butoxide, triethylamine, pyridine or the like) in an appropriate solvent (water, methanol, ethanol or the like, or a mixed solvent thereof) at 0°C to 100°C .

(W) The compound having unsubstituted carbamoyl on R^1 - R^6 among the objective compound [I] can be prepared by using any conventional method for converting cyano group into unsubstituted carbamoyl group. For example, the reaction can be preferably carried out by treating the compound having the corresponding cyano with a base (sodium hydroxide, potassium hydroxide, potassium tert-butoxide or the like) in an appropriate solvent (water, methanol, ethanol, isopropanol or the like, or a mixed solvent thereof) at -20°C to 100°C .

(X) The compound having tertiary alcohol on R^1 - R^6 among the objective compound [I] can be prepared by, for example, reacting the compound having the corresponding oxo under a condition of the above (U).

(Y) The preparation of the compound having optically-active secondary alcohol on R¹-R⁶ among the objective compound [I] can be carried out by using any conventional method for resolution of secondary alcohol compound in enzymatic transesterification. For example, the preparation can be preferably carried out by treating the corresponding racemic secondary alcohol with acyl donor (vinyl acetate or the like) in the presence of enzyme (lipase PS or the like) in an appropriate solvent (tert-butylmethyl ether, hexane, diisopropyl ether, THF, diethyl ether, water or the like) at -78°C to 100°C.

(Z) The preparation of the compound having alkyl on R¹-R⁶ among the objective compound [I] can be carried out by using so-called catalytic hydrogenation. For example, the compound can be preferably prepared by treating the compound having the corresponding alkenyl with a metal catalyst (palladium carbon, platinum dioxide or the like) under hydrogen in an appropriate solvent (methanol, ethanol, DMF, THF, acetic acid or the like, or a mixed solvent thereof) at 0°C to 200°C.

(AA) The preparation of the compound having 1,2-diol on R¹-R⁶ among the objective compound [I] can be preferably carried out by, for example, treating the compound having the corresponding alkenyl with an oxidizing agent (osmium tetroxide, ruthenium tetroxide, sodium periodate or the like) in an appropriate solvent (water, acetone, THF, acetonitrile, ethyl acetate or the like, or a mixed solvent thereof) at 0°C to 100°C.

(BB) The preparation of the compound having halogen atom on R¹-R⁶ among the objective compound [I] can be carried out by using any conventional method for halogenation of alcohol. For example, the preparation can be preferably carried out by treating the corresponding alcohol with carbon tetrabromide in the presence of triphenylphosphine in an appropriate solvent (methylene chloride, chloroform or the like) at 0°C to 100°C.

(CC) The preparation of the compound having unsubstituted and substituted alkylthio, heteroarylthio or arylthio on R¹, R², R⁵ or R⁶ among the objective compound [I] can be carried out by using any conventional method for coupling thiol with halogenated aryl, halogenated heteroaryl, aryl triflate or heteroaryl triflate. For example, the preparation can be preferably carried out by treating the compound having the corresponding haloaryl with thiol (hydroxyalkylthiol, dialkylaminoalkylthiol, or the like) in the presence of a metal catalyst (tetrakis(triphenylphosphine)palladium or the like) in an appropriate solvent (dioxane, toluene, THF, 1,2-dimethoxyethane or the like, or a mixed solvent thereof) in the presence or absence of a base (triethylamine, diisopropylamine or the like) at 0°C to 200°C.

(DD) The preparation of the compound having mono-substituted or di-substituted alkylamino on R¹-R⁶ among the objective compound [I] can be preferably carried out by, for example, treating the compound having the corresponding haloalkyl with mono-substituted or di-substituted alkylamine (dimethylamine, diethylamine, methylamine or the like) in an appropriate solvent (methanol, ethanol, dioxane, toluene, THF, 1,2-dimethoxyethane or the like) in the presence or absence of a base (triethylamine, diisopropylamine or the like) at 0°C to 200°C. Also, the compound having dimethylamino can be preferably prepared by treating the compound having the corresponding haloalkyl with N-(trimethylsilyl)dimethylamine in an appropriate solvent (methanol, ethanol, dioxane, toluene, THF, 1,2-dimethoxyethane or the like) at 0°C to 200°C.

(EE) The preparation of the compound having alkynyl on R¹, R², R⁵ or R⁶ among the objective compound [I] can be carried out by using any conventional method of so-called Sonogashira coupling reaction of halogenated aryl, halogenated heteroaryl, aryl triflate or heteroaryl triflate with the compound having alkyne. For example, the preparation

can be preferably carried out by treating the compound having the corresponding halogen with alkyne (propargyl alcohol, N,N-dimethylpropargylamine or the like) in the presence of a metal catalyst (tetrakis(triphenylphosphine)palladium or the like) in an appropriate solvent (dioxane, toluene, THF, 1,2-dimethoxyethane or the like) in the presence or absence of a base (triethylamine, diisopropylamine or the like) and/or copper salt (for example, cuprous iodide) at 0°C to 200°C.

(FF) The preparation of the compound having tetrazolyl on R¹-R⁶

among the objective compound [I] can be carried out by using any conventional method for converting cyano group into tetrazolyl group. For example, the preparation can be preferably carried out by treating the compound having the corresponding cyano with metal azide (sodium azide, tributyltin azide, trimethylsilyl azide) in an appropriate solvent (methanol, ethanol, DMF, dioxane, toluene, THF, 1,2-dimethoxyethane or the like) in the presence or absence of a base (triethylamine, diisopropylamine or the like) or a salt (triethylamine hydrochloride or the like) at 0°C to 200°C.

(GG) The preparation of the compound having O-

alkoxycarbonylhydroxyimine on R¹-R⁶ among the objective compound [I]

can be preferably carried out by treating the compound having the corresponding hydroxyimine with alkyl chlorocarbonate (ethyl chlorocarbonate or the like) in an appropriate solvent (DMF, dioxane, toluene, THF, 1,2-dimethoxyethane or the like) or as neat in the presence or absence of a base (pyridine, triethylamine or the like) at 0°C to 200°C.

(HH) The preparation of the compound having aryl or heteroaryl on R¹, R², R⁵ or R⁶ among the objective compound [I] can be carried out by using any conventional method of so-called Stille coupling or Suzuki coupling reaction.

For example, the preparation can be preferably carried out by treating the compound having the corresponding haloaryl

with aryltrialkyltin, heteroaryltrialkyltin, aryldihydroxyborane, heteroaryldihydroxyborane, arylcatecholborane, heteroarylcatecholborane or the like in the presence of a metal catalyst (for example, dichlorobis(triphenylphosphine)palladium, tetrakis-
5 (triphenylphosphine)palladium, tris(dibenzylideneacetone)dipalladium, dichloro[1,1'-bis(diphenylphosphino)ferrocene]palladium, palladium acetate or the like) in an appropriate solvent (dioxane, toluene, THF, 1,2-dimethoxyethane or the like, or a mixed solvent thereof) in the presence or absence of a base (triethylamine, diisopropylamine, sodium
10 tert-butoxide, sodium carbonate, cesium carbonate, potassium phosphate or the like) at 0°C to 200°C.

(II) In the above each reaction, a protecting group can be optionally introduced or removed to give the desired compound [I] finally. The method for introduction and removal of the protecting group can be
15 carried out according to the description of Protective Groups in Organic Synthesis Third Edition (Theodora W. Green and Peter G. Wuts).

(JJ) Alternatively, the compound [I] can be also synthesized by optionally carrying out any of the above reaction of (A) to (II) in the compound [II] to the compound [XIII] in an appropriate stage in each
20 process of (1) to (13).

Example of Experiment

A Glucokinase Activation Effect

(Method)

A glucokinase activity was examined by measuring the amount of
25 NADPH obtained in generating 6-phosphogluconic acid from glucose-6-phosphoric acid by a coupling enzyme glucose-6-phosphate dehydrogenase not by measuring directly the produced glucose-6-phosphoric acid. The glucokinase enzyme used in the examination is human-liver type GST-GK expressed in E.Coli. The measurement of GK
30 activity was carried out by the following procedures.

Twenty five mM HEPES buffer (pH7.4) containing 25 mM MgCl₂, 25 mM KCl, 1 mM DTT, 5 mM NADP (Roche), 16.64 µg/mL G6PDH (Roche 737-232 grade II from yeast) and 2.8 µg/mL GST-GK was prepared as a reaction solution. An evaluating compound dissolved in DMSO was added to the reaction solution to give final concentration of 0.001 to 100 µM (5% DMSO). Thereto was added glucose (final concentration of 5 mM) as a substrate and was added ATP (final concentration of 5 mM), and the reaction was started. The reaction temperature is 30°C and a generation of NADPH was monitored by changes of absorbance of 340 nm. An increasing in absorbance for 15 minutes from starting reaction was measured and the blank-corrected value was used as GK activity (mOD/min). EC₅₀ level was calculated by a GK activity level in at each concentration of an evaluating compound.

(Results)


EXAMPLE No.	EC ₅₀ (µM)
6	1.20
9	0.55
10	0.79
11	0.88
13-3	0.17
18-2	0.93
24-25	0.084
46-1	0.10
56	0.32
62-5	0.41
62-10	0.27
67-2	0.39
82-22	0.26
82-78	0.60
84-12	0.41
91-8	0.52
94-6	0.51
98-1	0.84
98-7	0.23
104-2	0.30
116	0.57
139-125	0.76
139-244	0.46

139-134	0.18
139-94	0.35
139-237	0.60
139-41	0.54
139-159	0.52
139-182	0.52
139-214	0.57
139-221	0.13

Another objective of the present invention is to provide an industrially advantageous method for preparing 5-substituted 2-aminothiazole and a salt thereof, and it has surprisingly been found that the desired 5-substituted 2-aminothiazole compound can be prepared in high yield by using 2-aminothiazole wherein the 5-position is not substituted as a starting material. The method of the present invention is an industrially very advantageous since the starting material, 2-aminothiazole, is commercially available at a low cost compared to 5-bromo-2-aminothiazole, which gives lowering of the production cost, and further various substituents can be introduced at 5-position of 2-aminothiazole.

Thus, the present invention includes the following embodiments of the method for preparing the desired compounds:

[1] A method for preparing 5-substituted 2-aminothiazole of the general formula:



wherein the symbols have the same meanings as mentioned above, or a salt thereof by treating 2-aminothiazole wherein the amino group may be protected, or a salt thereof with a base, followed by treating the resultant with an electrophile of the general formula:



wherein X is a leaving group, G is halogen atom, formyl, alkoxycarbonyl, alkylthio, alkylsulfinyl, alkylsulfonyl, aminosulfonyl, dialkylsulfonyl, alkylboryl or trialkylsilyl;

and removing the protecting group where the amino group is protected.

5 [2] A method for preparing of [1] wherein 2-amino-1,3-thiazole wherein the amino group may be protected is 2-amino-1,3-thiazole wherein the amino group is protected.

[3] A method for preparing of [1] or [2] wherein G is halogen atom or formyl.

10 [4] A method for preparing of [1], [2] or [3] wherein the base is alkyl lithium.

[5] A method for preparing of [1], [2], [3] or [4] wherein the base is used in two or more equivalents to one equivalent of 2-amino-1,3-thiazole or a salt thereof.

15 5-Substituted 2-aminothiazole or a salt thereof to be prepared by the method of the present invention is preferably a compound wherein the substituent G is halogen atom or formyl. A compound wherein G is fluorine atom or formyl, particularly fluorine atom, is more preferable.

A conventional protecting group can be used as a protecting
20 group of the amino group of the starting material, 2-aminothiazole. Said protecting group includes, for example, oxycarbonyl-type protecting group such as substituted or unsubstituted alkoxycarbonyl (for example, methoxycarbonyl, ethoxycarbonyl, 2,2,2-trichloroethoxycarbonyl, tert-butoxycarbonyl), substituted or unsubstituted
25 aralkyloxycarbonyl (for example, benzyloxycarbonyl), or substituted or unsubstituted aryloxycarbonyl (for example, phenoxycarbonyl); formyl; carbonyl-type protecting group such as substituted or unsubstituted alkanoyl (for example, trifluoroacetyl, tert-butanoyl), or substituted or unsubstituted arylcarbonyl (for example, benzoyl); or alkyl-type
30 protecting group such as substituted or unsubstituted alkyl (for

example, tert-butyl), or substituted or unsubstituted aralkyl (for example, benzyl, benzhydryl, trityl).

A preferable protecting group among them is oxycarbonyl-type protecting group, carbonyl-type protecting group, alkyl-type protecting group, more preferably oxycarbonyl-type protecting group, carbonyl-type protecting group, particularly oxycarbonyl-type protecting group.

A preferable oxycarbonyl-type protecting group is substituted or unsubstituted alkoxycarbonyl, substituted or unsubstituted aralkyloxycarbonyl, particularly substituted or unsubstituted alkoxycarbonyl. A preferable carbonyl-type protecting group is substituted or unsubstituted alkanoyl. A preferable alkyl-type protecting group is substituted or unsubstituted aralkyl.

A preferable substituted or unsubstituted alkoxycarbonyl is tert-butoxycarbonyl. A preferable substituted or unsubstituted aralkyloxycarbonyl is benzyloxycarbonyl. A preferable substituted or unsubstituted alkanoyl is trifluoroacetyl. A preferable substituted or unsubstituted aralkyl is benzhydryl.

A salt of 2-aminothiazole wherein the amino group may be protected includes a salt with an inorganic acid such as hydrochloride, hydrobromide, hydroiodide, sulfate, nitrate or phosphate; or a salt with an organic acid such as formate, acetate, propionate, oxalate, malonate, succinate, fumarate, maleate, lactate, malate, tartarate, citrate, methanesulfonate, ethanesulfonate, benzenesulfonate or toluenesulfonate.

A strong base can be preferably used as a base of base-treatment. Such a strong base includes lithium compound such as alkyllithium, cycloalkyllithium, aryllithium, lithium amide or lithium cyclyl-amide. Among them, using alkyllithium or cycloalkyllithium is preferable, most preferably alkyllithium in particular.

Alkyllithium includes n-butyllithium, tert-butyllithium, sec-

butyllithium or the like. Cycloalkyllithium includes cyclohexyllithium or the like. Aryllithium includes phenyllithium or the like. Lithium amide includes lithium dialkylamide (lithium diisopropylamide), lithium bis(trialkylsilyl)amide (lithium bis(trimethylsilyl)amide) or the like.

5 Lithium cyclyl-amide includes lithium 2,2,6,6-tetraalkylpiperidide (lithium 2,2,6,6-tetramethylpiperidide) or the like.

The base-treatment can be carried out in an appropriate solvent under cooling. Any of aliphatic hydrocarbon-type solvent, aromatic hydrocarbon-type solvent, ether-type solvent, phosphoric amide-type solvent, urea-type solvent, amine-type solvent or a mixed solvent thereof

10 can be preferably used as said solvent. A preferable solvent among them is ether-type solvent.

The aliphatic hydrocarbon-type solvent includes pentane, hexane, cyclohexane, preferably hexane or cyclohexane. The aromatic

15 hydrocarbon-type solvent includes toluene, xylene, preferably toluene. The ether-type solvent includes anisole, dimethyl ether, diethyl ether, diisopropyl ether, tert-butylmethyl ether, cyclopentyl methyl ether, THF, 1,2-dimethoxyethane, preferably diethyl ether, THF, 1,2-dimethoxyethane, particularly THF. The phosphoric amide-type solvent

20 includes hexaalkylphosphoric triamide, preferably hexamethylphosphoric triamide in particular. The urea-type solvent includes N,N'-dimethylpropyleneurea, N,N'-dimethylethyleneurea, preferably N,N'-dimethylpropyleneurea in particular. The amine-type solvent includes N,N,N',N'-tetramethylethylenediamine or the like.

25 The proceeding of the reaction can be promoted by adding a small portion of the phosphoric amide-type solvent, the urea-type solvent or the amine-type solvent as a co-solvent to the other solvent. For example, the co-solvent including hexamethylphosphoric triamide, N,N'-dimethylpropyleneurea, N,N'-dimethylethyleneurea or N,N,N',N'-

30 tetramethylethylene diamine, or a mixed solvent comprising one or

more kinds of these solvents can be added to the solvent including pentane, hexane, cyclohexane, toluene, xylene, anisole, dimethyl ether, diethyl ether, diisopropyl ether, tert-butylmethyl ether, cyclopentyl methyl ether, THF or 1,2-dimethoxyethane, or a mixed solvent comprising one or more kinds of these solvents. An amount of the co-solvent added in using in this way includes a range of 0.1% to 70%, preferably a range of 3% to 30% to an original solvent. In this case, the preferable original solvent among the above-mentioned is hexane, cyclohexane, toluene, diethyl ether, tert-butylmethyl ether, THF or 1,2-dimethoxyethane, or a mixed solvent comprising one or more kinds of these solvents, particularly hexane, toluene, diethyl ether, THF or 1,2-dimethoxyethane, or a mixed solvent comprising one or more kinds of these solvents. The most preferable one is THF.

The cooling condition in base-treatment includes a range of -100°C to 25°C, preferably a range of -78°C to 25°C. Particularly, a range of -78°C to 0°C is preferred.

The proceeding of the reaction can be promoted by using greater or equal to two equivalents of a base to one equivalent of 2-aminothiazole or a salt thereof in base-treatment.

An electrophile using in electrophile-treatment can include an electrophile of the general formula:



wherein the symbols have the same meanings as mentioned above.

Any conventional leaving group can be preferably used as X of the electrophile G-X. Therefore, G-X can be, for example, halide-type electrophile using halogen atom as X, ester-type electrophile using alkoxy or the like as X, or amine-type electrophile using substituted or unsubstituted amino group as X. Also, in case that G is halogen atom or alkylthio, G-X can be dimer-type electrophile of G (in case X = G).

Among them, amine-type electrophile is preferred.

The halide-type electrophile includes alkyl halocarbonate (ethyl chlorocarbonate, methyl chlorocarbonate or the like), alkylphosphoryl halide (ethylphosphoryl chloride or the like), trialkylsilyl halide (trialkylsilyl chloride, trialkylsilyl bromide or the like), alkylthio halide (methylthio chloride or the like), alkylsulfinyl halide (methylsulfinyl chloride or the like), or alkylsulfonyl chloride (methanesulfonyl chloride, ethanesulfonyl chloride or the like). Among them, alkylphosphoryl halide or trialkylsilyl halide is preferred, particularly ethylphosphoryl chloride or trimethylsilyl chloride is most preferred. The ester-type electrophile includes dialkyl carbonate (diethyl carbonate, dimethyl carbonate or the like) or trialkyl borate (trimethyl borate, triisopropyl borate or the like). Among them, trialkyl borate is preferred, particularly trimethyl borate is most preferred. The amine-type electrophile includes N-chlorosuccinimide, N-bromosuccinimide, N-iodosuccinimide, N-fluoropyridinium, 1-fluoropyridin-2-one, N-fluoroquinuclidium, 1-chloromethyl-4-fluoro-1,4-diazoniabicyclo[2.2.2]octane bistetrafluoroborate, N-fluoroperfluoropiperidine, N-fluorobenzenesulfonylimide, N-fluorotrifluoromethanesulfonylimide, N-fluoro-N-methyl-p-toluenesulfonylamide or 2,3-dihydro-3,3-dimethyl-2-fluoro-1,2-benzothiazole-1,1-dione. Among them, N-fluorobenzenesulfonylimide or N-fluoro-N-methyl-p-toluenesulfonylamide is preferred, particularly N-fluorobenzenesulfonylimide is most preferred. The dimer-type electrophile includes dialkyl disulfide (dimethyl disulfide, bis(trifluoromethyl)disulfide or the like) or halogen molecule (fluorine, chlorine, bromine or iodine). Among them, dialkyl disulfide is preferred, particularly dimethyl disulfide is most preferred.

The electrophile-treatment can be carried out in an appropriate solvent under cooling. The solvent can preferably include the solvent of the above-mentioned in the base-treatment.

The base-treatment and the electrophile-treatment can be carried

out in the same solvent by selecting an appropriate solvent. In this case, the electrophile-treatment can be carried out sequentially in the solvent used in the base-treatment. Such a solvent includes a mixed solvent of THF and hexane, a mixed solvent of diethyl ether and hexane, or the like.

The cooling condition of the electrophile-treatment includes a range of -100°C to 25°C ; preferably a range of -78°C to 25°C .

The protecting group which protects the amino group of 5-substituted 2-aminothiazole compound or a salt thereof can be removed by the conventional method. A removing method of such a protecting group includes, for example, acidolysis, acid hydrolysis, alkali hydrolysis, catalytic reduction or the like.

The acidolysis can be carried out by using an acid such as trifluoroacetic acid, hydrochloric acid, sulfuric acid, titanium tetrachloride or stannic chloride in an appropriate solvent (for example, methylene chloride, chloroform, toluene, methanol, ethanol, THF, water or the like). The acid hydrolysis can be carried out by using an acid such as hydrochloric acid or sulfuric acid in an appropriate solvent (for example, water, or a mixed solvent of methanol, ethanol, THF or the like with water). Also, the alkali hydrolysis can be carried out by using sodium hydroxide, potassium hydroxide, potassium carbonate, sodium carbonate, sodium bicarbonate or the like in a solvent such as water, methanol, ethanol or THF.

The resulting 5-substituted 2-aminothiazole compound or a salt thereof can be the desired salt by the conventional method. The salt of 5-substituted 2-aminothiazole compound includes a salt with an inorganic acid such as hydrochloride, hydrobromide, hydroiodide, sulfate, nitrate, phosphate; or a salt with an organic acid such as formate, acetate, propionate, oxalate, malonate, succinate, fumarate, maleate, lactate, malate, tartarate, citrate, methanesulfonate,

ethanesulfonate, benzenesulfonate, toluenesulfonate. Moreover, in case that substituent at 5-position has an acidic group, the salt includes a salt with an inorganic base such as alkali metal including lithium, sodium or potassium, alkali earth metal including calcium or magnesium, or other metal including zinc or aluminum; or a salt with an organic base such as ammonium, choline, diethanolamine, lysine, ethylenediamine, tert-butylamine, tert-octylamine, tris(hydroxymethyl)aminomethane, N-methylglucosamine, triethanolamine or dehydroabiethylamine.

In the above method for preparing of the present invention, aryl and aryl in aralkyl include monocyclic, bicyclic or tricyclic aryl having 6 to 14 carbons, preferably 6 to 10 carbons, specifically phenyl, naphthyl, phenanthryl, anthranyl or the like. Alkyl, alkoxy, alkanoyl, aryl or aralkyl may be substituted with one or more groups selected from halogen atom, alkyl, alkoxy or aryl.

The other groups are the same as the above-mentioned in the oxime derivative [I].

Meanwhile, thiazole includes 1,2-thiazole (isothiazole) and 1,3-thiazole, but in the above method for preparing of the present invention, it is described simply as "thiazole" in the meaning of 1,3-thiazole.

In the present specification, DMF represents N,N-dimethylformamide and THF represents tetrahydrofuran.

Effect of the Invention

The compound [I] of the present invention or a pharmaceutically acceptable salt thereof is useful for preventing or treating diseases involving glucokinase, for example, diabetes, particularly type 2 diabetes, or chronic complications associated with diabetes such as retinopathy, nephropathy, neuropathy, ischemic heart disease or arteriosclerosis, additionally obesity, because of its excellent glucokinase activation effect.

On the other hand, 5-substituted 2-aminothiazole compound or a salt thereof can be prepared in a good yield by the method for the preparing of the present invention. Also, the method for the preparing of the present invention is an industrially advantageous method which can introduce various substituents at 5-position of 2-aminothiazole depending on the kinds of the electrophile used. Additionally, it is an industrially very advantageous method because 2-aminothiazole, the starting material of the method of the present invention, is low in price compared to 5-bromo-2-aminothiazole, which gives lowering of production cost.

BEST MODE FOR CARRYING OUT THE INVENTION


The present invention is explained in more detail in the following EXAMPLES and REFERENCE EXAMPLES, but the invention is not limited to these explanations.

In EXAMPLES, APCI is atmospheric pressure chemical ionization mass spectrum and ESI is electrospray ionization mass spectrum.

EXAMPLES

EXAMPLE 1

66



- (1) To a solution of aluminum chloride (67.0 g, 503 mmol) in methylene chloride (380 ml) was added methyl chloroglyoxylate (48.9 g, 399 mmol) under ice-cooling, and the mixture was stirred for 30 minutes at the same temperature. To the mixture was added a solution of cyclopropyl phenyl sulfide (compound 1-A) (50 g, 333 mmol) in methylene chloride (60 ml), and then the ice-cooling bath was removed and the mixture was stirred at room temperature for 1.5 hours. The reaction mixture was poured onto ice, and then the methylene chloride layer was separated and concentrated in *vacuo*. The residue was dissolved in ethyl acetate and then washed sequentially with water, a saturated aqueous sodium bicarbonate solution and brine, followed by drying over sodium sulfate and concentrated in *vacuo*. The residue was recrystallized from hexane to give the compound (1-B) (69.5 g, yield 88%) as pale yellow crystals.

(2) To a solution of the above compound (57.0 g, 241 mmol) in methanol-THF (1:1) (1480 ml) was added dropwise an aqueous solution (513 ml) of oxone™ (177.9 g, 289 mmol) under ice-cooling over 1 hour, and then the mixture was stirred at room temperature for 12 hours after removing the ice bath. The insoluble materials were filtered off, and then the filtrate was concentrated in *vacuo*. The residue was dissolved in ethyl acetate, washed sequentially with water and brine, dried over sodium sulfate and concentrated in *vacuo*. The resulting residue was recrystallized from diethyl ether to give the compound (1-C) (44.3 g, yield 69%) as pale yellow crystals.

(3) To a solution of the above compound (65.0 g, 242 mmol) in methanol (450 ml) was added hydroxylamine hydrochloride (23.6 g, 339 mmol) at room temperature, and the mixture was heated to reflux for 3 hours. The reaction mixture was concentrated in *vacuo* and then the residue was dissolved in ethyl acetate and washed sequentially with water and brine, followed by drying over sodium sulfate and concentrated in *vacuo*. The residue was dissolved in trifluoroacetic acid (200 ml) and the mixture was stirred at room temperature for 12 hours. After concentration in *vacuo*, the residue was recrystallized from hexane-ethyl acetate to give the compound (1-D) (53.1 g, yield 78%) as colorless crystals.

(4)

(4-1) To a solution of the above compound (37.2 g, 130 mmol), triphenylphosphine (47.7 g, 182 mmol) and (S)-3-hydroxytetrahydrofuran (26.0 g, 294 mmol) in THF (400 ml) was added dropwise diisopropyl azodicarboxylate (36.8 g, 182 mmol) under ice-cooling, and the mixture was stirred at the same temperature for 3 hours and at room temperature for another 16 hours. The reaction mixture was ice-cooled again, and thereto were added water (55 ml) and a 5.4N sodium hydroxide solution (36 ml). The mixture was stirred at

the same temperature for 1 hour and concentrated. Thereto was added water, and the mixture was washed with ethyl acetate twice and then the aqueous layer was acidified with 10% hydrochloric acid to pH 2 to 3 and extracted with chloroform. The organic layer was separated, followed by washing sequentially with water and brine and drying over sodium sulfate, and concentrated in *vacuo*. The residue was recrystallized from ethyl acetate to give the compound (1-E) (31.6 g, yield 72%) as colorless crystals.

(4-2) The above compound (1-E) was also synthesized by using the following alternative method.

(4-2-1) To a solution of the compound (1-D) (68.1 g, 241 mmol) and potassium carbonate (66.5 g, 482 mmol) in DMF (1200 ml) was added (*S*)-3-tetrahydrofuranol *p*-toluenesulfonate (synthesized from (*S*)-3-hydroxytetrahydrofuran and *p*-toluenesulfonyl chloride) (69.9 g, 289 mmol) under ice-cooling, and the ice bath was removed. The mixture was stirred at room temperature overnight, and then diluted with ethyl acetate, washed sequentially with water and brine, dried over sodium sulfate and concentrated in *vacuo* to give the compound (1-D-1) (94.7 g, quantitatively).


(4-2-2) To a solution of the above compound (94.7 g) in water-methanol (1:3.3) (365 ml) was added an aqueous solution (55 ml) of sodium hydroxide (12.5 g, 312 mmol) under ice-cooling, and the mixture was stirred at the same temperature for 30 minutes. To the reaction mixture was added ethyl acetate, and the aqueous layer was separated and then acidified with 10% hydrochloric acid and extracted with chloroform. The organic layer was separated and then washed sequentially with water and brine, dried over sodium sulfate and concentrated in *vacuo* to give the compound (1-E) (58.5 g, yield 71%).

(5) To a solution of the compound (1-E) (41.9 g, 123 mmol), 2-amino-5-formylthiazole hydrochloride (30.4 g, 184 mmol) and *N,N*-

dimethylaminopyridine (22.5 g, 184 mmol) in methylene chloride (1270 ml) was added dropwise N-ethyl-N'-(3-dimethylaminopropyl)-carbodiimide (28.6 g, 184 mmol) at room temperature. The mixture was stirred at the same temperature for 12 hours and concentrated in *vacuo*.
5 The residue was purified by silica gel column chromatography (3% methanol-chloroform) to give the compound (1-F) (33.6 g, yield 61%) as pale yellow crystals.

MS (m/z) APCI: 450 [M+H]⁺

EXAMPLE 2




To a solution of the compound (1-F) (200 mg, 0.44 mmol) and (R)-2-methylpiperazine (223 mg, 2.65 mmol) in methylene chloride (4 ml) was added sodium triacetoxyborohydride (112 mg, 0.55 mmol) under ice-cooling, and the mixture was stirred at room temperature for 24
15 hours. To the reaction solution was added water, and the organic layer was separated, dried over magnesium sulfate and concentrated in *vacuo*. The residue was purified by silica gel column chromatography (NH-silica gel; 1 to 6% methanol-chloroform) to give the compound (2-A) (116.7 mg, yield 49%) as colorless crystals.




MS (m/z) APCI: 534 [M+H]⁺

EXAMPLES 3 to 10


Corresponding starting compounds were treated in the similar manner as EXAMPLE 2 to give the following compounds.



EXAMPLE No.	No	Structure	MS (m/z)
3			534 APCI [M+H] ⁺
4			548 APCI [M+H] ⁺
5			548 APCI [M+H] ⁺
6			534 APCI [M+H] ⁺
7			534 APCI [M+H] ⁺

8		606 APCI [M+H] ⁺
9		576 APCI [M+H] ⁺
10		562 APCI [M+H] ⁺


EXAMPLE 11



To a solution of the compound of EXAMPLE 4 (30 mg, 0.055 mmol) and diisopropylethylamine (21 mg, 0.165 mmol) in chloroform (1.5 ml) was added methoxyacetyl chloride (9.0 mg, 0.083 mmol) under ice-cooling, and the mixture was stirred at room temperature for 24 hours. To the reaction mixture was added an aqueous sodium bicarbonate solution. The organic layer was separated and concentrated in *vacuo*. The residue was purified by LC/MS (Xterra Prep MS C18 5 μ m, 30 x 50 mm; MeOH-10 mM (NH₄)₂CO₃aq, 40:60 to 70:30) to give the above compound (13 mg, yield 43%) as colorless crystals.

MS (m/z) ESI: 620 [M+H]⁺

EXAMPLE 12



(1) An enantiomer ((S)-isomer) of the compound (1-E) was synthesized by reacting in the similar manner as EXAMPLE 1-(4) using the corresponding antipode ((R)-isomer) as an alternative to (S)-3-hydroxytetrahydrofuranol used in EXAMPLE 1-(4) or a tosylate thereof.







(2) The titled compound was obtained by reacting the above compound in the similar manner as EXAMPLE 1-(5).


MS (m/z) APCI: 440 [M+H]⁺

EXAMPLE 13


Corresponding starting compounds were reacted in the similar manner as EXAMPLE 1-(5) to give the following compounds.

EXAMPLE No.	No	Structure	MS (m/z)
13.	1		489 APCI [M+H] ⁺
13	2		466 APCI [M+H] ⁺

13	3		560 APCI [M+H] ⁺
13	4		603 APCI [M+H] ⁺
13	5		546 APCI [M+H] ⁺
13	6		560 APCI [M+H] ⁺
13	7		574 APCI [M+H] ⁺
13	8		646 APCI [M+H] ⁺

13	9		632 APCI [M+H] ⁺
----	---	--	--------------------------------


EXAMPLE 14



The compound of EXAMPLE 13-(4) (2.69 g, 4.46 mmol) was dissolved in formic acid (50 ml). The mixture was stirred at room temperature for 20 hours and concentrated in *vacuo*. The residue was chased with toluene and solidified with ethyl acetate-hexane to give the above compound (2.46 g, quantitatively).

MS (m/z) ESI: 545 [M-H]⁻

EXAMPLE 15




To a suspension of the compound of EXAMPLE 13-(8) (151 mg, 0.234 mmol) in ethyl acetate (3 ml) was added a 4M hydrogen chloride solution in dioxane (6 ml, 24 mmol) at room temperature. The mixture was stirred for 16 hours at the same temperature and then diluted with

diethyl ether. The precipitated crystals were collected and dried to give the above compound (127 mg, yield 93%).

MS (m/z) APCI: 546 [M+H]⁺


EXAMPLE 16









The compound of EXAMPLE 13-(9) was treated in the similar manner as EXAMPLE 15 to give the above compound.







MS (m/z) APCI: 532 [M+H]⁺







EXAMPLE 17



To a solution of the carboxylic acid (1-E) of EXAMPLE 1 (100 mg, 0.295 mmol) and the amine of REFERENCE EXAMPLE 8 (68.5 mg, 0.324 mmol) in THF-N-methylpyrrolidone (1:1) (6 ml) was added 4-(4,6-dimethoxy-1,3,5-triazin-2-yl)-4-methylmorpholinium chloride (hereinafter called DMT-MM) (90 mg, 0.325 mmol) at room temperature. The mixture was stirred for 20 hours at the same temperature and diluted with diisopropyl ether-hexane. The resulting precipitates were collected and purified by LC/MS (Xterra Prep MS C18 5 μ m, 30 x 50 mm; MeOH-10 mM (NH₄)₂CO₃aq, 70:30) to give the above compound (8 mg, yield 6%) as colorless crystals.





18	6		588 APCI [M+H] ⁺
18	7		562 APCI [M+H] ⁺
18	8		562 APCI [M+H] ⁺
18	9		606 APCI [M+H] ⁺
18	10		507 APCI [M+H] ⁺
18	11		626 APCI [M+H] ⁺

18	12		634 APCI [M+H] ⁺
18	13		576 APCI [M+H] ⁺
18	14		548 APCI [M+H] ⁺
18	15		550 APCI [M+H] ⁺
18	16		536 APCI [M+H] ⁺
18	17		522 APCI [M+H] ⁺

18	18		548 APCI [M+H] ⁺
18	19		576 APCI [M+H] ⁺
18	20		523 APCI [M+H] ⁺
18	21		536 APCI [M+H] ⁺
18	22		550 APCI [M+H] ⁺
18	23		567 APCI [M+H] ⁺


18	24		537 APCI [M+H] ⁺
18	25		548 APCI [M+H] ⁺
18	26		523 APCI [M+H] ⁺
18	27		562 APCI [M+H] ⁺
18	28		564 APCI [M+H] ⁺
18	29		598 APCI [M+H] ⁺







18	30		612 APCI [M+H] ⁺
18	31		626 APCI [M+H] ⁺
18	32		597 APCI [M+H] ⁺
18	33		598 APCI [M+H] ⁺
18	34		597 APCI [M+H] ⁺
18	35		606 APCI [M+H] ⁺







18	36		560 APCI [M+H] ⁺
18	37		548 APCI [M+H] ⁺
18	38		563 APCI [M+H] ⁺
18	39		479 APCI [M+H] ⁺







EXAMPLE 19







Corresponding starting compounds were treated in the similar manner as EXAMPLE 11 to give the following compounds.






EXAMPLE No.	No	Structure	MS (m/z)
19	1		592 ESI ⁺ /UV [M+H] ⁺

19	2		576 ESI+/UV [M+H] ⁺
19	3		606 ESI+/UV [M+H] ⁺
19	4		590 ESI+/UV [M+H] ⁺
19	5		590 ESI+/UV [M+H] ⁺
19	6		620 ESI+/UV [M+H] ⁺
19	7		634 APCI [M+H] ⁺


19	8		576 APCI [M+H] ⁺
19	9		606 APCI [M+H] ⁺
19	10		592 APCI [M+H] ⁺
19	11		576 APCI [M+H] ⁺
19	12		606 APCI [M+H] ⁺
19	13		592 APCI [M+H] ⁺

19	14		606 APCI [M+H] ⁺
19	15		592 APCI [M+H] ⁺
19	16		606 APCI [M+H] ⁺
19	17		604 APCI [M+H] ⁺
19	18		590 APCI [M+H] ⁺
19	19		606 APCI [M+H] ⁺

19	20		604 APCI [M+H] ⁺
19	21		620 APCI [M+H] ⁺
19	22		626 APCI [M+H] ⁺
19	23		602 APCI [M+H] ⁺
19	24		612 APCI [M+H] ⁺
19	25		641 APCI [M+H] ⁺

19	26		619 APCI [M+H] ⁺
19	27		630 APCI [M+H] ⁺
19	28		602 APCI [M+H] ⁺
19	29		592 APCI [M+H] ⁺
19	30		605 APCI [M+H] ⁺

EXAMPLE 20






The compound of EXAMPLE (18-12) (640 mg, 1.01 mmol) was

dissolved in formic acid (10 ml). The mixture was stirred at room temperature for 24 hours, concentrated, neutralized with a saturated aqueous sodium carbonate solution and extracted with methylene chloride. The organic layer was washed sequentially with water and brine, dried over magnesium sulfate and concentrated in *vacuo*. The residue was purified by NH-silica gel column chromatography (5% methanol-chloroform) to give the above compound (385 mg, yield 72%) as a colorless solid.


MS (m/z) APCI: 534 [M+H]⁺

Corresponding starting compounds were converted in the similar manner as EXAMPLE 2 to the corresponding compounds having a tert-butoxycarbonyl group, and then the resulting starting compounds were treated in the similar manner as the above-mentioned to give the following compounds.

EXAMPLE No.	No	Structure	MS (m/z)
20	1		534 APCI [M+H] ⁺
20	2		534 APCI [M+H] ⁺
20	3		534 APCI [M+H] ⁺

*The compound (20-3) was isolated as a dihydrochloride salt.


EXAMPLE 21



The compound of EXAMPLE 19-(7) (2.7 g, 4.26 mmol) was dissolved in methanol (30 ml), and thereto was added potassium carbonate (600 mg, 4.26 mmol) at room temperature. The mixture was stirred at room temperature for 4 hours and concentrated, and thereto was added water. The mixture was extracted with ethyl acetate, and the organic layer was washed sequentially with water and brine, dried over sodium sulfate and concentrated in *vacuo*. The residue was purified by silica gel column chromatography (1.5 to 10% methanol-ethyl acetate) to give the above compound (1.7 g, yield 68%) as a colorless solid.

MS (m/z) APCI: 592[M+H]⁺

EXAMPLE 22







A solution of the compound of EXAMPLE 3 (80 mg, 0.15 mmol), difluoroacetic acid (0.028 ml, 0.45 mmol) and N-ethyl-N'-(3-dimethylaminopropyl)carbodiimide hydrochloride (116 mg, 0.60 mmol) in chloroform (3 ml) was stirred at room temperature for 7 hours. The

reaction mixture was poured into a saturated aqueous sodium bicarbonate solution and the mixture was extracted with ethyl acetate. The organic layer was dried over sodium sulfate, concentrated in *vacuo*, and the residue was purified by silica gel chromatography (0 to 5% methanol-chloroform) to give the above compound (85.6 mg, yield 93%) as a colorless solid.

MS (m/z) APCI: 612[M+H]⁺

Corresponding compounds were reacted in the similar manner as the above-mentioned to give the following compounds.

EXAMPLE No.	No	Structure	MS (m/z)
22	1		620 APCI [M+H] ⁺
22	2		646 APCI [M+H] ⁺
22	3		646 APCI [M+H] ⁺




The compound of EXAMPLE 3 (80 mg, 0.13 mmol) was dissolved in ethyl formate (3 ml), and the mixture was heated to reflux for 32 hours. The reaction mixture was concentrated and the residue was purified by silica gel column chromatography (0 to 8% methanol-chloroform) to give the above compound (72.4 mg, yield 98%) as a colorless solid.

MS (m/z) APCI: 562[M+H]⁺





Corresponding starting compounds were reacted in the similar manner as the above-mentioned to give the following compounds.







EXAMPLE No.	No	Structure	MS (m/z)
23	1		562 APCI [M+H] ⁺
23	2		562 APCI [M+H] ⁺
23	3		562 APCI [M+H] ⁺







23	4		562 APCI [M+H] ⁺
----	---	---	--------------------------------







EXAMPLE 24







Corresponding starting compounds were treated in the similar manner as EXAMPLE 1-(5) to give the following compounds.







EXAMPLE No.	No	Structure	MS (m/z)
24	1		440 APCI [M+H] ⁺
24	2		456/458 APCI [M+H] ⁺
24	3		436 APCI [M+H] ⁺
24	4		529 APCI [M+H] ⁺

24	5		423 APCI [M+H] ⁺
24	6		437 APCI [M+H] ⁺
24	7		416 APCI [M+H] ⁺
24	8		450/452 APCI [M+H] ⁺
24	9		417 APCI [M+H] ⁺
24	10		495/497 APCI [M+H] ⁺





24	11		507/509 APCI [M+H] ⁺
24	12		468 APCI [M+H] ⁺
24	13		447 APCI [M+H] ⁺
24	14		508 APCI [M+H] ⁺
24	15		584 APCI [M+H] ⁺
24	16		467 APCI [M+H] ⁺

24	17		472 APCI [M+H] ⁺
24	18		517 APCI [M+H] ⁺
24	19		502 APCI [M+H] ⁺
24	20		494/496 APCI [M+H] ⁺
24	21		430 APCI [M+H] ⁺
24	22		500/502 APCI [M+H] ⁺


24	23		490 APCI [M+H] ⁺
24	24		463 APCI [M+H] ⁺
24	25		503 APCI [M+H] ⁺
24	26		441 APCI [M+H] ⁺
24	27		469 APCI [M+H] ⁺
24	28		490 APCI [M+H] ⁺

24	29		434 APCI [M+H] ⁺
24	30		473 APCI [M+H] ⁺
24	31		494 APCI [M+H] ⁺
24	32		474 APCI [M+H] ⁺
24	33		451 APCI [M+NH ₄] ⁺
24	34		521 APCI [M+H] ⁺

24	35		507 APCI [M+H] ⁺
24	36		443 APCI [M+H] ⁺
24	37		451 APCI [M+H] ⁺
24	38		465 APCI [M+H] ⁺
24	39		457 APCI [M+H] ⁺
24	40		647 APCI [M+H] ⁺

24	41		471 APCI [M+H] ⁺
24	42		463 APCI [M+H] ⁺
24	43		522 APCI [M+NH ₄] ⁺
24	44	 Chiral	466 APCI [M+H] ⁺

EXAMPLE 25



- (1) To a solution of N-hydroxyphthalimide (142 g, 868 mmol), triphenylphosphine (252.9 g, 964 mmol) and (S)-3-hydroxytetrahydrofuran (70.7 g, 804 mmol) in THF (2800 ml) was added dropwise diisopropyl azodicarboxylate (195.0 g, 964 mmol) over 1.5 hours under ice-cooling. The mixture was stirred at room temperature for 16 hours and concentrated in *vacuo*. The residue was dissolved in ethanol (800 ml), and thereto was added hydrazine monohydrate (43.4 g, 867 mmol) at room temperature, and the mixture was heated to reflux for 4 hours and stirred at room temperature for another 40 hours. To the reaction mixture were added ethanol (500 ml) and a 4N hydrogen chloride solution in dioxane (300 ml, 1200 mmol). The precipitated crystals were filtered off. The filtrate was concentrated, and the residue was recrystallized from ethyl acetate to give the compound (25-A, monohydrochloride) (92.6 g, yield 83%) as colorless crystals.

MS (m/z) APCI: 104 [M+H]⁺

- (2) To a solution of the compound (1-C) (3.7 g, 13.8 mmol) in methanol (70 ml) was added sodium triacetoxyborohydride (7.33 g, 34.6

mmol) under ice-cooling. The mixture was stirred at the same temperature for 20 minutes, then the ice bath was removed. The reaction mixture was stirred at room temperature for another 3 hours, concentrated in *vacuo*, and then the residue was dissolved in ethyl acetate, washed sequentially with water and brine, dried over sodium sulfate and concentrated. The resulting crude ester (4.2 g) was dissolved in a mixed solvent of methanol (40 ml) and water (10 ml), and thereto was added a 2N aqueous sodium hydroxide solution (10.4 ml) under ice-cooling. The mixture was stirred at room temperature for 14 hours, concentrated, acidified with 10% hydrochloric acid and extracted with ethyl acetate. The organic layer was washed sequentially with water and brine, dried over sodium sulfate and concentrated. The residue was recrystallized from ethyl acetate-hexane to give the compound (25-B) (3.30 g, yield 93%) as colorless crystals.

MS (m/z) ESI: 255 [M-H]⁻

(3) A solution of the above compound (1.58 g, 6.15 mmol), 2-aminothiazole (1.23 g, 12.3 mmol) and N,N-dimethylaminopyridine (1.13 g, 9.25 mmol) in chloroform (30 ml) was ice-cooled, and thereto was added N-ethyl-N'-(3-dimethylaminopropyl)carbodiimide hydrochloride (1.77 g, 9.23 mmol). The reaction mixture was stirred at room temperature for 20 hours, diluted with ethyl acetate, washed sequentially with a 10% aqueous citric acid solution, brine, a saturated aqueous sodium bicarbonate solution and brine, dried over sodium sulfate and then concentrated in *vacuo*. The residue was purified by silica gel chromatography (0 to 10% methanol-chloroform) to give crude crystals. The resulting crude crystals were washed with diethyl ether to give the compound (25-C) (1.15 g, yield 55%) as colorless crystals.

MS (m/z) APCI: 339 [M+H]⁺

(4) To a solution of the above compound (957 mg, 2.83 mmol) in dimethylsulfoxide (30 ml) were added sequentially triethylamine (3.94


ml, 28.3 mmol) and sulfur trioxide-pyridine complex (2.25 g, 14.2 mmol) at room temperature, and the mixture was stirred at the same temperature for 1.5 hours. To the reaction mixture was added water, and the mixture was extracted with ethyl acetate. The organic layer
5 was washed sequentially with water and brine, dried over sodium sulfate and concentrated. The residue was solidified with diisopropyl ether to give the compound (25-D) (690 mg, yield 73%) as a colorless solid.

MS (m/z) APCI: 337 [M+H]⁺

10 (5) To a solution of the above compound (107 mg, 0.31 mmol) and the compound (25-A, monohydrochloride) (89 mg, 0.64 mmol) in methanol-THF (1:1) (4 ml) was added pyridine (0.068 ml, 0.80 mmol), and the mixture was stirred at room temperature for 16 hours and then heated to reflux for 2 hours. To the reaction mixture was added water,
15 and the mixture was extracted with ethyl acetate. The organic layer was washed sequentially with a 10% citric acid solution, water and brine, dried over sodium sulfate and concentrated. The residue was purified by silica gel column chromatography (0 to 10% methanol-chloroform) to give the compound (25-E) (76 mg, (E)-isomer, yield 32%)
20 and the corresponding (Z)-isomer (76 mg, yield 56%) as a colorless solid each.

MS (m/z) APCI: 422 [M+H]⁺

EXAMPLE 26




25 To a solution of the compound of EXAMPLE 1 (958 mg, 2.13 mmol) in methanol (40 ml) was added sodium borohydride (160 mg,

4.26 mmol) under ice-cooling, and the mixture was stirred for 2 hours at the same temperature. Acetone (1 ml) was added and the mixture was concentrated in *vacuo*. The resulting residue was purified by silica gel column chromatography (3 to 10% methanol-chloroform) to give the above compound (976 mg, yield 100%) as a colorless solid.

MS (m/z) APCI: 452 [M+H]⁺

EXAMPLE 27





To a solution of the compound of EXAMPLE 24-(12) (150 mg, 0.32 mmol) in methylene chloride (3 ml) was added m-chloroperbenzoic acid (110 mg, 0.48 mmol) under ice-cooling. The mixture was stirred for 2 hours at the same temperature and concentrated in *vacuo*. The resulting residue was purified by silica gel column chromatography (1 to 8% methanol-chloroform) to give the above compound (61 mg, yield 38%) as a colorless solid.

MS (m/z) APCI: 500 [M+H]⁺


EXAMPLE 28

The following compounds were synthesized by treating the compounds of EXAMPLE 24-(33) in the similar manner as EXAMPLE 2.

EXAMPLE No.	No	Structure	MS (m/z)
28	1		563 ESI+ [M+H] ⁺

28	2		549 ESI+ [M+H] ⁺
28	3		522 ESI [M+H] ⁺


EXAMPLE 29



To a solution of the compound of EXAMPLE 1 (1.28 g, 2.85 mmol) in THF (30 ml) was added a 3M solution of methyl magnesium bromide in diethyl ether (2 ml, 5.98 mmol) at -78°C , and then the mixture was warmed to 0°C and stirred for 1 hour at the same temperature. To the reaction mixture was added a saturated aqueous ammonium chloride solution, and the mixture was extracted with ethyl acetate. The organic layer was washed sequentially with water and brine, dried over magnesium sulfate and concentrated in *vacuo*. The resulting residue was purified by silica gel column chromatography (2 to 10% methanol-chloroform) to give the above compound (1.03 g, yield 78%) as a colorless solid.

MS (m/z) APCI: 466 [M+H]⁺


EXAMPLE 30



To a solution of the compound of EXAMPLE 24-(13) (100 mg, 0.224 mmol) in ethanol-water (1:1) (6 ml) were added sodium carbonate (40.4 mg, 0.38 mmol) and hydroxylamine hydrochloride (57.6 mg, 0.83 mmol) at room temperature. The reaction mixture was heated to reflux for 3 hours and extracted with chloroform, and the organic layer was washed sequentially with water and brine, dried over sodium sulfate and concentrated in *vacuo*. The resulting residue was purified by silica gel column chromatography (5 to 10% methanol-chloroform) to give the above compound (30 mg, yield 28%) as a colorless solid.

MS (m/z) APCI: 480 [M+H]⁺


EXAMPLE 31



To a solution of the compound of EXAMPLE 29 (100 mg, 0.21 mmol) in methylene chloride (7 ml) was added manganese dioxide (1 g) at room temperature. The mixture was stirred for 12 hours at the same temperature and filtered through Celite. The filtrate was concentrated in *vacuo*, and the resulting residue was purified by silica gel column chromatography (2 to 5% methanol-chloroform) to give the above compound (77.6 mg, yield 78%) as a colorless solid.

MS (m/z) APCI: 464 [M+H]⁺


EXAMPLE 32



To a solution of the compound of EXAMPLE 24-(13) (100 mg, 0.22 mmol) in acetone-water (4:1) (5 ml) were added potassium carbonate (31 mg, 0.22 mmol) and a 30% aqueous hydrogen peroxide solution (0.2 ml) under ice-cooling, and the mixture was stirred at room temperature for 38 hours. To the reaction mixture was added a 10% aqueous sodium sulfite solution, and then the mixture was extracted with a mixed solvent of 10% methanol-chloroform. The organic layer was dried over sodium sulfate and concentrated in *vacuo*, and the resulting residue was purified by silica gel column chromatography (10% methanol-chloroform) to give the above compound (74 mg, yield 71%) as a colorless solid.

MS (m/z) APCI: 465 [M+H]⁺


EXAMPLE 33



The above compound was obtained as a colorless solid by reacting the compound of EXAMPLE 31 in the similar manner as EXAMPLE 29.

MS (m/z) APCI: 480 [M+H]⁺


EXAMPLE 34



To a solution of the compound of EXAMPLE 29 (100 mg, 0.21 mmol) and vinyl acetate (0.4 ml, 4.30 mmol) in ethyl acetate (5 ml) was added Lipase PS (manufactured by Amano Pharmaceutical Co., Ltd.) (1.0 g) at room temperature, and the mixture was stirred at the same temperature for 3 days. The reaction mixture was filtered through Celite and the filtrate was concentrated in *vacuo*. The resulting residue was purified by silica gel column chromatography (0 to 5% methanol-ethyl acetate) to give the above compound (36.5 mg, yield 37%) as a colorless solid.

MS (m/z) APCI: 466 [M+H]⁺


EXAMPLE 35



To a solution of the compound of EXAMPLE 24-(14) (1.06 g, 2.09 mmol) in ethanol (30 ml) was added a 2N aqueous sodium hydroxide solution (2.09 ml, 4.18 mmol) under ice-cooling. The mixture was stirred at room temperature for 5 hours, acidified with 2N hydrochloric acid and extracted with methylene chloride. The organic layer was washed sequentially with water and brine, dried over magnesium sulfate and concentrated in *vacuo*. The resulting residue was solidified with diethyl ether to give the compound (920 mg, yield 92%) as a colorless solid.

MS (m/z) ESI: 478 [M-H]⁻


EXAMPLE 36



To a solution of the compound of EXAMPLE 24-(14) (100 mg, 0.20 mmol) in THF (4 ml) was added lithium borohydride (17.2 mg, 0.79 mmol) under ice-cooling, and then the mixture was stirred at room temperature for 24 hours. To the reaction mixture was added methanol (4 ml) and oxalic acid (100 mg), and the mixture was stirred at room temperature for another 24 hours, and thereto was added ethyl acetate. The organic layer was separated, washed sequentially with a saturated aqueous sodium carbonate solution, water and brine, dried over magnesium sulfate and concentrated in *vacuo*. The residue was purified by silica gel column chromatography (1 to 7% methanol-chloroform) to give the above compound (27.6 mg, yield 30%) as a colorless solid.

MS (m/z) APCI: 466 [M+H]⁺

EXAMPLE 37





To a solution of the compound of EXAMPLE 35 (200 mg, 0.42 mmol), dimethylamine hydrochloride (102 mg, 1.25 mmol) and 1-


hydroxybenzotriazole (169 mg, 1.25 mmol) in methylene chloride (6 ml) was added dropwise N-ethyl-N'-(3-diethylaminopropyl)carbodiimide (0.226 ml, 1.25 mmol) at room temperature. The mixture was stirred at the same temperature for 24 hours, diluted with methylene chloride, washed sequentially with a saturated aqueous sodium carbonate solution, water and brine, dried over magnesium sulfate and concentrated in *vacuo*. The residue was purified by NH-silica gel column chromatography (5% methanol-chloroform) to give the above compound (212 mg, yield 100%) as a colorless solid.

MS (m/z) APCI: 507 [M+H]⁺

The following compounds were synthesized by reacting the corresponding starting compounds in the similar manner as the above-mentioned.

EXAMPLE No.	No	Structure	MS (m/z)
37	1		493 APCI [M+H] ⁺
37	2		509 APCI [M+H] ⁺


EXAMPLE 38



To a solution of the compound of EXAMPLE 24-(36) (66 mg, 0.15 mmol) in ethanol-THF (1:1) (6 ml) was added 10% Pd/C (60 mg). The mixture was stirred at room temperature for 12 under hydrogen at normal pressure, filtered through Celite and then the filtrate was concentrated in *vacuo* to give the above compound (67 mg, yield 100%) as a colorless solid.

MS (m/z) APCI: 445 [M+H]⁺


EXAMPLE 39



To a solution of the compound of EXAMPLE 24-(22) (500 mg, 1.0 mmol) in DMF (15 ml) were added tributyl(2-methyl-1-propenyl)tin (690 mg, 2.0 mmol), diisopropylethylamine (0.87 ml, 5.0 mmol), lithium chloride (296 mg, 7.0 mmol) and tetrakis(triphenylphosphine)palladium (58 mg, 0.05 mmol) under argon. The mixture was stirred at 120°C for 4 hours, diluted with ethyl acetate, and thereto was added water and then filtered through Celite. The filtrate was washed sequentially with water and brine, dried over sodium sulfate and concentrated in *vacuo*. The resulting residue was purified by NH-silica gel column chromatography (50 to 100% chloroform-hexane) to give the above compound (268 mg, yield 56%) as a colorless solid.

MS (m/z) APCI: 476 [M+H]⁺

EXAMPLE 40




To a solution of the compound of EXAMPLE 39 (95 mg, 0.20 mmol) in acetone-acetonitrile-water (1:1:1) (6 ml) were added N-methylmorpholine N-oxide (59 mg, 0.50 mmol) and 10% microencapsulated osmium tetroxide (Osmium (VIII) Oxide, Microencapsulate™, Wako Pure Chemical Industries, Ltd., 102 mg, 0.04 mmol). The mixture was stirred at room temperature for 2 days, diluted with ethyl acetate and filtered through Celite, and then the filtrate was concentrated in *vacuo*. The residue was purified by silica gel column chromatography (3-10% methanol-chloroform) to give the above compound (96 mg, yield 94%) as a colorless solid.

MS (m/z) APCI: 510 [M+H]⁺

The following compound was synthesized by treating the corresponding starting compound in the similar manner as the above-mentioned.

EXAMPLE No.	No	Structure	MS (m/z)
40	1	<chem>CC(C)(O)C1=CN=C(NC(=O)N=C(N1)c2ccc(cc2)S(=O)(=O)C3CC3)S1</chem>	505 APCI [M+H] ⁺

EXAMPLE 41



(1) A corresponding starting compound and the compound (1-E) were treated in the similar manner as EXAMPLE 1-(5) to give the compound (41-A).

5 MS (m/z) APCI: 519 [M+H]⁺


(2) The above compound (72 mg, 0.14 mmol) was dissolved in formic acid (3 ml) at room temperature. The mixture was stirred at the same temperature for 24 hours and concentrated. The residue was purified by silica gel column chromatography (0 to 8% methanol-chloroform) to give the compound (41-B) (50.7 mg, yield 87%) as a colorless solid.

10 MS (m/z) APCI: 419 [M+H]⁺

The following compound was synthesized by treating the corresponding starting compound in the similar manner as the above-mentioned.

EXAMPLE No.	No	Structure	MS (m/z)
41	1		405 APCI [M+H] ⁺

15 EXAMPLE 42





To a solution of the compound of EXAMPLE 26 (300 mg, 0.66 mmol) in methylene chloride (10 ml) were added sequentially triethylamine (0.28 ml, 1.99 mmol) and acetic anhydride (0.095 ml, 1.0 mmol) at room temperature. The mixture was stirred at room temperature for 16 hours, concentrated in *vacuo* and then to the residue was added ethylene glycol (15 ml). The mixture was heated to reflux for 8 hours, diluted with ethyl acetate, and then washed sequentially with water and brine, dried over sodium sulfate and concentrated in *vacuo*. The resulting residue was purified by silica gel column chromatography (0 to 5% methanol-ethyl acetate) to the above compound (166 mg, yield 50%) as a colorless solid.

MS (m/z) APCI: 496 [M+H]⁺




The following compounds were synthesized by treating the corresponding starting compounds in the similar manner as the above-mentioned.







EXAMPLE No.	No	Structure	MS (m/z)
42	1		510 APCI [M+H] ⁺
42	2		510 APCI [M+H] ⁺
42	3		480 APCI [M+H] ⁺







42	4		494 APCI [M+H] ⁺
42	5		524 APCI [M+H] ⁺







EXAMPLE 43




Corresponding starting compounds were treated in the similar manner as EXAMPLE 1-(5) to give the following compounds.

EXAMPLE No.	No	Structure	MS (m/z)
43	1		431 APCI [M+H] ⁺
43	2		525 APCI [M+H] ⁺
43	3		535 APCI [M+H] ⁺


43	4		597 APCI [M+H] ⁺
43	5		463 APCI [M+H] ⁺
43	6		474 APCI [M+H] ⁺
43	7		550/552 ESI [M-H] ⁻
43	8		504 APCI [M+H] ⁺
43	9		474 APCI [M+H] ⁺

43	10		504 APCI [M+H] ⁺
43	11		516 APCI [M+H] ⁺
43	12		532 APCI [M+H] ⁺
43	13		554 APCI [M+H] ⁺
43	14		540 APCI [M+H] ⁺
43	15		491 APCI [M+H] ⁺

43	16		530 APCI [M+H] ⁺
43	17		571 APCI [M+H] ⁺
43	18		447 APCI [M+H] ⁺
43	19		460 APCI [M+H] ⁺
43	20		490 APCI [M+H] ⁺
43	21		450 APCI [M+H] ⁺

43	22		433 APCI [M+H] ⁺
43	23		467 APCI [M+H] ⁺
43	24		419 APCI [M+H] ⁺

EXAMPLE 44



- (1) To a solution of the compound (1-E) (200 mg, 0.59 mmol) and 2-(2-aminothiazol-5-ylsulfany)ethanol (104 mg, 0.59 mmol) in THF (4 ml) was added DMT-MM (4-(4,6-dimethoxy-1,3,5-triazin-2-yl)-4-methylmorpholinium chloride) (240 mg, 0.87 mmol) at room


temperature. The mixture was stirred at the same temperature for 20 hours, diluted with ethyl acetate, and then washed sequentially with water and brine, dried over magnesium sulfate and concentrated in *vacuo*. The residue was purified by NH-silica gel column chromatography (0 to 1% methanol-chloroform) to give the compound (44-A) (82 mg, yield 17%) as a pale yellow solid.

MS (m/z) APCI: 819 [M+H]⁺


(2) To a solution of the above compound (78 mg, 0.095 mmol) in THF-methanol-water (10:3:3) (1.6 ml) was added a 2N aqueous sodium hydroxide solution (0.12 ml, 0.24 mmol) under ice-cooling. The mixture was stirred at the same temperature for 2.5 hours, diluted with ethyl acetate, and then washed sequentially with water and brine, dried over magnesium sulfate and concentrated in *vacuo*. The residue was purified by silica gel chromatography (80 to 100% ethyl acetate-hexane) to give the compound (44-B) (31 mg, yield 66%).

MS (m/z) APCI: 498 [M+H]⁺

A corresponding starting compound was treated in the similar manner as the above-mentioned to give the following compound.

EXAMPLE No.	No	Structure	MS (m/z)
44	1		512 APCI [M+H] ⁺


EXAMPLE 45



To a solution of the compound of EXAMPLE 44-(1) (100 mg, 0.20 mmol) and carbon tetrabromide (321 mg, 0.97 mmol) in THF (4 ml) was added triphenylphosphine (256 mg, 0.97 mmol) at -10°C , and the mixture was stirred at room temperature for 38 hours, and then concentrated in *vacuo*. The residue was purified by NH-silica gel column chromatography (50 to 90% ethyl acetate-hexane) to give the above compound (176 mg, quantitatively).

MS (m/z) APCI: 574/576 $[\text{M}+\text{H}]^{+}$

EXAMPLE 46




To a solution of the compound of EXAMPLE 43-(14) (1800 mg, 3.34 mmol) in ethanol-THF (1:3) (26 ml) was added a 2N aqueous sodium hydroxide solution (5.0 ml, 10 mmol) under ice-cooling. The mixture was stirred at the same temperature for 2 hours, concentrated in *vacuo*, and then the residue was acidified with 2N hydrochloric acid. After diluting with ethyl acetate, the mixture was washed sequentially with water and brine, dried over sodium sulfate and concentrated in *vacuo*. The residue was solidified with ethyl acetate-hexane to give the above compound (1631 mg, yield 96%) as colorless crystals.


MS (m/z) APCI: 510 $[\text{M}-\text{H}]^{-}$

A corresponding starting compound was treated in the similar manner as the above-mentioned to give the following compound.

EXAMPLE No.	No	Structure	MS (m/z)
-------------	----	-----------	----------

46	1		524 ESI [M-H] ⁻
----	---	---	----------------------------


EXAMPLE 47




The compound of EXAMPLE 43-(12) (287 mg, 0.54 mmol) was dissolved in formic acid (3 ml). The mixture was stirred at room temperature for 24 hours at the same temperature and concentrated in *vacuo*. The residue was dissolved in chloroform, washed sequentially with a saturated aqueous sodium bicarbonate solution, water and brine, dried over sodium sulfate and concentrated in *vacuo*. The residue was purified by silica gel column chromatography (0 to 5% methanol-ethyl acetate) to give the above compound (180 mg, yield 77%) as a colorless solid.

MS (m/z).APCI: 432 [M+H]⁺





Corresponding starting compounds were treated in the similar manner as the above-mentioned to give the following compounds.



EXAMPLE No.	No	Structure	MS (m/z)
47	1		497 APCI [M+H] ⁺

47	2		511 APCI [M+H] ⁺
----	---	---	--------------------------------


EXAMPLE 48

Corresponding starting compounds were treated in the similar manner as EXAMPLE 37 to give the following compounds.

EXAMPLE No.	No	Structure	MS (m/z)
48	1		511 APCI [M+H] ⁺
48	2		525 APCI [M+H] ⁺
48	3		539 APCI [M+H] ⁺
48	4		525 APCI [M+H] ⁺

48	5		539 APCI [M+H] ⁺
48	6		553 APCI [M+H] ⁺

EXAMPLE 49




To a solution of the compound of EXAMPLE 24-(10) (248 mg, 0.50 mmol) in DMF (3 ml) were added tetrakis(triphenylphosphine)palladium (57 mg, 0.05 mmol) and 2-mercaptoethanol (98 mg, 1.25 mmol) under argon. The mixture was stirred at 120°C for 3 hours, diluted with ethyl acetate, and then washed sequentially with water and brine, dried over sodium sulfate and concentrated in *vacuo*. The residue was purified by silica gel column chromatography (50 to 100% ethyl acetate-hexane) to give the above compound (72.1 mg, yield 29%) as a colorless solid.


MS (m/z) APCI: 493 [M+H]⁺

A corresponding starting compound was treated in the similar manner as the above-mentioned to give the following compound.

EXAMPLE No.	No	Structure	MS (m/z)
-------------	----	-----------	----------

49	1		520 APCI [M+H] ⁺
----	---	---	--------------------------------


EXAMPLE 50



A corresponding starting compound was treated in the similar manner as EXAMPLE 1-(5) to give the above compound.

5 MS (m/z) APCI: 617 [M+H]⁺

EXAMPLE 51




To a solution of the compound of EXAMPLE 50 (810 mg, 1.31 mmol) in THF (10 ml) was added a 1.0 M solution of tetrabutylammonium fluoride in THF (3.9 ml, 3.9 mmol) under ice-cooling. The mixture was stirred at room temperature for 3 hours, diluted with ethyl acetate, washed sequentially with water and brine, dried over sodium sulfate and concentrated in *vacuo*. The residue was purified by silica gel column chromatography (0 to 8 % methanol-chloroform) to give the above compound (646 mg, yield 98%).

10

15

MS (m/z) APCI: 503 [M+H]⁺


EXAMPLE 52



To a solution of the compound of EXAMPLE 26 (296 mg, 0.66 mmol) in chloroform (3 ml) were added trifluoroacetic acid (2 ml) and sodium thiomethoxide (183 mg, 2.61 mmol). The mixture was stirred at 60°C in a sealed tube under microwave irradiation for 3 hours, diluted with ethyl acetate, washed sequentially with a saturated aqueous sodium bicarbonate solution, water and brine, dried over sodium sulfate and concentrated in *vacuo*. The residue was purified by silica gel column chromatography (ethyl acetate) to give the above compound (281 mg, yield 89%).

MS (m/z) APCI: 482 [M+H]⁺

EXAMPLE 53




To a solution of the compound of EXAMPLE 52 (176 mg, 0.365 mmol) in chloroform (5 ml) was added 70% m-chloroperbenzoic acid (225 mg, 0.913 mmol) under ice-cooling, and the mixture was stirred at the same temperature for 1 hour and at room temperature for another 3 hours. To the reaction mixture was added a 10% sodium sulfite solution, and the organic layer was separated, dried over sodium sulfate and concentrated in *vacuo*. The residue was purified by silica gel column chromatography (0 to 5% methanol-chloroform) to give the

above compound (148 mg, yield 79%).

MS (m/z) APCI: 514 [M+H]⁺


EXAMPLE 54



- 5 To a solution of the compound of EXAMPLE 45 (50 mg, 0.087 mmol) in methanol (1.5 ml) was added N,N-dimethyltrimethylsilylamine (459 mg, 3.92 mmol) under ice-cooling. The mixture was stirred at the same temperature for 30 minutes and at room temperature for another 18 hours, and concentrated in *vacuo*. The residue was purified by silica
10 gel column chromatography (0 to 6% methanol-chloroform) to give the above compound (41.2 mg, yield 88%).

MS (m/z) APCI: 539 [M+H]⁺

EXAMPLE 55




- 15 To a solution of the compound of EXAMPLE 26 (908 mg, 2.01 mmol) and 4-hydroxypiperidine (610 mg, 6.03 mmol) in toluene (60 ml) was added p-toluenesulfonic acid monohydrate (3.65 g, 19.2 mmol) at room temperature, and the mixture was heated to reflux for 1 hour using a Dean-Stark apparatus for azeotropic removal of the resulting
20 water. After standing to cool, to the reaction mixture was added a saturated aqueous sodium bicarbonate solution, and the mixture was


extracted with chloroform. The organic layer was dried over sodium sulfate and concentrated in *vacuo*. The residue was purified by silica gel column chromatography (aqueous ammonia-methanol-chloroform, 1:10:100) to give the above compound (461 mg, yield 43%).

5 MS (m/z) APCI: 535 [M+H]⁺

A corresponding starting compound was treated in the similar manner as the above-mentioned to give the following compound.

EXAMPLE No.	No	Structure	MS (m/z)
55	1		502 APCI [M+H] ⁺

EXAMPLE 56




10 To a solution of the compound of EXAMPLE 55 (107 mg, 0.2 mmol) and pyridine (0.081 ml, 1.0 mmol) in chloroform (5 ml) was added acetic anhydride (0.0284 ml, 0.3 mmol) at room temperature, and the mixture was stirred at the same temperature for 3 hours. To the reaction mixture was added a saturated aqueous sodium bicarbonate solution, and the mixture was extracted with chloroform.

15 The organic layer was dried over sodium sulfate, and concentrated in *vacuo*, and then the residue was purified by silica gel column chromatography (0 to 10% methanol-chloroform) to give the above


compound (105 mg, yield 91%).

MS (m/z) APCI: 577 [M+H]⁺

A corresponding starting compound was treated in the similar manner as the above-mentioned to give the following compound.

EXAMPLE No.	No	Structure	MS (m/z)
56	1		474 APCI [M+H] ⁺


5 EXAMPLE 57



To a mixture of the compound of EXAMPLE 55 (111 mg, 0.208 mmol) and a 38% aqueous formalin solution (1 ml) in chloroform (3 ml) was added sodium triacetoxyborohydride (132 mg, 0.603 mmol) under ice-cooling, and the mixture was stirred at room temperature for 3 hours. To the reaction mixture was added a saturated aqueous sodium bicarbonate solution, and the organic layer was separated, dried over magnesium sulfate and concentrated. The residue was purified by silica gel column chromatography (0 to 20% methanol-chloroform) to give the above compound (106 mg, yield 93%).

MS (m/z) APCI: 549 [M+H]⁺

EXAMPLE 58



(1) The compound of EXAMPLE 24-(36) was treated in the similar manner as EXAMPLE 40 to give the compound (58-A).

MS (m/z) APCI: 477 [M+H]⁺

5 (2) To a solution of the above compound (477 mg, 1.0 mmol) in acetone (10 ml) was added an aqueous solution of sodium periodate (235 mg, 1.10 mmol) (10 ml) under ice-cooling. The mixture was stirred at the same temperature for 2 hours, diluted with ethyl acetate, and then washed sequentially with a 1M aqueous sodium sulfite solution, 10 water and brine, dried over sodium sulfate and concentrated in *vacuo*. The resulting residue was purified by silica gel column chromatography (0 to 5% methanol-ethyl acetate) to give the compound (58-B) (401 mg, yield 90%).

MS (m/z) APCI: 445 [M+H]⁺

15 EXAMPLE 59


A corresponding starting compound was treated in the similar manner as EXAMPLE 29 to give the following compound.

EXAMPLE No.	No	Structure	MS (m/z)
59			461 APCI [M+H] ⁺

EXAMPLE 60



20 A corresponding starting compound was treated in the similar manner as EXAMPLE 31 to give the following compound.

EXAMPLE No.	No	Structure	MS (m/z)
-------------	----	-----------	----------


60			459 APCI [M+H] ⁺
----	--	---	--------------------------------

EXAMPLE 61

Corresponding starting compounds were treated in the similar manner as EXAMPLE 2 to give the following compounds.

EXAMPLE No.	No	Structure	MS (m/z)
61	1		529 APCI [M+H] ⁺
61	2		543 APCI [M+H] ⁺

EXAMPLE 62



5







The compound (1-D), 2-methoxyethanol and the corresponding starting compound were reacted in the similar manner as EXAMPLE 1 to give the above compound.







MS (m/z) APCI: 428 [M+H]⁺



10

Corresponding starting compounds were reacted in the similar manner as the above-mentioned to give the following compounds.


EXAMPLE	No	Structure	MS (m/z)
---------	----	-----------	----------

No.			
62	1		553 APCI [M+H] ⁺
62	2		553 APCI [M+H] ⁺
62	3		442 ESI ⁺ [M+H] ⁺
62	4		481 APCI [M+H] ⁺
62	5		462 APCI [M+H] ⁺
62	6		450 APCI [M+H] ⁺

62	7		472 APCI [M+H] ⁺
62	8		465 APCI [M+H] ⁺
62	9		505 APCI [M+H] ⁺
62	10		464 APCI [M+H] ⁺
62	11		558 APCI [M+H] ⁺
62	12		476 APCI [M+H] ⁺

62	13		467 APCI [M+H] ⁺
62	14		467 APCI [M+H] ⁺

EXAMPLE 63



(1) To a solution of the compound (1-D) (2.5 g, 8.8 mmol) and potassium carbonate (2.44 g, 17.7 mmol) in DMF (50 ml) was added bromoacetonitrile (0.737 ml, 10.6 mmol) at room temperature. The mixture was stirred at the same temperature for 15 hours, diluted with ethyl acetate, and then washed sequentially with water and brine, dried over sodium sulfate and concentrated in *vacuo*. The resulting residue was purified by silica gel column chromatography (40% ethyl acetate-hexane) to give the compound (63-A) (2.65 g, yield 93%).


MS (m/z) APCI: 340 [M+H]⁺

(2) To a solution of the above compound (2.8 g, 8.8 mmol) in methanol-THF (3:1) (20 ml) was added a 2N aqueous sodium hydroxide solution (4.4 ml, 8.8 mmol) under ice-cooling, and the ice bath was removed. The mixture was stirred at the same temperature for 30 minutes, and extracted with methylene chloride. The organic layer was washed sequentially with water and brine, dried over magnesium sulfate and concentrated in *vacuo* to give the compound (63-B) (2.8 g, quantitatively).

(3) A corresponding starting compound was reacted with the above compound in the similar manner as EXAMPLE 1-(5) to give the compound (63-C).

MS (m/z) APCI: 409 [M+H]⁺

EXAMPLE 64




(1) The compound of EXAMPLE 62-(1) (332 mg, 0.601 mmol) was dissolved in formic acid (9 ml). The mixture was stirred at room temperature for 24 hours and concentrated in *vacuo*. The residue was dissolved in ethyl acetate and thereto was added a 4N hydrogen chloride solution in dioxane, and the precipitated crystals were collected to give a crude amine (308 mg) as a monohydrochloride.

(2) To a solution of the above crude amine (60 mg) and pyridine (0.046 ml, 0.545 mmol) in chloroform (1 ml) was added acetic anhydride (0.015 ml, 0.163 mmol) at room temperature, and the mixture was stirred at the same temperature for 20 hours and then concentrated in


vacuo. The resulting residue was purified by NH-silica gel column chromatography (0 to 10% methanol-chloroform) to give the titled compound (50.4 mg, yield 87% in 2 steps).

MS (m/z) APCI: 495 [M+H]⁺

- 5 A corresponding starting compound was treated in the similar manner as the above-mentioned to give the following compound.

EXAMPLE No.	No	Structure	MS (m/z)
64	1		495 APCI [M+H] ⁺

EXAMPLE 65



- 10 (1) To a solution of the compound (1-C) (10 g, 37.2 mmol) and cyclopentylhydroxylamine (8.4 g, 83.0 mmol) in methanol (100 ml) was added pyridinium p-toluenesulfonate (10 mg, 0.04 mmol). The mixture was stirred at room temperature for 36 hours and concentrated in *vacuo*. The residue was purified by silica gel column chromatography (25 to 33% ethyl acetate-hexane) to give (E)-ester (65-A) (3.05 g, yield 23%)

and the corresponding (Z)-ester (6.17 g, yield 47%) as a colorless solid each.



(2) The above compound (65-A) was reacted in the similar manner as EXAMPLE 1-(4-2-2) to give the compound (65-B).

5 MS (m/z) ESI: 336 [M-H]⁻


(3) The above compound and the corresponding starting compound were reacted in the similar manner as EXAMPLE 1-(5) to give the compound (65-C).

MS (m/z) APCI: 438 [M+H]⁺

10 Corresponding starting compounds were treated in the similar manner as the above-mentioned to give the following compounds.

EXAMPLE No.	No	Structure	MS (m/z)
65	1		440 APCI [M+H] ⁺
65	2		460 APCI [M+H] ⁺

EXAMPLE 66



(1) To a solution of the compound of EXAMPLE 65-(2) (4.59 g, 9.99 mmol) in methylene chloride (200 ml) was added dropwise a 1.0M solution of boron tribromide in methylene chloride (50 ml, 50 mmol) over 30 minutes at -78°C , and then the mixture was stirred at the same temperature for 2 hours. To the reaction mixture was added water, and the mixture was warmed to room temperature and then extracted with chloroform. The organic layer was washed sequentially with water and brine, dried over sodium sulfate and concentrated in *vacuo*. The residue was solidified with a mixture of ethyl acetate-hexane (1:1) to give the compound (66-A) (2.97 g, yield 81%).

MS (m/z) APCI: 370 $[\text{M}+\text{H}]^{+}$

(2) To a solution of the above compound (500 mg, 1.35 mmol) in THF-N,N-dimethylacetamide (1:1) (4 ml) were added potassium tert-butoxide (379 mg, 3.38 mmol) and tert-butyl bromoacetate (0.22 ml, 1.49 mmol) under ice-cooling. The mixture was stirred at the same temperature for 40 minutes, diluted with ethyl acetate, and then washed sequentially with a saturated aqueous ammonium chloride solution, water and brine, dried over sodium sulfate and concentrated in *vacuo*. The residue was purified by silica gel column chromatography (20 to 50% ethyl acetate-hexane) to give the compound (66-B) (477 mg, yield 73%).

MS (m/z) APCI: 484 [M+H]⁺

(3) The above compound (448 mg, 0.927 mmol) was dissolved in formic acid (10 ml) at room temperature. The mixture was stirred at the same temperature for 70 hours and concentrated in *vacuo*, and then the residue was solidified with diisopropyl ether to give the carboxylic acid (66-C) (359 mg, yield 91%).


MS (m/z) ESI: 426 [M-H]⁻

(4) To a solution of the above compound (70 mg, 0.164 mmol) and 1-hydroxybenzotriazole (33.2 mg, 0.246 mmol) in DMF (2 ml) was added N-ethyl-N'-(3-dimethylaminopropyl)carbodiimide hydrochloride (63 mg, 0.328 mmol) under ice-cooling. The mixture was stirred at the same temperature for 1 hour. Then, thereto was added a 28% aqueous ammonia solution (1 ml), and the mixture was stirred at the same temperature for 30 minutes, diluted with ethyl acetate, washed sequentially with water and brine, dried over sodium sulfate and concentrated in *vacuo*. The residue was purified by silica gel column chromatography (0 to 7% methanol-chloroform) to give the compound (66-D) (24.7 mg, yield 35%).


MS (m/z) APCI: 427 [M+H]⁺

Corresponding starting compounds were treated in the similar manner as the above-mentioned to give the following compounds.

EXAMPLE No.	No	Structure	MS (m/z)
66	1		441 APCI [M+H] ⁺

66	2		455 APCI [M+H] ⁺
----	---	---	--------------------------------

EXAMPLE 67



(1) The compound (1-D), (S)-1-(tert-butyldimethylsilyloxy)-2-propanol and the corresponding starting compound were reacted in the similar manner as EXAMPLE 1-(4-1) to give the compound (67-A).

MS (m/z) APCI: 473 [M+NH₄]⁺

(2) The above compound was treated in the similar manner as EXAMPLE 1-(4-2-2) to give the compound (67-B).

MS (m/z) ESI: 440 [M-H]⁻

(3) The above compound was treated in the similar manner as EXAMPLE 1-(5) to give the compound (67-C).






MS (m/z) APCI: 542 [M+H]⁺





(4) To a solution of the above compound (73.7 mg, 0.136 mmol) in THF (3 ml) was added a 1.0M solution of tetrabutylammonium fluoride in THF (0.54 ml, 0.54 mmol) under ice-cooling. The mixture was stirred at room temperature for 24 hours, diluted with ethyl acetate, washed

sequentially with water and brine, dried over magnesium sulfate and concentrated in *vacuo*. The residue was purified by silica gel column chromatography (0 to 5% methanol-ethyl acetate) to give the compound (67-D) (38 mg, yield 65%).


5 MS (m/z) APCI: 428 [M+H]⁺

Corresponding starting compounds were treated in the similar manner as the above-mentioned to give the following compounds.

EXAMPLE No.	No	Structure	MS (m/z)
67	1		414 APCI [M+H] ⁺
67	2		428 APCI [M+H] ⁺
67	3		454 APCI [M+H] ⁺
67	4		454 APCI [M+H] ⁺
67	5		428 APCI [M+H] ⁺

67	6		454 APCI [M+H] ⁺
67	7		453 APCI [M+H] ⁺
67	8		468 APCI [M+H] ⁺
67	9		468 APCI [M+H] ⁺


EXAMPLE 68



The compound (66-A) and the corresponding starting compound were reacted in the similar manner as EXAMPLE 66-(2) to give the above compound.

MS (m/z) APCI: 464 [M+H]⁺

EXAMPLE 69



(1) To a solution of the compound (1-D) (858 mg, 3.03 mmol) in dimethylacetamide (5 ml) was added potassium tert-butoxide (374 mg, 3.33 mmol) under ice-cooling, and thereto was added (7*S*)-iodomethyl-(2*R*,3*R*)-diphenyl-1,4-dioxaspiro[4.4]nonane (WO2003095438) (1.40 g, 3.33 mmol). The mixture was stirred at room temperature overnight, diluted with ethyl acetate, washed sequentially with a 10% aqueous ammonium chloride solution, water and brine, dried over sodium sulfate and concentrated in *vacuo*. The residue was purified by silica gel column chromatography (30% ethyl acetate-hexane) to give the compound (69-A) (856 mg, yield 49%).

MS (*m/z*) APCI: 593 [*M*+NH₄]⁺

(2) To a solution of the above compound (1.05 g, 1.83 mmol) in dioxane (60 ml) was added 5N hydrochloric acid (30 ml) at room temperature. The mixture was stirred at the same temperature for 15 hours, diluted with ethyl acetate, washed sequentially with water and brine, dried over sodium sulfate and concentrated in *vacuo*. The residue was purified by silica gel column chromatography (50% ethyl acetate-hexane) to give the

compound (69-B) (512 mg, yield 74%).

MS (m/z) APCI: 380 [M+H]⁺


(3) The above compound was reacted in the similar manner as EXAMPLE 1-(4-2-2) to give the compound (69-C).

5 MS (m/z) ESI: 751 [2M+Na-2H]⁻

(4) The above compound and the corresponding starting compound were reacted in the similar manner as EXAMPLE 1-(5) to give the compound (69-D).

MS (m/z) APCI: 466 [M+H]⁺

10 EXAMPLE 70




(1) The compound (66-A) and the corresponding starting compound were reacted in the similar manner as EXAMPLE 66-(2) to give the compound (70-A).

15 MS (m/z) APCI: 534 [M+H]⁺

(2) The above compound (48.5 mg, 0.091 mmol) was dissolved in a mixed solvent of water-trifluoroacetic acid (1:3) (4 ml). The mixture was stirred at room temperature for 4 days, concentrated in *vacuo*, and then the residue was diluted with ethyl acetate, washed sequentially with a saturated aqueous sodium bicarbonate solution, water and brine, dried over sodium sulfate and concentrated in *vacuo*. The residue was purified by gel-filtration (column: JAIGEL, solvent: chloroform) to give the compound (70-B) (7 mg, yield 17%).

MS (m/z) APCI: 450 [M+H]⁺

25 EXAMPLE 71



The compound (69-D) (84.3 mg, 0.181 mmol) was dissolved in a mixture of methanol-THF (2:1) (6 ml), and thereto was added sodium borohydride (33.8 mg, 0.89 mmol) under ice-cooling. The mixture was stirred at room temperature for 90 minutes, diluted with ethyl acetate, washed sequentially with water and brine, dried over sodium sulfate and concentrated in *vacuo*. The residue was purified by silica gel column chromatography (0 to 7% methanol-chloroform) to give the above compound (79.8 mg, yield 94%).


MS (m/z) APCI: 468 [M+H]⁺

EXAMPLE 72

A corresponding starting compound was treated in the similar manner as EXAMPLE 62 and EXAMPLE 64 to give the following compound.

EXAMPLE No.	No	Structure	MS (m/z)
72			495 APCI [M+H] ⁺


EXAMPLE 73




To a solution of the compound of EXAMPLE 62-(7) (100 mg, 0.21 mmol) in methanol (3 ml) was added a 4N solution of hydrogen chloride in dioxane at room temperature, and the mixture was stirred for 3 days. To the reaction mixture was added a saturated aqueous sodium carbonate solution, and then the mixture was extracted with chloroform. The organic layer was washed sequentially with water and brine, dried over magnesium sulfate and concentrated in *vacuo*. The residue was purified by silica gel column chromatography (2 to 10% methanol-chloroform) to give the above compound (54.1 mg, yield 60%).

MS (m/z) APCI: 428 [M+H]⁺

A corresponding starting compound was treated in the similar manner as the above-mentioned to give the following compound.

EXAMPLE No.	No	Structure	MS (m/z)
73	1		428 APCI [M+H] ⁺

EXAMPLE 74



(1) To a solution of the compound (1-D) (1.0 g, 3.53 mmol) in THF (40 ml) was added sodium hydride (60%, 353 mg, 8.83mmol) under ice-cooling, and thereto was added (*R*)-2-bromopropionic acid (702 mg, 4.59 mmol) and the ice bath was removed. The mixture was stirred at room temperature for 3 hours, diluted with ethyl acetate, and then washed sequentially with 10% hydrochloric acid, water and brine, dried over magnesium sulfate and concentrated in *vacuo*. The resulting crude carboxylic acid (74-A) was used in the next reaction directly.

(2) To a solution of the above crude carboxylic acid in DMF (15 ml) were added sequentially 1-hydroxybenzotriazole (1.25 g, 9.29 mmol) and N-ethyl-N'-(3-dimethylaminopropyl)carbodiimide hydrochloride (2.97 g, 15.5 mmol) under ice-cooling, and the mixture was stirred at the same temperature for 4 hours. Then, thereto was added an aqueous solution of dimethylamine (50%) (5.0 ml) and the ice bath was removed and the mixture was stirred at room temperature for 30 minutes. The reaction mixture was diluted with ethyl acetate, and then washed sequentially with water and brine, dried over magnesium sulfate and concentrated in *vacuo*. The residue was purified by NH-silica gel column chromatography (50% ethyl acetate-hexane) to give the compound (74-B) (594 mg, yield 44%).


MS (m/z) APCI: 383 [M+H]⁺

(3) The above compound was treated in the similar manner as EXAMPLE 1-(4-2-2) to give the compound (74-C).

(4) The above compound and the corresponding starting compound were reacted in the similar manner as EXAMPLE 1-(5) to give the compound (74-D).

MS (m/z) APCI: 469 [M+H]⁺

EXAMPLE 75




To a solution of the compound of EXAMPLE 73 (100 mg, 0.23 mmol) and triethylamine (0.326 ml, 2.34 mmol) in dimethylsulfoxide (3 ml) was added sulfur trioxide-pyridine complex (186 mg, 1.17 mmol) at room temperature. The mixture was stirred at the same temperature for 18 hours, diluted with ethyl acetate, and then washed sequentially with water and brine, dried over magnesium sulfate and concentrated in *vacuo*. The residue was purified by silica gel column chromatography (30 to 100% ethyl acetate-hexane) to give the above compound (27.9 mg, yield 28%).

MS (m/z) APCI: 426 [M+H]⁺

A corresponding starting compound was treated in the similar manner as the above-mentioned to give the following compound.

EXAMPLE No.	No	Structure	MS (m/z)
75	1		466 APCI [M+H] ⁺

EXAMPLE 76



(1) To a solution of the compound (1-D) (1.0 g, 3.53 mmol) in THF (15 ml) were added sequentially potassium tert-butoxide (396 mg, 3.53 mmol) and then β -propiolactone (382 mg, 5.30 mmol) under ice-cooling, and the mixture was stirred at the same temperature for 2 hours and at room temperature for another 2 hours, and further at 50°C for 2 hours. To the reaction mixture was added 10% hydrochloric acid, and then the mixture was diluted with ethyl acetate, washed sequentially with water and brine, dried over magnesium sulfate and concentrated in *vacuo*. The resulting crude carboxylic acid (76-A) was used in the next reaction directly.

(2) The above crude carboxylic acid and the corresponding starting compound were reacted in the similar manner as EXAMPLE 74-(2) to give the compound (76-B).


MS (*m/z*) APCI: 383 [*M*+*H*]⁺

(3) The above compound was treated in the similar manner as EXAMPLE 1-(4-2-2) to give the compound (76-C).

(4) The above compound was reacted in the similar manner as EXAMPLE 1-(5) to give the compound (76-D).

MS (*m/z*) APCI: 469 [*M*+*H*]⁺

EXAMPLE 77



(1) To a solution of the compound (77-A) (1.77 g, 3.67 mmol) obtained from the compound (1-D) in the method of EXAMPLE 1-(4-2-1) in THF (10 ml) was added acetic acid (0.631 ml, 11.0 mmol) under ice-cooling, and thereto was added a 1.0M solution of tetrabutylammonium fluoride in THF (11.0 ml, 11.0 mmol); and the mixture was warmed to 50°C and then stirred for 3 hours, diluted with ethyl acetate, washed sequentially with water and brine, dried over magnesium sulfate and concentrated in *vacuo*. The residue was purified by silica gel column chromatography (50 to 80% ethyl acetate-hexane) to give the compound (77-B) (1.21 g, yield 90%).

MS (m/z) APCI: 368 [M+H]⁺

(2) To a solution of the above compound (1.05 g, 2.86 mmol) in DMF

(10 ml) were added silver (I) oxide (2.99 g, 12.7 mmol) and then methyl iodide (1.60 ml, 25.7 mmol) at room temperature, and the mixture was stirred at 40 to 50°C for 24 hours, diluted with ethyl acetate, and then filtered through Celite and the filtrate was washed sequentially with water and brine, dried over sodium sulfate and concentrated in *vacuo*. The residue was purified by silica gel column chromatography (30% ethyl acetate-hexane) to give the compound (77-C) (617 mg, yield 57%).

MS (m/z) APCI: 382 [M+H]⁺


(3) The above compound was reacted in the similar manner as EXAMPLE 1-(4-2-2) to give the compound (77-D).

MS (m/z) ESI: 366 [M-H]⁻

(4) The above compound was reacted in the similar manner as EXAMPLE 1-(5) to give the compound (77-E).

MS (m/z) APCI: 468 [M+H]⁺


EXAMPLE 78



(3-Chloro-4-methanesulfonylphenyl)oxoacetic acid ethyl ester and the corresponding starting compound were reacted in the similar manner as EXAMPLE 65 to give the above compound.

MS (m/z) APCI: 448/450 [M+H]⁺

EXAMPLE 79



(1) To a solution of 5-bromo-2-cyclopropylsulfanylp¹ridine 6.11g (26.6 mmol) in diethyl ether (200 ml) was added dropwise n-butyllithium (2.71M solution in hexane) (10.2 ml, 27.9 mmol) over 10 minutes at -78°C, and thereto was added diethyl oxalate (4.33 ml, 31.9 mmol) at the same temperature in one portion. To the reaction mixture was added a saturated aqueous ammonium chloride solution, and the mixture was extracted with ethyl acetate. The organic layer was separated and then washed sequentially with water and brine, dried over magnesium sulfate and concentrated in *vacuo*. The residue was purified by silica gel column chromatography (14% ethyl acetate-hexane) to give the above compound (79-A) (2.64 g, yield 40%).

MS (m/z) APCI: 252 [M+H]⁺

(2) The above compound and the corresponding starting compound were reacted in the similar manner as EXAMPLE 65-(1) and 27 to give the compound (79-B).

MS (m/z) APCI: 369 [M+H]⁺


(3) The above compound was reacted in the similar manner as EXAMPLE 1-(4-2-2) to give the compound (79-C).

MS (m/z) ESI: 339 [M-H]-

(4) The above compound was reacted in the similar manner as EXAMPLE 1-(5) to give the compound (79-D).

MS (m/z) APCI: 441 [M+H]⁺

EXAMPLE 80



(1) To a solution of thiophene-2-thiol (23.39 g, 200.4 mmol) in DMF (150 ml) was added potassium tert-butoxide (24.74 g, 220 mmol) under ice-cooling, and the mixture was stirred at the same temperature for 30 minutes and at room temperature for another 1 hour. To the above reaction mixture was added cyclopropyl bromide (17.8 ml, 222 mmol), and the mixture was stirred at 60°C for 30 hours and at 80°C for another 5 hours. The reaction mixture was diluted with ethyl acetate, washed sequentially with a saturated aqueous sodium bicarbonate solution, water and brine, dried over sodium sulfate and concentrated in *vacuo*. The residue was purified by NH-silica gel chromatography (ethyl acetate) and further purified by distillation under reduced pressure to give 2-cyclopropylsulfanyltrithiophene (80-A) (18.5g, yield 59%) as a colorless oil.

bp 70 to 100°C (19 mmHg), MS (m/z) APCI: 157 [M+H]⁺

(2) The above compound and the corresponding starting compound were reacted in the similar manner as EXAMPLE 1-(1) to give the above

compound (80-B).

MS (m/z) APCI: 257 [M+H]⁺


(3) The above compound and the corresponding starting compound
were reacted in the similar manner as EXAMPLE 1-(2) to (4) to give the
5 above compound (80-C).

MS (m/z) ESI: 344 [M-H]⁻

(4) The above compound and the corresponding starting compound
were reacted in the similar manner as EXAMPLE 1-(5) to give the above
compound (80-D).

10 MS (m/z) APCI: 446 [M+H]⁺

EXAMPLE 81



(1) (4-Nitrophenyl)oxoacetic acid ethyl ester and the corresponding
starting compounds were reacted in the similar manner as EXAMPLE 1-
15 (3) to (5) to give the compound (81-A).

MS (m/z) APCI: 381 [M+H]⁺

(2) To a solution of the above compound (3.0 g, 7.89 mmol) in
ethanol (150 ml) was added stannous chloride dihydrate (8.9 g, 39.4
mmol) at room temperature, and the mixture was stirred at the same



temperature for 16 hours. The reaction mixture was concentrated, and thereto was added ethyl acetate. The mixture was washed sequentially with a saturated aqueous sodium bicarbonate solution, water and brine, dried over magnesium sulfate and concentrated in *vacuo*. The residue was purified by silica gel chromatography (5% methanol-chloroform) to give the compound (81-B) 2.47 g, yield 89%).







MS (m/z) APCI: 351 [M+H]⁺

(3) To a solution of the above compound (100 mg, 0.29 mmol) and diisopropylethylamine (0.060 ml, 0.34 mmol) in THF (5 ml) was added p-nitrobenzoyl chloride (58 mg, 0.31 mmol) under ice-cooling, and then the ice bath was removed and the mixture was stirred at room temperature overnight. The reaction mixture was concentrated in *vacuo* and the residue was purified by NH-silica gel chromatography (0 to 3% methanol-chloroform) to give the compound (81-C) (160 mg, yield 57%).


MS (m/z) APCI: 500 [M+H]⁺

Corresponding starting compounds were treated in the similar manner as the above-mentioned to give the following compounds.

EXAMPLE No.	No	Structure	MS (m/z)
81	1		456 APCI [M+H] ⁺
81	2		475 APCI [M+H] ⁺

81	3		419 APCI [M+H] ⁺
81	4		456 APCI [M+H] ⁺
81	5		437 APCI [M+H] ⁺
81	6		455 APCI [M+H] ⁺
81	7		492 APCI [M+H] ⁺
81	8		471 APCI [M+H] ⁺

EXAMPLE 82









(1) The compound (81-B) (1.0 g, 2.9 mmol) was dissolved in a mixture of acetic acid (5 ml) and concentrated hydrochloric acid (14 ml), and thereto was added dropwise an aqueous solution (4ml) of sodium nitrite (217 mg, 3.14 mmol) under ice-cooling, and the mixture was stirred at the same temperature for 30 minutes. To the above reaction mixture were added sequentially copper (II) chloride dihydrate (243 mg, 1.43 mmol) and a solution of sodium bisulfite (4.45 g, 42.8 mmol) in 5.5M hydrochloric acid (22 ml), and the mixture was stirred at the same temperature for 10 minutes, and then the ice bath was removed. The mixture was stirred at room temperature for 3 hours and poured onto ice and extracted with ethyl acetate. The organic layer was dried over magnesium sulfate and concentrated in *vacuo* to give a crude sulfonyl chloride (82-A), which was used in the next reaction directly.







(2) To a solution of the above compound (142 mg) in THF (2 ml) was added a 28% aqueous ammonia solution (0.1 ml) at -5°C . The mixture was warmed to room temperature and stirred for 1 hour, diluted with ethyl acetate, washed sequentially with 2N hydrochloric acid, water and brine, dried over magnesium sulfate and concentrated in *vacuo*. The residue was purified by silica gel chromatography (40 to 60% ethyl acetate-hexane) to give the compound (82-B) (54 mg, yield 46% in 2 steps).







MS (m/z) APCI: 415 $[\text{M}+\text{H}]^{+}$







Corresponding starting compounds were treated in the similar manner as the above-mentioned to give the following compounds.







EXAMPLE	No	Structure	MS (m/z)
---------	----	-----------	----------







No.			
82	1		429 APCI [M+H] ⁺
82	2		443 APCI [M+H] ⁺
82	3		443 APCI [M+H] ⁺
82	4		457 APCI [M+H] ⁺
82	5		455 APCI [M+H] ⁺
82	6		455 APCI [M+H] ⁺







82	7		471 APCI [M+H] ⁺
82	8		485 APCI [M+H] ⁺
82	9		486 APCI [M+H] ⁺
82	10		500 APCI [M+H] ⁺
82	11		466 APCI [M+H] ⁺
82	12		457 APCI [M+H] ⁺

82	13		459 APCI [M+H] ⁺
82	14		473 APCI [M+H] ⁺
82	15		473 APCI [M+H] ⁺
82	16		487 APCI [M+H] ⁺
82	17		484 APCI [M+H] ⁺
82	18		498 APCI [M+H] ⁺







82	19		498 APCI [M+H] ⁺
82	20		526 APCI [M+H] ⁺
82	21		499 APCI [M+H] ⁺
82	22		499 APCI [M+H] ⁺
82	23		498 APCI [M+H] ⁺
82	24		512 APCI [M+H] ⁺







82	25		499 APCI [M+H] ⁺
82	26		499 APCI [M+H] ⁺
82	27		506 APCI [M+H] ⁺
82	28		520 APCI [M+H] ⁺
82	29		584 APCI [M+H] ⁺
82	30		512 APCI [M+H] ⁺







82	31		526 APCI [M+H] ⁺
82	32		555 APCI [M+H] ⁺
82	33		561 APCI [M+H] ⁺
82	34		562 APCI [M+H] ⁺
82	35		576 APCI [M+H] ⁺
82	36		590 APCI [M+H] ⁺







82	37		512 APCI [M+H] ⁺
82	38		540 APCI [M+H] ⁺
82	39		485 APCI [M+H] ⁺
82	40		485 APCI [M+H] ⁺
82	41		512 APCI [M+H] ⁺
82	42		512 APCI [M+H] ⁺







82	43		469 APCI [M+H] ⁺
82	44		487 APCI [M+H] ⁺
82	45		487 APCI [M+H] ⁺
82	46		501 APCI [M+H] ⁺
82	47		517 APCI [M+H] ⁺
82	48		533 APCI [M+H] ⁺







82	49		499 APCI [M+H] ⁺
82	50		561 APCI [M+H] ⁺
82	51		529 APCI [M+H] ⁺
82	52		560 APCI [M+NH ₄] ⁺
82	53		473 APCI [M+H] ⁺
82	54		473 APCI [M+H] ⁺

82	55		487 APCI [M+H] ⁺
82	56		487 APCI [M+H] ⁺
82	57		501 APCI [M+H] ⁺
82	58		501 APCI [M+H] ⁺
82	59		473 APCI [M+H] ⁺
82	60		473 APCI [M+H] ⁺







82	61		487 APCI [M+H] ⁺
82	62		487 APCI [M+H] ⁺
82	63		513 APCI [M+H] ⁺
82	64		526 APCI [M+H] ⁺
82	65		513 APCI [M+H] ⁺
82	66		513 APCI [M+H] ⁺

82	67		507 APCI [M+H] ⁺
82	68		509 APCI [M+H] ⁺
82	69		487 APCI [M+H] ⁺
82	70		501 APCI [M+H] ⁺
82	71		503 APCI [M+H] ⁺
82	72		531 APCI [M+H] ⁺

82	73		487 APCI [M+H] ⁺
82	74		487 APCI [M+H] ⁺
82	75		503 APCI [M+H] ⁺
82	76		499 APCI [M+H] ⁺
82	77		499 APCI [M+H] ⁺
82	78		461 APCI [M+H] ⁺

82	79		479 APCI [M+H] ⁺
82	80		489 APCI [M+H] ⁺
82	81		469 APCI [M+H] ⁺
82	82		514 APCI [M+H] ⁺
82	83		514 APCI [M+H] ⁺
82	84		528 APCI [M+H] ⁺


82	85		500 APCI [M+H] ⁺
82	86		472 APCI [M+H] ⁺
82	87		486 APCI [M+H] ⁺
82	88		500 APCI [M+H] ⁺
82	89		486 APCI [M+H] ⁺
82	90		528 APCI [M+H] ⁺

82	91		528 APCI [M+H] ⁺
82	92		542 APCI [M+H] ⁺
82	93		512 APCI [M+H] ⁺
82	94		512 APCI [M+H] ⁺
82	95		540 APCI [M+H] ⁺
82	96		512 APCI [M+H] ⁺

82	97	 SMILES: <chem>C1CCN(C1)S(=O)(=O)c2ccc(cc2CNC(=O)c3cc(NC(=O)c4nc(F)s4)ccc3OC5COCC5)</chem>	498 APCI [M+H] ⁺
82	98	 SMILES: <chem>C[C@H]1CCN(C1)S(=O)(=O)c2ccc(cc2CNC(=O)c3cc(NC(=O)c4nc(F)s4)ccc3OC5COCC5)</chem>	498 APCI [M+H] ⁺
82	99	 SMILES: <chem>C[C@H]1CC(O)[C@@H](C)CN1S(=O)(=O)c2ccc(cc2CNC(=O)c3cc(NC(=O)c4nc(F)s4)ccc3OC5COCC5)</chem>	513 APCI [M+H] ⁺
82	100	 SMILES: <chem>CN1CCCC1S(=O)(=O)c2ccc(cc2CNC(=O)c3cc(NC(=O)c4nc(F)s4)ccc3OC5COCC5)</chem>	526 APCI [M+H] ⁺
82	101	 SMILES: <chem>CCNS(=O)(=O)c1ccc(cc1CNC(=O)c2cc(NC(=O)c3nc(F)s3)ccc2OC4COCC4)</chem>	459 APCI [M+H] ⁺
82	102	 SMILES: <chem>CN(C)S(=O)(=O)c1ccc(cc1CNC(=O)c2cc(NC(=O)c3nc(F)s3)ccc2OC4COCC4)</chem>	459 APCI [M+H] ⁺

EXAMPLE 83

174



(1) To a solution of (4-methylthiophenyl)oxoacetic acid ethyl ester (44.9 g, 200 mmol) in chloroform (500 ml) was added 65% m-chloroperbenzoic acid (50 g, 188 mmol) over 30 minutes under ice-cooling, and the mixture was stirred at the same temperature for another 1 hour. The precipitated insoluble was filtered off, and then to the filtrate was added a saturated aqueous sodium bicarbonate solution, and the mixture was extracted with chloroform. The organic layer was dried over sodium sulfate and concentrated in *vacuo*. The residue was purified by silica gel chromatography (50 to 100% ethyl acetate-hexane) to give the compound (83-A) (41.1 g, yield 85%) as a pale yellow oil.

MS (m/z) APCI: 241 [M+H]⁺

(2) To a solution of the above compound (41.1 g, 171 mmol) in ethanol (400 ml) was added hydroxylamine hydrochloride (15.5 g, 222 mmol). The mixture was stirred at 50°C for 2 hours, concentrated in

vacuo, and then the residue was dissolved in ethyl acetate, washed sequentially with water and brine, dried over sodium sulfate and concentrated in *vacuo*. To a solution of the resulting crude oxime (45.7 g) and potassium carbonate (49.4 g, 358 mmol) in DMF (325 ml) was added 3-(*S*)-tetrahydrofuranol p-toluenesulfonate (56.1 g, 232 mmol) at room temperature. The mixture was stirred for 2 days, diluted with ethyl acetate, washed sequentially with water and brine, dried over sodium sulfate and concentrated in *vacuo*. The residue was purified by silica gel chromatography (ethyl acetate) to give the compound (83-B) (20.9 g, (*E*)-isomer:(*Z*)-isomer = 72:28).

MS (m/z) APCI: 326 [M+H]⁺

(3) The above compound (20.35 g) was dissolved in THF-ethanol-water (3:1:1) (350 ml), and thereto was added potassium carbonate (8.63 g, 62.5 mmol), and the mixture was stirred at room temperature for 23 hours. To the reaction mixture was added water, and the aqueous layer was washed with ethyl acetate, acidified with concentrated hydrochloric acid, saturated with sodium chloride and extracted with ethyl acetate several times. The extract was dried over sodium sulfate and concentrated in *vacuo* to give the compound (83-C, (*E*)-isomer) (10.9 g, yield 59%) as a pale yellow oil.

MS (m/z) APCI: 298 [M+H]⁺

(4) The above compound and the corresponding starting compound were reacted in the similar manner as EXAMPLE 1-(5) to give the compound (83-D).

MS (m/z) APCI: 398 [M+H]⁺



(5) To a solution of the above compound (2.85 g, 7.17 mmol) in chloroform (40 ml) was added trifluoroacetic anhydride (2.03 ml, 14.3 mmol) at room temperature, and the mixture was stirred at the same temperature for 3 hours and concentrated in *vacuo*. The residue was dissolved in methanol (15 ml), and thereto was added triethylamine (15







ml) at room temperature. The mixture was stirred at the same temperature for 30 minutes and concentrated in *vacuo* to give the crude thiol (83-E) (2.62 g).







(6) The above compound (150 mg, 0.41 mmol) was dissolved in DMF (2 ml), and thereto was added potassium tert-butoxide (36.6 mg, 0.33 mmol) under ice-cooling, and the mixture was stirred at the same temperature for 10 minutes. Then, thereto was added ethyl iodide (0.026 ml, 0.33 mmol). The mixture was stirred at the same temperature for 2 hours and at room temperature overnight, diluted with ethyl acetate, washed sequentially with water and brine, dried over sodium sulfate, concentrated in *vacuo*. The residue was purified by silica gel chromatography (25 to 75% ethyl acetate-hexane) to give the compound (83-F) (96.8 mg, yield 65% in 2 steps).




MS (m/z) APCI: 396 [M+H]⁺

Corresponding starting compounds were treated in the similar manner as the above-mentioned to give the following compounds.


EXAMPLE No.	No	Structure	MS (m/z)
83	1		426 APCI [M+H] ⁺
83	2		410 APCI [M+H] ⁺

83	3		422 APCI [M+H] ⁺
83	4		541 APCI [M+H] ⁺
83	5		499 APCI [M+NH ₄] ⁺
83	6		422 APCI [M+H] ⁺
83	7		438 APCI [M+H] ⁺
83	8		438 APCI [M+H] ⁺

83	9		456 APCI [M+H] ⁺
83	10		440 APCI [M+H] ⁺
83	11		426 APCI [M+H] ⁺
83	12		440 APCI [M+H] ⁺
83	13		440 APCI [M+H] ⁺
83	14		435 APCI [M+H] ⁺

83	15		421 APCI [M+H] ⁺
83	16		454 APCI [M+H] ⁺
83	17		454 APCI [M+H] ⁺







EXAMPLE 84















To a solution of the compound (83-F) (88.8 mg, 0.23 mmol) in chloroform (3 ml) was added 70% m-chloroperbenzoic acid (166 mg, 0.68 mmol) at room temperature, and the mixture was stirred at the same temperature for 6 hours and concentrated in *vacuo*. The residue was purified by NH-silica gel chromatography (0 to 5% methanol-chloroform) to give the above compound (76.8 mg, yield 80%).




MS (m/z) APCI: 428 [M+H]⁺

Corresponding starting compounds were reacted in the similar manner as EXAMPLE 83 and/or the above-mentioned to give the following compounds.


EXAMPLE No.	No	Structure	MS (m/z)
84	1		573 APCI [M+H] ⁺
84	2		514 APCI [M+H] ⁺
84	3		442 APCI [M+H] ⁺
84	4		442 APCI [M+H] ⁺
84	5		458 APCI [M+H] ⁺
84	6		454 APCI [M+H] ⁺

84	7		444 APCI [M+H] ⁺
84	8		456 APCI [M+H] ⁺
84	9		454 APCI [M+H] ⁺
84	10		470 APCI [M+H] ⁺
84	11		470 APCI [M+H] ⁺
84	12		494 APCI [M+H] ⁺

84	13		472 APCI [M+H] ⁺
84	14		458 APCI [M+H] ⁺
84	15		472 APCI [M+H] ⁺
84	16		472 APCI [M+H] ⁺
84	17		506 APCI [M+H] ⁺
84	18		467 APCI [M+H] ⁺

84	19		453 APCI [M+H] ⁺
84	20		486 APCI [M+H] ⁺
84	21		486 APCI [M+H] ⁺

EXAMPLE 85




(1) To a solution of the compound (83-E) (300 mg, 0.82 mmol) in chloroform (10 ml) were added sequentially triethylamine (0.17 ml, 1.2 mmol) and then tri-*n*-butyltin chloride (0.18ml, 0.65 mmol) under ice-cooling. The mixture was stirred at the same temperature for 1 hour and at room temperature for another 4 hours, diluted with ethyl acetate, washed sequentially with water and brine, dried over sodium sulfate and concentrated in *vacuo*. The residue was purified by silica gel chromatography (10 to 60% ethyl acetate-hexane) to give the compound (85-A) (294 mg, yield 55%).

MS (m/z) APCI: 654/656 [M+H]⁺

(2) To a solution of the above compound in toluene (3 ml) were added 2-bromopyridine (0.087 ml, 0.91 mmol) and tetrakis-(triphenylphosphine)palladium (71 mg, 0.061 mmol) under argon. The mixture was heated to reflux for 5 hours, diluted with ethyl acetate, washed sequentially with water and brine, dried over sodium sulfate and concentrated in *vacuo*. The residue was purified by silica gel chromatography (20 to 80% ethyl acetate-hexane) to give the compound (85-B) (76.7 mg, yield 79%).

MS (m/z) APCI: 445 [M+H]⁺

EXAMPLE 86




To a solution of the compound of EXAMPLE 84-(1) (264 mg, 0.46 mmol) in ethanol-THF (1:1) (12 ml) was added hydrazine hydrate (92.2 mg, 1.98 mmol) at room temperature. The mixture was stirred at the same temperature for 24 hours, and thereto was added a saturated aqueous sodium bicarbonate solution. The mixture was extracted with chloroform, and the organic layer was dried over sodium sulfate and concentrated in *vacuo*. The residue was dissolved in ethyl acetate, and thereto was added a 4N solution of hydrogen chloride in dioxane (0.5 ml) at room temperature, and the precipitated crystals were collected to give the above compound (242 mg, quantitatively) as a hydrochloride.

MS (m/z) APCI: 443 [M+H]⁺

EXAMPLE 87

185



(1) (4-Nitrophenyl)oxoacetic acid ethyl ester and the corresponding starting compound were reacted in the similar manner as EXAMPLE 1-(3) and 1-(4-1) to give the compound (87-A).

5 MS (m/z) ESI: 279 [M-H]⁻

(2) A solution of the above compound (21.5 g, 76.7 mmol) in DMF (358 ml) was ice-cooled and thereto were added potassium carbonate (15.9 g, 115 mmol) and methyl iodide (5.75 ml, 92.1 mmol). The mixture was stirred at room temperature for 4 hours, diluted with ethyl acetate, washed sequentially with water and brine, dried over sodium sulfate and concentrated in *vacuo* to give the compound (87-B) (23.9 g, quantitatively).

10 MS (m/z) APCI: 295 [M+H]⁺

(3) The above compound was treated in the similar manner as EXAMPLE 81-(2) to give the compound (87-C).

15

MS (m/z) APCI: 265 [M+H]⁺

(4) To a solution of the above compound (3.0 g, 11.4 mmol) in a mixture of concentrated hydrochloric acid (30 ml) and water (10 ml) was added an aqueous solution (8 ml) of sodium nitrite (875 mg, 12.5 mmol) under ice-cooling, and the mixture was stirred at the same temperature for 30 minutes. Then, thereto was added an aqueous solution (12 ml) of potassium iodide (5.68 g, 34.1 mmol), and the ice bath was removed. The mixture was stirred at room temperature for another 1 hour, poured onto ice and extracted with ethyl acetate. The organic layer was filtered through Celite, and then the filtrate was washed sequentially with water and brine, dried over magnesium sulfate and concentrated in *vacuo*. The residue was purified by silica gel chromatography (30% ethyl acetate-hexane) to give the compound (87-D) (2.57 g, yield 60%).

MS (m/z) APCI: 376 [M+H]⁺


(5) The above compound was reacted in the similar manner as EXAMPLE 1-(4-2-2) to give the compound (87-E).

MS (m/z) ESI: 360 [M-H]⁻

(6) The above compound and the corresponding starting compound were reacted in the similar manner as EXAMPLE 1-(5) to give the compound (87-F).

MS (m/z) APCI: 462 [M+H]⁺

EXAMPLE 88




To a suspension of the compound (87-F) (200 mg, 0.43 mmol), potassium tert-butoxide (53.5 mg, 0.48 mmol), 3-mercapto-4-methyl-4H-1,2,4-triazole (50 mg, 0.43 mmol) and bis(2-


diphenylphosphinophenyl) ether (23.4 mg, 0.043 mmol) in toluene (5 ml) was added tris(dibenzylideneacetone)dipalladium (20 mg, 0.022 mmol) at room temperature under argon. The mixture was heated to reflux for 2 hours, diluted with ethyl acetate, and then filtered through
 5 Celite and concentrated in *vacuo*. The residue was purified by NH-silica gel chromatography (0 to 3% methanol-chloroform) to give the above compound (79 mg, yield 41%).

MS (m/z) APCI: 449 [M+H]⁺

A corresponding starting compound was treated in the similar
 10 manner as the above-mentioned to give the following compound.

EXAMPLE No.	No	Structure	MS (m/z)
88	1		448 APCI [M+H] ⁺

EXAMPLE 89




To a solution of the compound of EXAMPLE 88 (80 mg, 0.18 mmol) in methylene chloride (3 ml) was added 65% m-chloroperbenzoic acid (189 mg, 0.71 mmol) under ice-cooling, and then the ice bath was removed and the mixture was stirred at room temperature for another 2 hours. To the reaction mixture was added a 10% aqueous sodium thiosulfate solution, and then the mixture was extracted with chloroform. The organic layer was dried over magnesium sulfate and
 15 concentrated in *vacuo*. The residue was purified by NH-silica gel
 20

chromatography (0 to 3% methanol-chloroform) to give the above compound (7.8 mg, yield 9%).

MS (m/z) APCI: 481 [M+H]⁺

EXAMPLE 90



(1) To a solution of 4-hydroxyphenyloxycetic acid ethyl ester (19.4 g, 108 mmol) in acetone (300 ml) were added sequentially potassium carbonate (44.1 g, 319 mmol) and then allyl bromide (15.4 g, 127 mmol) under ice-cooling, and then the ice bath was removed. The mixture was stirred at room temperature for 16 hours and heated to reflux for another 6 hours, filtered through Celite, and then concentrated in *vacuo*. The resulting residue was purified by silica gel chromatography (17% ethyl acetate-hexane) to give the compound (90-A) (3.38 g, yield 14%).



(2) The above compound was reacted in the similar manner as EXAMPLE 1-(3) and 1-(4-2) to give the compound (90-B).

MS (m/z) ESI: 290 [M-H]⁻


(3) The above compound and the corresponding starting compound were reacted in the similar manner as EXAMPLE 1-(5) to give the compound (90-C).

MS (m/z) APCI: 392 [M+H]⁺

Corresponding starting compounds were treated in the similar manner as the above-mentioned to give the following compounds.

EXAMPLE No.	No	Structure	MS (m/z)
90	1		394 APCI [M+H] ⁺
90	2		406 APCI [M+H] ⁺

EXAMPLE 91



- 5 (1) To a solution of cyclopentylamine (5.0 g, 49 mmol) and triethylamine (8.15 ml, 59 mmol) in THF (50 ml) was added dropwise a

solution of ethyl chloroglyoxylate (6.1 g, 45 mmol) in THF (25 ml) under ice-cooling, and the mixture was stirred at the same temperature for 30 minutes and at room temperature for another 1 hour. The insoluble was filtered off through Celite, and then the filtrate was concentrated in *vacuo*, and the resulting residue was purified by silica gel column chromatography (25% ethyl acetate-hexane) to give the compound (91-A) (7.97 g, yield 88%) as a pale yellow oil.

MS (m/z) APCI: 202 [M+H]⁺

(2) To a solution of the above compound (2.0 g, 10 mmol) in acetonitrile (80 ml) were added sequentially triphenylphosphine (3.93 g, 15 mmol) and carbon tetrabromide (4.97 g, 15 mmol) at room temperature, and the mixture was heated to reflux for 3 hours. After cooling to room temperature, the mixture was concentrated and the residue was purified by silica gel column chromatography (3% ethyl acetate-hexane) to give the compound (91-B) (1.87 g, yield 71%) as a colorless oil.

(3) To a solution of the above compound (5.28 g, 20 mmol), 4-methylthiophenylboronic acid (5.04 g, 30 mmol) in 1,2-dimethoxyethane (120 ml) were added a 1N aqueous sodium carbonate solution (60 ml) and dichlorobis(triphenylphosphine)palladium (1.4 g, 2.0 mmol) under argon, and the mixture was heated at 80°C for 30 minutes using a microwave reactor. After cooling to room temperature, the reaction mixture was diluted with ethyl acetate, washed sequentially with water and brine, dried over sodium sulfate and concentrated in *vacuo*. The residue was purified by silica gel column chromatography (2 to 3% ethyl acetate-hexane) to give the compound (91-C) (4.20 g, yield 68%) as a pale yellow oil.

(4) To a solution of the above compound (20 g, 65 mmol) in methylene chloride (320 ml) was added dropwise a solution of 75% m-chloroperbenzoic acid (32.9 g, 143 mmol) in methylene chloride (80 ml)

under ice-cooling, and then the mixture was stirred at the same temperature for 30 minutes and at room temperature for 2 hours. The precipitate was filtered off through Celite, and then to the filtrate was added a 10% aqueous sodium sulfite solution and the mixture was stirred for a while. The organic layer was separated, and then washed sequentially with a saturated aqueous sodium carbonate solution, water and brine, dried over sodium sulfate and concentrated in *vacuo*. The residue was purified by silica gel column chromatography (25% ethyl acetate-hexane) to give the compound (91-D) (19.35 g, yield 88%).


MS (m/z) APCI: 340[M+H]⁺







(5) The above compound was reacted in the similar manner as EXAMPLE 1-(4-2-2) to give the compound (91-E).

(6) The above compound and the corresponding starting compound were reacted in the similar manner as EXAMPLE 1-(5) to give the compound (91-F).







MS (m/z) APCI: 388 [M+H]⁺







The above compounds of EXAMPLE (91-C), (91-D) or (91-E) and the corresponding starting compounds were reacted in the similar manner as a combination of any or some of the above EXAMPLEs to give the following compounds.







EXAMPLE No.	No	Structure	MS (m/z)
91	1		394 APCI [M+H] ⁺







91	2		466 APCI [M+H] ⁺
91	3		436 ESI [M-H] ⁻
91	4		419 APCI [M+H] ⁺
91	5		424 APCI [M+H] ⁺
91	6		406 APCI [M+H] ⁺
91	7		479 ESI [M-H] ⁻

91	8		435 ESI [M-H] ⁻
91	9		422 APCI [M+H] ⁺
91	10		528 APCI [M+H] ⁺
91	11		428/430 APCI [M+H] ⁺
91	12		462 APCI [M+H] ⁺
91	13		412 APCI [M+H] ⁺

91	14		466 APCI [M+H] ⁺
91	15		424 APCI [M+H] ⁺
91	16		395 APCI [M+H] ⁺
91	17		480 APCI [M+H] ⁺
91	18		506 APCI [M+H] ⁺
91	19		493 APCI [M+H] ⁺

91	20		535 APCI [M+H] ⁺
91	21		477 APCI [M+H] ⁺
91	22		495 APCI [M+H] ⁺
91	23		507 APCI [M+H] ⁺
91	24		521 APCI [M+H] ⁺
91	25		520 APCI [M+H] ⁺

91	26		550 APCI [M+H] ⁺
91	27		521 APCI [M+H] ⁺
91	28		509 APCI [M+H] ⁺
91	29		508 APCI [M+H] ⁺
91	30		507 APCI [M+H] ⁺
91	31		528 APCI [M+H] ⁺


91	32		506 ESI+ [M+H] ⁺
91	33		500 ESI+ [M+H] ⁺
91	34		514 ESI+ [M+H] ⁺
91	35		528 ESI+ [M+H] ⁺
91	36		494 ESI+ [M+H] ⁺
91	37		480 ESI+ [M+H] ⁺

91	38		514 APCI [M+H] ⁺
91	39		451 APCI [M+H] ⁺
91	40		465 APCI [M+H] ⁺

EXAMPLE 92




Corresponding starting compounds were treated in the similar manner as EXAMPLE 65 to give the following compounds.

EXAMPLE No.	No	Structure	MS (m/z)
92	1		380 APCI [M+H] ⁺
92	2		444/446 APCI [M+H] ⁺


92	3		382 APCI [M+H] ⁺
----	---	---	--------------------------------

EXAMPLE 93

Corresponding starting compounds were treated in the similar manner as EXAMPLE 65 to give the following compounds.

EXAMPLE No.	No	Structure	MS (m/z)
93	1		428/430 APCI [M+H] ⁺
93	2		462/464 APCI [M+H] ⁺
93	3		430/432 APCI [M+H] ⁺

EXAMPLE 94



(1) The compound (1-C) was treated in the similar manner as EXAMPLE 65 to give the compound (94-A).

MS (m/z) APCI: 442 [M+H]⁺







5 (2) The above compound was reacted in the similar manner as EXAMPLE 66-(1) to give the compound (94-B).







MS (m/z) APCI: 352 [M+H]⁺







(3) To a solution of the above compound (95.0 mg, 0.27 mmol), triphenylphosphine (127 mg, 0.487 mmol) and cyclopropylmethanol (58.4 mg, 0.810 mmol) in THF (6 ml) was added dropwise diisopropyl azodicarboxylate (0.096 ml, 0.487 mmol) under ice-cooling. The mixture was stirred at the same temperature for 3 hours, concentrated in *vacuo*, and the residue was purified by gel-filtration (column: JAIGEL, solvent: chloroform) to give the compound (94-C) (82.8 mg, yield 78%) as a colorless solid.







MS (m/z) APCI: 406 [M+H]⁺







Corresponding starting compounds were treated in the similar manner as one or more combinations selected from the above methods, EXAMPLE 66-(2), EXAMPLE 64 and EXAMPLE 53 to give the following compounds.







EXAMPLE No.	No	Structure	MS (m/z)
94	1		442 APCI [M+H] ⁺
94	2		426 APCI [M+H] ⁺
94	3		410 APCI [M+H] ⁺
94	4		521 APCI [M+H] ⁺
94	5		467 APCI [M+H] ⁺
94	6		443 APCI [M+H] ⁺







94	7		456 APCI [M+H] ⁺
94	8		463 APCI [M+H] ⁺
94	9		447 APCI [M+H] ⁺
94	10		461 APCI [M+H] ⁺
94	11		467 APCI [M+H] ⁺
94	12		434 ESI ⁺ [M+H] ⁺




94	13		472 ESI+ [M+H] ⁺
94	14		443 ESI+ [M+H] ⁺
94	15		448 ESI+ [M+H] ⁺
94	16		424 ESI+ [M+H] ⁺
94	17		408 ESI+ [M+H] ⁺
94	18		394 ESI+ [M+H] ⁺

94	19		406 ESI+ [M+H] ⁺
94	20		449 APCI [M+H] ⁺
94	21		444 ESI+ [M+H] ⁺
94	22		476 ESI+ [M+H] ⁺
94	23		436 ESI+ [M+H] ⁺
94	24		436 APCI [M+H] ⁺


94	25		467 APCI [M+H] ⁺
94	26		460 ESI+ [M+H] ⁺
94	27		472 ESI+ [M+H] ⁺
94	28		472 ESI+ [M+H] ⁺
94	29		436 APCI [M+H] ⁺
94	30		424 APCI [M+H] ⁺

94	31		424 APCI [M+H] ⁺
94	32		463 APCI [M+H] ⁺
94	33		430 APCI [M+H] ⁺
94	34		449 APCI [M+H] ⁺
94	35		424 APCI [M+H] ⁺
94	36		457 APCI [M+H] ⁺

94	37		452 APCI [M+H] ⁺
94	38		484 APCI [M+H] ⁺
94	39		468 APCI [M+H] ⁺
94	40		444 APCI [M+H] ⁺
94	41		391 APCI [M+H] ⁺
94	42		410 ESI+ [M+H] ⁺

94	43		410 ESI+ [M+H] ⁺
94	44		535 APCI [M+H] ⁺
94	45		432 APCI [M+H] ⁺

EXAMPLE 95




(1) The compound (1-D) and the corresponding starting compound were reacted in the similar manner as EXAMPLE 1-(4-1) to give the compound (95-A).

MS (m/z) ESI: 362 [M-H]⁻

(2) The above compound and the corresponding starting compound were reacted in the similar manner as EXAMPLE 1-(5) to give the compound (95-B).



MS (m/z) APCI: 600 [M+H]⁺

A corresponding starting compound was treated in the similar manner as the above-mentioned to give the following compound.

EXAMPLE No.	No	Structure	MS (m/z)
95	1		490 APCI [M+H] ⁺

EXAMPLE 96


The compound (80-C) and the corresponding starting compounds were reacted in the similar manner as EXAMPLE 1-(5) to give the following compounds.







EXAMPLE No.	No	Structure	MS (m/z)
96	1		442 APCI [M+H] ⁺
96	2		479 APCI [M+H] ⁺


5

EXAMPLE 97


Corresponding starting compounds were treated in the similar manner as a combination of the method of EXAMPLE 91 and the above-mentioned method to give the following compounds.

EXAMPLE No.	No	Structure	MS (m/z)
97	1		450 APCI [M+H] ⁺

97	2		390 APCI [M+H] ⁺
97	3		362 APCI [M+H] ⁺
97	4		387 APCI [M+H] ⁺
97	5		392 APCI [M+H] ⁺
97	6		396/398 APCI [M+H] ⁺
97	7		449 APCI [M+H] ⁺

97	8		380 APCI [M+H] ⁺
----	---	---	--------------------------------

EXAMPLE 98



(1) 3-Methoxypropyl phenyl sulfide was reacted in the similar manner as EXAMPLE 1-(1) to (3) to give the compound (98-A).

5 MS (m/z) APCI: 316 [M+H]⁺

(2) The above compound and the corresponding starting compound were treated in the similar manner as EXAMPLE 1-(4-2-1) to give the compound (98-B).

MS (m/z) APCI: 386 [M+H]⁺


10 (3) The above compound was treated in the similar manner as EXAMPLE 1-(4-2-2) to give the compound (98-C).

MS (m/z) ESI: 741 [2M-H]⁻

(4) The above compound was treated in the similar manner as EXAMPLE 1-(5) to give the compound (98-D).

15 MS (m/z) APCI: 468 [M+H]⁺

The compound (98-B) was also synthesized in the following alternative method.




(1) A solution of the compound (83-C) (1.00 g, 3.36 mmol) in DMF (15 ml) was ice-cooled, and thereto were added potassium carbonate (697 mg, 5.04 mmol) and methyl iodide (0.0251 ml, 4.03 mmol). The mixture was stirred at room temperature for 3 hours, diluted with ethyl acetate, washed sequentially with water and brine, dried over sodium sulfate and concentrated in *vacuo*, and then the residue was purified by silica gel chromatography (0 to 10% methanol-ethyl acetate) to give the compound (98-E) (1.04 g, yield 99%) as a colorless oil.







MS (m/z) APCI: 312 [M+H]⁺

(2) The above compound (98-E) was reacted with the corresponding starting compound in the similar manner as EXAMPLE 83-(5), (6) and EXAMPLE 84 to give the compound (98-B).


MS (m/z) APCI: 386 [M+H]⁺

Corresponding starting compounds were treated in the similar manner as the above-mentioned to give the following compounds.

EXAMPLE No.	No	Structure	MS (m/z)
98	1		498 APCI [M+H] ⁺

98	2		608 APCI [M+H] ⁺
98	3		469 APCI [M+H] ⁺
98	4		535 APCI [M+H] ⁺
98	5		505 APCI [M+H] ⁺
98	6		592 APCI [M+H] ⁺
98	7		565 APCI [M+H] ⁺

EXAMPLE 99



(1) The compound (87-C) and the corresponding starting compound were treated in the similar manner as EXAMPLE 82 to give the compound (99-A).

5 MS (m/z) APCI: 387 [M+H]⁺

(2) The above compound was treated in the similar manner as EXAMPLE 63-(2) to give the compound (99-B).





MS (m/z) ESI: 765 [2M-2H+Na]⁻

(3) The above compound was treated in the similar manner as EXAMPLE 1-(5) to give the compound (99-C).


10 MS (m/z) APCI: 499 [M+H]⁺

Corresponding starting compounds were treated in the similar manner as the above-mentioned to give the following compounds.

EXAMPLE No.	No	Structure	MS (m/z)
99	1		470 APCI [M+H] ⁺

99	2		469 APCI [M+H] ⁺
99	3		609 APCI [M+H] ⁺
99	4		506 APCI [M+H] ⁺
99	5		536 APCI [M+H] ⁺

EXAMPLE 100



(1) The compound (87-C) and the corresponding starting compound were treated in the similar manner as EXAMPLE 82 to give the compound (100-A).

5 MS (m/z) APCI: 373 [M+H]⁺

(2) A solution of the above compound (2.89 g, 7.76mmol) and imidazole (1.28 g, 18.6 mmol) in DMF (48 ml) was ice-cooled, and thereto was added tert-butyldimethylchlorosilane (1.45 g, 9.3 mmol). The mixture was stirred at room temperature for 2 hours, diluted with ethyl acetate, and washed with water and brine, dried over sodium sulfate and concentrated in *vacuo*, and the resulting residue was purified by silica gel chromatography (33% ethyl acetate-hexane) to give the compound (100-B) (3.58 g, yield 95%) as a colorless oil.

10

MS (m/z) APCI: 487 [M+H]⁺

(3) The above compound was treated in the similar manner as EXAMPLE 63-(2) to give the compound (100-C).




(4) The above compound was treated in the similar manner as EXAMPLE 1-(5) to give the compound (100-D).

MS (m/z) APCI: 569 [M+H]⁺


(5) The above compound was treated in the similar manner as EXAMPLE 67-(4) to give the compound (100-E).

MS (m/z) APCI: 455 [M+H]⁺

Corresponding starting compounds were treated in the similar manner as the above-mentioned to give the following compounds.

EXAMPLE No.	No	Structure	MS (m/z)
100	1		485 APCI [M+H] ⁺
100	2		492 APCI [M+H] ⁺
100	3		522 APCI [M+H] ⁺

EXAMPLE 101



(1) The compound (87-C) was treated in the similar manner as EXAMPLE 82-(1) and 1-(4-2-2) to give the compound (101-A).



MS (m/z) ESI: 793 [2M-2H+Na]⁻

5 (2) The above compound was reacted in the similar manner as EXAMPLE 1-(5) to give the compound (101-B).

MS (m/z) APCI: 623 [M+H]⁺




Corresponding starting compounds were treated in the similar manner as the above-mentioned to give the following compounds.







EXAMPLE No.	No	Structure	MS (m/z)
101	1		483 APCI [M+H] ⁺







101	2		520 APCI [M+H] ⁺
101	3		550 APCI [M+H] ⁺







EXAMPLE 102






Corresponding starting compounds were treated in a combination of the above-mentioned methods to give the following compounds.

EXAMPLE No.	No	Structure	MS (m/z)
102	1		423 APCI [M+H] ⁺
102	2		422 APCI [M+H] ⁺
102	3		576 APCI [M+H] ⁺


102	4		606 APCI [M+H] ⁺
102	5		450 APCI [M+H] ⁺
102	6		576 APCI [M+H] ⁺
102	7		532 APCI [M+H] ⁺
102	8		560 APCI [M+H] ⁺
102	9		532 APCI [M+H] ⁺

102	10		519 APCI [M+H] ⁺
102	11		567 APCI [M+H] ⁺
102	12		448 APCI [M+H] ⁺
102	13		450 APCI [M+H] ⁺
102	14		551 APCI [M+H] ⁺
102	15		441 APCI [M+H] ⁺

102	16		469 APCI [M+H] ⁺
102	17		415 APCI [M+H] ⁺
102	18		478 APCI [M+H] ⁺
102	19		538 APCI [M+H] ⁺
102	20		552 APCI [M+H] ⁺
102	21		472 APCI [M+H] ⁺

102	22		518 APCI [M+H] ⁺
102	23		508 ESI [M-H] ⁻
102	24		493/495 APCI [M+H] ⁺
102	25		569 APCI [M+H] ⁺
102	26		366 ESI [M+H] ⁺

EXAMPLE 103



(1) The compound (1-D) and the corresponding starting compound were reacted in the similar manner as EXAMPLE 1-(4-1) to give the compound (103-A).

5 (2) The above compound and the corresponding starting compound were reacted in the similar manner as EXAMPLE 1-(5) to give the compound (103-B).



MS (m/z) APCI: 536/538 [M+H]⁺

10 (3) The above compound (107 mg, 0.21 mmol) was dissolved in methanol (2 ml), and thereto were added two drops of a 4N solution of hydrogen chloride in dioxane. The mixture was stirred at room temperature for 24 hours, diluted with ethyl acetate, washed with water and brine, dried over magnesium sulfate and concentrated in *vacuo*, and the resulting residue was purified by silica gel chromatography (0 to 3% methanol-ethyl acetate) to give the compound (103-C).

15 MS (m/z) APCI: 430/432 [M+H]⁺

Corresponding starting compounds were treated in the similar manner as the above-mentioned to give the following compounds.




EXAMPLE No.	No	Structure	MS (m/z)
-------------	----	-----------	----------







103	1		477 APCI [M+H] ⁺
103	2		410 APCI [M+H] ⁺






EXAMPLE 104

Corresponding starting compounds were treated in the similar manner as EXAMPLE 67 using the compound 67-B or the corresponding enantiomer synthesized in the method of EXAMPLE 67 to give the following compounds.

5


EXAMPLE No.	No	Structure	MS (m/z)
104	1		550 ESI+ [M+H] ⁺
104	2		444 ESI+ [M+H] ⁺
104	3		491 ESI+ [M+H] ⁺

104	4		444 ESI+ [M+H] ⁺
104	5		491 ESI+ [M+H] ⁺
104	6		478 APCI [M+H] ⁺
104	7		482 APCI [M+H] ⁺
104	8		461 APCI [M+H] ⁺
104	9		438/440 APCI [M+H] ⁺

104	10		418 APCI [M+H] ⁺
104	11		472 APCI [M+H] ⁺
104	12		424 APCI [M+H] ⁺
104	13		462 APCI [M+H] ⁺
104	14		454 APCI [M+H] ⁺

EXAMPLE 105

228



(1) To a solution of the compound (87-D) (300 mg, 0.80 mmol), 2-tributylstannylpyrazine (384 mg, 1.04 mmol) and copper (I) iodide (15.2 mg, 0.08 mmol) in THF (6 ml) was added tetrakis(triphenylphosphine)palladium (92.4 mg, 0.08 mmol) at room temperature under argon, and the mixture was heated to reflux for 6 hours. After cooling, thereto was added a 10% aqueous potassium fluoride solution and the mixture was diluted with ethyl acetate, and then the precipitated insoluble was filtered off through Celite. The filtrate was washed with brine, dried over magnesium sulfate and concentrated in *vacuo*. The residue was purified by silica gel chromatography (30 to 60% ethyl acetate-hexane) to give the compound (105-A) (206 mg, yield 79%).

MS (m/z) APCI: 328 [M+H]⁺

(2) The above compound was reacted in the similar manner as EXAMPLE 1-(4-2-2) to give the compound (105-B).


MS (m/z) ESI: 312 [M-H]⁻

(3) The above compound and the corresponding starting compound were reacted in the similar manner as EXAMPLE 1-(5) to give the

compound (105-C).

MS (m/z) APCI: 414 [M+H]⁺

EXAMPLE 106



- 5 (1) The compound (1-D) and the corresponding starting compound were reacted in the similar manner as EXAMPLE 1-(4-1) to give the compound (106-A).

MS (m/z) ESI: 326 [M-H]⁻


- 10 (2) The above compound and the corresponding starting compound were reacted in the similar manner as EXAMPLE 1-(5) to give the compound (103-B).

MS (m/z) ESI: 491 [M+H]⁺

Corresponding starting compounds were treated in the similar manner as the above-mentioned to give the following compounds.

EXAMPLE No.	No	Structure	MS (m/z)
106	1		444 ESI+ [M+H] ⁺
106	2		424 ESI+ [M+H] ⁺
106	3		454 APCI [M+H] ⁺

EXAMPLE 107



(1) The compound (1-D) and the corresponding starting compound were reacted in the similar manner as EXAMPLE 1-(4-1) to give the compound (107-A).

MS (m/z) ESI: 340 [M-H]⁻

(2) The above compound and the corresponding starting compound were reacted in the similar manner as EXAMPLE 1-(5) to give the compound (107-B).


MS (m/z) ESI: 458/460 [M+H]⁺

Corresponding starting compounds were treated in the similar manner as the above-mentioned to give the following compounds.

EXAMPLE No.	No	Structure	MS (m/z)
107	1		439 ESI+ [M+H] ⁺
107	2		564 ESI+ [M+H] ⁺

107	3		438 ESI+ [M+H] ⁺
107	4		505 APCI [M+H] ⁺
107	5		578 APCI [M+H] ⁺
107	6		468 APCI [M+H] ⁺

EXAMPLE 108



(1) The compound (1-D) and the corresponding starting compound were reacted in the similar manner as EXAMPLE 1-(4-2) to give the compound (108-A).

MS (m/z) ESI: 359 [M-H]⁻






(2) The above compound and the corresponding starting compound

were reacted in the similar manner as EXAMPLE 1-(5) to give the compound (108-B).

MS (m/z) APCI: 524 [M+H]⁺


Corresponding starting compounds were treated in the similar manner as the above-mentioned to give the following compounds.

5

EXAMPLE No.	No	Structure	MS (m/z)
108	1		477/479 APCI [M+H] ⁺
108	2		583 APCI [M+H] ⁺
108	3		457 APCI [M+H] ⁺
108	4		458 APCI [M+H] ⁺
108	5		597 APCI [M+H] ⁺

108	6		487 APCI [M+H] ⁺
108	7		461 APCI [M+H] ⁺

EXAMPLE 109



(1) The compound (1-D) and the corresponding starting compound were reacted in the similar manner as EXAMPLE 1-(4-2) to give the compound (109-A).







MS (m/z) ESI: 360 [M-H]⁻

(2) The above compound and the corresponding starting compound were reacted in the similar manner as EXAMPLE 1-(5) to give the compound (109-B).

MS (m/z) APCI: 525 [M+H]⁺

Corresponding starting compounds were treated in the similar manner as the above-mentioned to give the following compounds.



EXAMPLE No.	No	Structure	MS (m/z)
-------------	----	-----------	----------

109	1		478/480 APCI [M+H] ⁺
109	2		458 APCI [M+H] ⁺
109	3		584 APCI [M+H] ⁺
109	4		459 APCI [M+H] ⁺
109	5		598 APCI [M+H] ⁺
109	6		488 APCI [M+H] ⁺


EXAMPLE 110

Corresponding starting compounds were reacted in the similar

manner as EXAMPLE 1 using (R)-(-)-2,2-dimethyl-1,3-dioxolan-4-methanol or the corresponding (S)-isomer to give the following compounds.

EXAMPLE No.	No	Structure	MS (m/z)
110	1		547 APCI [M+H] ⁺
110	2		547 APCI [M+H] ⁺

EXAMPLE 111



5

To a solution of the compound of EXAMPLE 110-(1) (300 mg, 0.55 mmol) in THF (4 ml) was added 1N hydrochloric acid (2 ml, 2 mmol) at room temperature, and the mixture was stirred at the same temperature for 16 hours. The mixture was extracted with chloroform, and the organic layer was dried over sodium sulfate and concentrated in *vacuo*. The residue was purified by silica gel column chromatography (10% methanol-chloroform) to give the above compound (230 mg, yield 83%).


10

MS (m/z) APCI: 507 [M+H]⁺

15


The compound of EXAMPLE 110-(2) was reacted in the similar

manner as the above-mentioned to give the following compound.

EXAMPLE No.	No	Structure	MS (m/z)
111	1		507 APCI [M+H] ⁺

EXAMPLE 112

A corresponding starting compound was reacted in the similar manner as EXAMPLE 1-(5) to give the following compound.





5

MS (m/z) APCI: 462 [M+H]⁺

EXAMPLE 113


Corresponding starting compounds were reacted in the similar manner as EXAMPLE 40 to give the following compounds.

EXAMPLE No.	No	Structure	MS (m/z)
113	1		475 APCI [M+H] ⁺
113	2		503 APCI [M+H] ⁺

10

EXAMPLE 114

237



(1) To a solution of 2,5-dibromopyridine (23.7 g, 100 mmol) in diethyl ether (800 ml) was added dropwise a 2.6 M solution of n-butyllithium in hexane (39.2 ml, 102 mmol) over 15 minutes at -78°C under argon, and the mixture was stirred at the same temperature for 20 minutes. Then, thereto was added dropwise diethyl oxalate (16.4 ml, 120 mmol) over 15 minutes. The mixture was stirred at the same temperature for 30 minutes and warmed to 0°C over another 3 hours, poured into a saturated aqueous ammonium chloride solution, and thereto was added diethyl ether. The organic layer was separated, washed sequentially with water and brine, dried over magnesium sulfate and concentrated in *vacuo*. The residue was purified by silica gel chromatography (14% ethyl acetate-hexane) to give the compound (114-A) (11.6 g, yield 51%). MS (m/z) APCI: 276 $[\text{M}+\text{H}+\text{H}_2\text{O}]^{+}$

(2) To a solution of the above compound (11.6 g, 44.9 mmol) in DMF (200 ml) was added sodium thiomethoxide (3.15 g, 44.9 mmol) at room temperature. The mixture was stirred at the same temperature overnight, diluted with ethyl acetate, and then washed sequentially with water and brine, dried over magnesium sulfate and concentrated in *vacuo*. The residue was purified by silica gel chromatography (9% ethyl acetate-hexane) to give the compound (114-B) (11.2 g, yield 69%).

MS (m/z) APCI: 226 [M+H]⁺


(3) The above compound and the corresponding starting compound were reacted in the similar manner as EXAMPLE 65-(1) and 1-(5) to give the compound (114-C).

5 MS (m/z) APCI: 363 [M+H]⁺


(4) In this case, the compound (114-D) which was believed to be derived from (114-A) remaining unreacted in the reaction of the above (3) was also obtained.

MS (m/z) APCI: 395/397 [M+H]⁺

10 A corresponding starting compound was treated in the similar manner as the above-mentioned method and EXAMPLE 27 to synthesize the following compound.

EXAMPLE No.	No	Structure	MS (m/z)
114	1		429/431 APCI [M+H] ⁺

EXAMPLE 115



(1) To a solution of 4H-[1,2,4]triazole-3-carboxylic acid methyl ester (25.0 g, 197 mmol) in pyridine (350 ml) was added trityl chloride (65.8 g, 236 mmol) under ice-cooling. The mixture was stirred at room temperature 90 minutes and at 100°C for 2 hours and concentrated in *vacuo*, and then the residue was solidified with isopropanol to give the compound (115-A) (68.8 g, yield 95%).

(2) To a solution of the above compound (64.6 g, 175 mmol) in THF (1000 ml) was added gradually lithium aluminum hydride (10.8 g, 284 mmol) under ice-cooling, and the ice bath was removed. After stirring at room temperature for 3 hours, thereto was added a 10% aqueous sodium hydroxide solution, and the suspension was filtered. The filtrate was concentrated, and then the residue was solidified with diethyl ether to give the compound (115-B) (34.8 g, yield 58%).

MS (m/z) ESI: 364 [M+Na]⁺

(3) To a solution of the above compound (17.4 g, 50.8 mmol) and

diisopropylethylamine (9.9 g, 76.2 mmol) in methylene chloride (250 ml) was added methanesulfonyl chloride (7.76 g, 67.8 mmol) under ice-cooling, and the mixture was stirred at the same temperature for 30 minutes. To the reaction mixture was added water and the organic layer was separated, and dried over sodium sulfate, and concentrated in *vacuo* to give the crude methanesulfonate (115-C) (23.6 g).

(4) To a solution of the compound (1-D) (12.0 g, 42.4 mmol) and potassium carbonate (17.6 g, 127 mmol) in DMF (250 ml) was added dropwise a solution of the above sulfonate (115-C) in DMF (50 ml) under ice-cooling, and the ice bath was removed. After stirring at room temperature for 16 hours, the mixture was diluted with ethyl acetate, washed sequentially with water and brine, dried over sodium sulfate and concentrated in *vacuo* to give the crude ester (31.7 g, 115-D).

(5) The above compound was reacted in the similar manner as EXAMPLE 1-(4-2-2) to give the compound (115-E).

MS (m/z) ESI: 591 [M-H]⁻

(6) The above compound and the corresponding starting compound were reacted in the similar manner as EXAMPLE 1-(5) to give the compound (115-F).







MS (m/z) ESI: 689 [M+H]⁺


(7) The above compound (191 mg, 0.28 mmol) was dissolved in formic acid (3 ml). The mixture was stirred at room temperature for 18 hours and concentrated in *vacuo*, and then the residue was purified by silica gel chromatography (2 to 6% methanol-chloroform) to give the compound (115-G) (75 g, yield 61%).

MS (m/z) APCI: 447 [M+H]⁺


Corresponding starting compounds were treated in the similar manner as the above-mentioned to give the following compounds.

EXAMPLE No.	No	Structure	MS (m/z)
-------------	----	-----------	----------

115	1		448 APCI [M+H] ⁺
115	2		587 APCI [M+H] ⁺
115	3		484 APCI [M+H] ⁺
115	4		514 APCI [M+H] ⁺
115	5		462 APCI [M+H] ⁺
115	6		477 APCI [M+H] ⁺

115	8		451 APCI [M+H] ⁺
-----	---	---	--------------------------------

EXAMPLE 116



(1) The compound (1-D) and tert-butyldimethylsilyl-protected corresponding starting compound were reacted in the similar manner as EXAMPLE 67-(1), (2) to give the compound (116-A).

MS (m/z) ESI: 466 [M-H]⁻







(2) The above compound and the corresponding starting compound were reacted in the similar manner as EXAMPLE 1-(5) to give the compound (116-B).

(3) The above compound was reacted in the similar manner as EXAMPLE 67-(4) to give the compound (116-C).

MS (m/z) APCI: 451 [M+H]⁺


Corresponding starting compounds were treated in the similar manner as the above-mentioned to give the following compounds.

EXAMPLE No.	No	Structure	MS (m/z)
-------------	----	-----------	----------

116	1		487 APCI [M+H] ⁺
116	2		590 APCI [M+H] ⁺
116	3		480 APCI [M+H] ⁺
116	4		517 APCI [M+H] ⁺
116	5		574 APCI [M+H] ⁺
116	6		547 APCI [M+H] ⁺

EXAMPLE 117

244



(1) To a solution of 2-chloro-4-iodoaniline (25 g, 96.7 mmol) in THF (100 ml) was added acetic anhydride (47 ml, 473 mmol) under ice-cooling. The mixture was stirred at the same temperature for 10 minutes and at room temperature for another 18 hours, and concentrated in *vacuo*. The residue was recrystallized from a mixture of ethyl acetate-hexane (1:1) to give the compound (117-A) (28.6 g, quantitatively).

MS (m/z) APCI: 296/298 [M+H]⁺

(2) To a solution of the above compound (2.4 g, 8.1 mmol) and sodium azide (1.1 g, 16 mmol) in acetonitrile (40 ml) and methylene chloride (5 ml) was added trifluoromethanesulfonic anhydride (3.4 g, 12 mmol) under ice-cooling, and the ice bath was removed. The mixture was stirred at room temperature for 20 hours and concentrated in *vacuo*, and then to the residue was added ethyl acetate. The mixture

was washed sequentially with water and brine, dried over magnesium sulfate and concentrated in *vacuo*. The residue was purified by silica gel chromatography (20% ethyl acetate-hexane) to give the compound (117-B) (2.07 g, yield 80%).

5 MS (m/z) APCI: 321/323 [M+H]⁺

(3) To a solution of the above compound (4.8 g, 15 mmol), bis(tributyltin) (17.5 g, 30 mmol) in toluene (70 ml) was added dichlorobis(triphenylphosphine)palladium (1.06 g, 1.5 mmol) at room temperature under argon. The mixture was stirred at 100°C for 3 hours.

10 After air-cooling, thereto was added a 10% aqueous potassium fluoride solution, and the mixture was stirred at room temperature for 1 hour, and the precipitated insoluble was filtered off through Celite. The filtrate was dried over sodium sulfate and concentrated in *vacuo* and the residue was purified by silica gel chromatography (0 to 20% ethyl acetate-hexane) to give the compound (117-C) (4.8 g, yield 66%).

15 MS (m/z) APCI: 481/483/485 [M+H]⁺

(4) To a solution of the above compound (7.6 g, 16 mmol), diisopropylethylamine (3.6 ml, 20 mmol), potassium carbonate (130 mg, 0.94 mmol) and tris(dibenzylideneacetone)dipalladium (720 mg, 0.78

20 mmol) in THF (70 ml) was added ethyl chlorooxalate (2.64 ml, 24 mmol) at room temperature under argon, and the mixture was stirred at the same temperature for 3 hours. To the reaction mixture was added a 10% aqueous potassium fluoride solution, and the mixture was stirred at the same temperature for 3 hours and diluted with diethyl ether. The

25 insoluble was filtered off and the organic layer of the filtrate was separated, washed sequentially with water and brine, dried over magnesium sulfate and concentrated in *vacuo*. The residue was purified by silica gel chromatography (30% ethyl acetate-hexane) to give the compound (117-D) (1.04 g, yield 22%).



30 MS (m/z) APCI: 295/297 [M+H]⁺

(5) The above compound and the corresponding starting compound were reacted in the similar manner as EXAMPLE 1-(3), 65-(1), 1-(4-2-2) and (5) to give the compound (117-E).

MS (m/z) APCI: 432/434 [M+H]⁺


5 EXAMPLE 118


The compound (79-C) and the corresponding starting compounds were treated in the similar manner as EXAMPLE 1-(5) to give the following compounds.

EXAMPLE No.	No	Structure	MS (m/z)
118	1		504 APCI [M+H] ⁺
118	2		563 APCI [M+H] ⁺

EXAMPLE 119


10 Corresponding starting compounds were treated in the similar manner as EXAMPLE 1-(3), (4-1), (5) to give the following compounds.

EXAMPLE No.	No	Structure	MS (m/z)
119	1		498 APCI [M+H] ⁺

119	2		493 APCI [M+H] ⁺
-----	---	---	--------------------------------

EXAMPLE 120



A corresponding starting compound was treated in the similar manner as EXAMPLE 73 to give the following compound.



5 MS (m/z) APCI: 454 [M+H]⁺


EXAMPLE 121

Corresponding starting compounds were treated in the similar manner as EXAMPLE 1-(4-1), (5) to give the following compounds.

EXAMPLE No.	No	Structure	MS (m/z)
121	1		472 APCI [M+H] ⁺
121	2		502 APCI [M+H] ⁺

EXAMPLE 122

248



- (1) To a solution of 4-bromo-3-methylbenzenesulfonyl chloride (5.4 g, 200 mmol) in THF (120 ml) was added sodium borohydride (7.6 g, 200 mmol) at room temperature, and the mixture was heated to reflux for 16 hours. After cooling to room temperature, the reaction mixture was diluted with ethyl acetate, washed sequentially with 5N hydrochloric acid, water and brine, dried over sodium sulfate and concentrated in *vacuo* to give the crude thiol (5.2 g, quantitatively). A 2.0 g portion of the resulting crude thiol was dissolved in DMSO (50 ml), and thereto was added potassium tert-butoxide (1.2 g, 11 mmol), and the mixture was stirred at room temperature. At the mixture became homogeneous, cyclopropyl bromide (2.4 ml, 30 mmol) was added and the mixture was stirred at 80°C for 2 days. After cooling to room temperature, the reaction mixture was poured into water and the mixture was extracted with ethyl acetate (200 ml) twice. The organic layers were combined, washed sequentially with water and brine, dried over sodium sulfate, concentrated in *vacuo*, and then the residue was purified by silica gel

column chromatography (hexane) to give the compound (122-A) (155 mg, yield 8% in 2 steps) as an oil.

(2) To a solution of the above compound (150 mg, 0.62 mmol) in diethyl ether (15 ml) was added dropwise a 1.59 M solution (0.81 ml, 1.3 mmol) of tert-butyllithium in pentane at -70°C over 4 minutes, and thereto was added trimethyl borate (0.076 ml, 0.68 mmol) after 10 minutes in one portion, and the mixture was warmed to 0°C . Thereto was added a saturated aqueous ammonium chloride solution, and the mixture was stirred at room temperature for 15 minutes, and then was extracted with ethyl acetate (20 ml). The organic layer was dried over sodium sulfate and concentrated in *vacuo* to give the crude boronic acid (122-B).

(3) The above crude boronic acid was suspended in 1,2-dimethoxyethane (3 ml), and thereto were added the compound (122-C) (114 mg, 0.40 mmol), which was synthesized by reacting a corresponding starting compound in the similar manner as EXAMPLE 91, a 1M aqueous sodium carbonate solution (1.2 ml, 1.2 mmol) and dichlorobis(triphenylphosphine)palladium (28 mg, 0.04 mmol). After stirring at 80°C for 4 hours, the mixture was cooled to room temperature, poured into water and extracted with ethyl acetate (15 ml). The organic layer was washed with brine, dried over sodium sulfate and concentrated in *vacuo*. The resulting residue was purified by silica gel column chromatography (5% to 25% hexane-ethyl acetate) to give the compound (122-D) (30 mg, yield 20%) as an oil.

MS (m/z) APCI: 370 $[\text{M}+\text{H}]^{+}$

(4) The above compound was treated in the similar manner as EXAMPLE 1-(4-2-2) to give the compound (122-E).

MS (m/z) ESI: 681 $[2\text{M}-\text{H}]^{-}$

(5) The above compound and the corresponding starting compound were reacted in the similar manner as EXAMPLE 1-(5) to give the

compound (122-F).





MS (m/z) APCI: 442 [M+H]⁺


(6) The above compound was treated in the similar manner as EXAMPLE 84 to give the compound (122-G).

5 MS (m/z) APCI: 474 [M+H]⁺

EXAMPLE 123

Corresponding starting compounds were treated in a combination of the methods of EXAMPLE 1-(5), EXAMPLE 15, EXAMPLE 17 using the compound (77-D) to give the following compounds.

EXAMPLE No.	No	Structure	MS (m/z)
123	1		494 APCI [M+H] ⁺
123	2		604 APCI [M+H] ⁺
123	3		574 APCI [M+H] ⁺
123	4		561 APCI [M+H] ⁺




To a suspension of the compound of EXAMPLE 102-(24) (80 mg, 0.162 mmol), propargyl alcohol (0.047 ml, 0.81 mmol), copper (I) iodide (6.2 mg, 0.03 mmol) and triethylamine (0.5 ml, 3.6 mmol) in THF (2 ml) was added dichlorobis(triphenylphosphine)palladium (22.7 mg, 0.03 mmol) at room temperature under argon, and the mixture was heated to reflux for 8 hours. After cooling to room temperature, the mixture was concentrated in *vacuo* and the resulting residue was purified by silica gel chromatography (50 to 100% ethyl acetate-hexane) to give the above compound (25 mg, yield 33%).

MS (m/z) APCI: 469 [M+H]⁺

A corresponding starting compound was treated in the similar manner as the above-mentioned to give the following compound.

EXAMPLE No.	No	Structure	MS (m/z)
124	1		496 APCI [M+H] ⁺

EXAMPLE 125




15

A corresponding starting compound was treated in the similar

manner as EXAMPLE 58 to give the above compound.

MS (m/z) APCI: 443 [M+H]⁺


EXAMPLE 126



- 5 A corresponding starting compound was treated in the similar manner as EXAMPLE 29 to give the above compound.

MS (m/z) APCI: 459 [M+H]⁺


EXAMPLE 127



- 10 A corresponding starting compound was treated in the similar manner as EXAMPLE 26 to give the above compound.

MS (m/z) APCI: 445 [M+H]⁺


EXAMPLE 128



- 15 A corresponding starting compound was treated in the similar manner as EXAMPLE 83 and 84 to give the above compound.

MS (m/z) APCI: 498 [M+H]⁺

EXAMPLE 129




(1) The compound of EXAMPLE 97-(4) was reacted in the similar manner as EXAMPLE 83-(1) to give the compound (129-A).

MS (m/z) APCI: 403 [M+H]⁺

5 (2) To a solution of the above compound (100 mg, 0.25 mmol) in DMF (2 ml) were added sequentially ammonium chloride (134 mg, 2.5 mmol) and sodium azide (162 mg, 2.5 mmol), and the mixture was stirred at 115°C for 3 hours. After cooling to room temperature, the mixture was acidified with 10% hydrochloric acid, diluted with ethyl acetate, washed sequentially with water and brine, dried over
10 magnesium sulfate and concentrated in *vacuo* to give the compound (129-B).

MS (m/z) ESI: 444 [M-H]⁻


EXAMPLE 130



15 To a solution of the compound of EXAMPLE (134) (100 mg, 0.22 mmol) in pyridine (5 ml) was added ethyl chlorocarbonate (0.023 ml, 0.24 mmol), and the mixture was stirred at 120°C for 3 hours. After cooling to room temperature, the reaction mixture was concentrated in *vacuo*
20 and the residue was purified by silica gel chromatography (0 to 5% methanol-chloroform) to give the above compound (44 mg, yield 38%).

MS (m/z) ESI: 522 [M-H]⁻


EXAMPLE 131



To a solution of the compound of EXAMPLE 86 (58 mg, 0.11 mmol) and diisopropylethylamine (0.098 ml, 0.57mmol) in chloroform (2 ml) was added acetic anhydride (0.021 ml, 0.22 mmol) under ice-cooling, and the mixture was stirred at room temperature overnight. To the reaction mixture was added a saturated aqueous sodium bicarbonate solution, and the mixture was extrated with chloroform. The organic layer was washed sequentially with water and brine, dried over sodium sulfate and concentrated in *vacuo*. The resulting residue was purified by silica gel chromatography (0 to 5% methanol-chloroform) to give the above compound (42 mg, yield 77%).

MS (m/z) APCI: 485 [M+H]⁺

EXAMPLE 132




To a mixture of the compound of EXAMPLE 86 (59 mg, 0.11 mmol) and diisopropylethylamine (0.060 ml, 0.034mmol) in a 38% aqueous formalin solution (1 ml) and chloroform (2 ml) was added sodium triacetoxyborohydride (73 mg, 0.34 mmol) under ice-cooling, and the mixture was stirred at the same temperature for 1 hour and at room temperature for 3 hours. Then, to the reaction mixture was added

a saturated aqueous sodium bicarbonate solution, and the mixture was extracted with chloroform. The organic layer was washed sequentially with water and brine, dried over sodium sulfate and concentrated in *vacuo*. The resulting residue was purified by NH-silica gel chromatography (0 to 15% methanol-ethyl acetate) to give the above compound (29 mg, yield 54%).

MS (m/z) APCI: 471 [M+H]⁺

EXAMPLE 133




- 10 (1) The compound (83-E) was treated in the similar manner as
EXAMPLE 83 to give the compound (133-A).
- (2) The above compound was treated in the similar manner as
EXAMPLE 115-(7) to give the compound (133-B).
- MS (m/z) APCI: 449 [M+H]⁺
- 15 (3) The compound (133-A) was treated in the similar manner as
EXAMPLE 84 to give the compound (133-C).

(4) The above compound was treated in the similar manner as the above (2) to give the compound (133-D).

MS (m/z) APCI: 481 [M+H]⁺

EXAMPLE 134




5

The compound of EXAMPLE 91-(4) was treated in the similar manner as EXAMPLE 30 to give the above compound.

MS (m/z) APCI: 452 [M+H]⁺

EXAMPLE 135



10

(1) The compound of EXAMPLE 125 was treated in the similar manner as EXAMPLE 26 to give the compound (135-A).

MS (m/z) APCI: 445 [M+H]⁺

(2) To a solution of the compound obtained in the above (1) (78 mg, 0.18 mmol) and triphenylphosphine (92 mg, 0.35 mmol) in THF (2 ml) was added carbon tetrabromide (116 mg, 0.35 mmol) at room temperature, and the mixture was stirred at the same temperature for 6 hours and concentrated in *vacuo*. The residue was purified by silica gel column chromatography (30 to 80% ethyl acetate-hexane) to give the

15


compound (135-B) (73 mg, yield 80%).

MS (m/z) APCI: 507/509 [M+H]⁺

(3) To a solution of the compound obtained in the above (2) (70 mg, 0.14 mmol) in methanol (3 ml) was added silver (II) oxide (34 mg, 0.28 mmol) at room temperature. The mixture was stirred at the same temperature for 16 hours and filtered. The filtrate was concentrated in *vacuo*, and the resulting residue was purified by silica gel column chromatography (30 to 100% ethyl acetate-hexane) to give the compound (135-C) (10 mg, yield 16%).

MS (m/z) APCI: 459 [M+H]⁺

EXAMPLE 136



(1) A suspension of the compound (77-B) of EXAMPLE 77 (1.95 g, 5.31 mmol) and molecular sieves (4Å, powder) (1.95 g) in 1,2-dichloroethane (38 ml) was ice-cooled and thereto were added sequentially 1,8-bis(dimethylamino)naphthalene (3.35 g, 15.6 mmol) and triethyloxonium tetrafluoroborate (2.97 g, 15.6 mmol). The mixture was stirred at room temperature for 18 hours and filtered. To the filtrate was added ethyl acetate. The mixture was washed with brine, dried over sodium sulfate and then concentrated in *vacuo*, and the residue was purified by silica gel column chromatography (33 to 50%

ethyl acetate-hexane) to give the compound (136-A) (1.54 g, yield 73%).

MS (m/z) APCI: 396 [M+H]⁺


(2) The compound obtained in the above (1) was treated in the similar manner as EXAMPLE 1-(4-2-2) to give the compound (136-B).

5 MS (m/z) APCI: 783 [2M-2H+Na]⁻

(3) The compound obtained in the above (2) was treated in the similar manner as EXAMPLE 1-(5) to give the compound (136-C).

MS (m/z) APCI: 482 [M+H]⁺

EXAMPLE 137



10

To a solution of the compound of EXAMPLE 60 (49 mg, 0.11 mmol) and pyridine (0.50 ml) in methanol (1.0 ml) was added hydroxylamine hydrochloride (15 mg, 0.21 mmol), and the mixture was stirred at room temperature for 16 hours. To the reaction mixture was added chloroform, and the mixture was washed with water and brine, dried over sodium sulfate and concentrated in *vacuo*. The residue was purified by silica gel column chromatography (0 to 5% methanol-chloroform) to give the titled compound (52 mg, quantitatively).


15

MS (m/z) APCI: 474 [M+H]⁺

20

EXAMPLE 138

259



(1) The compound (1-E) of EXAMPLE 1 was treated in the similar manner as EXAMPLE 1-(5) to give the compound (138-A).

MS (m/z) APCI: 637 [M+H]⁺








(2) A solution of the compound obtained in the above (1) (3.20 g, 5.03 mmol) in chloroform-trifluoroacetic acid (1:1) (80 ml) was stirred at room temperature for 2 days. The reaction mixture was concentrated in *vacuo* and the residue was chased with toluene, and then to the residue was added a saturated aqueous sodium bicarbonate solution. The mixture was extracted with chloroform. The extract was dried over sodium sulfate and concentrated in *vacuo*. The residue was then purified by silica gel column chromatography (methanol-chloroform-ammonia water 200:10:1 to 100:10:1) to give the compound (138-B) (2.06 g, yield 77%).

MS (m/z) APCI: 537 [M+H]⁺








EXAMPLE 139








Corresponding starting compounds were treated in the similar manner as any of the above EXAMPLEs to give the following compounds.








EXAMPLE	No.	Structure	MS (m/z)
---------	-----	-----------	----------








No.			
139	1		582 APCI [M+H] ⁺
139	2		568 APCI [M+H] ⁺
139	3		430 APCI [M+H] ⁺
139	4		457 APCI [M+H] ⁺
139	5		462 ESI [M- H] ⁻
139	6		522 ESI [M- H] ⁻
139	7		456 ESI [M- H] ⁻








261

139	8		494 ESI [M-H] ⁻
139	9		504 ESI [M-H] ⁻
139	10		540/542 ESI [M-H] ⁻
139	11		476 ESI [M-H] ⁻
139	12		476 ESI [M-H] ⁻
139	13		ESI 512 [M-H] ⁻
139	14		530 ESI [M-H] ⁻







139	15		506 APCI [M+H] ⁺
139	16		506 APCI [M+H] ⁺
139	17		556/558 APCI [M+H] ⁺
139	18		506 APCI [M+H] ⁺
139	19		534 APCI [M+H] ⁺
139	20		542 APCI [M+H] ⁺
139	21		492 APCI [M+H] ⁺







139	22		524 APCI [M+H] ⁺
139	23		459 APCI [M+H] ⁺
139	24		502 ESI [M- H] ⁻
139	25		561 APCI [M+H] ⁺
139	26		601 APCI [M+H] ⁺
139	27		605 APCI [M+H] ⁺
139	28		611 APCI [M+H] ⁺







139	29		575 APCI [M+H] ⁺
139	30		475 APCI [M+H] ⁺
139	31		558 APCI [M+H] ⁺
139	32		572 APCI [M+H] ⁺
139	33		482 APCI [M+H] ⁺
139	34		508 APCI [M+H] ⁺
139	35		618 APCI [M+H] ⁺

139	36		575 APCI [M+H] ⁺
139	37	 HCl	588 APCI [M+H] ⁺
139	38		559 APCI [M+H] ⁺
139	39		546 APCI [M+H] ⁺
139	40		485 APCI [M+H] ⁺
139	41		475 APCI [M+H] ⁺
139	42		460 APCI [M+H] ⁺







139	43		489 APCI [M+H] ⁺
139	44		474 APCI [M+H] ⁺
139	45		516 APCI [M+H] ⁺
139	46		528 APCI [M+H] ⁺
139	47		602 APCI [M+H] ⁺
139	48		588 APCI [M+H] ⁺







139	49		450 APCI [M+H] ⁺
139	50		570 APCI [M+H] ⁺
139	51		498 APCI [M+H] ⁺
139	52		588 APCI [M+H] ⁺
139	53		588 APCI [M+H] ⁺
139	54		573 APCI [M+H] ⁺







139	55		474 APCI [M+H] ⁺
139	56		436 APCI [M+H] ⁺
139	57		478 APCI [M+H] ⁺
139	58		586 APCI [M+H] ⁺
139	59		544 APCI [M+H] ⁺
139	60		558 APCI [M+H] ⁺

139	61		574 APCI [M+H] ⁺
139	62		572 APCI [M+H] ⁺
139	63		560 APCI [M+H] ⁺
139	64		560 APCI [M+H] ⁺
139	65		558 APCI [M+H] ⁺
139	66		588 APCI [M+H] ⁺

270

139	67		574 APCI [M+H] ⁺
139	68		572 APCI [M+H] ⁺
139	69		558 APCI [M+H] ⁺
139	70		572 APCI [M+H] ⁺
139	71		559 APCI [M+H] ⁺
139	72		502 APCI [M+H] ⁺








139	73		532 APCI [M+H] ⁺
139	74		546 APCI [M+H] ⁺
139	75		559 APCI [M+H] ⁺
139	76		APCI 537 [M+H] ⁺
139	77		APCI 551 [M+H] ⁺
139	78		APCI 579 [M+H] ⁺

139	79		545 [M+H] ⁺
139	80		532 [M+H] ⁺
139	81		588 [M+H] ⁺
139	82		483 APCI [M+H] ⁺
139	83		484 APCI [M+H] ⁺
139	84		484 APCI [M+H] ⁺

139	85		513 APCI [M+H] ⁺
139	86		623 APCI [M+H] ⁺
139	87		550 APCI [M+H] ⁺
139	88		464 APCI [M+H] ⁺
139	89		514 APCI [M+H] ⁺
139	90		494 APCI [M+H] ⁺
139	91		508 APCI [M+H] ⁺








139	92		503 APCI [M+H] ⁺
139	93		522 APCI [M+H] ⁺
139	94		504 APCI [M+H] ⁺
139	95		609 APCI [M+H] ⁺
139	96		469 APCI [M+H] ⁺
139	97		506 APCI [M+H] ⁺
139	98		536 APCI [M+H] ⁺








275









139	99		579 APCI [M+H] ⁺
139	100		567 APCI [M+H] ⁺
139	101		470 APCI [M+H] ⁺
139	102		499 APCI [M+H] ⁺
139	103		490 APCI [M+H] ⁺
139	104		480 APCI [M+H] ⁺
139	105		593 ESI [M+H] ⁺









139	106		501 APCI [M+H] ⁺
139	107		497 APCI [M+H] ⁺
139	108		527 APCI [M+H] ⁺
139	109		498 APCI [M+H] ⁺
139	110		564 APCI [M+H] ⁺
139	111		508 APCI [M+H] ⁺
139	112		518 APCI [M+H] ⁺









139	113		567 APCI [M+H] ⁺
139	114		579 APCI [M+H] ⁺
139	115		480 APCI [M+H] ⁺
139	116		496 APCI [M+H] ⁺
139	117		512 APCI [M+H] ⁺
139	118		541 APCI [M+H] ⁺
139	119		651 APCI [M+H] ⁺








139	120		515 APCI [M+H] ⁺
139	121		511 APCI [M+H] ⁺
139	122		609 APCI [M+H] ⁺
139	123		532 APCI [M+H] ⁺
139	124		479 APCI [M+H] ⁺
139	125		581 APCI [M+H] ⁺
139	126		593 APCI [M+H] ⁺








139	127		484 APCI [M+H] ⁺
139	128		513 APCI [M+H] ⁺
139	129		494 APCI [M+H] ⁺
139	130		510 APCI [M+H] ⁺
139	131		491 APCI [M+H] ⁺
139	132		495 APCI [M+H] ⁺
139	133		490 APCI [M+H] ⁺








139	134		557 APCI [M+H] ⁺
139	135		520 APCI [M+H] ⁺
139	136		614 APCI [M+H] ⁺
139	137		630 APCI [M+H] ⁺
139	138		588 APCI [M+H] ⁺
139	139		501 APCI [M+H] ⁺
139	140		600 APCI [M+H] ⁺
139	141		470 APCI [M+H] ⁺








139	142		467 APCI [M+H] ⁺
139	143		496 APCI [M+H] ⁺
139	144		576 APCI [M+H] ⁺
139	145		563 APCI [M+H] ⁺
139	146		604 APCI [M+H] ⁺
139	147		590 APCI [M+H] ⁺
139	148		564 APCI [M+H] ⁺
139	149		603 APCI [M+H] ⁺







139	150		589 APCI [M+H] ⁺
139	151		532 APCI [M+H] ⁺
139	152		518 APCI [M+H] ⁺
139	153		562 APCI [M+H] ⁺
139	154		576 APCI [M+H] ⁺
139	155		589 APCI [M+H] ⁺
139	156		507 APCI [M+H] ⁺
139	157		518 APCI [M+H] ⁺







139	158		530 APCI [M+H] ⁺
139	159		489 APCI [M+H] ⁺
139	160		462 APCI [M+H] ⁺
139	161		491 APCI [M+H] ⁺
139	162		505 APCI [M+H] ⁺
139	163		521 APCI [M+H] ⁺
139	164		486 APCI [M+H] ⁺

139	166		493 APCI [M+H] ⁺
139	167		487 APCI [M+H] ⁺
139	168		590 APCI [M+H] ⁺
139	169		588 APCI [M+H] ⁺
139	170		602 APCI [M+H] ⁺
139	171		476 APCI [M+H] ⁺
139	172		608 APCI [M+H] ⁺

139	173		521 APCI [M+H] ⁺
139	174		506 APCI [M+H] ⁺
139	175		514 APCI [M+H] ⁺
139	176		511 APCI [M+H] ⁺
139	177		540 APCI [M+H] ⁺
139	178		607 APCI [M+H] ⁺
139	179		622 APCI [M+H] ⁺








139	180		580 APCI [M+H] ⁺
139	181		493 APCI [M+H] ⁺
139	182		482 APCI [M+H] ⁺
139	183		512 APCI [M+H] ⁺
139	184		578 APCI [M+H] ⁺
139	185		486 APCI [M+H] ⁺
139	186		483 APCI [M+H] ⁺








139	187		579 APCI [M+H] ⁺
139	188		592 APCI [M+H] ⁺
139	189		606 APCI [M+H] ⁺
139	190		519 APCI [M+H] ⁺
139	191		549 APCI [M+H] ⁺
139	192		618 APCI [M+H] ⁺








139	193		500 APCI [M+H] ⁺
139	194		496 APCI [M+H] ⁺
139	195		497 APCI [M+H] ⁺
139	196		526 APCI [M+H] ⁺
139	197		563 APCI [M+H] ⁺
139	199		593 APCI [M+H] ⁺

139	200		636 APCI [M+H] ⁺
139	201		594 APCI [M+H] ⁺
139	202		492 APCI [M+H] ⁺
139	203		507 APCI [M+H] ⁺
139	204		523 APCI [M+H] ⁺
139	205		517 APCI [M+H] ⁺
139	206		HCl 618 APCI [M+H] ⁺









290

139	207		632 APCI [M+H] ⁺
139	208		578 APCI [M+H] ⁺
139	209		566 APCI [M+H] ⁺
139	210		479 APCI [M+H] ⁺
139	211		464 APCI [M+H] ⁺
139	212		495 APCI [M+H] ⁺
139	213		577 APCI [M+H] ⁺

139	214		564 APCI [M+H] ⁺
139	215		536 APCI [M+H] ⁺
139	216		606 APCI [M+H] ⁺
139	217		592 APCI [M+H] ⁺
139	218		605 APCI [M+H] ⁺
139	219		591 APCI [M+H] ⁺
139	220		578 APCI [M+H] ⁺





139	221		534 APCI [M+H] ⁺
139	222		520 APCI [M+H] ⁺
139	223		564 APCI [M+H] ⁺
139	224		578 APCI [M+H] ⁺
139	225		591 APCI [M+H] ⁺
139	226		477 APCI [M+H] ⁺
139	227		592 APCI [M+H] ⁺

293

139	228		590 APCI [M+H] ⁺
139	229		604 APCI [M+H] ⁺
139	230		483 APCI [M+H] ⁺
139	231		479 APCI [M+H] ⁺
139	232		480 APCI [M+H] ⁺
139	233		509 APCI [M+H] ⁺
139	234		546 APCI [M+H] ⁺
139	235		577 APCI [M+H] ⁺



139	236		564 APCI [M+H] ⁺
139	237		606 APCI [M+H] ⁺
139	238		533 APCI [M+H] ⁺
139	239		563 APCI [M+H] ⁺
139	240		496 APCI [M+H] ⁺
139	241		462 APCI [M+H] ⁺
139	242		590 APCI [M+H] ⁺






295







139	243		595 APCI [M+H] ⁺
139	244		565 APCI [M+H] ⁺
139	245		553 APCI [M+H] ⁺
139	246		480 APCI [M+H] ⁺






EXAMPLE 140






Corresponding starting compounds are treated in the similar manner as any of the above EXAMPLES to give the following compounds.

EXAMPLE No.	No.	Structure
140	1	
140	2	






140	3	
140	4	
140	5	
140	6	
140	7	

140	8	
140	9	
140	10	
140	11	
140	12	
140	13	






140	14	
140	15	
140	16	
140	17	
140	18	

140	19	
140	20	
140	21	
140	22	
140	23	






300








140	24	
140	25	
140	26	
140	27	
140	28	








301






140	29	
140	30	
140	31	
140	32	
140	33	







302







140	34	
140	35	
140	36	
140	37	
140	38	






140	39	
140	40	
140	41	
140	42	
140	43	
140	44	
140	45	







140	46	
140	47	
140	48	
140	49	
140	50	
140	51	
140	52	







140	53	
140	54	
140	55	
140	56	
140	57	








140	58	
140	59	
140	60	
140	61	
140	62	
140	63	







140	64	
140	65	
140	66	
140	67	
140	68	
140	69	







140	70	
140	71	
140	72	
140	73	
140	74	







140	75	
140	76	
140	77	
140	78	
140	79	
140	80	








140	81	
140	82	
140	83	
140	84	
140	85	
140	86	








140	87	
140	88	
140	89	
140	90	
140	91	
140	92	
140	93	








140	94	
140	95	
140	96	
140	97	
140	98	
140	99	








140	100	
140	101	
140	102	
140	103	
140	104	
140	105	








140	106	
140	107	
140	108	
140	109	
140	110	
140	111	

140	112	
140	113	
140	114	
140	115	
140	116	
140	117	
140	118	








140	119	
140	120	
140	121	
140	122	
140	123	
140	124	
140	125	

140	126	
140	127	
140	128	
140	129	
140	130	
140	131	
140	132	








140	133	
140	134	
140	135	
140	136	
140	137	
140	138	
140	139	








140	140	
140	141	
140	142	
140	143	
140	144	
140	145	
140	146	







140	147	
140	148	
140	149	
140	150	
140	151	
140	152	
140	153	








140	154	
140	155	
140	156	
140	157	
140	158	
140	159	
140	160	








140	161	
140	162	
140	163	
140	164	
140	165	
140	166	
140	167	








140	168	
140	169	
140	170	
140	171	
140	172	
140	173	
140	174	







140	175	
140	176	
140	177	
140	178	
140	179	
140	180	
140	181	







140	182	
140	183	
140	184	
140	185	
140	186	
140	187	
140	188	







140	189	
140	190	
140	191	
140	192	
140	193	
140	194	
140	195	








140	196	
140	197	
140	198	
140	199	
140	200	
140	201	
140	202	







140	203	
140	204	
140	205	
140	206	
140	207	
140	208	
140	209	








140	210	
140	211	
140	212	
140	213	
140	214	
140	215	

140	216	
140	217	
140	218	
140	219	
140	220	
140	221	







140	222	
140	223	
140	224	
140	225	
140	226	
140	227	







140	228	
140	229	
140	230	
140	231	
140	232	
140	233	
140	234	








140	235	
140	236	
140	237	
140	238	
140	239	
140	240	








140	241	
140	242	
140	243	
140	244	
140	245	
140	246	
140	247	







335

140	248	
140	249	
140	250	
140	251	
140	252	
140	253	







140	254	
140	255	
140	256	
140	257	
140	258	
140	259	







140	260	
140	261	
140	262	
140	263	
140	264	
140	265	
140	266	







140	267	
140	268	
140	269	
140	270	
140	271	
140	272	
140	273	








140	274	
140	275	
140	276	
140	277	
140	278	
140	279	








140	280	
140	281	
140	282	
140	283	
140	284	
140	285	

140	286	
140	287	
140	288	
140	289	
140	290	
140	291	








140	292	
140	293	
140	294	
140	295	
140	296	
140	297	

140	298	
140	299	
140	300	
140	301	
140	302	
140	303	






140	304	
140	305	
140	306	
140	307	
140	308	
140	309	
140	310	







140	311	
140	312	
140	313	
140	314	
140	315	
140	316	
140	317	

140	318	
140	319	
140	320	
140	321	
140	322	
140	323	







140	324	
140	325	
140	326	
140	327	
140	328	
140	329	
140	330	







348

140	331	
140	332	
140	333	
140	334	
140	335	







140	336	
140	337	
140	338	
140	339	
140	340	
140	341	

350








140	342	
140	343	
140	344	
140	345	
140	346	
140	347	








140	348	
140	349	
140	350	
140	351	
140	352	
140	353	








352

140	354	
140	355	
140	356	
140	357	
140	358	
140	359	







353








140	360	
140	361	
140	362	
140	363	
140	364	
140	365	
140	366	








140	367	
140	368	
140	369	
140	370	
140	371	
140	372	
140	373	








140	374	
140	375	
140	376	
140	377	
140	378	
140	379	
140	380	








356








140	381	
140	382	
140	383	
140	384	
140	385	
140	386	








140	387	
140	388	
140	389	
140	390	
140	391	
140	392	
140	393	







140	394	
140	395	
140	396	
140	397	
140	398	
140	399	
140	400	







140	401	
140	402	
140	403	
140	404	
140	405	
140	406	
140	407	






140	408	
140	409	
140	410	
140	411	
140	412	
140	413	
140	414	







140	415	
140	416	
140	417	
140	418	
140	419	
140	420	
140	421	

140	422	
140	423	
140	424	
140	425	
140	426	
140	427	
140	428	








140	429	
140	430	
140	431	
140	432	
140	433	
140	434	








140	435	
140	436	
140	437	
140	438	
140	439	
140	440	







140	441	
140	442	
140	443	
140	444	
140	445	

140	446	
140	447	
140	448	
140	449	
140	450	
140	451	







367

140	452	
140	453	
140	454	
140	455	
140	456	
140	457	
140	458	

140	459	
140	460	
140	461	
140	462	
140	463	
140	464	
140	465	







140	466	
140	467	
140	468	
140	469	
140	470	
140	471	


370

140	472	
140	473	
140	474	
140	475	
140	476	
140	477	


140	478	
140	479	
140	480	
140	481	
140	482	
140	483	
140	484	

140	485	
140	486	
140	487	
140	488	
140	489	
140	490	
140	491	

140	492	
140	493	
140	494	
140	495	
140	496	
140	497	

140	498	
-----	-----	--

REFERENCE EXAMPLE 1



To a solution of 2-tert-butyloxycarbonylaminothiazole (88.0 g, 439 mmol) in THF (1760 ml) was added dropwise a 1.59M solution of n-butyllithium in hexane (729 ml, 1159 mmol) over 20 minutes at -78°C , and the mixture was warmed to -10°C over 1 hour. The mixture was cooled again to -78°C and thereto was added DMF (102 ml, 0.132 mmol) in one portion. The acetone-dry ice bath was removed. The mixture was stirred for 30 minutes, and then poured into cold water (1000 ml) and thereto was added ethyl acetate (2000 ml). The organic layer was washed sequentially with water and brine, dried over anhydrous magnesium sulfate and the solvents were removed. The residue was recrystallized from ethyl acetate to give 2-tert-butyloxycarbonylaminothiazole-5-carbaldehyde (74.8 g).

mp. 173 to 175°C

MS (m/z) APCI: 229 (M+H)⁺


To a solution of the above 2-tert-butyloxycarbonylaminothiazole-5-carbaldehyde (64.5 g, 282 mmol) in methylene chloride (322 ml) was added dropwise trifluoroacetic acid (322 ml) under ice-cooling over 20 minutes. The mixture was stirred at room temperature for 2 hours and

the solvent was removed by evaporation. Thereto was added chloroform (50 ml), and then added a 4N hydrogen chloride solution in dioxane (300 ml) dropwise under ice-cooling. After removing the solvents, the residue was washed with ethyl acetate and collected to give the titled compound (41.3 g) as monohydrochloride thereof.

mp. 190 to 194°C (decomposed)

MS (m/z) APCI: not detected

REFERENCE EXAMPLE 2



To a solution of 2-tert-butyloxycarbonylaminothiazole (60.0 g, 299 mmol) in THF (1200 ml) was added dropwise a 1.59M solution of n-butyllithium in hexane (428 ml, 659 mmol) at -78°C over 20 minutes, and the mixture was warmed to -10°C over 1 hour and cooled again to -78°C. Thereto was added N-fluorobenzenesulfonylimide (142 g, 449 mmol) in one portion. The acetone-dry ice bath was removed and the mixture was stirred for 30 minutes poured into cold water (1000 ml). Thereto was added ethyl acetate (1200 ml) and the organic layer was washed sequentially with 2N hydrochloric acid, water and brine, dried over anhydrous magnesium sulfate and concentrated. The residue was purified by silica gel column chromatography (15% ethyl acetate-hexane = 3:1), and then recrystallized from diethyl ether to give 2-tert-butyloxycarbonylamino-5-fluorothiazole (45.1 g).

mp. 157 to 159°C

MS (m/z) APCI: 219 (M+H)⁺


To a solution of the above 2-tert-butyloxycarbonylamino-5-fluorothiazole (38.0 g, 174 mmol) in methylene chloride (190 ml) was added dropwise trifluoroacetic acid (190 ml) over 20 minutes under ice-

cooling. The mixture was stirred at room temperature for 2.5 hours, and concentrated. Thereto was added chloroform (20 ml) and 4N hydrogen chloride solution in dioxane (180 ml) dropwise under ice-cooling. After concentration, the residue was washed with ethyl acetate and collected to give the titled compound (24.6 g) as monohydrochloride thereof.

mp. 142 to 144°C (decomposed)

MS (m/z) APCI: 119 (M+H)⁺

REFERENCE EXAMPLE 3




(1) To an aqueous solution (400 ml) of a compound of REFERENCE EXAMPLE 1 (80.0 g, 486 mmol) was added dropwise a 28% ammonia water (40 ml) at room temperature, and then the precipitated crystals were collected and dried to give the free aldehyde.

(2) The above product was suspended in methanol (600 ml) and thereto was added sodium borohydride (27.6 g, 729 mmol) under ice-cooling, and the mixture was stirred at room temperature for 1 hour. To the reaction solution were added acetone (30 ml) and water (50 ml), and then the mixture was concentrated in *vacuo*. The resulting residue was purified by silica gel column chromatography (20 to 25% methanol-chloroform) to give 2-amino-5-hydroxymethylthiazole (45.2 g, yield 71%). MS (m/z) APCI: 131 (M+H)⁺

(3) The above product (45.2 g) was dissolved in a 2M solution of hydrogen chloride in methanol and the mixture was stirred at 60°C for 5 hours. The reaction mixture was concentrated and then the residue was solidified with a mixture of methanol and diethyl ether to give the titled compound (49.8 g, yield 79%) as monohydrochloride thereof.

MS (m/z) APCI: 145 (M+H)⁺


REFERENCE EXAMPLE 4



5-Methoxy[1,3]thiazolo[5,4-b]pyridin-2-amine (5.0 g, 27.6 mmol) was dissolved in 30% hydrogen bromide in acetic acid (50 ml). The mixture was stirred at 130°C for 3 hours and cooled to room temperature, and then the solvents were removed in *vacuo* and the residue was solidified with diethyl ether to give 2-amino[1,3]thiazolo[5,4-b]pyridin-5-ol dihydrobromide (9.11 g, quantitatively) as colorless crystals.

MS (m/z) APCI: 168 (M+H)⁺


REFERENCE EXAMPLE 5



A solution of a compound of REFERENCE EXAMPLE 4 (1.18 g, 3.59 mmol), dimethylaminoethyl chloride monohydrochloride (569 mg, 3.95 mmol) and cesium carbonate (6.43 g, 19.75 mmol) in DMF (30 ml) was stirred at 60°C for 2 hours. After cooling to room temperature, thereto was added acetic acid (2.26ml, 39.5 mmol) and the mixture was diluted with water to a homogeneous solution and then concentrated in *vacuo*. The residue was purified by silica gel column chromatography (NH-silica gel; 50% ethyl acetate-hexane) to give the titled compound (273 mg, yield 32%) as pale yellow crystals.

MS (m/z) APCI: 239 (M+H)⁺


REFERENCE EXAMPLE 6



A solution of a compound of REFERENCE EXAMPLE 4 (1.54 g, 4.68 mmol), ethyl bromoacetate (0.571 ml, 5.15 mmol) and cesium carbonate (6.86g, 21.06 mmol) in DMF (40 ml) was stirred at room temperature for 30 minutes. Thereto was added acetic acid (2.47 ml, 43.2 mmol) and the mixture was diluted with water to a homogeneous solution and the solvents were removed in *vacuo*. The residue was purified by silica gel column chromatography (NH-silica gel; ethyl acetate) and solidified with diisopropyl ether to give the titled compound (862 mg, yield 67%) as colorless crystals.

MS (m/z) APCI: 254 (M+H)⁺


REFERENCE EXAMPLE 7



A corresponding starting compound was treated in the similar manner as REFERENCE EXAMPLE 5 to give the titled compound.

MS (m/z) APCI: 282 (M+H)⁺

REFERENCE EXAMPLE 8




To a solution of a compound of REFERENCE EXAMPLE 6 (862 mg, 3.40 mmol) in THF (20 ml) was added lithium borohydride (222 mg,

10.21 mmol) at room temperature, and the mixture was stirred for 24 hours. To the reaction mixture was added 10% hydrochloric acid for degradation of the excess reagents. Thereto was added a saturated aqueous sodium bicarbonate solution, and the mixture was extracted with 20% methanol-chloroform. The organic layer was separated and the solvents were removed. The resulting residue was purified by silica gel column chromatography (5 to 20% methanol-chloroform) to give the titled compound (369 mg, yield 51%) as colorless crystals.

MS (m/z) APCI: 212 (M+H)⁺

The above compound was also synthesized by the following alternative method.



(1) To a solution of the compound (8-a) (5.18 g, 32.7 mmol) and ethylene glycol (20.28 g, 327 mmol) in DMF (20 ml) was added potassium carbonate (13.55 g, 98.0 mmol). The mixture was stirred at room temperature for 2 hours, diluted with ethyl acetate, washed with water and brine, dried over sodium sulfate and concentrated in *vacuo*. The residue was recrystallized from ethyl acetate-hexane to give the compound (8-b) (5.59 g, yield 93%) as yellow crystals.

MS (APCI): 185 (M+H)⁺

(2) To a solution of the compound (8-b) (5.57 g, 30.25 mmol) in ethanol (50 ml) was added 10% Pd-C (0.50g). The mixture was stirred vigorously for 2 hours under hydrogen and concentrated in *vacuo* to give the crude amine.


(3) To a solution of the above amine in acetic acid (100 ml) was added potassium thiocyanate (17.64 g, 182 mmol), and thereto was added dropwise bromine (1.62 ml, 31.8 mmol) under cooling with ice

bath. The mixture was stirred at room temperature overnight, then concentrated in *vacuo*. The residue was chased with toluene, purified by silica gel chromatography (NH-silica gel; methanol-chloroform = 20:1 to 10:1) and triturated with ethyl acetate to give the compound (8-c)

5 (5.55 g, yield 87%) as colorless crystals.

MS (APCI): 212 (M+H)⁺

REFERENCE EXAMPLE 9




The compound of REFERENCE EXAMPLE 4 was treated with 2-bromoacetamide in the similar manner as REFERENCE EXAMPLE 5 to give the titled compound.

10

MS (m/z) APCI: 225 (M+H)⁺

REFERENCE EXAMPLE 10




The compound of REFERENCE EXAMPLE 4 was treated with 2-bromo-N-methylacetamide in the similar manner as REFERENCE EXAMPLE 5 to give the titled compound.

15

MS (m/z) APCI: 239 (M+H)⁺

REFERENCE EXAMPLE 11



(1) To a solution of the compound of REFERENCE EXAMPLE 9 (1380


20

mg, 3.02 mmol) in THF (40 ml) was added lithium aluminum hydride (1.17 g, 30.8 mmol) at room temperature, and the mixture was stirred at room temperature for 20 hours. After degrading excess reagents with a 30% aqueous ammonia solution, the mixture was filtered and the filtrate was concentrated in *vacuo*. The resulting residue was purified by silica gel column chromatography (30% aqueous ammonia-methanol-chloroform = 0.1:1:5) to give an amine (150 mg).

(2) To a suspension of the compound obtained in the above (1) (150 mg, 0.713 mmol) in THF (10 ml) was added a solution of di-tert-butylidicarbonate (187 mg, 0.856 mmol) in THF (5 ml) at room temperature, and the mixture was stirred at room temperature for 90 minutes. To the reaction mixture was added a 30% aqueous ammonia solution, and the mixture was stirred at room temperature for 30 minutes, and then diluted with ethyl acetate, washed with brine, dried over sodium sulfate and concentrated in *vacuo*. The resulting residue was purified by silica gel column chromatography (0 to 5% methanol-chloroform) to give the titled compound (177 mg, yield 9% in 2 steps) as colorless solids.

MS (m/z) APCI: 311 (M+H)⁺

The above compound was also synthesized by the following alternative method.



(1) To a solution of the compound (8-a) (15.0 g, 94.6 mmol) and N-(tert-butoxycarbonyl)ethanolamine (30.50 g, 189.2 mmol) in DMF (150 ml) was added portionwise potassium carbonate (26.15 g, 189.2 mmol), and the mixture was stirred at 50 to 60°C for 4 hours, cooled to room temperature, diluted with ethyl acetate, washed with water and brine,


and dried over sodium sulfate. After treating with activated charcoal, the mixture was concentrated in *vacuo* and the residue was crystallized from diisopropyl ether to give the compound (11-a) (11.86 g, yield 44%). MS (m/z) APCI: 284 (M+H)⁺

(2) To a solution of the above compound (1.08 g, 3.81 mmol) in ethyl acetate (10 ml) was added 10% Pd-C (128 mg), and the mixture was stirred vigorously at room temperature for 1 hour under hydrogen at atmospheric pressure. The reaction mixture was filtered and the filtrate was concentrated in *vacuo* to give the crude amine (1.09 g).

(3) To a solution of the above amine in acetic acid (20 ml) were added potassium acetate (1.87 g, 19.3 mmol) and potassium thiocyanate (2.22 g, 22.8 mmol), and thereto was added dropwise bromine (0.234 ml, 4.57 mmol) with cooling with an ice-water bath, and then the mixture was stirred at room temperature for 1 hour. The reaction mixture was diluted with ethyl acetate, washed with water, an aqueous sodium sulfite solution and brine, and dried over sodium sulfate. After treating with activated charcoal, the mixture was concentrated in *vacuo* and the residue was crystallized from ethyl acetate-hexane to give the above compound (516 mg, yield 44% through the above 2 steps).

MS (m/z) APCI: 311 (M+H)⁺

REFERENCE EXAMPLE 12




The compound of REFERENCE EXAMPLE 10 was treated in the similar manner as REFERENCE EXAMPLE 11 to give the titled compound as colorless crystals.

MS (m/z) APCI: 325 (M+H)⁺

The above compound was also synthesized by the following

alternative method.



(1) To a solution of the compound (8-a) (10.0 g, 63.08 mmol) and N-(tert-butoxycarbonyl)-N-methylethanolamine (16.55 g, 94.6 mmol) in dimethylsulfoxide (100 ml) was added portionwise potassium tert-butoxide (10.62 g, 94.6 mmol) with cooling with an ice-water bath. The mixture was stirred at room temperature for 50 minutes, diluted with ethyl acetate, washed with water and brine, and dried over sodium sulfate. After treating with activated charcoal, the mixture was concentrated in *vacuo* to give the compound (12-a) (21.76 g).

MS (m/z) APCI: 298 (M+H)⁺


(2) The compound obtained in the above (1) was reacted in the similar manner as the alternative method (2) of REFERENCE EXAMPLE 11 to give the crude amine.

MS (m/z) APCI: 268 (M+H)⁺

(3) The compound obtained in the above (2) was treated in the similar manner as the alternative method (3) of REFERENCE EXAMPLE 11 to give the above compound.

MS (m/z) APCI: 325 (M+H)⁺

REFERENCE EXAMPLE 13



(1) Chlorosulfuric acid (80.0 g, 687 mmol) was ice-cooled, and thereto was added 2-acetamidothiazole (20.00 g, 140.6 mmol) in several portions, and the mixture was stirred at 100°C overnight. The reaction solution was cooled to room temperature, poured into ice water and the


mixture was extracted with diethyl ether. The extract was washed with water and brine, dried over sodium sulfate, and then concentrated in *vacuo* to give the crude product (13-b) (9.41 g) as a yellow solid.

(2) A mixture of a 2M aqueous dimethylamine solution (7.2 ml, 14.3 mmol) and pyridine (3 ml) was ice-cooled, and thereto was added the compound (13-b) (1.50 g) in several portions. The mixture was stirred at the same temperature for 15 minutes and at room temperature overnight, concentrated in *vacuo* and the residue was purified by silica gel chromatography (0 to 10% methanol-chloroform) to give the crude product (13-c) (677 mg) as a yellow powder.

(3) To the above compound (447 mg) was added 6N hydrochloric acid. The mixture was stirred at 110°C for 1 hour, cooled to room temperature and concentrated in *vacuo*. To the residue were added water and ammonia water were to be basic, and the precipitated crystals were collected and dried to give the compound (13-d) (283 mg, yield 9% in 3 steps) as a yellow powder.

MS (APCI): 208 (M+H)⁺


REFERENCE EXAMPLE 14



Methylpiperazine was used in the similar manner as REFERENCE EXAMPLE 13 to give the titled compound (yield 3% in 3 steps).

MS (APCI): 263 (M+H)⁺

REFERENCE EXAMPLE 15



(1) To a solution of 2-tert-butyloxycarbonylaminothiazole-5-carbaldehyde of REFERENCE EXAMPLE 1 (4.37 g, 19.1 mmol) in THF (100 ml) were added diethylphosphonoacetic acid ethyl ester (9.14 ml, 45.9 mmol) and potassium tert-butoxide (5.16 g, 45.9 mmol) at room temperature, and the mixture was stirred at the same temperature overnight, and then heated to reflux for 5 hours. The reaction solution was cooled to room temperature, and thereto was added water, and the mixture was extracted with ethyl acetate. The organic layer was washed with water and brine, dried over sodium sulfate, treated with activated charcoal and then concentrated in *vacuo*, and the residue was crystallized from ethyl acetate-hexane (1:5) to give the compound (15-a) (4.13 g, yield 72%) as colorless crystals.

MS (APCI): 299 (M+H)⁺

(2) A mixture of the above compound (4.10 g, 13.74 mmol), 10% Pd/C (5.0 g), ethyl acetate (50 ml) and acetic acid (50 ml) was stirred vigorously at room temperature for 24 hours under hydrogen. The reaction mixture was filtered and the filtrate was concentrated in *vacuo*, and then chased with toluene several times to give the compound (15-b) (4.05 g, yield 98%) as colorless crystals.

MS (APCI): 301 (M+H)⁺

(3) A solution of the above compound (4.03 g, 13.42 mmol) in a mixed solvent of ethanol (20 ml), THF (40 ml) and water (20 ml) was ice-cooled, and thereto was added dropwise a 2N sodium hydroxide solution (16.1 ml, 32.2 mmol). The mixture was stirred at the same

temperature for 2 hours and then at room temperature for 3 hours, poured into a mixture of aqueous citric acid solution and ethyl acetate. The mixture was extracted with ethyl acetate, and the extract was washed with water and brine, dried over sodium sulfate, and concentrated in *vacuo*. The residue was crystallized from diisopropyl ether to give the compound (15-c) (3.16 g, yield 86%) as colorless crystals.

MS (ESI): 271 (M-H)⁻

(4) A solution of the above compound (589 mg, 2.16 mmol) and 1-hydroxybenzotriazole (584 mg, 4.33 mmol) in DMF (10 ml) was ice-cooled, and thereto was added N-ethyl-N'-(3-dimethylaminopropyl)-carbodiimide monohydrochloride (1.04 g, 5.40 mmol), and the mixture was stirred at the same temperature for 15 minutes and then at room temperature for 1 hour. The reaction mixture was ice-cooled again, and thereto was added a 50% aqueous methylamine solution (5 ml). The mixture was stirred at the same temperature for 10 minutes and at room temperature for 1 hour, poured into a mixture of ethyl acetate and brine, and the mixture was extracted with ethyl acetate. The extract was washed with a saturated aqueous sodium bicarbonate solution and brine, dried over sodium sulfate, and concentrated in *vacuo*. The residue was crystallized from ethyl acetate-hexane (1:10) to give the compound (15-d) (597 mg, yield 92%) as colorless crystals.


MS (APCI): 300 (M+H)⁺

(5) To a solution of the above compound (202 mg, 0.675 mmol) in formic acid (6 ml) was added a 4N solution of hydrogen chloride in dioxane (2 ml, 8 mmol), and the mixture was stirred at room temperature for 8 hours. To the mixture was added diethyl ether, and the mixture was stirred at room temperature. The resulting crystals were collected and washed with diethyl ether. To the crystals were added a 4N solution of hydrogen chloride in dioxane (2 ml, 8 mmol) and

diethyl ether (10 ml), and the mixture was stirred further at room temperature overnight. The crystals were collected and washed with diethyl ether to give the compound (15-e) (175 mg) as colorless crystals in quantitative yield.

5 MS (APCI): 200 (M+H)⁺

REFERENCE EXAMPLE 16




2 HCl salt

To the compound (15-e) of REFERENCE EXAMPLE 15 (173 mg, 0.734 mmol) was added ammonia water and brine, and the mixture was
10 extracted with chloroform several times. The extract was dried over sodium sulfate and concentrated in *vacuo* to give colorless crystals of the free compound (112 mg, yield 76%). A solution of the above compound (112 mg, 0.560 mmol) in THF (3 ml) was ice-cooled, and thereto was added lithium aluminum hydride (65 mg, 1.71 mmol), and
15 the mixture was stirred at the same temperature for 1 hour and at room temperature for 1 hour. To the reaction mixture was added ammonia water, and the mixture was stirred at room temperature overnight. Thereto was added sodium sulfate, and then the mixture was filtered. The filtrate was concentrated in *vacuo* and the resulting residue was
20 dissolved in ethyl acetate, and thereto were added a 4N solution of hydrogen chloride in dioxane (0.5 ml, 1 mmol) and diethyl ether, and the mixture was stirred at room temperature. The precipitate was collected and dried to give the titled compound (98 mg, yield 68%) as a colorless powder.

25 MS (APCI): 186 (M+H)⁺


REFERENCE EXAMPLE 17



A mixture of 2-amino-5-bromopyrazine (2.00 g, 11.5 mmol), cyclopropylboronic acid (1.28 g, 14.9 mmol), palladium acetate (258 mg, 1.15 mmol), tricyclohexylphosphine (644 mg, 2.30 mmol) and tripotassium phosphate (8.50 g, 40.23 mmol) in toluene-water (20:1) (53 ml) was heated to reflux for 18 hours. The mixture was cooled to room temperature, diluted with ethyl acetate, washed with water and brine, dried over magnesium sulfate, and then concentrated in *vacuo*. The resulting residue was purified by silica gel column chromatography (50 to 100% ethyl acetate-hexane) to give the titled compound (329 mg, yield 21%) as a colorless powder.

MS (APCI): 136 (M+H)⁺


REFERENCE EXAMPLE 18



To a solution of the compound of REFERENCE EXAMPLE 8 (176 mg, 0.833 mmol) and imidazole (188 mg, 2.76 mmol) in DMF (4 ml) was added dropwise a solution of tert-butyldimethylchlorosilane (188 mg, 1.25 mmol) in DMF (2 ml) under ice-cooling. The mixture was stirred at room temperature overnight, diluted with ethyl acetate, washed with water and brine, and then dried over sodium sulfate and concentrated in *vacuo*. Then, the residue was purified by silica gel chromatography (40 to 70% ethyl acetate-hexane) to give the titled compound (172 mg, yield 64%) as colorless crystals.

MS (APCI): 326 (M+H)⁺


REFERENCE EXAMPLE 19



To a solution of 2-amino-5-bromothiazole hydrobromide (3.00 g, 11.5 mmol) and 2-dimethylaminoethanethiol monohydrochloride (2.45 g, 17.31 mmol) in water (15 ml)-ethanol (20 ml) was added a 1N aqueous sodium hydroxide solution (23.5 ml, 23.5 mmol) at room temperature, and the mixture was heated to reflux for 2 hours. The reaction solution was cooled to room temperature and concentrated in *vacuo*, and thereto was added a saturated aqueous sodium bicarbonate solution and sodium chloride to saturation, and the mixture was extracted with ethyl acetate several times. The extract was dried over magnesium sulfate, concentrated in *vacuo*, and then the residue was purified by silica gel chromatography (NH-silica gel; methanol-chloroform = 49:1 to 19:1) to give the titled compound (2.10 g, yield 89%) as a brown solid.

MS (APCI): 204 (M+H)⁺


REFERENCE EXAMPLE 20



3-Mercapto-4-methyl-4H-1,2,4-triazole was used in the similar manner as REFERENCE EXAMPLE 19 to give the titled compound.

MS (APCI): 214 (M+H)⁺


REFERENCE EXAMPLE 21



2-Mercaptoethanol was used in the similar manner as REFERENCE EXAMPLE 19 to give the titled compound.

MS (APCI): 177 (M+H)⁺


REFERENCE EXAMPLE 22



3-Mercaptopropanol was used in the similar manner as
REFERENCE EXAMPLE 19 to give the titled compound.

MS (APCI): 191 (M+H)⁺


5 REFERENCE EXAMPLE 23



To a mixture of 2-amino-5-bromothiazole hydrobromide (5.50 g,
21.2 mmol), tert-butyl-N-(2-mercaptoethyl)carbamate (5.25 g, 29.6
mmol) and ethanol (80 ml) was added 1,8-diazabicyclo[5.4.0]-7-
undecene (7.73 g, 50.8 mmol) under ice-cooling, and the mixture was
10 stirred at room temperature for 7 hours. The reaction solution was
concentrated in *vacuo*, and the residue was diluted with ethyl acetate,
and then washed with water and brine, dried over magnesium sulfate
and concentrated in *vacuo*. Then, the residue was purified by silica gel
15 chromatography (ethyl acetate-hexane = 3:1) to give the titled
compound 5.72 g, yield 98%) as a colorless powder.

MS (APCI): 276 (M+H)⁺

REFERENCE EXAMPLE 24



20 (1) To a solution of the compound (24-a) synthesized according to the
known method (B. Koren et al., *Heterocycles*, 1987, 26(3) 689.) (1.00 g,
3.30 mmol) in DMF (40 ml) were added water (10 ml), sodium formate
(4.50 g, 66.2 mmol) and 10% Pd-C (200 mg), and the mixture was
stirred at 85°C for 14 hours. After cooling to room temperature, the
25 reaction mixture was filtered and the filtrate was concentrated in *vacuo*.


To the residue was added water, and the mixture was acidified with an aqueous citric acid solution, and then the precipitated crystals were collected and dried to give the compound (24-b) (720 mg, yield 97%) as a yellow powder.

5 MS (APCI): 225 (M+H)⁺

(2) To the above compound (700 mg, 3.12 mmol) was added a 10% aqueous sodium hydroxide solution (20 ml), and the mixture was stirred at 100°C for 4.5 hours. The reaction solution was cooled to room temperature, and then 10% hydrochloric acid was added to
10 neutralize and the precipitate was filtered off. The filtrate was extracted with ethyl acetate and the extract was concentrated in *vacuo*. The resulting residue was washed with diethyl ether to give the compound (24-c) (228 mg, yield 48%) as a yellow powder.

MS (APCI): 153 (M+H)⁺

15 REFERENCE EXAMPLE 25



(1) A mixture of the compound (25-a) synthesized according to the known method (B. Koren et al., *Heterocycles*, 1987, 26(3) 689.) (482 mg, 2.09 mmol), 2,5-hexanedione (0.490 ml, 4.17 ml), p-toluenesulfonic acid monohydrate (40 mg, 0.21 mmol) and toluene (5 ml) was heated to reflux for 3 hours with removing water by a Dean-Stark apparatus. The reaction mixture was cooled to room temperature, and then diluted with ethyl acetate and a small amount of methanol, washed with a saturated aqueous sodium bicarbonate solution and brine, and then dried over
20 sodium sulfate. After concentration in *vacuo*, the residue was purified by silica gel chromatography (NH-silica gel; ethyl acetate-hexane = 10:1, then ethyl acetate) to give the compound (25-b) (598 mg, yield 93%) as pale brown crystals.

MS (APCI): 309/311 (M+H)⁺


(2) A solution of the above compound (533 mg, 1.72 mmol) in methanol (15 ml)-DMF (15 ml) was ice-cooled, and thereto was added sodium methoxide (464 mg, 8.60 mmol). The mixture was stirred at room temperature for 42 hours, diluted with ethyl acetate, washed with water and brine, and then dried over sodium sulfate. After treatment with activated charcoal, the mixture was concentrated in *vacuo* to give the compound (25-c) (339 mg, yield 76%) as brown crystals.

MS (APCI): 261 (M+H)⁺

(3) A suspension of the above compound (330 mg, 1.27 mmol) in water (30 ml) was ice-cooled, and thereto was added trifluoroacetic acid (30 ml). The mixture was stirred at 60°C for 4 hours, then cooled to room temperature and concentrated in *vacuo*. The residue was chased with toluene, and purified by silica gel chromatography (NH-silica gel; methanol-chloroform = 1:20 to 1:5). The resulting crude crystals were washed with ethyl acetate-hexane (1:3) to give the compound (25-d) (154 mg, yield 67%) as reddish brown crystals.

MS (APCI): 183 (M+H)⁺

REFERENCE EXAMPLE 26




To a solution of potassium thiocyanate (6.10 g, 63 mmol) in acetic acid (25 ml) was added 5-aminopyrimidine (1.00 g, 10.5 mmol), and thereto was added dropwise a solution of bromine (1.08 ml, 21.0 mmol) in acetic acid (3 ml) with cooling by ice bath. The mixture was stirred at room temperature for 3 days and concentrated in *vacuo*. The residue was neutralized with a saturated aqueous sodium bicarbonate solution and then concentrated in *vacuo*. To the residue were added chloroform and THF, and the mixture was dried over magnesium sulfate and

filtered. The filtrate was concentrated in *vacuo* and the resulting residue was purified by silica gel chromatography (methanol-chloroform = 1:20 to 1:10) to give the titled compound (287 mg, yield 18%) as a yellow powder.

5 MS (APCI): 153 (M+H)⁺

REFERENCE EXAMPLE 27



(1) A suspension of the compound (27-a) synthesized according to the known method (T. Takahashi et al., *Chemical & Pharmaceutical Bulletin*, 1958, 6, 334.) (1.00 g, 4.62 mmol) in methanol (20 ml) was cooled to -40°C, and thereto was added a solution of 28% sodium methoxide in methanol (0.305 ml, 4.85 mmol), and the mixture was stirred at the same temperature for 6 hours. Water was added, and the mixture was extracted with ethyl acetate. The organic layer was dried over sodium sulfate and concentrated in *vacuo*. The residue was purified by silica gel chromatography (15 to 35% ethyl acetate-hexane) to give the compound (27-b) (194 mg, yield 20%) as a yellow powder.

(2) To a solution of the above compound (663 mg, 3.12 mmol) in acetic acid (6 ml) was added iron powder (678 mg, 12.2 mmol) at room temperature, and the mixture was stirred at 60°C for 4 hours. After cooling to room temperature, the mixture was diluted with ethyl acetate, and filtered through Celite. The filtrate was concentrated and then the residue was diluted with ethyl acetate, and thereto was added a saturated aqueous sodium bicarbonate solution, and then the insoluble was filtered off. The filtrate was extracted with ethyl acetate, and the organic layer was dried over sodium sulfate and concentrated in *vacuo*. The residue was purified by silica gel chromatography (ethyl acetate) to

give the compound (27-c) (231 mg, yield 41%) as a yellow powder.

MS (APCI): 183 (M+H)⁺

REFERENCE EXAMPLE 28



(1) A mixture of the compound (28-a) (3.05 g, 16.4 mmol), 2,5-hexanedione (3.85 ml, 32.9 mmol), p-toluenesulfonic acid monohydrate (313 mg, 1.64 mmol) in toluene (30 ml) was heated to reflux for 6 hours with removing water by a Dean-Stark apparatus. The reaction mixture was cooled to room temperature, and then diluted with ethyl acetate, washed with a saturated aqueous sodium bicarbonate solution and brine, and dried over sodium sulfate. After concentration in *vacuo*, the residue was purified by silica gel chromatography (ethyl acetate-hexane = 1:10) to give the compound (28-b) (3.84 g, yield 89%) as pale brown crystals.

MS (APCI): 264/266 (M+H)⁺

(2) To a suspension of the above compound (600 mg, 2.27 mmol), bis(dibenzylideneacetone)palladium (131 mg, 0.227 mmol), 2-dicyclohexylphosphono-2'-(N,N'-dimethylamino)biphenyl (89 mg, 0.226 mmol) and sodium tert-butoxide (437 mg, 4.55 mmol) in toluene (10 ml) was added a 2M solution of dimethylamine in THF (6.81 ml, 13.6 mmol) at room temperature under argon, and the mixture was stirred at the same temperature for 5 days. Water was added, and the mixture was extracted with ethyl acetate. The organic layer was washed with water and brine, dried over sodium sulfate and then concentrated in *vacuo*.


The residue was purified by silica gel chromatography (10 to 35% ethyl acetate-hexane) to give the compound (28-c) (419 mg, yield 68%) as brown crystals.

MS (APCI): 273 (M+H)⁺

(3) A suspension of the above compound (410 mg, 1.51 mmol) in water (30 ml) was ice-cooled, and thereto was added trifluoroacetic acid (30 ml). The mixture was stirred at 50°C for 1 hour, cooled to room temperature, and then concentrated in *vacuo*. The residue was
5 extracted with ethyl acetate. The organic layer was washed with a saturated aqueous sodium bicarbonate solution and brine, dried over sodium sulfate, and then treated with activated charcoal. After concentration in *vacuo*, the residue was crystallized from ethyl acetate-hexane (3:1) to give the compound (28-d) (194 mg, yield 66%) as a
10 colorless powder.

MS (APCI): 195 (M+H)⁺


REFERENCE EXAMPLE 29



A mixture of 2-amino-5-bromopyrazine (2.61 g, 15.0 mmol), tert-butylcarbamate (2.11 g, 18.0 mmol), copper (I) iodide (290 mg, 1.50 mmol), N,N'-dimethylethylenediamine (260 mg, 3.00 mmol), potassium carbonate (4.15 g, 30.0 mmol) in dioxane (80 ml) was heated to reflux
15 for 16 hours. The mixture was cooled to room temperature, poured into water and extracted with ethyl acetate. The extract was filtered through Celite and the filtrate was concentrated in *vacuo*, and the resulting
20 residue was purified by silica gel chromatography (30 to 80% ethyl acetate-hexane) to give the titled compound (yield 18%) as a colorless powder.

MS (APCI): 211 (M+H)⁺

REFERENCE EXAMPLE 30




(1) A suspension of the compound (28-b) (1.86 g, 7.05 mmol), tert-butyltrimethyl(tributylstannylmethoxy)silane (6.14 g, 14.1 mmol) and dichlorobis(triphenylphosphine)palladium (495 mg, 0.701 mmol) in dioxane (20 ml) was heated to reflux for 41 hours under argon. The reaction mixture was cooled to room temperature, and thereto was added activated charcoal. The mixture was filtered, and the filtrate was concentrated in *vacuo*. To the resulting residue was dissolved ethyl acetate, washed with a saturated aqueous sodium bicarbonate solution and brine, dried over sodium sulfate and concentrated in *vacuo* to give the compound (30-a) (10.44 g) as a yellow oil.

(2) A mixture of the above compound (10.4 g) in water (60 ml) was ice-cooled, and thereto was added trifluoroacetic acid (60 ml). The mixture was stirred at 50°C for 1 hour, cooled to room temperature, and then concentrated in *vacuo*. To the resulting residue was added a saturated aqueous sodium bicarbonate solution, and the mixture was saturated with sodium chloride and extracted with ethyl acetate. The extract was dried over sodium sulfate and then concentrated in *vacuo*, and the residue was chased with toluene and then crystallized from a mixed solvent of ethyl acetate-hexane to give the compound (30-b) (1.10 g) as a brown powder.

(3) To a solution of the above compound (1.10 g) and imidazole (1.54 g, 22.6 mmol) in DMF (30 ml) was added tert-butyltrimethylchlorosilane (1.70 g, 11.3 mmol) under ice-cooling. The mixture was stirred at room temperature for 1 hour, diluted with ethyl acetate, washed with a saturated aqueous sodium bicarbonate solution and brine, and dried over sodium sulfate. After concentration in *vacuo*, the resulting residue was purified by silica gel column chromatography (NH-silica gel; 60 to 100% ethyl acetate-hexane) to give the compound (30-c) (354 mg, yield 16% in 3 steps) as pale yellow crystals.

MS (APCI): 296 (M+H)⁺


REFERENCE EXAMPLE 31



A mixture of 2-amino-5-iodopyridine (1.00 g, 4.55 mmol), methyl
 thioglycolate (0.482 g, 4.55 mmol), tris(dibenzylideneacetone)-
 5 dipalladium (208 mg, 0.227 mmol), bis(2-diphenylphosphinophenyl)-
 ether (245 mg, 0.455 mmol) and potassium tert-butoxide (561 mg, 5.00
 mmol) in toluene (20 ml) was stirred at 100°C for 3 hours under argon.
 The reaction mixture was cooled to room temperature and filtered
 through Celite, and the filtrate was concentrated in *vacuo*. The residue
 10 was purified by silica gel chromatography (NH-silica gel, 0 to 3%
 methanol-chloroform) to give the titled compound (268 mg, yield 30%)
 as a pale yellow solid.

MS (APCI): 199 (M+H)⁺

REFERENCE EXAMPLE 32




(1) The compound (32-a), (S)-1-acetyl-2-methylpiperazine
 hydrochloride and diisopropylethylamine were treated in the similar
 manner as EXAMPLE 2 to give the compound (32-b).

MS (m/z) APCI: 349 (M+H)⁺

20 (2) The compound obtained in the above (1) was treated in the
 similar manner as EXAMPLE 138-(2) to give the compound (32-c).

MS (m/z) APCI: 249 (M+H)⁺


REFERENCE EXAMPLE 33






(1) A mixture of the compound (28-b) of REFERENCE EXAMPLE 28 (2.00 g, 7.58 mmol) and N,N,N'-trimethylethylenediamine (7.74 g, 75.8 mmol) was stirred at 125°C for 15 hours. The reaction mixture was cooled to room temperature and concentrated in *vacuo*. To the residue was added ethyl acetate and the insoluble materials were filtered off. The filtrate was concentrated in *vacuo* and the residue was purified by silica gel column chromatography (NH-silica gel; 15 to 35% ethyl acetate-hexane) to give the compound (33-a) (2.33 g, yield 93%). MS (m/z) APCI: 330 (M+H)⁺


(2) To a solution of the above compound (530 mg, 1.61 mmol) in methanol (2 ml) was added 10% hydrochloric acid (3.3 ml), and the mixture was stirred at 60°C for 19 hours, cooled to room temperature, neutralized with sodium bicarbonate and concentrated in *vacuo*. To the residue was added chloroform, and the mixture was dried over magnesium sulfate and filtered, and the filtrate was concentrated in *vacuo*. The resulting crude crystals were washed with ethyl acetate-diethyl ether to give the compound (33-b) (201 mg, yield 50%). MS (m/z) APCI: 252 (M+H)⁺

Corresponding starting compounds were treated in the similar manner as the above REFERENCE EXAMPLE 33, REFERENCE EXAMPLE 11-(2) or REFERENCE EXAMPLE 18 or an appropriate combination thereof to give the following compounds.

REFERENCE EXAMPLE No.	No	Structure	MS (m/z)
33	1		338 APCI [M+H] ⁺

33	2		238 APCI [M+H] ⁺
33	3		339 APCI [M+H] ⁺
33	4		325 APCI [M+H] ⁺

REFERENCE EXAMPLE 34



- (1) A mixture of the compound (28-b) (500 mg, 1.90 mmol), tert-butylcarbamate (333 mg, 2.84 mmol), cesium carbonate (1235 mg, 3.79 mmol), palladium (II) acetate (21.3 mg, 0.095 mmol), 4,5-bis(diphenylphosphino)-9,9-dimethylxanthene (87.7 mg, 0.152 mmol) and dioxane (5 ml) was stirred at 100°C for 15 hours under argon. The reaction mixture was cooled to room temperature, and thereto was added water, and the mixture was extracted with ethyl acetate. The extract was concentrated in *vacuo*. The residue was purified by silica

gel chromatography (5 to 15% ethyl acetate-hexane) to give the compound (34-a) (727 mg, quantitatively).

MS (m/z) APCI: 345 (M+H)⁺


(2) A solution of the compound obtained in the above (1) (633 mg, 1.84 mmol) in DMF (4 ml) was ice-cooled, and thereto were added sequentially methyl iodide (0.126 ml, 2.02 mmol) and 60% sodium hydride (110 mg, 2.76 mmol), and the mixture was stirred at the same temperature for 10 minutes, poured into water and extracted with ethyl acetate. The extract was concentrated in *vacuo*, and the residue was purified by silica gel chromatography (NH-silica gel, 0 to 10% ethyl acetate-hexane) to give the compound (34-b) (556 mg, yield 84%).

MS (m/z) APCI: 359 (M+H)⁺

(3) A mixture of the compound obtained in the above (2) (556 mg, 1.55 mmol), 10% hydrochloric acid (7 ml) and methanol (6 ml) was stirred at 75°C for 23 hours. The reaction mixture was cooled to room temperature, and thereto was added a saturated aqueous sodium bicarbonate solution and the mixture was extracted with chloroform. The extract was concentrated in *vacuo* to give the compound (34-c) (300 mg, quantitatively).

MS (m/z) APCI: 181 (M+H)⁺

REFERENCE EXAMPLE 35









The compound (34-a) of REFERENCE EXAMPLE 34 was treated in the similar manner as the method of REFERENCE EXAMPLE 34-(3) to give the titled compound.

MS (m/z) APCI: 167 (M+H)⁺


REFERENCE EXAMPLE 36

Corresponding starting compounds were reacted in the similar manner as the alternative method of REFERENCE EXAMPLE 11 or REFERENCE EXAMPLE 12 to give the following compounds.

REFERENCE EXAMPLE No.	No	Structure	MS (m/z)
36	1		351 APCI [M+H] ⁺
36	2		337 APCI [M+H] ⁺
36	3		337 APCI [M+H] ⁺
36	4		323 APCI [M+H] ⁺
36	5		339 APCI [M+H] ⁺

36	6		339 APCI [M+H] ⁺
----	---	--	--------------------------------


REFERENCE EXAMPLE 37



To a suspension of 2-amino-4-chloromethylthiazole hydrochloride (2.00 g, 10.8 mmol) in dioxane (15 ml) was added N-methylpiperazine (12 ml, 108 mmol), and the mixture was stirred at room temperature for 110 hours. To the reaction mixture was added a saturated aqueous sodium bicarbonate solution (5 ml) and the organic layer was separated by Chem Elut 1010® (manufactured by VARIAN) and the column was further eluted with chloroform. The eluate was concentrated in *vacuo* and the residue was purified by silica gel column chromatography (NH-silica gel; 2% methanol-chloroform) to give the above compound (1.18 g, yield 52%).

MS (m/z) APCI: 213 (M+H)⁺


REFERENCE EXAMPLE 38



A corresponding starting compound was treated in the similar manner as REFERENCE EXAMPLE 38 to give the above compound.

MS (m/z) APCI: 227 (M+H)⁺


REFERENCE EXAMPLE 39



To a solution of the compound of REFERENCE EXAMPLE 1 (10.00 g, 60.7 mmol), (S)-2-methyl-1-acetylpiperazine hydrochloride (15.0 g, 85.0 mmol) and diisopropylethylamine (26.0 ml, 149 mmol) in chloroform (300 ml) was added sodium triacetoxyborohydride (18.0 g, 85.0 mmol), and the mixture was stirred at room temperature for 17 hours. To the reaction mixture was added a saturated aqueous sodium bicarbonate solution, and the mixture was extracted with chloroform. The extract was dried over magnesium sulfate. After concentration in *vacuo* the residue was purified by silica gel column chromatography (6% methanol-chloroform) and crystallized from ethyl acetate-diethyl ether to give the above compound (10.99 g, yield 71%).

MS (m/z) APCI: 255 (M+H)⁺


REFERENCE EXAMPLE 40



A corresponding starting compound was treated in the similar manner as EXAMPLE 39 to give the above compound.

MS (m/z) APCI: 213 (M+H)⁺

REFERENCE EXAMPLE 41




A solution of the compound of REFERENCE EXAMPLE 29 (1.61 g, 7.70 mmol) in DMF (16 ml) was ice-cooled, and thereto was added potassium tert-butoxide (945 mg, 8.42 mmol). The mixture was stirred at room temperature for 10 minutes, ice-cooled again and thereto was added methyl iodide (572 ml, 9.19 mmol). The mixture was stirred at

room temperature for 6 hours. To the reaction mixture was added water, and the mixture was extracted with chloroform. The extract was washed sequentially with water and brine, dried over sodium sulfate and then concentrated in *vacuo*. The residue was purified by column chromatography (NH-silica gel, 10 to 35% ethyl acetate-hexane) to give the above compound (1.51 g, yield 88%).

MS (m/z) APCI: 225 (M+H)⁺


REFERENCE EXAMPLE 42



A mixture of 2-amino-5-bromopyrazine (5.00 g, 28.74 mmol), 2-pyrrolidinone (10.90 ml, 143.7 mmol), cuprous iodide (1.10 g, 5.75 mmol), (1*R*,2*R*)-(-)-1,2-diaminocyclohexane (1.38 ml, 11.50 mmol), potassium carbonate (7.94 g, 57.5 mmol) and dioxane (86 ml) was stirred at 120°C for 17 hours under argon. The reaction mixture was cooled to room temperature and thereto was added ethyl acetate-methanol (10:1), and the mixture was filtered through Celite. The filtrate was concentrated in *vacuo*, and the residue was purified by silica gel column chromatography (5% methanol-chloroform) and crystallized from diethyl ether to give the above compound (2.77 g, yield 54%).

MS (m/z) APCI: 179 (M+H)⁺


REFERENCE EXAMPLE 43



1-tert-Butoxycarbonyl-4-acetylthiopiperidine (4.60 g, 17.69 mmol) and 2-amino-5-bromothiazole hydrobromide were reacted in the similar manner as REFERENCE EXAMPLE 19 to give the above compound.

MS (m/z) APCI: 316 (M+H)⁺

REFERENCE EXAMPLE 44




To 2-methoxyethanol (55 ml) were added sequentially 60% sodium hydride (1.83 g, 45.8 mmol), 2-amino-5-bromopyrazine (7.00 g, 40.23 mmol) and copper powder (2.91 g, 53.9 mmol) under ice-cooling, and the mixture was stirred at 160°C for 20 hours in a sealed tube. The reaction mixture was cooled to room temperature, and thereto were added water, ammonia water and ethyl acetate, and the mixture was stirred and then filtered through Celite. The filtrate was extracted with ethyl acetate. The extract was dried over sodium sulfate, and after concentration in *vacuo* the residue was purified by silica gel column chromatography (40% ethyl acetate-hexane) to give the titled compound (2.74 g, yield 40%).

MS (m/z) APCI: 170 (M+H)⁺

Corresponding starting compounds were treated in the similar manner as the above REFERENCE EXAMPLE 44, REFERENCE EXAMPLE 11-(2) or REFERENCE EXAMPLE 18 or an appropriate combination thereof to give the following compounds.

REFERENCE EXAMPLE No.	No	Structure	MS (m/z)
44	1		270 APCI [M+H] ⁺
44	2		154 APCI [M+H] ⁺

REFERENCE EXAMPLE 45



(1) A solution of (S)-1-acetyl-2-methylpiperazine hydrochloride (0.800 g, 4.50 mmol) and diisopropylethylamine (1.81 ml, 10.4 mmol) in DMF (10 ml) was ice-cooled, and thereto was added the compound (45-a). The mixture was stirred at the same temperature for 3 hours, diluted with water, extracted with ethyl acetate and the extract was concentrated in *vacuo*. The residue was purified by silica gel column chromatography (ethyl acetate) to give the compound (45-b) (0.930 g, yield 76%).


MS (m/z) APCI: 350 (M+H)⁺


(2) The compound obtained in the above (1) was reacted in the similar manner as EXAMPLE 138-(2) to give the compound (45-c).

MS (m/z) APCI: 250 (M+H)⁺


REFERENCE EXAMPLE 46

Corresponding starting compounds were treated with 2-amino-6-hydroxybenzothiazole in the similar manner as REFERENCE EXAMPLE 5 to give the following compounds.

REFERENCE EXAMPLE No.	No	Structure	MS (m/z)
46	1		325 APCI [M+H] ⁺

46	2		324 APCI [M+H] ⁺
----	---	--	--------------------------------

REFERENCE EXAMPLE 47



(1) A solution of the compound (47-a) (1.50 g, 5.63 mmol) and N,N-dimethylethanolamine (535 mg, 6.00 mmol) in THF (20 ml) was ice-cooled, and thereto were added triphenylphosphine (1.57 g, 6.00 mmol) and diethyl azodicarboxylate (1.04 g, 6.00 mmol), and the mixture was stirred at the same temperature overnight. To the reaction mixture were added ethyl acetate and diluted hydrochloric acid. The aqueous layer was separated and basified with an aqueous sodium hydroxide solution. After extraction with ethyl acetate, the extract was dried over sodium sulfate and concentrated in *vacuo*, and the residue was purified by silica gel chromatography (40 to 60% acetone-chloroform) and then washed with diisopropyl ether to give the compound (47-b) (606 mg, yield 32%) as a colorless powder.

(2) To a solution of the compound (47-b) obtained in the above (1) (337 mg, 1.0 mmol) in methanol (5 ml) was added a 4 M hydrogen chloride solution in dioxane (2.5 ml) at room temperature, and the mixture was stirred at the same temperature overnight. The reaction mixture was concentrated in *vacuo* and the residue was washed with diethyl ether to give the compound (47-c) (302 mg, yield 97%) as a colorless powder.

MS (m/z) APCI: 238 (M+H)⁺


INDUSTRIAL APPLICABILITY

5 The compound of the present invention or a pharmaceutically acceptable salt thereof can provide an agent for preventing or treating diseases involving glucokinase because of having an excellent glucokinase activation effect.

Also, the method for preparing 5-substituted 2-aminothiazole compound of the present invention and a salt thereof is industrially advantageous.

CLAIMS


1. An oxime derivative of the general formula [I]:



wherein Ring A is aryl or heteroaryl;

Q is cycloalkyl, nonaromatic heterocycle, alkyl or alkenyl;

Ring T is heteroaryl or nonaromatic heterocycle of



- 10 R^1 is hydrogen atom;

R^2 is cycloalkylsulfonyl, substituted or unsubstituted aminosulfonyl, substituted or unsubstituted nonaromatic heterocyclylsulfonyl, substituted or unsubstituted heteroarylsulfonyl;

- 15 R^3 and R^4 are independently hydrogen atom, alkoxy, substituted or unsubstituted nonaromatic heterocycle, substituted or unsubstituted heteroaryl, alkoxyalkoxy, substituted or unsubstituted cycloalkyl, cyano, substituted or unsubstituted aryl, substituted or unsubstituted carbamoyl, hydroxy, alkanoyl, alkylthio, alkoxycarbonyl, substituted or unsubstituted aryloxy, halogen atom, oxo, or substituted or
- 20 unsubstituted arylcarbonyloxy;

R^5 is hydrogen atom, formyl, halogen atom, oxo, substituted or unsubstituted alkoxy, substituted or unsubstituted aminosulfonyl,


substituted or unsubstituted alkylthio, cyano, substituted or
 unsubstituted nonaromatic heterocyclyl-sulfonyl, nitro, substituted or
 unsubstituted cycloalkyl, alkoxycarbonyl, alkenyl, alkylsulfonyl,
 substituted or unsubstituted carbamoyl, substituted or unsubstituted
 5 heteroarylthio, substituted or unsubstituted amino, carboxyl,
 substituted or unsubstituted heteroaryl, substituted or unsubstituted
 alkynyl, substituted or unsubstituted nonaromatic heterocyclyl-
 carbonyl, substituted or unsubstituted nonaromatic heterocyclyl-oxy,
 substituted or unsubstituted nonaromatic heterocycle, substituted or
 10 unsubstituted nonaromatic heterocyclyl-thio, substituted or
 unsubstituted cycloalkyloxy, alkanoyl, or substituted or unsubstituted
 alkyl;

R^6 is hydrogen atom, substituted or unsubstituted alkyl, halogen
 atom, or carboxyl;

provided that Ring A is not thiazolyl or thiadiazolyl;
 or a pharmaceutically acceptable salt thereof.

2. The oxime derivative of claim 1, wherein R^2 is
 cycloalkylsulfonyl, or a pharmaceutically acceptable salt thereof.


3. An oxime derivative of the general formula [I]:



wherein Ring A is aryl or heteroaryl;

Q is cycloalkyl, nonaromatic heterocycle, alkyl or alkenyl;

Ring T is heteroaryl or nonaromatic heterocycle of



R¹ is hydrogen atom, halogen atom, cycloalkylsulfonyl, alkylsulfonyl, alkylsulfinyl, alkylthio, or substituted or unsubstituted tetrazolyl;

5 R² is hydrogen atom, halogen atom, cycloalkylsulfonyl, substituted or unsubstituted alkylsulfonyl, substituted or unsubstituted alkylthio, nitro, substituted or unsubstituted amino, substituted or unsubstituted aminosulfonyl, substituted or unsubstituted nonaromatic heterocyclyl-thio, substituted or
10 unsubstituted nonaromatic heterocyclyl-sulfinyl, substituted or unsubstituted nonaromatic heterocyclyl-sulfonyl, substituted or unsubstituted heteroarylsulfonyl, alkenyloxy, substituted or unsubstituted alkoxy, substituted or unsubstituted alkylsulfinyl, substituted or unsubstituted heteroaryl, substituted or unsubstituted
15 heteroarylthio;

 R³ and R⁴ are independently hydrogen atom, alkoxy, substituted or unsubstituted nonaromatic heterocycle, substituted or unsubstituted heteroaryl, alkoxyalkoxy, substituted or unsubstituted cycloalkyl, cyano, substituted or unsubstituted aryl, substituted or unsubstituted
20 carbamoyl, hydroxy, alkanoyl, alkylthio, alkoxycarbonyl, substituted or unsubstituted aryloxy, halogen atom, oxo, or substituted or unsubstituted arylcarbonyloxy;

 R⁵ is hydrogen atom, formyl, halogen atom, oxo, substituted or unsubstituted alkoxy, substituted or unsubstituted aminosulfonyl, substituted or unsubstituted alkylthio, cyano, substituted or
25 unsubstituted nonaromatic heterocyclyl-sulfonyl, nitro, substituted or unsubstituted cycloalkyl, alkoxycarbonyl, alkenyl, alkylsulfonyl, substituted or unsubstituted carbamoyl, substituted or unsubstituted

heteroarylthio, substituted or unsubstituted amino, carboxyl, substituted or unsubstituted heteroaryl, substituted or unsubstituted alkynyl, substituted or unsubstituted nonaromatic heterocyclyl-carbonyl, substituted or unsubstituted nonaromatic heterocyclyl-oxy, substituted or unsubstituted nonaromatic heterocycle, substituted or unsubstituted nonaromatic heterocyclyl-thio, substituted or unsubstituted cycloalkyloxy, alkanoyl, or substituted or unsubstituted alkyl;

R^6 is hydrogen atom, substituted or unsubstituted alkyl, halogen atom, or carboxyl;

provided that when Q is cycloalkyl, alkyl or alkenyl, then one of R^3 and R^4 is selected from substituted or unsubstituted nonaromatic heterocycle, substituted or unsubstituted heteroaryl, alkoxyalkoxy, substituted or unsubstituted cycloalkyl, substituted or unsubstituted carbamoyl, alkanoyl, substituted or unsubstituted aryloxy, oxo, or substituted or unsubstituted arylcarbonyloxy, and the other is selected from hydrogen atom, alkoxy, substituted or unsubstituted nonaromatic heterocycle, substituted or unsubstituted heteroaryl, alkoxyalkoxy, substituted or unsubstituted cycloalkyl, cyano, substituted or unsubstituted aryl, substituted or unsubstituted carbamoyl, hydroxy, alkanoyl, alkylthio, alkoxycarbonyl, substituted or unsubstituted aryloxy, halogen atom, oxo, or substituted or unsubstituted arylcarbonyloxy;

or a pharmaceutically acceptable salt thereof.

4. The oxime derivative of claim 3, wherein R^1 is hydrogen atom or halogen atom, or a pharmaceutically acceptable salt thereof.

5. The oxime derivative of claim 3, wherein R^1 is hydrogen atom, or a pharmaceutically acceptable salt thereof.

6. The oxime derivative of any one of claims 3 to 5, wherein R^2 is cycloalkylsulfonyl, substituted or unsubstituted alkylsulfonyl,

substituted or unsubstituted alkylthio, nitro, substituted or unsubstituted amino, substituted or unsubstituted aminosulfonyl, substituted or unsubstituted nonaromatic heterocyclyl-thio, substituted or unsubstituted nonaromatic heterocyclyl-sulfonyl, substituted or unsubstituted alkylsulfinyl, substituted or unsubstituted heteroarylsulfonyl, substituted or unsubstituted heteroarylthio, or substituted or unsubstituted heteroaryl, or a pharmaceutically acceptable salt thereof.

7. The oxime derivative of any one of claims 3 to 5, wherein R^2 is cycloalkylsulfonyl, substituted or unsubstituted alkylsulfonyl, substituted or unsubstituted aminosulfonyl, or substituted or unsubstituted nonaromatic heterocyclyl-sulfonyl, or a pharmaceutically acceptable salt thereof.

8. The oxime derivative of any one of claims 3 to 5, wherein R^2 is cycloalkylsulfonyl, substituted or unsubstituted alkylsulfonyl, or substituted or unsubstituted aminosulfonyl, or a pharmaceutically acceptable salt thereof.

9. The oxime derivative of any one of claims 3 to 5, wherein R^2 is cycloalkylsulfonyl, or a pharmaceutically acceptable salt thereof.

10. The oxime derivative of any one of claims 3 to 8, wherein the substituent of "substituted aminosulfonyl" in R^2 is substituted or unsubstituted alkyl, cycloalkyl, substituted or unsubstituted nonaromatic heterocycle, or alkoxy, or a pharmaceutically acceptable salt thereof.

11. The oxime derivative of any one of claims 3 to 8, wherein the substituent of "substituted alkylsulfonyl" in R^2 is alkoxy, or a pharmaceutically acceptable salt thereof.

12. The oxime derivative of any one of claims 3 to 11, wherein Ring A is aryl, or a pharmaceutically acceptable salt thereof.

13. The oxime derivative of any one of claims 3 to 11, wherein

Ring A is phenyl or pyridyl, or a pharmaceutically acceptable salt thereof.


14. The oxime derivative of any one of claims 3 to 13, wherein Q is cycloalkyl, nonaromatic heterocycle or alkyl, R³ and R⁴ are independently hydrogen atom, alkoxy, substituted or unsubstituted nonaromatic heterocycle, substituted or unsubstituted heteroaryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted aryl, substituted or unsubstituted carbamoyl, hydroxy, alkanoyl, alkylthio, substituted or unsubstituted aryloxy, halogen atom, oxo, or substituted or unsubstituted arylcarbonyloxy, provided that when Q is cycloalkyl or alkyl, then R³ and R⁴, each independently, are not hydrogen, alkoxy, substituted or unsubstituted aryl, hydroxy, alkylthio or halogen atom, or a pharmaceutically acceptable salt thereof.

15. The oxime derivative of any one of claims 3 to 13, wherein Q is cycloalkyl or nonaromatic heterocycle, R³ and R⁴ are independently hydrogen atom, alkoxy, substituted or unsubstituted heteroaryl, substituted or unsubstituted cycloalkyl, or hydroxy, provided that when Q is cycloalkyl, then R³ and R⁴, each independently, are not hydrogen, alkoxy, hydroxy, or a pharmaceutically acceptable salt thereof.

16. The oxime derivative of any one of claims 3 to 13, wherein the group of -Q(R³)(R⁴) is nonaromatic heterocycle, or alkyl substituted with 1 to 2 groups of substituted or unsubstituted heteroaryl, or a pharmaceutically acceptable salt thereof.

17. The oxime derivative of any one of claims 3 to 13, wherein Q is nonaromatic heterocycle and both of R³ and R⁴ are hydrogen atom, or a pharmaceutically acceptable salt thereof.

18. The oxime derivative of any one of claims 3 to 17, wherein Ring T is heteroaryl of



or a pharmaceutically acceptable salt thereof.

19. The oxime derivative of any one of claims 3 to 17, wherein Ring T is thiazolyl, thiazolopyridinyl, pyridyl, pyrazinyl, benzothiazolyl, quinolyl, thiadiazolyl, pyrazolyl, thiazolopyrazinyl, thiazolopyrimidinyl, cyclohexanothiazolyl or dihydrothiazolopyridinyl, or a pharmaceutically acceptable salt thereof.

20. The oxime derivative of any one of claims 3 to 17, wherein Ring T is thiazolyl, thiazolopyridinyl, pyridyl, pyrazinyl, benzothiazolyl, thiadiazolyl, thiazolopyrazinyl, thiazolopyrimidinyl, cyclohexanothiazolyl or dihydrothiazolopyridinyl, or a pharmaceutically acceptable salt thereof.

21. The oxime derivative of any one of claims 3 to 17, wherein Ring T is thiazolyl, thiazolopyridinyl, pyrazinyl, thiadiazolyl, thiazolopyrazinyl, thiazolopyrimidinyl, or a pharmaceutically acceptable salt thereof.

22. The oxime derivative of any one of claims 3 to 17, wherein Ring T is thiazolyl or thiazolopyridinyl, or a pharmaceutically acceptable salt thereof.

23. The oxime derivative of any one of claims 3 to 22, wherein R⁵ is hydrogen atom, formyl, halogen atom, oxo, substituted or unsubstituted alkoxy, substituted or unsubstituted aminosulfonyl, substituted or unsubstituted alkylthio, cyano, substituted or unsubstituted nonaromatic heterocyclyl-sulfonyl, nitro, substituted or unsubstituted cycloalkyl, alkoxycarbonyl, alkenyl, alkanoyl, substituted or unsubstituted carbamoyl, substituted or unsubstituted heteroarylthio, substituted or unsubstituted amino, substituted or unsubstituted heteroaryl, substituted or unsubstituted alkynyl,

substituted or unsubstituted nonaromatic heterocycl-
oxy, or substituted or unsubstituted alkyl, or a pharmaceutically acceptable salt thereof.

24. The oxime derivative of any one of claims 3 to 22, wherein
5 R⁵ is hydrogen atom, halogen atom, substituted or unsubstituted alkoxy, substituted or unsubstituted alkylthio, cyano, substituted or unsubstituted cycloalkyl, alkanoyl, substituted or unsubstituted carbamoyl, substituted or unsubstituted amino, substituted or unsubstituted nonaromatic heterocycl-
oxy, or substituted or
10 unsubstituted alkyl, or a pharmaceutically acceptable salt thereof.

25. The oxime derivative of any one of claims 3 to 22, wherein
R⁵ is halogen atom, substituted or unsubstituted alkoxy, substituted or unsubstituted alkylthio, substituted or unsubstituted amino, substituted or unsubstituted nonaromatic heterocycl-
oxy, or
15 substituted or unsubstituted alkyl, or a pharmaceutically acceptable salt thereof.

26. The oxime derivative of any one of claims 3 to 22, wherein
R⁵ is substituted or unsubstituted alkoxy, substituted or unsubstituted amino, substituted or unsubstituted nonaromatic heterocycl-
oxy, or
20 substituted or unsubstituted alkyl, or a pharmaceutically acceptable salt thereof.


27. The oxime derivative of any one of claims 3 to 22, wherein
the substituent of "substituted alkyl" in R⁵ is substituted or unsubstituted nonaromatic heterocycle, substituted or unsubstituted amino, substituted or unsubstituted alkoxy, substituted or
25 unsubstituted carbamoyl, hydroxy, trialkylsilyloxy, alkylthio, alkylsulfonyl, substituted or unsubstituted nonaromatic heterocycl-
oxy, heteroaryl, substituted or unsubstituted hydroxyimino, halogen atom, carboxyl, alkoxycarbonyl, or alkanoyloxy, or a pharmaceutically
30 acceptable salt thereof.

28. The oxime derivative of any one of claims 3 to 27, wherein R^6 is hydrogen atom, or substituted or unsubstituted alkyl, or a pharmaceutically acceptable salt thereof.


29. The oxime derivative of any one of claims 3 to 27, wherein R^6 is hydrogen atom, or a pharmaceutically acceptable salt thereof.

30. The oxime derivative of claim 3, wherein Ring A is phenyl, Q is 3-tetrahydrofuryl, Ring T is 2-thiazolyl, R^1 , R^3 , R^4 , and R^6 are hydrogen atom, R^2 is cyclopropylsulfonyl, and R^5 is piperazinyl-substituted alkyl being optionally substituted with 1 to 3 substituents selected from alkyl, oxo, alkanoyl and alkoxyalkanoyl.


31. A method for preparing an oxime derivative of either claim 1 or 3 of the general formula [I]:



wherein the symbols have the same meanings as defined in either claim 1 or 3, which comprises reacting a compound of the general formula [II]:




wherein Z^2 is hydrogen atom or alkyl and the other symbols have the same meanings as defined in either claim 1 or 3, with a compound of the general formula [III]:




wherein the symbols have the same meanings as defined in either claim 1 or 3.


32. A method for preparing an oxime derivative of either claim 1
5 or 3 of the general formula [I]:



wherein the symbols have the same meanings as defined in claim 31,
which comprises reacting a compound of the general formula [IX]:




- 10 wherein the symbols have the same meanings as defined in claim 31,
with a compound of the general formula [V]:




wherein Z^3 is hydroxy, halogen atom, arylsulfonyloxy or alkylsulfonyloxy
and the other symbols have the same meanings as defined in claim 31.


- 15 33. A method for preparing an oxime derivative of either claim 1
or 3 of the general formula [I]:



wherein the symbols have the same meanings as defined in claim 31,
which comprises reacting a compound of the general formula [X]:



- 5 wherein the symbols have the same meanings as defined in claim 31,
with a compound of the general formula [XI]:



wherein the symbols have the same meanings as defined in claim 31.