

## INCORPORATION OF EXOGENOUS LACTIC BACTERIA INTO THE ORAL MICROFLORA

5 The present invention relates to the incorporation of exogenous lactic bacteria into the oral microflora intended for the prophylaxis or the treatment of dental caries, dental plaque and periodontal infection.

### Background of the invention

10

The mouth (oral cavity) contains a resident and a non-resident microflora. The first includes microorganisms that are able to establish a more or less permanent residence on the oral surfaces. These bacteria are mainly localised on the tongue, the buccal mucosa and the teeth while the gingiva, lips, cheeks, palate and floor of the mouth only support a very sparse microflora.

20 On the tongue and the buccal mucosa, the natural resident microflora includes microorganisms selected from Streptococcus, Veillonella, Bacteroides and Haemophilus. On the teeth, Streptococci, Lactobacilli and Actinomyces predominate but a variety of Gram positive and negative cocci and rods can be found.

For example, Frandsen *et al.* showed that *S. sanguis* predominates on the buccal mucosa but its primary habitat is the surface of teeth, that *S. gordonii* grows in the mature supragingival plaque, that *S. oralis* and *S. mitis* grow in the initial dental plaque (Oral Microbiol. Immunol., 6, 129-133, 1991). Strains belonging to the mutans group are localised on teeth (*S. cricetus*, *S. downei*, *S. ferus*, *S. macacae*, *S. mutans*, *S. rattus*, *S. sobrinus*). Strains belonging to the *S. milleri* group predominate in dental abscesses (*S. anginosus*, *S. constellatus*, *S. intermedius*; Bentley *et al.*, Int. J. System. Bacter. 1991, 41, 487-494; Wood *et al.*, The Genera of Lactic Acid Bacteria, Blackie Academic and Professional, Chapman & Hall, W. H. eds., 1995).

35 Many of these microorganisms are innocuous commensal, but a lot of them have been recognised as the etiologic agent of quite a few diseases (Hill, M. J. and Marsh, P. D. eds. Human Microbial Ecology, 1990, CRC Press, Boca Raton Florida, USA)

The dental plaque is a film that forms on the surface of teeth consisting of bacterial cells in a matrix of extracellular polysaccharides and salivary products. Immediately after eruption, the teeth are covered with an amorphous layer of saliva, the acquired enamel pellicle (AEP) that is about 1.3  $\mu\text{m}$  thick and cannot be removed by normal tooth brushing. The deposition of bacteria on teeth follows immediately the formation of the AEP and plaque becomes evident in 8-12 hours as a multi-layered structure. The first layer consists of bacteria (earliest colonisers) that attach to teeth mainly via specific adhesin-receptor recognition; it forms a substratum for the second colonisers that adhere one to the other via analogous specific binding or via simple juxtaposition. Plaque cohesion is essentially guaranteed by three mechanisms: the presence of a salivary pellicle on the outer bacteria layer, the specific coaggregation among the different bacterial species and the glucans synthesised by the bacteria and that remain entrapped in the plaque matrix (Skopek *et al.*, Oral Microbiol. Immunol., 2, 19-24, 1994; Kolenbrander *et al.*, Meth. Enzymol., 253, 385-397, 1995; Hiroi *et al.*, FEMS Microbiol Lett., 96, 193-198, 1992; Gibbons *et al.*, Infect. Immun., 52, 555-561, 1986).

The organic acids produced by oral bacteria during the fermentation process directly cause dental caries. These acids attack the hard tissue of teeth with consequent release of ions such as calcium, phosphate, carbonate, magnesium, fluoride, sodium. When the pH in the oral cavity increases again around the neutrality, saliva becomes saturated with calcium and therefore its liberation from the tooth is prevented.

Among all the food residues found in the mouth, carbohydrates show the highest caries promoting effect being directly available for oral bacteria fermentation.

Potentially all micro-organisms fermenting sugars are cariogenic, but the primary etiological agents of coronal and root caries are the mutans streptococci because they are strong acid producers; lactobacilli, that are highly aciduric, can also be implicated. In humans, *S. mutans* and *S. sobrinus* are the more cariogenic strains, and live on teeth while not colonising the entire dentition. A decrease in their number from molar to anterior teeth was indeed demonstrated (Lindquist *et al.*, Dent. Res., 69, 1160-1166, 1990). Moreover in human approximal plaque, *S. mutans* and *S. sobrinus* preferentially colonise the most caries-prone site apical to the contact area (Ahmady *et al.*, Caries Res., 27, 135-139, 1993). A higher

prevalence of *S. sobrinus* was also found in the molar regions compared with that of *S. mutans* (Lindquist *et al.*, Caries Res., 25, 146-152, 1991).

*S. mutans* and *S. sobrinus* have been shown to attach to the pellicle of teeth  
5 mainly via specific adhesin-receptor. Gibbons *et al.* showed that *S. mutans* carries an adhesin which binds to salivary components in the pellicle, while *S. sobrinus* cells appear to possess an adhesin which binds to glucan in the pellicle (Infect. Immun., 52, 555-561, 1986).

10 The transient microflora comprises exogenous bacteria that can be occasionally present in the mouth, but that do not establish a permanent residence (even if repeated oral administrations of these bacteria are carried out). All the food bacteria, and in particular lactic acid bacteria, can be part of this transient microflora. These exogenous lactic bacteria have never been shown to be capable  
15 of directly adhering to the pellicle of teeth. Repeated administration of exogenous lactic bacteria may however lead to colonisation of the mouth on all the oral surfaces, such as the tongue, the buccal mucosa, the gingiva, lips, cheeks, palate, floor and the teeth. This colonisation may result from attachments via specific bindings to bacteria of the resident microflora (co-aggregation  
20 phenomena), or via entrapment in the matrix of polysaccharides produced by the resident bacteria, or via adhesion to saliva proteins (especially glycoproteins).

*Lactobacillus casei rhamnosus* GG (ATCC53103) has been reported to colonise the mouth, most probably on the epithelium of the buccal mucosa. This  
25 strain also adheres to the epithelium of the intestinal tract (US5032399, Gorbach *et al.*; Micr. Ecol. In Health and Dis., 7, 295-298, 1994). By contrast *L. rhamnosus* does not adhere to teeth.

Japanese patent n°4021633 (Cyconmedix KK) also reported colonisation of  
30 the mouth by *Lactobacillus acidophilus*, most probably on the epithelium of the buccal mucosa. Many *Lactobacillus acidophilus* are known to adhere also to the epithelium of the intestinal tract (EP577904; EP199535; Perdigon *et al.*, Medicina, 46, 751-754, 1986; Perdigon *et al.*, Immunology, 63, 17-23, 1988).

35 Exogenous bacteria can also produce factors that inhibit the growth of the resident microflora in the mouth. For example, EP759469 (Société des Produits Nestlé) described the use of a bacteriocin produced by *Micrococcus varians* for

inhibiting the development of the oral pathogens *S. sobrinus*, *S. sanguis*, *S. mutans* and *A. viscosus*.

Some strategies have been used to minimise the development of the resident  
5 microflora of the mouth, namely an administration of commensal bacteria of the  
resident microflora that are not cariogenic, such as *Streptococcus salivarius*  
and/or *Stomatococcus mucilaginosus*, and/or repeated administrations of  
exogenous lactic bacteria such as *L. casei*, *L. fermentum*, *L. acidophilus*, *L.*  
*crispatus*, *L. gasseri*, *L. salivarius*, *L. bulgaricus* and *S. salivarius* (Tanzer *et al.*,  
10 *Infec. and Immunity*, 48, 44-50, 1985; WO92/14475).

The application of bacteriocins is also one of the investigated strategies  
which have been set up to reduce tooth caries. These molecules have attracted  
interest as prospective anticaries agents and as factors important in modulating  
15 colonisation of the oral cavity. The anti-caries potential of applications of some  
bacteriocins comes from their potent and broad antibacterial activity against  
mutans streptococci and bacteria associated with dental plaque, and their natural  
occurrence in bacteria regarded as human safe (US5368845 of Colgate, and  
WO94/12150 of Smithkline Beecham).

The application of milk derivatives is also of interest for the health of the  
mouth. Indeed, US5427769 (Nestec S.A.) describes another alternative where  
dental caries are prevented by contacting teeth with an edible composition  
containing micellar casein in amount sufficient to inhibit colonisation by  
25 *Streptococcus sobrinus*. EP748591 (Société des Produits Nestlé S.A.) also  
reports the use of fluoridated micellar casein or its micellar subunits for treating  
dental caries or plaque. US4992420 (Nestec S.A.) describes treatment of the  
buccal cavity with kappa-caseino-glycomacropeptide derived from milk for  
eradicating plaque and caries.

Lactic bacteria that are not part of the resident microflora of the mouth have  
never been shown to be really capable of directly adhering to the pellicle of teeth.  
Thus, by colonising the surface of teeth, such lactic bacteria would exert an  
inhibitory activity against the growth of the resident microflora, including oral  
35 pathogens.



### Summary of the invention

The object of the present invention is to provide the use of lactic bacteria  
5 that is not part of the resident microflora of the mouth, that is low acidifying and  
that is capable of adhering directly to the pellicle of the teeth, for the preparation  
of a composition intended for the prophylaxis or the treatment of dental caries,  
dental plaque and periodontal infection.

10 Another object is the use of the lactic bacteria that has been genetically  
modified to increase its adherence to the pellicle of the teeth via adhesion factors  
and/ or genetically modified to be even less acidifying contributing to a pH in the  
oral cavity of about 5.5-7.

15 Another object is the use of lactic bacteria that is not part of the resident  
microflora of the mouth, for the preparation of a composition intended for the  
prophylaxis or the treatment of dental caries, dental plaque and periodontal  
infection, wherein the lactic bacteria is selected from the group consisting of :

- an acidifying lactic bacteria that adheres to the pellicle of the teeth and that  
20 has been genetically modified so that it is low acidifying;
- a non adherent lactic bacteria that is low acidifying and that has been  
genetically modified so that it adheres to the pellicle of the teeth;
- a non-adherent acidifying lactic bacteria that has been genetically modified  
so that it adheres to the pellicle of the teeth and genetically modified so that it is  
25 low acidifying.

Another object is to provide a composition for the health of the mouth  
comprising a lactic bacteria that is not part of the resident microflora, that is low  
acidifying and that is capable of adhering directly to the pellicle of the teeth.

30 Another object is to provide a composition for the health of the mouth  
comprising (1) at least a lactic bacteria that is not part of the resident microflora  
of the mouth, which is capable of adhering directly to the pellicle of the teeth and  
contributing to a pH in the oral cavity of above 5.5, and (2) any forms of caseino-  
glycomacropeptide, micellar casein, fluorinated micellar casein, renneted milk or  
35 bacteriocin.

This Sheet Has Been Replaced



Another object is to provide a composition for the health of the mouth comprising at least one lactic bacteria strain selected from the group consisting of the strains CNCM I-1984, CNCM I-1985, CNCM I-1986, CNCM I-1987 and LMG P-18997.

5

Another object is to provide a composition for the health of the mouth comprising at least one lactic bacteria that has been genetically modified to increase its adherence to the pellicle of the teeth via adhesion factors and/ or genetically modified to be even less acidifying and contributes to a pH in the oral cavity of about 5.5-7.

10

Another object is to provide a composition for the health of the mouth comprising lactic bacteria selected from the group consisting of:

15

- an acidifying lactic bacteria that adheres to the pellicle of the teeth and that has been genetically modified so that it is low acidifying;
- a non adherent lactic bacteria that is low acidifying and that has been genetically modified so that it adheres to the pellicle of the teeth;
- a non-adherent acidifying lactic bacteria that has been genetically modified so that it adheres to the pellicle of the teeth and genetically modified so that it is low acidifying.

20

In a last further aspect, the invention provides a method for screening lactic bacteria capable of adhering to tooth, comprising the steps of: (1) preparing monoclonal antibody recognising specific surface proteins of a lactic bacteria strain capable of adhering to the teeth, (2) screening any lactic bacteria strain by use of the monoclonal antibody of strain capable of adhering to the teeth.

25

#### Detailed description of the invention

30

Within the following description, the mouth defines the oral cavity of humans or animals such as pets, composed by the oral mucosa (gums, lips, cheeks, palate and floor of the mouth), the tongue and the teeth (including artificial structures).

35

The resident microflora of the mouth includes all microorganisms that naturally live in the mouth because they can establish a permanent residence on the oral surfaces. The resident microflora of the mouth also includes bacteria that live in the interfacial region between the dental hard and soft tissues (the junction

This Sheet Has Been Replaced



tooth-gingiva), even though the gingival crevice and the periodontal pocket are not present in a healthy mouth. This microflora includes microorganisms selected from *Streptococcus*, *Staphylococcus*, *Enterococcus*, *Micrococcus*, *Peptostreptococcus*, *Peptococcus*, *Lactobacillus*, *Corynebacterium*, *Actinomyces*,  
5 *Arachnia*, *Rothia*, *Alcaligenes*, *Eubacterium*, *Propionibacterium*, *Bifidobacterium*, *Bacillus*, *Clostridium*, *Neisseria/Branhamella*, *Veillonella*, *Enterobacteriaceae*, *Campylobacter*, *Eikenella*, *Actinobacillus*, *Capnocytophaga*, *Haemophilus*, *Simonsiella*, *Bacteroides*, *Fusobacterium*, *Porphyromonas*, *Prevotella*, *Leptotrichia*, *Wolinella/Selenomonas*, *Mycoplasma*, *Candida*,  
10 *Spirochaetes*, Protozoa.

The transient microflora comprises exogenous bacteria that can be occasionally present in the mouth, but that do not establish a permanent residence. This transient microflora may comprise all the food micro-organisms,  
15 such as the bifidobacteria (*B. infantis*, *B. adolescentis*, *B. breve* and *B. longum*); the lactococci (*Lactococcus lactis* subsp. *lactis*, *Lactococcus lactis* subsp. *cremoris*, and *Lactococcus lactis* subsp. *lactic* biovar *diacetylactis*); the streptococci (*Streptococcus thermophilus*, *S. lactis*, *S. lactis cremoris* and *S. lactis diacetylactis*); the lactobacilli (*Lactobacillus delbrueckii* subsp. *bulgaricus*,  
20 *Lactobacillus helveticus*, *Lactobacillus farciminis*, *Lactobacillus alimentarius*, *Lactobacillus casei* subsp. *casei*, *Lactobacillus delbrueckii* subsp. *lactis*, *Lactobacillus sake*, *Lactobacillus curvatus*, *Lactobacillus fermentum* and the acidophile group comprising *L. johnsonii*; see Fujisawa et al., Int. J. Syst. Bact., 42, 487-491, 1992); the pediococci (*Pediococcus pentosaceus*, *Pediococcus*  
25 *acidilactici* and *Pediococcus halophilus*); the enterococci; the staphylococci (*Staphylococcus xylosus* and *Staphylococcus carnosus*); the micrococci (*Micrococcus varians*); yeast of the genus *Debaromyces*, *Candida*, *Pichia*, *Torulopsis* and *Saccharomyces*; and mold of the genus *Aspergillus*, *Rhizopus*, *Mucor* and *Penicillium*.

30

With respect to the first object of the present invention, the use of lactic bacteria that is not part of the resident microflora of the mouth, that is low acidifying and that is capable of adhering directly to the pellicle of the teeth, for the preparation of a composition intended for the prophylaxis or the treatment of  
35 dental caries, dental plaque and periodontal infection, is concerned.

The lactic bacteria according to the invention are low acidifying and they are capable of adhering directly to the pellicle of the teeth so that the

compositions prepared with these lactic bacteria are intended for displacing pathogens of the teeth or preventing their attachment. Lactic bacteria according to the invention are "low acidifying", which means that they are less acidifying than pathogenic strains. Accordingly, they contribute to a pH in the oral cavity of  
5 about 5.5-7. They are preferably from dairy origin.

The lactic bacteria according to the invention adhere to the pellicle of the teeth via specific or unspecific interaction and/or adhesion factors. The specific adhesion factors are proteins or polysaccharides.  
10

At least one lactic bacteria is selected from the group consisting of *Streptococcus thermophilus*, *Lactococcus lactis subsp. lactis*, and *Lactococcus lactis subsp. lactis biovar diacetylactis* and particularly from the group consisting of the strains CNCM I-1984, CNCM I-1985, CNCM I-1986, CNCM I-1987 and  
15 LMG P-18997.

These strains have been selected among lactic bacteria strains for their capacity of adherence to the pellicle of the teeth, their optimal growth temperature is about 37°C, which is the temperature in the oral cavity. Moreover  
20 they are capable of fermenting glucose and sucrose and do not synthesise glucans, which are factors of pathogenicity of the cariogenic strains.

According to another object of the present invention, it is considering modifying genetically lactic bacteria so that it adheres to the pellicle of the teeth  
25 via adhesion factors. For lactic bacteria that already adheres to the pellicle of the teeth, this modification intends to make the strains more adherent to the surface of the teeth. In the same way, any non-adherent lactic acid bacteria (not *Lactobacilli*) can be genetically modified so that it adheres to the pellicle of the teeth.

This modification of the lactic bacteria can be achieved by insertion of  
30 the genes X17390, X14490 or X53657 (GenBank accession numbers), for example. These gene are responsible in *S. mutans* for the expression of the Antigen I/II that mediates adhesion to salivary glycoproteins.

According to the invention, it is also possible to genetically modify lactic  
35 bacteria so that they are low acidifying. For lactic bacteria that is already low acidifying this modification is intended to increase the said effect by decreasing its lactic acid production.

This modification can be achieved in many ways and preferably according to one the protocols described in the following documents : Boumerdassi *et al.*, Appl. Environ. Microbiol., 63, 2293-2299, 1997; Platteeuw *et al.*, Appl. Environ. Microbiol., 61, 3967-3971, 1995; Ito *et al.*, Biosci. Biotechnol. Biochem., 58, 1569-1573, 1994.

According to the present invention, at least one lactic bacteria genetically modified or not, is used in an "effective quantity" for the preparation of compositions intended for the prophylaxis or the treatment of dental caries, dental plaque and periodontal infection in humans or animals such as pets. This quantity is preferably comprised between  $10^4$  to  $10^9$  cfu/g.

It is also possible to use at least a lactic bacteria, in combination with milk derivatives, such as milk or fermented milk or milk derivatives selected from any forms of caseino-glycomacropeptide, micellar casein, fluorinated micellar casein, renneted milk or bacteriocin, for example.

## BIOCHEMICAL CHARACTERIZATION OF THE SELECTED STRAINS

**Fermentation patterns:** 49 simple sugars were tested with the api 50 CH bioMérieux strip test (bioMérieux SA, 69280 Marcy-l'Etoile, France) and the results are given in the table 1.

**Acidification curves:** the acidification curves were determined at 37°C in the following conditions:

- *S. sobrinus* OMZ 176: FUM sucrose 1% and FUM glucose 1%
- *S. thermophilus* CNCM I-1985: Belliker sucrose 1% and Belliker glucose 1%

Inoculation was always 5%; pH was recorded every 20 min.

*S. thermophilus* CNCM I-1985, from sucrose fermentation, lowers the pH to 4.5, while *S. sobrinus* OMZ 176 to 4.

| Sugar                        | <i>L. lactis</i><br>CNCM I-<br>1987 | <i>L. lactis</i><br>CNCM I-<br>1986 | <i>S. th.</i><br>CNCM I-<br>1984 | <i>S. th.</i><br>CNCM I-<br>1985 | <i>S. th.</i><br>LMG<br>18997 | P- |
|------------------------------|-------------------------------------|-------------------------------------|----------------------------------|----------------------------------|-------------------------------|----|
| Adonitol                     | +++                                 |                                     |                                  |                                  |                               |    |
| Aesculin                     | ++                                  | ++++                                |                                  |                                  |                               |    |
| Amygdalin                    | ++++                                |                                     |                                  |                                  |                               |    |
| D-Arabinose                  |                                     |                                     |                                  |                                  |                               |    |
| L-Arabinose                  |                                     |                                     |                                  |                                  |                               |    |
| D-Arabitol                   |                                     |                                     |                                  |                                  |                               |    |
| L-Arabitol                   | +++                                 |                                     |                                  |                                  |                               |    |
| Arbutin                      | +++                                 | +++                                 |                                  |                                  |                               |    |
| Cellobiose                   | +++                                 | ++++                                |                                  |                                  |                               |    |
| Dulcitol                     |                                     |                                     |                                  |                                  |                               |    |
| Erythritol                   |                                     |                                     |                                  |                                  |                               |    |
| D-Fructose                   | +                                   | ++++                                |                                  |                                  |                               |    |
| D-Fucose                     |                                     |                                     |                                  |                                  |                               |    |
| L-Fucose                     |                                     |                                     |                                  |                                  |                               |    |
| Galactose                    | ++                                  | ++++                                |                                  |                                  |                               |    |
| $\beta$ -Gentiobiose         |                                     | +++                                 |                                  |                                  |                               |    |
| Gluconate                    |                                     |                                     |                                  |                                  |                               |    |
| 2-keto-Gluconate             |                                     |                                     |                                  |                                  |                               |    |
| 5-keto-Gluconate             |                                     |                                     |                                  |                                  |                               |    |
| GlcNAc                       | +                                   | ++++                                |                                  |                                  |                               |    |
| D-Glucose                    | +                                   | ++++                                | +                                | ++                               | ++                            |    |
| Glycerol                     |                                     |                                     |                                  |                                  |                               |    |
| Glycogen                     |                                     |                                     |                                  |                                  |                               |    |
| Inositol                     |                                     |                                     |                                  |                                  |                               |    |
| Inulin                       |                                     |                                     |                                  |                                  |                               |    |
| Lactose                      | +                                   | ++++                                | +++                              | ++++                             | ++++                          |    |
| D-Lyxose                     |                                     |                                     |                                  |                                  |                               |    |
| Maltose                      | ++                                  |                                     |                                  |                                  |                               |    |
| Mannitol                     | +++                                 | ++                                  |                                  |                                  |                               |    |
| D-Mannose                    | +                                   | ++++                                |                                  |                                  |                               |    |
| Melezitose                   |                                     |                                     |                                  |                                  |                               |    |
| Melibiose                    |                                     |                                     |                                  |                                  |                               |    |
| $\alpha$ -Methyl-D-glucoside |                                     |                                     |                                  |                                  |                               |    |
| $\alpha$ -Methyl-D-mannoside |                                     |                                     |                                  |                                  |                               |    |

|                          |     |     |     |      |     |
|--------------------------|-----|-----|-----|------|-----|
| D-Raffinose              |     |     |     |      |     |
| Rhamnose                 |     |     |     |      |     |
| Ribose                   | ++  | ++  |     |      |     |
| Salicin                  | +++ | +++ |     |      |     |
| Sorbitol                 |     |     |     |      |     |
| L-Sorbose                |     |     |     |      |     |
| Starch                   |     |     |     |      |     |
| Sucrose                  |     |     | +++ | ++++ | +++ |
| D-Tagatose               |     |     |     |      |     |
| Trehalose                | ++  |     |     |      |     |
| D-Turanose               | ++  |     |     |      |     |
| Xylitol                  | +++ |     |     |      |     |
| D-Xylose                 |     |     |     |      |     |
| L-Xylose                 |     |     |     |      |     |
| $\beta$ -methyl-xyloside |     |     |     |      |     |

+, ++, +++, ++++ show if the fermentation begins after 3, 6, 24 or 48 hours.

**Table 1.** Sugar fermentation of *L. lactis* CNCM I- 1987, *L. lactis* CNCM I-1986, *S. thermophilus* CNCM I-1984, *S. thermophilus* CNCM I-1985 and , *S. thermophilus* LMG P-18997.

5

The second main object of the present invention concerns a composition for the health of the mouth comprising a lactic bacteria that is not part of the resident microflora of the mouth, that is low acidifying and that is capable of adhering directly to the pellicle of the teeth.

10

These compositions are in particular intended for the prophylaxis or the treatment of dental caries, dental plaque and periodontal infection.

The said lactic bacteria strain is selected from the group consisting of *Streptococcus thermophilus*, *Lactococcus lactis* subsp. *lactis*, and *Lactococcus lactis* subsp. *lactis* biovar *diacetylactis* and preferably from the group consisting of the strains CNCM I-1984, CNCM I-1985, LMG P-18997, CNCM I-1986 and CNCM I-1987.

In such compositions for the health of the mouth, lactic bacteria strains may be genetically modified as described above.

The said lactic bacteria strains may be included in a food, pet food, cosmetic or pharmaceutical composition, for example.

Accordingly, these compositions are preferably toothpaste, mouth rinse, gum, spray, beverage, candies, infant formula, ice cream, frozen dessert, sweet salad dressing, milk preparations, cheese, quark, yogurt, acidified milk, coffee cream or whipped cream, for example.

In these compositions lactic bacteria strains may be included alone or in combination with milk derivatives, for example, in order to obtain synergistic preparations. Accordingly, these compositions for the health of the mouth comprise:

- a lactic bacteria that is not part of the resident microflora of the mouth, which is capable of adhering directly to the pellicle of the teeth;
- any forms of lactic glycopeptides, renneted milk or bacteriocin.

15

The lactic glycopeptides are preferably caseino-glycomacropeptides (CGMP), fluorinated or not micellar casein (which can be obtained as described in the patents EP 0 604 802 and EP 0 748 591) or renneted milk may also be added. The caseino-glycomacropeptides are preferably added in a minimum amount of about 0.1%. It has also been shown that the caseino-glycomacropeptides do not prevent the said lactic bacteria from adhering to the teeth pellicle (fig. 2 and 3).

Synergistic compositions may also be prepared, adding at least one bacteriocin, which is active against Gram-positive oral bacteria. In that case, the oral hygiene compositions may comprise 0.00001 to 50%, and preferably from 0.00001 to 15% of purified bacteriocin, by weight of the composition. The bacteriocin is preferably variacin (EP 0 759 469).

In order to protect the composition from degradation, an oil-soluble antioxidant may also be included. Suitable antioxidants include the "tocopherols", butyl-hydroxyanisole (BHA), butyl-hydroxytoluene (BHT), and ascorbyl palmitate. The oil soluble antioxidant is present in amounts of from 0.005% to 0.5%, preferably 0.005% to 0.01% by weight of the composition.

35

Suitable abrasives for use in dentifrice compositions of the present invention include calcium carbonate, calcium aluminosilicate, alumina, hydrates

alumina, zinc orthophosphate, plastic particles, and silica, of which silica is the preferred abrasive.

5 Compositions according to the invention will have a pH which is orally acceptable and within which the activity of the said lactic bacteria is not compromised. The pH may be in the range 3.0-9.5, preferably in the range 3.5 to 6.5.

10 These compositions may be prepared by conventional processes comprising admixing the ingredients together in the appropriate relative amounts and finally, and if necessary, adjusting the pH to desired value.

15 According to the last aspect of the present invention, a method for screening lactic bacteria capable of adhering to tooth is concerned. This method comprises the steps of:

- (1) preparing monoclonal antibodies recognising specific surface proteins of a lactic bacteria strain capable of adhering to the teeth, and
- (2) screening any lactic bacteria strain by use of the monoclonal antibody of strain capable of adhering to the teeth.

20

The said monoclonal antibodies would be used as a tool to detect the said lactic bacteria strain among other strains growing nearby.

25 The present invention is not to be limited in scope by the specific embodiments described herein. Indeed, various modifications of the invention, in addition to those described herein, will become apparent to those skilled in the art from the foregoing description and accompanying figures. Such modifications are intended to fall within the scope of the claims. Various publications are cited herein, the disclosures of which are incorporated by reference in their entireties to the extent necessary for understanding the present invention. DNA  
30 manipulation, cloning and transformation of bacteria cells are, except where otherwise stated, carried out according to the textbook of Sambrook *et al.* (Sambrook *et al.*, Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Laboratory Press, U.S.A., 1989). These examples are preceded by a brief  
35 description of the plasmids, strains and the various media used, as well as the method for producing a monoclonal antibody.

---

This Sheet Has Been Replaced



The strains *S. thermophilus* S118 (NCC 1529), S123 (NCC 1561), *L. lactis subsp. lactis* 29 (NCC 2211), *L. lactis subsp. lactis biovar diacetylactis* 69 (NCC 2225) were deposited under the Budapest Treaty, at the Collection Nationale de Culture de Microorganismes (CNCM I-1984, CNCM I-1985, CNCM I-1986 and CNCM I-1987 respectively), 25 rue du docteur Roux, 75724 Paris, France, on March 3<sup>rd</sup>, 1998. The strain *S. thermophilus* BF11116 (CNBL1177) was deposited under the Budapest Treaty, at the Belgian Coordinated Collections of Microorganisms LMG P-18997, K.L. Ledeganckstraat 35, B-9000 Gent, Belgium, on July 5<sup>th</sup>, 1999. All restrictions as to the availability of these deposits will be withdrawn upon first publication of this application or another application which claims benefit of priority to this application.

### Figures

- Figures 1a, 1b and 1c represent respectively the adhesion saturation curves for *S. sobrinus* OMZ 176, *L. lactis* NCC2211 and *S. thermophilus* NCC1561.
- Figure 2 represents the curves obtained for the three strains by plotting the number of bound cells versus increasing amounts of CGMP.
- Figure 3 represents the curves obtained for the three strains by plotting the number of bound cells versus increasing amounts of As-CGMP.

### Example 1: STRAINS AND CULTURE CONDITIONS

More than 100 strains (belonging to the Nestlé culture collection) were screened for their ability to attach to saliva-coated hydroxyapatite beads, and in particular the following 23 strains: *S. thermophilus* YS4 (NCC 2284), *S. thermophilus* Sfi6 (NCC 1971), *S. thermophilus* Sfi13 (NCC 2008), *S. thermophilus* Sfi21 (NCC 2038), *S. thermophilus* Sfi39 (NCC 2130), *S. thermophilus* Sfi42 (NCC 2145), *S. thermophilus* Sfi47 (NCC 2172), *S. thermophilus* S118 (NCC 1529), *S. thermophilus* S119 (NCC 1536), *S. thermophilus* S122 (NCC 1554), *S. thermophilus* S123 (NCC 1561), *S. thermophilus* S126 (NCC 1587), *L. lactis subsp. cremoris* 15 (NCC 92), *L. lactis subsp. cremoris* 25 (NCC 1932), *L. lactis subsp. cremoris* 136 (NCC 2419), *L. lactis subsp. diacetylactis* 8 (NCC 1970), *L. lactis subsp. diacetylactis* 28 (NCC 2057), *L. lactis subsp. diacetylactis* 69 (NCC 2225), *L. lactis subsp. diacetylactis* 80 (NCC 2272), *L. lactis subsp. lactis* 29 (NCC 2211), *L. lactis subsp. lactis* 50 (NCC 2224), *L. lactis subsp. lactis* 54 (NCC 2228), *S. macedonicus* 216 (NCC 2484).

The 5 oral strains, *S. sobrinus* OMZ 176, *S. oralis* OMZ 607, *A. naeslundii* OMZ 745, *V. dispar* OMZ 493 and *F. nucleatum* OMZ 596 were obtained from the Institute für Orale Mikrobiologie und Allgemeine Immunologie, University of Zürich and they were cultured in FUM medium in anaerobiosis (GasPackSystem, BBL) at 37°C.

All the strains were stored in glycerol at -20°C and precultured for 14 hours prior to use at their specific optimal temperature; *S. sobrinus* OMZ 176 grew in FUM medium (Gmür *et al.*) lactococci and streptococci in M17 (Difco) except *S. thermophilus* NCC1529, S119, S122, NCC1561 and S126 that grew in Belliker (prepared by dissolution in 1 l water of 20 g tryptone, 5 g yeast extract, 2.5 g gelatine 5 g dextrose, 5 g sucrose, 5 g lactose, 4 g NaCl, 0.5 g Ascorbic acid, 10 g beef extract).

For the plate counting: *S. sobrinus* OMZ 176 was cultured in Mitis-Salivarius agar (Difco), *S. thermophilus* NCC1529, S119, S122, NCC1561, BF11116 and S126 in Belliker agar (prepared by adding to the liquid Belliker 15 g of Bacto agar, Difco), and the remaining lactic bacteria strains in M17 agar (Oxoid).

#### Example 2: PRODUCTION OF A MONOCLONAL ANTIBODY

A monoclonal antibody would be used as a tool to detect *L. lactis subsp. lactis* NCC2211 among 5 oral strains growing together on S-HA discs and forming a biofilm that simulates dental plaque. Therefore the monoclonal antibody was tested against these strains to verify there was no cross-reaction.

To this end, the monoclonal antibody is produced as described by Granato *et al.* "A mouse monoclonal IgE antibody anti-bovine milk lactoglobulin allows studies of allergy in the gastrointestinal tract., Clin. Exp. Immunol., 63, 703-710, 1986.

#### Example 3: SELECTION OF ADHERENT LACTIC BACTERIA

##### Attachment to saliva-coated hydroxyapatite beads (S-HA)

To select among the lactic bacteria dairy strains those able to attach to saliva-coated hydroxyapatite beads (S-HA), the procedure previously described by Neeser *et al.* (1994) was used with slight modifications: beads washings were

done with 150 µl volumes and Hyamine hydroxide was substituted with Benzethonium hydroxide (Sigma).

Briefly, all the strains were grown to the end of the log phase in FUM except *S. thermophilus* NCC1529, S119, S122, NCC1561 and S126 that were cultured in Belliker. *S. sobrinus* OMZ 176, *L. lactis subsp. lactis* NCC2211, 50 and 54, *S. thermophilus* NCC1529, S119, S122, NCC1561 and S126 grew at 37°C, the remaining lactococci at 30°C and the remaining streptococci at 42°C.

5 mg of hydroxyapatite beads (BDH Chemicals Ltd, Poole, England) were covered with 70 µl clarified saliva obtained from volunteers in the lab and prepared as previously explained (Neesser *et al.*, 1994). Saliva coated beads were kept overnight at 4°C, then washed (first with distilled water and after with HEPES buffer) and finally inoculated with 100 µl of the metabolically labelled bacterial suspension (bacteria had been grown in their medium, supplemented with 10 µCi/ml <sup>14</sup>C acetic acid). Adhesion took place during 45 min at 37°C, then unbound bacteria were washed away and the attached cells directly counted in a LKB scintillation counter (type 1219 Rackbeta).

The adhesion percentages are expressed as radioactivity bound to the beads on the total radioactivity added to each well. All measurements were done in triplicate. Table 2 reports the percentages of adhesion to saliva-coated hydroxyapatite beads obtained for several screened strains and for *S. sobrinus* OMZ176 (the reference strain).

| STRAIN                                              | % ADHESION (+/-SD) |
|-----------------------------------------------------|--------------------|
| <i>S. sobrinus</i> OMZ 176                          | 2.23 +/-0.49       |
| <i>S. thermophilus</i> Sfi42 (NCC 2145)             | 0.08 +/-0.02       |
| <i>S. thermophilus</i> Sfi47 (NCC 2172)             | 0.14 +/-0.04       |
| <i>S. thermophilus</i> NCC1529                      | 2.89 +/-0.60       |
| <i>S. thermophilus</i> S119 (NCC 1536)              | 0.15 +/-0.04       |
| <i>S. thermophilus</i> S122 (NCC 1554)              | 0.93 +/-0.17       |
| <i>S. thermophilus</i> NCC1561                      | 2.19 +/-0.50       |
| <i>S. thermophilus</i> S126 (NCC 1587)              | 1.19 +/-0.56       |
| <i>L. lactis subsp. diacetylactis</i> 28 (NCC 2057) | 1.59 +/-0.17       |
| <i>L. lactis subsp. diacetylactis</i> NCC2225       | 1.96 +/-0.40       |
| <i>L. lactis subsp. diacetylactis</i> 80 (NCC 2272) | 1.20 +/-0.35       |
| <i>L. lactis subsp. lactis</i> NCC2211              | 2.85 +/-0.85       |

**Table 2: percentages of adhesion to saliva-coated hydroxyapatite beads obtained for several screened strains**

Four strains, *S. thermophilus* NCC1529 (CNCM I-1984), *S. thermophilus* NCC1561 (CNCM I-1985), *L. lactis subsp. lactis* NCC2211 (CNCM I-1986) (hereinafter *L. lactis* NCC2211) and *L. lactis subsp. diacetylactis* NCC2225 (CNCM I-1987) showed values close to *S. sobrinus* OMZ 176.

*L. lactis* NCC2211 and *S. thermophilus* NCC1561 were chosen as the more promising candidates since they grow very well at 37°C, which is the temperature in the mouth, while *L. diacetylactis* NCC2225 has an optimal growth temperature of 30°C. In particular, *L. lactis* NCC2211 cannot grow on sucrose, but it can ferment a wide range of sugars, moreover other oral strain can provide glucose via their invertase.

#### Adhesion saturation curves

Curves of bound CFU versus CFU inoculated into the well were determined to verify if beads saturation could be obtained. The 50% saturation was directly drawn from the bending point of the curves obtained. The adhesion saturation curves for *S. sobrinus* OMZ 176, *L. lactis* NCC2211 and *S. thermophilus* NCC1561 were determined. They are shown in the Figure 1.

For the three strains the CFU number to be inoculated in the well to get the 50% beads saturation and the corresponding number of bound CFU were directly deduced from the bending point of the curves and are given in the table 3.

|                                | cfu/well | Bound cfu | % adhesion |
|--------------------------------|----------|-----------|------------|
| <i>S. sobrinus</i> OMZ 176     | 4.00E+07 | 4.00E+06  | 10%        |
| <i>L. lactis</i> NCC2211       | 1.00E+07 | 9.00E+05  | 9%         |
| <i>S. thermophilus</i> NCC1561 | 3.00E+07 | 2.00E+06  | 7%         |

**Table 3: Number of CFU to be inoculated per well to get the 50% beads saturation.**

#### Example 4: EFFECT OF CASEINOGLYCOMACROPEPTIDES

The influence of CGMP on the adhesion of *L. lactis* NCC2211 and *S. thermophilus* NCC1561 was studied to verify the possibility of using it to foster the predominance of one of this two strains on the pathogenic ones (namely *S. Sobrinus* OMZ 176. Caseino-glycopeptide (CGMP) and its desialylated

derivative (As-CGMP) were obtained from Nestec S. A., Lausanne (for their preparation see Neeser *et al.*, 1994).

5 The dose-response effect was studied on the adhesion to S-HA beads by inoculating in the well 100  $\mu$ l of bacterial suspension (CFU/ml corresponding to the 50% beads saturation previously calculated) which contained CGMP or As-CGMP in different concentrations and then performing the adhesion assay as usual. Concentrations in the range 0.05-3 mg/ml were tested. No previous incubation of the bacteria in presence of CGMP or As-CGMP was done.

10

In the figure 2, the curves obtained for the three strains by plotting the number of bound cells versus increasing amounts of CGMP are presented, the number of inoculated cells being the one corresponding to the 50% beads saturation formerly calculated for each strain. The strong inhibition observed in the case of *S. sobrinus* OMZ 176 confirms the previous results obtained by

15

As evidenced in the figure 2, 0.25 mg/ml produced 50% inhibition of the adhesion of *S. sobrinus* OMZ 176, while more than 2 mg/ml were necessary to have the same effect with *S. thermophilus* NCC1561. CGMP slightly enhances the adhesion of *L. lactis* NCC2211.

20

As in the case of CGMP, the desyalilated derivative inhibits the adhesion of *S. sobrinus* OMZ 176; only 0.05 mg/ml are needed to produce 50% decrease in the adhesion percentage. As-CGMP does not influence *L. lactis* NCC2211 adhesion, while it slightly fosters the one of *S. thermophilus* NCC1561 (Fig. 3).

25

#### Example 5: TOOTHPASTE

30 Toothpaste is prepared by adding  $10^5$  cfu/ml of at least one of the lactic bacteria strain CNCM I-1984, CNCM I-1985, CNCM I-1986, CNCM I-1987 or LMG P-18997 in a lyophilised form, to the following mixture containing:

|                                |       |
|--------------------------------|-------|
| Cetyl pyridinium chloride      | 1.65% |
| 35 Sorbitol (70% soln)         | 33.0% |
| Glycerin                       | 25.0% |
| Sodium carboxymethyl cellulose | 2.0%  |

|   |                               |       |
|---|-------------------------------|-------|
|   | Sodium fluoride               | 0.25% |
|   | Silica (RP 93)                | 26.3% |
|   | Thickening Silica (Sident 22) | 8.1%  |
| 5 | Sodium saccharine             | 0.5%  |
|   | Poloxamer (Pluronic F108)     | 3.2%, |

This toothpaste is intended for the prophylaxis or the treatment of dental caries, dental plaque and periodontal infection.

10

#### Example 6: ICE CREAM

A cream comprising 10.8% lactic fats, 13.5% milk solids (non fat), 0.3% Emulstab® SE30 et 0.3% Emulstab® foam (Grindsted, DK) is prepared, it is then pasteurised at 105°C for 20s, homogenised at 75°C and 300 bar, cooled to 38°C and inoculated with precultures in MRS medium, taken in exponential growth phase, at a rate of  $10^7$ - $10^8$  cfu/ml of at least one of the lactic bacteria strain CNCM I-1984, CNCM I-1985, CNCM I-1986, CNCM I-1987 or LMG P-18997. The cream is then fermented for 10 hours at 38°C up to a pH of about 4.5. At the end of the fermentation, sucrose and glucose syrup is added thereto. The composition of the cream is presented in table 4 below.

20

The mixture is then beaten, cooled to 4°C, stored at 4°C, chilled to a degree of expansion of 95°C by volume.

| Ingredients           | Composition (kg) | Fats (%) | Non-fat solids (%) | Sucrose (%) | Solids content (%) |
|-----------------------|------------------|----------|--------------------|-------------|--------------------|
| Cream (35%)           | 30,83            | 10,79    | 1,54               |             | 12,33              |
| Powdered skimmed milk | 12,45            |          | 11,95              |             | 11,95              |
| Emulstab® SE30        | 0,41             |          |                    |             | 0,37               |
| Emulstab® foam        | 0,41             |          |                    |             | 0,36               |
| Water                 | 55,91            |          |                    |             |                    |
| Total: cream base     | 100,00           | 10,79    | 13,49              | -           | 25,01              |
| Cream base            | 74,14            | 8,00     | 10,00              | -           | 18,54              |
| Sucrose               | 22,06            |          |                    | 15,00       | 15,00              |
| Glucose syrup         | 3,80             |          |                    |             | 3,00               |
| Fermented Ice cream   | 100,00           | 8,00     | 10,00              | 15,00       | 36,54              |

25 **Table 4**

**Example 7: YOGHURT**

5 5 l MRS culture medium are sterilised for 15 min at 121°C and then inoculated with 5% by volume of an active culture of at least one of the *S. Thermophilus* strain CNCM I-1984, CNCM I-1985 or LMG P-18997 containing approximately  $10^9$  cfu/ml. After incubation for 8 h at 41°C, a starter containing  $4.5 \cdot 10^8$  cfu/ml is obtained.

10 5 l reconstituted skimmed milk having a dry matter content of 10%, to which 0.1% yeast extract has been added, are sterilised for 15 min at 121°C and inoculated with 2% of an active culture of commercial thickening *Streptococcus thermophilus* containing approximately  $10^9$  cells/ml. After incubation for 4 h at 41°C, a starter containing  $4.5 \cdot 10^8$  cells/ml is obtained.

15 One batch of whole milk containing 3.7% fats strengthened with 2.5% skimmed milk powder and then pasteurised for 30 min at 90°C is then inoculated with 2% by volume of the starter of at least one of the strain CNCM I-1984, CNCM I-1985 or LMG P-18997 and 3% by volume of the starter of thickening *Streptococcus thermophilus*. The inoculated milk is stirred, poured into pots and incubated for 4 h at 41°C.

20

The yoghurt obtained has a good firm and smooth texture and is intended for the health of the mouth.

**Example 8: CHEWING GUM**

25

A chewing gum for preventing or treating dental caries, dental plaque or periodontal infection can be prepared adding an active culture of at least one of the *S. Thermophilus* strain CNCM I-1984, CNCM I-1985 or LMG P-18997 so that it contains approximately  $10^4$  to  $10^9$  cfu/g, to the following typical ingredients:

30

|    |                   |        |
|----|-------------------|--------|
|    | Xylitol           | 67.5 % |
|    | Gum base          | 20 %   |
|    | Calcium carbonate | 5 %    |
|    | Glycerin          | 3 %    |
| 35 | Pluronic F127     | 2 %    |
|    | Cellulose gum     | 1 %    |
|    | Balast compounds  | 0.5%   |
|    | Flavor            | 1 %    |

**Example 9: PET FOOD COMPOSITION**

5 A pet food for mouth health is obtained by preparing a feed mixture made up of corn, corn gluten chicken and fish meal, salts, vitamins and minerals. The feed mixture is fed into a preconditioner and moistened. The moistened feed leaving the preconditioner is then fed into an extruder-cooker and gelatinised. The gelatinised matrix leaving the extruder is forced through a die and extruded. The extrudate is cut into pieces suitable for feeding to dogs, dried at about 110°C  
10 for about 20 minutes and cooled to form pellets which have a water activity of about 0.6.

The pellets are sprayed with 3 coating mixtures. Each coating mixture contains active culture of at least one of the *S. Thermophilus* strain CNCM I-1984, CNCM I-1985 or LMG P-18997 but one coating mixture uses hydrogenated soy fat as a  
15 coating substrate, one coating mixture uses water as a coating substrate and one coating mixture uses protein digest as a coating substrate. The pellets contain approximately  $10^4$  to  $10^9$  cfu/g of said strains.

TRAITE DE BUDAPEST SUR LA RECONNAISSANCE  
INTERNATIONALE DU DEPOT DES MICRO-ORGANISMES  
AUX FINS DE LA PROCEDURE EN MATIERE DE BREVETS


## FORMULE INTERNATIONALE

## DESTINATAIRE :

**SOCIETE DES PRODUITS NESTLE S.A.**  
Patents Department  
Avenue Nestlé 55  
CH-1800 VEVEY

RECEPISSE EN CAS DE DEPOT INITIAL,  
délivré en vertu de la règle 7.1 par  
l'AUTORITE DE DEPOT INTERNATIONALE  
identifiée au bas de cette page

NOM ET ADRESSE  
DU DEPOSANT

|                                                                                                                                                                                                                                                                                        |                                                                                                                                                                                                                                                                                                                                         |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>I. IDENTIFICATION DU MICRO-ORGANISME</b>                                                                                                                                                                                                                                            |                                                                                                                                                                                                                                                                                                                                         |
| Référence d'identification donnée par le DEPOSANT :                                                                                                                                                                                                                                    | Numéro d'ordre attribué par l'AUTORITE DE DEPOT INTERNATIONALE :                                                                                                                                                                                                                                                                        |
| <b>NCC 1529</b>                                                                                                                                                                                                                                                                        | <b>I - 1984</b>                                                                                                                                                                                                                                                                                                                         |
| <b>II. DESCRIPTION SCIENTIFIQUE ET/OU DESIGNATION TAXONOMIQUE PROPOSEE</b>                                                                                                                                                                                                             |                                                                                                                                                                                                                                                                                                                                         |
| Le micro-organisme identifié sous chiffre I était accompagné :                                                                                                                                                                                                                         |                                                                                                                                                                                                                                                                                                                                         |
| <input checked="" type="checkbox"/> d'une description scientifique<br><input checked="" type="checkbox"/> d'une désignation taxonomique proposée<br>(Cocher ce qui convient)                                                                                                           |                                                                                                                                                                                                                                                                                                                                         |
| <b>III. RECEPTION ET ACCEPTATION</b>                                                                                                                                                                                                                                                   |                                                                                                                                                                                                                                                                                                                                         |
| La présente autorité de dépôt internationale accepte le micro-organisme identifié sous chiffre I, qu'elle a reçu le <b>03 MARS 1998</b> (date du dépôt initial) <sup>1</sup>                                                                                                           |                                                                                                                                                                                                                                                                                                                                         |
| <b>IV. RECEPTION D'UNE REQUETE EN CONVERSION</b>                                                                                                                                                                                                                                       |                                                                                                                                                                                                                                                                                                                                         |
| La présente autorité de dépôt internationale a reçu le micro-organisme identifié sous chiffre I le _____ (date du dépôt initial) et a reçu une requête en conversion du dépôt initial en dépôt conforme au Traité de Budapest le _____ (date de réception de la requête en conversion) |                                                                                                                                                                                                                                                                                                                                         |
| <b>V. AUTORITE DE DEPOT INTERNATIONALE</b>                                                                                                                                                                                                                                             |                                                                                                                                                                                                                                                                                                                                         |
| Nom : <b>CNCM</b><br>Collection Nationale<br>de Cultures de Microorganismes<br><br>Adresse : <b>INSTITUT PASTEUR</b><br>28, Rue du Docteur Roux<br>F-75724 PARIS CEDEX 15                                                                                                              | Signature(s) de la (des) personne(s) compétente(s) pour représenter l'autorité de dépôt internationale ou de l'(des) employé(s) autorisé(s) : <b>Mme Y. CERNIER</b><br>Directeur Administratif de la CNCM<br><br><br>Date : Paris, le 02 avril 1998 |

<sup>1</sup> En cas d'application de la règle 6.4.d), cette date est la date à laquelle le statut d'autorité de dépôt internationale a été acquis.

**TRAITE DE BUDAPEST SUR LA RECONNAISSANCE  
INTERNATIONALE DU DEPOT DES MICRO-ORGANISMES  
AUX FINS DE LA PROCEDURE EN MATIERE DE BREVETS**

**FORMULE INTERNATIONALE**

**DESTINATAIRE :**

**SOCIETE DES PRODUITS NESTLE S.A.**  
**Patents Department**  
**Avenue Nestlé 55**  
**CH-1800 VEVEY**

RECEPISSE EN CAS DE DEPOT INITIAL,  
délivré en vertu de la règle 7.1 par  
l'AUTORITE DE DEPOT INTERNATIONALE  
identifiée au bas de cette page

**NOM ET ADRESSE  
DU DEPOSANT**

|                                                                                                                                                                                                                                                                                     |                                                                                                                                                                                                                                                                                                                                                          |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>I. IDENTIFICATION DU MICRO-ORGANISME</b>                                                                                                                                                                                                                                         |                                                                                                                                                                                                                                                                                                                                                          |
| Référence d'identification donnée par le<br>DEPOSANT :<br><br><b>NCC 1561</b>                                                                                                                                                                                                       | Numéro d'ordre attribué par<br>l'AUTORITE DE DEPOT INTERNATIONALE :<br><br><b>I - 1985</b>                                                                                                                                                                                                                                                               |
| <b>II. DESCRIPTION SCIENTIFIQUE ET/OU DESIGNATION TAXONOMIQUE PROPOSEE</b>                                                                                                                                                                                                          |                                                                                                                                                                                                                                                                                                                                                          |
| Le micro-organisme identifié sous chiffre I était accompagné :                                                                                                                                                                                                                      |                                                                                                                                                                                                                                                                                                                                                          |
| <input checked="" type="checkbox"/> d'une description scientifique<br><input type="checkbox"/> d'une désignation taxonomique proposée<br><br>(Cocher ce qui convient)                                                                                                               |                                                                                                                                                                                                                                                                                                                                                          |
| <b>III. RECEPTION ET ACCEPTATION</b>                                                                                                                                                                                                                                                |                                                                                                                                                                                                                                                                                                                                                          |
| La présente autorité de dépôt internationale accepte le micro-organisme identifié sous<br>chiffre I, qu'elle a reçu le <b>03 MARS 1998</b> (date du dépôt initial) <sup>1</sup>                                                                                                     |                                                                                                                                                                                                                                                                                                                                                          |
| <b>IV. RECEPTION D'UNE REQUETE EN CONVERSION</b>                                                                                                                                                                                                                                    |                                                                                                                                                                                                                                                                                                                                                          |
| La présente autorité de dépôt internationale a reçu le micro-organisme identifié sous<br>chiffre I le (date du dépôt initial)<br>et a reçu une requête en conversion du dépôt initial en dépôt conforme au Traité de<br>Budapest le (date de réception de la requête en conversion) |                                                                                                                                                                                                                                                                                                                                                          |
| <b>V. AUTORITE DE DEPOT INTERNATIONALE</b>                                                                                                                                                                                                                                          |                                                                                                                                                                                                                                                                                                                                                          |
| Nom : <b>CNCM</b><br>Collection Nationale<br>de Cultures de Microorganismes<br><br>Adresse : <b>INSTITUT PASTEUR</b><br>28, Rue du Docteur Roux<br>F-75724 PARIS CEDEX 15                                                                                                           | Signature(s) de la (des) personne(s)<br>compétente(s) pour représenter l'autorité<br>de dépôt internationale ou de l'(des)<br>employé(s) autorisé(s) : <b>Mme Y. CERISIER</b><br>Directeur Administratif de la CNCM<br><br><br>Date : <b>Paris, le 02 avril 1998</b> |

<sup>1</sup> En cas d'application de la règle 6.4.d), cette date est la date à laquelle le statut d'autorité de dépôt internationale a été acquis.

TRAITE DE BUDAPEST SUR LA RECONNAISSANCE  
INTERNATIONALE DU DEPOT DES MICRO-ORGANISMES  
AUX FINS DE LA PROCEDURE EN MATIERE DE BREVETS

## FORMULE INTERNATIONALE

DESTINATAIRE :

**SOCIETE DES PRODUITS NESTLE S.A.**  
Patents Department  
Avenue Nestlé 55  
CH-1800 VEVEY

RECEPISSE EN CAS DE DEPOT INITIAL,  
délivré en vertu de la règle 7.1 par  
l'AUTORITE DE DEPOT INTERNATIONALE  
identifiée au bas de cette page

NOM ET ADRESSE  
DU DEPOSANT

|                                                                                                                                                                                                                                                                                        |                                                                                                                                                                                                                                                                                                                                     |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>I. IDENTIFICATION DU MICRO-ORGANISME</b>                                                                                                                                                                                                                                            |                                                                                                                                                                                                                                                                                                                                     |
| Référence d'identification donnée par le DEPOSANT :                                                                                                                                                                                                                                    | Numéro d'ordre attribué par l'AUTORITE DE DEPOT INTERNATIONALE :                                                                                                                                                                                                                                                                    |
| <b>NCC 2211</b>                                                                                                                                                                                                                                                                        | <b>I - 1986</b>                                                                                                                                                                                                                                                                                                                     |
| <b>II. DESCRIPTION SCIENTIFIQUE ET/OU DESIGNATION TAXONOMIQUE PROPOSEE</b>                                                                                                                                                                                                             |                                                                                                                                                                                                                                                                                                                                     |
| Le micro-organisme identifié sous chiffre I était accompagné :                                                                                                                                                                                                                         |                                                                                                                                                                                                                                                                                                                                     |
| <input type="checkbox"/> d'une description scientifique<br><input type="checkbox"/> d'une désignation taxonomique proposée<br>(Cocher ce qui convient)                                                                                                                                 |                                                                                                                                                                                                                                                                                                                                     |
| <b>III. RECEPTION ET ACCEPTATION</b>                                                                                                                                                                                                                                                   |                                                                                                                                                                                                                                                                                                                                     |
| La présente autorité de dépôt internationale accepte le micro-organisme identifié sous chiffre I, qu'elle a reçu le <b>03 MARS 1998</b> (date du dépôt initial) <sup>1</sup>                                                                                                           |                                                                                                                                                                                                                                                                                                                                     |
| <b>IV. RECEPTION D'UNE REQUETE EN CONVERSION</b>                                                                                                                                                                                                                                       |                                                                                                                                                                                                                                                                                                                                     |
| La présente autorité de dépôt internationale a reçu le micro-organisme identifié sous chiffre I le _____ (date du dépôt initial) et a reçu une requête en conversion du dépôt initial en dépôt conforme au Traité de Budapest le _____ (date de réception de la requête en conversion) |                                                                                                                                                                                                                                                                                                                                     |
| <b>V. AUTORITE DE DEPOT INTERNATIONALE</b>                                                                                                                                                                                                                                             |                                                                                                                                                                                                                                                                                                                                     |
| Nom : <b>CNCM</b><br>Collection Nationale<br>de Cultures de Microorganismes<br><br>Adresse : <b>INSTITUT PASTEUR</b><br>28, Rue du Docteur Roux<br>F-75724 PARIS CEDEX 15                                                                                                              | Signature(s) de la (des) personne(s) compétente(s) pour représenter l'autorité de dépôt internationale ou de l'(des) employé(s) autorisé(s) : <b>Mme Y. CERSIER</b><br>Directeur Administratif de la CNCM<br><br>Date : Paris, le 02 avril 1998 |

<sup>1</sup> En cas d'application de la règle 6.4.d), cette date est la date à laquelle le statut d'autorité de dépôt internationale a été acquis.

TRAITE DE BUDAPEST SUR LA RECONNAISSANCE  
INTERNATIONALE DU DEPOT DES MICRO-ORGANISMES  
AUX FINS DE LA PROCEDURE EN MATIERE DE BREVETS

## FORMULE INTERNATIONALE

DESTINATAIRE :

**SOCIETE DES PRODUITS NESTLE S.A.**  
**Patents Department**  
**Avenue Nestlé 55**  
**CH-1800 VEVEY**

RECEPISSE EN CAS DE DEPOT INITIAL,  
délivré en vertu de la règle 7.1 par  
l'AUTORITE DE DEPOT INTERNATIONALE  
identifiée au bas de cette page

NOM ET ADRESSE  
DU DEPOSANT

|                                                                                                                                                                                                                                                                            |                                                                                                                                                                                                                                                                                                                                                  |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>I. IDENTIFICATION DU MICRO-ORGANISME</b>                                                                                                                                                                                                                                |                                                                                                                                                                                                                                                                                                                                                  |
| Référence d'identification donnée par le DEPOSANT :                                                                                                                                                                                                                        | Numéro d'ordre attribué par l'AUTORITE DE DEPOT INTERNATIONALE :                                                                                                                                                                                                                                                                                 |
| <b>NCC 2225</b>                                                                                                                                                                                                                                                            | <b>I - 1987</b>                                                                                                                                                                                                                                                                                                                                  |
| <b>II. DESCRIPTION SCIENTIFIQUE ET/OU DESIGNATION TAXONOMIQUE PROPOSEE</b>                                                                                                                                                                                                 |                                                                                                                                                                                                                                                                                                                                                  |
| Le micro-organisme identifié sous chiffre I était accompagné :                                                                                                                                                                                                             |                                                                                                                                                                                                                                                                                                                                                  |
| <input checked="" type="checkbox"/> d'une description scientifique<br><input type="checkbox"/> d'une désignation taxonomique proposée<br>(Cocher ce qui convient)                                                                                                          |                                                                                                                                                                                                                                                                                                                                                  |
| <b>III. RECEPTION ET ACCEPTATION</b>                                                                                                                                                                                                                                       |                                                                                                                                                                                                                                                                                                                                                  |
| La présente autorité de dépôt internationale accepte le micro-organisme identifié sous chiffre I, qu'elle a reçu le <b>03 MARS 1998</b> (date du dépôt initial) <sup>1</sup>                                                                                               |                                                                                                                                                                                                                                                                                                                                                  |
| <b>IV. RECEPTION D'UNE REQUETE EN CONVERSION</b>                                                                                                                                                                                                                           |                                                                                                                                                                                                                                                                                                                                                  |
| La présente autorité de dépôt internationale a reçu le micro-organisme identifié sous chiffre I le (date du dépôt initial) et a reçu une requête en conversion du dépôt initial en dépôt conforme au Traité de Budapest le (date de réception de la requête en conversion) |                                                                                                                                                                                                                                                                                                                                                  |
| <b>V. AUTORITE DE DEPOT INTERNATIONALE</b>                                                                                                                                                                                                                                 |                                                                                                                                                                                                                                                                                                                                                  |
| Nom : <b>CNCM</b><br>Collection Nationale<br>de Cultures de Microorganismes<br><br>Adresse : <b>INSTITUT PASTEUR</b><br>28, Rue du Docteur Roux<br>F-75724 PARIS CEDEX 15                                                                                                  | Signature(s) de la (des) personne(s) compétente(s) pour représenter l'autorité de dépôt internationale ou de l'(des) employé(s) autorisé(s) : <b>Mme Y. CERISIER</b><br>Directeur Administratif de la CNCM<br><br><br>Date : <b>Paris, le 02 avril 1998</b> |

<sup>1</sup> En cas d'application de la règle 6.4.d), cette date est la date à laquelle le statut d'autorité de dépôt internationale a été acquis.

### Claims

1. The use of lactic bacteria that is not part of the resident microflora of the mouth, and that is low acidifying and is capable of adhering directly to the pellicle of the teeth, for the preparation of a composition intended for the prophylaxis or the treatment of dental caries, dental plaque and periodontal infection.
2. The use according to claim 1, wherein the composition is intended for displacing pathogens of the teeth or preventing their attachment.
3. The use according to claim 1 or 2, wherein the lactic bacteria is less acidifying than the pathogenic strains, contributing to a pH in the oral cavity of about 5.5-7.
4. The use according to any of preceding claims, wherein the lactic bacteria is from dairy origin.
5. The use according to any of preceding claims, wherein at least one lactic bacteria is selected from the group consisting of *Streptococcus thermophilus*, *Lactococcus lactis subsp. lactis*, and *Lactococcus lactis subsp. lactis biovar diacetylactis*.
6. The use according to any of preceding claims, wherein at least one lactic bacteria is selected from the group consisting of the strains CNCM I-1984, CNCM I-1985, CNCM I-1986, CNCM I-1987 and LMG P-18997.
7. The use according to any of preceding claims, wherein the lactic bacteria adheres to the pellicle of the teeth via adhesion factors.
8. The use according to any of preceding claims, wherein the lactic bacteria has been genetically modified to increase its adherence to the pellicle of the teeth and/ or genetically modified to be even less acidifying.
9. The use of lactic bacteria that is not part of the resident microflora of the mouth, for the preparation of a composition intended for the prophylaxis or the treatment of dental caries, dental plaque and periodontal infection, wherein the lactic bacteria is selected from the group consisting of :




This Sheet Has Been Replaced





- an acidifying lactic bacteria that adheres to the pellicle of the teeth and that has been genetically modified so that it is low acidifying;
  - a non adherent lactic bacteria that is low acidifying and that has been genetically modified so that it adheres to the pellicle of the teeth;
  - 5     - a non-adherent acidifying lactic bacteria that has been genetically modified so that it adheres to the pellicle of the teeth and genetically modified so that it is low acidifying.
- 10     10. The use according to claims 8 and 9, wherein the lactic bacteria has been genetically modified so that it adheres to the pellicle of the teeth via adhesion factors and contribute to a pH in the oral cavity of about 5.5-7.
- 15     11. The use according to any of preceding claims, wherein the composition is an edible composition comprising an effective quantity of lactic bacteria for the prophylaxis or the treatment of dental caries, dental plaque and periodontal infection.
- 20     12. The use according to any of preceding claims, wherein the composition contains at least  $10^4$ - $10^9$  cfu /g of lactic bacteria.
- 25     13. The use according to any of preceding claims, wherein the lactic bacteria is combined with milk, fermented milk, milk derivatives or bacteriocin.
- 30     14. The use according to claim 13, wherein the milk derivatives are selected from any form of caseino-glycomacropeptide, micellar casein, fluorinated micellar casein or renneted milk.
- 35     15. A composition for the health of the mouth comprising lactic bacteria that is not part of the resident microflora of the mouth and that is low acidifying and that is capable of adhering directly to the pellicle of the teeth.
16. A composition for the health of the mouth according to claim 15, wherein the lactic bacteria has been genetically modified to increase its adherence to the pellicle of the teeth and/ or genetically modified to be even less acidifying.
17. A composition for the health of the mouth according to claim 15 or 16, wherein the lactic bacteria is selected from the group consisting of :

This Sheet Has Been Replaced



- an acidifying lactic bacteria that adheres to the pellicle of the teeth and that has been genetically modified so that it is low acidifying;
- a non adherent lactic bacteria that is low acidifying and that has been genetically modified so that it adheres to the pellicle of the teeth;
- 5     - a non-adherent acidifying lactic bacteria that has been genetically modified so that it adheres to the pellicle of the teeth and genetically modified so that it is low acidifying.

10     18. A composition for the health of the mouth according to any of preceding claims, wherein the lactic bacteria has been genetically modified so that it adheres to the pellicle of the teeth via adhesion factors and contributes to a pH in the oral cavity of about 5.5-7.

15     19. A composition for the health of the mouth according to any of preceding claims, comprising an effective quantity of lactic bacteria for the prophylaxis or the treatment of dental caries, dental plaque and periodontal infection.

20     20. A composition for the health of the mouth according to any of preceding claims, containing at least  $10^4$ - $10^9$  cfu /g of lactic bacteria.

20     21. A composition for the health of the mouth comprising:

- (1) a lactic bacteria that is not part of the resident microflora of the mouth, which is capable of adhering directly to the pellicle of the teeth and contributing to a pH in the oral cavity of about 5.5-7;
- 25     (2) any forms of caseino-glycomacropeptide, micellar casein, fluorinated micellar casein or renneted milk or bacteriocin.

30     22. A composition for the health of the mouth according to any of preceding claims, comprising at least one lactic bacteria strain selected from the group consisting of *Streptococcus thermophilus*, *Lactococcus lactis subsp. lactis*, and *Lactococcus lactis subsp. lactis biovar diacetylactis*.

35     23. A composition for the health of the mouth according any of preceding claims, comprising at least one lactic bacteria strain selected from the group consisting of the strains CNCM I-1984, CNCM I-1985, CNCM I-1986, CNCM I-1987 and LMG P-18997.



24. A method for screening lactic bacteria capable of adhering the teeth, comprising the steps of:

- (1) preparing monoclonal antibodies recognising specific surface proteins of a lactic bacteria strain capable of adhering to the teeth, and
- 5 (2) screening any lactic bacteria strain by use of the monoclonal antibody of strain capable of adhering to the teeth.

10

