CLAIMS

1. Use of a therapeutically effective amount of a composition comprising pilocarpine and oxymetazoline in the manufacture of a medicament for treating an ocular condition in a subject; wherein the ocular condition is selected from the group consisting of presbyopia, mild hyperopia, irregular astigmatism, and hyperopic accommodative esotropia; and wherein the medicament is to be administered to the subject.

2. The use of claim 1, wherein ~~administering comprises administering~~ the medicament ~~composition~~ is to be administered to an eye of the subject.

3. The use of claim 1, wherein the medicament is to be administered to only one eye of the subject.

4. The use of claim 1, wherein the medicament increases refractive power of an eye of the subject by up to about 4.0 diopters.

5. The use of claim 1, wherein the ocular condition could alternatively be corrected with eye glasses having about +0.5D to about +1.0D lenses and wherein the therapeutically effective amount comprises from about 0.3% to about 1.0% pilocarpine.

6. The use of claim 1, wherein the ocular condition could alternatively be corrected with eye glasses having about +1.0D to about +1.5D lenses and wherein the therapeutically effective amount comprises from about 0.8% to about 1.6% pilocarpine.

7. The use of claim 1, wherein the ocular condition could alternatively be corrected with eye glasses having about +1.5D to about +2.0D lenses and wherein the therapeutically effective amount comprises from about 1.4 % to about 2.2% pilocarpine.

8. The use of claim 1, wherein the medicament comprises from about 0.01% to about 10% w/w pilocarpine and from about 0.01% to about 0.5% w/w oxymetazoline.

9. The use of claim 1, wherein the medicament comprises from about 0.5% to about 2% w/w pilocarpine and from about 0.02% to about 0.125% w/w oxymetazoline.

10. The use of claim 1, wherein administrationering ~~the medicament to a subject treats~~ of the medicament to a subject treats the ocular condition for at least four hours.

11. The use of claim 1, wherein administratrionering of the medicament to a subject treats the ocular condition for at least six hours.

12. The use of claim 1, wherein the ocular condition is presbyopia.

13. The use of claim 1, wherein the ocular condition is mild hyperopia.

14. The use of claim 1, wherein the ocular condition is irregular astigmatism.

15. The use of claim 1, wherein the ocular condition is hyperopic accommodative esotropia.