
atom or C_{1-6} alkyl, C_{3-8} cycloalkyl, C_{1-6} alkoxy or phenyl group, each optionally having a substituent selected from the group consisting of C_{1-6} alkyl, C_{3-8} cycloalkyl and $-Q^5-R^{36}$ (wherein Q^5 represents C_{1-3} alkyleneoxy group, and R^{36} represents a hydrogen atom, C_{1-3} alkyl, or cyclopropyl group), each optionally having a halogen atom;
or phenyl group optionally having one or two substituents selected from the group consisting of a halogen atom or C_{1-6} alkyl, phenyl, C_{1-6} alkoxy, pyrrolyl, pyrrolidinyl, pyrazolyl, oxazolyl, oxadiazolyl, imidazolyl, triazolyl, furanyl or morpholinyl group, each optionally having a substituent selected from the group consisting of a halogen atom, C_{1-6} alkyl, C_{3-8} cycloalkyl and $-Q^5-R^{36}$ (wherein Q^5 represents C_{1-3} alkyleneoxy group, and R^{36} represents a hydrogen atom, C_{1-3} alkyl, or cyclopropyl group):
R^4 and R^5 represent a hydrogen atom.

2. The nicotinamide derivative or a salt thereof according to claim 1, wherein R^{32} represents an alkyl group; an alkyl group substituted with a cycloalkyl group; a cycloalkyl group; or a cycloalkyl group substituted with an alkyl group, each containing 3 to 5 carbon atoms in total, or an alkoxyalkyl group containing 2 to 4 carbon atoms in total.

3. The nicotinamide derivative or a salt thereof according to claim 1, wherein R^{32} represents a methyl or ethyl group substituted with a substituent selected from the group consisting of C_{3-8} cycloalkyl and phenyl group.

4. The nicotinamide derivative or a salt thereof according to claim 1, which is represented by the following formula (I-1):
[Formula 5]



(I-1)

wherein R^{26} has the same definitions as those of R^2 described in claim 1, and R^3 has the same definitions as those described in claim 1.

5. The nicotinamide derivative or a salt thereof according to claim 1, wherein R^3 represents a pyridyl group represented by the following formula (VIII-3) or (VIII-4):
[Formula 7]