


10

in dimethylsulfoxide and sonicate prior to end of bombardment with aqueous [18]fluoride ion to a compound of formula



I

5 and, if desired, converting the compounds obtained into pharmaceutically acceptable acid addition salts.

In another aspect, there is provided a compound of formula I as defined herein for use in binding and imaging tau aggregates, beta-amyloid aggregates or alpha-synuclein aggregates.

10 In another aspect, there is provided a compound of formula I as defined herein for use in binding and imaging tau aggregates in Alzheimer patients.

In another aspect, there is provided a compound of formula I as defined herein for use in diagnostic imaging of tau-aggregates in the brain of a mammal.

In another aspect, there is provided a method of imaging tau-aggregate deposits, comprising


15 - detecting the compound of formula I that had associated with one or more tau-aggregate deposits following introduction of a detectable quantity of a preparation as defined herein into a mammal.

A further object of the embodiments disclosed herein is a pharmaceutical composition, containing compounds of formula I and pharmaceutically acceptable carriers, which may be used
20 for identifying potential patients.

In another aspect, there is provided a pharmaceutical preparation comprising a compound of formula I as defined herein and a pharmaceutical acceptable carrier.

Claims

1. A compound of formula



wherein

5 R is hydrogen or tritium;

F is fluoro or ¹⁸F;

wherein the compound of formula I does not have tritium and ¹⁸F simultaneously,
or a pharmaceutically acceptable acid addition salt.

10 2. A compounds of formula I according to claim 1, which compounds are


2-(6-fluoro-pyridin-3-yl)-9H-dipyrido[2,3-b;3',4'-d]pyrrole

³H-2-(6-fluoro-pyridin-3-yl)-9H-dipyrido[2,3-b;3',4'-d]pyrrole


and [18F]-2-(6-fluoro-pyridin-3-yl)-9H-dipyrido[2,3-b;3',4'-d]pyrrole.

15 3. A process for the manufacture of a compound of formula I as defined in any one of claims 1
or 2, which process comprises

a) coupling a compound of formula 2 (X = Cl, Br)




with suitable boronic acids or boronic acid esters of formula 3



20

wherein R' is hydrogen or lower alkyl,


to afford compounds of formula I




wherein R is tritium, and, if desired, converting the compounds obtained into pharmaceutically acceptable acid addition salts or

d) dissolving a compound of formula **10**

5



in dimethylsulfoxide and sonicate prior to end of bombardment with aqueous [18]fluoride ion to a compound of formula



10 and, if desired, converting the compounds obtained into pharmaceutically acceptable acid addition salts.

4. A compound of formula **I** according to any one of claims 1 or 2 for use in binding and imaging tau aggregates, beta-amyloid aggregates or alpha-synuclein aggregates.

15

5. A compound of formula **I** according to any one of claims 1 or 2 for use in binding and imaging tau aggregates in Alzheimer patients.

6. A compound of formula **I** according to any one of claims 1 or 2 for use in diagnostic imaging of tau-aggregates in the brain of a mammal.

20

7. A pharmaceutical preparation comprising a compound of formula **I** according to any one of claims 1 or 2 and a pharmaceutical acceptable carrier.

8. A method of imaging tau-aggregate deposits, comprising detecting the compound of formula **I** that had associated with one or more tau-aggregate deposits following introduction of a detectable quantity of a preparation according to claim 7 into a mammal.