BIOCONJUGATES MADE FROM RECOMBINANT N-GLYCOSYLATED PROTEINS FROM PROCARYOTIC CELLS

Field of the Invention

The present invention relates to bioconjugates, specifically bioconjugate vaccines, made from recombinant glycoproteins, namely N-glycosylated proteins. The invention comprises one or more introduced N-glycosylated proteins with optimized amino acid consensus sequence(s), nucleic acids encoding these proteins as well as corresponding vectors and host cells. In addition, the present invention is directed to the use of said proteins, nucleic acids, vectors and host cells for preparing bioconjugate vaccines. Furthermore, the present invention provides methods for producing bioconjugate vaccines.

Background of the Invention

Glycoproteins are proteins that have one or more covalently attached sugar polymers. N-linked protein glycosylation is an essential and conserved process occurring in the endoplasmic reticulum of eukarotic organisms. It is important for protein folding, oligomerization, stability, quality control, sorting and transport of secretory and membrane proteins (Helenius, A., and Aebi, M. (2004). Roles of N-linked glycans in the endoplasmic reticulum. Annu. Rev. Biochem. 73, 1019-1049).

Protein glycosylation has a profound influence on the antigenicity, the stability and the half-life of a protein. In addition, glycosylation can assist the purification of proteins by chromatography, e.g. affinity chromatography with lectin ligands bound to a solid phase interacting with glycosylated moieties of the protein. It is therefore established practice to produce many glycosylated proteins recombinantly in eukaryotic cells to provide biologically and pharmaceutically useful glycosylation patterns.

It has been demonstrated that a bacterium, the food-borne pathogen *Campylobacter jejuni,* can also N-glycosylate its proteins (Szymanski, et al. (1999). Evidence for a system of general protein glycosylation in *Campylobacter jejuni*. Mol. Microbiol. 32, 1022-1030). The machinery required for glycosylation is encoded by 12 genes that are clustered in the so-called *pgl* locus.

Disruption of N-gylcosylation affects invasion and pathogenesis of *C. jejuni* but is not lethal as in most eukaryotic organisms (Burda P. and M. Aebi, (1999). The dolichol pathway of N-linked glycosylation. Biochim Biophys Acta 1426(2):239-57). It is possible to reconstitute the N-glycosylation of *C. jejuni* proteins by recombinantly expressing the *pgl* locus and acceptor glycoprotein in *E. coli* at the same time (Wacker et al. (2002). N-linked glycosylation in *Campylobacter* jejuni and its functional transfer into *E. coli*. Science 298, 1790-1793).

Diarrheal illness is a major health problem associated with international travel in terms of frequency and economic impact. Traveller's diarrhea refers to an enteric illness acquired when a person travels from a developed to a developing country. Today, over 50 million people travel each year from developed countries to developing countries and up to 50% of these travelers report having diarrhea during the first 2 weeks of their week of their stay. There has been no significant decline in the incidence of traveller's diarrhea since the 1970s, despite efforts made by the tourism industry to improve local infrastructure.

Traveller's diarrhea is acquired through the ingestion of faecally contaminated food and less commonly water. Bacteria are the main cause of traveller diarrhea's, being responsible for up to 80% of the infections. Enterotoxigenic *E. coli* (ETEC) is the most frequently isolated bacterium in all parts of the world associated with traveler's diarrhea, followed by *Shigella* spp and *C. jejuni*.

Shigellosis remains a serious and common disease. In addition to causing watery diarrhea, *Shigellae* are a major cause of dysentery (fever, cramps, and blood and/or mucus in the stool). Man is the only natural host for this bacterium. The estimated number of Shigella infections is over 200 million annually. About 5 million of these cases need hospitalization and a million people die. Three serogroups are mostly responsible for the disease described as bacillary dysentery: *S. dysenteriae*, *S. flexneri* and *S. sonnei*.

S. dysenteriae and *S. flexneri* are responsible for most infections in the tropics, with case fatalities up to 20%. Shigellosis occurs both endemically and as epidemics. In many tropical countries, endemic infection is largely due to *S. flexneri* whereas major epidemics of *S. dysenteriae* have occurred in Central America, Central Africa and Southeast Asia. These


epidemics are major public-health risks. Infections, primarily due to *S. sonnei* and less frequently *S. flexneri* continue to occur in industrialized countries.

Conjugate vaccines have shown promising results against *Shigella* infections. O-specific polysaccharides of *S. dysenteriae* type 1 have been used to synthesize a conjugate vaccine that has elicited an immune response in mice. Such vaccines have been synthesized chemically and conjugated to human serum albumin or has been developed where the O-polysaccharide has been purified from *Shigella*. The O-specific polysaccharides of *S. sonnei* and *S. flexneri* also have been conjugated chemically to *P. aeruginosa* exotoxin and have elicited a significant immune response in mice. Additionally, they have been shown to be immunogenic and safe in humans. However, chemical conjugation is an expensive and time-consuming process that does not always yield reliable and reproducible vaccines. This leads to good manufacturing practices (GMP) problems when seeking to develop such bioconjugate vaccines on a commercial scale.

Summary of the Invention


In one aspect, the present invention is directed to a bioconjugate vaccine comprising: a protein carrier comprising an inserted consensus sequence, D/E – X – N – Z – S/T, wherein X and Z may be any natural amino acid except proline; at least one antigenic polysaccharide from at least one bacterium, linked to the protein carrier, wherein the at least one antigenic polysaccharide is at least one bacterial O-antigen from one or more strains of *Shigella*, *E. coli* or *Pseudomonas aeruginosa*; and, optionally, an adjuvant.

In another aspect, the present invention is directed to a *Shigella* bioconjugate vaccine comprising: a protein carrier comprising Exotoxin of *Pseudomonas aeruginosa* (EPA) that has been modified to contain at least one consensus sequence D/E – X – N – Z – S/T, wherein X and Z may be any natural amino acid except proline; at least one polysaccharide chain linked to the protein carrier and having the following structure:



; and, optionally, an adjuvant.

In yet another aspect, the present invention is directed to a *Shigella dysenteriae* O1 bioconjugate vaccine comprising: a protein carrier having the sequence provided in SEQ. ID NO.: 7; at least one polysaccharide chain linked to the protein carrier and having the following structure:



; and an adjuvant.

In yet additional aspects, the present invention is directed to: a plasmid comprising SEQ. ID NO. 5; a genetic sequence comprising SEQ. ID NO. 5; an amino acid sequence comprising SEQ. ID NO. 6; an amino acid sequence comprising SEQ. ID NO. 7; or vector pGVXN64.

In another aspect, the present invention is directed to an expression system for producing a bioconjugate vaccine against at least one bacterium comprising: a nucleotide sequence encoding an oligosaccharyl transferase (OST/OTase); a nucleotide sequence encoding a protein carrier; and at least one antigenic polysaccharide synthesis gene cluster from the at least one bacterium, wherein the antigenic polysaccharide is a bacterial O-antigen.

In still another aspect, the present invention is directed to an expression system for producing a bioconjugate vaccine against *Shigella dysenteriae* O1 comprising: a nucleotide sequence encoding PgIB having SEQ. ID NO. 2 ; a nucleotide sequence encoding a modified EPA having SEQ. ID NO. 6; and a polysaccharide synthesis gene cluster comprising SEQ. ID NO. 5.

In yet another aspect, the present invention contemplates a method of producing an O1-bioconjugate in a bioreactor comprising the steps: expressing in bacteria: modified EPA containing at least one consensus sequence, D/E – X – N – Z – S/T, wherein X and Z may be any natural amino acid except proline, or AcrA; PgIB; and one or more O1-polysaccharides; growing the bacteria for a period of time to produce an amount of the O1-bioconjugate comprising the AcrA or the modified EPA linked to the one more O1-polysaccharides; extracting periplasmic proteins; and separating the O1-bioconjugate from the extracted periplasmic proteins.

In an additional aspect, the present invention contemplates a method of producing an *S. dysenteriae* bioconjugate vaccine, said method comprising: assembling a polysaccharide of *S. dysenteriae* in a recombinant organism through the use of glycosyltransferases; linking said polysaccharide to an asparagine residue of one or more target proteins in said recombinant organism, wherein said one or more target proteins contain one or more T-cell epitopes.

In a further aspect, the present invention contemplates a method of producing an *S. dysenteriae* bioconjugate vaccine, said method comprising: introducing genetic information encoding for a metabolic apparatus that carries out N-glycosylation of a target protein into a prokaryotic organism to produce a modified prokaryotic organism, wherein the genetic information required for the expression of one or more recombinant target proteins is introduced into said prokaryotic organism, and wherein the metabolic apparatus comprises specific glycosyltransferases for the assembly of a polysaccharide of *S. dysenteriae* on a lipid carrier and an oligosaccharyltransferase, the oligosaccharyltransferase covalently linking the polysaccharide to an asparagine residue of the target protein, and the target protein containing at least one T-cell epitope; producing a culture of the modified prokaryotic organism; and obtaining glycosylated proteins from the culture medium.

Description of the Figures

Fig. 1 illustrates the N-glycosylation of Lip proteins derived from constructs A to C (see example 1). *E. coli* Top 10 cells carrying a functional *pgl* operon from *C. jejuni* (Wacker et al., 2002, supra) and a plasmid coding for constructs A (lane 2), B (lane 1), and C (lane 3) or a

mutant of construct C with the mutation D121A (lane 4). Proteins were expressed and purified from periplasmic extracts. Shown is the SDS-PAGE and Coomassie brilliant blue staining of the purified protein fractions.

Fig. 2 shows the N-glycosylation analysis of the different proteins that were analyzed for the sequence specific N-glycosylation by the *C. jejuni pgl* operon (Wacker et al., 2002, supra) in CLM24 cells (Feldman et al., (2005). Engineering N-linked protein glycosylation with diverse O antigen lipopolysaccharide structures in Escherichia coli. Proc. Natl. Acad. Sci. USA 102, 3016-3021) or Top10 cells (panel E lanes 1-6) or SCM7 cells (Alaimo, C, Catrein, I., Morf, L., Marolda, C. L., Callewaert, N., Valvano, M. A., Feldman, M. F., Aebi, M. (2006). Two distinct but interchangeable mechanisms for flipping of lipid-linked oligosaccharides. EMBO Journal 25, 967-976) (panel E, lanes 7, 8) expressing said proteins from a plasmid. Shown are SDS-PAGE separated periplasmic extracts that were transferred to a nitrocellulose membrane and visualized with specific antisera. In panels A-D, the top panels show immunoblots probed with anti AcrA antiserum (Wacker et al. 2002, supra; Nita-Lazar, M., Wacker, M., Schegg, B., Amber, S., and Aebi, M. (2005). The N-X- S/T consensus sequence is required but not sufficient for bacterial N-linked protein glycosylation. Glycobiology 15, 361-367), whereas the bottom panels show immunoblots probed with R12 antiserum (Wacker et al., 2002, *supra*). + and - indicate the presence of the functional or mutant *pgl* operon in the cells. Panel A contains samples of the soluble wildtype AcrA with the pelB signal sequence and the hexa histag (SEQ ID NO: 24) (lanes 1 , 2), AcrA- N273Q (lane 3, 4), and AcrA-D121A (lane 5). Panel B: AcrA (lanes 1, 2), AcrA-T145D (lane 3), AcrA-N123Q-N273Q-T145D (lanes 4, 5). Panel C: AcrA-F115D-T145D (lanes 1, 2), AcrA-N123Q-N273Q-N272D (lanes 3, 4). Panel D: AcrA-N273Q (lanes 1, 2), AcrA-N273Q-F122P (lanes 3, 4). Panel E: CtxB (lanes 1, 2), CtxB-W88D (lanes 3, 4), CtxB-Q56/DSNIT (lanes 5, 6), and CtxB-W88D-Q56/DSNIT. 'DSNIT' disclosed as SEQ ID NO: 25.

Fig. 3 shows the engineering of multiple glycosylation sites in OmpH1. The ΔwaaL strain SCM6 was co-transformed with plasmid pACYC*pgl* (encoding entire *pgl locus*) and plasmids expressing wild type OmpH1 (lane 1), $OmpH1^{N139S}$-myc (lane 2), $OmpH1^{KGN\rightarrow NIT,\ HFGDD\rightarrow DSNIT}$-myc (lane 3), $QmpH1^{RGD\rightarrow NIT,\ HFGDD\rightarrow DSNIT}$-myc (|lane 4), $OmpH1^{KGN\rightarrow NIT,\ RGD\rightarrow NIT}$-myc (lane 5)$^{KGN\text{-}NIT,\ RGD\rightarrow N,T.\ HFGDD\rightarrow DSNIT}$-myc (lane 6) or $OmpH1^{RGD\rightarrow NIT,\ V83T}$-myc (lane 7). 'HFGDD' and 'DSNIT' disclosed as SEQ ID NOS 26 and 25, respectively. The cells

were grown aerobically, induced with 0.5% arabinose for 3 hours prior to analysis. Whole cell lysates were TCA precipitated after equalizing the optical density of the cultures as described in the materials and methods section. The proteins were separated by 15% SDS-PAGE and transferred onto a PVDF membrane. *First panel*, immunoblot of whole cell lysates probed with anti-myc tag antobodies. *Bottom panel*, immunoblot of whole cell lysates probed with glycan-specific antiserum. The positions of unglycosylated- and glycosylated OmpH1 are indicated on the right.

Fig. 4. shows fluorescence microscopy of cells expressing various OmpH1 variants. Fluorescence microscopy was performed by the using an Axioplan2 microscope (Carl Zeiss). Images were combined by using Adobe Photoshop, version CS2. SCM6 cells expressing OmpH1 (panel A), OmpH1^{N139S} (panel B), OmpH1^{C20S} (panel C), OmpH1 $^{KGN\rightarrow NIT,\ HFGDD\rightarrow DSNIT}$ (panel D), OmpH1$^{RGD\rightarrow NIT,HFGDD\rightarrow DSNIT}$ (panel E), OmpH1$^{KGN\rightarrow NIT,\ RGD\rightarrow NIT}$ (panel F), OmpH1$^{V83T,KGN\rightarrow NIT}$ (panel G), and OmpH1$^{KGN\rightarrow NIT,RGD\rightarrow NIT,HFGDD\rightarrow DSNIT}$ (panel H). 'HFGDD' and 'DSNIT' disclosed as SEQ ID NOS 26 and 25, respectively. The first column is a merge of the pictures in columns 2, 3, and 4 represented in greytones on black background. Column 2: blue fluorescence in greytones from DAPI stain, column 3: green fluorescence from glycan specific fluorescence, column 4: red fluorescence from anti-myc staining.

Fig. 5A shows a schematic of the capsular polysaccharides and lipopolysacharides in Gram-positive and Gram-negative bacteria.

Fig. 5B shows genomic DNA, with integrated PglB and EPA, and plasmid DNA, which is interchangeable (i.e., exchangeable), encoding a polysaccharide synthesis gene cluster.

Fig. 6A shows the production process of conjugate vaccines using technology of the invention.

Fig. 6B shows the construction of the *Shigella dysenteriae* O1 antigen expression plasmid pGVXN64.

Figs. 7A and 7B show schematics of the protein glycosylation pathway utilized in the present invention.

Figs. 8A and 8B are schematics depicting expression platforms for bioconjugate production of the present invention.

Fig. 9 shows production of *Shigella* bioconjugates.

Fig. 10A shows polysaccharide biosynthesis of the O-antigen of *S. dysenteriae* Serotype O1 on undecaprenolpyrophosphate (UPP).

Fig. 10B shows a schematic of a carrier protein, such as EPA, onto which N-glycosylation sites can be designed.

Fig. 11 shows bicoonjugates that elicit an immune response against *Shigella* O1 polysaccharide in mice.

Fig. 12 shows results from the production of a *Shigella* O1-EPA bioconjugate (e.g., O1-EPA) in a bioreactor.

Fig. 13 shows purification of O1-EPA.

Fig. 14A shows a Western blot analysis of a series of specimens of *Shigella* O1-AcrA bioconjugates produced in an LB shake flask taken under various conditions.

Fig. 14B provides different serotypes of *Shigella* and the polysaccharide structure that defines their antigenicity (i.e., *Shigella* O-antigens).

Fig. 15 shows the expansion of the anomeric region of 1H NMR spectrum of an example of a *S. dysenteriae* Serotype O1 bioconjugate of the invention.

Fig. 16A shows protein samples of a *Shigella* O1 Bioconjugate (e.g., EPA-O1) normalized to biomass concentration (0.1 OD_{600nm} of cells/lane).

Fig. 16B shows the periplasmic extract from a *Shigella* O1 Bioconjugate production process which was loaded on a 7.5% SDS-PAGE, and stained with Coomasie to identify EPA and EPA-O1.

Fig. 17A shows protein fractions from 1. Source Q analyzed by SDS-PAGE and stained by Coomassie to identify the *Shigella* O1 bioconjugate.

Fig. 17B shows protein fractions from 2. Source Q column analyzed on SDS-PAGE and stained by Coomassie to identify the *Shigella* O1 bioconjugate.

Fig. 18A shows protein fractions from Superdex 200 column analyzed by SDS-PAGE and stained by Coomassie stained to identify the *Shigella* O1 bioconjugate.

Fig. 18B shows *Shigella* bioconjugates from different purification steps analyzed by SDS-PAGE and stained by Coomassie.

Detailed Description of the Invention

Introduction to Invention

The present invention provides a versatile in vivo glycosylation platform.

European Patent Application No. 03 702 276.1 (European Patent 1 481 057) teaches a procaryotic organism into which is introduced a nucleic acid encoding for (i) specific glycosyltransferases for the assembly of an oligosaccharide on a lipid carrier, (ii) a recombinant target protein comprising a consensus sequence "N - X - S/T", wherein X can be any amino acid except proline, and (iii) an oligosaccharyl transferase of *C. jejuni* (OTase) that covalently links said oligosaccharide to the consensus sequence of the target protein. Said procaryotic organism produces N-glycans with a specific structure which is defined by the type of the specific glycosyltransferases.

The known N-glycosylation consensus sequence in a protein allows for the N-glycosylation of recombinant target proteins in procaryotic organisms comprising the oligosaccharyl transferase (OTase) of *C. jejuni.*

The object of the present invention is to provide proteins as well as means and methods for producing such proteins having an optimized efficiency for N-glycosylation that can be produced in procaryotic organisms in vivo. Another object of the present invention aims at the

more efficient introduction of N-glycans into recombinant proteins for modifying antigenicity, stability, biological, prophylactic and/or therapeutic activity of said proteins. A further object is the provision of a host cell that efficiently displays recombinant N-glycosylated proteins of the present invention on its surface.

In a first aspect, the present invention provides a recombinant N-glycosylated protein, comprising one or more of the following N-glycosylated optimized amino acid sequence(s):

D/E - X - N - Z - S/T, (optimized consensus sequence)

wherein X and Z may be any natural amino acid except Pro, and wherein at least one of said N-glycosylated partial amino acid sequence(s) is introduced.

It was surprisingly found that the introduction of specific partial amino acid sequence(s) (optimized consensus sequence(s)) into proteins leads to proteins that are efficiently N-glycosylated by the oligosaccharyl transferase (OST, OTase) from *Campylobacter* spp., preferably *C. jejuni*, in these introduced positions.

The term "partial amino acid sequence(s)" as it is used in the context of the present invention will also be referred to as "optimized consensus sequence(s)" or "consensus sequence(s)". The optimized consensus sequence is N-glycosylated by the oligosaccharyl transferase (OST, OTase) from *Campylobacter* spp., preferably *C. jejuni*, much more efficiently than the regular consensus sequence "N - X - S/T" known in the prior art.

In general, the term "recombinant N-glycosylated protein" refers to any heterologous poly- or oligopeptide produced in a host cell that does not naturally comprise the nucleic acid encoding said protein. In the context of the present invention, this term refers to a protein produced recombinantly in any host cell, e.g. an eukaryotic or prokaryotic host cell, preferably a procaryotic host cell, e.g. *Escherichia* ssp., *Campylobacter* ssp., *Salmonella* ssp., *Shigella* ssp., *Helicobacter* ssp., *Pseudomonas* ssp., *Bacillus* ssp., more preferably *Escherichia coli, Campylobacter jejuni, Salmonella typhimurium* etc., wherein the nucleic acid encoding said protein has been introduced into said host cell and wherein the encoded protein is N-glycosylated by the OTase from *Campylobacter* spp., preferably *C. jejuni,* said transferase enzyme naturally occurring in or being introduced recombinantly into said host cell.

In accordance with the internationally accepted one letter code for amino acids the abbreviations D, E, N, S and T denote aspartic acid, glutamic acid, asparagine, serine, and threonine, respectively. Proteins according to the invention differ from natural or prior art proteins in that one or more of the optimized consensus sequence(s) D/E - X - N - Z - S/T is/are introduced and N-glycosylated. Hence, the proteins of the present invention differ from the naturally occurring *C. jejuni* proteins which also contain the optimized consensus sequence but do not comprise any additional (introduced) optimized consensus sequences.

The introduction of the optimized consensus sequence can be accomplished by the addition, deletion and/or substitution of one or more amino acids. The addition, deletion and/or substitution of one or more amino acids for the purpose of introducing the optimized consensus sequence can be accomplished by chemical synthetic strategies well known to those skilled in the art such as solid phase-assisted chemical peptide synthesis. Alternatively, and preferred for larger polypeptides, the proteins of the present invention can be prepared by standard recombinant techniques.

The proteins of the present invention have the advantage that they may be produced with high efficiency and in any procaryotic host comprising a functional *pgl* operon from *Campylobacter* spp., preferably *C. jejuni* . Preferred alternative OTases from *Campylobacter* spp. for practicing the aspects and embodiments of the present invention are *Campylobacter coli* and *Campylobacter lari* (see Szymanski, C.M. and Wren, B.W. (2005). Protein glycosylation in bacterial mucosal pathogens. Nat. Rev. Microbiol. 3:225-237). The functional *pgl* operon may be present naturally when said procaryotic host is *Campylobacter* spp., preferably *C. jejuni*. However, as demonstrated before in the art and mentioned above, the *pgl* operon can be transferred into cells and remain functional in said new cellular environment.

The term "functional *pgl* operon from *Campylobacter* spp., preferably *C. jejuni*" is meant to refer to the cluster of nucleic acids encoding the functional oligosaccharyl transferase (OTase) of *Campylobacter* spp., preferably *C. jejuni*, and one or more specific glycosyltransferases capable of assembling an oligosaccharide on a lipid carrier, and wherein said oligosaccharide can be transferred from the lipid carrier to the target protein having one or more optimized amino acid sequence(s): D/E - X N - Z - S/T by the OTase. It to be understood that the term "functional

pgl operon from *Campylobacter* spp., preferably *C. jejuni*' in the context of this invention does not necessarily refer to an operon as a singular transcriptional unit. The term merely requires the presence of the functional components for N-glycosylation of the recombinant protein in one host cell. These components may be transcribed as one or more separate mRNAs and may be regulated together or separately. For example, the term also encompasses functional components positioned in genomic DNA and plasmid(s) in one host cell. For the purpose of efficiency, it is preferred that all components of the functional *pgl* operon are regulated and expressed simultaneously.

It is important to realize that only the functional oligosaccharyl transferase (OTase) should originate from *Campylobacter* spp., preferably *C. jejuni*, and that the one or more specific glycosyltransferases capable of assembling an oligosaccharide on a lipid carrier may originate from the host cell or be introduced recombinantly into said host cell, the only functional limitation being that the oligosaccharide assembled by said glycosyltransferases can be transferred from the lipid carrier to the target protein having one or more optimized consensus sequences by the OTase. Hence, the selection of the host cell comprising specific glycosyltransferases naturally and/or incapacitating specific glycosyltransferases naturally present in said host as well as the introduction of heterologous specific glycosyltransferases will enable those skilled in the art to vary the N-glycans bound to the optimized N-glycosylation consensus site in the proteins of the present invention.

As a result of the above, the present invention provides for the individual design of N-glycan-patterns on the proteins of the present invention. The proteins can therefore be individualized in their N-glycan pattern to suit biological, pharmaceutical and purification needs.

In a preferred embodiment, the proteins of the present invention may comprise one but also more than one, preferably at least two, preferably at least 3, more preferably at least 5 of said N-glycosylated optimized amino acid sequences.

The presence of one or more N-glycosylated optimized amino acid sequence(s) in the proteins of the present invention can be of advantage for increasing their antigenicity, increasing their stability, affecting their biological activity, prolonging their biological half-life and/or simplifying their purification.

The optimized consensus sequence may include any amino acid except proline in position(s) X and Z. The term "any amino acids" is meant to encompass common and rare natural amino acids as well as synthetic amino acid derivatives and analogs that will still allow the optimized consensus sequence to be N-glycosylated by the OTase. Naturally occurring common and rare amino acids are preferred for X and Z. X and Z may be the same or different.

It is noted that X and Z may differ for each optimized consensus sequence in a protein according to the present invention.

The N-glycan bound to the optimized consensus sequence will be determined by the specific glycosyltransferases and their interaction when assembling the oligosaccharide on a lipid carrier for transfer by the OTase. Those skilled in the art can design the N-glycan by varying the type(s) and amount of the specific glycosyltransferases present in the desired host cell.

N-glycans are defined herein as mono-, oligo- or polysaccharides of variable compositions that are linked to an ε-amide nitrogen of an asparagine residue in a protein via an N-glycosidic linkage. Preferably, the N-glycans transferred by the OTase are assembled on an undecaprenol-pyrophosphate lipid-anchor that is present in the cytoplasmic membrane of gram-negative or positive bacteria. They are involved in the synthesis of O antigen, O polysaccharide and peptidoglycan (Bugg, T. D., and Brandish, P. E. (1994). From peptidoglycan to glycoproteins: common features of lipid-linked oligosaccharide biosynthesis. FEMS Microbiol Lett *119*, 255-262; Valvano, M. A. (2003). Export of O-specific lipopolysaccharide. Front Biosci *8*, s452-471).

In a preferred embodiment, the recombinant protein of the present invention comprises one or more N-glycans selected from the group of N-glycans from *Campylobacter* spp., preferably *C. jejuni*, the N-glycans derived from oligo- and polysaccharides transferred to O antigen forming O polysaccharide in Gram-negative bacteria or capsular polysaccharides from Gram-positive bacteria, preferably: P. aeruginosa 09, 011 ; *E. coli* 07, 09, 016, 0157 and *Shigella dysenteriae* O1 and engineered variants thereof obtained by inserting or deleting glycosyltransferases and epimerases affecting the polysaccharide structure.

In a further preferred embodiment, the recombinant protein of the present invention comprises two or more different N-glycans.

For example, different N-glycans on the same protein can prepared by controlling the timing of the expression of specific glycosyltransferases using early or late promoters or introducing factors for starting, silencing, enhancing and/or reducing the promoter activity of individual specific glycosyltransferases. Suitable promoters and factors governing their activity are routinely available to those in the art.

There is no limitation on the origin of the recombinant protein of the invention. Preferably said protein is derived from mammalian, bacterial, viral, fungal or plant proteins. More preferably, the protein is derived from mammalian, most preferably human proteins. For preparing antigenic recombinant proteins according to the invention, preferably for use as active components in vaccines, it is preferred that the recombinant protein is derived from a bacterial, viral or fungal protein.

In a further preferred embodiment, the present invention provides for recombinant proteins wherein either the protein and/or the N-glycan(s) is (are) therapeutically and/or prophylactically active. The introduction of at least one optimized and N-glycosylated consensus sequence can modify or even introduce therapeutic and/or prophylactic activity in a protein. In a more preferred embodiment, it is the protein and/or the N-glycan(s) that is (are) immunogenically active. In this case, the introduced N-glycosylation(s) may have a modifying effect on the proteins biological activity and/or introduce new antigenic sites and/or may mask the protein to evade degrading steps and/or increase the half-life.

The recombinant proteins of the present invention can be efficiently targeted to the outer membrane and/or surface of host cells, preferably bacteria, more preferably gram- negative bacteria. For assisting the surface display and/or outer membrane localization, it is preferred that the recombinant protein of the invention further comprise at least one polypeptide sequence capable of targeting said recombinant protein to the outer membrane and/or cell surface of a bacterium, preferably a gram-negative bacterium.

In a preferred embodiment, the recombinant protein of the invention is one, wherein said targeting polypeptide sequence is selected from the group consisting of type II signal peptides (Paetzel, M., Karla, A., Strynadka, N.C., and Dalbey, R. E. 2002. Signal peptidases. Chem Rev 102: 4549-4580.) or outer membrane proteins (reviewed in Wemerus, H., and Stahl, S. 2004. Biotechnological applications for surface-engineered bacteria. Biotechnol Appl Biochem 40: 209-228.), preferably selected from the group consisting of the full length protein or the signal peptides of OmpH1 from *C. jejuni*, JIpA from *C. jejuni*, outer membrane proteins from *E. coli*, preferably OmpS, OmpC, OmpA, OprF, PhoE, LamB, Lpp'OmpA (a fusion protein for surface display technology, see Francisco, JA_1 Earhart, C. F., and Georgiou, G. 1992. Transport and anchoring of beta- lactamase to the external surface of Escherichia coli. Proc Natl Acad Sci U S A 89: 2713- 2717.), and the lnp protein from *Pseudomonas aeruginosa*.

In a different aspect, the present invention relates to a nucleic acid encoding a recombinant protein according to the invention. Preferably, said nucleic acid is a mRNA, a DNA or a PNA, more preferably a mRNA or a DNA, most preferably a DNA. The nucleic acid may comprise the sequence coding for said protein and, in addition, other sequences such as regulatory sequences, e.g. promoters, enhancers, stop codons, start codons and genes required to regulate the expression of the recombinant protein via the mentioned regulatory sequences, etc. The term "nucleic acid encoding a recombinant protein according to the invention" is directed to a nucleic acid comprising said coding sequence and optionally any further nucleic acid sequences regardless of the sequence information as long as the nucleic acid is capable of producing the recombinant protein of the invention in a host cell containing a functional *pgl* operon from *Campylobacter* spp., preferably *C. jejuni*. More preferably, the present invention provides isolated and purified nucleic acids operably linked to a promoter, preferably linked to a promoter selected from the group consisting of known inducible and constitutive prokaryotic promoters, more preferably the tetracycline promoter, the arabinose promoter, the salicylate promoter, *lac-*, *trc-*, and *tac* promotors (Baneyx, F. (1999). Recombinant protein expression in Escherichia coli. Curr Opin Biotechnol *10,* 411-421 ; Billman-Jacobe, H. (1996). Expression in bacteria other than Escherichia coli. Curr Opin Biotechnol 7, 500-504.). Said operably linked nucleic acids can be used for, e.g. vaccination.

Furthermore, another aspect of the present invention relates to a host cell comprising a nucleic acid and/or a vector according to the present invention. The type of host cell is not limiting as long as it accommodates a functional *pgl* operon from *C. jejuni* and one or more nucleic acids coding for recombinant target protein(s) of the present invention. Preferred host cells are prokaryotic host cells, more preferably bacteria, most preferably those selected from the group consisting of *Escherichia* ssp., *Campylobacter* ssp., *Salmonella* ssp., *Shigella* ssp., *Helicobacter* ssp., *Pseudomonas* ssp., *Bacillus* ssp., preferably *Escherichia coli,* more preferably *E. coli* strains Top10, W3110, CLM24, BL21 , SCM6 and SCM7 (Feldman et al., (2005). Engineering N-linked protein glycosylation with diverse O antigen lipopolysaccharide structures in Escherichia coli. Proc. Natl. Acad. Sci. U S A *102*, 3016-3021 ; Alaimo, C, Catrein, I., Morf, L., Marolda, C. L., Callewaert, N., Valvano, M. A., Feldman, M. F., Aebi, M. (2006). Two distinct but interchangeable mechanisms for flipping of lipid-linked oligosaccharides. EMBO Journal 25, 967-976) and *S. enterica* strains SL3261 (*Salmonella enterica* sv. *Typhimurium* LT2 (delta) aroA, see Hoiseth, S. K., and Stocker, B.A. 1981 , Aromatic-dependent *Salmonella typhimurium* are non-virulent and effective as live vaccines. Nature 291 :238-239), SL3749 (*Salmonella enterica* sv. *Typhimurium* LT2 waaL, see Kaniuk et al., J. Biol. Chem. 279: 36470-36480) and SL3261 ΔwaaL.

In a more preferred embodiment, the host cell according to the invention is one that is useful for the targeting to the outer membrane and/or surface display of recombinant proteins according to the invention, preferably one, wherein said host cell is a recombinant gram-negative bacterium having:

i) a genotype comprising nucleotide sequences encoding for

a) at least one natural or recombinant specific glycosyltransferase for the assembly of an oligosaccharide on a lipid carrier,

b) at least one natural or recombinant prokaryotic oligosaccharyl transferase (OTase) from *Campylobacter* spp., preferably *C. jejuni*,

c) at least one recombinant protein according to the invention, preferably a protein further comprising a targeting polypeptide, and

ii) a phenotype comprising a recombinant N-glycosylated protein according to the invention that is located in and/or on the outer membrane of the gram-negative bacterium.

The host cell for the above embodiment is preferably selected from the group consisting of Escherichia ssp., *Campylobacter* ssp., *Shigella* ssp, *Helicobacter* ssp. and *Pseudomonas* ssp., *Salmonella* ssp., preferably *E. coli*, more preferably *E. coli* strains Top10, W3110, CLM24, BL21 , SCM6 and SCM7, and *S. enterica* strains SL3261 , SL3749 and SL326iδwaaL (see Hoiseth, S. K., and Stocker, B.A. 1981. Aromatic- dependent *Salmonella typhimurium* are non-virulent and effective as live vaccines. Nature 291 : 238-239), SL3749 (Kaniuk, N.A., Vinogradov, E., and Whitfield, C. 2004. Investigation of the structural requirements in the lipopolysaccharide core acceptor for ligation of O antigens in the genus *Salmonella*: WaaL "ligase" is not the sole determinant of acceptor specificity. J Biol Chem 279: 36470-36480).

Because preferred proteins of the present invention may have a therapeutic or prophylactic activity by themselves and/or due to the introduced N-glycosylation sites, they can be used for the preparation of a medicament. The type of protein for practicing the invention is not limited and, therefore, proteins of the invention such as EPO, IFN- alpha, TNFalpha, IgG, IgM, IgA, interleukins, cytokines, viral and bacterial proteins for vaccination like *C. jejuni* proteins such as HisJ (CjO734c), AcrA (CjO367c), OmpH1 (CjO982c), Diphteria toxin (CRM 197), Cholera toxin, P. aeruginosa exoprotein, to name just a few, and having introduced therein the optimized N-glycosylated consensus sequence are useful for preparing a medicament (Wyszynska, A., Raczko, A., Lis, M., and Jagusztyn-Krynicka, E. K. (2004). Oral immunization of chickens with avirulent *Salmonella* vaccine strain carrying *C. jejuni* 72Dz/92 cjaA gene elicits specific humoral immune response associated with protection against challenge with wild-type Campylobacter. Vaccine 22, 1379-1389).

In addition, the nucleic acids and/or vectors according to the invention are also useful for the preparation of a medicament, preferably for use in gene therapy.

Moreover, a host cell according to the invention, preferably one that has a phenotype comprising an N-glycosylated recombinant protein of the invention that is located in and/or on the outer membrane of a bacterium, preferably a gram-negative bacterium, more preferably one

of the above-listed gram-negative bacteria, is particularly useful for the preparation of a medicament.

More preferably, a protein of the invention is used for the preparation of a medicament for the therapeutic and/or prophylactic vaccination of a subject in need thereof.

In a more preferred embodiment, the present invention relates to the use of a nucleic acid and/or a vector according to the invention for the preparation of a medicament for the therapeutic and/or prophylactic vaccination of a subject in need thereof, preferably by gene therapy.

The host cells of the invention displaying said N-glycosylated recombinant proteins are particularly useful for preparing vaccines, because the displayed N-glycosylated proteins are abundantly present on the host cell's surface and well accessible by immune cells, in particular their hydrophilic N-glycans, and because the host cells have the added effect of an adjuvant, that, if alive, may even replicate to some extent and amplify its vaccination effects.

Preferably, the host cell for practicing the medical aspects of this invention is an attenuated or killed host cell.

Another advantage of the use of the inventive host cells for preparing medicaments, preferably vaccines, is that they induce IgA antibodies due to the cellular component.

Preferably, said host cells are used according to the invention for inducing IgA antibodies in an animal, preferably a mammal, a rodent, ovine, equine, canine, bovine or a human. It is preferred that said subject in need of vaccination is avian, mammalian or fish, preferably mammalian, more preferably a mammal selected from the group consisting of cattle, sheep, equines, dogs, cats, and humans, most preferably humans. Fowls are also preferred.

A further aspect of the present invention relates to a pharmaceutical composition, comprising at least one protein, at least one nucleic acid, a least one vector and/or at least one host cell according to the invention. The preparation of medicaments comprising proteins or host cells, preferably attenuated or killed host cells, and the preparation of medicaments comprising nucleic acids and/or vectors for gene therapy are well known in the art. The preparation scheme for the final pharmaceutical composition and the mode and details of its

administration will depend on the protein, the host cell, the nucleic acid and/or the vector employed.

In a preferred embodiment, the pharmaceutical composition of the invention comprises a pharmaceutically acceptable excipient, diluent and/or adjuvant.

The present invention provides for a pharmaceutical composition comprising at least one of the following, (i) a recombinant protein, a host cell, a nucleic acid and/or a recombinant vector being/encoding/expressing a recombinant protein according to the present invention, and (ii) a pharmaceutically acceptable excipient, diluent and/or adjuvant.

Suitable excipients, diluents and/or adjuvants are well-known in the art . An excipient or diluent may be a solid, semi-solid or liquid material which may serve as a vehicle or medium for the active ingredient. One of ordinary skill in the art in the field of preparing compositions can readily select the proper form and mode of administration depending upon the particular characteristics of the product selected, the disease or condition to be treated, the stage of the disease or condition, and other relevant circumstances (*Remington's Pharmaceutical Sciences*, Mack Publishing Co. (1990)). The proportion and nature of the pharmaceutically acceptable diluent or excipient are determined by the solubility and chemical properties of the pharmaceutically active compound selected, the chosen route of administration, and standard pharmaceutical practice. The pharmaceutical preparation may be adapted for oral, parenteral or topical use and may be administered to the patient in the form of tablets, capsules, suppositories, solution, suspensions, or the like. The pharmaceutically active compounds of the present invention, while effective themselves, can be formulated and administered in the form of their pharmaceutically acceptable salts, such as acid addition salts or base addition salts, for purposes of stability, convenience of crystallization, increased solubility, and the like.

A further aspect of the present invention is directed to a method for producing N-linked glycosylated proteins, comprising the steps of:

a) providing a recombinant organism, preferably a prokaryotic organism, comprising nucleic acids coding for

i) a functional *pgl* operon from *Campylobacter* spp., preferably *C. jejuni*, and

ii) at least one recombinant target protein comprising one or more of the following N-glycosylated optimized amino acid consensus sequence(s):

D/E - X - N - Z - S/T,

wherein X and Z may be any natural amino acid except Pro, and wherein at least one of said N-glycosylated optimized amino acid consensus sequence(s) is introduced, and

b) culturing the recombinant organism in a manner suitable for the production and N-glycosylation of the target protein(s).

Preferably, the target protein is one of the above described recombinant proteins according to the invention.

In a preferred method of the invention, the functional *pgl* operon from *Campylobacter* spp., *preferably C. jejuni*, comprises nucleic acids coding for

i) recombinant OTase from *Campylobacter* spp., *preferably C. jejuni*, and

ii) recombinant and/or natural specific glycosyltransferases from *Campylobacter* spp., preferably *C. jejuni*, and/or

iii) recombinant and/or natural specific glycosyltransferases from species other than *Campylobacter* spp.,

for the assembly of an oligosaccharide on a lipid carrier to be transferred to the target protein by the OTase.

Moreover, in a preferred embodiment the present invention relates to a method for preparing a host cell according to the invention comprising the steps of:

i) providing a gram-negative bacterium,

ii) introducing into said bacterium at least one nucleotide sequence encoding for

a) at least one recombinant specific glycosyltransferase for the assembly of an oligosaccharide on a lipid carrier, and/or

b) at least one recombinant oligosaccharyl transferase (OTase) from *Campylobacter* spp., preferably *C. jejuni*, and/or

c) at least one recombinant protein comprising one or more of the following N-glycosylated optimized amino acid consensus sequence(s):

D/E - X - N - Z - S/T,

wherein X and Z may be any natural amino acid except Pro, and wherein at least one of said N-glycosylated optimized amino acid consensus sequence(s) is introduced, and

iii) culturing said bacterium until at least one recombinant N-glycosylated protein coded by the nucleotide sequence of c) is located in and/or on the outer membrane of the gram-negative bacterium.

For practicing the preferred methods above, the recombinant procaryotic organism or host cell is preferably selected from the group of bacteria consisting of Escherichia ssp., *Campylobacter* ssp., *Salmonella* ssp., *Shigella* ssp., *Helicobacter* ssp., *Pseudomonas* ssp., Bacillus ssp., preferably Escherichia coli, preferably *E. coli* strains Top10, W3110, W3110ΔwaaL, BL21 , SCM6 and SCM7, and S. *enterica* strains SL3261 , SL3749 and SL3261ΔwaaL.

Another preferred method for producing, isolating and/or purifying a recombinant protein according to the invention comprises the steps of:

a) culturing a host cell,

b) removing the outer membrane of said recombinant gram-negative bacterium; and

c) recovering said recombinant protein.

Exemplary methods for removing the outer membrane of a cell, preferably a prokaryotic cell, more preferably a gram-negative bacterial cell, are suitable enzymatic treatment methods, osmotic shock detergent solubilisation and the French press method.

Most preferred, the present invention relates to a method, wherein recombinant or natural specific glycosyltransferases from species other than *Campylobacter* spp., preferably *C. jejuni*, are selected from the group of glycosyltransferases and epimerases originating from bacteria, archea, and/or eukaryota that can be functionally expressed in said host cell.

Bioconjugate Vaccines

An embodiment of the invention involves novel bioconjugate vaccines. A further embodiment of the invention involves a novel approach for producing such bioconjugate vaccines that uses recombinant bacterial cells that directly produce immunogenic or antigenic bioconjugates. In one embodiment, bioconjugate vaccines can be used to treat or prevent bacterial diseases, such as diarrhea, nosocomial infections and meningitis. In further embodiments, biooconjugate vaccines may have therapeutic and/or prophylactic potential for cancer or other diseases.

Conjugate vaccines can be administered to children to protect against bacterial infections and can provide a long lasting immune response to adults. Constructs of the invention have been found to generate an IgG response in animals. It has been found that an IgG response to a *Shigella* O-specific polysaccharide-protein conjugate vaccine in humans correllates with immune protection in humans. (Passwell, J.H. *et al.*, "Safety and Immunogenicity of Improved *Shigella* O-Specific Polysaccharide-Protein Conjugate Vaccines in Adults in Israel" Infection and Immunity, 69(3):1351-1357 (Mar. 2001).) It is believed that the polysaccharide (i.e. sugar residue) triggers a short-term immune response that is sugar-specific. Indeed, the human immune system generates a strong response to specific polysaccharide surface structures of bacteria, such as O-antigens and capsular polysaccharides. However, since the immune response to polysaccharides is IgM dependent, the immune system develops no memory. The protein carrier that carries the polysaccharide triggers an IgG response that is T-cell dependent and that provides long lasting protection since the immune system develops memory.

A typical vaccination dosage for humans is about 1 to 25 μg, preferably about 1 μg to about 10μg, most preferably about 10 μg. Optionally, a vaccine, such as a bioconjugate vaccine of the present invention, includes an adjuvant.

Synthesized complexes of polysaccharides (i.e., sugar residues) and proteins (i.e., protein carriers) can be used as conjugate vaccines to protect against a number of bacterial infections. In one aspect, the instant invention is directed to a novel bioengineering approach for producing immunogenic conjugate vaccines that provide advantages over classical chemical conjugation methods. In one embodiment, the approach involves *in vivo* production of glycoproteins in bacterial cells, for example, Gram-negative cells such as *E. coli*.

The biosynthesis of different polysaccharides is conserved in bacterial cells. The polysaccharides are assembled on carrier lipids from common precursors (activated sugar nucleotides) at the cytoplasmic membrane by different glycosyltransferases with defined specificity. Lipopolysaccharides (LPS) are provided in gram-negative bacteria only, e.g. *Shigella* spp., *Pseudomonas* spp. and *E. coli* (ExPEC, EHEC).

The synthesis of lipopolysaccharides (LPS) starts with the addition of a monosaccharide to the carrier lipid undecaprenyl phosphate at the cytoplasmic side of the membrane. The antigen is built up by sequential addition of monosaccharides from activated sugar nucleotides by different glycosyltransferases and the lipid-linked polysaccharide is flipped through the membrane by a flippase. The antigen-repeating unit is polymerized by an enzymatic reaction. The polysaccharide is then transferred to the Lipid A by the Ligase WaaL forming the LPS that is exported to the surface, whereas the capsular polysaccharide is released from the carrier lipid after polymerization and exported to the surface. The biosynthetic pathway of these polysaccharides enables the production of LPS bioconjugates *in vivo*, capturing the polysaccharides in the periplasm to a protein carrier. Bioconjugates, such as LPS bioconjugates, are preferred in the present invention.

As shown in Figure 5A, both Gram-positive and Gram-negative bacteria have a cell membrane that is surrounded by capsular polysaccharides. Gram-negative bacteria additionally have an outer membrane over the cell membrane, with a periplasmic space separating the two membranes. In addition, they contain lipopolysacharides at the surface. When Gram-negative

bacteria, such as *E. coli*, is used to produce a conjugate vaccine of the present invention, the glycoprotein used in the conjugate vaccine is assembled in the periplasmic space.

Conjugate vaccines have been successfully used to protect against bacterial infections. The conjugation of an antigenic polysaccharide to a protein carrier is required for protective memory response, as polysaccharides are T-cell independent antigens. Polysaccharides have been conjugated to protein carriers by different chemical methods, using activation reactive groups in the polysaccharide as well as the protein carrier.

Figure 6A shows the production process of conjugate vaccines using technology of the invention compared to the currently used process. Currently, conjugate vaccines are produced using two fermentation runs and after several purification and chemical cleavage steps, as schematically shown in the top panel. This current approach has a number of problems. First, large scale cultivation of pathogenic organisms is required. Second, the conjugation approach is dependent on the polysaccharide. Third, the approach has low reproducibility. Fourth, the approach has low homogeneity due to unspecific conjugation. Fifth, the approach also has low purity due to excess of polysaccharide in conjugation. Finally, the current approach provides yields of less than 20%.

As shown in the bottom panel of Figure 6A, in an embodiment, the innovative technology of the invention can be used to develop conjugate vaccines (e.g., bioconjugate vaccines) completely *in vivo* with non-pathogenic cells, avoiding chemical reactions and providing high purity after a few purification steps. This novel method also allows for the production of bioconjugate vaccines that are not feasible using current methods. Moreover, the conjugation and purification process is independent of the polysaccharide antigen that is used. As a result, bioconjugate vaccines can be engineered faster using novel glycan structures. The increased homogeneity of resulting conjugates and the improved reproducibility (i.e., no batch to batch variability) of such conjugates makes this a highly attractive process from quality control and regulatory perspectives. In addition, the novel method provides good yield (30-60 mg/L and up to 200 mg/L).

The present invention is directed to a novel conjugation process involving engineering bacterial cells to produce the final bioconjugate vaccines. One embodiment of the invention allows the production of bioconjugate vaccines *in vivo*, circumventing the chemical conjugation and therefore simplifying the production process. The technology includes a novel genetic/enzymatic mechanism for the *in vivo* synthesis of novel bioconjugates consisting of protein-linked saccharides.

The basis of one aspect of the invention includes the discovery that *Campylobacter jejuni* contains a general N-linked protein glycosylation system, an unusual feature for prokaryotic organisms. Various proteins of *C. jejuni* have been shown to be modified by a heptasaccharide. This heptasaccharide is assembled on undecaprenyl pyrophosphate, the carrier lipid, at the cytoplasmic side of the inner membrane by the stepwise addition of nucleotide activated monosaccharides catalyzed by specific glycosyltransferases. The lipid-linked oligosaccharide then flip-flops (diffuses transversely) into the periplasmic space by a flipppase, e.g., PglK. In the final step of N-linked protein glycosylation, the oligosaccharyltransferase (e.g., PglB) catalyzes the transfer of the oligosaccharide from the carrier lipid to Asn residues within the consensus sequence Asp/Glu-Xaa-Asn-Zaa-Ser/Thr (i.e., D/E – X – N – Z – S/T), where the Xaa and Zaa can be any amino acid except Pro (Figure 7A). We have successfully transferred the glycosylation cluster for the heptasaccharide into *E. coli* and were able to produce N-linked glycoproteins of *Campylobacter*.

We have been able to demonstrate that PglB does not have a strict specificity for the lipid-linked sugar substrate. The antigenic polysaccharides assembled on undecaprenyl pyrophosphate are captured by PglB in the periplasm and transferred to a protein carrier (Feldman, 2005; Wacker, M., *et al.*, Substrate specificity of bacterial oligosaccharyltransferase suggests a common transfer mechanism for the bacterial and eukaryotic systems. Proc Natl Acad Sci U S A, 2006. 103(18): p. 7088-93.) The enzyme will also transfer a diverse array of undecaprenyl pyrophosphate (UPP) linked oligosaccharides if they contain an N-acetylated hexosamine at the reducing terminus. The nucleotide sequence for *pglB* is provided at SEQ. ID NO. 1, whereas the amino acid sequence for PglB is provided at SEQ. ID. NO. 2.

Figures 7A and 7B show schematics of the protein glycosylation pathway (i.e., N-glycosylation system) of the present invention. In an embodiment, the protein glycosylation pathway of *C. jejuni* (e.g., including pgl operon) can be introduced into *E. coli*. In Figure 7A, an oligosaccharide, specifically a heptasaccharide made of five N-acetyl-D-galactosamine units, one glucose unit and one 2,4-diacetamido-2,4,6-trideoxy-D-glucose unit, is assembled onto a lipid carrier, undecaprenylpyrophosphate (UDP), using glycosyltransferases (e.g., pglA, pglC, pglH, J; I) at the cytoplasmic side of the inner membrane and is transferred to the periplasmic space by way of a flippase called PgIK. Separately, a carrier protein depicted as a spiral and containing consensus sequence D/E – X – N – Z – S/T (i.e., Asp/Glu–Xaa–Asn–Zaa–Ser/Thr) is translated in the cytoplasm and is secreted into the periplasmic space. In the final step, an oligosaccharyl transferase (OST or OTase) (e.g., PgIB) transfers the heptasaccharide to Asn residues within a consensus sequence of the carrier protein to produce a glycoprotein.

Figure 7B also shows biosynthesis of a polysaccharide (i.e., an antigenic polysaccharide or antigen) by stepwise action of glycosyltransferases, and transfer of the O-antigen to the periplasm by way of flippase, followed by polymerization into a polysaccharide using a polymerase (e.g., wzy). Separately, a carrier protein, such as EPA, is produced and secreted into the periplasm. An oligosaccharyl transferase (OST or OTase), such as PglB, has relaxed substrate specificity and transfers the polysaccharide from a lipid carrier to Asn in the consensus sequence within EPA.

Figure 8A shows a schematic depicting an embodiment of the expression platform for bioconjugate production of the present invention. The technology of the invention is versatile in that various existing carrier proteins can be employed, so long as the carrier protein contains or is modified to contain the consensus sequence, as discussed earlier. In particular, Figure 8A illustrates the construction of an expression host, such as an engineered *E. coli* bacterium in an embodiment of the invention. Such an *E. coli* contains the general components of a glycosylation system (i.e., an OST/OTase, e.g., PglB, and a protein carrier, e.g. EPA). Such components can be integrated into the genome of an *E. coli* strain. In addition, the Ligase WaaL as well as WecG are deleted. Additionally, specific components for polysaccharide antigen expression (i.e., a polysaccharide synthesis gene cluster containing, for example, glycosyl

transferase, polymerase, flippase, and sugar biosynthesis enzymes) can be provided by the addition of an exchangeable plasmid. This construction allows for specific glycosylation of the protein carrier with a polysaccharide of choice *in vivo*.

In an embodiment of the expression system for a bacterial bioconjugate that is compatible with Good Manufacturing Practices (GMP), DNA encoding the inducible oligosaccharyltransferase and carrier protein can be stably integrated into a bacterial (e.g., *E. coli*) genome such that genes for antibiotic selection can be omitted. For example, as shown in Figure 5B, PgIB and EPA is integrated into genomic DNA, whereas plasmid DNA, which is interchangeable (i.e., exchangeable), encodes a polysaccharide synthesis gene cluster.

In another embodiment, Figure 8B shows an expression system for a bacterial bioconjugate that includes three plasmids. A first plasmid codes for the carrier protein, e.g., AcrA from *Campylobacter jejuni*, which has two N-glycosylation sites and is directed to the periplasm by a PelB signal peptide. A second plasmid codes for the OST/OTase, e.g., PgIB from *C. jejuni*, which is membrane-bound. A third plasmid is a native plasmid that codes, e.g., for a polysaccharide (O antigen) synthesis cluster, such as that for *Shigella dysenteriae* O1.

In an embodiment, an expression plasmid for a bacterial O antigen, such as the *Shigella dysenteriae* O1 antigen, can be constructed as in pGVXN64 shown in Figure 6B. This plasmid encodes all enzymes necessary to synthesize the polysaccharides in the *Shigella dysenteriae* strain that make up the O1 serotype. These enzymes are listed in the left-hand column of Figure 6B. Vector pGVXN64 expressing the *Shigella dysenteriae* O1 antigen was constructed by digestion of pLARFR1 (Vanbleu, E. *et al.*, "Genetic and physical map of the pLAFR1 vector" DNA Seq. 15(3):225-227 (2004)) with EcoR1 followed by insertion of an oligonucleotide cassette (5`-AATTCTGCAGGATCCTCTAGAAGCTTGG (SEQ. ID NO. 3) and 5`-AATTCCAAGCTTCTAGAGGATCCTGCAG (SEQ. ID NO. 4). The BamH1 fragment of pSDM7 (Falt, I. *et al.*, "Construction of recombinant aroA salmonellae stably producing the Shigella Dysenteriae serotype 1 O-antigen and structural characterization of the Salmonella/Shigella hybrid LPS" Microb. Pathog. 20(1):11-30 (1996)) containing the rfb and rfp cluster of *Shigella dysenteriae* O1 was then cloned via the BamH1 site into the oligonucleotide

cassette containing pLAFR1. The complete nucleotide sequence encoding the *Shigella dysenteriae* O1 antigen in the pGVXN64 plasmid is set forth as SEQ. ID NO.: 5 in the Sequence Listing provided below.


The host organism for an expression system of the invention can be, e.g., an *Escherichia coli* strain such as *Escherichia coli* W31110ΔwaaL. The deletion of WaaL prevents the transfer of any polysaccharide to the lipid A core. The chromosomal copy of WaaL can also be replaced by PglB. The strain also contains mutation in wbbL, therefore it does not produce any *E. coli* O16 polysaccharide. To further increase the production of carrier lipid linked polysaccharide, wecG has been deleted to prevent the formation of ECA (Entero Common Antigen).

In one aspect, the instant invention is further directed to the development of bioconjugate vaccines, preferably LPS bioconjugate vaccines, against one or more *Shigella* species, which are invasive, gram-negative bacteria. *Shigella* species cause Shigellosis, a severe inflammation of the colon. There are 165 million cases in the world every year, with 70% of such cases being in children under 5 years of age. In developing countries, Shigellosis causes 1.1 million of deaths per year. This is a serious disease that is spread via the fecal-oral route and is highly transmissible. Potential groups that would benefit from immunization against *Shigella* species include, for example, children, travelers and people in refugee camps.

There are four different serogroups of *Shigella*, namely, *S. dysenteriae*, *S. flexneri*, *S. sonnei* and *S. boydii*. In embodiments of the present invention, immunogenic bioconjugates can be made against each of these different serogroups of *Shigella*. For example, Figure 14B provides different serotypes of *Shigella* and the polysaccharide structure that defines their antigenicity (i.e., *Shigella* O-antigens).

In further embodiments of the present invention, immunogenic LPS bioconjugates could be made against other bacteria using the teachings in this specification, including bacteria: (1) that cause nosocomial infections, such as *Pseudomonas aeruginosa*; and (2) that cause urinary tract infection, such as *Extraintestinal E. coli (ExPEC)*.

In an embodiment, the inventors have developed a *Shigella dysenteriae* O1 LPS bioconjugate vaccine (also referred to as a *S. dysenteriae* bioconjugate), using genetically engineered *E. coli* with simple fermentation and purification methods. Figure 9 shows production of *Shigella* bioconjugates. The top panel shows the synthesis of bioconjugates in *E. coli*. In an embodiment, the O-antigen repeating unit of *S. dysenteriae* O1 is assembled on the carrier lipid undecaprenyl pyrophosphate (UPP), flipped to the periplasmic space and polymerized. The structure of *S. dysenteriae* O1 is as follows and is also provided in the middle right of Figure 9:



PglB transfers the activated polysaccharide to Asn residues of protein carriers, forming the *Shigella* bioconjugates. The protein carrier can be, for example, AcrA or a protein carrier that has been modifed to contain the consensus sequence for protein glycosylation, i.e., D/E – X – N – Z – S/T, wherein X and Z can be any amino acid except proline (e.g., a modified Exotoxin *Pseudomonas aeruginosa* (EPA)). EPA has been used successfully in conjugate vaccines.

In an embodiment illustrated in Figure 9, periplasmic proteins of *E. coli* cells expressing the modified EPA in the presence of PglB and the O1 polysaccharide cluster were separated by SDS page and, after transfer to nitrocellulose, EPA was immunodetected with an antiserum that was raised against EPA (lane 2). In lane 1, periplasmic proteins of *E. coli* cells expressing the *Campylobacter* protein AcrA in the presence of PglB and the O1 polysaccharide cluster were separated and immunodetected with an antiserum that was raised against AcrA. Both proteins were glycosylated with the O1-polysaccharide cluster. In the lowest panel of Figure 9, the LPS from *E. coli* were separated by SDS-PAGE and visualized by Silver Staining. The left lane depicts the LPS extracted from a strain not expressing the WaaL, whereas the right lane shows the typical O1 LPS pattern. Both strains are expressing the polysaccharide biosynthesis cluster of *S. dysenteriae* O1.

The production of a bacterial bioconjugate, such as a *Shigella* bioconjugate, is described in an embodiment in further detail with reference to Figures 10A and 10B. Prior to assembly of the bacterial bioconjugate using a bacterial system, such as *E. coli*, it is necessary to introduce into the bacterial system certain genetic sequences coding for the various enzymes and proteins to be used, as discussed earlier with reference to Figures 8A and 8B. For example, this includes an OST/Otase, preferably from *C. jejuni* (e.g., PglB), a protein carrier (e.g. EPA) and a gene cluster directed to antigenic polysaccharide synthesis (e.g., the gene cluster for *S. dysenteriae* O1 polysaccharide synthesis).

Figure 10A shows Step 1 in the development of a bacterial bioconjugate, namely, the biosynthesis of a polysaccharide, such as the O-antigen of *S. dysenteriae* Serotype O1. In this step, the antigen is synthesized on the carrier lipid undecaprenyl pyrophosphate (UPP), and then transferred into the periplasm using a flippase. The antigen is polymerized by the polymerase Wzy and transferred to the lipid A core by the ligase WaaL. To transfer the polysaccharide to a protein carrier, the ligase is replaced by the oligosaccharyltransferase; PglB.

Step 2 in the production of a bacterial bioconjugate involves engineering a suitable protein carrier. Protein carriers that are useful preferably should have certain immunological and pharmacological features. From an immunological perspective, preferably, a protein carrier should: (1) have T-cell epitopes; (2) be capable of delivering an antigen to antigen presenting cells (APCs) in the immune system; (3) be potent and durable; and (4) be capable of generating an antigen-specific systemic IgG response. From a pharmacological perspective, preferably, a protein carrier should: (1) be non-toxic; and (2) be capable of delivering antigens efficiently across intact epithelial barriers. More preferably, in addition to these immunological and pharmacological features, a protein carrier suitable for the production of a bacterial bioconjugate should: (1) be easily secreted into the periplasmic space; and (2) be capable having antigen epitopes readily introduced as loops or linear sequences into it.

The inventors have found genetically detoxified *Pseudomonas aeruginosa* Exotoxin (EPA) and the *Campylobacter* protein AcrA to be suitable protein carriers, most preferably EPA. AcrA contains natural glycosylation sites whereas EPA needs to be modified to encode

glycosylation sites. Preferably, EPA is modifed to introduce two glycosylation sites directed to the *Shigella* O1 antigen. More preferably, two consensus sequences are introduced as discussed in Example 10.

The amino acid sequence of EPA, as modified in an embodiment of this invention to contain two glycosylation sites, is provided as SEQ. ID NO.: 6 (with signal sequence) and SEQ. ID NO.: 7 (without signal sequence) in the Sequence Listing provided below. The glycosylation sites in each of SEQ. ID NO.: 6 and SEQ. ID NO.: 7 are denoted with an underline.

Figure 10B shows a schematic of a carrier protein, such as EPA onto which N-glycosylation sites can be designed as Step 3 in the production of a bacterial bioconjugate. N-glycosylation sites require introduction of the consensus sequences discussed previously, namely, insertion of D/E – X – N – Z-S/T sequons, wherein X and Z may be any natural amino acid except proline. We have found that such consensus sequences preferably are introduced through surface loops, by insertion rather than mutation and considering using flanking residues to optimize the operation of the N-glycosylation site.

Figure 11 shows bicoonjugates that elicit an immune response against *Shigella* O1 polysaccharide in mice. O1-AcrA and O1-EPA was purified by affinity column and anionic exchange. The pure bioconjugate was injected into mice (n=10). Serum of mice that were immunized with O1-AcrA (top) or O1-EPA (bottom) three times (day 1, 21, 60) was pooled and analyzed by ELISA at day 70 for a sugar specific IgG response. The plates (Nunc, polysorb) were coated with LPS isolated from *S. dysenteriae* O1 and incubated with the serum and anti mouse Polyvalent-HRP. Mice that received either conjugate developed an IgG response against the polysaccharide, confirming the presence of T-cell epitopes on the two protein carriers.

Consequently, the bacterial bioconjugates of the present invention show *in vivo* immnogenicity. In an embodiment, bacterial bioconjugates are capable of exhibiting: (1) a carbohydrate specific response; and (2) a carrier specific response or a similar response irrespective of the carrier protein. Moreover, an IgG specific response shows T-cell dependency of the immune response, such that memory of the response is expected.

Figure 12 reflects production of a *Shigella* O1 bioconjugate, e.g., O1-EPA, in a bioreactor. *E. coli* cells expressing EPA, PgIB and the O1-polysaccharide were grown in a bioreactor to OD_{600} = 40 by two nutrient pulses. Expression of PgIB and EPA was induced and the cells were grown overnight by linear feed of nutrients. The growth curve is depicted in the top panel. Whole cell extracts were separated by SDS-PAGE and expression and glycosylation of EPA was analyzed by immunodetection using a polyclonal antierserum that was raised against EPA (bottom). The cells efficiently glycosylate EPA at high cell density. The process is reproducible and leads to a total optical density (OD) of 90, which is a 45-fold increase compared to the shake flask culture. Consequently, scale-up is possible using the fed-batch process.

Figure 13 shows an example of purification of O1-EPA. More specifically, Figure 13 shows the fractionation and chromatographic purification of *S. dysenteriae* O1 bioconjugate. *E. coli* cells expressing the O1-EPA were grown in the bioreactor to high cell density (See Figure 12). The cells were pelleted by centrifugation and periplasmic proteins were extracted by osmotic shock. Periplasmic proteins were separated by anionic exchange (Source Q). Fractions enriched for O1-EPA were further purified by a second column (Fluoroapatite). The different fractions were separated by SDS-PAGE and the proteins were visualized by Coomassie Blue. Lane 1 shows whole cells extracts, lane 2 periplasmic proteins after osmotic shock, lane 3 periplasmic proteins loaded on anionic exchange, lane 4 and 5 eluates from anionic exchange and lane 6 O1-EPA eluate after the second purification column. This process allows the purification of O1-EPA at large scale. In this embodiment, the purification process is: (1) efficient (at > 10 mg/L culture); (2) possible at large scale; and (3) compatible with Good Manufacturing Practices (GMP). Following such purification, the EPA-O1 yield for glycerol-LB fed-batch was up to 200 mg/L, which is substantially higher than the yield for LB shake flask, which was 0.6 mg/L.

Figure 14A shows a Western blot analysis of a series of specimens of AcrA-*Shigella* O1 bioconjugates produced in an LB shake flask taken under various conditions, including pre-induction, 4 hours after induction, and 4 hours and 19 hours after induction under oxygen-limited circumstances. After extraction and purification, periplasmic proteins were separated by SDS-PAGE. AcrA and AcrA-*Shigella* O1 bioconjugates were detected using anti-AcrA antibody and

chemiluminescent detection via a secondary antibody. Loaded samples were normalized to culture OD_{600} at time of sampling.

In summary, in one aspect, the technology of the present invention has been used to develop a vaccine against *S. dysenteriae* O1 infection. For example, the polysaccharide of *S. dysenteriae* O1 can be conjugated to EPA in *E. coli.* This is very beneficial since EPA previously has been successfully used in clinical trials with different conjugate vaccines. In the instant invention, the *S. dysenteriae* O1 bioconjugate was produced in a bioreactor at 3l scale. The cells were grown to high OD and the bioconjugate was extracted by osmotic shock. The bioconjugates were purified to 98% purity by anionic exchange and size exclusion chromatography. The bioconjugates were injected into different mice strains. After two as well as three injections, a sugar specific IgG response against the polysaccharide was detected using LPS from *Shigella dysenteriae* O1 for analysis (Figure 11). As expected, IgM specific response was elicited when the LPS was injected. The bioconjugates raised a specific IgG response against the polysaccharide isolated from *S. dysenteriae*. IgG response against the corresponding sugar antigen, which was chemically coupled to a carrier protein, has been shown to correlate with protection in humans.

These results strongly suggest that our inventive *E. coli* strain is suitable for the potential production of an antigenic bacterial vaccine, such as an antigenic *Shigella* vaccine. In an embodiment, the EPA-*Shigella* bioconjugate was characterized intensively by different methods, like NMR, HPLC and MS. Figure 15 shows the expansion of the anomeric region of 1H NMR spectrum of an example of a *S. dysenteriae* Serotype O1 bioconjugate of the invention. The bioconjugate contains a sugar/protein ratio of 0.15, with 13.2 repeating units of the antigen being linked to the protein, and 1-2 sites being glycosylated. Two consensus sequences for glycosylation were introduced into EPA and about 20% of the protein is fully glycoslyated. The polysaccharide is linked via the reducing end to the protein carrier; therefore, the antigen epitopes of the polysaccharide are unmodified. In addition, the *in vivo* conjugation method attaches just the O-antigen repeating units to the protein, but no monosaccharides of the lipid A core are attached.

Using this technology, bacterial bioconjugates can be produced that are immunogenic. Genetic modifications can be made allowing *in vivo* conjugation of bacterial polysaccharides in desired proteins and at desired positions. For example, in an embodiment and as discussed above, the antigenic polysaccharide of *S. dysenteriae* O1 can be expressed in *E. coli* and conjugated to two different protein carriers *in vivo* (i.e., EPA and AcrA). Both bioconjugates elicit a specific IgG response against the polysaccharide in mice. As another example, Table 1 below depicts different polysaccharide substrates for bacterial OSTs/OTases such as PgIB that can be used in the *in vivo* method of the present invention for conjugating a protein carrier with the polysaccharide.

Table1






Table 2 below depicts yet additional different LPS polysaccharide substrates that could be utilized in the present invention with respect to various strains of *Shigella* and *E. coli*., as well as of *Pseudomonas aeruginosa* O11 and *Francisella tularensis*.

Table 2


Shigella dysenteriae O1




S. flexneri 2a




S. flexneri 3a




S. flexneri 3b




S. flexneri 6




S. sonnei




E. coli O4:K52 (ExPEC)




E. coli O4:K6 (ExPEC)




E. coli O6:K2 (ExPEC)




E. coli O6:K54 (ExPEC)



E. coli O22 (ExPEC)



E. coli O75 (ExPEC)



E. coli O83 (ExPEC)






Table 2 (continued)


E. coli O7




E. coli O9




E. coli O16




E. coli O121




E. coli O157 (EHEC)



Pseudomonas aeruginosa O11



Francisella tularensis



Qui4NFm, 4,6-dideoxy-4-formamido-D-glucose
GalNAcAN, 2-acetamido-2-deoxy-D-galacturonamide
QuiNAc, 2-acetamido-2,6-dideoxy-D-glucose

For example, in a further embodiment of the invention, bioconjugate vaccines against *E. coli* can also be developed. *E. coli* is a well-known bacterial species. From a genetic and clinical perspective, *E. coli* strains of biological significance to humans can be broadly categorized as commensal strains, intestinal pathogenic strains and extraintestinal pathogenic *E. coli* (ExPEC). ExPEC strains can be part of the normal intestinal flora and are isolated in 11% of healthy individuals. They do not cause gastroenteritis in humans but their main feature is their capacity to colonize extraintestinal sites and to induce infections in diverse organs or anatomical sites. They are the main cause of urinary tract infections (UTI), are involved in septicemia, diverse abdominal infections and meningitis. Bacteremia can arise with a risk of severe sepsis. Severe sepsis due to ExPEC was associated with 41,000 estimated deaths in 2001. ExPEC strains have been susceptible to antibiotics; however more and more antibiotic resistant strains have evolved, both in hospital and in the community. This antimicrobial resistance is making the management of ExPEC infections more difficult; therefore, new vaccines would be an alternative strategy to prevent these infections.

In animal models, passive or active immunization against capsule, O-specific antigen and different outer membrane proteins have afforded protection against systemic infections and immunization with these different antigens are protective against urinary tract infections from ExPEC strains expressing these virulence factors. The serotypes O4, O6, O14, O22, O75 and O83 cause 75% of UTI. In one embodiment, the novel technology of the present invention can be used to develop a monovalent LPS bioconjugate including one antigen (e.g., serotype O6, one of the major serotypes) and even a multivalent LPS bioconjugate including these 6 antigens. For example, the gene cluster encoding for the enzymes that synthesize the O-antigen for ExPEC could be amplified and then expressed in the Shigella production strain.

The instant invention involves a highly efficient production process with high potential yields that can be used for industrial scale preparations in a cost-efficient process. This novel, cost efficient bioengineering approach to producing bioconjugate can be applied to other conjugates and for other applications. An additional feature of the invention involves a considerable simplification of the production of bacterial vaccines with high reproducibility and a potentially reduced risk of lot failures.

Process for Manufacturing Conjugate Vaccine

It is now possible to engineer bacterial expression systems so that specific bioconjugates are produced that are biologically active. For example, the O-specific polysaccharide of *S. dysenteriae* has been conjugated to different protein carriers and the resulting bioconjugate has elicited a specific IgG response against the polysaccharide in mice. In an embodiment, the technology of the invention makes use of an oligosaccharyl transferase, for example, PglB of *Campylobacter jejuni* to couple bacterial polysaccharides (O antigens) *in vivo* to simultaneously express recombinant carrier proteins, yielding highly immunogenic bioconjugate vaccines.

A production process has been established that can be used on an industrial scale. This opens up the possibility that a multitude of various conjugate vaccines can be developed and manufactured using simple bacterial fermentation. The process has several advantages compared to the *in vitro* conjugation method depicted in the top panel of Figure 6A. As it is a complete *in vivo* process, the cost and risk of failures are reduced significantly and the process is more reproducible. In addition, the consensus capture sequence allows the conjugation of polysaccharides to defined proteins at specific built-in sites, thereby facilitating regulatory acceptance and quality control. Finally, the development of conjugate vaccines is much faster since the process is simplified and requires only biotechnology tools. In addition, the *in vivo* conjugation process is suited for application where polysaccharide compositions prevent chemical cross-linking.

In an embodiment, the instant invention relates to the scaled-up production of recombinant glycosylated proteins in bacteria and factors determining glycosylation efficiency. For example, recombinant glycosylated proteins of the present invention can be made using the shakeflask process. Bioconjugates have previously been mainly produced in LB shake flask cultures. More preferably, in one aspect of the invention, a first fed-batch process can be used for the production of recombinant glycosylated proteins in bacteria. In a preferred manufacturing process, the aim is to achieve markedly increased final biomass concentrations while maintaining glycosylation efficiency and recombinant protein yield per cell and while maintaining simplicity and reproducibility in the process.

In one embodiment, bacterial bioconjugates of the present invention can be manufactured on a commercial scale by developing an optimized manufacturing method using typical *E. coli* production processes. First, one can use various types of feed strategies, such as batch, chemostat and fed-batch. Second, one can use a process that requires oxygen supply and one that does not require an oxygen supply. Third, one can vary the manner in which the induction occurs in the system to allow for maximum yield of product.

It has found been that, in contrast to expression of the carrier protein, the degree of N-linked glycosylation strongly reacts to changes in nutrient availability, type of carbon- and energy source, oxygen supply and time-point of induction. For example, in a fed-batch process, the addition of inducers to the batch and fed-batch cultures immediately leads to a 3-fold decrease in specific growth rate, indicating a high metabolic burden and/or stress due to synthesis of the carrier protein and membrane-bound oligosaccharyltransferase. Based on the inventors' finding of a recurring retardation of the appearance of glycosylated carrier protein compared to the non-glycosylated form after induction, it is concluded that glycosylation appears to be the rate-limiting step in bioconjugate biosynthesis.

Based on these results, in an example of an embodiment of the invention, the following process design for cultivation has been developed: fed-batch cultivation mode for achieving high cell densities; extended incubation after induction to facilitate maximal glycosylation; surplus nutrient supply (e.g., LB components yeast extract and tryptone) during biomass build-up until induction to provide a sufficient supply of building blocks for the production process; and glycerol as the main carbon and energy source to prevent catabolite repression and acetate formation. This bioprocess allows a 50-fold increase in yield compared to LB batch culture, paving the way towards a cost-effective production of conjugate vaccines in recombinant *Escherichia coli*. In this example, one can have oxic conditions throughout the production process, for example, achieved through oxygen-enriched aeration; however, low oxygen content is also feasible. Example 9 sets forth this example of a fed-batch process in greater detail. It should be recognized, however, that other processes may be used to produce the bacterial LPS bioconjugates of the present invention.

Consequently, in one embodiment of the invention, *E. coli* can be used for *in vivo* production of glycosylated proteins and is suitable for industrial production of glycosylated proteins.

The following examples serve to illustrate further the present invention and are not intended to limits its scope in any way.

Examples

Example 1: Selection of AcrA as model protein for optimizing N-glycosylation

To optimize the acceptor protein requirements for N-glycosylation detailed studies were performed on the *C. jejuni* glycoprotein AcrA (CjO367c). AcrA is a periplasmic lipoprotein of 350 amino acid residues. It has been shown that secretion to the periplasm but not lipid-anchoring is a prerequisite for glycosylation (Nita-Lazar et al., 2005, supra). The signal for export can either be the native AcrA signal sequence or the heterologous PelB signal when expressed in *E. coli*. Of the five potential λMinked glycosylation sequons (N117, N123, N147, N273, N274) the same two ones are used in *C. jejuni* and *E. coli* (N123 and N273 (Nita-Lazar et al., 2005, supra)). AcrA was chosen as model because it is the only periplasmic N-glycoprotein of *C. jejuni* for which detailed structural information is available. Recently, the crystal structure of an AcrA homologue, the MexA protein from the Gram-negative bacterium P. aeruginosa, was published (Higgins et al., (2004). Structure of the periplasmic component of a bacterial drug efflux pump. Proc. Natl. Acad. Sci. U S A 7Of$_1$ 9994-9999). Both proteins are members of the so-called periplasmic efflux pump proteins (PEP,(Johnson, J. M. and Church, G. M. (1999). Alignment and structure prediction of divergent protein families: periplasmic and outer membrane proteins of bacterial efflux pumps. J. Mol. Biol. 287, 695-715)). The elongated molecule contains three linearly arranged subdomains: an α-helical, anti-parallel coiled-coil which is held together at the base by a lipoyl domain, which is followed by a six-stranded β-barrel domain. The 23-28 residues at the *N*-terminus and 95-101 residues in the C- terminus are unstructured in the crystals. MexA and AcrA protein sequences are 29.3% identical and 50 % similar. Thus, the two proteins likely exhibit a similar overall fold.

Example 2: Elucidation of the primary peptide sequence that triggers glycosylation

It is known that lipoyl domains similar to MexA of P. aeruginosa and accordingly also in AcrA of *C. jejuni* form a compact protein that can be individually expressed in *E. coli* (reviewed by Berg, A., and de Kok, A. (1997). 2-Oxo acid dehydrogenase multienzyme complexes. The central role of the lipoyl domain. Biol. Chem. 378, 617-634). To check which acceptor peptide sequence was required for N-glycosylation by the *pgl* machinery in *E. coli* the lipoyl domain of AcrA was taken. It was used as a molecular scaffold to transport peptides of different lengths to the periplasm and present them to the *pgl* machinery *in vivo*.

Therefore, a plasmid coding for the lipoyl domain (Lip) was constructed and N-terminally fused to the signal sequence of OmpA (Choi, J. H., and Lee, S. Y. (2004). Secretory and extracellular production of recombinant proteins using Escherichia coli. Appl Microbiol Biotechnol 64, 625-635) and C-terminally to a hexa histag (SEQ ID NO: 24). Cloning was performed to place the gene expression under the control of the arabinose promoter. For the Lip domain borders amino acid positions were chosen that appeared at the same positions as the domain borders of the Lipoyl domain part in MexA. To test different peptides for their ability to accept an N-glycan stretches of the sequence were inserted between the two hammerhead-like parts of the Lip domain. The stretches consisted of sequences comprising the N-glycosylation site N 123 of *C. jejuni* AcrA. The resulting open reading frames consisted of the sequences coding for the OmpA signal sequence, the N-terminal hammerhead-like part of AcrA (D60-D95, the numbering of the amino acids refers to the mature AcrA polypeptide sequence numbering), the different stretches containing the native N 123 glycosylation site of AcrA (see below), the C-terminal hammerhead-like part of AcrA-Lip (L167-D210) and the C-terminal his-tag.

Construction of the plasmids was achieved by standard molecular biology techniques. Three stretches containing the native N123 glycosylation site of AcrA of different lengths were inserted between the two halves of Lip resulting in three different ORFs:

Construct A contains A118-S130 resulting in a protein sequence of:

MKKTAIAIAVALAGFATVAQADVIIKPQVSGVIVNKLFKAGDKVKKGQTLFIIEQDQASK **DFNRS**KALFSQLDHTEIKAPFDGTIGDALVNIGDYVSASTTELVRVTNLNPIYADGS*HHH HHH* (SEQ. ID NO. 8).

Construct B contains F122-E138 resulting in a protein sequence of:

MKKTAIAIAVALAGFATVAQADVIIKPQVSGVIVNKLFKAGDKVKKGQTLFIIEQDQ**FN RSK**ALFSQSAISQKELDHTEIKAPFDGTIGDALVNIGDYVSASTTELVRVTNLNPIYADGS *HHHHHH* (SEQ. ID NO. 9).

Construct C contains D121-A127 resulting in a protein sequence of:

MKKTAIAIAVALAGFATVAQADVIIKPQVSGVIVNKLFKAGDKVKKGQTLFIIEQDQDF **NRS**KALDHTEIKAPFDGTIGDALVNIGDYVSASTTELVRVTNLNPIYADGS*HHHHHH* (SEQ. ID. NO. 10).

The underlined stretches of sequence indicate the OmpA signal peptide, singly underlined residues were introduced for cloning reasons or to render the protein resistant to degradation. Bold: glycosylation site corresponding to N 123 of AcrA. Italics: hexa-histag (SEQ ID NO: 24). The corresponding genes were expressed under the control of the arabinose promoter in the backbone of the plasmid pEC415 (Schulz, H., Hennecke, H., and Thony-Meyer, L. (1998). Prototype of a heme chaperone essential for cytochrome c maturation. Science 281, 1197-1200).

To check which of the three stretches triggered glycosylation of the Lip proteins protein expression experiments were performed. *E. coli* Top10 cells (Invitrogen, Carlsbad, CA, USA) carrying pACYC*pgl* or pACYC*pgl*mut (Wacker et al., 2002, *supra*) and a plasmid coding constructs A_1 B or C were grown in LB medium containing ampicillin and chloramphenicol up to an OD of 0.5 at 37^0C. For induction 1/1000 volume 20% arabinose (w/v) solution was added and the cells were grown for another 2 hrs. The cells were then harvested by centrifugation and resuspended in 20 mM Tris/HCI, pH 8.5, 20% sucrose (w/v), 1 mM EDTA, 1 mM PMSF, and 1 g/l (w/v) lysozyme and incubated at 4^0C for 1 hr. Periplasmic extracts were obtained after pelletting of the spheroblasts and diluted with 1/9 volume (v/v) of 10x buffer A (3 M NaCI, 0.5 M Tris/HCI, pH 8.0 and 0.1 M imidazole) and $MgSO_4$ added to 2.5 mM. Ni-affinity purification

was performed on 1 ml Ni-Sepharose columns from Amersham Pharmacia Biotech (Uppsala, Sweden) in buffer A. Proteins were eluted in buffer A containing 0.25 M imidazole.

Fig. 1 shows Coomassie brilliant blue stained SDS-PAGE gel of the peak elution fractions from the Ni-purified periplasmic extracts. The expression analysis showed that construct B produced a prominent single protein species (Fig. 1 , lane 1). Constructs A and C both lead, in addition to the prominent protein, to a second protein band with slower electrophoretic mobility (Fig. 1 , lanes 2 and 3). That the heavier protein species was indeed glycosylated was proven by MALDI-TOF/TOF (not shown). The only amino acid missing in construct B but present in A and C was D121 , the aspartate residue 2 positions N-terminally to the glycosylated N123. This demonstrates that D121 plays an important role for glycosylation by the OTase. To verify that D121 is essential for glycosylation it was mutated to alanine in construct C. Expression analysis resulted in only one protein band (Figure 1, lane 4), thus showing that D121 is important for glycosylation. Furthermore, the fact that an artificial peptide display protein can be glycosylated shows that a short peptide of the D/E-X-N-Y-S/T type contains all information for *C. jejuni*-borne N-glycosylation to occur.

Example 3: Verification of Example 2; AcrA-D121A is not glycosylated at N 123

To confirm the findings from the peptide display approach an aspartate to alanine mutation was inserted at position 121 (D121A, i.e. 2 residues before the glycosylated N 123) in the full length soluble version of the AcrA protein and it was tested whether the site N123 could still be glycosylated in *E. coli*. In order to test this AcrA-D121A was expressed and its glycosylation status was analyzed. For the analysis an engineered AcrA was used. It differed from the original *C. jejuni* gene in that it contains the PelB signal sequence (Choi and Lee, 2004, supra) for secretion into the periplasm and a C- terminal hexa histag (SEQ ID NO: 24) for purification. It has been shown that this AcrA variant gets secreted, signal peptide-cleaved and glycosylated as the lipid anchored, native protein (Nita-Lazar et al., 2005, supra). The following is the amino acid sequence of the soluble AcrA protein:

MKYLLPTAAAGLLLLAAQPAMAMHMSKEEAPKIQMPPQPVTTMSAKSEDLPLSFTYPA
KLVSDYDVIIKPQVSGVIVNKLFKAGDKVKKGQTLFIIEQDKFKASVDSAYGQALMAKA
TFENASK**DFNRS**KALFSKSAISQKEYDSSLATFNNSKASLASARAQLANARIDLDHTEIK

APFDGTIGDALVNIGDYVSASTTELVRVTNLNPIYADFFISDTDKLNLVRNTQSGKWDLD
SIHANLNLNGETVQGKLYFIDSVIDANSGTVKAKAVF**DNNNS**TLLPGAFATITSEGFIQK
NGFKVPQIGVKQDQNDVYVLLVKNGKVEKSSVHISYQNNEYAIIDKGLQNGDKIILDNF
KKIQVGSEVKEIGAQLE*HHHHHH* (SEQ. ID NO. 11)

The underlined residues are the PelB signal peptide, italics the hexa-histag (SEQ ID NO: 24), and bold the two natural glycosylation sites at N 123 and N273. A plasmid containing the ORF for the above protein in the pEC415 plasmid (Schulz et al., 1998) was constructed to produce pAcrAper.

The assay to test the glycosylation status of AcrA and mutants thereof (see below) was as follows: expression of AcrA was induced with 0.02% arabinose in exponentially growing *E. coli* CLM24 (Feldman et al., 2005, *supra*) cells containing the plasmid-bome *pgl* operon in its active or inactive form (pACYCpg/ or pACYCpg/mut, see (Wacker et al., 2002, *supra*)) and a plasmid coding for AcrA (pAcrAper). After four hours of induction, periplasmic extracts were prepared as described above and analyzed by SDS-PAGE, electrotransfer and immunodetection with either anti-AcrA antiserum or R12 antiserum. The latter is specific for *C. jejuni* N-glycan containing proteins (Wacker et al., 2002, *supra*).

The first two lanes of Fig. 2A show AcrA in the absence and presence of a functional *pgl* operon. Only one band appears in the absence but three in the presence of the functional *pgl* operon (Fig. 2A, top panel). These correspond to unglycosylated AcrA (lane 1) and un-, mono- and diglycosylated AcrA (lane 2). That the two heavier proteins in lane 2 were glycosylated was confirmed by the R12 western blot (lane 2, bottom panel). When the mutant AcrA-N273Q was expressed the same way, only the monoglycosylated AcrA was detected in presence of the functional glycosylation *pgl* operon (lane 3). Unglycosylated AcrA was detected in absence of the functional *pgl* locus (lane 4). Analysis of the mutant AcrA-D121A produced only two bands, one of them glycosylated (lane 5) as observed with AcrA-N273Q in lane 3. This means that D121 is essential for efficient glycosylation at position 123-125.

Example 4: Introducing artificial glycosylation sites into AcrA

To test if the introduction of an aspartate residue could generate a glycosylation site, AcrA mutants were generated in which the residue in the -2 position of the not used glycosylation sites in positions N117 and N147 of soluble AcrA were exchanged for aspartate (F115D, T145D). It was then tested whether the modified glycosylation sites could be glycosylated by the same assay as described in example 2. Both mutations were individually inserted either into the wildtype sequence of the soluble version of AcrA or in the double mutant in which both used glycosylation sites were deleted (N123Q and N273Q). Periplasms extracts of cultures induced for 4 hrs were prepared, separated by SDS page and analyzed by Western blotting (Fig. 2B). As controls the samples of wildtype glycosylated and non glycosylated AcrA were run on the same gel (lanes 1 and 2). The T145D mutation affected the -2 position of the natively not used glycosylation sequon N147-S149. Upon expression of AcrA-T145D Western blotting with anti AcrA antiserum resulted in four bands, the highest of them with slower electrophoretic mobility than the doubly glycosylated protein in lane 2 (lane 3 in figure 2B). The R12 blot confirmed that the fourth band was a triply glycosylated AcrA. Despite the low intensity towards anti AcrA the heaviest band gave the strongest signal with the glycosylation specific R12 antiserum. When the same mutant AcrA-T145D was expressed in the absence of the native N-glycosylation sequence (AcrA-N123Q-N273Q-T145D), only monoglycosylated AcrA was detected in the presence of a functional *pgl* operon (Fig. 2B, lane 4), that was missing in absence of a functional *pgl* operon (lane 5). This demonstrates that the heavier band in lane 4 was glycosylated. Hence, by simply introducing the T145D mutation an optimized glycosylation site was generated (DFNNS) (SEQ ID NO: 27).

To further confirm that it is possible to introduce a glycosylation site by inserting an aspartate residue in the -2 position, the natively not used sites N117-S119 and N274-T276 were changed to optimize N-glycosylation. For this purpose further mutants were generated (Fig. 2C). Expression of AcrA-F115D-T145D in the above described system resulted in five protein species detected with the anti AcrA antiserum (lane 2). This is indicative for four glycosylates taking place on the same AcrA molecule. When the detection was performed with the *C. jejuni* N-glycan-specific R12 antiserum, a ladder of five bands was detected. The lowest faint band is unglycosylated AcrA because it is also present in the absence of glycosylation (lane 1), the highest results in a strong signal probably due to the five antigenic determinants in a fourfold glycosylated AcrA. Thus, the two introduced sites (at N117 and N 147) and the two natively

used sites (N123 and N273) are used and glycosylated by the *pgl* machinery. Expression of AcrA-N123Q-N273Q-N272D with and without the *pgl* operon demonstrated that a third artificially introduced glycosylation site, N274 (DNNST) (SEQ ID NO: 28), was also recognized by the *pgl* operon (Fig. 2C, lanes 3 and 4).

The above experiments confirm the finding that the bacterial N-glycosylation site recognized by the OTase of *C. jejuni* consists partly of the same consensus as the eukaryotic one (N - X - S/T, with X≠P) but, in addition, an aspartate in the -2 position is required for increasing efficiency. Furthermore, they demonstrate that it is possible to glycosylate a protein at a desired site by recombinantly introducing such an optimized consensus sequence.

Example 5: Verification of position -1 in the optimized N-glycosylation sequence

A further experiment was performed to test whether the -1 position in the bacterial glycosylation site exhibits the same restrictions as the +1 position in eukaryotes (Imperiali, B., and Shannon, K. L. (1991). Differences between Asn-Xaa-Thr-containing peptides: a comparison of solution conformation and substrate behaviour with oligosaccharyl-transferase. Biochemistry 30, 4374-4380; Rudd, P. M., and Dwek, R. A. (1997). Glycosylation: heterogeneity and the 3D structure of proteins. Crit. Rev. Biochem. Mol. Biol. 32, 1-100). A proline residue at +1 is thought to restrict the peptide in such a way that glycosylation is inhibited. To test if a similar effect could also be observed in the -1 position a proline residue was introduced at that position of the first natively used site in a point mutant that had the second native site knocked out (AcrA-N273Q-F122P). The control expression of AcrA-N273Q showed a monoglycosylated protein in the presence of a functional *pgl* operon (Fig. 2D, lane 1 and 2). However, AcrA-N273Q-F122P was not glycosylated (Fig. 2D, lanes 3 and 4). This indicates that proline inhibited bacterial N-glycosylation when it constitutes the residue between the asparagine and the negatively charged residue of the -2 position.

Sequence alignments of all the sites known to be glycosylated by the *C. jejuni pgl* machinery indicate that they all comprise a D or E in the -2 position (Nita-Lazar et al., 2005, supra; Wacker et al., 2002, supra; Young et al., (2002). Structure of the N-linked glycan present on multiple glycoproteins in the Gram-negative bacterium, Campylobacter jejuni. J. Biol. Chem. 277, 42530-42539). Thus, it was established that the glycosylation consensus sequence for

bacteria can be optimized by a negatively charged amino acid in the -2 position, resulting in D/E - X - N - Z - S/T, wherein X & Z ≠ P.

Example 6: N-glycosylation of a non-*C. jejuni* protein

To demonstrate that the primary sequence requirement (optimized consensus sequence) is sufficient for N-glycosylation in bacteria, it was tested whether a non-*C. jejuni* protein could be glycosylated by applying the above strategy. Cholera toxin B subunit (CtxB) was employed as a glycosylation target. The corresponding gene was amplified from *Vibrio cholerae* in such a way that it contained the coding sequence of the OmpA signal sequence on the N-terminus and a hexahistag (SEQ ID NO: 24) at the C-terminus, just the same as constructs A through C in example 1. The resulting DNA was cloned to replace construct A in the plasmids employed in example 1. A point mutation of W88 to D or a D insertion after W88 generated an optimized glycosylation site (DNNKT) (SEQ ID NO: 29). The wildtype and W88D CtxB proteins containing the signal sequence and his-tag were expressed in *E. coli* Top 10 and other cell types in the presence and absence of the functional *pgl* locus from *C. jejuni*. When periplasmic extracts from Top10 cells were analyzed by SDS-PAGE, electrotransfer and consecutive immunoblotting with a CtxB antiserum, only CtxB W88D produced a higher and thus glycosylated band in the *pgl* locus background (Fig. 2E, compare lanes 3 and 4). A consensus sequence (DSNIT) (SEQ ID NO: 25) was also inserted by replacing G54 or Q56 of CtxB (the latter is denoted CtxB-Q56/DSNIT) (SEQ ID NO: 25), i.e. in one of the loops that was reported to contribute to the ganglioside GM 1 binding activity of CtxB. Lanes 5 and 6 of Figure 2E demonstrate that the engineered protein (exemplified by the construct which contains the peptide sequence DSNIT (SEQ ID NO: 25) instead of Q56 expressed in Top10 cells) produced a lower mobility and thus glycosylated band in glycosylation competent but not glycosylation-deficient cells when analyzed in the same way as described above. It was also demonstrated that a CtxB containing two manipulations, i.e. the insertion of D after W88 as well as DSNIT (SEQ ID NO: 25) replacing Q56, was double-glycosylated in SCM7 cells (Alaimo et al., EMBO Journal 25: 967-976 (2006)) (panel E, lanes 7 and 8). The double- glycosylated protein CtxB shown in lane 7 was Ni^{2+} affinity-purified and analyzed by ESI- MS/MS after in-gel trypsinization according to standard protocols. The expected glycopeptides were detected confirming that bacterial N-glycosylation can also be directed to a non-*C. jejuni* protein by mutating or inserting the

optimized consensus sequence according to the invention for bacterial N-glycosylation (not shown). Examples of other suitable exemplary *E. coli* strains for practicing the present invention are W3110, CLM24, BL21 (Stratagene, La JoIla, CA, USA), SCM6 and SCM7.

The amino acid sequence of the CtxB protein used here is indicated below (recombinant OmpA signal sequence underlined, hexa-histag (SEQ ID NO: 24) italics, W88 bold):

MKKTAIAIAVALAGFATVAQATPQNITDLCAEYHNTQIHTLNDKIFSYTESLAGKREMAI ITFKNGATFQVEVPGSQHIDSQKKAIERMKDTLRIAYLTEAKVEKLCVWNNKTPHAIAAI SMANGS*HHHHHH* (SEQ. ID NO. 12)

Example 7: Introduction of artificial N-glycosylation sites into the *C. jejuni* outer membrane protein, OmpH1

A potential application of the N-glycosylation in bacteria is the display of the glycan on the surface of a bacterial host cell in order to link the pheno- to the genotype and thereby select for specific genetic mutations. To demonstrate that N-glycans can be presented on outer membrane proteins the OmpH1 protein was engineered in a way that it contained multiple optimized consensus sites according to the invention. The sites were engineered into loop regions of the protein as deduced from the known crystal structure (Muller, A., Thomas, G. H., Horler, R., Brannigan, J.A., Blagova, E., Levdikov, V.M., Fogg, M.J., Wilson, K.S., and Wilkinson, A.J. 2005. An ATP-binding cassette-type cysteine transporter in *Campylobacter* jejuni inferred from the structure of an extracytoplasmic solute receptor protein. Mol. Microbiol. 57: 143-155). Previous experiments showed that the best glycosylation sequons were generated by the mutations V83T, K59N-G601- N61T, R190N-G191I-D192T and H263D-F264S-G265N-D266I-D267T. For surface display it was desired to evaluate different combinations of those introduced sites in order to establish the most N-glycan-specific sample. The combinations were generated in a wild type OmpH1 encoding plasmid construct and tested in a similar manner as described for AcrA. Figure 3 shows the analysis of various OmpH1 variants harboring multiple glycosylation sequons in addition to the existing wild type sequon. OmpH1 variants were generated with three (lane 3, 4, 5 and 7) and four glycosylation sequons (lane 6). A wild type OmpH1 with only one glycosylation sequon and a mutant lacking the critical asparagine for glycosylation were also included in the experiment. All variants tested here did not only

demonstrate a high level of glycosylation efficiency but also that every glycosylation sequon was utilized. The results were confirmed with *Campylobacter N*-glycan specific immuneserum (Fig. 3 lower panel).

The following is the amino acid sequence of the OmpH1 protein of *Campylobacter* jejuni (strain 81-176) with attached myc tag in italics:

MKKILLSVLTTFVAVVLAACGGNSDSKTLNSLDKIKQNGWRIGVFGDKPPFGYVDEKG
NNQGYDIALAKRIAKELFGDENKVQFVLVEAANRVEFLKSNKVDIILANFTQTPERAEQ
VDFCLPYMKVALGVAVPK**DSNIT**SVEDLKDKTLLLNKGTTADAYFTQDYPNIKTLKYD
QNTETFAALMDKRGDALSHDNTLLFAWVKDHPDFKMGIKELGNKDVIAPAVKKGDKE
LKEFIDNLIIKLGQEQFFHKAYDETLKAHFGDDVKADDWIEGGKIL*EQKLISEEDL* (SEQ. ID NO. 13)

The native glycosylation site in the protein is bold, the signal sequence underlined.

Example 8: Surface display of N-glycans from *C. jejuni* on OmpH1 on the outer membrane of *E. coli* cells

In order to answer the question whether multiple glycosylated OmpH1 variants can be displayed on the surface of bacterial cells, immunofluorescence was performed on bacterial CLM24 or SCM6 (which is SCM7 ΔwaaL) cells expressing various OmpH1 variants. A wild type OmpH1 and a mutant lacking the critical asparagine for glycosylation were included in the experiment. In addition, a C20S mutant was constructed in order to retain the protein in the periplasm, thus serving as a control in the experiment. Immunostaining was carried out on the cells treated with paraformaldehyde. Paraformaldehyde fixes cells without destroying the cell structure or compartimentalization. The c-Myc- and *N*-glycan- specific immuneserum in combination with corresponding secondary antibodies conjugated to FITC and Cy3 were used to detect the protein (red fluorescence) and *N*-glycan (green) on the bacterial cell surface, respectively. Additionally, 4,6-diamino-2-phenylindole (DAPI, blue) was employed to stain for bacterial DNA to unambiguously differentiate between bacterial cells and cellular debris. When the cells expressing wild type OmpH1 were stained, immunofluorescence specific to the protein as well as the *N*-glycan was detected (Fig. 4 A). When a mutant lacking the critical asparagine

N139S was stained with both anti-Myc- and *N*-glycan- specific immuneserum only the protein but not glycan specific signals were obtained (panel 4 B) indicating specificity of the *N*-glycan-specific immune serum. When the protein was retained within the periplasm as in the C20S mutant, no protein specific, red immunofluorescence was detected indicating that the antibodies were unable to diffuse within the cell and were competent enough to detect any surface phenomenon (panel 4 C). Next, cells expressing multiple OmpH1 variants different in glycosylation were stained: OmpH1 $^{KGN\rightarrow NIT,,HFGDD\rightarrow DSNIT}$ (panel 4 D), OmpH1$^{RGD\rightarrow NIT,HFGDD\rightarrow DSNIT}$ (panel 4 E), OmpH1$^{KGN\rightarrow NIT,RGD\rightarrow NIT}$ (panel 4 F), OmpH1$^{V83T,KGN\rightarrow NIT}$ (panel 4 G) and OmpH1 $^{KGN\rightarrow NIT,RGD\rightarrow NIT,HFGDD\rightarrow DSNIT}$ (panel 4 H). 'HFGDD' and 'DSNIT' disclosed as SEQ ID NOS 26 and 25, respectively. All the OmpH1 variants were double-stained indicating the presence of glycosylated protein on the bacterial surface. Figure 4 is represented in grayscale, the first column is a merge picture of the other pictures of the same row.

Figure 4 shows fluorescence microscopy of cells expressing various OmpH1 variants. Cultures of *E. coli* strains CLM24 or SCM6 containing the expression plasmid for the wild type OmpH1 and its variants were equalized to OD_{600} of 0.25/ml. Cells were washed two times with phosphate-buffered saline (PBS), pH 7.4 and 100 µl cell suspensions was dropped onto gelatinized glass slides and incubated at room temperature (RT) for 30 min inside a humidified chamber. All subsequent steps in the whole-cell immunofluorescence labeling were done at room temperature inside a humidified chamber. The unbound cells were removed and rest was fixed with 4% paraformaldehyde containing PBS for 30 min at RT. Importantly, paraformaldehyde is considered not to permeabilize cells but keeping the compartimentalization by membranes intact. Fixed cells were washed two times with PBS and resuspended blocking buffer containing 5% BSA in PBS. After blocking, the cells were incubated with anti-*myc* monoclonal mouse IgG (1:50, Calbiochem) and/or anti-glycan antiserum (1:4000) for 1 h in 100 µl of PBS containing 5% BSA. The cells were washed three times with 100 µl of PBS for 5 min each and incubated with secondary anti-rabbit antibody conjugated to FITC (1:250, Jackson Immunoresearch Laboratories) and/or anti-mouse antibody conjugated to Cy3 (1:250, Jackson Immunoresearch Laboratories) for 1 h in 100 µl of PBS containing 5% BSA. If required, 4, 6-diamino-2-phenylindole (DAPI) (Sigma) (0.5 µg/ml) was added at the time of secondary antibody incubation to stain for bacterial DNA. The secondary antibody was rinsed from the

cells PBS_I and coverslips were mounted on slides by using vectashield (Vector Laboratories) mounting medium and sealed with nail polish. Fluorescence microscopy was performed by the using an Axioplan2 microscope (Carl Zeiss). Images were combined by using Adobe Photoshop, version CS2. SCM6 cells expressing OmpH1 (panel A), $OmpH1^{N139S}$ (panel B), $OmpH1^{C20S}$ (panel C), $OmpH1^{KGN\rightarrow NIT,\ HFGDD\rightarrow DSNIT}$ (panel D), $OmpH1^{RGD\rightarrow NIT,HFGDD\rightarrow DSNIT}$ (panel E), $OmpH1^{KGN\rightarrow NIT\ RGD\rightarrow NIT}$ (panel F), $OmpH1^{V83T,KGN\rightarrow NIT}$ (panel G), and $OmpH1^{KGN\rightarrow NIT,RGD\rightarrow NIT,HFGDD\rightarrow DSNIT}$ (panel H). 'HFGDD' and 'DSNIT' disclosed as SEQ ID NOS 26 and 25, respectively. The first column is a merge of the pictures in columns 2, 3, and 4 represented in greytones on black background. Column 2: blue fluorescence in greytones from DAPI stain, column 3: green fluorescence from glycan specific fluorescence, column 4: red fluorescence from anti-myc staining.

Example 9: An Example of a Production Process for *Shigella* O1 LPS Bioconjugate

This is an example of a production process; however, different conditions also lead to similar product formation.

A. Production process

E. coli strain W3110ΔwaaL containing three plasmids expressing PglB, EPA and the enzymes for the biosynthesis of the Shigella O1 polysaccharide was used for the production of the LPS bioconjugate. A single colony was inoculated in 50 ml LB medium and grown at 37°C O/N. The culture was used to inoculate a 1l culture in a 2l bioreactor. The bioreactor was stirred with 500 rpm, pH was kept at 7.0 by auto-controlled addition of either 2 M KOH or 20% H_3PO_4 and the cultivation temperature was set at 37°C. The level of dissolved oxygen (pO2) was kept between 0 and 10% oxygen. The cells were grown in a semi defined glycerol medium containing Kanamycin to an OD_{600} = 15. The medium contained the following ingredients:
330 mM Glycerol, 10 g Yeast extract, 20 g Tryptone, 34 mM K_2HPO_4, 22 mM KH_2PO_4, 38 mM $(NH_4)_2SO_4$, 2 mM $MgSO_4 \cdot 7\ H_2O$ and 5mM Citric acid. After an initial batch phase around 5h, a first nutrient pulse was added to sustain fast biomass build-up (glycerol, tryptone and yeast extract). After an additional 1.5 h the culture reached an OD_{600} = 30. At this timepoint a second nutrient pulse of glycerol and tryptone was added together with the required inducers 1% L-arabinose and 1 mM IPTG. In order to keep induction at maximum levels and supply further

amino acids for recombinant protein synthesis, a linear nutrient/inducer feed (28.8ml/h) was started with the addition this pulse. The feed was sustained until the end of the process. The bioreactor culture was harvested after a total of ≈ 24 h cultivation, when it should have reached an OD600 of ±80.

The production process was analyzed by Western blot as described previously (Wacker, M., *et al.*, N-linked glycosylation in *Campylobacter jejuni* and its functional transfer into *E. coli*. Science, 2002. 298(5599): p. 1790-3.). After being blotted on nitrocellulose membrane, the sample was immunostained with the specific anti-EPA (Wacker, M., *et al.*, N-linked glycosylation in *Campylobacter jejuni* and its functional transfer into *E. coli*. Science, 2002. 298(5599): p. 1790-3.). Anti-rabbit IgG-HRP (Biorad) was used as secondary antibody. Detection was carried out with ECL™ Western Blotting Detection Reagents (Amersham Biosciences, Little Chalfont Buchinghamshire).

Figure 16A shows proteins extracted of the *Shigella* O1 LPS Bioconjugate (i.e., EPA-O1) from a fed-batch process that were normalized to biomass concentration (0.1 OD_{600nm} of cells/lane). The proteins were separated by SDS-PAGE transferred to Nitrocellulose membrane and visualized by rabbit anti EPA antibody. The induction time for PglB and EPA expression was 1 h and O/N.

B. Periplasmic protein extraction

The cells were harvested by centrifugation for 20 min at 10,000 g and resuspended in 1 volume 0.9% NaCl. The cells were pelleted by centrifugation during 25-30 min at 7,000 g. The cells were resuspended in Suspension Buffer (25% Sucrose, 100 mM EDTA 200 mM Tris HCl pH 8.5, 250 OD/ml) and the suspension was incubated under stirring at 4-8°C during 30 min. The suspension was centrifuged at 4-8°C during 30 min at 7,000-10,000 g. The supernatant was discarded, the cells were resuspended in the same volume ice cold 20 mM Tris HCl pH 8.5 and incubated under stirring at 4-8°C during 30 min. The spheroblasts were centrifuged at 4-8°C during 25-30 min at 10,000 g, the supernatant was collected and passed through a 0.2 μ membrane.

As shown in Figure 16B, the periplasmic extract was loaded on a 7.5% SDS-PAGE, and stained with Coomasie to identify EPA and EPA-O1. EPA is a thick band that runs above the 70kDa marker. O1-EPA (i.e., EPA-O1) runs as a leader between 100 and 170 kDa.

C. Bioconjugate Purification

The supernatant containing periplasmic proteins obtained from 100,000 OD of cells was loaded on a Source Q anionic exchange column (XK 26/40 ≈ 180 ml bed material) equilibrated with buffer A(20 mM Tris HCl pH 8.0). After washing with 5 column volumes (CV) buffer A, the proteins were eluted with a linear gradient of 15CV to 50% buffer B (20mM Tris HCl + 1M NaCl pH 8.0) and then 2CV to 100% buffer B. Protein were analyzed by SDS-PAGE and stained by Coomassie. Fractions containing O1-EPA were pooled. Normally the bioconjugate eluted at conductivity between 6-17 mS. The sample was concentrated 10 times and the buffer was exchanged to 20 mM Tris HCl pH 8.0.

As shown in Figure 17A, protein fractions from 1. Source Q were analyzed by SDS-PAGE and stained by Coomassie. Fractions C1 to G9 contained O1 bioconjugate and were pooled.

The O1-Bioconjugate was loaded a second time on a Source Q column (XK 16/ 20 ≈ 28 ml bed material) that has been equilibrated with buffer A: 20 mM Tris HCl pH 8.0. The identical gradient that was used above was used to elute the bioconjugate. Protein were analyzed by SDS-PAGE and stained by Coomassie. Fractions containing O1-EPA were pooled. Normally the bioconjugate eluted at conductivity between 6-17 mS. The sample was concentrated 10 times and the buffer was exchanged to 20 mM Tris HCl pH 8.0.

As shown in Figure 17B, protein fractions from 2. Source Q column were analyzed on SDS-PAGE and stained by Coomassie. Fractions A11 to B3 containing O1 bioconjugate were pooled.

The O1-Biconjugate was loaded on Superdex 200 (Hi Load 26/60, prep grade) that was equilibrated with 20 mM Tris HCl pH 8.0.

As shown in Figure 18A, protein fractions from Superdex 200 column were analyzed by SDS-PAGE and stained by Coomassie stained. Fractions F1 to F11 were pooled.

As shown in Figure 18B, *Shigella* bioconjugate from different purification steps were analyzed by SDS-PAGE and stained by Coomassie. O1-EPA was purified to more than 98% purity using the process, showing that O1-EPA bioconjugate can be successfully produced using this technology.

Example 10: Engineering of Exotoxin A of *Pseudomonas aeruginosa* for glycosylation with antigenic carbohydrates

Exotoxin A of *Pseudomonas aeruginosa* (EPA) is a 67 kDa extracellulary secreted protein encoding mature 613 amino acids in its mature form and containing four disulfide bridges (C11-C15, C197-C214, C265-C287, C372-C379). To enable its glycosylation in *E. coli*, the protein must locate to the periplasmic space for glycosylation to occur. Therefore, a the signal peptide of the protein DsbA from *E. coli* was genetically fused to the N-terminus of the mature EPA sequence. A plasmid derived from pEC415 [Schulz, H., Hennecke, H., and Thony-Meyer, L., Prototype of a heme chaperone essential for cytochrome c maturation, *Science, 281*, 1197-1200, 1998] containing the DsbA signal peptide code followed by a RNase sequence was digested (NdeI to EcoRI) to keep the DsbA signal and remove the RNase insert. EPA was amplified using PCR (forward oligo was 5'-AAGCTAGCGCCGCCGAGGAAGCCTTCGACC (SEQ. ID NO. 14) and reverse oligo was 5'-AAGAATTCTCAGTGGTGGTGGTGGTGGTGCTTCAGGTCCTCGCGCGGCGG (SEQ. ID NO. 15)) and digested NheI/EcoRI and ligated to replace the RNase sequence removed previously. The resulting construct (pGVXN69) encoded a protein product with an DsbA signal peptide, the mature EPA sequence and a hexa-histag (SEQ ID NO: 24). Detoxification was achieved by mutating/deleting the catalytically essential residues L552VΔE553 according to [Lukac, M., Pier, G.B., and Collier, R.J., Toxoid of Pseudomonas aeruginosa exotoxin A generated by deletion of an active-site residue, Infect Immun, 56, 3095-3098, 1988] and [Ho, M.M., et al., Preclinical laboratory evaluation of a bivalent Staphylococcus aureus saccharide-

exotoxin A protein conjugate vaccine, Hum Vaccin, 2, 89-98, 2006] using quick change mutagenesis (Stratagene) and phosphorylated oligonucleotides 5'-GAAGGCGGGCGCGTGACCATTCTCGGC (SEQ. ID NO. 16) and 5'-GCCGAGAATGGTCACGCGCCCGCCTTC (SEQ. ID NO. 17) resulting in construct pGVXN70.

It is known that insertion of a pentapeptide sequence of the type D/E-Z-N-X-S/T into a suitable position results in glycosylation. To glycosylate EPA in *E. coli* cells, two different glycosylation sites were inserted into the previously described constructs according to the following description.

To insert a site at position 375, two steps were performed. First, quick change mutagenesis using oligos 5'-CCTGACCTGCCCCGGGGAATGCGCGG (SEQ. ID NO. 18) and 5'- CCGCGCATTCCCCGGGGCAGGTCAGG (SEQ. ID NO. 19) with pGVXN70 as a template resulted in a construct containing a single SmaI site at amino acid position 375 of EPA protein sequence by deleting three residues but otherwise keeping the starting protein sequence intact. In a second step, an insert composed of two complementary, phosphorylated oligonucleotides coding for (i) the previously deleted residues (when inserting the SmaI site), (ii) the pentapeptide glycosylation sequon and (iii) additional lysine residues flanking the consensus for optimization of glycosylation efficiency (as was found by further experiments) was ligated into this SmaI site (5'- GTCGCCAAAGATCAAAATAGAACTAAA (SEQ. ID NO. 20) and 5'-TTTAGTTCTATTTTGATCTTTGGCGAC (SEQ. ID NO. 21). The resulting construct was pGVXN137.

To insert an additional glycosylation site in the construct at amino acid 240, a one step procedure using quick change mutagenesis with oligonucleotides 5'-CATGACCTGGACATCAAGGATAATAATAATTCTACTCCCACGGTCATCAGTCATC (SEQ. ID NO. 22) and 5'-GATGACTGATGACCGTGGGAGTAGAATTATTATTATCCTTGATGTCCAGGTCATG (SEQ. ID NO. 23) was applied on construct pGVXN137. The resulting construct thus contained

various changes compared to the wild type EPA protein: two glycosylation sites, a DsbA signal peptide, detoxification mutation.

While this invention has been particularly shown and described with references to embodiments thereof, it will be understood by those skilled in the art that various changes in form and detail may be made therein without departing from the scope of the invention encompassed by the claims. Moreover, in instances in the specification where specific nucleotide or amino acid sequences are noted, it will be understood that the present invention encompasses homologous sequences that still embody the same functionality as the noted sequences. Preferably, such sequences are at least 85% homologous. More preferably, such sequences are at least 90% homologous. Most preferably, such sequences are at least 95% homologous.

Sequence Listing

SEQ. ID NO. 1: NUCLEOTIDE SEQUENCE FOR PglB

LOCUS pglb 2169 bp DNA SYN 17-FEB-2009
DEFINITION
ACCESSION pglb
KEYWORDS .
SOURCE .
ORGANISM C. jejuni 81116

REFERENCE 1 (bases 1 to 2169)
AUTHORS Self
JOURNAL Unpublished.
BASE COUNT 650 a 264 c 386 g 869 t
ORIGIN

```
   1 ATGTTGAAAA AAGAGTATTT AAAAAACCCT TATTTAGTTT TGTTTGCGAT GATTATATTA
  61 GCTTATGTTT TTAGTGTATT TTGCAGGTTT TATTGGGTTT GGTGGGCAAG TGAGTTTAAT
 121 GAGTATTTTT TCAATAATCA GTTAATGATC ATTTCAAATG ATGGCTATGC TTTTGCTGAG
 181 GGCGCAAGAG ATATGATAGC AGGTTTTCAT CAGCCTAATG ATTTGAGTTA TTATGGATCT
 241 TCTTTATCCG CGCTTACTTA TTGGCTTTAT AAAATCACAC CTTTTTCTTT TGAAAGTATC
 301 ATTTTATATA TGAGTACTTT TTTATCTTCT TTGGTGGTGA TTCCTACTAT TTTGCTAGCT
 361 AACGAATACA AACGTCCTTT AATGGGCTTT GTAGCTGCTC TTTTAGCAAG TATAGCAAAC
 421 AGTTATTATA ATCGCACTAT GAGTGGGTAT TATGATACGG ATATGCTGGT AATTGTTTTG
 481 CCTATGTTTA TTTTATTTTT TATGGTAAGA ATGATTTTAA AAAAAGACTT TTTTTCATTG
 541 ATTGCCTTGC CGTTATTTAT AGGAATTTAT CTTTGGTGGT ATCCTTCAAG TTATACTTTA
 601 AATGTAGCTT TAATTGGACT TTTTTTAATT TATACACTTA TTTTTCATAG AAAAGAAAAG
 661 ATTTTTTATA TAGCTGTGAT TTTGTCTTCT CTTACTCTTT CAAATATAGC ATGGTTTTAT
 721 CAAAGTGCCA TTATAGTAAT ACTTTTTGCT TTATTCGCCT TAGAGCAAAA ACGCTTAAAT
 781 TTTATGATTA TAGGAATTTT AGGTAGTGCA ACTTTGATAT TTTTGATTTT AAGTGGTGGG
 841 GTTGATCCTA TACTTTATCA GCTTAAATTT TATATTTTTA GAAGTGATGA AAGTGCGAAT
 901 TTAACGCAGG GCTTTATGTA TTTTAATGTC AATCAAACCA TACAAGAAGT TGAAAATGTA
 961 GATCTTAGCG AATTTATGCG AAGAATTAGT GGTAGTGAAA TTGTTTTTTT GTTTTCTTTG
1021 TTTGGTTTTG TATGGCTTTT GAGAAAACAT AAAAGTATGA TTATGGCTTT ACCTATATTG
1081 GTGCTTGGGT TTTTAGCCTT AAAAGGGGGG CTTAGATTTA CCATTTATTC TGTACCTGTA
1141 ATGGCCTTAG GATTTGGTTT TTTATTGAGC GAGTTTAAGG CTATAATGGT TAAAAAATAT
1201 AGCCAATTAA CTTCAAATGT TTGTATTGTT TTTGCAACTA TTTTGACTTT AGCTCCAGTA
1261 TTTATCCATA TTTACAACTA TAAAGCGCCA ACAGTTTTTT CTCAAAATGA AGCATCATTA
1321 TTAAATCAAT TAAAAAATAT AGCCAATAGA GAAGATTATG TGGTAACTTG GGCGGCTTAT
1381 GGTTATCCTG TGCGTTATTA TAGCGATGTG AAAACTTTAG TAGATGGTGG AAAGCATTTA
1441 GGTAAGGATA ATTTTTTCCC TTCTTTTGCT TTAAGCAAAG ATGAACAAGC TGCAGCTAAT
1501 ATGGCAAGAC TTAGTGTAGA ATATACAGAA AAAAGCTTTT ATGCTCCGCA AAATGATATT
1561 TTAAAAACAG ACATTTTGCA AGCCATGATG AAAGATTATA ATCAAAGCAA TGTGGATTTG
1621 TTTCTAGCTT CATTATCAAA ACCTGATTTT AAAATCGATA CGCCAAAAAC TCGTGATATT
1681 TATCTTTATA TGCCCGCTAG AATGTCTTTG ATTTTTTCTA CGGTGGCTAG TTTTTCTTTT
1741 ATTAATTTAG ATACAGGAGT TTTGGATAAA CCTTTTACCT TTAGCACAGC TTATCCACTT
1801 GATGTTAAAA ATGGAGAAAT TTATCTTAGC AACGGAGTGG TTTTAAGCGA TGATTTTAGA
1861 AGTTTTAAAA TAGGTGATAA TGTGGTTTCT GTAAATAGTA TCGTAGAGAT TAATTCTATT
1921 AAACAAGGTG AATACAAAAT CACTCCAATT GATGATAAGG CTCAGTTTTA TATTTTTTAT
1981 TTAAAGGATA GTGCTATTCC TTACGCACAA TTTATTTTAA TGGATAAAAC CATGTTTAAT
2041 AGTGCTTATG TGCAAATGTT TTTTTTAGGA AATTATGATA AGAATTTATT TGACTTGGTG
2101 ATTAATTCTA GAGATGCTAA GGTTTTTAAA CTTAAAATTT ACCCATACGA TGTTCCAGAT
2161 TACGCTTAA
```

SEQ. ID NO. 2: AMINO ACID SEQUENCE FOR PgIB

722 AA

MLKKEYLKNPYLVLFAMIILAYVFSVFCRFYWVWWASEFNEYFFNNQLMIISNDGYAFAE
GARDMIAGFHQPNDLSYYGSSLSALTYWLYKITPFSFESIILYMSTFLSSLVVIPTILLA
NEYKRPLMGFVAALLASIANSYYNRTMSGYYDTDMLVIVLPMFILFFMVRMILKKDFFSL
IALPLFIGIYLWWYPSSYTLNVALIGLFLIYTLIFHRKEKIFYIAVILSSLTLSNIAWFY
QSAIIVILFALFALEQKRLNFMIIGILGSATLIFLILSGGVDPILYQLKFYIFRSDESAN
LTQGFMYFNVNQTIQEVENVDLSEFMRRISGSEIVFLFSLFGFVWLLRKHKSMIMALPIL
VLGFLALKGGLRFTIYSVPVMALGFGFLLSEFKAIMVKKYSQLTSNVCIVFATILTLAPV
FIHIYNYKAPTVFSQNEASLLNQLKNIANREDYVVTWAAYGYPVRYYSDVKTLVDGGKHL
GKDNFFPSFALSKDEQAAANMARLSVEYTEKSFYAPQNDILKTDILQAMMKDYNQSNVDL
FLASLSKPDFKIDTPKTRDIYLYMPARMSLIFSTVASFSFINLDTGVLDKPFTFSTAYPL
DVKNGEIYLSNGVVLSDDFRSFKIGDNVVSVNSIVEINSIKQGEYKITPIDDKAQFYIFY
LKDSAIPYAQFILMDKTMFNSAYVQMFFLGNYDKNLFDLVINSRDAKVFKLKIYPYDVPD
YA

SEQ. ID NO. 3: OLIGONUCLEOTIDE CASSETTE IN pGVXN64 PLASMID

5`-AATTCTGCAGGATCCTCTAGAAGCTTGG

SEQ. ID NO. 4: OLIGONUCLEOTIDE CASSETTE IN pGVXN64 PLASMID

5`AATTCCAAGCTTCTAGAGGATCCTGCAG

SEQ. ID NO. 5: COMPLETE NUCLEOTIDE SEQUENCE ENCODING THE *SHIGELLA DYSENTERIAE* O1 ANTIGEN IN THE PGVXN64 PLASMID

LOCUS i-O1Cluste 10174 bp DNA SYN 23-JAN-2009
DEFINITION Inverted O1Cluster
ACCESSION i-O1Cluste
KEYWORDS .
SOURCE Unknown.
ORGANISM Unknown
Unclassified.
REFERENCE 1 (bases 1 to 10174)
AUTHORS Self
JOURNAL Unpublished.
BASE COUNT 3223 a 1581 c 2074 g 3296 t
ORIGIN

1 ATGAAAAACC ATAAGGTTAG TATTATCATC CCCTGTTTTA ATAACGGAAA AACTATAGAA
61 AGATGCGTTC TTTCCGCATT AAAACAAACC CATAGAAATA TAGAGATTAT TTGTATAAAT
121 GATGGACCGT CAGATAATAG CTGGAATATT TTAAAGAAAT TAAGCTCATT GTATGGAAAT
181 GTTTTTGCTT TTAATAATGA GGACAATTCA GGGCCGTCAT TTTCCAGGAT TAAAGGGGTT
241 TCACTATCTA CGGGGCATTT TTTGTCATTT TTGGATGCAG ATGATTACTG GCATCCAAAA
301 AAACTAGAAT TACAACTATC ATTTATTAAT GATGAAAACT TGGATTTTTT AGGTTCAACG
361 TGTTCCATTG GTGAGAAAAA TAACCAAGAA ATTAACCAAG GAATTAAAAA AGAACATTTA

421 AAATTAAAAA TAATTTCATT TAACATGATG TTGTTTAAGA ATTATTTCCA GACTCCAGCT
481 GTCATTATGA AAAGAGATAT TTTTATTCCA TTTAATGAGA ATCAGCGTTT TTCAGAGGAC
541 TACATGTCAT GGCTTGTTAT CGTTTATAAT AAAAAAACAA ATGTGGATTT AATATATGGA
601 AGGGATTTGG TTTTTCTCGA TAAATTTAAC TTTGGAGTGT CAGGGTTGAG TGGTAATTTA
661 TGGTTGATGG AGAAGTGGGA GTTAAAAAAT ATATTTAACT TCTTGTTGAA AGGTAAAATA
721 ATGGCAGTGC CTGCGATCTT GTTTTCTTTG ATAAAATATG AAAGAAGATG CGCTTTAACA
781 AAGAAAAATA AAGGTAAGGG TAATAAATAA TGAAGATCTC AATAATAGGG AACACAGCAA
841 ATGCTATGAT TTTGTTTAGA TTGGATTTAA TAAAAACACT AACCAACAAA GGGATTTCAG
901 TCTATGCTTT TGCTACTGAC TATAATGATT CATCCAAGGA AATAATAAAA AAAGCAGGCG
961 CCATTCCTGT TGATTATAAT TTAAGTCGCA GTGGTATTAA CCTTGCTGGT GATTTATGGA
1021 ATACTTACTT ATTAAGTAAA AAACTAAAGA AGATAAAACC AGATGCTATT TTATCTTTTT
1081 TTTCAAAGCC CTCTATCTTT GGATCGTTGG CTGGTATTTT TTCAGGCGTT AAAAATAATA
1141 ACGCTATGCT TGAGGGGTTA GGTTTTTTAT TTACAGAGCA GCCACATGGA ACTCCGTTAA
1201 AAACAAAGTT ACTTAAAAAT ATCCAGGTTC TCCTGTATAA AATAATATTT CCACATATCA
1261 ACTCATTAAT ACTCCTTAAC AAGGATGATT ATCATGATTT GATAGATAAA TACAAAATAA
1321 AATTAAAATC TTGCCATATT CTTGGTGGCA TTGGTTTAGA TATGAATAAT TACTGTAAAA
1381 GCACGCCACC AACAAATGAA ATATCATTCA TTTTTATAGC TCGTTTGCTA GCAGAAAAAG
1441 GAGTCAATGA GTTTGTTGCT GCCGCAAAAA AAATAAAAAA AACACATCCC AATGTTGAAT
1501 TTATTATACT TGGCGCTATA GATAAGGAAA ACCCCGGAGG GTTATCTGAA TCTGACGTAG
1561 ATACTTTAAT TAAATCAGGA GTTATTTCTT ATCCCGGATT TGTTTCTAAT GTGGCTGATT
1621 GGATTGAAAA ATCAAGCGTA TTTGTTCTTC CTTCCTATTA TCGAGAGGGA GTTCCTCGTA
1681 GTACACAAGA AGCGATGGCT ATGGGGAGGC CGATTTTAAC TACTAATTTA CCAGGCTGCA
1741 AAGAAACAAT TATTGATGGT GTGAATGGAT ATGTTGTAAA AAAATGGTCA CATGAAGATC
1801 TTGCAGAAAA AATGCTGAAG TTAATTAATA ATCCTGAAAA AATAATCAGT ATGGGAGAAG
1861 AAAGTTATAA GTTAGCAAGA GAAAGATTCG ATGCAAATGT AAATAATGTA AAGTTATTAA
1921 AAATACTAGG GATTCCTGAT TAATAAACGA AAAGCGGCTC TGATTCATTC GGAACTAAGA
1981 ACCTATCTCA ATAGGAGCTA AATTCATGAC CTTACCCAGC CATATCGACC TGCAGCCTGA
2041 CTGATGCCAT TGCCGAACTG GCGAAAAAAC AGTCCGTTGA TGCCATGCTG ATGACTGGAG
2101 ACAGCTACGA CTGCGGAAAA AAAATGGGCT ATATGCAGGC GTTTGTGAAG TATGGGCTGC
2161 GCAACCTCAA AGAAGGGGCG AAGTTCCGTA AAGGGATTGA GAAGCTGTTA AGCGAATAAT
2221 GAAAATCTGA CCGAATGTAA CGGTTGATAA GAAAATTATA ACGGCAGTGA AGATTCGTGG
2281 CGAAAGTAAT TTGTTGCGAA TATTCCTGCC GTTGTTTTAT ATAAACAATC AGAATAACAA
2341 AGAGTTAGCA ATAGGATTTT CGTCAAAGTT TTCCAGGATT TTCCTTGTTT CCAGAGCGGA
2401 TTGGTAAGAC AATTAGTGTT TGAATTTTTC GGGTTTAGCG CGAGTGGGTA ACGCTCGTCA
2461 CATCGTGGAC ATGTATGCAG TGCTCTGGTA GCTGTAAAGC CAGGGGCGGT AGCGTGCATT
2521 AATACCTCTA TTAATCAAAC TGAGAGCCGC TTATTTCACA GCATGCTCTG AAGTAATATG
2581 GAATAATAAA GTGAAGATAC TTGTTACTGG TGGCGCAGGA TTTATTGGTT CTGCTGTAGT
2641 TCGTCACATT ATAAATAATA CGCAGGATAG TGTTGTTAAT GTCGATAAAT TAACGTACGC
2701 CGGAAACCTG GAGTCACTTG CTGATGTTTC TGACTCTAAA CGCTATGTTT TTGAACATGC
2761 GGATATTTGC GATGCTGCTG CAATGGCGCG GATTTTTGCT CAGCATCAGC CGGATGCAGT
2821 GATGCACCTG GCTGCTGAAA GCCATGTGGA TCGTTCAATT ACAGGCCCTG CGGCATTTAT
2881 TGAAACCAAT ATTGTTGGTA CTTATGTCCT TTTGGAAGCG GCTCGCAATT ACTGGTCTGC
2941 TCTTGATGGC GACAAGAAAA ATAGCTTCCG TTTTCATCAT ATTTCTACTG ACGAAGTCTA
3001 TGGTGATTTG CCTCATCCTG ACGAAGTAAA TAATAAAGAA CAATTACCCC TCTTTACTGA
3061 GACGACAGCT TACGCGCCTA GTAGTCCTTA TTCCGCATCA AAAGCATCCA GCGATCATTT
3121 AGTCCGTGCG TGGAAACGTA CCTATGGTTT ACCGACCATT GTGACTAACT GTTCGAATAA
3181 CTACGGTCCT TATCACTTTC CGGAAAAATT GATTCCACTA GTAATTCTTA ATGCTCTGGA
3241 AGGTAAGGCA TTACCTATTT ATGGCAAAGG GGATCAAATT CGTGACTGGC TGTATGTTGA
3301 AGATCATGCG CGTGCGTTAT ATATCGTCGT AACCGAAGGT AAAGCGGGTG AAACTTATAA
3361 CATTGGTGGA CACAACGAAA AGAAAAACAT CGATGTAGTG CTCACTATTT GTGATTTGTT
3421 GGATGAGATT GTACCGAAAG AGAAATCTTA CCGCGAGCAA ATTACTTATG TTGCCGATCG
3481 CCCGGGACAC GATCGCCGTT ATGCGATTGA TGCAGAGAAG ATTAGCCGCG AATTGGGCTG
3541 GAAACCGCAG GAAACGTTTG AGAGCGGGAT TCGTAAAACG GTGGGATGGT ACCTCTCCAA
3601 TACAAAATGG GTTGATAATG TAAAAAGTGG TGCCTATCAA TCGTGGATTG AACAGAACTA
3661 TGAGGGCCGC CAGTAATGAA TATCCTCCTT TTCGGCAAAA CAGGGCAGGT AGGTTGGGAA
3721 CTACAGCGTG CTCTGGCACC TCTGGGTAAT TTGATTGCTC TTGATGTTCA CTCCACTGAT

3781 TACTGTGGTG ATTTTAGTAA TCCTGAAGGT GTAGCTGAAA CCGTAAGAAG CATTCGGCCT
3841 GATATTATTG TCAACGCAGC CGCTCACACC GCAGTAGACA AAGCAGAATC AGAACCGGAG
3901 TTTGCACAAT TACTTAACGC GACGAGTGTC GAAGCGATCG CGAAAGCAGC CAATGAAGTC
3961 GGCGCCTGGG TTATTCACTA CTCTACTGAC TACGTATTTC CGGGGACCGG TGAAATACCA
4021 TGGCAGGAGG CGGATGCAAC CGCACCGCTA AATGTTTACG GTGAAACCAA GTTAGCTGGA
4081 GAAAAAGCAT TACAAGAGCA TTGTGCGAAG CACCTAATTT TCCGTACAAG CTGGGTCTAT
4141 GCAGGTAAAG GAAATAACTT CGCCAAAACG ATGTTGCGTC TGGGAAAAGA GCGTGAAGAA
4201 TTAGCCGTTA TTAATGATCA GTTTGGTGCG CCAACAGGTG CTGAACTGCT GGCTGATTGT
4261 ACGGCACATG CAATTCGTGT GGCACTGAAT AAACCAGAAG TCGCAGGCTT GTACCATCTG
4321 GTAGCCACTG GTACCACAAC CTGGCACGAT TATGCTGCGC TGGTTTTTGA AGAGGCACGA
4381 AAAGCAGGTA TTCCCCTTGC ACTCAACAAG CTCAACGCAG TACCAACAAC AGCTTATCCT
4441 ACACCAGCTC GTCGTCCACA TAACTCTCGC CTTAATACAG AAAAATTTCA GCAAAATTTT
4501 GCGCTTGTTT TGCCTGACTG GCAGGTTGGC GTGAAACGAA TGCTCAACGA ATTATTTACG
4561 ACTACAGCAA TTTAATAGTT TTTGCATCTT GTTCGTGATG ATGGAGCAAG ATGAATTAAA
4621 AGGAATGATG TAATGAAAAC GCGTAAAGGT ATTATTTTAG CGGGTGGCTC TGGTACTCGT
4681 CTTTATCCTG TGACTATGGC TGTCAGTAAA CAGCTATTAC CTATTTATGA TAAGCCGATG
4741 ATCTATTACC CGCTCTCTAC ACTGATGTTG GCGGGTATTC GCGATATTCT GATTATTAGT
4801 ACGCCACAGG ATACTCCTCG TTTTCAACAA CTCCTGGGTG ATGGTAGCCA GTGGGGGTTA
4861 AATCTTCAGT ACAAAGTGCA ACCGAGTCCA GATGGTCTTG CGCAGGCATT TATCATCGGT
4921 GAAGAGTTTA TCGGTGGTGA TGATTGTGCT CTGGTTCTCG GTGATAATAT CTTCTACGGT
4981 CATGATCTGC CGAAGTTAAT GGATGTCGCT GTCAACAAAG AAAGTGGTGC AACGGTATTT
5041 GCCTATCACG TTAATGATCC TGAACGCTAC GGTGTTGTTG AGTTTGATAA AAACGGTACG
5101 GCAATCAGCC TGGAAGAAAA ACCGCTACAA CCAAAAAGTA ATTATGCGGT AACCGGGCTT
5161 TATTTCTATG ATAACGACGT TGTCGAAATG GCGAAAAACC TTAAGCCTTC TGCCCGTGGT
5221 GAACTGGAAA TTACCGATAT TAACCGTATT TATATGGAGC AGGGGCGTTT ATCCGTTGCC
5281 ATGATGGGAC GTGGTTATGC ATGGCTGGAC ACGGGGACAC ATCAAAGTCT TATTGAAGCA
5341 AGCAACTTCA TTGCAACAAT TGAAGAGCGC CAAGGGTTAA AGGTATCTTG CCTGGAAGAG
5401 ATTGCTTATC GTAAAGGCTT TATTGACGCA GAGCAGGTTA ATGTATTAGC CGAACCGCTA
5461 AAGAAAAATG CTTATGGTCA GTATCTGTTG AAAATGATTA AAGGTTATTA AAAATGAATG
5521 TAATTAAAAC TGAAATTCCA GATGTATTAA TTTTCGAGCC GAAAGTTTTT GGTGATGAAC
5581 GTGGTTTTTT TATGGAAAGC TTTAACCAGA AAGTTTTCGA AGAGGCTGTA GGGCGGAAGG
5641 TTGAATTTGT TCAGGATAAC CATTCTAAAT CAACTAAGGG TGTGTTACGC GGACTGCACT
5701 ATCAGTTGGA ACCTTATGCT CAAGGTAAAT TAGTTCGTTG TGTTGTCGGT GAAGTTTTTG
5761 ATGTAGCAGT TGATATTCGT AAATCGTCAC CTACATTTGG GAAATGGATT GGGGTGAATT
5821 TGTCTGCTGA GAATAAGCGT CAGTTGTGGA TACCTGAAGG ATTTGCGCAT GGATTTTTGG
5881 TGCTGAGTGA AACGGCTGAG TTTGTTTATA AAACAACAAA CTATTACAAT CCAAGTTTTG
5941 AAAAAAGTAT TTCATACTCA GATCCTACCA TTAAAATTCA GTGGCCCAAT TTACAGGATA
6001 TGCATTTTAA ATTATCAAAT AAGGATTTGA ATGCTAAGAA CTTTTTTAAT AACAATAGTT
6061 TAATGCAATG AAGAAAAATA TATTGCTCTT GTTCTTAGTA CATGGGGCAA ATTATTTGTT
6121 CCCGTTTATA GTTCTTCCAT ATCAAACTCG AATATTAAGC ATCGAGACAT TCGCAGATGT
6181 AGCAAAAATT CAAGCCGCTG TGATGCTTTT ATCTTTAATC GTAAATTATG GATATAACTT
6241 ATCAAGTACA AGAGCTATAG CTAGGGCCGT ATCTCAAGCA GAAATAAATA AGATCTATAG
6301 TGAGACTCTT ATTGTAAAAT TATTATTGGC AACCATTTGT CTTGCACTTG GTTGCGTACA
6361 TTTGATGTAT GTCAAAGAGT ACTCATTGAT ATATCCTTTT ATAATCAGTT CGATATATCT
6421 TTATGGTAGT GCATTATTTG CTACTTGGTT ATTCCAAGGA CTTGAGAAAA TGAAAGCGGT
6481 CGTTATAGCA ACAACAATCG CTAAACTGAC TGGTGTGATA CTTACTTTTA TTTTAGTTAA
6541 GTCTCCAAAT GATATAGTTG CAGCTCTTTT TACACAAAAC ATTGGGATGT TTATAAGTGG
6601 TATAATATCT ATTTATTTGG TAAGGAAAAA CAAATATGCA ACCGTAATAT GTTTTCGACT
6661 TAAAAATATT ATTGTAAGCT TAAAAGAAGC GTGGCCGTTT TTTTTATCAT TAGCTGCAAC
6721 AAGTGTATAT ACATATTTTA ATGTGATTTT ATTATCTTTT TATGCTGGCG ACTATGTTGT
6781 GGCAAATTTT AATGCTGCTG ATAAATTAAG AATGGCTGCT CAAGGGTTAC TTATTCCAAT
6841 AGGACAGGCT GTTTTCCCAC GATTATCTAA ACTAGAGGGC TATGAATATA GTTCTAAACT
6901 TAAAATTTAT GCAATAAGGT ATGCTATTTT TGGTGTTTGC ATTAGTGCGG GACTTGTATT
6961 TTTAGGTCCC ATGTTAACTA CTATTTATTT AGGCAAAGAA TATTCGTTGT CAGGAGAATA
7021 TCTTCAAAGT ATGTTTTTAC TACCTGCCAC TATTTCAATA TCGACTATAC TGAGTCAATG
7081 GATGTTGATA CCTCAAGGCA AAGAAAAAAT ATTAAGCAGA ATCTATATTC TAGGCGCCAT

7141 TGTCCATTTA TTATATGCAT TTCCTTTAGT TTACTATTAT GGGGCTTGGG GCATGGTAAT
7201 ATCAATTTTA TTTACTGAAG TCTTAATTGT ATTATTTATG CTTAAGGCTG TGAAATGACT
7261 TACTTTACTG GTTTTATTTT AATATTGTTT GCTATTATAA TTAAAAGATT AACTCCAAGT
7321 CAAAGCAAGA AAAATATTGT CTTAATAGCT AATGCGTTTT GGGGAATATT GTTGGTAGGT
7381 TATGCTTTCA ATGAACAATA TTTCGTACCA TTAAGTGCAA CAACCTTGTT TTTTATACTT
7441 GCATTCTTAT TTTTCTTTAG TATGACTTAT ATTTTAATTG CTAGGAGTGG AAGGGTTGTT
7501 TTTTCTTTCG GTACTGGTTT TATAGAAAGC AAATATATTT ACTGGTTTGC TGGGATGATT
7561 AATATTATTA GTATCTGCTT TGGCATTATC CTTTTATATA ATAATCATTT TTCTTTAAAA
7621 GTAATGAGAG AAGGAATTTT AGATGGTTCT ATTAGTGGGT TTGGATTGGG GATAAGTTTG
7681 CCACTTTCCT TCTGCTGTAT GTATTTAGCA AGACATGAGA ATAAAAAAAA TTATTTCTAT
7741 TGTTTTACAC TACTTTCATT CTTGCTTGCG GTGTTATCAA CTTCAAAGAT CTTCTTAATA
7801 TTATTCCTTG TATATATTGT TGGAATAAAT AGTTATGTAA GCAAAAAGAA ATTGCTTATT
7861 TATGGAGTGT TTGTATTTGG ACTGTTCGCT TTATCAAGTA TTATCTTGGG TAAGTTCTCT
7921 TCAGACCCTG AAGGCAAGAT TATTTCAGCA ATATTTGATA CGTTAAGGGT TTATCTTTTC
7981 TCGGGATTGG CAGCCTTTAA TCTTTATGTT GAAAAGAATG CCACGCTCCC CGAAAATTTA
8041 CTTTTGTATC CATTTAAGGA GGTTTGGGGG ACGACAAAAG ATATTCCCAA AACTGATATT
8101 TTGCCTTGGA TCAACATTGG TGTATGGGAC ACGAATGTAT ATACAGCTTT TGCACCATGG
8161 TATCAGTCAT TGGGATTATA TGCAGCTATA ATTATTGGTA TTCTCTTAGG GTTTTATTAC
8221 GGGATATGGT TTAGCTTTCG TCAAAATTTA GCTGTGGGTT TTTATCAAAC ATTTTTGTGT
8281 TTTCCTCTTT TAATGTTGTT TTTCCAGGAG CATTATTTGT TGTCATGGAA AATGCATTTT
8341 ATTTATTTTT TATGTGCAAT TTTATTAGCG ATGAGAAAAG CATTAGAGTA TGAATAAATA
8401 TTGTATCTTA GTACTATTTA ATCCAGATAT AAGTGTTTTT ATTGATAATG TCAAAAAGAT
8461 TTTATCTTTG GATGTAAGTT TATTTGTATA TGACAATTCA GCAAATAAAC ATGCATTCCT
8521 TGCTCTATCC TCACAAGAGC AAACAAAGAT AAATTACTTT TCGATATGTG AAAATATCGG
8581 ATTGTCGAAA GCTTATAATG AGACACTAAG GCATATTCTT GAATTTAATA AGAATGTGAA
8641 AAATAAAAGC ATTAATGATA GTGTGCTTTT TCTCGACCAA GACTCTGAAG TTGATTTAAA
8701 TTCCATCAAT ATTTTGTTTG AAACTATATC AGCAGCAGAG TCTAATGTGA TGATAGTCGC
8761 GGGGAATCCC ATAAGGAGAG ATGGACTACC GTATATAGAT TACCCCCACA CTGTAAACAA
8821 TGTAAAATTT GTAATTAGTA GTTATGCTGT GTATCGCTTA GACGCATTTA GAAACATCGG
8881 CTTGTTTCAA GAAGATTTTT TTATAGATCA TATCGATAGT GATTTTTGTT CAAGGCTGAT
8941 AAAAAGCAAT TACCAAATTC TCCTTAGAAA AGATGCCTTT TTTTATCAAC CAATAGGAAT
9001 AAAACCATTC AATCTCTGTG GTAGATATTT ATTCCCTATC CCATCACAAC ACCGAACATA
9061 TTTTCAAATT AGAAATGCTT TTTTAAGTTA CAGGCGCAAT GGTGTTACAT TTAATTTTTT
9121 ATTTAGGGAA ATTGTAAATA GATTGATTAT GAGTATATTC TCAGGCCTTA ACGAGAAAGA
9181 CTTATTGAAA CGATTGCATT TATATTTAAA AGGAATAAAA GATGGTCTTA AAATGTAATT
9241 CTTGGCTAGA AGTGGGGGCG TTGTGATTAA AAAAAAAGTG GCGGCGATAA TTATAACATA
9301 TAATCCAGAT CTAACAATTC TGCGAGAAAG TTATACGAGT CTATATAAGC AAGTCGATAA
9361 AATAATTCTT ATTGATAACA ACTCTACAAA CTATCAAGAA CTTAAGAAGT TATTCGAAAA
9421 AAAAGAAAAA ATAAAAATAG TGCCCTTGAG TGATAATATA GGACTAGCAG CAGCTCAAAA
9481 TTTAGGTTTG AACTTAGCTA TTAAAAATAA CTATACTTAT GCTATTTTAT TCGATCAGGA
9541 TAGCGTCTTA CAAGACAATG GAATTAACAG TTTCTTTTTT GAATTTGAGA AATTAGTTAG
9601 TGAAGAAAAA TTAAATATAG TTGCCATTGG GCCAAGTTTT TTTGACGAAA AGACAGGAAG
9661 ACGCTTTCGG CCTACAAAAT TTATCGGTCC CTTTTTATAT CCCTTTCGTA AAATAACCAC
9721 AAAAAATCCT CTAACAGAAG TTGACTTCTT GATTGCTTCT GGTTGTTTCA TAAAATTGGA
9781 GTGTATTAAA TCAGCCGGAA TGATGACTGA ATCGTTATTC ATCGATTATA TTGATGTTGA
9841 ATGGTCATAT CGTATGCGTT CGTATGGCTA TAAGCTATAT ATTCATAATG ATATTCACAT
9901 GAGTCATTTA GTGGGAGAAT CTCGAGTTAA TTTAGGATTG AAAACTATTT CTTTACATGG
9961 GCCGCTAAGA CGATATTACT TATTTAGGAA TTATATTTCA ATTTTAAAAG TGAGATATAT
10021 ACCGTTAGGA TATAAAATAC GTGAGGGTTT TTTTAATATC GGAAGATTTT TGGTAAGTAT
10081 GATTATAACT AAAAATAGAA AAACTTTAAT TTTATACACT ATAAAAGCAA TTAAGGACGG
10141 AATAAATAAT GAAATGGGGA AATATAAAGG CTAA

SEQ. ID NO. 6: EPA AA SEQUENCE (643 AA), CONTAINING TWO GLYCOSYLATION SITES, WITH SIGNAL SEQUENCE

MKKIWLALAGLVLAFSASAAEEAFDLWNECAKACVLDLKDGVRSSRMSVDPAIADTN
GQGVLHYSMVLEGGNDALKLAIDNALSITSDGLTIRLEGGVEPNKPVRYSYTRQARGSW
SLNWLVPIGHEKPSNIKVFIHELNAGNQLSHMSPIYTIEMGDELLAKLARDATFFVRAHE
SNEMQPTLAISHAGVSVVMAQAQPRREKRWSEWASGKVLCLLDPLDGVYNYLAQQRC
NLDDTWEGKIYRVLAGNPAKHDLDIKDNNNSTPTVISHRLHFPEGGSLAALTAHQACHL
PLEAFTRHRQPRGWEQLEQCGYPVQRLVALYLAARLSWNQVDQVIRNALASPGSGGDL
GEAIREQPEQARLALTLAAAESERFVRQGTGNDEAGAASADVVSLTCPVAKDQNRTKG
ECAGPADSGDALLERNYPTGAEFLGDGGDVSFSTRGTQNWTVERLLQAHRQLEERGYV
FVGYHGTFLEAAQSIVFGGVRARSQDLDAIWRGFYIAGDPALAYGYAQDQEPDARGRIR
NGALLRVYVPRWSLPGFYRTGLTLAAPEAAGEVERLIGHPLPLRLDAITGPEEEGGRVTI
LGWPLAERTVVIPSAIPTDPRNVGGDLDPSSIPDKEQAISALPDYASQPGKPPREDLK

SEQ. ID NO. 7: PROCESSED EPA AA SEQUENCE (624 AA), CONTAINING TWO GLYCOSYLATION SITES

AEEAFDLWNECAKACVLDLKDGVRSSRMSVDPAIADTNGQGVLHYSMVLEGGNDALK
LAIDNALSITSDGLTIRLEGGVEPNKPVRYSYTRQARGSWSLNWLVPIGHEKPSNIKVFIH
ELNAGNQLSHMSPIYTIEMGDELLAKLARDATFFVRAHESNEMQPTLAISHAGVSVVMA
QAQPRREKRWSEWASGKVLCLLDPLDGVYNYLAQQRCNLDDTWEGKIYRVLAGNPAK
HDLDIKDNNNSTPTVISHRLHFPEGGSLAALTAHQACHLPLEAFTRHRQPRGWEQLEQC
GYPVQRLVALYLAARLSWNQVDQVIRNALASPGSGGDLGEAIREQPEQARLALTLAAA
ESERFVRQGTGNDEAGAASADVVSLTCPVAKDQNRTKGECAGPADSGDALLERNYPTG
AEFLGDGGDVSFSTRGTQNWTVERLLQAHRQLEERGYVFVGYHGTFLEAAQSIVFGGV
RARSQDLDAIWRGFYIAGDPALAYGYAQDQEPDARGRIRNGALLRVYVPRWSLPGFYR
TGLTLAAPEAAGEVERLIGHPLPLRLDAITGPEEEGGRVTILGWPLAERTVVIPSAIPTDPR
NVGGDLDPSSIPDKEQAISALPDYASQPGKPPREDLK

SEQ. ID NO. 8: CONSTRUCT A CONTAINING A118-S130 FROM EXAMPLE 2

MKKTAIAIAVALAGFATVAQADVIIKPQVSGVIVNKLFKAGDKVKKGQTLFIIEQDQASK
DFNRSKALFSQLDHTEIKAPFDGTIGDALVNIGDYVSASTTELVRVTNLNPIYADGSHHH
HHH

SEQ. ID NO. 9: CONSTRUCT B CONTAINING F122-E138 FROM EXAMPLE 2

MKKTAIAIAVALAGFATVAQADVIIKPQVSGVIVNKLFKAGDKVKKGQTLFIIEQDQ**FN**
RSKALFSQSAISQKELDHTEIKAPFDGTIGDALVNIGDYVSASTTELVRVTNLNPIYADGS
HHHHHH

SEQ. ID NO. 10: CONSTRUCT C CONTAINING D121-A127 FROM EXAMPLE 2

MKKTAIAIAVALAGFATVAQADVIIKPQVSGVIVNKLFKAGDKVKKGQTLFIIEQDQDF
NRSKALDHTEIKAPFDGTIGDALVNIGDYVSASTTELVRVTNLNPIYADGSHHHHHH

SEQ. ID NO. 11: AMINO ACID SEQUENCE OF THE SOLUBLE ACRA PROTEIN FROM EXAMPLE 3

MKYLLPTAAAGLLLLAAQPAMAMHMSKEEAPKIQMPPQPVTTMSAKSEDLPLS/TYPA
KLVSDYDVIIKPQVSGVIVNKLFKAGDKVKKGQTLFIIEQDKFKASVDSAYGQALMAKA
TFENASK**DFNRS**KALFSKSAISQKEYDSSLATFNNSKASLASARAQLANARIDLDHTEIK
APFDGTIGDALVNIGDYVSASTTELVRVTNLNPIYADFFISDTDKLNLVRNTQSGKWDLD
SIHANLNLNGETVQGKLYFIDSVIDANSGTVKAKAVF**DNNNS**TLLPGAFATITSEGFIQK
NGFKVPQIGVKQDQNDVYVLLVKNGKVEKSSVHISYQNNEYAIIDKGLQNGDKIILDNF
KKIQVGSEVKEIGAQLEHHHHHH

SEQ. ID NO. 12: AMINO ACID SEQUENCE OF THE CTXB PROTEIN FROM EXAMPLE 6

MKKTAIAIAVALAGFATVAQATPQNITDLCAEYHNTQIHTLNDKIFSYTESLAGKREMAI
ITFKNGATFQVEVPGSQHIDSQKKAIERMKDTLRIAYLTEAKVEKLCVWNNKTPHAIAAI
SMANGSHHHHHH

SEQ. ID NO. 13: AMINO ACID SEQUENCE OF THE OMPH1 PROTEIN OF *CAMPYLOBACTER JEJUNI* (STRAIN 81-176) FROM EXAMPLE 7

MKKILLSVLTTFVAVVLAACGGNSDSKTLNSLDKIKQNGWRIGVFGDKPPFGYVDEKG
NNQGYDIALAKRIAKELFGDENKVQFVLVEAANRVEFLKSNKVDIILANFTQTPERAEQ
VDFCLPYMKVALGVAVPK**DSNIT**SVEDLKDKTLLLNKGTTADAYFTQDYPNIKTLKYD
QNTETFAALMDKRGDALSHDNTLLFAWVKDHPDFKMGIKELGNKDVIAPAVKKGDKE
LKEFIDNLIIKLGQEQFFHKAYDETLKAHFGDDVKADDWIEGGKILEQKL/SEEDL

SEQ. ID NO. 14: FORWARD OLIGO FOR AMPLIFYING EPA USING PCR IN EXAMPLE 10

5'-AAGCTAGCGCCGCCGAGGAAGCCTTCGACC

SEQ. ID NO. 15: REVERSE OLIGO FOR AMPLIFYING EPA USING PCR IN EXAMPLE 10

5'-AAGAATTCTCAGTGGTGGTGGTGGTGGTGCTTCAGGTCCTCGCGCGGCGG

SEQ. ID NO. 16: PHOSPHORYLATED OLIGONUCLEOTIDE USED TO CONSTRUCT pGVXN70 IN EXAMPLE 10

5'-GAAGGCGGGCGCGTGACCATTCTCGGC

SEQ. ID NO. 17: PHOSPHORYLATED OLIGONUCLEOTIDE USED TO CONSTRUCT pGVXN70 IN EXAMPLE 10

5'-GCCGAGAATGGTCACGCGCCCGCCTTC

SEQ. ID NO. 18: OLIGO FOR QUICK CHANGE MUTAGENESIS USING pGVXN70 AS A TEMPLATE IN EXAMPLE 10

5'-CCTGACCTGCCCCGGGGAATGCGCGG

SEQ. ID NO. 19: OLIGO FOR QUICK CHANGE MUTAGENESIS USING pGVXN70 AS A TEMPLATE IN EXAMPLE 10

5'-CCGCGCATTCCCCGGGGCAGGTCAGG

SEQ. ID NO. 20: OLIGO FOR PRODUCTION OF pGVXN137 IN EXAMPLE 10

5'- GTCGCCAAAGATCAAAATAGAACTAAA

SEQ. ID NO. 21: OLIGO FOR PRODUCTION OF pGVXN137 IN EXAMPLE 10

5'- TTTAGTTCTATTTTGATCTTTGGCGAC

SEQ. ID NO. 22: OLIGO FOR INSERTION OF ADDITIONAL GLYCOSYLATION SITE INTO pGVXN137 IN EXAMPLE 10

5'-CATGACCTGGACATCAAGGATAATAATAATTCTACTCCCACGGTCATCAGTCATC

SEQ. ID NO. 23: OLIGO FOR INSERTION OF ADDITIONAL GLYCOSYLATION SITE INTO pGVXN137 IN EXAMPLE 10

5'-GATGACTGATGACCGTGGGAGTAGAATTATTATTATCCTTGATGTCCAGGTCATG

CLAIMS

1. A method of producing a composition comprising *Shigella* bioconjugates in a bioreactor comprising the steps:

(i) growing bacteria that comprises (a) modified exotoxin of *Pseudomonas aeruginosa* (EPA) containing at least one consensus sequence, D/E – X – N – Z – S/T, wherein X and Z may be any natural amino acid except proline; (b) an oligosaccharyl transferase; and (c) one or more *Shigella* polysaccharides for a period of time to produce an amount of *Shigella* bioconjugates comprising the modified EPA linked to the one more *Shigella* polysaccharides;

(ii) extracting periplasmic proteins; and


(iii) separating the *Shigella* bioconjugates from the extracted periplasmic proteins by anionic exchange.

2. The method of claim 1, wherein said modified EPA has an amino acid sequence comprising SEQ. ID NO.: 6.

3. The method of claim 1, wherein said modified EPA has an amino acid sequence comprising SEQ. ID NO.: 7.

4. The method of any one of the preceding claims, wherein said *Shigella* polysaccharide is from *Shigella dysenteriae* **O1**.

5. The method of any one of the preceding claims, wherein said *Shigella* polysaccharide comprises the following structure:



6. The method of any one of the preceding claims, wherein said bacteria comprises a nucleotide sequence comprising SEQ ID NO: 5.

7. The method of any one of the preceding claims, wherein the oligosaccharyl transferase is from *Campylobacter* spp.

8. The method of any one of the preceding claims, wherein the oligosaccharyl transferase is from *Campylobacter jejuni*.

9. Use of the method of any one of the preceding claims for the production of a bioconjugate vaccine.

10. The method of any one of the preceding claims, wherein said bacteria is *E. coli*.