

**ENHANCED FORMULATIONS FOR NEUTRALIZATION OF
CHEMICAL, BIOLOGICAL AND INDUSTRIAL TOXANTS**

5 **CROSS-REFERENCE TO RELATED APPLICATIONS**

 This application is a continuation-in-part application of U.S. Patent Application Serial
No. 09/607,586, entitled "Formulations for Neutralization of Chemical and Biological Toxants", filed
on June 29, 2000, now U.S. Patent No. 6,566,574, which was a continuation-in-part application of
U.S. Patent Application Serial No. 09/108,235, entitled "Aqueous Foams for Mitigation and
10 Decontamination of Chemical and Biological Weapons Agents", filed on June 30, 1998, now
abandoned, and which claimed the benefit of the filing of U.S. Provisional Patent Application Serial
No. 60/146,432, filed on July 29, 1999, and the specifications thereof are incorporated herein by
reference.

15 This application is also a continuation-in-part application of U.S. Patent Application Serial
No. 09/952,940, now Patent Publication No. 2003/060517, entitled "Concentrated Formulations and
Methods for Neutralizing Chemical and Biological Toxants", filed on September 14, 2001, which was
a continuation-in-part application of U.S. Patent Application Serial No. 09/607,586, entitled
"Formulations for Neutralization of Chemical and Biological Toxants", filed on June 29, 2000, and the
20 specifications thereof are incorporated herein by reference.

 This application claims the benefit of the filing of U.S. Provisional Patent Application Serial
No. 60/326,508, entitled "DF-200 – An Enhanced Formulation for Decontamination and Mitigation of
CBW Agents and Biological Pathogens", filed on October 1, 2001, and of U.S. Provisional Patent
25 Application Serial No. 60/334,271, entitled "Configurations for the Rapid Deployment of DF-200",
filed on November 30, 2001, and of U.S. Provisional Patent Application Serial No. 60/387,104,
entitled "Decontamination Formulations", filed on June 7, 2002, and the specifications thereof are
incorporated herein by reference.

WO 03/028429

PCT/US02/29886

-2-

GOVERNMENT RIGHTS

The Government has rights to this invention pursuant to Contract No. DE-AC04-94AL85000 awarded by the U.S. Department of Energy.

5

BACKGROUND OF THE INVENTIONField of the Invention (Technical Field):

The present invention relates to formulations for neutralization of chemical, biological and industrial toxants.

10 Background:

The present invention is directed to materials and methods for neutralization of toxic chemical, biological and industrial compounds or agents, especially chemical and biological weapons agents, and methods of making same. In particular, the present invention is directed to materials containing solubilizing compounds, reactive compounds and bleaching activators that can be delivered as foams, sprays, liquids, fogs and aerosols to enhance the rate of reactions leading to neutralization of chemical compounds, and other additives which serve to kill or attenuate certain biological compounds or agents.

15

20

Terrorist threats, potentially involving weapons of mass destruction, are increasing both in the United States and abroad. The use, and threat of use, of chemical and biological agents in the context of weapons of mass destruction are of paramount concern both to national defense as well as to state and local law enforcement.

25

Certain chemical warfare ("CW") agents known to pose a threat by terrorists share chemical characteristics that present an opportunity for the development of countermeasures. The chemical agents sarin, soman, and tabun (G-agents) are all examples of phosphorus-containing compounds which, when altered chemically, can lose their toxicity. Mustard, which is an example of the H-agents, and VX, which is an example of the V-agents, can also be altered chemically and rendered harmless. In addition, certain of the known BW agents include botulinum toxin, anthrax and other

WO 03/028429

PCT/US02/29886

-3-

spore-forming bacteria, vegetative bacteria, including plague and various viruses can also be deactivated chemically.

5 A CW or biological warfare ("BW") attack can involve either local placement or wide dispersal of the agent or agents so as to affect a population of human individuals. Because of the flexibility with which CW and BW ("CBW") agents can be deployed, respondents might encounter the agents in a variety of physical states including bulk, aerosol and vapors.

10 An effective, rapid, and safe (non-toxic and non-corrosive) decontamination technology is required for the restoration of civilian facilities in the event of a domestic terrorist attack. Ideally, this technology should be applicable to a variety of scenarios such as the decontamination of open, semi-enclosed, and enclosed facilities as well as sensitive equipment. Examples of types of facilities where the decontamination formulation may be utilized include a stadium (open), an underground subway station (semi-enclosed), and an airport terminal or office building (enclosed). A foaming
15 version is useful for extending the contact time of the formulation on vertical surface.

Decontamination of chemical compounds have focused primarily on chemical warfare agents, particularly on the nerve agents (such as G agents and V agents) and on the blistering agents (such as mustard gas, or simply, mustard). Reactions involved in detoxification of chemical agents can be
20 divided into substitution and oxidation reactions. Decontamination of biological agents is primarily focused on bacterial spores (e.g., anthrax), which are considered to be the most difficult of all microorganisms to kill. Additional background is discussed in U.S. Patent Applications Serial Nos. 09/607,586 and 09/952,940.

25 A need also exists for rapid, safe, and effective neutralization of toxic industrial chemicals, such as Malathion, Hydrogen Cyanide, Sodium Cyanide, Butyl Isocyanate, Carbon Disulfide, and Phosgene gas.

WO 03/028429

PCT/US02/29886

-4-

U.S. Patent Application Serial No. 09/607,586 is related generally to an aqueous-based decontamination technology ("DF-100") that rapidly neutralizes chemical and biological warfare ("CBW") agents. The formulation:

- is effective for neutralizing both chemical and biological agents;
- 5 • is environmentally benign (i.e., non-toxic and non-corrosive);
- works on a number of anticipated material surfaces; and
- can be incorporated into a wide variety of carriers (e.g., foams, liquid sprays, fogs)

that satisfy a wide variety of operational objectives.

10 A major interest for the use of the technology was from the civilian first responder (e.g., fire departments, police departments, and HazMat units who would be the first to arrive at the scene of an attack utilizing CBW agents) followed by a secondary interest in use of the formulation for facility restoration. Technical issues exist with DF-100 that make use of the formulation by the civilian first responder less than optimal. These technical problems include: (1) The pH of the DF-100 must be
15 adjusted to optimally decontaminate each specific chemical and biological agent. In other words, a different formulation may be required to neutralize each specific agent. Although it is relatively simple to adjust the pH of the formulation in the laboratory, this is more difficult in the field and is generally unsuitable for the primary users of the technology (i.e., the civilian first responder). (2) The reaction rate for one chemical agent, Mustard, is rather slow as compared to the reaction rates for
20 other chemical agents.

These technical problems limit the effectiveness of DF-100 in actual use. A modified formulation, DF-100A, disclosed in U.S. Patent Application Serial No. 09/952,940, addressed the requirement to adjust the pH for each specific agent (i.e., the first technical problem described
25 above). However, while DF-100A does improve upon the performance of the formulation at a single pH, it does not completely solve this problem, and does not even address the second technical problem (i.e., the relatively slow reaction rate with Mustard). Additionally, some versions of DF-100/100A can use short-chain alcohols (e.g., isobutanol, isopropanol), which can cause flammability problems if the formulation is packaged in a concentrated form. Also, some versions of

WO 03/028429

PCT/US02/29886

-5-

DF-100/100A can use diethylene glycol monobutyl ether (DEGMBE), which can cause false alarms on some chemical agent sensors and detectors (especially older sensors that are used in some military settings).

As a point of comparison, the following is an example of a preferred formulation for DF-100:

- 5 DF-100
- 2.6% Variquat 80MC (*cationic surfactant*)
 - 3.3% Adogen 477 (*cationic hydrotrope*)
 - 0.8% 1-Dodecanol (*fatty alcohol*)
 - 0.5% Isobutanol (*short chain alcohol*)
 - 10 1.6% Isopropanol (*short chain alcohol*)
 - 0.1% Jaguar 8000 (*cationic polymer*)
 - 1.6% Diethylene Glycol Monobutyl Ether (*solvent*)
 - 4% Sodium Bicarbonate (*buffer and peroxide activator*)
 - 4% Hydrogen Peroxide (*liquid oxidant*)
 - 15 75% Water

This formulation can be adjusted to a pH value of 8 for optimal decontamination of Mustard and Anthrax spores; and can be adjusted to a pH value of 10.5 for optimal decontamination of VX.

Decontamination of chemical agents is generally effective anywhere between pH 8 and 10.

As a further point of comparison, the following is an example of a preferred formulation for DF-100A:

- 20 DF-100A
- 5.3% Variquat 80MC (*cationic surfactant*)
 - 2.8% Adogen 477 (*cationic hydrotrope*)
 - 0.65% 1-Dodecanol (*fatty alcohol*)
 - 0.6% Isobutanol (*short chain alcohol*)
 - 25 0.1% Jaguar 8000 (*cationic polymer*)
 - 1.35% Diethylene Glycol Monobutyl Ether (*solvent*)
 - 4% Potassium Bicarbonate (*buffer and peroxide activator*)
 - 4% Hydrogen Peroxide (*oxidant*)
 - 81% Water

WO 03/028429

PCT/US02/29886

-6-

This formulation can be adjusted to a pH value of 8 for optimal decontamination of Mustard and Anthrax spores; and can be to a pH value of 10 for optimal decontamination of VX. Decontamination of chemical agents is generally effective anywhere between pH 8 and 10. Also, it can be adjusted to a pH of 9.2 for less than optimal decontamination of all agents.

5

In both of the examples shown above for DF-100 and DF-100A, the hydrogen peroxide and bicarbonate salt react to produce a highly reactive negatively charged nucleophilic species, hydroperoxycarbonate (HCO_4^-), which is a strong oxidant. Other negatively-charged nucleophiles are formed by the use of hydrogen peroxide, including: hydroxyl ions (OH^-) and hydroperoxide ions (OOH^-). The function of the other components in these formulations is discussed extensively in U.S. Patent Applications Serial Nos. 09/607,586 and 09/952,940.

10

The present invention presents enhanced decontamination formulations (generically designated "DF-200") that include bleaching activators, which leads to faster reaction kinetics, improved performance, and the elimination of the need for pH adjustment. Although bleaching activators are commonly used in (anionic) laundry detergents, the present invention can employ them with cationic surfactants, and where good solubility in water of the activator is useful for achieving quick reaction times. A desirable bleaching activator for use in the present invention is preferably water-soluble, non-toxic, non-flammable, and low-cost.

15

20

Glucose pentaacetate is an O-acetyl peroxide activator that has been used as an activation agent in a sterilizing composition comprising a cationic surface-active agent and inorganic peroxide (See Japanese Laid-Open Patent Publication No. 62-155203, entitled "Sterilizing Composition for Cattle Shed" (1987)). Glucose pentaacetate is a solid at room temperature (i.e., melting point 110 degrees C), and is insoluble in water. In an aqueous solution containing peroxide, it dissolves very slowly in water as it reacts with the peroxide. At a glucose pentaacetate concentration of about 2%, it takes several hours to dissolve. This makes its use less than desirable for rapid deployment configurations.

25

SUMMARY OF THE INVENTION (DISCLOSURE OF THE INVENTION)

The present inventions is of a formulation for use in neutralization of at least one toxant, comprising: at least two solubilizing compounds, wherein at least one solubilizing compound is a cationic surfactant and at least one solubilizing compound is a cationic hydrotrope; at least one
5 reactive compound, wherein the at least one reactive compound is selected from the group consisting of hydrogen peroxide, urea hydrogen peroxide, hydroperoxycarbonate, peracetic acid, sodium perborate, sodium peroxyphosphate, sodium peroxysilicate, and sodium percarbonate; and at least one bleaching activator selected from the group consisting of O-acetyl, N-acetyl, and nitrile group bleaching activators; wherein the at least two solubilizing compounds, the at least one
10 reactive compound, and the at least one bleaching activator, when mixed with water and exposed to the at least one toxant, neutralizes the at least one toxant. In the preferred embodiment, the cationic surfactant comprises a quaternary ammonium salt, most preferably cetyltrimethyl ammonium bromide, benzalkonium chloride, benzethonium chloride, cetylpyridinium chloride, alkyldimethylbenzylammonium salt, tetrabutyl ammonium bromide, or WITCO VARIQUAT 80MC™,
15 or a combination thereof. The formulation may further comprise water-soluble polymer, preferably polyvinyl alcohol, guar gum, (cationic or non-ionic) polydiallyl dimethyl ammonium chloride, polyacrylamide, poly (ethylene oxide), glycerol, polyethylene glycol 8000 (PEG 8000), or JAGUAR 8000™ (Guar Gum 2-hydroxypropyl ether), or a combination thereof. The formulation may also further comprise a fatty alcohol comprising from 8 to 20 carbon atoms per molecule, a solvent
20 (preferably Di(propylene glycol) methyl ether or diethylene glycol monobutyl ether or a combination thereof), and/or a carbonate salt (preferably potassium bicarbonate, sodium bicarbonate, ammonium bicarbonate, ammonium hydrogen bicarbonate, lithium bicarbonate, ammonium carbonate, or potassium carbonate, or a combination thereof). The bleaching activator is preferably water-soluble, most preferably acetylcholine chloride, monoacetin (glycerol monoacetate), diacetin (glycerol
25 diacetate), 4-cyanobenzoic acid, ethylene glycol diacetate, propylene glycol monomethyl ether acetate, methyl acetate, dimethyl glutarate, diethylene glycol monoethyl ether acetate, glycerol diacetate (Diacetin), glycerol monoacetate, glycerol triacetate, or propylene glycol diacetate, or a combination thereof. Alternatively, the bleaching activator may be water-insoluble, preferably tetraacetyl ethylenediamine (TAED), n-nonanoyloxybenzenesulfonate (NOBS), or N-acetyl

WO 03/028429

PCT/US02/29886

-8-

glucosamine, or a combination thereof. The formulation, when mixed with water, preferably has a pH value between about 9.6 and about 9.8. One embodiment of the invention consists essentially of 1-10% Benzalkonium Chloride, 1-8% Propylene Glycol Diacetate; 1-16% Hydrogen Peroxide; and 2-8% Potassium Bicarbonate.

5

The invention is also of a formulation for use in neutralization of at least one toxant, comprising: at least one cationic surfactant; at least one reactive compound, wherein the at least one reactive compound is selected from the group consisting of hydrogen peroxide, urea hydrogen peroxide, hydroperoxycarbonate, peracetic acid, sodium perborate, sodium peroxyphosphate, sodium peroxydicarbonate, and sodium percarbonate; at least one bleaching activator selected from the group consisting of O-acetyl, N-acetyl, and nitrile group bleaching activators; and at least one carbonate salt not one of the at least one reactive compounds; wherein the at least one surfactant, the at least one reactive compound, the at least one bleaching activator, and the at least one carbonate salt, when mixed with water and exposed to the at least one toxant, neutralizes the at least one toxant. In the preferred embodiment, the formulation, when mixed with water, has a pH value between about 9.6 and about 9.8 and the cationic surfactant comprises a quaternary ammonium salt, most preferably benzalkonium chloride. The at least one carbonate salt is preferably potassium bicarbonate, sodium bicarbonate, ammonium bicarbonate, ammonium hydrogen bicarbonate, lithium bicarbonate, ammonium carbonate, or potassium carbonate, or a combination thereof. One embodiment consists substantially of the at least one cationic surfactant, the at least one reactive compound, the at least one bleaching activator, and the at least one carbonate salt. The bleaching activator is preferably water-soluble, most preferably acetylcholine chloride, monoacetin (glycerol monoacetate), diacetin (glycerol diacetate), 4-cyanobenzoic acid, ethylene glycol diacetate, propylene glycol monomethyl ether acetate, methyl acetate, dimethyl glutarate, diethylene glycol monoethyl ether acetate, glycerol diacetate (Diacetin), glycerol monoacetate, glycerol triacetate, or propylene glycol diacetate, or a combination thereof.

25

WO 03/028429

PCT/US02/29886

-9-

The invention is further of a formulation for use in neutralization of at least one toxant, comprising: at least one cationic surfactant; at least one reactive compound, wherein the at least one reactive compound is selected from the group consisting of hydrogen peroxide, urea hydrogen peroxide, hydroperoxycarbonate, peracetic acid, sodium perborate, sodium peroxyphosphate, sodium peroxydicarbonate, and sodium percarbonate; and at least one water-soluble bleaching activator selected from the group consisting of O-acetyl, N-acetyl, and nitrile group bleaching activators; wherein the at least one surfactant, the at least one reactive compound, and the at least one water-soluble bleaching activator, when mixed with water and exposed to the at least one toxant, neutralizes the at least one toxant. In the preferred embodiment, the at least one water-soluble bleaching activator is acetylcholine chloride, monoacetin (glycerol monoacetate), diacetin (glycerol diacetate), 4-cyanobenzoic acid, ethylene glycol diacetate, propylene glycol monomethyl ether acetate, methyl acetate, dimethyl glutarate, diethylene glycol monoethyl ether acetate, glycerol diacetate (Diacetin), glycerol monoacetate, glycerol triacetate, or propylene glycol diacetate, or a combination thereof. The cationic surfactant preferably comprises a quaternary ammonium salt, most preferably benzalkonium chloride. The formulation preferably further comprises at least one carbonate salt, most preferably potassium bicarbonate, sodium bicarbonate, ammonium bicarbonate, ammonium hydrogen bicarbonate, lithium bicarbonate, ammonium carbonate, or potassium carbonate, or a combination thereof. The formulation, when mixed with water, has a pH value between about 9.6 and about 9.8. One embodiment consists substantially of the at least one cationic surfactant, the at least one reactive compound, and the at least one water-soluble bleaching activator.

The invention is yet further of a formulation for use in neutralization of at least one toxant, the formulation comprising: at least one solubilizing compound, selected from the group consisting of a cationic hydrotrope and a fatty alcohol comprising from 8 to 20 carbon atoms per molecule; at least one reactive compound, wherein the at least one reactive compound is selected from the group consisting of hydrogen peroxide, urea hydrogen peroxide, hydroperoxycarbonate, peracetic acid, sodium perborate, sodium peroxyphosphate, sodium peroxydicarbonate, and sodium percarbonate; and at least one bleaching activator selected from the group consisting of O-acetyl, N-acetyl, and

WO 03/028429

PCT/US02/29886

-10-

nitrile group bleaching activators; wherein the at least one solubilizing compound, the at least one reactive compound, and the at least one bleaching activator, when mixed with water and exposed to the at least one toxant, neutralizes the at least one toxant.

- 5 The invention is also of any of the above formulations packaged as a kit. In a first embodiment, the kit comprises: a premixed component comprising at least two solubilizing agents; a first component comprising at least one bleaching activator; and a second component comprising at least one reactive compound. The first embodiment may further comprise water and a base, the premixed component may additionally comprise a water-soluble polymer, the at least one bleaching
- 10 activator is preferably propylene glycol diacetate, glycerol diacetate, and/or TAED, the premixed component may additionally comprise a fatty alcohol comprising from 8 to 20 carbon atoms per molecule, the at least one reactive compound may be urea hydrogen peroxide with the second component comprising the at least one reactive compound additionally comprising sodium percarbonate, and the premixed component may additionally comprise a short-chained alcohol. A
- 15 second embodiment comprises: a first premixed component comprising at least two solubilizing agents and water; and a second premixed component comprising at least one bleaching activator and the at least one reactive compound, wherein the at least one bleaching activator is in solid form. In the second embodiment, the first premixed component preferably additionally comprises an acid, the at least one bleaching activator preferably comprises acetylcholine chloride, the at least one
- 20 reactive compound comprises urea hydrogen peroxide and the at least one bleaching activator comprises TAED, and the at least one bleaching activator is encapsulated to prevent premature reaction with the at least one reactive compound. A third embodiment comprises: a premixed component comprising at least two solubilizing agents and at least one bleaching activator; and a component comprising at least one reactive compound. In the third embodiment, the premixed
- 25 component may additionally comprise water and an acid, the component comprising the at least one reactive compound may comprise sodium percarbonate and additionally an acid, and the at least one reactive compound may comprise urea hydrogen peroxide with the component comprising the at least one reactive compound additionally comprising a mixture of potassium carbonate and potassium bicarbonate.

Objects, advantages and novel features, and further scope of applicability of the present invention will be set forth in part in the detailed description to follow, taken in conjunction with the accompanying drawings, and in part will become apparent to those skilled in the art upon
5 examination of the following, or may be learned by practice of the invention. The objects and advantages of the invention may be realized and attained by means of the instrumentalities and combinations particularly pointed out in the appended claims.

BRIEF DESCRIPTION OF THE DRAWINGS

10 The accompanying drawings, which are incorporated into and form a part of the specification, illustrate one or more embodiments of the present invention and, together with the description, serve to explain the principles of the invention. The drawings are only for the purpose of illustrating one or more preferred embodiments of the invention and are not to be construed as limiting the invention. In the drawings:

15 Fig. 1 is a graph of the effect of DF-200 components on *Bacillus globigii* (anthrax simulant) spore kill;

Fig. 2 is a schematic diagram of a preferred mixing procedure for a first rapid deployment
20 configuration of the present invention ("DF-200HF Rapid Deployment"); and

Fig. 3 is a schematic diagram of a preferred mixing procedure for a second rapid deployment configuration of the present invention ("DF-200HF Slurry Rapid Deployment").

DESCRIPTION OF THE PREFERRED EMBODIMENTS

(BEST MODES FOR CARRYING OUT THE INVENTION)

25 The present invention addresses the need for a general formulation that neutralizes the adverse effects of either or both chemical and biological toxants, where a toxant is defined as any

WO 03/028429

PCT/US02/29886

-12-

chemical or biological compound, constituent, species, or agent that through its chemical or biological action on life processes can, if left untreated, cause death, temporary incapacitation, or permanent harm to humans or animals. This includes all such chemicals or biological agents, regardless of their origin or of their method of production, and regardless of whether they are
5 produced in facilities, in munitions or elsewhere. Neutralization is defined as the mitigation, de-toxification, decontamination, or otherwise destruction of toxants to the extent that the toxants no longer cause acute adverse effects to humans or animals. The formulation and described variations of the present invention can neutralize, and does not itself contain or produce, infection, significant adverse health effects, or even fatality in animals.

10

An important subset of chemical and biological compounds that the present invention addresses is that of chemical warfare ("CW") and biological warfare ("BW") agents. However, the present invention also addresses toxants that can cause potential adverse health effects to animals, including humans, where such adverse health effects include infections, acute and chronic health
15 effects, and fatalities. Such toxants can be found in an agricultural facility, animal or dairy farm, or food products processing or packaging facility. Additionally, the present invention addresses the need for such a formulation that is itself non-toxic and non-corrosive, and that can be delivered by a variety of means and in different phases. Certain embodiments are discussed in U.S. Patent Applications Serial Nos. 09/607,586 and 09/952,940. The present invention presents additional
20 embodiments that have substantial differences over both the prior art and the earlier embodiments, as will be described below.

25

The word "formulation" is defined herein as the activated product or solution (e.g., aqueous solution) that is applied to a surface or body for the purpose of neutralization, with or without the addition of a gas (e.g., air) to create a foam. Unless otherwise specifically stated, the concentrations, constituents, or components listed herein are relative to the weight percentage of the overall activated solution. The word "water" is defined herein to broadly include: pure water, tap water, deionized water, demineralized water, saltwater, or any other liquid consisting primarily of H₂O.

-13-

One example of a minimum set of constituents for a DF-200 formulation that can achieve a significant rate of spore kill comprises four components:

- (1) a solubilizing agent selected from the group consisting of a cationic surfactant (e.g., Variquat 80MC), a cationic hydrotrope (e.g., Adogen 477), and a fatty alcohol (e.g., 1-Dodecanol);
 - (2) a bleaching activator selected from the group consisting of O-acetyl, N-acetyl, and nitrile group peroxide activators (e.g., propylene glycol diacetate);
 - (3) a reactive compound (e.g., hydrogen peroxide, peracetic acid); and
 - (4) water.
- 10 The solubilizing agent serves to effectively render the toxant susceptible to attack, while the reactive compound serves to attack and neutralize the toxant, and the bleaching activator enhances the process.

Examples of suitable cationic surfactants include: quaternary ammonium salts and polymeric quaternary salts. Examples of suitable quaternary ammonium salts include: cetyltrimethyl ammonium bromide, benzalkonium chloride, benzethonium chloride, cetylpyridinium chloride, alkyldimethylbenzylammonium salt, and tetrabutyl ammonium bromide. A preferred cationic surfactant is WITCO VARIQUAT 80MC™, which is a mixture of benzyl (C12-C16) alkyldimethylammonium chlorides. The concentration of quaternary ammonium salt used in DF-200 formulations is preferably no more than about 10%, because at higher concentrations the quaternary ammonium salt becomes significantly toxic to humans and the environment.

Examples of suitable cationic hydrotropes include: tetrapentyl ammonium bromide, triacetyl methyl ammonium bromide, and tetrabutyl ammonium bromide. A preferred cationic hydrotrope is WITCO ADOGEN 477™, which is a pentamethyltallow alkyltrimethylenediammonium dichloride.

Examples of suitable fatty alcohols include alcohols having 8-20 carbon atoms per molecule, such as: 1-dodecanol, pure dodecanol, hexadecanol, and 1-tetradecanol.

Examples of suitable bleaching activators are discussed subsequently.

Examples of suitable reactive compounds include: peroxide compounds; hydrogen peroxide; urea hydrogen peroxide; sodium perborate; sodium percarbonate; sodium carbonate perhydrate; sodium peroxyphosphate; sodium peroxydicarbonate; peroxide adducts of pyrophosphates; citrates; sodium sulfate; urea; and sodium silicate; an activated peroxide compound (e.g., hydrogen peroxide + bicarbonate); peracetic acid; oximates (e.g., butane-2,3-dione, monooximate ion, and benzohydroxamate); alkoxides (e.g., methoxide and ethoxide); aryloxides (e.g., aryl substituted benzenesulfonates); aldehydes (e.g., glutaraldehyde); peroxydisulfate; Fenton's reagent (a mixture of iron and peroxide); and sodium hypochlorite. Use of these reactive compounds in DF-200 formulations can produce a variety of negatively-charged nucleophiles, e.g., hydroxyl ions (OH^-) and hydroperoxide ions (OOH^-) produced when using hydrogen peroxide; and/or hydroperoxycarbonate ions (HCO_4^-) produced when hydrogen peroxide is combined with a carbonate salt. Hydroperoxycarbonate ions (HCO_4^-) are a much stronger oxidant than hydroxyl ions (OH^-) or hydroperoxide ions (OOH^-), and are especially effective in reacting with biological toxants. When using hydrogen peroxide in DF-200 formulations, its concentration is preferably less than about 10% because higher concentrations are significantly corrosive, especially in the range of 30-50% hydrogen peroxide concentration.

To achieve very high rates of spore kill, a carbonate salt (e.g., sodium bicarbonate or potassium bicarbonate) is preferably added to the minimum set of constituents for DF-200 formulations described above. When using a peroxide compound (e.g., hydrogen peroxide) as the reactive compound for DF-200, the added carbonate salt combines with, e.g., hydrogen peroxide to form the highly reactive hydroperoxycarbonate species (HCO_4^-). Addition of carbonate salts can also buffer the formulation to optimize the pH.

Hence, a minimum set of constituents for DF-200 formulations that can achieve a very high rate of spore kill can comprise five components:

-15-

- (1) a solubilizing agent selected from the group consisting of a cationic surfactant (e.g., Variquat 80MC), a cationic hydrotrope (e.g., Adogen 477), and a fatty alcohol (e.g., 1-Dodecanol);
- (2) a bleaching activator selected from the group consisting of O-acetyl, N-acetyl, and nitrile group peroxide activators (e.g., propylene glycol diacetate);
- (3) a reactive component (e.g., hydrogen peroxide, peracetic acid, etc.);
- (4) a carbonate salt (e.g., sodium bicarbonate); and
- (5) water.

Examples of suitable carbonate salts include: potassium bicarbonate, sodium bicarbonate, ammonium bicarbonate, ammonium hydrogen bicarbonate, lithium bicarbonate, ammonium carbonate, and potassium carbonate.

Figure 1 shows the results of decontamination tests on *Bacillus globigii* spores (initial concentration 10^7 spores/ml). The spores were exposed to four different sub-combinations of the various components of DF-200 formulations for 1 hour (the pH of the formulation was 9.8). The degree of spore kill was determined by culturing surviving organisms. As shown in Figure 3, significant spore kill was achieved by using two different combinations: (1) an aqueous solution of 2% Variquat (cationic surfactant), 2% propylene glycol diacetate (bleaching activator), and 2% hydrogen peroxide (oxidant) and (2) an aqueous solution of 2% Variquat (a cationic surfactant), 5% potassium bicarbonate (buffer and peroxide activator) and 2% hydrogen peroxide (oxidant). However, very high spore kill was achieved by using a third combination, comprising (3) an aqueous solution of 2% Variquat (a cationic surfactant), 2% propylene glycol diacetate (bleaching activator), 5% potassium bicarbonate (buffer and peroxide activator) and 2% hydrogen peroxide (oxidant).

Next, a variety of alternative embodiments and configurations of DF-200 formulations will be presented.

DF-200HF (High Foam)

The present invention provides an enhanced decontamination formulation for high foam applications ("DF-200HF"). An example of a formulation for DF-200HF comprises:

DF-200HF (High Foam)

- 1-4 % (preferably 2%) Variquat 80MC (*cationic surfactant*)
- 0.5-3% (preferably 1%) Adogen 477 (*cationic hydrotrope*)
- 0.2-0.8% (preferably 0.4%) 1-Dodecanol (*fatty alcohol*)
- 0.05-0.1% Jaguar 8000 (*cationic water-soluble polymer*)
- 0.5% Di(propylene glycol) Methyl Ether (*solvent*)
- 0.1-10% (preferably 1-4%) Hydrogen Peroxide (*oxidant*)
- 0.1-10% (preferably 2-8%) Bicarbonate salt (*buffer and peroxide activator*)
- 1-4% Propylene Glycol Diacetate (*bleaching activator*)
- 67-97% Water

This formulation is effective at a pH value between 7.5 and 10.5. This formulation can be adjusted to a pH value between 9.6 and 9.8 for optimal decontamination of all target agents. This "high-foam" formulation includes a cationic water-soluble polymer (e.g., Jaguar 8000™), which increases the bulk viscosity of the solution and produces a more stable foam.

Examples of suitable non-anionic water-soluble polymers include: polyvinyl alcohol, guar gum, (cationic or non-ionic) polydiallyl dimethyl ammonium chloride, polyacrylamide, polyethylene glycol 8000 (PEG 8000), and JAGUAR 8000™ (Guar Gum 2-hydroxypropyl ether). A cationic polymer is preferred over a non-ionic polymer; an anionic polymer does not work well. The fatty alcohol 1-dodecanol serves to increase the surface viscosity of the foam lamellae to also increase foam stability against drainage and bubble collapse.

The term 'High Foam' refers to the ability of a formulation to form a highly stable and persistent foam, whereas a 'Low Foam' formulation forms a much less stable foam. The following tables show the improved performance of DF-200HF and DF-200LF as compared to DF-100A. The notation "ND" refers to a concentration below detectable limits, and "PGD" refers to propylene glycol diacetate (a preferred bleaching activator).

WO 03/028429

PCT/US02/29886

-18-

Table 1: Summary of the reaction rates for Mustard simulant (2-Chloroethyl phenyl sulfide).

Formulation	Mustard Simulant (% Decontaminated)		
	1 Minute	15 Minutes	60 Minutes
DF-100A(pH 8)	18	42	81
DF-100A (pH 9.2)	16	38	83
DF-200HF (2% PGD / 3% H ₂ O ₂ / 4.5% Bicarb)	42	62	ND
DF-200HF (2% PGD / 3.5% H ₂ O ₂ /4% Bicarb)	94	98	ND
DF-200LF (2.5% PGD / 3% H ₂ O ₂ / 4.5% Bicarb)	55	91	ND

5

Table 2: Summary of the reaction rates for VX simulant (O-Ethyl S-Ethyl Phenylphosphonothioate).

Formulation	VX Simulant (% Decontaminated)		
	1 Minute	15 Minutes	60 Minutes
DF-100A (pH 10)	45	99	ND
DF-100A (pH 9.2)	33	71	93
DF-200HF (2% PGD / 3% H ₂ O ₂ / 4.5% Bicarb)	63	98	ND
DF-200HF (2% PGD / 3.5% H ₂ O ₂ / 4% Bicarb)	66	99	ND
DF-200LF (2.5% PGD / 3% H ₂ O ₂ / 4.5 Bicarb)	79	98	ND

10

Table 3: Summary of the reaction rates for G Agent simulant (Diphenyl chlorophosphate).

Formulated	G Agent Simulant (% Decontaminated)		
	1 Minute	15 Minutes	60 Minutes
DF-100A (pH 8)	53	ND	ND
DF-100A (pH 9.2)	ND	ND	ND
DF-200 HF (2% PGD / 3% H ₂ O ₂ /4.5 Bicarb)	ND	ND	ND
DF-200HF (2% PGD / 3.5% H ₂ O ₂ 4% Bicarb)	ND	ND	ND
DF-200LF (2.5% PGD / 3% H ₂ O ₂ 4.5% Bicarb)	ND	ND	ND

15

Table 4: Summary of the kill rates for Anthrax simulant (Bacillus globigii spores)

Formulation	Anthrax Simulant	Anthrax Simulant
	% Kill after 30 Minutes	% Kill after 60 Minutes
DF-100A (pH 8)	99.99	99.99999
DF-100A (pH 9.2)	90	99.9
DF-200HF (2% PGD / 3% H ₂ O ₂ / 4.5 Bicarb)	99.99999	99.99999
DF-200LF (2.5% PGD / 3% H ₂ O ₂ / 4.5 Bicarb)	99.99999	99.99999

WO 03/028429

PCT/US02/29886

-19-

Differences between formulations of DF-200 and DF-100/100A include:

- DF-200 is active against all agents at a single pH. The formulation is effective at pH values between about 7.5 and 10.5; is more effective at pH values between about 9.2 and 9.8; and is most effective at pH values between about pH 9.6 and 9.8;
- 5 • DF-200 has better performance against Mustard;
- DF-200 has better performance against bacterial spores;
- DF-200 has lower concentrations of both the cationic surfactant and/or the cationic hydrotrope, which further lowers the (already low) toxicity and corrosivity properties of the formulation;
- 10 • DF-200 has a lower concentration of the foam stability component, 1-Dodecanol;
- DF-200 doesn't use a short-chained alcohol (e.g., isobutanol, isopropanol), which causes flammability problems when the formulation is packaged in a concentrated form;
- DF-200 doesn't use Diethylene Glycol MonoButyl Ether (DEGMBE), which can cause false alarms on some chemical agent sensors and detectors (especially older sensors which are
- 15 used in some military settings); and
- DF-200 can contain a lower concentration of hydrogen peroxide, which also reduces the (already low) toxicity and corrosivity properties of the formulation.

Additional differences between DF-200 and DF-100A include:


- 20 • DF-200 performs optimally at a higher pH (about 9.6 to 9.8) as compared to DF-100A. However, note that this is the typical pH value for common household products such as laundry detergents, shampoos, and dishwashing detergents; and
- DF-200 has more individual components that should be stored separately (e.g., the hydrogen peroxide and the bleaching activator) from the bulk formulation until use, as compared to
- 25 DF-100A (where only one component, hydrogen peroxide, should be stored separately). This will be discussed in greater detail below.

One reason for the better performance of DF-200 formulations (e.g., DF-200HF and DF-200LF) over DF-100 and DF-100A formulations is the addition of a bleaching activator (e.g.,

-20-

propylene glycol diacetate). Bleaching activators can be compounds with O- or N- bounded acetyl groups that react with the strongly nucleophilic hydroperoxy anion (OOH^-) to yield peroxygenated species, which are more efficient oxidizers than hydrogen peroxide alone.

5



10

Since the 1950's, a number of different bleaching activators have been used in commercial laundry detergents, as well as other commercial products. The most common activators are tetraacetyl ethylenediamine (TAED), which is primarily used in Europe and Asia; and n-nonanoyloxybenzenesulfonate (NOBS), which is primarily used in the United States. NOBS is a proprietary chemical of the Proctor and Gamble company. In a laundry detergent, hydrogen peroxide is provided in a solid form (usually as sodium perborate, which reacts in water to form the hydroperoxy anion). The addition of a bleaching activator greatly enhances the ability of a laundry detergent to remove stains from clothing.

15

It should be noted that TAED and NOBS bleaching activators are extremely insoluble in water (e.g., TAED is only 0.1% soluble at 25°C). To get around this problem in a laundry detergent, the solid TAED or NOBS particles are kept in suspension by the agitating action of the washing machine, where they slowly react with the hydrogen peroxide in the detergent. However, agitation in the field of DF-200 formulations presents practical problems; hence, a water-soluble bleaching activator is preferred.


20

25

Useful water-soluble bleaching activators include short-chained organic compounds that contain an ester bond, e.g., ethylene glycol diacetate, propylene glycol monomethyl ether acetate, methyl acetate, dimethyl glutarate, diethylene glycol monoethyl ether acetate, glycerol diacetate (Diacetin), glycerol monoacetate, glycerol triacetate, and propylene glycol diacetate. A preferred water-soluble bleaching activator is propylene glycol diacetate (PGD), which is shown below.

30

-21-



- 5 This molecule reacts with hydroperoxy anions (OOH^-), giving up the ester bonds to form two peroxygenated molecules.

Propylene glycol diacetate also acts as an organic solvent that is highly effective in solubilizing insoluble organic molecules (e.g., chemical warfare agents, as well as foam stabilizers/boosters (such as 1-dodecanol and Lauramide DEA)). Therefore, an added function of this compound is that it can be used to supplement the diethylene glycol monobutyl ether (DEGMBE) solvent that is used in DF-100 and DF-100A, or to supplement the Di(propylene glycol) methyl ether solvent used in some DF-200 formulations, thereby allowing the propylene glycol diacetate to serve a dual purpose (i.e., solvent and bleaching activator).

15

Bleaching activators are generally not stable in water for long periods of time. This is especially true when the aqueous solution is at a high pH (>10). Therefore, for long shelf life, the propylene glycol diacetate (or other bleaching activator) is preferably stored separate from the aqueous solution until use. This is not unlike other products that utilize bleach activators (e.g., laundry detergents), where all the components of the formulation are kept dry and separated until use (in the case of laundry detergent, the bleaching activator is encapsulated to prevent it from reacting with the peroxide component until both components are mixed in water).

20

Another example of a water-soluble bleaching activator is ethylene glycol diacetate, which works well in DF-200 formulations. However, when ethylene glycol diacetate reacts with hydrogen peroxide, it forms ethylene glycol (i.e., anti-freeze), which is a relatively toxic byproduct. Propylene glycol diacetate, on the other hand, does not form this relatively toxic byproduct.

25

DF-200NF (Non-Foaming)

30

The present invention is also of a non-foaming embodiment ("DF-200NF") that may be used for specific applications, e.g., the kill of biological organisms, batch processing (such as in chemical

WO 03/028429

PCT/US02/29886

-22-

agent demilitarization neutralization processes, i.e., in a bath of solution), or spray applications. A preferred example of this formulation comprises (amounts illustrative):

DF-200NF (Non-Foaming)

- 5 1-10% (preferably 2.5%) Benzalkonium Chloride (*cationic surfactant*)
 1-8% Propylene Glycol Diacetate (*bleaching activator*)
 1-16% Hydrogen Peroxide (*oxidant*)
 2-8% Potassium Bicarbonate (*buffer and peroxide activator*)
 65.5-93.5% Water

10

The formulation can be adjusted to a pH value between about 9.6 and 9.8 for optimum performance, and is effective for decontamination of all target agents.

DF-100E

- 15 The present invention is also of an enhanced version of DF-100A that utilizes the propylene glycol diacetate bleaching activator. A preferred embodiment of this enhanced formulation, ("DF-100E") comprises (amounts illustrative):

20

WO 03/028429

PCT/US02/29886

-23-

DF-100E

5.3% Variquat 80MC

2.8% Adogen 477

0.65% 1-Dodecanol

5 0.5% Isobutanol

0.1% Jaguar 8000

1.35% Diethylene Glycol Monobutyl Ether

2-8% Bicarbonate Salt

1-4% Hydrogen Peroxide

10 1-4% Propylene Glycol Diacetate

73-85% Water

This formulation can be adjusted to a pH value between about 9.6 and 9.8 for optimal performance against all agents. The performance of DF-100E (2% PGD / 3.00% H₂O₂ / 3.75% Bicarbonate salt)

15 against chemical simulants is shown below in Table 5:

Table 5: Summary of the reaction rates for the DF-100E formulation in kinetic testing.

Simulant	% Decontaminated		
	1 Minute	15 Minutes	60 Minutes
Mustard (HD)	83	92	ND
G Agents	ND	ND	ND
VX	66	96	ND

20

Tests against the anthrax spore simulant (*Bacillus globigii* spores) demonstrated 99.9999% (7-Log) kill after a 30 minute exposure to DF-100E.

Other bleaching activators (such as water-insoluble NOBS or TAED) can be used in place of Propylene Glycol Diacetate in DF-100E. However, as noted above, this produces a slurry mixture instead of a true liquid solution.

25

The following table summarizes some differences between DF-100, DF-100A, DF-100E, DF-200HF, DF-200LF, DF-200NF, and DF-200HF Slurry:

Table 6: Comparison of Various Decontamination Formulations

5

Formulation	Requires pH Adjustment	Improved Kinetics	Reduction of Flammable Constituents	Forms Highly Stable Foam	Can Use Saltwater for Makeup Water
DF-100	Y	N	N	Y	N
DF-100A	N	N	N	Y	N
DF-100E	N	Y	N	Y	Y
DF-200HF	N	Y	Y	Y	Y
DF-200LF	N	Y	Y	N	Y
DF-200NF	N	Y	Y	N	Y
DF-200HF-Slurry	N	Y	Y	Y	Y

Kit Configurations

10 In the following sections, various examples of 2-part, 3-part, and 4-part "kit" configurations are shown for the different embodiments of DF-200 formulations. In general, the 2-part and 3-part kits have the bulk of the water already "pre-packaged" in one of the two (or three) containers (usually the foam component). This allows for rapid deployment of the decontamination solution, the use of small-scale units (e.g., backpacks), and doesn't require any extra water to be provided in the field.

15 Conversely, the 4-part kits generally require that the make-up water is added in the field at, or near, the site of contamination. This allows the "package" containing the other three parts to be much lighter, which makes it easier to ship and store. However, a source of make-up is required in the field (which can be saltwater).

WO 03/028429

PCT/US02/29886

-25-

In general, the DF-200 formulations can be configured either way, with the bulk of the water "pre-packaged", or without, depending on the application.

5 DF-200HF (Kit Configuration)

The DF-200HF formulation can be configured as a 4-part kit, and then prepared for field use as follows (amounts illustrative):

DF-200 HF (4-part kit)

10 Part A (Foam Concentrate):

20 g Variquat 80MC

10 g Adogen 477

4 g b-Dodecanol

1 g Jaguar 8000 Polymer

15 5 g Di(propylene glycol) methyl ether

7.5 g Potassium Bicarbonate

141 g Water

Part B (Solid Component):

50 g Sodium Percarbonate

20 50 g Urea Hydrogen Peroxide

Part C (Bleaching Activator):

20 g Propylene Glycol Diacetate

Part D (Make-up Water):

800 g Water

25

In this example of a 4-part configuration, the bulk of the water is not included in the "package" (i.e., Parts A, B, and C), which minimizes the weight of the package for shipping and storage. Here, the make-up water (Part D) would be supplied in the field at, or near, the site of contamination. The pH of the formulation can be adjusted to be between about 9.6 and 9.8 for

optimal performance. The formulation as described above will produce 1 liter of "high" foam solution. In this example, sodium percarbonate supplies a portion of the hydrogen peroxide, a portion of the bicarbonate, and a base for buffering the solution. The remainder of the hydrogen peroxide is supplied by the urea hydrogen peroxide. The total hydrogen peroxide concentration is approximately 5 3% in this example. The viscosity of the formulation can be adjusted to be between about 4-9 mm²/s.

A variety of different methods can be used in the field to mix the DF-200HF formulation configured as a 4-part kit, for example:

10 Method 1: Supply the make-up water (Part D). Then, mix Part B into Part D. Then, add Parts C and A to Part B+D. Use, preferably, within 8 hours.

Method 2: Mix Part C into Part A. Supply the make-up water (Part D). Then, mix Part B into Part D. Keep separate until use. When use is required, mix Part A+C into Part B+D. Use, 15 preferably, within 8 hours of first mixing Parts A+C into Parts B+D.

In general, activated DF-200 formulations are used preferably within 8 hours after mixing, however, they still can be effective for up to 24 hours and longer. DF-200HF (High Foam) can be applied to a surface for a long period of time, and then rinsed off. However, DF-200LF (Low Foam) 20 can be used in a different manner than the DF-100/100A and DF-200HF formulations. Instead of leaving DF-200LF on a surface for long periods of time, it can be applied to a surface, left for a relatively short period of time (e.g., 15-60 minutes), and then rinsed off with a high pressure freshwater or salt water spray. This will minimize corrosion of the material to which it is applied, which will make it especially useful for decontaminating aircraft and other equipment where corrosion 25 is a concern. It will also minimize the time required for decontamination, which is especially advantageous for military use (on the battlefield or at fixed sites).

WO 03/028429

PCT/US02/29886

-27-

Saltwater can also be effectively used as the make-up water (Part D) for DF-200 formulations. The table below shows the results of kinetic tests using DF-200HF (2% PGD / 3.50% H₂O₂ / 4.0% Bicarbonate salt) with saltwater (~35,000 ppm total dissolved solids):

- 5 Table 7: Summary of the reaction rates for DF-200HF formulation (2% PGD / 3.5% H₂O₂ / 4.0% Bicarbonate salt) with saltwater used as the make-up water (Part D).

Simulant	% Decontaminated		
	1 Minute	15 Minutes	60 Minutes
Mustard	24	42	89
G Agents	ND	ND	ND
VX	62	96	>99

10

Tests against the anthrax spore simulant (*Bacillus globigii* spores) demonstrated 99.9999% (7-Log) kill after a one hour exposure to DF-200HF with saltwater used as the make-up water.

- Surface testing was conducted with DF-200HF against the Mustard and VX simulant. For this test, 8 mg of simulant was applied to a 2" diameter test coupon made of CARC (Chemical Agent Resistant Coating). CARC is a material commonly used to paint military vehicles to protect them against chemical agent attack. After waiting one hour, the test coupon was placed in a horizontal position and covered with 1.0 g of DF-200HF (2% PGD / 3.5% H₂O₂ / 4.0% Bicarbonate salt). After 60 minutes, the test coupon was immersed in 25 ml of acetonitrile for 15 minutes to extract unreacted simulant from the surface. The extraction solvent (acetonitrile) was then analyzed for the unreacted simulant. Results (shown in Table 8) demonstrate more effective decontamination of the test coupon as compared to DF-100A.
- 15
- 20

WO 03/028429

PCT/US02/29886

-28-

Table 8: Results of Surface Testing of DF-200HF on CARC.

<u>Decon Formulation</u>	<u>mg VX Simulant (Unreacted simulant on surface after 60 minutes)</u>	<u>Mg Mustard Simulant (Unreacted simulant on surface after 60 minutes)</u>
Control	8.0 $\pm$ 0.3	8.0 $\pm$ 0.3
DF-100A (pH 9.2)	2.9	4.5
DF-200HF (2% PGD / 3.5% H ₂ O ₂ / 4.0% Bicarb)	1.4	2.5

5

DF-200HF Slurry (Kit Configuration)

Insoluble bleach activators (such as TAED, NOBS, and N-acetyl glucosamine) can be utilized in place of the (water-soluble) propylene glycol diacetate for DF-200 formulations. However, in this case, the formulation results in a slurry when mixed with water, instead of a true aqueous solution.

10

The present invention also provides a method to utilize a water-insoluble solid bleach activator (e.g., TAED) to produce a reactive slurry (wherein a slurry is defined as a watery mixture that includes insoluble, undissolved matter). This embodiment, designated "DF-200HF Slurry", is a modification of the DF-200HF formulation. An example of a 4-part kit configuration is shown below (amounts illustrative):

15

WO 03/028429

PCT/US02/29886

-29-

DF-200HF Slurry (4-part kit)Part A (Foam Concentrate):

20 g Variquat 80MC
10 g Adogen 477
5 4 g 1-Dodecanol
1 g Jaguar 8000 Polymer
5 g Di(propylene glycol) methyl ether
7.5 g Potassium Bicarbonate
161 g Water

Part B (Solid Component):

50 g Sodium Percarbonate
50 g Urea Hydrogen Peroxide

Part C (Bleaching Activator):

10 g TAED (*preferably encapsulated TAED, such as Warwick B637*)

Part D (Make-up Water):

800 g Water (*can be freshwater or saltwater supplied at the site where use is to occur*)

The formulation as described above will produce 1 liter of foam solution. The pH of the final
20 formulation can be adjusted to be between about 9.6 and 9.8 for optimal performance. The following
mixing procedure can be used: Mix Part B into Part D. Then, add Parts C and A to Parts B+D. Use,
preferably, within 8 hours.

The performance against each chemical agent simulant for DF-200HF Slurry (1% TAED / 3%
25 H₂O₂ / 4% Bicarbonate salt) is shown below in Table 9:

Table 9: Summary of the reaction rates for the DF-200HF Slurry formulation in kinetic testing. Note that improved performance can be achieved by using higher concentrations of TAED (e.g., 2% TAED, instead of 1% TAED).

5

Simulant	% Decontaminated		
	1 Minute	15 Minutes	60 Minutes
Mustard (HD)	67	96	ND
G Agents	ND	ND	ND
VX	33	95	ND

The above examples of different embodiments of DF-200 would typically be used in large-scale operations where dedicated deployment equipment and a source of make-up water is readily available (e.g., for use by the military to decontaminate 'fixed sites' such as airbases and seaports), or used to minimize the volume of 'pre-packaged' water in order to minimize the weight of the formulation that needs to be shipped and stored.

DF-200 Rapid Deployment Configurations

The present invention is also of configurations emphasizing the rapid deployment of DF-200 formulations, and/or its deployment using small-scale deployment equipment (such as hand-held units, backpack units, or units mounted on small dollies). For these applications, all of the water is 'pre-packaged' into the formulation, so that no extra water is required in the field. A first example of a 3-part kit configuration for a Rapid Deployment version of DF-200HF, "DF-200HF Rapid Deployment #1", comprises (amounts illustrative):

WO 03/028429

PCT/US02/29886

-31-

DF-200HF Rapid Deployment #1 (3-part kit)Part A (Liquid Foam Component):

20 g Variquat 80MC

10 g Adogen 477

5 4 g 1-Dodecanol

5 g Poly (Ethylene Oxide)

8 g Diethylene Glycol Monobutyl Ether

5 g Isobutanol

45 g Potassium Bicarbonate

10 approx. 19 g Potassium Hydroxide (*the pH of Part A should be approximately 10.2*)

933 g Water

Part B (Solid Oxidant Component):

97 g Urea Hydrogen Peroxide

Part C (Liquid Bleaching Activator):

15 20 g Propylene Glycol Diacetate

This configuration will produce 1 liter of foam solution. The pH of the final formulation can be adjusted to be between about 9.6 and 9.8 for optimal performance. The following mixing procedure can be used: Mix Part B into Part A. After dissolution of the urea hydrogen peroxide, add Part C to Part A+B. Use, preferably, within 8 hours. The performance of DF-200HF Rapid Deployment against chemical agent simulants is shown below in Table 10:

Table 10: Reaction rates from kinetic testing of DF-200HF Rapid Deployment #1 configuration.

Simulant	% Decontaminated		
	1 Minute	15 Minutes	60 Minutes
Mustard (HD)	48	82	ND
G Agents	ND	ND	ND
VX	71	97	>99

WO 03/028429

PCT/US02/29886

-32-

Tests against the anthrax spore simulant (*Bacillus globigii* spores) demonstrated 99.9999% (7-Log) kill after a 30 minute exposure to DF-200HF Rapid Deployment.

5 A schematic example of a preferred mixing procedure for this first example of a rapid deployment configuration of DF-200HF is shown in Figure 2. Urea hydrogen peroxide dissolves rapidly in water. Therefore, the formulation can be prepared and deployed in a very short time at the scene of an incident involving chemical or biological warfare agents, making it ideal for use by civilian first responders (firefighters, HazMat units, police officers, and others who would be the first to arrive
10 at the location of a CBW attack), and/or the military.

However, the particular bleaching activator (propylene glycol diacetate) used in this formulation is not stable in an aqueous solution where the pH is greater than approximately 9.9. Therefore, it is important to mix the right components in the correct order. For example, if Part C is
15 mixed into Part A before the addition of Part B, there may be some loss of activity in DF-200HF since the propylene glycol diacetate is exposed to a solution having a pH value >9.9. This is not true, however, if Part B is added to Part A before the addition of Part C, since the addition of Part B to Part A brings the pH of the Part A+B mixture to a value below about 9.9.

20 The solvent, diethylene glycol monobutyl ether, used in Part A (the foam solution) of the first example shown above for DF-200HF Rapid Deployment #1 is different than the solvent that was used in the previously described DF-200HF formulation (Di(propylene glycol) methyl ether), because Di(propylene glycol) methyl ether is not stable in the high pH environment required for the foam component (Part A) in the rapid deployment configuration. Also, note that a short-chained alcohol
25 (i.e., isobutanol) has been added to the foam component (Part A) in the rapid deployment configuration #1 of DF-200HF. While this low molecular weight alcohol can cause flammability problems in highly concentrated configurations of DF-200HF, it is not a problem in the less concentrated configurations described herein. The use of isobutanol also helps solubilize the 1-dodecanol in Part A, and improves the kinetics (chemical reactivity) of the formulation. In addition,

-33-

the formulation preferably uses a different polymer, poly (ethylene oxide), than the polymer used in the other earlier described DF-200 formulations (i.e., Jaguar 8000). This alternative polymer is used because Jaguar 8000 is also not stable in the high pH environment of the liquid foam portion (Part A) of the rapid deployment formulation. Accordingly, a preferred formulation for DF-200HF Rapid

5 Deployment #1 comprises:

DF-200HF Rapid Deployment #1

1-4% (preferably 2%) Variquat 80MC (*cationic surfactant*)

0.5-3% (preferably 1%) Adogen 477 (*cationic hydrotrope*)

10 0.2-0.8% (preferably 0.4%) 1-Dodecanol (*fatty alcohol*)

0.5-8% (preferably 0.5%) Poly (Ethylene Glycol) (*polymer*)

0.6-1.2% (preferably 0.8%) Diethylene Glycol Monobutyl Ether (*solvent*)

0-1% (preferably 0.5%) Isobutanol (*short-chained alcohol*)

0.1-10% (preferably 2-8%) Bicarbonate salt (*buffer and peroxide activator*)

15 0.1-10% (preferably 1-4%) Hydrogen Peroxide (*oxidant*)

0.1-10% (preferably 1-4%) Propylene Glycol Diacetate (*bleaching activator*)

52-97% Water

The formulation can be adjusted to a pH value between about 9.6 and 9.8 for optimal performance,
20 and is effective for decontamination of all target agents.

A second example of a 3-part kit configuration for a Rapid Deployment version of DF-200HF, "DF-200HF Rapid Deployment #2", comprises (amounts illustrative):

WO 03/028429

PCT/US02/29886

-34-

DF-200HF Rapid Deployment #2 (3-part kit)Part A (Liquid Foam Component):

20 g Variquat 80MC

10 g Adogen 477

5 4 g 1-Dodecanol

20 g Polyethylene Glycol 8000 polymer

8 g Diethylene Glycol Monobutyl Ether

5 g Isobutanol

50 g Potassium Bicarbonate

10 approx. 25 g Potassium Hydroxide (*the pH of Part A should be approximately 10.2*)

933 g Water

Part B (Solid Oxidant Component):

97 g Urea Hydrogen Peroxide

Part C (Liquid Bleaching Activator):

15 20 g Propylene Glycol Diacetate

In this second example, Polyethylene Glycol 8000 polymer replaced the poly (Ethylene Oxide) polymer used in DF-200HF Rapid Deployment #1.

20 A third example of a 3-part kit configuration for a Rapid Deployment version of DF-200HF, "DF-200HF Rapid Deployment #3", comprises (amounts illustrative):

WO 03/028429

PCT/US02/29886

-35-

DF-200HF Rapid Deployment #3 (3-part kit)Part A (Liquid Foam Component):

20 g Variquat 80MC

10 g Adogen 477

5 4 g 1-Dodecanol

20 g Polyethylene Glycol 8000 polymer

10 g Hexylene Glycol

45 g Potassium Carbonate

5 g Potassium Bicarbonate

10 700 g Water

Part B (Solid Oxidant Component):

83 g Urea Hydrogen Peroxide

Part C (Liquid Bleaching Activator):

20 g Glycol Diacetate (i.e., Diacetin)

15

In this third example, Polyethylene Glycol 8000 polymer replaced the poly (Ethylene Oxide) polymer used in DF-200HF Rapid Deployment #1 as a water-soluble polymer. Also, Hexylene Glycol replaced Diethylene Glycol Monobutyl Ether and Isobutanol used as a solvents. Finally, Glycol Diacetate (i.e., Diacetin) replaced Propylene Glycol Diacetate used as the bleaching activator.

20 These changes in the third example were made to reduce or eliminate the use of short-chained alcohols and/or high vapor-pressure solvents to prevent possible problems with very long-term (months to years) shelf life of the liquid foam component (Part A), especially at high ambient storage temperatures, due to evaporation of the most-volatile components. Note that the combination of 45 grams of potassium carbonate and the 5 grams of potassium bicarbonate were chosen to supply

25 both the right amount of carbonate/bicarbonate, and to adjust the pH appropriately. Alternatively, 50 grams of potassium bicarbonate could have been used (with no potassium carbonate), and then the right amount of potassium hydroxide (base) could have been added to increase the pH to the desired value, as is well-known in the art.

-36-

DF-200HF Slurry Rapid Deployment

The present invention is also of a 2-part kit configuration of a rapid deployment embodiment of the DF-200HF Slurry embodiment ("DF-200HF Slurry Rapid Deployment"), in which TAED (or other solid peroxide activator) is utilized as the bleaching activator. This example of a rapid deployment configuration also requires no additional water to be added in the field (amounts illustrative):

DF-200HF-Slurry Rapid Deployment (2-part kit)Part A (Liquid Foam Component):

10 20 g Variquat 80MC
 10 g Adogen 477
 4 g 1-Dodecanol
 5 g Poly (Ethylene Glycol)
 8 g Diethylene Glycol Monobutyl Ether
15 5 g Isobutanol
 50 g Potassium Bicarbonate
 28 g Potassium Hydroxide (*the pH of Part A should be approximately*
10.4)
 953 g Water

20 Part B (Solid Oxidant and Bleaching Activator):

 97 g Urea Hydrogen Peroxide
 30 g TAED (preferably in encapsulated form, such as Warwick
International B637)

25 This formulation will produce 1 liter of foam solution. The pH of the final
formulation can be adjusted to be between 9.6 and 9.8 for optimal performance. The
following procedure can be used to mix the formulation: Mix Part B into Part A.
Then, wait for at least one minute before use to allow time for the TAED to react with
the hydrogen peroxide. Use, preferably, within 8 hours. It is useful to note that TAED
30 will not immediately dissolve in water, but will remain as solid particles for at least
15-20

WO 03/028429

PCT/US02/29886

-37-

minutes. Therefore, a filter or screen may be required so that the undissolved TAED particles will not damage or clog any pumps or other components of the deployment device. However, the formulation is ready for use approximately 1 minute after addition of the TAED particles in Part B to Part A.

5

It is useful to employ an encapsulated form of TAED in this configuration for two reasons. First, the protective coating (which slowly dissolves in water) will protect the TAED so that it will not react with the urea hydrogen peroxide during storage. Second, the coating will protect the TAED from the high pH conditions in Part A until the urea hydrogen peroxide dissolves and lowers the pH of the formulation to a value below approximately 9.9. TAED should be used in a similar manner as propylene glycol diacetate with respect to protecting the activator against exposure to high pH solutions. TAED will lose much of its effectiveness as a bleaching activator if it is exposed to solutions with a pH greater than 9.9. Therefore, the use of an encapsulated form of TAED will minimize its exposure to the high pH conditions immediately after Part B is mixed into Part A. The performance of DF-200HF Slurry Rapid Deployment against chemical agent simulants is given in Table 11 below.

15

Table 11: Reaction rates in kinetic testing for the DF-200HF Slurry Rapid Deployment formulation.

20

Simulant	% Decontaminated		
	1 Minute	15 Minutes	60 Minutes
Mustard (HD)	76	97	ND
G Agents	ND	ND	ND
VX	57	97	ND

Tests against the anthrax spore simulant (*Bacillus globigii* spores) demonstrated 99.9999% (7-Log) kill after a 30 minute exposure to DF-200HF Slurry Rapid Deployment.

25

A schematic illustrating an example of a process for field mixing DF-200HF Slurry Rapid Deployment is shown in Figure 3.

The above are only a few examples of rapid deployment configurations for DF-200 formulations. As understood by one of ordinary skill in the art, other rapid deployment configurations are possible by utilizing the fundamental considerations presented in this application.

5

The present invention is also of a method for preparing the foam component (Part A) of the rapid deployment configurations. The following is an example of such a method:

1. Place the appropriate mass of water in a mixing vessel.
2. While stirring the water in the mixing vessel, add the bicarbonate salt. Stir until
10 completely dissolved.
3. Slowly add the Poly (Ethylene Glycol) polymer to the mixing vessel while rapidly stirring. Be careful to avoid lumps. Stir for approximately 30 minutes, at a minimum.
4. While stirring the foam solution in the mixing vessel, add the Variquat 80MC. Stir
until completely mixed.
- 15 5. While stirring the foam solution in the mixing vessel, add the Adogen 477. Stir until completely mixed.
6. Mix the diethylene glycol monobutyl ether, isobutanol, and 1-dodecanol in a separate vessel. Slowly add this mixture to the foam solution while stirring.
7. While stirring the foam solution in the mixing vessel, slowly add solid KOH until the
20 pH reaches the appropriate value.

Alternative DF-200 Formulations

The invention is also of the following alternative DF-200 formulations:

1. An alternative formulation that includes *propylene glycol* to lower the freezing point of
25 the solution;
2. An alternative formulation that utilizes *sodium percarbonate* as a solid source of hydrogen peroxide;
3. An alternative formulation that includes a *corrosion inhibitor*;

WO 03/028429

PCT/US02/29886

-39-

4. An alternative formulation that includes *glycerol* as a viscosity builder for operations such as skin decontamination;

5. An alternative formulation that utilizes *O-acetyl* bleaching activators, including one which is available in solid form; and

5 6. An alternative formulation that utilizes a bleaching activator containing a *nitrile group*.

DF-200 with Propylene Glycol

The following is a first example of a 2-part kit configuration for DF-200HF that includes propylene glycol as a freezing point depressant, and where all of the water is 'pre-packaged' in Part

10 A, comprising (amounts illustrative):

DF-200HF Rapid Deployment with Propylene Glycol, first example (2-part kit)

Part A (Liquid Foam Component):

20 g Variquat 80MC

15 10 g Adogen 477

20 g Poly (Ethylene Glycol) (*MW 8000*)

8 g Diethylene Glycol Monobutyl Ether

5 g Isobutanol

4 g 1-Dodecanol

20 20 g Propylene Glycol Diacetate

150 g Propylene Glycol (*freeze-point depressant*)

approx. 6 g of 10% HCl Solution (*sufficient to give a final pH of 2.5 in Part A*)

777 g Water

Part B (Solid Additive):

25 97 g Urea Hydrogen Peroxide

12 g Potassium Bicarbonate

38 g Potassium Carbonate (*buffer, to adjust final pH*)

This formulation will produce 1 liter of foam solution. The pH of the final formulation can be adjusted to be between about 9.6 and 9.8 for optimal performance. A person of ordinary skill in the art will understand that the ratio of potassium carbonate to potassium bicarbonate used in Part B can be adjusted to achieve the desired final pH of the formulation (preferably about 9.6 to about 9.8). Hence, in this example, the potassium carbonate serves as both a base and a source of carbonate/bicarbonate. To prepare this formulation, mix Part B into Part A. Use, preferably, within 8 hours. The performance of this first example of DF-200HF with propylene glycol against chemical agent simulants is shown in Table 12.

Table 12: Reaction rates from kinetic testing for DF-200HF with propylene glycol (first example).

Simulant	% Decontaminated		
	1 Minute	15 Minutes	60 Minutes
Mustard (HD)	16	80	ND
G Agents	ND	ND	ND
VX	66	90	>99

Tests against the anthrax spore simulant (*Bacillus globigii* spores) demonstrated 99.9999% (7-Log) kill after a 30 minute exposure to DF-200HF with propylene glycol (first example).

When all of the water is "pre-packaged" in Part A, the mixing of the formulation for use can be accomplished in a very short time since it only consists of two parts. Therefore, it could be deployed very rapidly at the scene of an incident involving chemical and biological warfare agents. This configuration is ideal for use the civilian first responder (firefighter, HazMat units, police officers, and others who would be the first to arrive at the location of a CBW attack). However, it is heavier to carry than other configurations that add water in the field.

This configuration also incorporates the bleaching activator, propylene glycol diacetate, into the foam component Part A (rather than storing it as a separate, third component). This is possible because the pH of the foam component is less than 3. Propylene glycol diacetate will hydrolyze in

WO 03/028429

PCT/US02/29886

-41-

solutions of pH greater than 3, but is hydrolytically stable in solutions of pH less than 3. This configuration also uses the polyethylene glycol polymer (PEG 8000) for viscosity enhancement. This polymer is used in many cosmetics and is extremely soluble and stable in water. In addition, it is easier to mix into solution than Jaguar 8000 or a high molecular weight poly(ethylene oxide), since it
5 does not have the tendency to clump.

This configuration includes propylene glycol as a freeze-point depressant. Propylene glycol is considered to be an environmentally friendly antifreeze. In this case, the concentration is approximately 15% by weight, which lowers the freezing point of Part A to approximately -20°C. This
10 configuration has also been tested with good results with propylene glycol concentrations as high as 40% by weight.

An alternative to the first example of DF-200HF with Propylene Glycol shown above is to use sodium percarbonate as the source of the bicarbonate and as a portion of the peroxide in Part B,
15 instead of using urea hydrogen peroxide. This substitution is useful because sodium percarbonate is much less expensive than urea hydrogen peroxide. This second example of DF-200HF with Propylene Glycol is shown below (amounts illustrative):

DF-200HF Rapid Deployment with Propylene Glycol, second example (2-part kit)Part A (Liquid Foam Component):

20 g Variquat 80MC

10 g Adogen 477

5 20 g Poly (Ethylene Glycol) (MW 8000)

8 g Diethylene Glycol Monobutyl Ether

5 g Isobutanol

4 g 1-Dodecanol

20 g Propylene Glycol Diacetate

10 150 g Propylene Glycol (*freeze-point depressant*)approx. 6 g of 10% HCl Solution (*sufficient to give a final pH of 2.5 in Part A*)

777 g Water

Part B (Solid Additive):

90 g Sodium Percarbonate

15 15 g Citric Acid (*buffer, to adjust final pH*)

This formulation will produce 1 liter of foam solution. The pH of the final formulation can be adjusted to be between about 9.6 and 9.8 for optimal performance. The following mixing procedure can be used: Mix Part B into Part A. Use, preferably, within 8 hours. Alternatively, sodium bisulfate (a common pool conditioning chemical), or other acid, can be used in place of citric acid to adjust the pH. The performance of this second example of DF-200HF with Propylene Glycol (utilizing sodium percarbonate) against chemical agent simulants is shown in Table 13.

Table 13: Reaction rates from kinetic testing for the second example of DF-200HF with propylene glycol (utilizing sodium percarbonate).

Simulant	% Decontaminated		
	1 Minute	15 Minutes	60 Minutes
Mustard (HD)	80	ND	ND
VX	76	96	>99

In general, sodium percarbonate dissolves much more slowly than urea hydrogen peroxide after it has been added to Part A. However, to increase the dissolution velocity, sodium percarbonate can be milled to approximately a 100 mesh size for use in this configuration. The time
5 to dissolve the sodium percarbonate was decreased from approximately 30 minutes to about 2 minutes when milled sodium percarbonate was used.

DF-200 with Corrosion Inhibitor

Corrosion inhibitors can be added to DF-200 formulations to reduce their corrosivity. A
10 preferred corrosion inhibitor for use in DF-200 formulations is N,N-dimethyl ethanolamine. However, other corrosion inhibitors, such as triethanolamine, ethanolamine salts of C9, C10, and C12 diacid mixtures, dicyclohexyl amine nitrite, and N,N-dibenzylamine, can be used. The Corrosion inhibitors added to DF-200 formulations can serve multiple purposes:

- 15
1. a corrosion inhibitor,
 2. a pH buffer,
 3. a solvent to keep 1-dodecanol in solution, and
 4. a co-solvent to solubilize insoluble chemical agents, such as sarin or mustard.

20 An example of a 3-part kit configuration of DF-200HF with a corrosion inhibitor comprises (amounts illustrative):

WO 03/028429

PCT/US02/29886

-44-

DF-200HF Rapid Deployment with Corrosion Inhibitor (3-part kit)Part A (Liquid Foam Component):

20 g Variquat 80MC

10 g Adogen 477

5 4 g 1-Dodecanol

5 g Poly (Ethylene Glycol)

10 g N,N-dimethyl ethanolamine (*corrosion inhibitor*)

50 g Potassium Bicarbonate

approx. 18 g Potassium Hydroxide (*sufficient to give a final pH of 10.2 in Part A*)

10 936 g Water

Part B (Solid Oxidant Component):

97 g Urea Hydrogen Peroxide

Part C (Liquid Bleaching Activator):

20 g Propylene Glycol Diacetate

15

This formulation will produce 1 liter of foam solution. The pH of the final formulation can be adjusted to be between about 9.6 and 9.8 for optimal performance. The following mixing procedure can be used: Mix Part B into Part A. Then, after dissolution of the urea hydrogen peroxide, add Part C to Part A+B. Use, preferably, within 8 hours. The performance of DF-200HF with corrosion inhibitor is shown below against chemical agent simulants is given in Table 14.

20

Table 14: Reaction rates in kinetic testing for DF-200HF with a corrosion inhibitor.

Simulant	% Decontaminated		
	1 Minute	15 Minutes	60 Minutes
Mustard (HD)	7	41	79
VX	58	94	99

25

Tests against the anthrax spore simulant (*Bacillus globigii* spores) demonstrated 99.9999% (7-Log) kill after a 60 minute exposure to DF-200HF with a corrosion inhibitor. The addition of the

-45-

corrosion inhibitor has a detrimental effect on the performance of DF-200 against chemical agents, but has no measured effect on the performance of DF-200HF against biological agents. Similar results were obtained when an alternative corrosion inhibitor, 1% triethanolamine, was used.

5 DF-200 with Glycerol

In another embodiment of a DF-200 formulation, glycerol may be employed as a viscosity builder in place of Jaguar 8000, poly (ethylene oxide), or polyethylene glycol. Glycerol is a common ingredient in cosmetics, where it is used as a viscosity builder, as well as a solvent, humectant and emollient. Thus, the use of glycerol in DF-200 formulations can serve multiple purposes:

- 10 1. Viscosity builder,
2. a humectant (i.e., a substance which moisturizes the skin),
3. a solvent to keep 1-dodecanol in solution, and
4. a co-solvent to solubilize insoluble chemical agents, such as sarin or mustard.

- 15 An example of a 3-part kit configuration of DF-200HF with glycerol comprises (amounts illustrative):

DF-200HF Rapid Deployment with Glycerol (3-part kit)

Part A (Liquid Foam Component):

- | | |
|----|---|
| 20 | 20 g Variquat 80MC |
| | 10 g Adogen 477 |
| | 4 g 1-Dodecanol |
| | 40 g Glycerol (<i>viscosity builder</i>) |
| | 40 g Potassium Bicarbonate |
| 25 | approx. 17 g Potassium Hydroxide (<i>sufficient to give a final pH of 10.2 in Part A</i>) |
| | 906 g Water |

WO 03/028429

PCT/US02/29886

-46-

Part B (Solid Oxidant Component):

97 g Urea Hydrogen Peroxide

Part C (Liquid Bleaching Activator):

5 20 g Propylene Glycol Diacetate

This formulation will produce 1 liter of foam solution. The pH of the final formulation can be adjusted to be between about 9.6 and 9.8 for optimal performance. The following mixing procedure can be used: Mix Part B into Part A. Then after dissolution of the urea hydrogen peroxide, add Part C to
10 Part +/B. Use, preferably, within 8 hours. The performance of DF-200HF with glycerol against chemical agent simulants is given in Table 15.

Table 15: Reaction rates in kinetic testing for DF-200HF with glycerol.

15

Simulant	% Decontaminated		
	1 Minute	15 Minutes	60 Minutes
Mustard (HD)	63	96	ND
G Agents	ND	ND	ND
VX	76	99	ND

Tests against the anthrax spore simulant (*Bacillus globigii* spores) demonstrated 99.9999% (7-Log) kill after a 30 minute exposure to DF-200HF with glycerol.

20

This formulation can be used for direct application to humans because the glycerol will act as a humectant. This formulation could also be utilized, e.g., as a spray or shower, by removing foaming constituents (such as 1-dodecanol and Adogen 477), and by reducing the concentration of peroxide. However, a drawback to the use of glycerol is that it is solid at a fairly high temperature (below about 10°C). Therefore, it would preferably be used in controlled temperature conditions (i.e.,
25 warm temperature conditions).

WO 03/028429


PCT/US02/29886

-47-

Propylene glycol diacetate, a bleaching activator used in many of the previously described DF-200 configurations is not presently available in solid form. However, other bleaching activators are available in solid form.

5 DF-200 with Acetylcholine Chloride

Solid O-acetyl bleaching activators (e.g., acetylcholine chloride, which is often used in eyedrop solutions) can be used in DF-200 formulations in place of (liquid) propylene glycol diacetate. The chemical structure of this O-acetyl bleaching activator is shown below. As can be seen, the molecule contains an O-acetyl group that can activate peroxide, and it is a quaternary compound,
 10 which is very compatible with DF-200 formulations. Acetylcholine chloride is also soluble in water and is very hygroscopic.



An example of a 2-part kit configuration of DF-200HF using acetylcholine chloride comprises
 15 (amounts illustrative):

WO 03/028429

PCT/US02/29886

-48-

DF-200HF Rapid Deployment using Acetylcholine Chloride (2-part kit)Part A (Liquid Foam Component):

20 g Variquat 80MC

10 g Adogen 477

5 30 g Poly (Ethylene Glycol) (MW 8000)

8 g Diethylene Glycol Monobutyl Ether

5 g Isobutanol

4 g 1-Dodecanol

150 g Propylene Glycol

10 50 g Potassium Bicarbonate

approx. 17 g Potassium Hydroxide (*sufficient to give a final pH of 10.2 in Part A*)

803 g Water

Part B (Solid Additive):

97 g Urea Hydrogen Peroxide

15 25 g Acetylcholine Chloride (*solid bleaching activator*)

This formulation will produce approximately 1 liter of foam solution. The pH of the final formulation can be adjusted to be between about 9.6 and 9.8 for optimal performance. To use this formulation, mix Part B into Part A. Use, preferably, within 8 hours. The performance of DF-200HF using acetylcholine chloride against chemical agent simulants is shown in Table 16.

Table 16: Reaction rates from kinetic testing for the DF-200HF using acetylcholine chloride as an activator.

Simulant	% Decontaminated		
	1 Minute	15 Minutes	60 Minutes
Mustard (HD)	60	98	ND
VX	10	85	>99

Tests against the anthrax spore simulant (*Bacillus globigii* spores) demonstrated 99.9999% (7-Log) kill after a 30 minute exposure to DF-200HF using acetylcholine chloride.

Two other O-acetyl bleaching activators, monoacetin (glycerol monoacetate) and diacetin (glycerol diacetate), have also been tested for their effectiveness in DF-200 formulations. Both of these compounds have also proven to be extremely effective bleaching activators. These compounds are water-soluble liquids.

Experiments have also shown that the peroxide in DF-200 formulations is also effectively activated by a nitrile-containing compound, such as 4-cyanobenzoic acid (which is water-soluble), at a concentration of, for example, 2%, for the neutralization of both chemical agent and biological agent simulants.

DF-200 using Peracetic Acid

Tests were conducted using peracetic acid as the oxidant in DF-200, instead of hydrogen peroxide. The following formulation was used:

2% Variquat 80MC (*cationic surfactant*)
2% peracetic acid (*oxidant*)
5% potassium bicarbonate (*buffer and activator*)
91% water

The pH was adjusted to 9.8 with solid KOH and the formulation was tested against the simulants for mustard, VX, and anthrax spores. The performance of this formulation is shown in Table 17 against chemical agent simulants.

WO 03/028429

PCT/US02/29886

-50-

Table 17: Reaction rates in kinetic testing for DF-200 with 2% peracetic acid.

Simulant	% Decontaminated		
	1 Minute	15 Minutes	60 Minutes
Mustard (HD)	27	58	68
VX	68	76	95

5 Tests against the anthrax spore simulant (*Bacillus globigii* spores) demonstrated 99.9999% (7-Log) kill after a 30 minute exposure to DF-200 with 2% peracetic acid.

Tests were also conducted for DF-200 using a higher concentration of peracetic acid (3.5%) in the following formulation:

10

2% Variquat 80MC (*cationic surfactant*)

3.5% peracetic acid (*oxidant*)

5% potassium bicarbonate (*buffer and activator*)

89.5% water

15

The pH was adjusted to 9.8 with solid KOH and the formulation was tested against the simulants for mustard, VX, and anthrax spores. The performance of this formulation is shown in Table 18 against chemical agent simulants.

20

Table 18: Reaction rates in kinetic testing for DF-200 with 3.5% peracetic acid.

Simulant	% Decontaminated		
	1 Minute	15 Minutes	60 Minutes
Mustard (HD)	40	94	ND
VX	74	96	98

25

The results show that use of peracetic acid as an alternative oxidant is effective against chemical agent simulants, but is not as effective as DF-200 formulations using activated hydrogen

WO 03/028429

PCT/US02/29886

-51-

peroxide (i.e., the combination of hydrogen peroxide, bicarbonate, and propylene glycol diacetate) as the oxidant. However, the DF-200 formulations with 2-3.5% peracetic acid are very effective for spore kill. Nevertheless, use of this oxidant is not as attractive as hydrogen peroxide because peracetic acid is not presently available in a safe, convenient solid form, and the shelf life of the liquid form is rather short.

Tests were also conducted to determine the minimum constituents required for spore kill in a DF-200 formulation which utilizes peracetic acid as an oxidant. These results indicate that only three constituents, i.e., peracetic acid, bicarbonate and the cationic surfactant, are necessary to achieve high rates of spore kill.

Live Agent Tests Using DF-200HF

Live agent tests on three chemical agents (soman ("GD"), VX, and mustard ("HD")) and two biological agents (anthrax spores and *Yersinia pestis*) were conducted. The results of kinetic testing of DF-200HF (using a three-part configuration) on the chemical agents is shown in Table 19.

Table 19: Reaction rates in kinetic testing for DF-200HF against chemical agents.

Chemical Agent	% Destruction of Chemical Agent at Time Interval		
	1 minute	15 minutes	60 minutes
GD	99.98 $\pm$ 0.01	99.97 $\pm$ 0.01	99.98 $\pm$ 0.01
VX	91.20 $\pm$ 8.56	99.80 $\pm$ 0.08	99.88 $\pm$ 0.04
HD	78.13 $\pm$ 10.53	98.46 $\pm$ 1.43	99.84 $\pm$ 0.32

After exposure of GD to DF-200HF, methylphosphonic acid (MPA) and pinacolyl methylphosphonic acid (PMPA) were identified as byproducts. After exposure of VX to DF-200HF, ethyl methylphosphonic acid (EMPA) and MPA were identified as byproducts. This indicated that the destruction of the VX followed the more desirable path to the phosphonic acids, rather than to EA2192 (a toxic byproduct which can also be produced during VX degradation). Lastly, after exposure of HD to DF-200HF, the initial degradation products for HD comprised a mixture of the

WO 03/028429

PCT/US02/29886

-52-

sulfoxide and sulfone byproducts, followed later by nearly complete disappearance of each of these byproducts after 60 minutes.

Results of tests against anthrax spores is shown in Tables 20 and 21 and against *Yersinia* pestis (i.e., the plague bacterium) are shown in Table 22 (NG refers to 'no growth'). The detection limit for these tests were 10 CFU/ml. Note that the 'error bars' in the '% Reduction' column takes into account this detection limit.

Table 20: Kill rates for *B. anthracis* AMES-RID spores in a solution of DF-200HF.

<i>B. anthracis</i> AMES-RID	Average CFU/ml	Log Reduction	% Reduction
Control	1.21E+07	0	0.00
15 min contact	NG	7	100±.00004
30 min contact	NG	7	100±.00004
60 min contact	NG	7	100±.00004

Table 21: Kill rates for *B. anthracis* ANR-1 spores in a solution of DF-200HF.

<i>B. anthracis</i> ANR-1	Average CFU/ml	Log Reduction	% Reduction
Control	6.42E+07	0	0/00
15 min contact	NG	7	100±.00004
30 min contact	NG	7	100±.00004
60 min contact	NG	7	100±.00004

Table 22: Kill rates for *Y. pestis* cells in a solution of DF-200HF.

<i>Y. pestis</i> (ATCC 11953)	Average CFU/ml	Log Reduction	%Reduction
Control	1.33E+07	0	0.00
15 min contact	NG	7	100±.00004
30 min contact	NG	7	100±.00004
60 min contact	NG	7	100±.00004

The petri dishes used for cell growth on each of these tests were saved for 21 days following the tests to verify that DF-200HF had actually killed the spores, rather than just inhibiting their growth. No growth on any of the petri dishes was observed after the 21-day period.

5 Toxic Industrial Chemicals Tests Using DF-200HF

Several toxic industrial chemicals (TICs) have also been tested against DF-200HF. A summary of the TICs tested to date and the results of those tests is shown in Table 23. Note that the results for malathion, butyl isocyanate, sodium cyanide, and carbon disulfide were obtained by analyzing for the unreacted chemical in foam solution, while the results for phosgene was obtained
10 by analyzing for the chemical in the headspace above a foam solution. These results demonstrate very effective neutralization of these toxic industrial chemicals.

Table 23: Summary of Toxic Industrial Chemical (TIC) Neutralization Tests with DF-200HF.

15

TIC	% Decontaminated		
	1 minute	15 minutes	60 minutes
Malathion (liquid)	89	95	Below Detection
Hydrogen Cyanide (gas)	>99	>99	>99
Sodium Cyanide (solid)	93	98	>99
Butyl Isocyanate (liquid)	99	Below Detection	Below Detection
Carbon Disulfide (liquid)	>99	>99	Below Detection
Phosgene (gas)	98	>99	>99

20

The preceding examples can be repeated with similar success by substituting the generically or specifically described reactants and/or operating conditions of this invention for those used in the preceding examples.

Although the invention has been described in detail with particular reference to these preferred embodiments, other embodiments can achieve the same results. Variations and modifications of the present invention will be obvious to those skilled in the art and it is intended to

17 FEB 2005

WO 03/028429

PCT/US02/29886

-54-

cover in the appended claims all such modifications and equivalents. The entire disclosures of all references, applications, patents, and publications cited above are hereby incorporated by reference.

-55-

CLAIMS

What is claimed is:

- 5 1. A formulation for use in neutralization of at least one toxant, said formulation comprising:
- at least two solubilizing compounds, wherein at least one solubilizing compound is a cationic surfactant and at least one solubilizing compound is a cationic hydrotrope;
- at least one reactive compound, wherein said at least one reactive compound
- 10 is selected from the group consisting of hydrogen peroxide, urea hydrogen peroxide, hydroperoxycarbonate, peracetic acid, sodium perborate, sodium peroxyphosphate, sodium peroxysulfate, and sodium percarbonate; and
- at least one bleaching activator selected from the group consisting of O-acetyl, N-acetyl, and nitrile group bleaching activators;
- 15 wherein said at least two solubilizing compounds, said at least one reactive compound, and said at least one bleaching activator, when mixed with water and exposed to the at least one toxant, neutralizes the at least one toxant.
- 20 2. The formulation according to claim 1 wherein said cationic surfactant comprises a quaternary ammonium salt.
3. The formulation according to claim 2 wherein said quaternary ammonium salt is selected from the group consisting of cetyltrimethyl ammonium bromide, benzalkonium chloride, benzethonium chloride, cetylpyridinium chloride, alkyldimethylbenzylammonium salt, tetrabutyl
- 25 ammonium bromide, and a mixture of benzyl (C-12-C16) alkyldimethylammonium chlorides.
4. The formulation according to claim 1 further comprising a water-soluble polymer.

-56-

5. The formulation according to claim 4 wherein the water-soluble polymer is selected from the group consisting of polyvinyl alcohol, guar gum, (cationic or non-ionic) polydiallyl dimethyl ammonium chloride, polyacrylamide, poly (ethylene oxide), glycerol, polyethylene glycol 8000 (PEG 8000), and JAGUAR 8000™ (Guar Gum 2-hydroxypropyl ether), and combinations thereof.

5

6. The formulation according to claim 1 further comprising a fatty alcohol comprising from 8 to 20 carbon atoms per molecule.

7. The formulation according to claim 1 further comprising a solvent.

10

8. The formulation according to claim 7 wherein the solvent comprises a member of the group consisting of Di(propylene glycol) methyl ether and diethylene glycol monobutyl ether and a combination thereof.

15

9. The formulation according to claim 1 further comprising a carbonate salt.

10. The formulation according to claim 9 wherein the carbonate salt is selected from the group consisting of potassium bicarbonate, sodium bicarbonate, ammonium bicarbonate, ammonium hydrogen bicarbonate, lithium bicarbonate, ammonium carbonate, and potassium carbonate, and combinations thereof.

20

11. The formulation according to claim 1 wherein said at least one bleaching activator is water-soluble.

-57-

12. The formulation according to claim 11 wherein said at least one water-soluble bleaching activator is selected from the group consisting of acetylcholine chloride, monoacetin (glycerol monoacetate), diacetin (glycerol diacetate), 4-cyanobenzoic acid, ethylene glycol diacetate, propylene glycol monomethyl ether acetate, methyl acetate, dimethyl glutarate, diethylene glycol monoethyl ether acetate, glycerol diacetate (Diacetin), glycerol monoacetate, glycerol triacetate, and propylene glycol diacetate, and combinations thereof.

13. The formulation according to claim 1 wherein said at least one bleaching activator is water-insoluble.

14. The formulation according to claim 13 wherein said at least one water-insoluble peroxide activator is selected from the group consisting of tetraacetyl ethylenediamine (TAED), n-nonanoyloxybenzenesulfonate (NOBS), and N-acetyl glucosamine, and combinations thereof.

15. The formulation according to claim 1 wherein the formulation, when mixed with water, has a pH value between about 9.6 and about 9.8.

-58-

16. A formulation for use in neutralization of at least one toxant, said formulation comprising:

at least one cationic surfactant;

at least one reactive compound, wherein said at least one reactive compound

5 is selected from the group consisting of hydrogen peroxide, urea hydrogen peroxide, hydroperoxycarbonate, peracetic acid, sodium perborate, sodium peroxyphosphate, sodium peroxysilicate, and sodium percarbonate;

at least one bleaching activator selected from the group consisting of O-acetyl, N-acetyl, and nitrile group bleaching activators; and

10 at least one carbonate salt not one of the at least one reactive compounds; wherein said at least one surfactant, said at least one reactive compound, said at least one bleaching activator, and said at least one carbonate salt, when mixed with water and exposed to the at least one toxant, neutralizes the at least one toxant.

15 17. The formulation according to claim 16 wherein the formulation, when mixed with water, has a pH value between about 9.6 and about 9.8.

18. The formulation according to claim 16 wherein said cationic surfactant comprises a quaternary ammonium salt.

20

19. The formulation according to claim 18 wherein said quaternary ammonium salt comprises benzalkonium chloride.

20. The formulation according to claim 16 wherein said at least one carbonate salt is
25 selected from the group consisting potassium bicarbonate, sodium bicarbonate, ammonium bicarbonate, ammonium hydrogen bicarbonate, lithium bicarbonate, ammonium carbonate, and potassium carbonate, and combinations thereof.

21. The formulation according to claim 16 consisting substantially of said at least one cationic surfactant, said at least one reactive compound, said at least one bleaching activator, and said at least one carbonate salt.

5 22. The formulation according to claim 16 wherein said at least one bleaching activator is water-soluble.

23. The formulation according to claim 22 wherein said at least one water-soluble bleaching activator is selected from the group consisting of acetylcholine chloride, monoacetin
10 (glycerol monoacetate), diacetin (glycerol diacetate), 4-cyanobenzoic acid, ethylene glycol diacetate, propylene glycol monomethyl ether acetate, methyl acetate, dimethyl glutarate, diethylene glycol monoethyl ether acetate, glycerol diacetate (Diacetin), glycerol monoacetate, glycerol triacetate, and propylene glycol diacetate, and combinations thereof.

15 24. The formulation according to claim 1 packaged in a kit configuration.

25. The formulation according to claim 24 wherein said kit configuration comprises:
a premixed component comprising said at least two solubilizing agents;
a first component comprising said at least one bleaching activator; and
20 a second component comprising said at least one reactive compound.

26. The formulation according to claim 25 further comprising water and a base.

27. The formulation according to claim 25 wherein said premixed component additionally
25 comprises a water-soluble polymer.

28. The formulation according to claim 25 wherein said at least one bleaching activator is selected from the group consisting of propylene glycol diacetate, glycerol diacetate, and TAED, and combinations thereof.

-60-

29. The formulation according to claim 25 wherein said premixed component additionally comprises a fatty alcohol comprising from 8 to 20 carbon atoms per molecule.

5 30. The formulation according to claim 25 wherein said at least one reactive compound comprises urea hydrogen peroxide and wherein said second component comprising said at least one reactive compound additionally comprises sodium percarbonate.

31 The formulation according to claim 25 wherein said premixed component additionally
10 comprises a short-chained alcohol.

32. The formulation according to claim 24 wherein said kit configuration comprises:
a first premixed component comprising said at least two solubilizing agents
and water; and
15 a second premixed component comprising said at least one bleaching
activator and said at least one reactive compound, wherein said at least one bleaching activator is in
solid form.

33. The formulation according to claim 32 wherein said first premixed component
20 additionally comprises an acid.

34. The formulation according to claim 32 wherein said at least one bleaching activator
comprises acetylcholine chloride.

25 35. The formulation according to claim 32 wherein said at least one reactive compound
comprises urea hydrogen peroxide and wherein said at least one bleaching activator comprises
TAED.

WO 03/028429

PCT/US02/29886

-61-

36. The formulation according to claim 32 wherein said at least one bleaching activator is encapsulated to prevent premature reaction with said at least one reactive compound.

37. The formulation according to claim 24 wherein said kit configuration comprises:
5 a premixed component comprising said at least two solubilizing agents and
said at least one bleaching activator; and
a component comprising said at least one reactive compound.

38. The formulation according to claim 37 wherein said premixed component additionally
10 comprises water and an acid.

39. The formulation according to claim 37 wherein said component comprising said at least one reactive compound comprises sodium percarbonate and additionally comprises an acid.

15 40. The formulation according to claim 37 wherein said at least one reactive compound comprises urea hydrogen peroxide and wherein said component comprising said at least one reactive compound additionally comprises a mixture of potassium carbonate and potassium bicarbonate.

-62-

41. A formulation for use in neutralization of at least one toxant, said formulation comprising:

at least one cationic surfactant;

at least one reactive compound, wherein said at least one reactive

5 compound is selected from the group consisting of hydrogen peroxide, urea hydrogen peroxide, hydroperoxycarbonate, peracetic acid, sodium peroxyphosphate, sodium peroxydicarbonate, and sodium percarbonate;

at least one water-soluble bleaching activator selected from the group consisting of O-acetyl, N-acetyl, nitrile group bleaching activators; , acetylcholine chloride,

10 monoacetin (glycerol monoacetate), diacetin (glycerol diacetate), 4-cyanobenzoic acid, ethylene glycol diacetate, propylene glycol monomethyl ether acetate, methyl acetate, dimethyl glutarate, diethylene glycol monoethyl ether acetate, glycerol acetate (Diacetin), glycerol monoacetate, glycerol triacetate, and propylene glycol diacetate, and combinations thereof; and

wherein said at least one surfactant, said at least one reactive compound
15 and said at least one water-soluble bleaching activator, when mixed with water and exposed to the at least one toxant, neutralizes the at least one toxant.

42. Canceled

20 43. The formulation according to claim 41 wherein said cationic surfactant comprises a quaternary ammonium salt.

44. The formulation according to claim 43 wherein said quaternary ammonium salt comprises benzalkonium chloride.

25

45. The formulation according to claim 41 further comprising at least one carbonate salt.

-63-

46. The formulation according to claim 45 wherein said at least one carbonate salt is selected from the group consisting potassium bicarbonate, sodium bicarbonate, ammonium bicarbonate, ammonium hydrogen bicarbonate, lithium bicarbonate, ammonium carbonate, and potassium carbonate, and combinations thereof.

5

47. The formulation according to claim 41 wherein the formulation, when mixed with water, has a pH value between about 9.6 and about 9.8.

48. The formulation according to claim 41 consisting substantially of said at least one cationic surfactant, said at least one reactive compound, and said at least one water-soluble bleaching activator.

10

49. The formulation according to claim 1 consisting substantially of:

1-10% benzalkonium chloride;

1-8% propylene glycol diacetate;

1-16% hydrogen peroxide; and

2-8% potassium bicarbonate.

15

WO 03/028429

PCT/US02/29886

-64-

50. A formulation for use in neutralization of at least one toxant, said formulation comprising:

at least one solubilizing compound, selected from the group consisting of a cationic hydrotrope and a fatty alcohol comprising from 8 to 20 carbon atoms per molecule;

5 at least one reactive compound, wherein said at least one reactive compound is selected from the group consisting of hydrogen peroxide, urea hydrogen peroxide, hydroperoxycarbonate, peracetic acid, sodium perborate, sodium peroxyphosphate, sodium peroxysilicate, and sodium percarbonate; and

10 at least one bleaching activator selected from the group consisting of O-acetyl, N-acetyl, and nitrile group bleaching activators;

wherein said at least one solubilizing compound, said at least one reactive compound, and said at least one bleaching activator, when mixed with water and exposed to the at least one toxant, neutralizes the at least one toxant.