

Claims

1. A pharmaceutical composition comprising 1) an amorphous solid dispersion comprising pralsetinib or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable hydrophilic polymer; and 2) an effervescent couple.
- 5 2. A pharmaceutical composition comprising 1) an amorphous solid dispersion comprising pralsetinib or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable amphiphilic polymer; and 2) an effervescent couple.
3. The composition of claim 1 or 2, wherein the polymer is selected from the group consisting of hydroxypropyl methylcellulose (hypromellose), hydroxypropyl methylcellulose acetate succinate (HPMC-AS), vinylpyrrolidone-vinyl acetate copolymer (KOLLIDON
10 VA64 or KOLLIDON K30), dimethylaminoethyl methacrylate-copolymer (EUDRAGIT EPO), poly(ethylene) oxide (POLYOX), and polyvinyl caprolactam–polyvinyl acetate–polyethylene glycol graft copolymer (SOLUPLUS).
4. The composition of any one of claims 1-3 wherein the polymer is hydroxypropyl
15 methylcellulose.
5. The composition of any one of claims 1-4, wherein the polymer is hydroxypropylmethylcellulose E3 (HPMC-E3) or hydroxypropylmethylcellulose E5 (HPMC-E5).
6. The composition of any one of claims 1, 3, and 4, wherein the pralsetinib, or an
20 equivalent amount of a pharmaceutically acceptable salt thereof, and the hydrophilic polymer are in a weight percent ratio of about 1:1.
7. The composition of any one of claims 2-4, wherein the pralsetinib, or an equivalent amount of a pharmaceutically acceptable salt thereof, and the amphiphilic polymer are in a weight percent ratio of about 1:1.
- 25 8. The composition of any one of claims 1, 3, and 4, wherein the pralsetinib, or an equivalent amount of a pharmaceutically acceptable salt thereof, and the hydrophilic polymer are in a weight percent ratio of from about 1:5 to about 5:1.
9. The composition of any one of claims 2-4, wherein the pralsetinib, or an equivalent amount of a pharmaceutically acceptable salt thereof, and the amphiphilic polymer are in a
30 weight percent ratio of from about 1:5 to about 5:1.
10. The composition of any one of claims 1 to 9, wherein the composition comprises the amorphous solid dispersion from about 25% to about 65% by weight of the composition, based on the total weight of the composition.

11. The composition of any one of claims 1 to 10, wherein the composition comprises the amorphous solid dispersion from about 50% to about 60% by weight of the composition, based on the total weight of the composition.
12. The composition of any one of claims 1 to 11, wherein the amorphous solid dispersion
5 is prepared by hot melt extrusion, lyophilization, spray drying, solvent casting, or melt quenching.
13. The composition of any one of claims 1 to 12, wherein the effervescent couple comprises a water-soluble acid and a water-soluble base.
14. The composition of any one of claims 1 to 12, wherein the effervescent couple
10 comprises a water-soluble base.
15. The composition of claim 13, wherein the water-soluble acid is selected from the group consisting of citric acid, tartaric acid, fumaric acid, adipic acid, succinic acid, malonic acid, benzoic acid, oxalic acid, malic acid, and glutaric acid.
16. The composition of any one of claims 13-15, wherein the water-soluble base is selected
15 from the group consisting of sodium bicarbonate, sodium carbonate, potassium bicarbonate, potassium carbonate and magnesium carbonate.
17. The composition of claim 13, wherein the water-soluble acid is citric acid and the water-soluble base is sodium bicarbonate.
18. The composition of any one of claims 13 and 15-17, wherein the composition
20 comprises the water-soluble acid from about 1% to about 20% by weight of the composition, based on the total weight of the composition.
19. The composition of any one of claims 13-18, wherein the composition comprises the water-soluble base from about 1% to about 35% by weight of the composition, based on the total weight of the composition.
20. The composition of any one of claims 1 to 19, wherein the composition further
25 comprises a moisture scavenger.
21. The composition of claim 20, wherein the moisture scavenger is a starch.
22. The composition of claim 21, wherein the starch is pregelatinized starch.
23. The composition of any one of claims 20 to 22, wherein the composition comprises the
30 moisture scavenger from about 0.5% to about 30% by weight of the composition, based on the total weight of the composition.
24. The composition of any one of claims 1 to 23, wherein the composition further comprises a diluent.
25. The composition of claim 24, wherein the diluent is a cellulose.

26. The composition of claim 25, wherein the cellulose is microcrystalline cellulose.
27. The composition of any one of claims 24 to 26, wherein the composition comprises the diluent from about 5% to about 60% by weight of the composition, based on the total weight of the composition.
- 5 28. The composition of any one of claims 1 to 27, wherein the composition further comprises a lubricant.
29. The composition of claim 28, wherein the lubricant is magnesium stearate.
30. The composition of any one of claims 28 or 29, wherein the composition comprises the lubricant from about 0.1% to about 5% by weight of the composition, based on the total
- 10 weight of the composition.
31. The composition of any one of claims 1 to 30, wherein the composition is prepared in an oral dosage form.
32. The composition of claim 31, wherein the oral dosage form is a capsule.
33. The composition of claim 31, wherein the oral dosage form is a tablet.
- 15 34. The composition of any one of claims 1 to 33, wherein the composition is an immediate release composition.
35. The composition of any one of claims 1 to 34, wherein the composition comprises about 10 mg, about 20 mg, about 30 mg, about 40 mg, about 50 mg, about 60 mg, about 70 mg, about 80 mg, about 90 mg, about 100 mg, or about 200 mg of pralsetinib, or an equivalent
- 20 amount of a pharmaceutically acceptable salt thereof.
36. An oral dosage form comprising a) an amorphous solid dispersion comprising pralsetinib, or a pharmaceutically acceptable salt thereof and hydroxypropyl methylcellulose (HPMC); b) microcrystalline cellulose (MCC); c) pregelatinized starch; d) sodium bicarbonate; e) citric acid; and f) magnesium stearate.
- 25 37. The oral dosage form of claim 36, wherein the oral dosage form comprises the amorphous solid dispersion from about 25% to about 65% percent by weight of the oral dosage form, based on the total weight of the oral dosage form.
38. The oral dosage form of claim 36 or 37, wherein the oral dosage form comprises the microcrystalline cellulose (MCC) from about 5% to about 60% percent by weight of the oral
- 30 dosage form, based on the total weight of the oral dosage form.
39. The oral dosage form of any one of claims 36 to 38, wherein the oral dosage form comprises the pregelatinized starch from about 0.5% to about 30% by weight of the oral dosage form, based on the total weight of the oral dosage form.

40. The oral dosage form of any one of claims 36 to 39, wherein the oral dosage form comprises the sodium bicarbonate from about 1% to about 35% percent by weight of the oral dosage form, based on the total weight of the oral dosage form.
41. The oral dosage form of any one of claims 36 to 40, wherein the oral dosage form
5 comprises the citric acid from about 1% to about 20% by weight of the oral dosage form, based on the total weight of the oral dosage form.
42. The oral dosage form of any one of claims 36 to 41, wherein the oral dosage form comprises the magnesium stearate from about 0.1% to about 5% by weight of the oral dosage form, based on the total weight of the oral dosage form.
- 10 43. The oral dosage form of any one of claims 36 to 42 wherein the pralsetinib, or an equivalent amount of a pharmaceutically acceptable salt thereof, and the hydrophilic polymer are in a weight percent ratio of about 1:1.
44. The oral dosage form of any one of claims 36 to 43, wherein the amorphous solid dispersion is prepared by hot melt extrusion, lyophilization, spray drying, solvent casting, or
15 melt quenching.
45. The oral dosage form of any one of claims 36 to 44, wherein the oral dosage form is a capsule.
46. The oral dosage form of claim 32 or 45, wherein the size of the capsule is 0.
47. The oral dosage form of any one of claims 36-44, wherein the oral dosage form is a
20 tablet.
48. The oral dosage form of any one of claims 36 to 47, wherein the oral dosage form comprises about 10 mg, about 20 mg, about 30 mg, about 40 mg, about 50 mg, about 60 mg, about 70 mg, about 80 mg, about 90 mg, or about 100 mg, or about 200 mg of pralsetinib, or an equivalent amount of a pharmaceutically acceptable salt thereof.
- 25 49. The oral dosage form of any one of claims 36 to 48, wherein the oral dosage form is an immediate release oral dosage form.
50. An immediate release oral dosage form comprising:
- a) an amorphous solid dispersion comprising: pralsetinib, or a pharmaceutically acceptable salt thereof, and hydroxypropyl methylcellulose E3, wherein the
30 pralsetinib or an equivalent amount of pharmaceutically acceptable salt thereof and the hydroxypropyl methyl cellulose E3 are in about a 1:1 weight ratio;
- b) an effervescent couple comprising about 3 to about 13 w/w% citric acid and about 7 to about 30 w/w% sodium bicarbonate; wherein w/w% is based on the total weight of the oral dosage form;

c) a diluent; and optionally

d) moisture scavenger and/or a lubricant.

51. The oral dosage form of claim 50, wherein the amorphous solid dispersion comprises about 30 mg, about 50 mg, about 60 mg or about 100 mg of pralsetinib or an equivalent amount of a pharmaceutically acceptable salt thereof.
52. The composition or oral dosage form of any one of claims 1-51, wherein at least 80% of the pralsetinib is released in 45 minutes using USP <711> with a Type 2 Apparatus, media containing 900 mL 0.1 M HCl, and a paddle speed of 100 rpm.
53. The composition or oral dosage form of any one of claims 1-51, wherein the dosage form is a capsule, wherein the capsule disintegrates in about 7 to 15 minutes using USP <701>, with Basket Type A and a disc with a maintained temperature at $37^{\circ}\text{C} \pm 2^{\circ}\text{C}$.
54. The composition or oral dosage form of any one of claims 1-51, wherein the dosage form is a capsule, wherein at least 80% of the pralsetinib is released in about 120 minutes using USP II apparatus with a media containing 900 mL pH 6.8 sodium phosphate buffer with 0.5% CTAB and a paddle speed of $75 \text{ rpm} \pm 2 \text{ rpm}$.
55. A method for preparing the amorphous solid dispersion according to any one of claims 1 to 54, comprising: mixing the pralsetinib or the pharmaceutically acceptable salt thereof, with the hydrophilic polymer in about a 1:1 ratio; adding a solvent, and removing the solvent by heating.
56. A method of treating a RET-altered cancer comprising administering to a patient in need thereof a therapeutically effective amount of the composition according to any one of claims 1 to 35 or an oral dosage form of any one of claims 36 to 54.
57. A method of treating a patient with rearranged during transfection (RET)-positive locally advanced or metastatic non-small cell lung cancer (NSCLC) comprising administering to a patient in need thereof a therapeutically effective amount of the composition according to any one of claims 1 to 35 or an oral dosage form of any one of claims 36 to 54.
58. The method of claim 57, wherein the (RET)-positive locally advanced or metastatic non-small cell lung cancer (NSCLC) is detected by an FDA approved test.
59. A method of treating a patient with RET-mutation positive locally advanced or metastatic medullary thyroid cancer (MTC) comprising administering to a patient in need thereof a therapeutically effective amount of the composition according to any one of claims 1 to 35 or an oral dosage form of any one of claims 36 to 54.
60. A method of treating a patient with RET-fusion positive locally advanced or metastatic thyroid cancer who require systemic therapy and have no satisfactory alternative treatment

options comprising administering to a patient in need thereof a therapeutically effective amount of the composition according to any one of claims 1 to 35 or an oral dosage form of any one of claims 36 to 54.

61. Crystalline Form A of pralsetinib.

5 62. The crystalline Form A of claim 61, wherein the crystalline form is characterized by at least three, at least four, or at least five x-ray powder diffraction peaks at 2-theta angles selected from $5.0\pm0.2^\circ$, $9.7\pm0.2^\circ$, $12.7\pm0.2^\circ$, $13.6\pm0.2^\circ$, and $16.1\pm0.2^\circ$.

63. The crystalline Form A of claim 61, wherein the crystalline form is characterized by at least three, at least four, at least five, at least six, at least seven, at least eight, or at least nine
10 x-ray powder diffraction peaks at 2-theta angles selected from $5.0\pm0.2^\circ$, $6.8\pm0.2^\circ$, $9.7\pm0.2^\circ$, $12.7\pm0.2^\circ$, $13.6\pm0.2^\circ$, $16.1\pm0.2^\circ$, $19.2\pm0.2^\circ$, $19.5\pm0.2^\circ$, and $23.5\pm0.2^\circ$.

64. The crystalline Form A of claim 61, wherein the crystalline form is characterized by at least three, at least four, at least five, at least six, at least seven, at least eight, at least nine, or at least ten x-ray powder diffraction peaks at 2-theta angles selected from $5.0\pm0.2^\circ$, $6.8\pm0.2^\circ$,
15 $9.7\pm0.2^\circ$, $12.7\pm0.2^\circ$, $13.6\pm0.2^\circ$, $14.8\pm0.2^\circ$, $16.1\pm0.2^\circ$, $17.2\pm0.2^\circ$, $17.8\pm0.2^\circ$, $19.2\pm0.2^\circ$, $19.5\pm0.2^\circ$, $20.5\pm0.2^\circ$, $21.6\pm0.2^\circ$, $23.1\pm0.2^\circ$, $23.5\pm0.2^\circ$, $24.8\pm0.2^\circ$, $25.6\pm0.2^\circ$, $26.0\pm0.2^\circ$, $27.9\pm0.2^\circ$, and $29.4\pm0.2^\circ$.

65. The crystalline Form A of any one of claims 61 to 64, wherein the peaks have a relative intensity of at least 10%, of at least 15%, of at least 20%, or of at least 25%.

20 66. The crystalline Form A of claim 61, wherein the crystalline form has an XRPD pattern that is substantially the same XRPD pattern shown in FIG. 1A.

67. The crystalline Form A of claim 61, wherein the crystalline form has an XRPD pattern that substantially includes the peaks in Table 1A.

68. The crystalline Form A of any one of claims 61 to 67, wherein the crystalline form has
25 a DSC pattern that is substantially the same DSC pattern shown in FIG. 1B.

69. The crystalline Form A of any one of claims 61 to 68, wherein an endothermic event at about $205^\circ\text{C}\pm 2^\circ\text{C}$ is observed in the DSC thermogram.

70. The crystalline Form A of any one of claims 61 to 69, wherein the crystalline form has a TGA pattern that is substantially the same TGA pattern shown in FIG. 1B.

30 71. The crystalline Form A of any one of claims 61 to 70, wherein the crystalline form has a DVS pattern that is substantially the same DVS pattern shown in FIG. 1C.

72. The crystalline Form A of any one of claims 61 to 71, wherein the crystalline form is characterized by a reversible mass change of about 10% by DVS between 2-95% relative humidity.

73. A process for preparing the crystalline Form A of any one of claims 61 to 72, wherein the process comprises the step selected from the group consisting of (a) slurrying in alcohols, acetone, or ACN; (b) evaporative crystallization and cooling crystallization in IPA and 1-propanol; (c) recrystallization in acetone:water; (d) heating a sample of Form B to at least about 190 °C; or (e) heating a sample of pralsetinib Form C to at least about 190 °C under suitable conditions to yield Form A.
74. Crystalline Form B of pralsetinib.
75. The crystalline Form B of claim 74, wherein the crystalline form is characterized by at least three, at least four, or at least five x-ray powder diffraction peaks at 2-theta angles selected from $5.9 \pm 0.2^\circ$, $8.8 \pm 0.2^\circ$, $11.6 \pm 0.2^\circ$, $14.7 \pm 0.2^\circ$, and $19.5 \pm 0.2^\circ$.
76. The crystalline Form B of claim 74, wherein the crystalline form is characterized by at least three, at least four, at least five, at least six, at least seven, or at least eight x-ray powder diffraction peaks at 2-theta angles selected from $5.9 \pm 0.2^\circ$, $8.8 \pm 0.2^\circ$, $11.6 \pm 0.2^\circ$, $14.7 \pm 0.2^\circ$, $17.0 \pm 0.2^\circ$, $17.6 \pm 0.2^\circ$, $19.5 \pm 0.2^\circ$, and $22.2 \pm 0.2^\circ$.
77. The crystalline Form B of any one of claims 74 to 76, wherein the peaks have a relative intensity of at least 10%, of at least 15%, of at least 20%, or of at least 25%.
78. The crystalline Form B of claim 74, wherein the crystalline form has an XRPD pattern that is substantially the same XRPD pattern shown in FIG. 1A.
79. The crystalline Form B of claim 74, wherein the crystalline form has an XRPD pattern that substantially includes the peaks in Table 2A.
80. The crystalline Form B of any one of claims 74 to 79, wherein the crystalline form has a DSC pattern that is substantially the same DSC pattern shown in FIG. 2B.
81. The crystalline Form B of any one of claims 74 to 80, when crystalline Form B is characterized by DSC: an endotherm with onset at $149^\circ\text{C} \pm 2^\circ\text{C}$, an exotherm with onset at $162^\circ\text{C} \pm 2^\circ\text{C}$, and melting with onset $205^\circ\text{C} \pm 2^\circ\text{C}$.
82. The crystalline Form B of any one of claims 74 to 81, wherein the crystalline form has a TGA pattern that is substantially the same TGA pattern shown in FIG. 2B.
83. The crystalline Form B of any one of claims 74 to 82, wherein the crystalline form has a TGA pattern showing that the mass loss was 0.5%.
84. The crystalline Form B of any one of claims 74 to 83, wherein the crystalline form has a DVS pattern that is substantially the same DVS pattern shown in FIG. 2C.
85. The crystalline Form B of any one of claims 74 to 84, wherein the crystalline form showed a total mass change of 1.4 wt.% between 2% and 95% relative humidity.

86. A process for preparing the crystalline Form B of any one of claims 74 to 85, wherein the process comprises the step of heating a sample of Form C to about 150 °C.
87. Crystalline Form C of pralsetinib.
88. The crystalline Form C of claim 87, wherein the crystalline form is characterized by at least three, at least four, or at least five x-ray powder diffraction peaks at 2-theta angles selected from $5.8 \pm 0.2^\circ$, $8.7 \pm 0.2^\circ$, $11.0 \pm 0.2^\circ$, $13.6 \pm 0.2^\circ$, and $20.2 \pm 0.2^\circ$.
89. The crystalline Form C of claim 87, wherein the crystalline form is characterized by at least three, at least four, at least five, at least six, at least seven, at least eight, or at least nine x-ray powder diffraction peaks at 2-theta angles selected from $5.8 \pm 0.2^\circ$, $8.7 \pm 0.2^\circ$, $11.0 \pm 0.2^\circ$, $11.6 \pm 0.2^\circ$, $13.6 \pm 0.2^\circ$, $14.5 \pm 0.2^\circ$, $20.2 \pm 0.2^\circ$, $22.2 \pm 0.2^\circ$, and $23.2 \pm 0.2^\circ$.
90. The crystalline Form C of claim 87, wherein the crystalline form is characterized by at least three, at least four, at least five, at least six, at least seven, at least eight, at least nine, or at least ten x-ray powder diffraction peaks at 2-theta angles selected from $5.8 \pm 0.2^\circ$, $8.7 \pm 0.2^\circ$, $11.0 \pm 0.2^\circ$, $11.6 \pm 0.2^\circ$, $12.0 \pm 0.2^\circ$, $13.6 \pm 0.2^\circ$, $14.5 \pm 0.2^\circ$, $17.1 \pm 0.2^\circ$, $18.2 \pm 0.2^\circ$, $19.5 \pm 0.2^\circ$, $20.2 \pm 0.2^\circ$, $20.6 \pm 0.2^\circ$, $21.3 \pm 0.2^\circ$, $22.2 \pm 0.2^\circ$, $22.6 \pm 0.2^\circ$, $23.2 \pm 0.2^\circ$, $24.2 \pm 0.2^\circ$, $24.5 \pm 0.2^\circ$, $26.0 \pm 0.2^\circ$, $26.8 \pm 0.2^\circ$, and $28.1 \pm 0.2^\circ$.
91. The crystalline Form C of claim 87, wherein the peaks have a relative intensity of at least 10%, of at least 15%, of at least 20%, or of at least 25%.
92. The crystalline Form C of claim 87, wherein the crystalline form has an XRPD pattern that is substantially the same XRPD pattern shown in FIG. 3A.
93. The crystalline Form C of claim 87, wherein the crystalline form has an XRPD pattern that substantially includes the peaks in Table 3A.
94. The crystalline Form C of any one of claims 87 to 93, wherein the crystalline form has a DSC pattern that is substantially the same DSC pattern shown in FIG. 1B.
95. The crystalline Form C of any one of claims 87 to 94, wherein onsets occurring at $122^\circ \pm 2^\circ\text{C}$, $127^\circ \pm 2^\circ\text{C}$, and $206^\circ \pm 2^\circ\text{C}$ are seen in the DSC thermogram.
96. The crystalline Form C of any one of claims 87 to 95, wherein the crystalline form has a TGA pattern that is substantially the same TGA pattern shown in FIG. 3B.
97. The crystalline Form C of any one of claims 87 to 96, wherein a mass loss of about 3 wt.% is observed in the Form C TGA thermogram.
98. The crystalline Form C of any one of claims 87 to 97, wherein the crystalline form has a DVS pattern that is substantially the same DVS pattern shown in FIG. 3C.

99. The crystalline Form C of any one of claims 87 to 98, wherein the DVS of the crystalline form showed a total mass change of 1.4 wt.% between 2% and 95% relative humidity.
100. A process for preparing the crystalline Form C of any one of claims 87 to 99, wherein
5 the process comprises the step selected from the group consisting of a) recrystallization in various water containing solvent systems (acetone:water, MeOH:water, IPA:water, DMAc:water, THF:water); b) conversion from Form A during competitive slurry experiments in methanol:water at high ratios of water to methanol and lower temperatures.
101. Crystalline pralsetinib HCl salt Form I.
102. The crystalline Form I of claim 101, wherein the crystalline form is characterized by at
10 least three, at least four, or at least five x-ray powder diffraction peaks at 2-theta angles selected from $.0\pm0.2^\circ$, $6.1\pm0.2^\circ$, $9.1\pm0.2^\circ$, $9.9\pm0.2^\circ$, and $14.7\pm0.2^\circ$.
103. The crystalline Form I of claim 101, wherein the crystalline form is characterized by at
15 at least three, at least four, at least five, at least six, at least seven, at least eight, at least nine, or at least ten x-ray powder diffraction peaks at 2-theta angles selected from $5.0^\circ\pm0.2$, $6.1^\circ\pm0.2$, $9.1^\circ\pm0.2$, $9.9^\circ\pm0.2$, $13.8^\circ\pm0.2$, $14.7^\circ\pm0.2$, $15.3^\circ\pm0.2$, $17.2^\circ\pm0.2$, $18.1^\circ\pm0.2$, $19.6^\circ\pm0.2$, $20.3^\circ\pm0.2$, $20.7^\circ\pm0.2$, $21.8^\circ\pm0.2$, $24.2^\circ\pm0.2$, $25.6^\circ\pm0.2$, and $26.3^\circ\pm0.2$.
104. The crystalline Form I of any one of claims 101 to 103, wherein the peaks have a relative intensity of at least 10%, of at least 15%, of at least 20%, or of at least 25%.
- 20 105. The crystalline Form I of claim 101, wherein the crystalline form has an XRPD pattern that is substantially the same XRPD pattern shown in FIG. 4A.
106. The crystalline Form I of claim 101, wherein the crystalline form has an XRPD pattern that substantially includes the peaks in Table 4A.
107. The crystalline Form I of any one of claims 101 to 106, wherein the crystalline form has
25 a DSC pattern that is substantially the same DSC pattern shown in FIG. 4B.
108. A process for preparing the crystalline Form I of any one of claims 101 to 107, wherein the process comprises the step of isolating the solid from the slurry of the HCl salt in EtOH or IPA:water (9:1 Vol).
109. Crystalline pralsetinib HCl salt Form II.
- 30 110. The crystalline Form II of claim 109, wherein the crystalline form is characterized by at least three, at least four, or at least five x-ray powder diffraction peaks at 2-theta angles selected from $6.1\pm0.2^\circ$, $8.9\pm0.2^\circ$, $9.5\pm0.2^\circ$, $15.0\pm0.2^\circ$, $16.6\pm0.2^\circ$.
111. The crystalline Form II of claim 109, wherein the crystalline form is characterized by at least three, at least four, at least five, at least six, at least seven, at least eight, or at least nine

- x-ray powder diffraction peaks at 2-theta angles selected from $6.1 \pm 0.2^\circ$, $8.9 \pm 0.2^\circ$, $9.5 \pm 0.2^\circ$, $15.0 \pm 0.2^\circ$, $16.6 \pm 0.2^\circ$, $17.2 \pm 0.2^\circ$, $17.9 \pm 0.2^\circ$, $18.4 \pm 0.2^\circ$, $19.8 \pm 0.2^\circ$, $25.8 \pm 0.2^\circ$, and $26.8 \pm 0.2^\circ$.
112. The crystalline Form II of any one of claims 109 to 111, wherein the peaks have a relative intensity of at least 10%, of at least 15%, of at least 20%, or of at least 25%.
- 5 113. The crystalline Form II of claim 109, wherein the crystalline form has an XRPD pattern that is substantially the same XRPD pattern shown in FIG. 5A.
114. The crystalline Form II of claim 109, wherein the crystalline form has an XRPD pattern that substantially includes the peaks in Table 5A.
115. The crystalline Form II of any one of claims 109 to 114, wherein the crystalline form
- 10 has a DSC pattern that is substantially the same DSC pattern shown in FIG. 5B.
116. The crystalline Form II of any one of claims 109 to 115, wherein the crystalline form was observed to have a broad endotherm with an onset of $88.7^\circ\text{C} \pm 2^\circ\text{C}$ and a melt which had an onset of $244.2^\circ\text{C} \pm 2^\circ\text{C}$ in the DSC thermogram.
117. The crystalline Form II of any one of claims 109 to 116, wherein the crystalline form
- 15 has a TGA pattern that is substantially the same TGA pattern shown in FIG. 5B.
118. The crystalline Form II of any one of claims 109 to 117, wherein an initial mass loss of 3.4 wt. % associated with a broad endotherm with an onset of $94.4^\circ\text{C} \pm 2^\circ\text{C}$ and a second mass loss event of 6.7 wt. % was observed from the end of the first broad endotherm to the end of the melt which had an onset of $244.2^\circ\text{C} \pm 2^\circ\text{C}$ was observed in the TGA thermogram.
- 20 119. A process for preparing the crystalline Form II of any one of claims 109 to 118, wherein the process comprises the step of isolating the solid from EtOAc and IPA:water (9:1 vol).
120. Crystalline pralsetinib HCl salt Form III.
121. The crystalline Form III of claim 120, wherein the crystalline form is characterized by at least three, at least four, or at least five x-ray powder diffraction peaks at 2-theta angles
- 25 selected from $6.4 \pm 0.2^\circ$, $8.5 \pm 0.2^\circ$, $8.9 \pm 0.2^\circ$, $9.6 \pm 0.2^\circ$, and $17.3 \pm 0.2^\circ$.
122. The crystalline Form III of claim 120, wherein the crystalline form is characterized by at least three, at least four, at least five, at least six, at least seven, at least eight, or at least nine x-ray powder diffraction peaks at 2-theta angles selected from $6.4 \pm 0.2^\circ$, $8.5 \pm 0.2^\circ$, $8.9 \pm 0.2^\circ$, $9.6 \pm 0.2^\circ$, $11.5 \pm 0.2^\circ$, $16.7 \pm 0.2^\circ$, $17.3 \pm 0.2^\circ$, and $19.2 \pm 0.2^\circ$.
- 30 123. The crystalline Form III of claim 120, wherein the crystalline form is characterized by at least three, at least four, at least five, at least six, at least seven, at least eight, at least nine, or at least ten x-ray powder diffraction peaks at 2-theta angles selected from $6.0 \pm 0.2^\circ$, $6.4 \pm 0.2^\circ$, $8.5 \pm 0.2^\circ$, $8.9 \pm 0.2^\circ$, $9.6 \pm 0.2^\circ$, $11.5 \pm 0.2^\circ$, $12.7 \pm 0.2^\circ$, $15.9 \pm 0.2^\circ$, $16.7 \pm 0.2^\circ$, $17.3 \pm 0.2^\circ$, $19.2 \pm 0.2^\circ$, $21.0 \pm 0.2^\circ$, and $26.9 \pm 0.2^\circ$.

124. The crystalline Form III of any one of claims 120 to 123, wherein the peaks have a relative intensity of at least 10%, of at least 15%, of at least 20%, or of at least 25%.
125. The crystalline Form III of claim 120, wherein the crystalline form has an XRPD pattern that is substantially the same XRPD pattern shown in FIG. 6A.
- 5 126. The crystalline Form III of claim 120, wherein the crystalline form has an XRPD pattern that substantially includes the peaks in Table 6A.
127. The crystalline Form III of any one of claims 120 to 126, wherein the crystalline form has a DSC pattern that is substantially the same DSC pattern shown in FIG. 6B.
128. The crystalline Form III of any one of claims 120 to 127, wherein the crystalline form
10 had observed DSC onsets of $86.8\text{ }^{\circ}\text{C} \pm 2^{\circ}\text{C}$, $224.1\text{ }^{\circ}\text{C} \pm 2^{\circ}\text{C}$, and $241.7\text{ }^{\circ}\text{C} \pm 2^{\circ}\text{C}$.
129. The crystalline Form III of any one of claims 120 to 128, wherein the crystalline form has a TGA pattern that is substantially the same TGA pattern shown in FIG. 6B.
130. The crystalline Form III of any one of claims 120 to 129, wherein an initial mass loss of 3.4 wt. % and a second mass loss event of 2 wt. % was observed in the TGA thermogram.
- 15 131. A process for preparing the crystalline Form III of any one of Claims 120 to 130, wherein the process comprises the step of drying the isolated Pralsetinib HCl salt Form II.
132. A pharmaceutically acceptable hydrochloride salt of Compound (I):

