

DESCRIPTION
PROPHYLACTIC OR THERAPEUTIC AGENT FOR ATTENTION
DEFICIT/HYPERACTIVITY DISORDER

Technical Field

5 [0001]

The present invention relates to a prophylactic or therapeutic agent for attention deficit/hyperactivity disorder.

[0002]

(Background of the Invention)

10 Attention deficit/hyperactivity disorder (ADHD) is said to be one of the developmental disorders characterized by symptoms such as hyperactivity, inattention and impulsivity. For example, non-patent document 1 describes them in detail. The disease is often found before and after entering
15 elementary school where social rules such as staying still and the like increase. It is generally considered that the disease is of genetic etiology, and includes cases with no suitable diagnosis but with similar symptoms. Although a therapeutic method has not been established at present, improvement of
20 daily life is sought by prescription of medicament that enhances attention. For example, while a central nervous stimulant is used for increasing the arousal level to prevent symptoms, it is a symptomatic therapy and is not a curative treatment.

25 Therefore, a therapeutic drug effective for ADHD (particularly, medicament effective for hyperactivity/impulsivity disorder) is awaited in the medical practice.

non-patent document 1: Barkley, R. (2006). "Attention-
30 Deficit Hyperactivity Disorder, Third Edition: A Handbook for Diagnosis and Treatment"

Disclosure of the Invention

Problems to be Solved by the Invention

[0003]

35 An object of the present invention is to provide a novel

prophylactic or therapeutic agent for attention deficit/hyperactivity disorder.


Means of Solving the Problems

[0004]

5 The present inventors have conducted intensive studies and found that a compound represented by the following formula (I) or a salt thereof, particularly, (S)-N-[2-(1,6,7,8-tetrahydro-2H-indeno[5,4-b]furan-8-yl)ethyl]propionamide (general name: ramelteon; hereinafter sometimes to be
10 abbreviated as compound A) is effective for the prophylaxis or treatment of attention deficit/hyperactivity disorder, which resulted in the completion of the present invention.

Accordingly, the present invention relates to
[1] a prophylactic or therapeutic agent for attention
15 deficit/hyperactivity disorder, comprising a compound represented by the formula (I)

[0005]



[0006]

20 wherein R^1 is a hydrocarbon group optionally having substituent(s), an amino group optionally having substituent(s) or a heterocyclic group optionally having substituent(s),
 R^2 is a hydrogen atom or a hydrocarbon group optionally having
25 substituent(s),
 R^3 is a hydrogen atom, a hydrocarbon group optionally having substituent(s) or a heterocyclic group optionally having substituent(s),

X is CHR^4 , NR^4 , CO, O or S wherein R^4 is a hydrogen atom, a hydrocarbon group optionally having substituent(s) or a hydroxyl group,

Y is C, CH or N,

5 [0007]

[0008]

is a single bond or a double bond,

ring A is a 5- to 7-membered heterocycle containing an oxygen
10 atom and optionally having substituent(s),

ring B is a benzene ring optionally having substituent(s), and
m is an integer of 1 to 4, (hereinafter to be also referred to
as "compound (I)"), or a salt thereof as an active ingredient;

[2] the agent of the above-mentioned [1], wherein compound (I)
15 or a salt thereof is (S)-N-[2-(1,6,7,8-tetrahydro-2H-
indeno[5,4-b]furan-8-yl)ethyl]propionamide;

[3] a prophylactic or therapeutic agent for attention
deficit/hyperactivity disorder, comprising, as an active
ingredient, compound (I) or a salt thereof, and one or more
20 kinds of medicaments selected from other prophylactic or
therapeutic agents for attention deficit/hyperactivity
disorder in combination;

[4] the agent of the above-mentioned [3], wherein compound (I)
or a salt thereof is (S)-N-[2-(1,6,7,8-tetrahydro-2H-
25 indeno[5,4-b]furan-8-yl)ethyl]propionamide;

[5] a method for the prophylaxis or treatment of attention
deficit/hyperactivity disorder, comprising administering an
effective amount of compound (I) or a salt thereof to a mammal
(including human: hereinafter the same);

30 [6] the method of the above-mentioned [5], wherein compound
(I) or a salt thereof is (S)-N-[2-(1,6,7,8-tetrahydro-2H-
indeno[5,4-b]furan-8-yl)ethyl]propionamide;

[7] a method for the prophylaxis or treatment of attention
deficit/hyperactivity disorder, comprising administering an

effective amount of compound (I) or a salt thereof, and one or more kinds of medicaments selected from other prophylactic or therapeutic agents for attention deficit/hyperactivity disorder in combination to a mammal;

5 [8] the method of the above-mentioned [7], wherein compound (I) or a salt thereof is (S)-N-[2-(1,6,7,8-tetrahydro-2H-indeno[5,4-b]furan-8-yl)ethyl]propionamide;

[9] compound (I) or a salt thereof for use as a prophylactic or therapeutic agent for attention deficit/hyperactivity

10 disorder;

[10] (S)-N-[2-(1,6,7,8-tetrahydro-2H-indeno[5,4-b]furan-8-yl)ethyl]propionamide for use as a prophylactic or therapeutic agent for attention deficit/hyperactivity disorder;

[11] compound (I) or a salt thereof for use as a prophylactic
15 or therapeutic agent for attention deficit/hyperactivity disorder, in combination with one or more kinds of medicaments selected from other prophylactic or therapeutic agents for attention deficit/hyperactivity disorder;

[12] (S)-N-[2-(1,6,7,8-tetrahydro-2H-indeno[5,4-b]furan-8-
20 yl)ethyl]propionamide for use as a prophylactic or therapeutic agent for attention deficit/hyperactivity disorder, in combination with one or more kinds of medicaments selected from other prophylactic or therapeutic agents for attention deficit/hyperactivity disorder;

25 [13] use of compound (I) or a salt thereof for the production of a prophylactic or therapeutic agent for attention deficit/hyperactivity disorder;

[14] the use of the above-mentioned [13], wherein compound (I) or a salt thereof is (S)-N-[2-(1,6,7,8-tetrahydro-2H-
30 indeno[5,4-b]furan-8-yl)ethyl]propionamide;

[15] use of compound (I) or a salt thereof for the production of a prophylactic or therapeutic agent for attention deficit/hyperactivity disorder to be combined with one or more kinds of medicaments selected from other prophylactic or
35 therapeutic agents for attention deficit/hyperactivity

disorder;

[16] the use of the above-mentioned [15], wherein compound (I) or a salt thereof is (S)-N-[2-(1,6,7,8-tetrahydro-2H-indeno[5,4-b]furan-8-yl)ethyl]propionamide;

5 and the like.

Effect of the Invention

[0009]

The present invention provides a novel prophylactic or therapeutic agent for attention deficit/hyperactivity disorder,
10 and the like.

Brief Description of the Drawings

[0010]

Fig. 1 is a schematic presentation showing lamps on the wall and the position of the lever of the test apparatus,
15 wherein A is a house lamp, B_L and B_R are cue lamps, C_L and C_R are retractable levers, and D is a reward delivery opening.

Fig. 2 is a graph showing transition of total number of reaction during trials (comparison between SHR-SP rat (control group) and WKY rat).

20 Fig. 3 is a graph showing transition of the number of correct answers (comparison between SHR-SP rat (control group) and WKY rat).

Fig. 4 is a graph showing transition of correct answer latency (comparison between SHR-SP rat (control group) and WKY
25 rat).

Fig. 5 is a graph showing transition of the total number of reaction during trials (comparison between SHR-SP rat (control group) and SHR-SP rat (positive drug group)).

Fig. 6 is a graph showing transition of the number of
30 correct answers (comparison between SHR-SP rat (control group) and SHR-SP rat (positive drug group)).

Fig. 7 is a graph showing transition of correct answer latency (comparison between SHR-SP rat (control group) and SHR-SP rat (positive drug group)).

35 [0011]

(Detailed Description of the Invention)

Compound (I) and a salt thereof are described in detail in the following.

In the above-mentioned formula (I), R^1 is a hydrocarbon group optionally having substituent(s), an amino group optionally having substituent(s) or a heterocyclic group optionally having substituent(s),

R^2 is a hydrogen atom or a hydrocarbon group optionally having substituent(s),

R^3 is a hydrogen atom, a hydrocarbon group optionally having substituent(s) or a heterocyclic group optionally having substituent(s),

X is CHR^4 , NR^4 , CO, O or S wherein R^4 is a hydrogen atom, a hydrocarbon group optionally having substituent(s) or a hydroxyl group,

Y is C, CH or N,

[0012]

[0013]

is a single bond or a double bond,
ring A is a 5- to 7-membered heterocycle containing an oxygen atom and optionally having substituent(s),
ring B is a benzene ring optionally having substituent(s), and
m is an integer of 1 to 4.

Preferably, when X is CH_2 , then Y is C or CH.

[0014]

In the present specification, examples of the "hydrocarbon group" of the "hydrocarbon group optionally having substituent(s)" include an aliphatic hydrocarbon group, a monocyclic saturated hydrocarbon group, an aromatic hydrocarbon group and the like, with preference given to those having a carbon number of 1 to 16. Specifically, for example, an alkyl group, an alkenyl group, an alkynyl group, a cycloalkyl group, an aryl group and the like are used.

Preferable examples of the "alkyl group" include a lower alkyl group and the like and, for example, a C₁₋₆ alkyl group such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl and tert-butyl, pentyl, hexyl etc., and the like are
5 widely used.

Preferable examples of the "alkenyl group" include a lower alkenyl group and the like and, for example, a C₂₋₆ alkenyl group such as vinyl, 1-propenyl, allyl, isopropenyl, butenyl and isobutenyl etc., and the like are widely used.

10 Preferable examples of the "alkynyl group" include a lower alkynyl group and the like and, for example, a C₂₋₆ alkynyl group such as ethynyl, propargyl, 1-propynyl etc., and the like are widely used.

Preferable examples of the "cycloalkyl group" include a
15 lower cycloalkyl group and the like and, for example, a C₃₋₆ cycloalkyl group such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl etc., and the like are widely used.

Preferable examples of the "aryl group" include a C₆₋₁₄ aryl group such as phenyl, 1-naphthyl, 2-naphthyl, biphenyl, 2-anthryl etc., and the like and, for example, a phenyl group
20 and the like are widely used.

[0015]

As the substituent that the "hydrocarbon group" of the "hydrocarbon group optionally having substituent(s)"
25 optionally has, a halogen atom (e.g., fluorine, chlorine, bromine, iodine etc.), a nitro group, a cyano group, a hydroxyl group, an optionally halogenated lower alkyl group (e.g., an optionally halogenated C₁₋₆ alkyl group such as methyl, chloromethyl, difluoromethyl, trichloromethyl, trifluoromethyl,
30 ethyl, 2-bromoethyl, 2,2,2-trifluoroethyl, pentafluoroethyl, propyl, 3,3,3-trifluoropropyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, 4,4,4-trifluorobutyl, pentyl, isopentyl, neopentyl, 5,5,5-trifluoropentyl, hexyl, 6,6,6-trifluorohexyl etc.), a lower alkoxy group (e.g., a C₁₋₆ alkoxy group such as
35 methoxy, ethoxy, propoxy, isopropoxy, butoxy, isobutoxy,

5 pentyloxy, hexyloxy etc., and the like), an amino group, a mono-lower alkylamino group (e.g., a mono-C₁₋₆ alkylamino group such as methylamino, ethylamino etc., and the like), a di-lower alkylamino group (e.g., a di-C₁₋₆ alkylamino group such as
10 dimethylamino, diethylamino etc., and the like), a carboxyl group, a lower alkylcarbonyl group (e.g., a C₁₋₆ alkyl-carbonyl group such as acetyl, propionyl etc., and the like), a lower alkoxy carbonyl group (e.g., a C₁₋₆ alkoxy-carbonyl group such as methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl,
15 butoxycarbonyl etc., and the like), a carbamoyl group, a mono-lower alkylcarbamoyl group (e.g., a mono-C₁₋₆ alkyl-carbamoyl group such as methylcarbamoyl, ethylcarbamoyl etc., and the like), a di-lower alkylcarbamoyl group (e.g., a di-C₁₋₆ alkyl-carbamoyl group such as dimethylcarbamoyl, diethylcarbamoyl
20 etc., and the like), an arylcarbamoyl group (e.g., a C₆₋₁₀ aryl-carbamoyl group such as phenylcarbamoyl, naphthylcarbamoyl etc.), an aryl group (e.g., a C₆₋₁₀ aryl group such as phenyl, naphthyl etc.), an aryloxy group (e.g., a C₆₋₁₀ aryloxy group such as phenyloxy, naphthyloxy etc.), an optionally
25 halogenated lower alkylcarbonylamino group (e.g., an optionally halogenated C₁₋₆ alkyl-carbonylamino group such as acetylamino, trifluoroacetylamino etc., and the like), an oxo group and the like can be used.

The "hydrocarbon group" of the "hydrocarbon group
25 optionally having substituent(s)" may have 1 to 5, preferably 1 to 3, of the aforementioned substituents at substitutable position(s) of the hydrocarbon group. When the number of the substituents is two or more, the respective substituents may be the same or different.

30 [0016]

In the present specification, examples of the "heterocyclic group" of the "heterocyclic group optionally having substituent(s)" include a 5- to 14-membered (preferably 5- to 10-membered) (monocyclic to tricyclic, preferably
35 monocyclic or bicyclic) heterocyclic group containing, besides

carbon atom, one or two kinds of 1 to 4 (preferably 1 to 3) hetero atoms selected from a nitrogen atom, an oxygen atom and a sulfur atom and the like. For example, a 5-membered ring group containing, besides carbon atom, 1 to 4 hetero atoms
5 selected from an oxygen atom, a sulfur atom and a nitrogen atom, such as 2- or 3-thienyl, 2- or 3-furyl, 1-, 2- or 3-pyrrolyl, 1-, 2- or 3-pyrrolidinyl, 2-, 4- or 5-oxazolyl, 3-, 4- or 5-isoxazolyl, 2-, 4- or 5-thiazolyl, 3-, 4- or 5-isothiazolyl, 3-, 4- or 5-pyrazolyl, 2-, 3- or 4-pyrazolidinyl,
10 2-, 4- or 5-imidazolyl, 1,2,3-triazolyl, 1,2,4-triazolyl, 1H- or 2H-tetrazolyl and the like; a 6-membered ring group containing, besides carbon atom, 1 to 4 hetero atoms selected from an oxygen atom, a sulfur atom and a nitrogen atom, such as 2-, 3- or 4-pyridyl, N-oxide-2-, 3- or 4-pyridyl, 2-, 4- or
15 5-pyrimidinyl, N-oxide-2-, 4- or 5-pyrimidinyl, thiomorpholinyl, morpholinyl, piperidino, 2-, 3- or 4-piperidyl, thiopyranyl, 1,4-oxazinyl, 1,4-thiazinyl, 1,3-thiazinyl, piperazinyl, triazinyl, 3- or 4-pyridazinyl, pyrazinyl, N-oxide-3- or 4-pyridazinyl and the like; a
20 bicyclic or tricyclic fused ring group containing, besides carbon atom, 1 to 4 hetero atoms selected from an oxygen atom, a sulfur atom and a nitrogen atom (preferably, a group formed by condensation of the aforementioned 5- or 6-membered ring and one or two 5- or 6-membered ring groups optionally
25 containing, besides carbon atom, 1 to 4 hetero atoms selected from an oxygen atom, a sulfur atom and a nitrogen atom), such as indolyl, benzofuryl, benzothiazolyl, benzoxazolyl, benzimidazolyl, quinolyl, isoquinolyl, phthalazinyl, quinazolinyl, quinoxalinyl, indolizinyl, quinolizinyl, 1,8-
30 naphthyridinyl, dibenzofuranyl, carbazolyl, acrydiny, phenanthridinyl, chromanyl, phenothiazinyl, phenoxazinyl and the like; and the like can be used. Of these, a 5- to 7-membered (preferably 5- or 6-membered) heterocyclic group containing, besides carbon atom, 1 to 3 hetero atoms selected
35 from an oxygen atom, a sulfur atom and a nitrogen atom is

preferable.

[0017]

As the substituent that the "heterocyclic group" of the "heterocyclic group optionally having substituent(s)"
5 optionally has, a halogen atom (e.g., fluorine, chlorine, bromine, iodine etc.), a lower alkyl group (e.g., a C₁₋₆ alkyl group such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, hexyl etc., and the like), a cycloalkyl group (e.g., a C₃₋₆ cycloalkyl group such as
10 cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl etc., and the like), a lower alkynyl group (e.g., a C₂₋₆ alkynyl group such as ethynyl, 1-propynyl, propargyl etc., and the like), a lower alkenyl group (e.g., a C₂₋₆ alkenyl group such as vinyl, allyl, isopropenyl, butenyl, isobutenyl etc., and the like), an
15 aralkyl group (e.g., a C₇₋₁₁ aralkyl group such as benzyl, α -methylbenzyl, phenethyl etc., and the like), an aryl group (e.g., C₆₋₁₀ aryl group such as phenyl, naphthyl etc., and the like, preferably a phenyl group), a lower alkoxy group (e.g., a C₁₋₆ alkoxy group such as methoxy, ethoxy, propoxy, isopropoxy, butoxy, isobutoxy, sec-butoxy, tert-butoxy etc., and the like),
20 an aryloxy group (e.g., C₆₋₁₀ aryloxy group such as phenoxy etc., and the like), a lower alkanoyl group (e.g., formyl; a C₁₋₆ alkyl-carbonyl group such as acetyl, propionyl, butyryl, isobutyryl etc., and the like), arylcarbonyl (e.g., a C₆₋₁₀
25 aryl-carbonyl group such as a benzoyl group, a naphthoyl group etc., and the like), a lower alkanoyloxy group (e.g., formyloxy; a C₁₋₆ alkyl-carbonyloxy group such as acetyloxy, propionyloxy, butyryloxy, isobutyryloxy etc., and the like), an arylcarbonyloxy group (e.g., a C₆₋₁₀ aryl-carbonyloxy group
30 such as benzoyloxy, naphthoyloxy etc., and the like), a carboxyl group, a lower alkoxy-carbonyl group (e.g., a C₁₋₆ alkoxy-carbonyl group such as methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, isopropoxycarbonyl, butoxycarbonyl, isobutoxycarbonyl, tert-butoxycarbonyl etc., and the like),
35 aralkyloxycarbonyl (e.g., a C₇₋₁₁ aralkyloxycarbonyl group such

as benzyloxycarbonyl etc., and the like), a carbamoyl group, a mono-, di- or tri-halogeno-lower alkyl group (e.g., a mono-, di- or tri-halogeno-C₁₋₄ alkyl group such as chloromethyl, dichloromethyl, trifluoromethyl, 2,2,2-trifluoroethyl etc., and the like), an oxo group, an amidino group, an imino group, an amino group, a mono-lower alkylamino group (e.g., a mono-C₁₋₄ alkylamino group such as methylamino, ethylamino, propylamino, isopropylamino, butylamino etc., and the like), a di-lower alkylamino group (e.g., a di-C₁₋₄ alkylamino group such as dimethylamino, diethylamino, dipropylamino, diisopropylamino, dibutylamino, methylethylamino etc., and the like), a 3- to 6-membered cyclic amino group optionally containing, besides carbon atom and one nitrogen atom, 1 to 3 hetero atoms selected from an oxygen atom, a sulfur atom and a nitrogen atom (e.g., a 3- to 6-membered cyclic amino group such as aziridinyl, azetidiny, pyrrolidinyl, pyrrolinyl, pyrrolyl, imidazolyl, pyrazolyl, imidazolidinyl, piperidyl, morpholinyl, dihydropyridyl, pyridyl, N-methylpiperazinyl, N-ethylpiperazinyl etc., and the like), an alkylenedioxy group (e.g., a C₁₋₃ alkylenedioxy group such as methylenedioxy, ethylenedioxy etc., and the like), a hydroxyl group, a nitro group, a cyano group, a mercapto group, a sulfo group, a sulfinio group, a phosphono group, a sulfamoyl group, a monoalkylsulfamoyl group (e.g., a mono-C₁₋₆ alkylsulfamoyl group such as N-methylsulfamoyl, N-ethylsulfamoyl, N-propylsulfamoyl, N-isopropylsulfamoyl, N-butylsulfamoyl etc., and the like), a dialkylsulfamoyl group (e.g., a di-C₁₋₆ alkylsulfamoyl group such as N,N-dimethylsulfamoyl, N,N-diethylsulfamoyl, N,N-dipropylsulfamoyl, N,N-dibutylsulfamoyl etc., and the like), an alkylthio group (e.g., a C₁₋₆ alkylthio group such as methylthio, ethylthio, propylthio, isopropylthio, butylthio, sec-butylthio, tert-butylthio etc., and the like), an arylthio group (e.g., a C₆₋₁₀ arylthio group such as phenylthio, naphthylthio etc., and the like), a lower alkylsulfinyl group (e.g., a C₁₋₆ alkylsulfinyl group such as methylsulfinyl,

ethylsulfinyl, propylsulfinyl, butylsulfinyl etc., and the like), an arylsulfinyl group (e.g., a C₆₋₁₀ arylsulfinyl group such as phenylsulfinyl, naphthylsulfinyl etc., and the like), a lower alkylsulfonyl group (e.g., a C₁₋₆ alkylsulfonyl group
5 such as methylsulfonyl, ethylsulfonyl, propylsulfonyl, butylsulfonyl etc., and the like), an arylsulfonyl group (e.g., a C₆₋₁₀ arylsulfonyl group such as phenylsulfonyl, naphthylsulfonyl etc., and the like) and the like can be used.

The "heterocyclic group" of the "heterocyclic group
10 optionally having substituent(s)" may have 1 to 5, preferably 1 to 3, substituents from the aforementioned substituents at substitutable position(s) of the heterocyclic group. When the number of the substituents is two or more, the respective substituents may be the same or different.

15 [0018]

In the present specification, examples of the "amino group optionally having substituent(s)" include an amino group optionally having, as substituents, 1 or 2 substituents such as the aforementioned "hydrocarbon group optionally having
20 substituent(s)" etc., and the like. Preferable examples of the substituent that the "amino group" optionally has include a C₁₋₆ alkyl group optionally having substituent(s), a C₆₋₁₀ aryl group optionally having substituent(s) and the like. As the substituents that the "C₁₋₆ alkyl group" and the "C₆₋₁₀ aryl
25 group" optionally have, those similar to the substituents that the aforementioned "hydrocarbon group" optionally has can be used.

[0019]

In the present specification, the "lower alkyl group" of
30 the "lower alkyl group optionally having substituent(s)" is, for example, a C₁₋₆ alkyl group such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl and tert-butyl etc., and the like, and may have, as substituents, for example, 1 to 3 substituents that the aforementioned "hydrocarbon group"
35 optionally has, and the like.

In the present specification, the "lower alkoxy group" of the "lower alkoxy group optionally having substituent(s)" is, for example, a C₁₋₆ alkoxy group such as methoxy, ethoxy, propoxy, isopropoxy, butoxy, isobutoxy, sec-butoxy, tert-butoxy etc., and the like, and may have, as substituents, for example, 1 to 3 substituents that the aforementioned "hydrocarbon group" optionally has, and the like.

[0020]

In the present specification, the "benzene ring optionally having substituent(s)" is a benzene ring optionally having 1 or 2 substituents selected from, for example, a halogen atom (e.g., fluorine, chlorine, bromine, iodine etc.), a hydrocarbon group optionally having substituent(s), an amino group optionally having substituent(s), an amide group (e.g., a C₁₋₃ acylamino group such as formamide, acetamide etc., and the like), a hydroxyl group, a lower alkoxy group optionally having substituent(s), a lower alkylenedioxy group (e.g., a C₁₋₃ alkylenedioxy group such as methylenedioxy, ethylenedioxy etc., and the like), and the like at substitutable position(s).

As these "hydrocarbon group optionally having substituent(s)", "amino group optionally having substituent(s)" and "lower alkoxy group optionally having substituent(s)", those similar to the groups exemplified above can be used. When the number of the substituents that the "hydrocarbon group", "amino group" and "lower alkoxy group" have is two or more, the respective substituents may be the same or different.

As the "benzene ring optionally having substituent(s)", a benzene ring optionally substituted by 1 or 2 substituents selected from a halogen atom (e.g., fluorine, chlorine etc.), a C₁₋₆ alkyl group (e.g., methyl, ethyl etc.), a hydroxyl group and a mono-C₁₋₆ alkylamino group, and the like are preferable.

[0021]

In the aforementioned formula (I), R¹ is a hydrocarbon group optionally having substituent(s), an amino group

optionally having substituent(s) or a heterocyclic group optionally having substituent(s).

Preferable examples of the "hydrocarbon group" of the "hydrocarbon group optionally having substituent(s)" for R¹ include an alkyl group (e.g., a C₁₋₆ alkyl group such as methyl, ethyl, propyl, isopropyl etc., and the like), an alkenyl group (e.g., a C₂₋₆ alkenyl group such as vinyl etc., and the like), an alkynyl group (e.g., a C₂₋₆ alkynyl group such as ethynyl etc., and the like), a cycloalkyl group (e.g., a C₃₋₆ cycloalkyl group such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl etc., and the like) and an aryl group (e.g., a C₆₋₁₄ aryl group such as phenyl etc., and the like) and the like. Particularly, an alkyl group (e.g., a C₁₋₆ alkyl group such as methyl etc., and the like), a cycloalkyl group (e.g., a C₃₋₆ cyclopropyl group such as cyclopropyl etc., and the like) and the like are widely used. The "alkyl group", "alkenyl group", "alkynyl group", "cycloalkyl group" and "aryl group" may have, for example, 1 to 5, preferably 1 to 3, substituents that the aforementioned "hydrocarbon group" optionally has (preferably, a halogen atom such as fluorine etc., and the like) and the like.

[0022]

Preferable examples of the substituent of the "amino group optionally having substituent(s)" for R¹ include 1 or 2 from a lower alkyl group optionally having substituent(s), an aryl group optionally having substituent(s) and the like. Particularly, one from a lower alkyl group optionally having substituent(s) and the like is used. As the "lower alkyl group", for example, a C₁₋₆ alkyl group such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl etc., and the like are used. The "lower alkyl group" may have 1 to 3 substituents from, for example, those optionally possessed by the aforementioned "hydrocarbon group", and the like. As the "aryl group", for example, a C₆₋₁₀ aryl group such as a phenyl group etc., and the like are used. The "aryl group" may have 1

to 5, preferably 1 to 3, substituents from, for example, those optionally possessed by the aforementioned "hydrocarbon group" (preferably, a halogen atom such as fluorine, chlorine etc., a C₁₋₆ alkoxy group such as methoxy, ethoxy etc., and the like).

5 As the "amino group optionally having substituent(s)", a phenylamino group substituted by 1 to 3 lower alkoxy groups (e.g., a C₁₋₄ alkoxy group such as methoxy etc., and the like), a monoalkylamino group substituted by a lower alkyl group (e.g., a C₁₋₄ alkyl group such as methyl, ethyl, propyl, butyl, 10 tert-butyl etc., and the like) and the like are widely used.
[0023]

Preferable examples of the "heterocyclic group" of the "heterocyclic group optionally having substituent(s)" for R¹ include a 5- or 6-membered heterocyclic group containing, 15 besides carbon atom, 1 to 3 hetero atoms selected from a nitrogen atom, an oxygen atom and a sulfur atom, and the like. Specific examples include 1-, 2- or 3-pyrrolidinyl, 2- or 4-imidazoliny, 2-, 3- or 4-pyrazolidinyl, piperidino, 2-, 3- or 4-piperidyl, 1- or 2-piperazinyl, morpholinyl, 2- or 3-thienyl, 20 2-, 3- or 4-pyridyl, 2-furyl or 3-furyl, pyrazinyl, 2-pyrimidinyl, 3-pyrrolyl, 3-pyridazinyl, 3-isothiazolyl, 3-isoxazolyl and the like. Particularly preferably, a 6-membered nitrogen-containing heterocyclic group (e.g., pyridyl etc.) and the like are used.

25 Preferable examples of the substituent of the "heterocyclic group optionally having substituent(s)" for R¹ include a halogen atom (e.g., chlorine, fluorine etc.), a C₁₋₆ alkyl group (e.g., methyl, ethyl etc.), a C₁₋₆ alkoxy group (e.g., methoxy, ethoxy etc.), an aralkyloxycarbonyl group (e.g., C₇₋₁₂ aralkyloxy-carbonyl such as benzyloxycarbonyl etc., 30 and the like), and the like.

[0024]

For example, R¹ is preferably (i) a lower alkyl group optionally having substituent(s), (ii) a lower cycloalkyl 35 group optionally having substituent(s), (iii) a lower alkenyl

group optionally having substituent(s), (iv) an aryl group optionally having substituent(s), (v) a mono- or di-lower alkylamino group optionally having substituent(s), (vi) an arylamino group optionally having substituent(s), (vii) a 5-
5 or 6-membered nitrogen-containing heterocyclic group optionally having substituent(s) or the like.

The aforementioned "lower alkyl group" is preferably, for example, a C₁₋₆ alkyl group such as methyl, ethyl, propyl, isopropyl, butyl, pentyl, hexyl etc., and the like. The "lower
10 cycloalkyl group" is preferably, for example, a C₃₋₆ cycloalkyl group such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl etc., and the like. The "lower alkenyl group" is preferably, for example, a C₂₋₆ alkenyl group such as vinyl, 1-propenyl, butenyl etc., and the like. The "aryl group" is preferably,
15 for example, C₆₋₁₀ aryl group such as phenyl, 1-naphthyl, 2-naphthyl etc., and the like. The "lower alkylamino group" is preferably, for example, a mono- or di-C₁₋₆ alkylamino group such as methylamino, ethylamino, propylamino, isopropylamino, butylamino, tert-butylamino, dimethylamino, diethylamino,
20 methylethylamino etc., and the like. The "arylamino group" is preferably a C₆₋₁₀ arylamino group such as phenylamino etc., and the like. The "5- or 6-membered nitrogen-containing heterocyclic group" is preferably, for example, a 5- or 6-membered nitrogen-containing heterocyclic group such as 2-, 3-
25 or 4-pyridyl etc., and the like. As the substituents that these groups optionally have, 1 to 5 substituents that the aforementioned "hydrocarbon group" may have and the like are used.

In the present specification,
30 [0025]

[0026]

is a single bond or a double bond.

[0027]

More preferable examples of R^1 include (i) a C_{1-6} alkyl group optionally substituted by 1 to 4 each of halogen, a hydroxyl group and a C_{1-6} alkoxy group, (ii) a C_{3-6} cycloalkyl group, (iii) a C_{2-6} alkenyl group, (iv) a C_{6-10} aryl group optionally substituted by 1 to 4 each of C_{1-6} alkoxy, nitro, halogeno C_{1-6} alkyl-carbonylamino and a halogen atom, (v) a mono- or di- C_{1-6} alkylamino group, (vi) a C_{6-10} arylamino group optionally substituted by 1 to 3 C_{1-6} alkoxy groups or (vii) a 6-membered nitrogen-containing heterocyclic group optionally substituted by 1 or 2 C_{7-11} aralkyloxycarbonyl groups and the like. Particularly, an optionally halogenated C_{1-6} alkyl group (e.g., methyl, chloromethyl, difluoromethyl, trichloromethyl, trifluoromethyl, ethyl, 2-bromoethyl, 2,2,2-trifluoroethyl, pentafluoroethyl, propyl, 3,3,3-trifluoropropyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, 4,4,4-trifluorobutyl, pentyl, isopentyl, neopentyl, 5,5,5-trifluoropentyl, hexyl, 6,6,6-trifluorohexyl etc.), a C_{3-6} cycloalkyl group (e.g., cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl etc.), a mono- C_{1-6} alkylamino group (e.g., methylamino, ethylamino, propylamino, isopropylamino, butylamino, tert-butylamino etc.) and the like are widely used. Among these, an optionally halogenated C_{1-6} alkyl group or a mono- C_{1-6} alkylamino group, particularly an optionally halogenated C_{1-6} alkyl group, among others a C_{1-3} alkyl group (e.g., methyl, ethyl, propyl etc.), are preferable.

[0028]

In the aforementioned formula (I), R^2 is a hydrogen atom or a hydrocarbon group optionally having substituent(s).

As R^2 , a hydrogen atom or a lower (C_{1-6}) alkyl group optionally having substituent(s) is preferably used, a hydrogen atom or a lower (C_{1-6}) alkyl group is more preferably used, and particularly, a hydrogen atom is widely used.

[0029]

In the aforementioned formula (I), R^3 is a hydrogen atom, a hydrocarbon group optionally having substituent(s) or a

heterocyclic group optionally having substituent(s).

Preferable examples of the "hydrocarbon group" of the "hydrocarbon group optionally having substituent(s)" for R³ include an alkyl group (e.g., a C₁₋₆ alkyl group such as methyl, ethyl, propyl, isopropyl etc., and the like), an alkenyl group (e.g., a C₂₋₆ alkenyl group such as vinyl etc., and the like), an alkynyl group (e.g., a C₂₋₆ alkynyl group such as ethynyl etc., and the like), a cycloalkyl group (e.g., a C₃₋₆ cycloalkyl group such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl etc., and the like) and an aryl group (e.g., a C₆₋₁₄ aryl group such as phenyl etc., and the like) and the like, particularly an alkyl group (e.g., a C₁₋₆ alkyl group such as methyl etc., and the like), an aryl group (e.g., a C₆₋₁₄ aryl group such as phenyl etc., and the like) and the like are widely used. The "alkyl group", "alkenyl group", "alkynyl group", "cycloalkyl group" and "aryl group" may have 1 to 5, preferably 1 to 3, for example, substituents optionally possessed by the aforementioned "hydrocarbon group" (preferably, a halogen atom such as fluorine etc., and the like) and the like.

[0030]

Preferable examples of the "heterocyclic group" of the "heterocyclic group optionally having substituent(s)" for R³ include a 5- or 6-membered heterocyclic group containing, besides carbon atom, 1 to 3 hetero atoms selected from a nitrogen atom, an oxygen atom and a sulfur atom, and the like. Specific examples include 1-, 2- or 3-pyrrolidinyl, 2- or 4-imidazolyl, 2-, 3- or 4-pyrazolidinyl, piperidino, 2-, 3- or 4-piperidyl, 1- or 2-piperazinyl, morpholinyl, 2- or 3-thienyl, 2-, 3- or 4-pyridyl, 2- or 3-furyl, pyrazinyl, 2-pyrimidinyl, 3-pyrrolyl, 3-pyridazinyl, 3-isothiazolyl, 3-isoxazolyl and the like. Particularly preferably, a 6-membered nitrogen-containing heterocyclic group (e.g., pyridyl etc.) and the like are used.

[0031]

Preferable examples of the substituent of the

"heterocyclic group optionally having substituent(s)" for R^3 include a halogen atom (e.g., chlorine, fluorine etc.), a C_{1-6} alkyl group (e.g., methyl, ethyl etc.), a C_{1-6} alkoxy group (e.g., methoxy, ethoxy etc.), an aralkyloxycarbonyl group (e.g., C_{7-12} aralkyloxy-carbonyl such as benzyloxycarbonyl etc., and the like), an amino group, a mono- C_{1-6} alkylamino group (e.g., methylamino, ethylamino etc.), a di- C_{1-6} alkylamino group (e.g., dimethylamino, diethylamino etc.) and the like.

[0032]

For example, R^3 is preferably (i) a hydrogen atom, (ii) a lower alkyl group optionally having substituent(s), (iii) an aryl group optionally having substituent(s), (iv) a 5- or 6-membered heterocyclic group optionally having substituent(s) or the like. Furthermore, for example, (i) a hydrogen atom, (ii) a lower alkyl group, (iii) a C_{6-10} aryl group optionally having substituent(s), (iv) a 6-membered nitrogen-containing heterocyclic group optionally having substituent(s) and the like are preferable. Examples of the substituent include a halogen atom, a C_{1-6} alkyl group, a C_{1-6} alkoxy group, an amino group, a mono- C_{1-6} alkylamino group, a di- C_{1-6} alkylamino group and the like. More preferably, R^3 is a hydrogen atom, a phenyl group, or a 2-, 3- or 4-pyridyl group. Particularly preferred is a hydrogen atom.

[0033]

In the aforementioned formula (I), X is CHR^4 , NR^4 , CO, O or S wherein R^4 is a hydrogen atom, a hydroxyl group or a hydrocarbon group optionally having substituent(s).

As R^4 , a hydrogen atom, a hydroxyl group or a lower (C_{1-6}) alkyl group optionally having substituent(s) is preferable, and a hydrogen atom is widely used.

X is preferably CHR^4 wherein R^4 is as defined above, CO, O or S. Alternatively, X is preferably CHR^4 or NR^4 wherein R^4 is as defined above.


[0034]

In the aforementioned formula (I), ring A is a 5- to 7-

membered heterocycle containing an oxygen atom and optionally having substituent(s).

Examples of the "5- to 7-membered heterocycle containing an oxygen atom" include a 5- to 7-membered (preferably 5- or 6-membered) heterocycle containing, besides carbon atom and oxygen atom, 1 to 3 (preferably 1 or 2) of one or two kinds selected from a nitrogen atom, an oxygen atom and a sulfur atom and the like. As the ring, a ring represented by the formula

10 [0035]



[0036]


wherein E is (i) CH_2CH_2 , (ii) $\text{CH}=\text{CH}$, (iii) CH_2O , (iv) OCH_2 , (v) $\text{CH}_2\text{S}(\text{O})_{q'}$, wherein q' is an integer of 0 to 2, (vi) $\text{S}(\text{O})_{q'}\text{CH}_2$,
 15 wherein q' is as defined above, (vii) CH_2NH , (viii) NHCH_2 , (ix) $\text{N}=\text{N}$, (x) $\text{CH}=\text{N}$, (xi) $\text{N}=\text{CH}$ or (xii) CONH , and n' is an integer of 0 to 2, is preferable.

E is preferably (i) CH_2CH_2 , (ii) $\text{CH}=\text{CH}$, (iii) CH_2O , (iv) OCH_2 , (v) CH_2NH , (vi) NHCH_2 , (vii) $\text{N}=\text{N}$, (viii) $\text{CH}=\text{N}$ or (ix) $\text{N}=\text{CH}$,
 20 particularly preferably (i) CH_2CH_2 or (ii) $\text{CH}=\text{CH}$.

Specifically, for example, 5-membered heterocycle containing oxygen atom(s) such as 2,3-dihydrofuran, furan, 1,3-dioxole, oxazoline, isoxazole, 1,2,3-oxadiazole, oxazole etc., 6-membered heterocycle containing oxygen atom(s) such as
 25 2H-3,4-dihydropyran, 2H-pyran, 2,3-dehydro-1,4-dioxane, 2,3-dehydromorpholine etc., and the like are preferable.

More preferred is a ring represented by the formula

[0037]



[0038]

wherein n is an integer of 0 to 2, and

[0039]

5 [0040]

is a single bond or a double bond.

Specifically, for example, 2,3-dihydrofuran, furan, 2H-3,4-dihdropyran and 2H-pyran are widely used.

[0041]

10 As the substituent of ring A, a halogen atom (e.g., fluorine, chlorine, bromine, iodine etc.), a lower alkyl group optionally having substituent(s), a cycloalkyl group optionally having substituent(s), a lower alkynyl group optionally having substituent(s), a lower alkenyl group
15 optionally having substituent(s), an aryl group optionally having substituent(s), a lower alkoxy group (e.g., a C₁₋₆ alkoxy group such as methoxy, ethoxy, propoxy, isopropoxy, butoxy, isobutoxy, sec-butoxy, tert-butoxy etc., and the like), an aryloxy group (e.g., a C₆₋₁₀ aryloxy group such as phenoxy etc.,
20 and the like), a lower alkanoyl group (e.g., formyl; a C₁₋₆ alkyl-carbonyl group such as acetyl, propionyl, butyryl, isobutyryl etc., and the like), an arylcarbonyl group (e.g., a C₆₋₁₀ aryl-carbonyl group such as a benzoyl group, a naphthoyl group etc., and the like), a lower alkanoyloxy group (e.g.,
25 formyloxy; a C₁₋₆ alkyl-carbonyloxy group such as acetyloxy, propionyloxy, butyryloxy, isobutyryloxy etc., and the like), an arylcarbonyloxy group (e.g., a C₆₋₁₀ aryl-carbonyloxy group such as benzoyloxy, naphthoyloxy etc., and the like), a carboxyl group, a lower alkoxycarbonyl group (e.g., a C₁₋₆
30 alkoxy-carbonyl group such as methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, isopropoxycarbonyl, butoxycarbonyl, isobutoxycarbonyl, tert-butoxycarbonyl etc., and the like), aralkyloxy-carbonyl (e.g., a C₇₋₁₁ aralkyloxy-carbonyl group such as benzyloxy-carbonyl etc., and the like), a carbamoyl group, a

mono-, di- or tri-halogeno-lower alkyl group (e.g., a mono-, di- or tri-halogeno-C₁₋₄ alkyl group such as chloromethyl, dichloromethyl, trifluoromethyl, 2,2,2-trifluoroethyl etc., and the like), an oxo group, an amidino group, an imino group, 5 an amino group, a mono-lower alkylamino group (e.g., a mono-C₁₋₄ alkylamino group such as methylamino, ethylamino, propylamino, isopropylamino, butylamino etc., and the like), a di-lower alkylamino group (e.g., a di-C₁₋₄ alkylamino group such as dimethylamino, diethylamino, dipropylamino, diisopropylamino, 10 dibutylamino, methylethylamino etc., and the like), a 3- to 6-membered cyclic amino group containing, besides carbon atom and one nitrogen atom, 1 to 3 hetero atoms selected from an oxygen atom, a sulfur atom and a nitrogen atom (e.g., a 3- to 6-membered cyclic amino group such as aziridinyl, azetidiny, 15 pyrrolidinyl, pyrrolinyl, pyrrolyl, imidazolyl, pyrazolyl, imidazolidinyl, piperidyl, morpholinyl, dihydropyridyl, pyridyl, N-methylpiperazinyl, N-ethylpiperazinyl etc., and the like), an alkylenedioxy group (e.g., a C₁₋₃ alkylenedioxy group such as methylenedioxy, ethylenedioxy etc., and the like), a 20 hydroxyl group, a nitro group, a cyano group, a mercapto group, a sulfo group, a sulfinio group, a phosphono group, a sulfamoyl group, a monoalkylsulfamoyl group (e.g., a mono-C₁₋₆ alkylsulfamoyl group such as N-methylsulfamoyl, N-ethylsulfamoyl, N-propylsulfamoyl, N-isopropylsulfamoyl, N- 25 butylsulfamoyl etc., and the like), a dialkylsulfamoyl group (e.g., a di-C₁₋₆ alkylsulfamoyl group such as N,N-dimethylsulfamoyl, N,N-diethylsulfamoyl, N,N-dipropylsulfamoyl, N,N-dibutylsulfamoyl etc., and the like), an alkylthio group (e.g., a C₁₋₆ alkylthio group such as methylthio, ethylthio, 30 propylthio, isopropylthio, butylthio, sec-butylthio, tert-butylthio etc., and the like), an arylthio group (e.g., a C₆₋₁₀ arylthio group such as phenylthio, naphthylthio etc., and the like), a lower alkylsulfinyl group (e.g., a C₁₋₆ alkylsulfinyl group such as methylsulfinyl, ethylsulfinyl, propylsulfinyl, 35 butylsulfinyl etc., and the like), an arylsulfinyl group (e.g.,

a C₆₋₁₀ arylsulfinyl group such as phenylsulfinyl, naphthylsulfinyl etc., and the like), a lower alkylsulfonyl group (e.g., a C₁₋₆ alkylsulfonyl group such as methylsulfonyl, ethylsulfonyl, propylsulfonyl, butylsulfonyl etc., and the like), an arylsulfonyl group (e.g., a C₆₋₁₀ arylsulfonyl group such as phenylsulfonyl, naphthylsulfonyl etc., and the like) and the like are used.

The "lower alkyl group", "lower alkenyl group", "lower alkynyl group", "lower cycloalkyl group" and "aryl group" may have 1 to 5, preferably 1 to 3, for example, substituents optionally possessed by the aforementioned "hydrocarbon group", and the like.


[0042]

Preferable examples of the substituent of ring A include a halogen atom, a C₁₋₆ alkyl group optionally having substituent(s), a C₁₋₆ alkoxy group optionally having substituent(s), a hydroxyl group, a nitro group, a cyano group, an amino group optionally having substituent(s), an oxo group and the like. As the "substituent" of the "C₁₋₆ alkyl group optionally having substituent(s)", "C₁₋₆ alkoxy group optionally having substituent(s)" and "amino group optionally having substituent(s)", for example, substituents optionally possessed by the aforementioned "hydrocarbon group" can be mentioned.

Ring A optionally contains 1 to 4, preferably 1 or 2, substituents from the aforementioned substituents at substitutable position(s) according to the size of the ring. When the number of the substituents is two or more, the respective substituents may be the same or different.

Examples of the ring A include

[0043]



[0044]

wherein R^5 is a hydrogen atom or 1 or 2 substituents from the above-mentioned "preferable substituents for ring A", and other symbols are as defined above, and the like. Of these, one wherein R^5 is a hydrogen atom, a hydroxyl group or a C_{1-6} alkyl group optionally having substituent(s), particularly one wherein R^5 is a hydrogen atom (unsubstituted ring A), are widely used.

In the formula, n is an integer of 0 to 2 and n is preferably an integer of 0 or 1.

Particularly, n is preferably 0.

[0045]


In the aforementioned formula (I), ring B is a benzene ring optionally having substituent(s).

Examples of the substituent of ring B include the "substituent" of the above-mentioned "benzene ring optionally having substituent(s)". Among these, a halogen atom, a hydroxyl group or a lower (C_{1-6}) alkyl group optionally having substituent(s) is preferable, and particularly, a halogen atom, a hydroxyl group or a lower (C_{1-6}) alkyl group (preferably methyl) is widely used. The "substituent" of the above-mentioned "lower (C_{1-6}) alkyl group optionally having substituent(s)" is, for example, the substituent optionally possessed by the aforementioned "hydrocarbon group".

Ring B optionally has 1 or 2, preferably 1, substituent at substitutable position(s). When the number of the substituents is two or more, the respective substituents may be the same or different.

As ring B, for example,

[0046]



[0047]


wherein R^6 is a hydrogen atom, a hydroxyl group, a halogen atom, a lower (C_{1-6}) alkyl group optionally having substituent(s) or a lower (C_{1-6}) alkoxy group optionally having substituent(s), and the like are preferable. For example, R^6 is preferably a hydrogen atom, a hydroxyl group, a halogen atom or a lower (C_{1-6}) alkyl group (preferably methyl). More preferred is a hydrogen atom.

[0048]

In the aforementioned formula (I), m is an integer of 1 to 4, preferably an integer of 1 to 3. Furthermore, m is preferably 2 or 3, particularly preferably 2.

Preferable examples of the part represented by the formula


[0049]



[0050]

include

[0051]




20 [0052]

wherein $R^{4'}$ is a hydrocarbon group optionally having substituent(s) or a hydroxyl group, and other symbols are each as defined above, and the like.

$R^{4'}$ is preferably lower (C_{1-3}) alkyl optionally having
5 substituent(s).

Other preferable examples of the part represented by the formula


[0053]



10 [0054]

include


[0055]



[0056]

15 wherein each symbol is as defined above, and the like. Of these,

[0057]




[0058]

20 wherein each symbol is as defined above, and the like are

preferable.

Among these,


[0059]



5 [0060]

wherein each symbol is as defined above, and the like are preferable. Particularly preferred is

[0061]



10 [0062]

wherein each symbol is as defined above.

[0063]

Preferable examples of compound (I) include a compound wherein

- 15 R¹ is (i) a lower alkyl group optionally having substituent(s),
(ii) a lower cycloalkyl group optionally having substituent(s),
(iii) a lower alkenyl group optionally having substituent(s),
(iv) an aryl group optionally having substituent(s), (v) a
mono- or di-lower alkylamino group optionally having
20 substituent(s), (vi) an arylamino group optionally having
substituent(s) or (vii) a 5- or 6-membered nitrogen-containing
heterocyclic group optionally having substituent(s),
R² is a hydrogen atom or a lower (C₁₋₆) alkyl group optionally
having substituent(s),
25 R³ is (i) a hydrogen atom, (ii) a lower alkyl group optionally
having substituent(s) or (iii) an aryl group optionally having
substituent(s),
X is CHR⁴ or NR⁴ wherein R⁴ is a hydrogen atom or a lower (C₁₋₆)
alkyl group optionally substituted by an oxo group),

Y is C, CH or N (when X is CH₂, then Y is C or CH),
[0064]

[0065]

5 is a single bond or a double bond,
ring A is a 5- to 7-membered heterocycle containing an oxygen
atom and optionally having substituent(s),
ring B is a benzene ring optionally having substituent(s), and
m is 1 or 2, and the like.


10 [0066]

More preferable examples include a compound wherein
R¹ is (i) a C₁₋₆ alkyl group optionally substituted by 1 to 4
each of halogen, a hydroxyl group and a C₁₋₆ alkoxy group, (ii)
a C₃₋₆ cycloalkyl group, (iii) a C₂₋₆ alkenyl group, (iv) a C₆₋₁₀
15 aryl group optionally substituted by 1 to 4 each of C₁₋₆ alkoxy,
nitro, halogeno C₁₋₆ alkyl-carbonylamino and a halogen atom, (v)
a mono- or di-C₁₋₆ alkylamino group, (vi) a C₆₋₁₀ arylamino group
optionally substituted by 1 to 3 C₁₋₆ alkoxy groups or (vii) a
6-membered nitrogen-containing heterocyclic group optionally
20 substituted by 1 or 2 C₇₋₁₁ aralkyloxy-carbonyl groups, R² is a
hydrogen atom or a lower (C₁₋₆) alkyl group,
R³ is (i) a hydrogen atom, (ii) a lower (C₁₋₆) alkyl group or
(iii) a C₆₋₁₄ aryl group,
X is CHR⁴ or NR⁴ wherein R⁴ is a hydrogen atom or a lower (C₁₋₆)
25 alkyl group optionally substituted by an oxo group,
Y is C, CH or N (when X is CH₂, Y is C or CH,
[0067]

[0068]

30 is a single bond or a double bond,
ring A is


[0069]



[0070]

wherein each symbol is as defined above,
ring B is

5 [0071]



[0072]


wherein R^{6a} is a hydrogen atom, a halogen atom or a lower (C_{1-6})
alkyl group, and

10 m is 1 or 2, and the like.

[0073]

Among them, a compound represented by the formula


[0074]



15 [0075]

wherein R^{1b} is a C_{1-6} alkyl group optionally having a hydroxyl
group, R^{6b} is a hydrogen atom or a halogen atom, n is 0 or 1,

[0076]



20 [0077]

is a single bond or a double bond, when X^b is CH_2 , then

[0078]

a

[0079]

is a single bond or a double bond, when X^b is NH, then

[0080]


a

5

is a single bond, or a salt thereof is preferable.

In addition, a compound represented by the formula

[0081]



10 [0082]

wherein R^{1b} is a C_{1-6} alkyl group optionally having a hydroxyl group,

X' is CH_2 , NH or NCHO,

R^{3a} is a hydrogen atom or a phenyl group,

15 [0083]

[0084]

is a single bond or a double bond,

E^a is CH_2CH_2 , $CH=CH$, CH_2O , $CH=N$, CONH or

20 CH_2NH ,

n^a is 0 or 1,

ring A'' is a 5- or 6-membered heterocycle containing an oxygen atom and optionally having 1 or 2 C_{1-6} alkyl groups optionally substituted by a hydroxyl group, and

25 ring B' is a benzene ring optionally substituted by a halogen atom, or a salt thereof is also preferable. Among them, a compound wherein when X' is CH_2 or NCHO, then

[0085]

[0086]

is a single bond or a double bond, and when X' is NH, then

5 [0087]

[0088]

is a single bond, or a salt thereof is also preferable.

[0089]

10 Preferable examples of compound (I) include N-[2-(1,6,7,8-tetrahydro-2H-indeno[5,4-b]furan-8-yl)ethyl]acetamide, N-[2-(1,6,7,8-tetrahydro-2H-indeno[5,4-b]furan-8-yl)ethyl]butylamide, N-[2-(1,6,7,8-tetrahydro-2H-indeno[5,4-b]furan-8-yl)ethyl]propionamide,
15 N-[2-(3,7,8,9-tetrahydropyrano[3,2-e]indol-1-yl)ethyl]propionamide, N-[2-(5-fluoro-3,7,8,9-tetrahydrocyclopenta[f][1]benzopyran-9-yl)ethyl]propionamide,
20 N-[2-(5-fluoro-1,2,3,7,8,9-hexahydrocyclopenta[f][1]benzopyran-9-yl)ethyl]propionamide, N-[2-(3,7,8,9-tetrahydropyrano[3,2-e]indol-1-yl)ethyl]butylamide, N-[2-(1,2,3,7,8,9-hexahydropyrano[3,2-e]indol-1-yl)ethyl]propionamide,
25 N-[2-(1,2,3,7,8,9-hexahydropyrano[3,2-e]indol-1-yl)ethyl]butylamide, N-[2-(4-fluoro-1,6,7,8-tetrahydro-2H-indeno[5,4-b]furan-8-yl)ethyl]butylamide,
30 N-[2-(4-fluoro-1,6,7,8-tetrahydro-2H-indeno[5,4-b]furan-8-yl)ethyl]propionamide,
(S)-N-[2-(1,6,7,8-tetrahydro-2H-indeno[5,4-b]furan-8-yl)ethyl]propionamide,

(R)-N-[2-(1,6,7,8-tetrahydro-2H-indeno[5,4-b]furan-8-yl)ethyl]propionamide,
 N-[2-(1,6,7,8-tetrahydro-2H-indeno[5,4-b]furan-8-yl)ethyl]butylamide,
 5 N-[2-(1,6-dihydro-2H-indeno[5,4-b]furan-8-yl)ethyl]acetamide,
 N-[2-(1,6-dihydro-2H-indeno[5,4-b]furan-8-yl)ethyl]propionamide,
 N-[2-(1,6-dihydro-2H-indeno[5,4-b]furan-8-yl)ethyl]butylamide,
 N-[2-(7,8-dihydro-6H-indeno[4,5-d]-1,3-dioxol-8-yl)ethyl]propionamide,
 10 N-[2-(7,8-dihydro-6H-indeno[4,5-d]-1,3-dioxol-8-yl)ethyl]butylamide,
 N-[2-(2,3,8,9-tetrahydro-7H-indeno[4,5-b]-1,4-dioxin-9-yl)ethyl]propionamide,
 15 N-[2-(2,3,8,9-tetrahydro-7H-indeno[4,5-b]-1,4-dioxin-9-yl)ethyl]butylamide,
 N-[2-(1,6,7,8-tetrahydro-2H-furo[3,2-e]indol-8-yl)ethyl]propionamide,
 N-[2-(1,6,7,8-tetrahydro-2H-furo[3,2-e]indol-8-yl)ethyl]butylamide,
 20 N-[2-(7-phenyl-1,6-dihydro-2H-indeno[5,4-b]furan-8-yl)ethyl]propionamide,
 N-[2-(7-phenyl-1,6-dihydro-2H-indeno[5,4-b]furan-8-yl)ethyl]butylamide,
 25 (2S)-2-hydroxy-N-[2-[(8S)-1,6,7,8-tetrahydro-2H-indeno[5,4-b]furan-8-yl)ethyl]propionamide,
 N-[2-[(8R)-6-oxo-1,6,7,8-tetrahydro-2H-indeno[5,4-b]furan-8-yl)ethyl]propionamide,
 2-hydroxy-N-[2-[(8R)-6-oxo-1,6,7,8-tetrahydro-2H-indeno[5,4-b]furan-8-yl)ethyl]propionamide,
 30 N-[2-[(8S)-1-hydroxy-1,6,7,8-tetrahydro-2H-indeno[5,4-b]furan-8-yl)ethyl]propionamide,
 N-[2-[(8R)-1-hydroxy-6-oxo-1,6,7,8-tetrahydro-2H-indeno[5,4-b]furan-8-yl)ethyl]propionamide,
 35 N-[2-[(8R)-6-hydroxy-1,6,7,8-tetrahydro-2H-indeno[5,4-b]furan-

8-yl]ethyl]propionamide,
2-hydroxy-N-[2-[(8R)-6-hydroxy-1,6,7,8-tetrahydro-2H-
indeno[5,4-b]furan-8-yl]ethyl]propionamide,
N-[2-[(8S)-4-hydroxy-1,6,7,8-tetrahydro-2H-indeno[5,4-b]furan-
5 8-yl]ethyl]propionamide and the like.

[0090]

More preferred are N-[2-(1,6,7,8-tetrahydro-2H-
indeno[5,4-b]furan-8-yl)ethyl]acetamide,
N-[2-(1,6,7,8-tetrahydro-2H-indeno[5,4-b]furan-8-
10 yl)ethyl]propionamide,
N-[2-(5-fluoro-3,7,8,9-tetrahydrocyclopenta[f][1]benzopyran-9-
yl)ethyl]propionamide,
N-[2-(5-fluoro-1,2,3,7,8,9-
hexahydrocyclopenta[f][1]benzopyran-9-yl)ethyl]propionamide,
15 (S)-N-[2-(1,6,7,8-tetrahydro-2H-indeno[5,4-b]furan-8-
yl)ethyl]propionamide,
(R)-N-[2-(1,6,7,8-tetrahydro-2H-indeno[5,4-b]furan-8-
yl)ethyl]propionamide,
N-[2-(1,6,7,8-tetrahydro-2H-indeno[5,4-b]furan-8-
20 yl)ethyl]butylamide,
N-[2-(1,6-dihydro-2H-indeno[5,4-b]furan-8-yl)ethyl]acetamide,
N-[2-(1,6-dihydro-2H-indeno[5,4-b]furan-8-
yl)ethyl]propionamide,
N-[2-(1,6-dihydro-2H-indeno[5,4-b]furan-8-yl)ethyl]butylamide,
25 N-[2-(1,6,7,8-tetrahydro-2H-furo[3,2-e]indol-8-
yl)ethyl]propionamide,
N-[2-(1,6,7,8-tetrahydro-2H-furo[3,2-e]indol-8-
yl)ethyl]butylamide,
N-[2-(7-phenyl-1,6-dihydro-2H-indeno[5,4-b]furan-8-
30 yl)ethyl]propionamide,
N-[2-(7-phenyl-1,6-dihydro-2H-indeno[5,4-b]furan-8-
yl)ethyl]butylamide,
(2S)-2-hydroxy-N-[2-[(8S)-1,6,7,8-tetrahydro-2H-indeno[5,4-
b]furan-8-yl]ethyl]propionamide, and
35 N-[2-[(8R)-6-oxo-1,6,7,8-tetrahydro-2H-indeno[5,4-b]furan-8-

yl)ethyl]propionamide.

[0091]

Particularly preferred are (S)-N-[2-(1,6,7,8-tetrahydro-
2H-indeno[5,4-b]furan-8-yl)ethyl]propionamide,
5 N-[2-(1,6,7,8-tetrahydro-2H-furo[3,2-e]indol-8-
yl)ethyl]propionamide,
N-[2-(1,6,7,8-tetrahydro-2H-furo[3,2-e]indol-8-
yl)ethyl]butylamide,
N-[2-(7-phenyl-1,6-dihydro-2H-indeno[5,4-b]furan-8-
10 yl)ethyl]propionamide,
N-[2-(7-phenyl-1,6-dihydro-2H-indeno[5,4-b]furan-8-
yl)ethyl]butylamide,
(2S)-2-hydroxy-N-[2-[(8S)-1,6,7,8-tetrahydro-2H-indeno[5,4-
b]furan-8-yl)ethyl]propionamide, and
15 N-[2-[(8R)-6-oxo-1,6,7,8-tetrahydro-2H-indeno[5,4-b]furan-8-
yl)ethyl]propionamide.

[0092]

As the salt of compound (I), for example, a
pharmacologically acceptable salt and the like are used. For
20 example, a salt with inorganic base, a salt with organic base,
a salt with inorganic acid, a salt with organic acid, a salt
with basic or acidic amino acid and the like can be mentioned.
Preferable examples of the salt with inorganic base include
alkali metal salts such as sodium salt, potassium salt and the
25 like, alkaline earth metal salts such as calcium salt,
magnesium salt and the like, aluminum salt, ammonium salt and
the like. Preferable examples of the salt with organic base
include a salt with trimethylamine, triethylamine, pyridine,
picoline, 2,6-lutidine, ethanolamine, diethanolamine,
30 triethanolamine, cyclohexylamine, dicyclohexylamine, N,N'-
dibenzylethylenediamine and the like. Preferable examples of
the salt with inorganic acid include a salt with hydrochloric
acid, hydrobromic acid, nitric acid, sulfuric acid, phosphoric
acid and the like. Preferable examples of the salt with
35 organic acid include a salt with formic acid, acetic acid,

trifluoroacetic acid, phthalic acid, fumaric acid, oxalic acid, tartaric acid, maleic acid, citric acid, succinic acid, malic acid, methanesulfonic acid, benzenesulfonic acid, p-toluenesulfonic acid and the like. Preferable examples of the
5 salt with basic amino acid include a salt with arginine, lysine, ornithine and the like. Preferable examples of the salt with acidic amino acid include a salt with aspartic acid, glutamic acid and the like.

Of the above-mentioned salts, a pharmacologically
10 acceptable salt is preferable. When compound (I) has a basic functional group, for example, a salt with inorganic acid such as hydrochloric acid, hydrobromic acid, nitric acid, sulfuric acid, phosphoric acid and the like, a salt with organic acid such as acetic acid, phthalic acid, fumaric acid, tartaric
15 acid, maleic acid, citric acid, succinic acid, methanesulfonic acid, p-toluenesulfonic acid and the like can be mentioned, when it has an acidic functional group, for example, alkali metal salts such as sodium salt, potassium salt and the like, alkaline earth metal salts such as calcium salt, magnesium
20 salt and the like, ammonium salt and the like can be mentioned.

In addition, compound (I) may be a hydrate or a non-hydrate.

[0093]

As compound (I) or a salt thereof, particularly preferred
25 is (S)-N-[2-(1,6,7,8-tetrahydro-2H-indeno[5,4-b]furan-8-yl)ethyl]propionamide (general name: ramelteon).

[0094]

Compound (I) or a salt thereof [e.g., (S)-N-[2-(1,6,7,8-tetrahydro-2H-indeno[5,4-b]furan-8-yl)ethyl]propionamide] is a
30 known therapeutic agent for sleep disorder, which is disclosed in US Patent No. 6034239 and the like, and can be produced by a known method such as the method described in the US Patent and the like.

In addition, compound (I) can be synthesized according to
35 the production method described in JP-A-10-287665 or a method

analogous thereto and, when desired, said method in combination with a conventional oxidation method, and the like.
[0095]

Since compound (I) or a salt thereof, particularly
5 compound A, has an action to improve attentional deficit and suppress impulsivity, it can be used for the prophylaxis or treatment of attention deficit/hyperactivity disorder.

In addition, compound (I) or a salt thereof, particularly compound A, is expected to be effective for the prophylaxis or
10 treatment of bipolar disorder (particularly manic state).
[0096]

When compound (I) or a salt thereof (particularly, compound A) is administered to a patient, compound (I) or a salt thereof (particularly, compound A) can be safely
15 administered singly, or as a pharmaceutical composition containing pharmacologically acceptable carrier, such as tablet (including sugar-coated tablet, film-coated tablet), powder, granule, capsule, liquid, emulsion, suspension, injection, suppository, sustained-release preparation, plaster
20 and the like, orally or parenterally (e.g., topical, rectal or intravenous administration etc.).

The content of compound (I) or a salt thereof (particularly, compound A) is generally about 0.01 - 100 wt% of the whole composition.

25 While the dose of compound (I) or a salt thereof (particularly, compound A) varies depending on the subject of administration, administration route, disease and the like, for example, it is about 0.0005 - 2 mg/kg body weight, preferably about 0.001 - 1 mg/kg body weight, more preferably
30 about 0.001 - 0.5 mg/kg body weight, of compound (I) or a salt thereof (e.g., compound A) as an active ingredient in the form of an oral preparation for an adult, which can be administered in one to several portions a day.

[0097]

35 Since compound A shows extremely low toxicity, it can be

combined with other prophylactic or therapeutic agent for attention deficit/hyperactivity disorder and used for the prophylaxis or treatment of attention deficit/hyperactivity disorder, and a superior prophylactic or therapeutic effect
5 can be expected by such combined use with other medicament. It is further expected that such combination therapy decreases the dose of other prophylactic or therapeutic agent for attention deficit/hyperactivity disorder, and reduces side effects it has.

10 Examples of "other prophylactic or therapeutic agent for attention deficit/hyperactivity disorder" to be used in combination with compound A include central nervous stimulant (methylphenidate, atomoxetine, phemodafinil, armodafinil etc.).
[0098]

15 These "other prophylactic or therapeutic agent for attention deficit/hyperactivity disorder" may be in a free form or a pharmaceutically acceptable salt. Examples of such salt include, when the drug has an acidic functional group, for example, inorganic salts such as alkali metal salts (e.g.,
20 sodium salt, potassium salt etc.), alkaline earth metal salts (e.g., calcium salt, magnesium salt, barium salt etc.) and the like, ammonium salt and the like can be mentioned, and when it has a basic functional group, for example, a salt with organic acid such as hydrochloric acid, hydrobromic acid, nitric acid,
25 sulfuric acid, phosphoric acid and the like, a salt with organic acid such as acetic acid, phthalic acid, fumaric acid, oxalic acid, tartaric acid, maleic acid, citric acid, succinic acid, methanesulfonic acid, p-toluenesulfonic acid and the like can be mentioned. The "other prophylactic or therapeutic
30 agent for attention deficit/hyperactivity disorder" exemplified here is easily obtained as a commercially available product, or can be produced according to a known method.

[0099]

35 When compound (I) or a salt thereof (particularly,

compound A) and "other prophylactic or therapeutic agent for attention deficit/hyperactivity disorder" are used in combination, examples of the administration form include

- (1) administration of a single preparation obtained by simultaneously processing compound (I) or a salt thereof (particularly, compound A) and "other prophylactic or therapeutic agent for attention deficit/hyperactivity disorder",
- (2) simultaneous administration of two kinds of preparations of compound (I) or a salt thereof (particularly, compound A) and "other prophylactic or therapeutic agent for attention deficit/hyperactivity disorder", which have been separately produced, by the same administration route,
- (3) administration of two kinds of preparations of compound (I) or a salt thereof (particularly, compound A) and "other prophylactic or therapeutic agent for attention deficit/hyperactivity disorder", which have been separately produced, by the same administration route in a staggered manner,
- (4) simultaneous administration of two kinds of preparations of compound (I) or a salt thereof (particularly, compound A) and "other prophylactic or therapeutic agent for attention deficit/hyperactivity disorder", which have been separately produced, by different administration routes,
- (5) administration of two kinds of preparations of compound (I) or a salt thereof (particularly, compound A) and "other prophylactic or therapeutic agent for attention deficit/hyperactivity disorder", which have been separately produced, by different administration routes in a staggered manner (for example, administration in the order of compound (I) or a salt thereof (particularly, compound A) and "other prophylactic or therapeutic agent for attention deficit/hyperactivity disorder", or in the reverse order) and the like.

From the aspect of convenience for patients,

administration of a single preparation obtained by simultaneously processing compound (I) or a salt thereof (particularly, compound A) and "other prophylactic or therapeutic agent for attention deficit/hyperactivity disorder" is a particularly preferable embodiment.

[0100]

The dose of the "other prophylactic or therapeutic agent for attention deficit/hyperactivity disorder" can be appropriately determined based on the clinically employed dose as the standard. In addition, the mixing ratio of compound (I) or a salt thereof (particularly, compound A) and the "other prophylactic or therapeutic agent for attention deficit/hyperactivity disorder" can be appropriately determined according to the subject of administration, administration route, condition, the kind of combination drug to be used, and the like. Generally, it can be determined based on the general dose of the "other prophylactic or therapeutic agent for attention deficit/hyperactivity disorder" as the standard. When the subject of administration is human, for example, 0.01 - 100 parts by weight of the "other prophylactic or therapeutic agent for attention deficit/hyperactivity disorder" can be used per 1 part by weight of compound (I) or a salt thereof (particularly, compound A).

[0101]

The present invention is further explained in detail by the following Example and Formulation Examples. These Examples are mere Examples or Formulation Examples and do not limit the present invention.

Examples

[0102]

Example 1

A. Test method

(1) While spontaneously hypertensive rat (SHR) is a hypertension model rat widely used in research, this lineage

not only develops hypertension but also behaves similarly in the juvenile stage to patients with attention deficit/hyperactivity disorder (attention-deficit/hyperactivity disorder: ADHD), as evidenced by
5 disrupted retention of attention and the like (Sagvolden T, Russell VA, Aase H, Johansen EB, Farshbaf M. (2005): Rodent models of attention-deficit/hyperactivity disorder. Biol Psychiatry. 57(11): 1239-47). Particularly, SHR-SP (subtype of SHR) in the juvenile stage is recently recognized as an animal
10 model of developmental disorder such as ADHD by those of ordinary skill in the art (Ueno KI, Togashi H, Mori K, Matsumoto M, Ohashi S, Hoshino A, Fujita T, Saito H, Minami M, Yoshioka M. (2002): Behavioural and pharmacological relevance of stroke-prone spontaneously hypertensive rats as an animal
15 model of a developmental disorder. Behav Pharmacol. 13(1): 1-13).

SHR-SP is known to show not only enhanced spontaneous motility (hyperactivity) but also impairment of short-term memory due to attentional deficit and low anxiety condition
20 considered to be an index of impulsivity. In addition, methylphenidate, which is a first choice drug for ADHD treatment, shows an improving effect on these abnormal behaviors of SHR-SP at a low dose. Thus, those of ordinary skill in the art consider that SHR-SP is suitable as an ADHD
25 model.

Therefore, the inventor employed the SHR-SP rats and evaluated the effect of compound A on ADHD.

(2) The test was performed using 24 SHR-SP rats and 12 WKY rats (both 5-week-old; provided by CLEA Japan, Inc.). They
30 were reared at 12 hr light-dark cycle (light was on at 7:00 am) and allowed free access to water. The feeding was restricted such that the body weight was 85% of that by free food ingestion. All rats were subjected to handling before the test.

35 As a test apparatus, 4 Skinner boxes each equipped with a

45 mg feed pellet feeding tray (reward delivery opening) in the center of the wall and a cue lamp and a retractable lever set on the sides of the tray were set (see Fig. 1).

(3) The rats were trained for 2 days for lever pressing -
5 feeding associative learning. One session consisted of 30 trials or 900 sec. Two levers were presented, pressing either lever affords feed pellets (reinforcer), and a buzzer sound indicating correctness was made. Solely in the training on the initial day, feed pellets were automatically given with a
10 buzzer every 45 seconds. All animals reached the criteria obtaining 30 feed pellets within 300 sec.

(4) In the test session, a 2-lever discriminative reaction task was performed. In the 2-lever discriminative reaction task, 1 session consisted of 30 sets of ITI (inter-trial
15 interval) for 30 sec and reaction latency for 30 sec maximum. The house lamp was turned on at the start of the session to indicate both retractable levers. The leverpressing reaction of the animal was sequentially recorded while indicating the levers until completion of 30 trials. After completion of each
20 ITI, either cue lamp was turned on for 30 sec maximum in a pseudo-random order and when the animal responded to the lever on the side where the cue lamp was on, a reward was given along with a sound of the buzzer. In the contrary case, the cue lamp was turned off and the next ITI began. The levers
25 were retracted after completion of all 30 trials, and the house lamp was turned off 3 sec later, with which the session was ended. The one session of the experiment was performed once every week day except Saturdays and Sundays. To eliminate an influence of the apparatus such as house lamp illuminance,
30 sound volume, lever sensitivity and the like, the Skinner box used for the test was in rotation for each session.

(5) Drug administration

SHR-SP rats were divided into the positive drug group and the control group (12 per group) based on the total number of
35 reaction during trials in the first session of the 2-lever

discriminative reaction task.

The positive drug group was orally administered with compound A (1.0 mg/kg body weight), and the control group was orally administered with a solvent (0.5% methylcellulose injection water) alone at around 18:00. They were prepared immediately before administration every day and administered for 14 consecutive days up to session 11.

[0103]

B. Test results

(1) The transition of average values of three parameters of the number of correct answers, correct answer latency and total number of reaction during trials was compared. For comparison of SHR-SP rat and WKY rat, repeated measurement ANOVA was performed for the transition of the number of correct answers, correct answer latency and total number of reaction during trials. For comparison of the positive drug group and the control group, t-test was performed for the values of the last session.

(2) The leaning curves of the WKY rat and SHR-SP rat (control group) were compared. After progress of 11 session, both groups did not show much difference in the transition of "the number of correct answers (number of reward obtained)", which is an "index of attentional deficit", and the transition of the "correct answer latency (reaction latency when correct answer was selected)", which is an "index of motivation, sedation and the like" (see Figs. 3 and 4). However, in the transition of the "total number of reaction during trials", which is an "index of impulsivity", SHR-SP rats showed a significantly greater total number of reaction during trials ($p < 0.05$) (see Fig. 2). From these results, the usefulness of the SHR-SP rat as an evaluation model of impulsivity similar to ADHD was confirmed.

(3) Simultaneously, the SHR-SP rat was studied for the effect of continuous administration of compound A for the learning of the 2-lever discriminative reaction task. The transition of

the "total number of reaction during trials", which is an "index of impulsivity", was similar up to session 4 between the positive drug group and the control group. However, in session 11, the total number of reaction during trials in the positive drug group tended to decrease ($p = 0.09$) (see Fig. 5).

Therefrom it was clarified that compound A is effective for the suppression of impulsivity.

On the other hand, "the number of correct answers (number of reward obtained)", which is considered an "index of attentional deficit", tended to increase in the positive drug group ($p = 0.09$) (see Fig. 6). On the other hand, in the "correct answer latency (reaction latency when correct answer was selected)", which is an "index of motivation, sedation and the like", the positive drug group and the control group did not show much difference (see Fig. 7).

From the above findings, it is understood that compound (I) including compound A or a salt thereof improves attentional deficit and impulsivity.

[0104]

Formulation Example 1

Compound A (160 g), lactose (4064 g) and cornstarch (640 g) are uniformly mixed in a fluid bed dryer granulator, and the mixture is granulated while spraying an aqueous solution of hydroxypropylcellulose (160 g) in the granulator, and dried in the granulator. The obtained granulation product is milled in a POWER MILL using a 1.5 mmφ punching screen to give a sieved powder. The sieved powder (3894 g) is taken, cornstarch (124 g) and magnesium stearate (12.4 g) are added thereto and they are mixed to give granules for tableting. The granules are tableted by a tableting machine using a 7.0 mmφ punch to give a core tablet weighing 130 mg. A dispersion of titanium oxide and yellow ferric oxide in hydroxypropylmethylcellulose 2910 and copolividon is sprayed on the obtained core tablets in a film coating machine to give film-coated tablets (about 25000 tablets) containing 4 mg of compound A per tablet and

the formulation shown in Table 1.

[0105]

Table 1

composition	amount (mg)
compound A	4.0
lactose	101.6
cornstarch	20.0
hydroxypropylcellulose	4.0
magnesium stearate	0.4
core tablet	130.0
hydroxypropylmethylcellulose 2910	3.74
copovidon	0.75
titanium oxide	0.5
yellow ferric oxide	0.01
total	135.0

5 [0106]

This application is based on patent application No. 2008-022059 filed in Japan, the contents of which are encompassed in full herein.

Claims

[1] A prophylactic or therapeutic agent for attention deficit/hyperactivity disorder, comprising (S)-N-[2-(1,6,7,8-tetrahydro-2H-indeno[5,4-b]furan-8-yl)ethyl]propionamide as an active ingredient.

[2] A prophylactic or therapeutic agent for attention deficit/hyperactivity disorder, comprising (S)-N-[2-(1,6,7,8-tetrahydro-2H-indeno[5,4-b]furan-8-yl)ethyl]propionamide and one or more kinds of medicaments selected from other prophylactic or therapeutic agents for attention deficit/hyperactivity disorder in combination.

[3] A method for the prophylaxis or treatment of attention deficit/hyperactivity disorder, comprising administering an effective amount of (S)-N-[2-(1,6,7,8-tetrahydro-2H-indeno[5,4-b]furan-8-yl)ethyl]propionamide to a mammal.

[4] A method for the prophylaxis or treatment of attention deficit/hyperactivity disorder, comprising administering, to a mammal, an effective amount of (S)-N-[2-(1,6,7,8-tetrahydro-2H-indeno[5,4-b]furan-8-yl)ethyl]propionamide, and one or more kinds of medicaments selected from other prophylactic or therapeutic agents for attention deficit/hyperactivity disorder in combination.

[5] (S)-N-[2-(1,6,7,8-Tetrahydro-2H-indeno[5,4-b]furan-8-yl)ethyl]propionamide for use as a prophylactic or therapeutic agent for attention deficit/hyperactivity disorder.

[6] (S)-N-[2-(1,6,7,8-Tetrahydro-2H-indeno[5,4-b]furan-8-yl)ethyl]propionamide for use as a prophylactic or therapeutic agent for attention deficit/hyperactivity disorder in combination with one or more kinds of medicaments selected

from other prophylactic or therapeutic agents for attention deficit/hyperactivity disorder.

[7] Use of (S)-N-[2-(1,6,7,8-tetrahydro-2H-indeno[5,4-b]furan-
5 8-yl)ethyl]propionamide for the production of a prophylactic or therapeutic agent for attention deficit/hyperactivity disorder.

[8] Use of (S)-N-[2-(1,6,7,8-tetrahydro-2H-indeno[5,4-b]furan-
10 8-yl)ethyl]propionamide for the production of a prophylactic or therapeutic agent for attention deficit/hyperactivity disorder, in combination with one or more kinds of medicaments selected from other prophylactic or therapeutic agents for attention deficit/hyperactivity disorder.