

DESCRIPTION

TITLE OF THE INVENTION: TRANS-2-DECENOIC ACID DERIVATIVE AND DRUG CONTAINING SAME

5

TECHNICAL FIELD

[0001] The present invention relates to a novel trans-2-decenoic acid derivative or a pharmaceutically acceptable salt thereof, and a pharmaceutical agent containing the compound (in the present invention, in a case where it is simply referred to as a "compound," it may include a pharmaceutically acceptable salt thereof) as an active ingredient. More specifically, the present invention relates to a trans-2-decenoic acid derivative or a pharmaceutically acceptable salt thereof, having prophylactic or therapeutic actions for peripheral nerve disorders induced by administration of anti-cancer agents, neurotrophic factor-like activity such as nerve growth factor (NGF) or brain-derived neurotrophic factor (BDNF), and a pharmacological action such as analgesic action, and a pharmaceutical agent containing the compound as an active ingredient.

10

15

20

BACKGROUND ART

[0002] To date, in cancer (malignant tumor) treatment, surgery, irradiation or chemotherapy is used alone or in any combination thereof as required. Among them, anti-cancer agents (anti-malignant-tumor agents) used in the chemotherapy inherently have cytotoxicity and damage not only the cancer cells but also human normal cells to cause side effects. Thus, it is

25

important that the anti-cancer agents are administered to patients so as to prevent or treat such side effects as far as possible and to provide sufficient anti-cancer effects.

[0003] Examples of the side effects induced by administration of anti-cancer agents include variously blood disorders, gastrointestinal disorders, nerve disorders, etc. and, in particular, acute or chronic nerve disorders have increased as a recent trend. This trend is considered to be caused by the following factors: frequent occurrence of nerve disorders as a main side effect of new anti-cancer agents providing remarkable anticancer effects, the effects of multiple drug therapy as recent main therapy, and a relatively improved tendency of side effects such as blood disorders and gastrointestinal disorders. In this manner, no effective countermeasures against the nerve disorders, which are a main side effect caused by the current cancer chemotherapy, are available once the disorders have developed, due to the difficulty of nerve cell regeneration. Therefore, serious symptoms or irreversible disorders may be developed so that the administration of anti-cancer agents cannot be continued in some cases. Accordingly, the nerve disorders are an important therapeutic problem for cancer.

[0004] The nerve disorders induced by administration of anti-cancer agents are observed in, besides the central nervous system, the autonomic nervous system, and the peripheral nervous system, the sense organs such as the sense of taste. Among them, nerve disorders in the peripheral nervous system that occur in a comparatively high frequency to be problems are pains such as a stinging pain and burning pain, paresthesia such as

numbness of limb extremities and a burning sensation, hyperesthesia such as hypersensitivity to cold stimuli, dysesthesia such as sensory loss, sensory paralysis and sense of discomfort, sensory ataxia, muscle weakness or the like. In the present invention the anti-cancer agent inducing the peripheral nerve disorder may be any kinds of them, and the lesions in the peripheral nervous system induced by administration of anti-cancer agents are considered to be mainly due to neuroaxonal degeneration. Microtubules in the axon play an important role in maintaining the normal function of cells, for example, forming a spindle during cell division, and involving in placing the subcellular organelle and transporting substances. A taxane drug such as paclitaxel and docetaxel and a vinca alkaloid drug such as vincristine, vinblastine, vindesine and vinorelbine target the microtubules to inhibit the proliferation of cancer cells. Thus, it is considered that the microtubules in normal nerve cells are also more likely to be damaged to cause nerve disorders. Furthermore, it is considered that a platinum drug such as oxaliplatin, carboplatin, cisplatin and nedaplatin directly damage nerve cells, and consequently secondarily leading to axonopathy, so that it is considered to cause much of the nerve disorders.

[0005] In spite of the above situation, the neurotoxicity of the anti-cancer agents has not been studied in detail and sufficient prophylactic and therapeutic methods for the peripheral nerve disorder which is a side effect caused by taxane anti-cancer drugs, etc. as described above have yet to be established. Therefore, at present, for relieving numbness symptoms, vitamin B₁₂ preparations such as mecobalamin and a Chinese herbal medicine, Gosha-jinki-gan, are used. For pains, an antidepressant

(amitriptyline hydrochloride), an antiepileptic agent (carbamazepine), an antiarrhythmic agent (mexiletine hydrochloride), adrenocorticosteroid and the like are used. However, these therapies have limited effectiveness.

Stopping the administration of anti-cancer agent is the only reliable method for preventing the development of the peripheral nerve disorder, but even after stopping the administration, the peripheral nerve disorder may continue or get worse. It would require the administration of anti-cancer agents for treatment of cancer, but a significant peripheral nerve disorder can make it difficult to continue the administration of important anti-cancer agents which are highly effective against cancer and it becomes a serious problem for treatment of cancer. In view of the above problems, more effective prophylactic or therapeutic agents against a peripheral nerve disorder induced by anti-cancer agents have been strongly required in clinical practices.

[0006] In addition, a nerve cell is a cell having an information transmitting function, and its damage emerges as a serious loss of a cranial nerve function. The regeneration of the axon can hardly be expected in central nerves of the brain and the spinal cords, and thus, when nerve cells are found to have damages or degeneration, it is necessary to protect and activate the nerve cells. As the biological protective mechanism mentioned above, the roles of neurotrophic factors in differentiation of nerve cells, survival sustenance, increase in synapse functions, and regeneration or repair of the damaged nerve axon are indispensable.

[0007] Among neurotrophic factors, a nervous growth factor (NGF), a brain-derived neurotrophic factor (BDNF), neurotrophin-3 (NT-3),

neurotrophin-4/5 (NT-4/5), and the like construct a neurotrophin family having 50% or more of sequence homology with NGF as a prototype.

When the extracellularly secreted neurotrophin is bound to high-affinity receptor (Trks) on a nerve cell membrane, signals are transmitted in three

5 directions in the nerve cells. A transcription factor CREB (cAMP-response element binding protein) is activated through the activation of MAP kinase (mitogen-activated protein (MAP) kinases/extracellular signal-regulated protein kinases 1/2 (ERK1/2)) information transmission pathway, including the activation (phosphorylation) of the MAP kinase,
10 which is one of the high-affinity receptors, thereby regulating numerous gene expressions. Therefore, once signal transmissions through a MAP kinase information transmission pathway can be activated, there is a possibility of clinical applications to a nervous disorder caused by degeneration of nerve cells or cellular death. Further, there is a report
15 about associations between BDNF and some diseases.

[0008] According to the studies on gene polymorphisms of BDNF, it is reported that particular polymorphisms of the BDNF are associated with Parkinson's disease, Alzheimer's disease, depression, bipolar depression, anxiety disorders, autistic disorder spectrum, glaucoma, and the like.

20 Furthermore, there are some reports that decrease in a synapse function of a genetically converted mouse having Huntington's disease is cured with administration of a BDNF, and that administration of a MAP kinase phosphorylation inhibitor provokes an antidepressant condition.

[0009] Neurotrophic factors, as is seen in examples of BDNF mentioned
25 above, show therapeutic effects against particular nerve diseases, and have

some actions of germinating the axon and extending its length. However, since neurotrophic factors are high-molecular weight proteins, they have a problem of having difficulty in reaching the brain since they cannot pass through a blood-brain barrier even if administered from a peripheral. Thus, it has been tried to search for a pharmaceutical agent having a neurotrophic factor-like activity that activates nerve cells with a low-molecular weight compound and a pharmaceutical agent promoting the production and secretion of neurotrophic factors.

[0010] Patent Publication 1 discloses that a fatty acid or a fatty acid ester having 8 carbon atoms or 10 to 12 carbon atoms has a neurotrophic factor-like activity. However, the compounds described in Patent Publication 1 is an ordinary fatty acid or an ester thereof, clearly differing in the structure in the carboxylic acid moiety from the compounds of the present invention. In addition, the compounds of the present invention, 1-((*E*)-2-decenoyl)pyrrolidine (Compound 1), 1-((*E*)-2-decenoyl)piperidine (Compound 5) and 1-((*E*)-2-decenoyl)azepane (Compound 32), are disclosed in Patent Publication 2, and the compound of the present invention, (4-((*E*)-2-decenoyl) morpholine (Compound 17), is disclosed in Patent Publication 3 or Non-Patent Publication 1, respectively. However, Patent Publication 2 relates to a toxic action against molluskans, and Patent Publication 3 relates to growth inhibitory actions against disease germs or molds, and Non-Patent Publication 1 is a publication relating to a repellent against houseflies. In conclusion, none of these publications include pharmaceutical applications of the compounds of the present invention and any descriptions suggesting the applications.

PRIOR ART REFERENCES

PATENT PUBLICATIONS

- [0011] Patent Publication 1: WO 2009/038110
Patent Publication 2: WO 2010/123894
Patent Publication 3: U.S. Patent No. 3,294,794

NON-PATENT PUBLICATIONS

- [0012] Non-Patent Publication 1: *Journal of Economic Entomology*, 1970,
63(6), 1752-1755

SUMMARY OF THE INVENTION

PROBLEMS TO BE SOLVED BY THE INVENTION


- [0013] An object of the present invention is to provide a novel trans-2-decenoic acid derivative or a pharmaceutically acceptable salt thereof, and a pharmaceutical agent containing the compound as an active ingredient, and more specifically to provide a pharmaceutical agent having a prophylactic or therapeutic action against a peripheral nerve disorder induced by an anti-cancer agent, a neurotrophic factor-like activity or a pharmacological action such as analgesic action, wherein in the present invention, the "peripheral nerve disorder" can be replaced with a synonym "neuropathy"; in addition, "prevention" and "treatment" include "improvement" or "alleviation," and "prophylactic agent" and "therapeutic agent" include "improvement agent" or "alleviating agent."

MEANS TO SOLVE THE PROBLEMS

[0014] As a result of intensively studies in order to solve the above problems, the present inventors have found that a trans-2-decenoic acid derivative represented by the general formula (I) defined below or a pharmaceutically acceptable salt thereof has an excellent pharmacological action such as a prophylactic or therapeutic action for peripheral nerve disorders induced by anti-cancer agents, a neurotrophic factor-like activity, or an analgesic action. The present inventors have made further studies on the basis of the above findings, and whereby the present invention has been accomplished.

[0015] Specifically, the present invention provides the following compounds and pharmaceutical agents containing the compounds.

(1) A trans-2-decenoic acid derivative represented by the following general formula (I') or a pharmaceutically acceptable salt thereof:



wherein X' is:

- (a) a 1-pyrrolidyl substituted with carboxyl or alkoxycarbonyl,
- (b) a 3-thiazolidyl,
- (c) a piperidino substituted with alkyl, oxo, hydroxy, alkoxy, carboxyl, alkoxycarbonyl, alkylamino, alkylaminoalkyl, phenyl, carboxyalkyl, alkoxycarbonylalkyl, cyano, or halogenophenyl,
- (d) a thiomorpholino,
- (e) a 1-piperazyl which may be substituted with alkyl, carboxyalkyl,

alkoxycarbonylalkyl, alkylaminoalkyl, cycloalkyl, piperidinoalkyl,
phenylalkyl, pyridyl, pyrimidyl, carboxyphenylalkyl or
alkoxycarbonylphenylalkyl,

(f) a 1-piperazyl substituted with a phenyl which may be substituted
with alkylamino, halogen, alkoxy, alkyl, hydroxy, carboxyalkoxy or
alkoxycarbonylalkoxy,

(g) a 1,4-diazepanyl which may be substituted with alkyl or
alkylaminoalkyl, or

(h) a carboxymorpholino.

(2) The trans-2-decenoic acid derivative or a pharmaceutically
acceptable salt thereof according to the above (1), wherein X' is a 1-
pyrrolidyl substituted with carboxyl or alkoxycarbonyl.

(3) The trans-2-decenoic acid derivative or a pharmaceutically
acceptable salt thereof according to the above (1), wherein X' is a 3-
thiazolidyl.

(4) The trans-2-decenoic acid derivative or a pharmaceutically
acceptable salt thereof according to the above (1), wherein X' is a
piperidino substituted with alkyl, oxo, hydroxy, alkoxy, carboxyl,
alkoxycarbonyl, alkylamino, alkylaminoalkyl, phenyl, carboxyalkyl,
alkoxycarbonylalkyl, cyano or halogenophenyl.

(5) The trans-2-decenoic acid derivative or a pharmaceutically
acceptable salt thereof according to the above (1), wherein X' is a
thiomorpholino.

(6) The trans-2-decenoic acid derivative or a pharmaceutically
acceptable salt thereof according to the above (1), wherein X' is a 1-


piperazyl which may be substituted with alkyl, carboxyalkyl, alkoxycarbonylalkyl, alkylaminoalkyl, cycloalkyl, piperidinoalkyl, phenylalkyl, pyridyl, pyrimidyl, carboxyphenylalkyl or alkoxycarbonylphenylalkyl.

5 (7) The trans-2-decenoic acid derivative or a pharmaceutically acceptable salt thereof according to the above (1), wherein X' is a 1-piperazyl substituted with a phenyl which may be substituted with alkylamino, halogen, alkoxy, alkyl, hydroxy, carboxyalkoxy or alkoxycarbonylalkoxy.

10 (8) The trans-2-decenoic acid derivative or a pharmaceutically acceptable salt thereof according to the above (1), wherein X' is a 1,4-diazepanyl which may be substituted with alkyl or alkylaminoalkyl.

(9) The trans-2-decenoic acid derivative or a pharmaceutically acceptable salt thereof according to the above (1), wherein X' is a
15 carboxymorpholino.

[0016] (10) A pharmaceutical agent containing, as an active ingredient, at least one member of a trans-2-decenoic acid derivative represented by the following general formula (I) and a pharmaceutically acceptable salt thereof:



20

wherein X is:

- (a) a 1-pyrrolidyl substituted with carboxyl or alkoxycarbonyl,
- (b) a 3-thiazolidyl,

(c) a piperidino substituted with alkyl, oxo, hydroxy, alkoxy, carboxyl, alkoxy carbonyl, alkylamino, alkylaminoalkyl, phenyl, carboxyalkyl, alkoxy carbonylalkyl, cyano, or halogenophenyl,

(d) a thiomorpholino,

5 (e) a 1-piperazyl which may be substituted with alkyl, carboxyalkyl, alkoxy carbonylalkyl, alkylaminoalkyl, cycloalkyl, piperidinoalkyl, phenylalkyl, pyridyl, pyrimidyl, carboxyphenylalkyl or alkoxy carbonylphenylalkyl,

10 (f) a 1-piperazyl substituted with a phenyl which may be substituted with alkylamino, halogen, alkoxy, alkyl, hydroxy, carboxyalkoxy or alkoxy carbonylalkoxy,

(g) a 1,4-diazepanyl which may be substituted with alkyl or alkylaminoalkyl, or

(h) a carboxymorpholino.

15 (11) The pharmaceutical agent according to the above (10), wherein X is a 1-pyrrolidyl substituted with carboxyl or alkoxy carbonyl.

(12) The pharmaceutical agent according to the above (10), wherein X is a 3-thiazolidyl.

20 (13) The pharmaceutical agent according to the above (10), wherein X is a piperidino substituted with alkyl, oxo, hydroxy, alkoxy, carboxyl, alkoxy carbonyl, alkylamino, alkylaminoalkyl, phenyl, carboxyalkyl, alkoxy carbonylalkyl, cyano, or halogenophenyl.

(14) The pharmaceutical agent according to the above (10), wherein X is a thiomorpholino.

25 (15) The pharmaceutical agent according to the above (10), wherein X is

a 1-piperazyl which may be substituted with alkyl, carboxyalkyl, alkoxycarbonylalkyl, alkylaminoalkyl, cycloalkyl, piperidinoalkyl, phenylalkyl, pyridyl, pyrimidyl, carboxyphenylalkyl or alkoxycarbonylphenylalkyl.

5 (16) The pharmaceutical agent according to the above (10), wherein X is a 1-piperazyl substituted with a phenyl which may be substituted with alkylamino, halogen, alkoxy, alkyl, hydroxy, carboxyalkoxy or alkoxycarbonylalkoxy.

(17) The pharmaceutical agent according to the above (10), wherein X is
10 a 1,4-diazepanyl which may be substituted with alkyl or alkylaminoalkyl.

(18) The pharmaceutical agent according to the above (10), wherein X is a carboxymorpholino.

[0017] (19) The pharmaceutical agent according to any one of the above (10) to (18), wherein the pharmaceutical agent is a prophylactic agent or
15 therapeutic agent for peripheral nerve disorders induced by administration of a drug containing at least one anti-cancer agent.

(20) The pharmaceutical agent according to any one of the above (10) to (18), wherein the pharmaceutical agent is an analgesic against pain diseases.

20 [0018] (21) The pharmaceutical agent according to any one of the above (10) to (18), wherein the pharmaceutical agent is an injectable preparation.

(22) The pharmaceutical agent according to any one of the above (10) to (18), wherein the pharmaceutical agent is an oral preparation.

(23) The pharmaceutical agent according to any one of the above (10) to
25 (18), wherein the pharmaceutical agent is an external preparation.

EFFECTS OF THE INVENTION

[0019] The compound of the present invention is effective as a drug for prevention or treatment of nerve disorders of the peripheral nerve system induced by anti-cancer agents such as paresthesia such as numbness of limb extremities of human and animals, and hyperalgesia such as pains.

[0020] In addition, the compound of the present invention has an excellent neurotrophic factor-like activity, so that the compound is usable as an agent having a neurotrophic factor-like activity. This agent having a neurotrophic factor-like activity is useful as prophylactic agents or therapeutic agents for nerve disorders because signal transmissions are activated via MAP kinase information transmission pathway with a neurotrophic factor-like activity. Among the nerve disorders, the agent is especially useful as prophylactic agents or therapeutic agents for a neurodegenerative disease, such as dementia, Alzheimer's disease, Parkinson's disease, amyotrophic lateral sclerosis (ALS), Huntington's disease, progressive supranuclear palsy (PSP), diabetic neuropathy, or glaucoma. In addition, among the nerve disorders, the agent is also useful as an improving agent for prevention of a mental disease. Among the mental diseases, the agent is especially useful as prophylactic agents or improving agents for depression, anxiety disorders (neuroses), or autistic disorder spectrum, and particularly useful as prophylactic agents or therapeutic agents for depression and anxiety disorders (neuroses).


[0021] In addition, the compound of the present invention is useful as a therapeutic agent for spinal cord injury (in the case of spinal cord injury, it

is sometimes called as “repairing agent”).

Further, the compound of the present invention is a compound showing an excellent analgesic action, which is useful as a drug for prevention or treatment of various pain diseases such as pains caused by arthralgia such as orthoarthritis.

MODES FOR CARRYING OUT THE INVENTION

[0022] The present invention relates to a trans-2-decenoic acid derivative represented by the following general formula (I') or a pharmaceutically acceptable salt thereof:



wherein X' is:


- (a) a 1-pyrrolidyl substituted with carboxyl or alkoxycarbonyl,
- (b) a 3-thiazolidyl,
- (c) a piperidino substituted with alkyl, oxo, hydroxy, alkoxy, carboxyl, alkoxycarbonyl, alkylamino, alkylaminoalkyl, phenyl, carboxyalkyl, alkoxycarbonylalkyl, cyano, or halogenophenyl,
- (d) a thiomorpholino,
- (e) a 1-piperazyl which may be substituted with alkyl, carboxyalkyl, alkoxycarbonylalkyl, alkylaminoalkyl, cycloalkyl, piperidinoalkyl, phenylalkyl, pyridyl, pyrimidyl, carboxyphenylalkyl or alkoxycarbonylphenylalkyl,
- (f) a 1-piperazyl substituted with a phenyl which may be substituted

with alkylamino, halogen, alkoxy, alkyl, hydroxy, carboxyalkoxy or
alkoxycarbonylalkoxy,

(g) a 1,4-diazepanyl which may be substituted with alkyl or
alkylaminoalkyl, or

5 (h) a carboxylmorpholino.

[0023] In addition, the present invention relates to a pharmaceutical agent,
such as a prophylactic agent or a therapeutic agent of a peripheral nerve
disorder induced by an anti-cancer agent, an agent having a neurotrophic
factor-like activity, or an analgesic, containing, as an active ingredient, at
10 least one member of a trans-2-decenoic acid derivative represented by the
following general formula (I) and a pharmaceutically acceptable salt
thereof:



wherein X is:

- 15 (a) a 1-pyrrolidyl which may be substituted with carboxyl or
alkoxycarbonyl,
- (b) a 3-thiazolidyl,
- (c) a piperidino which may be substituted with alkyl, oxo, hydroxy,
alkoxy, carboxyl, alkoxycarbonyl, alkylamino, alkylaminoalkyl, phenyl,
20 carboxyalkyl, alkoxycarbonylalkyl, cyano, or halogenophenyl,
- (d) a morpholino which may be substituted with carboxyl,
- (e) a thiomorpholino,
- (f) a 1-piperazyl which may be substituted with alkyl, carboxyalkyl,

alkoxycarbonylalkyl, alkylaminoalkyl, cycloalkyl, piperidinoalkyl, phenylalkyl, pyridyl, pyrimidyl, carboxyphenylalkyl or alkoxycarbonylphenylalkyl,

(g) a 1-piperazyl substituted with a phenyl which may be substituted with alkylamino, halogen, alkoxy, alkyl, hydroxy, carboxyalkoxy or alkoxycarbonylalkoxy,

(h) a 1-azepanyl, or

(i) a 1,4-diazepanyl which may be substituted with alkyl or alkylaminoalkyl. The compound represented by the general formula (I)

embraces the compound represented by the general formula (I') mentioned above.

[0024] In the substituents of the general formulae (I) and (I') mentioned above, the alkyl (including "alkyl" in alkylamino, alkylaminoalkyl, carboxyalkyl, alkoxycarbonylalkyl, piperidinoalkyl, phenylalkyl, carboxyphenylalkyl, or alkoxycarbonylphenylalkyl) may be any ones, and the alkyl is linear or branched alkyl groups having from 1 to 4 carbon atoms, preferably including methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, t-butyl, etc. Also, the alkylamino (including "alkylamino" in alkylaminoalkyl) refers to an amino group substituted with 1 or 2 alkyls.


[0025] The alkoxy (including "alkoxy" in alkoxycarbonyl, carboxyalkoxy, or alkoxycarbonylalkoxy) may be any ones, and the alkoxy is a linear or branched alkoxy group having from 1 to 4 carbon atoms, preferably including methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, isobutoxy, sec-butoxy, t-butoxy, etc.

The cycloalkyl may be any ones, and the cycloalkyl is a cycloalkyl group having from 3 to 8 carbon atoms, preferably including cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, etc., and more preferably a cycloalkyl group having 5 or 6 carbon atoms.

5 The halogen (including "halogeno" in the halogenophenyl) is fluorine, chlorine, bromine, iodine, etc.

[0026] The compound represented by the general formula (I) (including the compound represented by the formula (I'); hereinafter referred to the same) of the present invention can be produced using trans-2-decenoic acid as a raw material in the manner, for example, as shown in the following reaction formula:

Reaction Formula



wherein X is as defined above.

[0027] The compound represented by the general formula (I) can be produced by subjecting a compound represented by the general formula (II) and a compound represented by the general formula (III) having an active hydrogen to dehydration condensation. As the dehydration condensation reaction, a known method can be employed. For example, a compound represented by the general formula (II) and a compound represented by the general formula (III) can be allowed to react in the

presence of an appropriate condensing agent, including, for example, dicyclohexylcarbodiimide (DCC) or N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide•HCl. The reaction may be usually carried out in a solvent, including, for example, dichloromethane. The amount of the compound represented by the general formula (III) used is usually from 0.5 to 2 mol, and preferably from 1 to 1.5 mol, per one mol of the compound represented by the general formula (II).

[0028] Alternatively, for example, the compound represented by the general formula (II) is once converted to a carboxylic halide, and thereafter allowed to react with a compound represented by the general formula (3) in the presence or absence of a base. The conversion to the carboxylic halide may be carried out, for example, using a halogenating agent such as thionyl chloride, sulfuryl chloride, phosphorus trichloride, phosphorus pentachloride, oxalyl chloride or phosphoric acid trichloride. Examples of the base include triethylamine and pyridine. The amount of the compound represented by the general formula (III) used is usually from 0.5 to 2 mol, and preferably from 1 to 1.5 mol, per one mol of the compound represented by the general formula (II). When a base is used, the amount of the base used is usually from 1 to 5 mol or so, per one mol of the compound represented by the general formula (II).

After the termination of the above reaction, known purification and isolation procedures, including, for example, extraction, chromatography, distillation, or recrystallization, are employed, whereby an intended compound can be obtained.












[0029] Examples of the compound produced as such are shown in Tables 1

to 5. When each compound is referred to hereinafter, the compound number mentioned in the tables is used.

[Table 1]











No.	Compound Name	Structural Formula
1	1-((<i>E</i>)-2-Decenoyl)pyrrolidine (Reference compound)	
2	(<i>S</i>)-1-((<i>E</i>)-2-Decenoyl)pyrrolidine-2-carboxylic acid	
3	Methyl (<i>S</i>)-1-((<i>E</i>)-2-Decenoyl)-pyrrolidine-2-carboxylate	
4	3-((<i>E</i>)-2-Decenoyl)thiazolidine	
5	1-((<i>E</i>)-2-Decenoyl)piperidine (Reference compound)	
6	1-((<i>E</i>)-2-Decenoyl)-4-methylpiperidine	
7	1-((<i>E</i>)-2-Decenoyl)-piperidin-4-one	
8	1-((<i>E</i>)-2-Decenoyl)-4-hydroxypiperidine	
9	1-((<i>E</i>)-2-Decenoyl)-4-methoxypiperidine	
10	1-((<i>E</i>)-2-Decenoyl)piperidine-4-carboxylic acid	
11	Ethyl 1-((<i>E</i>)-2-Decenoyl)-piperidine-4-carboxylate	
12	1-((<i>E</i>)-2-Decenoyl)-4-dimethylaminopiperidine	
13	1-((<i>E</i>)-2-Decenoyl)-4-diethylaminopiperidine	

[0030] [Table 2]

No.	Compound Name	Structural Formula
14	1-((<i>E</i>)-2-Decenoyl)-4-diethylaminomethylpiperidine	
15	1-((<i>E</i>)-2-Decenoyl)-4-(2-dimethylaminoethyl)piperidine	
16	1-((<i>E</i>)-2-Decenoyl)-4-phenylpiperidine	
17	4-((<i>E</i>)-2-Decenoyl)morpholine (Reference compound)	
18	4-((<i>E</i>)-2-Decenoyl)-thiomorpholine	
19	1-((<i>E</i>)-2-Decenoyl)-4-methylpiperazine	
20	1-((<i>E</i>)-2-Decenoyl)-4-isopropylpiperazine	
21	3-[4-((<i>E</i>)-2-Decenoyl)piperazin-1-yl]propionic acid	
22	1-((<i>E</i>)-2-Decenoyl)-4-[2-(dimethylamino)ethyl]piperazine	
23	4-Cyclohexyl-1-((<i>E</i>)-2-decenoyl)piperazine	
24	1-((<i>E</i>)-2-Decenoyl)-4-(2-piperidin-1-ylethyl)piperazine	

[0031]

[Table 3]






No.	Compound Name	Structural Formula
25	1-((<i>E</i>)-2-Decenoyl)-4-phenylpiperazine	
26	4-Benzyl-1-((<i>E</i>)-2-decenoyl)-piperazine	
27	1-((<i>E</i>)-2-Decenoyl)-4-(2-phenylethyl)piperazine	
28	1-((<i>E</i>)-2-Decenoyl)-4-(4-dimethylaminophenyl)-piperazine	
29	1-((<i>E</i>)-2-Decenoyl)-4-(pyridin-4-yl)piperazine	
30	1-((<i>E</i>)-2-Decenoyl)-4-(pyridin-2-yl)piperazine	
31	1-((<i>E</i>)-2-Decenoyl)-4-(pyrimidin-2-yl)piperazine	
32	1-((<i>E</i>)-2-Decenoyl)azepane (Reference compound)	
33	1-((<i>E</i>)-2-Decenoyl)-4-methyl-[1,4]diazepane	
34	1-((<i>E</i>)-2-Decenoyl)-4-(2-dimethylaminoethyl)-[1,4]diazepane	

[0032]

[Table 4]

No.	Compound Name	Structural Formula
35	2-(1-((<i>E</i>)-2-Decenoyl)-4-piperidyl)acetic acid	
36	3-(1-((<i>E</i>)-2-Decenoyl)-4-piperidyl)propanoic acid	
37	1-((<i>E</i>)-2-Decenoyl)-4-cyanopiperidine	
38	1-((<i>E</i>)-2-Decenoyl)-4-(4-chlorophenyl)piperidine	
39	[4-[(<i>E</i>)-2-Decenoyl]-piperazin-1-yl]acetic acid hydrochloride	
40	4-[4-[(<i>E</i>)-2-Decenoyl]-piperazin-1-yl]butanoic acid hydrochloride	
41	1-((<i>E</i>)-2-Decenoyl)-4-(4-chlorophenyl)piperazine	
42	1-((<i>E</i>)-2-Decenoyl)-4-(4-methoxyphenyl)piperazine	
43	1-((<i>E</i>)-2-Decenoyl)-4-(4-methylphenyl)piperazine	
44	1-((<i>E</i>)-2-Decenoyl)-4-(4-fluorophenyl)piperazine	

[0033] [Table 5]

No.	Compound Name	Structural Formula
45	1-((E)-2-Decenoyl)-4-(2-chlorophenyl)piperazine	
46	1-((E)-2-Decenoyl)-4-(4-hydroxyphenyl)piperazine	
47	2-[4-[4-((E)-2-Decenoyl)piperazin-1-yl]-phenoxy]acetic acid	
48	4-[4-[[4-((E)-2-Decenoyl)piperazin-1-yl]-methyl]phenyl]benzoic acid	
49	4-[(E)-2-Decenoyl]morpholin-3-carboxylic acid	

[0034] When a salt of the compound represented by the general formula (I) of the present invention is formed, in a case where a compound is used as a pharmaceutical agent, the form of a pharmaceutically acceptable salt is preferred. Examples of the salts include addition salts with an acid such as phosphoric acid, hydrochloric acid, sulfuric acid, nitric acid, hydrobromic acid, perchloric acid, thiocyanic acid, boric acid, formic acid, acetic acid, haloacetic acid, propionic acid, citric acid, tartaric acid, lactic acid, glycolic acid, succinic acid, gluconic acid, malonic acid, fumaric acid, anthranilic acid, benzoic acid, cinnamic acid, p-toluenesulfonic acid,

naphthalenesulfonic acid, or sulfanilic acid. Also, the salts include salts with a metal, such as an alkali metal such as sodium or potassium, an alkaline earth metal such as calcium or magnesium, or aluminum; and salts with a base such as ammonia or an organic amine. These salts can be produced from each compound in a free form, or converted reversibly, in accordance with a known method. In addition, in a case where the compound represented by the general formula (I) of the present invention contains an asymmetric carbon, the compound also embraces various isomers such as optical isomers, racemic mixtures, and diastereomers. In a case where the compound of the present invention is formed into crystals, the crystals also embrace various kinds of crystalline forms (crystal polymorphisms) which can be formed thereby.

[0035] The compound represented by the general formula (I) of the present invention is effective as an agent for having prophylactic or therapeutic action for a peripheral nerve disorder induced by administration of anti-cancer agents. One of the anti-cancer agents developing the peripheral nerve disorder is an anti-cancer agent that damages microtubules to induce the peripheral nerve disorder. Examples of such medicinal agent include taxane drug such as paclitaxel or docetaxel, and vinca alkaloid drug such as vincristine, vinplastine, vindesine or vinorelbine. Another anti-cancer agent is an anti-cancer agent that damages nerve cells to cause axonopathy and then induces the peripheral nerve disorder. Examples of such agents include platinum formulation such as oxaliplatin, carboplatin, cisplatin or nedaplatin.

[0036] Examples of the peripheral nerve disorder induced by these anti-

cancer agents include pains such as a stinging pain and burning pain, numbness of limb extremities, paresthesia such as a burning sensation, hyperesthesia such as hypersensitivity to cold stimuli, dysesthesia such as sensory loss, sensory paralysis and sense of discomfort, sensory ataxia and muscle weakness. The peripheral nerve disorder induced by anti-cancer agents targeted for the prevention or treatment in the compounds of the present invention includes a peripheral nerve disorder induced by monotherapy using one of anti-cancer agents as well as a peripheral nerve disorder induced by multiple drug therapy in which a plurality of medicinal agents having various action mechanisms are together administered or by biochemical modulation in which a combination of medicinal agents and an administration method are designed such that the medicinal agents having various action mechanisms can provide the maximum effectiveness.

[0037] Since the compound represented by the general formula (I) of the present invention has a neurotrophic factor-like activity, the compound is useful as an agent having a neurotrophic factor-like activity, and is useful in the prevention and treatment of nervous disorders. The nervous disorder refers to a clinical condition of damaging functions of nerve cells due to degeneration or cell death of the nerve cells, and includes neurodegenerative diseases and mental diseases. The neurodegenerative diseases refer to dementia, Alzheimer's disease, Parkinson's disease, amyotrophic lateral sclerosis (ALS), Huntington's disease, progressive supranuclear palsy (PSP), diabetic neuropathy, glaucoma of optic nerve diseases etc. The mental diseases refer to depression (including bipolar depression), anxiety disorder (neurosis), integration disorder syndrome,

autistic disorder spectrum etc. When the agent is used for depression, it takes at least 3 to 4 weeks until there appear effects of conventionally existing depression therapeutic agents such as a tricyclic antidepressant, a tetracyclic antidepressant, a selective serotonin reuptake inhibitor (SSRI), a serotonin noradrenalin reuptake inhibitor (SNRI), and the like, and these agents had to be taken periodically during this term; however, the pharmaceutical agent containing the compound of the present invention can be expected to have immediate affectivity as compared to existing medical drugs.

10 **[0038]** The compound represented by the general formula (I) is useful as a therapeutic agent (repairing agent) for spinal cord injury. There are no effective methods of treatment for spinal cord injury suffering physical damages caused by traffic accidents, athletic sports accidents, or compressive fractures of elder people, and various studies on methods of treatment by regeneration therapy have been made. It is expected that the pharmaceutical agent containing the compound of the present invention can treat (repair) spinal cord injury with administrations by injections, internal medication, and external applications.

15 **[0039]** In addition, the compound represented by the general formula (I) of the present invention is useful as prophylactic agents or therapeutic agents for various pain diseases. Examples of the pain diseases include arthralgia such as the pains by osteoarthritis (e.g., knee osteoarthritis and hip osteoarthritis) or the pains by rheumatoid arthritis.

20 **[0040]** The compounds of the present invention can be made into a pharmaceutical preparation in various dosage forms (such as oral,

25

injectable and external preparations) by appropriately combining with an appropriate pharmaceutical carrier or diluent. The pharmaceutical agent of the present invention may be a formulation in which the compound of the present invention is used together with other pharmaceutically active ingredient(s). Further, the pharmaceutical agent of the present invention may be formed into a preparation as an inclusion complex with cyclodextrin etc. As a result, there may be the cases where enhancement of pharmacological activity, improvement in stability, prolonged action, easy handling, etc. can be obtained. The inclusion complex can be formed by, for example, mixing the compound of the present invention with α -, β -, or γ -cyclodextrin in accordance with an ordinary method.

[0041] When the compound of the present invention is made into an oral preparation, a tablet, powder, granule or capsule preparation can be made by formulation of proper combinations of appropriate additives, including, for example, an excipient, a binder, a disintegrator, a lubricant, an extender, a wetting agent, a buffer, a preservative and a flavoring. When made into an injectable preparation, it is possible to make into an injectable preparation by addition of a stabilizer, a preservative, an isotonic agent or the like to a solution or suspension containing the ester of C10 fatty acid. When made into an external preparation, it is possible to make into an external preparation such as patch preparation, gel preparation, ointment, cream preparation or the like. Thus, the compound of the present invention is, for example, mixed with, melted in or emulsified in an appropriate base to make a preparation, and in the case of a patch preparation, the above is spread and applied onto a support. In the case of

a patch preparation, a gel preparation or the like, it can be made, for example, into a composition using an organogelling agent. Incidentally, depending upon the dosage form of each external preparation, a commonly used preservative, antioxidant, flavoring agent, adhesive or the like may be appropriately selected and added to a formulation.

[0042] A desired dose of the compound of the present invention may be appropriately increased or decreased by taking dose regimen, age, sex, symptom in a patient, etc. into consideration, and the compound can be usually administered in an amount of from 1 to 1,000 mg, and preferably 5 to 300 mg, for adult, at once or in several divided administrations per day.

EXAMPLES

[0043] The present invention will be hereinafter explained by means of Examples.

[0044] Example 1

Production of 1-((*E*)-2-Decenoyl)pyrrolidine [Compound 1] (Reference compound)

A tetrahydrofuran solution (20 mL) of pyrrolidine (0.71 g, 0.01 mol) containing pyridine (0.79 g, 0.01 mol) was added to a tetrahydrofuran solution (20 mL) of (*E*)-2-decenoic acid chloride (1.9 g, 0.01 mol), synthesized using (*E*)-2-decenoic acid and thionyl chloride, and the mixture was heated under refluxing in a warm water bath for 3 hours. After distilling off excess tetrahydrofuran, water was added to a liquid reaction mixture, and the liquid mixture was extracted with ethyl acetate, and washed with water, and ethyl acetate was distilled off. The residue

was purified with silica gel column chromatography (developing solvent: hexane : ethyl acetate = 1:3), to give an intended compound (1.4 g) in the form of a colorless oily substance.

Colorless oily substance, $C_{14}H_{25}NO$ MW 223, ESIMS (positive ion mode: rel. int): m/z 246.184 $[M+Na]^+$ (Calcd for $C_{14}H_{25}NONa$, 246.1834), 1H -NMR (400 MHz, $CDCl_3$) δ : 0.88 (3H, t, $J = 6.8$ Hz), 1.25_1.32 (8H, m), 1.45 (2H, m), 1.86 (2H, m), 1.96 (2H, m), 2.20 (2H, m), 3.518 (2H, t, $J = 7.2$ Hz), 3.524 (2H, t, $J = 7.2$ Hz), 6.09 (1H, d, $J = 15.4$ Hz), 6.91 (1H, dt, $J = 15.4, 6.8$ Hz).

[0045] Example 2

Production of (*S*)-1-((*E*)-2-Decenoyl)pyrrolidine-2-carboxylic acid methyl ester [Compound 3]

Triethylamine (0.84 mL, 6 mmol) and *l*-ethyl-3-(3-dimethylaminopropyl) carbodiimide hydrochloride (WSC·HCl) (1.05 g, 5.5 mmol) were added to a dichloromethane solution (50 mL) containing (*E*)-2-decenoic acid (0.92 mL, 5 mmol) and L-proline methyl ester hydrochloride (0.91 g, 5.5 mmol) at room temperature, and the mixture was stirred for 24 hours. The obtained reaction solution was washed with water and saturated sodium chloride solution, and then dried over anhydrous sodium sulfate. The residue was purified with silica gel column chromatography (hexane : ethyl acetate = 7:3), to give an intended compound (1.23 g, 87%) in the form of a colorless oily substance. Here, in a case where the starting raw material amine is not a hydrochloride, triethylamine may not be added.

Colorless oily substance, $C_{16}H_{27}NO_3$ MW 281.4, EIMS: m/z 282 $[M+H]^+$,

$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 0.86 (t, $J = 7.0$ Hz, 3H), 1.21-1.31 (m, 8H), 1.34-1.44 (m, 2H), 1.70-1.97 (m, 3H), 2.02-2.27 (m, 3H), 3.38-3.65 (m, 5H), 4.31-4.35 and 4.70-4.74 (m, 1H), 5.97 and 6.25 (m, 1H), 6.61-6.71 (m, 1H).

5 [0046] Example 3

Production of (*S*)-1-((*E*)-2-Decenoyl)pyrrolidine-2-carboxylic acid
[Compound 2]

Compound 3 (0.84 g, 3 mmol) was dissolved in methanol (40 mL), and a 1 mol/L aqueous sodium hydroxide solution (4 mL, NaOH 4 mmol) was added thereto at room temperature, and the mixture was stirred for 20 hours. The solvent was distilled off under a reduced pressure, and the residue was then dissolved in water, and a 10% aqueous citric acid solution was added thereto to adjust its pH to about 4. The liquid mixture was extracted with dichloromethane, an organic layer was washed with water and saturated sodium chloride solution, and the washed extract was then dried over anhydrous sodium sulfate, to give an intended compound (0.72 g, 90%) in the form of a colorless oily substance.

Colorless oily substance, $\text{C}_{15}\text{H}_{25}\text{NO}_3$ MW 267.4, EIMS: m/z 268 $[\text{M}+\text{H}]^+$, $^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 0.86 (t, $J = 7.0$ Hz, 3H), 1.20-1.45 (m, 10H), 1.70-2.27 (m, 6H), 3.37-3.63 (m, 2H), 4.23-4.28 and 4.55-4.59 (m, 1H), 6.25 and 5.99 (m, 1H), 6.62-6.71 (m, 1H), 12.30-12.70 (br, 1H).

[0047] Example 4

Production of 3-((*E*)-2-Decenoyl)thiazolidine [Compound 4]

The same procedures as in Example 1 were carried out using (*E*)-2-decenoic acid and thiazolidine as starting raw materials, to produce an

intended compound.

Colorless oily substance, $C_{13}H_{23}NO$ MW 241, HR-ESIMS (positive ion mode): m/z 264.1443 $[M+Na]^+$ (Calcd for $C_{13}H_{23}NOSNa$, 264.1398),

1H -NMR (400 MHz, $CDCl_3$) δ : 0.88 (3H, t, $J = 6.8$ Hz), 1.22-1.34 (8H, m), 1.46 (2H, m), 2.22 (2H, m), 3.01 (1H, t, $J = 6.1$ Hz), 3.11 (1H, t, $J = 6.1$ Hz), 3.83 (1H, t, $J = 6.1$ Hz), 3.91 (1H, brt), 4.58 (1H, s), 4.65 (1H, s), 6.11 (1H, d, $J = 15.0$ Hz), 6.96 (1H, dt, $J = 15.0, 7.3$ Hz).

[0048] Example 5

Production of 1-((*E*)-2-Decenoyl)piperidine [Compound 5] (Reference compound)

The same procedures as in Example 1 were carried out using (*E*)-2-decenoic acid and piperidine as starting raw materials, to produce an intended compound.

Colorless oily substance, $C_{15}H_{27}NO$ MW 237, EIMS: m/z 237 $[M]^+$,

1H -NMR (500 MHz, $CDCl_3$) δ : 0.88 (3H, t, $J = 7.0$ Hz), 1.22-1.34 (8H, m), 1.42-1.47 (2H, m), 1.54-1.59 (4H, m), 1.62-1.68 (2H, m), 2.16-2.21 (2H, m), 3.44-3.65 (4H, m), 6.22-6.26 (1H, m), 6.83 (1H, dt, $J = 15.1, 7.1$ Hz)

[0049] Example 6

Production of 1-((*E*)-2-Decenoyl)-4-methylpiperidine [Compound 6]

The same procedures as in Example 2 were carried out using (*E*)-2-decenoic acid and 4-methylpiperidine as starting raw materials, to produce an intended compound.

Colorless oily substance, $C_{16}H_{29}NO$ MW 251.4, EIMS: m/z 252 $[M+H]^+$,

1H -NMR (500 MHz, $DMSO-d_6$) δ : 0.93-1.11 (m, 8H), 1.20-1.32 (m, 8H), 1.35-1.44 (m, 2H), 1.54-1.67 (m, 3H), 2.12-2.19 (m, 2H), 2.51-2.60 (m,

1H), 2.92-3.11 (m, 1H), 3.96-4.15 (m, 1H), 4.32-4.41 (m, 1H), 6.43 (d, J = 15.0 Hz, 1H), 6.61 (dt, J = 15.0, 7.0 Hz, 1H).

[0050] Example 7

Production of 1-((*E*)-2-decenoyl)piperidin-4-one [Compound 7]

5 The same procedures as in Example 1 were carried out using (*E*)-2-decenoic acid and piperidin-4-one as starting raw materials, to produce an intended compound.

Pale yellow oily substance, C₁₅H₂₅NO₂ MW 251, HR-ESIMS (positive ion mode): m/z 274.1824 [M+Na]⁺ (Calcd for C₁₅H₂₅NO₂Na, 274.1783),

10 ¹H-NMR (400 MHz, CDCl₃) δ: 0.88 (3H, t, J = 6.8 Hz), 1.22-1.35 (8H, m), 1.47 (2H, m), 2.23 (2H, m), 2.51 (4H, t, J = 6.4 Hz) 3.82-3.97 (4H, m), 6.31 (1H, d, J = 15.2 Hz), 6.97 (1H, dt, J = 15.2, 7.3 Hz).

[0051] Example 8

Production of 1-((*E*)-2-Decenoyl)-4-hydroxypiperidine [Compound 8]

15 The same procedures as in Example 2 were carried out using (*E*)-2-decenoic acid and 4-hydroxypiperidine as starting raw materials, to produce an intended compound.

Colorless oily substance, C₁₅H₂₇NO₂ MW 253.4, EIMS: m/z 254 [M+H]⁺,

20 ¹H-NMR (500 MHz, DMSO-d₆) δ: 0.86 (t, J = 7.0 Hz, 3H), 1.20-1.32 (m, 10H), 1.36-1.44 (m, 2H), 1.66-1.76 (m, 2H), 2.13-2.20 (m, 2H), 3.00-3.09 (m, 1H), 3.16-3.25 (m, 1H), 3.65-3.73 (m, 1H), 3.76-3.84 (m, 1H), 3.88-3.96 (m, 1H), 4.74 (d, J = 4.1 Hz, 1H), 6.45 (d, J = 15.0 Hz, 1H), 6.63 (dt, J = 15.0, 7.0 Hz, 1H).

[0052] Example 9

25 Production of 1-((*E*)-2-Decenoyl)-4-methoxypiperidine [Compound 9]

Compound 8 (0.62 g, 2.4 mmol) was dissolved in THF (10 mL), and sodium hydride (60% in mineral oil) (0.10 g, 2.6 mmol) was added thereto under ice-cooling. After mixing for 15 minutes under ice-cooling, methyl iodide (0.16 mL, 2.6 mmol) was added thereto, and stirred at room temperature for 20 hours. Water was added to the liquid reaction mixture, the mixture was extracted with ethyl acetate, an organic layer was washed with water and saturated sodium chloride solution, and the washed extract was then dried over anhydrous sodium sulfate. The residue was purified with silica gel column chromatography (developing solvent: hexane : ethyl acetate = 3:2), to give an intended compound (0.42 g, 65%) in the form of a pale yellow oily substance.

Pale yellow oily substance, $C_{16}H_{29}NO_2$ MW 267.4, EIMS:

m/z 268 $[M+H]^+$, 1H -NMR (500 MHz, DMSO- d_6) δ : 0.86 (t, J = 7.0 Hz, 3H), 1.20-1.43 (m, 12H), 1.75-1.85 (m, 2H), 2.12-2.19 (m, 2H), 3.08-3.17 (m, 1H), 3.22-3.30 (m, 1H), 3.25 (s, 3H), 3.33-3.42 (m, 1H), 3.70-3.88 (m, 2H), 6.45 (d, J = 15.0 Hz, 1H), 6.63 (dt, J = 15.0, 7.0 Hz, 1H).

[0053] Example 10

Production of 1-((*E*)-2-Decenoyl)piperidine-4-carboxylic acid [Compound 10]

The same procedures as in Example 3 were carried out using Compound 11 as a starting raw material, to produce an intended compound. White crystals, mp 88°-89°C, $C_{16}H_{27}NO_3$ MW 281.4, EIMS: m/z 281 $[M]^+$, 1H -NMR (500 MHz, DMSO- d_6) δ : 0.86 (t, J = 7.0 Hz, 3H), 1.19-1.31 (m, 8H), 1.32-1.46 (m, 4H), 1.77-1.87 (m, 2H), 2.12-2.20 (m, 2H), 2.46-2.54 (m, 1H), 2.71-2.82 (m, 1H), 3.03-3.16 (m, 1H), 3.90-4.00 (m, 1H),

4.18-4.28 (m, 1H), 6.44 (d, $J = 15.0$ Hz, 1H), 6.63 (dt, $J = 15.0, 7.0$ Hz, 1H), 12.28 (brs, 1H).

[0054] Example 11

Production of 1-((*E*)-2-Decenoyl)piperidine-4-carboxylic acid ethyl ester
[Compound 11]

The same procedures as in Example 2 were carried out using (*E*)-2-decenoic acid and ethyl isonipecotate as starting raw materials, to produce an intended compound.

Colorless oily substance, $C_{18}H_{31}NO_3$ MW 309.4, EIMS: m/z 309 $[M]^+$,
 1H -NMR (500 MHz, DMSO- d_6) δ : 0.86 (t, $J = 7.0$ Hz, 3H), 1.18 (t, $J = 7.0$ Hz, 3H), 1.20-1.32 (m, 8H), 1.33-1.49 (m, 4H), 1.80-1.88 (m, 2H), 2.13-2.19 (m, 2H), 2.56-2.64 (m, 1H), 2.71-2.81 (m, 1H), 3.05-3.16 (m, 1H), 3.92-4.07 (m, 1H), 4.07 (q, $J = 7.0$ Hz, 2H), 4.20-4.28 (m, 1H), 6.45 (d, $J = 15.0$ Hz, 1H), 6.63 (dt, $J = 15.0, 7.0$ Hz, 1H).

[0055] Example 12

Production of 1-((*E*)-2-decenoyl)-4-dimethylaminopiperidine [Compound 12]

The same procedures as in Example 2 were carried out using (*E*)-2-decenoic acid and 4-dimethylaminopiperidine as starting raw materials, to produce an intended compound.

Pale yellow oily substance, $C_{17}H_{32}N_2O$ MW 280.5, EIMS: m/z 281 $[M+H]^+$, 1H -NMR (500 MHz, DMSO- d_6) δ : 0.86 (t, $J = 7.0$ Hz, 3H), 1.13-1.31 (m, 10H), 1.36-1.44 (m, 2H), 1.70-1.79 (m, 2H), 2.12-2.20 (m, 7H), 2.26-2.34 (m, 1H), 2.56-2.65 (m, 1H), 2.93-3.04 (m, 1H), 3.99-4.07 (m, 1H), 4.33-4.41 (m, 1H), 6.45 (d, $J = 15.0$ Hz, 1H), 6.63 (dt,

$J = 15.0, 7.0 \text{ Hz, } 1\text{H}$).

In addition, an oily product Compound 12 was dissolved in methylene chloride, and treated with hydrogen chloride-dioxane, to produce 1-((*E*)-2-decenoyl)-4-dimethylaminopiperidine hydrochloride
 5 (crystals, mp 185°-188°C).

[0056] Example 13

Production of 1-((*E*)-2-Decenoyl)-4-diethylaminopiperidine [Compound 13]

The same procedures as in Example 2 were carried out using (*E*)-2-decenoic acid and 4-diethylaminopiperidine as starting raw materials, to
 10 produce an intended compound.

Pale yellow oily substance, $\text{C}_{19}\text{H}_{36}\text{N}_2\text{O}$ MW 308.5, EIMS:

m/z 309 $[\text{M}+\text{H}]^+$, $^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 0.86 (t, $J = 7.0 \text{ Hz}$, 3H), 0.95 (t, $J = 7.0 \text{ Hz}$, 6H), 1.15-1.31 (m, 10H), 1.36-1.43 (m, 2H), 1.63-
 15 1.72 (m, 2H), 2.13-2.20 (m, 2H), 2.45 (q, $J = 7.0 \text{ Hz}$, 4H), 2.50-2.60 (m, 1H), 2.65-2.73 (m, 1H), 2.90-3.02 (m, 1H), 4.01-4.10 (m, 1H), 4.39-4.47 (m, 1H), 6.44 (d, $J = 15.0 \text{ Hz}$, 1H), 6.61 (dt, $J = 15.0, 7.0 \text{ Hz}$, 1H).

[0057] Example 14

Production of 1-((*E*)-2-Decenoyl)-4-diethylaminomethylpiperidine [Compound 14]

The same procedures as in Example 2 were carried out using (*E*)-2-decenoic acid and 4-(diethylaminomethyl)piperidine as starting raw materials, to produce an intended compound.

Pale red oily substance, $\text{C}_{20}\text{H}_{38}\text{N}_2\text{O}$ MW 322.5, EIMS: m/z 323 $[\text{M}+\text{H}]^+$,

25 $^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 0.86 (t, $J = 7.0 \text{ Hz}$, 3H), 0.93 (t,

$J = 7.0$ Hz, 6H), 0.90-0.97 (m, 2H), 1.20-1.30 (m, 8H), 1.35-1.43 (m, 2H), 1.60-1.80 (m, 3H), 2.10-2.19 (m, 4H), 2.41 (q, $J = 7.0$ Hz, 4H), 2.51-2.60 (m, 1H), 2.92-3.02 (m, 1H), 3.97-4.06 (m, 1H), 4.33-4.42 (m, 1H), 6.43 (d, $J = 15.0$ Hz, 1H), 6.61 (dt, $J = 15.0, 7.0$ Hz, 1H).

5 **[0058]** Example 15

Production of 1-((*E*)-2-Decenoyl)-4-(2-dimethylaminoethyl)piperidine
[Compound 15]

10 The same procedures as in Example 2 were carried out using (*E*)-2-decenoic acid and 4-(2-dimethylaminoethyl)piperidine as starting raw materials, to produce an intended compound.

Pale yellow oily substance, $C_{19}H_{36}N_2O$ MW 308.5, EIMS:

15 m/z 309 $[M+H]^+$, 1H -NMR (500 MHz, DMSO- d_6) δ : 0.86 (t, $J = 7.0$ Hz, 3H), 0.90-1.15 (m, 2H), 1.20-1.34 (m, 10H), 1.35-1.43 (m, 2H), 1.48-1.59 (m, 1H), 1.62-1.71 (m, 2H), 2.09 (s, 6H), 2.13-2.22 (m, 4H), 2.50-2.58 (m, 1H), 2.90-3.01 (m, 1H), 3.95-4.05 (m, 1H), 4.32-4.41 (m, 1H), 6.43 (d, $J = 15.0$ Hz, 1H), 6.60 (dt, $J = 15.0, 7.0$ Hz, 1H).

[0059] Example 16

Production of 1-((*E*)-2-Decenoyl)-4-phenyl piperidine [Compound 16]

20 The same procedures as in Example 1 were carried out using (*E*)-2-decenoic acid and 4-phenylpiperidine as starting raw materials, to produce an intended compound.

Colorless oily substance, $C_{21}H_{31}NO$ MW 313, HR-ESIMS (positive ion mode): m/z 336.2330 $[M+Na]^+$ (Calcd for $C_{21}H_{31}NONa$, 336.2303),

25 1H -NMR (400 MHz, $CDCl_3$) δ : 0.88 (3H, t, $J = 7.2$ Hz), 1.22-1.34 (8H, m), 1.46 (2H, m), 1.66 (2H, m), 1.91 (2H, brd), 2.21 (2H, m), 2.66-2.80 (2H,

m), 3.15 (1H, brt), 4.14 (1H, brd), 4.84 (1H, brd), 6.29 (1H, d, $J = 15.0$ Hz), 6.89 (1H, dt, $J = 15.0, 7.3$ Hz), 7.18-7.24 (3H, m), 7.32 (2H, t, $J = 7.4$ Hz).

[0060] Example 17

Production of 4-((*E*)-2-Decenoyl)morpholine [Compound 17] (Reference compound)

The same procedures as in Example 1 were carried out using (*E*)-2-decenoic acid and morpholine as starting raw materials, to produce an intended compound.

Colorless oily substance, $C_{14}H_{25}NO_2$ MW 239, HR-ESIMS (positive ion

mode): m/z 262.1827 $[M+Na]^+$ (Calcd for $C_{14}H_{25}NO_2Na$, 262.1783),

1H -NMR (400 MHz, $CDCl_3$) δ : 0.88 (3H, t, $J = 7.2$ Hz), 1.24-1.33 (8H, m),

1.45 (2H, m), 2.20 (2H, m), 3.57 (2H, brs), 3.69 (6H, brs), 6.20 (1H, d,

$J = 15.2$ Hz), 6.91 (1H, dt, $J = 15.2, 7.1$ Hz).

[0061] Example 18

Production of 4-((*E*)-2-decenoyl)thiomorpholine [Compound 18]

The same procedures as in Example 1 were carried out using (*E*)-2-decenoic acid and thiomorpholine as starting raw materials, to produce an intended compound.

Colorless oily substance, $C_{14}H_{25}NOS$ MW 255, HR-ESIMS (positive ion

mode): m/z 278.1575 $[M+Na]^+$ (Calcd for $C_{14}H_{25}NOSNa$, 278.1555),

1H -NMR (400 MHz, $CDCl_3$) δ : 0.88 (3H, t, $J = 7.2$ Hz), 1.23-1.33 (8H, m),

1.45 (2H, m), 2.20 (2H, m), 2.63 (4H, brs), 3.83 (2H, brs), 3.92 (2H, brs),

6.20 (1H, d, $J = 15.2$ Hz), 6.87 (1H, dt, $J = 15.2, 7.3$ Hz).

[0062] Example 19

Production of 1-((*E*)-2-Decenoyl)-4-methylpiperazine [Compound 19]

The same procedures as in Example 2 were carried out using (*E*)-2-decenoic acid and 1-methylpiperazine as starting raw materials, to produce an intended compound.

Pale yellow oily substance, C₁₅H₂₈N₂O MW 252.4, EIMS:

5 m/z 253 [M+H]⁺, ¹H-NMR (500 MHz, DMSO-d₆) δ: 0.86 (t, J = 7.0 Hz, 3H), 1.20-1.31 (m, 8H), 1.35-1.44 (m, 2H), 2.13-2.19 (m, 2H), 2.17 (s, 3H), 2.22-2.30 (m, 4H), 3.45-3.55 (m, 4H), 6.44 (dt, J_d = 14.0 Hz, J_t = 1.1 Hz, 1H), 6.65 (dt, J = 14.0, 7.2 Hz, 1H).

[0063] Example 20

10 Production of 1-((*E*)-2-Decenoyl)-4-isopropylpiperazine [Compound 20]

The same procedures as in Example 2 were carried out using (*E*)-2-decenoic acid and 1-isopropylpiperazine as starting raw materials, to produce an intended compound.

15 Pale yellow oily substance, C₁₇H₃₂N₂O MW 280.5, EIMS: m/z 280 [M]⁺, ¹H-NMR (500 MHz, DMSO-d₆) δ: 0.86 (t, J = 7.0 Hz, 3H), 0.96 (d, J = 6.5 Hz, 6H), 1.20-1.31 (m, 8H), 1.36-1.42 (m, 2H), 2.11-2.20 (m, 2H), 2.35-2.42 (m, 4H), 2.65 (qq, J = 6.5, 6.5 Hz, 1H), 3.44-3.54 (m, 4H), 6.43 (dt, J = 15.0, 1.1 Hz, 1H), 6.63 (dt, J = 15.0, 7.0 Hz, 1H).

[0064] Example 21

20 Production of 3-[4-((*E*)-2-decenoyl)piperazin-1-yl]propionic acid [Compound 21]

The same procedures as in Example 2 were carried out using (*E*)-2-decenoic acid and ethyl 3-piperazin-1-yl propionate dihydrochloride (ethyl ester of Compound 21) as starting raw materials, to produce ethyl 3-[4-
25 ((*E*)-2-decenoyl)piperazin-1-yl]propionate, and subjected to alkali

saponification, to obtain an intended compound as white crystals.

White crystals, mp 53°-55°C, C₁₇H₃₀N₂O₃ MW 310.4, EIMS:

m/z 311 [M+H]⁺, ¹H-NMR (500 MHz, DMSO-d₆) δ: 0.86 (t, J = 7.0 Hz, 3H), 1.19-1.31 (m, 8H), 1.35-1.44 (m, 2H), 2.12-2.20 (m, 2H), 2.32-
 2.43 (m, 6H), 2.55-2.61 (m, 2H), 3.43-3.57 (m, 4H), 6.45 (d, J = 15.0 Hz, 1H), 6.65 (dt, J = 15.0, 7.0 Hz, 1H), 11.70-12.70 (brs, 1H).

In addition, the low-melting point crystals Compound 21 were dissolved in methylene chloride, and treated with hydrogen chloride-dioxane, to produce 3-[4-((*E*)-2-decenoyl)piperazin-1-yl]propionic acid hydrochloride (crystals, mp 201°-203°C).

[0065] Example 22

Production of 1-((*E*)-2-Decenoyl)-4-[2-(dimethylamino)ethyl]piperazine [Compound 22]

The same procedures as in Example 1 were carried out using (*E*)-2-decenoic acid and 1-[2-(dimethylamino)ethyl]piperazine as starting raw materials, to produce an intended compound.

Oily substance, C₁₈H₃₅N₃O MW 309, HR-ESIMS (positive ion mode):

m/z 310.2868 [M+H]⁺ (calcd for C₁₈H₃₆N₃O, 310.2858),

¹H-NMR (500 MHz, CDCl₃) δ: 0.88 (3H, t, J = 7.4 Hz), 1.27_1.31 (8H, m), 1.45 (2H, br t, J = 7.5 Hz), 2.19 (2H, dt, J = 6.9, 8.0 Hz), 2.32 (6H, s), 2.47 (4H, br s), 2.53 (4H, s), 3.56 (2H, br s), 3.68 (2H, br s), 6.21 (1H, d, J = 15.2 Hz), 6.86 (1H, dt, J = 6.9, 15.2 Hz).

[0066] Example 23

Production of 4-Cyclohexyl-1-((*E*)-2-decenoyl)piperazine [Compound 23]

The same procedures as in Example 2 were carried out using (*E*)-2-

decenoic acid and 1-cyclohexylpiperazine as starting raw materials, to produce an intended compound.

White crystals, mp 33°-34°C, C₂₀H₃₆N₂O MW 320.5, EIMS: m/z 320 [M]⁺,
¹H-NMR (500 MHz, DMSO-d₆) δ: 0.85 (t, J = 7.0 Hz, 3H), 1.01-1.11 (m, 1H), 1.11-1.21 (m, 4H), 1.21-1.31 (m, 8H), 1.36-1.43 (m, 2H), 1.53-1.59 (m, 1H), 1.68-1.76 (m, 4H), 2.13-2.28 (m, 3H), 2.40-2.48 (m, 4H), 3.41-3.53 (m, 4H), 6.43 (d, J = 15.0 Hz, 1H), 6.64 (dt, J = 15.0, 7.0 Hz, 1H).

[0067] Example 24

Production of 1-((E)-2-Decenoyl)-4-(2-piperidin-1-ylethyl)piperazine [Compound 24]

The same procedures as in Example 2 were carried out using (E)-2-decenoic acid and 1-[2-(1-piperidinyl)ethyl]piperazine as starting raw materials, to produce an intended compound.

White crystals, C₂₁H₃₉N₃O MW 349.6, EIMS: m/z 350 [M+H]⁺,
¹H-NMR (500 MHz, DMSO-d₆) δ: 0.86 (t, J = 6.7 Hz, 3H), 1.20-1.31 (m, 8H), 1.32-1.42 (m, 4H), 1.43-1.50 (m, 4H), 2.11-2.20 (m, 2H), 2.27-2.43 (m, 12H), 3.42-3.53 (m, 4H), 6.43 (d, J = 15.0 Hz, 1H), 6.64 (dt, J = 15.0, 7.0 Hz, 1H).

[0068] Example 25

Production of 1-((E)-2-Decenoyl)-4-phenylpiperazine [Compound 25]

The same procedures as in Example 1 were carried out using (E)-2-decenoic acid and 1-phenylpiperazine as starting raw materials, to produce an intended compound.

Pink oily substance, C₂₀H₃₀N₂O MW 314, HR-ESIMS (positive ion mode):

m/z 337.2244 [M+Na]⁺ (Calcd for C₂₀H₃₀N₂ONa, 337.2256),
¹H-NMR (400 MHz, CDCl₃) δ: 0.88 (3H, t, J = 6.8 Hz), 1.21-1.36 (8H, m),
 1.46 (2H, m), 2.22 (2H, m), 3.18 (4H, t, J = 5.4 Hz), 3.71 (2H, brs),
 3.83 (2H, brs), 6.27 (1H, d, J = 15.6 Hz), 6.88-6.96 (4H, m), 7.26-
 7.30 (2H, m).

[0069] Example 26

Production of 4-Benzyl-1-((E)-2-decenoyl)piperazine [Compound 26]

The same procedures as in Example 2 were carried out using (E)-2-decenoic acid and 1-benzylpiperazine dihydrochloride as starting raw materials, to produce an intended compound.

White crystals, mp 62°-63°C, C₂₁H₃₂N₂O MW 328.5, EIMS:
 m/z 2329 [M+H]⁺, ¹H-NMR (500 MHz, DMSO-d₆) δ: 0.86 (t, J = 7.0 Hz, 3H), 1.19-1.31 (m, 8H), 1.35-1.42 (m, 2H), 2.11-2.19 (m, 2H), 2.28-2.38 (m, 4H), 3.48 (s, 2H), 3.48-3.57 (m, 4H), 6.43 (d, J = 15.0 Hz, 1H), 6.65 (dt, J = 15.0, 7.0 Hz, 1H), 7.22-7.36 (m, 5H).

[0070] Example 27

Production of 1-((E)-2-Decenoyl)-4-(2-phenylethyl)piperazine [Compound 27]

The same procedures as in Example 2 were carried out using (E)-2-decenoic acid and 1-(2-phenetyl)piperazine dihydrochloride as starting raw materials, to produce an intended compound.

White crystals, mp 33°-34°C, C₂₂H₃₄N₂O MW 342.5, EIMS:
 m/z 343 [M+H]⁺, ¹H-NMR (500 MHz, DMSO-d₆) δ: 0.86 (t, J = 7.0 Hz, 3H), 1.19-1.31 (m, 8H), 1.36-1.44 (m, 2H), 2.11-2.21 (m, 2H), 2.36-2.45 (m, 4H), 2.52 (dd, J = 6.5, 8.3 Hz, 2H), 2.74 (dd, J = 6.5, 8.3 Hz, 2H),

3.46-3.56 (m, 4H), 6.45 (d, $J = 15.0$ Hz, 1H), 6.66 (dt, $J = 15.0, 7.0$ Hz, 1H), 7.15-7.31 (m, 5H).

[0071] Example 28

Production of 1-((*E*)-2-Decenoyl)-4-(4-dimethylaminophenyl)piperazine
[Compound 28]

The same procedures as in Example 2 were carried out using (*E*)-2-decenoic acid and 1-(4-dimethylaminophenyl)piperazine dihydrochloride as starting raw materials, to produce an intended compound.

Pale brown crystals, mp 73°-74°C, $C_{22}H_{35}N_3O$ MW 357.5, EIMS:

m/z 357 $[M]^+$, 1H -NMR (500 MHz, DMSO- d_6) δ : 0.85 (t, $J = 7.0$ Hz, 3H), 1.20-1.31 (m, 8H), 1.36-1.46 (m, 2H), 2.15-2.21 (m, 2H), 2.78 (s, 6H), 2.87-2.96 (m, 4H), 3.55-3.74 (m, 4H), 6.49 (d, $J = 15.0$ Hz, 1H), 6.64-6.72 (m, 3H), 6.86 (d, $J = 8.7$ Hz, 2H).

[0072] Example 29

Production of 1-((*E*)-2-Decenoyl)-4-(pyridin-4-yl)piperazine [Compound 29]

The same procedures as in Example 1 were carried out using (*E*)-2-decenoic acid and 1-(4-pyridyl)piperazine as starting raw materials, to produce an intended compound.

Pale brown oily substance, $C_{19}H_{29}N_3O$ MW 315, HR-ESIMS (positive ion mode): m/z 316.2349 $[M+H]^+$ (Calcd for $C_{19}H_{30}N_3O$, 316.2383),

1H -NMR (400 MHz, $CDCl_3$) δ : 0.89 (3H, t, $J = 7.2$ Hz), 1.19-1.35 (10H, m), 1.47 (2H, m), 2.23 (2H, m), 2.49 (1H, brs), 3.39 (4H, t, $J = 5.4$ Hz), 3.70-3.88 (4H, m), 6.25 (1H, d, $J = 15.0$ Hz), 6.67 (2H, d, $J = 6.4$ Hz), 6.95 (1H, dt, $J = 15.0, 7.3$ Hz), 8.31 (2H, d, $J = 6.4$ Hz)

[0073] Example 30Production of 1-((E)-2-Decenoyl)-4-(pyridin-2-yl)piperazine [Compound 30]

The same procedures as in Example 1 were carried out using (E)-2-decenoic acid and 1-(2-pyridyl)piperazine as starting raw materials, to produce an intended compound.

Colorless crystals, mp 59°-61°C, C₁₉H₂₉N₃O MW 315,

HR-ESIMS (positive ion mode): m/z 316.2346 [M+H]⁺ (Calcd for

C₁₉H₂₉N₃O, 316.2383), ¹H-NMR (400 MHz, CDCl₃) δ: 0.88 (3H, t,

J = 7.2 Hz), 1.22-1.35 (8H, m), 1.47 (2H, m), 2.22 (2H, m), 3.53 (2H, brs),

3.60-3.71 (4H, m), 3.81 (2H, brs), 6.27 (1H, d, J = 15.2 Hz), 6.64-6.69 (2H,

t, m), 6.93 (1H, dt, J = 15.2, 7.3 Hz), 7.51 (1H, m), 8.20 (1H, m)

[0074] Example 31Production of 1-((E)-2-Decenoyl)-4-(pyrimidin-2-yl)piperazine[Compound 31]

The same procedures as in Example 1 were carried out using (E)-2-decenoic acid and 1-(2-pyrimidyl)piperazine as starting raw materials, to produce an intended compound.

Colorless crystals, mp 93°-94°C, C₁₈H₂₈N₄O MW 316,

HR-ESIMS (positive ion mode): m/z 339.2155 [M+Na]⁺ (Calcd for

C₁₈H₂₉N₄ONa, 339.2161), ¹H-NMR (400 MHz, CDCl₃) δ: 0.88 (3H, t,

J = 7.2 Hz), 1.22-1.35 (8H, m), 1.47 (2H, m), 2.23 (2H, m), 3.64 (2H, brs),

3.76 (2H, brs), 3.82-3.88 (4H, m), 6.28 (1H, d, J = 15.2 Hz), 6.54 (1H, t,

J = 4.9 Hz), 6.93 (1H, dt, J = 15.2, 7.4 Hz), 8.33 (2H, d, J = 4.9 Hz).

[0075] Example 32

Production of 1-((*E*)-2-Decenoyl)azepane [Compound 32] (Reference compound)

The same procedures as in Example 1 were carried out using (*E*)-2-decenoic acid and hexamethyleneimine as starting raw materials, to produce an intended compound.

Pale brown oily substance, C₁₆H₂₉NO MW 251, HR-ESIMS (positive ion mode): m/z 274.2145 [M+Na]⁺ (Calcd for C₁₆H₂₉NONa, 274.2147), ¹H-NMR (400 MHz, CDCl₃) δ : 0.88 (3H, t, J = 6.8 Hz), 1.24-1.33 (8H, m), 1.45 (2H, m), 1.53-1.60 (4H, m), 1.69-1.77 (4H, m), 2.20 (2H, m), 3.50 (2H, t, J = 6.0 Hz), 3.58 (2H, t, J = 6.2 Hz), 6.22 (1H, d, J = 15.6 Hz), 6.90 (1H, dt, J = 15.6, 7.4 Hz).

[0076] Example 33

Production of 1-((*E*)-2-Decenoyl)-4-methyl-[1,4]diazepane [Compound 33]

The same procedures as in Example 2 were carried out using (*E*)-2-decenoic acid and 1-methyl-1,4-diazepane as starting raw materials, to produce an intended compound.

Pale yellow oily substance, C₁₆H₃₀N₂O MW 266.4, EIMS: m/z 267 [M+H]⁺, ¹H-NMR (500 MHz, DMSO-d₆) δ : 0.86 (t, J = 7.0 Hz, 3H), 1.19-1.31 (m, 8H), 1.36-1.45 (m, 2H), 1.71-1.81 (m, 2H), 2.13-2.20 (m, 2H), 2.23 and 2.24 (s x 2, 3H), 2.39-2.55 (m, 4H), 3.46-3.58 (m, 4H), 6.34-6.40 (m, 1H), 6.61-6.69 (m, 1H).

[0077] Example 34

Production of 1-((*E*)-2-Decenoyl)-4-(2-dimethylaminoethyl)-[1,4]diazepane [Compound 34]

The same procedures as in Example 2 were carried out using (*E*)-2-decenoic acid and 1-(2-dimethylaminoethyl)-1,4-diazepane as starting raw materials, to produce an intended compound.

Pale yellow oily substance, C₁₉H₃₇N₃O MW 323.5, EIMS:

5 m/z 324 [M+H]⁺, ¹H-NMR (500 MHz, DMSO-d₆) δ: 0.86 (t, J = 7.0 Hz, 3H), 1.20-1.30 (m, 8H), 1.36-1.44 (m, 2H), 1.67-1.75 (m, 2H), 2.11 (s, 6H), 2.13-2.20 (m, 2H), 2.26-2.31 (m, 2H), 2.48-2.59 (m, 4H), 2.60-2.67 (m, 2H), 3.44-3.57 (m, 4H), 6.33-6.40 (m, 1H), 6.60-6.69 (m, 1H).

[0078] Example 35

10 Production of 2-(1-((*E*)-2-Decenoyl)-4-piperidyl)acetic acid [Compound 35]

(1) The same procedures as in Example 2 were carried out using (*E*)-2-decenoic acid and ethyl 2-(4-piperidyl)acetate hydrochloride as starting raw materials, to produce ethyl 2-(1-((*E*)-2-decenoyl)-4-piperidyl)acetate
15 (ethyl ester of Compound 35).

(2) The ethyl 2-(1-((*E*)-2-decenoyl)-4-piperidyl)acetate was subjected to alkali saponification in the same manner as in Example 3, to produce an intended compound.

White crystals, mp 65°-66°C, C₁₇H₂₉NO₃ MW 295.4, EIMS:

20 m/z 296 [M+H]⁺, ¹H-NMR (500 MHz, DMSO-d₆) δ: 0.87 (t, J = 7.1 Hz, 3H), 0.96-1.10 (m, 2H), 1.20-1.31 (m, 8H), 1.36-1.43 (m, 2H), 1.63-1.72 (m, 2H), 1.85-1.94 (m, 1H), 2.11-2.18 (m, 4H), 2.53-2.62 (m, 1H), 2.95-3.05 (m, 1H), 3.98-4.05 (m, 1H), 4.32-4.41 (m, 1H), 6.43 (d, J = 15.0 Hz, 1G), 6.61 (dt, J_d = 15.0 Hz, J_t = 7.0 Hz, 1H).

25 [0079] Example 36

Production of 3-(1-((*E*)-2-Decenoyl)-4-piperidyl)propanoic acid

[Compound 36]

(1) The same procedures as in Example 2 were carried out using (*E*)-2-decenoic acid and ethyl 3-(4-piperidyl)propionate hydrochloride as starting raw materials, to produce ethyl 3-(1-((*E*)-2-decenoyl)-4-piperidyl)propionate (ethyl ester of Compound 36).

(2) The ethyl 3-(1-((*E*)-2-decenoyl)-4-piperidyl)propionate was subjected to alkali saponification in the same manner as in Example 3, to produce an intended compound.

Colorless oily substance, $C_{18}H_{31}NO_3$ MW 309.4, EIMS: m/z 310 $[M+H]^+$, 1H -NMR (500 MHz, DMSO- d_6) δ : 0.86 (t, $J = 6.5$ Hz, 3H), 0.90-1.02 (m, 2H), 1.20-1.31 (m, 8H), 1.36-1.51 (m, 5H), 1.63-1.71 (m, 2H), 2.13-2.18 (m, 2H), 2.23 (t, $J = 7.5$ Hz, 2H), 2.50-2.58 (m, 1H), 2.90-3.00 (m, 1H), 3.98-4.06 (m, 1H), 4.35-4.43 (m, 1H), 6.43 (d, $J = 15.0$ Hz, 1H), 6.61 (dt, $J_d = 15.0$ Hz, $J_t = 7.0$ Hz, 1H), 12.03 (br, 1H).

[0080] Example 37

Production of 1-((*E*)-2-Decenoyl)-4-cyanopiperidine [Compound 37]

The same procedures as in Example 2 were carried out using (*E*)-2-decenoic acid and piperidine-4-carbonitrile as starting raw materials, to produce an intended compound.

Colorless oily substance, $C_{16}H_{26}N_2O$ MW 262.4, EIMS: m/z 263 $[M+H]^+$, 1H -NMR (500 MHz, DMSO- d_6) δ : 0.87 (t, $J = 6.5$ Hz, 3H), 1.20-1.23 (m, 8H), 1.37-1.43 (m, 2H), 1.56-1.69 (m, 2H), 1.81-1.90 (m, 2H), 2.14-2.19 (m, 2H), 3.08-3.14 (m, 1H), 3.20-3.28 (m, 1H), 3.35-3.43 (m, 1H), 3.70-3.83 (m, 2H), 6.45 (d, $J = 15.0$ Hz, 1H), 6.65 (dt, $J_d = 15.0$ Hz,

$J_t = 7.0$ Hz, 1H).

[0081] Example 38

Production of 1-((*E*)-2-Decenoyl)-4-(4-chlorophenyl)piperidine

[Compound 38]

5 The same procedures as in Example 2 were carried out using (*E*)-2-decenoic acid and 4-(4-chlorophenyl)piperidine as starting raw materials, to produce an intended compound.

Pale yellow oily substance, $C_{21}H_{30}ClNO$ MW 347.9, EIMS:

10 m/z 348 $[M+H]^+$, 1H -NMR (500 MHz, DMSO- d_6) δ : 0.86 (t, $J = 7.0$ Hz, 3H), 1.22-1.32 (m, 8H), 1.32-1.54 (m, 4H), 1.74-1.83 (m, 2H), 2.15-2.21 (m, 2H), 2.26-2.69 (m, 1H), 2.75-2.83 (m, 1H), 3.05-3.15 (m, 1H), 4.12-4.21 (m, 1H), 4.54-4.62 (m, 1H), 6.48 (d, $J = 15.0$ Hz, 1H), 6.65 (dt, $J_d = 15.0$ Hz, $J_t = 7.0$ Hz, 1H), 7.27 (d, $J = 8.5$ Hz, 2H), 7.34 (d, $J = 8.5$ Hz, 2H).

15 [0082] Example 39

Production of [4-[(*E*)-2-Decenoyl]piperazin-1-yl]acetic acid hydrochloride

[Compound 39]

20 (1) The same procedures as in Example 2 were carried out using (*E*)-2-decenoic acid and ethyl 2-(1-piperazinyl)acetate dihydrochloride as starting raw materials, to produce ethyl [4-[(*E*)-2-decenoyl]piperazin-1-yl]acetate (ethyl ester of a free form of Compound 39).

(2) The ethyl [4-[(*E*)-2-decenoyl]piperazin-1-yl]acetate was subjected to alkali saponification in the same manner as in Example 3, to produce [4-[(*E*)-2-decenoyl]piperazin-1-yl]acetic acid (a free form of Compound 39).

25 (3) The compound obtained was dissolved in methylene chloride, and

then treated with hydrogen chloride-dioxane, to produce an intended compound.

White crystals, mp 195°C (decomposition), $C_{16}H_{29}ClN_2O_3$ MW 332.9, EIMS: m/z 297 $[M+H]^+$, 1H -NMR (500 MHz, DMSO- d_6) δ : 0.87 (t, $J = 7.0$ Hz, 3H), 1.20-1.33 (m, 8H), 1.38-1.46 (m, 2H), 2.15-2.21 (m, 2H), 3.20-3.53 (m, 4H), 3.70-4.10 (m, 4H), 4.12 (s, 2H), 6.48 (d, $J = 15.0$ Hz, 6.73 (dt, $J_d = 15.0$ Hz, $J_t = 7.0$ Hz, 1H).

[0083] Example 40

Production of 4-[4-((*E*)-2-Decenoyl)piperazin-1-yl]butanoic acid hydrochloride [Compound 40]

(1) The same procedures as in Example 2 were carried out using (*E*)-2-decenoic acid and ethyl 4-(1-piperazinyl)butyrate dihydrochloride as starting raw materials, to produce ethyl 4-[4-((*E*)-2-decenoyl)piperazin-1-yl]butanoate (ethyl ester of a free form of Compound 40).

(2) The ethyl 4-[4-((*E*)-2-decenoyl)piperazin-1-yl]butanoate was subjected to alkali saponification in the same manner as in Example 3, to produce 4-[4-((*E*)-2-decenoyl)piperazin-1-yl]butanoic acid (a free form of Compound 40).

(3) The compound obtained was dissolved in methylene chloride, and then treated with hydrogen chloride-dioxane, to produce an intended compound.

White crystals, mp 203°-205°C, $C_{18}H_{33}ClN_2O_3$ MW 360.9, EIMS: m/z 325 $[M+H]^+$, 1H -NMR (500 MHz, DMSO- d_6) δ : 0.86 (t, $J = 7.0$ Hz, 3H), 1.20-1.31 (m, 8H), 1.35-1.43 (m, 2H), 1.61-1.69 (m, 2H), 2.13-2.18 (m, 2H), 2.23 (t, $J = 7.2$ Hz, 2H), 2.26-2.35 (m, 6H), 3.45-3.55 (m,

4H), 6.43 (d, $J = 15.0$ Hz, 1H), 6.64 (dt, $J_d = 15.0$ Hz, $J_t = 7.0$ Hz, 1H).

[0084] Example 41

Production of 1-((*E*)-2-Decenoyl)-4-(4-chlorophenyl)piperazine

[Compound 41]

5 The same procedures as in Example 2 were carried out using (*E*)-2-decenoic acid and 1-(4-chlorophenyl)piperazine as starting raw materials, to produce an intended compound.

Pale yellow oily substance, $C_{20}H_{29}ClN_2O$ MW 348.9, EIMS: m/z 348 $[M]^+$,
 1H -NMR (500 MHz, $DMSO-d_6$) δ : 0.86 (t, $J = 6.5$ Hz, 3H), 1.20-1.31 (m,
 10 8H), 1.37-1.45 (m, 2H), 2.15-2.22 (m, 2H), 3.07-3.18 (m, 4H), 3.60-3.70 (m, 4H), 6.51 (d, $J = 15.0$ Hz, 1H), 6.70 (dt, $J_d = 15.0$ Hz, $J_t = 7.0$ Hz, 1H), 6.90 (d, $J = 9.0$ Hz, 2H), 7.25 ($J = 9.0$ Hz, 2H).

[0085] Example 42

Production of 1-((*E*)-2-Decenoyl)-4-(4-methoxyphenyl)piperazine

[Compound 42]

15 The same procedures as in Example 2 were carried out using (*E*)-2-decenoic acid and 1-(4-methoxyphenyl)piperazine as starting raw materials, to produce an intended compound.

White crystals, mp 50° - 52° C, $C_{21}H_{32}N_2O_2$ MW 344.5, EIMS:

20 m/z 344 $[M]^+$, 1H -NMR (500 MHz, $DMSO-d_6$) δ : 0.86 (t, $J = 6.8$ Hz, 3H), 1.20-1.31 (m, 8H), 1.38-1.45 (m, 2H), 2.15-2.21 (m, 2H), 2.94-3.01 (m, 4H), 3.60-3.67 (m, 4H), 3.68 (s, 3H), 6.50 (d, $J = 15.0$ Hz, 1H), 6.68 (dt, $J_d = 15.0$ Hz, $J_t = 7.0$ Hz, 1H), 6.83 (d, $J = 9.1$ Hz, 2H), 6.91 (d, $J = 9.1$ Hz, 2H).

25 [0086] Example 43

Production of 1-((*E*)-2-Decenoyl)-4-(4-methylphenyl)piperazine

[Compound 43]

The same procedures as in Example 2 were carried out using (*E*)-2-decenoic acid and 1-(*p*-tolyl)piperazine as starting raw materials, to produce an intended compound.

White crystals, mp 45°-46°C, C₂₁H₃₂N₂O MW 328.5, EIMS: m/z 328 [M]⁺, ¹H-NMR (500 MHz, DMSO-d₆) δ: 0.85 (t, J = 6.9 Hz, 3H), 1.20-1.30 (m, 8H), 1.36-1.45 (m, 2H), 2.20 (s, 3H), 2.15-2.23 (m, 2H), 3.00-3.08 (m, 4H), 3.60-3.70 (m, 4H), 6.50 (d, J = 15.0 Hz, 1H), 6.69 (dt, J_d = 15.0 Hz, J_t = 7.0 Hz, 1H), 6.85 (d, J = 8.5 Hz, 2H), 7.03 (d, J = 8.5 Hz, 2H).

[0087] Example 44

Production of 1-((*E*)-2-Decenoyl)-4-(4-fluorophenyl)piperazine

[Compound 44]

The same procedures as in Example 2 were carried out using (*E*)-2-decenoic acid and 1-(4-fluorophenyl)piperazine as starting raw materials, to produce an intended compound.

Pale yellow oily substance, C₂₀H₂₉FN₂O MW 332.5, EIMS: m/z 332 [M]⁺, ¹H-NMR (500 MHz, DMSO-d₆) δ: 0.86 (t, J = 6.5 Hz, 3H), 1.20-1.31 (m, 8H), 1.37-1.45 (m, 2H), 2.16-2.22 (m, 2H), 3.00-3.09 (m, 4H), 3.62-3.72 (m, 4H), 6.50 (d, J = 15.0 Hz, 1H), 6.69 (dt, J_d = 15.0 Hz, J_t = 7.0 Hz, 1H), 6.95-7.00 (m, 2H), 7.03-7.09 (m, 2H).

[0088] Example 45

Production of 1-((*E*)-2-Decenoyl)-4-(2-chlorophenyl)piperazine

[Compound 45]

The same procedures as in Example 2 were carried out using (*E*)-2-

decenoic acid and 1-(2-chlorophenyl)piperazine as starting raw materials, to produce an intended compound.

Pale yellow oily substance, $C_{20}H_{29}ClN_2O$ MW 348.9, EIMS: m/z 348 $[M]^+$, 1H -NMR (500 MHz, DMSO- d_6) δ : 0.86 (t, J = 6.5 Hz, 3H), 1.21-1.31 (m, 8H), 1.38-1.45 (m, 2H), 2.15-2.21 (m, 2H), 2.90-2.98 (m, 4H), 3.65-3.75 (m, 4H), 6.50 (d, J = 15.0 Hz, 1H), 6.70 (dt, J_d = 15.0 Hz, J_t = 7.0 Hz, 1H), 7.04-7.08 (m, 1H), 7.13-7.17 (m, 1H), 7.28-7.32 (m, 1H), 7.41-7.45 (m, 1H).

[0089] Example 46

Production of 1-((*E*)-2-Decenoyl)-4-(4-hydroxyphenyl)piperazine
[Compound 46]

The same procedures as in Example 2 were carried out using (*E*)-2-decenoic acid and 4-(1-piperazinyl)phenol as starting raw materials, to produce an intended compound.

Pale red crystals, mp 85°-87°C, $C_{20}H_{30}N_2O_2$ MW 330.5, EIMS: m/z 330 $[M+H]^+$, 1H -NMR (500 MHz, DMSO- d_6) δ : 0.86 (t, J = 7.0 Hz, 3H), 1.20-1.30 (m, 8H), 1.38-1.40 (m, 2H), 2.15-2.21 (m, 2H), 2.87-2.96 (m, 4H), 3.59-3.69 (m, 4H), 6.49 (d, J = 15.0 Hz, 1H), 6.63-6.71 (m, 3H), 6.80 (d, J = 6.8 Hz, 2H), 8.88 (s, 1H).

[0090] Example 47

Production of 2-[4-[4-((*E*)-2-Decenoyl)piperazin-1-yl]phenoxy]acetic acid
[Compound 47]

(1) The same procedures as in Example 2 were carried out using (*E*)-2-decenoic acid and ethyl 2-[4-(1-piperazinyl)phenoxy]acetate dihydrochloride as starting raw materials, to produce ethyl 2-[4-[4-((*E*)-2-

decenoyl)piperazin-1-yl]phenoxy]acetate (ethyl ester of Compound 47).

(2) The ethyl 2-[4-[4-((*E*)-2-decenoyl)piperazin-1-yl]phenoxy]acetate was subjected to alkali saponification in the same manner as in Example 3, to produce an intended compound.

White crystals, mp 210°C (decomposition), C₂₂H₃₂N₂O₄ MW 388.5, EIMS: m/z 388 [M]⁺, ¹H-NMR (500 MHz, DMSO-d₆) δ: 0.86 (t, J = 7.0 Hz, 3H), 1.21-1.32 (m, 8H), 1.37-1.45 (m, 2H), 2.15-2.20 (m, 2H), 2.91-3.00 (m, 4H), 3.60-3.71 (m, 4H), 6.50 (d, J = 15.0 Hz, 1H), 6.68 (dt, J_d = 15.0 Hz, J_t = 7.0 Hz, 1H), 6.76 (d, J = 7.1 Hz, 2H), 6.87 (d, J = 7.1 Hz, 2H).

[0091] Example 48

Production of 4-[[4-((*E*)-2-Decenoyl)piperazin-1-yl]methyl]benzoic acid [Compound 48]

(1) The same procedures as in Example 2 were carried out using (*E*)-2-decenoic acid and methyl 4-[1-piperazinylmethyl]benzoate dihydrochloride as starting raw materials, to produce methyl 4-[[4-((*E*)-2-decenoyl)piperazin-1-yl]methyl]benzoate (methyl ester of Compound 48).

(2) The methyl 4-[[4-((*E*)-2-decenoyl)piperazin-1-yl]methyl]benzoate was subjected to alkali saponification in the same manner as in Example 3, to produce an intended compound.

White crystals, mp 114°-116°C, C₂₂H₃₂N₂O₃ MW 372.5, EIMS: m/z 373 [M+H]⁺, ¹H-NMR (500 MHz, DMSO-d₆) δ: 0.86 (t, J = 7.0 Hz, 3H), 1.20-1.30 (m, 8H), 1.35-1.43 (m, 2H), 2.12-2.18 (m, 2H), 2.30-2.38 (m, 4H), 3.45-3.55 (m, 4H), 3.55 (s, sH), 6.43 (d, J = 15.0 Hz, 1H), 6.64 (dt, J_d = 15.0 Hz, J_t = 7.0 Hz, 1H), 7.44 (d, J = 8.2 Hz, 2H), 12.8 (br,

1H).

[0092] Example 49

Production of 4-[(E)-2-Decenoyl]morpholine-3-carboxylic acid

[Compound 49]

(1) The same procedures as in Example 2 were carried out using (E)-2-decenoic acid and methyl morpholine-3-carboxylate hydrochloride as starting raw materials, to produce methyl 4-[(E)-2-decenoyl]morpholine-3-carboxylate (methyl ester of Compound 49).

(2) The methyl 4-[(E)-2-decenoyl]morpholine-3-carboxylate was subjected to alkali saponification in the same manner as in Example 3, to produce an intended compound.

Pale yellow oily substance, $C_{15}H_{25}NO_4$ MW 283.4, EIMS:

m/z 284 $[M+H]^+$, 1H -NMR (500 MHz, $CDCl_3$) δ : 0.88 (t, $J = 7.0$ Hz, 3H), 1.22-1.35 (m, 8H), 1.40-1.50 (m, 2H), 2.17-2.26 (m, 2H), 3.10-4.00 (m, 5H), 4.45-4.40 and 5.15-5.18 (m, 2H), 6.05 and 6.23 (d x 2, $J = 15.0$ Hz, 1H), 6.88-7.02 (m, 1H), 9.00 (br, 1H).

[0093] Test Example 1 (Evaluation for Action Against Rat Peripheral Nerve Disorder Induced by Paclitaxel)

The effects of the compounds of the present invention were evaluated for a peripheral nerve disorder which is a side effect induced by the administration of an anti-cancer agent paclitaxel, such as hyperesthesia including allodynia induced by mechanical stimuli, which is a severe pain induced by such tactile stimuli which usually cause no pains. The compound of the present invention was intraperitoneally administered to rats as a test drug to conduct a von Frey test.

[0094] (1) Preparation of Rats with Paclitaxel-Induced Peripheral Nerve Disorder and Administration of Test Drugs

Male SD rats of six weeks age (six rats per group) were used as experimental animals, and paclitaxel (2 mg/kg) was intraperitoneally administered every other day for four times in total to prepare rats with paclitaxel-induced peripheral nerve disorder. During the period of 18 to 25 days or during the period of 20 to 27 days after the initiation of paclitaxel administration, the test drug was intraperitoneally administered in a single dose of 300 μ /kg, and the following von Frey test was conducted.

[0095] (2) von Frey test

The rats of the above (1) were placed in a transparent acrylic cage with a wire-meshed floor and habituated for about three minutes, and the 50% reaction threshold values to the mechanical stimulus of right hind limb were measured before administration of the test drug and after 1 and 5 hours from the initiation of the administration.

The measurement was conducted using von Frey filaments (manufactured by North Coast Medical Inc.) in accordance with the methods of Chaplan, et al. (*Journal of Neuroscience Methods*, 53(1), 55-63, 1994) and Dixon, et al. (*Annual Review of Pharmacology and Toxicology*, 20, 441-462, 1980). Of eight filaments [stimulus loads (g): 0.4, 0.6, 1.0, 2.0, 4.0, 6.0, 8.0 and 15.0], the test was started as from the filament of 2.0 g, the filament was vertically attached to the sole for 2 to 3 seconds with such a force that the filament was lightly bent and the case where the hind limb showed an escape reaction was called a positive reaction. The case where the rat escaped at the instance of removing the filament was

also called positive. When the positive reaction was found, stimulus was applied similarly with a filament of one rank weaker, and in the absence of the reaction, stimulus was similarly applied with a filament of one rank stronger, and a point where the reaction changed from negative to positive or from positive to negative was called the first two reactions.

Subsequently, stimuli were applied for continuously four times by the same up-down method. A 50% reaction threshold value to the mechanical stimulus was measured using the reaction to the six stimuli in total and then (mean value) $\pm$ (standard error) for each group was calculated.

Incidentally, when stimulus reached to a load of 15.0 g without positive reaction, or when positive reaction continued to a load of 0.4 g, 15.0 g or 0.25 g was adopted as each of 50% reaction threshold values. With regard to the higher 50% reaction threshold value between the 50% threshold values after 1 hour and 5 hours from administration of a test drug, a recovery rate (%) of the 50% reaction threshold value was calculated by the following expression in which 15 was taken as a normal threshold value. One example of the above test results is shown in Tables 6 and 7.

Recovery rate (%) of 50% reaction threshold value = $\frac{[(50\% \text{ reaction threshold value after 1 hour or 5 hours from administration of test drug}) - (50\% \text{ reaction threshold value before administration of test drug})]}{[(\text{Normal threshold value}) - (50\% \text{ reaction threshold value before administration of test drug})]} \times 100$

[0096]

[Table 6]

Test Drug	Recovery Rate of 50% Reaction Threshold Value (%)
Compound 1 (Reference compound)	58.2
Compound 2	23.5
Compound 4	43.7
Compound 5 (Reference compound)	41.6
Compound 6	33.8
Compound 7	45.2
Compound 8	49.7
Compound 9	35.6
Compound 10	44.6
Compound 12	49.7
Compound 13	23.2
Compound 14	26.0
Compound 15	34.2
Compound 16	38.6
Compound 17 (Reference compound)	30.6
Compound 18	48.9
Compound 19	19.8
Compound 20	26.9
Compound 21	34.8
Compound 22	29.7
Compound 23	33.0
Compound 24	49.7
Compound 25	55.3
Compound 26	24.1
Compound 27	69.1
Compound 28	20.3
Compound 29	42.8
Compound 30	22.6
Compound 31	22.9
Compound 32 (Reference compound)	51.1
Compound 33	33.4
Compound 34	35.5

[0097] [Table 7]

Test Drug	Recovery Rate of 50% Reaction Threshold Value (%)
Compound 35	41.2
Compound 36	42.0
Compound 37	22.4
Compound 38	55.5
Compound 39	16.6
Compound 40	61.6
Compound 41	81.5
Compound 42	45.8
Compound 43	31.3
Compound 44	42.6
Compound 45	33.3
Compound 46	53.7
Compound 47	54.5
Compound 48	56.4

[0098] Tables 6 and 7 show that the compounds of the present invention exhibits an excellent improving effect for hyperesthesia induced by administration of paclitaxel, and thus have an alleviating action for the side effects induced by administration of an anti-cancer agent. In addition, the compounds of the present invention also show an excellent improving action to hyperesthesia when oxaliplatin of platinum drugs, is used in the same manner as that when paclitaxel of taxane drugs is used.

[0099] Test Example 2 (Evaluation of Activation (Phosphorylation) of MAP Kinase)

With regard to the compounds of the present invention, the activation of MAP kinase (ERK 1/2) was measured in accordance with a Western immunoblotting in the following manner.

Nerve cells were dispersed from cerebral cortex of a fetal rat of 17 days age, and the nerve cells were cultured for one day in a Dulbecco's modified Eagle's medium (DMEM) containing 5% fetal bovine serum. The culture medium was exchanged with a serum-free medium (B27 supplemented Neurobasal; Invitrogen), and the nerve cells were cultured at the density of 20,000 to 40,000 cells/cm² in a polyornithine-coated culture dish.

[0100] After three days, the compound of the present invention was added, and the culture was continued for 30 minutes. Thereafter, the cells were recovered on ice using a solution containing a phosphatase inhibitor where Tris-HCl buffer was used as a base. The protein concentration of the resulting cell extract was quantified with a BCA Protein Assay Kit (Takara Bio Co., Ltd.), and a certain amount of the protein (3 µg for the measurement of MAP kinase and 5 µg for the measurement of phosphorylated MAP kinase) was subjected to polyacrylamide gel electrophoresis. The protein was transferred from the electrophoresed gel to a PVDF membrane, and Western immunoblotting was carried out using each of an anti-MAP kinase antibody (Cell Signaling Technology) and an anti-phosphorylated MAP kinase antibody (Cell Signaling Technology) of the primary antibody.

[0101] Subsequently, the reaction with an alkaline phosphatase-labeled anti-rabbit IgG antibody (Promega) of the secondary antibody was carried out, and the enzymatic activity was allowed to develop color, to measure MAP kinase (MAPK) and the phosphorylated MAP kinase (pMAPK).

Incidentally, each of the compounds of the present invention was

dissolved in 0.1% DMSO to adjust its concentration to 250 µg/ml. As to the control, 0.1% DMSO was added.

[0102] The intensity of the concentration of the bands of eletrophoresed the gel was calculated and quantified using Image J (K. K. Bioarts). The numerical values for the MAPK when using the compounds of the present invention were each divided by the numerical value for the MAPK of the control, and the numerical values for the pMAPK when using the compounds of the present invention were divided by the numerical value for the pMAPK of the control, whereby a ratio of the MAPK when using each of the compounds of the present invention to the control, and a ratio of the pMAPK when using each of the compounds of the present invention to the control were obtained. Subsequently, the ratio of the pMAPK to the control obtained was divided by the ratio of the MAPK to the control to obtain a ratio of the pMAPK to the MAPK. One example of the results are shown in Tables 8 to 10.

[0103]

[Table 8]

	Ratio of pMAPK to MAPK
Control	1.00
Compound 4	1.47
Compound 7	1.62
Compound 18	1.28

[0104] [Table 9]

	Ratio of pMAPK to MAPK
Control	1.00
Compound 6	1.71
Compound 14	2.08
Compound 20	1.27
Compound 23	1.30
Compound 24	2.58

[0105] [Table 10]

	Ratio of pMAPK to MAPK
Control	1.00
Compound 2	1.28
Compound 13	1.63
Compound 21	1.54
Compound 26	1.48
Compound 27	1.44

[0106] Tables 8 to 10 show that the compounds of the present invention show high activation (phosphorylation) of the MAP kinase, as compared to the control, suggesting to have a neurotrophic factor-like activity.

[0107] Test Example 3 Evaluation of Activation (Phosphorylation) of MAP Kinase under Mild Stress Loads

As to the compounds of the present invention, the activation level of the MAP kinase (ERK1/2) was measured using chronic mild stress-induced depression model mice in the following manner.

(1) Preparation of Chronic Mild Stress-Induced Depression Model Mice and Administration of Test Drug

Female ddY mice of seven weeks age ($n = 8$ to 12) were subjected to the following procedures: (A) The mice were subjected to forced swimming for 15 minutes, and then to normal breeding for two days.

(B) The mice were subjected to breeding in an inclined cage for two days,

and then to normal breeding for one day. (C) The mice were subjected to breeding for one day in a cage of which floor mat was made wet and then to normal breeding for one day. (D) The mice were subjected to breeding for one day with a cage rotating at a rate of 180 rotations/minute, and then to normal breeding for one day. Further, after one cycle of (A) to (D) was

carried out, (B) to (D) were repeated for two cycles. As such, stress was applied over three weeks in total to prepare chronic mild stress-induced depression model mice. During that time, the compounds of the present invention each dissolved in PBS or in a PBS solvent containing DMSO, etc. were orally administered once every day over a period of three weeks.

[0108] (2) Measurements of MAP Kinase and Phosphorylated MAP Kinase

After the termination of the behavioral analyses after the above chronic mild stress loads (tail suspension test and light-darkness test), hippocampus was excised from the brain. The hippocampus and a disruption (20 mM Tris-HCl buffer (pH 7.4): RIPA buffer containing 1% Nonidet P40[registered trademark], 1% sodium deoxycholate, 2 mM EDTA, 0.1% sodium dodecyl sulfate (SDS), 0.15 M NaCl, 10 mg/mL aprotinin, 10 mg/mL leupeptin, 50 mM NaF, 1 mM sodium orthovanadate, 1 mM phenylmethanesulfonyl fluoride (PMSF)) in an amount 19 times the wet weight of the hippocampus were placed in a microtube (1.5 mL), and

the mixture was sonically disrupted. Thereafter, the microtube was allowed to stand on ice for 30 minutes, centrifuged at $1400 \times g$ for 15 minutes, to provide the supernatant as a protein extract. To the protein extract were added a sample buffer for electrophoresis (0.2 M Tris-HCl buffer (pH 7.2), 8% SDS, 40% glycerol, bromophenol blue (BPB)) in an amount 1/3 that of the protein extract, and 2-mercaptoethanol in an amount 1/10 that of the protein extract. The mixture was heat-treated at 95°C for 5 minutes, and subjected to SDS electrophoresis with 10% polyacrylamide gel.

[0109] Each of the MAP kinase (MAPK) and the phosphorylated MAP kinase (pMAPK) was subjected to immunoblot analysis, and 2 μ g or 5 μ g each of tissue-extracted proteins was subjected to electrophoresis. Thereafter, the protein was transferred from the gel to a PVDF membrane. The transferred PVDF membrane was blocked with a solution containing 5% skim milk, and allowed to react with each of 1,000-fold dilutions of an anti-MAPK antibody and an anti-pMAPK antibody overnight at 4°C. Next, the reaction mixture was allowed to react with a secondary antibody (alkali phosphatase-labeled anti-mouse IgG antibody (5,000-fold dilution)), and finally incubated in a solution supplemented with a substrate of the alkali phosphatase for several minutes to allow color development. The staining intensity of each band was quantified, and a ratio of pMAPK to MAPK was obtained ($n = 4$ to 7). Here, the significance test was conducted in accordance with Dunnett's test.

[0110] As a result of the above test, the ratio of pMAPK to MAPK of the normal animal control group was 1.26 ± 0.11 , and the same ratio of the

mild stress-loaded animal control group was significantly lowered to 0.56 ± 0.04 . The same ratios in the group administered with the compounds of the present invention were such that the group administered with Compound 10 (1,000 $\mu\text{g/kg}$) was 1.00 ± 0.04 , and that the group administered with Compound 12 (500 $\mu\text{g/kg}$) was 1.09 ± 0.04 , thereby showing that the compounds of the present invention significantly increased the phosphorylation of MAPK that was lowered in the mild stress-loaded animal control group, and suggesting to have a neurotrophic factor-like activity.

[0111] Test Example 4 Evaluation of Activation (Phosphorylation) of CREB under Mild Stress Loads

With respect to the compounds of the present invention, the activation level of transcription factor CREB (a ratio of phosphorylated CREB (pCREB) to CREB), which plays a key role in the functions of nerve cells and memorization and learning abilities, of which signal transmission pathway is located downstream of MAPK (ERK 1/2) was measured using the chronic mild stress-induced depression model mice, in the same manner as in Test Example 3 mentioned above in accordance with Western blotting ($n = 3$ to 8). However, the primary antibodies (anti-CREB antibody and anti-pCREB antibody) were allowed to react in 1,000-fold dilutions, and the secondary antibodies were allowed to react in 10,000-fold dilutions.

[0112] As a result of the above test, the ratio of pCREB to CREB of the normal animal control group was 0.95 ± 0.09 , and the same ratio of the mild stress-loaded animal control group was significantly lowered to

0.50 ± 0.07. The same ratios in the group administered with the compounds of the present invention were such that the group administered with Compound 10 (1,000 µg/kg) was 0.94 ± 0.10, and that the group administered with Compound 12 (500 µg/kg) was 1.08 ± 0.14, thereby showing that the compounds of the present invention significantly increased the ratio of pCREB that was lowered in the mild stress-loaded animal control group in the same manner as the case of the phosphorylation of MAPK, and suggesting to have a neurotrophic factor-like activity.

[0113] Test Example 5 Evaluation of Suppressive Effects of Depression Symptoms Accompanying Mild Stress Loads

The compounds of the present invention were administered as a test drug to the chronic mild stress-induced depression model mice of Test Example 3(1) mentioned above, and the model mice were then subjected to tail suspension test to evaluate suppressive effects of depression symptoms. Specifically, mice were gripped at a location 1 cm away from the tip end of the mouse tail with a hand, and kept at a height 10 cm from the floor. The observations were made for 6 minutes, and the length of the immobility time, serving as an index of depression symptoms, was measured. Here, the significance test was conducted in accordance with Dunnett's test.

[0114] As a result of the above-mentioned tail suspension test, the immobility time of the normal animal control group was 73.16 ± 9.20 seconds, and the immobility time of the mild stress-loaded animal control was significantly extended to 157.01 ± 11.97 seconds. The immobility time of the group administered with Compound 12 of the present invention

at a dose of 500 $\mu\text{g/kg}$ was 69.26 ± 16.41 seconds, thereby significantly showing suppressive effects on depression symptoms. In addition, the immobility time of the group administered with Compound 10 of the present invention at a dose of 1,000 $\mu\text{g/kg}$ was 96.03 ± 25.24 seconds, so that some suppression of depression symptoms was found.

[0115] Test Example 6 Evaluation of Suppressive Effects of Anxiety Symptoms Accompanying Mild Stress Loads

The compounds of the present invention were administered as a test drug to the chronic mild stress-induced depression model mice of Test Example 3(1) mentioned above, and the model mice were then subjected to light-dark test to evaluate suppressive effects of anxiety symptoms.

Specifically, a wooden rectangular apparatus composed of a light chamber and a dark chamber having a height of 50 cm, a length of 50 cm, a width of 25 cm was used, a mouse was placed in a dark chamber and allowed to freely search, and then the number of times invading into a light chamber, and the time staying in the light chamber were recorded for 5 minutes.

This test is a test for evaluating anxiety level of mouse utilizing its characteristic of the tendency of the mice of liking dark places, and less likely to stay in light places, but might invade into a light chamber from a searching behavior out of curiosity. Here, the significance test was carried out in accordance with Dunnett's test.

[0116] As a result of the above light-dark test, contrary to the staying time in the light chamber of the normal animal control group of 110.74 ± 5.29 seconds, the staying time in the light chamber of the mild stress-loaded control group was significantly lowered to 74.72 ± 4.09 seconds. The

staying time in the light chamber of the group administered with Compound 12 of the present invention at a dose of 500 µg/kg was 108.70 ± 12.06 seconds, showing significant suppressive effects of anxiety symptoms. In addition, the staying time in the light chamber of the group administered with Compound 10 of the present invention at a dose of 1,000 µg/kg was 107.05 ± 8.29 seconds, so that inhibition of anxiety symptoms was found.

[0117] Test Example 7 Evaluation of Analgesic Action Against Osteoarthritis Model Rats

The following experiments were conducted for examining the analgesic action of the analgesic of the present invention using the osteoarthritis (OA) rats induced by sodium monoiodoacetate (MIA) which were model animals for OA.

(1) Preparation of MIA-Induced OA Rats

A 50% reaction threshold value to the mechanical stimulus of male Wistar rats of six weeks age was measured (the measurement method will be mentioned later), and a normal control group was selected. MIA prepared with saline was administered in a single dose of 300 µg/50 µL into the right knee joint of the rats except for the normal control group, while 50 µL of saline was administered into the left knee joint, whereby MIA-induced OA rats were prepared. To the normal control group, 50 µL of saline was administered into the joints of both the knees.

[0118] (2) Grouping

With regard to the male Wistar rats of six weeks age used as experimental animals except the normal control group, their 50% reaction

threshold values to the mechanical stimulus (the measuring method will be mentioned later) and body weights were measured after 24 days from the administration of MIA mentioned in (1) and then 3 groups, i.e. a normal control group, an onset control group, and a test drug-administered group, six rats per group, were organized.

[0119] (3) Administration of Test Drug

A test drug solution (100 µg/mL) using each of the compounds of the present invention as a test drug was prepared using a phosphate buffered saline (PBS) containing 0.1 vol% dimethyl sulfoxide (DMSO).

Immediately after the grouping (after 14 days from the administration of MIA), a test drug solution was intraperitoneally administered in a single dose of 500 µg/kg to a test drug-administered group. Also, to the normal control group and the onset control group, PBS containing 0.1 vol% DMSO was intraperitoneally administered in a single dose.

[0120] (4) Results of the Measurements of 50% Reaction Threshold Values to Mechanical Stimulus (von Frey Test)

The von Frey test was conducted in the same manner as in Test Example 1 (2), and a recovery rate (%) of the 50% reaction threshold values against hyperalgesia of OA induced by the administration of MIA was calculated. One example of the results is shown in Table 11.

[0121] [Table 11]

Test Drug	Recovery Rate of 50% Reaction Threshold (%)
Compound 1 (Reference compound)	24.2
Compound 5 (Reference compound)	35.8
Compound 22	13.2
Compound 25	18.9

[0122] As is clear from Table 11, the compounds of the present invention showed suppressive effects against hyperalgesia of OA induced by the administration of MIA, thereby suggesting their usefulness as an analgesic.

5

INDUSTRIAL APPLICABILITY

[0123] As is clear from the above-mentioned pharmacological test results, it was found that the compounds of the present invention had excellent therapeutic effects against hyperalgesia against mechanical stimulus induced in rats by the administration of anti-cancer agent. Therefore, the compounds of the present invention are effective as an agent for preventing or treating peripheral nerve disorders induced by anti-cancer agents, such as such as paresthesia such as numbness of limb extremities of human and animals, and hyperalgesia such as pains, so that the compounds are very highly useful.

10

15


In addition, in the evaluations using the rat cerebral cortex-cultured nerve cells and the hippocampus of the chronic mild stress-induced depression model mice, the compounds showed excellent MAP kinase

phosphorylation action and CREB phosphorylation action (neurotrophic factor-like activity). Further, in various tests using the chronic mild stress-induced depression model mice, actions of improving depression symptoms or anxiety symptoms were shown. Therefore, it is expected that the compounds of the present invention are useful as prophylactic or therapeutic agents for dementia, Alzheimer's disease, Parkinson's disease, diabetic neuropathy, depression, glaucoma, autistic disorder spectrum, or the like, and repairing agents for spinal cord injury.

In addition, the compounds of the present invention have excellent analgesic effects and suppressive effects for hyperalgesia, in the animal experiments using MIA-induced OA rats, which are OA models. Therefore, the compounds of the present invention are highly useful as prophylactic or therapeutic agent for various pain diseases, including, for example, pains induced by OA or the like.

CLAIMS

[Claim 1] A trans-2-decenoic acid derivative represented by the following general formula (I') or a pharmaceutically acceptable salt thereof:



wherein X' is:

- (a) a 1-pyrrolidyl substituted with carboxyl or alkoxycarbonyl,
- (b) a 3-thiazolidyl,
- (c) a piperidino substituted with alkyl, oxo, hydroxy, alkoxy, carboxyl, alkoxycarbonyl, alkylamino, alkylaminoalkyl, phenyl, carboxyalkyl, alkoxycarbonylalkyl, cyano, or halogenophenyl,
- (d) a thiomorpholino,
- (e) a 1-piperazyl which may be substituted with alkyl, carboxyalkyl, alkoxycarbonylalkyl, alkylaminoalkyl, cycloalkyl, piperidinoalkyl, phenylalkyl, pyridyl, pyrimidyl, carboxyphenylalkyl or alkoxycarbonylphenylalkyl,
- (f) a 1-piperazyl substituted with a phenyl which may be substituted with alkylamino, halogen, alkoxy, alkyl, hydroxy, carboxyalkoxy or alkoxycarbonylalkoxy,
- (g) a 1,4-diazepanyl which may be substituted with alkyl or alkylaminoalkyl, or
- (h) a carboxymorpholino.

[Claim 2] The trans-2-decenoic acid derivative or a pharmaceutically

acceptable salt thereof according to claim 1, wherein X' is a 1-pyrrolidyl substituted with carboxyl or alkoxycarbonyl.

[Claim 3] The trans-2-decenoic acid derivative or a pharmaceutically acceptable salt thereof according to claim 1, wherein X' is a 3-thiazolidyl, a thiomorpholino or a carboxymorpholino.

[Claim 4] The trans-2-decenoic acid derivative or a pharmaceutically acceptable salt thereof according to claim 1, wherein X' is a piperidino substituted with alkyl, oxo, hydroxy, alkoxy, carboxyl, alkoxycarbonyl, alkylamino, alkylaminoalkyl, phenyl, carboxyalkyl, alkoxycarbonylalkyl, cyano or halogenophenyl.


[Claim 5] The trans-2-decenoic acid derivative or a pharmaceutically acceptable salt thereof according to claim 1, wherein X' is a 1-piperazyl which may be substituted with alkyl, carboxyalkyl, alkoxycarbonylalkyl, alkylaminoalkyl, cycloalkyl, piperidinoalkyl, phenylalkyl, pyridyl, pyrimidyl, carboxyphenylalkyl or alkoxycarbonylphenylalkyl.

[Claim 6] The trans-2-decenoic acid derivative or a pharmaceutically acceptable salt thereof according to claim 1, wherein X' is a 1-piperazyl substituted with a phenyl which may be substituted with alkylamino, halogen, alkoxy, alkyl, hydroxy, carboxyalkoxy or alkoxycarbonylalkoxy.

[Claim 7] The trans-2-decenoic acid derivative or a pharmaceutically acceptable salt thereof according to claim 1, wherein X' is a 1,4-diazepanyl which may be substituted with alkyl or alkylaminoalkyl.

[Claim 8] A pharmaceutical agent comprising, as an active ingredient, at least one member of a trans-2-decenoic acid derivative represented by the following general formula (I) and a pharmaceutically acceptable salt

thereof:



wherein X is:

- (a) a 1-pyrrolidyl substituted with carboxyl or alkoxycarbonyl,
- 5 (b) a 3-thiazolidyl,
- (c) a piperidino substituted with alkyl, oxo, hydroxy, alkoxy, carboxyl, alkoxycarbonyl, alkylamino, alkylaminoalkyl, phenyl, carboxyalkyl, alkoxycarbonylalkyl, cyano, or halogenophenyl,
- (d) a thiomorpholino,
- 10 (e) a 1-piperazyl which may be substituted with alkyl, carboxyalkyl, alkoxycarbonylalkyl, alkylaminoalkyl, cycloalkyl, piperidinoalkyl, phenylalkyl, pyridyl, pyrimidyl, carboxyphenylalkyl or alkoxycarbonylphenylalkyl,
- (f) a 1-piperazyl substituted with a phenyl which may be substituted
- 15 with alkylamino, halogen, alkoxy, alkyl, hydroxy, carboxyalkoxy or alkoxycarbonylalkoxy,
- (g) a 1,4-diazepanyl which may be substituted with alkyl or alkylaminoalkyl, or
- (h) a carboxymorpholino.

20 [Claim 9] The pharmaceutical agent according to claim 8, wherein the pharmaceutical agent is a prophylactic agent or therapeutic agent for peripheral nerve disorders induced by administration of a drug comprising at least one anti-cancer agent.

[Claim 10] The pharmaceutical agent according to claim 8, wherein the pharmaceutical agent is a prophylactic agent or therapeutic agent for neurodegenerative diseases.

5 [Claim 11] The pharmaceutical agent according to claim 8, wherein the pharmaceutical agent is a prophylactic agent or therapeutic agent for mental diseases.

[Claim 12] The pharmaceutical agent according to claim 11, wherein the mental diseases are depression, anxiety disorders (neuroses), or autistic disorder spectrum.

10 [Claim 13] The pharmaceutical agent according to claim 8, wherein the pharmaceutical agent is an analgesic against pain diseases.

[Claim 14] The pharmaceutical agent according to any one of claims 8 to 13, wherein the pharmaceutical agent is an injectable preparation.

15 [Claim 15] The pharmaceutical agent according to any one of claims 8 to 13, wherein the pharmaceutical agent is an oral preparation.

[Claim 16] The pharmaceutical agent according to any one of claims 8 to 13, wherein the pharmaceutical agent is an external agent.