

ANTIBODY AGAINST GLYPICAN-3 AND APPLICATION THEREOF

Technical field

The present invention relates to the field of immunotherapy or diagnosis of tumor, and
5 in particular, to antibodies that specifically recognize glypican-3 (GPC3) and uses thereof.

Background

At present, adoptive immunotherapy based on immune effector cells has achieved
some effects in some tumors, and this immunotherapy method can overcome the defects of
10 antibody treatment, however the therapeutic effects in most tumors are still unsatisfactory
[Grupp SA, et al. Adoptive cellular therapy. Curr Top Microbiol Immunol., 2011; 344:
149-72.]. In recent years, it was discovered that the recognition specificity of cytotoxic
lymphocytes (CTLs) to target cells depends on T cell receptors (TCRs), scFv of antibodies
against tumor cell associated antigens and intracellular signal-activating motifs of T
15 lymphocyte receptor CD3 ζ or Fc ϵ RI γ were fused to a chimeric antigen receptor (CAR),
and T lymphocyte was genetically modified by the chimeric antigen receptors on its surface
by means of, for example, lentivirus infection. Such CAR T lymphocytes are capable of
selectively targeting T lymphocytes to tumor cells and specifically killing the tumor in a
non-limiting manner by Major Histocompatibility Complex (MHC). CAR T lymphocyte is
20 a new immunotherapy strategy in the field of tumor immunotherapy. When designing
CAR-modified immune effector cells, especially T cells, the targeted antigen genes are in
fact a crucial choice. Given the complexity of gene expressions *in vivo* and various
uncontrollable factors, selection for suitable genes for a CAR is very difficult. Moreover,
for many tumor-specific antigens, it is difficult to find a specific molecule directed against
25 them and suitable for constructing CAR-modified immune effector cells.

Glypican-3 (GPC3, also known as DGSX, GTR2-2, MXR7, OCI-5, SDYS, SGB, SGBS
or SGBS1) is a cell surface protein belonging to heparan sulfate proteoglycan family. GPC3
gene encodes a precursor core protein of about 70-kDa which is cleaved by furin to produce
a soluble, blood-accessible, amino-terminal (N-terminal) peptide of about 40-kDa and a
30 membrane bound carboxy-terminal (C-terminal) peptide containing about 2 heparan sulfate
(HS) sugar chains of about 30-kDa. GPC3 proteins attach to the cell membrane via
glycosylphosphatidylinositol (GPI) anchors.

GPC3 is highly expressed in fetal liver, while not in normal adult liver tissue, however
its expression is restored in hepatocellular carcinoma and closely related to the development

and progression of hepatocellular carcinoma. GPC3 is not only highly detected in early stage of hepatocellular carcinoma, but the detection rate is also increased with the development of liver cancer. GPC3 expression was not detected in liver adenocarcinoma, cholangiocarcinoma, liver metastases, and 12 common solid tumors and 21 non-hepatoma cell lines. In addition, GPC3 is also expressed in tumors such as melanoma, ovarian clear cell carcinoma, yolk sac tumor and neuroblastoma. GPC3 is considered as a candidate target for tumor immunotherapy in view of the high specific expression of GPC3 in tumors such as hepatocellular carcinoma and melanoma.

Although protocols for the detection of liver cancer using anti-GPC3 antibodies and antibody-dependent (ADCC) or complement-dependent (CDC) cytotoxicity assays using anti-GPC3 antibodies have been reported, there are currently no available anti-GPC3 antibodies. Only GC33 antibody disclosed in CN200580000807.4 was in phase I of clinical research, while clinical application remains to unclear.

In addition, anti-GPC3 antibodies can be used not only as therapeutic antibodies but also as immunotherapies such as T lymphocytes for diagnosis and chimeric antigen receptor modification (CAR). Current research shows that different antibodies may have different immunogenicity, and such immunogenicity will lead to the therapeutic effects and side effects of an antibody or its derived therapeutic agent. For example, CAR T cells based on mouse anti-mesothelin antibodies can cause allergic reactions that affect their long-term survival in human body. In view of the above considerations, there is a need in the art for further optimization and preparation of new anti-GPC3 antibodies that have good tumor-killing activity and excellent clinical application prospects when applied in solid tumors, so that GPC3 antibodies can be effectively used in tumor diagnosis and treatment, and benefit patients.

Summary of the invention

It is an object of the present invention to provide anti-glypican-3 antibodies and uses thereof.

In the first aspect of the present invention, an antibody that specifically recognizes glypican-3 (GPC3) is provided, and the antibody has a light chain variable region and a heavy chain variable region, and

CDR1 of the light chain variable region has an amino acid sequence selected from a group consisting of SEQ ID NO: 16, SEQ ID NO: 67, SEQ ID NO: 69;

CDR2 of the light chain variable region has an amino acid sequence selected from a

group consisting of SEQ ID NO: 18, SEQ ID NO: 68, SEQ ID NO: 70;

CDR3 of the light chain variable region has the amino acid sequence: SEQ ID NO: 20;

CDR1 of the heavy chain variable region has an amino acid sequence selected from a group consisting of SEQ ID NO: 10, SEQ ID NO: 60, SEQ ID NO: 62, SEQ ID NO: 64;

5 CDR2 of the heavy chain variable region has an amino acid sequence selected from a group consisting of SEQ ID NO: 12, SEQ ID NO: 61, SEQ ID NO: 63, SEQ ID NO: 65, SEQ ID NO: 66, SEQ ID NO: 71;

CDR3 of the heavy chain variable region has an amino acid sequence selected from a group consisting of SEQ ID NO: 14, SEQ ID NO: 72.

10 In a preferred embodiment, the antibody comprises:

antibody (a) (P7D4), and its light chain variable region has CDR1 of SEQ ID NO: 16, CDR2 of SEQ ID NO: 18 and CDR3 of SEQ ID NO: 20, or its heavy chain variable region has CDR1 of SEQ ID NO: 10, CDR2 of SEQ ID NO: 12 and CDR3 of SEQ ID NO: 14;

15 antibody (b) (am4), and its light chain variable region has CDR1 of SEQ ID NO: 16, CDR2 of SEQ ID NO: 18 and CDR3 of SEQ ID NO: 20; or its heavy chain variable region has CDR1 of SEQ ID NO: 60, CDR2 of SEQ ID NO: 61 and CDR3 of SEQ ID NO: 14;

antibody (c) (am14), and its light chain variable region has CDR1 of SEQ ID NO: 16, CDR2 of SEQ ID NO: 18 and CDR3 of SEQ ID NO: 20, or its heavy chain variable region has CDR1 of SEQ ID NO: 62, CDR2 of SEQ ID NO: 63 and CDR3 of SEQ ID NO: 14;

20 antibody (d) (am20), and its light chain variable region has CDR1 of SEQ ID NO: 16, CDR2 of SEQ ID NO: 18 and CDR3 of SEQ ID NO: 20; or its heavy chain variable region has CDR1 of SEQ ID NO: 64, CDR2 of SEQ ID NO: 65 and CDR3 of SEQ ID NO: 14;

antibody (e) (am35), and its light chain variable region has CDR1 of SEQ ID NO: 67, CDR2 of SEQ ID NO: 68 and CDR3 of SEQ ID NO: 20; or its heavy chain variable region has CDR1 of SEQ ID NO: 10, CDR2 of SEQ ID NO: 66 and CDR3 of SEQ ID NO: 14;

25 antibody (f) (am42), and its light chain variable region has CDR1 of SEQ ID NO: 69, CDR2 of SEQ ID NO: 70 and CDR3 of SEQ ID NO: 20, or its heavy chain variable region has CDR1 of SEQ ID NO: 10, CDR2 of SEQ ID NO: 66 and CDR3 of SEQ ID NO: 14;

30 antibody (g) (T2-23), and its light chain variable region has CDR1 of SEQ ID NO: 16, CDR2 of SEQ ID NO: 70 and CDR3 of SEQ ID NO: 20, or its heavy chain variable region has CDR1 of SEQ ID NO: 10, CDR2 of SEQ ID NO: 71 and CDR3 of SEQ ID NO: 72;

antibody (h), which recognizes the same antigenic determinant as that recognized by the antibody according to any one of (a) to (g).

In another preferred embodiment, the antibody specifically recognizing glypican-3

(GPC3) may be single chain antibody (scFV), monoclonal antibody, domain antibody, Fab fragment, Fd fragment, Fv fragment, F (ab')₂ fragment and a derivative thereof, or other forms of antibody; preferably single chain antibody.

In another preferred embodiment, the antibody that specifically recognizes glypican-3 (GPC3) is humanized or fully fully humanized; preferably, fully fully humanized.

In another preferred embodiment, the amino acid sequence of the heavy chain variable region of antibody (a) (P7D4) is shown in positions 1-121 of SEQ ID NO: 4; or the amino acid sequence of the light chain variable region of the antibody is shown in positions 137-247 of SEQ ID NO: 4;

The amino acid sequence of the heavy chain variable region of antibody (b) (am4) is shown in positions 1-121 of SEQ ID NO: 25; or the amino acid sequence of the light chain variable region of the antibody is shown in positions 137-247 of SEQ ID NO: 25;

The amino acid sequence of the heavy chain variable region of antibody (c) (am14) is shown in positions 1-121 of SEQ ID NO: 27; or the amino acid sequence of the light chain variable region of the antibody is shown in positions 137-247 of SEQ ID NO: 27;

The amino acid sequence of the heavy chain variable region of the antibody (d) (am20) is shown in positions 1-121 of SEQ ID NO: 29; or the amino acid sequence of the light chain variable region of the antibody is shown in positions 137-247 of SEQ ID NO: 29;

The amino acid sequence of the heavy chain variable region of the antibody (e) (am35) is shown in positions 1-121 of SEQ ID NO: 31; or the amino acid sequence of the light chain variable region of the antibody is shown in positions 137-247 of SEQ ID NO: 31;

The amino acid sequence of the heavy chain variable region of the antibody (f) (am42) is shown in positions 1-121 of SEQ ID NO: 33; or the amino acid sequence of the light chain variable region of the antibody is shown in positions 137-247 of SEQ ID NO: 33; or

The amino acid sequence of the heavy chain variable region of antibody (g) (T2-23) is shown in positions 1-121 of SEQ ID NO: 35; or the amino acid sequence of the light chain variable region of the antibody is shown in positions 137-247 of SEQ ID NO: 35.

In another aspect of the present invention, an antibody that specifically recognizes glypican-3 (GPC3) is provided, which is a humanized monoclonal antibody comprising a heavy chain variable region and a light chain variable region. Its heavy chain variable region comprises: CDR1 of the amino acid sequence of SEQ ID NO: 73, CDR2 of the amino acid sequence of SEQ ID NO: 74 and CDR3 of the amino acid sequence of SEQ ID NO: 75; or its light chain variable region comprises CDR1 of the amino acid sequence of SEQ ID NO: 76, CDR2 of the amino acid sequence of SEQ ID NO: 77 and CDR3 of the amino acid sequence

of SEQ ID NO: 78.

In a preferred embodiment, the antibody has a heavy chain variable region of the amino acid sequence of SEQ ID NO: 58 or a light chain variable region of the amino acid sequence of SEQ ID NO: 59.

5 In another aspect of the invention, a nucleic acid encoding any one of the preceding antibodies is provided.

In another aspect of the present invention, an expression vector comprising the nucleic acid is provided.

10 In another aspect of the present invention, a host cell comprising the expression vector or having the nucleic acid integrated into its genome is provided.

In another aspect of the invention, use of any one of the antibodies described above is provided for the preparation of a targeted drug, antibody-drug conjugate, or a polyfunctional antibody that specifically targets tumor cells expressing glypican-3; or for the preparation of a reagent for diagnosing a tumor expressing Glypican-3; or for the preparation of chimeric
15 antigen receptor-modified immune cells.

In another aspect of the present invention, a multi-functional immunoconjugate is provided, comprising: any one of the antibodies described above; and a functional molecule linked thereto (including covalently linked, conjugated, attached, adsorbed); and the functional molecule is selected from a group consisting of a molecule targeting a tumor
20 surface marker, a tumor-suppressing molecule, a molecule targeting a surface marker of an immune cell, or a detectable label.

In a preferred embodiment, in the multifunctional immunoconjugate, the molecule that targets a tumor surface marker is an antibody or ligand that binds to a tumor surface marker; or the tumor-suppressing molecule is an anti-tumor cytokine or anti-tumor toxin; and
25 preferably, the cytokine includes, but are not limited to: IL-12, IL-15, IFN-beta and TNF-alpha.

In another preferred embodiment, the detectable label in the multi-functional immunoconjugate includes a fluorescent label and a chromogenic label.

30 In another preferred embodiment, in the multifunctional immunoconjugate, the antibody that binds to a tumor surface marker refers to an antibody that recognizes antigens other than glypican-3, wherein the other antigens include: EGFR, EGFRvIII, mesothelin, HER2, EphA2, Her3, EpCAM, MUC1, MUC16, CEA, Claudin 18.2, folate receptor, Claudin 6, WT1, NY-ESO- 1, MAGE 3, ASGPR1 or CDH16.

In another preferred embodiment, in the multifunctional immunoconjugate, the

molecule that targets the surface marker of the immune cell is an antibody that binds to T cell surface marker, which can form a T-cell-engaging bifunctional antibody with any one of the above described antibodies (bispecific T cell engager, BiTE).

5 In another preferred embodiment, in the multifunctional immunoconjugate, the antibody that binds to the immune cell surface marker is an anti-CD3 antibody.

In another preferred embodiment, the anti-CD3 antibody is a single chain antibody (scFV), a monoclonal antibody, a Fab fragment, an Fd fragment, an Fv fragment, an F(ab')₂ fragment and a derivative thereof, or an antibody of other form; preferably single chain antibody.

10 In another preferred embodiment, the anti-CD3 antibody is humanized, chimeric, fully human or murine.

In another preferred embodiment, the multifunctional immunoconjugate is a fusion peptide, and further comprises a linker peptide (linker) between any one of the above described antibodies and the functional molecule linked thereto.

15 In another preferred embodiment, the linker peptide has the sequence (GlyGlyGlyGlySer)_n, wherein n is an integer from 1 to 5; more preferably, n = 3.

In another preferred embodiment, the multi-functional immunoconjugate is administered in a form of polypeptide or in the manner of gene administration.

In another aspect of the invention, a nucleic acid encoding any one of the multi-functional immunoconjugates described above is provided.

20 In another aspect of the present invention, use of any one of the above described multi-functional immunoconjugates is provided, for the preparation of an antineoplastic agent or an agent for diagnosis of tumors that express Glypican-3; or for the preparation of chimeric antigen receptor-modified immune cells. Preferably, the immune cells include T lymphocyte, NK cell or NKT lymphocyte.

25 In another aspect of the present invention, a chimeric antigen receptor (CAR) comprising any one of the above described antibodies is provided, and the chimeric antigen receptor comprises: any one of the above described antibodies, a transmembrane region and an intracellular signal region, which are sequentially linked; and the intracellular signal region is selected from a group consisting of intracellular signal region sequences of CD3ζ, FcεRIγ, CD27, CD28, CD137, CD134, MyD88, CD40 or a combination thereof.
30

In another preferred embodiment, the transmembrane region in the chimeric antigen receptor comprises a transmembrane region of CD8 or CD28.

In another preferred embodiment, in the chimeric antigen receptor, the chimeric antigen receptor comprises the following sequentially connected antibody, a transmembrane region

and an intracellular signal region: any one of the above described antibodies, CD8, and CD3 ζ ; any one of the above described antibodies, CD8, CD137 and CD3 ζ ; any one of the above described antibodies, the transmembrane region of CD28 molecule, the intracellular signaling region of CD28 molecule and CD3 ζ ; or any one of the above described antibodies, the transmembrane region of CD28 molecule, the intracellular signaling region of CD28 molecule, CD137 and CD3 ζ .

In another preferred embodiment, in the chimeric antigen receptor, the antibody is a single chain antibody or a domain antibody. In another preferred embodiment, the chimeric antigen receptor comprises:

- 10 SEQ ID NO: 49 or the amino acid sequence shown in positions 22-346 thereof;
- SEQ ID NO: 50 or the amino acid sequence shown in positions 22-447 thereof;
- SEQ ID NO: 51 or the amino acid sequence shown in positions 22-491 thereof;
- SEQ ID NO: 52 or the amino acid sequence shown in positions 22-494 thereof;
- SEQ ID NO: 53 or the amino acid sequence shown in positions 22-536 thereof;
- 15 The amino acid sequence of SEQ ID NO: 85;
- The amino acid sequence of SEQ ID NO: 86; or
- The amino acid sequence of SEQ ID NO: 87.

In another aspect of the invention, a nucleic acid encoding the chimeric antigen receptor is provided. In another preferred embodiment, the nucleic acid encoding the chimeric antigen receptor comprises:

- 20 SEQ ID NO: 44 or the nucleotide sequence shown in positions 380-1420 or 443-1420;
- SEQ ID NO: 45 or the nucleotide sequence shown in positions 380-1723 or 443-1723;
- SEQ ID NO: 46 or the nucleotide sequence shown in positions 380-1855 or 443-1855;
- SEQ ID NO: 47 or the nucleotide sequence shown in positions 380-1864 or 443-1864;
- 25 SEQ ID NO: 48 or the nucleotide sequence shown in positions 380-1990 or 443-1990;
- The nucleotide sequence of SEQ ID NO: 85;
- The nucleotide sequence of SEQ ID NO: 87; or
- The nucleotide sequence of SEQ ID NO: 89.

In another aspect of the present invention, an expression vector comprising a nucleic acid encoding the chimeric antigen receptor is provided. In another preferred embodiment, the expression vector is derived from lentiviral plasmid pWPT (or pWPT-eGFP).

In another aspect of the present invention, a virus comprising said vector is provided.

Use of the chimeric antigen receptor, or a encoding nucleic acid thereof, or a expression vector or virus comprising the nucleic acid for the preparation of genetically

modified immune cells targeting the tumor that expresses Glypican-3 is provided.

In another preferred embodiment, the tumor expressing glypican-3 includes (but is not limited to) liver cancer, melanoma, ovarian clear cell carcinoma, yolk sac tumor and neuroblastoma.

5 In another aspect of the present invention, a genetically modified immune cell is provided, which is transduced with a nucleic acid encoding the chimeric antigen receptor or an expression vector or a virus containing the nucleic acid; or has the chimeric antigen receptor expressed on its surface.

10 In another preferred embodiment, the immune cells further carry an exogenous encoding sequence for cytokines; and preferably, the cytokines include IL-12, IL-15 or IL-21.

In another preferred embodiment, the immune cell also expresses another chimeric antigen receptor which does not contain CD3 ζ but contains the intracellular signaling domain of CD28, the intracellular signaling domain of CD137, or a combination of both.

15 In another preferred embodiment, the immune cell further expresses a chemokine receptor; and preferably, the chemokine receptor includes: CCR2.

In another preferred embodiment, the immune cell further expresses siRNA which can reduce expression of PD-1 or a protein which blocks PD-L1.

20 In another preferred embodiment, the immune cell further expresses a safety switch; and preferably, the safety switch includes iCaspase-9, Truncated EGFR or RQR8.

In another preferred embodiment, the immune cells include T lymphocytes, NK cells or NKT cells.

25 In another aspect of the invention, use of the genetically modified immune cells is provided for the preparation of a tumor-inhibiting drug, and the tumor is a tumor expressing glypican-3.

In another aspect of the invention, a pharmaceutical composition (including a medicament or diagnostic reagent) is provided, comprising:

any one of the above described antibodies or a nucleic acid encoding the antibody; or

30 any one of the above described immunoconjugates or a nucleic acid encoding the conjugate; or

any one of the above described chimeric antigen receptors or a nucleic acid encoding the chimeric antigen receptor; or

any one of the above described genetically modified immune cells.

Other aspects of the invention will be apparent to those skilled in the art from the

disclosure herein.

Description of drawings

Figure 1 shows binding of antibodies P1B12E and P7D4 to human GPC3 and control BSA in a single phage ELISA assay. The values for antibodies P1B12E and P7D4 to human GPC3 and negative control BSA demonstrate that the two selected antibodies can specifically bind to human GPC3.

Figure 2 shows purification electrophoresis of antibodies to human GPC3.

Figure 3 shows binding activity analysis of antibodies scFv-P1B12E-Fc and scFv-P7D4-Fc to GPC3-expression positive HepG2 cells.

Figure 4 shows SPR analysis of binding ability for P7D4 series antibodies to GPC3.

Figure 5 shows P7D4 series antibodies specifically binding to recombinant human GPC3.

Figure 6 shows SDS PAGE electrophoresis of expressed and purified GPC3 protein.

Figure 7 shows kinetic analysis of binding of antibody Y035 to human GPC3.

Figure 8 shows kinetic analysis of binding of antibody 5A5 to human GPC3.

Figure 9 shows antibody Y035 specifically binding to GPC3-expression positive HepG cells.

Modes for carrying out the invention

After intensive research and screening, the inventors obtained antibodies that specifically recognize GPC3, including single chain antibodies and humanized antibodies. The antibody of the present invention can be used to prepare various targeting-antitumor drugs and drugs for diagnosing tumors.

Anti-GPC3 antibody

Specific antibodies with good binding properties to GPC3 were screened and obtained in all-human natural antibody libraries by the present inventors, and further subjected to amino acid mutations to obtain anti-GPC3 antibodies with significantly increased affinity, and key CDR regions for them to exert their binding properties were also found by the inventors.

The present inventors also obtained a mouse antibody against GPC3 using hybridoma technique, humanized it, and obtained a humanized anti-GPC3 antibody through repeated comparison with extremely excellent binding property to GPC3 and key

CDR regions for its binding performance were also found.

Antibodies of the invention may be intact immunoglobulin molecules or antigen-binding fragments including but not limited to Fab fragments, Fd fragments, Fv fragments, F(ab')₂ fragments, complementarity determining region (CDR) fragments, single chain antibody (scFv), domain antibody, bivalent single chain antibody, single chain phage antibody, bispecific diabody, triple chain antibody, quadruple chain antibody.

The antigen-binding properties of an antibody can be described by three specific regions located in variable regions of the heavy and light chains, termed complementarity determining regions (CDRs), which divide the variable regions into four framework regions (FR), and the amino acid sequences of four FRs are relatively conservative, not directly involved in binding reaction. These CDRs form a loop structure, in which β -folds formed by the FRs are located close to each other in space and the antigen binding site of the antibody is constituted by CDRs on the heavy chain and CDRs on the corresponding light chain. It is possible to determine which amino acids make up FR or CDR regions by comparing the amino acid sequences of the same type of antibody. The CDR regions are sequences of immunologically interesting proteins and the CDR regions of the antibodies of the invention are brand new. The antibody may comprise two, three, four, five, or all six of the CDR regions disclosed herein.

Another aspect of the invention includes functional variants of the antibodies described herein. If the variant is capable of competing with the parental antibody for specific binding to GPC3 and its ability to recognize GPC3 expressed on the surface of tumor cells is close to that of the specific antibodies provided in Examples of the present invention. The functional variants may have conservative sequence modifications, including nucleotide and amino acid substitutions, additions and deletions. These modifications can be introduced by standard techniques known in the art, such as directed mutagenesis and random PCR-mediated mutagenesis, and can include both natural and non-natural nucleotides and amino acids. Preferably, modification of the sequence occurs on a region outside the CDR region of the antibody.

Immunoconjugate

In the present invention, a multifunctional immunoconjugate is also provided, comprising the antibodies described herein and further comprising at least one functional molecule of other type. The functional molecule is selected from, but not limited to, a

molecule that targets a tumor surface marker, a tumor-suppressing molecule, a molecule that targets a surface marker of an immune cell, or a detectable label. The antibody and the functional molecule may form a conjugate by covalent attachment, coupling, attachment, cross-linking, or the like.

5 As a preferred mode, the immunoconjugate may comprise an antibody of the invention and at least one molecule that targets a tumor surface marker or a tumor-suppressing molecule. The tumor-suppressing molecule may be anti-tumor cytokines or anti-tumor toxins. Preferably, the cytokines include but are not limited to IL-12, IL-15, IFN-beta, TNF-alpha. The molecules that target tumor surface markers, for example, can act synergistically with
10 the antibodies of the invention to more precisely target tumor cells.

As a preferred mode, the immunoconjugate may comprise an antibody of the present invention and a detectable label. Such detectable labels include, but are not limited to, fluorescent labels, chromogenic labels such as enzymes, prosthetic groups, fluorescent
15 materials, luminescent materials, bioluminescent materials, radioactive materials, positron-emitting metals and non-radioactive paramagnetic metal ion. More than one marker can also be included. The label used to label the antibody for the purpose of detection and / or analysis and / or diagnosis depends on the used particular detection / analysis / diagnosis technique and / or method, eg, immunohistochemical staining (tissue) samples, flow
20 cytometry, and the like. Suitable labels for detection / analysis / diagnosis techniques and / or methods known in the art are well known to those skilled in the art.

As a preferred mode, the immunoconjugate may comprise an antibody of the invention as well as a molecule that targets a surface marker of an immune cell. The molecule that targets surface markers of immune cells can recognize immune cells and carry the antibodies
25 of the invention to the immune cells, so that the antibodies of the invention can target the immune cells to the tumor cells and thus trigger immunocyte for specifically killing tumor.

As a means of chemically generating an immunoconjugate by conjugation, either directly or indirectly (eg, by a linker), the immunoconjugate can be produced as a fusion protein comprising an antibody of the invention and other suitable proteins. The fusion
30 protein can be produced by a method known in the art, for example recombinantly produced by constructing and subsequently expressing the nucleic acid molecule which comprises the nucleotide sequence encoding the antibody in frame with a nucleotide sequence encoding a suitable label.

In another aspect of the invention, a nucleic acid molecule encoding at least one

antibody of the invention, a functional variant, or an immunoconjugate thereof is provided. Once obtaining the relevant sequence, the recombination method can be used to obtain the relevant sequence in large quantities. This is usually done by cloning it into a vector, transferring it to a cell, and then isolating the relevant sequence from the proliferating host cells by conventional methods.

The present invention also relates to vectors comprising the appropriate DNA sequences described above as well as appropriate promoters or control sequences. These vectors can be used to transform an appropriate host cell to enable expression of the protein. The host cell may be a prokaryotic cell, such as a bacterial cell; or a lower eukaryotic cell, such as a yeast cell; or a higher eukaryotic cell, such as a mammalian cell.

Chimeric antigen receptor and genetically modified immune cell

In the present invention, a chimeric antigen receptor expressed on the surface of an immune effector cell (immune cell) is provided, wherein the chimeric antigen receptor comprises sequentially linked: extracellular binding region, transmembrane region and intracellular signal region, and the extracellular binding region comprises the antibody of the invention. By expressing the chimeric antigen receptor on the surface of immune effector cells, immune effector cells can have a highly specific cytotoxic effect on tumor cells that express GPC3.

As used herein, "immune cells" and "immune effector cells" are used interchangeably and include: or NKT cells, and the like; and preferably, NK cells and T lymphocytes.

As a preferred embodiment of the present invention, the antibody contained in the chimeric antigen receptor is a single chain antibody, which is connected to CD8 or the transmembrane region of CD28 through the hinge region of CD8, and the transmembrane region is immediately followed by the intracellular signal region.

The invention also includes nucleic acids encoding the chimeric antigen receptors. The present invention also relates to variants of the above described polynucleotides, which encode a polypeptide, or a fragment, analog and derivative of the polypeptide having the same amino acid sequence as the present invention.

The transmembrane region of the chimeric antigen receptor may be selected from the transmembrane region of a protein such as CD8 or CD28. The human CD8 protein is a heterodimer composed of two chains, $\alpha\beta$ or $\gamma\delta$. In one embodiment of the invention, the transmembrane region is selected from the transmembrane region of CD8 α or CD28. In addition, the CD8 α hinge is a flexible region so that CD8 or CD28 and the transmembrane region as well as the hinge region are used to connect the target recognition domain scFv of the chimeric antigen receptor CAR to the

intracellular signal region.

The intracellular signal region may be selected from a group consisting of intracellular signal region of CD3 ζ , Fc ϵ RI γ , CD27, CD28, CD137, CD134, MyD88, CD4 protein, and combinations thereof. The CD3 molecule consists of five subunits, in which CD3 ζ subunit (also known as CD3 zeta, abbreviated as Z) contains 3 ITAM motifs that are important signal transduction regions in TCR-CD3 complex. CD3 δ Z is a truncated CD3 ζ sequence without ITAM motif and is generally constructed in the present invention as a negative control. Fc ϵ RI γ is mainly distributed on the surface of mast cells and basophils, which contains an ITAM motif, which is similar to CD3 ζ in structure, distribution and function. In addition, as mentioned above, CD28, CD137 and CD134 are co-stimulatory signaling molecules. The co-stimulatory effect of their intracellular signaling segments upon binding to the respective ligands results in the continued proliferation of immune effector cells, primarily T lymphocytes, and increase in the level of cytokines such as IL-2 and IFN- γ secreted by immune effector cells, and the survival period and anti-tumor effect of CAR immune effector cells *in vivo* are increased.

The chimeric antigen receptor of the present invention can be sequentially linked as follows:

The antibody of the invention, CD8 and CD3 ζ ;

The antibody of the invention, CD8, CD137 and CD3 ζ ;

The antibody of the invention, the transmembrane region of CD28 molecule, the intracellular signal region of CD28 molecule and CD3 ζ ; or

The antibodies of the invention, the transmembrane region of CD28 molecule, the intracellular signal region of CD28 molecule, CD137 and CD3 ζ .

And combinations thereof, wherein CD28a in the relevant chimeric antigen receptor protein represents the transmembrane region of CD28 molecule and CD28b represents the intracellular signal region of CD28 molecule. The various chimeric antigen receptors described above are collectively referred to as scFv (GPC3)-CAR.

The present invention also provides a vector comprising the above-mentioned nucleic acid encoding a chimeric antigen receptor protein expressed on the surface of an immune effector cell. In a specific embodiment, the vector used in the present invention is a lentiviral plasmid vector pWPT-eGFP. This plasmid belongs to the third generation of self-inactivating lentiviral vector system. The system has three plasmids, packaging plasmid psPAX2 encoding protein Gag / Pol, encoding Rev protein; envelope plasmid PMD2.G encoding VSV-G protein; and empty vector pWPT-eGFP, which can be used for recombinant introduction of a nucleic acid sequence of interest, i.e., a nucleic acid encoding CAR. In the empty vector pWPT-eGFP, the expression of enhanced

green fluorescent protein (eGFP) is regulated by elongation factor-1 α (EF-1 α) promoter. While in the recombinant expression vector pWPT-eGFP-F2A-CAR containing the nucleic acid sequence encoding CAR, co-expression of eGFP and CAR is achieved by ribosomal skipping sequence 2A (abbreviated as F2A) from food-and-mouth disease virus (FMDV).

5 The invention also includes viruses comprising the vectors described above. The viruses of the invention include packaged infectious viruses as well as viruses to be packaged that contain the necessary components for packaging into infectious viruses. Other viruses known in the art that can be used to transduce exogenous genes into immune effector cells and their corresponding plasmid vectors are also useful in the present invention.

10 The present invention further includes a genetically modified T lymphocyte, which is transduced with a nucleic acid of the present invention or transduced with the above-mentioned recombinant plasmid containing the nucleic acid of the present invention or a viral system containing the plasmid. Conventional nucleic acid transduction methods in the art, including non-viral and viral transduction methods, can be used in the present invention. Non-viral
15 transduction methods include electroporation and transposon methods. Recently, nucleofector nuclear transfection instrument developed by Amaxa can directly introduce foreign genes into nucleus to achieve highly efficient transduction of target genes. In addition, compared with conventional electroporation, the transduction efficiency of transposon system based on Sleeping Beauty system or PiggyBac transposon was significantly improved. The combination of
20 nucleofector transfection instrument and SB Sleeping Beauty transposon system has been reported [Davies JK., et al. Combining CD19 redirection and alloanergization to generate tumor-specific human T cells for allogeneic cell therapy of B-cell malignancies. Cancer Res, 2010, 70(10): OF1-10.], and high transduction efficiency and site-directed integration of target genes can be achieved by this method. In one embodiment of the invention, the transduction method of a T
25 lymphocyte modified by a chimeric antigen receptor gene is a transduction method based on a virus such as a retrovirus or a lentivirus. The method has the advantages of high transduction efficiency and stable expression of exogenous gene, and the time for in vitro culturing T lymphocytes to clinical level can be shorten. The transduced nucleic acid is expressed on the surface of the transgenic T lymphocytes by transcription, translation. In vitro cytotoxicity assay
30 performed on various cultured tumor cells demonstrated that the immune effector cells of the present invention have highly specific tumor cell killing effects (also known as cytotoxicity). Therefore, the nucleic acid encoding a chimeric antigen receptor protein of the present invention, a plasmid comprising the nucleic acid, a virus comprising the plasmid, and a transgenic immune effector cells transfected with the nucleic acid, plasmid or virus described above can be effectively

used in tumor immunotherapy.

The immune cells of the present invention may also carry exogenous encoding sequences for cytokines, including but not limited to IL-12, IL-15 or IL-21. These cytokines have immunomodulatory or antitumor activity, enhance the function of effector T cells and activated NK cells, or directly exert anti-tumor effects. Therefore, those skilled in the art will understand that the use of these cytokines will help the immune cells to function better.

In addition to the chimeric antigen receptor described above, the immune cells of the present invention may also express another chimeric antigen receptor, which does not contain CD3 ζ , but contains intracellular signaling domain of CD28 and intracellular signal domain of CD137, or a combination of both.

The immune cells of the present invention may also express chemokine receptors; the chemokine receptors include, but are not limited to, CCR2. A skilled person will understand that the CCR2 chemokine receptor can competitively bind CCR2 in the body and is beneficial for blocking the metastasis of the tumor.

The immune cells of the present invention may also express siRNAs that can reduce PD-1 expression or PD-L1-blocking proteins. A skilled person will understand that competitive blocking of the interaction between PD-L1 and its receptor PD-1 will facilitate the recovery of anti-tumor T-cell responses, thereby inhibiting tumor growth.

The immune cells of the present invention may also express a safety switch; preferably, the safety switch includes iCaspase-9, Truncated EGFR or RQR8.

Pharmaceutical composition

The antibodies, immunoconjugates comprising the antibodies, and genetically modified immune cells of the present invention can be used in the preparation of a pharmaceutical composition or diagnostic reagent. In addition to an effective amount of the antibody, immunological conjugate, or immune cell, the composition may further comprise a pharmaceutically acceptable carrier. The term "pharmaceutically acceptable" means that when the molecular entities and compositions are properly administered to animals or humans, they do not cause adverse, allergic or other untoward reactions.

Specific examples of some of the substances which may be used as pharmaceutically acceptable carriers or components thereof are sugars, such as lactose, dextrose and sucrose; starches, such as corn starch and potato starch; cellulose and its derivatives, such as carboxymethylcellulose sodium, ethylcellulose and methylcellulose; gum tragacanth; malt; gelatin; talc; solid lubricants such as stearic acid and magnesium stearate; calcium sulfate; vegetable oils, such as peanut oil, cottonseed oil, sesame oil, olive oil, corn oil and cocoa butter; polyhydric

alcohols such as propylene glycol, glycerin, sorbitol, mannitol and polyethylene glycol; alginic acid; emulsifiers such as Tween®; wetting agents such as sodium lauryl sulfate; coloring agents; flavoring agents; tablets, stabilizers; antioxidants; preservatives; pyrogen-free water; isotonic saline solutions; and phosphate buffers and the like.

5 The composition of the present invention can be prepared into various dosage forms as needed, and the dosage to be administered to a patient can be determined by a physician according to factors, such as type, age, body weight, and general disease condition of a patient, mode of administration, and the like. For example, injection or other treatment may be used.

10 The present invention is further described below with reference to specific embodiments. It should be understood that these examples are only for illustrating the present invention and are not intended to limit the scope of the present invention. Experimental procedures in the following examples where no specific conditions are indicated are generally carried out in accordance with the conditions described in customary conditions such as those compiled by J. Sambrook et al.,
15 Molecular Cloning Experiments Guide, Third Edition, Science Press, 2002, or according to the manufacturer Suggested conditions.

Example 1. Preparation of a specific single chain antibody (scFv) binding to human GPC3

1.1 Screening of GPC3-specific binding antibodies based on phage display

20 Using phage display technology, human GPC3 (hereinafter referred to as huGPC3) specific antibody was screened from the all-human natural antibodies library. For this purpose, glycerol bacteria (purchased from Shanghai Rui Jin Biotechnology Co., Ltd.) from the natural library of phage-displayed all-human single-chain antibody were inoculated in
25 400 ml of 2 × YT/ampicillin medium so that the cell density reached $OD_{600} = 0.1$, and incubated at 37°C and 200 rpm until cell density reached $OD_{600} = 0.5$. Cells were infected with 10^{12} pfu of M13KO7 helper phage (purchased from Invitrogen) and incubated at 30°C and 50 rpm for 30 minutes. After 50 mg/L kanamycin was added and shaking-culture was performed at 37°C and 200 rpm for 30 minutes, the pellet was separated by centrifugation
30 (15 minutes, $1600 \times g$, 4°C) and resuspended in 400 ml of 2 × YT/Penicillin/kanamycin medium and shaken for 16 hours at 37°C and 200 rpm. Finally, the pellet was separated by centrifugation (5000 rpm, 4°C for 20 minutes) and discarded. The supernatant was filtered through a 0.45 μm filter and 1/4 volume of 20% (w/v) PEG 8000, 2.5 M NaCl solution was added and incubated in an ice bath for 1 hour to precipitate bacteriophage pellets. The pellet

was then precipitated by centrifugation (20 min, $8000 \times g$, 4°C) and the supernatant discarded. The phage were resuspended in 25 ml of prechilled PBS (137 mM NaCl, 2.7 mM KCl, 8 mM Na_2HPO_4 , 2 mM KH_2PO_4) and centrifuged (5 minutes, $20000 \times g$, 4°C). 1/4 volume of 20% (w/v) PEG8000, 2.5 M NaCl solution was added to the supernatant and incubated in an ice bath for 30 minutes to precipitate phage particles again. The pellets were centrifuged (30 min, $20000 \times g$, 4°C) and the phage pellets were resuspended in 2 ml of prechilled PBS again, kept on ice for 30 min and centrifuged (30 min, $17000 \times g$, 4°C). Supernatants were mixed with 4% (w/v) BSA in PBS at 1: 1, placed on a rotary mixer and incubated for 30 minutes at room temperature before being directly used for screening.

Using the above phage antibody library, four rounds of directional screening were performed on biotinylated human GPC3 recombinant protein (purchased from Shanghai Rui Jin Biotechnology Co., Ltd.) with the following scheme: The phage antibody library was incubated with biotin-labeled antigen GPC3 at room temperature for 2 hours and then incubated with streptavidin magnetic beads MyOne Cl (from Invitrogen) blocked with 2% (w/v) BSA (bovine serum albumin, purchased from Shanghai Bioengineering) at room temperature for 30 minutes. The beads were then washed with PBST (containing 0.1% Tween-20) buffer to remove phages which were not specifically bound or with weak binding capacities. Strongly-binding phages were then eluted from magnetic beads with glycine-HCl (pH 2.2), neutralized with Tris neutralizing solution (pH 9.1), and used to infect *E. coli* ER2738 in the mid-logarithmic growth phase and for the next round of screening. In the four rounds of screening, the beads were used in an amount of 50 μl , 20 μl , 10 μl and 10 μl , and the concentrations of biotin-labeled antigen GPC3 were 200 nM, 10 nM, 5 nM and 1 nM, respectively, and the time for PBST-washing was 10, 10, 15 and 20, respectively.

1.2 Identification of GPC3-specific binding antibodies

96 clones were randomly selected in the clones obtained from the fourth round of screening and their binding capability to human GPC3 was analyzed by single phage ELISA (enzyme-linked immunosorbent assay). For this purpose, each single colony was inoculated in 300 μl of $2 \times \text{YT}$ /ampicillin medium (containing 2% glucose) in a 96-well deep-well plate and cultured with shaking at 37°C and 250 rpm for 16 hours. 20 μl of culture was inoculated into 500 μl of $2 \times \text{YT}$ /ampicillin medium (containing 0.1% glucose) and shaken at 37°C and 250 rpm for 1.5 hours. To prepare the helper phage solution, 75 μl of M13KO7 (titer of 3×10^{12} pfu/ml) was taken and mixed into 15 ml of $2 \times \text{YT}$ medium and added into a culture plate at 50 μl /well, and incubated at 37°C and 150 rpm for 30 minutes. And then prepared

kanamycin solution (180 μ l of 50 mg/ml kanamycin was taken and added into 15 ml of 2 $\times$ YT medium) was added at 50 μ l/well and incubated with shaking for 16 hours at 37°C and 250 rpm. Finally, cells were precipitated by centrifugation (30 mins, 5000 $\times$ g, 4°C) and the supernatant was transferred to a new 96-well deep-well plate.

5 For single phage ELISA, 100 ng/well of antigen GPC3 and negative control protein BSA (100 μ l/well) were used in a 96-well MediSorp ELISA plate (purchased from Nunc) and coated overnight at 4°C. Each well was blocked with PBST containing 2% BSA (w/v). The wells were then washed with PBST for three times and PBST was discarded. Then, each phage solution prepared above was added into each well of the plate at 100 μ l/well. After
10 incubated at 37°C for 2 hours, the plate was washed for three times with PBST. To detect bound phage, anti-M13 antibody peroxidase conjugate (purchased from GE Healthcare) was diluted at 1: 5000 in PBST and 100 μ l was taken and added into each well. After incubated at 37°C for 1 hour, the wells were rinsed for three times with PBST and then rinsed for three times with PBS. Finally, 50 μ l of TMB substrate was pipetted into the wells
15 and developed for 10 minutes at room temperature, followed by addition of 50 μ l of 2M H₂SO₄ per well to quench the color reaction. Extinction values were measured at 450 nm with an enzyme-linked immunosorbent (Bio-Rad).

Two different single chain antibodies P1B12E (SEQ ID NO: 1 (nucleotide), 2 (amino acid)) and P7D4 (SEQ ID NO: 3 (nucleotide), 4 (amino acid)) were observed with
20 sequencing analysis, which exhibited significantly stronger binding signal to human GPC3 (huGPC3) in ELISA assay, while not binding to BSA (Figure 1).

Example 2. Expression and Purification of Anti-GPC3 Single-chain Antibody

According to the standard protocol, scFv-P1B12E fragment was amplified from the
25 plasmid (pCantab 5E-P1B12E) of screened clone P1B12E using primer pair V5-P1B12E-F (SEQ ID NO: 5) and V5-P1B12E-R (SEQ ID NO: 6), and scFv-P7D4 fragment was amplified from the plasmid (pCantab 5E-P7D4) of screened clone p7D4 (pCantab 5E-P7D4) using primer pair V5-P7D4-F (SEQ ID NO: 7) and V5-P7D4-R (SEQ ID NO: 8), digested with NheI/BamHI (purchased from NEB) and connected to vector plasmid pCMV-V5-Fc (Fc
30 fragment of human antibody IgG1 was fused and expressed downstream to the multiple cloning site of this vector, hereinafter referred to as V5-Fc, purchased from Shanghai Rui Jin Biotechnology Co., Ltd.) digested with NheI / BamHI using T4 DNA ligase (purchased from NEB) and was transformed into host strain TOP10. Clones were picked out and positive clones were identified by PCR and confirmed by sequencing, thereby obtaining

V5-scFv-P1B12E-Fc and V5-scFv-P7D4-Fc eukaryotic expression plasmids, respectively.

The above expression plasmids were respectively transfected into well-grown HEK-293F cells, which were cultured at 37°C, 5% CO₂, 125 rpm for 7 days and centrifuged at 4000rpm for 10 mins. Pellets were removed, and the supernatant was collected and filtered with 0.45 μm membrane. Processed samples were affinity-purified with protein A (from GE) affinity column, and purified antibody-Fc fusion protein scFv-P1B12E-Fc and scFv-P7D4-Fc were finally obtained. The identification results are shown in FIG. 2, and their molecular weights are around 50 kD.

Example 3. Binding of each cell line to anti-GPC3 single chain antibody analyzed through Flow cytometry

The ability of each of antibodies scFv-P1B12E-Fc and scFv-P7D4-Fc to bind to GPC3-positive hepatocarcinoma HepG2 cell line (ATCC) was analyzed by fluorescence activated cell sorter (FACS) (BD FACSCalibur).

Specific methods are as follows:

1) inoculating HepG2 hepatocarcinoma cell line in logarithmic growth phase into a 6 cm dish at inoculation cell density of about 90%, and incubating overnight at 37°C in an incubator.

2) digesting cells with 10 mM EDTA, collecting cells through centrifugation at 200 g × 5 mins, and resuspending cells in 1% phosphate buffered saline (NBS PBS) containing calf serum at 1×10^6 to 1×10^7 /mL into a flow-specific tube in an amount of 100 μl per tube.

3) centrifuging at 200g × 5min, and discarding the supernatant.

4) adding antibodies scFv-P1B12E-Fc and scFv-P7D4-Fc to be tested, respectively, while using PBS as a negative control. The final concentration of antibody is 10 μg/ml. 100 μl was added to each tube, and incubated in an ice bath for 45 minutes.

5). Adding 2 ml of 1% NBS PBS to each tube and centrifuging at 200 g × 5 min for two times.

6) Discarding the supernatant and adding FITC fluorescently labeled goat anti-human antibody (Shanghai Kangcheng Bioengineering Co., Ltd.) at a dilution of 1:50 with 100ul being added to each tube, incubating in an ice bath for 45 minutes.

7). Adding 2 ml of 1% NBS PBS into each tube, centrifuging at 200 g × 5 min for two times.

8) Discarding the supernatant, resuspending in 300 ul of 1% NBS PBS and detecting by flow cytometry.

9). analyzing the data using flow cytometry data analysis software WinMDI 2.9.

The results are shown in Fig. 3, and the results of flow cytometry showed that antibody scFv-P7D4-Fc can specifically recognize GPC3-expressing HepG2 cells, and antibody scFv-P1B12E-Fc did not bind to HepG2 cells.

5

Example 4. Screening and preparation of P7D4 single chain antibody mutants with increased GPC3 binding capacity

For enhancing the ability of P7D4 single-chain antibody to bind to GPC3, some amino acids in its heavy chain CDR1 and CDR2 regions, or light chain CDR1 and CDR2 were randomly mutated and the corresponding affinity mature libraries H12 and L12 were constructed.

4.1 CDR $\times$ P7D4 light and heavy chains and CDR regions thereof

The nucleotide sequence of P7D4 scFv is shown as follows (SEQ ID NO: 3; wherein positions 76-105 is heavy chain CDR1, positions 148-198 is the heavy chain CDR2, positions 295-330 is heavy chain CDR3, positions 475-516 is light chain CDR1; positions 562-582 is light chain CDR2, positions 679-708 is light chain CDR3; wherein positions 1-363 is heavy chain nucleotide sequence, 409-741 is light chain nucleotide sequence, positions 364-408 is (Gly₄Ser)₃ linker).

CAGGTGCAGCTGCAGGAGTCCGGGGGAGGCTTAGTTCAGCCTGGGAGGTCCCTGAGACTCTCC
20 TGTGCAGCCTCTGGATTACCTTCAGTAGCTATGCTATGCACITGGGTCCGCCAGGCTCCAGGCA
AGGGGCTGGAGTGGGTCTCAGCTATTAGTGGTAGTGGTGGTAGCACATACTACGCAGACTCCG
TGAAGGGCGCGTTCACCATCTCCAGAGACAATTCCAAGAACACGCTGTATCTGCAAATGAACA
GCCTGAGAGCCGAGGACACGGCCGTATATTACTGTGCGAAAATCGACGAGGGAGCCACGCT
GATGCTTTTGATGTCITGGGGCCAAGGAACCTGGTCACCGTCTCGAGTGGTGGAGGCGGTTTCAG
25 GCGGAGGTGGTTCTGGCGGTGGCGGATCGCAGTCTGCCCTGACTCAGCCTCCCTCCGCGTCCGG
GTCTCCTGGACAGTCAGTCACCATCTCCTGCCTGGAACCAGCAGTGACGTTGGTGGTTATAAC
TATGTCTCCITGGTACCAACAGTACCCAGGCAAAGCCCCAACTCCTCATCTATGGTAACAGC
AATCGGCCCTCAGGGGTCCCTGACCGATTCTCTGGCTCCAAGTCTGGCACCTCAGCCTCCCTGG
CCATCACTGGGCTCCAGGCTGAGGATGGGGCTGATTATTACTGCAGTCCTATGACAGCAGCC
30 TCGCTGTGGTAITTCGGCGGAGGGACCAAGGTCACCGTCCTAGGT

P7D4 scFv amino acid sequence is shown as follows (SEQ ID NO: 4):

QVQLQESGGGLVQPGRSRLRLSCAASGFTFSSYAMHWVRQAPGKGLEWVSAISGSGGSTYYA
DSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAKDRRGSHADAFDVWGQGLTVTVSSGGGG
35 SGGGSGGGGSQSALTQPPSASGSPGQSVTISCTGTSSDVGGYNYVSWYQQYPGKAPKLLIYGNSN
RPSGVPDRFSGSKSGTSASLAITGLQAEDGADYYCQSYDSSLRVVFGGGTKVTVLG

P7D4 VH CDR1:

nucleotide sequence: GGATTCACCTTCAGTAGCTATGCTATGCAC (SEQ ID NO: 9);
amino acid sequence: GFTFSSYAMH (SEQ ID NO: 10).

P7D4 VH CDR2:

nucleotide sequence: gctattagtggtagtggtggtagcacatactacgcagactccgtgaagggc (SEQ ID NO: 11);
amino acid sequence: AISGSGGSTYYADSVKG (SEQ ID NO: 12).

P7D4 VH CDR3:

nucleotide sequence: gatcgacgagggagccacgctgatgctttgatgtc (SEQ ID NO: 13);
amino acid sequence: DRRGSHADAFDV (SEQ ID NO: 14).

P7D4 VL CDR1:

nucleotide sequence: actggaaccagcagtgacgttggtggttataactatgtctcc (SEQ ID NO: 15);
amino acid sequence: TGTSSDVGGYNYVS (SEQ ID NO: 16).

P7D4 VL CDR2:

nucleotide sequence: ggtaacagcaatcggccctca (SEQ ID NO: 17);
amino acid sequence: GNSNRPS (SEQ ID NO: 18).

P7D4 VL CDR3:

nucleotide sequence: cagtcctatgacagcagcctgcgtgtggta (SEQ ID NO: 19);
amino acid sequence: QSYDSSLRVV (SEQ ID NO: 20).

4.2 Construction of Affinity Mature Library of H12

By sequence alignment and analysis of P7D4 single-chain antibody, part of amino acids in the first and second CDR regions of P7D4 heavy chain were selected and randomized mutations were introduced by primers to construct a heavy chain affinity mature library.

To prepare a DNA fragment encoding the P7D4 mutant library, two DNA fragments were respectively obtained by PCR using plasmid pCantab P7D4 as a template, followed by splicing through bypass PCR method. Specifically, the following procedure was used: for synthesizing genes, PCR reactions were performed in a volume of 50 μ l each using plasmid pCantab P7D4 as a template with a final concentration of 0.2 μ M for each primer and 5 μ l of 10 $\times$ KOD Plus buffer, 4 μ l dNTPs (dATP, dCTP, dGTP and dTTP, 2 mM each), 2 μ l 25

mM MgSO₄ and 1 U KOD Plus (from Takara) were added and the PCR procedure was started in a thermal cycler after making up the volume with water. The reaction was firstly heated to 94°C for 5 minutes and then incubated for 25 cycles of 94°C for 30 seconds, 56°C for 30 seconds and 68°C for 30 seconds, and finally, at 68°C for 10 minutes. The first
5 fragment was amplified using primers S1 (SEQ ID NO: 21, CAACGTGAAAAATTATTATTCGC) and 74H12F1r (SEQ ID NO: 22, CCAGCCCCTTGCCCTGGAGCCTGGCGGACCCAMNNCATAGCATAMNNACTGAAGGTGAATCCAG) and the second fragment was amplified using primer 74H12F2f (SEQ ID NO: 23,
10 GCTCCAGGCAAGGGGCTGGAGTGGGTCTCANNKATTAGTNNKNNKGNTNNKNNKA
CATACTACGCAGACTCC) and S6 (SEQ ID NO: 21, GTAAATGAATTTTCTGTATGAGG).

Expected PCR products were identified by analytical agarose gel electrophoresis and purified from samples by Wizard SV Gel and PCR Clean-up Kit (available from Promega). The two
15 fragments were added in equimolar ratio to a second round of bridge PCR as a template and the reaction system still used KOD Plus system mentioned above. The reaction was firstly heated to 94°C for 5 minutes and then incubated for 10 cycles, each cycling reaction conditions were 94°C for 30 seconds, 60°C for 30 seconds and 68°C for 30 seconds, and finally, at 68°C for 10 minutes. Subsequently, primers S1 and S6 were directly added to the reaction system at a final
20 concentration of 0.2 μM, and the PCR program was started. The reaction was firstly heated to 94°C for 5 minutes and then incubated for 25 cycles of 94°C for 30 seconds, 56°C for 30 seconds and 68°C for 30 seconds, and finally, at 68°C for 10 minutes. The expected PCR products were separated by preparative agarose gel electrophoresis and purified by Wizard SV Gel and PCR Clean-up kits according to the manufacturer's instructions.

25 In the library, complete DNA fragments contained *sfiI* and *NotI* restriction enzyme recognition sites at each end, and was digested by restriction endonuclease *sfiI* / *NotI* for restriction digestion and inserted into phagemid vector pCANTAB 5E digested by the same two enzymes. Ligation products were isolated and desalted using Wizard SV Gel and PCR Clean-up Kit for electrotransformation. For electrotransformation, a home-made competent *E. coli* ER2738
30 (available from NEB) was used with electroporation cuvette and electroporation instrument Gene Pulser II (from Bio-Rad). A library containing 8.9×10^9 mutants was finally confirmed.

4.3 Construction of Affinity Mature Library of L12

By sequence alignment and analysis of P7D4 single-chain antibody, part of amino acids

in the first and second CDR regions of P7D4 light chain were selected and randomized mutations were introduced by primers to construct a light chain affinity mature mutant library.

To prepare a DNA fragment encoding the P7D4 mutant library, two DNA fragments were respectively obtained by PCR using plasmid pCantab P7D4 as a template, followed by splicing through bypass PCR method. Specifically, the following procedure was used: for synthesizing genes, PCR reactions were performed in a volume of 50 μ l each using plasmid pCantab P7D4 as a template with a final concentration of 0.2 μ M for each primer and 5 μ l of 10 $\times$ KOD Plus buffer, 4 μ l dNTPs (dATP, dCTP, dGTP and dTTP, 2 mM each), 2 μ l 25 mM MgSO₄ and 1 U KOD Plus were added and the PCR procedure was started in a thermal cycler after making up the volume with water. The reaction was firstly heated to 94°C for 5 minutes and then incubated for 25 cycles of 94°C for 30 seconds, 56°C for 30 seconds and 68°C for 30 seconds, and finally, at 68°C for 10 minutes. The first fragment was amplified using primers S1 and 74L12F1r (SEQ ID NO: 83, GTTTGGGGGCTTTGCCTGGGTACTGTTGGTACCAGGAGACMNNAHNMNNAHNACC AACGTCACTGCTG) and the second fragment was amplified using primer 74L12F2f (SEQ ID NO: 84, ACCCAGGCAAAGCCCCAAACTCCTCATCTATNNKNNKNNKNN KCGGCCCTCAGGGGTC) and S6.

Expected PCR products were identified by analytical agarose gel electrophoresis and purified from samples by Wizard SV Gel and PCR Clean-up Kit. The two fragments were added in equimolar ratio to a second round of bridge PCR as a template and the reaction system still used KOD Plus system mentioned above. In the absence of primers, the reaction was firstly heated to 94°C for 5 minutes and then incubated for 10 cycles, each cycling reaction conditions were 94°C for 30 seconds, 60°C for 30 seconds and 68°C for 30 seconds, and finally, at 68°C for 10 minutes. Subsequently, primers S1 and S6 were directly added to the reaction system at a final concentration of 0.2 μ M, and the PCR program was started. The reaction was firstly heated to 94°C for 5 minutes and then incubated for 25 cycles of 94°C for 30 seconds, 56°C for 30 seconds and 68°C for 30 seconds, and finally, at 68°C for 10 minutes. The expected PCR products were separated by preparative agarose gel electrophoresis and purified by Wizard SV Gel and PCR Clean-up kits according to the manufacturer's instructions.

In the library, complete DNA fragments contained *sfi*I and *Not*I restriction enzyme recognition sites at each end, and was digested by restriction endonuclease *sfi*I / *Not*I for restriction digestion and inserted into phagemid vector pCANTAB 5E digested by the same two enzymes. Ligation products were isolated and desalted using Wizard SV Gel and PCR Clean-up Kit for

electrotransformation. For electrotransformation, a home-made competent *E. coli* ER2738 was used with electroporation cuvette and electroporation instrument Gene Pulser II. A library containing 1.1×10^{10} mutants was finally confirmed.

In addition, the inventors adopted the method of error prone PCR to randomly mutate the whole P7D4 fragment to construct a library T2 with a capacity of 7.9×10^9 , in which the amplification primer pair is S1 and S6, and the manner for cloning and construction is the same as that for the above described H12 and L12.

The screening of the above three affinity mature libraries is consistent with the screening procedure in Example 1. The inventors identified six high-affinity P7D4 series of mutant clones, am4, am14, am20, am35, am42 and T2-23 by screening, sequence information of which is shown as follows:

am4 nucleotide sequence (SEQ ID NO: 24, wherein positions 76-105 is the heavy chain CDR1, positions 148-198 is the heavy chain CDR2, positions 295-330 is the heavy chain CDR3, and positions 475-516 is light chain CDR1; positions 562-582 is light chain CDR2 and positions 679-708 is light chain CDR3; wherein positions 1-363 is heavy chain nucleotide sequence and positions 409-741 is light chain nucleotide sequence, and positions 364-408 is (Gly₄Ser)₃ linker):

```
CAGGTGCAGCTGCAGGAGTCCGGGGGAGGCTTAGTTCAGCCTGGGAGGTCCCTGAGACTC
TCCTGTGCAGCCTCTGGATTCACCTTCAGTACGTATGCTATGACGTTGGGTCCGCCAGGCT
CCAGGCAAGGGGCTGGAGTGGGTCTCACTATTAGTAGTAGTGGTGAAAGTACATACTAC
GCAGACTCCGTGAAGGGCGGTTCAACATCTCCAGAGACAATCCAAGAACACGCTGTAT
CTGCAATGAACAGCCTGAGAGCCGAGGACACGGCCGTATATTACTGTGCGAAA GATCGA
CGAGGGAGCCACGCTGATGCTTTTGATGCTTGGGGCCAAGGAACCCTGGTCACCGTCTCG
AGTGGTGGAGGCGGTTGAGGCGGAGGTGGTCTGGCGGTGGCGGATCGCAGTCTGCCCTG
ACTCAGCCTCCCTCCGCGTCCGGGTCTCCTGGACAGTCAGTCACCATCTCCTGC ACTGGA
ACCAGCAGTGACGTTGGTGGTTATACTATGTCTCC TGGTACCAACAGTACCCAGGCAAA
GCCCCAAACTCCTCATCTATGGTAACAGCAATCGGCCCTCA GGGGTCCCTGACCGATTC
TCTGGCTCCAAGTCTGGCACCTCAGCCTCCCTGGCCATCACTGGGCTCCAGGCTGAGGAT
GGGGCTGATTATTACTGC CAGTCCATGACAGCAGCCTGCGTGTGGTA TCGGCGGAGGG
ACCAAGGTCACCGTCCTAGGT
```

am4 amino acid sequence (SEQ ID NO: 25):

```
QVQLQESGGGLVQGRSLRLSCAASGFTFSTYAMTWVRQAPGKGLEWVSSISSSGESTYY
ADSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCAKDRRGSHADAFDVWGQGLVTVS
SGGGGSGGGGSGGGGQSALTQPPSASGSPGQSVTISC TGTSSDVGGYNYVS WYQQYPGK
APKLLIYGNSNRPSGVPDRFSGSKSGTSASLAITGLQAEDGADYYC QSYDSSLRVV FGGG
TKVTVLG
```

Heavy chain CDR1: GFTFSTYAMT (SEQ ID NO: 60)

Heavy chain CDR2: SISSSGESTYYADSVKG (SEQ ID NO: 61)

Heavy chain CDR3: DRRGSHADAFDV (SEQ ID NO: 14)

Light chain CDR1: TGTSSDVGGYNYVS (SEQ ID NO: 16)

Light chain CDR2: GNSNRPS (SEQ ID NO: 18)

Light chain CDR3: QSYDSSLRVV (SEQ ID NO: 20)

Wherein positions 1-121 is heavy chain amino acid sequence, positions 137-247 is light chain amino acid sequence, and positions 122-136 is (Gly₄Ser)₃ linker sequence.

5 am14 nucleotide sequence (SEQ ID NO: 26, wherein, positions 76-105 is heavy chain CDR1, positions 148-198th is heavy chain CDR2, positions 295-330 is heavy chain CDR3, positions 475-516 is light chain CDR1, positions 562-582 is light chain CDR2 and positions 679-708 is light chain CDR3; wherein positions 1-363 is heavy chain nucleotide sequence and positions 409-741 is light chain nucleotide sequence, positions 364-408 is (Gly₄Ser)₃ linker):

CAGGTGCAGCTGCAGGAGTCCGGGGGAGGCTTAGTTCAGCCTGGGAGGTCCCTGAGACTC
TCCTGTGCAGCCTCTGGATTACCTTCAGTACTTATGCTATGGCTTGGGTCCGCCAGGCT
CCAGGCAAGGGGCTGGAGTGGGTCTCAGAAATTAGTAGTTCTGGTAGTAGGACATACTAC
GCAGACTCCGTGAAGGGCGGTTACCATCTCCAGAGACAATTCCAAGAACACGCTGTAT
15 CTGCAAATGAACAGCCTGAGAGCCGAGGACACGGCCGTATATTACTGTGCGAAA GATCGA
CGAGGGAGCCACGCTGATGCTTTTGATGTC TGGGGCCAAGGAACCCTGGTCACCGTCTCG
AGTGGTGGAGGCGGTT CAGGCGGAGGTGGTCTGGCGGTGGCGGATCGCAGTCTGCCCTG
ACTCAGCCTCCCTCCGCGTCCGGGTCTCCTGGACAGTCAGTCACCATCTCCTGC ACTGGA
ACCAGCAGTGACGTTGGTGGTTATACTATGTCTCC TGGTACCAACAGTACCCAGGCAAA
20 GCCCCAAACTCCTCATCTAT GGTAACAGCAATCGGCCCTCA GGGGTCCCTGACCGATTC
TCTGGCTCCAAGTCTGGCACCTCAGCCTCCCTGGCCATCACTGGGCTCCAGGCTGAGGAT
GGGGCTGATTATTACTGC CAGTCCTATGACAGCAGCCTGCGTGTGGTA TCGGCGGAGGG
ACCAAGGTCACCGTCCTAGGT

am14 amino acid sequence (SEQ ID NO: 27):

25 QVQLQESGGGLVQGRSLRLSCAAS GFTFSTYAMA WVRQAPGKGLEWVS EISSSGSRTYY
ADSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCAK DRRGSHADAFDV WGQGLVTVS
SGGGSGGGSGGGGSQSALTQPPSASGSPGQSVTISC TGTSSDVGGYNYVS WYQQYPGK
APKLLIY GNSNRPS GVPDRFSGSKSGTSASLAITGLQAEDGADYYC QSYDSSLRVV FGGG
TKVTVLG

30 Heavy chain CDR1: GFTFSTYAMA (SEQ ID NO: 62)

Heavy chain CDR2: EISSSGSRTYYADSVKG (SEQ ID NO: 63)

Heavy chain CDR3: DRRGSHADAFDV (SEQ ID NO: 14)

Light chain CDR1: TGTSSDVGGYNYVS (SEQ ID NO: 16)

Light chain CDR2: GNSNRPS (SEQ ID NO: 18)

35 Light chain CDR3: QSYDSSLRVV (SEQ ID NO: 20)

Wherein positions 1-121 is heavy chain amino acid sequence and positions 137-247 is light chain amino acid sequence, and positions 122-136 is (Gly₄Ser)₃ linking sequence.

am20 nucleotide sequence (SEQ ID NO: 28, positions 76-105 is heavy chain CDR1,
40 positions 148-198 is heavy chain CDR2, positions 295-330 is heavy chain CDR3, positions

475-516 is light chain CDR1, positions 562-582 is light chain CDR2, positions 679-708 is light chain CDR3; wherein positions 1-363 is heavy chain nucleotide sequence, and positions 409-741 is light chain nucleotide sequence, positions 364-408 is (Gly₄Ser)₃ linking sequence):

5 CAGGTGCAGCTGCAGGAGTCCGGGGGAGGCTTAGTTCAGCCTGGGAGGTCCCTGAGACTC
 TCCTGTGCAGCCTCTGGATTACCTTCAGTACGTATGCTATGAATTGGGTCCGCCAGGCT
 CCAGGCAAGGGGCTGGAGTGGGTCTCAGCGATTAGTATGTCTGGTGAATCTACATACTAC
 GCAGACTCCGTGAAGGGCCGGTTCACCATCTCCAGAGACAATTCCAAGAACACGCTGTAT
 CTGCAAATGAACAGCCTGAGAGCCGAGGACACGGCCGTATATTACTGTGCGAAA GATCGA
 10 CGAGGGAGCCACGCTGATGCTTTGATGTC TGGGGCCAAGGAACCCTGGTCACCGTCTCG
 AGTGGTGGAGGCGGTT CAGGCGGAGGTGGTTCTGGCGGTGGCGGATCGCAGTCTGCCCTG
 ACTCAGCCTCCCTCCGCGTCCGGGTCTCCTGGACAGTCAGTCACCATCTCCTGC ACTGGA
 ACCAGCAGTGACGTTGGTGGTTATAACTATGTCTCC TGGTACCAACAGTACCCAGGCAAA
 15 GCCCCAAACTCCTCATCTATGGTAACAGCAATCGGCCCTCA GGGGTCCCTGACCGATTCT
 TCTGGCTCCAAGTCTGGCACCTCAGCCTCCCTGGCCATCACTGGGCTCCAGGCTGAGGAT
 GGGGCTGATTATTACTGC CAGTCTATGACAGCAGCCTGCGTGTGGTA TTCGGCGGAGGG
 ACCAAGGTCACCGTCCTAGGT

am20 amino sequence (SEQ ID NO: 29):

20 QVQLQESGGGLVQPGRSLRLSCAASGFTFSTYAMNVWRQAPGKGLEWVS AISMSGESTYY
 ADSVKG RFTISRDN SKNTLYLQMNSLRAEDTAVYYCAK DRRGSHADAFDV WGQGLTVTVS
 SGGGSGGGGSGGGGQSALTQPPSASGSPGQSVTISC TGTSSDVGGYNYVS WYQQYPGK
 APKLLIY GNSNRPS GVPDRFSGSKSGTSASLAITGLQAEDGADYYC QSYDSSLRVV FGGG
 TKVTVLG

Heavy chain CDR1: GFTFSTYAMA (SEQ ID NO: 64)

25 Heavy chain CDR2: AISMSGESTYYADSVKG (SEQ ID NO: 65)

Heavy chain CDR3: DRRGSHADAFDV (SEQ ID NO: 14)

Light chain CDR1: TGTSSDVGGYNYVS (SEQ ID NO: 16)

Light chain CDR2: GNSNRPS (SEQ ID NO: 18)

Light chain CDR3: QSYDSSLRVV (SEQ ID NO: 20)

30 Wherein positions 1-121 is heavy chain amino acid sequence and positions 137-247 is light chain amino acid sequence, and positions 122-136 is (Gly₄Ser)₃ linking sequence.

am35 nucleotide sequence (SEQ ID NO: 30, positions 76-105 is heavy chain CDR1, positions 148-198 is heavy chain CDR2, positions 295-330 is heavy chain CDR3, positions 475-516 is light chain CDR1, positions 562-582 is light chain CDR2, positions 679-708 is light chain CDR3; wherein positions 1-363 is heavy chain nucleotide sequence, and positions 409-741 is light chain nucleotide sequence, positions 364-408 is (Gly₄Ser)₃ linking sequence):

40 CAGGTGCAGCTGCAGGAGTCCGGGGGAGGCTTAGTTCAGCCTGGGAGGTCCCTGAGACTC
 TCCTGTGCAGCCTCTGGATTACCTTCAGTAGCTATGCTATGCAC TGGGTCCGCCAGGCT
 CCAGGCAAGGGGCTGGAGTGGGTCTCAGCTATTAGTAGTAGTGGTGGTAGCACATACTAC

5 GCAGACTCCGTGAAGGGC CGGTTCACCATCTCCAGAGACAATTCCAAGAACACGCTGTAT
 CTGCAAATGAACAGCCTGAGAGCCGAGGACACGGCCGTATATTACTGTGCGAAA GATCGA
 CGAGGGAGCCACGCTGATGCTTTTGATGTC TGGGGCCAAGGAACCCCTGGTCACCGTCTCG
 AGTGGTGGAGGCGGTT CAGGCGGAGGTGGTCTGGCGGTGGCGGATCG CAGTCTGCCCTG
 ACTCAGCCTCCCTCCGCGTCCGGGTCTCCTGGACAGTCAGTCACCATCTCCTGC ACTGGA
 10 ACCAGCAGTGACGTTGGTCATAAGTTTCCTGTCTCC TGGTACCAACAGTACCCAGGCAAA
 GCCCCAAACTCCTCATCTAT AAGAATCTTTTGGCGCCCTCA GGGGTCCCTGACCGATTG
 TCTGGCTCCAAGTCTGGCACCTCAGCCTCCCTGGCCATCACTGGGCTCCAGGCTGAGGAT
 GGGGCTGATTATTACTGC CAGTCCTATGACAGCAGCCTGCGTGTGGTA TTCGGCGGAGGG
 ACCAAGGTCACCGTCCTAGGT

am35 amino acid sequence (SEQ ID NO: 31):
 QVQLQESGGGLVQPGRSLRLSCAAS GFTFSSYAMH WVRQAPGKGLEWVS AISSSGGSTYY
 ADSVKG RFTISRDNSKNTLYLQMNSLRAEDTAVYYCAK DRRGSHADAFDV WGQGLTVTS
 15 SGGGSGGGGSGGGGSQSALTQPPSASGSPGQSVTISC TGTSSDVGHKFPVS WYQYYPGK
 APKLLIY KNLLRPS GVPDRFSGSKSGTSASLAITGLQAEDGADYYC QSYDSSLRVV FGGG
 TKVTVLG

Heavy chain CDR1: GFTFSSYAMH (SEQ ID NO: 10)
 Heavy chain CDR2: AISSSGGSTYYADSVKG (SEQ ID NO: 66)
 Heavy chain CDR3: DRRGSHADAFDV (SEQ ID NO: 14)
 20 Light chain CDR1: TGTSSDVGHKFPVS (SEQ ID NO: 67)
 Light chain CDR2: KNLLRPS (SEQ ID NO: 68)
 Light chain CDR3: QSYDSSLRVV (SEQ ID NO: 20)

25 Wherein positions 1-121 is heavy chain amino acid sequence and positions 137-247 is
 light chain amino acid sequence, and positions 122-136 is (Gly₄Ser)₃ linking sequence.

am42 nucleotide sequence (SEQ ID NO: 32, positions 76-105 is heavy chain CDR1,
 positions 148-198 is heavy chain CDR2, positions 295-330 is heavy chain CDR3, positions
 475-516 is light chain CDR1, positions 562-582 is light chain CDR2, positions 679-708 is
 light chain CDR3; wherein positions 1-363 is heavy chain nucleotide sequence, and positions
 30 409-741 is light chain nucleotide sequence, positions 364-408 is (Gly₄Ser)₃ linking
 sequence):

35 CAGGTGCAGCTGCAGGAGTCCGGGGGAGGCTTAGTTCAGCCTGGGAGGTCCCTGAGACTC
 TCCTGTGCAGCCTCT GGATTCACCTTCAGTAGCTATGCTATGCAC TGGGTCCGCCAGGCT
 CCAGGCAAGGGGCTGGAGTGGGTCTCA GCTATTAGTAGTAGTGGTGGTAGCACATACTAC
 GCAGACTCCGTGAAGGGC CGGTTCACCATCTCCAGAGACAATTCCAAGAACACGCTGTAT
 CTGCAAATGAACAGCCTGAGAGCCGAGGACACGGCCGTATATTACTGTGCGAAA GATCGA
 CGAGGGAGCCACGCTGATGCTTTTGATGTC TGGGGCCAAGGAACCCCTGGTCACCGTCTCG
 AGTGGTGGAGGCGGTT CAGGCGGAGGTGGTCTGGCGGTGGCGGATCG CAGTCTGCCCTG
 ACTCAGCCTCCCTCCGCGTCCGGGTCTCCTGGACAGTCAGTCACCATCTCCTGC ACTGGA
 40 ACCAGCAGTGACGTTGGTCTTATGCATAATGTCTCC TGGTACCAACAGTACCCAGGCAAA
 GCCCCAAACTCCTCATCTAT AAGTCTTCGTCTCGGCCCTCA GGGGTCCCTGACCGATTG
 TCTGGCTCCAAGTCTGGCACCTCAGCCTCCCTGGCCATCACTGGGCTCCAGGCTGAGGAT
 GGGGCTGATTATTACTGC CAGTCCTATGACAGCAGCCTGCGTGTGGTA TTCGGCGGAGGG

ACCAAGGTCACCGTCCTAGGT

am42 amino acid sequence (SEQ ID NO: 33):

QVQLQESGGGLVQPGRLRLSCAAS[GFTFSSYAMH]WVRQAPGKGLEWVS[AISSSGGSTYY]
[ADSVKG]RFTISRDNKNTLYLQMNSLRAEDTAVYYCAK[DRRGSHADAFDV]WGQGLVTVS
5 SGGGSGGGGSGGGGQSALTQPPSASGSPGQSVTISC[TGTSSDVGLMHNVS]WYQQYPGK
APKLLIY[KSSSRPS]GVPDRFSGSKSGTSASLAITGLQAEDGADYYC[QSYDSSLRVV]FGGG
TKVTVLG

Heavy chain CDR1: GFTFSSYAMH (SEQ ID NO: 10)

Heavy chain CDR2: AISSSGGSTYYADSVKG (SEQ ID NO: 66)

10 Heavy chain CDR3: DRRGSHADAFDV (SEQ ID NO: 14)

Light chain CDR1: TGTSSDVGLMHNVS (SEQ ID NO: 69)

Light chain CDR2: KSSSRPS (SEQ ID NO: 70)

Light chain CDR3: QSYDSSLRVV (SEQ ID NO: 20)

Wherein positions 1-121 is heavy chain amino acid sequence and positions 137-247 is
15 light chain amino acid sequence, and positions 122-136 is (Gly₄Ser)₃ linking sequence.

T2-23 nucleotide sequence (SEQ ID NO: 34, positions 76-105 is heavy chain CDR1,
positions 148-198 is heavy chain CDR2, positions 295-330 is heavy chain CDR3, positions
475-516 is light chain CDR1, positions 562-582 is light chain CDR2, positions 679-708 is
20 light chain CDR3; wherein positions 1-363 is heavy chain nucleotide sequence, and positions
409-741 is light chain nucleotide sequence, positions 364-408 is (Gly₄Ser)₃ linking
sequence):

CAGGTGCAGCTGCAGGAGTCCGGGGGAGGCTTAGTTCAGCCTGGGAGGTCCCTGAGACTC
TCCTGTGCAGCCTCT[GGATTACCTTCAGTAGCTATGCTATGCAC]TGGGTCCGCCAGGCT
25 CCAGGCAAGGGGCTGGAGTGGGTCTCA[GCTATTAGTAGTAGTGGTCGTAGCACATACTAC]
[GCAGACTCCGTGGAGGGG]CGGTTCACCATCTCCAGAGACAATTCCAAGAACACGCTGTAT
CTGCAAATGAACAGCCTGAGAGCCGAGGACACGGCCGTATATTACTGTGCGAAA[GATCGA]
[CGAGGGAGCCACGCTGATGCTTTAAATGTC]TGGGGCCAAGGAACCCTGGTCACCGTCTCG
AGTGGTGGAGGCGGTT[CAGGCGGAGGTGGTTCGGCGGTGGCGGATCG]CAGTCTGCCCTG
30 ACTCAGCCTCCCTCCGCGTCCGGGTCTCCTGGACAGTCAGTCACCATCTCCTGC[ACTGGA]
[ACCAGCAGTGACGTTGGTGGTTATACTATGTCTCC]TGGTACCAACAGTACCCAGGCAAA
GCCCCAAACTCCTCATCTAT[GGTAACAGCAATCGGCCCTCA]GGGGTCCCTGACCGATTC
TCTGGCTCCAAGTCTGGCACCTCAGCCTCCCTGGCCATCACTGGGCTCCAGGCTGAGGAT
GGGGCTGATTATTACTGC[CAGTCCTATGACAGCAGCCTGCGTGTGGTA]TTCGGCGGAGGG
35 ACCAAGGTCACCGTCCTAGGT

T2-23 amino acid sequence (SEQ ID NO: 35):

QVQLQESGGGLVQPGRLRLSCAAS[GFTFSSYAMH]WVRQAPGKGLEWVS[AISSGRSTYY]
[ADSVKG]RFTISRDNKNTLYLQMNSLRAEDTAVYYCAK[DRRGSHADALNV]WGQGLVTVS
40 SGGGSGGGGSGGGGQSALTQPPSASGSPGQSVTISC[TGTSSDVGGYNYVS]WYQQYPGK
APKLLIY[GNSNRPS]GVPDRFSGSKSGTSASLAITGLQAEDGADYYC[QSYDSSLRVV]FGGG
TKVTVLG

Heavy chain CDR1: GFTFSSYAMH (SEQ ID NO: 10)

Heavy chain CDR2: AISSSGRSTYYADSVEG (SEQ ID NO: 71)

Heavy chain CDR3: DRRGSHADALNV (SEQ ID NO: 72)

Light chain CDR1: TGTSSDVGGYNYVS (SEQ ID NO: 16)

Light chain CDR2: KSSSRPS (SEQ ID NO: 70)

5 Light chain CDR3: QSYDSSLRVV (SEQ ID NO: 20)

Wherein positions 1-121 is heavy chain amino acid sequence and positions 137-247 is light chain amino acid sequence, and positions 122-136 is (Gly₄Ser)₃ linking sequence.

4.4 SPR analysis of the binding ability of P7D4 series antibodies to GPC3

10 To quantitatively analyze the binding of P7D4 series antibodies to GPC3, the affinity and kinetic parameters of P7D4 series single-chain antibodies were measured by capture method using Biacore T200 system (from GE). An anti-human IgG (Fc) antibody (purchased from GE) was coupled to carboxymethyl dextran surface of sensor chip CM5 through primary amino with NHS / EDC coupling according to the manufacturer's instructions.

15 Measurements were performed in 1 × HBS-EP + working buffer at 25°C, 30 µl/min, and regeneration condition was 3 M MgCl₂, 10 µl/min for 30 seconds. In each round of the testing cycle, the antibody to be tested is firstly captured onto the chip. GPC3 of a certain concentration flowed over the chip surface. The interaction between human GPC3 and the captured antibody caused the change of the molecular concentration on the surface of the

20 sensor chip which was measured according to changes in SPR Signal and expressed in resonance units (RU). Time vs resonance unit (RU) was plotted. The resulting sensorgram documents the entire reaction including the binding and dissociation processes (Figure 4). In all single-cycle kinetics, GPC3 concentrations were 5 nM, 10 nM, 20 nM, 40 nM and 80 nM, respectively. The resulting curves were evaluated using Biacore T200 evaluation software

25 and the affinity KD values were calculated. The binding data for all P7D4 single chain antibodies to GPC3 are summarized in Table 1 below.

Table 1. Binding data for P7D4 series single chain antibodies to GPC3

Antibody sample	ka (1/Ms)	kd (1/s)	KD (M)
P7D4	9.31E+04	6.17E-03	6.64E-08
am4	5.39E+05	4.08E-05	7.56E-11
am14	6.73E+05	4.43E-05	6.46E-11
am20	2.12E+05	6.47E-05	3.06E-10
am35	7.38E+05	5.2 E-05	7.14E-11

am42	8.16E+05	6.70E-05	8.21E-11
T2-23	8.86E+05	5.19E-05	5.85E-11

It can be seen from Table 1 that P7D4 series mutant single chain antibodies obtained by modification exhibited good and significant increased affinity relative to P7D4 single-chain antibody.

4.5 Analysis of Identification Specificity of P7D4 series Antibodies

In order to analyze the binding specificity of P7D4 series antibody to GPC3 protein, binding activity of P7D4 series single chain antibody to human GPC family members, including recombinant human GPC1 (rhGPC1), GPC2 (rhGPC2), GPC3 (rhGPC3), GPC5 (rhGPC5) and GPC6 (rhGPC6) (purchased from Andy Biosciences (Shanghai) Co., Ltd.) was detected by ELISA, respectively.

For this purpose, the above 5 antigens were diluted with 0.1 M NaHCO₃ (pH 9.6) coating solution and each well was coated with 100 ng at 50 µl/well overnight at 4°C and blocked with 2% (w/v) BSA in PBST for 2 hours at room temperature. The plate was then rinsed for three times with PBST and PBST was removed. Subsequently, 100 ng of each antibody protein in PBST was added to each well plate, and the assay for each sample was performed in duplicate wells. After incubated for 2 hours at 37°C, the plates were rinsed for three times with PBST followed by addition of 100 µl/well HRP-labeled rabbit anti-human Fc antibody (purchased from Shanghai Rui Jin Biotechnology Co., Ltd.) diluted at 1: 20000 and incubated at 37°C for 1 hour. For detection, wells were rinsed for three times with PBST followed by three times with PBS and finally with TMB for development for 10 min. The chromogenic reaction was stopped with 50 µl of 2 M H₂SO₄ per well, and extinction value was measured at 450 nm with enzyme-linked immunodetector (Bio-Rad).

Results are shown in Figure 5, and all P7D4 series single chain antibodies only specifically bind to human GPC3 and do not bind to human GPC1, GPC2, GPC5 and GPC6.

Example 5. Preparation of Human GPC3 Antigen and Humanized Monoclonal Antibodies against the Antigen

5.1 Expression and purification of Human GPC3 in eukaryotic expression system

5.1.1 Construction and identification of GPC3 vector

PCR amplification was performed using human hepatoma cell line Huh-7 cDNA as a template with following primers GPC3-F: GATCGCTAGCACAGCCCCCGCCGCCGC (SEQ ID NO: 54), GPC3-R: GTACGGATCCTTCAGCGGGGAATGAACGTTC (SEQ ID

NO: 55) to obtain GPC3 fragment (1.6 kb) with NheI / BamHI site at both ends. The obtained PCR fragment was double digested with endonuclease NheI / BamHI (purchased from Fermentas), and the plasmid vector V5H (purchased from Raygene) was double digested with endonuclease NheI / BamHI. The vector and inserted fragment were recovered by agarose gel electrophoresis and ligated by T4 DNA ligase (purchased from NEB) and transformed into host strain TOP10 (purchased from LIFE Co.). The host strains were screened by Ampicillin-resistance, clones were pick out, plasmids were extracted and double digested by NheI / BamHI (purchased from Fermentas), and positive clones containing the inserted fragment were identified and verified by sequencing to obtain eukaryotic expression plasmid V5H- GPC3 containing the correct human GPC3 gene sequence.

5.1.2 Expression and purification of human GPC3 protein

5.1.2.1. Liposome transfection and culture of V5H-GPC3 plasmid

Well-grown HEK293F cells (HEK293F, purchased from LIFE Co., Ltd.) were seeded in a cell culture flask at a density of 1×10^6 cells/ml and incubated overnight at 37°C and 120 rpm for future use. The plasmid V5H-GPC3 obtained in the above step and lipofectamine 293Fectin (purchased from LIFE) was diluted with DMEM and mixed gently, incubated at room temperature for 20 min. The incubated DNA-liposome complex was added to 293F cells and cultured at 37°C and 120 rpm for 120 h. The cell culture was collected and centrifuged at 4500g for 15min. Cells were removed and the supernatant was kept.

5.1.2.2. Purification of GPC3 protein

1 ml of Ni-NTA Agarose affinity filler was loaded and the Ni-NTA affinity column was equilibrated with 10 column volumes of equilibration buffer (50 mM PB, 0.3 M NaCl, 10 mM imidazole, pH 8.0). After centrifugation, the supernatant of cell culture was passed through a Ni-NTA affinity column at 1 ml/min and the flow-through was collected and stored at 4°C. 10 column volumes of wash buffer 1 (50 mM PB, 0.3 M NaCl, 20 mM imidazole, pH 8.0) was used for washing and the flow-through was collected at 4°C. 4-5 column volumes of elution buffer (50 mM PB, 0.3 M NaCl, 250 mM imidazole, pH 8.0) were used for elution and the eluate was collected and dialysed in dialysis fluid (50 mM PB, pH 7.8, 0.3 M NaCl, Glycerol) overnight at 4°C to obtain GPC3(H) protein, and a small amount was used in SDS PAGE electrophoresis (FIG. 6).

5.2 Human GPC3 antigen immunization

Recombinant protein immunization: 1 ml of the purified human GPC3 protein GPC3(H) (1.0 mg/mL) obtained in the above Example 5.1 as an antigen was sufficiently emulsified and

mixed with 1 mL of complete Freund's adjuvant (Sigma-Aldrich Co., Ltd.) for subcutaneously immunizing BALB/c mice (6-8 weeks old, 100 µg of human GPC3(H) protein antigen per mouse). Four weeks later, the human GPC3 antigen was emulsified and mixed with incomplete Freund's adjuvant. The mice were immunized through intraperitoneal injection (50 µg for each mouse), followed by an interval of 2 weeks and then 50 µg of antigen was intraperitoneally administered for booster immunization. One week after the fourth booster immunization, GPC3(H) protein was used for coating. The mouse antiserum titer was detected by ELISA, and the booster immunization was continued until the antiserum titer reached $> 10^5$ in mice.

Three weeks after the last booster immunization, 20 µg of human GPC3(H) protein was immunized in spleen for future use.

5.3 Establishment of Anti-human GPC3 hybridoma cell lines

Four days after boosting in the spleen of mouse, the spleens were taken aseptically and the lymphocytes were isolated by filtration through a 100-mesh filter, fused with myeloma cell line SP2/0, selectively cultured with hypoxanthine, aminopterin and thymidine (HAT) for three days, and then HT medium was added and the culture was continued for one week.

GPC3(H) antigen was used for coating, and the positive clones were screened by ELISA and subcloned for three times by limiting dilution. The cells were cultured for another 2 months. Finally, stable hybridoma cell lines (clone numbers 5A5, 7C9 and 11D3) were obtained.

5.4 Production of ascites and purification of antibody

F1 mice of 8-10 weeks old were injected intraperitoneally with 100 µL of pristane (purchased from Sigma-Aldrich), and 1 week later, hybridoma cell clones of the above Example 5.4 were taken and intraperitoneally injected into the mice at 5×10^5 cells/mouse to prepare ascites. After 7 to 10 days, ascites was collected and centrifuged at 10000g for 10 mins, and the supernatant was taken for future use. Protein G affinity column (purchased from GE) was returned to room temperature and 5 column volumes of PBS (0.01 M PB, 0.15 M NaCl, pH 7.4) was used for equilibration. The supernatant of ascites was mixed with an equal volume of PBS (0.01 M PB, 0.15 M NaCl, pH 7.4), filtered through a 0.22 µm filter and the filtered ascites supernatant was loaded on a Protein G affinity column and washed with 5 volumes of PBS. The column was eluted with elution buffer (0.1M Glycine HCl, pH 2.7) and the elution was neutralized with 1/10 volume of neutralization buffer (1M NaH₂PO₄,

pH 9.0). The solution was dialyzed against PBS (0.01 M PB, 0.15 M NaCl, pH 7.4), during which PBS was changed for two times and the interval between two changes was longer than 5 hours. The dialyzed solution was centrifuged at 10000g for 10min and the supernatant was filtered through a 0.22um filter to obtain a solution of the purified anti-human GPC3 monoclonal antibody produced by each clone. According to analysis, 5A5 exhibited good antigen binding ability.

5.5 Humanization of GPC3 clone 5A5

5.5.1 Sequence analysis of 5A5 antibody

5A5 exhibited good antigen binding ability. The variable region sequences of VL and VH of 5A5 antibody were determined. According to the naming scheme of CDRs of Kabat, Chothia and IMGT, the sequence of six CDRs of the light and heavy chain of the antibody were determined.

The nucleotide sequence of 5A5 heavy chain is shown as follows (SEQ ID NO: 56, where the three underlined segments are successively CDR1, CDR2, CDR3):

CAGGTTCAACTGCAGCAGTCTGGGACTGAGCTGGTGAGGCCTGGGGCTTCA
GTGAAGCTGTCCTGCAAGGCTTTGGGCTACACATTTACTGACTATGAAATGC
ACTGGGTGAAGCAGACACCTGTGCATGGCCTGGAGTGGATTGGAGCTATTC
ATCCAGGAAGTGGTGATACTGCCTACAATCAGAGGTTCAAGGGCAAGGCCA
CACTGACTGCAGACAAATCTTCCAGCACAGCCTACATGGAGTACAGCAGCC
TGACATCTGAGGACTCTGCTGTCTATTACTGTACAAGATTTTATTCCTATGCT
TACTGGGGCCAAGGGACTCTGGTCACTGTCTCTGCA

The nucleotide sequence of 5A5 light chain is shown as follows (SEQ ID NO: 57, where the three underlined segments are successively CDR1, CDR2, CDR3):

GATGTTGTGATGACCCAACTCCACTCTCCCTGCCTGTCAGTCTTGGAGATCAAGCCTCCATCT
CTTGCAGATCTAGTCAGAGCCTTGTACACAGTAATGGAAACACCTATTTACAGTGGTACCTGCA
GAAGCCAGGCCAGTCTCCAAAGCTCCTGATCTACAAAAGTTTCCAATCGATTTTCTGGGGTCCCA
GACAGGTTCACTGGCAGAGGATCAGGGACAGATTTACACTCAAGATCAGCAGAGTGGAGGCT
GAGGATCTGGGAGTTTATTTCTGCTCTCAAAGTATATATGTTCCGTACACGTTCCGAGGAGGGA
CCAAGCTGGAAATAAAACGG

Amino acid sequence of 5A5 heavy chain (SEQ ID NO: 81, where the three underlined segments are successively CDR1, CDR2, CDR3):

QVQLQQSGTELVRPGASVKLSCKALGYTFTDYEMHWVVKQTPVHGLEWIGAIHPGSGDTAYNQRF
KGKATLTADKSSSTAYMEYSSLTSEDSAVYYCTRFYSYAYWGQGLTVTSA

Amino acid sequence of 5A5 light chain (SEQ ID NO: 82, where the three underlined

segments are successively CDR1, CDR2, CDR3)

DVVMQTQTPLSLPVSLGDQASISCRSSQSLVHSNGNTY¹LQWYLQKPGQSPKLLIYK²VS³NRFS⁴GV⁵PD⁶RF⁷SG⁸R
GSGTDFTLKISRVEAEDLGVYFCS⁹QSIYVPY¹⁰TFGGG¹¹TKLEIKR

5 5.5.2 Selection of antibody template

There are four major factors in the selection of antibody template framework regions: immunogenicity, antigen binding properties, expression, and antibody stability.

(1) Determination of the amino acid residues supporting loop structure of an antibody in the light and heavy chain variable regions of the antibody and the amino acid residues of the light and heavy chain binding regions according to WO2008021156.

(2) Alignment of the full length of the 5A5 light chain or heavy chain variable region with antibodies from IMGT, V BASE or NCBI.

(3) Removing amino acid residues associated with supporting loop within CDR regions of 5A5 light or heavy chain variable region and then sequence alignment with antibodies from IMGT, V BASE or NCBI.

(4) Removing amino acid residues in CDR regions in the light chain or heavy chain variable region of 5A5 and then sequence alignment with antibodies from IMGT, V BASE or NCBI.

(5) Only keeping amino acid residues associated with supporting loop in the light or heavy chain sequence of 5A5, and then sequence similarity alignment with antibodies from IMGT, V BASE or NCBI.

(6) Only keeping amino acid residues associated with supporting loop in light chains or heavy chain frameworks of 5A5 are retained, and then sequence similarity alignment with antibodies from IMGT, V BASE or NCBI.

(7) Selecting the antibody with the highest similarity as the antibody template according to results from the alignment of sequence similarity.

(8) Results from alignment of sequence similarity of heavy chain: Based on the similarity, VH1_69 * 06 (IMGT, Accession numbers: L22583) was selected as the antibody template for 5A5 heavy chain.

(9) Results from alignment of sequence similarity of light chain: based on the similarity, VK2D_29 * 02 (IMGT, Accession numbers: U41644) was selected as the antibody template for 5A5 light chain.

5.5.3 CDR transplantation

The light chain or heavy chain CDR regions of 5A5 antibody are substituted for the CDR

regions of the antibody template. Humanized antibody (Y035) was finally identified. Its amino acid sequence is shown as follows:

Humanized Y035 heavy chain (SEQ ID NO: 58):

EVQLVQSGAEVKKPGASVKVSCKASGYTFSDIYEMHWVRQAPGQGLEWMGAIHPGSGDTAYNQ

5 RFKGRVTITADKSTSTAYMELSSLRSEDTAVYYCARFYSYAYWGQGLTVSA

Heavy chain CDR1 amino acid sequence is: Asp Tyr Glu Met His (SEQ ID NO: 73);

Heavy chain CDR2 amino acid sequence is: Ala Ile His Pro Gly Ser Gly Asp Thr Ala Tyr Asn
Gln Arg Phe Lys Gly (SEQ ID NO: 74);

Heavy chain CDR3 amino acid sequence is: Phe Tyr Ser Tyr Ala Tyr (SEQ ID NO: 75).

10

Humanized Y035 light chain (SEQ ID NO: 59):

DIVMTQTPLSLPVTPGEPASISCRSSQSLVHSNGNTYLQWYLQKPGQSPQLLIYKVSNRFSGVPDRF
SGSGSGTDFTLKISRVEAEDVGVYYCSQSIYVPYTFGQGTKLEIKR

15 Light chain CDR1 amino acid sequence is: Arg Ser Ser Gln Ser Leu Val His Ser Asn Gly Asn
Thr Tyr Leu Gln (SEQ ID NO: 76);

Light chain CDR2 amino acid sequence is: Lys Val Ser Asn Arg Phe Ser (SEQ ID NO: 77);

Light chain CDR3 amino acid sequence is: Ser Gln Ser Ile Tyr Val Pro Tyr Thr (SEQ ID NO:
78).

20 5.5.4 Expression and purification of humanized antibody

(1) Based on the amino acid sequence of humanized antibody (Y035), the nucleotide sequence was designed and synthesized.

Synthesized light chain nucleotide sequence (SEQ ID NO: 79, including signal peptide encoding sequence):

25 GGATCGATATCCACCATGGACATGATGGTGCTGGCCCAGTTCCTGGCCTTCCTGCTGCTGTGGTTCC
CAGGCGCTAGATGCGACATCGTGATGACCCAGACCCCCCTGAGCCTGCCCGTGACCCCCGGCGA
GCCCCGCCAGCATCAGCTGCCGGAGCAGCCAGAGCCTGGTGCACAGCAACGGCAACACCTACC
TGCAGTGGTACCTGCAGAAGCCCGGCCAGAGCCCCCAGCTGCTGATCTACAAGGTGAGCAAC
30 CGGTTACAGCGGCGTGCCCGACCGGTTACAGCGGCAGCGGCAGCGGCACCGACTTCACCCTGA
AGATCAGCCGGGTGGAGGCCGAGGACGTGGGCGTGTTACTACTGCAGCCAGAGCATCTACGTG
CCCTACACCTTCGGCCAGGGCACCAAGCTGGAGATCAAACGTACGGTGGCT

Synthesized heavy chain nucleotide sequence (SEQ ID NO: 80, including signal peptide encoding sequence):

35 GGATCGATATCTGCGGCCTATCTAGCCACCATGCGGGTGCTGATCCTGCTGTGGCTGTTTACCGCCT
TCCCCGGCTTCCTGAGCGAGGTGCAGCTGGTGCAGAGCGGCGCCGAGGTGAAGAAGCCCGGCG
CCAGCGTGAAGGTGAGCTGCAAGGCCAGCGGCTACACCTTCAGCGACTACGAGATGCACTGG
GTGCGGCAGGCCCCCGGCCAGGGCCTGGAGTGGATGGGCGCCATCCACCCCGGCAGCGGCG

ACACCGCCTACAACCAGCGGTTCAAGGGCCGGGTGACCATCACCGCCGACAAGAGCACCAGC
ACCGCCTACATGGAGCTGAGCAGCCTGCGGAGCGAGGACACCGCCGTGTACTACTGCGCCCG
GTTCTACAGCTACGCCTACTGGGGCCAGGGCACCTGGTGACCGTGAGCGCCGCTAGCACCAAA

(2) The synthesized antibody nucleotide sequence (signal peptide: positions 16-81 (light chain), positions 31-84 (heavy chain)) was inserted into mammalian cell expression vector, so as to construct antibody expression vector containing heavy chain and light chain, respectively, and the sequence of the antibody was identified by sequencing. In particular:

The synthesized light chain and pIK-hu12L (purchased from Shanghai Rui Jin Biotechnology Co., Ltd.) were double digested with EcoRV and BsiWI, and the light chain was inserted into pIK-hu12L.

The synthesized heavy chain and the modified pIH-hu12H (purchased from Shanghai Rui-Jin Biotechnology Co., Ltd., an NheI restriction site was added by synonymous mutation on CH1) were double digested with EcoRV and NheI, and the heavy chain was inserted into pIH-hu12H.

The digested and linked vector was transformed into TOP10 competent cells, and the positive clones were identified by PCR. The correct clones were confirmed by sequencing. The vector was transiently transfected into 293F cells by 293Fectin and expressed. The 293F culture supernatant was collected by centrifugation, filtered through a 0.45 μ m filter and passed through Protein A column for affinity chromatography. A280 was determined by spectrophotometer to determine the concentration of the purified antibody.

5.6 SPR analysis of the binding ability of humanized antibody Y035 to human GPC3

To quantitatively analyze the binding of antibody Y035 to antigen human GPC3, the affinity and kinetic parameters of antibody Y035 were measured by multi-cycle kinetic assay using Biacore T200 system (from GE). Antigen human GPC3 (purchased from Shanghai Rui-Jin Biotechnology Co., Ltd.) was coupled to carboxymethyl dextran surface of sensor chip CM5 through primary amino with NHS / EDC coupling according to the manufacturer's instructions, and the final amount of the conjugated ligand was 305RU. Kinetic measurements were performed in 1 $\times$ HBS-EP + working buffer at 25°C, 30 μ l/min, and regeneration condition was 10 mM Glycine-HCl (pH2.5), 10 μ l/min for 25 seconds. In each round of the testing cycle, antibody Y035 of a certain concentration flowed over the chip surface. The interaction between the antibody and the antigen protein human GPC3 fixed on the chip caused the change of the molecular concentration on the surface of the sensor chip which was measured according to changes in SPR Signal and expressed in resonance units (RU). Time vs resonance unit (RU) was plotted and after RU value of the reference channel

without fixed antigen was subtracted, the resulting sensorgram documents the entire reaction including the binding and dissociation processes. Kinetic analysis of the binding of antibody Y035 and antibody 5A5 to human GPC3 are shown in Figure 7-8. In different cycles of kinetic assay, the concentration of antibody Y035 was 6.25 nM, 12.5 nM, 25 nM, 50 nM and 100 nM, respectively. The resulting curves were evaluated using Biacore T200 evaluation software and the affinity KD values were calculated. Using the same method, the affinity and kinetic parameters of the parental murine monoclonal antibody 5A5 prior to humanization of Y035 were also assayed. The binding data of antibodies Y035 and 5A5 to human GPC3 proteins, respectively, are summarized in Table 2, and humanized antibody Y035 binds human GPC3 protein much better than mouse 5A5 after humanization.

Table 2. Binding kinetic parameter of antibody Y035 and 5A5 to human GPC3

Antibody	ka (1/Ms)	kd (1/s)	KD (M)
Y035	3.96E+05	2.51E-05	6.34E-11
5A5	2.80E+05	2.30E-05	8.19E-11

5.7 FACS analysis of Cell-binding activity of antibody Y035

Antibody Y035 was analyzed for binding ability to GPC3-positive hepatocarcinoma HepG2 cell line (ATCC) by Fluorescence Activated Cell Sorter (FACS) using negative 293T (ATCC) cells expressing GPC3 as a negative control.

Specific method is as follows:

1) inoculating HepG2 cells and 297T in logarithmic growth phase into a 6 cm dish respectively at inoculation cell density of about 90%, and incubating overnight at 37°C in an incubator.

2) digesting cells with 10 mM EDTA, collecting cells through centrifugation at 200 g × 5 mins, and resuspending cells in 1% phosphate buffered saline (NBS PBS) containing calf serum at 1×10^6 to 1×10^7 /mL into a flow-specific tube in an amount of 100 µl per tube.

3) centrifuging at 200g × 5min, and discarding the supernatant.

4) adding antibodies Y035 to be tested into the experimental groups of both cell lines, respectively, while using PBS without antibody as a blank control. The final concentration of antibody is 20 µg/ml. 100 µl was added to each tube, and incubated in an ice bath for 45 minutes.

5). Adding 2 ml of 1% NBS PBS to each tube and centrifuging at 200 g × 5 min for two times.

6) Discarding the supernatant and adding goat anti-human antibody-FITC (Shanghai

Karrie Biotech Co., Ltd.) at a dilution of 1: 100 with 100ul being added to each tube, incubating in an ice bath for 45 minutes.

7). Adding 2 ml of 1% NBS PBS into each tube, centrifuging at $200\text{ g} \times 5\text{ min}$ for two times.

8) Discarding the supernatant, resuspending in 300 ul of 1% NBS PBS and detecting by flow cytometry.

Data was analyzed by flow cytometry data analysis software Flowjo7.6, and the results are shown in Figure 9. Results from flow cytometry show that antibody Y035 can specifically recognize HepG2 cells expressing GPC3, and not bind to GPC3-negative 293T cells.

Example 6. Construction of CAR expression vector based on P7D4 single chain antibody and preparation of lentivirus

By way of example, the vector system used for the lentiviral plasmid vectors constructed below belongs to self-inactivating lentiviral vector system of the third generation, which has 4 plasmids, namely, packaging plasmid pMDLg RRE encoding protein Gag/Pol (from addgene), packaging plasmid pRSV-REV encoding Rev protein (from addgene); envelope plasmid pCMV-VSV-G encoding VSV-G protein (from addgene); and the recombinant expression vector encoding the gene of interest CAR based on empty vector pRRLSIN-cPPT.PGK-GFP.WPRE (from addgene), which can effectively reduce the risk of forming replicable lentivirus particles.

In the present system, the present inventor firstly modified the empty vector pRRLSIN-cPPT.PGK-GFP.WPRE by the conventional technique of molecular cloning, and replaced the original promoter in the vector with the promoter of elongation factor-1 α (EF-1 α) and added MluI restriction site between the promoter and CD8 α sp signal peptide. Specifically, the vector pWPT-EGFP (purchased from Addgene) was double-digested with ClaI/SalI (purchased from NEB), and a 1.1 kb of DNA fragment was recovered and ligated with T4 DNA ligase into ClaI/SalI digested vector pRRLSIN- cPPT.PGK-GFP.WPRE and transformed into host strain TOP10. The positive clones were picked out by colony PCR and confirmed by sequencing. The recombinant plasmid pRRLSIN-cPPT.EF-1 α -EGFP.WPRE was obtained.

Chimeric antigen receptors were prepared using the previously optimized P7D4 scFv. Table 3 explains the connection order of the parts of the chimeric antigen receptor exemplified in the present invention.

Table 3

Chimeric antigen receptor	Extracellular binding region - transmembrane region - intracellular signal region 1 - intracellular signal region 2 and the like	Description
P7D4- δ Z	P7D4 scFv-CD8-CD3 δ zeta	Negative control
P7D4-Z	P7D4 scFv-CD8-CD3 zeta	1 st generation
P7D4-BBZ	P7D4 scFv-CD8-CD137-CD3 zeta	2 nd generation
P7D4-28Z	P7D4 scFv-CD28a-CD28b-CD3 zeta	2 nd generation
P7D4-28BBZ	P7D4 scFv-CD28a-CD28b-CD137-CD3 zeta	3 rd generation

Note: CD28a represents the transmembrane region of CD28 molecule and CD28b represents the intracellular signaling region of CD28 molecule.

1. Amplification of nucleic acid fragments

(1) Amplification of scFv sequences

P7D4 scFv fragment was obtained by PCR using V5-scFv-P7D4-Fc plasmid as a template with a forward primer P7D4fd (SEQ ID NO: 36, ctccacgccgccaggccgcaggtgcagctgcaggag, comprising part of the sequence of CD8 signal peptide) and a reverse primer P7D4re (SEQ ID NO: 37, CGGCGCTGGCGTTCGTGGTACCTAGGACGGTGACCTTGG, comprising part of the sequence of CD8 hinge).

(2) Nucleic acid sequences of other parts of the chimeric antigen receptor

The nucleic acid sequences of other parts of the anti-GPC3 chimeric antigen receptor protein except for P7D4 scFv were respectively obtained by PCR using the sequences SEQ ID NO: 26, 27, 28, 29 and 30 disclosed in Patent Application No. 201310164725.X as templates.

Specifically, the fragment F1 containing the CD8 α sp sequence was obtained by PCR amplification with the primer pair PWXLF (SEQ ID NO: 38, gcaggggaaagaatagtagaca) and PRRL-CD8SP-R1 (SEQ ID NO: 39, CGGCCTGGCGGCGTGAG) using the plasmid pRRLSIN-cPPT.EF-1 α -EGFP.WPRE constructed in this example as a template.

Fragment F3- δ Z containing CD8-CD3 δ zeta (δ Z) was obtained by PCR amplification with the primer pair PRRL-CD8hinge (SEQ ID NO: 40, accacgacgccagcgccg) and δ Zre (SEQ ID NO: 41, GAGGTCGACCTACGCGGGGGCGTCTGCGCTCCTGCTGAACTTCACTCT) using the plasmid SEQ ID NO: 26 in 201310164725.X as a template.

Fragment F3-Z comprising CD8-CD3 zeta(Z), fragment F3-BBZ comprising

CD8-CD137-CD3 zeta(BBZ), fragment F3-28Z comprising CD28a-CD28b-CD3 zeta(28Z) and fragment F3-28BBZ comprising CD28a-CD28b-CD137-CD3 zeta(28BBZ) was obtained by PCR amplification with the primer pair PRRL-CD8hinge (SEQ ID NO: 42, accacgacgccagcgccg) and R3R (SEQ ID NO: 43, aatccagagggttgattgtcgacctagcgagggggcagggcctgc) using the plasmids corresponding to SEQ ID NO: 27, SEQ ID NO: 28, SEQ ID NO: 29 and SEQ ID NO: 30 in the application No. 201310164725.X, respectively.

2. Splicing of nucleic acid fragments

F1 nucleic acid fragment obtained as described above and equimolar P7D4 scFv nucleic acid fragment and equimolar F3- δ Z or F3-Z or F3-BBZ or F3-28Z or F3-28BBZ nucleic acid fragments were subjected to three-segment splicing and PCR as shown in Figure 9 under the following conditions: Pre-denaturation: 94°C for 4 min; denaturation: 94°C for 40 s; annealing: 60°C for 40 s; extension: 68°C for 140 s for 5 cycles and then total extension 68°C for 10 min. DNA polymerase and forward primer PWXLF and reverse primer R3R (reverse primer corresponding to CD8-CD3 δ zeta was δ Z re, and other is R3R) were supplemented, and then PCR was performed for 30 cycles with the amplification conditions: Pre-denaturation: 94°C for 4 min; denaturation: 94°C for 40 s; annealing: 60°C for 40 s; extension: 68°C for 140 s for 30 cycles and then total extension 68°C for 10 min. The fragment consisting of F1 fragment, P7D4 scFv or P7D4 mutant scFv, and various F3 fragments constructed in this Example may be simply referred to as CAR fragment, and the amplified fragments were referred to as:

P7D4 scFv- δ Z (SEQ ID NO: 44),
P7D4 scFv-Z (SEQ ID NO: 45),
P7D4 scFv-BBZ (SEQ ID NO: 46),
P7D4 scFv-28Z (SEQ ID NO: 47),
P7D4 scFv-28BBZ (SEQ ID NO: 48).

3. Construction of lentiviral plasmid vector

In this example, target gene consisting of F1 fragment, P7D4 scFv and F3 fragment components (abbreviated as CAR) was double-digested by MluI and SalI restriction enzymes and ligated into the same double digested pRRLSIN-cPPT.EF-1 α -EGFP.WPRE vector to construct a lentiviral vector expressing each chimeric antigen receptor. The constructed vector was identified by MluI and SalI digestion and sequenced correctly, which was ready for lentivirus packaging. As mentioned above, the relevant CAR gene was transcribed and translated into one peptide chain,

where the anti-GPC3 chimeric antigen receptor will be localized under the guidance of CD8 α signal peptide on the cell membrane.

5 CAR polypeptide sequences were respectively obtained through the above construction, which are named as:

- P7D4- δ Z (SEQ ID NO: 49);
- P7D4-Z (SEQ ID NO: 50);
- P7D4-BBZ (SEQ ID NO: 51);
- P7D4-28Z (SEQ ID NO: 52);
- P7D4-28BBZ (SEQ ID NO: 53).

4. Plasmid-transfected 293T packaging lentivirus

HEK-293T cells (ATCC: CRL-11268) cultured at passage 6 to passage 10 were seeded at a density of 6×10^6 in 10 cm dishes and cultured overnight at 37°C in 5% CO₂ for transfection. The medium was DMEM containing 10% fetal bovine serum (purchased from Life).

Transfection steps are as follows:

4.1 Preparation of liquid A: dissolving 5.2 μ g of each of the desired gene plasmids pRRLSIN-cPPT.EF-1 α -CAR with 6.2 μ g of packaging plasmid pMDLg RRE and pRSV-REV and 2.4 μ g of envelope plasmid pCMV-VSV-G into 800 μ L of serum-free DMEM medium and mixing well.

4.2 Preparation of liquid B: dissolving 60 μ g PEI (polyethylenimine 1 μ g/ μ l, purchased from Polysciences) in 800 μ L serum-free DMEM medium, mixing gently and incubating at room temperature for 5min.

4.3 Formation of transfection complex: adding liquid A into liquid B and gently mixing, vortexing or gently mixing immediately after addition, incubating at room temperature for 20min.

4.4 Adding 1.6 ml of the transfection complex into HEK-293T cells dropwise, and after 4-5 h, changing to DMEM with 2% FBS for transfected 293T cells.

After 72 h of transfection, the virus was collected by filtration using a 0.45 μ m filter (available from Millipore Corporation) and centrifuged at 28,000 rpm using a Beckman Optima L-100XP ultracentrifuge for 2 hours at 4°C. The supernatant was discarded and the resulting pellet was centrifuged at 1/10 ~ 1/50 stock solution of AIM-V (purchased from Life) and resuspend at 100 μ L/tube in -80°C for virus titration or infection of T lymphocytes.

Example 7. T cells infected by Recombinant lentivirus

Human peripheral blood mononuclear cells were obtained from healthy human peripheral

blood by density gradient centrifugation (supplied by Shanghai Blood Center) and added in AIM-V lymphocyte medium (purchased from Invitrogen) at a density of about 2×10^6 / mL and added. The magnetic beads coated with anti-CD3 and CD28 antibodies (Invitrogen) were added in a 1: 1 ratio of cells to magnetic beads, and recombinant human IL-2 (purchased from Shanghai Huaxin Biotechnology Co., Ltd.) at a final concentration of 300U/mL was added for stimulation and culture for 48 h. And then T cells were infected with the above recombinant lentivirus (MOI ≈ 10). The infected cells were passaged every other day at a density of 5×10^5 /mL and recombinant human IL-2 at a final concentration of 300 U / mL was supplemented in the lymphocyte culture medium.

Infected CTL cells were detected by flow cytometry on day 8 of culture for the expression of different chimeric antigen receptors. FACS assays were performed using His-tagged human GPC3 recombinant protein (purchased from Shanghai Rui Jin Biotech Co., Ltd.); the cells were firstly incubated with the protein (50 ug / ml) for 1 hr, washed twice with D-PBS, and Anti-His-tag antibody (1:50 dilution, purchased from Shanghai Reagent Biotechnology Co., Ltd.) was added and incubated for 1 hr and then washed twice with D-PBS and then FITC labeled goat anti-mouse secondary antibody (Shanghai Kangcheng Biotechnology Co., Ltd.), incubated for 50 min, washed with D-PBS for three times, and tested by FACS. Uninfected T lymphocytes was used as a negative control, the positive rates of virus-infected T cells expressing different chimeric antigen receptors are shown in Table 4. The positive rate results show that a certain positive rate of CAR⁺T cells can be obtained by lentivirus infection.

Table 4

T cells transfected by following CARs	Positive rate of T cells transfection
P7D4- δ Z (negative control)	59%
P7D4-Z	65%
P7D4-BBZ	53%
P7D4-28Z	61%
P7D4-28BBZ	54%

T cells were infected with viruses that had different chimeric antigen receptors packaged, respectively, and then subcultured at a cell density of 5×10^5 / ml quaque die alterna, counted, and supplemented with IL-2 (final concentration of 300 U / ml). On the 11th day of culture, about 100 ~ 1000 times of amplification was obtained, indicating that the T cells expressing different chimeric antigen receptors can be expanded in a certain amount *in vitro*, which ensures subsequent

in vitro toxicity tests and *in vivo* experiments.

Example 8. *in vitro* toxicity test of T lymphocytes expressing chimeric antigen receptors

5 *In vitro* toxicity experiments used the following materials:

GPC3-positive hepatoma cells (HepG2 and Huh-7) and GPC3-negative hepatoma cells (SK-HEP-1) as shown in Table 5 were used as target cells and effector cells were CTL cultured for 12 days *in vitro*, which were verified in Example 7 and detected chimeric antigen receptor-expression positive by FACS. Effective target ratios were 3: 1, 1: 1 and 1: 3, respectively.

10 The number of target cells was 10000/well, and effector cells corresponded to different effective target ratio. Each group had 5 replicate wells, average of 5 wells was calculated, and detection time was 18h.

Each experimental group and each control group are listed as follows:

15 Each experimental group: each target cell + CTL expressing different chimeric antigen receptors;

Control group 1: target cells with maximum LDH release;

Control group 2: target cells with spontaneous LDH release;

Control group 3: effector cells with spontaneous LDH release.

20 Detection method: CytoTox 96® Non-Radioactive Cytotoxicity Assay (Promega) is used, which is a colorimetric based assay that can replace 51Cr release assay. CytoTox 96® Assay measures lactate dehydrogenase (LDH) quantitatively. LDH is a stable cytosolic enzyme that is released upon lysis of cells and is released in the same way as radioactive 51Cr is released. The supernatant with released LDH medium can be detected by a 30-minute coupled enzyme reaction in which LDH converts a tetrazolium salt (INT) to a red formazan.

25 The amount of red product produced is proportional to the number of lysed cells. Details can be found in instructions of CytoTox 96 non-radioactive cytotoxicity detection kit.

Cytotoxicity is calculated as:

$$\text{Cytotoxicity \%} = [(\text{experiment group} - \text{control group 2} - \text{control group 3}) / (\text{control group 1} - \text{control group 2})] \times 100$$

30 Specifically, as shown in Table 5, the CAR of the P7D4 single chain antibody of the present invention exhibited significant killing activity on GPC3-positive hepatoma cells, and the second and third generations of P7D4 CAR T cells were slightly more potent than the antitumor activity of the first generation. In addition, all CAR T cells showed no cytotoxic activity on GPC3-negative SK-HEP-1 positive hepatoma cells. These results indicate that

P7D4 can selectively target GPC3-positive hepatoma cells and kill them effectively. In addition, the first, second and third generation of CART expressing P7D4 of the present invention exhibited a effector target ratio gradient dependency, that is, the higher the effector target ratio, the stronger the cytotoxic effects.

5

Table 5. Cytotoxicity of CAR T cells expressing the single chain antibody P7D4

Cytotoxicity (%)	P7D4-28BBZ			P7D4-BBZ			P7D4-28Z			P7D4-Z			P7D4-δZ		
	Different effector target ratio			Different effector target ratio			Different effector target ratio			Different effector target ratio			Different effector target ratio		
	3:1	1:1	1:3	3:1	1:1	1:3	3:1	1:1	1:3	3:1	1:1	1:3	3:1	1:1	1:3
HuH-7	75.3	48.4	17.2	69.2	32.2	10.2	62.4	31.3	7.8	42.9	20.2	5.6	3.2	1.3	5.3
HepG2	85.8	59.1	21.9	71.3	38.6	13.9	68.2	35.6	6.5	48.5	25.9	7.2	4.8	2.5	3.2
SK-HEP-1	2.5	3.2	3.9	4.5	2.1	3.8	1.9	2.4	3.6	2.3	3.7	4.8	3.1	2.3	1.2

Example 9. Preparation and comparison of CAR-T cells based on Y035 and GC33

Two types of CAR-T cells, Y035-BBZ, Y035-28Z and Y035-28BBZ, were prepared based on the light chain variable region and the heavy chain variable region of Y035, respectively, according to the procedures of Examples 6 and 7; and two CAR-T cells, GC33-28Z and GC33-28BBZ, were prepared based on light chain variable region and heavy chain variable region of GC33, as a control. Parts of the chimeric antigen receptor are connected as shown in Table 6.

15

Table 6

Chimeric antigen receptor	Extracellular binding region - transmembrane region - intracellular signal region 1 - intracellular signal region 2 and the like
Y035-BBZ	Y035 scFv-CD8-CD137-CD3 zeta
Y035-28Z	Y035 scFv-CD28a-CD28b-CD3 zeta
Y035-28BBZ	Y035 scFv-CD28a-CD28b-CD137-CD3 zeta
GC33-28Z	GC33 scFv-CD28a-CD28b-CD3 zeta

Wherein nucleotide sequence of Y035-BBZ is show as follows (SEQ ID NO: 85):

gagggtgcagctggtgcagagcggcgccgaggtgaagaagcccggcgccagcgtgaaggtgagctgcaaggccagcggc
 tacaccttcagcgactacgagatgcactgggtgcggcaggccccggccagggcctggagtggatgggcgccatccaccccgga
 gcggcgacaccgcctacaaccagcgggtcaagggccgggtgaccatcaccgccgacaagagcaccagcaccgcctacatggagc
 5 tgagcagcctgcggagcagagacaccgccgtgtactactgcgcccgggttctacagctacgcctactggggccagggcaccctggtg
 accgtgagcgcgggtggaggcgggtcaggcgagggtggttctggcggtggcggtatcgacatcgtgatgaccagacccccctga
 gcctgcccgtgacccccggcgagccccccagcatcagctgcggagcagccagagcctggtgcacagcaacggcaacacctacc
 tgagtggttacctgcagaagcccggccagagccccagctgctgatctacaaggtgagcaaccggttcagcggcggtgcccgaccg
 gttcagcggcgagcggcgagcggcaccgacttcaccctgaagatcagccgggtggaggccgaggacgtgggcgtgtactactgcagc
 10 cagagcatctacgtgccctacaccttcggccagggcaccaagctggagatcaaacgtaccacgacgccagcgcgcgaccacca
 caccggcgcccaccatcgctgcagccccctgtccctgcgcccagaggcgtgcggccagcggcgggggggcgagtgacacg
 aggggggctggacttcgctgtgatactacatctggcgcccttggccgggacttgggggtccttctcctgtcactggttatcaccttt
 actgaaacggggcagaagaagaactcctgtatatattcaaacaccatttatgagaccagtacaaactactcaagaggaagatggctgt
 agctgccgatttccagaagaagaaggaggtgtaactgagagtgaagttcagcaggagcgcagacgccccgcgtacaagc
 15 agggccagaaccagctctataacgagctcaatctaggacgaagagaggagtacgatgttttgacaagagacgtggccgggaccct
 gagatggggggaaagccgagaaggaagaaccctcaggaaggcctgtacaatgaactgcagaagataagatggcgagggcctac
 agtgagattgggatgaaaggcgagcgcggagggggaaggggcacgatggcctttaccaggggtcagtagcaccaccaaggac
 acctacgacgcccttcacatgcaggccctgccccctcgc

amino acid sequence of Y035-BBZ is shown as follows (SEQ ID NO: 86)

EVQLVQSGAEVKKPGASVKVSCKASGYTFS DYEMHWVRQAPGQGLEWMGAIH
 PGSGDTAYNQRFKGRVTITADKSTSTAYMELSSLRSED TAVYYCARFYSYAYWGQGT
 LVTVSAGGGGSGGGGSGGGGSDIVMTQTPLSLPVTPEPASISCRSSQSLVHSNGNTYL
 QWYLQKPGQSPQLLIYKVS NRFSGVPDRFSGSGSTDFTLKISRVEAEDVGVYYCSQSI
 YVPYTFGQGTKLEIKRTTTPAPRPPTPAPTIASQPLSLRPEACRPAAGGAVHTRGLDFAC
 25 DIYIWAPLAGTCGVLLLSLVITLYCKRGRKKLLYIFKQPFMRPVQTTQEEDGCSCRFPE
 EEEGGCELRVKFSRSADAPAYKQGQNQLYNELNLGRREEYDVLDRRGRDPPEMGGK
 PRRKNPQEGLYNELQKDKMAEAYSEIGMKGERRRGKGHDGLYQGLSTATKDTYDAL
 HMQALPPR

nucleotide sequence of Y035-28Z is shown as follows (SEQ ID NO: 87):

gagggtgcagctggtgcagagcggcgccgaggtgaagaagcccggcgccagcgtgaaggtgagctgcaaggccagcggc
 tacaccttcagcgactacgagatgcactgggtgcggcaggccccggccagggcctggagtggatgggcgccatccaccccgga
 gcggcgacaccgcctacaaccagcgggtcaagggccgggtgaccatcaccgccgacaagagcaccagcaccgcctacatggagc
 5 tgagcagcctgcggagcagagacaccgccgtgtactactgcgcccgggttctacagctacgcctactggggccagggcaccctggtg
 accgtgagcgcgggtggaggcgggtcaggcgagggtggttctggcggtggcggtatcgacatcgtgatgaccagacccccctga
 gcctgcccgtgacccccggcgagccccccagcatcagctgcggagcagccagagcctggtgcacagcaacggcaacacctacc
 35 tgagtggttacctgcagaagcccggccagagccccagctgctgatctacaaggtgagcaaccggttcagcggcggtgcccgaccg
 gttcagcggcgagcggcgagcggcaccgacttcaccctgaagatcagccgggtggaggccgaggacgtgggcgtgtactactgcagc
 cagagcatctacgtgccctacaccttcggccagggcaccaagctggagatcaaacgtaccacgacgccagcgcgcgaccacca
 caccggcgcccaccatcgctgcagccccctgtccctgcgcccagaggcgtgcggccagcggcgggggggcgagtgacacg
 40 aggggggctggacttcgctgtgatttttgggtgctggtggtggtggtgagtcctggcttgctatagcttgtagtaacagtggccttat
 tattttctgggtgaggagtaagaggagcaggctcctgcacagtactacatgaacatgactccccgccgccccgggccaaccgca
 gcattaccagccctatgccccaccacgcgacttcgcagcctatcgctccagagtgaagttcagcaggagcgcagacgccccgcgt
 accagcaggggccagaaccagctctataacgagctcaatctaggacgaagagaggagtacgatgttttgacaagagacgtggccgg
 45 gaccctgagatggggggaaagccgcagagaaggaagaaccctcaggaaggcctgtacaatgaactgcagaagataagatggcg
 gaggcctacagtgagattgggatgaaaggcgagcgcggagggggaaggggcacgatggcctttaccaggggtcagtagcaccacca
 ccaaggacacctacgacgcccttcacatgcaggccctgccccctcgc

amino acid sequence of Y035-28Z is shown as follows (SEQ ID NO: 88):

EVQLVQSGAEVKKPGASVKVSCKASGYTFS DYEMHWVRQAPGQGLEWMGAIH
PGSGDTAYNQRFKGRVTITADKSTSTAYMELSSLRSED TAVYYCARFYSYAYWGQGT
LVTVSAGGGGSGGGGSGGGGSDIVMTQTPLSLPVT PGEPA SISRSSQSLVHSNGNTYL
QWYLQKPGQSPQLLIYKVS NRFSGV PDRFSGSGSGTDFTLKISRVEAEDVGVYYCSQSI
5 YVPYTFGQGTKLEIKRTTTPAPRPPTPAPT IASQPLSLRPEACRPAAGGAVHTRGLDFAC
DFWVLVVVGVLACYSLLVTVAFIIFWVRSKR S RLLHSDYMNMTPRRPGPTRKH YQP
YAPPRDFAAYRSRVKFSRSADAPAYQQGQNQLYNELNLGRREEYDVLDKRRGRDPE
MGGKPQRRKNPQEGLYNELQKDKMAEAYSEIGMKGERRRGKGHDGLYQGLSTATK
DTYDALHMQALPPR

10 nucleotide sequence of Y035-28BBZ is shown as follows (SEQ ID NO: 89):

gaggtgcagctggtgcagagcggcgccgaggtgaagaagccccggcgccagcgtgaaggtgagctgcaaggccagcggc
tacaccttcagcgactacgagatgcactgggtgcggcaggccccggccagggcctggagtggtggcgccatccacccccggca
gcggcgacaccgcctacaaccagcgggtcaagggccgggtgacctaccgccgacaagagcaccagcaccgcctacatggagc
tgagcagcctgcggagcgaggacaccgcccgtgtactactgcgcccgggttctacagctacgcctactggggccagggcaccctggtg
15 accgtgagcgccgggtggagggcggttcaggcgggaggtggttctggcggtggcggtatcgacatcgatgacccagacccccctga
gcctggcgtgacccccggcgagccccgcagcatcagctgccggagcagccagagcctggtgcacagcaacgggaacacctacc
tgagtggttacctgcagaagccccggccagagccccagctgctgatctacaaggtgagcaaccgggttcagcggcgtgcccgaccg
gttcagcggcgagcggcgagcggcaccgacttcacccctaagatcagccgggtggaggccgaggacgtggggcgtgtactactgcagc
cagagcatctacgtgccctacaccttcggccagggcaccaagctggagatcaaactaccacgacgccagcgccgcgaccaccaa
20 caccggcgcccaccatcgctgcagcccctgtcctgcgcccagaggcgtgccggccagcggcgggggcgagtgacacg
agggggcgtggacttcgctgtgattttgggtgctggtggtggtggtgagtcctggcttgctatagcttgtagtaacagtggccttat
tattttctgggtgaggagtaagaggagcaggctcctgcacagtgactacatgaacatgactccccgccgccccgggccaacccgcaa
gcattaccagccctatgccccaccacgcgacttcgcagcctatcgctccaaacggggcagaaagaaactcctgtatatattcaacaa
ccatttatgagaccagtacaaactactcaagaggaagatggctgtagctgccgatttcagaagaagaagaaggaggtgtgaactg
25 agagtgaagttcagcaggagcgagacgcccccgctaccagcagggccagaaccagctctataacgagctcaatctaggacgaa
gagaggagtacgatgttttgacaagagacgtggccgggaccctgagatgggggggaaagccgcagagaaggaagaaccctcagg
aaggcctgtacaatgaactgcagaaagataagatggcgaggcctacagtgagattgggatgaaaggcgagcgccggaggggca
aggggcacgatggcctttaccagggtctcagtagccaccaaggacacctacgacgcccttcacatgcaggccctgccccctcgc

amino acid sequence of Y035-28BBZ is shown as follows (SEQ ID NO: 90)

30 EVQLVQSGAEVKKPGASVKVSCKASGYTFS DYEMHWVRQAPGQGLEWMGAIH
PGSGDTAYNQRFKGRVTITADKSTSTAYMELSSLRSED TAVYYCARFYSYAYWGQGT
LVTVSAGGGGSGGGGSGGGGSDIVMTQTPLSLPVT PGEPA SISRSSQSLVHSNGNTYL
QWYLQKPGQSPQLLIYKVS NRFSGV PDRFSGSGSGTDFTLKISRVEAEDVGVYYCSQSI
35 YVPYTFGQGTKLEIKRTTTPAPRPPTPAPT IASQPLSLRPEACRPAAGGAVHTRGLDFAC
DFWVLVVVGVLACYSLLVTVAFIIFWVRSKR S RLLHSDYMNMTPRRPGPTRKH YQP
YAPPRDFAAYRSKRGRKLLYIFKQPFMRPVQTTQEEDGCSCRFPEEEEGGCEL RVKF
SRSADAPAYQQGQNQLYNELNLGRREEYDVLDKRRGRDPEMGGKPQRRKNPQEGLY
NELQKDKMAEAYSEIGMKGERRRGKGHDGLYQGLSTATKDTYDALHMQALPPR

GPC3-positive hepatoma Huh-7 cells were used as target cells and *in vitro* toxicity test
40 was conducted according to the *in vitro* toxicity test method of Example 8. When the effector
target ratio was 1: 3, the killing rates Y035-28Z and Y035-28BBZ on Huh-7 can be over 35%,
the killing rate of GC33-28Z was only 6%, and the killing rate of GC33-28BZZ was only
23%. When the effector target ratio was 1: 1, the killing rate of Y035-28Z on Huh-7 can be
over 60% %, While the killing rate of GC33-28Z was only 22%. The above results show that
45 the cell-killing activity of Y035-CAR T cells of the present application was significantly
better than that of GC33-CAR T cells.

Example 10. Preparation of Bispecific Antibody BiTE

(1) Y035 单链抗体核苷酸序列的制备 Preparation of nucleotide sequence of Y035 single-chain antibody

The hybridoma cell line secreting the monoclonal antibody was cultured to logarithmic growth phase, and cells were counted, and 1×10^7 hybridoma cells were taken. Total RNA was extracted from the cell pellet according to instruction of TRIzol® Plus RNA Purification kit (Invitrogen, 12183-555).

a. cDNA obtained by Reverse transcription

Total RNAs of hybridoma cells were used as a template and cDNA was synthesized by reverse transcription using High capacity RNA to cDNA kit (Invitrogen, 4387406).

b. Amplification of antibody variable region by 5'-RACE method

The cDNA was used as a template and amplified with primers tgcttggtttccaggtgcaagatgtgaggtgcagctggtgcaga (SEQ ID NO: 91) and primer TATCGGATCCACCACCTCCACGTTTGATCTCCAGCTTGGTG (SEQ ID NO: 92) using 5'-Full RACE kit (TAKARA, D315). The PCR product was separated by 1.5% agarose gel electrophoresis, and the obtained PCR product was Y035 single-chain antibody nucleic acid.

(2) Construction of CD3 single chain antibody nucleic acid sequence

PCR was performed with primer (SEQ ID NO: 93, gatatcaaactgcagcagtcag) and primer (SEQ ID NO: 94, GAGAGGGAGTACTCACCCCAAC) using the anti-EpCAM/CD3 bispecific antibody disclosed in 201310025302.X as a template with reference to KOD PLUS Enzyme Manual. The gel was recovered and purified, and the obtained PCR product was a nucleic acid of CD3 single-chain antibody.

(3) Introduction of NheI restriction sites

PCR was performed with primers (ggctaactagagaacccactgc, SEQ ID NO: 95) and PH-R (ACATCTTGACCTGGAAACCAAAGC, SEQ ID NO: 96) using vector PIH as a template, and specific reaction can refer to KOD PLUS Enzyme Manual. Gel was recovered and the PCR product was purified, thereby obtaining a short fragment with NheI restriction site.

(4) Construction of coding sequence nucleic acid of Y035/CD3 bispecific antibody

Overlap PCR was performed using the PCR products obtained in steps (1), (2) and (3), and PCR was performed with primer (SEQ ID NO: 95) and primer (SEQ ID NO: 92). Gel was recovered and the PCR product was purified, thereby obtaining Y035/CD3 bispecific antibody coding sequence (SEQ ID NO: 97) with the following sequence:

gaggtgcagctggtgcagagcggcgccgaggtgaagaagcccgccagcgtaaggtgagctgcaaggccagcggc
tacaccttcagcgactacgagatgcactgggtgcggcaggcccccggccagggcctggagtggatgggcgccatccaccccgca

gctggcgacaccgcctacaaccagcgggtcaagggccgggtgacctaccgcccagaagagcaccagcaccgcctacatggagc
tgagcagcctgcggagcgaggacaccgccgtgtactactgcgcccgggttctacagctacgcctactggggccagggcacccctgggtg
accgtgagcgcgggtggaggcggttcaggcgagggtggttctggcggtggcggtatcgacatcgatgacccagacccccctga
gcctgcccgtgacccccggcgagccccccagcatcagctgccggagcagccagagcctgggtgcacagcaacggcaacacctacc
5 tgcagtgggtacgtgcagaagcccggccagagccccagctgctgatctacaaggtgagcaaccgggttcagcggcgtgcccga ccg
gttcagcggcagcggcagcggcaccgacttcaccctgaagatcagccgggtggaggccgaggacgtgggcgtgtactactgcagc
cagagcatctacgtgccctacaccttcggccagggcaccaagctggagatcaaacgtggaggtggatccgatataaactgcag
cagtcaggggctgaactggcgaagacctggggcctcagtgaaagtgtcctgcaagacttctggctacacctttactaggtacacgatgc
actgggtaaaacagaggcctggacaggggtctggaatggattggataattaatcctagccgtgggtataactaattacaatcagaagttca
10 aggacaaggccacattgactacagacaaatcctccagcacagcctacatgcaactgagcagcctgacatctgaggactctgcagtct
attactgtgcaagatattatgatgatcattactgccttgactactggggccaaggcaccactctcacagtctcctcagtcgaaggtggaa
gtggaggttctgggtggaagtggaggttcaggtggagtcgacgacattcagctgacccagctccagcaatcatgtctgcattctccagg
ggagaaggtcacctgacctgcagagccagttcaagtgttaagtacatgaactggtaccagcagaagtcaggcacctccccaaaaag
atggatttatgacacatccaaagtggcttctggagtccttatcgcttcagtgagggtctgggacctcatactctctcacaatcagc
15 agcatggagggtgaagatgctgccacttattactgccaacagtgaggagtagtaacccgctcacgttcgggtgctgggaccaagctggag
ctgaaa

(5) Expression and Purification of Bispecific Antibodies Targeting Y035 / CD3

Plasmids of Y035 / CD3 BiTE were transiently transfected into eukaryotic cells 293F and cell transfection was performed with reference to 293fectin instruction. The culture
20 supernatant was purified by NiSepharose™ 6 Fast Flow (purchased from GE Healthcare Bio-Science), a metal affinity chromatography column, to give the bispecific antibody of Y035/CD3 (SEQ ID NO: 98) with the following sequence:

EVQLVQSGAEVKKPGASVKVSCKASGYTFSDYEMHWVRQAPGQGLEWMGAIH
PGSGDTAYNQRFKGRVTITADKSTSTAYMELSSLRSEDTAVYYCARFYSYAYWGQGT
25 LVTVSAGGGGSGGGGSGGGGSDIVMTQTPLSLPVTPGEPASISCRSSQSLVHSNGNTYL
QWYLQKPGQSPQLLIYKVSNRFSGVPDRFSGSGSGTDFTLKISRVEAEDVGVYYCSQSI
YVPYTFGQGTKLEIKRGGGSDIKLQQSGAELARPGASVKMSCKTSGYTFTRYTMHW
VKQRPQGQLEWIGYINPSRGYTNYNQKFKDKATLTDDKSSSTAYMQLSSLTSEDSAVY
YCARYYDDHYCLDYWGQGTTLTVSSVEGGSGGSGGSGGSGGVDDIQLTQSPAIMSAS
30 PGEKVTMTCRASSSVSYMNWYQQKSGTSPKRWIYDTSKVASGVPYRFSGSGSGTSYS
LTISSMEAEDAATYYCQWSSNPLTFGAGTKLELK

GPC3-positive HepG2 and SK-Hep-1 GPC3 cells were selected for studying *in vitro* cytotoxicity of Y035/CD3 BiTE. Results shows that the cytotoxicity of Y035/CD3 BiTE at a
concentration of 0.1 ng/ml on HepG2 cells can reach about 60% and the cytotoxicity on
35 SK-Hep-1 GPC3 can reach about 40%. Compared with GPC3/CD3 BiTE described in CN103833852A, cytotoxicity of Y035/CD3 BiTE is significantly increased.

All references mentioned in the present application are incorporated herein by reference, as if each reference was individually incorporated by reference. In addition, it should be
40 understood that after reading the above teachings of the present invention, those skilled in the art can make various modifications or changes to the present invention, and such equivalent forms also fall within the scope of the appended claims of the present application.

Claims

1. An antibody that specifically recognizes glypican-3, wherein the antibody has a light chain variable region and a heavy chain variable region, and

5 CDR1 of the light chain variable region has an amino acid sequence selected from a group consisting of SEQ ID NO: 16, SEQ ID NO: 67, SEQ ID NO: 69;

CDR2 of the light chain variable region has an amino acid sequence selected from a group consisting of SEQ ID NO: 18, SEQ ID NO: 68, SEQ ID NO: 70;

CDR3 of the light chain variable region has the amino acid sequence: SEQ ID NO: 20;

10 CDR1 of the heavy chain variable region has an amino acid sequence selected from a group consisting of SEQ ID NO: 10, SEQ ID NO: 60, SEQ ID NO: 62, SEQ ID NO: 64;

CDR2 of the heavy chain variable region has an amino acid sequence selected from a group consisting of SEQ ID NO: 12, SEQ ID NO: 61, SEQ ID NO: 63, SEQ ID NO: 65, SEQ ID NO: 66, SEQ ID NO: 71;

15 CDR3 of the heavy chain variable region has an amino acid sequence selected from a group consisting of SEQ ID NO: 14, SEQ ID NO: 72.

2. The antibody of claim 1, wherein the antibody comprises:

antibody (a) (P7D4), and its light chain variable region has CDR1 of SEQ ID NO: 16, CDR2 of SEQ ID NO: 18 and CDR3 of SEQ ID NO: 20, or its heavy chain variable region
20 has CDR1 of SEQ ID NO: 10, CDR2 of SEQ ID NO: 12 and CDR3 of SEQ ID NO: 14;

antibody (b) (am4), and its light chain variable region has CDR1 of SEQ ID NO: 16, CDR2 of SEQ ID NO: 18 and CDR3 of SEQ ID NO: 20; or its heavy chain variable region has CDR1 of SEQ ID NO: 60, CDR2 of SEQ ID NO: 61 and CDR3 of SEQ ID NO: 14;

antibody (c) (am14), and its light chain variable region has CDR1 of SEQ ID NO: 16,
25 CDR2 of SEQ ID NO: 18 and CDR3 of SEQ ID NO: 20, or its heavy chain variable region has CDR1 of SEQ ID NO: 62, CDR2 of SEQ ID NO: 63 and CDR3 of SEQ ID NO: 14;

antibody (d) (am20), and its light chain variable region has CDR1 of SEQ ID NO: 16, CDR2 of SEQ ID NO: 18 and CDR3 of SEQ ID NO: 20; or its heavy chain variable region has CDR1 of SEQ ID NO: 64, CDR2 of SEQ ID NO: 65 and CDR3 of SEQ ID NO: 14;

30 antibody (e) (am35), and its light chain variable region has CDR1 of SEQ ID NO: 67, CDR2 of SEQ ID NO: 68 and CDR3 of SEQ ID NO: 20; or its heavy chain variable region has CDR1 of SEQ ID NO: 10, CDR2 of SEQ ID NO: 66 and CDR3 of SEQ ID NO: 14;

antibody (f) (am42), and its light chain variable region has CDR1 of SEQ ID NO: 69, CDR2 of SEQ ID NO: 70 and CDR3 of SEQ ID NO: 20, or its heavy chain variable region

has CDR1 of SEQ ID NO: 10, CDR2 of SEQ ID NO: 66 and CDR3 of SEQ ID NO: 14;

antibody (g) (T2-23), and its light chain variable region has CDR1 of SEQ ID NO: 16, CDR2 of SEQ ID NO: 70 and CDR3 of SEQ ID NO: 20, or its heavy chain variable region has CDR1 of SEQ ID NO: 10, CDR2 of SEQ ID NO: 71 and CDR3 of SEQ ID NO: 72;

5 antibody (h), which recognizes the same antigenic determinant as that recognized by the antibody according to any one of (a) to (g).

3. The antibody of claim 2, wherein the amino acid sequence of the heavy chain variable region of antibody (a) (P7D4) is shown in positions 1-121 of SEQ ID NO: 4; or the amino acid sequence of the light chain variable region of the antibody is shown in positions
10 137-247 of SEQ ID NO: 4;

The amino acid sequence of the heavy chain variable region of antibody (b) (am4) is shown in positions 1-121 of SEQ ID NO: 25; or the amino acid sequence of the light chain variable region of the antibody is shown in positions 137-247 of SEQ ID NO: 25;

The amino acid sequence of the heavy chain variable region of antibody (c) (am14) is
15 shown in positions 1-121 of SEQ ID NO: 27; or the amino acid sequence of the light chain variable region of the antibody is shown in positions 137-247 of SEQ ID NO: 27;

The amino acid sequence of the heavy chain variable region of the antibody (d) (am20) is shown in positions 1-121 of SEQ ID NO: 29; or the amino acid sequence of the light chain variable region of the antibody is shown in positions 137-247 of SEQ ID NO: 29;

20 The amino acid sequence of the heavy chain variable region of the antibody (e) (am35) is shown in positions 1-121 of SEQ ID NO: 31; or the amino acid sequence of the light chain variable region of the antibody is shown in positions 137-247 of SEQ ID NO: 31;

The amino acid sequence of the heavy chain variable region of the antibody (f) (am42) is shown in positions 1-121 of SEQ ID NO: 33; or the amino acid sequence of the light chain
25 variable region of the antibody is shown in positions 137-247 of SEQ ID NO: 33; or

The amino acid sequence of the heavy chain variable region of antibody (g) (T2-23) is shown in positions 1-121 of SEQ ID NO: 35; or the amino acid sequence of the light chain variable region of the antibody is shown in positions 137-247 of SEQ ID NO: 35.

4. An antibody that specifically recognizes glypican-3 (GPC3) is provided, which is a
30 humanized monoclonal antibody comprising a heavy chain variable region and a light chain variable region. Its heavy chain variable region comprises: CDR1 of the amino acid sequence of SEQ ID NO: 73, CDR2 of the amino acid sequence of SEQ ID NO: 74 and CDR3 of the amino acid sequence of SEQ ID NO: 75; or its light chain variable region comprises CDR1 of the amino acid sequence of SEQ ID NO: 76, CDR2 of the amino acid sequence of SEQ ID

NO: 77 and CDR3 of the amino acid sequence of SEQ ID NO: 78.

5. The antibody of claim 4, wherein the antibody has a heavy chain variable region of the amino acid sequence of SEQ ID NO: 58 or a light chain variable region of the amino acid sequence of SEQ ID NO: 59.

5 6. A nucleic acid encoding the antibody of any one of claims 1-5.

7. An expression vector comprising the nucleic acid of claim 6.

8. A host cell comprising the expression vector of claim 7 or having the nucleic acid of claim 6 integrated into its genome.

9. Use of the antibody of any one of claims 1-5 for the preparation of a targeted drug,
10 antibody-drug conjugate, or a polyfunctional antibody that specifically targets tumor cells expressing glypican-3; or

for the preparation of a reagent for diagnosing a tumor expressing Glypican-3; or

for the preparation of chimeric antigen receptor-modified immune cells.

10. A multi-functional immunoconjugate, comprising:

15 the antibody of any one of claims 1-5; and

a functional molecule linked thereto; and the functional molecule is selected from a group consisting of a molecule targeting a tumor surface marker, a tumor-suppressing molecule, a molecule targeting a surface marker of an immune cell, or a detectable label.

11. The multifunctional immunoconjugate of claim 9, wherein the molecule that targets
20 a tumor surface marker is an antibody or ligand that binds to a tumor surface marker; or

the tumor-suppressing molecule is an anti-tumor cytokine or anti-tumor toxin; and preferably, the cytokine includes: IL-12, IL-15, IFN-beta and TNF-alpha.

12. The multifunctional immunoconjugate of claim 10, wherein the detectable label includes a fluorescent label or a chromogenic label.

25 13. The multifunctional immunoconjugate of claim 11, wherein the antibody that binds to a tumor surface marker refers to an antibody that recognizes antigens other than glypican-3, wherein the other antigens include: EGFR, EGFRvIII, mesothelin, HER2, EphA2, Her3, EpCAM, MUC1, MUC16, CEA, Claudin 18.2, folate receptor, Claudin 6, WT1, NY-ESO- 1, MAGE 3, ASGPR1 or CDH16.

30 14. The multifunctional immunoconjugate of claim 10, wherein the molecule that targets the surface marker of the immune cell is an antibody that binds to T cell surface marker, which can form a T-cell-engaging bifunctional antibody with the antibody of any one of claims 1-5.

15. The multifunctional immunoconjugate of claim 14, wherein the antibody that binds

to the immune cell surface marker is an anti-CD3 antibody.

16. The multifunctional immunoconjugate of claim 15, wherein the multifunctional immunoconjugate is a fusion peptide, and further comprises a linker peptide between the antibody of any one of claims 1-5 and the functional molecule linked thereto.

17. A nucleic acid encoding the multi-functional immunoconjugate of any one of claims 10-16.

18. Use of the multifunctional immunoconjugate of any one of claims 10-16, for the preparation of an antineoplastic agent, or

for the preparation of an agent for diagnosis of tumors that express Glypican-3; or

for the preparation of chimeric antigen receptor-modified immune cells; preferably, the immune cells include T lymphocyte, NK cell or NKT lymphocyte.

19. A chimeric antigen receptor comprising the antibody of any one of claims 1-5, and the chimeric antigen receptor comprises: the antibody of any one of claims 1-5, a transmembrane region and an intracellular signal region, which are sequentially linked; and the intracellular signal region is selected from a group consisting of intracellular signal region sequences of CD3 ζ , Fc ϵ RI γ , CD27, CD28, CD137, CD134, MyD88, CD40 or a combination thereof.

20. The chimeric antigen receptor of claim 19, wherein the transmembrane region comprises a transmembrane region of CD8 or CD28.

21. The chimeric antigen receptor of claim 19, wherein the chimeric antigen receptor comprises the following sequentially connected antibody, a transmembrane region and an intracellular signal region:

The antibody of any one of claims 1-5, CD8, and CD3 ζ ;

The antibody of any one of claims 1-5, CD8, CD137 and CD3 ζ ;

The antibody of any one of claims 1-5, the transmembrane region of CD28 molecule, the intracellular signaling region of CD28 molecule and CD3 ζ ; or

The antibody of any one of claims 1-5, the transmembrane region of CD28 molecule, the intracellular signaling region of CD28 molecule, CD137 and CD3 ζ .

22. The chimeric antigen receptor of claim 19, wherein the antibody is a single chain antibody or a domain antibody.

23. The chimeric antigen receptor of claim 19, wherein the chimeric antigen receptor comprises:

SEQ ID NO: 49 or the amino acid sequence shown in positions 22-346 thereof;

SEQ ID NO: 50 or the amino acid sequence shown in positions 22-447 thereof;

SEQ ID NO: 51 or the amino acid sequence shown in positions 22-491 thereof;

SEQ ID NO: 52 or the amino acid sequence shown in positions 22-494 thereof;

SEQ ID NO: 53 or the amino acid sequence shown in positions 22-536 thereof;

The amino acid sequence of SEQ ID NO: 85;

5 The amino acid sequence of SEQ ID NO: 86; or

The amino acid sequence of SEQ ID NO: 87.

24. A nucleic acid encoding the chimeric antigen receptor of any one of claims 19-23.

25. An expression vector comprising the nucleic acid of claim 24.

26. A virus comprising the vector of claim 25.

10 27. Use of the chimeric antigen receptor of any one of claims 19-23, or the nucleic acid of claim 24, or the expression vector of claim 25 or the virus of claim 26 for the preparation of genetically modified immune cells targeting the tumor that expresses Glypican-3.

28. The use of claim 27, wherein the tumor expressing glypican-3 includes (but is not limited to) liver cancer, melanoma, ovarian clear cell carcinoma, yolk sac tumor and
15 neuroblastoma.

29. A genetically modified immune cell, which is transduced with the nucleic acid of claim 24 or the expression vector of claim 25 or the virus of claim 26; or

has the chimeric antigen receptor of any one of claims 19-23 expressed on its surface.

30. The immune cell of claim 29, wherein it further carry an exogenous encoding
20 sequence for cytokines; and preferably, the cytokines include IL-12, IL-15 or IL-21.

31. The immune cell of claim 29, wherein it also expresses another chimeric antigen receptor which does not contain CD3 ζ but contains the intracellular signaling domain of CD28, the intracellular signaling domain of CD137, or a combination of both.

32. The immune cell of claim 29, wherein it further expresses a chemokine receptor;
25 and preferably, the chemokine receptor includes: CCR2.

33. The immune cell of claim 29, wherein it further expresses siRNA which can reduce expression of PD-1 or a protein which blocks PD-L1.

34. The immune cell of claim 29, wherein it further expresses a safety switch; and preferably, the safety switch includes iCaspase-9, Truncated EGFR or RQR8.

30 35. The immune cell of claim 29, wherein the immune cells include T lymphocytes, NK cells or NKT cells.

36. Use of the genetically modified immune cell of any one of claims 29-35 for the preparation of a tumor-inhibiting drug, and the tumor is a tumor expressing glypican-3.

37. A pharmaceutical composition comprising:

The antibody of any one of claims 1-5 or a nucleic acid encoding the antibody; or

The immunoconjugate of any one of claims 10-16 or a nucleic acid encoding the conjugate; or

5 The chimeric antigen receptor of any one of claims 19-23 or a nucleic acid encoding the chimeric antigen receptor; or

The genetically modified immune cell of any one of claims 29-35.