

Description

This is a continuation-in-part of United States Application Serial No. 158,439, filed February 22, 1988.

5 BACKGROUND OF THE INVENTION

United States Patent 4,647,576, which is herein incorporated by reference, discloses certain trans-6-[2-(substituted-pyrrol-1-yl)alkyl]-pyran-2-ones.

10 United States Patent 4,681,893, which is herein incorporated by reference, discloses certain trans-6-[2-(3- or 4-carboxamido-substituted pyrrol-1-yl)alkyl]-4-hydroxy-pyran-2-ones.


The compounds disclosed in the above United States patents are useful as inhibitors of the enzyme 3-hydroxy-3-methylglutaryl-coenzyme A reductase (HMG-CoA reductase) and are thus useful hypolipidemic and hypocholesterolemic agents. Particularly valuable as hypolipidemic and hypocholesterolemic agents are trans(±) 5-(4-fluorophenyl)-2-(1-methylethyl)-N,4-diphenyl-1-[2-(tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)-ethyl]-1H-pyrrole-3-carboxamide and (2R-trans)-5-(4-fluorophenyl)-2-(1-methylethyl)-N,4-diphenyl-1-[2-(tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)ethyl]-1H-pyrrole-3-carboxamide. The aforementioned compounds have been prepared by a linear synthetic route which employed two reactions conducted at low temperatures (-78°C) under carefully controlled conditions. The two reactions included the addition of the dianion of ethyl acetoacetate to an aldehyde and the reduction of the hydroxy ketone produced in this reaction with sodium borohydride and a trialkylborane. Although these reactions provide the target compounds in high diastereomeric excess, they are difficult to conduct on large-scale and use expensive reagents which are difficult to handle. They also do not produce enantiomerically pure products. The materials produced by the earlier methods can be separated into enantiomerically pure products but the process is very expensive, time-consuming, and results in the loss of more than 50% of the starting material.

The object of the present invention is an improved process for preparing the compounds described above by using a novel synthesis.

Further, we have unexpectedly found that the particularly valuable hypolipidemic and hypocholesterolemic agents trans(±)5-(4-fluorophenyl)-2-(1-methylethyl)-N,4-diphenyl-1-[2-(tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)ethyl]-1H-pyrrole-3-carboxamide and (2R-trans)-5-(4-fluorophenyl)-2-(1-methylethyl)-N,4-diphenyl-1-[2-(tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)ethyl]-1H-pyrrole-3-carboxamide can be prepared from a novel intermediate in fewer steps and higher yields than the previous methods. Moreover, the present method proceeds from inexpensive starting materials and is amenable to large-scale synthesis.

35 SUMMARY OF THE INVENTION

Accordingly, a first aspect of the present invention is an improved process for the preparation of a compound of Formula I




50 and a dihydroxy acid and pharmaceutically acceptable salts thereof, corresponding to the opened lactone ring of a compound of Formula I

wherein R₁ is

1-naphthyl,
2-naphthyl,
55 cyclohexyl,
cyclohexylmethyl,
norbornenyl,
phenyl,

- phenyl substituted with
fluorine,
chlorine,
bromine,
5 hydroxyl,
trifluoromethyl,
alkyl of from one to four carbon atoms,
alkoxy of from one to four carbon atoms, or
alkanoyloxy of from two to eight carbon atoms,
10 benzyl,
2-, 3-, or 4-pyridinyl, or
2-, 3-, or 4-pyridinyl-N-oxide;
R₂ or R₃ is independently
hydrogen,
15 alkyl of from one to six carbon atoms,
cyclopropyl,
cyclobutyl,
cyclopentyl,
cyclohexyl,
20 phenyl,
phenyl substituted with
fluorine,
chlorine,
bromine,
25 hydroxyl,
trifluoromethyl,
alkyl of from one to four carbon atoms, or
alkoxy of from one to four carbon atoms, cyano,
trifluoromethyl, or -CONR₅R₆ where R₅ and R₆ are independently
30 hydrogen,
alkyl of from one to six carbon atoms,
phenyl,
phenyl substituted with
fluorine,
35 chlorine,
bromine,
cyano, or
trifluoromethyl;
R₄ is
40 alkyl of from one to six carbon atoms,
cyclopropyl,
cyclobutyl,
cyclopentyl,
cyclohexyl, or
45 trifluoromethyl;
which comprises:
(a) reacting 1,6-heptadien-4-ol with an
(1) alkyl lithium
(2) followed by iodine and carbon dioxide and
50 (3) treating the resulting iodocarbonate intermediate in situ with a base in an aqueous alcohol at about
0 ° C to about 40 ° C to afford a compound of Formula IX;




IX

(b) treating the compound of Formula IX with

(1) an alkali cyanide at about 0°C to about 40°C and

(2) reacting the resulting diol intermediate in situ with a ketal-forming reagent in the presence of an acid to afford a compound of Formula VIII




VIII

wherein R₇ and R₈ are independently hydrogen, alkyl of from one to three carbon atoms, phenyl or R₇ and R₈ taken together as -(CH₂)_n-, wherein n is 4 or 5;

(c) treating the compound of Formula VIII with

(1) ozone in an inert solvent and


(2) reacting the resulting intermediate in situ with oxygen and triphenylphosphine at about -20°C to about -78°C to afford a compound of Formula VII



VII

wherein R₇ and R₈ are as defined above;

(d) treating the compound of Formula VII with an oxidizing reagent at about 0°C to afford a compound of Formula VI




VI

wherein R_7 and R_8 are as defined above;

(e) treating the compound of Formula VI with a compound of Formula

Hal-R_{9a}

wherein Hal is halogen and R_{9a} is alkyl of from one to eight carbon atoms or a three- to six-membered cycloalkyl group, in the presence of a base to afford a compound of Formula V_a


 V_a

wherein R_7 and R_8 , and R_{9a} are as defined above:

or treating the compound of Formula VI with a compound of Formula


HO-R_{9b}

wherein R_{9b} is tertiary butyl, tertiary amyl, or α , α -dimethylbenzyl in the presence of an activating agent, a catalytic amount of a base and an inert solvent to afford a compound of Formula V_b


 V_b

wherein R_7 and R_8 , and R_{9b} are as defined above;


(f) treating the compound of Formula V_a with hydrogen in the presence of a catalyst and an acid at about 0°C to about 70°C to afford a compound of Formula IV_a

IV_a

wherein R₇ and R₈, and R_{9a} are as defined above,
or treating the compound of Formula V_b with hydrogen in the presence of a catalyst and an acid or a
catalyst and a base at about 0°C to about 70°C to afford a compound of Formula IV_b


IV_b

wherein R₇ and R₈, and R_{9b} are as defined above;
(g) treating the compound of Formula IV_a with a compound of Formula III




III

wherein R₁, R₂, R₃, and R₄ are as defined above in an inert solvent to afford a compound of Formula II_a

II_a

wherein R₁, R₂, R₃, R₄, R₇, R₈, and R_{9a} are as defined above, or treating the compound of Formula IV_b
with a compound of Formula III in an inert solvent to afford a compound of Formula II_b

II_b

wherein R₁, R₂, R₃, R₄, R₇, R₈, and R_{9b} are as defined above;

(h) and finally treating a compound of Formula II_a with


- (1) an acid in the presence of an inert solvent
- (2) followed by hydrolysis with a base
- (3) followed by neutralization with an acid and
- (4) dissolution and/or heating in an inert solvent with concomitant removal of water to give a compound of Formula I,

or treating a compound of Formula II_b with

- (1) an acid in the presence of an inert solvent
- (2) followed by addition of a base
- (3) followed by neutralization with an acid and
- (4) dissolution and/or heating in an inert solvent with concomitant removal of water to give a compound of Formula I;


(i) and if desired converting the resulting compound of Formula I to a dihydroxy acid corresponding to the opened lactone ring of structural Formula I by conventional hydrolysis and further, if desired converting the dihydroxy acid to a corresponding pharmaceutically acceptable salt by conventional means, and if so desired converting the corresponding pharmaceutically acceptable salt to a dihydroxy acid by conventional means, and if so desired converting the dihydroxy acid to a compound of Formula I by heating in an inert solvent.

A second aspect of the present invention is an improved process for the preparation of the compound of Formula I_a

I_a


and the hydroxy acid and pharmaceutically acceptable salts thereof, corresponding to the opened lactone ring of the compound of Formula I_a which comprises:

(a) reacting the compound of Formula XVII




XVII I

with a compound of Formula




wherein R_7 and R_8 are independently hydrogen, alkyl of from one to three carbon atoms, phenyl or R_7 and R_8 are taken together as $-(CH_2)_n-$ wherein n is 4 or 5 and R_{13} is hydrogen or



in an inert solvent and treating the resulting intermediate with an acid to afford the compound of Formula I_a .


(b) and if desired, converting the resulting compound of Formula I_a to a hydroxy acid corresponding to the opened lactone ring of structural Formula I_a by conventional hydrolysis and further, if desired, converting the hydroxy acid to a corresponding pharmaceutically acceptable salt by conventional means, and if so desired, converting the hydroxy acid to a compound of Formula I_a by dissolution and/or heating in an inert solvent with concomitant removal of water.

A third aspect of the present invention is an improved process for the preparation of the compound of Formula I_a .




and the hydroxy acid and pharmaceutically acceptable salts thereof, corresponding to the opened lactone ring of the compound of Formula I_a which comprises


(a) reacting 4-methyl-3-oxo-N-phenylpentanamide with benzaldehyde in the presence of a catalyst and an inert solvent to afford the compounds of Formula XVIII




(b) reacting the compounds of Formula XVIII with 4-fluorobenzaldehyde in the presence of a catalyst, a base, and an inert solvent to afford the compound of Formula XVII




(c) reacting the compound of Formula XVII with a compound of Formula



wherein R_{10} and R_{11} are alkyl of one to eight carbon atoms or R_{10} and R_{11} together are




$-\text{CH}_2-\text{CH}_2-$ or $-\text{CH}_2-\text{CH}_2-\text{CH}_2-$ in the presence of a catalyst and an inert solvent to afford a compound of Formula XVI



XVI

wherein R_{10} and R_{11} are alkyl of one to eight carbon atoms or R_{10} and R_{11} together are




$-\text{CH}_2\text{CH}_2-$ or $-\text{CH}_2-\text{CH}_2-\text{CH}_2-$,

(d) and finally converting a compound of Formula XVI in a conventional manner to afford a compound of Formula I_a

(e) and if desired, converting the resulting compound of Formula I_a to a hydroxy acid corresponding to the opened lactone ring of structural Formula I_a by conventional hydrolysis and further, if desired, converting the hydroxy acid to a corresponding pharmaceutically acceptable salt by conventional means, and if so desired, converting the hydroxy acid to a compound of Formula I_a by dissolution and/or heating in an inert solvent with concomitant removal of water.

A fourth aspect of the present invention is a novel intermediate of Formula II




II

wherein R_9 is alkyl of from one to eight carbon atoms, a three to six-membered cycloalkyl group or α,α -dimethylbenzyl and R_1, R_2, R_3, R_4, R_7 , and R_8 are as defined above, which is useful in the preparation of inhibitors of cholesterol biosynthesis of Formula I.


Preferred are compounds of formula II, wherein R_7 and R_8 are methyl, R_9 is tertiary butyl, and the two optically active centers are R and compounds wherein R_9 is alkyl of from one to eight carbon atoms, a three to six-membered cycloalkyl group, or α,α -dimethylbenzyl.

A fifth aspect of the present invention is a novel intermediate of Formula XVI




XVI

wherein R_{10} and R_{11} are alkyl of one to eight carbon atoms or R_{10} and R_{11} together are



or $-CH_2-CH_2-CH_2-$ which is useful in the preparation of the inhibitor of cholesterol biosynthesis of Formula I_a.


A sixth aspect of the present invention is a novel intermediate of Formula IV



IV

wherein R_7 , R_8 , and R_9 are as defined above, which is useful in the preparation of a compound of Formula II, which in turn is useful in the preparation of inhibitors of cholesterol biosynthesis of Formula I.


A seventh aspect of the present invention is a novel intermediate of Formula XXI



XXI

wherein R_7 and R_8 are independently hydrogen, alkyl of from one to three carbon atoms, phenyl or R_7 and R_8 are taken together as $-(\text{CH}_2)_n$, wherein n is 4 or 5, which is useful in the preparation of the inhibitor of cholesterol biosynthesis of Formula Ia.


An eighth aspect of the present invention is a novel intermediate of Formula V



V

wherein R_7 , R_8 , and R_9 are as defined above, which is useful in the preparation of a compound of Formula IV, which in turn is useful in the preparation of a compound of Formula II, which in turn is useful in the preparation of inhibitors of cholesterol biosynthesis of Formula I.


A ninth aspect of the present invention is a novel intermediate of Formula VI



VI

wherein R_7 and R_8 are as defined above, which is useful in the preparation of a compound of Formula V, which in turn is useful in the preparation of a compound of Formula IV, which in turn is useful in the preparation of inhibitors of cholesterol biosynthesis of Formula I.


A tenth aspect of the present invention is a novel intermediate of Formula VII



VII

wherein R_7 and R_8 are as defined above, which is useful in the preparation of a compound of Formula VI, which in turn is useful in the preparation of a compound of Formula V, which in turn is useful in the preparation of a compound of Formula IV, which in turn is useful in the preparation of inhibitors of cholesterol biosynthesis of Formula I.


An eleventh aspect of the present invention is a novel intermediate of Formula VIII



VIII

wherein R_7 and R_8 are as defined above, which is useful in the preparation of a compound of Formula VII, which in turn is useful in the preparation of a compound of Formula VI, which in turn is useful in the preparation of a compound of Formula V, which in turn is useful in the preparation of inhibitors of cholesterol biosynthesis of Formula I.


A twelfth aspect of the present invention is the novel intermediate of Formula XVII



XVII

which is useful in the preparation of a compound of Formula XVI, which in turn is useful in the preparation of the inhibitor of cholesterol biosynthesis of Formula Ia.

A thirteenth aspect of the present invention is the novel intermediates of Formula XVIII



XVIII

which are useful in the preparation of the compound of Formula XVII, which, in turn, is useful in the preparation of a compound of Formula XVI, which, in turn, is useful in the preparation of the inhibitor of cholesterol biosynthesis of Formula Ia.

DETAILED DESCRIPTION OF THE INVENTION

In this invention, the term "alkyl" means a straight or branched hydrocarbon group having from one to eight carbon atoms and includes, for example, methyl, ethyl, *n*-propyl, isopropyl, *n*-butyl, isobutyl, tertiary-butyl, *n*-pentyl, tertiary-amyl, *n*-hexyl, *n*-heptyl, *n*-octyl, and the like.

"Cycloalkyl" refers to a three- to six-membered saturated hydrocarbon ring and includes, for example, cyclobutyl, cyclopentyl, cyclohexyl, and the like.

"Alkoxy" is O-alkyl in which alkyl is as defined above.

"Alkanoyloxy" is an alkyl group, as defined above, attached to a carbonyl group and thence, through an oxygen atom, to the parent molecular residue.

"Carboalkoxy" is an alkyl group, as defined above, attached to an oxygen atom and thence, through a carbonyl group, to the parent molecular residue.

"Norbornenyl" is a group derived by the removal of a hydrogen atom (other than at a bridgehead carbon atom) from bicyclo[2.2.1]hept-2-ene.

"Halogen" is iodine, bromine, and chlorine.

"Alkali metal" is a metal in Group IA of the periodic table and includes, for example, lithium, sodium, potassium, and the like.

"Alkaline-earth metal" is a metal in Group IIA of the periodic table and includes, for example, calcium, barium, strontium, and the like.

"Noble metal" is platinum, palladium, rhodium, ruthenium, and the like.

A preferred compound of Formula I prepared by the improved process of the present invention is one wherein R₁ is 1-naphthyl, norbornenyl, phenyl, or phenyl substituted with fluorine, chlorine,

- 5 bromine, hydroxyl, trifluoromethyl, alkyl of from one to four carbon atoms, alkoxy of from one to four carbon atoms, or
10 alkanoyloxy of from two to eight carbon atoms.

Also preferred is a compound of Formula I prepared by the improved process of the present invention wherein R₄ is alkyl of from one to six carbon atoms, cyclopropyl, or trifluoromethyl.

Particularly preferred compounds of Formula I prepared by the improved process of the present invention are the following:

- 15 trans-6-[2-[2-(4-fluorophenyl)-5-(trifluoromethyl)-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one;
trans-6[2-[2-(4-fluorophenyl)-5-methyl-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one;
trans-6-[2-[2-(4-fluorophenyl)-5-(1-methylethyl)-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one;
20 trans-6-[2-[2-cyclopropyl-5-(4-fluorophenyl)-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one;
trans-6-[2-[2-(1,1-dimethylethyl)-5-(4-fluorophenyl)-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one;
trans-tetrahydro-4-hydroxy-6-[2-[2-(2-methoxyphenyl)-5-methyl-1H-pyrrol-1-yl]ethyl]-2H-2-one;
trans-tetrahydro-4-hydroxy-6-[2-[2-(2-methoxyphenyl)-5-(1-methylethyl)-1H-pyrrol-1-yl]ethyl]-2H-pyran-2-one;
25 one;
trans-tetrahydro-4-hydroxy-6-[2-[2-methyl-5-(1-naphthalenyl)-1H-pyrrol-1-yl]ethyl]-2H-pyran-2-one;
trans-6-[2-(2-bicyclo[2.2.1]hept-5-en-2-yl-5-methyl-1H-pyrrol-1-yl)ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one;
trans(±)-5-(4-fluorophenyl)-2-(1-methylethyl)-N,4-diphenyl-1-[2-(tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)ethyl]-1H-pyrrole-3-carboxamide;
30 (2R)-trans-5-(4-fluorophenyl)-2-(1-methylethyl)-N,4-diphenyl-1-[2-(tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)ethyl]-1H-pyrrole-3-carboxamide;
trans-2-(4-fluorophenyl)-N,4-diphenyl-1-[2-(tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)ethyl]-5-trifluoromethyl-1H-pyrrole-3-carboxamide;
35 trans-5-(4-fluorophenyl)-N,4-diphenyl-1-[2-(tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)ethyl]-2-trifluoromethyl-1H-pyrrole-3-carboxamide; and

a dihydroxy acid and pharmaceutically acceptable salts thereof, corresponding to the opened lactone ring of compounds of structural Formula I.

- As previously described, the compounds of Formula I are useful as inhibitors of the enzyme 3-hydroxy-40 3-methylglutaryl-coenzyme A reductase (HMG CoA reductase) and are thus useful as hypolipidemic or hypocholesterolemic agents.

The process of the present invention in its first aspect is a new, improved, economical, and commercially feasible method for preparing HMG CoA reductase inhibitors of Formula 1. The process of the present invention in its first aspect is outlined in Scheme 1:

SCHEME I

5

10

15

20

25

30


35

40


45

50

55



SCHEME I (continued)



A compound of Formula IX is prepared from the known 1,6-heptadien-4-ol (XI) using the methodology described by Bongini, A., et al, *Journal of Organic Chemistry*, **47**, pp 4626-4633 (1982) and Majewski, M., et al, *Tetrahedron Letters*, **25**, pp 2101-2104 (1984). Thus, the homoallylic alcohol (XI) is reacted with an alkyl lithium such as, for example, *n*-butyllithium followed by iodine and carbon dioxide to give the iodocarbonate (X) at -35°C to -20°C which is not isolated but treated in situ with a base such as an alkali or alkaline-earth metal hydroxide or carbonate, for example, sodium hydroxide, potassium hydroxide, calcium hydroxide, sodium carbonate, potassium carbonate, calcium carbonate, and the like, in an aqueous alcohol of from one to three carbon atoms such as, for example, methanol, ethanol, isopropanol, and the like, at about 0°C to about 40°C , to afford the epoxide of Formula IX. Preferably the reaction is carried out with potassium carbonate in aqueous methanol at about 0°C to about 40°C , preferably 0°C . The epoxide ring of Formula IX is opened with either potassium or sodium cyanide in an aqueous alcohol such as, for example, methanol, ethanol, tertiary butanol, isopropanol, and the like, at about 0°C to about 40°C . Preferably the reaction is carried out with potassium cyanide in aqueous isopropanol at about 25°C . The resulting diol intermediate is not isolated but treated in situ with a ketal forming reagent such as, for example, acetone, dimethoxypropane, 2-methoxypropene, benzaldehyde, cyclopentane, cyclohexanone, 1,1-dimethoxycyclopentane, 1,1-dimethoxycyclohexane, and the like, in the presence of an acid such as, for example, camphorsulfonic acid, *para*-toluenesulfonic acid, and the like, in the presence of excess reagent or in an inert solvent such as, for example, dichloromethane, and the like, at about 0°C to the reflux temperature of

the reagent or solvent to afford a compound of Formula VIII, wherein R_7 and R_8 are independently hydrogen, alkyl of from one to three carbon atoms, phenyl or R_7 and R_8 are taken together as $-(CH_2)_n-$, wherein n is 4 or 5. A compound of Formula VIII is treated with ozone in an inert solvent such as, for example, dichloromethane and the like, and the resulting intermediate ozonide which is not isolated is flushed in situ with oxygen to remove the ozone and then treated with triphenylphosphine or dimethyl sulfide at about -20°C to about -78°C , preferably about -78°C , to afford a compound of Formula VII, wherein R_7 and R_8 are as defined above. A compound of Formula VII is treated with an oxidizing reagent such as, for example, chromium trioxide-sulfuric acid-water, and the like, at about 0°C to afford a compound of Formula VI, wherein R_7 and R_8 are as defined above. A compound of Formula VI is treated with a compound of Formula




wherein Hal is halogen such as, for example, iodine, chlorine, bromine, and R_{9a} is alkyl of from one to eight carbon atoms, or a three- to six-membered cycloalkyl group, preferably isopropyl, isobutyl, and the like in the presence of a base such as, for example, 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU), and the like to afford a compound of Formula V_a , wherein R_7 , R_8 , and R_{9a} are as defined above. Additionally, treating the compound of Formula VI with a compound of Formula



wherein R_{9b} is tertiary butyl, tertiary amyl, or α,α -dimethylbenzyl in the presence of an activating agent such as, for example, dicyclohexylcarbodiimide, 1,1'-carbonyldiimidazole and the like in the presence of a base such as, for example, 4-dimethylaminopyridine and the like in an inert solvent such as, for example, dichloromethane, tetrahydrofuran, and the like to afford a compound of Formula V_b , wherein R_7 , R_8 , and R_{9b} are as defined above. A compound of Formula V_a is treated with hydrogen in the presence of a catalyst such as a noble metal, for example, platinum, palladium, rhodium, ruthenium, derivatives thereof, and the like, or Raney nickel, preferably platinum dioxide, and an acid, such as, for example, acetic acid, propanoic acid and the like, preferably acetic acid, at about 0°C to about 70°C and about 1×10^5 to about 7×10^5 Pa (14 to 100 psi) pressure to afford a compound of Formula IV_a , wherein R_7 , R_8 , and R_{9a} are as defined above. Additionally, a compound of Formula V_b is treated with hydrogen in the presence of a catalyst such as a noble metal, for example, platinum, palladium, rhodium, ruthenium, derivatives thereof, and the like, and an acid such as, for example, acetic acid, propanoic acid and the like, or a catalyst such as, for example, Raney nickel, Raney cobalt and the like, in an inert solvent such as, for example, methanol, ethanol, isopropanol, tetrahydrofuran and the like, saturated with anhydrous ammonia or saturated with aqueous ammonium hydroxide or aqueous sodium hydroxide, preferably the reaction is carried out with Raney nickel in methanol saturated with anhydrous ammonia at about 0°C to about 70°C and about 1×10^5 Pa to about 7×10^5 Pa (about 14 to about 100 psi) pressure to afford a compound formula IV_b , wherein R_7 , R_8 , and R_{9b} are as defined above. Raney nickel and Raney cobalt as described above are finely divided forms of nickel and cobalt.

A compound of Formula XXI is prepared by treating a compound of Formula VI




VI

wherein R_7 and R_8 are as defined above with hydrogen in the presence of a catalyst such as, for example, Raney nickel, Raney cobalt and the like in finely divided form, in an inert solvent such as, for example, methanol, ethanol, isopropanol, tetrahydrofuran and the like, saturated with anhydrous ammonia or saturated with aqueous ammonium hydroxide solution or a catalyst such as, for example, platinum, palladium and the like, in an inert solvent such as, for example, methanol, ethanol, isopropanol, tetrahydrofuran and the like in


the presence of an acid such as, for example, acetic acid, propanoic acid and the like, at about 0°C to about 70°C and about 1×10^5 to about 7×10^5 Pa (about 14 to about 100 psi) pressure to afford a compound of Formula XXI.

A compound of Formula XXIII is prepared from a compound of Formula XXV



XXV

wherein R_7 and R_8 are as defined above using the methodology previously described for preparing a compound of Formula XXI from a compound of Formula VI. A compound of Formula XXV is prepared by treating a compound of Formula VI




VI

wherein R_7 and R_8 are as defined above and tertiary butyl alcohol with a coupling reagent such as, for example, dicyclohexylcarbodiimide and the like in the presence of a base such as, for example, 4-dimethylaminopyridine and the like in an inert solvent such as, for example dichloromethane and the like to afford a compound of Formula XXV.

An optically active (R) compound of Formula XXIII_a is prepared as outlined in Scheme II. The starting material (R)-4-cyano-3-[[1,1-dimethylethyl]dimethylsilyl]oxy]butanoic acid of Formula XXIX is synthesized starting from isoascorbic acid using syntheses well known to practitioners of the art. This chemistry is identical to that disclosed in United States Patent 4,611,067 (Merck & Co. Inc.) using ascorbic acid which is herein incorporated by reference which produces (S)-4-cyano-3-[[1,1-dimethylethyl]dimethylsilyl]oxy]butanoic acid.¹

¹U.S. Patent 4,611,067 apparently incorrectly assigned the configuration R to the product of this sequence of reactions starting with ascorbic acid.

SCHEME II



Thus, the optically active compounds are prepared from the known isoascorbic acid using the methodology described by Volante R. P. et al, in United States Patent 4,611,067 but in that case starting with ascorbic acid. This establishes the optically active centers desired in Formula XXVa and Formula XXIIIa as R. Thus, the (R)-4-cyano-3-[(1,1-dimethylethyl)dimethylsilyl]oxybutanoic acid of Formula XXIX is treated with carbonyldiimidazole in tetrahydrofuran at 0°C to -40°C, preferably -20°C, warmed to 25°C and the activated acid derivative is not isolated but the solution is added to a suspension of a salt of 1,1-dimethylethyl malonic acid such as, for example, the potassium salt of 1,1-dimethylethyl malonic acid (half ester, half salt) anhydrous magnesium chloride, and an amine such as, for example, diisopropylethylamine in acetonitrile at -10°C to 20°C preferably at 5°C. The mixture is poured into a mixture of 1N hydrochloric acid and ethyl acetate to afford the (R)-1,1-dimethylethyl 6-cyano-5-[(1,1-dimethylethyl)dimethylsilyl]oxy-3-oxohexanoate of Formula XXVIII. The ketone of Formula XXVIII is treated with fluoride ion at 0°C to 65°C, preferably 25°C to afford the (R)-1,1-dimethylethyl 6-cyano-5-hydroxy-3-oxo-hexanoate of Formula XXVII. The ketone of Formula XXVII is treated with triethylborane and air (or methoxydiethylborane without air) followed by sodium borohydride and methanol in tetrahydrofuran at -78°C to -110°C, preferably -100°C to afford [R-(R*,R*)]-1,1-dimethylethyl 6-cyano-3,5-hydroxyhexanoate of Formula XXVI. The diol of Formula XXVI is treated with a ketal forming reagent such as, for example, acetone, dimethoxypropane, 2-methoxypropene, benzaldehyde, cyclopentanone, cyclohexanone, 1,1-dimethoxycyclopentane, 1,1-dimethoxycyclohexane and the like, in the presence of an acid such as, for example, camphorsulfonic acid, para-toluenesulfonic acid, and the like, in the presence of excess reagent or in an inert solvent such as, for example, dichloromethane, and the like, at 0°C to the reflux temperature of the reagent or solvent to afford a compound of Formula XXVa wherein R_7 and R_8 are independently hydrogen, alkyl of from one to three carbon atoms, phenyl or R_7 and R_8 are taken together as $-(\text{CH}_2)_n-$, wherein n is 4 or 5.


A compound of Formula XXVa is treated with hydrogen gas in an alcohol such as methanol saturated with anhydrous ammonia or aqueous ammonium hydroxide in the presence of a catalyst such as Raney nickel or Raney cobalt or with a noble metal catalyst such as platinum oxide in the presence of an alkanolic

acid such as acetic acid to afford a compound of Formula XXIII_a, wherein R₇ and R₈ as defined above.

Additionally, an optically active compound of Formula IV_a or Formula IV_b may be prepared starting from the optically active epoxide of Formula IX. The preparation of the optically active epoxide of Formula IX is described by Kocienski, P. J., et al, Journal of the Chemical Society Perkin Transaction I, pp 2183-2187 (1987).

The process of preparing a compound of Formula I is outlined in Scheme III:

SCHEME III



A compound of Formula IV_a or Formula IV_b is reacted with a compound of Formula III, wherein R₁ is

- 1-naphthyl,
- 2-naphthyl,
- 5 cyclohexyl,
- cyclohexylmethyl,
- norbornenyl,
- phenyl,
- phenyl substituted with
- 10 fluorine,
- chlorine,
- bromine,
- hydroxyl,
- trifluoromethyl,
- 15 alkyl of from one to four carbon atoms,
- alkoxy of from one to four carbon atoms, or
- alkanoyloxy of from two to eight carbon atoms,
- benzyl,
- 2-, 3-, or 4-pyridinyl, or
- 20 2-, 3-, or 4-pyridinyl-N-oxide;

R₂ or R₃ is independently

- hydrogen
- alkyl of from one to six carbon atoms,
- cyclopropyl,
- 25 cyclobutyl,
- cyclopentyl,
- cyclohexyl,
- phenyl,
- phenyl substituted with
- 30 fluorine,
- chlorine,
- bromine,
- hydroxyl,
- trifluoromethyl,
- 35 alkyl of from one to four carbon atoms, or
- alkoxy of from one to four carbon atoms, cyano,
- trifluoromethyl or -CONR₅R₆ where R₅ and R₆ are independently
- hydrogen,
- alkyl of from one to six carbon atoms,
- 40 phenyl,
- phenyl substituted with
- fluorine,
- chlorine,
- bromine,
- 45 cyano, or
- trifluoromethyl;

R₄ is

- alkyl of from one to six carbon atoms,
- cyclopropyl,
- 50 cyclobutyl,
- cyclopentyl,
- cyclohexyl, or
- trifluoromethyl;

in an inert solvent such as, for example, toluene, and the like, at the reflux temperature of the solvent to give a compound of Formula II_a, or Formula II_b, wherein R₁, R₂, R₃, R₄, R₇, R₈, R_{9a} and R_{9b} are as defined above. Finally, a compound of Formula II_a is treated with an acid such as, for example, aqueous hydrochloric acid, and the like, in an inert solvent such as, for example, tetrahydrofuran, followed by hydrolysis with a base such as, for example sodium hydroxide. The reaction is neutralized with an acid

such as, for example, aqueous hydrochloric acid and dissolved and/or heated in an inert solvent such as, for example, toluene, and the like, with concomitant removal of water to give a compound of Formula I, wherein R_1 , R_2 , R_3 , and R_4 are as defined above.

15 Additionally, a compound of Formula II_b is treated with an acid such as, for example, aqueous hydrochloric acid, and the like, in an inert solvent such as, for example, tetrahydrofuran and the like for about 15 hours, followed by the addition of a base such as, for example, sodium hydroxide and the like and stirred for about 30 hours. The reaction is reacidified with an acid such as, for example, aqueous hydrochloric acid and dissolved and/or heated in an inert solvent such as, for example, toluene and the like, with concomitant removal of water to give a compound of Formula I, wherein R_1 , R_2 , R_3 , and R_4 are as defined above.

20 The process of the present invention in its second aspect is a new, improved, economical, and commercially feasible method for preparing the HMG CoA reductase inhibitor of Formula I_a. The process of the present invention in its second aspect is outlined in Scheme IV:

15

20

25

30

35

40

45


50

55

SCHEME IV


5

10




XXI

15




XVII

20




XXIII

+




XVII

25



XXII

30



XXIV


35

40

45


50

55



Ia


The compound of Formula XVII is reacted with a compound of Formula XXI



XXI

wherein R_7 and R_8 are independently hydrogen, alkyl of from one to three carbon atoms, phenyl or R_7 and R_8 are taken together as $-(\text{CH}_2)_n-$ wherein n is 4 or 5 in the presence of an inert solvent such as, for example, dimethyl sulfoxide and the like for about 15 hours at about 105°C with the removal of water to afford the intermediate derivative (XXII). Preferably the reaction is carried out in dimethyl sulfoxide for about 15 hours at about 105°C . The intermediate derivative XXII, which is not isolated, is treated with an acid such as, for example, concentrated hydrochloric acid and the like in an inert solvent such as, for example, ethyl acetate and the like to afford the compound of Formula I_a. Preferably, the reaction is carried out with concentrated hydrochloric acid in ethyl acetate.


Additionally, the compound of Formula XVII is reacted with a compound of Formula XXIII



XXIII


wherein R_7 and R_8 are independently hydrogen, alkyl of from one to three carbon atoms, phenyl or R_7 and R_8 are taken together as $-(\text{CH}_2)_n-$ wherein n is 4 or 5 in the presence of an inert solvent or solvents such as, for example, hexane, toluene and the like for about 24 hours at about the reflux temperature of the solvent or solvents. The intermediate derivative XXIV, which is not isolated, is treated with an acid such as, for example, a 10% aqueous solution of hydrochloric acid for about 15 hours, followed by the addition of a base such as, for example, sodium hydroxide and the like and reacidification with an acid for about 30 hours to afford the compound of Formula I_a.

The process of the present invention in its third aspect is a new, improved, economical, and commercially feasible method for preparing the HMG CoA reductase inhibitor of Formula I_a. The process of the present invention in its third aspect is outlined in Scheme V:




4-Methyl-3-oxo-N-phenylpentanamide (XIX) is obtained by heating a mixture of methyl 4-methyl-3-oxopentanoate (XX), aniline and ethylene diamine in toluene. 4-Methyl-3-oxo-N-phenylpentanamide (XIX) is subsequently reacted with benzaldehyde in the presence of a catalyst such as, for example, piperidine and glacial acetic acid, ethylene diamine and glacial acetic acid, β -alanine and glacial acetic acid, and the like in an inert solvent such as, for example, toluene, heptane, hexane, and the like for about 24 to about 36 hours at about 60° to about 120° C with the removal of water to afford 4-methyl-3-oxo-N-phenyl-2-(phenylmethylene)pentanamide (XVIII). Preferably the reaction is carried out with β -alanine and glacial acetic acid at reflux for about 24 hours in hexane. The 4-methyl-3-oxo-N-phenyl-2-(phenylmethylene)pentanamide (XVIII) is reacted with 4-fluorobenzaldehyde in the presence of a catalyst such as, for example, 3-benzyl-5-(2-hydroxyethyl)-4-methylthiazolium chloride, 3,4-dimethyl-5-(2-hydroxyethyl)thiazolium iodide, 3-ethyl-5-(2-

hydroxyethyl)-4-methylthiazolium bromide, thiamine hydrochloride, and the like, and a base such as, for example, N,N-diisopropylethylamine, pyridine, N,N-dimethylaniline, triethylamine, 1,8-diazabicyclo[5.4.0]-undec-7-ene (DBU), 1,4-diazabicyclo[2.2.2]octane (DABCO), 4-dimethylaminopyridine (DMAP), N,N,N',N'-tetramethylethylenediamine (TMEDA) and the like, either neat or in the presence of a solvent such as, for example, tetrahydrofuran, tertiary-butyl methyl ether, ethanol, dimethylformamide, dimethylsulfoxide, N-methylpyrrolidone, acetonitrile, methylisobutyl ketone, ethyl acetate, isopropanol, pyridine and the like for about 20 to about 30 hours under anhydrous conditions at about room temperature to about the reflux temperature of the solvent to afford the compound of Formula XVII. Preferably the reaction is carried out in the presence of 3-ethyl-5-(2-hydroxyethyl)-4-methylthiazolium bromide and triethylamine in ethanol at about 75-80 °C for about 24 hours. The compound of Formula XVII is reacted with a compound of Formula



wherein R₁₀ and R₁₁ are alkyl of one to eight carbon atoms or R₁₀ and R₁₁ together are




-CH₂CH₂- or -CH₂CH₂CH₂- in the presence of a catalyst of Formula



wherein R₁₂ is CH₃, CF₃, ClCH₂-, C₆H₅CH₂CH₂-, C₆H₅CH₂-, HO₂CCH₂-, HO₂CCH₂CH₂-, C₆H₅-, para-Cl-C₆H₄-, ClCH₂CH₂-, meta-H₃C-C₆H₄-, para-H₃C-C₆H₄-, tertiary-C₄H₉- or triethylamine hydrochloride and a solvent or mixtures thereof such as, for example, tetrahydrofuran, hexane, toluene, ethanol, tertiary-butyl acetate, ethyl acetate, 1,2-dichloroethane, isopropanol, dimethyl sulfoxide and the like for about 24 to about 48 hours at about 5 °C to about the reflux temperature of the solvent with the removal of water to afford a compound of Formula XVI. Preferably, the reaction is carried out in the presence of pivalic acid and a mixture of toluene and heptane at reflux for about 48 hours with the removal of water. A compound of Formula XVI is converted in a conventional manner using the methodology disclosed in United States Patent 4,681,893 to the compound of Formula Ia.

Certain of the compounds of Formula III are either known or capable of being prepared by methods known in the art. The ring-opened dihydroxy acids of formula



wherein R₁, R₂, R₃, and R₄ are as defined above may be prepared from the lactone compounds of Formula I by conventional hydrolysis such as, for example, sodium hydroxide in methanol, sodium hydroxide in tetrahydrofuran-water, and the like, of the lactone compounds of Formula I.

The ring-opened dihydroxy acid of Formula XII may be produced from the lactone compound of Formula Ia by conventional hydrolysis of the lactone compound of Formula Ia.

In the ring-opened dihydroxy acid form, compounds of the present invention react to form salts with pharmaceutically acceptable metal and amine cations formed from organic and inorganic bases. The term

"pharmaceutically acceptable metal salt" contemplates salts formed with the sodium, potassium, calcium, magnesium, aluminum, iron, and zinc ions. The term "pharmaceutically acceptable amine salt" contemplates salts with ammonia and organic nitrogenous bases strong enough to form salts with carboxylic acids. Bases useful for the formation of pharmaceutically acceptable nontoxic base addition salts of the compound of the present invention form a class whose limits are readily understood by those skilled in the art.

The dihydroxy free acid form of the compounds of the invention may be regenerated from the salt form, if desired, by contacting the salt with a dilute aqueous solution of an acid such as hydrochloric acid.

The ring closed lactone form of the compounds of the invention may be regenerated by dissolution of the dihydroxy acid form of the compounds of the invention in an inert solvent such as, for example, toluene, benzene, ethyl acetate, and the like, at about 0°C to about the boiling point of the solvent usually but not necessarily with concomitant removal of the resulting water and usually but not necessarily with strong acid catalysis such as, for example, concentrated hydrochloric acid and the like.

The base addition salts may differ from the free acid forms of the compounds of this invention in such physical characteristics as solubility and melting point, but are otherwise considered equivalent to the free acid form for the purposes of this invention.

The compounds of the present invention may exist in solvated or unsolvated form and such forms are equivalent to the unsolvated form for the purposes of this invention.

The compounds of structural Formulas I, I_a, and XII above possesses two asymmetric carbon centers, one at the 4-hydroxy position of the pyran-2-one ring, and the other at the 6-position of the pyran-2-one ring where the alkylpyrrole group is attached. This asymmetry gives rise to four possible isomers, two of which are the 4R,6S and 4S,6R-isomers and the other two of which are the 4R,6R and 4S,6S-isomers. The preferred isomer in this invention is the 4R,6R-isomer of the compounds of Formulas I, I_a and XII above.

The following nonlimiting examples illustrates the inventors' preferred method for preparing the compounds of the invention.

EXAMPLE 1

Trans-6-[2-[2-ethyl-5-(4-fluorophenyl)-1H-pyrrol-1-yl]-ethyl]-tetrahydro-4-hydroxy-2H-pyran-2-one.

Step A: Preparation of (R*,R*)-α-2-propenyloxirane-ethanol.

n-Butyllithium, 129 mL (200 mmol), is added dropwise to a 0°C solution of 1,6-heptadien-4-ol, 22.4 g (0.2 mol), in 200 mL of anhydrous tetrahydrofuran until the triphenylmethane indicator turned red. Carbon dioxide is then bubbled in for 30 minutes (lecture bottle carbon dioxide passed through drierite) and the light yellow solution is stirred for 30 minutes under a balloon of carbon dioxide. To this solution is added iodine, 101.4 g (0.4 mol), dissolved in ~200 mL of anhydrous tetrahydrofuran over 60 minutes. The mixture is allowed to warm to room temperature overnight, diluted with ethyl acetate, washed with 10% sodium bisulfite solution, saturated solution of sodium bicarbonate, brine, and dried (magnesium sulfate). The crude product is dissolved in 200 mL of methanol and 20 mL of water, cooled to 0°C and 0.5 g of solid potassium carbonate is added. The mixture is vigorously stirred for six hours, filtered, concentrated, and partitioned between ethyl acetate and brine. After extracting the aqueous layer 2x with ethyl acetate, the combined organics are washed with brine and dried (magnesium sulfate). Flash chromatography (4:1 hexane-ethyl acetate) provides, after concentration in vacuo, 18 g of (R*,R*)-α-2-propenyloxiraneethanol. 200 MHz NMR (CDCl₃) δ 1.5-1.65 (m, 1H), 1.90 (dt, 1H, J = 15, 4 Hz), 2.2 (m, 3H), 2.53 (m, 1H), 2.79 (m, 1H), 3.12 (m, 1H), 3.94 (m, 1H), 5.19 (m, 2H), 5.80 (m, 1H). IR (film) 3400, 3077, 2980, 2925, 1643, 1412, 1260, 918, 827 cm⁻¹.

Step B: Preparation of (±)-cis-2,2-dimethyl-6-(2-propenyl)-1,3-dioxane-4-acetonitrile.

5 Potassium cyanide, 1.3 g (20 mmol), is added to a room temperature solution of (R*,R*)-α-2-propenyloxiraneethanol, 2.56 g (20 mmol), in 25 mL of 4:1 isopropanol-water. The solution is stirred for 20 hours at ambient temperature, concentrated, and partitioned between ethyl acetate and brine. The aqueous layer is extracted 2x with ethyl acetate and the combined ethyl acetate extracts are washed with brine and dried
10 (magnesium sulfate). The crude product is dissolved in 20 mL of 2,2-dimethoxypropane, camphorsulfonic acid is added and the solution stirred for 18 hours at room temperature. Concentration and flash chromatography provides 1.30 g of (±)-cis-2,2-dimethyl-6-(2-propenyl)-1,3-dioxane-4-acetonitrile. 200 MHz NMR (CDCl₃) δ 1.35 (m, 1H), 1.40 (s, 3H), 1.45 (s, 3H), 1.67 (m, 1H), 2.20 (m, 1H), 2.33 (m, 1H), 2.50 (m, 2H), 3.89 (m, 1H), 4.10 (m, 1H), 5.10 (m, 2H), 5.7-5.9 (m, 1H).
15 IR (film) 2995, 2944, 2255, 1644, 1334 cm⁻¹.

Step C: Preparation of (±)-cis-6-(2-oxoethyl)-2,2-dimethyl-1,3-dioxane-4-acetonitrile.

20 A solution of (±)-cis-2,2-dimethyl-6-(2-propenyl)-1,3-dioxane-4-acetonitrile, 3 g (15.36 mmol), in 100 mL of dichloromethane is cooled to -78 °C under nitrogen. Ozone (Welsbach generator, flow rate 0.1, voltage = 90V) is then passed through a fritted gas inlet tube into the solution until the blue color of ozone appears.
25 The current is turned off, and oxygen bubbled through until the blue color is discharged. Triphenylphosphine, 4.2 g (16 mmol), is added and the colorless solution allowed to warm to room temperature. Flash chromatography provides, after concentration in vacuo, 2.5 g of pure (±)-cis-6-(2-oxoethyl)-2,2-dimethyl-1,3-dioxane-4-acetonitrile. 200 MHz NMR (CDCl₃) δ 1.30 (m, 1H), 1.39 (s, 3H), 1.48 (s, 3H), 1.78 (m, 1H), 2.46-2.75 (m, 4H), 4.2 (m, 1H), 4.40 (m, 1H), 9.79 (t, 1H, J = 1.6 Hz).
30 IR (film) 2250, 1720 cm⁻¹.

Step D: Preparation of (±)-cis-6-(cyanomethyl)-2,2-dimethyl-1,3-dioxane-4-acetic acid.

35 Jones reagent (chromium trioxide-sulfuric acid-water), 3.8 mL (7.6 mmol), is added dropwise to a 0 °C solution of (±)-cis-6-(2-oxoethyl)-2,2-dimethyl-1,3-dioxane-4-acetonitrile, 1.50 g (7.6 mmol), dissolved in 50 mL of acetone until the orange color is not discharged. After stirring a further 15 minutes, the mixture is poured into 300 mL of diethyl ether and washed with brine until the aqueous washes are colorless. The diethyl ether layer is dried (magnesium sulfate), filtered, and concentrated to provide 1.2 g of the acid which solidifies on standing. Trituration with isopropyl ether provides (±)-cis-6-(cyanomethyl)-2,2-dimethyl-1,3-dioxane-4-acetic acid as a colorless solid; mp 92-95 °C. A second trituration/recrystallization from isopropyl
45 ether provides material of mp 98-103 °C. 200 MHz NMR (CDCl₃) δ 1.35 (m, 1H), 1.42 (s, 3H), 1.49 (s, 3H), 1.82 (m, 1H), 2.4-2.7 (m, 4H), 4.18 (m, 1H), 4.35 (m, 1H). ¹³C-NMR(d₆-acetone, 50 MHz) δ 19.95, 24.91, 30.17, 35.88, 41.34, 65.79, 66.35, 99.70, 117.77, 171.83. IR (KBr) broad OH 3500-2400, 2254, 1711, 940 cm⁻¹.
50

Step E: Preparation of (±)-cis-1-methylethyl 6-(cyanomethyl)-2,2-dimethyl-1,3-dioxane-4-acetate.

55 To a solution of (±)-cis-6-(cyanomethyl)-2,2-dimethyl-1,3-dioxane-4-acetic acid, 0.6 g (3 mmol), in acetonitrile, 2 mL, is added 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU), 0.45 mL (3 mmol), and 2-iodopropane, 0.33 mL (3.3 mmol). The solution is stirred overnight at room temperature, diluted with diethyl

ether, washed with brine, and dried (magnesium sulfate). Flash chromatography provides 0.55 g of ($\pm$)-cis-1-methylethyl 6-(cyanomethyl)-2,2-dimethyl-1,3-dioxane-4-acetate.

90 MHz NMR (CDCl_3) δ 1.22 (d, 6H, J = 7 Hz), 1.3 (m, 1H), 1.35 (s, 3H), 1.40 (s, 3H), 1.75 (m, 1H), 2.2-2.7 (m, 4H), 3.9-4.4 (m, 2H), 4.95 (septet, 1H, J = 7 Hz).

5

Step F: Preparation of ($\pm$)-cis-1-methylethyl 6-(2-aminoethyl)-2,2-dimethyl-1,3-dioxane-4-acetate.

10

A mixture of ($\pm$)-cis-1-methylethyl 6-(cyanomethyl)-2,2-dimethyl-1,3-dioxane-4-acetate, 0.55 g, in glacial acetic acid is hydrogenated with platinum dioxide at 3.45×10^5 Pa (50 psi). Concentration dilution with ethyl acetate and bicarbonate wash, followed by washing with brine and drying provides 250 mg of ($\pm$)-cis-1-methylethyl 6-(2-aminoethyl)-2,2-dimethyl-1,3-dioxane-4-acetate. MS 260.1, 244.1 200 MHz NMR (CDCl_3) δ 1.25 (d, 6H, J = 7 Hz), 1.32 (m, 1H), 1.36 (s, 3H), 1.45 (s, 3H), 1.60 (m, 1H), 2.33 (dd, 1H, J = 15, 6 Hz), 2.49 (dd, 1H, J = 15, 6 Hz), 2.85 (br t, 2H, J = 6 Hz), 3.40 (br s, 2H), 4.00 (m, 1H), 4.29 (m, 1H), 12.03 (septet, 1H, J = 7 Hz). IR (film) 1734, 1387 cm^{-1} .

15

20

Step G: Preparation of ($\pm$)-cis-1-methylethyl 6-[2-[2-ethyl-5-(4-fluorophenyl)-1H-pyrrol-1-yl]ethyl]-2,2-dimethyl-1,3-dioxane-4-acetate.

25

A solution of ($\pm$)-cis-1-methylethyl 6-(2-aminoethyl)-2,2-dimethyl-1,3-dioxane-4-acetate, 0.15 g (0.58 mmol), and 1-(4-fluorophenyl)-1,4-hexanedione (Example A), 0.125 g (0.6 mmol), in 5 mL of toluene is stirred and heated at reflux overnight. The cooled solution is concentrated and the highly UV active pyrrole is separated from starting material by preparative thin layer chromatography, eluting 2x with 4:1 hexane-ethyl acetate. This provides 130 mg of pure ($\pm$)-cis-1-methylethyl 6-[2-[2-ethyl-5-(4-fluorophenyl)-1H-pyrrol-1-yl]ethyl]-2,2-dimethyl-1,3-dioxane-4-acetate.

30

200 MHz NMR (CDCl_3) δ 1.51 (m, 1H), 1.23 (d, 6H, J = 6 Hz), 1.33 (m, 9H), 1.5-1.6 (m, 3H), 2.27 (dd, 1H, J = 15.3, 6 Hz), 2.44 (dd, 1H, J = 15.3, 6 Hz), 2.66 (g, 2H, J = 7.5 Hz), 3.62 (m, 1H), 3.8-4.2 (m, 3H), 5.03 (septet, 1H, J = 6 Hz), 5.97 (d, 1H, J = 3.5 Hz), 6.11 (d, 1H, J = 3.5 Hz), 7.0-7.4 (m, 4H).

35

A solution of ($\pm$)-cis-1-methylethyl 6-[2-[2-ethyl-5-(4-fluorophenyl)-1H-pyrrol-1-yl]ethyl]-2,2-dimethyl-1,3-dioxane-4-acetate, 0.13 g (0.3 mmol), in 12 mL of 1:2 2M hydrochloric acid-tetrahydrofuran is stirred overnight at room temperature. To this is added sufficient 2M sodium hydroxide to bring the pH to 10. Stirring is continued for 30 minutes, water, 30 mL, is added and the mixture is extracted with diethyl ether. The aqueous layer is acidified with ice cold 6N hydrochloric acid and extracted with ethyl acetate (2x). The organic layer is washed with brine and dried (magnesium sulfate). The residue which remains on filtration and concentration is dissolved in toluene (50 mL) and heated at reflux with azeotropic removal of water (6 hours). The cooled solution is concentrated and flash chromatographed to provide 60 mg of trans-6-[2-[2-ethyl-5-(4-fluorophenyl)-1H-pyrrol-1-yl]ethyl]-tetrahydro-4-hydroxy-2H-pyran-2-one (elution with 2:1 hexane-ethyl acetate).

40

90 MHz NMR (CDCl_3) δ 1.25 (d, 6H, J = 7 Hz) 1.3-1.8 (m, 4H), 2.3 (br s, 1H, -OH), 2.55 (m, 2H), 2.65 (q, 2H, J = 7Hz), 3.9-4.5 (m, 4H), 5.90 (d, 1H, J = 3.5 Hz), 6.05 (d, 1H, J = 3.5 Hz), 6.9-7.4 (m, 4H).

45

50

55

EXAMPLE 2

Trans-(±)-5-(4-fluorophenyl)-2-(1-methylethyl)-N,4-
diphenyl-1-[2-(tetrahydro-4-hydroxy-6-oxo-2H-pyran-
2-yl)ethyl]-1H-pyrrole-3-carboxamide.

METHOD A

Step A: Preparation of 4-Methyl-3-oxo-N-phenyl-2-(phenylmethylene)pentanamide.

- 15 A suspension of 100 kg of 4-methyl-3-oxo-N-phenylpentanamide (Example B) in 660 kg of hexanes is treated with agitation under nitrogen with 8 kg of β -alanine, 47 kg of benzaldehyde, and 13 kg of glacial acetic acid. The resulting suspension is heated to reflux with removal of water for 20 hours. An additional 396 kg of hexanes and 3 kg of glacial acetic acid is added and reflux continued with water removal for one hour. The reaction mixture is cooled to 20-25 °C and the product is isolated by filtration. The product is
- 20 purified by slurrying in hexanes at 50-60 °C, cooling, and filtration. The product is slurried twice with water at 20-25 °C, filtered, and dried in vacuo to yield 110 kg of 4-methyl-3-oxo-N-phenyl-2-(phenylmethylene)-pentanamide, mp 143.7-154.4 °C.
- Vapor Phase Chromatography (VPC): 30 meter DB 5 capillary column 50-270 °C at 15 °C/min. 19.33 min, 99.7% (area).
- 25 Gas Chromatography/Mass Spectrometry (GC/MC): M/Z 293 [M]⁺.
 Nuclear Magnetic Resonance (NMR): (CDCl₃) δ 1.16 (6H, d), 3.30 (1H, quin.), 7.09 (1H, m), 7.28 (5H, m), 7.49 (5H, m), 8.01 (1H, brs).

Step B: Preparation of (±)4-Fluoro- α -[2-methyl-1-oxopropyl]- γ -oxo-N, β -diphenylbenzenebutaneamide mixture of [R-(R*,R*)], [R-(R*,S*)], [S-(R*,R*)] and [S-(R*,S*)] isomers.

- 30 A solution of 17.5 kg of 3-ethyl-5-(2-hydroxyethyl)-4-methylthiazolium bromide in 300 L of anhydrous ethanol is concentrated by distillation of 275 L of the ethanol. Under an argon atmosphere, 100 kg (340 mol) of 4-methyl-3-oxo-N-phenyl-2-(phenylmethylene)pentanamide, 47.5 L (340 mol) of triethylamine, and 40 L (375 mol) of 4-fluorobenzaldehyde are added. The resulting solution is stirred and heated at 75-80 °C for 23 hours. The product begins to form as a solid after approximately 1.5 hours but approximately 24 hours is required for essentially complete conversion. The slurry is dissolved in 600 L of isopropanol at 80 °C. The resulting solution is slowly cooled and the (±)4-fluoro- α -[2-methyl-1-oxopropyl]- γ -oxo-N, β -diphenylbenzenebutaneamide mixture of [R-(R*,R*)], [R-(R*,S*)], [S-(R*,R*)] and [S-(R*,S*)] isomers isolated by filtration.
- 40 Washing the precipitate with isopropanol and drying in vacuo yielded 99 kg of (±)4-fluoro- α -[2-methyl-1-oxopropyl]- γ -oxo-N, β -diphenylbenzenebutaneamide mixture of [R-(R*,R*)], [R-(R*,S*)], [S-(R*,R*)] and [S-(R*,S*)] isomers; mp 206.8-207.6 °C.
- NMR: (CDCl₃) δ 1.03 (3H, d), 1.22 (3H, d), 2.98 (1H, quin.), 4.91 (1H, d, J = 11 Hz), 5.51 (1H, d, J = 11 Hz), 6.98-7.43 (12H, m), 8.17 (2H, dd), 9.41 (1H, brs).
- 45 High Pressure Liquid Chromatography (HPLC): (Acetonitrile:tetrahydrofuran:water) (40:25:55) Econosil C₁₈ 5 μ 25 cm 1.0 mL/min. 254 nm 16.77 min. 99.2% (area).

Step C: Preparation of 1-(3,3-Diethoxypropyl)-5-(4-fluorophenyl)-2-(1-methylethyl)-N,4-diphenyl-1H-pyrrole-3-carboxamide.

- 50 To a nitrogen purged flask equipped with a mechanical stirrer is added 130 kg (311 mol) of (±)4-fluoro- α -[2-methyl-1-oxopropyl]- γ -oxo-N, β -diphenylbenzenebutaneamide mixture of [R-(R*,R*)], [R-(R*,S*)], [S-(R*,R*)] and [S-(R*,S*)] isomers, 540 L of heptanes and 60 L of toluene, 59 kg (400 mol) of 3-amino-1,1-diethoxypropane, and 22.3 kg (218 mol) of pivalic acid. The mixture is stirred and heated to reflux, removing water with a Dean Stark trap. The mixture is refluxed 32 hours and slowly cooled to 60-65 °C, diluted with 500 L of 2-propanol-water (3:2), seeded, and cooled to 20-25 °C. The product is isolated by filtration, washed with 300 L of 2-propanol, and dried in vacuo to yield 133.5 kg of 1-(3,3-diethoxypropyl)-5-(4-fluorophenyl)-2-(1-methylethyl)-N,4-diphenyl-1H-pyrrole-3-carboxamide; mp 125.1-127.7 °C after recrystal-

lization from ethanol.

HPLC: (acetonitrile:tetrahydrofuran:water) (40:25:55) 1.5 mL/min 254 nm Econosil C₁₈ 5 μ 25 cm R.T. = 37.70 min 99.4% (area)

NMR: ((CD₃)₂CO) δ 1.04 (6H, m, t), 1.47 (6H, d), 1.82 (2H, m), 3.40 (5H, m), 3.99 (2H, m), 4.43 (1H, brt), 6.90-7.50 (14H, m), 8.26 (1H, brs)

In a process analogous to Step C using appropriate starting materials, the following compounds of Formula XVI are prepared:

1-(3,3-Dimethoxypropyl)-5-(4-fluorophenyl)-2-(1-methylethyl)-N,4-diphenyl-1H-pyrrole-3-carboxamide; mp 167-168.2 °C.

5-(4-Fluorophenyl)-1-[2-(4-methyl-1,3-dioxolan-2-yl)ethyl]-2-(1-methylethyl)-N,4-diphenyl-1H-pyrrole-3-carboxamide; bp 141.5-145.9 °C.

Step D: Preparation of 5-(4-Fluorophenyl)-2-(1-methylethyl)-1-(3-oxopropyl)-N,4-diphenyl-1H-pyrrole-3-carboxamide.

To a nitrogen purged flask fitted with an overhead stirrer, a thermometer, and a condenser is added 20 kg (37.8 mol) of 1-(3,3-diethoxypropyl)-5-(4-fluorophenyl)-2-(1-methylethyl)-N,4-diphenyl-1H-pyrrole-3-carboxamide along with 200 L of acetone. The solution is stirred and 100 L of 2N hydrochloric acid solution is added. The mixture is heated to reflux for four hours then cooled to 50 °C $\pm$ 5 °C, seeded, and cooled to 0 °C $\pm$ 5 °C. The product is collected by filtration, washed with 100 L 2-propanol-water (1:1) and dried in vacuo at 50 °C for 64 hours to yield 16.2 kg of 5-(4-fluorophenyl)-2-(1-methylethyl)-1-(3-oxopropyl)-N,4-diphenyl-1H-pyrrole-3-carboxamide as an off-white solid.

Step E: Preparation of 2-(4-Fluorophenyl)-6-hydroxy-5-(1-methylethyl)- β -oxo-3-phenyl-4-[(phenyl-amino)carbonyl]-1H-pyrrole-1-heptanoic acid, methyl ester.

A 22-L three-necked flask fitted with an overhead stirrer, a low temperature thermometer, and a 2-L calibrated addition funnel is dried with a nitrogen purge and 78 g (1.95 mol) 60% sodium hydride in mineral oil is added, followed by 248 mL (1.76 mol) diisopropylamine and 8 L tetrahydrofuran. The reaction is cooled to -10 ° to 0 °C, a significant nitrogen purge was introduced through the flask, and 212 mL (1.92 mol) methyl acetoacetate is added in a slow stream over a 10 to 30-minute period. Stirring is continued at -10 ° to 10 °C for an additional 10 to 30 minutes. After cooling to -15 to -5 °C, 2.2 L of 1.6 M *n*-butyllithium in hexanes is added over a 30 to 60-minute period while maintaining the temperature below 0 °C. Stirring is continued for an additional 1 to 1.5 hours at -15 ° to 0 °C and the mixture cooled to -35 ° to -15 °C.

In a separate 5-L flask, 0.7 kg (1.54 mol) 5-(4-fluorophenyl)-2-(1-methylethyl)-1-(3-oxopropyl)-N,4-diphenyl-1H-pyrrole-3-carboxamide is dissolved with 2.0 L of dry tetrahydrofuran, cooled to 0 ° to -5 °C, and added to the anion solution over a 30 to 45 minute period. The reaction is stirred at -20 ° to -15 °C for 30 to 45 minutes, then quenched by the addition of 4 L aqueous 2 N hydrochloric acid solution over 5 to 15 seconds while stirring rapidly. After stopping the agitation, the lower layer is separated and the remaining organic layer washed with 4 L saturated aqueous sodium chloride.

Step F: Preparation of cis-(4-Fluorophenyl)- β , δ -dihydroxy-5-(1-methylethyl)-3-phenyl-4-[(phenylamino)carbonyl]-1H-pyrrole-1-heptanoic acid, methyl ester.

The reaction solution obtained from Step E contained in a 50-L jacketed glass still is concentrated by vacuum distillation to a thick oil, then dissolved with 19 L tetrahydrofuran and cooled to 0°C under an air atmosphere. Triethylborane, a 1 molar solution in hexanes, (3.20 L, 1.4 equivalents based on Step E) is added over a 10 minute period, the atmosphere on the flask is switched to nitrogen, and the flask cooled to -105°C over 3.5 hours. During this period two liters of methanol are added when the temperature reaches -67°C. Powdered sodium borohydride (184 g, 4.8 mol) is added in 20 to 50-g portions over 1.5 hours, and the reaction maintained between -95° and -106°C for 13 hours, then between -60° and -100°C for 10 hours. Unreacted sodium borohydride is quenched by the addition of 750 mL (12.7 mol) acetic acid in 50-mL portions over a 45 minute period with a substantial amount of gas evolution and with a temperature rise from -60° to -40°C. Further quenching is accomplished by the addition of a solution of 1.0 L 30% hydrogen peroxide (9.7 mol), 3.0 L water, and 100 g dihydrogen sodium phosphate over a 15 minute period and is accompanied by a temperature rise from -40° to 0°C. The reaction is allowed to warm to room temperature overnight, then the lower layer separated off and the upper layer washed with 4.0 L of saturated aqueous sodium chloride solution.

Variation of Step F: Preparation of cis-2-(4-Fluorophenyl)- β , δ -dihydroxy-5-(1-methylethyl)-3-phenyl-4-(phenylamino)carbonyl-1H-pyrrole-1-heptanoic acid, methyl ester.

The reaction solution obtained from Step E is concentrated under vacuum to a volume of 5 to 8 L, then dissolved in 20 L tetrahydrofuran and 4 L methanol under a nitrogen atmosphere. This solution is cooled to -85°C and 2.1 L of a 15% solution of methoxydiethylborane in tetrahydrofuran (2.1 mol, 1.0 equivalent based on Step E) is added. The reaction is cooled to -97°C over one hour and 144 g (3.78 mol) of sodium borohydride added in 20 to 50-g portions over 1.5 hours. The reaction is maintained between -93° and -97°C for 13 hours and allowed to warm to room temperature and stand for 60 hours under a nitrogen atmosphere.

The reaction is quenched by the addition of 460 mL (7.9 mol) acetic acid and concentrated by vacuum distillation to an oil. The residue is dissolved with 8 L methanol, concentrated by vacuum distillation, redissolved with 8 L methanol, and reconstituted by vacuum distillation to a volume of 6 L. The solution is diluted with 8 L tetrahydrofuran, 4 L hexanes, and carried into the next step.

Step G: Preparation of trans-($\pm$)-5-(4-Fluorophenyl)-2-(1-methylethyl)-N,4-diphenyl-1-[2-(tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)ethyl]-1H-pyrrole-3-carboxamide.

The crude reaction mixture of Step F is cooled to 8°C, 8.0 L of 2.0 N aqueous sodium hydroxide is added, and the reaction stirred for 2 hours at 15° to 18°C. The reaction is diluted with 12 L water and the upper layer removed. The remaining aqueous layer is washed with 8 L hexanes then 8 L ethyl acetate is added followed by 1 L concentrated aqueous hydrochloric acid solution. The well-stirred mixture is allowed to separate, the lower layer discarded, and the upper layer washed four times with 4 L each of 2 N aqueous hydrochloric acid solution.

The ethyl acetate layer is concentrated to a foamy syrup by vacuum distillation, and the residue dissolved in 8 L toluene. The toluene is concentrated by vacuum distillation to a volume of 6 L, then allowed to stand at room temperature for 16 hours. The resulting thick slurry is filtered on a Buchner funnel, washed with 1 L of cool toluene, washed with 2 L of hexanes, and dried in a vacuum oven for 24 hours at room temperature, resulting in 686 g of trans-(±)-5-(4-fluorophenyl)-2-(1-methylethyl)-N,4-diphenyl-1-[2-(tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)ethyl]-1H-pyrrole-3-carboxamide. The filtrates are washed with 2 N aqueous hydrochloric acid solution and concentrated in vacuo to a volume of 2 L, then allowed to stand at room temperature for three days. The resulting solid is filtered, washed, and dried as before, resulting in 157 g of trans-(±)-5-(4-fluorophenyl)-2-(1-methylethyl)-N,4-diphenyl-1-[2-(tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)ethyl]-1H-pyrrole-3-carboxamide. HPLC of the solids indicates 95% trans with 1.3% cis lactone for the first crop and 95% trans with 2.3% cis lactone for the second crop. The two crops of solid are dissolved in 8 L ethyl acetate by heating to 50 to 60 °C, then filtered through a Buchner funnel along with 8 L of hexanes which has been warmed to 50 °C. The solution is allowed to cool to room temperature over 16 hours, the resulting slurry filtered through a Buchner funnel, and the solid washed with 2 L hexanes. The resulting solid is dried in a vacuum oven for 24 hours at room temperature, resulting in 720 g of trans-(±)-5-(4-fluorophenyl)-2-(1-methylethyl)-N,4-diphenyl-1-[2-(tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)ethyl]-1H-pyrrole-3-carboxamide with a 98%:0.9% trans:cis HPLC assay. The second crop obtained by concentration as before is approximately 100 g.

METHOD B

Step A: Preparation of (±)-cis-6-(2-aminoethyl)-2,2-dimethyl-1,3-dioxane-4-acetic acid.

A solution of 1.04 g (4.88 mmol) of (±)-cis-6-(cyanomethyl)-2,2-dimethyl-1,3-dioxane-4-acetic acid in 100 mL of methanol saturated with anhydrous ammonia is added to a Parr shaker bottle containing 0.53 g of water wet Raney nickel #30. The solution is heated at 45 °C and 3.45 x 10⁵ Pa (50 psig) hydrogen pressure for 17 hours. The suspension is cooled and filtered to remove the Raney nickel through filter aid and the precipitate washed with methanol. The filtrate is concentrated at reduced pressure. The residue is dissolved in methanol saturated with anhydrous ammonia treated with decolorizing charcoal, filtered through filter aid and evaporated to give 0.56 g of (±)-cis-6-(2-aminoethyl)-2,2-dimethyl-1,3-dioxane-4-acetic acid, mp 165 °C with decomposition at 169 °C.

Fourier Transform Infrared (FTIR) (KBr): 1201.1, 1399.2, 1561.2, 2924.4, 3569.9 cm⁻¹.

¹H-NMR (D₂O, 200 MHz) δ 0.84 (m, 1H), 0.96 (s, 3H), 1.11 (s, 3H), 1.2-1.5 (m, 3H), 1.84 (dd, 1H, J = 14.0 Hz, J = 6.6 Hz), 1.99 (dd, 1H, J = 14.0 Hz, J = 6.8 Hz), 2.68 (t, 2H, J = 7.2 Hz), 3.6-3.85 (m, 1H), 3.85-4.15 (m, 1H).

¹³C-NMR (D₂O, 50 MHz) δ 19.64, 29.32, 32.94, 35.86, 36.95, 44.73, 68.16, 68.25, 100.18, 178.56.

Mass Spectrum (GC/MS), m/z 202, 198, 173, 142, 138, 120, 97, 82, 59, 43.

Step B: Preparation of Trans-(±)-5-(4-fluorophenyl)-2-(1-methylethyl)-N,4-diphenyl-1-[2-(tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)ethyl]-1H-pyrrole-3-carboxamide.

A solution of 0.26 g (1.21 mmol) of (±)-cis-6-(2-aminoethyl)-2,2-dimethyl-1,3-dioxane-4-acetic acid and 0.504 g (1.20 mmol) of (±)4-fluoro-α-[2-methyl-1-oxopropyl]-γ-oxo-N,β-diphenylbenzenebutaneamide mixture of [R-(R*,R*)], [R-(R*,S*)], [S-(R*,R*)] and [S-(R*,S*)] isomers in 5 mL of dimethyl sulfoxide is heated at 105 °C for 15 hours. The solution is cooled and poured into 100 mL of diethyl ether and 50 mL of saturated ammonium chloride in water. The layers are separated and the organic layer washed with water (2 x 50 mL) and 5% sodium hydroxide solution (2 x 100 mL - to extract the intermediate from unreacted diketone). The aqueous layer is acidified with dilute hydrochloric acid solution and extracted with 30 mL of ethyl acetate. A drop of concentrated hydrochloric acid is added to the ethyl acetate solution and allowed to stand 18 hours.

The solution is concentrated in vacuo and the concentrate is redissolved in 30 mL of ethyl acetate and treated with one drop of concentrated hydrochloric acid. The solution is stirred two hours, concentrated in vacuo, and dissolved in 6 mL of toluene. Trans-(±)-5-(4-fluorophenyl)-2-(1-methylethyl)-N,4-diphenyl-1-[2-(tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)ethyl]-1H-pyrrole-3-carboxamide crystallizes and is isolated by filtration. A total of 0.16 g of trans-(±)-5-(4-fluorophenyl)-2-(1-methylethyl)-N,4-diphenyl-1-[2-(tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)ethyl]-1H-pyrrole-3-carboxamide is isolated in two crops.

METHOD C

10 Step A: Preparation of either (±)-(2α,4β,6β) or (±)-(2α,4α,6β)-2-phenyl-6-(2-propenyl)-1,3-dioxane-4-acetonitrile.

Potassium cyanide, 1.3 g (20 mmol), is added to a room temperature solution of (R*,R*)-α-2-propenyloxiraneethanol, 2.56 g (20 mmol), in 25 mL of 4:1 isopropanol:water. The solution is stirred for 20 hours at ambient temperature, concentrated, and partitioned between ethyl acetate and brine. The aqueous layer is extracted 2x with ethyl acetate and the combined ethyl acetate extracts are washed with brine and dried (magnesium sulfate). The crude product is dissolved in 20 mL of benzaldehyde dimethyl acetal, camphor-sulfonic acid is added, and the solution is stirred for 18 hours at room temperature. Concentration and flash chromatography after concentration in vacuo provides 1.30 g of either (±)-(2α,4β,6β) or (±)-(2α,4α,6β)-2-phenyl-6-(2-propenyl)-1,3-dioxane-4-acetonitrile

200 MHz NMR (CDCl₃) δ 1.48 (m, 1H), 1.71 (m, 1H), 2.41 (m, 2H), 2.58 (m, 2H), 3.87 (m, 1H), 4.03 (m, 1H), 5.1 - 5.2 (m, 2H), 5.53 (s, 1H), 5.87 (m, 1H), 7.3 - 7.6 (m, 5H)

¹³C-NMR (CDCl₃, 50 MHz) δ 24.23, 35.07, 39.79, 71.57, 75.48, 100.44, 116.37, 117.53, 125.89, 127.91, 128.61, 133.05, 137.71.

GC/MS m/e 243 (M⁺), 242, 203, 202, 120, 107, 105, 79, 75, 41.

FTIR (neat) 699.6, 758.7, 920.8, 1028.8, 1051.9, 1121.4, 1345.2, 1383.7, 1401.7, 2253.1, 2916.6 cm⁻¹.

30 Step B: Preparation of either (±)-(2α,4α,6α) or (±)-(2α,4β,6β)-6-(2-oxoethyl)-2-phenyl-1,3-dioxane-4-acetonitrile.

A solution of either (±)-(2α,4β,6β) or (±)-(2α,4α,6α)-2-phenyl-6-(2-propenyl)-1,3-dioxane-4-acetonitrile, 4.11 g (16.89 mmol), in 100 mL of dichloromethane is cooled to -78°C under nitrogen. Ozone (Welsbach generator, flow rate 0.1, voltage = 90V) is then passed through a fritted gas inlet tube into the solution until the blue color of ozone appears. The current is turned off, and oxygen bubbled through until the blue color is discharged. Triphenylphosphine, 4.87 g (18.58 mmol), is added and the colorless solution is allowed to warm to room temperature. Flash chromatography provides after concentration in vacuo 3.75 g of pure either (±)-(2α,4α,6α) or (±)-(2α,4β,6β)-6-(2-oxoethyl)-2-phenyl-1,3-dioxane-4-acetonitrile.

GC/MS m/e 245 (M⁺), 244, 123, 105, 95, 94, 77, 51, 41.

40 Step C: Preparation of either (±)-(2α,4α,6α) or (±)-(2α,4β,6β)-6-(cyanomethyl)-2-phenyl-1,3-dioxane-4-acetic acid.

Jones reagent (chromium trioxide-sulfuric acid-water), 3.8 mL (97.6 mmol), is added dropwise to a 0°C solution of either (±)-(2α,4α,6α) or (±)-(2α,4β,6β)-6-(2-oxoethyl)-2-phenyl-1,3-dioxane-4-acetonitrile, 1.86 g (7.6 mmol), dissolved in 50 mL of acetone until the orange color is not discharged. After stirring a further 15 minutes, the mixture is poured into 300 mL of diethyl ether and washed with brine until the aqueous washes are colorless. The diethyl ether layer is dried (magnesium sulfate), filtered, and concentrated to provide 1.84 g of either (±)-(2α,4α,6α) or (±)-(2α,4β,6β)-6-(cyanomethyl)-2-phenyl-1,3-dioxane-4-acetic acid as a yellow gum.

200 MHz NMR (CDCl₃) δ 1.61 (m, 1H), 2.04 (m, 1H), 2.6 - 2.8 (m, 3H) 2.82 (dd, 1H, J = 15.9 Hz, J = 7.1 Hz), 4.20 (m, 1H), 4.40 (m, 1H), 5.60 (s, 1H), 7.2 - 7.5 (m, 5H).

55 Step D: Preparation of either (±)-(2α,4α,6α) or (±)-(2α,4β,6β)-6-(2-aminoethyl)-2-phenyl-1,3-dioxane-4-acetic acid.

A solution of 0.16 g (0.61 mmol) of either (±)-(2α,4α,6α) or (±)-(2α,4β,6β)-6-(cyanomethyl)-2-phenyl-1,3-dioxane-4-acetic acid in 50 mL of methanol saturated with anhydrous ammonia is added to a Parr shaker bottle containing 0.2 g of water wet Raney nickel #30. The solution is heated at 40°C and 3.45 x 10⁵ Pa (50

psig) hydrogen pressure for 6 hours. The suspension is cooled and filtered to remove the Raney nickel through filter aid and the precipitate washed with methanol. The filtrate is concentrated at reduced pressure. The residue is dissolved in methanol saturated with anhydrous ammonia treated with decolorizing charcoal, filtered through filter aid and evaporated to give 0.11 g of either (±)-(2α,4α,6α) or (±)-(2α,4β,6β)-6-(2-aminoethyl)-2-phenyl-1,3-dioxane-4-acetic acid; mp 215.9 - 217.9 °C with decomposition.

200 MHz NMR (D₂O) δ 1.4 - 1.9 (m, 4H), 2.39 (dd, 1H, J = 14.8 Hz, J = 6.7 Hz), 2.55 (dd, 1H, J = 14.8 Hz, J = 7.4 Hz), 2.73 (t, 2H, J = 7.2 Hz) 4.11 (m, 1H), 4.33 (m, 1H), 5.70 (s, 1H), 7.4 - 7.6 (m, 5H).

¹³C-NMR (D₂O, 50 MHz) δ 39.20, 39.78, 40.83, 47.11, 78.34, 78.73, 104.06, 129.15, 131.56, 132.38, 140.51, 181.89.

Step E: Preparation of Trans-(±)-5-(4-fluorophenyl)-2-(1-methylethyl)-N,4-diphenyl-1-[2-(tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)ethyl]-1H-pyrrole-3-carboxamide.

A solution of 0.31 g (1.21 mmol) of either (±)-(2α,4α,6α) or (±)-(2α,4β,6β)-6-(2-aminoethyl)-2-phenyl-1,3-dioxane-4-acetic acid and 0.504 g (1.20 mmol) of (±)-4-fluoro-α-[2-methyl-1-oxopropyl]-γ-oxo-N,β-diphenylbenzenebutaneamide mixture of [R-(R*,R*)], [R-(R*,S*)], [S-(R*,R*)] and [S-(R*,S*)] isomers in 5 mL of dimethyl sulfoxide is heated at 105 °C for 15 hours. The solution is cooled and poured into 100 mL of diethyl ether and 50 mL of saturated ammonium chloride in water. The layers are separated and the organic layer washed with water (2 x 50 mL) and 5% sodium hydroxide solution (2 x 100 mL - to extract the intermediate from unreacted diketone). The aqueous layer is acidified with dilute hydrochloric acid solution, stirred for three hours and extracted with 30 mL of ethyl acetate. A drop of concentrated hydrochloric acid is added to the ethyl acetate solution, the solution is stirred and allowed to stand 18 hours. The solution is concentrated in vacuo and the concentrate is redissolved in 30 mL of ethyl acetate and treated with one drop of concentrated hydrochloric acid. The solution is stirred two hours, concentrated in vacuo, and dissolved in 6 mL of toluene. Trans-(±)-5-(4-fluorophenyl)-2-(1-methylethyl)-N,4-diphenyl-1-[2-(tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)ethyl]-1H-pyrrole-3-carboxamide crystallizes and is isolated by filtration. A total of 0.16 g of trans-(±)-5-(4-fluorophenyl)-2-(1-methylethyl)-N,4-diphenyl-1-[2-(tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)ethyl]-1H-pyrrole-3-carboxamide is isolated in two crops.

METHOD D

Step A: Preparation of (±)-cis-9-(2-propenyl)-6,10-dioxaspiro[4.5]decane-7-acetonitrile.

Potassium cyanide, 1.3 g (20 mmol), is added to a room temperature solution of (R*,R*)-α-2-propenyloxiraneethanol, 2.56 g (20 mmol), in 25 mL of 4:1 isopropanol:water. The solution is stirred for 20 hours at ambient temperature, concentrated, and partitioned between ethyl acetate and brine. The aqueous layer is extracted 2x with ethyl acetate and the combined ethyl acetate extracts are washed with brine and dried (magnesium sulfate). The crude product is dissolved in 20 mL of 1,1-dimethoxycyclopentane, camphorsulfonic acid is added, and the solution is stirred for 18 hours at room temperature. Concentration and flash chromatography after concentration in vacuo provides 1.40 g of (±)-cis-9-(2-propenyl)-6,10-dioxaspiro[4.5]decane-7-acetonitrile.

Step B: Preparation of (±)-cis-9-(2-oxoethyl)-6,10-dioxaspiro[4.5]decane-7-acetonitrile.

A solution of ($\pm$)-cis-9-(2-propenyl)-6,10-dioxaspiro[4.5]decane-7-acetonitrile, 3.4 g (15.36 mmol), in 100 mL of dichloromethane is cooled to -78°C under nitrogen. Ozone (Welsbach generator, flow rate 0.1, voltage = 90V) is then passed through a fritted gas inlet tube into the solution until the blue color of ozone appears. The current is turned off, and oxygen bubbled through until the blue color is discharged.
 5 Triphenylphosphine, 4.2 g (16 mmol), is added and the colorless solution is allowed to warm to room temperature. Flash chromatography provides after concentration in vacuo 2.5 g of pure ($\pm$)-cis-9-(2-oxoethyl)-6,10-dioxaspiro[4.5]decane-7-acetonitrile.

10 **Step C: Preparation of ($\pm$)-cis-9-(cyanomethyl)-6,10-dioxaspiro[4.5]decane-7-acetic acid.**

Jones reagent (chromium trioxide-sulfuric acid-water), 3.8 mL (97.6 mmol), is added dropwise to a 0°C solution of ($\pm$)-cis-9-(2-oxoethyl)-6,10-dioxaspiro[4.5]decane-2-acetonitrile 1.70 (7.6 mmol), dissolved in 50 mL of acetone until the orange color is not discharged. After stirring a further 15 minutes, the mixture is poured into 300 mL of diethyl ether and washed with brine until the aqueous washes are colorless. The diethyl ether layer is dried (magnesium sulfate), filtered, and concentrated to provide 1.20 g of ($\pm$)-cis-9-(cyanomethyl)-6,10-dioxaspiro[4.5]decane-7-acetic acid as a colorless solid.

20

Step D: Preparation of ($\pm$)-cis-9-(2-aminoethyl)-6,10-dioxaspiro[4.5]decane-7-acetic acid.

25

A solution of 1.17 g (4.88 mmol) of ($\pm$)-cis-9-(cyanomethyl)-6,10-dioxaspiro[4.5]decane-7-acetic acid in 100 mL of methanol saturated with anhydrous ammonia is added to a Parr shaker bottle containing 0.53 g of water wet Raney nickel #30. The solution is heated at 45°C and 3.45×10^5 Pa (50 psig) hydrogen pressure for 17 hours. The suspension is cooled and filtered to remove the Raney nickel through filter aid and the precipitate is washed with methanol. The filtrate is concentrated at reduced pressure. The residue is dissolved in methanol saturated with anhydrous ammonia treated with decolorizing charcoal, filtered through filter aid and evaporated to give 0.56 g of ($\pm$)-cis-9-(2-aminoethyl)-6,10-dioxaspiro[4.5]decane-7-acetic acid.

30

35 **Step E: Preparation of Trans-($\pm$)-5-(4-fluorophenyl)-2-(1-methylethyl)-N,4-diphenyl-1-[2-(tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)ethyl]-1H-pyrrole-3-carboxamide.**

40

A solution of 0.295 g (1.21 mmol) of ($\pm$)-cis-9-(2-aminoethyl)-6,10-dioxaspiro[4.5]decane-7-acetic acid and 0.504 g (1.20 mmol) of ($\pm$)-4-fluoro- α -[2-methyl-1-oxopropyl]- γ -oxo-N,8-diphenylbenzenebutaneamide mixture of [R-(R*,R*)], [R-(R*,S*)], [S-(R*,R*)] and [S-(R*,S*)] isomers in 5 mL of dimethyl sulfoxide is heated at 105°C for 15 hours. The solution is cooled and poured into 100 mL of diethyl ether and 50 mL of saturated ammonium chloride in water. The layers are separated and the organic layer washed with water (2 x 50 mL) and 5% sodium hydroxide solution (2 x 100 mL - to extract the intermediate protected acid from unreacted diketone). The aqueous layer is acidified with dilute hydrochloric acid solution, stirred for three hours and extracted with 30 mL of ethyl acetate. A drop of concentrated hydrochloric acid is added to the ethyl acetate solution and allowed to stand 18 hours. The solution is concentrated in vacuo and the concentrate is redissolved in 30 mL of ethyl acetate and treated with one drop of concentrated hydrochloric acid. The solution is stirred two hours, concentrated in vacuo, and dissolved in 6 mL of toluene. Trans-($\pm$)-5-(4-fluorophenyl)-2-(1-methylethyl)-N,4-diphenyl-1-[2-(tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)ethyl]-1H-pyrrole-3-carboxamide crystallizes and is isolated by filtration. A total of 0.16 g of trans-($\pm$)-5-(4-fluorophenyl)-2-(1-methylethyl)-N,4-diphenyl-1-[2-(tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)ethyl]-1H-pyrrole-3-carboxamide is isolated in two crops.

55

METHOD E

Step A: Preparation of (±)-cis-4-(2-propenyl)-1,5-dioxaspiro[5.5]undecane-2-acetonitrile.

Potassium cyanide, 1.3 g (20 mmol), is added to a room temperature solution of (R*,R*)-α-2-propenyloxiraneethanol, 2.56 g (20 mmol), in 25 mL of 4:1 isopropanol:water. The solution is stirred for 20 hours at ambient temperature, concentrated, and partitioned between ethyl acetate and brine. The aqueous layer is extracted 2x with ethyl acetate and the combined ethyl acetate extracts are washed with brine and dried (magnesium sulfate). The crude product is dissolved in 20 mL of 2,2-dimethoxycyclohexane, camphorsulfonic acid is added, and the solution is stirred for 18 hours at room temperature. Concentration and flash chromatography after concentration in vacuo provides 1.50 g of (±)-cis-4-(2-propenyl)-1,5-dioxaspiro[5.5]undecane-2-acetonitrile.

200 MHz NMR (CDCl₃) δ 1.26 (m, 1H), 1.4 - 1.7 (m, 8H), 1.87 (m, 2H), 2.1 - 2.3 (m, 2H), 2.51 (d, 2H, J = 6.05 Hz), 3.95 (m, 1H), 4.15 (m, 1H), 5.0 - 5.2 (m, 2H), 5.83 (m, 1H).

¹³C-NMR (CDCl₃, 50 MHz) δ 22.16, 24.71, 25.42, 28.20, 35.47, 38.33, 40.30, 64.08, 66.87, 98.84, 116.53, 116.83, 113.46.

GC/MS m/e 235 (m⁺), 206, 192, 120, 99, 93, 79, 69, 55, 41.

FTIR (film) 964.5, 1121.4, 1160.0, 2253.1, 2937.2 cm⁻¹.

Step B: Preparation of (±)-cis-4-(2-oxoethyl)-1,5-dioxaspiro[5.5]undecane-2-acetonitrile.

A solution of (±)-cis-4-(2-propenyl)-1,5-dioxaspiro[5.5]undecane-2-acetonitrile 4.26 g (19.42 mmol), in 100 mL of dichloromethane is cooled to -78 °C under nitrogen. Ozone (Welsbach generator, flow rate 0.1, voltage = 90V) is then passed through a fritted gas inlet tube into the solution until the blue color of ozone appears. The current is turned off, and oxygen bubbled through until the blue color is discharged. Triphenylphosphine, 5.6 g (21.36 mmol), is added and the colorless solution is allowed to warm to room temperature. Flash chromatography provides after concentration in vacuo 4.04 g of pure (±)-cis-4-(2-oxoethyl)-1,5-dioxaspiro[5.5]undecane-2-acetonitrile.

200 MHz NMR (CDCl₃) δ 1.3 - 2.0 (m, 12H), 2.5 - 2.7 (m, 4H), 4.20 (m, 1H), 4.36 (m, 1H), 9.81 (t, 1H, J = 1.74 Hz).

¹³C-NMR (CDCl₃, 50 MHz) δ 22.39, 22.44, 24.97, 25.59, 28.44, 35.82, 38.48, 49.54, 63.25, 64.17, 99.66, 116.57, 199.82.

GC/MS m/e 237 (m⁺), 208, 194, 122, 94, 81, 55, 42, 41.

FTIR (film) 969.6, 1069.9, 1126.5, 1160.0, 1368.3, 1386.3, 1728.4, 2934.6 cm⁻¹.

Step C: Preparation of (±)-cis-4-(cyanomethyl)-1,5-dioxaspiro[5.5]undecane-2-acetic acid.

Jones reagent (chromium trioxide-sulfuric acid-water), 4.4 mL (8.85 mmol), is added dropwise to a 0 °C solution of (±)-cis-4-(2-oxoethyl)-1,5-dioxaspiro[5.5]undecane-2-acetonitrile, 3.62 g (12.6 mmol), dissolved in 30 mL of acetone until the orange color is not discharged. After stirring a further 15 minutes, the mixture is poured into 300 mL of diethyl ether and washed with brine until the aqueous washes are colorless. The diethyl ether layer is dried (magnesium sulfate), filtered, and concentrated to provide 3.65 g of (±)-cis-4-(cyanomethyl)-1,5-dioxaspiro[5.5]undecane-2-acetic acid as a yellow solid.

200 MHz NMR (Pyridine) δ 1.2 - 2.0 (m, 12H), 2.5 - 2.9 (m, 4H), 4.19 (m, 1H), 4.50 (m, 1H).

Step D: Preparation of (±)-cis-4-(2-aminoethyl)-1,5-dioxaspiro[5.5]undecane-2-acetic acid.

5

A solution of 0.13 g of (±)-cis-4-(cyanomethyl)-1,5-dioxaspiro[5.5]undecane-2-acetic acid in 20 mL of methanol saturated with anhydrous ammonia is added to a Parr shaker bottle containing 0.2 g of water wet Raney nickel #30. The solution is heated at 40°C and 3.45×10^5 Pa (50 psig) hydrogen pressure for 17 hours. The suspension is cooled and filtered to remove the Raney nickel through filter aid and the precipitate is washed with methanol. The filtrate is concentrated at reduced pressure. The residue is dissolved in methanol saturated with anhydrous ammonia, treated with decolorizing charcoal, filtered through filter aid and evaporated to give 0.13 g of (±)-cis-4-(2-aminoethyl)-1,5-dioxaspiro[5.5]undecane-2-acetic acid.

200 MHz NMR (D_2O) δ 1.1 - 1.7 (m, 10H), 1.75 - 2.1 (m, 4H), 2.19 (dd, 1H, $J = 14.6$ Hz, $J = 6.7$ Hz), 2.31 (dd, 1H, $J = 14.6$ Hz, $J = 7.3$ Hz), 2.69 (t, 2H, $J = 7.1$ Hz), 4.09 (m, 1H), 4.34 (m, 1H).

^{13}C -NMR (D_2O , 50 MHz) δ 24.50, 24.73, 27.55, 30.80, 38.91, 39.39, 39.63, 40.48, 47.03, 69.37, 69.54, 102.74, 181.33.

Step E: Preparation of Trans-(±)-5-(4-fluorophenyl)-2-(1-methylethyl)-N,4-diphenyl-1-[2-(tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)ethyl]-1H-pyrrole-3-carboxamide.

25

A solution of 0.31 g (1.21 mmol) of (±)-cis-4-(2-aminoethyl)-1,5-dioxaspiro[5.5]undecane-2-acetic acid and 0.504 g (1.20 mmol) of (±)-4-fluoro- α -[2-methyl-1-oxopropyl]- γ -oxo-N, β -diphenylbenzenebutaneamide mixture of [R-(R*,R*)], [R-(R*,S*)], [S-(R*,R*)] and [S-(R*,S*)] isomers in 5 mL of dimethyl sulfoxide is heated at 105°C for 15 hours. The solution is cooled and poured into 100 mL of diethyl ether and 50 mL of saturated ammonium chloride in water. The layers are separated and the organic layer washed with water (2 x 50 mL) and 5% sodium hydroxide solution (2 x 100 mL - to extract the intermediate acid from unreacted diketone). The aqueous layer is acidified with dilute hydrochloric acid solution, stirred for three hours and extracted with 30 mL of ethyl acetate to remove the protecting group before lactonization. The extract is concentrated and dissolved in 30 mL of ethyl acetate. A drop of concentrated hydrochloric acid is added to the ethyl acetate solution and allowed to stand 18 hours. The solution is concentrated in vacuo and the concentrate is redissolved in 30 mL of ethyl acetate and treated with one drop of concentrated hydrochloric acid. The solution is stirred two hours, concentrated in vacuo, and dissolved in 6 mL of toluene. Trans-(±)-5-(4-fluoro-phenyl)-2-(1-methylethyl)-N,4-diphenyl-1-[2-(tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)ethyl]-1H-pyrrole-3-carboxamide crystallizes and is isolated by filtration. A total of 0.155 g of trans-(±)-5-(4-fluorophenyl)-2-(1-methylethyl)-N,4-diphenyl-1-[2-(tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)ethyl]-1H-pyrrole-3-carboxamide is isolated in two crops.

METHOD F

45

Step A: Preparation of (±)-cis-6-(2-propenyl)-1,3-dioxane-4-acetonitrile

50

Potassium cyanide, 1.3 g (20 mmol), is added to a room temperature solution of (R*,R*)- α -2-propenyloxiraneethanol, 2.56 g (20 mmol), in 25 mL of 4:1 isopropanol:water. The solution is stirred for 20 hours at ambient temperature, concentrated, and partitioned between ethyl acetate and brine. The aqueous layer is extracted 2x with ethyl acetate and the combined ethyl acetate extracts are washed with brine and dried (magnesium sulfate). The crude product is dissolved in 20 mL of dimethoxymethane, camphorsulfonic acid is added, and the solution is stirred for 18 hours at room temperature. Concentration and flash chromatography after concentration in vacuo provides 1.20 g of (±)-cis-6-(2-propenyl)-1,3-dioxane-4-acetonitrile.

Step B: Preparation of (±)-cis-6-(2-oxoethyl)-1,3-dioxane-4-acetonitrile.

5

A solution of (±)-cis-6-(2-propenyl)-1,3-dioxane-4-acetonitrile, 2.57 g (15.36 mmol), in 100 mL of dichloromethane is cooled to -78°C under nitrogen. Ozone (Welsbach generator, flow rate 0.1, voltage = 90V) is then passed through a fritted gas inlet tube into the solution until the blue color of ozone appears. The current is turned off, and oxygen bubbled through until the blue color is discharged. Triphenylphosphine 4.87 g (28.58 mmol), is added and the colorless solution is allowed to warm to room temperature. Flash chromatography provides after concentration in vacuo 2.3 g of pure (±)-cis-6-(2-oxoethyl)-1,3-dioxane-4-acetonitrile.

15

Step C: Preparation of (±)-cis-6-(cyanomethyl)-1,3-dioxane-4-acetic acid.

Jones reagent (chromium trioxide-sulfuric acid-water), 3.8 mL (97.6 mmol), is added dropwise to a 0°C solution of (±)-cis-6-(2-oxoethyl)-1,3-dioxane-4-acetonitrile, 1.29 g (7.6 mmol), dissolved in 50 mL of acetone until the orange color is not discharged. After stirring a further 15 minutes, the mixture is poured into 300 mL of diethyl ether and washed with brine until the aqueous washes are colorless. The diethyl ether layer is dried (magnesium sulfate), filtered, and concentrated to provide 1.2 g of (±)-cis-6-(cyanomethyl)-1,3-dioxane-4-acetic acid as a colorless solid.

25

Step D: Preparation of (±)-cis-6-(2-aminoethyl)-1,3-dioxane-4-acetic acid.

30

A solution of 1.04 g (4.88 mmol) of (±)-cis-6-(cyanomethyl)-1,3-dioxane-4-acetic acid in 100 mL of methanol saturated with anhydrous ammonia is added to a Parr shaker bottle containing 0.53 g of water wet Raney nickel #30. The solution is heated at 45°C and 3,45 x 10⁵ Pa (50 psig) hydrogen pressure for 17 hours. The suspension is cooled and filtered to remove the Raney nickel through filter aid and the precipitate washed with methanol. The filtrate is concentrated at reduced pressure. The residue is dissolved in methanol saturated with anhydrous ammonia, treated with decolorizing charcoal, filtered through filter aid and evaporated to give 0.56 g of (±)-cis-6-(2-aminoethyl)-1,3-dioxane-4-acetic acid.

40

Step E: Preparation of Trans-(±)-5-(4-fluorophenyl)-2-(1-methylethyl)-N,4-diphenyl-1-[2-(tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)ethyl]-1H-pyrrole-3-carboxamide.

45

A solution of 0.26 g (1.21 mmol) of (±)-cis-6-(2-aminoethyl)-1,3-dioxane-4-acetic acid and 0.504 g (1.20 mmol) of (±)-4-fluoro-α-[2-methyl-1-oxopropyl]-γ-oxo-N,β-diphenylbenzenebutaneamide in 5 mL of dimethyl sulfoxide is heated at 105°C for 15 hours. The solution is cooled and poured into 100 mL of diethyl ether and 50 mL of saturated ammonium chloride in water. The layers are separated and the organic layer washed with water (2 x 50 mL) and 5% sodium hydroxide solution (2 x 100 mL - to extract the intermediate acid from unreacted diketone). The aqueous layer is acidified with dilute hydrochloric acid solution and extracted with 30 mL of ethyl acetate. A drop of concentrated hydrochloric acid is added to the ethyl acetate solution and allowed to stand 18 hours. The solution is concentrated in vacuo and the concentrate is redissolved in 30 mL of ethyl acetate and treated with one drop of concentrated hydrochloric acid. The solution is stirred two hours, concentrated in vacuo, and dissolved in 6 mL of toluene. Trans-(±)-5-(4-fluorophenyl)-2-(1-methylethyl)-N,4-diphenyl-1-[2-(tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)ethyl]-1H-pyrrole-3-carboxamide crystallizes and is isolated by filtration. A total of 0.15 g of trans-(±)-5-(4-

fluorophenyl)-2-(1-methylethyl)-N,4-diphenyl-1-[2-(tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)ethyl]-1H-pyrrole-3-carboxamide is isolated in two crops.

METHOD G

5

Step A: Preparation of either $(\pm)$ -(2 α ,4 β ,6 β) or $(\pm)$ -(2 α ,4 α ,6 α)-2-methyl-6-(2-propenyl)-1,3-dioxane-4-acetonitrile.

Potassium cyanide, 1.3 g (20 mmol), is added to a room temperature solution of (R*,R*)- α -2-propenyloxiraneethanol, 2.56 g (20 mmol), in 25 mL of 4:1 isopropanol:water. The solution is stirred for 20 hours at ambient temperature, concentrated, and partitioned between ethyl acetate and brine. The aqueous layer is extracted 2x with ethyl acetate and the combined ethyl acetate extracts are washed with brine and dried (magnesium sulfate). The crude product is dissolved in 20 mL of 1,1-dimethoxyethane, camphorsulfonic acid is added, and the solution is stirred for 18 hours at room temperature. Concentration and flash chromatography after concentration in vacuo provides 1.30 g of either $(\pm)$ -(2 α ,4 β ,6 β) or $(\pm)$ -(2 α ,4 α ,6 α)-2-methyl-6-(2-propenyl)-1,3-dioxane-4-acetonitrile.

Step B: Preparation of either $(\pm)$ -(2 α ,4 α ,6 α) or $(\pm)$ -(2 α ,4 β ,6 β)-2-methyl-6-(2-oxoethyl)-1,3-dioxane-4-acetonitrile.

20

A solution of either $(\pm)$ -(2 α ,4 β ,6 β) or $(\pm)$ -(2 α ,4 α ,6 α)-2-methyl-6-(2-propenyl)-1,3-dioxane-4-acetonitrile, 2.78 g (15.36 mmol), in 100 mL of dichloromethane is cooled to -78 °C under nitrogen. Ozone (Welsbach generator, flow rate 0.1, voltage = 90V) is then passed through a fritted gas inlet tube into the solution until the blue color of ozone appears. The current is turned off, and oxygen bubbled through until the blue color is discharged. Triphenylphosphine, 4.2 g (16 mmol), is added and the colorless solution is allowed to warm to room temperature. Flash chromatography provides after concentration in vacuo 2.5 g of pure either $(\pm)$ -(2 α ,4 α ,6 α) or $(\pm)$ -(2 α ,4 β ,6 β)-2-methyl-6-(2-oxoethyl)-1,3-dioxane-4-acetonitrile.

Step C: Preparation of either $(\pm)$ -(2 α ,4 α ,6 α) or $(\pm)$ -(2 α ,4 β ,6 β)-6-(cyanomethyl)-2-methyl-1,3-dioxane-4-acetic acid.

Jones reagent (chromium trioxide-sulfuric acid-water), 3.8 mL (7.6 mmol), is added dropwise to a 0 °C solution of either $(\pm)$ -(2 α ,4 α ,6 α) or $(\pm)$ -(2 α ,4 β ,6 β)-2-methyl-6-(2-oxoethyl)-1,3-dioxane-4-acetonitrile, 1.40 g (7.6 mmol), dissolved in 50 mL of acetone until the orange color is not discharged. After stirring a further 15 minutes, the mixture is poured into 300 mL of diethyl ether and washed with brine until the aqueous washes are colorless. The diethyl ether layer is dried (magnesium sulfate), filtered, and concentrated to provide 1.01 g of either $(\pm)$ -(2 α ,4 α ,6 α) or $(\pm)$ -(2 α ,4 β ,6 β)-6-(cyanomethyl)-2-methyl-1,3-dioxane-4-acetic acid as a colorless solid.

Step D: Preparation of either $(\pm)$ -(2 α ,4 α ,6 α) or $(\pm)$ -(2 α ,4 β ,6 β)-6-(2-aminoethyl)-2-methyl-1,3-dioxane-4-acetic acid.

A solution of 0.97 g (4.88 mmol) of either $(\pm)$ -(2 α ,4 α ,6 α) or $(\pm)$ -(2 α ,4 β ,6 β)-6-(cyanomethyl)-2-methyl-1,3-dioxane-4-acetic acid in 100 mL of methanol saturated with anhydrous ammonia is added to a Parr shaker bottle containing 0.53 g of water wet Raney nickel #30. The solution is heated at 45 °C and 3,45 x 10⁵ Pa (50 psig) hydrogen pressure for 17 hours. The suspension is cooled and filtered to remove the Raney nickel through filter aid and the precipitate washed with methanol. The filtrate is concentrated at reduced pressure. The residue is dissolved in methanol saturated with anhydrous ammonia treated with decolorizing charcoal, filtered through filter aid and evaporated to give 0.50 g of either $(\pm)$ -(2 α ,4 α ,6 α) or $(\pm)$ -(2 α ,4 β ,6 β)-6-(2-aminoethyl)-2-methyl-1,3-dioxane-4-acetic acid.

Step E: Preparation of Trans-($\pm$)-5-(4-fluorophenyl)-2-(1-methylethyl)-N,4-diphenyl-1-[2-(tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)ethyl]-1H-pyrrole-3-carboxamide.

A solution of 0.31 g of either ($\pm$)-(2 α ,4 α ,6 α) or ($\pm$)-(2 α ,4 β ,6 β)-6-(2-aminoethyl)-2-methyl-1,3-dioxane-4-acetic acid and 0.504 g (1.20 mmol) of ($\pm$)-4-fluoro- α -[2-methyl-1-oxopropyl]- γ -oxo-N, β -diphenylbenzenebutaneamide in 5 mL of dimethyl sulfoxide is heated at 105 °C for 15 hours. The solution is cooled and poured into 100 mL of diethyl ether and 50 mL of saturated ammonium chloride in water. The layers are separated and the organic layer washed with water (2 x 50 mL) and 5% sodium hydroxide solution (2 x 100 mL - to extract the intermediate acid from unreacted diketone). The aqueous layer is acidified with dilute hydrochloric acid solution, stirred for three hours and extracted with 30 mL of ethyl acetate. A drop of concentrated hydrochloric acid is added to the ethyl acetate solution and allowed to stand 18 hours. The solution is concentrated in vacuo and the concentrate is redissolved in 30 mL of ethyl acetate and treated with one drop of concentrated hydrochloric acid. The solution is stirred two hours, concentrated in vacuo, and dissolved in 6 mL of toluene. Trans-($\pm$)-5-(4-fluorophenyl)-2-(1-methylethyl)-N,4-diphenyl-1-[2-(tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)ethyl]-1H-pyrrole-3-carboxamide crystallizes and is isolated by filtration. A total of 0.14 g of trans-($\pm$)-5-(4-fluorophenyl)-2-(1-methylethyl)-N,4-diphenyl-1-[2-(tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)ethyl]-1H-pyrrole-3-carboxamide is isolated in two crops.

METHOD H

Step A: Preparation of ($\pm$)-cis-1,1-dimethylethyl 6-(cyanomethyl)-2,2-dimethyl-1,3-dioxane-4-acetate.

($\pm$)-cis-6-(cyanomethyl)-2,2-dimethyl-1,3-dioxane-4-acetic acid (3.72 g, 17.44 mmol) is dissolved in 20 mL of dichloromethane, cooled to 0 °C and 0.2 g of 4-dimethylaminopyridine (DMAP) is added, followed by t-butyl alcohol and followed by 4.32 g of dicyclohexylcarbodiimide (DCC). This solution is allowed to slowly warm to room temperature over a 76.5-hour period. Thin layer chromatography (TLC) shows mainly product, and some slightly lower R_f by-products. The mixture is stirred one hour and 50 mL of dichloromethane is added and stirring continues five hours. An additional 100 mL of diethyl ether is added and the mixture filtered. The precipitate is washed with diethyl ether. The filtrate is concentrated to an oil. The crude product is chromatographed on silica gel eluting with 4:1 hexane:ethyl acetate. The eluate is concentrated to yield ($\pm$)-cis-1,1-dimethylethyl 6-(cyanomethyl)-2,2-dimethyl-1,3-dioxane-4-acetate. 200 MHz NMR (CDCl₃) δ 1.36 (m, 1H), 1.42 (s, 3H), 1.49 (s, 9H), 1.50 (s, 3H), 1.79 (dt, 1H, J = 2.5 Hz, J = 12.1 Hz), 2.40 (dd, 1H, J = 6.2 Hz, J = 15.4 Hz), 2.5 - 2.7 (m, 1H), 2.55 (d, 2H, J = 6.1 Hz) 4.18 (m, 1H), 4.32 (m, 1H). ¹³C-NMR (CDCl₃, 50 MHz) δ 19.80, 25.16, 28.30, 29.94, 35.66, 42.56, 65.27, 65.87, 80.96, 99.57, 116.72, 169.83. GCMS m/e 254, 199, 198, 154, 138, 59, 57, 43, 41. FTIR (neat) 954.2, 987.6, 1152.3, 1201.1, 1257.7, 1316.9, 1368.3, 1383.7, 1728.4, 2253.1, 2942.4, 2983.5 cm⁻¹.

Step B: Preparation of ($\pm$)-cis-1,1-dimethylethyl 6-(2-aminoethyl)-2,2-dimethyl-1,3-dioxane-4-acetate.

A solution of 6.75 g of ($\pm$)-cis-1,1-dimethyl ethyl 6-(cyanomethyl)-2,2-dimethyl-1,3-dioxane-4-acetate in 80 mL of methanol saturated with gaseous ammonia is treated with 0.7 g of Raney nickel #30 and hydrogen gas in a shaker at 3,45 x 10⁵ Pa (50 psig) and 40 °C. After 10 hours, thin layer chromatography indicates no starting nitrile present. The suspension is cooled, filtered through filter aid, and concentrated to an oil. This crude oil is purified by flash chromatography on silica gel with 30:20:1 (ethyl acetate:methanol-ammonium hydroxide) as eluent to give 5.48 g of ($\pm$)-cis-1,1-dimethylethyl 6-(2-aminoethyl)-2,2-dimethyl-1,3-dioxane-4-acetate (98.2 area %) as a clear oil which hardens after time. 200 MHz NMR (CDCl₃) δ 1.0 - 1.2 (m, 1H), 1.22 (s, 3H), 1.31 (s, 12H), 1.35 - 1.45 (m, 1H), 1.77 (brs, 2H), 2.15 (dd, 1H, J = 15.1 Hz, J = 6.2 Hz), 2.29 (dd, 1H, J = 15.1 Hz, J = 7.0 Hz), 2.66 (t, 2H, J = 6.6 Hz), 3.82 (m, 1H), 4.12 (m, 1H). ¹³C-NMR (CDCl₃, 50 MHz) δ 19.56, 27.92, 29.96, 36.43, 38.18, 39.65, 42.55, 66.03, 67.16, 80.19, 98.32, 169.80. GC/MS m/e 258, 216, 215, 202, 200, 142, 113, 100, 99, 72, 57, 43. FTIR (neat) 951.6, 1157.4, 1201.1, 1260.3, 1314.3, 1368.3, 1381.2, 1728.4, 2361.1, 2939.8, 2980.9 cm⁻¹.

Step C: Preparation of Trans-(±)-5-(4-fluorophenyl)-2-(1-methylethyl)-N,4-diphenyl-1-[2-(tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)ethyl]-1H-pyrrole-3-carboxamide.

A solution of 0.79 g (2.89 mmol) of (±)-cis-1,1-dimethylethyl 6-(2-aminoethyl)-2,2-dimethyl-1,3-dioxane-4-acetate and 1.00 g (2.41 mmol) of (±)-4-fluoro-α-[2-methyl-1-oxopropyl]-γ-oxo-N,β-diphenylbenzenebutaneamide mixture of [R-(R*,R*)], [R-(R*,S*)], [S-(R*,R*)] and [S-(R*,S*)] isomers in 15 mL of heptane:toluene (9:1) is heated at reflux for 24 hours. The solution is cooled and poured into 100 mL of tetrahydrofuran and 50 mL of saturated ammonium chloride in water. The layers are separated and the organic layer washed with brine. To the organic layer is added 5 mL of 10% hydrochloric acid solution and the solution is stirred for 15 hours. To this solution is added 1.2 g of sodium hydroxide and the mixture is stirred for 30 hours. The reaction is stopped by adding 50 mL of water, 30 mL of hexane, and separating the layers. The aqueous layer is acidified with dilute hydrochloric acid solution, stirred for three hours and extracted with 50 mL of ethyl acetate. A drop of concentrated hydrochloric acid is added to the ethyl acetate solution and the solution is allowed to stand 18 hours. The solution is concentrated in vacuo and the concentrate is redissolved in 50 mL of ethyl acetate and treated with one drop of concentrated hydrochloric acid. The solution is stirred two hours, concentrated in vacuo, and dissolved in 10 mL of toluene. Trans-(±)-5-(4-fluorophenyl)-2-(1-methylethyl)-N,4-diphenyl-1-[2-(tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)ethyl]-1H-pyrrole-3-carboxamide is isolated in two crops.

METHOD I

Step A: Preparation of either (±)-(2α,4α,6α) or (±)-(2α,4β,6β)-1,1-dimethylethyl 6-(2-cyanomethyl)-2-phenyl-1,3-dioxane-4-acetate.

Either (±)-(2α,4α,6α) or (±)-(2α,4β,6β)-6-(cyanomethyl)-2-phenyl-1,3-dioxane-4-acetic acid (3.07 g, 11.75 mmol) is dissolved in 15 mL of dichloromethane, cooled to 0 °C and 0.1 g of 4-dimethylaminopyridine (DMAP) added, followed by t-butyl alcohol, followed by 2.91 g of dicyclohexylcarbodiimide (DCC). This solution is allowed to slowly warm to room temperature and stirred over a 16.5-hour period. Thin layer chromatography (TLC) shows mainly product, and some slightly lower R_f by-products. The mixture is stirred one hour and 50 mL of dichloromethane is added, and stirring is continued for five hours. An additional 100 mL of diethyl ether is added and then filtered. The precipitate is washed with diethyl ether. The filtrate is concentrated to an oil. The crude product is chromatographed on silica gel eluting with 4:1 hexane:ethyl acetate. The eluate is concentrated to yield either (±)-(2α,4α,6α) or (±)-(2α,4β,6β)-1,1-dimethylethyl 6-(2-cyanomethyl)-2-phenyl-1,3-dioxane-4-acetate.

GC/MS m/e 260, 244, 202, 138, 107, 105, 77, 57, 41.

Step B: Preparation of either (±)-(2α,4α,6α) or (±)-(2α, 4β, 6β) 1,1-dimethylethyl 6-(2-aminoethyl)-2-phenyl-1,3-dioxane-4-acetate.

A solution of 1.72 g of either (±)-(2α,4α,6α) or (±)-(2α, 4β, 6β)-1,1-dimethylethyl 6-(cyanomethyl)-2-phenyl-1,3-dioxane-4-acetate in 30 mL of methanol saturated with gaseous ammonia is treated with 0.3 g of Raney nickel #30 and hydrogen gas in a shaker at 3.45 x 10⁵ Pa (50 psig) and 40 °C. After 10 hours, thin layer chromatography indicates no starting nitrile present. The suspension is cooled, filtered through filter aid, and concentrated to an oil. This crude oil is purified by flash chromatography on silica gel with 30:20:1 (ethyl acetate:methanol: ammonium hydroxide) as eluent to give 1.56 g of either (±)-(2α,4α,6α) or (±)-(2α,4β,6β)-1,1-dimethylethyl 6-(2-aminoethyl)-2-phenyl-1,3-dioxane-4-acetate (98.2 area %) as a clear oil which hardens after time.

200 MHz NMR (CDCl₃) δ 1.2 - 1.9 (m, 4H) 1.44 (s, 9H), 2.03, (br.s, 2H), 2.42 (dd, 1H, J = 15.3 Hz, J = 6.3 Hz), 2.63 (dd, 1H, J = 15.3 Hz, J = 7.0 Hz), 2.89 (t, 2H, J = 6.8 Hz), 3.97 (m, 1H), 4.26 (m, 1H), 5.56 (s, 1H), 7.3 - 7.4 (m, 3H), 7.4 - 7.5 (m, 2H).

¹³C NMR (CDCl₃, 50 MHz) δ 28.07, 36.57, 38.23, 39.25, 42.17, 73.47, 74.87, 80.60, 100.36, 125.82, 127.88, 128.34, 138.45, 169.73.

GC/MS, m/e 321, 320, 248, 215, 174, 142, 105, 57.

FTIR (film) 699.6, 756.2, 1026.2, 1116.2, 1149.7, 1368.3, 1394.0, 1718.1, 1733.5, 2872.9, 2932.1 cm^{-1} .

Step C: Preparation of Trans-(±)-5-(4-fluorophenyl)-2-(1-methylethyl)-N,4-diphenyl-1-[2-(tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)ethyl]-1H-pyrrole-3-carboxamide.

In a process analogous to Method H by substituting (±)-(2 α ,4 α ,6 α) or (±)-(2 α ,4 β ,6 β) 1,1-dimethylethyl-6-(2-aminoethyl)-2-phenyl-1,3-dioxane-4-acetate for (±)-cis-1,1-dimethylethyl 6-(2-aminoethyl)-2,2-dimethyl-1,3-dioxane-4-acetate one obtains the title compound.

METHOD J

Step A: Preparation of (±)-cis-1,1-dimethylethyl 4-(cyanomethyl)-1,5-dioxaspiro[5.5]undecane-2-acetate.

(±)-cis-4-(cyanomethyl)-1,5-dioxaspiro[5.5]undecane-2-acetic acid 3.32 g (13.12 mmol), is dissolved in 15 mL of dichloromethane, cooled to 0°C and 0.1 g of 4-dimethylaminopyridine (DMAP) added, followed by t-butyl alcohol, and followed by 3.25 g of dicyclohexylcarbodiimide (DCC). This solution is stirred and allowed to slowly warm to room temperature over a 16.5-hour period. TLC shows mainly product, and some slightly lower R_f by-product. The mixture is stirred one hour and 50 mL of dichloromethane is added and then stirring continued four hours. One hundred mL of diethyl ether is added and then filtered. The filtrate is concentrated at reduced pressure. This crude concentrate is chromatographed on silica gel and eluted with 4:1 hexane:ethyl acetate to yield (±)-cis-1,1-dimethylethyl 4-(cyanomethyl)-1,5-dioxaspiro[5.5]undecane-2-acetate.

200 MHz NMR (CDCl_3) δ 1.1 - 2.0 (m, 12H) 1.43 (s, 9H), 2.36 (m, 2H), 2.48 (m, 2H), 4.1 - 4.4 (m, 2H).
 ^{13}C -NMR (CDCl_3 , 50 MHz) 22.37, 22.45, 25.08, 28.15, 28.55, 35.80, 38.57, 42.59, 64.31, 64.92, 80.76, 99.56, 116.65, 169.82.
 GC/MS m/e 309 (m^+), 266, 224, 210, 138, 120, 99, 57, 55.
 FTIR (KBr) 964.5, 1149.7, 1157.4, 1332.3, 1368.3, 1712.9, 2939.8 cm^{-1} .

Step B: Preparation of (±)-cis-1,1-dimethylethyl 4-(2-aminoethyl)-1,5-dioxaspiro[5.5]undecane-2-acetate.

A solution of 1.19 g of (±)-cis-1,1-dimethylethyl 4-(cyanomethyl)-1,5-dioxaspiro[5.5]undecane-2-acetate in 30 mL of methanol saturated with gaseous ammonia is treated with 0.3 g of Raney nickel #30 and hydrogen gas in a shaker at 3.45×10^5 Pa (50 psig) and 40°C. After 22 hours, thin layer chromatography indicates no starting nitrile present. The suspension is cooled, filtered through filter aid, and concentrated to an oil. This crude oil is purified by silica gel flash chromatography (30:20:1; ethyl acetate:methanol:ammonium hydroxide) to give 1.18 g of (±)-cis-1,1-dimethylethyl 4-(2-aminoethyl)-1,5-dioxaspiro[5.5]undecane-2-acetate as a clear oil which solidifies upon standing.

200 MHz NMR (CDCl_3) δ 1.2 - 2.0 (m, 12H), 1.43 (s, 9H), 2.34, (m, 2H), 2.50 (br.s, 2H), 2.84 (t, 2H, J = 7 Hz), 3.99 (m, 1H), 4.28 (m, 1H).
 GC/MS, m/e 313, 270, 214, 185, 144, 142, 99.
 FTIR (film) 961.9, 1098.2, 1154.8, 1368.3, 1725.8, 2934.6 cm^{-1} .

Step C: Preparation of Trans-(±)-5-(4-fluorophenyl)-2-(1-methylethyl)-N,4-diphenyl-1-[2-(tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)ethyl]-1H-pyrrole-3-carboxamide.

In a process analogous to Method H by substituting (±)-cis-1,1-dimethylethyl 4-(2-aminoethyl)-1,5-dioxaspiro[5.5]undecane-2-acetate for (±)-cis-1,1-dimethylethyl 6-(2-aminoethyl)-2,2-dimethyl-1,3-dioxane-4-acetate one obtains the title compound.

EXAMPLE 3

(2R-Trans-5-(4-fluorophenyl)-2-(1-methylethyl)-N,4-diphenyl-1-[2-(tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)ethyl]-1H-pyrrole-3-carboxamide.

METHOD A

Step A: Preparation of (R)-1,1-dimethylethyl 6-cyano-5-[[[(1,1-dimethylethyl)dimethylsilyl]oxy]-3-oxohexanoate.

(R)-4-cyano-3-[[[(1,1-dimethylethyl)dimethylsilyl]oxy]butanoic acid, 32 g (0.132 mol), is dissolved in 300 mL of tetrahydrofuran. The solution is cooled to -20°C and carbonyldiimidazole, 27 g (0.165 mol), is added. The solution is stirred and allowed to warm to 25°C over two hours. The solution is added to a slurry of potassium 1,1-dimethylethyl malonate (half ester, half salt), 60 g (0.3 mol), anhydrous magnesium chloride, 27.2 g (0.246 mol), diisopropylethylamine, 53 mL (0.3 mol) in 700 mL of dry acetonitrile. The mixture is stirred at 5°C for 18 hours and at 15°C for 108 hours. The mixture is poured into a mixture of 1 L of 1N hydrochloric acid and 1 L of ethyl acetate and the resulting two-phase system is stirred for 15 minutes. The layers are separated. The organic layer is washed with 500 mL of saturated salt solution and concentrated to yield an oil. The oil consists of (R)-1,1-dimethylethyl 6-cyano-5-[[[(1,1-dimethylethyl)dimethylsilyl]oxy]-3-oxohexanoate and some 1,1-dimethylethyl malonate (half ester, half acid) that is used directly in Step B. The oil has acceptable NMR spectra after subtracting the recovered malonate spectra.

Step B: Preparation of (R)-1,1-dimethylethyl 6-cyano-5-hydroxy-3-oxo-hexanoate.

A solution of crude (R)-1,1-dimethylethyl 6-cyano-5-[[[(1,1-dimethylethyl)dimethylsilyl]oxy]-3-oxohexanoate, 43 g (0.126 mol) in 350 mL of tetrahydrofuran is treated with 213 mL of tetrabutylammonium fluoride solution (1.0 M in hexane). The resulting mixture is stirred for five hours, at 25°C. The mixture is treated with 500 mL of water, 300 mL of diethyl ether is added, and the layers separated. The organic layer is dried (magnesium sulfate) and then filtered through a plug of silica gel with the aid of anhydrous diethyl ether. The solvent is removed under a vacuum to obtain 21 g of crude (R)-1,1-dimethylethyl 6-cyano-5-hydroxy-3-oxohexanoate with acceptable NMR, MS and IR spectra.

200 MHz NMR (CDCl₃) δ 1.48 (s, 9H), 2.62 (m, 2H), 2.89 (d, 2H, J = 6.1), 3.43 (s, 2H), 4.41 (pentet, 2H, J = 6.1 Hz)

¹³C-NMR (CDCl₃, 50 MHz) δ 25.05, 27.86, 48.03, 50.81, 63.39, 82.43, 117.03, 165.84, 202.03.

MS (Chemical ionization) m/e 228, 200, 172, 154.

FTIR (KBr) 1144.5, 1327.2, 1370.9, 1715.5, 1733.5, 2253.1, 2934.6, 2980.9, 3459.3 cm⁻¹.

Step C: Preparation of [R-(R*,R*)]-1,1-dimethylethyl 6-cyano-3,5-dihydroxyhexanoate.

Crude (R)-1,1-dimethylethyl 6-cyano-5-hydroxy-3-oxohexanoate, 21 g (0.0924 mol), is dissolved in 940 mL of tetrahydrofuran and 190 mL of methanol under a nitrogen atmosphere. This solution is cooled to -85°C and 95 mL of a 15% solution of methoxydiethylborane in tetrahydrofuran is added. The reaction is cooled to -97°C and 6.5 g (0.172 mol) of sodium borohydride is added in 0.5 g portions over 1.5 hours. The reaction is maintained between -93°C and -97°C for 13 hours and allowed to warm to room temperature and stand for 60 hours under a nitrogen atmosphere. The reaction is quenched by the addition of 25 mL (0.395 mol) acetic acid and concentrated by vacuum distillation to an oil. The residue is dissolved with 500 mL methanol, concentrated by vacuum distillation, redissolved with 500 mL methanol and reconcentrated by vacuum distillation to give a dark brown oil. This oil is taken up in 500 mL of ethyl acetate and filtered through a plug of silica gel with the aid of 250 mL of ethyl acetate. The solution is evaporated to give 15 g of crude [R-(R*,R*)]-1,1-dimethylethyl 6-cyano-3,5-dihydroxyhexanoate which is used without further purification.

Step D: Preparation of (4R-cis)-1,1-dimethylethyl 6-cyanomethyl-2,2-dimethyl-1,3-dioxane-4-acetate.

Crude [R-(R*,R*)]-1,1-dimethylethyl 6-cyano-3,5-dihydroxyhexanoate, 15 g (61 mol), is dissolved in 150 mL of 2,2-dimethoxypropane, camphorsulfonic acid is added, and the solution is stirred for 18 hours at room temperature. Concentration and flash chromatography after concentration in vacuo provides 11.8 g of (4R-cis)-1,1-dimethylethyl 6-cyanomethyl-2,2-dimethyl-1,3-dioxane-4-acetate as an off-white solid, mp 64.7-68 °C with acceptable IR, NMR, C-NMR and analysis.

200 MHz NMR (CDCl₃) δ 1.36 (m, 1H), 1.42 (s, 3H), 1.49 (s, 9H), 1.50 (s, 3H), 1.79 (dt, 1H, J = 2.5 Hz, J = 12.1 Hz), 2.40 (dd, 1H, J = 6.2 Hz, J = 15.4 Hz), 2.5 - 2.7 (m, 1H), 2.55 (d, 2H, J = 6.1 Hz), 4.18 (m, 1H), 4.32 (m, 1H).

¹³C-NMR (CDCl₃, 50 MHz) δ 19.74, 25.09, 28.24, 29.88, 35.58, 42.50, 65.20, 65.81, 80.87, 99.48, 116.68, 169.75.

GC/MS m/e 254, 198, 154, 138, 120, 59, 57, 43, 41.

FTIR (KBr) 941.4, 1116.2, 1154.8, 1188.3, 1257.7, 1293.7, 1309.1, 1368.3, 1725.8, 2361.1, 2983.5, 2996.4 cm⁻¹.

Step E: Preparation of (4R-cis)-1,1-dimethylethyl 6-(2-aminoethyl)-2,2-dimethyl-1,3-dioxane-4-acetate.

A solution of (4R-cis)-1,1-dimethylethyl 6-cyanomethyl-2,2-dimethyl-1,3-dioxane-4-acetate, 5.63 g (0.048 mol), in 100 mL of methanol saturated with gaseous ammonia is treated with 0.5 g of Raney nickel #30 and hydrogen gas in a shaker at 3.45 x 10⁵ Pa (50 psi) and 40 °C. After 16 hours, thin layer chromatography indicates no starting nitrile present. The suspension is cooled, filtered through filter aid, and concentrated to an oil. This crude oil is purified by flash chromatography on silica gel with 30:20:1 (ethyl acetate:methanol:ammonium hydroxide) as eluent to give 4.93 g of (4R-cis)-1,1-dimethylethyl 6-(2-aminoethyl)-2,2-dimethyl-1,3-dioxane-4-acetate (98.2 area %) as a clear oil with acceptable IR, NMR, C-NMR and MS spectra.

200 MHz NMR (CDCl₃) 1.0 - 1.2 (m, 1H), 1.22 (s, 3H), 1.31 (s, 12 H), 1.35 - 1.45 (m, 3H), 2.15 (dd, 1H, J = 15.1 Hz, J = 6.2 Hz), 2.29 (dd, 1H, J = 15.1 Hz, J = 7.0 Hz), 2.66 (t, 2H, J = 6.6 Hz), 3.82 (m, 1H), 4.12 (m, 1H).

¹³C-NMR (CDCl₃, 50 MHz) δ 19.60, 27.96, 30.00, 36.50, 38.25, 39.79, 42.61, 66.08, 67.18, 80.21, 98.35, 169.82.

GC/MS m/e 202, 200, 173, 158, 142, 140, 114, 113, 100, 99, 97, 72, 57.

FTIR (neat) 951.6, 1159.9, 1201.1, 1260.3, 1314.3, 1368.3, 1381.2, 1731.0, 2870.3, 2939.8, 2980.9, 3382.2 cm⁻¹.

Step F: Preparation of (4R-cis)-1,1-dimethylethyl 6-[2[2-(4-fluorophenyl)-5-(1-methylethyl)-3-phenyl-4-[(phenylamino)carbonyl]-1H-pyrrol-1-yl]ethyl]-2,2-dimethyl-1,3-dioxane-4-acetate.

A solution of (4R-cis)-1,1-dimethylethyl 6-(2-aminoethyl)-2,2-dimethyl-1,3-dioxane-4-acetate, 1.36 g (4.97 mol), and (±)-4-fluoro-α-[2-methyl-1-oxopropyl]-γ-oxo-N,β-diphenylbenzenebutaneamide mixture of [R-(R*,R*)], [R-(R*,S*)], [S-(R*,R*)] and [S-(R*,S*)] isomers, 1.60 g (3.83 mol), in 50 mL of heptane:toluene (9:1) is heated at reflux for 24 hours. The solution is cooled slightly and 15 mL of 2-propanol added. The mixture is allowed to cool to 25 °C and filtered to give 1.86 g of (4R-cis)-1,1-dimethylethyl 6-[2[2-(4-fluorophenyl)-5-(1-methylethyl)-3-phenyl-4-[(phenylamino)carbonyl]-1H-pyrrol-1-yl]ethyl]-2,2-dimethyl-1,3-dioxane-4-acetate as a yellow solid with acceptable PNMR & C-NMR spectra.

¹H-NMR (CDCl₃, 200 MHz) δ 1 - 1.7 (m, 5H), 1.30 (s, 3H), 1.36 (s, 3H), 1.43 (s, 9H), 1.53 (d, 6H, J = 7.1 Hz), 2.23 (dd, 1H, J = 15.3 Hz, J = 6.3 Hz), 2.39 (dd, 1H, J = 15.3 Hz, J = 6.3 Hz), 3.5 - 3.9 (m, 3H), 4.0 - 4.2 (m, 2H), 6.8 - 7.3 (m, 14H).

¹³C-NMR (CDCl₃, 50 MHz) δ 19.69, 21.60, 21.74, 26.12, 27.04, 28.12, 29.95, 36.05, 38.10, 40.89, 42.54, 65.92, 66.46, 80.59, 98.61, 115.00, 115.34, 115.42, 119.52, 121.78, 123.36, 126.44, 128.21, 128.31, 128.52, 128.75, 130.43, 133.01, 133.17, 134.69, 138.38, 141.47, 159.72, 164.64, 169.96.

**Step G: Preparation of (2R-trans)-5-(4-fluorophenyl)-
2-(1-methylethyl)-N,4-diphenyl-1-[2-(tetra-
hydro-4-hydroxy-6-oxo-2H-pyran-2-yl)ethyl]-
1H-pyrrole-3-carboxamide.**

(4R-cis)-1,1-dimethylethyl 6-[2[2-(4-fluorophenyl)-5-(1-methylethyl)-3-phenyl-4-[(phenylamino)carbonyl]-
1H-pyrrol-1-yl]ethyl]-2,2-dimethyl-1,3-dioxane-4-acetate, 4.37 g (6.68 mol), is dissolved in 200 mL of
tetrahydrofuran and 15 mL of 10% hydrochloric acid solution is added, and the solution is stirred for 15
hours. To this solution is added sodium hydroxide (3.6 g) and the mixture is stirred for 30 hours. The
reaction is stopped by adding 150 mL of water, 90 mL of hexane, and separating the layers. The aqueous
layer is acidified with dilute hydrochloric acid solution, stirred for three hours and extracted with 150 mL of
ethyl acetate. A drop of concentrated hydrochloric acid is added to the ethyl acetate solution and the
solution is allowed to stand 18 hours. The solution is concentrated in vacuo and the concentrate is
redissolved in 50 mL of ethyl acetate and treated with one drop of concentrated hydrochloric acid. The
solution is stirred two hours, concentrated in vacuo, and dissolved in 3.0 mL of toluene. (2R-trans)-5-(4-
fluorophenyl)-2-(1-methylethyl)-N,4-diphenyl-1-[2-(tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)ethyl]-1H-
pyrrole-3-carboxamide (3.01 g) is isolated in two crops.

METHOD B

A solution of (4R-cis)-1,1-dimethylethyl 6-(2-aminoethyl)-2,2-dimethyl-1,3-dioxane-4-acetate; 2.56 g (9.36
mol), and (±)-4-fluoro-α-[2-methyl-1-oxopropyl]-γ-oxo-N,β-diphenylbenzenebutaneamide mixture of [R-
(R*,R*)], [R-(R*,S*)], [S-(R*,R*)] and [S-(R*,S*)] isomers, 3.00 g (7.20 mol), in 60 mL of heptane:toluene (9:1) is
heated at reflux for 24 hours. The solution is cooled and poured into 300 mL of tetrahydrofuran and 150 mL
of saturated ammonium chloride in water. The layers are separated and the organic layer is added to 15 mL
of 10% hydrochloric acid solution and the solution is stirred for 15 hours. To this solution is added sodium
hydroxide (3.6 g) and the mixture is stirred for 30 hours. The reaction is stopped by adding 150 mL of
water, 90 mL of hexane, and separating the layers. The aqueous layer is acidified with dilute hydrochloric
acid solution, stirred for three hours and extracted with 150 mL of ethyl acetate. A drop of concentrated
hydrochloric acid is added to the ethyl acetate solution and the solution is allowed to stand 18 hours. The
solution is concentrated in vacuo and the concentrate is redissolved in 50 mL of ethyl acetate and treated
with one drop of concentrated hydrochloric acid. The solution is stirred two hours, concentrated in vacuo,
and dissolved in 3.0 mL of toluene. (2R-trans)-5-(4-fluorophenyl)-2-(1-methylethyl)-N,4-diphenyl-1-[2-
(tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)ethyl]-1H-pyrrole-3-carboxamide (2.92 g) is isolated in two
crops.

PREPARATION OF STARTING MATERIALS

EXAMPLE A

1-(4-Fluorophenyl)1,4-hexanedione.

A mixture of 36.61 g (295 mmol) of 4-fluorobenzaldehyde, 25 g (297.2 mmol) of ethyl vinyl ketone, 29
mL (206.9 mmol) of triethylamine and 11.94 g (44.25 mmol) of 3-benzyl-5-(2-hydroxyethyl)-4-methyl-
thiazolium chloride is stirred and heated at 70 °C for six hours. The cooled solution is diluted with 2 liters of
diethylether and washed with 2 x 300 mL of water, 2 x 100 mL of 2M hydrochloric acid, 100 mL of water,
200 mL of a saturated solution of sodium bicarbonate and brine. The organic layer is separated, dried
(magnesium sulfate), filtered, and concentrated to provide 55 g of 1-(4-fluorophenyl)-1,4-hexanedione after
recrystallization from methanol; mp 56-57 °C.

EXAMPLE B


4-Methyl-3-oxo-N-phenylpentanamide

A three-necked, 12-L round-bottom flask equipped with a mechanical stirrer, a thermometer and set up for distillation is charged with 2.6 L of toluene, 1.73 kg (12 mol) of methyl 4-methyl-3-oxopentanoate and 72 g (1.18 mol) of ethylene diamine. The mixture is heated to 80°C and charged with 0.49 kg of aniline. The mixture is brought to reflux and distillation started. After 40 minutes a further 0.245 kg of aniline is charged and at 40 minute intervals a further two portions of aniline (0.245 and 0.25 kg) are charged. Distillation is continued for a further one to five hours until a total of 985 mL of solvent is removed. The solution is stirred at room temperature for 16 hours and a further 550 mL of solvent is removed by vacuum distillation (using approximately 2.67×10^3 Pa (20 mm Hg)). The mixture is cooled and 2 L of water is charged to provide an oil. The mixture is warmed to 40°C and a further 1.0 L of water is charged. Seven hundred milliliters of toluene-water mixture is removed by vacuum distillation (using approximately 2.67×10^3 Pa (20 mm Hg)). Two liters of water is charged and the mixture is allowed to stand for 10 days. The product is isolated by filtration and washed with three portions of hexane. Drying in vacuo gives 1.7 kg of 4-methyl-3-oxo-N-phenylpentanamide as a hydrate; mp 46.5-58.8°C.

HPLC: 98.8% - retention time 3.56 min. 65/35 acetonitrile/water on a dry basis.
VPC: 87.6% - retention time 12.43 min. also 10.8% aniline (decomposition).

Claims

1. A process for the preparation of a compound of Formula I



I

and a dihydroxy acid and pharmaceutically acceptable salts thereof, corresponding to the opened lactone ring of a compound of Formula I
wherein R: is

- 1-naphthyl,
- 2-naphthyl,
- cyclohexyl,
- cyclohexylmethyl,
- norbornenyl,
- phenyl,
- phenyl substituted with
- fluorine,
- chlorine,
- bromine,
- hydroxyl,
- trifluoromethyl,
- alkyl of from one to four carbon atoms,
- alkoxy of from one to four carbon atoms, or
- alkanoyloxy of from two to eight carbon atoms,
- benzyl,
- 2-, 3-, or 4-pyridinyl, or




EXAMPLE B

4-Methyl-3-oxo-N-phenylpentanamide

- 5 A three-necked, 12-L round-bottom flask equipped with a mechanical stirrer, a thermometer and set up for distillation is charged with 2.6 L of toluene, 1.73 kg (12 mol) of methyl 4-methyl-3-oxopentanoate and 72 g (1.18 mol) of ethylene diamine. The mixture is heated to 80 °C and charged with 0.49 kg of aniline. The mixture is brought to reflux and distillation started. After 40 minutes a further 0.245 kg of aniline is charged and at 40 minute intervals a further two portions of aniline (0.245 and 0.25 kg) are charged. Distillation is
 10 continued for a further one to five hours until a total of 985 mL of solvent is removed. The solution is stirred at room temperature for 16 hours and a further 550 mL of solvent is removed by vacuum distillation (using approximately 2.67×10^3 Pa (20 mm Hg)). The mixture is cooled and 2 L of water is charged to provide an oil. The mixture is warmed to 40 °C and a further 1.0 L of water is charged. Seven hundred milliliters of toluene-water mixture is removed by vacuum distillation (using approximately 2.67×10^3 Pa (20 mm Hg)).
 15 Two liters of water is charged and the mixture is allowed to stand for 10 days. The product is isolated by filtration and washed with three portions of hexane. Drying in vacuo gives 1.7 kg of 4-methyl-3-oxo-N-phenylpentanamide as a hydrate; mp 46.5-58.8 °C.
 HPLC: 98.8% - retention time 3.56 min. 65/35 acetonitrile/water on a dry basis.
 VPC: 87.6% - retention time 12.43 min. also 10.8% aniline (decomposition).

Claims

1. A process for the preparation of a compound of Formula I



I

- and a dihydroxy acid and pharmaceutically acceptable salts thereof, corresponding to the opened lactone ring of a compound of Formula I
 40 wherein R: is

- 1-naphthyl,
 2-naphthyl,
 cyclohexyl,
 45 cyclohexylmethyl,
 norbornenyl,
 phenyl,
 phenyl substituted with
 fluorine,
 50 chlorine,
 bromine,
 hydroxyl,
 trifluoromethyl,
 alkyl of from one to four carbon atoms,
 55 alkoxy of from one to four carbon atoms, or
 alkanoyloxy of from two to eight carbon atoms,
 benzyl,
 2-, 3-, or 4-pyridinyl, or



2-, 3-, or 4-pyridinyl-N-oxide;

R₂ or R₃ is independently

hydrogen,
alkyl of from one to six carbon atoms,
cyclopropyl,
cyclobutyl,
cyclopentyl,
cyclohexyl,
phenyl,
phenyl substituted with
fluorine,
chlorine,
bromine,
hydroxyl,
trifluoromethyl,
alkyl of from one to four carbon atoms, or
alkoxy of from one to four carbon atoms, cyano,
trifluoromethyl, or -CONR₅R₆ where R₅ and R₆ are
independently

hydrogen,
alkyl of from one to six carbon atoms,
phenyl,
phenyl substituted with
fluorine,
chlorine,
bromine,
cyano, or
trifluoromethyl;

R₄ is

alkyl of from one to six carbon atoms,
cyclopropyl,
cyclobutyl,
cyclopentyl,
cyclohexyl, or
trifluoromethyl;


which comprises:

(a) reacting 1,6-heptadien-4-ol with an

(1) alkyl lithium

(2) followed by iodine and carbon dioxide and

(3) treating the resulting iodocarbonate intermediate in situ with a base in an aqueous alcohol at
about 0 °C to about 40 °C to afford a compound of Formula IX;




IX

(b) treating the compound of Formula IX with

(1) an alkali cyanide at about 0 °C to about 40 °C and

(2) reacting the resulting diol intermediate in situ with a ketal-forming reagent in the presence of
an acid to afford a compound of Formula VIII




VIII

wherein R_7 and R_8 are independently hydrogen, alkyl of from one to three carbon atoms, phenyl or R_7 and R_8 taken together as $-(\text{CH}_2)_n-$, wherein n is 4 or 5;

(c) treating the compound of Formula VIII with

(1) ozone in an inert solvent and


(2) reacting the resulting intermediate in situ with oxygen and triphenylphosphine at about -20°C to about -78°C to afford a compound of Formula VII



VII

wherein R_7 and R_8 are as defined above;

(d) treating the compound of Formula VII with an oxidizing reagent at about 0°C to afford a compound of Formula VI




VI

wherein R_7 and R_8 are as defined above;

(e) treating the compound of Formula VI with a compound of Formula

Hal-R_{9a}


wherein Hal is halogen and R_{9a} is alkyl of from one to eight carbon atoms or a three- to six-membered cycloalkyl group, in the presence of a base to afford a compound of Formula V_a

 V_a

wherein R_7 , R_8 , and R_{9a} are as defined above; or treating the compound of Formula VI with a compound of Formula


$\text{HO}-R_{9b}$

wherein R_{9b} is tertiary butyl, tertiary amyl, or α , α -dimethylbenzyl in the presence of an activating agent, a catalytic amount of a base and an inert solvent to afford a compound of Formula V_b


 V_b

wherein R_7 , R_8 , and R_{9b} are as defined above;

(f) treating the compound of Formula V_a with hydrogen in the presence of a catalyst and an acid at about 0°C to about 70°C to afford a compound of Formula IV_a


 IV_a

wherein R_7 , R_8 , and R_{9a} are as defined above, or treating the compound of Formula V_b with hydrogen in the presence of a catalyst and an acid or a catalyst and a base at about 0°C to about 70°C to afford a compound of Formula IV_b

IV_b


wherein R₇, R₈, and R_{9b} are as defined above;

(g) treating the compound of Formula IV_a with a compound of Formula III




III

wherein R₁, R₂, R₃, and R₄ are as defined above in an inert solvent to afford a compound of Formula II_a

II_a

wherein R₁, R₂, R₃, R₄, R₇, R₈, and R_{9a} are as defined above, or treating the compound of Formula IV_b with a compound of Formula III in an inert solvent to afford a compound of Formula II_b.

II_b

wherein R₁, R₂, R₃, R₄, R₇, R₈, and R_{9b} are as defined above;

(h) and finally treating a compound of Formula II_a with



- (1) an acid in the presence of an inert solvent
 - (2) followed by hydrolysis with a base
 - (3) followed by neutralization with an acid and
 - (4) dissolution and/or heating in an inert solvent with concomitant removal of water to give a compound of Formula I, or treating a compound of Formula II_b with
 - (1) an acid in the presence of an inert solvent
 - (2) followed by addition of a base
 - (3) followed by neutralization with an acid and
 - (4) dissolution and/or heating in an inert solvent with concomitant removal of water to give a compound of Formula I;
 - (i) and if desired converting the resulting compound of Formula I to a dihydroxy acid corresponding to the opened lactone ring of structural Formula I by conventional hydrolysis and further, if desired converting the dihydroxy acid to a corresponding pharmaceutically acceptable salt by conventional means, and if so desired converting the corresponding pharmaceutically acceptable salt to a dihydroxy acid by conventional means, and if so desired converting the dihydroxy acid to a compound of Formula I by heating in an inert solvent.
2. A process according to Claim 1 wherein the alkyl lithium in step (a) is n-butyllithium.
 3. A process according to Claim 1 wherein the alkali cyanide in step (b) is potassium cyanide.
 4. A process according to Claim 1 wherein the oxidizing reagent in step (d) is chromium trioxide-sulfuric acid-water.
 5. A process according to Claim 1 wherein the catalyst in step (f) for preparing a compound of Formula IV_a is platinum dioxide.
 6. A process according to Claim 1 wherein the catalyst and base in Step (f) for preparing a compound of Formula IV_b are Raney nickel and anhydrous ammonia, respectively.
 7. A process according to Claim 1 wherein the acid in step (h) is hydrochloric acid.
 8. A process according to Claim 1 wherein the base in step (h) is sodium hydroxide.
 9. A process according to Claim 1 wherein R₁ is 1-naphthyl, norbornenyl, phenyl, or phenyl substituted with fluorine, chlorine, bromine, hydroxyl, trifluoromethyl, alkyl of from one to four carbon atoms, alkoxy of from one to four carbon atoms, or alkanoyloxy of from two to eight carbon atoms.
 10. A process according to Claim 9 wherein R₄ is alkyl of from one to six carbon atoms, cyclopropyl, or trifluoromethyl.
 11. A process according to Claim 1 and for the preparation of a compound selected from the group consisting of:
 - trans-6-[2-[2-(4-fluorophenyl)-5-(trifluoromethyl)-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one;
 - trans-6-[2-[2-(4-fluorophenyl)-5-methyl-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one;
 - trans-6-[2-[2-(4-fluorophenyl)-5-(1-methylethyl)-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one;
 - trans-6-[2-[2-cyclopropyl-5-(4-fluorophenyl)-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one;
 - trans-6-[2-[2-(1,1-dimethylethyl)-5-(4-fluorophenyl)-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one;
 - trans-tetrahydro-4-hydroxy-6-[2-[2-(2-methoxyphenyl)-5-methyl-1H-pyrrol-1-yl]ethyl]-2H-pyran-2-one;
 - trans-tetrahydro-4-hydroxy-6-[2-[2-(2-methoxyphenyl)-5-(1-methylethyl)-1H-pyrrol-1-yl]ethyl]-2H-pyran-2-one;
 - trans-tetrahydro-4-hydroxy-6-[2-[2-methyl-5-(1-naphthalenyl)-1H-pyrrol-1-yl]ethyl]-2H-pyran-2-one;



trans-6-[2-(2-bicyclo[2.2.1]hept-5-en-2-yl-5-methyl-1H-pyrrol-1-yl)ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one;


trans(±)-5-(4-fluorophenyl)-2-(1-methylethyl)-N,4-diphenyl-1-[2-(tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)ethyl]-1H-pyrrole-3-carboxamide;

(2R-trans)-5-(4-fluorophenyl)-2-(1-methylethyl)-N,4-diphenyl-1-[2-(tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)ethyl]-1H-pyrrole-3-carboxamide;

trans-2-(4-fluorophenyl)-N,4-diphenyl-1-[2-(tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)ethyl]-5-trifluoromethyl-1H-pyrrole-3-carboxamide; and

trans-5-(4-fluorophenyl)-N,4-diphenyl-1-[2-(tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)ethyl]-2-trifluoromethyl-1H-pyrrole-3-carboxamide.


12. A process according to claim 1 for the preparation of the compound of Formula I_a



I_a


and the hydroxy acid and pharmaceutically acceptable salts thereof, corresponding to the opened lactone ring of the compound of Formula I_a which comprises:

(a) reacting the compound of Formula XVII




XVII

with a compound of Formula



wherein R_7 and R_8 are independently hydrogen alkyl of from one to three carbon atoms, phenyl or R_7 and R_8 are taken together as $-(\text{CH}_2)_n$ -wherein n is 4 or 5, and R_{13} is hydrogen or




in an inert solvent and treating the resulting intermediate with an acid to afford the compound of Formula I_a ;

(b) and if desired, converting the resulting compound of Formula I_a to a hydroxy acid corresponding to the opened lactone ring of structural Formula I_a by conventional hydrolysis and further, if desired, converting the hydroxy acid to a corresponding pharmaceutically acceptable salt by conventional means, and if so desired, converting the hydroxy acid to a compound of Formula I_a by heating in an inert solvent.

13. A process according to Claim 12 wherein the acid in Step (a) is hydrochloric acid.

14. A process according to Claim 12 and for the preparation of (2R-trans)-5-(4-fluorophenyl)-2-(1-methylethyl)-N,4-diphenyl-1-[2-(tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)ethyl]-1H-pyrrole-3-carboxamide.


15. A process for the preparation of the compound of Formula I_a



I_a


and the hydroxy acid and pharmaceutically acceptable salts thereof, corresponding to the opened lactone ring of the compound of Formula I_a which comprises

(a) reacting 4-methyl-3-oxo-N-phenylpentanamide with benzaldehyde in the presence of a catalyst and an inert solvent to afford the compound of Formula XVIII




XVIII

(b) reacting the compound of Formula XVIII with 4-fluorobenzaldehyde in the presence of a catalyst, a base, and an inert solvent to afford the compound of Formula XVII




XVII


(c) reacting the compound of Formula XVII with a compound of Formula



wherein R_{10} and R_{11} are alkyl of one to eight carbon atoms or R_{10} and R_{11} together are




$-\text{CH}_2-\text{CH}_2-$ or $-\text{CH}_2\text{CH}_2\text{CH}_2-$ in the presence of a catalyst and an inert solvent to afford a compound of Formula XVI.



XVI

wherein R_{10} and R_{11} are alkyl of one to eight carbon atoms or R_{10} and R_{11} together are



-CH₂-CH₂- or -CH₂-CH₂-CH₂-

(d) and finally converting a compound of Formula XVI in a conventional manner to afford the compound of Formula I_a.


(e) and if desired, converting the resulting compound of Formula I_a to a hydroxy acid corresponding to the opened lactone ring of structural Formula I_a by conventional hydrolysis and further, if desired, converting the hydroxy acid to a corresponding pharmaceutically acceptable salt by conventional means, and if so desired, converting the hydroxy acid to a compound of Formula I_a by heating in an inert solvent.

16. A process according to Claim 15 wherein the catalyst in step (a) is selected from the group consisting of piperidine and glacial acetic acid, ethylene diamine and glacial acetic acid, and β -alanine and glacial acetic acid.
17. A process according to Claim 16 wherein the catalyst is β -alanine and glacial acetic acid.
18. A process according to Claim 15 wherein the catalyst in step (b) is selected from the group consisting of 3-benzyl-5-(2-hydroxyethyl)-4-methylthiazolium chloride, 3,4-dimethyl-5-(2-hydroxyethyl)thiazolium iodide, 3-ethyl-5-(2-hydroxyethyl)-4-methyl-thiazolium bromide and thiamine hydrochloride.
19. A process according to Claim 18 wherein the catalyst is 3-ethyl-5-(2-hydroxyethyl)-4-methylthiazolium bromide.
20. A process according to Claim 15 wherein the catalyst in step (c) is selected from the group consisting of $R_{12}CO_2H$ wherein R_{12} is CH_3 , CF_3 , $ClCH_2-$, $C_6H_5CH_2CH_2-$, $C_6H_5CH_2-$, HO_2CCH_2- , $HO_2CCH_2CH_2-$, C_6H_5- , para-Cl- C_6H_5- , $ClCH_2CH_2-$, meta- $H_3C-C_6H_5-$, para- $H_3C-C_6H_5-$, or tertiary- C_4H_9- and triethylamine hydrochloride.
21. A process according to Claim 20 wherein the catalyst is tertiary- $C_4H_9CO_2H$.
22. A process according to Claim 15 wherein the base in step (b) is selected from the group consisting of N,N-diisopropylethylamine, pyridine, N,N-dimethylaniline, triethylamine, 1,8-diazabicyclo[5.4.0]undec-7-ene, 1,4-diazabicyclo[2.2.2]octane, 4-dimethylaminopyridine, and N,N,N',N'-tetramethylethylenediamine.
23. A process according to Claim 22 wherein the base is triethylamine.



24. A process according to Claim 15 and for the preparation of (2R-trans-5-(4-fluorophenyl)-2-(1-methylethyl)-N,4-diphenyl-1-[2-(tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)ethyl]-1H-pyrrole-3-carboxamide.

25. A compound of Formula II



II

wherein R_1 , R_2 , R_3 , R_4 , R_7 and R_8 are as defined in claim 1 and R_9 is alkyl of from one to eight carbon atoms, a three- to six-membered cycloalkyl group, or α,α -dimethylbenzyl.


26. A compound according to Claim 25 wherein R_7 and R_8 are methyl, R_9 is tertiary butyl, and the two optically active centers are R.

27. A compound according to Claim 25 wherein R_1 is 4-fluorophenyl, R_2 and R_3 are hydrogen, R_4 is ethyl, R_7 and R_8 are methyl, and R_9 is isopropyl.

28. A compound according to Claim 25 wherein R_1 is 4-fluorophenyl, R_2 is phenyl, R_3 is C_6H_5NHCO- , R_4 is isopropyl, R_7 and R_8 are methyl, and R_9 is tertiary butyl.


29. A compound according to Claim 28 wherein the two optically active centers are R.

30. A compound of Formula XVI



XVI


wherein R_{10} and R_{11} are alkyl of one to eight carbon atoms or R_{10} and R_{11} together are






or $-\text{CH}_2-\text{CH}_2-\text{CH}_2-$.

31. A compound according to Claim 30 wherein R_{10} and R_{11} are methyl or ethyl or R_{10} and R_{11} together are




32. A compound of Formula IV



IV

wherein R_7 and R_8 are independently hydrogen, alkyl of from one to three carbon atoms, phenyl or R_7 and R_8 are taken together as $-(\text{CH}_2)_n-$, wherein n is 4 or 5, and R_9 is alkyl of from one to eight carbon atoms, a three- to six-membered cycloalkyl group, or α,α -dimethylbenzyl.

33. A compound according to Claim 32 wherein R_9 is




34. A compound according to Claim 32 wherein R_7 and R_8 are methyl, and R_9 is isopropyl.

35. A compound according to Claim 32 wherein R_7 and R_8 are methyl, and R_9 is tertiary butyl.

36. A compound according to Claim 32 selected from the group consisting of (4R-cis)-1,1-dimethylethyl 6-(2-aminoethyl)-2,2-dimethyl-1,3-dioxane-4-acetate; [2R-(2 α ,4 α ,6 α)]-1,1-dimethylethyl 6-(2-aminoethyl)-2-phenyl-1,3-dioxane-4-acetate; [2S-(2 α ,4 β ,6 β)]-1,1-dimethylethyl 6-(2-aminoethyl)-2-phenyl-1,3-dioxane-4-acetate; (7R-cis)-1,1-dimethylethyl 9-(2-aminoethyl)-6,10-dioxaspiro[4.5]decane-7-acetate; (2R-cis)-1,1-dimethylethyl 4-(2-aminoethyl)-1,5-dioxaspiro[5.5]undecane-2-acetate; (4R-cis)-1,1-dimethylethyl 4-(2-aminoethyl)-1,3-dioxane-4-acetate; [2R-(2 α ,4 α ,6 α)]-1,1-dimethylethyl 6-(2-aminoethyl)-2-methyl-1,3-dioxane-4-acetate; and [2S-(2 α ,4 β ,6 β)]-1,1-dimethylethyl 6-(2-aminoethyl)-2-methyl-1,3-dioxane-4-acetate.



37. A compound of Formula XXI




XXI

wherein R_7 and R_8 are independently hydrogen alkyl of from one to three carbon atoms, phenyl or R_7 and R_8 are taken together as $-(CH_2)_n-$, wherein n is 4 or 5.

38. A compound according to Claim 37 selected from the group consisting of (4R-cis)-6-(2-aminoethyl)-2,2-dimethyl-1,3-dioxane-4-acetic acid; [2R-(2 α , 4 α , 6 α)]-6-(2-aminoethyl)-2-phenyl-1, 3-dioxane-4-acetic acid; [2S-(2 α , 4 β , 6 β)]-6-(2-aminoethyl)-2-phenyl-1,3-dioxane-4-acetic acid; (7R-cis)-9-(2-aminoethyl)-6,10-dioxaspiro [4.5]decane-7-acetic acid; (2R-cis)-4-(2-aminoethyl)-1,5-dioxaspiro[5.5]undecane-2-acetic acid; (4R-cis)-4-(2-aminoethyl)-1,3-dioxane-4-acetic acid; [2R-(2 α , 4 α , 6 α)]-6-(2-aminoethyl)-2-methyl-1,3-dioxane-4-acetic acid; and [2S-(2 α , 4 β , 6 β)]-6-(2-aminoethyl)-2-methyl-1,3-dioxane-4-acetic acid.

39. A compound of Formula V



V


wherein R_7 and R_8 are independently hydrogen, alkyl of from one to three carbon atoms, phenyl or R_7 and R_8 are taken together as $-(CH_2)_n-$, wherein n is 4 or 5, and R_9 is alkyl of from one to eight carbon atoms, a three- to six-membered cycloalkyl group, or α, α -dimethylbenzyl.

40. A compound according to Claim 39 wherein R_7 and R_8 are methyl, and R_9 is isopropyl.

41. A compound according to Claim 39 wherein R_7 and R_8 are methyl, R_9 is tertiary butyl, and the two optically active centers are R.



42. A compound of Formula VI




VI

wherein R_7 and R_8 are independently hydrogen, alkyl of from one to three carbon atoms, phenyl or R_7 and R_8 are taken together as $-(\text{CH}_2)_n-$, wherein n is 4 or 5.

43. A compound according to Claim 42 wherein R_7 and R_8 are methyl and the two optically active centers are R.

44. A compound of Formula VII




VII

wherein R_7 and R_8 are independently hydrogen, alkyl of from one to three carbon atoms, phenyl or R_7 and R_8 are taken together as $-(\text{CH}_2)_n-$, wherein n is 4 or 5.

45. A compound according to Claim 44 wherein R_7 and R_8 are methyl and the two optically active centers are R.

46. A compound of Formula VIII



VIII


wherein R_7 and R_8 are independently hydrogen, alkyl of from one to three carbon atoms, phenyl or R_7



and R_8 are taken together as $-(CH_2)_n-$, wherein n is 4 or 5.


47. A compound according to Claim 46 wherein R_7 and R_8 are methyl and the two optically active centers are R.

48. A compound of Formula XVII



XVII


49. A mixture of compounds of Formula XVIII



XVIII

Patentansprüche

1. Verfahren zur Herstellung einer Verbindung der Formel I



I

und einer Dihydroxysäure und pharmazeutisch akzeptabler Salze hiervon, entsprechend dem geöffne-



ten Lactonring einer Verbindung der Formel I,
worin bedeuten:

R₁ 1-Naphthyl; 2-Naphthyl; Cyclohexyl; Cyclohexylmethyl; Norbornenyl; Phenyl; Phenyl, substituiert durch Fluor, Chlor, Brom, Hydroxyl, Trifluormethyl, Alkyl mit 1 bis 4 Kohlenstoffatom(en), Alkoxy mit 1 bis 4 Kohlenstoffatom(en) oder Alkanoyloxy mit 2 bis 8 Kohlenstoffatomen; Benzyl; 2-, 3- oder 4-Pyridinyl oder 2-, 3- oder 4-Pyridinyl-N-oxid;

R₂ oder R₃ unabhängig voneinander Wasserstoff; Alkyl mit 1 bis 6 Kohlenstoffatom(en); Cyclopropyl; Cyclobutyl; Cyclopentyl; Cyclohexyl; Phenyl; Phenyl, substituiert durch Fluor, Chlor, Brom, Hydroxyl, Trifluormethyl, Alkyl mit 1 bis 4 Kohlenstoffatom(en) oder Alkoxy mit 1 bis 4 Kohlenstoffatom(en); Cyano; Trifluormethyl oder -CONR₅R₆ mit R₅ und R₆ unabhängig voneinander jeweils gleich Wasserstoff, Alkyl mit 1 bis 6 Kohlenstoffatom(en), Phenyl, Phenyl, substituiert durch Fluor, Chlor, Brom, Cyano oder Trifluormethyl;

R₄ Alkyl mit 1 bis 6 Kohlenstoffatom(en); Cyclopropyl; Cyclobutyl; Cyclopentyl; Cyclohexyl oder Trifluormethyl;


durch

(a) Umsetzen von 1,6-Heptadien-4-ol mit einem

(1) Alkylolithium und anschließend

(2) Jod und Kohlendioxid und

(3) Behandeln des gebildeten Jodcarbonatzwischenprodukts in situ mit einer Base in einem wäßrigen Alkohol bei etwa 0 °C bis etwa 40 °C zur Bildung einer Verbindung der Formel IX




IX

(b) Behandeln der Verbindung der Formel IX mit

(1) einem Alkylcyanid bei etwa 0 °C bis etwa 40 °C und

(2) Umsetzen des gebildeten Diolzwischenprodukts in situ mit einem Ketalbildner in Gegenwart einer Säure zur Bildung einer Verbindung der Formel VIII




VIII

worin R₇ und R₈ unabhängig voneinander für Wasserstoff, Alkyl mit 1 bis 3 Kohlenstoffatom(en) oder Phenyl oder R₇ und R₈ zusammen -(CH₂)_n- mit n gleich 4 oder 5 stehen;

(c) Behandeln der Verbindung der Formel VIII mit

(1) Ozon in einem inerten Lösungsmittel und


(2) Umsetzen des gebildeten Zwischenprodukts in situ mit Sauerstoff und Triphenylphosphin bei etwa -20 °C bis etwa -78 °C zur Bildung einer Verbindung der Formel VII



VII

worin R_7 und R_8 die zuvor angegebene Bedeutung besitzen;

(d) Behandeln der Verbindung der Formel VII mit einem Oxidationsmittel bei etwa 0°C zur Bildung einer Verbindung der Formel VI



VI


worin R_7 und R_8 die zuvor angegebene Bedeutung besitzen;

(e) Behandeln der Verbindung der Formel VI mit einer Verbindung der Formel:



worin Hal für Halogen steht und R_{9a} Alkyl mit 1 bis 8 Kohlenstoffatom(en) oder eine 3- bis 6-gliedrige Cycloalkylgruppe bedeutet,


in Gegenwart einer Base zur Bildung einer Verbindung der Formel V_a

 V_a

worin R_7 , R_8 und R_{9a} die zuvor angegebene Bedeutung besitzen, oder Behandeln der Verbindung der Formel VI mit einer Verbindung der Formel:




worin R_{9b} für tert.-Butyl, tert.-Amyl oder α,α -Dimethylbenzyl steht, in Gegenwart eines Aktivierungsmittels, einer katalytischen Menge einer Base und eines inerten Lösungsmittels zur Bildung einer Verbindung der Formel V_b


V_b

worin R₇, R₈ und R_{9b} die zuvor angegebene Bedeutung besitzen;

(f) Behandeln der Verbindung der Formel V_a mit Wasserstoff in Gegenwart eines Katalysators und einer Säure bei etwa 0°C bis etwa 70°C zur Bildung einer Verbindung der Formel IV_a


IV_a

worin R₇, R₈ und R_{9b} die zuvor angegebene Bedeutung besitzen, oder Behandeln der Verbindung der Formel V_b mit Wasserstoff in Gegenwart eines Katalysators und einer Säure oder eines Katalysators und einer Base bei etwa 0°C bis etwa 70°C zur Bildung einer Verbindung der Formel IV_b

IV_b


worin R₇, R₈ und R_{9b} die zuvor angegebene Bedeutung besitzen;

(g) Behandeln der Verbindung der Formel IV_a mit einer Verbindung der Formel III




III

worin R₁, R₂, R₃ und R₄ die zuvor angegebene Bedeutung besitzen, in einem inerten Lösungsmittel zur Bildung einer Verbindung der Formel II_a



II a-

worin R₁, R₂, R₃, R₄, R₇, R₈ und R_{9b} die zuvor angegebene Bedeutung besitzen, oder Behandeln der Verbindung der Formel IV_b mit einer Verbindung der Formel III in einem inerten Lösungsmittel zur Bildung einer Verbindung der Formel II_b

II_b

worin $R_1, R_2, R_3, R_4, R_7, R_8$ und R_{gh} die zuvor angegebene Bedeutung besitzen; und

(h) schließlich Behandeln einer Verbindung der Formel II_g mit

- (1) einer Säure in Gegenwart eines inerten Lösungsmittels,
- (2) anschließende Hydrolyse mit einer Base,
- (3) anschließende Neutralisation mit einer Säure und

(4) Auflösen und/oder Erwärmen in einem inerten Lösungsmittel unter gleichzeitiger Entfernung von Wasser zur Bildung einer Verbindung der Formel I,

oder Behandeln einer Verbindung der Formel II_b mit

- (1) einer Säure in Gegenwart eines inerten Lösungsmittels,
- (2) anschließende Zugabe einer Base,
- (3) anschließende Neutralisation mit einer Säure und
- (4) Auflösen und/oder Erwärmen in einem inerten Lösungsmittel unter gleichzeitiger Entfernung von Wasser zur Bildung einer Verbindung der Formel I;

(i) und gewünschtenfalls Umwandeln der erhaltenen Verbindung der Formel I in eine Dihydroxysäure entsprechend dem geöffneten Lactonring der Strukturformel I durch übliche Hydrolyse und ferner, gewünschtenfalls, Umwandeln der Dihydroxysäure in ein entsprechendes pharmazeutisch akzeptables Salz in üblicher bekannter Weise und gewünschtenfalls Umwandeln des entsprechenden pharmazeutisch akzeptablen Salzes in eine Dihydroxysäure in üblicher bekannter Weise und gewünschtenfalls Umwandeln der Dihydroxysäure in eine Verbindung der Formel I durch Erwärmen in einem inerten Lösungsmittel.

- 2. Verfahren nach Anspruch 1, wobei das Alkylolithium in Stufe (a) aus n-Butyllithium besteht.**

- 3. Verfahren nach Anspruch 1, wobei das Alkalicyanid in Stufe (b) aus Kaliumcyanid besteht.**


- 4. Verfahren nach Anspruch 1, wobei das Oxidationsmittel in Stufe (d) aus Chromtrioxid/Schwefelsäure/Wasser besteht.**



5. Verfahren nach Anspruch 1, wobei der Katalysator in Stufe (f) zur Herstellung einer Verbindung der Formel IV_a aus Platindioxid besteht.
6. Verfahren nach Anspruch 1, wobei der Katalysator bzw. die Base in Stufe (f) zur Herstellung einer Verbindung der Formel IV_b aus Raney-Nickel bzw. wasserfreiem Ammoniak besteht.
7. Verfahren nach Anspruch 1, wobei die Säure in Stufe (h) aus Chlorwasserstoffsäure besteht.
8. Verfahren nach Anspruch 1, wobei die Base in Stufe (h) aus Natriumhydroxid besteht.
9. Verfahren nach Anspruch 1, wobei R₁ für 1-Naphthyl, Norbornenyl, Phenyl oder Phenyl, substituiert durch Fluor, Chlor, Brom, Hydroxyl, Trifluormethyl, Alkyl mit 1 bis 4 Kohlenstoffatom(en), Alkoxy mit 1 bis 4 Kohlenstoffatom(en) oder Alkanoyloxy mit 2 bis 8 Kohlenstoffatomen steht.
10. Verfahren nach Anspruch 9, wobei R₄ für Alkyl mit 1 bis 6 Kohlenstoffatom(en), Cyclopropyl oder Trifluormethyl steht.
11. Verfahren nach Anspruch 1 zur Herstellung einer Verbindung, ausgewählt aus der Gruppe:
 - trans-6-[2-[2-(4-Fluorphenyl)-5-(trifluormethyl)-1H-pyrrol-1-yl]-ethyl]-tetrahydro-4-hydroxy-2H-pyran-2-on;
 - trans-6-[2-[2-(4-Fluorphenyl)-5-methyl-1H-pyrrol-1-yl]-ethyl]-tetrahydro-4-hydroxy-2H-pyran-2-on;
 - trans-6-[2-[2-(4-Fluorphenyl)-5-(1-methylethyl)-1H-pyrrol-1-yl]-ethyl]-tetrahydro-4-hydroxy-2H-pyran-2-on;
 - trans-6-[2-[2-Cyclopropyl-5-(4-fluorphenyl)-1H-pyrrol-1-yl]-ethyl]-tetrahydro-4-hydroxy-2H-pyran-2-on;
 - trans-6-[2-[2-(1,1-Dimethylethyl)-5-(4-fluorphenyl)-1H-pyrrol-1-yl]-ethyl]-tetrahydro-4-hydroxy-2H-pyran-2-on;
 - trans-Tetrahydro-4-hydroxy-6-[2-[2-(2-methoxyphenyl)-5-methyl-1H-pyrrol-1-yl]-ethyl]-2H-pyran-2-on;
 - trans-Tetrahydro-4-hydroxy-6-[2-[2-(2-methoxyphenyl)-5-(1-methylethyl)-1H-pyrrol-1-yl]-ethyl]-2H-pyran-2-on;
 - trans-Tetrahydro-4-hydroxy-6-[2-[2-methyl-5-(1-naphthalenyl)-1H-pyrrol-1-yl]-ethyl]-2H-pyran-2-on;
 - trans-6-[2-(2-Bicyclo[2.2.1]hept-5-en-2-yl)-5-methyl-1H-pyrrol-1-yl]-ethyl]-tetrahydro-4-hydroxy-2H-pyran-2-on;
 - trans(±)-5-(4-Fluorphenyl)-2-(1-methylethyl)-N,4-diphenyl-1-[2-(tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)-ethyl]-1H-pyrrol-3-carboxamid;
 - (2R-trans)-5-(4-Fluorphenyl)-2-(1-methylethyl)-N,4-diphenyl-1-[2-(tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)-ethyl]-1H-pyrrol-3-carboxamid;
 - trans-2-(4-Fluorphenyl)-N,4-diphenyl-1-[2-(tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)-ethyl]-5-trifluormethyl-1H-pyrrol-3-carboxamid und
 - trans-5-(4-Fluorphenyl)-N,4-diphenyl-1-[2-(tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)-ethyl]-5-trifluormethyl-1H-pyrrol-3-carboxamid.




12. Verfahren nach Anspruch 1 zur Herstellung einer Verbindung der Formel I_a




und der Hydroxysäure und pharmazeutisch akzeptabler Salze hiervon, entsprechend dem geöffneten Lactonring der Verbindung der Formel I_a durch:


(a) Umsetzen der Verbindung der Formel XVII



mit einer Verbindung der Formel



worin R₇ und R₈ unabhängig voneinander für Wasserstoff, Alkyl mit 1 bis 3 Kohlenstoffatom(en) oder Phenyl stehen oder R₇ und R₈ zusammen -(CH₂)_n- mit n = 4 oder 5 bilden und R₁₃ Wasserstoff oder



5

darstellt, in einem inerten Lösungsmittel und Behandeln des gebildeten Zwischenprodukts mit einer Säure zur Bildung einer Verbindung der Formel I_a;

10

(b) und gewünschtenfalls Umwandeln der gebildeten Verbindung der Formel I_a in eine Hydroxysäure entsprechend dem geöffneten Lactonring der Strukturformel I_a durch übliche Hydrolyse und ferner gewünschtenfalls Umwandeln der Hydroxysäure in ein entsprechendes pharmazeutisch akzeptables Salz in üblicher bekannter Weise und gewünschtenfalls Umwandeln der Hydroxysäure in eine Verbindung der Formel I_a durch Erwärmen in einem inerten Lösungsmittel.

15

13. Verfahren nach Anspruch 12, wobei die Säure in Stufe (a) aus Chlorwasserstoffsäure besteht.

14. Verfahren nach Anspruch 12 zur Herstellung von (2R-trans)-5-(4-Fluorphenyl)-2-(1-methylethyl)-N,4-diphenyl-1-[2-(tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)-ethyl]-1H-pyrrol-3-carboxamid.


20

15. Verfahren zur Herstellung der Verbindung der Formel I_a

25

30

35

I_a

40


und der Hydroxysäure und pharmazeutisch akzeptabler Salze hiervon, entsprechend dem geöffneten Lactonring der Verbindung der Formel I_a durch

(a) Umsetzen von 4-Methyl-3-oxo-N-phenylpentanamid mit Benzaldehyd in Gegenwart eines Katalysators und eines inerten Lösungsmittels zu einer Verbindung der Formel XVIII

45


50

55




XVIII

(b) Umsetzen der Verbindung der Formel XVIII mit 4-Fluorbenzaldehyd in Gegenwart eines Katalysators, einer Base und eines inerten Lösungsmittels zur Bildung der Verbindung der Formel XVII




XVII

(c) Umsetzen der Verbindung der Formel XVII mit einer Verbindung der Formel:



worin R_{10} und R_{11} für Alkyl mit 1 bis 8 Kohlenstoffatom(en) stehen oder R_{10} und R_{11} zusammen




$-\text{CH}_2-\text{CH}_2-$ oder $-\text{CH}_2\text{CH}_2\text{CH}_2-$ bilden, in Gegenwart eines Katalysators und eines inerten Lösungsmittels zur Bildung einer Verbindung der Formel XVI



5

10


15



XVI

worin R_{10} und R_{11} für Alkyl mit 1 bis 8 Kohlenstoffatom(en) stehen oder R_{10} und R_{11} zusammen

20



25

$-\text{CH}_2-\text{CH}_2-$ oder $-\text{CH}_2-\text{CH}_2-\text{CH}_2-$ bilden;

(d) und schließlich übliches bekanntes Umwandeln einer Verbindung der Formel XVI in die Verbindung der Formel Ia;

(e) und gewünschtenfalls Umwandeln der gebildeten Verbindung der Formel Ia in eine Hydroxysäure entsprechend dem geöffneten Lactonring der Strukturformel Ia durch übliche Hydrolyse und ferner gewünschtenfalls Umwandeln der Hydroxysäure in ein entsprechendes pharmazeutisch akzeptables Salz in üblicher bekannter Weise und gewünschtenfalls Umwandeln der Hydroxysäure in eine Verbindung der Formel Ia durch Erwärmen in einem inerten Lösungsmittel.

30

35

16. Verfahren nach Anspruch 15, wobei der Katalysator in Stufe (a) aus der Gruppe Piperidin und Eisessig, Ethylendiamin und Eisessig und β -Alanin und Eisessig ausgewählt ist.

17. Verfahren nach Anspruch 16, wobei der Katalysator aus β -Alanin und Eisessig besteht.

40

18. Verfahren nach Anspruch 15, wobei der Katalysator in Stufe (b) aus der Gruppe 3-Benzyl-5-(2-hydroxyethyl)-4-methylthiazoliumchlorid, 3,4-Dimethyl-5-(2-hydroxyethyl)-thiazoliumjodid, 3-Ethyl-5-(2-hydroxyethyl)-4-methyl-thiazoliumbromid und Thiaminhydrochlorid ausgewählt ist.

19. Verfahren nach Anspruch 18, wobei der Katalysator aus 3-Ethyl-5-(2-hydroxyethyl)-4-methylthiazoliumbromid besteht.

45

20. Verfahren nach Anspruch 15, wobei der Katalysator in Stufe (c) aus der Gruppe $R_{12}\text{CO}_2\text{H}$ mit R_{12} gleich CH_3 , CF_3 , ClCH_2- , $\text{C}_6\text{H}_5\text{CH}_2\text{CH}_2-$, $\text{C}_6\text{H}_5\text{CH}_2-$, HO_2CCH_2- , $\text{HO}_2\text{CCH}_2\text{CH}_2-$, C_6H_5- , $p\text{-Cl-C}_6\text{H}_5-$, $\text{ClCH}_2\text{CH}_2-$, $m\text{-H}_3\text{C-C}_6\text{H}_5-$, $p\text{-H}_3\text{C-C}_6\text{H}_5-$ oder $\text{tert.-C}_4\text{H}_9-$ und Triethylaminhydrochlorid ausgewählt ist.

50

21. Verfahren nach Anspruch 20, wobei der Katalysator aus $\text{tert.-C}_4\text{H}_9\text{CO}_2\text{H}$ besteht.

22. Verfahren nach Anspruch 15, wobei die Base in Stufe (b) aus der Gruppe N,N-Diisopropylethylamin, Pyridin, N,N-Dimethylanilin, Triethylamin, 1,8-Diazabicyclo[5.4.0]undec-7-en, 1,4-Diazabicyclo[2.2.2]octan, 4-Dimethylaminopyridin und N,N,N',N'-Tetramethylethylendiamin ausgewählt ist.


55

23. Verfahren nach Anspruch 22, wobei die Base aus Triethylamin besteht.



24. Verfahren nach Anspruch 15 zur Herstellung von (2R-trans-5-(4-Fluorphenyl)-2-(1-methylethyl)-N,4-diphenyl-1-[2-(tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)-ethyl]-1H-pyrrol-3-carboxamid.

25. Verbindung der Formel II



II

worin R_1 , R_2 , R_3 , R_4 , R_7 und R_8 die in Anspruch 1 angegebene Bedeutung besitzen und R_9 für Alkyl mit 1 bis 8 Kohlenstoffatom(en), eine 3- bis 6-gliedrige Cycloalkylgruppe oder α,α -Dimethylbenzyl steht.

26. Verbindung nach Anspruch 25, worin R_7 und R_8 für Methyl stehen, R_9 tert.-Butyl bedeutet und die beiden optisch aktiven Zentren R sind.

27. Verbindung nach Anspruch 25, wobei bedeuten:

R_1 4-Fluorphenyl;
 R_2 und R_3 Wasserstoff;
 R_4 Ethyl;
 R_7 und R_8 Methyl und
 R_9 Isopropyl.

28. Verbindung nach Anspruch 25, wobei bedeuten:

R_1 4-Fluorphenyl;
 R_2 Phenyl;
 R_3 C_6H_5NHCO ;
 R_4 Isopropyl;
 R_7 und R_8 Methyl und
 R_9 tert.-Butyl.

29. Verbindung nach Anspruch 28, wobei die beiden optisch aktiven Zentren R sind.




30. Verbindung der Formel XVI

5

10

15




XVI

20

worin R_{10} und R_{11} für Alkyl mit 1 bis 8 Kohlenstoffatom(en) stehen oder R_{10} und R_{11} zusammen

25




oder $-\text{CH}_2-\text{CH}_2-\text{CH}_2-$ bilden.

30

31. Verbindung nach Anspruch 30, worin R_{10} und R_{11} für Methyl oder Ethyl stehen oder R_{10} und R_{11} zusammen

35




bilden.

40

32. Verbindung der Formel IV

45



50

IV


worin bedeuten:

55

R_7 und R_8 unabhängig voneinander Wasserstoff, Alkyl mit 1 bis 3 Kohlenstoffatom(en) oder Phenyl oder R_7 und R_8 zusammen $-(\text{CH}_2)_n-$ mit n gleich 4 oder 5 und R_9 Alkyl mit 1 bis 8 Kohlenstoffatom(en), eine 3- bis 6-gliedrige Cycloalkylgruppe oder α,α -Dimethylbenzyl.



33. Verbindung nach Anspruch 32, worin R₉ für




steht.

34. Verbindung nach Anspruch 32, worin bedeuten: R₇ und R₈ Methyl und R₉ Isopropyl.

35. Verbindung nach Anspruch 32, worin bedeuten: R₇ und R₈ Methyl und R₉ tert.-Butyl.

36. Verbindung nach Anspruch 32, ausgewählt aus der Gruppe
 (4R-cis)-1,1-Dimethylethyl-6-(2-aminoethyl)-2,2-dimethyl-1,3-dioxan-4-acetat;
 [2R-(2 α ,4 α ,6 α)]-1,1-Dimethylethyl-6-(2-aminoethyl)-2-phenyl-1,3-dioxan-4-acetat;
 [2S-(2 α ,4 β ,6 β)]-1,1-Dimethylethyl-6-(2-aminoethyl)-2-phenyl-1,3-dioxan-4-acetat;
 (7R-cis)-1,1-Dimethylethyl-9-(2-aminoethyl)-6,10-dioxaspiro[4.5]-decan-7-acetat;
 (2R-cis)-1,1-Dimethylethyl-4-(2-aminoethyl)-1,5-dioxaspiro[5.5]-undecan-2-acetat;
 (4R-cis)-1,1-Dimethylethyl-4-(2-aminoethyl)-1,3-dioxan-4-acetat;
 [2R-(2 α ,4 α ,6 α)]-1,1-Dimethylethyl-6-(2-aminoethyl)-2-methyl-1,3-dioxan-4-acetat und
 [2S-(2 α ,4 β ,6 β)]-1,1-Dimethylethyl-6-(2-aminoethyl)-2-methyl-1,3-dioxan-4-acetat.


37. Verbindung der Formel XXI




XXI

worin R₇ und R₈ unabhängig voneinander für Wasserstoff, Alkyl mit 1 bis 3 Kohlenstoffatom(en) oder Phenyl stehen oder R₇ und R₈ zusammen -(CH₂)_n- mit n gleich 4 oder 5 bilden.

38. Verbindung nach Anspruch 37, ausgewählt aus der Gruppe
 (4R-cis)-6-(2-Aminoethyl)-2,2-dimethyl-1,3-dioxan-4-essigsäure;
 [2R-(2 α ,4 α ,6 α)]-6-(2-Aminoethyl)-2-phenyl-1,3-dioxan-4-essigsäure;
 [2S-(2 α ,4 β ,6 β)]-6-(2-Aminoethyl)-2-phenyl-1,3-dioxan-4-essigsäure;
 (7R-cis)-9-(2-Aminoethyl)-6,10-dioxaspiro[4.5]decan-7-essigsäure;
 (2R-cis)-4-(2-Aminoethyl)-1,5-dioxaspiro[5.5]undecan-2-essigsäure;
 (4R-cis)-4-(2-Aminoethyl)-1,3-dioxan-4-essigsäure;
 [2R-(2 α ,4 α ,6 α)]-6-(2-Aminoethyl)-2-methyl-1,3-dioxan-4-essigsäure und
 [2S-(2 α ,4 β ,6 β)]-6-(2-Aminoethyl)-2-methyl-1,3-dioxan-4-essigsäure.



39. Verbindung der Formel V



V

worin R_7 und R_8 unabhängig voneinander für Wasserstoff, Alkyl mit 1 - 3 Kohlenstoffatom(en) oder Phenyl stehen oder R_7 und R_8 zusammen $-(\text{CH}_2)_n-$ mit n gleich 4 oder 5 bilden und R_9 Alkyl mit 1 bis 8 Kohlenstoffatom(en), eine 3- bis 6-gliedrige Cycloalkylgruppe oder α, α -Dimethylbenzyl bedeutet.


40. Verbindung nach Anspruch 39, wobei bedeuten:

R_7 und R_8 Methyl und

R_9 Isopropyl.

41. Verbindung nach Anspruch 39, wobei R_7 und R_8 für Methyl stehen; R_9 tert.-Butyl bedeutet und die beiden optisch aktiven Zentren R sind.

42. Verbindung der Formel VI




VI

worin R_7 und R_8 unabhängig voneinander für Wasserstoff, Alkyl mit 1 bis 3 Kohlenstoffatom(en) oder Phenyl stehen oder R_7 und R_8 zusammen $-(\text{CH}_2)_n-$ mit n gleich 4 oder 5 bilden.

43. Verbindung nach Anspruch 42, wobei R_7 und R_8 für Methyl stehen und die beiden optisch aktiven Zentren R sind.



44. Verbindung der Formel VII




VII

worin R_7 und R_8 unabhängig voneinander für Wasserstoff, Alkyl mit 1 bis 3 Kohlenstoffatom(en) oder Phenyl stehen oder R_7 und R_8 zusammen $-(\text{CH}_2)_n-$ mit n gleich 4 oder 5 bilden.

45. Verbindung nach Anspruch 44, wobei R_7 und R_8 für Methyl stehen und die beiden optisch aktiven Zentren R sind.

46. Verbindung der Formel VIII




VIII

worin R_7 und R_8 unabhängig voneinander für Wasserstoff, Alkyl mit 1 bis 3 Kohlenstoffatom(en) oder Phenyl stehen und R_7 und R_8 zusammen $-(\text{CH}_2)_n-$ mit n gleich 4 oder 5 bilden.

47. Verbindung nach Anspruch 46, wobei R_7 und R_8 für Methyl stehen und die beiden optisch aktiven Zentren R sind.


48. Verbindung der Formel XVII



XVII




49. Mischung von Verbindungen der Formel XVIII



XVIII

Revendications

1. Un procédé de préparation d'un dérivé de formule I:



I

et d'un dihydroxyacide et des sels pharmaceutiquement acceptables de ce dernier, correspondant au cycle lactone ouvert du dérivé de formule I, dans laquelle:

R_1 représente un radical 1-naphtyle, 2-naphtyle, cyclohexyle, cyclohexylméthyle, norbornényle, phényle, un radical phényle substitué par:

un atome de fluor, de chlore, de brome, un groupe hydroxyle, trifluorométhyle, un radical alkyle comportant de 1 à 4 atomes de carbone, un radical alcoxy comportant de 1 à 4 atomes de carbone, ou alcanoyloxy comportant de 2 à 8 atomes de carbone, benzyle, 2-, 3-, ou 4-pyridinyle, ou 2-, 3-, ou 4-pyridinyl-N-oxyde;

R_2 ou R_3 représente indépendamment un atome d'hydrogène, un radical alkyle comportant de 1 à 6 atomes de carbone, un radical cyclopropyle, cyclobutyle, cyclopentyle, cyclohexyle, phényle, un radical phényle substitué par un atome de:

fluor, chlore, brome, un groupe hydroxyle, trifluorométhyle, un radical alkyle comportant de 1 à 4 atomes de carbone, ou alcoxy comportant de 1 à 4 atomes de carbone,

cyano, trifluorométhyle, ou $-CONR_5R_6$, dans lequel:

R_5 et R_6 sont indépendamment: un atome d'hydrogène, un radical alkyle comportant de 1 à 6 atomes de carbone, un radical phényle, un radical phényle substitué par un atome de fluor, de chlore, de brome, un groupe cyano ou un radical trifluorométhyle;

R_4 représente un radical alkyle comportant de 1 à 6 atomes de carbone, cyclopropyle, cyclobutyle, cyclopentyle, cyclohexyle ou trifluorométhyle;

qui comprend:


(a) la réaction de 1,6-heptadiène-4-ol avec:

(1) un alkyllithium,

(2) puis avec de l'iode et du dioxyde de carbone, et



(3) le traitement de l'iodocarbonate intermédiaire résultant in situ avec une base dans un alcool aqueux à environ 0 °C à environ 40 °C, de façon à obtenir un composé de formule IX:




IX

(b) le traitement du dérivé de formule IX avec:

(1) un cyanure alcalin à environ 0 °C à environ 40 °C, et

(2) la réaction du diol intermédiaire résultant in situ avec un réactif formant acétal en présence d'un acide pour conduire au dérivé de formule VIII:



VIII

dans laquelle:


R_7 et R_8 représentent indépendamment un atome d'hydrogène, un radical alkyle comportant de 1 à 3 atomes de carbone, un radical phényle ou R_7 et R_8 forment ensemble $-(\text{CH}_2)_n-$, dans lequel:

n est égal à 4 ou 5;

(c) le traitement du dérivé de formule VIII avec:

(1) l'ozone dans un solvant inerte, et

(2) la réaction de l'intermédiaire résultant in situ avec de l'oxygène et de la triphénylphosphine à environ -20 °C à -78 °C pour obtenir un dérivé de formule VII:




VII

dans laquelle:

R_7 et R_8 sont tels que définis ci-dessus;

(d) le traitement du dérivé de formule VII avec un réactif oxydant à environ 0 °C pour conduire au dérivé de formule VI:



VI

dans laquelle:

R_7 et R_8 sont tels que définis ci-dessus;


(e) le traitement du dérivé de formule VI avec un dérivé de formule:

Hal-R_{9a}

dans laquelle:

Hal est un atome d'halogène, et

R_{9a} est un radical alkyle comportant de 1 à 8 atomes de carbone, ou un groupe cycloalkyle comportant de 3 à 6 membres, en présence d'une base pour conduire au dérivé de formule V_a :

 V_a


dans laquelle:

R_7 , R_8 et R_{9a} sont tels que définis ci-dessus; ou le traitement du dérivé de formule VI avec un dérivé de formule:

HO-R_{9b}

dans laquelle:

R_{9b} représente un radical butyle tertiaire, amyle tertiaire ou α, α -diméthylbenzyle, en présence d'un agent d'activation, d'une quantité catalytique d'une base et d'un solvant inerte pour conduire au dérivé de formule V_b :


 V_b



dans laquelle:

R_7 , R_8 et R_{9a} sont tels que définis ci-dessus;

(f) le traitement du dérivé de formule V_a avec de l'hydrogène en présence d'un catalyseur et d'un acide à environ 0°C à environ 70°C , pour conduire à un dérivé de formule IV_a :




IV_a

dans laquelle:

R_7 , R_8 et R_{9a} sont tels que définis ci-dessus;

ou le traitement du dérivé de formule V_b avec de l'hydrogène en présence d'un catalyseur et d'un acide ou d'un catalyseur et d'une base à environ 0°C à environ 70°C pour conduire à un dérivé de formule IV_b :




IV_b

dans laquelle:

R_7 , R_8 et R_{9b} sont tels que définis ci-dessus;

(g) le traitement du dérivé de formule IV_a avec un dérivé de formule III:




III

dans laquelle:


R_1 , R_2 , R_3 et R_4 sont tels que définis ci-dessus;

dans un solvant inerte pour conduire au dérivé de formule II_a :

II_a

dans laquelle:

R₁, R₂, R₃, R₄, R₇, R₈ et R_{9a} sont tels que définis ci-dessus,
ou le traitement du dérivé de formule IV_b avec un dérivé de formule III dans un solvant inerte pour
conduire au dérivé de formule II_b:

II_b

dans laquelle:

R₁, R₂, R₃, R₄, R₇, R₈ et R_{9b} sont tels que définis ci-dessus;

(h) et finalement le traitement d'un dérivé de formule II_a avec:

- (1) un acide en présence d'un solvant inerte,
- (2) suivi de l'hydrolyse à l'aide d'une base,
- (3) suivi d'une étape de neutralisation avec un acide, et
- (4) dissolution et/ou chauffage dans un solvant inerte avec élimination concomitant de l'eau pour
obtenir un dérivé de formule I,

ou traitement du dérivé de formule II_b avec:

- (1) un acide en présence d'un solvant inerte,
- (2) suivi de l'addition d'une base,
- (3) suivi d'une étape de neutralisation avec un acide, et
- (4) dissolution et/ou chauffage dans un solvant inerte avec élimination concomitant de l'eau pour
obtenir un dérivé de formule I;

(i) et, si cela est nécessaire, la conversion du dérivé de formule I résultant en un dihydroxyacide
correspondant au cycle lactone ouvert de formule structurale I par une hydrolyse classique et de
plus, si cela est nécessaire, la conversion du dihydroxyacide en un sel pharmaceutiquement
acceptable correspondant par des moyens classiques et, si cela est nécessaire, la conversion du sel
pharmaceutiquement acceptable correspondant en un dihydroxyacide par des techniques classiques
et, si cela est nécessaire, la conversion du dihydroxyacide en un dérivé de formule I par chauffage
dans un solvant inerte.

2. Un procédé selon la revendication 1, caractérisé en ce que l'alkyllithium dans l'étape (a) est le n-
butyllithium.


3. Un procédé selon la revendication 1, caractérisé en ce que le cyanure alcalin dans l'étape (b) est du
cyanure de potassium.



4. Un procédé selon la revendication 1, caractérisé en ce que le réactif oxydant dans l'étape (d) est le trioxyde de chrome/acide sulfurique/eau.
5. Un procédé selon la revendication 1, caractérisé en ce que le catalyseur dans l'étape (f) pour la préparation du dérivé de formule IV_a est du dioxyde de platine.
6. Un procédé selon la revendication 1, caractérisé en ce que le catalyseur et la base dans l'étape (f) pour la préparation du dérivé de formule IV_b sont respectivement le nickel de Raney et l'ammoniac anhydre.
7. Un procédé selon la revendication 1, caractérisé en ce que l'acide dans l'étape (h) est de l'acide chlorhydrique.
8. Un procédé selon la revendication 1, caractérisé en ce que la base dans l'étape (h) est de l'hydroxyde de sodium.
9. Un procédé selon la revendication 1, caractérisé en ce que R₁ représente un radical 1-naphtyle, norbornényle, phényle ou phényle substitué par un atome de fluor, chlore, brome, un groupe hydroxyle, trifluorométhyle, un radical alkyle comportant de 1 à 4 atomes de carbone, alcoxy comportant de 1 à 4 atomes de carbone, ou alcanoyloxy comportant de 2 à 8 atomes de carbone.
10. Un procédé selon la revendication 9, caractérisé en ce que R₄ représente un radical alkyle comportant de 1 à 6 atomes de carbone, cyclopropyle ou trifluorométhyle.
11. Un procédé selon la revendication 1 et pour la préparation d'un dérivé sélectionné dans le groupe consistant en:
 - trans-6-[2-[2-(4-fluorophényl)-5-(trifluorométhyl)-1H-pyrrol-1-yl]éthyl]tétrahydro-4-hydroxy-2H-pyran-2-one;
 - trans-6-[2-[2-(4-fluorophényl)-5-méthyl-1H-pyrrol-1-yl]éthyl]tétrahydro-4-hydroxy-2H-pyran-2-one;
 - trans-6-[2-[2-(4-fluorophényl)-5-(1-méthyléthyl)-1H-pyrrol-1-yl]éthyl]tétrahydro-4-hydroxy-2H-pyran-2-one;
 - trans-6-[2-[2-cyclopropyl-5-(4-fluorophényl)-1H-pyrrol-1-yl]éthyl]tétrahydro-4-hydroxy-2H-pyran-2-one;
 - trans-6-[2-[2-(1,1-diméthyléthyl)-5-(4-fluorophényl)-1H-pyrrol-1-yl]éthyl]tétrahydro-4-hydroxy-2H-pyran-2-one;
 - trans-tétrahydro-4-hydroxy-6-[2-[2-(2-méthoxyphényl)-5-méthyl-1H-pyrrol-1-yl]éthyl]-2H-pyran-2-one;
 - trans-tétrahydro-4-hydroxy-6-[2-[2-(2-méthoxyphényl)-5-(1-méthyléthyl)-1H-pyrrol-1-yl]éthyl]-2H-pyran-2-one;
 - trans-tétrahydro-4-hydroxy-6-[2-[2-méthyl-5-(1-naphtalényle)-1H-pyrrol-1-yl]éthyl]-2H-pyran-2-one;
 - trans-6-[2-(2-bicyclo[2,2,1]hept-5-èn-2-yl-5-méthyl-1H-pyrrol-1-yl)éthyl]tétrahydro-4-hydroxy-2H-pyran-2-one;
 - trans-(±)-5-(4-fluorophényl)-2-(1-méthyléthyl)-N,4-diphényl-1-[2-(tétrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)éthyl]-1H-pyrrole-3-carboxamide;
 - (2R-trans)-5-(4-fluorophényl)-2-(1-méthyléthyl)-N,4-diphényl-1-[2-(tétrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)éthyl]-1H-pyrrole-3-carboxamide;
 - (trans)-2-(4-fluorophényl)-N,4-diphényl-1-[2-(tétrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)éthyl]-5-trifluorométhyl-1H-pyrrole-3-carboxamide; et
 - (trans)-5-(4-fluorophényl)-N,4-diphényl-1-[2-(tétrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)éthyl]-2-trifluorométhyl-1H-pyrrole-3-carboxamide.




12. Un procédé selon la revendication 1, pour la préparation d'un dérivé de formule I_a:




et l'hydroxy-acide ainsi que des sels pharmaceutiquement acceptables de ce dernier correspondant à un cycle lactone ouvert du composé de formule I_a, qui comprend:

(a) la réaction du dérivé de formule XVII:



avec un dérivé de formule:




dans laquelle:

R₇ et R₈ représentent indépendamment un atome d'hydrogène, un radical alkyle comportant de 1 à 3 atomes de carbone, un radical phényle, ou **R₇ et R₈** forment ensemble $-(CH_2)_n-$, dans laquelle:

n est égal à 4 ou 5, et

R₁₃ représente un atome d'hydrogène ou



dans un solvant inerte,


et le traitement de l'intermédiaire résultant avec un acide pour conduire au dérivé de formule I_a;

(b) et si cela est nécessaire la conversion du dérivé de formule I_a résultant en un hydroxyacide correspondant au cycle lactone ouvert de formule structurelle I_a par hydrolyse classique et de plus, si cela est nécessaire, la conversion de l'hydroxyacide en un sel pharmaceutiquement acceptable correspondant par les moyens classiques et, si cela est nécessaire, la conversion de l'hydroxyacide en un dérivé de formule I_a par chauffage dans un solvant inerte.

13. Un procédé selon la revendication 12, caractérisé en ce que l'acide dans l'étape (a) correspond à l'acide chlorhydrique.

14. Un procédé selon la revendication 12 et pour la préparation de (2R-trans)-5-(4-fluorophényl)-2-(1-méthyléthyl)-N,4-diphényl-1-[2-(tétrahydro-4-hydroxy-6-oxo-2H-pyran-2yl)éthyl]-1H-pyrrole-3-carboxamide.


15. Un procédé de préparation d'un dérivé de formule I_a:



I_a


et de l'hydroxy-acide et de sels pharmaceutiquement acceptables de ce dernier, correspondant au cycle lactone ouvert du dérivé de formule I_a, qui comprend:

(a) la réaction du 4-méthyl-3-oxo-N-phénylpentanamide avec du benzaldéhyde en présence d'un catalyseur et d'un solvant inerte pour conduire au dérivé de formule XVIII:




XVIII

(b) la réaction du dérivé de formule XVIII avec du 4-fluorobenzaldéhyde en présence d'un catalyseur, d'une base, et d'un solvant inerte pour conduire au dérivé de formule XVII:



XVII


(c) la réaction du dérivé de formule XVII avec un dérivé de formule:




dans laquelle:

R_{10} et R_{11}

représentent des radicaux alkyles comportant de 1 à 8 atomes de carbone, ou R_{10} et R_{11} forment ensemble:



$-\text{CH}_2 - \text{CH}_2 -$ ou $-\text{CH}_2 - \text{CH}_2 - \text{CH}_2 -$ en présence d'un catalyseur et d'un solvant inerte pour conduire au dérivé de formule XVI:




XVI

dans laquelle:

R_{10} et R_{11}

sont des radicaux alkyles comportant de 1 à 8 atomes de carbone, ou R_{10} et R_{11} forment ensemble:



$-\text{CH}_2-\text{CH}_2-$ ou $-\text{CH}_2-\text{CH}_2-\text{CH}_2-$

(d) et finalement la conversion d'un dérivé de formule XVI selon un procédé classique pour conduire au dérivé de formule Ia;

(e) et, si cela est nécessaire, la conversion du dérivé de formule Ia résultant en un hydroxyacide correspondant au cycle lactone ouvert de formule structurelle Ia par une hydrolyse classique, et de plus, si cela est nécessaire, la conversion de l'hydroxyacide en un sel pharmaceutiquement acceptable correspondant par des moyens classiques et, si cela est nécessaire, la conversion de l'hydroxyacide en un dérivé de formule Ia par chauffage dans un solvant inerte.

16. Un procédé selon la revendication 15, caractérisé en ce que le catalyseur dans l'étape (a) est sélectionné dans le groupe consistant en pipéridine et acide acétique glacial, éthylènediamine et acide acétique glacial, et β -alanine et acide acétique glacial.

17. Un procédé selon la revendication 16, caractérisé en ce que le catalyseur correspond à β -alanine et à l'acide acétique glacial.

18. Un procédé selon la revendication 15, caractérisé en ce que le catalyseur dans l'étape (b) est sélectionné dans le groupe consistant en: chlorure de 3-benzyl-5-(2-hydroxyéthyl)-4-méthylthiazolium; iodure de 3,4-diméthyl-5-(2-hydroxyéthyl)thiazolium; bromure de 3-éthyl-5-(2-hydroxyéthyl)-4-méthylthiazolium et chlorhydrate de thiamine.

19. Un procédé selon la revendication 18, caractérisé en ce que le catalyseur est le bromure de 3-éthyl-5-(2-hydroxyéthyl)-4-méthylthiazolium.

20. Un procédé selon la revendication 15, caractérisé en ce que le catalyseur dans l'étape (c) est sélectionné dans le groupe consistant en $R_{12}\text{CO}_2\text{H}$, dans lequel R_{12} représente CH_3 , CF_3 , ClCH_2- , $\text{C}_6\text{H}_5\text{CH}_2\text{CH}_2-$, $\text{C}_6\text{H}_5\text{CH}_2-$, HO_2CCH_2- , $\text{HO}_2\text{CCH}_2\text{CH}_2-$, C_6H_5- , para- $\text{Cl-C}_6\text{H}_5-$, $\text{ClCH}_2\text{CH}_2-$, meta- $\text{H}_3\text{C-C}_6\text{H}_5-$, para- $\text{H}_3\text{C-C}_6\text{H}_5-$, ou t- C_4H_9- et chlorhydrate de triéthylamine.

21. Un procédé selon la revendication 20, caractérisé en ce que le catalyseur est t- $\text{C}_4\text{H}_9\text{CO}_2\text{H}$.

22. Un procédé selon la revendication 15, caractérisé en ce que la base dans l'étape (b) est sélectionnée dans le groupe consistant en N,N-diisopropyléthylamine, pyridine, N,N-diméthylaniline, triéthylamine,




1,8-diaza-bicyclo[5,4,0]undec-7-ène, 1,4-diazabicyclo[2,2,2]octane, 4-diméthylaminopyridine et N,N,N',N'-tétraméthyléthylènediamine.

23. Un procédé selon la revendication 22, caractérisé en ce que la base correspond à la triéthylamine.

24. Un procédé selon la revendication 15, pour la préparation de (2R-trans)-5-(4-fluorophényl)-2-(1-méthyléthyl)-N,4-diphényl-1-[2-(tétrahydro-4-hydroxy-6-oxo-2H-pyran-2yl)-éthyl]-1H-pyrrole-3-carboxamide.

25. Un dérivé de formule II:



II

dans laquelle:

R_1, R_2, R_3, R_4, R_7 et R_8 sont tels que définis dans la revendication 1, et R_9 représente un radical alkyle comportant de 1 à 8 atomes de carbone, un groupe cycloalkyle comportant de 3 à 6 membres, ou un groupe α, α -diméthylbenzyle.


26. Un dérivé selon la revendication 25, caractérisé en ce que R_7 et R_8 représentent un radical méthyle, R_9 représente un radical t-butyle et les deux centres optiquement actifs correspondent à R.

27. Un dérivé selon la revendication 25, caractérisé en ce que R_1 représente un radical 4-fluorophényle, R_2 et R_3 représentent un atome d'hydrogène, R_4 représente un radical éthyle, R_7 et R_8 représentent un radical méthyle et R_9 représente un radical isopropyle.

28. Un dérivé selon la revendication 25, caractérisé en ce que R_1 représente un radical 4-fluorophényle, R_2 représente un radical phényle, R_3 représente C_6H_5NHCO- , R_4 représente un radical isopropyle, R_7 et R_8 représentent des radicaux méthyles et R_9 représente un radical t-butyle.

29. Un dérivé selon la revendication 28, caractérisé en ce que les deux centres optiquement actifs sont R.

30. Un dérivé représenté par la formule XVI:



XVI




dans laquelle:

R_{10} et R_{11}

sont des radicaux alkyles comportant de 1 à 8 atomes de carbone, ou R_{10} et R_{11} forment ensemble:

5




10

ou $-\text{CH}_2-\text{CH}_2-\text{CH}_2-$

31. Un dérivé selon la revendication 30, caractérisé en ce que R_{10} et R_{11} représentent un radical méthyle ou éthyle, ou R_{10} et R_{11} forment ensemble:


15



20

32. Un dérivé de formule IV:

25



30

IV

dans laquelle:

35

R_7 et R_8

représentent indépendamment un atome d'hydrogène, un radical alkyle comportant de 1 à 3 atomes de carbone, un radical phényle ou R_7 et R_8 forment ensemble $-(\text{CH}_2)_n-$ dans laquelle:

n

est égal à 4 ou 5, et


R_9

représente un radical alkyle comportant de 1 à 8 atomes de carbone, un groupe cycloalkyle comportant de 3 à 6 membres ou un groupe α,α -diméthylbenzyle.

40

33. Un dérivé selon la revendication 32, caractérisé en ce que R_9 représente:

45



50

34. Un dérivé selon la revendication 32, caractérisé en ce que R_7 et R_8 sont des radicaux méthyles et R_9 est un radical isopropyle.

35. Un dérivé selon la revendication 32, caractérisé en ce que R_7 et R_8 sont des radicaux méthyles et R_9 est un radical t-butyle.


55

36. Un dérivé selon la revendication 32, sélectionné dans le groupe consistant en (4R-cis)-1,1-diméthyléthyl-6-(2-aminoéthyl)-2,2-diméthyl-1,3-dioxane-4-acétate; [2R-(2 α , 4 α , 6 α)]-1,1-diméthyléthyl-6-(2-aminoéthyl)-2,2-diméthyl-1,3-dioxane-4-acétate.



noéthyl)-2-phényl-1,3-dioxane-4-acétate; [2S-(2 α , 4 β , 6 β)]-1,1-diméthyléthyl-6-(2-aminoéthyl)-2-phényl-1,3-dioxane-4-acétate; (7R-cis)-1,1-diméthyléthyl-9-(2-aminoéthyl)-6,10-dioxaspiro[4,5]-décane-7-acétate; (2R-cis)-1,1-diméthyléthyl-4-(2-aminoéthyl)-1,5-dioxaspiro[5,5]-undécane-2-acétate; (4R-cis)-1,1-diméthyléthyl-4-(2-aminoéthyl)-1,3-dioxane-4-acétate; [2R-(2 α , 4 α , 6 α)]-1,1-diméthyléthyl-6-(2-aminoéthyl)-2-méthyl-1,3-dioxane-4-acétate; et [2S-(2 α , 4 β , 6 β)]-1,1-diméthyléthyl-6-(2-aminoéthyl)-2-méthyl-1,3-dioxane-4-acétate.

37. Un dérivé de formule XXI:




XXI

dans laquelle:

R₇ et R₈ représentent indépendamment un atome d'hydrogène, un radical alkyle comportant de 1 à 3 atomes de carbone, un radical phényle ou R₇ et R₈ forment ensemble -(CH₂)_n- dans lequel:
n est égal à 4 ou 5.

38. Un dérivé selon la revendication 37, sélectionné dans le groupe consistant en: acide (4R-cis)-6-(2-aminoéthyl)-2,2-diméthyl-1,3-dioxane-4-acétique; acide [2R-(2 α , 4 α , 6 α)]-6-(2-aminoéthyl)-2-phényl-1,3-dioxane-4-acétique; acide [2S-(2 α , 4 β , 6 β)]-6-(2-aminoéthyl)-2-phényl-1,3-dioxane-4-acétique; acide (7R-cis)-9-(2-aminoéthyl)-6,10-dioxaspiro[4,5]décane-7-acétique; acide (2R-cis)-4-(2-aminoéthyl)-1,5-dioxaspiro[5,5]undécane-2-acétique; acide (4R-cis)-4-(2-aminoéthyl)-1,3-dioxane-4-acétique; acide [2R-(2 α , 4 α , 6 α)]-6-(2-aminoéthyl)-2-méthyl-1,3-dioxane-4-acétique et acide [2S-(2 α , 4 β , 6 β)]-6-(2-aminoéthyl)-2-méthyl-1,3-dioxane-4-acétique.

39. Un dérivé représenté par la formule V:



V

dans laquelle:


R₇ et R₈ représentent indépendamment un atome d'hydrogène, un radical alkyle comportant de 1 à 3 atomes de carbone, un radical phényle ou R₇ et R₈ forment ensemble -(CH₂)_n- dans lequel:
n est égal à 4 ou 5, et
R₉ représente un radical alkyle comportant de 1 à 8 atomes de carbone, un groupe cycloalkyle comportant de 3 à 6 membres ou un groupe α,α -diméthylbenzyle.

40. Un dérivé selon la revendication 39, caractérisé en ce que R₇ et R₈ sont des radicaux méthyle, et R₉ est un radical isopropyle.



41. Un dérivé selon la revendication 39, caractérisé en ce que R_7 et R_8 sont des radicaux méthyle, R_9 est un radical t-butyle et les deux centres optiquement actifs représentent R.

42. Un dérivé représenté par la formule VI:




VI

dans laquelle:

R_7 et R_8 représentent indépendamment un atome d'hydrogène, un radical alkyle comportant de 1 à 3 atomes de carbone, un radical phényle ou R_7 et R_8 forment ensemble $-(CH_2)_n-$ dans lequel: n est égal à 4 ou 5.

43. Un dérivé selon la revendication 42, caractérisé en ce que R_7 et R_8 sont des radicaux méthyles et les deux centres optiquement actifs représentent R.

44. Un dérivé représenté par la formule VII:



VII


dans laquelle:

R_7 et R_8 représentent indépendamment un atome d'hydrogène, un radical alkyle comportant de 1 à 3 atomes de carbone, un radical phényle ou R_7 et R_8 forment ensemble $-(CH_2)_n-$ dans lequel: n est égal à 4 ou 5.

45. Un dérivé selon la revendication 44, caractérisé en ce que R_7 et R_8 représentent des radicaux méthyles et les deux centres optiquement actifs représentent R.



46. Un dérivé représenté par la formule VIII:




VIII

dans laquelle:

R_7 et R_8 représentent indépendamment un atome d'hydrogène, un radical alkyle comportant de 1 à 3 atomes de carbone, un radical phényle ou R_7 et R_8 forment ensemble $-(CH_2)_n-$ dans lequel:
 n est égal à 4 ou 5.


47. Un dérivé selon la revendication 46, caractérisé en ce que R_7 et R_8 sont des radicaux méthyles et les deux centres optiquement actifs correspondent à R.

48. Un dérivé représenté par la formule XVII:



XVII

49. Un mélange des dérivés représentés par la formule XVIII:



XVIII